## Complexity of Charged Dynamical Spherical System in Modified Gravity

M. Sharif<sup>1</sup> \*and K. Hassan<sup>2</sup>  $^{\dagger}$ 

 <sup>1</sup> Department of Mathematics and Statistics, The University of Lahore, 1-KM Defence Road Lahore, Pakistan.
 <sup>2</sup> Department of Mathematics, University of the Punjab,

Quaid-e-Azam Campus, Lahore-54590, Pakistan.

#### Abstract

In this paper, we consider the effect of electromagnetic field to the definition of complexity in the context of f(G,T) gravity, where G and T express the Gauss-Bonnet term and energy-momentum tensor, respectively. The physical parameters such as anisotropic pressure, charge, energy density inhomogeneity, heat dissipation and correction terms are found responsible to induce complexity within the self-gravitating objects. The scalar functions are determined using Herrera's orthogonal splitting approach, which results in a complexity factor that includes all of the system's essential features. Furthermore, we investigate the dynamics of charged spherical distribution by choosing homologous mode as the simplest evolutionary pattern. Dissipative and non-dissipative cases associated with complexity free and homologous conditions are also discussed. Finally, we study the components responsible for producing complexity during the evolution process. We deduce that the inclusion of extra curvature terms and charge in f(G,T) gravity enhances the complexity of the selfgravitating structure.

Keywords: Complexity factor; Non-minimal coupled gravity; Self-gravitating systems; Anisotropic fluid.

**PACS:** 98.62.Gq; 04.40.Dg;04.40.-b

<sup>\*</sup>msharif.math@pu.edu.pk

<sup>&</sup>lt;sup>†</sup>komalhassan3@gmail.com

### 1 Introduction

The cosmos is made up of both microscopic and immense scale entities that range from tiny planets to galaxies (comprising billions of stars). These cosmic structures help in forming the basis for cosmological studies and persuade the universe's progression. Einstein believed that the cosmos is static but Hubble used a relationship between the distance and recession velocity of galaxies to prove that the universe is in an expanding phase. The rapid expansion of the cosmos is believed to be caused by the mysterious quantity termed as dark energy. Numerous astronomical phenomena like supernovae and microwave background radiations [1, 2] indicate the accelerated expanding universe. The  $\Lambda$ CDM model describes the best paradigm for explaining the present accelerated expanding universe but it faces two serious issues: cosmic coincidence and fine-tuning. Consequently, modified theories of gravity are introduced as alternative to discuss the cosmic rapid expansion.

Nojiri and Odintsov [3] added the function of Gauss-Bonnet (GB) invariant in the Einstein-Hilbert action and proposed modified GB gravity, also known as f(G) theory, where  $G = -4R^{\vartheta\tau}R_{\psi\chi} + R^{\vartheta\tau\alpha\delta}R_{\vartheta\tau\alpha\delta} + R^2$ . This theory discusses various astrophysical and cosmological events happening in the universe. Bamba et al. [4] chose a scale factor as a conjunction of two exponential functions and reconstructed the f(G) model to analyze the earlytime bounce and late-time cosmic expansion. Abbas et al. [5] investigated the possibility of constructing celestial objects and used the power-law model together with Krori-Barua metric to inspect their physical features in this theory. Sharif and Ikram [6] studied the dynamics of inflationary phase of homogeneous and isotropic cosmos utilizing viable f(G) model and found various metric coefficients that correspond to well-known inflationary models. Sharif and Ramzan [7] used the embedding class-1 approach to develop the anisotropic stellar structures and their salient physical aspects.

Sharif and Ikram [8] formulated f(G, T) theory by incorporating trace of the energy-momentum tensor in f(G) action. They remodeled the powerlaw and de Sitter ansatz to discuss energy conditions for the FRW cosmos. The same authors [9] discussed the stable behavior of the Einstein universe through anisotropic homogeneous perturbations. Hossienkhani et al. [10] analyzed energy constraints of anisotropic universe and found a direct relation between weak energy condition and anisotropy. Sharif and Naeem [11] studied the viable and stable behavior of certain stellar quantities by formulating a new solution. Shamir [12] determined the favorable bouncing solutions for the chosen equation of state with respect to two f(G, T) gravity models.

The electromagnetic field plays an important role (due to the suppression of gravitational force) in exploring the stability and evolution of massive bodies. To maintain the stable character of astrophysical objects, a substantial quantity of charge is essential to overcome the gravitational pull. Ivanov [13] investigated three distinct classes of regular and general solutions to the field equations for the charged celestial body. Esculpi and Aloma [14] used a linear equation of state that connects tangential and radial pressures to study the impact of anisotropic factor on charged compact objects. Sharif and Naz [15] inspected the collapse of a cylindrical charged configuration using dynamical field equations in f(G) gravity. We have figured out the essential characteristics of uncharged/charged and charged anisotropic celestial objects through minimal and extended geometric deformation scheme, respectively, in f(G, T) gravity [16, 17, 18].

The complexity of astrophysical entities predominantly relies on the internal attributes (heat, pressure, temperature, energy density, etc.) of selfgravitating objects. In order to calculate the complexity of cosmic objects, a mathematical formula that incorporates all important physical variables must be applied. López-Ruiz and his collaborators [19] proposed the idea of complexity in terms of information and entropy. The ideal gas and perfect crystal were the first two physical quantities on which this notion was used. The perfect crystal has no entropy due to the symmetry of molecules while the ideal gas exhibits maximal entropy due to the random distribution of its molecules. The probability distribution of perfect crystal does not reveal much new information, thus a tiny segment describes all of its important features. In contrast to perfect crystal, we attain a large amount of information during the study of small portion of ideal gas. One can notice that both the entities are at their extremes in behavior, so zero complexity is allotted to them. A new concept of complexity was developed based on how various probabilistic states deviate from the equiprobable distribution of the system [20, 21]. According to this notion, perfect crystal and ideal gas also have zero complexity.

Herrera [22] revised the concept of complexity in such a way that it incorporates all the essential state determinants (pressure anisotropy, inhomogeneous energy density, Tolman mass) for the static symmetric anisotropic matter distribution. This technique works on splitting the Riemann tensor which gives rise to the formulation of various scalar functions. Thus, we call a factor as complexity producing source that includes all the abovementioned quantities. Sharif and Butt [23] discussed the complexity factor for the charged sphere. This notion of complexity was then extended to non-static dissipative fluid by Herrera et al [24] and discussed the simplest evolutionary mode. The application of this concept has also been found in axially symmetric spacetime by the same authors [25]. Sharif and Butt [26] employed the Herrera's strategy to gauge the complexity of static cylindrically anisotropic source.

The analysis of complexity has also been investigated in modified theories. Sharif et al. [27] discussed complexity through  $\mathcal{Y}_{TF}$  (the complexity factor) for spherical system in f(G) gravity. Yousaf et al. [28, 29] studied the the complexity producing factor for the charged and uncharged spherical geometry in f(G,T) gravity. We have also figured out the impact of complexity on static uncharged/charged cylindrical source and non-static (spherical and cylindrical) structures in f(G,T) gravity through the orthogonal splitting of the Riemann tensor [30, 31, 32, 33]. In the background of different modified theories, there is a substantial body of work on the assessment of complexity in diverse geometries [34, 35, 36, 37, 38, 39, 40].

This paper explores the complexity producing factor for the dynamical charged spherical configuration as well as evolutionary patterns in f(G,T) theory. The paper is arranged in the following format. The fundamental properties of the matter source and the field equations characterizing the evolution of the system are given in section **2**. The Riemann tensor is orthogonally decomposed into its structure scalars in section **3**. In section **4**, two patterns of evolution, i.e., homologous and homogeneous are investigated. In section **5**, we derive kinematical and dynamical quantities to examine possible solutions for dissipative and non-dissipative systems. The conditions under which self-gravitating objects deviate from zero complexity condition are observed in section **6**. The summary of the obtained results is presented in section **7**.

### **2** f(G,T) Gravity and Matter Determinants

The generic f(G,T) function along with Ricci scalar in Einstein-Maxwell action takes the form

$$\mathbf{S}_{f(G,T)} = \frac{1}{2k^2} \int d^4x [f(G,T) + R] \sqrt{-g} + \int \sqrt{-g} (\mathbb{L}_m + \mathbb{L}_E) d^4x, \quad (1)$$

where the Lagrangian densities for the normal matter and electromagnetic field are represented by  $\mathbb{L}_m$  and  $\mathbb{L}_E$ , respectively, and  $k^2$  denotes the coupling constant. The stress-energy tensor and Lagrangian density are interlinked through the relation

$$T_{\vartheta\tau} = g_{\vartheta\tau} \mathbb{L}_m - \frac{2\partial \mathbb{L}_m}{\partial g^{\vartheta\tau}}.$$
 (2)

Here, the variation of action (1) with the metric tensor  $(g_{\vartheta\tau})$  formulates the modified field equations as

$$G_{\vartheta\tau} = 8\pi (T_{\vartheta\tau} + \mathbf{S}_{\vartheta\tau}) - (\Theta_{\vartheta\tau} + T_{\vartheta\tau}) f_T(G, T) + \frac{1}{2} g_{\vartheta\tau} f(G, T) + (4R_{\mu\tau} R^{\mu}_{\vartheta} + 4R^{\mu\nu} R_{\vartheta\mu\tau\nu} - 2RR_{\vartheta\tau} - 2R^{\mu\nu\gamma}_{\vartheta} R_{\tau\mu\nu\gamma}) f_G(G, T) + (4g_{\vartheta\tau} R^{\mu\nu} \nabla_{\mu} \nabla_{\nu} + 2R \nabla_{\vartheta} \nabla_{\tau} + 4R_{\vartheta\tau} \nabla^2 - 2g_{\vartheta\tau} R \nabla^2 - 4R^{\mu}_{\vartheta} \nabla_{\tau} \nabla_{\mu} - 4R^{\mu}_{\tau} \nabla_{\vartheta} \nabla_{\mu} - 4R_{\vartheta\mu\tau\nu} \nabla^{\mu} \nabla^{\nu}) f_G(G, T),$$
(3)

where d' Alembert operator is described by  $\nabla^2 = \nabla^l \nabla_l = \Box$  and  $\Theta_{\vartheta \tau} = -2T_{\vartheta \tau} + \mathfrak{p}g_{\vartheta \tau}$ . The Einstein tensor is indicated by  $G_{\vartheta \tau} = R_{\vartheta \tau} - \frac{1}{2}Rg_{\vartheta \tau}$ ,  $f_G(G,T)$  and  $f_T(G,T)$  are the partial derivatives of f(G,T) (arbitrary function) with respect to G and T, respectively.

In this theory, the matter components are coupled with the geometrical terms that lead to the non-conservation of the stress-energy tensor. Due to this, an extra force is induced that forces the particles in the gravitational field to deviate from the geodesic trajectory, therefore, the covariant differentiation of Eq.(3) reads

$$\nabla^{\vartheta} T_{\vartheta\tau} = \frac{f_T(G,T)}{k^2 - f_T(G,T)} \left[ -\frac{1}{2} g_{\vartheta\tau} \nabla^{\vartheta} T + (\Theta_{\vartheta\tau} + T_{\vartheta\tau}) \nabla^{\vartheta} (\ln f_T(G,T)) + \nabla^{\vartheta} \Theta_{\vartheta\tau} \right].$$
(4)

Equation (3) can also be written as

$$G_{\vartheta\tau} = 8\pi T_{\vartheta\tau}^{(\text{tot})} = 8\pi (T_{\vartheta\tau}^{(M)} + \mathbf{S}_{\vartheta\tau} + T_{\vartheta\tau}^{(\text{GT})}),$$
(5)

where  $T_{\vartheta\tau}^{(\mathsf{GT})}$  are extra curvature terms of f(G,T) given as

$$T_{\vartheta\tau}^{(\mathsf{GT})} = \frac{1}{8\pi} [\{(\mathfrak{u} + \mathfrak{p})v_{\vartheta}v_{\tau} + \Pi_{\vartheta\tau} + \zeta(v_{\vartheta}\chi_{\tau} + \chi_{\vartheta}v_{\tau})\}f_{T}(G, T) + (4R_{\mu\tau}R_{\vartheta}^{\mu} + 4R^{\mu\nu}R_{\vartheta\mu\tau\nu} - 2RR_{\vartheta\tau} - 2R_{\vartheta}^{\mu\nu\gamma}R_{\tau\mu\nu\gamma})f_{G}(G, T) + (4g_{\vartheta\tau}R^{\mu\nu}\nabla_{\nu}\nabla_{\mu} - 4R_{\vartheta\mu\tau\nu}\nabla^{\nu}\nabla^{\mu} - 4R_{\vartheta}^{\mu}\nabla_{\tau}\nabla_{\mu} - 2g_{\vartheta\tau}R\nabla^{2}]$$

$$- 4R^{\mu}_{\tau}\nabla_{\vartheta}\nabla_{\mu} + 2R\nabla_{\vartheta}\nabla_{\tau} + 4R_{\vartheta\tau}\nabla^{2})f_{G}(G,T) + \frac{g_{\vartheta\tau}f(G,T)}{2}.$$
 (6)

The normal matter distribution possessing anisotropy and dissipation due to heat flux is followed by the energy-momentum tensor

$$T^{(M)}_{\vartheta\tau} = \mathfrak{u} v_\vartheta v_\tau + \mathfrak{p} h_{\vartheta\tau} + \Pi_{\vartheta\tau} + \zeta (v_\vartheta \chi_\tau + \chi_\vartheta v_\tau), \tag{7}$$

where  $h_{\vartheta\tau}$  denotes the projection tensor. The radial four-vector, four velocity and heat flux are expressed as  $\chi^{\vartheta} = (0, \mathsf{B}^{-1}, 0, 0), v^{\vartheta} = (\mathsf{A}^{-1}, 0, 0, 0)$  and  $\zeta^{\vartheta} = (0, \zeta \mathsf{B}^{-1}, 0, 0)$ , respectively. These quantities satisfy the following relations

$$\chi^{\vartheta}\chi_{\vartheta} = 1, \quad \chi^{\vartheta}v_{\vartheta} = 0, \quad v^{\vartheta}\zeta_{\vartheta} = 0, \quad v^{\vartheta}v_{\vartheta} = -1.$$
(8)

The remaining entities are delineated as

$$\Pi_{\vartheta\tau} = \Pi\left(\chi_{\vartheta}\chi_{\tau} - \frac{h_{\vartheta\tau}}{3}\right), \quad \Pi = \mathfrak{p}_r - \mathfrak{p}_{\perp}, \tag{9}$$

$$\mathfrak{p} = \frac{\mathfrak{p}_r + 2\mathfrak{p}_\perp}{3}, \quad h_{\vartheta\tau} = g_{\vartheta\tau} + v_\vartheta v_\tau. \tag{10}$$

The stress-energy tensor in the electromagnetic field reads

$$\mathbf{S}_{\vartheta\tau} = \frac{1}{4\pi} \left( \mathcal{F}_{\vartheta}^{l} \mathcal{F}_{\tau l} - \frac{1}{4} g_{\vartheta\tau} \mathcal{F}_{lm} \mathcal{F}^{lm} \right), \tag{11}$$

where  $\mathcal{F}_{\vartheta\tau}$  signifies the Maxwell field tensor. In tensorial form, the Maxwell field equations are described as

$$\mathcal{F}_{[\vartheta\tau;l]} = 0, \quad \mathcal{F}^{\vartheta\tau}_{;\tau} = 4\pi \mathcal{J}^{\vartheta},$$

where  $\mathcal{J}^{\vartheta} = \varepsilon v^{\vartheta}$  is the four current and  $\varepsilon$  indicates the charge density.

The line element representing the non-static geometrical structure (surrounded by a hypersurface  $\Sigma$ ) involving dissipation is provided by

$$ds^{2} = -\mathsf{A}^{2}dt^{2} + \mathsf{B}^{2}dr^{2} + \mathsf{C}^{2}d\theta^{2} + \mathsf{C}^{2}\sin^{2}\theta d\phi^{2}, \qquad (12)$$

where B=B(t,r), A=A(t,r) and C = C(t,r). The modified field equations for the non-static realm are computed as

$$8\pi(\mathsf{A}^{2}\mathfrak{u} + \frac{\mathbf{s}^{2}\mathsf{A}}{8\pi\mathsf{C}^{4}} + T_{00}^{(\mathsf{GT})}) = \frac{-\mathsf{A}^{2}\left[\frac{2\mathsf{C}''}{\mathsf{C}} + \frac{\mathsf{C}'^{2}}{\mathsf{C}^{2}} - \frac{\mathsf{B}^{2}}{\mathsf{C}^{2}} - \frac{2\mathsf{C}'\mathsf{B}'}{\mathsf{C}\mathsf{B}}\right]}{\mathsf{B}^{2}} + \frac{\dot{\mathsf{C}}(\frac{2\dot{\mathsf{B}}}{\mathsf{B}} + \frac{\dot{\mathsf{C}}}{\mathsf{C}})}{\mathsf{C}}, (13)$$

$$8\pi(-\zeta \mathsf{A}\mathsf{B} + T_{01}^{(\mathsf{GT})}) = \frac{2\mathsf{A}'\dot{\mathsf{C}}}{\mathsf{A}\mathsf{C}} + \frac{2\mathsf{C}'\dot{\mathsf{B}}}{\mathsf{C}\mathsf{B}} - \frac{2\dot{\mathsf{C}}'}{\mathsf{C}}, \qquad (14)$$

$$8\pi (\mathsf{B}^{2}\mathfrak{p}_{r} - \frac{\mathsf{s}^{2}\mathsf{B}^{2}}{8\pi\mathsf{C}^{4}} + T_{11}^{(\mathsf{GT})}) = \frac{-\mathsf{B}^{2} \left[-\frac{\mathsf{C}}{\mathsf{C}} \left(\frac{2\mathsf{A}}{\mathsf{A}} - \frac{\mathsf{C}}{\mathsf{C}}\right) + \frac{2\mathsf{C}}{\mathsf{C}}\right]}{\mathsf{A}^{2}} - \frac{\mathsf{B}^{2}}{\mathsf{C}^{2}} + \left(\frac{2\mathsf{A}'\mathsf{C}'}{\mathsf{A}\mathsf{C}} + \frac{\mathsf{C}'^{2}}{\mathsf{C}^{2}}\right),$$
(15)

$$8\pi(\mathsf{C}^{2}\mathfrak{p}_{\perp} + \frac{\mathbf{s}^{2}}{8\pi\mathsf{C}^{2}} + T_{22}^{(\mathsf{GT})}) = \frac{-\mathsf{C}^{2}}{\mathsf{A}^{2}} \left[ \frac{\dot{\mathsf{B}}\dot{\mathsf{C}}}{\mathsf{B}\mathsf{C}} + \frac{\ddot{\mathsf{B}}}{\mathsf{B}} + \frac{\ddot{\mathsf{C}}}{\mathsf{C}} - \frac{\dot{\mathsf{A}}}{\mathsf{A}}(\frac{\dot{\mathsf{C}}}{\mathsf{C}} + \frac{\dot{\mathsf{B}}}{\mathsf{B}}) \right] \\ + \frac{\mathsf{C}^{2}}{\mathsf{B}^{2}} \left[ (\frac{\mathsf{A}'}{\mathsf{A}} - \frac{\mathsf{B}'}{\mathsf{B}}) \frac{\mathsf{C}'}{\mathsf{C}} - \frac{\mathsf{A}'\mathsf{B}'}{\mathsf{A}\mathsf{B}} + \frac{\mathsf{A}''}{\mathsf{A}} + \frac{\mathsf{C}''}{\mathsf{C}} \right], \quad (16)$$

where temporal and radial partial derivatives are denoted by . and  $\prime$ , respectively. The expressions for modified terms  $(T_{00}^{(GT)}, T_{01}^{(GT)}, T_{11}^{(GT)}, T_{22}^{(GT)})$  are presented in Eqs.(A1)-(A4) of Appendix **A**. Using Eq.(4), the non-zero components of the Bianchi identities for the present setup are derived as

$$T_{;\tau}^{\vartheta\tau}v_{\vartheta} = \frac{-1}{\mathsf{A}}\left\{\dot{\mathfrak{u}} + \frac{\dot{\mathsf{B}}}{\mathsf{B}}\left(\mathfrak{u} + \mathfrak{p}_{r}\right) + 2\frac{\dot{\mathsf{C}}}{\mathsf{C}}\left(\mathfrak{u} + \mathfrak{p}_{\perp}\right)\right\} - \frac{1}{\mathsf{B}}\left\{\zeta' + 2\zeta\left(\frac{\mathsf{A}'}{\mathsf{A}} + \frac{\mathsf{C}'}{\mathsf{C}}\right)\right\} = \mathsf{Z}_{1}$$

$$(17)$$

$$T_{;\tau}^{\vartheta\tau}\chi_{\vartheta} = \frac{1}{\mathsf{A}}\left\{\dot{\zeta} + 2\zeta\left(\frac{\dot{\mathsf{B}}}{\mathsf{B}} + \frac{\dot{\mathsf{C}}}{\mathsf{C}}\right)\right\} + \frac{1}{\mathsf{B}}\left\{\mathfrak{p}_{r}' + \left(\mathfrak{u} + \mathfrak{p}_{r}\right)\frac{\mathsf{A}'}{\mathsf{A}} - \frac{\mathsf{ss}'}{4\pi\mathsf{C}^{4}} + \left(2\mathfrak{p}_{r} - 2\mathfrak{p}_{\perp}\right)\right\}$$

$$\times \frac{\mathsf{C}'}{\mathsf{C}}\right\} = \mathsf{Z}_{2},$$

$$(18)$$

where the extra curvature terms  $Z_1$  and  $Z_2$  are prescribed in Eqs.(A5) and (A6) of Appendix **A**.

The non-zero elements of shear tensor, four acceleration and expansion scalar, respectively, for the present distribution are defined as

$$\sigma_{11} = \frac{2}{3}\mathsf{B}^2\sigma, \quad \sigma_{22} = \frac{\sigma_{33}}{\sin^2\theta} = -\frac{1}{3}\mathsf{C}^2\sigma,$$
 (19)

$$\sigma^{\vartheta\tau}\sigma_{\vartheta\tau} = \frac{2}{3}\sigma^2, \ \sigma = \left(\frac{\dot{\mathsf{B}}}{\mathsf{B}} - \frac{\dot{\mathsf{C}}}{\mathsf{C}}\right)\frac{1}{\mathsf{A}},\tag{20}$$

$$a = \sqrt{a^{\vartheta}a_{\vartheta}} = \frac{\mathsf{A}'}{\mathsf{B}\mathsf{A}}, \quad a_1 = \frac{\mathsf{A}'}{\mathsf{A}},$$
 (21)

$$\Theta = \left(2\frac{\dot{\mathsf{C}}}{\mathsf{C}} + \frac{\dot{\mathsf{B}}}{\mathsf{B}}\right)\frac{1}{\mathsf{A}}.$$
(22)

The influence of shear and expansion scalars on the fluid matter source is investigated through Eq.(14) in the following form

$$4\pi \left(\zeta \mathsf{B} - \frac{T_{01}^{(\mathsf{GT})}}{\mathsf{A}}\right) = -\sigma \frac{\mathsf{C}'}{\mathsf{C}} + \frac{1}{3} \left(\Theta - \sigma\right)' = \frac{\mathsf{C}'}{\mathsf{B}} \left[\frac{1}{3} D_{\mathsf{C}} \left(\Theta - \sigma\right) - \frac{\sigma}{\mathsf{C}}\right], \quad (23)$$

where proper radial differentiation is indicated by  $D_{\mathsf{C}} = \frac{1}{\mathsf{C}} \frac{\partial}{\partial r}$ . The mass of inner spherical object can be employed by Misner and Sharp [41] defined as

$$m = \frac{1}{2} \mathsf{C}^3 R_{232}^3 = \left[ 1 - \left(\frac{\mathsf{C}'}{\mathsf{B}}\right)^2 + \left(\frac{\dot{\mathsf{C}}}{\mathsf{A}}\right)^2 \right] \frac{\mathsf{C}}{2} + \frac{\mathsf{s}^2}{2\mathsf{C}}.$$
 (24)

Here, the dynamics of self-gravitating structure is examined by introducing proper time derivative, i.e.,  $D_T = \frac{1}{A} \frac{\partial}{\partial t}$ . During collapse, the outward pressure is suppressed by gravity which results in the decrement of radius of any celestial object. Due to this, the velocity (U) of the collapsing fluid in terms of areal radius (C) becomes negative as

$$U = D_T \mathsf{C} < 0. \tag{25}$$

The mass and velocity of the charged matter source are interlinked as follows

$$E \equiv \frac{\mathsf{C}'}{\mathsf{B}} = \left(1 - \frac{2m}{\mathsf{C}} + \frac{\mathbf{s}^2}{2\mathsf{C}} + U^2\right)^{\frac{1}{2}}.$$
 (26)

In view of time and radial proper derivatives, the mass function takes the form

$$D_T m = -4\pi \left[ \left( \mathfrak{p}_r - \frac{\mathbf{s}^2}{8\pi \mathsf{C}^4} + \frac{T_{11}^{(\mathsf{GT})}}{\mathsf{B}^2} \right) U + \left( \zeta - \frac{T_{01}^{(\mathsf{GT})}}{\mathsf{A}\mathsf{B}} \right) E \right] \mathsf{C}^2 + \frac{\mathbf{s}\dot{\mathbf{s}}}{\mathsf{A}\mathsf{C}} - \frac{\mathbf{s}^2\dot{\mathsf{C}}}{2\mathsf{A}\mathsf{C}^2},$$
(27)
$$(27)$$

$$D_{\mathsf{C}}m = 4\pi \left[ \left( \mathfrak{u} + \frac{\mathbf{s}^2}{8\pi\mathsf{C}^4} + \frac{T_{00}^{(\mathsf{G}\,\mathsf{I}\,\mathsf{I}\,\mathsf{I}}}{\mathsf{A}^2} \right) + \left( \zeta - \frac{T_{01}^{(\mathsf{G}\,\mathsf{I}\,\mathsf{I}\,\mathsf{I}}}{\mathsf{A}\mathsf{B}} \right) \frac{U}{E} \right] \mathsf{C}^2 + \frac{\mathbf{s}\mathbf{s}'}{\mathsf{C}\mathsf{C}'} - \frac{\mathbf{s}^2}{2\mathsf{C}^2}, \quad (28)$$
which yield

which yield

$$\frac{3m}{\mathsf{C}^3} = 4\pi \left( \mathfrak{u} + \frac{\mathbf{s}^2}{8\pi\mathsf{C}^4} + \frac{T_{00}^{(\mathsf{GT})}}{\mathsf{A}^2} \right) - \frac{4\pi}{\mathsf{C}^3} \int_0^r \mathsf{C}^3 \left[ D_\mathsf{C} \left( \mathfrak{u} + \frac{\mathbf{s}^2}{8\pi\mathsf{C}^4} + \frac{T_{00}^{(\mathsf{GT})}}{\mathsf{A}^2} \right) \right]$$

$$- 3\left(\zeta - \frac{T_{01}^{(\mathsf{GT})}}{\mathsf{AB}}\right)\frac{U}{\mathsf{C}E}\right]\mathsf{C}'dr + \frac{3\mathbf{s}^2}{2\mathsf{C}^4}.$$
(29)

The deformation induced in a self-gravitating body (as a result of varying gravitational field) of nearby object is estimated by the Weyl tensor  $(C^{\lambda}_{\vartheta \tau u})$ . This tensor is split into two components, i.e., electric and magnetic. For the spherical geometry, the magnetic component appears to be zero while electric part takes the following form

$$E_{\vartheta\tau} = \mathsf{C}_{\vartheta\mathfrak{u}\tau\nu} v^{\mathfrak{u}} v^{\nu} = \epsilon \left(-\frac{h_{\vartheta\tau}}{3} + \chi_{\vartheta}\chi_{\tau}\right),\tag{30}$$

where the Weyl scalar reads

$$\epsilon = \frac{1}{2A^2} \left[ \frac{\ddot{C}}{C} - \frac{\ddot{B}}{B} - \left( \frac{\dot{C}}{C} + \frac{\dot{A}}{A} \right) \left( \frac{\dot{C}}{C} - \frac{\dot{B}}{B} \right) \right] - \frac{1}{2C^2} + \frac{1}{2B^2} \left[ -\frac{C''}{C} + \frac{A''}{A} + \left( \frac{C'}{C} - \frac{A'}{A} \right) \left( \frac{B'}{B} + \frac{C'}{C} \right) \right].$$
(31)

The following relation explains the influence of tidal force on the charged matter distribution as

$$\frac{3m}{\mathsf{C}^3} = 4\pi \left[ -\Pi^{(\text{tot})} + \left( \mathfrak{u} + \frac{3\mathbf{s}^2}{8\pi\mathsf{C}^4} + \frac{T_{00}^{(\mathsf{GT})}}{\mathsf{A}^2} \right) \right] + \frac{3\mathbf{s}^2}{2\mathsf{C}^4} - \epsilon, \qquad (32)$$

where  $\Pi^{(tot)} = \Pi^{(GT)} + \Pi$ .

### **3** Structure Scalars

The concept of orthogonal splitting of the Riemann tenor was introduced by Bel [42]. Following this procedure, Herrera [43] developed different scalar functions to devise complexity of the system named as structure scalars. The Ricci tensor, Weyl tensor and Ricci scalar are defined in terms of the Riemann tensor as

$$R^{\rho}_{\vartheta\tau\mu} = \mathsf{C}^{\rho}_{\vartheta\tau\mu} + \frac{1}{2}R_{\vartheta\mu}\delta^{\rho}_{\tau} + \frac{1}{2}R^{\rho}_{\tau}g_{\vartheta\mu} - \frac{1}{2}R_{\vartheta\tau}\delta^{\rho}_{\mu} - \frac{1}{2}R^{\rho}_{\mu}g_{\vartheta\tau} - \frac{1}{6}R\left(\delta^{\rho}_{\tau}g_{\vartheta\mu} - g_{\vartheta\tau}\delta^{\rho}_{\mu}\right),\tag{33}$$

which can also be expressed in the form of matter source as

$$R_{\tau\delta}^{\vartheta\gamma} = \mathsf{C}_{\tau\delta}^{\vartheta\gamma} + 16\pi T_{[\tau}^{(\mathsf{tot})[\vartheta}\delta_{\delta]}^{\gamma]} + 8\pi T^{(\mathsf{tot})} \left(\frac{1}{3}\delta_{[\tau}^{\vartheta}\delta_{\delta]}^{\gamma} - \delta_{[\tau}^{[\vartheta}\delta_{\delta]}^{\gamma]}\right). \tag{34}$$

Here, we propose two tensors  $X_{\vartheta\tau}$  and  $Y_{\vartheta\tau}$  defined as

$$X_{\vartheta\tau} = {}^{*}R^{*}_{\vartheta\gamma\tau\delta}v^{\gamma}v^{\delta} = \frac{1}{2}\eta^{\epsilon\mu}_{\vartheta\gamma}R^{*}_{\epsilon\mu\tau\delta}v^{\gamma}v^{\delta}, \qquad (35)$$

$$Y_{\vartheta\tau} = R_{\vartheta\gamma\tau\delta}v^{\gamma}v^{\delta}, \qquad (36)$$

where  $R^*_{\vartheta\tau\gamma\delta} = \frac{1}{2}\eta_{\epsilon\mu\gamma\delta}R^{\epsilon\mu}_{\vartheta\tau}$  and  $\eta^{\epsilon\mu}_{\vartheta\gamma}$  indicates the Levi-Civita symbol. One can express these tensors into four scalars, i.e., a combination of trace  $(\mathcal{Y}_T, \mathcal{X}_T)$ and trace-free parts  $(\mathcal{Y}_{TF}, \mathcal{X}_{TF})$  as

$$Y_{\vartheta\tau} = (\chi_{\vartheta}\chi_{\tau} - \frac{h_{\vartheta\tau}}{3})\mathcal{Y}_{TF} + \frac{h_{\vartheta\tau}\mathcal{Y}_{T}}{3}, \qquad (37)$$

$$X_{\vartheta\tau} = (\chi_{\vartheta}\chi_{\tau} - \frac{h_{\vartheta\tau}}{3})\mathcal{X}_{TF} + \frac{h_{\vartheta\tau}\mathcal{X}_{T}}{3}.$$
 (38)

For the considered setup, the four structure scalars turn out to be

$$\mathcal{Y}_T = \frac{\mathbf{s}^2}{\mathsf{C}^4} + 4\pi \left( \mathfrak{u} + 3\mathfrak{p}_r - 2\Pi \right) + \frac{\left( \mathfrak{u} + \mathfrak{p} \right) f_T}{2} + \mathsf{M}^{(\mathsf{GT})}, \qquad (39)$$

$$\mathcal{Y}_{TF} = \epsilon - 4\pi \Pi - \frac{\mathbf{s}^2}{\mathsf{C}^4} + \frac{\Pi}{2} f_T + \mathsf{L}^{(\mathsf{GT})}, \qquad (40)$$

$$\mathcal{X}_T = 8\pi(\mathfrak{u} + \frac{\mathbf{s}^2}{8\pi \mathbf{C}^4}) + \mathbf{Q}^{(\mathsf{GT})}, \qquad (41)$$

$$\mathcal{X}_{TF} = -\epsilon - 4\pi\Pi - \frac{\mathbf{s}^2}{\mathsf{C}^4} + \frac{\Pi}{2}f_T, \qquad (42)$$

where  $\mathsf{L}^{(\mathsf{GT})} = \frac{\mathsf{J}_{\vartheta\tau}^{(\mathsf{GT})}}{\chi_{\vartheta\chi\tau} - \frac{1}{3}h_{\vartheta\tau}}$ . The extra terms in scalar functions  $(\mathsf{M}^{(\mathsf{GT})}, \mathsf{J}_{\vartheta\tau}^{(\mathsf{GT})}, \mathsf{Q}^{(\mathsf{GT})})$  are given in Appendix **B**. The contribution of energy density and charge in the system is attributed to the scalar  $\mathcal{X}_T$  whereas the anisotropic stresses along with electromagnetic effects are controlled by  $\mathcal{Y}_T$ . The scalar function  $\mathcal{Y}_{TF}$  in terms of all physical parameters is obtained using Eqs.(31) and (40) as

$$\mathcal{Y}_{TF} = \frac{4\pi}{\mathsf{C}^3} \int \mathsf{C}^3 \left[ D_\mathsf{C} \left( \mathfrak{u} + \frac{\mathbf{s}^2}{8\pi\mathsf{C}^4} + \frac{T_{00}^{(\mathsf{GT})}}{\mathsf{A}^2} \right) - 3 \left( \zeta - \frac{T_{01}^{(\mathsf{GT})}}{\mathsf{AB}} \right) \frac{U}{\mathsf{C}E} \right] \mathsf{C}' dr$$

$$- 4\pi\Pi - 4\pi\Pi^{(\mathsf{GT})} - \mathsf{L}^{(\mathsf{GT})} - \frac{\mathbf{s}^2}{2\mathsf{C}^4} + \frac{\Pi}{2}f_T.$$
(43)

One can notice from the above equation that the scalar  $\mathcal{Y}_{TF}$  is composed of heat flux, inhomogeneous energy density, modified terms, charge terms and pressure anisotropy. On the other hand,  $\mathcal{X}_{TF}$  includes the involvement of inhomogeneity due to energy density together with additional terms and charge as

$$\mathcal{X}_{TF} = 4\pi\Pi^{(\mathsf{GT})} - \frac{2\mathbf{s}^2}{\mathsf{C}^4} - \frac{4\pi}{\mathsf{C}^3} \int \mathsf{C}^3 \left[ D_\mathsf{C} \left( \mathfrak{u} + \frac{\mathbf{s}^2}{8\pi\mathsf{C}^4} + \frac{T_{00}^{(\mathsf{GT})}}{\mathsf{A}^2} \right) - 3 \left( \zeta - \frac{T_{01}^{(\mathsf{GT})}}{\mathsf{AB}} \right) \frac{U}{\mathsf{C}E} \right] \mathsf{C}' dr$$
(44)

### 4 Evolution Modes

The physical parameters (pressure, charge and energy density) have a significant impact on how the cosmic structures behave. The scalar  $\mathcal{Y}_{TF}$  involves energy density inhomogeneity, heat flux and pressure anisotropy along with f(G,T) corrections. The complexity factor for the anisotropic dissipative massive source in a non-static geometry is thus identified as  $\mathcal{Y}_{TF}$ . To obtain the complexity-free structure, we substitute  $\mathcal{Y}_{TF} = 0$  which is regarded as complexity-free condition. As the fluid under consideration is evolving with time, thus it is necessary to investigate the patterns of evolution. In the subsequent sections, two evolutionary modes, i.e., homologous evolution as well as homogeneous expansion are examined. We also determine the simplest mode that can reduce complexity in the evolutionary process.

#### 4.1 Homologous Evolution

The homologous refers to the phenomenon in which the whole system depicts a similar pattern. The collapse happens when all the substance of any astronomical object descends towards the center in the interior region. A direct relationship between radial distance and velocity is observed in homologous collapse, implying that during the collapse, all material of the astrophysical entity falls into its core at the same rate. In contrast to homologously collapsing objects, those structures whose cores collapse first will thus generate less gravitational radiations. Equation (23) can also be written as

$$D_{\mathsf{C}}\left(\frac{U}{\mathsf{C}}\right) = \frac{\sigma}{\mathsf{C}} + \frac{4\pi}{E}\left(\zeta - \frac{T_{01}^{(\mathsf{GT})}}{\mathsf{AB}}\right),\tag{45}$$

and the integration of the above equation leads to

$$U = \mathsf{C} \int_0^r \left[ \frac{\sigma}{\mathsf{C}} + \frac{4\pi}{E} \left( \zeta - \frac{T_{01}^{(\mathsf{GT})}}{\mathsf{AB}} \right) \right] \mathsf{C}' dr + h(t) \mathsf{C}, \tag{46}$$

where integration constant is represented by h(t) and its value at the boundary gives

$$U = \frac{U_{\Sigma}}{\mathsf{C}_{\Sigma}}\mathsf{C} - \mathsf{C} \int_{r}^{r_{\Sigma}} \left[ \frac{\sigma}{\mathsf{C}} + \frac{4\pi}{E} \left( \zeta - \frac{T_{01}^{(\mathsf{GT})}}{\mathsf{AB}} \right) \right] \mathsf{C}' dr.$$
(47)

The factors responsible for the deviation of matter source from the homologous mode are heat dissipation and shear scalar. To observe the necessary condition for homologous evolution [44, 45], i.e.,  $U \sim \mathsf{C}$  the integrand in the above equation must cancel the effect of each other. Further, it yields  $U = h(t)\mathsf{C}$  where  $h(t) = \frac{U_{\Sigma}}{\mathsf{C}_{\Sigma}}$ . In f(G,T) gravity, the homologous condition is of the form

$$\frac{\sigma}{\mathsf{C}} + \frac{4\pi\mathsf{B}}{\mathsf{C}'}\left(\zeta - \frac{T_{01}^{(\mathsf{GT})}}{\mathsf{AB}}\right) = 0.$$
(48)

#### 4.2 Homogeneous Expansion

Another mode under discussion is homogeneous expansion for which the required condition is  $\Theta' = 0$ . In contrast to preceding mode, the homogeneous contraction or expansion occurs when its rate is independent of r. The combination of this condition on Eq.(23) becomes

$$4\pi \left(\zeta - \frac{T_{01}^{(\mathsf{GT})}}{\mathsf{AB}}\right) = -\frac{\mathsf{C}'}{\mathsf{B}} \left[\frac{\sigma}{\mathsf{C}} + \frac{1}{3}D_{\mathsf{C}}(\sigma)\right].$$
(49)

Utilizing the homologous condition in Eq.(49) provides  $D_{\mathsf{C}}(\sigma) = 0$ . The regularity conditions at the core give rise to  $\sigma = 0$ , and its implication in the above equation yields

$$\zeta = \frac{T_{01}^{(\text{GT})}}{\text{AB}}.$$
(50)

This shows that the fluid incorporates the influence of heat dissipation. In the framework of general relativity (GR), the matter distribution was nondissipative as well as shear-free during homogeneous evolution.

### 5 Kinematical and Dynamical Quantities

Here, we investigate the behavior of some physical entities to select the simplest evolutionary mode. For our convenience, we assume that the metric coefficient C is a separable function of t and r. Employing homologous condition (48) in Eq.(23), we obtain

$$(\Theta - \sigma)' = \left(\frac{3\dot{\mathsf{C}}}{\mathsf{A}\mathsf{C}}\right)' = 0.$$
(51)

This equation indicates the geodesic nature of matter configuration as A' = 0 (a = 0) implying that A = 1 without any loss of generality. Combining the values of  $\sigma$  and  $\Theta$  together with the geodesic condition leads to

$$\Theta - \sigma = \frac{3\dot{\mathsf{C}}}{\mathsf{C}}.\tag{52}$$

The homologous condition is recovered by taking the successive derivatives of Eq.(52). One can observe that the necessary and sufficient condition for the dynamical sphere to evolve homologously is that the fluid must obey the geodesic path and vice-versa. In GR, the unavailability of heat flux ( $\zeta = 0$ ) corresponds to a shear-free matter distribution, while in this modified theory its effect can be seen as

$$\sigma = 4\pi \frac{T_{01}^{(\mathsf{GT})}\mathsf{C}}{\mathsf{C}'}.$$
(53)

If the matter source follows the homogeneous collapse with no dissipation then we have  $T_{01}^{(GT)} = 0$ , while the shear scalar from Eq.(49) becomes

$$\sigma = \frac{12\pi}{\mathsf{C}^3} \int \frac{\mathsf{C}^3 T_{01}^{(\mathsf{GT})}}{\mathsf{A}} dr + \frac{b(t)}{\mathsf{C}^3},\tag{54}$$

where b(t) stands for the integration function. Thus, one can say that when we neglect the additional terms and heat dissipation, the homogeneous expansion implies homologous condition as  $\sigma = 0 \Rightarrow U \sim \mathsf{C}$ . This result shows that homologous evolution is the simplest mode. During collapse, the mass and velocity undergoing homologous evolution are interlinked as

$$D_T U = \frac{\mathbf{s}^2}{2\mathbf{C}^3} - \frac{m}{\mathbf{C}^2} - 4\pi \left( \mathbf{p}_r - \frac{\mathbf{s}^2}{8\pi\mathbf{C}^4} + \frac{T_{11}^{(\mathsf{GT})}}{\mathbf{B}^2} \right) \mathsf{C}.$$
 (55)

In terms of  $\mathcal{Y}_{TF}$ , the above expression can be obtained as

$$\frac{3D_T U}{\mathsf{C}} = \mathcal{Y}_{TF} - \frac{\Pi}{2} f_T - \mathsf{L}^{(\mathsf{GT})} + \frac{\mathbf{s}^2}{\mathsf{C}^4} + 4\pi \Pi^{(\mathsf{GT})} - 4\pi \left[ \left( \mathfrak{u} + \frac{\mathbf{s}^2}{8\pi \mathsf{C}^4} + \frac{T_{00}^{(\mathsf{GT})}}{\mathsf{A}^2} \right) + 3 \left( \mathfrak{p}_r - \frac{\mathbf{s}^2}{8\pi \mathsf{C}^4} + \frac{T_{11}^{(\mathsf{GT})}}{\mathsf{B}^2} \right) - 2\Pi \right].$$
(56)

After some manipulations, we have

$$\frac{\ddot{\mathsf{C}}}{\mathsf{C}} - \frac{\ddot{\mathsf{B}}}{\mathsf{B}} = \mathcal{Y}_{TF} - \mathsf{L}^{(\mathsf{GT})} + \frac{\mathbf{s}^2}{\mathsf{C}^4} - \frac{\Pi}{2} f_T + 4\pi \Pi^{(\mathsf{GT})}.$$
(57)

This shows that charged spherical distribution gets complexity-free when

$$\frac{\ddot{\mathsf{C}}}{\mathsf{C}} - 4\pi\Pi^{(\mathsf{GT})} - \frac{\mathbf{s}^2}{\mathsf{C}^4} - \frac{\ddot{\mathsf{B}}}{\mathsf{B}} + L^{(\mathsf{GT})} + \frac{\Pi}{2}f_T = 0.$$
(58)

Here, we find feasible solutions corresponding to non-dissipative and dissipative scenarios by using zero complexity and homologous conditions. The system under consideration is non-static and modified field equations contain non-linear terms, thus, a linear f(G,T) model of the form  $f(G,T) = \gamma G^n + \omega T$  [46, 47] is chosen for the sake of simplicity. Moreover,  $\gamma$  and  $\omega$ denote the real numbers and n > 0, thus we can take  $\gamma = n = 1$ . There are three metric potentials (A(t,r), B(t,r), C(t,r)) to be calculated in the current setup. As we have selected A = 1, so we are left with only two unknowns. It means that two equations are required to find the unknowns. For this reason, the zero complexity and homologous conditions are utilized. These constraints, in the case of non-dissipative fluid, are calculated as

$$\frac{1}{\mathsf{B}^{3}\mathsf{C}^{2}(\omega+8\pi)}\left[\left(-2\mathsf{C}\left(5\ddot{\mathsf{B}}\dot{\mathsf{C}}'+\mathsf{C}'\dot{\mathsf{B}}\right)+\dot{\mathsf{B}}\mathsf{C}'\ddot{\mathsf{C}}\right)\mathsf{B}+10\dot{\mathsf{B}}\ddot{\mathsf{B}}\mathsf{C}\mathsf{C}'\right.\\\left.+2\mathsf{B}^{2}\dot{\mathsf{C}}'\left(\mathsf{C}-4\ddot{\mathsf{C}}\right)\right]=0,\tag{59}$$

$$\frac{1}{2\mathsf{B}^{4}\mathsf{C}^{4}}\left\{2\mathsf{B}^{3}\dot{\mathsf{B}}\mathsf{C}^{2}\dot{\mathsf{C}}\left(-4\ddot{\mathsf{C}}+\mathsf{C}\right)+\mathsf{B}\mathsf{C}^{2}\mathsf{C}'\left(-2\mathsf{B}'\mathsf{C}+\left(\dot{\mathsf{B}}^{2}+3\ddot{\mathsf{B}}\right)\mathsf{C}'\right.\\\left.+32\dot{\mathsf{B}}\dot{\mathsf{C}'}^{2}\right)-4\dot{\mathsf{B}}^{2}\mathsf{C}^{2}\mathsf{C}'^{2}+\mathsf{B}^{4}\left(\mathsf{s}^{2}-2\mathsf{C}^{2}\left(\dot{\mathsf{C}}^{2}+1\right)\right)+\mathsf{B}^{2}\mathsf{C}^{2}\left(2\mathsf{C}'^{2}+16\dot{\mathsf{C}'}^{2}-2\mathsf{C}\dot{\mathsf{C}}+\mathsf{s}^{2}\right)\right\}=0.$$
(60)

For the second case (dissipative fluid), the complexity-free condition remains the same as for non-dissipative source while the homologous condition turns out to be

$$\zeta = \frac{1}{16\pi^{2}\mathsf{C}^{4}\mathsf{B}^{5}(\omega+1)} \big[\mathsf{C}'\big(-\mathsf{B}\dot{\mathsf{C}}+\dot{\mathsf{B}}\mathsf{C}\big)\big(\mathsf{B}\big(\mathsf{C}\big(40\pi\dot{\mathsf{C}}'\ddot{\mathsf{B}}+\dot{\mathsf{B}}\mathsf{C}'\big)-4\pi\ddot{\mathsf{C}}\dot{\mathsf{B}}\mathsf{C}'\big) - 40\pi\dot{\mathsf{B}}\ddot{\mathsf{B}}\mathsf{C}\mathsf{C}'+\mathsf{B}^{2}\big(-\dot{\mathsf{C}}'\big)\big(\mathsf{C}-32\pi\ddot{\mathsf{C}}\big)\big)\big].$$
(61)

### 6 Stability of Zero Complexity Condition

It is possible that the system has zero complexity at first and then becomes complicated due to the presence of some factors as it evolves. These factors are determined by working on the evolution equation for the scalar  $\mathcal{Y}_{TF}$ following the technique [43] together Eqs.(17), (40) and (42) as

$$-4\pi \left(\mathfrak{p}_{r} + \frac{T^{11(\text{GT})}}{\text{B}^{2}} + \mathfrak{u} + T^{00(\text{GT})}\right)\sigma + \frac{\mathbf{s}^{2}}{\text{C}^{4}} + \frac{\mathbf{s}^{2}\mathbf{s}'}{\text{C}^{4}\text{B}} - \frac{4\pi}{\text{B}}\left[\zeta' - (\frac{T^{00(\text{GT})}}{\text{B}^{2}})'\text{B}\right] \\ -(\zeta - \frac{T^{01(\text{GT})}}{\text{B}^{2}})\frac{\text{C}'}{\text{C}} - \dot{\mathcal{Y}}_{TF} + \frac{3\mathbf{s}\dot{\mathbf{s}}}{\text{C}^{4}} - \dot{\textbf{L}}^{(\text{GT})} - 8\pi\dot{\Pi} - 4\pi\dot{\Pi}^{(\text{GT})} - 4\pi \left(T^{(\text{GT})}_{00}\right)'$$
$$-3\frac{\dot{\textbf{C}}}{\textbf{C}}\left(\mathcal{Y}_{TF} - \textbf{L}^{(\text{GT})}\right) - 12\pi \left(T^{00(\text{GT})} + \frac{\mathbf{s}^{2}}{8\pi\text{C}^{4}}\right)\frac{\dot{\textbf{C}}}{\textbf{C}} - 4\pi\text{Z}_{1} + \frac{4\pi T^{00(\text{GT})}\text{B}'}{\text{B}^{2}} \\ + \left(12\pi T^{(\text{GT})}_{00} - 12\pi\Pi^{(\text{GT})} + \frac{12\pi T^{(\text{GT})}_{11}}{\text{B}^{2}}\right)\frac{\dot{\textbf{C}}}{\textbf{C}} - \frac{16\pi\Pi^{(\text{tot})}\dot{\textbf{C}}}{\textbf{C}} - \frac{5\mathbf{s}^{2}\dot{\textbf{C}}}{2\mathbf{C}^{5}} = 0. \quad (62)$$

For the non-dissipative source, we substitute  $\mathcal{Y}_{TF} = \Pi^{(\text{tot})} = \zeta = \sigma = 0$  in the above equation at the initial time (t = 0), so that it becomes

$$-\dot{\mathcal{Y}}_{TF} - 4\pi \left(T_{00}^{(\mathsf{GT})}\right) - \dot{\mathsf{L}}^{(\mathsf{GT})} + 4\pi \left(\frac{T^{00(\mathsf{GT})}}{\mathsf{B}^{2}}\right)' - 4\pi \dot{\Pi}^{(\mathsf{GT})} - 8\pi \dot{\Pi} + 3\mathsf{L}^{(\mathsf{GT})}$$
$$\times \frac{\dot{\mathsf{C}}}{\mathsf{C}} - 4\pi\mathsf{Z}_{1} - 12\pi \left(T^{00(\mathsf{GT})} + \frac{\mathsf{s}^{2}}{8\pi\mathsf{C}^{4}}\right) \frac{\dot{\mathsf{C}}}{\mathsf{C}} + \frac{4\pi T^{00(\mathsf{GT})}\mathsf{B}'}{\mathsf{B}^{2}} + \left(12\pi T_{00}^{(\mathsf{GT})} + \frac{12\pi T_{11}^{(\mathsf{GT})}}{\mathsf{B}^{2}}\right)$$

$$\times \frac{\dot{\mathsf{C}}}{\mathsf{C}} + \frac{\mathbf{s}^2}{\mathsf{C}^4} + \frac{\mathbf{s}^2 \mathbf{s}'}{\mathsf{C}^4 \mathsf{B}} + \frac{3\mathbf{s}\dot{\mathbf{s}}}{\mathsf{C}^4} - \frac{5\mathbf{s}^2 \dot{\mathsf{C}}}{2\mathsf{C}^5} = 0.$$

Making use of the above constraint with the derivative of Eq.(43) at t = 0 provides

$$\frac{4\pi}{\mathsf{C}^{3}} \int_{0}^{r} \left[ \left(\mathfrak{u} + \frac{\mathbf{s}^{2}}{8\pi\mathsf{C}^{4}} + \frac{T_{00}^{(\mathsf{GT})}}{\mathsf{A}^{2}}\right) \cdot \right]' = 4\pi \left(\frac{T^{00(\mathsf{GT})}}{\mathsf{B}^{2}}\right)' - 4\pi \left(T_{00}^{(\mathsf{GT})}\right) \cdot + \frac{4\pi T^{00(\mathsf{GT})}\mathsf{B}'}{\mathsf{B}^{2}} - 12\pi \left(T^{00(\mathsf{GT})} + \frac{\mathbf{s}^{2}}{8\pi\mathsf{C}^{4}}\right) \frac{\dot{\mathsf{C}}}{\mathsf{C}} + 3L^{(\mathsf{GT})} \frac{\dot{\mathsf{C}}}{\mathsf{C}} - 4\pi\mathsf{Z}_{1} + \left(12\pi T_{00}^{(\mathsf{GT})} + \frac{12\pi T_{11}^{(\mathsf{GT})}}{\mathsf{B}^{2}}\right) \times \frac{\dot{\mathsf{C}}}{\mathsf{C}} + \frac{\mathbf{s}^{2}}{\mathsf{C}^{4}} + \frac{\mathbf{s}^{2}\mathbf{s}'}{\mathsf{C}^{4}} + \frac{4\mathbf{s}\dot{\mathsf{s}}}{\mathsf{C}^{4}} - \frac{\mathbf{s}^{2}\dot{\mathsf{C}}}{2\mathsf{C}^{5}}.$$
(63)

The stability of the complexity-free condition is associated with the pressure and energy density. The above equation demonstrates that the contributions of pressure anisotropy, charge, modified terms and energy density inhomogeneity in the inner matter source forced the system to deviate from the stable position. This depicts that the stability of the system is disturbed by the presence of electromagnetic field. Further, using the conditions  $\mathcal{Y}_{TF} = \sigma = 0$  in conjunction with the derivative of Eq.(62) with respect to t yields

$$\begin{aligned} &-\ddot{\mathcal{Y}}_{TF} - 4\pi\ddot{\Pi}^{(\mathsf{GT})} + \left[4\pi\left(\frac{T^{00(\mathsf{GT})}}{\mathsf{B}^{2}}\right)'\right]^{\cdot} - \ddot{\mathsf{L}}^{(\mathsf{GT})} - 4\pi\left(T^{(\mathsf{GT})}_{00}\right)^{\cdot} + \dot{\mathsf{L}}^{(\mathsf{GT})} - 4\pi\dot{Z}_{1} \\ &- 12\pi\left\{\left(\frac{\mathbf{s}^{2}}{8\pi\mathsf{C}^{4}} + T^{00(\mathsf{GT})}\right)\frac{\dot{\mathsf{C}}}{\mathsf{C}}\right\}^{\cdot} + \frac{3\dot{\mathsf{C}}}{\mathsf{C}}\left[4\pi\left(T^{(\mathsf{GT})}_{00}\right)^{\cdot} - 4\pi\left(\frac{T^{00(\mathsf{GT})}}{\mathsf{B}^{2}}\right)' + 4\pi\dot{\Pi}^{(\mathsf{GT})} \\ &+ 4\pi\mathsf{Z}_{1} + \frac{3\mathbf{s}\dot{\mathbf{s}}}{\mathsf{C}^{4}} - 4\pi\left(\frac{\mathsf{B}'T^{00(\mathsf{GT})}}{\mathsf{B}^{2}}\right) - 3L^{(\mathsf{GT})}\frac{\dot{\mathsf{C}}}{\mathsf{C}} + 12\pi\left(\frac{T^{00(\mathsf{GT})}\dot{\mathsf{C}}}{\mathsf{C}}\right) + 8\pi\dot{\Pi} - \frac{5\mathsf{s}^{2}\dot{\mathsf{C}}}{2\mathsf{C}^{5}} \\ &- \left(12\pi T^{(\mathsf{GT})}_{00} + \frac{12\pi T^{(\mathsf{GT})}_{11}}{\mathsf{B}^{2}}\right)\frac{\dot{\mathsf{C}}}{\mathsf{C}} + \frac{\mathsf{s}^{2}}{\mathsf{C}^{4}} + \frac{\mathsf{s}^{2}\mathsf{s}'}{\mathsf{C}^{4}\mathsf{B}}\right] + 3\left(\frac{L^{(\mathsf{GT})}\dot{\mathsf{C}}}{\mathsf{C}}\right)^{\cdot} - \frac{16\pi\Pi^{(\mathsf{tot})}\dot{\mathsf{C}}}{\mathsf{C}} - 8\pi\ddot{\Pi} \\ &+ \left[(12\pi T^{(\mathsf{GT})}_{00} + \frac{12\pi T^{(\mathsf{GT})}_{11}}{\mathsf{B}^{2}})\frac{\dot{\mathsf{C}}}{\mathsf{C}}\right]^{\cdot} + 4\pi\left(\frac{\mathsf{B}'T^{00(\mathsf{GT})}}{\mathsf{B}^{2}}\right)^{\cdot} + \left(\frac{\mathsf{s}^{2}}{\mathsf{C}^{4}} + \frac{\mathsf{s}^{2}\mathsf{s}'}{\mathsf{C}^{4}\mathsf{B}} + \frac{3\mathsf{s}\dot{\mathsf{s}}}{\mathsf{C}^{4}} - \frac{5\mathsf{s}^{2}\dot{\mathsf{C}}}{2\mathsf{C}^{5}}\right)^{\cdot} = 0. \end{aligned}$$

It is worth mentioning here that for the dissipative case, the heat flux works as an additional factor in disturbing the vanishing complexity constraint.

### 7 Concluding Remarks

In this paper, we have checked the influence of non-minimal theory to calculate the complexity of charged dynamical structure. The interior region was filled with energy density inhomogeneity, anisotropic pressure, heat flux and charge. Using Herrera's method, we have divided the Riemann tensor into four scalars, each of which is associated with a certain set of physical attributes. Among all the four scalars, we have considered  $\mathcal{Y}_{TF}$  to produce complexity for the reasons stated below.

- 1. In GR and f(G,T) gravity, this factor has already been worked as complexity factor for the static spacetime. Thus, we can reobtain this complexity-producing scalar for the static spherical system from the non-static geometry by using Eq.(40).
- 2. All the physical parameters (anisotropic pressure, energy density inhomogeneity, heat flux, charge and correction terms) are incorporated in this factor that might increase the complexity of the system.

We have looked at two different types of evolutionary patterns, homologous and homogenous. By using the criterion of zero complexity and selecting homologous mode as the most basic pattern of evolution, we are able to obtain the results for both dissipative and non-dissipative scenarios. The variables that lead the system to depart from zero complexity criterion throughout the evolution process have also been covered.

The fundamental properties of the charged complex sphere are strongly influenced by the state determinants and additional curvature factors in f(G,T) gravity. As a result, the charged dynamical sphere becomes complicated due to the inclusion of higher order curvature factors. Since the homologous fluid is assumed to be geodesic in nature (A = 1), this mode was proposed as the most basic pattern of evolution. Further, the homologous condition contains the extra terms of this theory. In GR, the conditions  $\mathfrak{u} = \Pi = \zeta = 0$  are responsible for the formation of a complex free system  $(\mathcal{Y}_{TF} = 0)$ , whereas in this theory, the structure corresponds to complexityfree only when the constraint  $L^{(\text{GT})} - \frac{s^2}{c^4} + \frac{\Pi}{2}f_T - 4\pi\Pi^{(\text{GT})} = 0$  and the aforementioned factors are satisfied. We have seen that the presence of correction terms causes the shear to be established even in the non-dissipative case. We have employed the modified complexity free and homologous conditions to assess the feasible solutions representing dissipative and non-dissipative scenarios. The stability criterion for the vanishing complexity condition has also been analyzed. Finally, we have addressed the factors responsible for the deviation of the system from zero complexity. We have also found that the contribution of charge and extra curvature terms of this modified theory make the system more complex. It is worth mentioning here that the constraint f(G,T) = 0 reduces all the obtained results to GR.

## Appendix A

The additional terms present in the field equations of f(G,T) theory are described as

$$\begin{split} T_{00}^{(\text{GT})} &= \frac{1}{8\pi} \left[ (\mathfrak{u} + \mathfrak{p}) A^2 f_T - \frac{A^2}{2} f + \left( \frac{8\dot{B}\dot{C}\ddot{C}}{A^2BC^2} - \frac{16\dot{A}'\dot{C}\dot{C}'}{AB^2C^2} + \frac{8AC'A'C''}{B^4C^2} \right. \\ &+ \frac{4\ddot{B}}{BC^2} + \frac{8\dot{A}'\dot{B}\dot{C}\dot{C}'}{AB^3C^2} - \frac{8\dot{A}B'C'\dot{C}}{AB^3C^2} + \frac{8\dot{A}\dot{C}C''}{AB^2C^2} + \frac{4\dot{C}^2\dot{A}'}{AC^2} - \frac{12\dot{A}\dot{B}\dot{C}^2}{A^3BC^2} \\ &\times \frac{B'}{B^3} + \frac{4\dot{A}\dot{C}'^2\dot{B}}{AB^3C^2} - \frac{12AA'B'C'^2}{B^5C^2} + \frac{4AC'^2A''}{B^4C^2} + \frac{4\dot{C}^2\ddot{B}}{A^2BC^2} + \frac{8\dot{C}'^2}{B^2C^2} \\ &+ \frac{8B'\ddot{C}C'}{B^3C^2} - \frac{16\dot{B}\dot{C}'\dot{C}'}{C^2B^3} - \frac{4\dot{A}\dot{B}}{AC^2B} - \frac{4\dot{C}^2\ddot{A}'}{B^2AC^2} + \frac{8\dot{C}^{2}A'^2}{B^2A^2C^2} + \frac{8C'^2\dot{B}^2}{C^2B^4} \\ &- \frac{8\ddot{C}C''}{C^2B^2} - \frac{4AA''}{C^2B^2} + \frac{B'AA'}{B^3C^2} - \frac{4C'^2\ddot{B}}{B^3C^2} \right) f_G + \left( \frac{8\dot{B}\dot{C}C'}{B^2C^2} - \frac{8B'C'\dot{C}}{B^3C^2} \right) \\ &- \frac{4\dot{B}}{BC^2} - \frac{12\dot{B}\dot{C}^2}{A^2BC^2} + \frac{4\dot{B}\dot{C}'^2}{B^3C^2} \right) \dot{f}_G + \left( \frac{8\dot{B}\dot{C}C'}{B^2C^2} - \frac{4A^2B'}{B^3C^2} - \frac{4A^2B'}{B^3C^2} \right) \\ &+ \frac{12A^2B'C'^2}{B^5C^2} - \frac{8A^2C'C''}{B^4C^2} \right) f_G' + \left( \frac{4\dot{C}^2}{B^2C^2} + \frac{4A^2}{B^2C^2} - \frac{4A^2C'^2}{B^3C^2} \right) \\ &\times f''_G], \end{split} \tag{A1}$$

$$\begin{aligned} & - \frac{A'\dot{C}B'\dot{C}'}{AB^{3}CC^{2}} + \frac{8A'\ddot{C}\ddot{C}}{A^{3}C^{2}} - \frac{8A'\dot{A}\dot{C}^{2}}{A^{4}C^{2}} + \frac{8\dot{A}\dot{C}\dot{C}'}{A^{3}CC^{2}} - \frac{10\ddot{B}\dot{C}'}{A^{2}BC} + \frac{10A''\dot{C}'}{AB^{2}C} \\ & - \frac{8\ddot{C}\dot{C}'}{A^{3}C^{2}} \int f_{G} + \left( -\frac{8\dot{A}\dot{B}\dot{A}'}{A^{4}B} - \frac{4A'}{AC^{2}} + \frac{8\dot{C}\dot{C}'}{A^{2}C^{2}} + \frac{4A'C'^{2}}{AB^{2}C^{2}} + \frac{8A'^{2}B'}{A^{2}B^{3}} \right) \\ & - \frac{12A'\dot{C}^{2}}{A^{3}C^{2}} - \frac{8A'A''}{A^{2}B^{2}} + \frac{8A'\ddot{B}}{A^{3}B} \right) \dot{f}_{G} \left( \frac{8A'C'\dot{C}}{AB^{2}C^{2}} + \frac{8A'B'\dot{B}}{AB^{4}} - \frac{4\ddot{B}\dot{C}^{2}}{A^{2}BC^{2}} \right) \\ & - \frac{8\dot{A}\dot{B}}{A^{3}B^{2}} + \frac{12\dot{B}\dot{C}'^{2}}{B^{3}C^{2}} + \frac{8\ddot{B}\dot{B}}{A^{2}B^{2}} - \frac{8A''\dot{B}}{AB^{3}} - \frac{8C'\dot{C}'}{B^{2}C^{2}} - \frac{4\dot{B}}{BC^{2}} \right) f'_{G} + (4) \\ & \times \frac{\dot{C}^{2}}{A^{2}C^{2}} + \frac{4}{C^{2}} - \frac{4C'^{2}}{B^{2}C^{2}} + \frac{8A''}{AB^{2}} - \frac{8\ddot{B}}{AB^{3}} - \frac{8A'B'}{AB^{3}} + \frac{8\dot{A}\dot{B}}{A^{3}B} \right) \dot{f}'_{G} \right], \quad (A2) \\ T_{11}^{(GT)} &= \frac{1}{8\pi} \left[ \frac{2}{3}\Pi B^{2}f_{T} + \frac{B^{2}}{2}f + \left( \frac{32\dot{B}C'\dot{C}'^{2}}{A^{2}BC^{2}} - \frac{8A'C'C''}{AB^{2}C^{2}} - \frac{8B\dot{B}\dot{C}\ddot{C}}{A^{2}C^{4}} - 8 \right) \\ & \times \frac{B'\ddot{C}C'}{BC^{2}A^{2}} + \frac{\ddot{B}C'^{2}}{A^{2}BC^{2}} - \frac{4A''C'^{2}}{AB^{2}C^{2}} - \frac{16\dot{C}'^{2}}{A^{2}C^{2}} - \frac{8\dot{A}\dot{C}C''}{A^{3}C^{2}} - \frac{8B\dot{B}\dot{C}\ddot{C}}{A^{3}C^{2}} - 24A''\dot{C}^{2}}{A^{3}C^{2}} - \frac{4B\ddot{B}\dot{A}}{A^{3}C^{2}} - \frac{4A'B'}{A^{3}C^{2}} - 24A''\dot{C}^{2}}{A^{3}C^{2}} - 24A''\dot{C}^{2} - 24A''\dot{C}^{2} + \frac{4A''C'^{2}}{A^{3}C^{2}} - 24A''\dot{C}^{2} + \frac{4A''C'^{2}}{A^{3}C^{2}} - 24A''\dot{A}^{2}\dot{C}^{2} - 24A''\dot{A}^{2}\dot{A}^{2}\dot{C}^{2} - 24A''\dot{A}^{2}\dot{A}^{2}\dot{C}^{2} - 24A''\dot{A}^{2}\dot{A}^{2}\dot{C}^{2} + \frac{4\dot{A}B\dot{C}'}{A^{3}C^{2}} - 24A''\dot{C}^{2}\dot{A}^{2}\dot{A}^{2}\dot{C}^{2} + \frac{4\dot{A}B\dot{C}}{A^{3}C^{2}} - \frac{24\dot{A}'C'^{2}}{A^{3}C^{2}} - 24A''\dot{A}^{2}\dot{C}^{2} + \frac{4\dot{A}B''}{A^{3}BC^{2}} - \frac{24\dot{A}'C'^{2}}{A^{3}C^{2}} - 24A''\dot{A}^{2}\dot{A}^{2}\dot{C}^{2} + \frac{4\dot{A}C'^{2}}{A^{3}C^{2}} + \frac{4\dot{A}C'^{2}}{A^{3}C^{2}} + \frac{4\dot{A}C'^{2}}{A^{3}C^{2}} - \frac{4\dot{A}\dot{A}C'^{2}}{A^{3}C^{2}} + \frac{4\dot{A}\dot{A}C'^{2}}{A^{3}C^{2}} + \frac{4\dot{A}C'^{2}}{A^{3}C^{2}} + \frac{4\dot{A}C'^{2}}{A^{3}C^{2}} + \frac{4\dot{A}C'^{2}}{A^{3}C^{$$

$$\times \frac{C''}{B^2} - \frac{4A'B'}{AB^3} - \frac{8\dot{B}\dot{C}\ddot{C}}{A^4B} - \frac{8C'B'\ddot{C}}{B^3A^2} + \frac{8A'\dot{C}\dot{C}'}{A^3B^2} + \frac{4\dot{A}\dot{B}}{A^3B} - \frac{4C'^2A''}{AB^4} \\ - \frac{4A'^2\dot{C}^2}{A^4B^2} - \frac{4\ddot{B}\dot{C}^2}{A^4B} + \frac{4\ddot{B}C'^2}{A^2B^3} + \frac{4A''\dot{C}^2}{A^3B^2} + \frac{8\ddot{C}C''}{A^2B^2} - \frac{4\dot{B}^2C'^2}{A^2B^4} \right) f_G \\ + \left(\frac{4\ddot{B}\dot{C}\dot{C}}{A^4B} - \frac{12\dot{A}\dot{B}\dot{C}\dot{C}}{A^5B} - \frac{4\dot{A}B'CC'}{A^3B^3} + \frac{8A'C'\dot{B}}{A^3B^3} + \frac{4A'B'\dot{C}\dot{C}}{A^3B^3} - \frac{4}{A^3} \right) f_G \\ \times \frac{A''\dot{C}\dot{C}}{B^2} + \frac{4\ddot{B}\dot{C}\ddot{C}}{A^4B} - \frac{12A'\dot{C}\dot{C}'}{A^3B^2} + \frac{4\dot{A}\dot{C}C''}{A^3B^2} + \frac{12\dot{C}\dot{C}A'^2}{A^4B^2} \right) \dot{f}_G + (8 \\ \times \frac{A'\dot{C}\dot{B}\dot{C}}{A^3B^3} + \frac{4B'C\ddot{C}}{B^3A^2} + \frac{4\dot{C}\dot{A}\dot{C}\dot{B}}{A^3B^3} - \frac{12C'CA'B'}{A^3B^2} - \frac{4\dot{C}\dot{C}B'\dot{A}}{A^3B^3} + \frac{4}{B^4} \\ \times \frac{CC'A''}{A} + \frac{8\dot{B}\dot{C}\dot{C}\dot{C}}{A^2B^3} - \frac{4\ddot{B}\dot{C}'C}{B^3A^2} - \frac{12\dot{B}\dot{C}'C}{B^3A^2} + \frac{4\dot{C}\dot{A}\dot{C}C''}{B^4A} + \frac{12A'^2}{A^2} \\ \times \frac{\dot{C}\dot{C}}{B^4} f_G' + \left(\frac{4\dot{A}\dot{C}\dot{C}}{A^3B^2} + \frac{4A'C'C}{AB^4} - \frac{4\ddot{C}\dot{C}}{A^2B^2} \right) f_G' + \left(\frac{12\dot{C}\dot{C}'}{A^2B^2} - \frac{12\dot{C}}{A^2} \\ \times \frac{\dot{B}C'}{B^3} - \frac{12A'\dot{C}\dot{C}}{A^3B^2} \dot{F}_G' + \left(\frac{4\dot{B}\dot{C}C}{A^3B^2} + \frac{4B'C'C}{AB^4} - \frac{4\ddot{C}C}{A^2B^2} \dot{F}_G' + \frac{12\dot{C}\dot{C}}{A^2B^2} - \frac{12\dot{C}}{A^2} \\ \times \frac{\dot{B}C'}{B^3} - \frac{12A'\dot{C}\dot{C}}{A^3B^2} \dot{F}_G' + \left(\frac{4\dot{B}\dot{C}C}{A^4B} + \frac{4B'C'C}{A^2B^3} - \frac{4\dot{C}C''}{A^2B^2} \dot{F}_G' \right) \dot{F}_G' \right].$$
(A4)

The modified expressions of  $Z_1 \mbox{ and } Z_2 \mbox{ are given as }$ 

$$Z_{1} = \frac{f_{T}}{8\pi - f_{T}} \left[ \left( \ln f_{T} \right)^{\cdot} \left( \frac{\mathfrak{u}}{\mathsf{A}} + \frac{\mathfrak{s}^{2}}{8\pi\mathsf{C}^{4}\mathsf{A}} + \frac{T^{00(\mathsf{GT})}}{\mathsf{A}^{3}} + \frac{\mathfrak{p}}{\mathsf{A}} \right) + 2 \left( \frac{\zeta}{\mathsf{B}} - \frac{T^{01(\mathsf{GT})}}{\mathsf{A}\mathsf{B}^{2}} \right) \right] \\ + \frac{1}{2\mathsf{A}} \left( \mathfrak{u} + 3\mathfrak{p} \right)^{\cdot} + \left( \ln f_{T} \right)^{\prime} \left( \frac{\zeta}{\mathsf{B}} - \frac{T^{01(\mathsf{GT})}}{\mathsf{A}\mathsf{B}^{2}} \right) + \left( \frac{2\mathfrak{u}}{\mathsf{A}} + \frac{\mathfrak{s}^{2}}{4\pi\mathsf{C}^{4}\mathsf{A}} + \frac{2T^{00(\mathsf{GT})}}{\mathsf{A}^{3}} \right) \\ + \frac{\mathfrak{p}}{\mathsf{A}} \right)^{\cdot} \right],$$

$$Z_{2} = \frac{f_{T}}{8\pi - f_{T}} \left[ \left( \ln f_{T} \right)^{\prime} \left( \frac{-\mathfrak{p}_{r}}{\mathsf{B}} - \frac{\mathfrak{s}^{2}}{8\pi\mathsf{C}^{4}\mathsf{B}} + \frac{T^{11(\mathsf{GT})}}{\mathsf{B}^{3}} + \frac{\mathfrak{p}}{\mathsf{B}} \right) + 2 \left( \frac{T^{01(\mathsf{GT})}}{\mathsf{A}^{2}\mathsf{B}} - \frac{\zeta}{\mathsf{A}} \right) \\ - \frac{1}{2} \frac{\left( 3\mathfrak{p} + \mathfrak{u} \right)^{\prime}}{\mathsf{B}} + \left( \frac{-2\mathfrak{p}_{r}}{\mathsf{B}} - \frac{\mathfrak{s}^{2}}{4\pi\mathsf{C}^{4}\mathsf{B}} - \frac{2T^{11(\mathsf{GT})}}{\mathsf{B}^{3}} + \frac{\mathfrak{p}}{\mathsf{B}} \right)^{\prime} + \left( \frac{-\zeta}{\mathsf{A}} + \frac{T^{01(\mathsf{GT})}}{\mathsf{A}^{2}\mathsf{B}} \right) \\ \times \left( \ln f_{T} \right)^{\cdot} \right].$$
(A6)

# Appendix B

In scalar functions, the extra curvature terms are described as follows

$$\begin{split} \mathsf{M}^{(\mathsf{GT})} &= 2 \left[ R_{\mu\tau} R^{\mu}_{\vartheta} f_G + R^{\mu\nu} R_{\mu\tau\nu\vartheta} f_G - \frac{1}{2} R R_{\vartheta\tau} f_G - \frac{1}{2} R_{\tau\mu\nun} R^{\mu\nun}_{\vartheta} f_G \right. \\ &+ \frac{1}{2} R \nabla_{\vartheta} \nabla_{\tau} f_G + R_{\vartheta\tau} \Box f_G - R^{\mu}_{\vartheta} \nabla_{\tau} \nabla_{\mu} f_G - R^{\mu\nu} \nabla_{\vartheta} \nabla_{\mu} f_G \\ &- R_{\mu\tau\nu\vartheta} \nabla^{\mu} \nabla^{\nu} f_G \right] g^{\vartheta\tau} + 2 \left[ -R_{\mu\delta} R^{\mu}_{\vartheta} f_G - R^{\mu\nu} R_{\mu\delta\nu\vartheta} f_G + \frac{1}{2} R R_{\vartheta\delta} f_G \right. \\ &+ \frac{1}{2} R_{\delta\mu\nun} R^{\mu\nun}_{\vartheta} f_G - R_{\delta\vartheta} \Box f_G + R_{\mu\delta\nu\vartheta} \nabla^{\nu} \nabla^{\mu} f_G + R^{\mu}_{\vartheta} \nabla_{\mu} \nabla_{\delta} f_G \\ &+ R^{\mu}_{\delta} \nabla_{\vartheta} \nabla_{\mu} f_G - \frac{1}{2} R \nabla_{\vartheta} \nabla_{\delta} f_G \right] v_{\tau} v^{\delta} g^{\vartheta\tau} + 2 \left[ -R_{\mu\tau} R^{\mu\gamma} f_G \right. \\ &+ \frac{1}{2} R_{\tau\mu\nun} R^{\mu\nun\gamma} f_G - \frac{1}{2} R R^{\gamma}_{\tau} f_G R^{\mu\nu} R^{\gamma}_{\mu\tau\nu} f_G + R^{\mu\gamma} \nabla_{\tau} \nabla_{\mu} f_G \\ &- \frac{1}{2} R \nabla^{\gamma} \nabla_{\tau} f_G + R^{\gamma}_{\mu\tau\nu} \nabla^{\nu} \nabla^{\mu} f_G - R^{\gamma}_{\tau} \Box f_G + R^{\mu}_{\tau} \nabla_{\mu} \nabla^{\gamma} f_G \right] v_{\vartheta} v_{\gamma} g^{\vartheta\tau} \\ &+ 2 \left[ R^{\mu\gamma} R_{\mu\delta} f_G + R^{\mu\nu} R^{\gamma}_{\mu\delta\nu} f_G + \frac{1}{2} R \nabla_{\delta} \nabla^{\gamma} f_G - \frac{1}{2} R^{\mu\nun\gamma} R_{\delta\mu\nun} f_G \\ &+ R^{\gamma}_{\delta} \Box f_G - R^{\mu}_{\delta} \nabla_{\mu} \nabla^{\gamma} f_G - R^{\mu\gamma} \nabla_{\mu} \nabla_{\delta} f_G - R^{\gamma}_{\mu\delta\nu} \nabla^{\nu} \nabla^{\mu} f_G - \frac{1}{2} R R^{\gamma}_{\delta} f_G \right] \\ &\times g_{\vartheta\tau} v_{\gamma} v^{\delta} g^{\vartheta\tau} - \left[ 4 R_{\mu m} R^{\mu m} f_G - 2 R^2 f_G + 4 R^{m}_{\mu m\nu} R^{\mu\nu} f_G \\ &- 2 R^{\mu\nu n}_{\mu m\nu} \nabla^{\nu} \nabla^{\mu} f_G - 4 R^{\nu l} \nabla_{\nu} \nabla_{l} f_G \right] - 6 R \Box f_G - \frac{1}{2} f + 12 R^{\mu\nu} \nabla_{\mu} \nabla_{\nu} f_G, \end{aligned}$$

$$\begin{aligned} \mathsf{J}_{(\vartheta\tau)}^{(\mathsf{GT})} &= \left[ 2R_{\mu d}R_{c}^{\mu}f_{G} + 2R^{\mu\nu}R_{\mu d\nu c}f_{G} - RR_{cd}f_{G} - R_{d\mu\nu n}R_{c}^{\mu\nu n}f_{G} + 2R_{cd}\Box f_{G} \right. \\ &+ R\nabla_{d}\nabla_{c}f_{G} - 2R_{d}^{\mu}\nabla_{\mu}\nabla_{c}f_{G} - 2R_{c}^{\mu}\nabla_{\mu}\nabla_{d}f_{G} - 2R_{\mu d\nu c}\nabla^{\mu}\nabla^{\nu}f_{G} \right]h_{\vartheta}^{c}h_{\tau}^{d} \\ &+ 2\left[ R^{\mu\gamma}R_{\mu\delta}f_{G} + R_{\mu\delta\nu}^{\gamma}R^{\mu\nu}f_{G} - \frac{1}{2}RR_{\delta}^{\gamma}f_{G} - \frac{1}{2}R_{\delta\mu\nu n}R^{\mu\nu n\gamma}f_{G} \right. \\ &+ R_{\delta}^{\gamma}\Box f_{G} - R^{\mu\gamma}\nabla_{\mu}\nabla_{\delta}f_{G} + \frac{1}{2}R\nabla_{\delta}\nabla^{\gamma}f_{G} - R_{\delta}^{\mu}\nabla^{\gamma}\nabla_{\mu}f_{G} \\ &- R_{\mu\delta\nu}^{\gamma}\nabla^{\mu}\nabla^{\nu}f_{G} \right]h_{\vartheta\tau}v_{\gamma}v^{\delta} - 2\left[ R_{\mu\tau}R_{\vartheta}^{\mu}f_{G} + R^{\mu\nu}R_{\mu\tau\nu\vartheta}f_{G} - \frac{1}{2}RR_{\vartheta\tau}f_{G} \right] \end{aligned}$$

$$- \frac{1}{2}R_{\tau\mu\nu n}R_{\vartheta}^{\mu\nu n}f_{G} + R_{\vartheta\tau}\Box f_{G} + \frac{1}{2}R\nabla_{\vartheta}\nabla_{\tau}f_{G} - R_{\vartheta}^{\mu}\nabla_{\tau}\nabla_{\mu}f_{G} - R_{\tau}^{\mu}\nabla_{\vartheta}\nabla_{\mu}f_{G} - R_{\mu\tau\nu\vartheta}\nabla^{\mu}\nabla^{\nu}f_{G}] - 2\left[-R_{\mu\delta}R_{\vartheta}^{\mu}f_{G} - R^{\mu\nu}R_{\mu\delta\nu\vartheta}f_{G} + \frac{1}{2}RR_{\vartheta\delta}f_{G} + \frac{1}{2}R_{\delta\mu\nu n}R_{\vartheta}^{\mu\nu n}f_{G} - R_{\delta\vartheta}\Box f_{G} + R_{\mu\delta\nu\vartheta}\nabla^{\mu}\nabla^{\nu}f_{G} - \frac{1}{2}R\nabla_{\vartheta}\nabla_{\delta}f_{G} + R_{\vartheta}^{\mu}\nabla_{\delta}\nabla_{\mu}f_{G} + R_{\delta}^{\mu}\nabla_{\vartheta}\nabla_{\mu}f_{G}\right]v_{\tau}v^{\delta} - 2\left[-R_{\mu\tau}R^{\mu\gamma}f_{G} - R^{\mu\nu}R_{\mu\tau\nu}^{\gamma}f_{G} + \frac{1}{2}RR_{\tau}^{\gamma}f_{G} + \frac{1}{2}R_{\tau\mu\nu n}R^{\mu\nu n\gamma}f_{G} - R_{\tau}^{\gamma}\Box f_{G} + R^{\mu\gamma}\nabla_{\mu}\nabla_{\tau}f_{G} + R_{\tau}^{\mu}\nabla_{\mu}\nabla^{\gamma}f_{G} - \frac{1}{2}R\nabla^{\gamma}\nabla_{\tau}f_{G} + R_{\mu\tau\nu}^{\gamma}\nabla^{\mu}\nabla^{\nu}f_{G}\right]v_{\vartheta}v_{\gamma} - 2\left[R_{\mu\delta\nu}^{\gamma}R^{\mu\nu}f_{G} + R_{\mu\delta}R^{\mu\gamma}f_{G} - \frac{1}{2}RR_{\delta}^{\gamma}f_{G} - \frac{1}{2}R^{\mu\nu n\gamma}R_{\delta\mu\nu n}f_{G} + R_{\delta}^{\gamma}\Box f_{G} + \frac{1}{2}R\nabla_{\delta}\nabla^{\gamma}f_{G} - R_{\delta}^{\mu}\nabla_{\mu}\nabla^{\gamma}f_{G} - R^{\mu\gamma}\nabla_{\mu}\nabla_{\delta}f_{G} - R_{\mu\delta\nu}^{\gamma}\nabla^{\nu}\nabla^{\mu}f_{G}\right]v_{\gamma}v^{\delta}g_{\vartheta\tau},$$

$$\begin{aligned} \mathsf{Q}^{(\mathsf{GT})} &= \left[ \frac{1}{2} R_{\mu\epsilon} R^{\mu\rho} f_G + \frac{1}{2} R^{\mu\nu} R^p_{\mu\epsilon\nu} f_G - \frac{1}{4} R R^p_{\epsilon} f_G - \frac{1}{4} R_{\epsilon\mu\nun} R^{\mu\nunp} f_G \right. \\ &+ \left. \frac{1}{2} R^p_{\epsilon} \Box f_G + \frac{1}{4} R \nabla^p \nabla_{\epsilon} f_G - \frac{1}{4} R^{\mu\rho} \nabla_{\epsilon} \nabla_{\mu} f_G - \frac{1}{2} R^p_{\epsilon} \nabla^p \nabla_{\mu} f_G \right. \\ &- \left. \frac{1}{2} R^p_{\mu\epsilon\nu} \nabla^{\mu} \nabla^{\nu} f_G \right] g^{\vartheta\tau} \epsilon_{p\delta\tau} \epsilon^{\epsilon\delta}_{\vartheta} + \left[ -\frac{1}{2} R_{\mu\delta} R^{\mu\rho} f_G - \frac{1}{2} R^{\mu\nu} R^p_{\mu\delta\nu} f_G \right. \\ &+ \left. \frac{1}{4} R^p_{\delta} R f_G - \frac{1}{2} R^p_{\delta} \Box f_G + \frac{1}{4} R^{\mu\nunp} R_{\delta\mu\nun} f_G - \frac{1}{4} R \nabla^p \nabla_{\delta} f_G \right. \\ &+ \left. \frac{1}{2} R^p_{\delta} \nabla^p \nabla_{\mu} f_G + \frac{1}{4} R^{\mu\rho} \nabla_{\delta} \nabla_{\mu} f_G + \frac{1}{2} R^p_{\mu\delta\nu} \nabla^{\mu} \nabla^{\nu} f_G \right] g^{\vartheta\tau} \epsilon_{\rho\epsilon\tau} \epsilon^{\epsilon\delta}_{\vartheta} \\ &+ \left[ -\frac{1}{2} R_{\mu\epsilon} R^{\mu\gamma} f_G - \frac{1}{2} R^{\mu\nu} R^{\gamma}_{\mu\epsilon\nu} f_G + \frac{1}{4} R R^{\epsilon}_{\epsilon} f_G + \frac{1}{4} R_{\epsilon\mu\nun} R^{\mu\nun\gamma} f_G \right. \\ &- \left. \frac{1}{2} R^{\gamma}_{\epsilon} \Box f_G + \frac{1}{4} R^{\mu\gamma} \nabla_{\epsilon} \nabla_{\mu} f_G - \frac{1}{4} R \nabla^{\gamma} \nabla_{\epsilon} f_G + \frac{1}{2} R^p_{\epsilon} \nabla_{\mu} \nabla^{\gamma} f_G \right. \\ &+ \left. \frac{1}{2} R^{\gamma}_{\mu\epsilon\nu} \nabla^{\nu} \nabla^{\mu} f_G \right] g^{\vartheta\tau} \epsilon_{\delta\gamma\tau} \epsilon^{\epsilon\delta}_{\vartheta} + \left[ \frac{1}{2} R^{\mu\nu} R^{\gamma}_{\mu\delta\nu} f_G + \frac{1}{2} R^{\mu\gamma}_{\epsilon} R_{\mu\delta} f_G \right. \\ &- \left. \frac{1}{4} R R^{\gamma}_{\delta} f_G + \frac{1}{2} R^{\gamma}_{\delta} \Box f_G - \frac{1}{4} R^{\mu\nun\gamma}_{\epsilon} R_{\delta\mu\nun} f_G + \frac{1}{4} R \nabla_{\delta} \nabla^{\gamma} f_G \right. \end{aligned}$$

$$- \frac{1}{4}R^{\mu\gamma}\nabla_{\delta}\nabla_{\mu}f_{G} - \frac{1}{2}R^{\gamma}_{\mu\delta\nu}\nabla^{\mu}\nabla^{\nu}f_{G} - \frac{1}{2}R^{\mu}_{\delta}\nabla^{\gamma}\nabla_{\mu}f_{G} \right]g^{\vartheta\tau}\epsilon_{\epsilon\gamma\tau}\epsilon^{\epsilon\delta}_{\vartheta}$$

$$- 12R^{\mu\nu}\nabla_{\nu}\nabla_{\mu}f_{G} + 6R\Box f_{G} + [4R_{\mu\nu}R^{\mu\nu}f_{G} + (\mu + \mathfrak{p})f_{T}$$

$$+ 4R^{\mu\nu}R^{m}_{\mu\num}f_{G} - 2R^{l}_{\mu\nun}R^{\mu\nun}_{l}f_{G} - 2R^{2}f_{G} - 4R\Box f_{G}$$

$$+ 16R^{\mu\nu}\nabla_{\nu}\nabla_{\mu}f_{G} - 4R^{\mu l}\nabla_{l}\nabla_{\mu}f_{G} - 4R^{\mu m}\nabla_{\mu}\nabla_{m}f_{G}$$

$$- 4R^{m}_{\mu m\nu}\nabla^{\nu}\nabla^{\mu}f_{G} \right] + \frac{f}{2}.$$

Data Availability Statement: This manuscript has no associated data.

### References

- L. Bergström, Non-baryonic dark matter: observational evidence and detection methods, Rep. Prog. Phys. 63 (2000) 793.
- [2] D. Pietrobon, A. Balbi, D. Marinucci, Integrated Sachs-Wolfe effect from the cross correlation of WMAP 3 year and the NRAO VLA sky survey data: New results and constraints on dark energy, Phys. Rev. D 74 (2006) 043524.
- [3] S. Nojiri, S.D. Odintsov, Modified Gauss-Bonnet theory as gravitational alternative for dark energy, Phys. Lett. B 631 (2005) 1-6.
- [4] K. Bamba, A.N. Makarenko, A.N. Myagky, S.D. Odintsov, Bouncing cosmology in modified Gauss-Bonnet gravity, Phys. Lett. B 732 (2014) 349-55.
- [5] G. Abbas, D. Momeni, M.A. Ali, R. Myrzakulov, S. Qaisar, Anisotropic compact stars in f(G) gravity, Astrophys. Space Sci. 357 (2015) 1-11.
- [6] M. Sharif, A. Ikram, Inflationary dynamics in f(G) gravity, Int. J. Mod. Phys. D 26 (2017) 1750030.
- [7] M. Sharif, A. Ramzan, Anisotropic compact stellar objects in modified Gauss-Bonnet gravity, Phys. Dark Universe 30 (2020) 100737.
- [8] M. Sharif, A. Ikram, Energy conditions in f(G, T) gravity, Eur. Phys. J. C 76 (2016) 1-13.

- [9] M. Sharif, A. Ikram, Anisotropic perturbations and stability of a static universe in f(G, T) gravity, Eur. Phys. J. Plus 132 (2017) 1-9.
- [10] H. Hossienkhani, V. Fayaz, A. Jafari, Energy conditions and modified gravity in anisotropic universe, Can. J. Phys. 96 (2018) 225-232.
- [11] M. Sharif, A. Naeem, Anisotropic solution for compact objects in f(G,T) gravity, Int. J. Mod. Phys. A 35 (2020) 2050121.
- [12] M.F. Shamir, Bouncing universe in f(G, T) gravity, Phys. Dark Universe 32 (2021) 100794.
- [13] B.V. Ivanov, Rigidly rotating cylinders of charged dust, Class. Quantum Grav. 19 (2002) 5131.
- [14] M. Esculpi, E. Aloma, Conformal anisotropic relativistic charged fluid spheres with a linear equation of state, Eur. Phys. J. C 7 (2010) 521-532.
- [15] M. Sharif, S. Naz, Charged cylindrical collapse in f(G) gravity, Mod. Phys. Lett. A 35 (2020) 1950340.
- [16] M. Sharif, K. Hassan, Influence of charge on decoupled anisotropic spheres in f(G, T) gravity, Eur. Phys. J. Plus 137 (2022) 997.
- [17] M. Sharif, K. Hassan, Anisotropic decoupled spheres in f(G, T) gravity, Int. J. Geom. Methods Mod. Phys. 19 (2022) 2250150.
- [18] M. Sharif, K. Hassan, Charged Anisotropic Solutions through Decoupling in f(G,T) Gravity, Int. J. Geom. Methods Mod. Phys (doi: 10.1142/S0219887823501001).
- [19] R. López-Ruiz, H.L. Mancini, X. Calbet, A statistical measure of complexity, Phys. Lett. A 209 (1995) 321-326.
- [20] X. Calbet, R. López-Ruiz, Tendency towards maximum complexity in a nonequilibrium isolated system, Phys. Rev. E 63 (2001) 066116.
- [21] R.G. Catalán, J. Garay, R. López-Ruiz, Features of the extension of a statistical measure of complexity to continuous systems, Phys. Rev. E 66 (2002) 011102.

- [22] L. Herrera, New definition of complexity for self-gravitating fluid distributions: The spherically symmetric, static case, Phys. Rev. D 97 (2018) 044010.
- [23] M. Sharif, I.I. Butt, Complexity factor for charged spherical system, Eur. Phys. J. C 78 (2018) 688.
- [24] L. Herrera, A. Di Prisco, J. Ospino, Definition of complexity for dynamical spherically symmetric dissipative self-gravitating fluid distributions, Phys. Rev. D 98 (2018) 104059.
- [25] L. Herrera, A. Di Prisco, J. Ospino, Complexity factors for axially symmetric static sources, Phys. Rev. D 99 (2019) 044049.
- [26] M. Sharif, I.I. Butt, Complexity factor for static cylindrical system, Eur. Phys. J. C 78 (2018) 850.
- [27] M. Sharif, A. Majid, M.M.M. Nasir, Complexity factor for selfgravitating system in modified Gauss-Bonnet gravity Int. J. Mod. Phys. A 34 (2019) 1950210.
- [28] Z. Yousaf, M.Z. Bhatti, K. Hassan, Complexity for self-gravitating fluid distributions in f(G, T) gravity, Eur. Phys. J. Plus 135 (2020) 397.
- [29] Z. Yousaf, K. Bamba, M.Z. Bhatti and K. Hassan, Measure of complexity in self-gravitating systems using structure scalars, New Astron. 84 (2021) 101541.
- [30] M. Sharif, K. Hassan, Complexity factor for static cylindrical objects in f(G,T) gravity, Pramana 96 (2022) 50.
- [31] M. Sharif, K. Hassan, Electromagnetic effects on the complexity of static cylindrical object in f(G, T) gravity, Eur. Phys. J. Plus 137 (2022) 1380.
- [32] M. Sharif, K. Hassan, Complexity of dynamical cylindrical system in f(G,T) gravity, Mod. Phys. Lett. A 37 (2022) 2250027.
- [33] M. Sharif, K. Hassan, Complexity for dynamical anisotropic sphere in f(G,T) gravity, Chin. J. Phys 77 (2022) 1479-1492.
- [34] G. Abbas, R. Ahmed, Complexity factor for a class of compact stars in f(R,T) gravity, Astrophys. Space Sci. 364 (2019) 1-9.

- [35] M. Sharif, A. Majid, Complexity factor for static sphere in selfinteracting Brans-Dicke gravity, Chin. J. Phys. 61 (2019) 38-46.
- [36] M. Sharif, A. Majid, Complexity of dynamical sphere in self-interacting Brans-Dicke gravity, Eur. Phys. J. C 80 (2020) 1-9.
- [37] M. Sharif, A. Majid, Complexity factor for cylindrical system in Brans-Dicke gravity, Indian J. Phys. 95 (2021) 769-777.
- [38] M. Zubair, H. Azmat, Complexity analysis of cylindrically symmetric self-gravitating dynamical system in f(R,T) theory of gravity, Phys. Dark Universe 28 (2020) 100531.
- [39] M. Sharif, T. Naseer, Complexity analysis of charged dynamical dissipative cylindrical structure in modified gravity, Eur. Phys. J. Plus 137 (2022) 1-16.
- [40] M. Sharif, T. Naseer, Complexity of a dynamical dissipative cylindrical system in non-minimally coupled theory, Chin. J. Phys. 77 (2022) 2655-2667.
- [41] C.W. Misner, D.H. Sharp, Relativistic equations for adiabatic, spherically symmetric gravitational collapse, Phys. Rev. 136 (1964) B571.
- [42] L. Bel, Electromagnetic and gravitational inductions, Annales de l'institut Henri Poincaré 17 (1961) 37-57.
- [43] L. Herrera, J Ospino, A. Di Prisco, E. Fuenmayor, O. Troconis, Structure and evolution of self-gravitating objects and the orthogonal splitting of the Riemann tensor, Phys.Rev. D. 79 (2009) 064025.
- [44] R. Kippenhahn, A. Weigert, A. Weiss, Stellar Structure and Evolution, Springer, 1990.
- [45] C.J. Hansen, S.D. Kawaler, V. Trimble, Stellar Interiors: Physical Principles, Structure and Evolution, Springer, New York, 1994.
- [46] M.F. Shamir, M. Ahmad, Stellar hydrostatic equilibrium compact structures in f(G, T) gravity, Mod. Phys. Lett. A 34 (2019) 1950038.
- [47] M.F. Shamir, N. Uzair, Spherically symmetric spacetime with radiating fluids in f(G, T) gravity, Mod. Phys. Lett. A 34 (2019)1950215.