

Width of Convex Bodies in Hyperbolic Space

Marek Lassak

Abstract. For every hyperplane H supporting a convex body C in the hyperbolic space \mathbb{H}^d we define the width of C determined by H as the distance between H and a most distant ultraparallel hyperplane supporting C . We define bodies of constant width in \mathbb{H}^d in the standard way as bodies whose all widths are equal. We show that every body of constant width is strictly convex. The minimum width of C over all supporting H is called the thickness $\Delta(C)$ of C . A convex body $R \subset \mathbb{H}^d$ is said to be reduced if $\Delta(Z) < \Delta(R)$ for every convex body Z properly contained in R . We show that regular tetrahedra in \mathbb{H}^3 are not reduced. Similarly as in the Euclidean and spherical spaces, we introduce complete bodies and bodies of constant diameter also in \mathbb{H}^d . We show that every body of constant width δ is a body of constant diameter δ and a complete body of diameter δ . Moreover, the two last conditions are equivalent.

Mathematical Subject Classification (2010). 52A55.

Keywords. Hyperbolic geometry, convex body, width, constant width, thickness, diameter, reduced body, complete body.

1 Introduction

Here we present a sketch of the content of this paper.

In Section 2 we introduce some notions and present four lemmas.

Section 3 is devoted to the notion of width of a convex body C in the hyperbolic space \mathbb{H}^d . For any hyperplane H supporting C we define $\text{width}_H(C)$ of C determined by H as the distance between H and any farthest ultraparallel hyperplane supporting C . Proposition 1 shows that $\text{width}_H(C)$ equals to the maximum distance between H and a point of C , and Proposition 2 claims that $\text{width}_H(C)$ equals to the distance between H and the nearest equidistant surface E to H such that C is a subset of the equidistant strip bounded by H and E .

Section 4 is about the diameter and thickness. Theorem 1 says that the maximum width of C equals to the diameter of C . The thickness $\Delta(C)$ of a convex body $C \in \mathbb{H}^d$ is defined as the infimum of $\text{width}_H(C)$ over all hyperplanes H supporting C . In Theorem 2 we assume that $\text{width}_H(C) = \Delta(C)$ for a hyperplane H supporting C and that there exists exactly one farthest point j of C from H . The thesis is that the projection of j onto H belongs to $C \cap H$. Finally, we find a formula for the width of the regular simplex determined

by the hyperplane containing its facet.

Section 5 concerns bodies of constant width and reduced bodies. We define the notion of a body of constant width as a body whose all widths are equal. In Proposition 3 we show that every body of constant width is strictly convex. By a reduced body we mean a convex body $R \subset \mathbb{H}^d$ such that $\Delta(Z) < \Delta(R)$ for every convex body $Z \subset R$ different from R . Proposition 4 says that every body of constant width is a reduced body. In Remark 2 we explain why every regular tetrahedron in \mathbb{H}^3 is not reduced.

In Section 6 we consider complete bodies and bodies of constant diameter in \mathbb{H}^d . Theorem 3 says that every body of constant width δ is a body of constant diameter δ , and that a body is of constant diameter δ if and only if it is a complete body of diameter δ . We conjecture that the three kinds of bodies in \mathbb{H}^d coincide.

In our considerations it is convenient to work with the hyperboloid model of \mathbb{H}^d , so about the model on the upper sheet $x_{d+1} = \sqrt{x_1^2 + \dots + x_d^2 + 1}$ of the two-sheeted hyperboloid. This model permits a reasonable comparison of the obtained results with the analogous ones in the Euclidean space \mathbb{E}^d and the spherical space \mathbb{S}^d . Our figures show the orthogonal look to the sheet from the above.

2 A few notions and lemmas

The geodesic between two different points $a, b \in \mathbb{H}^d$ is called the **segment** ab . By a **ball of radius** ρ we mean the set of points of \mathbb{H}^d which are at a distance at most ρ from a fixed point called the **center** of this ball. If $d = 2$, the ball is called a **disk**, and its boundary a **circle**. We say that a set $C \subset \mathbb{H}^d$ is **convex** if together with every two points a, b it contains the whole segment ab . By a **convex body** in \mathbb{H}^d we mean a closed bounded convex set with non-empty interior. Of course, the intersection of any family of convex sets is also convex. Thus for every set $Q \subset \mathbb{H}^d$ there exists the unique smallest convex set containing Q . It is called **the convex hull of** Q .

If a hyperplane G of \mathbb{H}^d has a common point with a convex body $C \subset \mathbb{H}^d$ and if its intersection with the interior of C is empty, we say that G **supports** C . If at every boundary point of C exactly one hyperplane supports C , the body is said to be **smooth**.

Let $A, B \subset \mathbb{H}^d$ be two non-empty sets. The symbol $\text{dist}(A, B)$ denotes the infimum of distances $|ab|$ over all $a \in A$ and $b \in B$. We call it the distance between A and B . If A is a one point set $\{c\}$, we simply write $\text{dist}(c, B)$.

We omit an easy proof of the following lemma.

Lemma 1. *For the projection h of a point g onto a hyperplane H we have $\text{dist}(g, H) = |gh|$. Moreover, $|gk| > |gh|$ for every $k \in H$ different from h .*

Proof. There are $h \in H$ and $j \in J$ such that hj is orthogonal to H and J . Clearly, $\text{dist}(H, J) = |hj|$.

Imagine that $J \cap C$ has more than one point. Then there is at least one point $f \in J \cap C$ different from j (see Fig. 2). Denote by e the projection of f onto H . Clearly, $|ef| > |hj|$.

Let F be the hyperplane through f orthogonal to the segment ef . By Lemma 1 we have $\text{dist}(F, H) = |ef| > |hj| = \text{dist}(J, H)$.

If F supports C , then it is a farthest ultraparallel hyperplane to H supporting C than J . This contradicts the assumption that J is a farthest ultraparallel hyperplane to H supporting C .

If F does not support C , then F passes through the interior of C . There exists a supporting hyperplane G of C orthogonal to the straight line containing ef such that F is a subset of the interior of the strip between H and G . By g denote the point of intersection of G with this line.

Of course, $g \neq f$. Thus by $f \in eg$ we obtain $|ef| < |eg|$. So $\text{dist}(G, H) = |ge|$ and $\text{dist}(F, H) = |fe|$ imply $\text{dist}(G, H) > \text{dist}(F, H)$. Thus from the earlier established inequality $\text{dist}(F, H) > \text{dist}(J, H)$ we get $\text{dist}(G, H) > \text{dist}(J, H)$. This contradicts the description of J as the most distant ultraparallel to H hyperplane supporting C . The obtained contradiction implies that $J \cap C$ is a one-point set. \square

Assume that a hyperplane H supports a convex body $C \in \mathbb{H}^d$ and J is a farthest from H ultraparallel hyperplane supporting C . Then a most distant from J ultraparallel hyperplane K supporting C does not have to be H (differently than in \mathbb{E}^d). A simple example showing this is the regular simplex and the hyperplane H containing a facet of it.

Let p be a point out of a hyperplane H . Recall that the **equidistant surface to H through p** is the set of all points in the distance $\text{dist}(p, H)$ from H which are in the half-space bounded by H and containing p .

3 Width of a convex body

Let H be a hyperplane supporting a convex body $C \subset \mathbb{H}^d$. We define the **width of C determined by H** as the distance between H and any farthest ultraparallel hyperplane supporting C (by compactness arguments there exists at least one such a most distant one, sometimes there are a finite or even infinitely many of them). The symbol $\text{width}_H(C)$ denotes this width of C determined by H .

Proposition 1. *Let $C \subset \mathbb{H}^d$ be a convex body and H be any supporting hyperplane of C . Then $\text{width}_H(C)$ equals to the maximum distance between H and a point of C .*

Proof. Since C is compact, there exists a point $j \in C$ in the maximum distance from H . Denote by h the projection of j onto H . Provide the hyperplane J through j which is orthogonal to hj . Clearly, H and J are ultraparallel. By Lemma 3 we conclude that J supports C at j (see Fig. 3).




Fig. 3. Illustration to the proof of Proposition 1

Let us show that J is a farthest ultraparallel hyperplane to H supporting C . Imagine the opposite. Then there is a more distant from H ultraparallel hyperplane J' supporting C . Denote by j' its closest to H point of support. Its projection onto H is denoted by h' . By $\text{dist}(H, J') > \text{dist}(H, J)$ we have $|h'j'| > |hj|$. This contradicts the choice of j .

We have shown that J is a farthest ultraparallel hyperplane to H supporting C . Thus by the definition of $\text{width}_H(C)$ we have $\text{width}_H(C) = \text{dist}(H, J)$. Since $h \in H$, $j \in J$ and hj is orthogonal to both these ultraparallel hyperplanes, we have $\text{dist}(H, J) = \text{dist}(H, j)$. Hence $\text{width}_H(C) = \text{dist}(H, j)$, which is our thesis. \square

In connection with this proposition, pay attention that in all the geometries \mathbb{E}^d , \mathbb{S}^d and \mathbb{H}^d , we may define the $\text{width}_H(C)$ as the maximum distance between H and a point of C . Recall here that $\text{width}_H(C)$ in \mathbb{E}^d is the width of C in the direction orthogonal to H , and $\text{width}_H(C)$ for \mathbb{S}^d is defined in [10].

We omit an easy proof of the following proposition.

Proposition 2. *Let $C \subset \mathbb{H}^d$ be a convex body and H any supporting hyperplane of C . Then $\text{width}_H(C)$ equals to the distance between H and the nearest equidistant surface E to H such that C is a subset of the equidistant strip being the convex hull of $H \cup E$.*

Each of the above propositions may be regarded as a different definition of $\text{width}_H(C)$.

Let us add that a few different notions of width of a convex body $C \subset \mathbb{H}^d$ are known. For a comparison of them see the paper [7] by Horváth.

4 Diameter and thickness of a convex body

We put the material on the diameter and the thickness in one section since they are of analogous nature. Just we see this from the below Theorem 1 characterizing the diameter of a convex body $C \subset \mathbb{H}^d$ and the below definition of the thickness of C : they are the maximum and the minimum values of $\text{width}_H(C)$ over all hyperplanes H supporting C .

By the **diameter** $\text{diam}(A)$ of a set $A \subset \mathbb{H}^d$ we mean the supremum of the distances between pairs of points of A . Clearly, if A a convex body, by its compactness, $\text{diam}(A)$ is realized for at least one pair of points of A .

Theorem 1. *For every convex body $C \subset \mathbb{H}^d$ we have*

$$\max\{\text{width}_H(C); H \text{ is a supporting hyperplane of } C\} = \text{diam}(C).$$

Proof. Denote the left side of this equality by $\max \text{width}(C)$.

By the compactness of C there exists a supporting hyperplane K of C such that $\text{width}_K(C) = \max \text{width}(C)$. By Proposition 1 the value $\text{width}_K(C)$ equals to the maximum distance of a point $c \in C$ from K . For the projection k of c onto K we have $|kc| = \text{width}_K(C)$. Since K supports C , there exists a point $f \in K \cap C$. Of course $|cf| \geq |ck|$. Hence $\text{diam}(C) \geq \max \text{width}(C)$.

There are h, j in $\text{bd}(C)$ such that $|hj| = \text{diam}(C)$. Provide the hyperplane H through h orthogonal to hj . Observe that H supports C (still if not, then there exists a point $y \in C$ which is in the strictly opposite side of H than j , which means that $|yj| > |hj|$ in contradiction to $|hj| = \text{diam}(C)$). Since H supports C and the point j is in C , by Proposition 1 we get $\text{width}_H(C) \geq |hj|$. Hence $\max \text{width}(C) \geq |hj|$. This and $|hj| = \text{diam}(C)$ lead to $\text{diam}(C) \leq \max \text{width}(C)$.

From the above two paragraphs we obtain the thesis of our theorem. \square

Claim 1. *Assume that $\text{diam}(C) = |ab|$ for some points a, b of a convex body $C \subset \mathbb{H}^d$. Denote by S the ultraparallel strip whose both bounding hyperplanes are orthogonal to the segment ab . We have $C \subset S$.*

Proof. Denote by A the hyperplane through a bounding S and by B the hyperplane through b bounding S . Imagine that $C \not\subset S$. Then a point $c \in C$ is out of S . It is strictly separated from S by A or B . For instance, let it be separated by A . Denote by a' the intersection of bc with A . Of course, $a' \in C$. From the right triangle baa' we conclude that $|a'b| > |ab|$. This contradicts $\text{diam}(C) = |ab|$. Consequently, $C \subset S$. \square

By the **thickness** $\Delta(C)$ of a convex body $C \subset \mathbb{H}^d$ we mean the infimum of $\text{width}_H(C)$ over all hyperplanes H supporting C . By compactness arguments, this infimum is realized, so $\Delta(C)$ is the minimum of the numbers $\text{width}_H(C)$.

Remark 1. In \mathbb{E}^d the following statement holds true. *Let C be a convex body. Let H be a supporting hyperplane of C such that $\text{width}_H(C) = \Delta(C)$. Denote by j any farthest point of C from H . Then the projection of j onto H belongs to $H \cap C$.* It follows from the property formulated by Eggleston at the bottom of page 77 of [4]. An analogous statement holds true in \mathbb{S}^d as a consequence of Claim 2 of [10] (see also Corollary 2.13 of [12]). Submitting the first version of this paper, the author expected that such a statement holds also in \mathbb{H}^d . But it is not true as it results from the following example shown by the referee.

Example 1. Let H be a line and E be an equidistant curve to H in a distance ρ . Take a point $a \in H$ and the orthogonal line ℓ to H through a . Let b and c be points of E symmetric with respect to ℓ such that the angle $\angle bac$ is right or obtuse. Define C as the convex hull of the set consisting of the point a and all the points of E between b and c . Clearly, $\text{width}_H(C) = \rho = \Delta(C)$. The projections of all the points of E between bc besides its middle onto H do not belong to $H \cap C$.

Here is a weaker form of the (not true in \mathbb{H}^d) statement from Remark 1.

Theorem 2. *Let $C \subset \mathbb{H}^d$ be a convex body and let H be a supporting hyperplane of C such that $\text{width}_H(C) = \Delta(C)$. Assume that there exists a unique most distant point $j \in C$ from H . Then the projection of j onto H belongs to $H \cap C$.*

Proof. Imagine the opposite to the thesis, this is that the projection h of j onto H does not belong to $H \cap C$.

Since $H \cap C$ is convex and compact, there exists exactly one point $g \in H \cap C$ in the minimum distance from h . The above uniqueness of g results from the fact that g is the intersection of the $(d-1)$ -dimensional ball $B \subset H$ centered at h which touches $H \cap C$. We do not lose the generality assuming that the vector hg is directed into the positive orientation on the straight line containing the segment gh .

Denote by m the midpoint of gh . Take the $(d-2)$ -dimensional hyperplane M of H through m orthogonal to gh .

Provide the hyperplane G orthogonal to gm through g . Let G^+ be the closed halfspace of \mathbb{H}^d bounded by G which contains j , and let G^- be the opposite closed halfspace.

Since j is the unique point of C in the maximum distance from H and by the compactness of $C \cap G^-$ there is a positive number $\kappa^- < \Delta(C)$ for which every point of $C \cap G^-$ is in a distance at most κ^- from H . Hence we can rotate H around M , in the positive orientation, up to a position H' for which there is positive number λ^- strictly between κ^- and $\Delta(C)$ such that every point of $C \cap G^-$ is in a distance at most λ^- from H' .

Clearly, our H' can be simultaneously chosen so close to H that it has empty intersection with $C \cap G^+$, and thus with C . Hence there exists a λ^+ such that every point of

$C \cap G^+$ is in a distance at most λ^+ .

From the two preceding paragraphs we conclude that every point of C is in a distance at most $\lambda = \max\{\lambda^-, \lambda^+\}$ from H' . Since $\lambda < \text{width}_H(C)$, by Proposition 1 applied to H' we conclude that $\text{width}_{H'}(C) < \text{width}_H(C)$. This contradicts the assumption that $\text{width}_H(C) = \Delta(C)$. Hence h must be in $C \cap H$. \square

Example 2. In order to find the height η_x of the regular triangle T of side $2x$ we take a vertex a of T and the midpoint m of the opposite side bc . Look at the triangle amb . Clearly, at m there is the right angle. Thus by the Pythagorean theorem for \mathbb{H}^2 we obtain $\cosh |ab| = \cosh |mb| \cdot \cosh |ma|$, this is $\cosh 2x = \cosh x \cdot \cosh \eta_x$. Hence $\eta_x = \text{arccosh} \frac{\cosh 2x}{\cosh x}$.

Example 3. Denote by S_{2x} the regular tetrahedron $uvwz$ whose edges are of length $2x$ and by p the projection of z onto the facet uvw . By Lemma 1 of Kellerhals [8] for \mathbb{H}^2 we know that the circumradius of the regular triangle of side $2x$ is $\text{arcsinh}(\sqrt{4/3} \cdot \sinh x)$. So this is $|up|$. Since the triangle zpu has the right angle at p , by the Pythagorean theorem for \mathbb{H}^2 we obtain $\cosh 2x = \cosh[\text{arcsinh}(\sqrt{4/3} \cdot \sinh x)] \cdot \cosh |zp|$. This and $|zp| = \text{width}_G(S_{2x})$ for the hyperplane G containing uvw lead to

$$\text{width}_G(S_{2x}) = \text{arccosh} \frac{\cosh 2x}{\cosh[\text{arcsinh}(\sqrt{4/3} \sinh x)]}. \quad (1)$$

5 Bodies of constant width and reduced bodies

If for every hyperplane supporting a convex body $W \subset \mathbb{H}^d$ the width of W determined by this hyperplane is δ , we say that W is a **body of constant width** δ . This definition is analogous to the definitions in \mathbb{E}^d and \mathbb{S}^d (for instance see [3] and [14]). In all the three cases the concept of width of C with respect to a supporting hyperplane (hemisphere in \mathbb{S}^d) is analogous. By the way, there are different definitions of a body of constant width in \mathbb{H}^d . Some of them apply different notions of width, as for instance Santaló in [18]. Some other take into account a property of a convex body in \mathbb{E}^d equivalent to the definition of a body of constant body there, for instance see the recent paper by Böröczki and Sagemester [2]. A task would be to check the relationships of all such definitions of a body of constant width in \mathbb{H}^d , analogously as this is done for \mathbb{S}^d in the last section of [12].

It is easy to check that balls in \mathbb{H}^d and every Reuleaux polygon in \mathbb{H}^2 (defined as in \mathbb{E}^2) are bodies of constant width.

From Theorem 1 we obtain the following claim.

Claim 2. *For any body W of constant width δ we have $\text{diam}(W) = \delta$.*

Proposition 3. *Every body of constant width is strictly convex.*

Proof. Imagine the opposite to the thesis, i.e., that there exists a body W of constant width δ which is not strictly convex. Then a hyperplane H supporting W contains more than one point. There is a farthest from H ultraparallel hyperplane J supporting C . By Theorem 2 and the sentence just before it there is exactly one point $j \in W \cap J$ and its projection h onto H belongs to $C \cap H$. Clearly, $|jh| = \delta$. Recall that H contains a point t different from h . By Lemma 1 we have $|jt| > |jh|$ which implies that $\text{diam}(W) > \delta$. On the other hand, by Claim 2 we get $\text{width}_H(W) = \text{diam}(W)$. This contradiction shows that our proposition holds true. \square

A few different notions of a body of constant width in \mathbb{H}^d exist. Some information on them are given in pages 240–242 of the book [16] by Martini, Montejano and Oliveros.

In analogy to the definition of reduced bodies in Euclidean space \mathbb{E}^d (see [6] and the survey article [13]) and on the sphere S^d (see [10] and the survey article [12]) we define reduced convex bodies in \mathbb{H}^d . We say that a convex body $R \subset \mathbb{H}^d$ is **reduced** if $\Delta(Z) < \Delta(R)$ for every convex body $Z \subset R$ different from R .

From Zorn's lemma we see that *every convex body $C \subset \mathbb{H}^d$ contains a reduced body whose thickness is $\Delta(C)$.*

It is easy to show that all regular odd-gons in \mathbb{H}^2 are reduced bodies.

Proposition 4. *Every body of constant width in \mathbb{H}^d is reduced.*

Proof. Let $W \subset \mathbb{H}^d$ be a body of constant width. In order to confirm that W is reduced take an arbitrary convex body $Z \subset W$ different from W ; we intend to show that $\Delta(Z) < \Delta(W)$.

There exists a hyperplane K supporting Z which has non-empty intersection with the interior of W . Take any point $q \in Z$ farthest from K , which by Proposition 1 means that $\text{dist}(q, K) = \text{width}_K(Z)$. Provide an ultraparallel hyperplane L to K which supports W as far as it is possible from K on the opposite side of K than Z is situated. By Lemma 1 we have $\text{dist}(q, K) < \text{dist}(q, L)$. By Proposition 1 applied to $\text{width}_L(W)$ we see that $\text{width}_L(W) \geq \text{dist}(q, L)$ (still each most far point of W from L is in a distance at least $\text{dist}(q, L)$). Since W is of constant width, every its width is equal to $\Delta(W)$. In particular, $\text{width}_L(W) = \Delta(W)$.

By the above observations we obtain $\Delta(W) = \text{width}_L(W) \geq \text{dist}(q, L) > \text{dist}(q, K) = \text{width}_K(Z) \geq \Delta(Z)$. From the obtained inequality $\Delta(Z) < \Delta(W)$ we conclude that W is a reduced body. \square

Take a reduced body in \mathbb{H}^2 with an axis of symmetry and rotate it in \mathbb{H}^3 around this axis. It appears that the obtained 3-dimensional body is reduced (see Fig. 4 for rotating the Reuleaux triangle around its axis of symmetry). Similarly as in \mathbb{E}^3 and \mathbb{S}^3 .




Fig. 4. A rotational reduced body




Fig. 5. A $\frac{1}{8}$ -th part of a ball

Dissect a ball in \mathbb{H}^d by d orthogonal hyperplanes through its center into 2^d closed parts called $\frac{1}{2^d}$ -**th parts of a ball**. Clearly, the thickness of each of them is equal to the radius of the above ball. Observe that every $\frac{1}{2^d}$ -th part of a ball is a reduced body (similarly as in \mathbb{E}^d and \mathbb{S}^d , see [9] and [10]). The 3-dimensional case is seen in Fig. 5.

Example 4. Take S_{2x} from Example 3. Provide the hyperplane H containing the edge uv and orthogonal to mn , where m is the midpoint of uv and n is the midpoint of wz (see Fig. 6). By the Pythagorean theorem for the rectangular triangle nmv we obtain $\cosh |nv| = \cosh |mn| \cdot \cosh |mv|$ which means that $\cosh |mn| = \frac{\cosh |nv|}{\cosh |mv|}$, i.e., that $|mn| = \operatorname{arccosh} \frac{\cosh |nv|}{\cosh |mv|}$. Hence $|mn| = \operatorname{arccosh} \frac{\cosh |nv|}{\cosh |mv|}$. Still $|mv| = x$ and $\cosh |nv| = \cosh \eta_x = \frac{\cosh 2x}{\cosh x}$ as shown Example 2. By Proposition 1 the width of S_{2x} determined by H equals to the distance of a farthest point of S_{2x} from H . Such a point is z . By h denote the projection of z onto H . Then $\operatorname{width}_H(C) = |hz|$. Observe that points m, n, h and z are in a hyperplane. Consider the Lambert quadrilateral $nmhz$. The right angles are at n, m and h . Thus $\sinh |zh| = \sinh |mn| \cdot \cosh |nz|$. Hence $|zh| = \operatorname{arcsinh} (\sinh |mn| \cdot \cosh |nz|)$. By Proposition 1 we have $|zh| = \operatorname{width}_H(T)$. Moreover, taking into account $|nz| = x$ and the established earlier formula for $|mn|$ we obtain

$$\operatorname{width}_H(S_{2x}) = \operatorname{arcsinh} \left[\sinh \left(\operatorname{arccosh} \frac{\cosh 2x}{\cosh^2 x} \right) \cdot \cosh x \right]. \quad (2)$$

Remark 2. In \mathbb{E}^d for $d \geq 3$ every regular simplex is not reduced (see [17]). The author tried to check the situation in \mathbb{H}^3 expecting that sufficiently large tetrahedra are reduced. It appears that the opposite is true. We show this by applying the formulas $\cosh(\operatorname{arccosh} a) = \sqrt{1 - a^2}$, $\sinh(\operatorname{arccosh} a) = \sqrt{a^2 - 1}$, $\cosh 2x = 2 \cosh^2 x - 1$ and $\sinh x = \sqrt{\cosh^2 x - 1}$ for the difference of (1) and (2), which after substituting $\lambda = \cosh 2x$ leads to the inequality




Fig. 6. Illustration to Example 4

$29\lambda^2 - 36\lambda + 7 < 0$. It is never fulfilled for $\lambda > 1$, so never for any positive x . So every S_{2x} is not reduced (the author expects that the situation is analogous for every $d > 3$). By the way, $\text{width}_H(S_{2x})/\text{width}_G(S_{2x})$ tends to 1 as $x \rightarrow \infty$.

We conjecture that for any reduced convex body $R \subset \mathbb{H}^d$ and any supporting hyperplane H such that $\text{width}_H(C) = \Delta(R)$ there are unique $a \in R \cap H$ and $b \in \text{bd}(R)$ such that $ab \perp H$ and $|ab| = \Delta(R)$.

Is it true that every smooth reduced body in \mathbb{H}^d is of constant width? Similarly as in \mathbb{E}^d , which is proved by Groemer [5] (compare Corollary 1 from [9]) and also as in the version for \mathbb{S}^d shown in Theorem 5 from [10].

6 Complete and constant diameter bodies

We say that a convex body $D \subset \mathbb{H}^d$ is of **constant diameter** δ provided $\text{diam}(D) = \delta$ and for every $p \in \text{bd}(D)$ there is a point $p' \in \text{bd}(D)$ with $|pp'| = \delta$. This definition is analogous to the one for \mathbb{S}^d from Section 4 of [14].

Similarly to the traditional notion of a complete set in \mathbb{E}^d (for instance, see [1], [3] and [4]) we say that a set $C \subset \mathbb{H}^d$ of diameter δ is **complete** provided $\text{diam}(C \cup \{x\}) > \delta$ for every $x \notin C$.

We omit the proof of the following two claims since they are similar to the proof by Lebesgue [15] in \mathbb{E}^d , which is recalled in Section 64 of [1].

Claim 3. *Every set of a diameter δ in \mathbb{H}^d is a subset of a complete set of diameter δ .*

Claim 4. *Every complete body C of a diameter δ in \mathbb{H}^d coincides with the intersection of all balls of radius δ centered at points of C .*

This claim permits to use the term a **complete convex body** for a complete set.

Proposition 5. *If $C \subset H^d$ is a complete body of diameter δ , then for every $p \in \text{bd}(C)$ there exists $p' \in C$ such that $|pp'| = \delta$.*

Proof. Imagine that the thesis is not true. Then $|pp'| < \delta$ for a point $p \in \text{bd}(C)$ and for every point $p' \in C$. Since C is compact, there is an $\omega < \delta$ such that $|pp'| \leq \omega$ for every $p' \in C$. Take a ball B of radius $\delta - \omega$ centered at p . Of course, there is a supporting hyperplane H supporting C at p . Observe that the open halfspace bounded by H and containing the interior of C contains the relative interior of the segment pp' . Consequently, the straight line through p' and p contains a point $x \in B$, but out of C , in the distance $\delta - \omega$ from p . Thus $|p'x| = \delta$. We get $\text{diam}(C \cup \{x\}) \geq \delta$. Since $x \notin C$, this contradicts the assumption that C is a complete body of diameter δ . Hence our proposition holds true. \square

This proposition is analogous the Lemma 2 of [11] and holds true also in E^d . Just this proof can be repeated also for every complete body in E^d .

Theorem 3. *Let $C \subset \mathbb{H}^d$. We have (a) \implies (b) \iff (c), where*

- (a) *C is a body of constant width δ ,*
- (b) *C is a body of constant diameter δ ,*
- (c) *C is complete body of diameter δ .*

Proof. Show that (a) implies (b).

Assume that C is of constant width δ . By Theorem 1 we have $\text{diam}(C) = \delta$, which is the first requirement of the definition of a body of constant diameter.

Take an arbitrary point $p \in \text{bd}(C)$. Let P be any hyperplane supporting C at p . Since C is of constant width δ , we have $\text{width}_P(C) = \delta$. Let p' be a farthest point of C from P . By Proposition 1 its distance from P is $\text{width}_P(C)$. This and $p \in P$ imply $|pp'| \geq \delta$. Hence by $\text{diam}(C) = \delta$ we get $|pp'| = \text{width}_P(C)$. We conclude that p' fulfills the second requirement of the definition of a body of constant diameter.

From the above two paragraphs we see that C is a body of constant diameter δ .

Show that (b) and (c) are equivalent.

Consider a body $C \subset \mathbb{H}^d$ of constant diameter δ . Take any $r \notin C$. Let B be the largest ball whose interior is disjoint with C . Denote by ρ the radius of this ball and by p the common point of B and C . Since C is of constant diameter δ , there is $p' \in C$ in the

distance δ from p . Clearly, a unique hyperplane supports B and C at p . Hence $\angle rpp' \geq \frac{\pi}{2}$. Thus by the hyperbolic law of cosines for the triangle rpp' we get $|rp'| > |pp'|$ which means that $|rp'| > \delta$. Consequently, $\text{diam}(C \cup \{r\}) > \text{diam}(C)$ for every $r \notin C$. So C is complete.

On the other side, if $C \subset \mathbb{H}^d$ is a complete body of diameter δ , then by Proposition 5 we conclude that C is of constant diameter. \square

We conjecture that (a), (b) and (c) are equivalent. In order to confirm this, it is sufficient to show that (b) or (c) imply (a).

References

- [1] Bonnesen, T., Fenchel, T. W.: *Theorie der konvexen Körper*. Springer, Berlin (1934) (English translation: *Theory of Convex Bodies*, BCS Associates, Moscow, Idaho, (1987))
- [2] Böröczky, K, Sagemester, Á.: *Convex bodies of constant width in spaces of constant curvature and the extremal area of Reuleaux triangles*, *Studia Scientiarum Mathematicarum Hungarica* **59**, 244–273 (2022)
- [3] Chakerian G. D., Groemer, H.: *Convex bodies of constant width*, In *Convexity and its applications*, pp. 49–96, Birkhäuser, Basel (1983)
- [4] Eggleston, E. G.: *Convexity*, vol. 47. Cambridge Tracts in Mathematics and Mathematical Physics, Cambridge University Press, New York (1958)
- [5] Groemer, H.: *Extremal convex sets*, *Monatsh. Math.* **96** 29–39 (1983)
- [6] Heil, E.: *Kleinste konvexe Körper gegebener Dicke*, Preprint No. 453, Fachbereich Mathematik der TH Darmstadt, (1978)
- [7] Horváth, Á.: *Diameter, width and thickness in the hyperbolic plane*, *J. Geom.* **112**, No. 47, 29p. (2021)
- [8] Kellerhals, R.: *Regular simplices and lower volume bounds for hyperbolic n -manifolds*, *Annals of Global Analysis and Geometry* **13**, 377–392 (1995)
- [9] Lassak, M.: *Reduced convex bodies in the plane*, *Israel J. Math.* **70**, No. 3, 365–379 (1990)
- [10] Lassak, M.: *Width of spherical convex bodies*, *Aequationes Math.* **89**, 555–567 (2015)
- [11] Lassak, M.: *Complete spherical convex bodies*, *J. Geom.* **111**(2), No. 35, 6p. (2020)
- [12] Lassak, M.: *Spherical Geometry – A Survey on Width and Thickness of Convex bodies* in *Surveys in geometry I*. Papadopoulos, Athanase (ed.), Cham: Springer. 7–47 (2022)
- [13] Lassak, M., Martini, H.: *Reduced convex bodies in Euclidean space – a survey*, *Results Math.* **66**, 405–426 (2014)
- [14] Lassak, M., Musielak M.: *Spherical convex bodies of constant width*, *Aequationes Math.* **92** (2018), 627–640.

- [15] Lebesgue, H.: *Sur quelques questions de minimum, relatives aux courbes orbiformes, et leurs rapports avec le calcul des variations*, J. Math. Pures Appl. **8**, 67–96 (1921)
- [16] Martini, M., Montejano, L., Oliveros, D.: *Bodies of constant width. An introduction to convex geometry with applications*, Springer Nature Switzerland AG, 2019
- [17] Martini, H., Swanepoel, K.: *Non-planar simplices are not reduced*, Publ. Math. Debrecen **64**, 101–106 (2004)
- [18] Santaló, L. A.: *Note on convex curves on the hyperbolic plane*, Bull. Amer. Math. Soc. **51**, 405–412 (1945)

Marek Lassak
University of Science and Technology
85-789 Bydgoszcz, Poland
e-mail: lassak@pbs.edu.pl