From dense to sparse design: Optimal rates under the supremum norm for estimating the mean function in functional data analysis

Max Berger, Philipp Hermann and Hajo Holzmann* Department of Mathematics and Computer Science Philipps-Universität Marburg {mberger, herm, holzmann}@mathematik.uni-marburg.de

May 9, 2024

Abstract

We derive optimal rates of convergence in the supremum norm for estimating the Hölder-smooth mean function of a stochastic process which is repeatedly and discretely observed with additional errors at fixed, multivariate, synchronous design points, the typical scenario for machine recorded functional data. Similarly to the optimal rates in L_2 obtained in Cai and Yuan (2011), for sparse design a discretization term dominates, while in the dense case the parametric \sqrt{n} rate can be achieved as if the n processes were continuously observed without errors. The supremum norm is of practical interest since it corresponds to the visualization of the estimation error, and forms the basis for the construction uniform confidence bands. We show that in contrast to the analysis in L_2 , there is an intermediate regime between the sparse and dense cases dominated by the contribution of the observation errors. Furthermore, under the supremum norm interpolation estimators which suffice in L_2 turn out to be sub-optimal in the dense setting, which helps to explain their poor empirical performance. In contrast to previous contributions involving the supremum norm, we discuss optimality even in the multivariate setting, and for dense design obtain the \sqrt{n} rate of convergence without additional logarithmic factors. We also obtain a central limit theorem in the supremum norm, and provide simulations and real data applications to illustrate our results.

Keywords. asymptotic confidence sets, dense and sparse design, functional data, minimax optimality, supremum norm

1 Introduction

Functional data analysis (FDA) refers to situations where the data involve functions or images, potentially as features with additional observations in regression and classification problems.

^{*}Corresponding author. Prof. Dr. Hajo Holzmann, Department of Mathematics and Computer Science, Philipps-Universität Marburg, Hans-Meerweinstr., 35043 Marburg, Germany

The field was popularized by Ramsay and Silvermann (1998), and since then has attracted the attention of many theoretical and applied statisticians. Wang et al. (2016), Morris (2015) and Cuevas (2014) contribute overviews of various aspects of the field.

Replication and regularization are two central notions in FDA (Morris, 2015; Ramsay and Silvermann, 1998). By replication one makes use of information across the repeated observations of the functions, while regularization takes advantage of information within functions, which in practice are only observed at discrete design points in time or space.

Fixed, non-random points which are equal across functions are referred to as a deterministic, synchronous design. This typically arises for machine recorded data, such as temperature and further weather data recorded at regular time intervals at weather stations. In this paper, for such designs we clarify even in a multivariate setting the phase transition from dense to sparse case, by using minimax optimality over Hölder smoothness classes and the supremum norm as loss function.

The multivariate and in particular the bivariate settings which we comprehensively deal with in the present paper evidently are of great interest, for example for repeatedly observed images. Much less work is available here than for the univariate case, in which the covariate typically refers to time. A notable exception is Wang et al. (2020), who consider images, focus on the dense case and work under a more complex set of assumptions then is considered in the present paper.

The analysis of estimators in FDA has traditionally been conducted for L_2 and pointwise errors (Hall et al., 2006). Arguably the supremum norm is of more or at least equal practical interest, since it corresponds to the visualization of the estimation error. Further it constitutes the basis for the construction uniform confidence bands, which allow for the assessment of the precision of the estimates. Therefore in recent years much research effort has been devoted to an analysis of estimation errors in the supremum norm. Notable contributions in the random, asynchronous design are Yao et al. (2005); Li and Hsing (2010); Zhang and Wang (2016). Rates in the supremum norm and asymptotic confidence bands in the deterministic, synchronous design that we focus on have been investigated by Degras (2011); Chang et al. (2017); Xiao (2020); Kalogridis and Van Aelst (2022).

A precise assessment of the performance of methods can however only be given by providing optimality results, which we do in terms of minimax rates under the supremum norm in this paper, and which have not been previously derived in the literature.

More technically speaking, we show that in the sparse regime, a discretization term dominates rather than the estimation error arising from the errors as in the random design case. This is in line with results in Cai and Yuan (2011) for the L_2 -loss and strong smoothness assumptions on the paths of the processes. In particular, they require that the sample paths have the same smoothness as the target mean function, a much stronger assumption than what is imposed in our work. Notably and in contrast to the L_2 norm there exists a transition phase between the sparse and dense designs when considering the supremum norm in which the contribution of the errors dominates. Our analysis also shows that simple interpolation estimators without smoothing, which according to Cai and Yuan (2011) suffice to achieve optimal rates in L_2 for dense design, in case of observational errors actually deteriorate with respect to the supremum norm for a large number of row wise observations. Also note that the rates of convergence presented in Xiao (2020); Kalogridis and Van Aelst (2022) by using penalized spline estimators are suboptimal. In particular, the parametric \sqrt{n} -rate in the dense design is not attained, which is however very desirable for asymptotic inference. The parametric \sqrt{n} -rate in the dense setting allows us to provide a central limit theorem in the supremum norm, which is the basis for the construction of uniform confidence bands, e.g. by bootstrap methods (Degras, 2011; Wang et al., 2020) or by using bounds based on the Kac-Rice formula as recently proposed in Liebl and Reimherr (2023). As real-data illustrations we consider the biomechanical data set from Liebl and Reimherr (2023) as well as daily temperature curves at Nuremberg, Germany. The effect of coarser discretization on the estimation quality is determined by using a confidence band for the estimate based on the finest discretization, and investigating whether estimates with coarser design still lie inside this band.

The paper is organized as follows. In Section 2 we introduce the model and the linear estimators that we shall consider. We work with generic assumptions on the weights, and in Section 4 verify them for weights of local polynomial estimates. Section 3 contains the main results on rates of estimation of the mean function in the supremum norm. Upper bounds and matching lower bounds are presented. We also show convergence in distribution to a Gaussian process. In Section 5 we illustrate our theoretical results in a simulation study. In Section 6 we provide real-data illustrations. The concluding Section 7 describes how to transfer the results to more general designs, and points to open problems for the sparse, asynchronous case. Proofs of the main results are given in Section 8, while some technical proofs as well as additional simulations are deferred to an appendix.

Notation

In the paper we use the following notation. For $\boldsymbol{a} = (a_1, \ldots, a_d)^{\top}$ and $\boldsymbol{b} = (b_1, \ldots, b_d)^{\top} \in \mathbb{R}^d$ write $\boldsymbol{a} \leq \boldsymbol{b}$ if $a_i \leq b_i$ for all $i = 1, \ldots, d$, $\boldsymbol{a}^{\boldsymbol{b}} := a_1^{b_1} \cdots a_d^{b_d}$, $|\boldsymbol{a}| = a_1 + \ldots + a_d$, $\boldsymbol{a}! := a_1! \cdots a_d!$, $\boldsymbol{a}_{\min} = \min_{1 \leq r \leq d} a_r$ and $\partial^{\boldsymbol{a}} := \partial_1^{a_1} \ldots \partial_d^{a_d}$ if the coordinates are integers. Further set $\boldsymbol{1} := (1, \ldots, 1)^{\top} \in \{1\}^d$. Given $\boldsymbol{p} := (p_1, \ldots, p_d)^{\top}$ denote by $\{\boldsymbol{j} \in \mathbb{N}^d \mid \boldsymbol{1} \leq \boldsymbol{j} \leq \boldsymbol{p}\}$ the set of $\boldsymbol{j} \in \mathbb{N}^d$ that are component wise between $\boldsymbol{1}$ and \boldsymbol{p} , and denote $\sum_{\boldsymbol{j=1}}^{\boldsymbol{p}} := \sum_{j_1=1}^{p_1} \cdots \sum_{j_d=1}^{p_d}$. If for a vector the indices are written in bold letters this indexes vectors such as $\boldsymbol{p}_{\boldsymbol{j}} := (p_{j_1}, \ldots, p_{j_d})^{\top}$. If only the indices are written bold this indexes a scalar like $\varepsilon_{i,\boldsymbol{j}} := \varepsilon_{i,j_1,\ldots,j_d}$ or $Y_{\boldsymbol{j}} := Y_{j_1,\ldots,j_d}$. For sequences $(p_n), (q_n)$ tending to infinity, we write $p_n \leq q_n$, if $p_n = \mathcal{O}(q_n), p_n \simeq q_n$ if $p_n \leq q_n$ and $q_n \leq p_n$, and $p_n \ll q_n$, if $p_n = o(q_n)$.

The notation $\mathcal{O}_{\mathbb{P},h\in(h_1,h_0]}$ stands for stochastic dominance uniformly for $h\in(h_1,h_0]$. Finally, $\|\cdot\|_{\infty}$ denotes the supremum norm on $[0,1]^d$ (sup-norm in the following).

2 The model, smoothness classes and linear estimators

Assume that we have data $(Y_{i,j}, x_j)$ according to the model

$$Y_{i,j} = \mu(\boldsymbol{x}_j) + Z_i(\boldsymbol{x}_j) + \varepsilon_{i,j}, \quad i = 1, \dots, n, \quad \boldsymbol{j} = (j_1, \dots, j_d), \quad (1)$$

where $Y_{i,j}$ are real-valued response variables and $x_j \in T \subset \mathbb{R}^d$ the design points. The set T is assumed to be a hypercube and we take $T = [0, 1]^d$ in the following. The mean function $\mu: T \to \mathbb{R}$ is to be estimated, $\varepsilon_{i,j}$ are independently distributed errors with mean zero and Z, Z_1, \ldots, Z_n are centered, independent and identically distributed, square integrable random fields on T distributed as the generic Z. We shall assume a Cartesian product structure $x_j = (x_{1,j_1}, \ldots, x_{d,j_d}), 1 \leq j_k \leq p_k, k = 1, \ldots, d$, and $x_{k,l} < x_{k,l+1}, 1 \leq l \leq p_k - 1$, for the covariates. The number $p^1 = p^1(n) = \prod_{k=1}^d p_k(n)$ of design points and the design points x_j depend on n, which is often suppressed in the notation.

Given $\alpha > 0$ we set $\lfloor \alpha \rfloor = \max\{k \in \mathbb{N}_0 \mid k < \alpha\}$. A function $f: T \to \mathbb{R}$ is Hölder-smooth with index α if for all indices $\boldsymbol{\beta} = (\beta_1, \ldots, \beta_d)^\top$ with $|\boldsymbol{\beta}| \leq \lfloor \alpha \rfloor$ the derivatives $\partial^{\boldsymbol{\beta}} f(\boldsymbol{x})$ exist and the Hölder-norm given by

$$\|f\|_{\mathcal{H},\alpha} \coloneqq \max_{|\boldsymbol{\beta}| \leq \lfloor \alpha \rfloor} \sup_{\boldsymbol{x} \in T} |\partial^{\boldsymbol{\beta}} f(\boldsymbol{x})| + \max_{|\boldsymbol{\beta}| = \lfloor \alpha \rfloor} \sup_{\boldsymbol{x}, \boldsymbol{y} \in T, \, \boldsymbol{x} \neq \boldsymbol{y}} \frac{|\partial^{\boldsymbol{\beta}} f(\boldsymbol{x}) - \partial^{\boldsymbol{\beta}} f(\boldsymbol{y})|}{\|\boldsymbol{x} - \boldsymbol{y}\|_{\infty}^{\alpha - \lfloor \alpha \rfloor}}$$

is finite. The Hölder class on T with parameters $\alpha > 0$ and L > 0 is defined by

$$\mathcal{H}_T(\alpha, L) = \left\{ f \colon T \to \mathbb{R} \mid \|f\|_{\mathcal{H}, \alpha} \le L \right\}.$$
(2)

For the distribution of the processes Z we assume that $\mathbb{E}[Z(\mathbf{0})^2] < \infty$ and that there exists a random variable $M = M_Z > 0$ with $\mathbb{E}[M^2] < \infty$ such that

$$|Z(\boldsymbol{x}) - Z(\boldsymbol{y})| \le M \|\boldsymbol{x} - \boldsymbol{y}\|_{\infty}^{\beta}$$
 almost surely (3)

for all $x, y \in T$, where $0 < \beta \leq 1$ is a constant. Given $C_Z > 0$ and $0 < \beta_0 \leq 1$, we consider the class of processes

$$\mathcal{P}_T(\beta_0, C_Z) = \left\{ Z : T \to \mathbb{R} \text{ random field} \mid \exists \beta \in [\beta_0, 1] \text{ and } M \text{ s.th.} \\ \mathbb{E}[M^2] + \mathbb{E}[Z(\mathbf{0})^2] \le C_Z \text{ and } (3) \text{ holds} \right\}.$$
(4)

Remark 1. Note that Hölder continuity of the process Z in (3) is a natural assumption when deriving the parametric $1/\sqrt{n}$ -rate and CLTs in the supremum norm, in view of the Jain Markus Theorem (van der Vaart and Wellner, 1996, Example 2.11.13). Various contributions which work without direct smootheness assumptions on the paths (Li and Hsing, 2010; Zhang and Wang, 2016; Xiao, 2020) only achieve a rate of $(\log(n)/n)^{1/2}$ in the dense setting. While this does not constitute a big difference in terms of rates, for the asymptotics one requires the exact parametric rate. Further, it is of interest when reconstruction can be conducted as if the processes were continuously observed without errors. Let us point out that in contrast to Cai and Yuan (2011), we however do not assume that the paths of the process Z are as smooth as the mean function. Typically μ will be in a Hölder class of smoothness $\alpha > 0$ which will be larger than β in (3).

Assumption 1 (Distributional assumptions). The random variables $\{\varepsilon_{i,j} \mid 1 \leq i \leq n, 1 \leq j \leq p\}$ are independent and independent of the processes Z_1, \ldots, Z_n . Further we assume that the distribution of $\varepsilon_{i,j}$ is sub-Gaussian, and setting $\sigma_{ij}^2 := \mathbb{E}[\varepsilon_{i,j}^2]$ we have that $\sigma^2 := \sup_n \max_{1 \leq j \leq p} \sigma_j^2 < \infty$ and that there exists $\zeta > 0$ such that $\zeta^2 \sigma_{i,j}^2$ is an upper bound for the sub-Gaussian norm of $\varepsilon_{i,j}$.

Remark 2. In the case where the distribution of the additional errors $\varepsilon_{i,j}$ is not a sub-Gaussian distribution a Gaussian approximation yields similar rates of convergence under the assumption that $\mathbb{E}|\varepsilon_{i,j}|^q < \infty$ for some $q \ge 4$ and a additional assumption on the weights of the estimator. The resulting rate of convergence and the proof and the discussion of the dominating rates of convergence for the one-dimensional case d = 1 can be found in the Appendix D.

For the mean function μ , we consider linear estimators of the form

$$\widehat{\mu}_{n,\boldsymbol{p},h}(\boldsymbol{x}) = \sum_{\boldsymbol{j=1}}^{\boldsymbol{p}} w_{\boldsymbol{j},\boldsymbol{p}}(\boldsymbol{x};h;\boldsymbol{x_1},\ldots,\boldsymbol{x_p}) \, \bar{Y}_{\boldsymbol{j}} \,, \qquad \text{where} \quad \bar{Y}_{\boldsymbol{j}} = \frac{1}{n} \sum_{i=1}^{n} Y_{i,\boldsymbol{j}}, \tag{5}$$

where $\boldsymbol{x} \in T$, $w_{\boldsymbol{j},\boldsymbol{p}}(\boldsymbol{x};h;\boldsymbol{x}_1,\ldots,\boldsymbol{x}_p)$ are deterministic weights depending on the design points and on a bandwidth parameter h > 0. We often abbreviate the notation by writing $w_{\boldsymbol{j},\boldsymbol{p}}(\boldsymbol{x};h) :=$ $w_{\boldsymbol{j},\boldsymbol{p}}(\boldsymbol{x};h;\boldsymbol{x}_1,\ldots,\boldsymbol{x}_p)$.

While the following assumptions on the weights are standard, note that we require them to hold also for h of order $1/p_{\min}$, a regime which is not relevant in traditional nonparametric regression. Here recall the notation $p_{\min} = \min_k p_k$ for $p = (p_1, \ldots, p_d)$.

Assumption 2 (Weights of linear estimator). There are a c > 0 and a $h_0 > 0$ such that for sufficiently large p_{\min} , the following holds for all $h \in (c/p_{\min}, h_0]$ for suitable constants $C_1, C_2 > 0$ which are independent of n, p, h and x.

(W1) The weights reproduce polynomials of a suitable degree $\zeta \geq 1$, that is for $x \in T$,

$$\sum_{1 \le j \le p} w_{j,p}(\boldsymbol{x},h) = 1, \quad \sum_{1 \le j \le p} (\boldsymbol{x}_j - \boldsymbol{x})^{\boldsymbol{r}} w_{j,p}(\boldsymbol{x};h) = 0, \quad \boldsymbol{r} \in \mathbb{N}^d \text{ s.t. } |\boldsymbol{r}| = 1, \dots, \zeta.$$

(W2) We have $w_{j,p}(\boldsymbol{x};h) = 0$ if $\|\boldsymbol{x}_j - \boldsymbol{x}\|_{\infty} > h$ with $\boldsymbol{x} \in T$.

(W3) For the absolute values of the weights, $\max_{1 \le j \le p} |w_{j,p}(\boldsymbol{x};h)| \le \frac{C_1}{p^1 h^d}, \, \boldsymbol{x} \in T.$

(W4) For a constant $C_2 > 0$ it holds that

$$|w_{\boldsymbol{j},\boldsymbol{p}}(\boldsymbol{x};h) - w_{\boldsymbol{j},\boldsymbol{p}}(\boldsymbol{y};h)| \leq \frac{C_2}{\boldsymbol{p}^1 h^d} \left(\frac{\|\boldsymbol{x}-\boldsymbol{y}\|_{\infty}}{h} \wedge 1\right), \qquad \boldsymbol{x},\boldsymbol{y} \in T.$$

Moreover, we require the following design assumption.

Assumption 3 (Design Assumption). There is a constant $C_3 > 0$ such that for each $x \in T$ and h > 0 we have that

card
$$\{ \boldsymbol{j} \in \{\boldsymbol{1},\ldots,\boldsymbol{p}\} \mid \boldsymbol{x} - (h,\ldots,h) \leq \boldsymbol{x}_{\boldsymbol{j}} \leq \boldsymbol{x} + (h,\ldots,h) \} \leq C_3 h^d \boldsymbol{p}^1.$$

Note that for a product design, we could formulate Assumption 3 coordinatewise. However, in the present formulation it also applies to more general designs. In case of design densities, Assumption 3 is satisfied, and we check the above assumptions on the weights of local polynomial estimators in Section 4.

3 Optimal rates of estimation for the mean function

Upper bounds

We start by deriving upper bounds on the rate of convergence of the linear estimator in (5) under Assumption 1. Bounding the terms in the error decomposition

$$\widehat{\mu}_{n,p,h}(\boldsymbol{x}) - \mu(\boldsymbol{x}) = \sum_{\boldsymbol{k}=1}^{\boldsymbol{p}} w_{\boldsymbol{k},\boldsymbol{p}}(\boldsymbol{x};h) \left(\mu(\boldsymbol{x}_{\boldsymbol{k}}) - \mu(\boldsymbol{x})\right) + \sum_{\boldsymbol{k}=1}^{\boldsymbol{p}} w_{\boldsymbol{k},\boldsymbol{p}}(\boldsymbol{x};h) \,\overline{\varepsilon}_{\boldsymbol{k}} + \sum_{\boldsymbol{k}=1}^{\boldsymbol{p}} w_{\boldsymbol{k},\boldsymbol{p}}(\boldsymbol{x};h) \,\overline{Z}_{n}(\boldsymbol{x}_{\boldsymbol{k}}) \\ =: I_{1}^{\boldsymbol{p},h}(\boldsymbol{x}) + I_{2}^{n,\boldsymbol{p},h}(\boldsymbol{x}) + I_{3}^{n,\boldsymbol{p},h}(\boldsymbol{x}) , \qquad \boldsymbol{x} \in [0,1]^{d} , \qquad (6)$$

where

$$ar{arepsilon_j} = rac{1}{n}\sum_{i=1}^n arepsilon_{i,j}$$
 and $ar{Z}_n(oldsymbol{x}) = rac{1}{n}\sum_{i=1}^n Z_i(oldsymbol{x})$,

leads to the following result.

Theorem 1. In model (1) under Assumptions 1 and 3, consider the linear estimator $\hat{\mu}_{n,\boldsymbol{p},h}$ in (5) with weights satisfying Assumption 2 with $\gamma = \lfloor \alpha \rfloor$ for given Hölder smoothness $\alpha > 0$. Then for sufficiently large \boldsymbol{p}_{\min} and n,

$$\sup_{h \in (c/\boldsymbol{p}_{\min},h_0]} \sup_{Z \in \mathcal{P}_T(\beta_0,C_Z), \mu \in \mathcal{H}(\alpha;M)} a_{n,\boldsymbol{p},h}^{-1} \mathbb{E}_{\mu,Z} \Big[\big\| \widehat{\mu}_{n,\boldsymbol{p},h} - \mu \big\|_{\infty} \Big] = \mathcal{O}(1),$$
(7)

where

$$a_{n,\boldsymbol{p},h} = \max\left(h^{\alpha}, \left(\frac{\log(h^{-1})}{n\boldsymbol{p}^{1}h^{d}}\right)^{1/2}, n^{-\frac{1}{2}}\right).$$
 (8)

Therefore, setting

$$h^{\star} = \max\left(c/\boldsymbol{p}_{\min}, \left(\frac{\log(n\,\boldsymbol{p}^{1})}{n\,\boldsymbol{p}^{1}}\right)^{\frac{1}{2\alpha+d}}\right),\tag{9}$$

 $we \ obtain$

$$\sup_{Z \in \mathcal{P}_T(\beta_0, C_Z), \mu \in \mathcal{H}(\alpha; M)} \mathbb{E}_{\mu, Z} \Big[\big\| \widehat{\mu}_{n, \boldsymbol{p}, h^\star} - \mu \big\|_{\infty} \Big] = \mathcal{O} \Big(\max \Big(\boldsymbol{p}_{\min}^{-\alpha}, \Big(\frac{\log(n \, \boldsymbol{p}^1)}{n \, \boldsymbol{p}^1} \Big)^{\frac{\alpha}{2\alpha + d}}, n^{-\frac{1}{2}} \Big) \Big).$$
(10)

Here the bound h^{α} in (8) arises from the bias term $I_1^{\boldsymbol{p},h}(\boldsymbol{x})$ in (6) by standard bias estimates using property (W1) of the weights, the bound $n^{-1/2}$ follows from the stochastic process term $I_3^{n,\boldsymbol{p},h}(\boldsymbol{x})$ by using $\mathbb{E}[\|\bar{Z}_n\|_{\infty}] = \mathcal{O}(n^{-1/2})$ and boundedness of sums of absolute values of the weights, and the intermediate term in (8) from $I_2^{n,\boldsymbol{p},h}(\boldsymbol{x})$ upon using Dudley's entropy bound (van der Vaart and Wellner, 1996, Corollary 2.2.8). We provide the details in Section 8.1.

Lower bounds

Next we show optimality of the upper bounds in Section 3 by deriving corresponding lower bounds of estimation in the minimax sense (see Tsybakov (2004, Section 2)).

Theorem 2. Assume that in model (1) the errors $\varepsilon_{i,j}$ are *i.i.d.* $\mathcal{N}(0, \sigma_0^2)$ - distributed, $\sigma_0^2 > 0$. The design points $\mathbf{x_1}, \ldots, \mathbf{x_p}$ shall satisfy Assumption 5. Then setting

$$a_{n,\boldsymbol{p}} = \max\left\{\boldsymbol{p}_{\min}^{-\alpha}, \left(\frac{\log(n\,\boldsymbol{p}^{1})}{n\boldsymbol{p}^{1}}\right)^{\frac{\alpha}{2\alpha+d}}, n^{-\frac{1}{2}}\right\}$$

we have that

$$\liminf_{n,\boldsymbol{p}\to\infty}\inf_{\hat{\mu}_{n,\boldsymbol{p}}}\sup_{\mu\in\mathcal{H}(\alpha,L),\ Z\in\mathcal{P}_{T}(\beta_{0},C_{Z})}\mathbb{E}_{\mu,Z}\left[a_{n,\boldsymbol{p}}^{-1}\left\|\hat{\mu}_{n,\boldsymbol{p}}-\mu\right\|_{\infty}\right]\geq c>0,$$

where the infimum is taken over all estimators $\hat{\mu}_{n,p}$ of μ .

The proof is given in Section 8.2.

Remark 3. We point out that the lower bound does not require the design Assumption 3. The argument and hence the result can be adapted to more general distributions of the errors. What is required in our argument is that the Kullback-Leibler divergence in the location model associated with the distribution of the errors is quadratic in the location parameter, at least locally around 0. See for example Tsybakov (2004), Assumption B in Section 2.3 and Section 2.5. Moreover, intuitively speaking Gaussian errors are quite regular, and we show that these already suffice to achieve a lower bound slow enough to match the upper bound. The upper bound holds more generally, so slower minimax rates cannot occur, and when moving away from Gaussian errors, one does not expect faster rates except for degenerate errors. Thus, it appears reasonable to consider Gaussian errors in the lower bounds.

Discussion

Remark 4 (Regimes in the rate). Let us discuss the minimax rates given in Theorems 1 and 2 in terms of p and n and relate them to results previously obtained in the literature. For simplicity, we concentrate on the setting where all coordinates of p are of the same order p, which is assumed to be polynomial in n.

If $p \leq (n/\log n)^{1/(2\alpha)}$, the resulting rate in (10) is $p^{-\alpha}$. The discretization term dominates as observed for the error analysis in L_2 in Cai and Yuan (2011). Note that for the asynchronous, random design considered in Li and Hsing (2010); Wang et al. (2016) this term does not arise.

If in contrast $p \gtrsim (\log n)^{1/d} n^{1/(2\alpha)}$ (10) reduces to $n^{-1/2}$, without the additional logarithmic factors that arise e.g. in Li and Hsing (2010); Wang et al. (2016); Xiao (2020). In between the rate is $(\log(n)/(np^d))^{\alpha/(2\alpha+d)}$, a regime which does not occur in the error

In between the rate is $(\log(n)/(n p^a))^{\alpha/(2a+a)}$, a regime which does not occur in the error analysis in L_2 in Cai and Yuan (2011) but has been obtained for the asynchronous, random design in Li and Hsing (2010, Corollary 3.2, (a)), see also Wang et al. (2016, Corollary 4.2, (1)).

Remark 5 (Choice of the bandwidth). Consider the setup of Theorem 1 in the case that $\max\left((\log(p)/p^1)^{1/d}, p_{\min}^{-1}\right) \leq n^{-\frac{1}{2\alpha}}$ and define the interval

$$\bar{H}_n := \left[c_1 \max\left(\left(\log(p)/\boldsymbol{p}^1\right)^{1/d}, \boldsymbol{p}_{\min}\right), c_2 n^{-\frac{1}{2\alpha}}\right],$$

with constants $c_1, c_2 > 0$. Then the result from Theorem 1 specializes to

$$\sup_{h \in \bar{H}_n} \sup_{Z \in \mathcal{P}_T(\beta_0, C_Z), \mu \in \mathcal{H}(\alpha; M)} \sqrt{n} \mathbb{E}_{\mu, Z} \left[\left\| \widehat{\mu}_{n, p, h} - \mu \right\|_{\infty} \right] = \mathcal{O}(1) \,. \tag{11}$$

Therefore any choice of $h_n \in \overline{H}_n$ is sufficient in order to achieve the \sqrt{n} -rate. Consider for simplicity the setting where all coordinates in p are of the same order p what reduces the assumption on p to $p \gtrsim (\log n)^{(\zeta+\delta)/d} n^{1/(2\alpha)}$ for some $\delta \ge 1$ and $\zeta \ge 0$. In this case taking $h \simeq (\log n)^{\delta/d}/p$ already yields the $n^{-1/2}$ -rate. If ζ has to be 0 then the bias-term is also of order $n^{-1/2}$, and if $\delta = 1$ then the error term is of order $n^{-1/2}$. Therefore asymptotic inference, see Theorem 3, is only possible for $\delta > 1$ and $\zeta > 0$. Further data-driven bandwidths which are contained in such intervals with probability converging to 1 are valid in order to obtain the \sqrt{n} -rate.

Remark 6 (Interpolation estimators). Note that the interpolation estimator as suggested in Cai and Yuan (2011) for the dense regime has an error contribution of the form

$$\max_{\boldsymbol{j}} |\bar{\varepsilon}_{\boldsymbol{j}}|$$

which for independent and identically distributed Gaussian random errors is of order $\sqrt{\log p^1}/\sqrt{n}$. If the coordinates of p are polynomial in n this is of order $\sqrt{\log n}/\sqrt{n}$, thus slower than $1/\sqrt{n}$.

4 Asymptotic normality and local polynomial estimators

Asymptotic normality

In case the term $I_3^{n,p,h}$ in the decomposition (6) dominates, we can apply the functional central limit theorem to obtain asymptotic normality of the linear estimator in (5), thereby generalizing Degras (2011, Theorem 1) to multiple dimensions and general linear estimators. We require the following additional assumption on the process Z.

Assumption 4. The covariance kernel $\Gamma: [0,1]^d \times [0,1]^d \to \mathbb{R}$ of the process $Z: \Omega \times [0,1]^d \to \mathbb{R}$ belongs to the Hölder-class with smoothness $\gamma \in (0,1]$, that is $\Gamma \in \mathcal{H}_{[0,1]^{2d}}(\gamma, \tilde{L})$.

Theorem 3. In model (1) under Assumptions 1 and 4, consider the linear estimator $\hat{\mu}_{n,\boldsymbol{p},h}$ in (5) with weights satisfying Assumption 2 with $\zeta = \lfloor \alpha \rfloor$. If for some $\delta > 1$ we have that $\log(\boldsymbol{p}_{\min})^{\delta/d} / (\min((\boldsymbol{p}^1)^{1/d}, \boldsymbol{p}_{\min})) \lesssim n^{-\frac{1}{2\alpha}} \log(\boldsymbol{p}_{\min})^{-\delta}$, then for all sequences of smoothing parameters $h_n = h$ in $H_n := [c_1 \log(\boldsymbol{p}_{\min})^{\delta/d} / (\min((\boldsymbol{p}^1)^{1/d}, \boldsymbol{p}_{\min})), c_2 n^{-\frac{1}{2\alpha}} \log(\boldsymbol{p}_{\min})^{-\delta}]$ with constants $c_1, c_2 > 0$, we obtain the convergence in distribution

$$\sqrt{n} (\widehat{\mu}_{n,\boldsymbol{p},h} - \mu) \xrightarrow{D} \mathcal{G}(0,\Gamma),$$

where \mathcal{G} is a real-valued Gaussian process on $[0,1]^d$ with covariance kernel Γ of Z.

Remark 7. The assumption $\log(\boldsymbol{p}_{\min})^{\delta} / (\min((\boldsymbol{p}^1)^{1/d}, \boldsymbol{p}_{\min})) \leq n^{-\frac{1}{2\alpha}} \log(\boldsymbol{p}_{\min})^{-\delta}$ for a $\delta > 1$ guarantees that the intervals H_n are not empty for n being large enough. The rate of the upper bounds yields

$$h^{lpha} \lesssim n^{-1/2} \log(\boldsymbol{p}_{\min})^{-lpha \, \delta} \ll n^{-1/2}$$

Therefore the discretization error is negligible. Note that instead of $\log(p_{\min})^{-\delta}$ any factor that would make up for a strictly faster rate of the discretization error would be permissible. The rate of the lower bound guarantees

$$\left(\frac{\log(1/h)}{n\,\boldsymbol{p}^{1}h^{d}}\right)^{1/2} \lesssim \left(\frac{\log(\boldsymbol{p}_{\min})^{1-\delta}}{n\,\boldsymbol{p}^{1}\min(\boldsymbol{p}^{1},\boldsymbol{p}_{\min}^{d})^{-1}}\right)^{1/2} \lesssim \left(\frac{\log(\boldsymbol{p}_{\min})^{1-\delta}}{n}\right)^{1/2} \lesssim \frac{1}{\sqrt{n}}\,.$$

Here $\delta > 1$ is required.

Local polynomial estimators

We discuss local polynomial estimators and show that in case of design densities and mild assumptions on the kernel, the weights satisfy the properties in Assumption 2.

Assumption 5. Assume that there exist Lipschitz continuous densities $f_k: [0,1] \to \mathbb{R}$ bounded by $0 < f_{k\min} \leq f_k(t) \leq f_{k\max} < \infty, t \in [0,1]$ and $1 \leq k \leq d$, such that the design points $x_{k,l}, 1 \leq l \leq p_k$

$$\int_0^{x_{k,l}} f_k(t) \, \mathrm{d}t = \frac{l - 0.5}{p_k}, \qquad l = 1, \dots, p_k, \quad k = 1, \dots, d.$$

Lemma 7 in the Appendix A shows that Assumption 5 implies Assumption 3.

Let $N_{l,d} := \binom{d+l}{d}$ let $\psi_l: \{1, \ldots, N_{l,d-1}\} \to \{\mathbf{r} \in \{0, 1, \ldots, l\}^d \mid |\mathbf{r}| = l\}$ be an enumeration of the set $\{\mathbf{r} \in \{0, 1, \ldots, l\}^d \mid |\mathbf{r}| = l\}$ and let $U_m: \mathbb{R}^d \to \mathbb{R}^{N_{m,d}}$ be defined as $U_m(\mathbf{u}) := (1, P_{\psi_1(1)}(\mathbf{u}), \ldots, P_{\psi_1(d)}(\mathbf{u}), \ldots, P_{\psi_m(1)}(\mathbf{u}), \ldots, P_{\psi_m(N_{m,d-1})}(\mathbf{u}))$ for the monomial $P_{\psi_l(j)}(\mathbf{u}) := \frac{\mathbf{u}^{\psi_l(j)}}{\psi_l(j)!}$, for all $j = 1, \ldots, N_{l,d-1}$ and $l = 1, \ldots, m$. As a result of the properties of the binomial coefficient we note that $U_m(\mathbf{u}) \in \mathbb{R}^{N_{m,d}}$. Choose a non-negative kernel function $K: \mathbb{R}^d \to [0, \infty)$ and a bandwidth h > 0, set $K_h(\mathbf{x}) := K(\mathbf{x}/h)$ and $U_h(\mathbf{x}) = U_m(\mathbf{x}/h)$, and given an order $m \in \mathbb{N}$ define

$$\widehat{\boldsymbol{\vartheta}}_{n,\boldsymbol{p},h}(\boldsymbol{x}) := \operatorname*{argmin}_{\boldsymbol{\vartheta} \in \mathbb{R}^{N_{m,d}}} \sum_{\boldsymbol{j}=\boldsymbol{1}}^{\boldsymbol{p}} \left(\bar{Y}_{\boldsymbol{j}} - \boldsymbol{\vartheta}^{\top} U_{h}(\boldsymbol{x}_{\boldsymbol{j}} - \boldsymbol{x}) \right)^{2} K_{h}(\boldsymbol{x}_{\boldsymbol{j}} - \boldsymbol{x}), \quad \boldsymbol{x} \in T.$$
(12)

The local polynomial estimator of μ of order m at x is given as

$$\widehat{\boldsymbol{\mu}}_{n,\boldsymbol{p},h}^{\mathrm{LP},m}(\boldsymbol{x}) := \left(\widehat{\boldsymbol{\vartheta}}_{n,\boldsymbol{p},h}(\boldsymbol{x})\right)_{1}, \quad \boldsymbol{x} \in T.$$
(13)

Lemma 1. Let the kernel function K have compact support in $[-1,1]^d$, be Lipschitz continuous and satisfy

$$K_{\min}\mathbb{1}_{[-\boldsymbol{\Delta},\boldsymbol{\Delta}]}(\boldsymbol{u}) \le K(\boldsymbol{u}) \le K_{\max} \qquad \boldsymbol{u} \in \mathbb{R}^d,$$
(14)

for positive constants $\Delta, K_{\min}, K_{\max} > 0$, where $\mathbf{\Delta} = (\Delta, \dots, \Delta)^{\top} \in \mathbb{R}^d$. Then under the design Assumption 5, there exist $p_0 \in \mathbb{N}$ and $h_0, c > 0$ such that for $\mathbf{p} = (p_1, \dots, p_d)^{\top}$ with $\mathbf{p}_{\min} \geq p_0$ and $h \in (c/\mathbf{p}_{\min}, h_0]$, the local polynomial estimator is uniquely defined by (13), and is a linear estimator with weights satisfying Assumption 2 with $\gamma = m$.

The proof is given in Section A in the appendix.



Figure 1: Mean function μ_0 (solid line), n = 10 observational rows with p = 21 design points. Estimator based on all (dashed line) as well as on p = 11 observations (dotted line).

5 Simulations

We illustrate the rates in Theorem 1 in a simulation. We consider dimension d = 1, and for the mean function choose

$$\mu_0(x) = \sin(3\pi(2x-1))\exp(-2|2x-1|), \quad x \in [0,1].$$
(15)

We take independent standard Gaussian errors $\varepsilon_{i,j}$ and for the individual deviations we use independent Brownian motions. Simulations are performed in R, and the local polynomial estimator is computed using the R-package locpol with the function locPolWeights. Figure 1 contains plots of μ_0 , n = 10 observations at p = 21 design points as well as the local polynomial estimator of degree 2 based on all as well as only on p = 11 observations.

In the following, since we have synchronous design points we directly sample the averaged errors and processes, resulting in $\bar{\varepsilon}_{j,n} \sim \mathcal{N}(0, 1/n)$ and for the individual deviation $\bar{Z}_{j,n} \sim n^{-1/2}B(t), t \in [0, 1]$ for a Brownian motion $(B(t))_{t \in [0, 1]}$.

To determine the sensitivity on the bandwidth, in Figure 2, for sample size n = 600 and number of design points $p \in \{65, 115, 175, 275, 400, 550\}$, based on N = 1000 iterations we simulate the supremum norm errors on a grid of bandwidths of step size 0.005, starting at 3/p. Again we use the local polynomial estimator of degree 2 and complement the results with an interpolation estimator. For large but also very small bandwidths - depending on p- the estimation error in the supremum norm starts again to increase.

Figure 3 contains boxplots of bandwidths selected by *leave-one-curve-out cross validation*. Comparing the values to Figure 2 we see that overall reasonable values area selected, and in the majority of cases the resulting bandwidth seems to result in minimal overall supremum norm error.



Figure 2: Sup-norm error for n = 600 and various sizes of p for bandwidths on a 0.005 grid starting at 3/p, as well as error of interpolation estimator. The values with sup.err = 0 indicate the bandwidth chosen by leaveone-curve-out cross validation.



Figure 3: Boxplots of N = 200 iterations of chosen bandwidths chosen by *leave-one-curve-out cross validation* in the setup of Figure 2.

Figure 4 contains the sup-norm error of the local polynomial estimator with optimally tuned bandwidth, of a penalized splines estimator computed using the function $mooth_spline$ from package stats and optimally tuned smoothing parameter, as well as an interpolation estimator for n = 600 and a range of values of p. We again use N = 1000 repetitions to determine the expected sup-norm error. While for local polynomial and smoothing splines the sup-norm error first decreases and then stabilizes for increasing p, for the interpolation estimator it actually increases with increasing number of design points p. Taking a closer look at the contributions of the terms in the error decomposition in (6) in Figure 5, we observe



Figure 4: Effect of increasing p on overall sup-norm error for local polynomial estimator (green dotted line), smoothing spline (blue dashed line) and interpolation estimator (solid red line).

Figure 5: Effect of increasing p on contribution of bias (red curves), process terms Z (blue curves) as well as errors (green curves) on the the sup-norm error of the three methods.

that while the contributions from the processes B_i are comparable for all three methods, and the bias of the interpolation estimators is quite small, the contribution from the observation errors increases for interpolation as also observed in Remark 6.

6 Real data illustrations

In this section we investigate the effect of a coarser discretization in two data examples. Smaller values of p result in fewer data points, which may be advantageous for storage and also for energy efficiency in the measurement process, as long as estimation quality is not too adversely affected.

To quantify the effect on the sup-norm error, we proceed as follows. We use the full data set to compute the estimator as well as the fast and fair confidence bands proposed in Liebl and Reimherr (2023). Here we use the package ffscb from Liebl and Reimherr (2023) with two intervals in the time spans in our examples. Then we compute the estimator for coarser discretizations, that is smaller values of p using only subsets of the full data, and investigate whether the resulting estimator is still fully contained in the confidence bands.

6.1 Temperature series

First we consider daily temperatures (per month) for the years 2000 until 2022 in Nuremberg, Germany. The data were obtained from the *Deutschen Wetter Dienst (DWD)* at [Link]. To remove serial dependence, only the 1, 4, 8, 12, 15, 18, 22, 25 and 29th days of each month were used. Data are recorded every 10 minutes, which results in p = 144. Some NA's had to be removed which results in varying n for the different months, see Table 1.

Figure 6 contains plots of the observed daily temperature curves of the 1, 8, 15, 22 and 29th day of the month, together with a local polynomial fit. The bandwidth is chosen by leave-one-curve-out cross validation, the results are displayed in Table 1.

Month	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
n	178	165	178	178	175	187	184	186	184	184	183	188
h	72.0	72.0	58.4	64.8	43.2	64.8	57.6	57.6	57.6	57.6	136.8	43.2

Table 1: Number of observations n and bandwidth h for each month.

Next, we use the full data to compute the fast and fair confidence bands proposed in Liebl and Reimherr (2023) with two time intervals. Finally, we compute the estimator for a coarser discretization for one- and two hour intervals, and investigate whether these curves are still contained in the confidence bands. The results for January and August are displayed in Figure 7. While in January, both curves are fully contained in the confidence band, this is not the case in August due to higher temperature variations in day- and night times. The corresponding plots for all months are given in Figure 10 in the appendix.



Figure 6: Daily temperature curves in Nuremberg, Germany, per month from 2000 until 2022, together with a local polynomial estimate of the mean curve (yellow curve).



Figure 7: LP estimator with p = 144 (10-minute data, red line), p = 24 (one hour data, blue dotted/dashed line) and p = 12 (two hour data, green dashed line). Confidence bands for the p = 144 curve from Liebl and Reimherr (2023) for January and August.





Figure 8: Curves of pairwise differences of torque curves with different shoes for each runner, together with an local polynomial estimate of the mean function (solid blue curve)

Figure 9: Estimated mean difference curves using p = 200 (solid red), p = 50 (dotted green) and p = 20 (dashed blue) design points, and a uniform confidence band computed from the full data

6.2 Biomechanics

Second we revisit a data set of a sports biomechanics experiment from Liebl and Reimherr (2023). It contains n = 18 pairs of torque curves in Newton metres (N/m) standardized by the bodyweight (kg) for runners who wear first, extra cushioned running shoes and second, certain normal cushioned running shoes. The time scale is normalized to $t \in [0, 100]$ with p = 200 grid points. See Liebl and Reimherr (2023) for further details. Figure 8 contains the curves of pairwise difference for each runner together with a local polynomial estimate of the corresponding mean curve. The bandwidth is selected by leave-one-curve-out cross validation, resulting in h = 5.5.

Figure 9 contains the confidence band computed with the methods from Liebl and Reimherr (2023) using two intervals in the time span, together with mean curve estimates based on p = 50 as well as p = 20 data points in each row. While the fit with p = 50 is hardly distinguishable from that based on the full data (p = 200), the estimate based on only p = 20 is outside the confidence band in the critical starting phase at about t = 6.

7 Discussion

Let us indicate how to extend our results to more general design assumptions. First, we note that if the design satisfies Assumption 3 and the weights satisfy Assumption 2, the upper bound (7) in Theorem 1 follows. Here p^1 is interpreted as the total number of design points. While the range of admissible bandwidths has to be adapted according to the design under investigation, no Cartesian product structure is required.

Further, the upper bounds in Theorem 1 and even the CLT in Theorem 3 can be extended to asynchronous design (e.g. design depending on the observation row i) if the number of points in each row is of the same order. Here we use the linear estimator (5) in each row,

$$\widehat{\mu}_{n,\boldsymbol{p},h,i}(\boldsymbol{x}) = \sum_{\boldsymbol{j}=1}^{\boldsymbol{p}_i} w_{\boldsymbol{j},\boldsymbol{p}_i,i}(\boldsymbol{x};h;\boldsymbol{x}_{1,i},\ldots,\boldsymbol{x}_{\boldsymbol{p}_i,i}) Y_{\boldsymbol{j}}, \qquad (16)$$

and average to obtain the estimator of μ , potentially using a weighted average (Zhang and Wang, 2016). If the weights $w_{j,p_i,i}$ in each row satisfy Assumption 2, with constants $C_1, C_2 > 0$ and Assumption 3 with $C_3 > 0$, which may be chosen independently of *i*, we may obtain results similar to Theorem 1 and Theorem 3, where the $n^{-1/2}$ - rate in the dense setting in Theorem 1 follows from the more elaborate manageability argument used in the proof of Theorem 3.

The smooth first then average approach discussed above for the asynchronous setting cannot work in the truly sparse case, where the coordinates of p remain bounded. Here an alternative is pooling, that is, forming a single linear estimator from all observations potentially with weighting. This has been worked out in detail in (Li and Hsing, 2010; Zhang and Wang, 2016). In the dense case, however, only a $\log n/\sqrt{n}$ - rate is obtained. Thus, the parametric rate together with the central limit theorem in supremum norm for dense designs seem not to be developed for pooling estimators yet.

8 Proofs

8.1 Proof of Theorem 1

From the error decomposition (6) we obtain the bound

$$\sup_{\mu \in \mathcal{H}(\alpha,L), \ Z \in \mathcal{P}_{T}(\beta_{0},C_{Z})} \mathbb{E}_{\mu} \left[\left\| \widehat{\mu}_{n,\boldsymbol{p},h} - \mu \right\|_{\infty} \right]$$

$$\leq \sup_{\mu \in \mathcal{H}(\alpha,L)} \left\| I_{1}^{\boldsymbol{p},h} \right\|_{\infty} + \mathbb{E} \left[\left\| I_{2}^{n,\boldsymbol{p},h} \right\|_{\infty} \right] + \sup_{Z \in \mathcal{P}_{T}(\beta_{0},C_{Z})} \mathbb{E} \left[\left\| I_{3}^{n,\boldsymbol{p},h} \right\|_{\infty} \right].$$
(17)

Theorem 1 follows from (17) together with Lemma 2 below. Note that property (W3) in Assumption 2 together with Assumption 3 imply the following.

(W5) We have
$$\sum_{1 \leq j \leq p} |w_{j,p}(\boldsymbol{x};h)| \leq C_4, \, \boldsymbol{x} \in T.$$

Lemma 2. In model (1) under Assumption 1, consider the linear estimator $\hat{\mu}_{n,\mathbf{p},h}$ in (5) with error decomposition in (17).

i) If the weights satisfy (W1) with $\gamma = |\alpha|$, (W2) of Assumption 2 as well as (W5), then

$$\sup_{\mu \in \mathcal{H}(\alpha,L)} \left\| I_1^{\boldsymbol{p},h} \right\|_{\infty} = \mathcal{O}\left(h^{\alpha}\right).$$

ii) If the weights satisfy (W3), (W4) of Assumption 2 and Assumption 3, then

$$\mathbb{E}\left[\left\|I_{2}^{n,\boldsymbol{p},h}\right\|_{\infty}\right] = \mathcal{O}\left(\sqrt{\frac{\log(h^{-1})}{n\boldsymbol{p}^{1}h^{d}}}\right)$$

iii) If the weights satisfy (W5) then

$$\limsup_{n, \boldsymbol{p} \to \infty} \sup_{h \in (\boldsymbol{c}/\boldsymbol{p}, h_0]} \sup_{Z \in \mathcal{P}_T(\beta_0, C_Z)} \mathbb{E} \Big[\left\| \sqrt{n} I_3^{n, \boldsymbol{p}, h} \right\|_{\infty}^2 \Big] < \infty \,.$$

In particular,

$$\left\|I_{3}^{n,\boldsymbol{p},h}\right\|_{\infty}=\mathcal{O}_{\mathbb{P},h\in(c/\boldsymbol{p},h_{0}]}\left(n^{-\frac{1}{2}}\right).$$

Proof of Lemma 2. i): This is a well-known result in nonparametric curve estimation by using Taylor expansion and the properties of the weights. The proof can be found in the Appendix, Section B.

iii): For processes Z in the class $\mathcal{P}_T(\beta_0, C_Z)$, we have that $\frac{1}{\sqrt{n}} |\sum_{i=1}^n Z_i(\boldsymbol{x})|$ is uniformly bounded in expected value. For example, by using the inequality (25) in the appendix, which is Pollard (1990, Section 7, display (7.10)), we explicitly have that

$$\mathbb{E}\Big[\sup_{\boldsymbol{x}\in[0,1]^d} \frac{1}{\sqrt{n}} \left|\sum_{i=1}^n Z_i(\boldsymbol{x})\right|\Big] \le \operatorname{const}\left(C_Z/\beta_0\right)^{1/2},\tag{18}$$

see Lemma 5 below. Therefore, by using property (W5) of the weights we obtain

$$\mathbb{E}\left[\left\|\sqrt{n} I_{3}^{n,\boldsymbol{p},\boldsymbol{h}}\right\|_{\infty}\right] = \mathbb{E}\left[\sup_{\boldsymbol{x}\in[0,1]^{d}}\sqrt{n}\left|\sum_{\boldsymbol{j=1}}^{\boldsymbol{p}} w_{\boldsymbol{j},\boldsymbol{p}}(\boldsymbol{x};\boldsymbol{h})\bar{Z}_{n}(\boldsymbol{x}_{\boldsymbol{j}})\right|\right]$$
$$\leq C_{4} \mathbb{E}\left[\sup_{\boldsymbol{x}\in[0,1]^{d}}\left|\frac{1}{\sqrt{n}}\sum_{i=1}^{n} Z_{i}(\boldsymbol{x})\right|\right] \qquad (by (W5))$$
$$\leq \operatorname{const}\left(C_{Z}/\beta_{0}\right)^{1/2} < \infty. \qquad (by (18))$$

ii): We shall apply Dudley's entropy bound (van der Vaart and Wellner, 1996, Corollary 8). To this end, note that by Assumption 1,
$$\bar{\varepsilon}_1, \ldots, \bar{\varepsilon}_n$$
 are independent, centered, sub-

2.2.8). To this end, note that by Assumption 1, $\bar{\varepsilon}_1, \ldots, \bar{\varepsilon}_p$ are independent, centered, sub-Gaussian random variables with parameters bounded by $\zeta^2 \sigma^2/n > 0$ and $\mathbb{E}[\bar{\varepsilon}_{j,n}^2] = \sigma_j^2/n$, $j = 1, \ldots, p$. Therefore, the process

$$S_{n,\boldsymbol{p},h}(\boldsymbol{x}) = \sqrt{n \boldsymbol{p}^{\mathbf{1}} h^d} I_2^{n,\boldsymbol{p},h}(\boldsymbol{x})$$

is a sub-Gaussian process w.r.t. the semimetric

$$d_{S}^{2}(\boldsymbol{x},\boldsymbol{y}) := \zeta^{2} \sigma^{2} \boldsymbol{p}^{1} h^{d} \sum_{\boldsymbol{j=1}}^{\boldsymbol{p}} \left(w_{\boldsymbol{j},\boldsymbol{p}}(\boldsymbol{x};h) - w_{\boldsymbol{j},\boldsymbol{p}}(\boldsymbol{y};h) \right)^{2}$$
$$\leq M^{2} \left(\frac{\|\boldsymbol{x} - \boldsymbol{y}\|_{\infty}}{h} \wedge 1 \right)^{2} \leq M^{2}, \qquad (19)$$

where $M = \sqrt{2C_3}C_2 \zeta \sigma$, and we used the Lipschitz continuity of the weights, (W4). Now, given $0 < \delta < M$ to bound the covering number $N([0,1]^d, \delta; d_S)$ and hence the packing number $D([0,1]^d, \delta; d_S)$ of $[0,1]^d$ w.r.t. d_S , note that from (19),

$$\mathrm{d}_S(\boldsymbol{x}, \boldsymbol{ au}) \leq \delta \qquad \Longrightarrow \qquad \left\| \boldsymbol{ au}_j - \boldsymbol{x} \right\|_\infty \leq \frac{\delta h}{M}, \qquad \boldsymbol{x}, \boldsymbol{ au} \in [0, 1]^d.$$

We obtain

$$D([0,1]^d,\delta;\mathbf{d}_S) \le N([0,1]^d,\delta/2;\mathbf{d}_S) \le \left(\frac{2M}{\delta h}\right)^d.$$
(20)

By observing from (19) that the diameter of $[0, 1]^d$ under d_S is upper bounded by 2M, Dudley's entropy bound implies that

$$\mathbb{E}\Big[\sup_{\boldsymbol{x}\in[0,1]^d} \left|S_{n,\boldsymbol{p},h}(\boldsymbol{x})\right|\Big] \le \mathbb{E}\Big[\left|S_{n,\boldsymbol{p},h}(\boldsymbol{x}_0)\right|\Big] + C_4 \int_0^{2M} \sqrt{\log\left(D\big([0,1]^d,\delta;\mathbf{d}_S\big)\right)} \,\mathrm{d}\delta \tag{21}$$

for fixed $x_0 \in [0,1]^d$ and a universal constant $C_4 > 0$. Using (20), the integral in the second term is bounded by

$$\int_{0}^{2M} \sqrt{\log\left((2M/(\delta h)^{d})\right)} d\delta = \frac{2M d}{h} \int_{0}^{h} \sqrt{\log(\delta^{-1})} d\delta$$
$$\leq 2M d\left(\sqrt{-\log(h)} - \frac{1}{2\sqrt{-\log(h)}}\right) = \mathcal{O}\left(\sqrt{-\log(h)}\right).$$

For the first term in (21), apply Jensen's inequality to obtain

$$\mathbb{E}\Big[\big|S_{\boldsymbol{p}}(\boldsymbol{x}_{0})\big|\Big]^{2} \leq \mathbb{E}\Big[\widetilde{S}_{\boldsymbol{p}}^{2}(\boldsymbol{x}_{0})\Big] \leq \sigma^{2}\boldsymbol{p}^{1}h^{d}\sum_{\boldsymbol{j}=\boldsymbol{1}}^{\boldsymbol{p}}w_{\boldsymbol{j},\boldsymbol{p}}(\boldsymbol{x}_{0};h)^{2} \leq \sigma^{2}C_{4}C_{1} < \infty$$

by (W3) and (W5). This concludes the proof of the theorem.

8.2 Proof of Theorem 2 (optimality)

Proof of Theorem 2. In the proof we rely on the reduction to hypothesis testing as presented e.g. in Tsybakov (2004, section 2).

For the lower bound $p_{\min}^{-\alpha}$, using the method of two sequences of hypotheses functions we set $\mu_0 = 0$ and construct $\mu_1 = \mu_{1,p}$ such that $\|\mu_0 - \mu_{1,p}\|_{\infty} \ge c p_{\min}^{-\alpha}$ for some constant c > 0 and that $\mu_{1,p}(\boldsymbol{x_j}) = 0$ at all design points $\boldsymbol{x_j}$, so that the distribution of the observations for μ_0 and μ_1 coincide.

If r is such that $p_r = \mathbf{p}_{\min}$, observing Assumption 5 for a constant $\tilde{L} > 0$ to be specified we set

$$g_{\boldsymbol{p}}(\boldsymbol{x}) = g(\boldsymbol{x}) = \tilde{L}\left(\frac{1}{p_r f_{\max r}}\right)^{\alpha} \exp\left(-\frac{1}{1-x_r^2}\right) \mathbb{1}_{\{|x_r|<1\}}, \qquad \boldsymbol{x} = (x_1, \dots, x_d),$$

and for some fixed $1 \le l \le p_r - 1$ let

$$\mu_1(\boldsymbol{x}) = g((2p_r f_{\max r}) (x_r - (x_{r,l} + x_{r,l+1}) / 2)).$$

For the design points $x_{r,l}$ in the r^{th} coordinate, by Assumption 5 the distance is at least $p_r f_{\max r}$, and therefore by the definition of g and the Cartesian product structure of the design it follows that $\mu_{1,p}(x_j) = 0$ at all design points. Further, if the r^{th} coordinate is $(x_{r,l} + x_{r,l+1})/2$, $\mu_{1,p}$ takes the value $p_r^{-\alpha} \tilde{L} e^{-1}/f_{\max r}^{\alpha}$, so that $\|\mu_0 - \mu_{1,p}\|_{\infty} \ge c p_{\min}^{-\alpha}$ holds true. Finally, using the chain rule and the fact that all derivatives of the bump function in the

definition of g are uniformly bounded, μ_1 is α - Hölder smooth (indeed all partial derivatives except in direction r vanish) with constant proportional to \tilde{L} , which can be adjusted to yield the Hölder norm L. This concludes the proof for the lower bound $p_{\min}^{-\alpha}$.

Let us turn to the lower bound of order $(\log(n p^1)/(np^1))^{\alpha/(2\alpha+d)}$, which corresponds to the minimax rate in d dimensions in the conventional nonparametric regression model with np^1 design points. Taking Z = 0 ($\in \mathcal{P}_T(\beta_0, C_Z)$) we can adapt the argument in Tsybakov (2004, Theorem 2.10). Indeed, by sufficiency and taking averages using the normal distribution we could reduce the analysis to a single row with errors distributed as $\mathcal{N}(0, \sigma_0^2/n)$. For convenience we provide the details, and do not use sufficiency, so that the argument can be extended to further error distributions (Tsybakov, 2004, Assumption B, Section 2.3). For sufficiently small c_i , i = 0, 1 to be specified set

$$N_{n,\boldsymbol{p}} = \left\lceil c_0 \left(\frac{n \boldsymbol{p}^1}{\log(n \boldsymbol{p}^1)} \right)^{\frac{1}{2\alpha+d}} \right\rceil, \qquad h_{n,\boldsymbol{p}} = N_{n,\boldsymbol{p}}^{-1}, \qquad s_{n,\boldsymbol{p}} := c_1 h_{n,\boldsymbol{p}}^{\alpha}.$$

For \tilde{L} to be specified let

$$\tilde{g}(\boldsymbol{x}) = \tilde{L}h_{n,\boldsymbol{p}}^{\alpha} \exp\left(-\left(1 - \|\boldsymbol{x}\|_{2}^{2}\right)^{-1}\right) \mathbb{1}_{\{\|\boldsymbol{x}\|_{2} < 1\}}$$

and for

$$\boldsymbol{l} = (l_1, \dots, l_d) \in \{1, \dots, N_{n, \boldsymbol{p}}\}^d$$

 set

$$\mu_{l}(\boldsymbol{x}) = \tilde{g}(2(\boldsymbol{x} - \boldsymbol{z}_{l})/h_{n,\boldsymbol{p}}), \qquad \boldsymbol{z}_{l} = (l_{1} - 1/2, \dots, l_{d} - 1/2)/N_{n,\boldsymbol{p}}.$$

Using the chain rule one checks that $\mu_{l} \in \mathcal{H}(\alpha, L)$ for suitable choice of \tilde{L} depending on Land c_{0} . Further, by construction the μ_{l} have disjoint supports in $[0, 1]^{d}$, so that $\|\mu_{l} - \mu_{r}\|_{\infty} \geq 2 s_{n,p}$ for all $l \neq r$, and c_{1} sufficiently small depending on c_{0} .

To apply Tsybakov (2004, Proposition 2.3) and thus to obtain $s_{n,p}$ as lower bound for the rate of convergence, it remains to show for a $\kappa \in (0, 1/8)$ and n and p large enough that

$$\frac{1}{N_{n,\boldsymbol{p}}^{d}} \sum_{\boldsymbol{l}} \operatorname{KL}(\mathbb{P}_{\mu_{\boldsymbol{l}}}, \mathbb{P}_{0}) \leq d \kappa \log(N_{n,\boldsymbol{p}}) \quad \text{for each } \boldsymbol{l},$$
(22)

where \mathbb{P}_{μ_l} is the (joint normal) distribution of the observations for the mean function μ_l . To this end, since Kullback-Leibler divergence for product measures is additive, we can estimate

$$\frac{1}{N_{n,\boldsymbol{p}}^{d}} \sum_{\boldsymbol{l}} \operatorname{KL}(\mathbb{P}_{\mu_{\boldsymbol{l}}}, \mathbb{P}_{0}) = \frac{1}{N_{n,\boldsymbol{p}}^{d}} \frac{n}{2\sigma_{0}^{2}} \sum_{\boldsymbol{l}} \sum_{\boldsymbol{l} \leq \boldsymbol{j} \leq \boldsymbol{p}} \mu_{\boldsymbol{l}}^{2}(\boldsymbol{x}_{\boldsymbol{j}}) \qquad \text{(normal distr.)}$$

$$\leq \frac{n}{2\sigma_{0}^{2}} \frac{\tilde{L}^{2}}{e^{2}} h_{n,\boldsymbol{p}}^{2\alpha+d} \sum_{\boldsymbol{l} \leq \boldsymbol{j} \leq \boldsymbol{p}} \sum_{\boldsymbol{l}} \mathbbm{1}_{\{2 \| \boldsymbol{x}_{\boldsymbol{j}} - \boldsymbol{z}_{\boldsymbol{l}} \|_{2} \leq h_{n,\boldsymbol{p}}\}} \qquad \text{(constr. of } \mu_{\boldsymbol{l}})$$

$$\leq \frac{n}{2\sigma_{0}^{2}} \frac{\tilde{L}^{2}}{e^{2}} \frac{h_{n,\boldsymbol{p}}^{2\alpha+d}}{2^{2\alpha-d}} \boldsymbol{p}^{1}$$

$$\leq \text{const. } \log(n \, \boldsymbol{p}^{1}), \qquad \text{(choice of } h_{n,\boldsymbol{p}} = N_{n,\boldsymbol{p}}^{-1})$$

which implies (22) and completes the proof for the lower bound $(\log(n p^1)/(np^1))^{\alpha/(2\alpha+d)}$.

Finally for the lower bound of order $n^{-1/2}$ we may follow the argument in Cai and Yuan (2011): Take μ to be a constant function, and Z_i to be a random constant, which is centered normally distributed with variance $\leq C_Z$. Since the Kullback - divergence between two normal distributions with different mean and equal variance only decreases under convolution with a further centered normal distribution, it suffices to obtain the lower bound $n^{-1/2}$ in the model without errors $\varepsilon_{i,j}$. But then it follows as the known rate of convergence for estimating a normal mean in a sample of size n. This concludes the proof of Theorem 2.

8.3 Manageability and a maximal inequality from Pollard (1990)

We recall the concept of manageability of a triangular array of processes from Pollard (1990, Section 7), which we require to prove (18) as well as Theorem 3.

For a bounded subset $B \subset \mathbb{R}^n$, $n \in \mathbb{N}$, we call a vector $\mathbf{\Phi} = (\Phi_1, \dots, \Phi_n)^\top \in \mathbb{R}^n$ an envelope for B if $|b_i| \leq \Phi_i$, for all $\mathbf{b} = (b_1, \dots, b_n)^\top \in B$. For a triangular array $(X_{n,i}(x))_{n \in \mathbb{N}, 1 \leq i \leq k_n} : \Omega \times [0, 1]^d \to \mathbb{R}$ of stochastic processes with independent rows we call the sequence $\mathbf{\Phi}_n$ of k_n dimensional random vectors a sequence of envelopes of $(X_{n,i})$ if $\mathbf{\Phi}_n(\omega)$ is an envelope of the set $F_n(\omega) := \{(X_{n,1}(\omega, \mathbf{x}), \dots, X_{n,k_n}(\omega, \mathbf{x}))^\top \mid \mathbf{x} \in [0, 1]^d\}$ for $n \in \mathbb{N}$ and $\omega \in \Omega$.

Further we call a triangular array of processes $(X_{n,i}(\boldsymbol{x})), i = 1, \ldots, k_n, \boldsymbol{x} \in [0, 1]^d$ manageable (with respect to its envelope Φ_n) (Pollard, 1990, see Definition 7.9, p. 38) if there exists a deterministic function $\lambda \colon [0, 1] \to \mathbb{R}$, the capacity bound, for which

- i) $\int_0^1 \sqrt{\log(\lambda(x))} \mathrm{d}\varepsilon < \infty$,
- ii) $D(\varepsilon \| \boldsymbol{\alpha} \circ \Phi_n(\omega) \|_2, \boldsymbol{\alpha} \circ F_n(\omega)) \leq \lambda(\varepsilon)$ pointwise for $\varepsilon \in [0, 1]$ and all $\boldsymbol{\alpha} \in \mathbb{R}^n_+$, $n \in \mathbb{N}$ and all $\omega \in \Omega$

where $D(\varepsilon \| \boldsymbol{\alpha} \circ \Phi_n(\omega) \|_2, \boldsymbol{\alpha} \circ F_n(\omega))$ is the packing number and $\boldsymbol{x} \circ \boldsymbol{y}$ denotes the Hadamard (component-wise) product of two vectors of equal dimension. A sequence (X_i) of processes manageable if the array defined by $X_{n,i} = X_i$ for $i \leq n$ is manageable. To check manageability we shall use the following Lemma.

Lemma 3. Consider the triangular array $(X_{n,i})_{n \in \mathbb{N}, 1 \leq i \leq k_n}$ of stochastic processes on $[0,1]^d$ with a suiting sequence of envelopes $(\mathbf{\Phi}_n)_{n \in \mathbb{N}}$. If there are constants $K_1, K_2, \kappa \in \mathbb{R}^+$ such that for all $\mathbf{x}, \mathbf{x}' \in [0,1]^d, n \in \mathbb{N}$ and $\epsilon > 0$ it holds that

$$\|\boldsymbol{x} - \boldsymbol{x}'\|_{2} \le K_{1} \epsilon^{\kappa} \Rightarrow \left|X_{n,i}(\boldsymbol{x}) - X_{n,i}(\boldsymbol{x}')\right| \le \epsilon K_{2} \Phi_{n,i},$$
(23)

where $i = 1, ..., k_n$, and $\mathbf{\Phi}_n = (\Phi_{n,1}, ..., \Phi_{n,k_n})^{\top}$, then we have for the covering number that

$$N(\epsilon \|\boldsymbol{\alpha} \circ \boldsymbol{\Phi}_n\|_2, \boldsymbol{\alpha} \circ F_n) \le \left(\frac{K_2^{\kappa}}{2K_1}\epsilon^{-\kappa} + 2\right)^d =: \lambda_{\kappa}(\varepsilon)$$

Further it holds that

$$\Lambda_{\kappa} := \int_0^1 \sqrt{\log(\lambda_{\kappa}(\varepsilon))} \,\mathrm{d}\epsilon < \infty \tag{24}$$

Proof of Lemma 3. Let $\tau_1, \ldots, \tau_N \in [0, 1]^d$ be such that for $\boldsymbol{x} \in [0, 1]^d$ there exists τ_m such that $\|\boldsymbol{x} - \tau_m\|_2^2 \leq \frac{K_1}{K_2^\kappa} \epsilon^{\kappa}$. For such \boldsymbol{x} and τ_m , from (23) we obtain

 $|X_{n,i}(x) - X_{n,i}(\tau_m)| \le \epsilon \Phi_n \quad \text{and hence} \quad \|\alpha \circ X_n(x) - \alpha \circ X_{n,i}(\tau_m)\|_2 \le \epsilon \|\alpha \circ \Phi_n\|_2$

for all $\alpha \in \mathbb{R}^d$. Therefore the amount of balls with radius $\epsilon \|\alpha \circ \Phi_n\|_2$ at points $T_m = \alpha \circ X_n(\tau_m)$ needed to cover $\alpha \circ F_n$ is upper bounded by the number of balls with radius $K_1 \epsilon^{\kappa} K_2^{-\kappa}$ needed to cover $[0, 1]^d$, which in turn is upper bounded by

$$\lambda_{\kappa}^{d}(\epsilon) := \left(\frac{K_{2}^{\kappa}}{2K_{1}\epsilon^{\kappa}} + 2\right)^{d}.$$

The integral in (24) is finite since

$$\begin{split} \int_{0}^{1} \sqrt{\log(\lambda_{\kappa}^{d}(\epsilon))} \mathrm{d}\epsilon &\leq \sqrt{d} \int_{0}^{1} \left(\sqrt{\log\left(\frac{K_{2}^{\kappa}}{2 K_{1}}\right)} + \sqrt{-\kappa \log(\epsilon)} + \sqrt{\log\left(1 + \frac{K_{1}}{K_{2}^{\kappa}}\right)} \right) \mathrm{d}\epsilon \\ &\leq \sqrt{\frac{d \kappa \pi}{2}} + \mathrm{const} < \infty \,. \end{split}$$

Let us state the maximal inequality Pollard (1990, Section 7, display (7.10)).

Lemma 4. Let $(X_{n,i})$ be a triangular array of stochastic processes with a suitable sequence of envelopes (Φ_n) . Further let $\lambda : [0,1] \to \mathbb{R}$ be the capacity bound such that $(X_{n,i})$ is manageable with respect to (Φ_n) and λ . We write

$$S_n(\boldsymbol{x}) := \sum_{i=1}^{k_n} X_{n,i}(\boldsymbol{x}) \qquad and \qquad M_n(\boldsymbol{x}) := \mathbb{E}\big[S_n(\boldsymbol{x})\big],$$

then for all $1 \leq p < \infty$ there exists a constant $\tilde{C}_p > 0$ such that

$$\mathbb{E}\left[\|S_n - M_n\|_{\infty}^p\right] \le \tilde{C}_p\left(\int_0^1 \sqrt{\log(\lambda(\varepsilon))} \mathrm{d}\varepsilon\right)^p \mathbb{E}\left[\|\boldsymbol{\Phi}_n\|_2^p\right].$$
(25)

From Lemmas 3 and 4 we may conclude the bound (18), which we state in the following lemma.

Lemma 5. For Z, Z_1, \ldots, Z_n i.i.d. stochastic processes on $[0,1]^d$ with $Z \in \mathcal{P}_T(\beta_0, C_Z)$ we have the bound

$$\mathbb{E}\Big[\sup_{\boldsymbol{x}\in[0,1]^d}\frac{1}{\sqrt{n}}\left|\sum_{i=1}^n Z_i(\boldsymbol{x})\right|\Big] \le \left(2\,\tilde{C}_2\,C_Z\,\Lambda_{1/\beta_0}\right)^{1/2} \tag{26}$$

where Λ_{1/β_0} is given in (24), and \tilde{C}_2 in (25).

Proof. From the Hölder continuity of the paths (3), Lemma 3 implies manageability of each process $n^{-1/2}Z_i$ w.r.t. to the envelope $n^{-1/2}(M_i + |Z_i(\mathbf{0})|), i = 1, ..., n$, where we may take $K_1 = K_2 = 1$ and $\kappa = 1/\beta_0$ uniformly over $\mathcal{P}_T(\beta_0, C_Z)$. The bound follows from Jensen's inequality and (25) with p = 2.

8.4 Proof of Theorem 3 (asymptotic normality)

Proof of Theorem 3. In view of the error decomposition (6), Lemma 1 together with the assumptions on p and n and the choice of the smoothing parameter h imply

$$\sup_{\mu \in \mathcal{H}(\alpha,L)} \sqrt{n} \left\| I_1^{\boldsymbol{p},h} \right\|_{\infty} \to 0 \quad \text{and} \quad \sqrt{n} \left\| I_2^{n,\boldsymbol{p},h} \right\|_{\infty} \xrightarrow{\mathbb{P}} 0$$

We shall apply the functional central limit Theorem in Pollard (1990, Theorem (10.6)) to the triangular array of processes $(X_{ni})_{n \in \mathbb{N}, 1 \leq i \leq n}$ given by

$$X_{ni}(x) = \frac{1}{\sqrt{n}} \sum_{j=1}^{p} w_{j,p}(x;h) Z_i(x_j) \text{ and } S_n(x) := \sum_{i=1}^{n} X_{ni}(x), \quad x \in [0,1]^d,$$

to obtain

$$\sqrt{n} I_3^{n,\boldsymbol{p},h} = S_n(\boldsymbol{x}) \stackrel{D}{\longrightarrow} \mathcal{G}(\boldsymbol{0},c) \,. \tag{27}$$

With Slutsky's Theorem (van der Vaart and Wellner, 1996, Example 1.4.7) we then obtain the assertion of Theorem 3.

For (27) we have to check the conditions i) - v) of Pollard (1990, Theorem (10.6)).

i). We show manageability of the processes $X_n = (X_{n1}, \ldots, X_{nn})$ with respect to $\Phi_n := (\Phi_{n1}, \ldots, \Phi_{nn})$, with

$$\Phi_{ni} := \frac{C_4}{\sqrt{n}} \left(|Z_i(\mathbf{0})| + M_i \right),$$

where the random variables M_i are given (3). We apply Lemma 3 and distinguish the cases $\epsilon > h^{\beta}$ and $\epsilon \le h^{\beta}$. For $\epsilon > h^{\beta}$, using (W1) we get for $\|\boldsymbol{x} - \boldsymbol{y}\|_2 \le \epsilon^{1/\beta}$ that

$$\begin{split} \sqrt{n} |X_{ni}(\boldsymbol{x}) - X_{ni}(\boldsymbol{y})| &\leq \bigg| \sum_{\boldsymbol{j=1}}^{\boldsymbol{p}} w_{\boldsymbol{j},\boldsymbol{p}}(\boldsymbol{x};h) \big(Z_i(\boldsymbol{x}_{\boldsymbol{j}}) - Z_i(\boldsymbol{x}) \big) \bigg| + \big| Z_i(\boldsymbol{x}) - Z_i(\boldsymbol{y}) \big| \\ &+ \bigg| \sum_{\boldsymbol{j=1}}^{\boldsymbol{p}} w_{\boldsymbol{j},\boldsymbol{p}}(\boldsymbol{y};h) \big(Z_i(\boldsymbol{y}) - Z_i(\boldsymbol{x}_{\boldsymbol{j}}) \bigg|. \end{split}$$

The second term is bounded by

$$|Z_i(\boldsymbol{x}) - Z_i(\boldsymbol{y})| \le M_i \|\boldsymbol{x} - \boldsymbol{y}\|_{\infty}^{\beta} \le M_i \|\boldsymbol{x} - \boldsymbol{y}\|_2^{\beta} \le M_i \epsilon$$

and the first and third terms by

$$\left|\sum_{j=1}^{P} w_{j,p}(\boldsymbol{x};h) \left(Z_{i}(\boldsymbol{x}_{j}) - Z_{i}(\boldsymbol{x}) \right) \right| \leq C_{4} \left| Z_{i}(\boldsymbol{x}_{j}) - Z_{i}(\boldsymbol{x}) \right| \mathbf{1}_{\|\boldsymbol{x}-\boldsymbol{x}_{j}\|_{\infty} \leq h}$$
$$\leq C_{4} M_{i} h^{\beta} \leq C_{4} M_{i} \epsilon \,.$$

Thus (23) in Lemma 3 follows with $K_1 = 1, K_2 = 3$ and $\kappa = 1/\beta$. For the second case $\epsilon \leq h^{\beta}$ let $\|\boldsymbol{x} - \boldsymbol{y}\|_2 \leq (C_2 C_3)^{-1} \epsilon^{1+1/\beta}$. Then

$$\begin{aligned} \left| X_{ni}(\boldsymbol{x}) - X_{ni}(\boldsymbol{y}) \right| &\leq \Phi_{ni} \sum_{\boldsymbol{j=1}}^{\boldsymbol{p}} \left| w_{\boldsymbol{j},\boldsymbol{p}}(\boldsymbol{x};h) - w_{\boldsymbol{j},\boldsymbol{p}}(\boldsymbol{y};h) \right| \\ &\leq \Phi_{ni} C_2 C_3 \left(\frac{\|\boldsymbol{x} - \boldsymbol{y}\|}{h} \wedge 1 \right) \leq \Phi_{ni} C_2 C_3 \epsilon. \end{aligned}$$

In this case (23) in Lemma 3 follows for $K_1 = (H C_2 C_3)^{-1}, K_2 = 1$ and $\kappa = 1 + 1/\beta$. Summarizing we may bound the capacity number by

$$N(\epsilon \| \alpha \circ \Phi_n \|, \alpha \circ F_n) \le \left(\frac{\max(3^{1/\beta}, C_2 C_3)}{2} \,\epsilon^{-(1+1/\beta)} + 2\right)^d =: \lambda_\beta^d(\epsilon)$$

which is integrable, see Lemma 3.

For ii) we calculate $\lim_{n\to\infty} \mathbb{E}[S_n(\boldsymbol{x})S_n(\boldsymbol{x}')]$. It holds

$$\mathbb{E}[S_n(\boldsymbol{x})S_n(\boldsymbol{x}')] = \sum_{\boldsymbol{j},\boldsymbol{k}=\boldsymbol{1}}^{\boldsymbol{p}} w_{\boldsymbol{j},\boldsymbol{p}}(\boldsymbol{x};h)w_{\boldsymbol{j},\boldsymbol{p}}(\boldsymbol{x}';h)\Gamma(\boldsymbol{x}_{\boldsymbol{j}},\boldsymbol{x}_{\boldsymbol{k}}) \stackrel{n \to \infty}{\longrightarrow} \Gamma(\boldsymbol{x},\boldsymbol{x}'),$$

uniformly for all $\boldsymbol{x}, \boldsymbol{x}' \in [0, 1]^d$ since $c \in \mathcal{H}_{[0,1]^{2d}}(\gamma, \tilde{L})$ as stated in Assumption 4.

For iii) we make use of the fact that the processes Z_i have finite second moment

$$\sum_{i=1}^{n} \mathbb{E}[\phi_{n,i}^2] \le 2 C_4^2 \mathbb{E}[Z_1^2(\mathbf{0}) + M_1^2] < \infty.$$

For iv) it holds for all fixed $\epsilon > 0$ that

$$\sum_{i=1}^{n} \mathbb{E}[\Phi_{n,i} \mathbf{1}_{\{\Phi_{n,i} > \epsilon\}}] \leq 2 C_4 \mathbb{E}[(Z_1^2(\mathbf{0}) + M_1^2) \mathbf{1}_{\{\Phi_{n,i} > \epsilon\}}] \xrightarrow{n \to \infty} 0$$

by the monotone convergence Theorem.

For v) we calculate

$$\rho_n^2(\boldsymbol{x}, \boldsymbol{x}') = \sum_{\boldsymbol{j}, \boldsymbol{k} = 1}^{\boldsymbol{p}} w_{\boldsymbol{j}, \boldsymbol{p}}(\boldsymbol{x}; h) w_{\boldsymbol{k}, \boldsymbol{p}}(\boldsymbol{x}; h) \Gamma(\boldsymbol{x}_{\boldsymbol{j}}, \boldsymbol{x}_{\boldsymbol{k}}) - 2 \sum_{\boldsymbol{j}, \boldsymbol{k} = 1}^{\boldsymbol{p}} w_{\boldsymbol{j}, \boldsymbol{p}}(\boldsymbol{x}'; h) w_{\boldsymbol{k}, \boldsymbol{p}}(\boldsymbol{x}; h) \Gamma(\boldsymbol{x}_{\boldsymbol{j}}, \boldsymbol{x}_{\boldsymbol{k}}) + \sum_{\boldsymbol{j}, \boldsymbol{k} = 1}^{\boldsymbol{p}} w_{\boldsymbol{j}, \boldsymbol{p}}(\boldsymbol{x}'; h) w_{\boldsymbol{k}, \boldsymbol{p}}(\boldsymbol{x}'; h) \Gamma(\boldsymbol{x}_{\boldsymbol{j}}, \boldsymbol{x}_{\boldsymbol{k}}) \xrightarrow{n \to \infty} \Gamma(\boldsymbol{x}, \boldsymbol{x}) - \Gamma(\boldsymbol{x}, \boldsymbol{x}') + \Gamma(\boldsymbol{x}', \boldsymbol{x}')$$

uniformly for all $\boldsymbol{x}, \boldsymbol{x}' \in [0, 1]^d$. Further let $(\boldsymbol{x}_n)_{n \in \mathbb{N}}, (\boldsymbol{y}_n)_{n \in \mathbb{N}} \subset [0, 1]^d$ be two deterministic sequences that suffice $\rho(\boldsymbol{x}_n, \boldsymbol{y}_n) \to 0, n \to \infty$. Then it holds

$$0 \le \rho_n(\boldsymbol{x}_n, \boldsymbol{y}_n) \le \sup_{\boldsymbol{x}, \boldsymbol{x}' \in [0, 1]^d} \left| \rho_n(\boldsymbol{x}, \boldsymbol{x}') - \rho(\boldsymbol{x}, \boldsymbol{x}') \right| + \rho(\boldsymbol{x}_n, \boldsymbol{y}_n) \stackrel{n \to \infty}{\longrightarrow} 0.$$

Thus the functional central limit Theorem is applicable and yields the stated result. \Box

References

- Cai, T. T. and M. Yuan (2011). Optimal estimation of the mean function based on discretely sampled functional data: Phase transition. *Ann. Statist.* 39(5) 2330–2355.
- Chang, C. X. Lin, and R. T. Ogden (2017). Simultaneous confidence bands for functional regression models. J. Statist. Plann. Inference 188 67–81.

- Cuevas, A. (2014). A partial overview of the theory of statistics with functional data. J. Statist. Plann. Inference 147 1–23.
- Degras, D. A. (2011). Simultaneous confidence bands for nonparametric regression with functional data. *Statist. Sinica* 1735–1765.
- Hall, P. H.-G. Müller, and J.-L. Wang (2006). Properties of principal component methods for functional and longitudinal data analysis. *Ann. Statist.* 1493–1517.
- Kalogridis, I. and S. Van Aelst (2022). Robust optimal estimation of location from discretely sampled functional data. *Scand. J. Stat.*.
- Li, Y. and T. Hsing (2010). Uniform convergence rates for nonparametric regression and principal component analysis in functional/longitudinal data. Ann. Statist. 38(6) 3321–3351.
- Liebl, D. and M. Reimherr (2023). Fast and fair simultaneous confidence bands for functional parameters. J. R. Stat. Soc. Ser. B. Stat. Methodol. 85(3) 842–868.
- Morris, J. S. (2015). Functional regression. Annu. Rev. Stat. Appl. 2 321–359.
- Pollard, D. (1990). Empirical processes: theory and applications. In NSF-CBMS regional conference series in probability and statistics pp. i–86. JSTOR.
- Ramsay, J. and B. Silvermann (1998). Functional data analysis. springer series in statistics.
- Rosenthal, H. P. (1970). On the subspaces of L_p (p > 2) spanned by sequences of independent random variables. *Israel J. Math.* 8 273–303.
- Sakhanenko, A. I. (1991). On the accuracy of normal approximation in the invariance principle. Siberian Adv. Math. 1 58–91.
- Tsybakov, A. B. (2004). Introduction to nonparametric estimation, 2009 Volume 9.
- van der Vaart, A. and J. Wellner (1996). Weak convergence and empirical processes: with applications to statistics. Springer Science & Business Media.
- Wang, J.-L. J.-M. Chiou, and H.-G. Müller (2016). Functional data analysis. Annu. Rev. Stat. Appl. 3 257–295.
- Wang, Y. G. Wang L. Wang, and R. T. Ogden (2020). Simultaneous confidence corridors for mean functions in functional data analysis of imaging data. *Biometrics* 76(2) 427–437.
- Xiao, L. (2020). Asymptotic properties of penalized splines for functional data. Bernoulli 26(4) 2847–2875.
- Yao, F. H.-G. Müller, and J.-L. Wang (2005). Functional data analysis for sparse longitudinal data. J. Amer. Statist. Assoc. 100(470) 577–590.
- Zhang, X. and J.-L. Wang (2016). From sparse to dense functional data and beyond. Ann. Statist. 44(5) 2281–2321.

A Proof of Lemma 1

For the proof of Lemma 1 we start with an auxiliary Lemma concerning the approximate localisation of the design points on each axis and their approximate distance.

Lemma 6. Let Assumption 5 be satisfied. Then for all $j = 1, ..., p_k$ and $1 \le k \le d$ it follows that

$$\frac{j - 0.5}{f_{k_{\max}} p_k} \le x_{k,j} \le \frac{j - 0.5}{f_{k_{\min}} p_k} \quad and \quad 1 - \frac{p_k - j + 0.5}{f_{k_{\min}} p_k} \le x_{k,j} \le 1 - \frac{p_k - j + 0.5}{f_{k_{\max}} p_k}, \tag{28}$$

and for all $1 \leq j < l \leq p_k$ that

$$\frac{l-j}{f_{k\max} p_k} \le x_{k,l} - x_{k,j} \le \frac{l-j}{f_{k\min} p_k} \,.$$
(29)

Proof of Lemma 6. The first assertion follows by the equivalence

$$\int_0^{x_{k,j}} f_{k\min} \,\mathrm{d}t \le \int_0^{x_{k,j}} f_k(t) \,\mathrm{d}t \le \int_0^{x_{k,j}} f_{k\max} \,\mathrm{d}t \quad \Leftrightarrow \quad x_{k,j} \,f_{k\min} \le \frac{j-0.5}{p_k} \le x_{k,j} \,f_{k\max} \,.$$

Similar equivalences and transformations with

$$\int_{x_{k,j}}^{1} f_{k\min} \, \mathrm{d}t \le \int_{x_{k,j}}^{1} f_k(t) \, \mathrm{d}t = 1 - \int_0^{x_{k,j}} f_k(t) \, \mathrm{d}t \le \int_{x_{k,j}}^{1} f_{k\max} \, \mathrm{d}t$$

and

$$\int_{x_{k,j}}^{x_{k,l}} f_{k\min} \, \mathrm{d}t \le \int_{x_{k,j}}^{x_{k,l}} f_k(t) \, \mathrm{d}t = \int_0^{x_{k,l}} f_k(t) \, \mathrm{d}t - \int_0^{x_{k,j}} f_k(t) \, \mathrm{d}t \le \int_{x_{k,j}}^{x_{k,l}} f_{k\max} \, \mathrm{d}t$$

show the other two assertations.

Lemma 7. Suppose that Assumption 5 is satisfied. Then we obtain for all subsets defined by intervals $I_1, \ldots, I_d \subseteq [0, 1]$ via $I = I_1 \times \ldots \times I_d \subseteq [0, 1]^d$ the estimate

$$\sum_{\mathbf{1} \leq \boldsymbol{j} \leq \boldsymbol{p}} \mathbb{1}_{\{\boldsymbol{x}_{\boldsymbol{j}} \in I\}} \leq 2^{d} \boldsymbol{f}_{\max}^{\mathbf{1}} \prod_{k=1}^{d} \max\left(p_{k} \nu(I_{k}), 1\right),$$

where $\nu(I_k)$ is the Lebesgue measure of I_k . In particular this means that Assumption 3 is satisfied.

Proof of Lemma 7. Without loss of generality we set $I = [a_1, b_1] \times \cdots \times [a_d, b_d] \subseteq [0, 1]^d$. We distinguish two cases with respect to the length of the k-interval $[a_k, b_k]$. Let $x_{k,1}, \ldots, x_{k,p_k} \in [0, 1]$ be the grid points on k^{th} axis. If $b_k - a_k < 1/p_k$, we can bound the number of design points in the interval $[a_k, b_k]$ by $\lceil f_{k_{\max}} \rceil = \min\{z \in \mathbb{N} \mid z > f_{k_{\max}}\}$. Otherwise, if we assume that $x_{k,j}$ is the smallest point larger than or equal to a_k , the lower bound in (29) would yield to the contradiction

$$x_{k,j+\lceil f_{k_{\max}}\rceil} - x_{k,j} \ge \frac{\lceil f_{k_{\max}}\rceil}{f_{k_{\max}} p_k} > \frac{1}{p_k}.$$

Hence,

$$\sum_{j=1}^{p_k} \mathbb{1}_{\{x_{k,j}\in[a_k,b_k]\}} \le \lceil f_{k\max} \rceil \le \frac{2f_{k\max}}{p_k} = 2f_{k\max}\max\left(p_k\left(b_k - a_k\right), 1\right)$$

follows. Conversely, if $b_k - a_k \ge 1/p_k$, the number of points in die corresponding interval is bounded by $\lceil (b_k - a_k) f_{k\max} p_k \rceil$ since otherwise we would obtain

$$x_{k,j+\lceil (b_k-a_k)f_{k_{\max}}p_k\rceil} - x_{k,j} \ge \frac{\lceil (b_k-a_k)f_{k_{\max}}p_k\rceil}{f_{k_{\max}}p_k} > b_k - a_k$$

just like before. Again this leads to

$$\sum_{j=1}^{p_k} \mathbb{1}_{\{x_{k,j} \in [a_k, b_k]\}} \le \lceil (b_k - a_k) f_{k_{\max}} p_k \rceil \le f_{k_{\max}}(b_k - a_k) + 1 \le 2f_{k_{\max}} \max\left(p_k \left(b_k - a_k \right), 1 \right).$$

Together with $\sum_{j=1}^{p} \mathbb{1}_{\{x_j \in I\}} = \prod_{k=1}^{d} \sum_{j=1}^{p_k} \mathbb{1}_{x_{k,j} \in [a_k, b_k]}$ the two cases yield the first claim. Assumption 3 follows with $C_4 = 4^d f_{\max}^1$ by

$$\sum_{j=1}^{p} \mathbf{1}_{x_{j} \in [x-(j,\dots,h),x+(h,\dots,h)]} \leq 2^{d} f_{\max}^{1} \prod_{k=1}^{d} \max(2h p_{k}, 1) = 4^{d} f_{\max}^{1} h^{d} p^{1}.$$

Construction of the weights of the linear estimator.

In order to derive the form of the weights $w_{j,p}(\boldsymbol{x};h)$ of the linear estimator $\hat{\mu}_{n,p,h}^{\text{LP},m}$ in (13) we set

$$B_{\boldsymbol{p},h}(\boldsymbol{x}) := \frac{1}{\boldsymbol{p}^{1}h^{d}} \sum_{\boldsymbol{j}=1}^{\boldsymbol{p}} U_{h}(\boldsymbol{x}_{\boldsymbol{j}}-\boldsymbol{x}) U_{h}^{\top}(\boldsymbol{x}_{\boldsymbol{j}}-\boldsymbol{x}) K_{h}(\boldsymbol{x}_{\boldsymbol{j}}-\boldsymbol{x}) \quad \in \mathbb{R}^{N_{m,d} \times N_{m,d}}$$
(30)

and

$$\boldsymbol{a}_{\boldsymbol{p},h}(\boldsymbol{x}) := \frac{1}{\boldsymbol{p}^{1}h^{d}} \sum_{\boldsymbol{j}=1}^{\boldsymbol{p}} U_{h}(\boldsymbol{x}_{\boldsymbol{j}} - \boldsymbol{x}) K_{h}(\boldsymbol{x}_{\boldsymbol{j}} - \boldsymbol{x}) \bar{Y}_{\boldsymbol{j}} \quad \in \mathbb{R}^{N_{m,d}}$$

for $\boldsymbol{x} \in T$, then $\widehat{\boldsymbol{\vartheta}}_{n,\boldsymbol{p},h}(\boldsymbol{x})$ as in (12) is the solution to the weighted least squares problem

$$\widehat{\boldsymbol{\vartheta}}_{n,\boldsymbol{p},h}(\boldsymbol{x}) = \operatorname*{argmin}_{\boldsymbol{\vartheta} \in \mathbb{R}^{N_{m,d}}} \left(-2 \, \boldsymbol{\vartheta}^\top \boldsymbol{a}_{\boldsymbol{p},h}(\boldsymbol{x}) + \boldsymbol{\vartheta}^\top B_{\boldsymbol{p},h}(\boldsymbol{x}) \, \boldsymbol{\vartheta} \right).$$

The solution is determined by the normal equations

$$B_{\boldsymbol{p},h}(\boldsymbol{x})\,\widehat{\boldsymbol{\vartheta}}_{n,\boldsymbol{p},h}(\boldsymbol{x}) = \boldsymbol{a}_{\boldsymbol{p},h}(\boldsymbol{x})\,,$$

where the matrix $B_{p,h}(\boldsymbol{x})$ is positive semidefinite. In particular, if $B_{p,h}(\boldsymbol{x})$ is positive definite, the solution in (12) is uniquely determined and we obtain

$$\widehat{\mu}_{n,\boldsymbol{p},h}^{\mathrm{LP},m}(\boldsymbol{x}) = \sum_{\boldsymbol{j}=1}^{\boldsymbol{p}} w_{\boldsymbol{j},\boldsymbol{p}}(\boldsymbol{x};h) \, \bar{Y}_{\boldsymbol{j}}$$

with

$$w_{\boldsymbol{j},\boldsymbol{p}}(\boldsymbol{x};h) := \frac{1}{\boldsymbol{p}^{1}h^{d}} U_{m}^{\top}(\boldsymbol{0}) B_{\boldsymbol{p},h}^{-1}(\boldsymbol{x}) U_{h}(\boldsymbol{x}_{\boldsymbol{j}}-\boldsymbol{x}) K_{h}(\boldsymbol{x}_{\boldsymbol{j}}-\boldsymbol{x}) , \qquad (31)$$

so that the local polynomial estimator is a linear estimator. In the following Lemma we show a result that implicates that $B_{p,h}(x)$ is positive definite.

Lemma 8. Suppose that the kernel K satisfies (14) in Lemma 1. Then there exist a sufficiently large $p_0 \in \mathbb{N}$ and a sufficiently small $h_0 \in \mathbb{R}_+$ such that for all \boldsymbol{p} with $\boldsymbol{p}_{\min} \geq p_0$ and $h \in (c/\boldsymbol{p}_{\min}, h_0]$, where $c \in \mathbb{R}_+$ is a sufficiently large constant, the smallest eigenvalues $\lambda_{\min}(B_{\boldsymbol{p},h}(\boldsymbol{x}))$ of the matrices $B_{\boldsymbol{p},h}(\boldsymbol{x})$, which are given in (30), are bounded below by a universal positive constant $\lambda_0 > 0$ for any $\boldsymbol{x} \in [0, 1]^d$.

An immediate consequence of Lemma 8 is the invertibility of $B_{p,h}(\boldsymbol{x})$ for all \boldsymbol{p} with $\boldsymbol{p}_{\min} \geq \boldsymbol{p}_0, h \in (c/\boldsymbol{p}_{\min}, h_0]$ and $\boldsymbol{x} \in [0, 1]^d$, and hence also the uniqueness of the local polynomial estimator for these \boldsymbol{p} and h. In Tsybakov (2004, Lemma 1.5) the lower bound for the smallest eigenvalues has only be shown for a fixed sequence h_p (for d = 1) of bandwidths which satisfies $h_p \to 0$ and $p h_p \to \infty$. In contrast, we allow an uniformly choice of h which results, in particular, in the findings of Section 3.

Proof of Lemma 8. In the following let $\boldsymbol{v} \in \mathbb{R}^{N_{m,d}}$. We show that there exist a sufficiently large $p_0 \in \mathbb{N}$ and a component wise sufficiently small $h_0 \in \mathbb{R}_+$ such that the estimate

$$\inf_{\boldsymbol{p} \ge \boldsymbol{p}_0} \inf_{h \in (c/\boldsymbol{p}_{\min},h_0]} \inf_{\boldsymbol{x} \in [0,1]^d} \inf_{\|\boldsymbol{v}\|_2 = 1} \boldsymbol{v}^\top B_{\boldsymbol{p},h}(\boldsymbol{x}) \, \boldsymbol{v} \ge \lambda_0 \tag{32}$$

is satisfied. Then we obtain for these \boldsymbol{p} and h, and any $\boldsymbol{x} \in [0,1]^d$ also

$$\lambda_{\min} ig(B_{\boldsymbol{p},h}(\boldsymbol{x}) ig) = \boldsymbol{e}_{\min}^{ op} ig(B_{\boldsymbol{p},h}(\boldsymbol{x}) ig) B_{\boldsymbol{p},h}(\boldsymbol{x}) \, \boldsymbol{e}_{\min} ig(B_{\boldsymbol{p},h}(\boldsymbol{x}) ig) \ge \inf_{\|\boldsymbol{v}\|_2 = 1} \boldsymbol{v}^{ op} B_{\boldsymbol{p},h}(\boldsymbol{x}) \, \boldsymbol{v} \ge \lambda_0 \,,$$

where $\boldsymbol{e}_{\min}(B_{\boldsymbol{p},h}(\boldsymbol{x})) \in \mathbb{R}^{m+1}$ is a normalized eigenvector of $\lambda_{\min}(B_{\boldsymbol{p},h}(\boldsymbol{x}))$. Let $\Psi: \{0,1\}^d \mapsto \{1,\ldots,2^d\}$ be an enumeration of the subsets of $\{0,1\}^d$ with $\Psi(\mathbf{0}) = 1$ and $E^i := J_{(\Psi^{-1}(i))_1} \times \cdots \times J_{(\Psi^{-1}(i))_d}$ for all $i = 1,\ldots,2^d$, whereby $J_0 := [0,\Delta)$ and $J_1 := (-\Delta,0]$ with the constants $\Delta \in \mathbb{R}_+$ and $K_{\min} > 0$ given in (14). We set

$$\lambda_i(\boldsymbol{v}) \coloneqq \boldsymbol{f}_{\min}^{\boldsymbol{1}} K_{\min} \int_{E^i} \left(\boldsymbol{v}^\top U_m(\boldsymbol{z}) \right)^2 \mathrm{d}\boldsymbol{z} \,, \qquad \lambda_i \coloneqq \inf_{\|\boldsymbol{v}\|_2 = 1} \lambda_i(\boldsymbol{v})$$

for all $i = 1, ..., 2^d$ and $f_{\min}^1 := \prod_{k=1}^d f_{k\min}$ given in Assumption 5. Applying Tsybakov (2004, Lemma 1.4) with $K(\mathbf{z}) = \mathbb{1}_{E^i}(\mathbf{z})$ leads to

$$\lambda_i(\boldsymbol{v}) \geq \lambda_i > 0$$
.

Therefore we find a $\lambda_0 > 0$ such that $\min(\lambda_1, \ldots, \lambda_{2^d}) > \lambda_0 > 0$, e.g. $\lambda_0 := \min(\lambda_1, \ldots, \lambda_{2^d})/2$. Now we want to specify a partition $S_1 \cup \ldots \cup S_{2^d} = [0, 1]^d$ and functions $A_{p,h}^{(1)}(\boldsymbol{x}, \boldsymbol{v}), \ldots, A_{p,h}^{(2^d)}(\boldsymbol{x}, \boldsymbol{v})$ such that

$$\mathbf{v}^{\top} B_{\mathbf{p},h}(\mathbf{x}) \, \mathbf{v} \ge A_{\mathbf{p},h}^{(i)}(\mathbf{x},\mathbf{v}) \,, \quad \|\mathbf{v}\|_2 = 1, \quad \mathbf{x} \in S_i \,, \quad i = 1, \dots, 2^d \,,$$
 (33)

and

$$\sup_{\boldsymbol{x}\in S_i} \sup_{\|\boldsymbol{v}\|_2=1} \left| A_{\boldsymbol{p},h}^{(i)}(\boldsymbol{x},\boldsymbol{v}) - \lambda_i(\boldsymbol{v}) \right| \le \sum_{k=1}^d \frac{c_1}{p_k h}, \quad i = 1,\dots,2$$
(34)

hold true for a positive constant $c_1 > 0$. Setting $c := dc_1/(\min(\lambda_1, \ldots, \lambda_{2^d}) - \lambda_0)$ yields $\sum_{k=1}^d c_1/(p_k h) < dc_1/c \leq \lambda_i - \lambda_0$ for all $i = 1, \ldots, 2^d$, since $h \in (c/p_{\min}, h_0]$. Hence, it follows by (33) and (34) that

$$\begin{split} \inf_{\boldsymbol{x}\in S_i} \inf_{\|\boldsymbol{v}\|_2=1} \boldsymbol{v}^\top B_{\boldsymbol{p},h}(\boldsymbol{x}) \, \boldsymbol{v} &= \inf_{\boldsymbol{x}\in S_i} \inf_{\|\boldsymbol{v}\|_2=1} \left(\lambda_i(\boldsymbol{v}) + \boldsymbol{v}^\top B_{\boldsymbol{p},h}(\boldsymbol{x}) \, \boldsymbol{v} - \lambda_i(\boldsymbol{v})\right) \\ &\geq \lambda_i + \inf_{\boldsymbol{x}\in S_i} \inf_{\|\boldsymbol{v}\|_2=1} \left(A_{p,h}^{(i)}(\boldsymbol{x},\boldsymbol{v}) - \lambda_i(\boldsymbol{v})\right) \\ &= \lambda_i - \sup_{\boldsymbol{x}\in S_i} \sup_{\|\boldsymbol{v}\|_2=1} \left(\lambda_i(\boldsymbol{v}) - A_{p,h}^{(i)}(\boldsymbol{x},\boldsymbol{v})\right) \\ &\geq \lambda_i - \sup_{\boldsymbol{x}\in S_i} \sup_{\|\boldsymbol{v}\|_2=1} \left|A_{p,h}^{(i)}(\boldsymbol{x},\boldsymbol{v}) - \lambda_i(\boldsymbol{v})\right| \\ &\geq \lambda_i - \sum_{k=1}^d \frac{c_1}{p_k \, h} \geq \lambda_0 \,, \end{split}$$

which leads to (32) because of

$$\begin{split} \inf_{\boldsymbol{p} \geq \boldsymbol{p}_0} \inf_{h \in (c/\boldsymbol{p}_{\min},h_0]} \inf_{\boldsymbol{x} \in [0,1]^d} \inf_{\|\boldsymbol{v}\|_2 = 1} \boldsymbol{v}^\top B_{\boldsymbol{p},h}(\boldsymbol{x}) \, \boldsymbol{v} \\ &= \inf_{\boldsymbol{p} \geq \boldsymbol{p}_0} \inf_{h \in (c/\boldsymbol{p}_{\min},h_0]} \min_{i=1,...,2^d} \left(\inf_{\boldsymbol{x} \in S_i} \inf_{\|\boldsymbol{v}\|_2 = 1} \boldsymbol{v}^\top B_{\boldsymbol{p},h}(\boldsymbol{x}) \, \boldsymbol{v} \right) \geq \lambda_0 \, . \end{split}$$

Now we show (33) and (34). Let $I_0^k := [0, 1 - \Delta h - 1/(2f_{k\min}p_k)]$ and $I_1^k := [1 - \Delta h - 1/(2f_{k\min}p_k), 1]$. We define $S_i := I_{(\Psi^{-1}(i))_1}^1 \times \ldots \times I_{(\Psi^{-1}(i))_d}^d$. It is clear that we get 2^d different subsets and $\bigcup_{i=1}^{2^d} S_i = [0, 1]^d$. We indicate the design points on the k^{th} axis by $x_{k,1}, \ldots, x_{k,p_k}$ and define $y_{k,j} := (x_{k,j} - x_k)/h$, $x_k \in [0, 1]$ for all $1 \leq j \leq p_k$ and $1 \leq k \leq d$ and let $y_j := (y_{1,j_1}, \ldots, y_{d,j_d}) \in \mathbb{R}^d$. We have to differentiate in two different cases. First let $x_k \in I_0^k$

$$y_{k,1} \leq \frac{1}{2f_{k\min} p_k h} - \frac{x_k}{h} \leq \frac{1}{2f_{k\min} p_k h},$$

$$y_{k,p_k} \geq \frac{1 - \frac{1}{2f_{k\min} p_k} - x_k}{h} = \frac{1 - x_k}{h} - \frac{1}{2f_{k\min} p_k h}$$

$$\geq \frac{1 - \left(1 - \Delta h - (2f_{k\min} p_k)^{-1}\right)}{h} - \frac{1}{2f_{k\min} p_k h} = \Delta$$

by (28) in Lemma 6. For appropriate \boldsymbol{p} and $h \in (c/\boldsymbol{p}_{\min}, h_0]$ the quantity $(2f_{k\min} p_k h)^{-1}$ gets small if c is chosen large enough. Consequently, the points $y_{k,1}, \ldots, y_{k,p_k}$ form a grid which covers an interval containing $[1/(2f_{k\min} p_k h), \Delta]$. Hence, there exist $1 \leq j_{k*} := j_{k*}(x_k) < j_k^* := j_k^*(x_k) \leq p_k$ such that $y_{k,j_{k*}} \geq 0 \land (j_{k*} = 1 \lor y_{k,j_{k*}-1} < 0)$ and $y_{k,j_k^*} \leq \Delta \land y_{k,j_{k}^*+1} > \Delta$. Here \wedge denotes the logical and, and \vee the logical or. For the second case let $x_k \in I_1^k$. Then we obtain the estimates

$$y_{k,1} \leq \frac{1}{2f_{k\min}p_k h} - \frac{x_k}{h} \leq \Delta - \frac{1}{h} \leq -\Delta,$$
$$y_{k,p_k} \geq -\frac{1}{2f_{k\min}p_k h}$$

since h has to be sufficiently small and p_k sufficiently large. Consequently, the points $y_{k,1}, \ldots, y_{k,p_k}$ form a grid which covers an interval containing $[-\Delta, -1/(2f_{k\min}p_k h)]$. Therefore we could define \tilde{j}_{k*} and \tilde{j}_{k}^* analogously as before but in the following we only examine the case for $S_1 = I_0^1 \times \ldots \times I_0^d$. The arguments for the other subsets are basically the same after the right case distinctions. We set $g(\boldsymbol{x}) := (\boldsymbol{v}^\top U_m(\boldsymbol{x}))^2$ to save some space in the upcoming calculations. By Assumption 5, inequality (29) in Lemma 6 for the $y_{k,j}$ and the presentation of $B_{\boldsymbol{p},h}(\boldsymbol{x})$ in (30) we get for $\boldsymbol{x} \in S_1$

$$\begin{split} \boldsymbol{v}^{\top} B_{\boldsymbol{p},h}(\boldsymbol{x}) \, \boldsymbol{v} &= \frac{1}{\boldsymbol{p}^{1} h^{d}} \boldsymbol{v}^{\top} \bigg(\sum_{\boldsymbol{j}=1}^{\boldsymbol{p}} U_{m}(\boldsymbol{y}_{\boldsymbol{j}}) \, U_{m}^{\top}(\boldsymbol{y}_{\boldsymbol{j}}) \, K(\boldsymbol{y}_{\boldsymbol{j}}(\boldsymbol{x})) \bigg) \boldsymbol{v} \\ &\geq \frac{\boldsymbol{f}_{\min}^{1} K_{\min}}{\boldsymbol{f}_{\min}^{1} \boldsymbol{p}^{1} h^{d}} \sum_{\boldsymbol{j}=1}^{\boldsymbol{p}} \Big(\boldsymbol{v}^{\top} U_{m}(\boldsymbol{y}_{\boldsymbol{j}}) \Big)^{2} \mathbb{1}_{[-\boldsymbol{\Delta},\boldsymbol{\Delta}]}(\boldsymbol{y}_{\boldsymbol{j}}) \\ &\geq \boldsymbol{f}_{\min}^{1} K_{\min} \sum_{\boldsymbol{j}_{1}=1}^{p_{1}-1} \cdots \sum_{\boldsymbol{j}_{d}=1}^{p_{d}-1} g(\boldsymbol{y}_{\boldsymbol{j}})(y_{1,j_{1}+1}-y_{1,j_{1}}) \cdots (y_{d,j_{d}+1}-y_{d,j_{d}}) \mathbb{1}_{[\boldsymbol{0},\boldsymbol{\Delta})}(\boldsymbol{y}_{(j_{1},\dots,j_{d})}) \\ &\geq \boldsymbol{f}_{\min}^{1} K_{\min} \sum_{\boldsymbol{j}_{1}=j_{1*}}^{j_{1}^{*}} \cdots \sum_{\boldsymbol{j}_{d}=j_{d*}}^{j_{d}^{*}} g(\boldsymbol{y}_{\boldsymbol{j}})(y_{1,j_{1}+1}-y_{1,j_{1}}) \cdots (y_{d,j_{d}+1}-y_{d,j_{d}}) =: A_{\boldsymbol{p},h}^{(1)}(\boldsymbol{x},\boldsymbol{v}) \end{split}$$

Inserting this function in (34), dropping the scalar $f_{\min}^1 K_{\min}$ and oppressing the sups yields

$$\left| \sum_{j_{1}=j_{1*}}^{j_{1}^{*}} \cdots \sum_{j_{d}=j_{d*}}^{j_{d}^{*}} g(\boldsymbol{y}_{j})(y_{1,j_{1}+1} - y_{1,j_{1}}) \cdots (y_{d,j_{d}+1} - y_{d,j_{d}}) - \int_{E^{1}} g(\boldsymbol{z}) \, \mathrm{d}z_{1} \cdots \mathrm{d}z_{d} \right| \\
\leq \sum_{k=1}^{d} \left| \int_{0}^{\Delta_{1}} \cdots \int_{0}^{y_{k,k*}} \cdots \int_{0}^{\Delta_{d}} g(\boldsymbol{z}) \, \mathrm{d}z_{1} \cdots \mathrm{d}z_{d} \right| \\
+ \sum_{j_{1}=j_{1*}}^{j_{1}^{*}-1} \cdots \sum_{j_{k}=j_{k*}}^{j_{k}^{*}-1} \left| \int_{y_{1,j_{1}}}^{y_{1,j_{1}+1}} \cdots \int_{y_{d,j_{d}}}^{y_{d,j_{d}+1}} \left(g(\boldsymbol{y}_{j}) - g(\boldsymbol{z}) \right) \, \mathrm{d}z_{1} \cdots \mathrm{d}z_{d} \right| \quad (35) \\
+ \Delta^{d-1} \sum_{k=1}^{d} \left| g(\boldsymbol{y}_{(k_{1}^{*},\ldots,k_{d}^{*})})(y_{k,j_{k}^{*}+1} - y_{k,j_{k}^{*}}) \right| \\
+ \sum_{k=1}^{d} \left| \int_{0}^{\Delta_{1}} \cdots \int_{y_{k,j_{k}^{*}}}^{\Delta_{d}} \cdots \int_{0}^{\Delta_{d}} g(\boldsymbol{z}) \, \mathrm{d}z_{1} \cdots \mathrm{d}z_{d} \right|.$$

We want to bound the order of these terms separately. Firstly, note that $g(\boldsymbol{z}) = (\boldsymbol{v}^{\top} U_m(\boldsymbol{z}))^2$ for all coordinate wise bounded \boldsymbol{z} . Further $g(\boldsymbol{z})$ is LIPSCHITZ-continuous since it is a multivariate polynomial. Therefore it holds that $|g(\boldsymbol{y}_j) - g(\boldsymbol{z})| = \mathcal{O}((\boldsymbol{p}_{\min}h)^{-1})$. Together with the fact that $|y_{k,j_{k*}}|, |\Delta - y_{k,j_k^*}|, |y_{k,j_k+1} - y_{k,j_k}| = \mathcal{O}((p_k h)^{-1})$ and $\sum_{j_k=j_{k*}}^{j_k^*-1} 1 = \mathcal{O}(p_k h)$ by Lemma 7 we get that

$$(35) = \mathcal{O}\left(\frac{d}{\boldsymbol{p}_{\min}\,h}\right).$$

Now we can prove Lemma 1.

Proof of Lemma 1. By Lemma 8 there exist a sufficiently large $p_0 \in \mathbb{N}$ and a sufficiently small $h_0 \in \mathbb{R}_+$ such that for all $\mathbf{p} = (p_1, \ldots, p_d)^\top$ with $\mathbf{p}_{\min} \geq \mathbf{p}_0$ and $h \in (c/\mathbf{p}_{\min}, h_0]$, where c > 0 is a sufficiently large constant, the local polynomial estimator in (13) with any order $m \in \mathbb{N}$ is unique and a linear estimator with weights given in (31). Further, the mentioned Lemma together with Lemma 7, Tsybakov (2004, Lemma 1.3) and (14) imply (W2) and (W3) of Assumption 2 for the weights of the local polynomial estimator. Moreover, Lemma 8 and Tsybakov (2004, Proposition 1.12) lead to (W1) with $\gamma = m$. Note that the mentioned Lemma has to be generalized to the *d*-dimensional case. In the following we make use of

$$\left\|B_{\boldsymbol{p},h}^{-1}(\boldsymbol{x})\right\|_{M,2} \le \frac{1}{\lambda_0} \tag{36}$$

for all $\boldsymbol{p} \geq \boldsymbol{p}_0$, $h \in (c/\boldsymbol{p}_{\min}, h_0]$ and $\boldsymbol{x} \in [0, 1]^d$, where $\|M\|_{M, 2}$ is the spectral norm of a symmetric matrix $M \in \mathbb{R}^{N_{m,d} \times N_{m,d}}$ what is an immediate consequence of Lemma 8.

We continue with the LIPSCHITZ-condition (W4), and divide the proof in three cases with respect to the fact whether the weights vanish or not. In the following let $1 \leq j \leq p$ and $x, y \in [0, 1]^d$.

Let min $(\|\boldsymbol{x} - \boldsymbol{x}_{\boldsymbol{j}}\|_{\infty}, \|\boldsymbol{y} - \boldsymbol{x}_{\boldsymbol{j}}\|_{\infty}) > h$, then by (W2) both weights $w_{\boldsymbol{j},\boldsymbol{p}}(\boldsymbol{x};h)$ and $w_{\boldsymbol{j},\boldsymbol{p}}(\boldsymbol{y};h)$ vanish, and hence (W4) is clear.

Let $\max(\|\boldsymbol{x} - \boldsymbol{x}_{\boldsymbol{j}}\|_{\infty}, \|\boldsymbol{y} - \boldsymbol{x}_{\boldsymbol{j}}\|_{\infty}) > h$. We assume $\|\boldsymbol{y} - \boldsymbol{x}_{\boldsymbol{j}}\|_{\infty} > h$ and $\|\boldsymbol{x} - \boldsymbol{x}_{\boldsymbol{j}}\|_{\infty} \leq h$ without loss of generality. Once again (W2) leads to $w_{\boldsymbol{j},\boldsymbol{p}}(\boldsymbol{y};h) = 0$, and hence the CAUCHY-SCHWARZ inequality, $\|U_m(\mathbf{0})\|_2 = 1$ and (36) imply

$$\begin{split} \left| w_{\boldsymbol{j},\boldsymbol{p}}(\boldsymbol{x};h) - w_{\boldsymbol{j},\boldsymbol{p}}(\boldsymbol{y};h) \right| &= \frac{1}{\boldsymbol{p}^{1}h^{d}} \left| U_{m}^{\top}(\boldsymbol{0}) B_{\boldsymbol{p},h}^{-1}(\boldsymbol{x}) U_{h}(\boldsymbol{x}_{\boldsymbol{j}} - \boldsymbol{x}) \right| \left| K_{h}(\boldsymbol{x}_{\boldsymbol{j}} - \boldsymbol{x}) \right| \\ &\leq \frac{1}{\boldsymbol{p}^{1}h^{d}} \left\| U_{m}(\boldsymbol{0}) \right\|_{2} \left\| B_{\boldsymbol{p},h}^{-1}(\boldsymbol{x}) U_{h}(\boldsymbol{x}_{\boldsymbol{j}} - \boldsymbol{x}) \right\|_{2} \left| K_{h}(\boldsymbol{x}_{\boldsymbol{j}} - \boldsymbol{x}) \right| \\ &\leq \frac{1}{\boldsymbol{p}^{1}h^{d}} \left\| B_{\boldsymbol{p},h}^{-1}(\boldsymbol{x}) \right\|_{M,2} \left\| U_{h}(\boldsymbol{x}_{\boldsymbol{j}} - \boldsymbol{x}) \right\|_{2} \left| K_{h}(\boldsymbol{x}_{\boldsymbol{j}} - \boldsymbol{x}) \right| \\ &\leq \frac{1}{\lambda_{0} \boldsymbol{p}^{1}h^{d}} \left| K_{h}(\boldsymbol{x}_{\boldsymbol{j}} - \boldsymbol{x}) \right| \left(\sum_{|\boldsymbol{r}|=0}^{m} \left(\frac{(\boldsymbol{x}_{\boldsymbol{j}} - \boldsymbol{x})^{\boldsymbol{r}}}{\boldsymbol{h}^{\boldsymbol{r}} \boldsymbol{r}!} \right)^{2} \right)^{\frac{1}{2}} \\ &\leq \frac{c_{1}}{\lambda_{0} \boldsymbol{p}^{1}h^{d}} \left| K_{h}(\boldsymbol{x}_{\boldsymbol{j}} - \boldsymbol{x}) \right| \end{split}$$

for a positive constant $c_1 > 0$. In the last step we used the fact that the sum can't get arbitrarily large, also for component wise small h, because the kernel K has compact support in $[-1,1]^d$. If $\|\boldsymbol{x} - \boldsymbol{y}\|_{\infty} > h$, we use the upper bound of the kernel function in (14) and obtain

$$|w_{\boldsymbol{j},\boldsymbol{p}}(\boldsymbol{x};h) - w_{\boldsymbol{j},\boldsymbol{p}}(\boldsymbol{y};h)| \leq \frac{c_1 K_{\max}}{\lambda_0 \, \boldsymbol{p}^1 h^d}$$

Conversely, if $\|\boldsymbol{x} - \boldsymbol{y}\|_{\infty} \leq h$, we add $K_h(\boldsymbol{x_j} - \boldsymbol{y}) = 0$ and estimate

$$\left|w_{\boldsymbol{j},\boldsymbol{p}}(\boldsymbol{x};h) - w_{\boldsymbol{j},\boldsymbol{p}}(\boldsymbol{y};h)\right| \leq \frac{c_1}{\lambda_0 \, \boldsymbol{p}^1 h^d} \left|K_h(\boldsymbol{x}_{\boldsymbol{j}} - \boldsymbol{x}) - K_h(\boldsymbol{x}_{\boldsymbol{j}} - \boldsymbol{y})\right| \leq \frac{c_1 \, L_K}{\lambda_0 \, \boldsymbol{p}^1 h^d} \cdot \frac{\|\boldsymbol{x} - \boldsymbol{y}\|_{\infty}}{h},$$

because of the LIPSCHITZ-continuity of the kernel. So in total we get

$$w_{\boldsymbol{j},\boldsymbol{p}}(\boldsymbol{x};h) - w_{\boldsymbol{j},\boldsymbol{p}}(\boldsymbol{y};h) \bigg| \leq \frac{c_1 \max(K_{\max}, L_K)}{\lambda_0 \, \boldsymbol{p}^1 h^d} \left(\frac{\|\boldsymbol{x} - \boldsymbol{y}\|_{\infty}}{h} \wedge 1 \right).$$
(37)

Let $\max(\|\boldsymbol{x} - \boldsymbol{x}_{j}\|_{\infty}, \|\boldsymbol{y} - \boldsymbol{x}_{j}\|_{\infty}) \leq h$, then both weights don't vanish and we have to show a proper LIPSCHITZ-property for

$$w_{\boldsymbol{j},\boldsymbol{p}}(\boldsymbol{x};h) = \frac{1}{\boldsymbol{p}^{\mathbf{1}}h^{d}} U_{m}^{\top}(\boldsymbol{0}) B_{\boldsymbol{p},h}^{-1}(\boldsymbol{x}) U_{h}(\boldsymbol{x}_{\boldsymbol{j}}-\boldsymbol{x}) K_{h}(\boldsymbol{x}_{\boldsymbol{j}}-\boldsymbol{x})$$

The kernel K and polynomials on compact intervals are LIPSCHITZ-continuous, hence it suffices to show that $B_{p,h}^{-1}(x)$ has this property as well. Then the weights are products of bounded LIPSCHITZ-continuous functions and, thus, also LIPSCHITZ-continuous. The entries of the matrix $B_{p,h}(x)$, which is defined in (30), considered as functions from [0, 1] to \mathbb{R} are LIPSCHITZ-continuous. Indeed they are of order one by using Assumption 3. Hence, the row sum norm $||B_{p,h}(x) - B_{p,h}(y)||_{M,\infty}$ is a sum of these LIPSCHITZ-continuous functions and, in consequence, there exists a positive constant $L_{\infty} > 0$ such that

$$\|B_{\boldsymbol{p},h}(\boldsymbol{x}) - B_{\boldsymbol{p},h}(\boldsymbol{y})\|_{\mathcal{M},\infty} \le (N_{m,d}+1)L_{\infty} \ \frac{\|\boldsymbol{x} - \boldsymbol{y}\|_{\infty}}{h}$$

is satisfied. Since the matrices are symmetric the column sum norm is equal to the row sum norm and we obtain

$$\|B_{p,h}(\boldsymbol{x}) - B_{p,h}(\boldsymbol{y})\|_{M,2} \le \|B_{p,h}(\boldsymbol{x}) - B_{p,h}(\boldsymbol{y})\|_{M,\infty} \le (N_{m,d}+1)L_{\infty} \frac{\|\boldsymbol{x} - \boldsymbol{y}\|_{\infty}}{h}.$$

This leads together with (36) and the submultiplicativity of the spectral norm to

$$\begin{split} \left\| B_{\boldsymbol{p},h}^{-1}(x) - B_{\boldsymbol{p},h}^{-1}(y) \right\|_{M,2} &= \left\| B_{\boldsymbol{p},h}^{-1}(y) \left(B_{\boldsymbol{p},h}(\boldsymbol{y}) - B_{\boldsymbol{p},h}(\boldsymbol{x}) \right) B_{\boldsymbol{p},h}^{-1}(x) \right\|_{M,2} \\ &\leq \left\| B_{\boldsymbol{p},h}^{-1}(y) \right\|_{M,2} \left\| B_{\boldsymbol{p},h}(\boldsymbol{y}) - B_{\boldsymbol{p},h}(\boldsymbol{x}) \right\|_{M,2} \left\| B_{\boldsymbol{p},h}^{-1}(x) \right\|_{M,2} \\ &\leq \frac{(N_{m,d}+1)L_{\infty}}{\lambda_0^2} \left\| \frac{\|\boldsymbol{x} - \boldsymbol{y}\|_{\infty}}{h} \right\|_{M,2} \end{split}$$

which is the LIPSCHITZ-continuity of $B_{p,h}^{-1}(x)$ with respect to the spectral norm. So finally there exists a positive constant $c_2 > 0$ such that

$$\left|w_{\boldsymbol{j},\boldsymbol{p}}(\boldsymbol{x};h) - w_{\boldsymbol{j},\boldsymbol{p}}(\boldsymbol{y};h)\right| \leq \frac{c_2}{2\boldsymbol{p}^{\mathbf{1}}h^d} \ \frac{\|\boldsymbol{x} - \boldsymbol{y}\|_{\infty}}{h} \leq \frac{c_2}{\boldsymbol{p}^{\mathbf{1}}h^d}$$

is satisfied. Here we used in the last step that $\max(\|\boldsymbol{x} - \boldsymbol{x}_{\boldsymbol{j}}\|_{\infty}, \|\boldsymbol{y} - \boldsymbol{x}_{\boldsymbol{j}}\|_{\infty}) \leq h$ implies $\|\boldsymbol{x} - \boldsymbol{y}\|_{\infty} \leq 2h$.

Finally we choose $C_3 \ge \max(c_1 \max(K_{\max}, L_K)/\lambda_0, c_2)$ and obtain with (37) finally Assumption (W4).

B Proof of Lemma 2

Proof of Lemma 2 i). For the first part we want to use Taylor expansion and that the weights of the estimator reproduce polynomials of a certain degree. For certain $\theta_j \in [0, 1], 1 \leq j \leq p$, s.t. $\tau_j := \mathbf{x} + \theta_j (\mathbf{x}_j - \mathbf{x}) \in [0, 1]^d$ it holds

$$\begin{split} \sum_{j=1}^{p} w_{j,p}(\boldsymbol{x};h) \mu(\boldsymbol{x}_{j}) - \mu(\boldsymbol{x}) &= \sum_{j=1}^{p} w_{j,p}(\boldsymbol{x};h)(\mu(\boldsymbol{x}_{j}) - \mu(\boldsymbol{x})) \\ &= \sum_{j=1}^{p} \Big(\sum_{|\boldsymbol{k}|=1}^{\gamma-1} D^{\boldsymbol{k}} \mu(\boldsymbol{x}) \frac{(\boldsymbol{x}_{j}-\boldsymbol{x})^{\boldsymbol{k}}}{\boldsymbol{k}!} + \sum_{|\boldsymbol{k}|=\gamma} D^{\boldsymbol{k}} \mu(\boldsymbol{\tau}_{j}) \frac{(\boldsymbol{x}_{j}-\boldsymbol{x})^{\boldsymbol{k}}}{\boldsymbol{k}!} \Big) w_{j,p}(\boldsymbol{x};h) \\ &= \sum_{j=1}^{p} \sum_{|\boldsymbol{k}|=\gamma} D^{\boldsymbol{k}} \mu(\boldsymbol{\tau}_{j}) \frac{(\boldsymbol{x}_{j}-\boldsymbol{x})^{\boldsymbol{k}}}{\boldsymbol{k}!} w_{j,p}(\boldsymbol{x};h) \qquad \text{(Taylor and (W1))} \\ &= \sum_{j=1}^{p} \left(\sum_{|\boldsymbol{k}|=\gamma} \left(D^{\boldsymbol{k}} \mu(\boldsymbol{\tau}_{j}) - D^{\boldsymbol{k}} \mu(\boldsymbol{x}) \right) \frac{(\boldsymbol{x}_{j}-\boldsymbol{x})^{\boldsymbol{k}}}{\boldsymbol{k}!} \right) w_{j,p}(\boldsymbol{x};h) \qquad \text{(by (W1))} \\ &\leq \sum_{j=1}^{p} \left(\sum_{|\boldsymbol{k}|=\gamma} \| \theta(\boldsymbol{x}_{j}-\boldsymbol{x}) \|_{\infty}^{\alpha-\gamma} \frac{(\boldsymbol{x}_{j}-\boldsymbol{x})^{\boldsymbol{k}}}{\boldsymbol{k}!} \right) w_{j,p}(\boldsymbol{x};h) \\ &\leq \sum_{j=1}^{p} \| \boldsymbol{x}_{j} - \boldsymbol{x} \|_{\infty}^{\alpha} \left(\sum_{|\boldsymbol{k}|=\gamma} \frac{1}{\boldsymbol{k}!} \right) w_{j,p}(\boldsymbol{x};h) \\ &\leq \| \boldsymbol{h} \|_{\infty}^{\alpha} \sum_{|\boldsymbol{k}|=\gamma} \frac{C_{1}}{\boldsymbol{k}!}. \end{split}$$



C Additional numerical results

Figure 10: LP estimator with p = 144 (10-minute data, red line), p = 24 (one houre data, blue dotted/dashed line) and p = 12 (tow hour data, green dashed line). Confidence bands for the p = 144 curve from Liebl and Reimherr (2023).

D Gaussian approximation for non sub-Gaussian errors

In the following we determine the stochastic rate of convergence for the error term $I_2^{n,p,h}$ for errors which only satisfy mild moment assumptions. We restrict ourselves to one-dimensions: d = 1. To his end we consider the situation of Assumption 2. For the Gaussian approximation we need a further property for the weights. Let $C_5 > 0$ be a constant not depending on n, pand h. Then for $x \in [0, 1]$ and $j = 1, \ldots, p - 1$,

(W6)
$$|w_j(x;h) - w_{j+1}(x;h)| \le \frac{C_5}{(p\,h)^2} \mathbf{1}_{[x_j-h,x_j+h]\cup[x_{j+1}-h,x_{j+1}+h]}(x).$$

In Lemma 10 we show that in one dimensions that the weights of the local polynomial estimator satisfy this assumption. Further we replace Assumption 1 as follows. Assumption 6. The random variables $\{\varepsilon_{i,j} \mid 1 \leq i \leq n, 1 \leq j \leq p\}$ are centered, independent and independent of the processes Z_1, \ldots, Z_n . Further we assume that for a $q \geq 4$ it holds $\sup_{i=1,\ldots,n, j=1,\ldots,p} \mathbb{E}[|\varepsilon_{i,j}|^q] \leq C < \infty$ for a constant C > 0.

Proposition 4. In model (1) with d = 1, under Assumption 3 let the errors $\epsilon_{i,j}$ satisfy Assumption 6. Then

$$I^{n,p,h}(x) = \sum_{j=1}^{p} w_j(x;h) \frac{1}{n} \sum_{i=1}^{n} \epsilon_{i,j}, \quad x \in [0,1],$$
(38)

with weights $w_i(x;h)$ satisfying Assumption 2 and (W6) satisfies

$$\left\|I^{n,p,h}\right\|_{\infty} = \mathcal{O}_{\mathbb{P}}\left(\sqrt{\frac{\log(1/h)}{n\,p\,h}} + \frac{p^{1/q}}{\sqrt{n\,p\,h}}\right). \tag{39}$$

Remark 8. Comparing the two rates yields that rate of the Gaussian approximation is slower then the initial rate if $h \leq p^{2/q}/(p \log(1/h)) \sim p^{2/q}/(p \log(p))$ due to

$$\frac{p^{1/q}}{\sqrt{n}\,p\,h} \gtrsim \frac{p^{1/q}}{\sqrt{n}\,p\,h} \cdot \frac{\sqrt{\log(1/h)}}{p^{1/q}} \simeq \sqrt{\frac{\log(1/h)}{n\,p\,h}}$$

and vice versa. Therefore we need to compare the optimal choice for the rate of convergence h^* in (9) with $p^{2/q}/(p \log(1/p))$. This results in

$$p\gtrsim \left(\frac{n}{\log(n)}\right)^{\frac{q}{2(\alpha(q-2)+1)}}$$

If this condition on p is satisfied the \sqrt{n} -rate and the asymptotic normality are still guaranteed. For the worst case covered by the theory, q = 4, this threshold for p is larger then $n^{1/(2\alpha)} \log(n)$ and therefore the \sqrt{n} -rate can only be obtained for ratewise larger p then before. However for large q we have

$$\left(\frac{n}{\log(n)}\right)^{\frac{q}{2(\alpha(q-2)-1)}} \stackrel{q\to\infty}{\longrightarrow} \left(\frac{n}{\log(n)}\right)^{\frac{1}{2\alpha}}$$

and in consequence the initial thresholds from Remark 4 remain valid.

Table 2 contains the quotient of the error term $I_2^{n,p,h}$ with normal (nominator) and standardized t_5 distributed errors (denominator) for different $n, p \in \mathbb{N}$. Even for very small sample sizes, the smoothing already suffices for both terms to behave very similarly. Differences can however be seen without smoothing, which gives a further argument against the use of interpolation estimators.

p n	5	10	25	50
1	1.011	1.018	0.999	1.024
5	1.000	1.022	1.006	1.020
10	1.010	1.007	1.005	0.996
25	0.999	1.013	1.014	1.009
50	1.002	1.013	1.001	0.993

p n	5	10	25	50
1	0.975	0.929	0.858	0.799
5	0.992	0.972	0.933	0.904
10	0.998	0.979	0.962	0.946
25	0.993	0.996	0.979	0.973
50	0.995	1.002	0.990	0.980

Table 2: Quotient of $\left\|\sum_{j=1}^{p} w_{j}(\cdot; h) \bar{\epsilon}_{j,n}\right\|_{\infty}$ Table 3: Quotient of $\max_{j} |\bar{\epsilon}_{j,n}|$ divided by $\max_{j} \left|\frac{1}{n} \sum_{i=1}^{n} t_{i,j}\right|$ with $\bar{\epsilon}_{j,n} \sim \mathcal{N}(0, n^{-1})$ and $\lim_{j \to \infty} w_{j}(\cdot; h) \frac{1}{n} \sum_{i=1}^{n} t_{i,j}\|_{\infty}$ with $t_{i,j} \sim \sqrt{3/5} t_{5}$. $t_{i,j} \sim \sqrt{3/5} t_5.$

Lemma 9. In model (1) under Assumption 3 let X_1, \ldots, X_p be independent random variables which satisfy $\mathbb{E}[X_1] = 0$, $\mathbb{E}[X_1^2] = 1$ and $\mathbb{E}[|X_j|^q] \leq C < \infty$, as required in Assumption 6. Consider the term

$$E^{p,h}(x) = \sum_{j=1}^{p} w_j(x;h) X_j, \quad x \in [0,1],$$
(40)

where the weights $w_i(x;h)$ satisfy Assumption 2 and (W6). This term becomes small with the rate

$$\left\|E^{p,h}\right\|_{\infty} = \mathcal{O}_{\mathbb{P}}\left(\sqrt{\frac{\log(1/h)}{p\,h}} + \frac{p^{1/q}}{p\,h}\right). \tag{41}$$

Proof of Lemma 9. Consider T > 0 and $q \ge 4$. Then from Sakhanenko (1991, Corollary 4, §5) there exists a Brownian motion $(B_t)_{t\in[0,\infty)}$ such that

$$\mathbb{P}\Big(\max_{k=1,\dots,p} \left|\sum_{j=1}^{k} X_{j} - B_{k}\right| > q T p^{1/q} \Big) \leq (p T^{q})^{-1} \sum_{j=1}^{p} \mathbb{E}\big[|X_{j}|^{q}\big] + \mathbb{P}\Big(\max_{j=1,\dots,p} |X_{j}| > T q^{1/q}\Big) \\
\leq \frac{C}{T^{q}} + \mathbb{P}\Big(\max_{j=1,\dots,p} |X_{j}| > T p^{1/q}\Big).$$

Now, by using the independence and the existence of the fourth moment of the errors, we get

$$\mathbb{P}\left(\max_{j=1,\dots,p}|X_j| > T p^{1/q}\right) = 1 - \prod_{j=1}^p \mathbb{P}\left(|X_j| \le T p^{1/q}\right)$$
$$\leq 1 - \prod_{j=1}^p \left(1 - \frac{\mathbb{E}[|X_j|^q]}{p T^q}\right)$$
$$\leq 1 - \left(1 - \frac{C T^{-q}}{p}\right)^p \stackrel{p \to \infty}{\to} 1 - \exp\left(-C T^{-q}\right)$$

Therefore we get

$$\max_{k=1,...,p} \left| \sum_{j=1}^{k} X_j - B_k \right| = \mathcal{O}_{\mathbb{P}}(p^{1/q}).$$
(42)

Now set $N_j := B_j - B_{j-1}, j = 1, ..., p$ with $B_0 = 0$ by writing $\eta_{j,n} := X_j - N_j$ we have $\eta_{j,n} = \sum_{i=1}^j \eta_{i,n} - \sum_{i=1}^{j-1} \eta_{i,n}$. By setting $\sum_{i=1}^0 \eta_{i,n} = 0$ we get

$$\left|\sum_{j=1}^{p} \eta_{j}\right| = \left|\sum_{j=1}^{p} X_{j} - B_{p}\right|.$$
(43)

With the properties of the weights (W3) and (W6) and the previous statement we get

$$\begin{split} \sup_{x \in [0,1]} \left| E^{p,h}(x) - \sum_{j=1}^{p} w_j(x;h) N_j \right| \\ &= \sup_{x \in [0,1]} \left| w_p(x;h) \sum_{j=1}^{p} \eta_j + \sum_{j=1}^{p-1} \left(w_j(x;h) - w_{j+1}(x;h) \right) \sum_{i=1}^{j} \eta_i \right| \\ &\leq \sup_{x \in [0,1]} \max_{j=1,\dots,p} \left| w_j(x;h) \right| \left| \sum_{j=1}^{p} \eta_j \right| \\ &+ \max_{j=1,\dots,p} \left| \sum_{i=1}^{j} \eta_i \right| \sup_{x \in [0,1]} \left| \sum_{j=1}^{p-1} \left(w_j(x;h) - w_{j+1}(x;h) \right) \right| \\ &\leq \sup_{x \in [0,1]} \frac{C_1}{ph} \left| \sum_{j=1}^{p} X_j - B(p) \right| \qquad (\text{with (W3) and (43)}) \\ &+ \max_{j=1,\dots,p} \left| \sum_{i=1}^{j} X_i - B(j) \right| \sup_{x \in [0,1]} \left| \sum_{j=1}^{p-1} \left(w_j(x;h) - w_{j+1}(x;h) \right) \right| \\ &\leq \left(\frac{C_1}{ph} + \frac{C_3 C_5}{ph} \right) \cdot \mathcal{O}_{\mathbb{P}}(p^{1/q}) \quad (\text{with Assumption 3, (42), (W2) and (W6)}) \\ &= \mathcal{O}_{\mathbb{P}}\left(\frac{p^{1/q}}{ph} \right). \end{split}$$

Applying Lemma 2 ii) to the term $\sum_{j=1}^{p} w_j(x;h) N_j$ for the case n = 1 and d = 1 we get that the rate of this term is bounded by $\sqrt{\log(1/h)/(ph)}$ which concludes the proof.

Proof of Proposition 4. Since $\sigma \geq \sigma_j$ for all $j = 1, \ldots, p$, we have

$$\left\|I^{n,p,h}\right\|_{\infty} \leq \frac{\sigma}{\sqrt{n}} \sup_{x \in [0,1]} \left|\sum_{j=1}^{p} w_j(x;h) \frac{1}{\sigma_j \sqrt{n}} \sum_{i=1}^{n} \epsilon_{i,j}\right|.$$

Setting $\tilde{\epsilon}_j = (\sigma_j \sqrt{n})^{-1} \sum_{i=1}^n \epsilon_{i,j}$ we have that $\mathbb{E}[\tilde{\epsilon}_j] = 0$ and $\mathbb{E}[\tilde{\epsilon}_j^2] = 1$ by by Assumption 6. By using the Rosenthal inequality (Rosenthal, 1970) we further get for independent, centered random variables $\mathbb{E}[|\tilde{\epsilon}_j|^q] \leq K_q \max\{C n^{1-q/2}, 1\}$, where $K_q > 0$ is a constant only depending on q. Now $\tilde{\epsilon}_j$ suffice Assumption 6 and therefore the claim follows from Lemma 9.

Lemma 10. Consider the same setup as in Lemma 1 in the one dimensional case. Then the weights the local polynomial estimator also suffice (W6).

Proof. With the upper bound for $||B_{p,h}^{-1}(x)||_{M,2} \le \lambda_0^{-1}$ from (36) and the Lipschitz continuity of $U_m(x)K(x)$ we get

$$\begin{aligned} \left| w_{j}(x;h) - w_{j+1}(x,h) \right| &\leq \frac{1}{ph} \left\| B_{p,h}^{-1}(x) \left(U_{h}(x_{j}-x) K_{h}(x_{j}-x) - U_{h}(x_{j+1}-x) K_{h}(x_{j+1}-x) \right) \right\|_{2} \\ &\leq \frac{L}{\lambda_{0} p h} \frac{|x_{j}-x_{j+1}|}{h} \mathbf{1}_{[x_{j}-h,x_{j}+h] \cup [x_{j+1}-h,x_{j+1}+h]}(x) \\ &\leq \frac{L}{\lambda_{0} f_{\min} p^{2} h^{2}} \mathbf{1}_{[x_{j}-h,x_{j}+h] \cup [x_{j+1}-h,x_{j+1}+h]}(x) \,, \end{aligned}$$

due to the Cauchy-Schwarz inequality, $||U(0)||_2 = 1$ and (29). Setting $C_5 = L/(\lambda_0 f_{\min})$ yields the claim.