

Higgs boson rapidity distribution in bottom annihilation at NNLL and beyond

Goutam Das

*Institut für Theoretische Teilchenphysik und Kosmologie,
RWTH Aachen University, D-52056 Aachen, Germany*

E-mail: goutam@physik.rwth-aachen.de

ABSTRACT: We present precise resummed predictions for Higgs boson rapidity distribution through bottom quark annihilation at next-to-next-to-leading logarithmic (NNLL) accuracy matched to next-to-next-to-leading order (NNLO) and at next-to-next-to-next-to-leading logarithmic (N3LL) accuracy matched to next-to-next-to-next-to-leading order soft-virtual (N3LO_{sv}) in the strong coupling. Exploiting the universal behavior of soft radiation near the threshold, we determine the analytic expressions for the process-dependent and universal perturbative ingredients for threshold resummation in double singular limits of partonic threshold variables z_1, z_2 . Subsequently, the threshold resummation is performed in the double Mellin space within the standard QCD framework. The new third-order process-dependent non-logarithmic coefficients are determined using three-loop bottom quark form factor and third-order quark soft distribution function in rapidity distribution. The effect of these new resummed coefficients are studied at the 13 TeV LHC. We observe a better perturbative convergence in the resummed predictions on the Higgs rapidity spectrum in bottom quark annihilation. We also find that the NNLL and N3LL corrections are sizeable which typically are of the order of -2.5% and -1.5% over the respective available fixed orders with the scale uncertainty remaining at the same level as the fixed order.

KEYWORDS: Resummation, Perturbative QCD

Contents

1	Introduction	1
2	Theoretical Framework	3
3	Numerical Results	6
4	Conclusions	10
A	Anomalous Dimensions	11
B	Yukawa running	12
C	The process dependent coefficient $g_{a,0}^b$	13
D	Soft-virtual coefficients in double Mellin space	14

1 Introduction

The Standard Model (SM) Higgs boson is one of the important fundamental particles to study at colliders like the Large Hadron Collider (LHC). Understanding the Higgs properties is crucial to know the SM well and is also critical to the search of new physics beyond the SM (BSM) where new physics might couple to the Higgs sector. Testing the Higgs properties and understanding its interactions with other fundamental particles are indeed main tasks at the LHC in the upcoming runs. Precision calculations play a prominent role in these studies by calculating the higher-order contributions in the perturbation theory and improving the predictions of Higgs boson properties to a very high accuracy.

The dominant mode of Higgs production at the LHC is through gluon fusion. On the other hand, the Higgs production in bottom annihilation channel, despite being subdominant is also interesting to study. Firstly, Higgs dominantly decays to bottom quarks which can give direct access to the Higgs Yukawa coupling. This purely hadronic final state, however, is challenging to measure [1, 2]. This also requires production of Higgs boson at the first place, through bottom annihilation channel along with the dominant gluon fusion. Secondly, it also gives access to the Higgs Yukawa coupling to bottom quarks even when Higgs is decayed through cleaner channels like di-photons [3, 4] or four-leptons [5, 6] productions. Although in the SM the Higgs Yukawa coupling to bottom quark is suppressed by small bottom mass, in the extensions of SM e.g. two-Higgs doublet models, or minimal supersymmetric standard model the coupling could be enhanced and is crucial to the search for new physics. Therefore, a precise understanding of the SM contribution

will be beneficial in BSM analyses. Thirdly, the Higgs production through bottom annihilation is also interesting on how bottom quark is treated: as whether it is a part of the proton, taking bottom quark as a massless parton except in the Yukawa coupling which is done in the 5 flavor scheme (5FS), or whether it is taken as a massive quark throughout and excluded from the proton structure as is done in 4 flavor scheme (4FS). While in 4FS a massive bottom quark is produced from gluon splitting from proton, in the 5FS scheme, massless bottom has its own parton distributions.

Due to its high importance, Higgs inclusive cross-section is now available theoretically to a very high accuracy to next-to-next-to-next-to leading order (N3LO) [7, 8] in Higgs effective field theory (HEFT) in the gluon fusion channel providing a correction of around 2% with scale uncertainty around 3% reducing from 9% at NNLO [9–11]. It is also known up to N3LO in the vector boson fusion [12–15] where the correction already stabilizes, and the scale uncertainty is found to be below 0.2% at N3LO. Beyond the fixed order, efforts were made to study the dominant threshold contributions by studying the soft-virtual (SV) corrections at fourth order as well as partial subleading logarithmic effects in gluon fusion [16, 17] with a further enhancement of the cross-section ranging from 0.2%-2.7% depending on the scale choices. The fixed order cross-section is further improved by performing threshold resummation at the next-to-next-to-leading logarithmic (NNLL) accuracy [18–21] and to the third logarithmic accuracy (N3LL) [22, 23] as well as resummation of π^2 terms [24] arising from time-like Sudakov form factor. The finite top mass effect is also known at NLO [25, 26], partially at NNLO with top mass expansion [27, 28, 28–30] and recently to exact NNLO [31] where an increment of 0.6% is observed compared to the HEFT approximation. The electroweak corrections are also known to NLO [32, 33] which amount to a positive correction of about 5% compared to the NNLO QCD. Higgs production through bottom annihilation, despite being the subdominant channel, has also been studied extensively in the literature. Due to the availability of third order form factor [34], the soft distributions [35], and the relevant splitting functions [36, 37], the inclusive cross-section is known up to N3LO [38–42] in the 5FS where the residual scale uncertainties are found to be around 5% and to NLO [43–45] in the 4FS. There have been several studies to combine the 5FS and 4FS prediction through different matching prescriptions [46–54]. Further a complete N3LL resummation is performed in [55, 56] where the scale uncertainty at this order reduces to about 4.9%. The pure QED and mixed QCD-QED effects are studied in [57] where the corrections are found to be below 0.03% of the LO.

Differential measurement like the rapidity distribution of the Higgs boson is important to understand Higgs interaction within the SM. Similar to the total production cross-section, rapidity is also inclusive to extra radiations. While this sheds light on the spin of the particle itself, it is also useful to constrain the parton distribution functions (PDFs). In particular, the region with large momentum fraction is not well-constrained where the resummed results could play an important role. Rapidity distribution has been known to NNLO for the Higgs production through gluon fusion [58, 59], as well as in bottom quark annihilation [60, 61]. The accuracy is further extended beyond the NNLO level by studying the threshold contributions up to the third order [62, 63] in gluon fusion as well in to bottom annihilation [64]. It was observed there that the threshold contribu-

tions play a prominent role in the rapidity distributions for these processes. Recently even complete N3LO corrections are also obtained for Higgs [65] and Drell-Yan (DY) [66] rapidities using q_T subtractions [67–70] where the corrections are shown to be around 3% and –2% respectively over NNLO in the central rapidity range. However, unlike the DY case, the uncertainty band for the Higgs rapidity distribution at N3LO show a nice perturbative convergence, reducing the scale uncertainty below 5% and residing within the NNLO uncertainty band. Beyond NNLO, large threshold logarithms are also included at next-to-next-to-leading logarithmic (NNLL) accuracy [71] in the gluon fusion channel. Efforts [72] are also made to resum partial subleading logarithms in the gluon fusion.

In this article we focus on the bottom induced rapidity distribution which is known to NNLO [60, 61] for quite some time. We aim to improve this by including the threshold effects at NNLL and beyond. Within the traditional QCD resummation framework, a formalism [71, 73–77] has been already developed to resum the large threshold logarithms in rapidity distribution for colorless particles. Originally the formalism was proposed for the x_F distribution in the seminal work [73] by Catani and Trentadue. Later it was extended for rapidity¹ following the framework developed in [62, 78]. The idea is to identify proper scaling variables z_1, z_2 corresponding to partonic threshold (z) and rapidity (y_p). One then resums large rapidity logarithms by resumming these scaling variables simultaneously going to the threshold limit $z_1, z_2 \rightarrow 1$. This was termed [71, 75, 76] as the Mellin-Mellin (M-M) approach as the resummation is performed in the double Mellin space corresponding to z_1, z_2 . Essentially in this approach one resums all the double singular terms arising from the delta function $\delta(\bar{z}_i)$ and plus-distributions $\left[\frac{\ln^n \bar{z}_i}{\bar{z}_i}\right]_+$ where $\bar{z}_i = 1 - z_i$. This is also consistent with the generalized threshold resummation approach [79] employing soft-collinear effective theory (SCET) in the double singular limit. A recent comparison for different approaches up to the next-to-leading power (beyond double soft) level can be found in [80].

In this article we follow the standard M-M approach at the leading power within the traditional QCD resummation framework and study the impact of these threshold logarithms for the Higgs boson rapidity distribution in the bottom quark annihilation channel. We organize the paper as follows: in sec. (2) we lay out the theoretical framework for the resummation of rapidity distributions in the M-M approach, in sec. (3) we present the results relevant at the 13 TeV LHC, and finally we conclude in sec. (4) and collect all the analytical results required upto N3LL in the appendices (A - D).

2 Theoretical Framework

The effective Lagrangian for the interaction of a scalar Higgs boson with the bottom quark is given as,

$$\mathcal{L}_{\text{int}}^{(\text{S})} = -\lambda \bar{\psi}_b(x) \psi_b(x) \phi(x), \quad (2.1)$$

where $\psi_b(x)$ and $\phi(x)$ are the bottom quark field and the Higgs field respectively. Here λ is the Yukawa interaction which is given by $\lambda = m_b/v$, with v being the vacuum expectation

¹Note that the threshold behavior is same for both x_F and rapidity.

value (VEV), and m_b being mass of the bottom quark. We follow the 5FS where we use non-zero mass of the bottom quark only in the Yukawa coupling, elsewhere it is treated as a massless quark. The rapidity distribution of the Higgs boson at proton-proton collider takes the following form,

$$\frac{d\sigma(\tau, y)}{dy} = \sum_{i,j=q,\bar{q},g} \int_0^1 dx_1 \int_0^1 dx_2 f_i(x_1, \mu_f) f_j(x_2, \mu_f) \times \int_0^1 dz_1 \int_0^1 dz_2 \delta(x_1^0 - x_1 z_1) \delta(x_2^0 - x_2 z_2) \hat{\sigma}_{d,ij}(z_1, z_2, \mu_f, \mu_r). \quad (2.2)$$

Here $\tau = m_H^2/S = x_1^0 x_2^0$ with S being the hadronic center of mass energy. The hadronic rapidity is defined as $y = \frac{1}{2} \ln(x_1^0/x_2^0)$. The rapidity-dependent partonic coefficient function ($\hat{\sigma}_{d,ij}$) can be decomposed in terms of singular SV piece consisting of plus-distributions and delta function in partonic threshold variables (z_1, z_2) and non-singular or regular piece as,

$$\hat{\sigma}_{d,ij}(z_1, z_2, \mu_f, \mu_r) = \sigma_0(\mu_r) \left(\Delta_{d,ij}^{\text{sv}}(z_1, z_2, \mu_f, \mu_r) + \Delta_{d,ij}^{\text{reg}}(z_1, z_2, \mu_f, \mu_r) \right). \quad (2.3)$$

The singular SV ($\Delta_{d,ij}^{\text{sv}}$) part gets contributions only from the diagonal channel i.e. in the present case $i, j = b, \bar{b}$ for the SV part [81]. On the other hand, the regular terms ($\Delta_{d,ij}^{\text{reg}}$) are subdominant at the threshold and gets contributions from all partonic channels. The overall Born normalization factor $\sigma_0(\mu_r)$ takes the following form,

$$\sigma_0(\mu_r) = \frac{\pi \lambda^2(\mu_r) \tau}{6m_H^2} \equiv \frac{\pi m_b^2(\mu_r) \tau}{6m_H^2 v^2}. \quad (2.4)$$

Note that the μ_r dependence in the Born factor above comes only through the running of bottom mass ($m_b(\mu_r)$) or equivalently the Yukawa ($\lambda(\mu_r)$). The Yukawa running is performed using the mass anomalous dimensions γ_m which we collect in the appendix (B) up to the third order.

Resummation is conveniently performed in the Mellin space where the double Mellin transformation is performed taking Mellin transformation on both partonic threshold variables z_1, z_2 in the following way (suppressing the μ_r, μ_f dependence),

$$\Delta_{d,b\bar{b}}(N_1, N_2) = \int_0^1 dz_1 z_1^{N_1-1} \int_0^1 dz_2 z_2^{N_2-1} \Delta_{d,b\bar{b}}^{\text{sv}}(z_1, z_2). \quad (2.5)$$

Here N_i is the Mellin variable corresponding to partonic threshold z_i . The singular SV contribution can be resummed through the integral form in terms of universal cusp anomalous dimensions A^b and rapidity-dependent threshold non-cusp anomalous dimension D_d^b as well as process-dependent coefficients g_0^b . In the double-Mellin space this can be written in the following integral form,

$$\begin{aligned} \tilde{\Delta}_{d,b\bar{b}}(N_1, N_2) = g_{d,0}^b(a_s) \exp \left(\int_0^1 dz_1 z_1^{N_1-1} \int_0^1 dz_2 z_2^{N_2-1} \right. \\ \left. \times \left(\delta(\bar{z}_2) \left[\frac{1}{\bar{z}_1} \left\{ \int_{\mu_f^2}^{M_H^2 \bar{z}_1} \frac{d\eta^2}{\eta^2} A^b(a_s(\eta^2)) + D_d^b(a_s(M_H^2 \bar{z}_1)) \right\} \right] \right) \right)_+ \end{aligned}$$

$$+ \left[\frac{1}{2\bar{z}_1\bar{z}_2} \left\{ A^b(a_s(z_{12})) + \frac{dD_d^b(a_s(z_{12}))}{d \ln z_{12}} \right\} \right]_+ + (z_1 \leftrightarrow z_2) \Big) \Big), \quad (2.6)$$

where $z_{12} = M_H^2 z_1 z_2$ and $\bar{z}_i = 1 - z_i$. Expansion of the above expression at a fixed order in strong coupling will reproduce the Mellin space fixed order results corresponding to eq. (2.5). Performing the Mellin integration above will produce some non-logarithmic constants in double Mellin space which can be combined with the process-dependent prefactor ($g_{d,0}'^b$) and one can finally organize the large threshold logarithms in the Mellin space in the following form,

$$\tilde{\Delta}_{d,b\bar{b}}(N_1, N_2) = g_{d,0}^b(a_s, \mu_r, \mu_f) \exp \left(G_d^b(a_s, \bar{\omega}, \mu_r, \mu_f) \right). \quad (2.7)$$

The non-logarithmic coefficient ($g_{d,0}^b$) has the following perturbative expansion

$$g_{d,0}^b = 1 + \sum_{i=1}^{\infty} a_s^i g_{d,0}^{b,(i)}. \quad (2.8)$$

To achieve N3LL accuracy, they are needed up to the third order in strong coupling. We have obtained these using the third order $b\bar{b}H$ form factor [34] and third order soft distribution function [35], and we present these in appendix (C). The exponent is universal and resums the large logarithms (which now appear in $N_i \rightarrow \infty$ limit) to all orders. It can be expanded in the strong coupling and the inclusion of successive terms defines the resummed order,²

$$G_d^b(a_s, \bar{\omega}) = g_{d,1}^b(\bar{\omega}) \ln(\bar{N}_1 \bar{N}_2) + \sum_{i=0}^{\infty} a_s^i g_{d,i+2}^b(\bar{\omega}), \quad (2.9)$$

where $\bar{\omega} = a_s \beta_0 \ln(\bar{N}_1 \bar{N}_2)$, with $\bar{N}_i = e^{\gamma_E} N_i, i = 1, 2$. These process-independent resummed exponent are same as the quark-initiated Drell-Yan process and up to N3LL accuracy these can be found e.g. in [82].

The resummed expression in eq. (2.7) only resums the leading singular terms which appear through the large logarithms of N_1 and N_2 in Mellin space. This lacks the subleading regular pieces which can be included through the available fixed order results to improve the resummed predictions. However, one can not simply add them as the fixed order also contains the same logarithmic contributions up to a certain order which are already taken into account in the resummed expression. Therefore, a matching procedure has to be invoked removing these logarithms which also appear in the fixed order. This is done through the following all order matched expression,

$$\begin{aligned} \frac{d\sigma^{\text{res}}}{dy} &= \frac{d\sigma^{\text{f.o.}}}{dy} + \sigma_0(\mu_r) \sum_{k,l=b,\bar{b}} \int_{c_1-i\infty}^{c_1+i\infty} \frac{dN_1}{2\pi i} \int_{c_2-i\infty}^{c_2+i\infty} \frac{dN_2}{2\pi i} e^{y(N_2-N_1)} (\sqrt{\tau})^{-N_1-N_2} \\ &\quad \times \tilde{f}_k(N_1) \tilde{f}_l(N_2) \left[\tilde{\Delta}_{d,b\bar{b}}(N_1, N_2) - \tilde{\Delta}_{d,b\bar{b}}^{\text{f.o.}}(N_1, N_2) \right]. \end{aligned} \quad (2.10)$$

²Note that in the threshold region $\ln(\bar{N}_1 \bar{N}_2) \sim 1/a_s$.

The first term on the right-hand side of the equality is the fixed order contribution containing singular and regular contribution up to a fixed order in strong coupling. The first term inside the square bracket organizes the resummed series up to a certain logarithmic accuracy provided by the knowledge of cusp and rapidity anomalous dimensions as well as the process-dependent $g_{d,0}^b$ coefficients. The symbol ‘f.o.’ in the second term inside the square bracket means the function is truncated to a fixed order in order to avoid double counting of singular terms already present in the fixed order ($\frac{d\sigma^{\text{f.o.}}}{dy}$). Here $\tilde{f}_j(N_i) \equiv \int_0^1 dz_i z_i^{N_i-1} f_j(z_i)$ are the PDFs in the Mellin space. In practice, we use the x -space PDF through the LHAPDF6 [83] interface using the derivatives of PDF as described in [84].

The Mellin inversion in eq. (2.10) is not straightforward as the resummed expression diverges when $\bar{\omega} = 1$. This corresponds to the Landau pole where strong coupling diverges. The perturbative formalism thus break down in this region. One way to proceed is by choosing the contour of the Mellin inversion according to the *minimal prescription* (MP) [85]. The basic idea is to choose the contour in such a way that all the poles remain at the left of the contour except for the Landau pole which remains far right of the contour. In double Mellin space this is little involved as the Landau pole is now a function of two Mellin variables. Typically, one needs to project the complex integration on real variables r_i and chose the contour accordingly. One can still fix the contour of one of the Mellin variable ($N_1 = c_1 + r_1 \exp(i\phi_1)$) according to MP. Once one fixes the contour for N_1 through c_1 and ϕ_1 , the Landau pole is not constrained anymore on the real axis of the second Mellin variable ($N_2 = c_2 + r_2 \exp(i\phi_2)$) and in fact, it now depends on the first Mellin variable (N_1). In order to satisfy the MP, a reasonable choice [74, 82] of the contour for the second Mellin variable as $\phi_2 = \max\left(\pi/2 - 1/2 \arg(\frac{1}{N_1} \exp(\frac{1}{a_S \beta_0} - 2\gamma_E)), 3\pi/4\right)$.

3 Numerical Results

With the setup introduced in the previous section, we now focus on the numerical impact of the threshold logarithms in the rapidity distribution. We focus on 13 TeV LHC with CT14 as our default PDF choice which is used through the LHAPDF6 [83] interface.³ The fixed order results are obtained from [61] using the N-jettiness slicing method [86, 87] as implemented in MCFM [88–90]. The strong coupling is taken through the corresponding PDF sets. At the third order we evolve the strong coupling using four-loop QCD beta function [91–102] for which we set the initial condition as $\alpha_S(m_Z) = 0.118$ where $m_Z = 91.1876$ GeV is the Z boson mass. We set the Higgs mass to be $m_H = 125$ GeV. The central scale choice for this process is taken as $(\mu_r^c, \mu_f^c) = (1, 1/4)m_H$ GeV. The choice of the low μ_f^c scale is done following the observation in [40] to minimize the effect of large collinear logarithms which appear at $\mu_f^c \sim m_H/4$. The Yukawa coupling is also evolved through renormalisation group equation using 4-loop mass anomalous dimensions with bottom mass $m_b(m_b) = 4.18$ GeV. The required mass anomalous dimensions are collected in the appendix (B). To have an estimation of the residual scale uncertainties we follow the standard seven-point scale variations around the central scale choice stated above, with the

³The fixed order results for $b\bar{b}H$ rapidity at NNLO are available with CT14 PDF set from [61].

restriction $1/2 \leq (\mu_r/\mu_r^c)/(\mu_f/\mu_f^c) \leq 2$. This amounts to seven configurations for the scale (μ_r, μ_f) . For each bin, the uncertainty envelope is obtained by considering the maximum and minimum deviations from the central scale choice. Note that the scale variation also includes the scale dependence as arising from the Yukawa running in the $\overline{\text{MS}}$ scheme. The double Mellin inversion in eq. (2.10) is performed with an in house code which we also interface to LHAPDF6 as well as to Cuba [103, 104] for the final integration. Accordingly we chose the contour in eq. (2.10) as $c_1 = c_2 = 1.9$ and $\phi_1 = 3\pi/4$ and ϕ_2 given in the previous section.

We further define the following perturbative quantities in order to assess the higher order effects. We define the ratios K -factor and R -factor [105–107] corresponding to the fixed order and resummed order respectively as,

$$K_{ij} = \left[\frac{d\sigma}{dy} \right]_{\text{NiLO}} / \left[\frac{d\sigma}{dy} \right]_{\text{NjLO}}, R_{ij} = \left[\frac{d\sigma}{dy} \right]_{\text{NiLO+NiLL}} / \left[\frac{d\sigma}{dy} \right]_{\text{NjLO+NjLL}}. \quad (3.1)$$

Further we define the RF_{ij} ratios to estimate the resummed contributions over the fixed orders as,

$$RF_{ij} = \left[\frac{d\sigma}{dy} \right]_{\text{NiLO+NiLL}} / \left[\frac{d\sigma}{dy} \right]_{\text{NjLO}}. \quad (3.2)$$

For such ratios, we calculate the correlated error by taking both the numerator and the denominator at the same scale and obtain the seven-point uncertainties from seven such ratios. In order to shorten the notation, we denote +LL, +NLL, +NNLL to indicate the complete matched resummed results, meaning the resummed corrections are matched to the FO results according to eq. (2.10). At the third order similar notation is adopted where the complete resummed N3LL results are matched to the N3LOsv results and are denoted as +N3LLsv.

In fig. (1), we present the rapidity distribution of the Higgs boson in bottom quark annihilation up to NNLO (left panel) and to +NNLL (right panel). The asymmetric band is obtained by taking the envelope of maximum(minimum) deviation from the central scale according to the seven-point scale variation. On the fixed order side, the NLO gets a correction of similar size to LO, which gets further increased at NNLO by 9.3% compared to NLO in the central rapidity region (at $y = 0 - 1.6$). In the higher rapidity region (at $y = 2 - 4$), the behavior is different where NLO gets negative correction up to 50% compared to LO, whereas on the other hand, the NNLO gets positive correction up to 27% of NLO. The corresponding scale uncertainties are respectively $^{+62.9\%}_{-53.9\%}$ at LO, $^{+22.0\%}_{-31.1\%}$ at NLO, and $^{+1.7\%}_{-20.0\%}$ at NNLO in the central rapidity region. In the resummed case, the convergence is faster with +NLL getting a correction 73% compared to +LL whereas +NNLL gets a further increment of 10.1% compared to +NLL in the central region. Corresponding corrections at the higher rapidity ($y = 3.2$) region are -31.7% and 8.6% respectively. The scale uncertainty does not improve compared to the FO, a behavior which is also seen in the neutral and charged DY productions [82]. In the central region, the asymmetric scale uncertainties are $^{+63.4\%}_{-54.0\%}$ at +LL, $^{+27.1\%}_{-32.2\%}$ at +NLL, $^{+3.8\%}_{-20.4\%}$ at +NNLL respectively. At $y = 3.2$ the corresponding uncertainties are $^{+52.9\%}_{-50.4\%}$ at +LL, $^{+19.8\%}_{-27.6\%}$ at +NLL, $^{+11.9\%}_{-5.4\%}$




Figure 1. Higgs rapidity distribution in bottom annihilation for 13 TeV LHC. The left figure compares different fixed order and matched resummed orders. The notation $+Nk\text{LL}$ indicates that the matched resummed results are obtained by matching with the corresponding FO results. The right figure compares the SV results at the third order against the matched result at the same order where the matching is done with corresponding SV results.

at $+NNLL$ respectively. We notice that while $+LL$ gets a positive contribution ranging from 13.7% to 22.8% over LO, the $+NLL$ gets -3.2% to -2.3% corrections over NLO and $+NNLL$ gets relatively flat correction ranging -2.4% to -3.1% going from central rapidity to higher rapidity. On the right panel of fig. (1), we present the new third order SV results which is further matched with the third order resummed results. While SV results at the third order gets a increment up to 0.5% compared to the NNLO, the scale uncertainty is not improved. The third order matched results however gets a flat correction of about 1.8% throughout the rapidity region compared to $+NNLL$ with the scale uncertainties are at the similar level as the fixed order.




Figure 2. RF factor (as defined in eq. (3.2)) along with correlated errors up to the third order. For the third order, resummed N3LL results are matched to the N3LO_{sv} results.

It is also possible to match the resummed result in a multiplicative way instead of the

additive matching in eq. (2.10). In order to estimate the effect of such a procedure, we followed the prescription⁴ as presented in [108]. We find that the +NLL gets an increment of around 84% at $y = 0$ compared to +LL whereas at +NNLL the correction is less than 1% of +NLL. Similarly, at $y = 3.2$, the corrections are -31% and 5% respectively. Thus, we observe a faster convergence for multiplicative matching compared to the additive matching. The corresponding asymmetric scale uncertainties at $y = 3.2$ are $^{+53.9}_{-50.8}\%$ at +LL, $^{+25.0}_{-33.8}\%$ at +NLL, $^{+6.4}_{-2.9}\%$ at +NNLL respectively. Up to +NLL, the uncertainties remain similar compared to the additive matching, at +NNLL, the uncertainties get better at higher rapidities. On the other hand we see similar scale uncertainties as the additive case at $y = 0$ at +NNLL level. In the rest of the article we follow the additive matching as provided by eq. (2.10).




Figure 3. K, R factors (as defined in eq. (3.1)) along with correlated errors upto the second order (left) and the same at the third order (right). The third order results are up to SV accuracy and same for matched.

A better way to visualize the higher order effects is through the ratios *viz.* the K , R , and RF factors as defined in eq. (3.1)-eq. (3.2) which are presented in fig. (3) and fig. (2). In the left panel we show these ratios up to the second order. It is clear that NLO (or +NLL) corrections shape the rapidity distributions very well, whereas the corrections from NNLO (or +NNLL) are rather flat over a large rapidity region. Compared to the K_{21} factor we observe slight increment of the central scale in the case of R_{21} in the higher rapidity region. On the right panel of fig. (3), we present these ratios at the third order. Again we observe a relatively flat QCD correction over the large rapidity range at N3LOsv amounting to about 8% uncertainty at the higher rapidities. The matched result becomes almost flat even in the higher rapidity region and the correlated scale uncertainty reduces to 4%.

To estimate the intrinsic PDF uncertainty, we define the quantity $\delta\text{PDF} = 1 \pm \delta[\frac{d\sigma}{dy}]/[\frac{d\sigma}{dy}]_0 \times 100\%$, where the numerator $\delta[\frac{d\sigma}{dy}]$ is the intrinsic PDF uncertainty and the denominator $[\frac{d\sigma}{dy}]_0$ is the central prediction. We study only the SV part of the fixed order

⁴See Eq. (4.3) of [108].




Figure 4. Effects of matched NNLL results to NNLOsv are presented and compared against the NNLOsv for CT14 PDF at 13 TeV LHC.

which might be improved by resummed results. This is shown in fig. (4). Notice that the central predictions are different for FO and resummed cases. Indeed, we observe a -2.5% to -2.1% change in the central predictions for resummed case compared to FO SV while going from $y = 0$ to $y = 4$. The PDF uncertainties at the second order (SV) results are about 8% in the central rapidity ($y = 0$) which gets reduced to about 4.5% at $y = -2$. At the higher rapidity region the uncertainties further increases. This is a known behavior which is also observed in the case of DY rapidities [82]. The matched resummed results show a slight improvement (below 0.1%) over the fixed order PDF uncertainty in all rapidity region.

4 Conclusions

At the LHC, soft gluons play an important role in predicting observables correctly in different phase space corners. Particularly in the threshold region their contribution dominates and hence to have a reliable prediction one needs to resum them and match them to the available fixed order to better predict an observable. For rapidity distribution in the threshold region both threshold variables corresponding to partonic threshold and rapidity become equally important and one needs to resum both of them consistently order by order in the perturbation theory. In QCD, this is achieved by the M-M approach, and we exploit this to resum large threshold logarithms to NNLL accuracy matched to available NNLO result. The threshold effects at fixed order amount to -3.2% enhancement over the NLO distribution over a large range of rapidity. On the other hand, the resummed corrections at NNLO+NNLL amount to -2.5% enhancement over the fixed order NNLO. In general, we find a better perturbative convergence in the resummed spectrum which combines merit of both resummed logarithms and non-singular contributions from fixed order which are not captured in purely resummed predictions. While the third order analytic ingredients presented in this article will be useful to match to the third order fixed order results once it will be available, we present the first predictions at the third order coming from the dominant threshold logarithms and also match them to N3LL. We observe

a relatively flat correction in the third order matched results for all rapidities with reduced scale uncertainties to below 4% particularly in the higher rapidities, a region dominated by the large threshold logarithms. Our resummed results can be useful in constraining the bottom quark PDFs at the large momentum fraction.

Acknowledgments

G.D. is grateful to C. Williams for providing the NNLO results. G.D. also thanks V. Ravindran for encouraging to work on this project. Part of the analytical computation has been performed using symbolic manipulation system FORM [109, 110]. Simulations were performed with computing resources granted by RWTH Aachen University under project `rwth1298`. The research of G.D. was supported by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) under grant 396021762 - TRR 257 (*Particle Physics Phenomenology after Higgs discovery*).

A Anomalous Dimensions

The cusp anomalous dimensions have the following perturbative expansion in strong coupling,

$$A^b = \sum_{i=1}^{\infty} a_S^i A_i^b. \quad (\text{A.1})$$

We collect here the coefficients up to fourth order [111–113] needed at N3LL,

$$\begin{aligned} A_1^b &= 4C_F, \\ A_2^b &= 8C_F C_A \left(\frac{67}{18} - \zeta_2 \right) + 8C_F n_f \left(-\frac{5}{9} \right), \\ A_3^b &= 16C_F C_A^2 \left(\frac{245}{24} - \frac{67}{9}\zeta_2 + \frac{11}{6}\zeta_3 + \frac{11}{5}\zeta_2^2 \right) + 16C_F^2 n_f \left(-\frac{55}{24} + 2\zeta_3 \right) + 16C_F C_A n_f \left(-\frac{209}{108} + \frac{10}{9}\zeta_2 - \frac{7}{3}\zeta_3 \right) - 16C_F n_f^2 \left(\frac{1}{27} \right), \\ A_4^b &= C_F n_f^3 \left(-\frac{32}{81} + \frac{64}{27}\zeta_3 \right) + C_F^2 n_f^2 \left(\frac{2392}{81} - \frac{640}{9}\zeta_3 + \frac{64}{5}\zeta_2^2 \right) + C_A C_F n_f^2 \left(\frac{923}{81} + \frac{2240}{27}\zeta_3 - \frac{608}{81}\zeta_2 - \frac{224}{15}\zeta_2^2 \right) + C_F^3 n_f \left(\frac{572}{9} - 320\zeta_5 + \frac{592}{3}\zeta_3 \right) + C_A^2 C_F n_f \left(-\frac{24137}{81} + \frac{2096}{9}\zeta_5 - \frac{23104}{27}\zeta_3 + \frac{20320}{81}\zeta_2 + \frac{448}{3}\zeta_2\zeta_3 - \frac{352}{15}\zeta_2^2 \right) + C_A C_F^2 n_f \left(-\frac{34066}{81} + 160\zeta_5 + \frac{3712}{9}\zeta_3 + \frac{440}{3}\zeta_2 - 128\zeta_2\zeta_3 - \frac{352}{5}\zeta_2^2 \right) + C_A^3 C_F \left(\frac{84278}{81} - \frac{3608}{9}\zeta_5 + \frac{20944}{27}\zeta_3 - 16\zeta_3^2 - \frac{88400}{81}\zeta_2 - \frac{352}{3}\zeta_2\zeta_3 + \frac{3608}{5}\zeta_2^2 - \frac{20032}{105}\zeta_2^3 \right) + n_f \frac{d_{FF}^{(4)}}{N_F} \left(-\frac{1280}{3}\zeta_5 - \frac{256}{3}\zeta_3 + 256\zeta_2 \right) + \frac{d_{FA}^{(4)}}{N_F} \left(\frac{3520}{3}\zeta_5 + \frac{128}{3}\zeta_3 - 384\zeta_3^2 - 128\zeta_2 - \frac{7936}{35}\zeta_2^3 \right). \end{aligned} \quad (\text{A.2})$$

The quartic casimirs appearing above are defined as

$$\begin{aligned}\frac{d_{FA}^{(4)}}{N_F} &\equiv \frac{d_F^{abcd} d_A^{abcd}}{n_c} = \frac{(n_c^2 - 1)(n_c^2 + 6)}{48} = \frac{5}{2}, \\ \frac{d_{FF}^{(4)}}{N_F} &\equiv \frac{d_F^{abcd} d_F^{abcd}}{n_c} = \frac{(n_c^2 - 1)(n_c^4 - 6n_c^2 + 18)}{96n_c^3} = \frac{5}{36},\end{aligned}\tag{A.3}$$

with $N_F = n_c = 3$ for QCD. The threshold non-cusp anomalous dimensions D_d^b has the following perturbative expansion in the strong coupling,

$$D_d^b = \sum_{i=1}^{\infty} a_S^i D_{d,i}^b.\tag{A.4}$$

The coefficients are the same as the quark ones (see e.g. [82]). Up to the N3LL accuracy, these are needed up to third order [62] which we collect below,

$$\begin{aligned}D_{d,1}^b &= 0, \\ D_{d,2}^b &= C_F n_f \left(\frac{112}{27} - \frac{8}{3} \zeta_2 \right) + C_A C_F \left(-\frac{808}{27} + 28 \zeta_3 + \frac{44}{3} \zeta_2 \right), \\ D_{d,3}^b &= C_F n_f^2 \left(-\frac{1856}{729} - \frac{32}{27} \zeta_3 + \frac{160}{27} \zeta_2 \right) + C_A C_F n_f \left(\frac{62626}{729} + \frac{208}{15} \zeta_2^2 - \frac{536}{9} \zeta_3 - \frac{7760}{81} \zeta_2 \right) \\ &\quad + C_F^2 n_f \left(\frac{1711}{27} - \frac{32}{5} \zeta_2^2 - \frac{304}{9} \zeta_3 - 8 \zeta_2 \right) + C_A^2 C_F \left(-\frac{297029}{729} - \frac{616}{15} \zeta_2^2 - 192 \zeta_5 \right. \\ &\quad \left. + \frac{14264}{27} \zeta_3 + \frac{27752}{81} \zeta_2 - \frac{176}{3} \zeta_2 \zeta_3 \right).\end{aligned}\tag{A.5}$$

B Yukawa running

The Higgs Yukawa coupling with bottom quark is given as $\lambda = m_b/v$. Here $m_b(\mu_r)$ is the $\overline{\text{MS}}$ running mass of the bottom quark. Thus, the running of Yukawa goes through the running of bottom mass as,

$$\mu_r^2 \frac{d}{d\mu_r^2} \lambda(\mu_r) = \gamma_m(a_S) \lambda(\mu_r).\tag{B.1}$$

The mass anomalous dimension (γ_m) has the following perturbative expansion:

$$\gamma_m = \sum_{i=0}^{\infty} a_S^{i+1} \gamma_m^{(i)},\tag{B.2}$$

where the coefficients are known up to four loops [99, 114–117]. We collect all the coefficients up to four-loop order,

$$\gamma_m^{(0)} = C_F \left(3 \right),\tag{B.3}$$

$$\gamma_m^{(1)} = C_F^2 \left(\frac{3}{2} \right) + C_A C_F \left(\frac{97}{6} \right) + n_f C_F \left(-\frac{5}{3} \right), \quad (\text{B.4})$$

$$\begin{aligned} \gamma_m^{(2)} = & C_F^3 \left(\frac{129}{2} \right) + C_A C_F^2 \left(-\frac{129}{4} \right) + C_A^2 C_F \left(\frac{11413}{108} \right) + n_f C_F^2 \left(-23 + 24\zeta_3 \right) \\ & + n_f C_A C_F \left(-\frac{278}{27} - 24\zeta_3 \right) + n_f^2 C_F \left(-\frac{35}{27} \right), \end{aligned} \quad (\text{B.5})$$

$$\begin{aligned} \gamma_m^{(3)} = & \frac{d_{FA}}{N_c} \left(\frac{69383}{21} - \frac{10560}{7} \zeta_5 + \frac{16384}{21} \zeta_3 \right) + C_F^4 \left(-\frac{1261}{8} - 336\zeta_3 \right) + C_A C_F^3 \left(\frac{15349}{12} \right. \\ & + 316\zeta_3 \left. \right) + C_A^2 C_F^2 \left(\frac{182015}{252} - \frac{2200}{7} \zeta_5 + \frac{2480}{21} \zeta_3 \right) + n_f C_F^3 \left(\frac{148}{3} - 240\zeta_5 - 444\zeta_3 \right) \\ & + n_f C_A C_F^2 \left(-\frac{13139}{54} + 40\zeta_5 + 784\zeta_3 - \frac{264}{5} \zeta_2^2 \right) + n_f C_A^2 C_F \left(-\frac{59843}{324} + 200\zeta_5 \right. \\ & - \frac{1732}{3} \zeta_3 + \frac{264}{5} \zeta_2^2 \left. \right) + n_f^2 C_F^2 \left(\frac{76}{27} - 40\zeta_3 + \frac{48}{5} \zeta_2^2 \right) + n_f^2 C_A C_F \left(\frac{671}{162} + 40\zeta_3 \right. \\ & - \frac{48}{5} \zeta_2^2 \left. \right) + n_f^3 C_F \left(-\frac{83}{81} + \frac{16}{9} \zeta_3 \right). \end{aligned} \quad (\text{B.6})$$

C The process dependent coefficient $g_{d,0}^b$

Below we present the new process dependent coefficients ($g_{d,0i}^b$) up to N3LL accuracy,

$$g_{d,0}^{b(1)} = C_F \left\{ -4 + 16\zeta_2 + \left(-6 \right) L_{fr} \right\}, \quad (\text{C.1})$$

$$\begin{aligned} g_{d,0}^{b(2)} = & C_F n_f \left\{ \frac{8}{9} + \frac{8}{9} \zeta_3 - \frac{40}{3} \zeta_2 + \left(\frac{2}{3} + \frac{16}{3} \zeta_2 \right) L_{fr} + \left(\frac{16}{3} \zeta_2 \right) L_{qr} + \left(-2 \right) L_{fr}^2 \right\} + C_F^2 \left\{ 16 \right. \\ & - 60\zeta_3 - 32\zeta_2 + \frac{552}{5} \zeta_2^2 + \left(21 - 48\zeta_3 - 72\zeta_2 \right) L_{fr} + \left(48\zeta_3 - 24\zeta_2 \right) L_{qr} \\ & + \left(18 \right) L_{fr}^2 \left. \right\} + C_A C_F \left\{ \frac{166}{9} + \frac{280}{9} \zeta_3 + \frac{256}{3} \zeta_2 - \frac{92}{5} \zeta_2^2 + \left(-12 - 24\zeta_3 \right. \right. \\ & - \frac{88}{3} \zeta_2 \left. \right) L_{qr} + \left(-\frac{17}{3} + 24\zeta_3 - \frac{88}{3} \zeta_2 \right) L_{fr} + \left(11 \right) L_{fr}^2 \left. \right\}, \end{aligned} \quad (\text{C.2})$$

$$\begin{aligned} g_{d,0}^{b(3)} = & C_F n_f^2 \left\{ \frac{16}{27} + \frac{160}{81} \zeta_3 + \frac{256}{27} \zeta_2 + \frac{448}{135} \zeta_2^2 + \left(-\frac{8}{9} \right) L_{fr}^3 + \left(\frac{4}{9} + \frac{32}{9} \zeta_2 \right) L_{fr}^2 + \left(\frac{34}{9} \right. \right. \\ & + \frac{32}{9} \zeta_3 - \frac{160}{27} \zeta_2 \left. \right) L_{fr} + \left(-\frac{64}{27} \zeta_3 - \frac{320}{27} \zeta_2 \right) L_{qr} + \left(\frac{32}{9} \zeta_2 \right) L_{qr}^2 \left. \right\} + C_F^2 n_f \left\{ -\frac{70}{9} \right. \\ & - \frac{608}{9} \zeta_5 + \frac{8872}{27} \zeta_3 - \frac{3428}{27} \zeta_2 - \frac{256}{3} \zeta_2 \zeta_3 - \frac{15688}{135} \zeta_2^2 + \left(-4 - 32\zeta_3 - 48\zeta_2 \right) L_{fr}^2 \\ & + \left(\frac{8}{3} - \frac{656}{3} \zeta_3 + \frac{200}{3} \zeta_2 + \frac{464}{5} \zeta_2^2 \right) L_{qr} + \left(38 + \frac{256}{3} \zeta_3 + 56\zeta_2 + \frac{272}{5} \zeta_2^2 \right) L_{fr} \\ & + \left(32\zeta_3 - 16\zeta_2 \right) L_{qr}^2 + \left(-32\zeta_2 \right) L_{qr} L_{fr} + \left(12 \right) L_{fr}^3 \left. \right\} + C_F^3 \left\{ \frac{1078}{3} + 848\zeta_5 \right. \\ & - 1188\zeta_3 + 32\zeta_3^2 - \frac{166}{3} \zeta_2 - 544\zeta_2 \zeta_3 - \frac{744}{5} \zeta_2^2 + \frac{169504}{315} \zeta_2^3 + \left(-113 + 480\zeta_5 \right. \end{aligned}$$

$$\begin{aligned}
& + 416\zeta_3 + 12\zeta_2 - 704\zeta_2\zeta_3 - \frac{1968}{5}\zeta_2^2) L_{fr} + \left(-100 - 480\zeta_5 - 56\zeta_3 + 132\zeta_2 \right. \\
& + 704\zeta_2\zeta_3 - \frac{1344}{5}\zeta_2^2) L_{qr} + \left(-54 + 288\zeta_3 + 144\zeta_2 \right) L_{fr}^2 + \left(-288\zeta_3 \right. \\
& + 144\zeta_2) L_{qr} L_{fr} + \left(-36 \right) L_{fr}^3 \Big\} + C_A C_F n_f \left\{ -\frac{11540}{81} - 8\zeta_5 - \frac{15944}{81}\zeta_3 \right. \\
& - \frac{13040}{81}\zeta_2 + \frac{880}{9}\zeta_2\zeta_3 + \frac{184}{135}\zeta_2^2 + \left(-40 - \frac{400}{9}\zeta_3 + \frac{2672}{27}\zeta_2 - \frac{8}{5}\zeta_2^2 \right) L_{fr} + \left(-\frac{146}{9} \right. \\
& + 16\zeta_3 - \frac{352}{9}\zeta_2) L_{fr}^2 + \left(-8 - 16\zeta_3 - \frac{352}{9}\zeta_2 \right) L_{qr}^2 + \left(\frac{88}{9} \right) L_{fr}^3 + \left(\frac{196}{3} + \frac{3440}{27}\zeta_3 \right. \\
& + \frac{4480}{27}\zeta_2 - \frac{344}{15}\zeta_2^2) L_{qr} \Big\} + C_A C_F^2 \left\{ -\frac{982}{3} - \frac{3352}{9}\zeta_5 - \frac{11188}{27}\zeta_3 + \frac{592}{3}\zeta_3^2 \right. \\
& + \frac{19658}{27}\zeta_2 + \frac{2528}{3}\zeta_2\zeta_3 + \frac{25676}{27}\zeta_2^2 - \frac{123632}{315}\zeta_2^3 + \left(-\frac{327}{2} - 240\zeta_5 - \frac{2536}{3}\zeta_3 \right. \\
& - 212\zeta_2 + 352\zeta_2\zeta_3 - \frac{1136}{5}\zeta_2^2) L_{fr} + \left(1 + 32\zeta_3 + 264\zeta_2 \right) L_{fr}^2 + \left(72 + 144\zeta_3 \right. \\
& + 176\zeta_2) L_{qr} L_{fr} + \left(\frac{388}{3} + 240\zeta_5 + \frac{3296}{3}\zeta_3 - \frac{1748}{3}\zeta_2 - 352\zeta_2\zeta_3 - 472\zeta_2^2 \right) L_{qr} + \left(\right. \\
& - 176\zeta_3 + 88\zeta_2) L_{qr}^2 + \left(-66 \right) L_{fr}^3 \Big\} + C_A^2 C_F \left\{ \frac{68990}{81} - 84\zeta_5 + \frac{42748}{81}\zeta_3 - \frac{400}{3}\zeta_3^2 \right. \\
& + \frac{39980}{81}\zeta_2 - \frac{7768}{9}\zeta_2\zeta_3 - \frac{25328}{135}\zeta_2^2 + \frac{7088}{63}\zeta_2^3 + \left(-\frac{1180}{3} + 80\zeta_5 - \frac{15472}{27}\zeta_3 \right. \\
& - \frac{12800}{27}\zeta_2 + \frac{1964}{15}\zeta_2^2) L_{qr} + \left(-\frac{242}{9} \right) L_{fr}^3 + \left(44 + 88\zeta_3 + \frac{968}{9}\zeta_2 \right) L_{qr}^2 + \left(\frac{493}{9} \right. \\
& - 88\zeta_3 + \frac{968}{9}\zeta_2) L_{fr}^2 + \left(\frac{1657}{18} - 80\zeta_5 + \frac{3104}{9}\zeta_3 - \frac{8992}{27}\zeta_2 + 4\zeta_2^2 \right) L_{fr} \Big\}. \quad (C.3)
\end{aligned}$$

D Soft-virtual coefficients in double Mellin space

For completeness, here we collect all the singular SV coefficients [81] up to third order in the double Mellin space as defined in eq. (2.5). The perturbative expansion in Mellin space takes the following form,

$$\Delta_{d,b\bar{b}}(N_1, N_2) \equiv \tilde{\Delta}_{d,b\bar{b}}^{\text{f.o.}}(N_1, N_2) = 1 + \sum_{i=1}^{\infty} a_S^i \Delta_{d,b\bar{b}}^{(i)}. \quad (D.1)$$

Defining $\bar{L} \equiv \ln(\bar{N}_1 \bar{N}_2)$, the coefficients up to third order take the form,

$$\begin{aligned}
\Delta_{d,b\bar{b}}^{(1)} &= \bar{L}^2 \left\{ \binom{2}{2} C_F \right\} + \bar{L} \left\{ \left(\binom{-4}{2} L_{qr} + \binom{4}{2} L_{fr} \right) C_F \right\} + g_{d,0}^{b(1)}, \\
\Delta_{d,b\bar{b}}^{(2)} &= \bar{L}^4 \left\{ \binom{2}{2} C_F^2 \right\} + \bar{L}^3 \left\{ \left(\binom{-8}{2} L_{qr} + \binom{8}{2} L_{fr} \right) C_F^2 + \left(-\frac{4}{9} \right) C_F n_f \right. \\
&\quad \left. + \left(\frac{22}{9} \right) C_A C_F \right\} + \bar{L}^2 \left\{ \left(-4\zeta_2 + \left(-\frac{22}{3} \right) L_{qr} + \frac{134}{9} \right) C_A C_F + \left(32\zeta_2 \right. \right.
\end{aligned} \quad (D.2)$$

$$\begin{aligned}
& + \left(-16 \right) L_{qr} L_{fr} + \left(-12 \right) L_{fr} + \left(8 \right) L_{qr}^2 + \left(8 \right) L_{fr}^2 - 8 \Big) C_F^2 + \left(\left(\frac{4}{3} \right) L_{qr} \right. \\
& - \left. \frac{20}{9} \right) C_F n_f \Big\} + \overline{L} \left\{ \left(-28\zeta_3 + \left(-8\zeta_2 + \frac{268}{9} \right) L_{fr} + \left(8\zeta_2 - \frac{268}{9} \right) L_{qr} + \left(\right. \right. \right. \\
& - \left. \left. \frac{22}{3} \right) L_{fr}^2 + \left(\frac{22}{3} \right) L_{qr}^2 + \frac{808}{27} \right) C_A C_F + \left(\left(-64\zeta_2 + 16 \right) L_{qr} + \left(64\zeta_2 - 16 \right) L_{fr} \right. \\
& + \left(-24 \right) L_{fr}^2 + \left(24 \right) L_{qr} L_{fr} \Big) C_F^2 + \left(\left(-\frac{40}{9} \right) L_{fr} + \left(-\frac{4}{3} \right) L_{qr}^2 + \left(\frac{4}{3} \right) L_{fr}^2 \right. \\
& + \left. \left. \left(\frac{40}{9} \right) L_{qr} - \frac{112}{27} \right) C_F n_f \right\} + g_{d,0}^{b(2)}, \tag{D.3}
\end{aligned}$$

$$\begin{aligned}
\Delta_{d,b\bar{b}}^{(3)} = & \overline{L}^6 \left\{ \left(\frac{4}{3} \right) C_F^3 \right\} + \overline{L}^5 \left\{ \left(\left(-8 \right) L_{qr} + \left(8 \right) L_{fr} \right) C_F^3 + \left(-\frac{8}{9} \right) C_F^2 n_f \right. \\
& + \left. \left(\frac{44}{9} \right) C_A C_F^2 \right\} + \overline{L}^4 \left\{ \left(-8\zeta_2 + \left(-\frac{220}{9} \right) L_{qr} + \left(\frac{88}{9} \right) L_{fr} + \frac{268}{9} \right) C_A C_F^2 \right. \\
& + \left(32\zeta_2 + \left(-32 \right) L_{qr} L_{fr} + \left(-12 \right) L_{fr} + \left(16 \right) L_{qr}^2 + \left(16 \right) L_{fr}^2 - 8 \right) C_F^3 + \left(\left(\right. \right. \\
& - \left. \left. \frac{16}{9} \right) L_{fr} + \left(\frac{40}{9} \right) L_{qr} - \frac{40}{9} \right) C_F^2 n_f + \left(-\frac{44}{27} \right) C_A C_F n_f + \left(\frac{4}{27} \right) C_F n_f^2 \\
& + \left(\frac{121}{27} \right) C_A^2 C_F \Big\} + \overline{L}^3 \left\{ \left(-\frac{88}{9} \zeta_2 + \left(-\frac{484}{27} \right) L_{qr} + \frac{3560}{81} \right) C_A^2 C_F + \left(-\frac{64}{9} \zeta_2 \right. \right. \\
& + \left(-\frac{136}{9} \right) L_{fr} + \left(\frac{8}{3} \right) L_{fr}^2 + \left(\frac{16}{3} \right) L_{qr} L_{fr} + \left(\frac{160}{9} \right) L_{qr} + \left(-8 \right) L_{qr}^2 \\
& - \frac{212}{27} \Big) C_F^2 n_f + \left(\frac{16}{9} \zeta_2 + \left(\frac{176}{27} \right) L_{qr} - \frac{1156}{81} \right) C_A C_F n_f + \left(\frac{352}{9} \zeta_2 - 56\zeta_3 + \left(\right. \right. \\
& - \left. \left. 32\zeta_2 + \frac{940}{9} \right) L_{fr} + \left(32\zeta_2 - \frac{1072}{9} \right) L_{qr} + \left(-\frac{88}{3} \right) L_{qr} L_{fr} + \left(-\frac{44}{3} \right) L_{fr}^2 \\
& + \left(44 \right) L_{qr}^2 + \frac{1352}{27} \Big) C_A C_F^2 + \left(\left(-128\zeta_2 + 32 \right) L_{qr} + \left(128\zeta_2 - 32 \right) L_{fr} \right. \\
& + \left(\left(-32 \right) L_{fr} + \left(32 \right) L_{qr} + 48 \right) L_{qr} L_{fr} + \left(-\frac{32}{3} \right) L_{qr}^3 + \left(\frac{32}{3} \right) L_{fr}^3 \\
& + \left(-48 \right) L_{fr}^2 \Big) C_F^3 + \left(\left(-\frac{16}{27} \right) L_{qr} + \frac{80}{81} \right) C_F n_f^2 \Big\} + \overline{L}^2 \left\{ \left(-\frac{504}{5} \zeta_2^2 + \frac{3824}{9} \zeta_2 \right. \right. \\
& + \frac{560}{9} \zeta_3 + \left(-176\zeta_2 + 64\zeta_3 - \frac{3088}{27} \right) L_{qr} + \left(-\frac{104}{3} \zeta_2 - 64\zeta_3 + \frac{514}{27} \right) L_{fr} + \left(\right. \\
& - \left. 32\zeta_2 + \frac{1072}{9} \right) L_{qr}^2 + \left(-32\zeta_2 + \frac{1270}{9} \right) L_{fr}^2 + \left(64\zeta_2 + \left(\frac{88}{3} \right) L_{qr} + \left(\frac{88}{3} \right) L_{fr} \right. \\
& - \left. \frac{1748}{9} \right) L_{qr} L_{fr} + \left(-\frac{88}{3} \right) L_{qr}^3 + \left(-\frac{88}{3} \right) L_{fr}^3 - \frac{68}{3} \Big) C_A C_F^2 + \left(\frac{88}{5} \zeta_2^2 - \frac{536}{9} \zeta_2 \right. \\
& - 88\zeta_3 + \left(\frac{88}{3} \zeta_2 - \frac{3560}{27} \right) L_{qr} + \left(\frac{242}{9} \right) L_{qr}^2 + \frac{15503}{81} \Big) C_A^2 C_F + \left(\frac{1104}{5} \zeta_2^2 - 64\zeta_2 \right. \\
& - \left. 120\zeta_3 + \left(-256\zeta_2 + \left(-48 \right) L_{qr} + \left(96 \right) L_{fr} + 64 \right) L_{qr} L_{fr} + \left(-144\zeta_2 - 96\zeta_3 \right. \right.
\end{aligned}$$

$$\begin{aligned}
& + 42) L_{fr} + \left(-48\zeta_2 + 96\zeta_3 \right) L_{qr} + \left(128\zeta_2 - 32 \right) L_{qr}^2 + \left(128\zeta_2 + 4 \right) L_{fr}^2 \\
& + \left(-48 \right) L_{fr}^3 + 32) C_F^3 + \left(-\frac{560}{9}\zeta_2 + \frac{160}{9}\zeta_3 + \left(\frac{32}{3}\zeta_2 - \frac{52}{27} \right) L_{fr} + \left(32\zeta_2 \right. \right. \\
& + \left. \frac{412}{27} \right) L_{qr} + \left(\left(-\frac{16}{3} \right) L_{qr} + \left(-\frac{16}{3} \right) L_{fr} + \frac{248}{9} \right) L_{qr} L_{fr} + \left(-\frac{196}{9} \right) L_{fr}^2 + \left(\right. \\
& - \left. \frac{160}{9} \right) L_{qr}^2 + \left(\frac{16}{3} \right) L_{qr}^3 + \left(\frac{16}{3} \right) L_{fr}^3 - \frac{23}{3} \Big) C_F^2 n_f + \left(\frac{80}{9}\zeta_2 + \left(-\frac{16}{3}\zeta_2 \right. \right. \\
& + \left. \frac{1156}{27} \right) L_{qr} + \left(-\frac{88}{9} \right) L_{qr}^2 - \frac{4102}{81} \Big) C_A C_F n_f + \left(\left(-\frac{80}{27} \right) L_{qr} + \left(\frac{8}{9} \right) L_{qr}^2 \right. \\
& + \left. \frac{200}{81} \right) C_F n_f^2 \Big\} + \bar{L} \left\{ \left(-\frac{88}{5}\zeta_2^2 + \frac{176}{3}\zeta_2\zeta_3 - \frac{6392}{81}\zeta_2 - \frac{12328}{27}\zeta_3 + 192\zeta_5 + \left(\right. \right. \right. \\
& - \left. \frac{176}{5}\zeta_2^2 + \frac{1072}{9}\zeta_2 + 176\zeta_3 - \frac{31006}{81} \right) L_{qr} + \left(\frac{176}{5}\zeta_2^2 - \frac{1072}{9}\zeta_2 + \frac{88}{3}\zeta_3 + \frac{490}{3} \right) L_{fr} \\
& + \left(-\frac{88}{3}\zeta_2 + \frac{3560}{27} \right) L_{qr}^2 + \left(\frac{88}{3}\zeta_2 - \frac{3560}{27} \right) L_{fr}^2 + \left(-\frac{484}{27} \right) L_{qr}^3 + \left(\frac{484}{27} \right) L_{fr}^3 \\
& + \frac{297029}{729} \Big) C_A^2 C_F + \left(-\frac{16}{5}\zeta_2^2 + \frac{824}{81}\zeta_2 + \frac{904}{27}\zeta_3 + \left(-\frac{160}{9}\zeta_2 + \frac{8204}{81} \right) L_{qr} + \left(\right. \right. \\
& - \left. \frac{16}{3}\zeta_2 + \frac{1156}{27} \right) L_{fr}^2 + \left(\frac{16}{3}\zeta_2 - \frac{1156}{27} \right) L_{qr}^2 + \left(\frac{160}{9}\zeta_2 - \frac{112}{3}\zeta_3 - \frac{836}{27} \right) L_{fr} + \left(\right. \\
& - \left. \frac{176}{27} \right) L_{fr}^3 + \left(\frac{176}{27} \right) L_{qr}^3 - \frac{62626}{729} \Big) C_A C_F n_f + \left(\frac{32}{5}\zeta_2^2 - \frac{1792}{27}\zeta_2 + \frac{304}{9}\zeta_3 + \left(\right. \right. \\
& - \left. \frac{1120}{9}\zeta_2 + \frac{320}{9}\zeta_3 + \frac{86}{9} \right) L_{fr} + \left(-\frac{128}{3}\zeta_2 + \frac{4}{3} \right) L_{qr}^2 + \left(\frac{128}{3}\zeta_2 + 28 \right) L_{fr}^2 \\
& + \left(\frac{1120}{9}\zeta_2 - \frac{320}{9}\zeta_3 + \frac{46}{3} \right) L_{qr} + \left(\left(8 \right) L_{qr} + \left(8 \right) L_{fr} - \frac{88}{3} \right) L_{qr} L_{fr} \\
& + \left(-16 \right) L_{fr}^3 - \frac{421}{9} \Big) C_F^2 n_f + \left(-448\zeta_2\zeta_3 + \frac{12928}{27}\zeta_2 + 112\zeta_3 + \left(-\frac{1008}{5}\zeta_2^2 \right. \right. \\
& + \frac{7648}{9}\zeta_2 + \frac{2632}{9}\zeta_3 - \frac{2024}{9} \Big) L_{fr} + \left(\frac{1008}{5}\zeta_2^2 - \frac{7648}{9}\zeta_2 - \frac{1120}{9}\zeta_3 + \frac{136}{3} \right) L_{qr} + \left(\right. \\
& - \left. \frac{560}{3}\zeta_2 + 96\zeta_3 - 172 \right) L_{fr}^2 + \left(-48\zeta_2 - 192\zeta_3 + \left(-44 \right) L_{qr} + \left(-44 \right) L_{fr} \right. \\
& + \left. \frac{460}{3} \right) L_{qr} L_{fr} + \left(\frac{704}{3}\zeta_2 + 96\zeta_3 + \frac{56}{3} \right) L_{qr}^2 + \left(88 \right) L_{fr}^3 - \frac{3232}{27} \Big) C_A C_F^2 + \left(\frac{32}{9}\zeta_3 \right. \\
& + \left(-\frac{400}{81} \right) L_{qr} + \left(-\frac{80}{27} \right) L_{fr}^2 + \left(-\frac{16}{27} \right) L_{qr}^3 + \left(-\frac{16}{27} \right) L_{fr} + \left(\frac{16}{27} \right) L_{fr}^3 \\
& + \left(\frac{80}{27} \right) L_{qr}^2 + \frac{1856}{729} \Big) C_F n_f^2 + \left(\left(-\frac{2208}{5}\zeta_2^2 + 128\zeta_2 + 240\zeta_3 - 64 \right) L_{qr} \right. \\
& + \left(\frac{2208}{5}\zeta_2^2 - 128\zeta_2 - 240\zeta_3 + 64 \right) L_{fr} + \left(-288\zeta_2 - 192\zeta_3 + 84 \right) L_{fr}^2 + \left(96\zeta_2 \right. \\
& - \left. 192\zeta_3 \right) L_{qr}^2 + \left(192\zeta_2 + 384\zeta_3 + \left(-72 \right) L_{fr} - 84 \right) L_{qr} L_{fr} + \left(72 \right) L_{fr}^3 \Big) C_F^3 \Big\}
\end{aligned}$$

$$+ g_{d,0}^{b(3)} . \tag{D.4}$$

References

- [1] CMS collaboration, A. M. Sirunyan et al., *Observation of Higgs boson decay to bottom quarks*, *Phys. Rev. Lett.* **121** (2018) 121801 [[1808.08242](#)].
- [2] ATLAS collaboration, M. Aaboud et al., *Observation of $H \rightarrow b\bar{b}$ decays and VH production with the ATLAS detector*, *Phys. Lett. B* **786** (2018) 59 [[1808.08238](#)].
- [3] ATLAS collaboration, M. Aaboud et al., *Measurements of Higgs boson properties in the diphoton decay channel with 36 fb^{-1} of pp collision data at $\sqrt{s} = 13 \text{ TeV}$ with the ATLAS detector*, *Phys. Rev. D* **98** (2018) 052005 [[1802.04146](#)].
- [4] CMS collaboration, A. M. Sirunyan et al., *A measurement of the Higgs boson mass in the diphoton decay channel*, *Phys. Lett. B* **805** (2020) 135425 [[2002.06398](#)].
- [5] ATLAS collaboration, M. Aaboud et al., *Measurement of the four-lepton invariant mass spectrum in 13 TeV proton-proton collisions with the ATLAS detector*, *JHEP* **04** (2019) 048 [[1902.05892](#)].
- [6] CMS collaboration, A. M. Sirunyan et al., *Measurements of production cross sections of the Higgs boson in the four-lepton final state in proton-proton collisions at $\sqrt{s} = 13 \text{ TeV}$* , *Eur. Phys. J. C* **81** (2021) 488 [[2103.04956](#)].
- [7] C. Anastasiou, C. Duhr, F. Dulat, F. Herzog and B. Mistlberger, *Higgs Boson Gluon-Fusion Production in QCD at Three Loops*, *Phys. Rev. Lett.* **114** (2015) 212001 [[1503.06056](#)].
- [8] B. Mistlberger, *Higgs boson production at hadron colliders at $N^3\text{LO}$ in QCD*, *JHEP* **05** (2018) 028 [[1802.00833](#)].
- [9] R. V. Harlander and W. B. Kilgore, *Next-to-next-to-leading order Higgs production at hadron colliders*, *Phys. Rev. Lett.* **88** (2002) 201801 [[hep-ph/0201206](#)].
- [10] C. Anastasiou and K. Melnikov, *Higgs boson production at hadron colliders in NNLO QCD*, *Nucl. Phys. B* **646** (2002) 220 [[hep-ph/0207004](#)].
- [11] V. Ravindran, J. Smith and W. L. van Neerven, *NNLO corrections to the total cross-section for Higgs boson production in hadron hadron collisions*, *Nucl. Phys. B* **665** (2003) 325 [[hep-ph/0302135](#)].
- [12] P. Bolzoni, F. Maltoni, S. Moch and M. Zaro, *Higgs production via vector-boson fusion at NNLO in QCD*, *Phys. Rev. Lett.* **105** (2010) 011801 [[1003.4451](#)].
- [13] P. Bolzoni, F. Maltoni, S. Moch and M. Zaro, *Vector boson fusion at NNLO in QCD: SM Higgs and beyond*, *Phys. Rev. D* **85** (2012) 035002 [[1109.3717](#)].
- [14] F. A. Dreyer and A. Karlberg, *Vector-Boson Fusion Higgs Production at Three Loops in QCD*, *Phys. Rev. Lett.* **117** (2016) 072001 [[1606.00840](#)].
- [15] A. Buckley et al., *A comparative study of Higgs boson production from vector-boson fusion*, *JHEP* **11** (2021) 108 [[2105.11399](#)].
- [16] G. Das, S. Moch and A. Vogt, *Soft corrections to inclusive deep-inelastic scattering at four loops and beyond*, *JHEP* **03** (2020) 116 [[1912.12920](#)].
- [17] G. Das, S. Moch and A. Vogt, *Approximate four-loop QCD corrections to the Higgs-boson production cross section*, *Phys. Lett. B* **807** (2020) 135546 [[2004.00563](#)].

- [18] S. Catani, D. de Florian, M. Grazzini and P. Nason, *Soft gluon resummation for Higgs boson production at hadron colliders*, *JHEP* **07** (2003) 028 [[hep-ph/0306211](#)].
- [19] S. Moch and A. Vogt, *Higher-order soft corrections to lepton pair and Higgs boson production*, *Phys. Lett. B* **631** (2005) 48 [[hep-ph/0508265](#)].
- [20] E. Laenen and L. Magnea, *Threshold resummation for electroweak annihilation from DIS data*, *Phys. Lett. B* **632** (2006) 270 [[hep-ph/0508284](#)].
- [21] A. Idilbi, X.-d. Ji, J.-P. Ma and F. Yuan, *Threshold resummation for Higgs production in effective field theory*, *Phys. Rev. D* **73** (2006) 077501 [[hep-ph/0509294](#)].
- [22] M. Bonvini and S. Marzani, *Resummed Higgs cross section at N^3LL* , *JHEP* **09** (2014) 007 [[1405.3654](#)].
- [23] M. Bonvini, S. Marzani, C. Muselli and L. Rottoli, *On the Higgs cross section at N^3LO+N^3LL and its uncertainty*, *JHEP* **08** (2016) 105 [[1603.08000](#)].
- [24] M. A. Ebert, J. K. L. Michel and F. J. Tackmann, *Resummation Improved Rapidity Spectrum for Gluon Fusion Higgs Production*, *JHEP* **05** (2017) 088 [[1702.00794](#)].
- [25] D. Graudenz, M. Spira and P. M. Zerwas, *QCD corrections to Higgs boson production at proton proton colliders*, *Phys. Rev. Lett.* **70** (1993) 1372.
- [26] M. Spira, A. Djouadi, D. Graudenz and P. M. Zerwas, *Higgs boson production at the LHC*, *Nucl. Phys. B* **453** (1995) 17 [[hep-ph/9504378](#)].
- [27] S. Marzani, R. D. Ball, V. Del Duca, S. Forte and A. Vicini, *Higgs production via gluon-gluon fusion with finite top mass beyond next-to-leading order*, *Nucl. Phys. B* **800** (2008) 127 [[0801.2544](#)].
- [28] A. Pak, M. Rogal and M. Steinhauser, *Finite top quark mass effects in NNLO Higgs boson production at LHC*, *JHEP* **02** (2010) 025 [[0911.4662](#)].
- [29] R. V. Harlander, H. Mantler, S. Marzani and K. J. Ozeren, *Higgs production in gluon fusion at next-to-next-to-leading order QCD for finite top mass*, *Eur. Phys. J. C* **66** (2010) 359 [[0912.2104](#)].
- [30] R. V. Harlander and K. J. Ozeren, *Finite top mass effects for hadronic Higgs production at next-to-next-to-leading order*, *JHEP* **11** (2009) 088 [[0909.3420](#)].
- [31] M. Czakon, R. V. Harlander, J. Klappert and M. Niggetiedt, *Exact Top-Quark Mass Dependence in Hadronic Higgs Production*, *Phys. Rev. Lett.* **127** (2021) 162002 [[2105.04436](#)].
- [32] U. Aglietti, R. Bonciani, G. Degrossi and A. Vicini, *Two loop light fermion contribution to Higgs production and decays*, *Phys. Lett. B* **595** (2004) 432 [[hep-ph/0404071](#)].
- [33] S. Actis, G. Passarino, C. Sturm and S. Uccirati, *NLO Electroweak Corrections to Higgs Boson Production at Hadron Colliders*, *Phys. Lett. B* **670** (2008) 12 [[0809.1301](#)].
- [34] T. Gehrmann and D. Kara, *The $Hb\bar{b}$ form factor to three loops in QCD*, *JHEP* **09** (2014) 174 [[1407.8114](#)].
- [35] C. Anastasiou, C. Duhr, F. Dulat, E. Furlan, T. Gehrmann, F. Herzog et al., *Higgs boson gluon-fusion production at threshold in N^3LO QCD*, *Phys. Lett. B* **737** (2014) 325 [[1403.4616](#)].
- [36] A. Vogt, S. Moch and J. A. M. Vermaseren, *The Three-loop splitting functions in QCD: The Singlet case*, *Nucl. Phys. B* **691** (2004) 129 [[hep-ph/0404111](#)].

- [37] S. Moch, J. A. M. Vermaseren and A. Vogt, *The Three loop splitting functions in QCD: The Nonsinglet case*, *Nucl. Phys. B* **688** (2004) 101 [[hep-ph/0403192](#)].
- [38] D. Dicus, T. Stelzer, Z. Sullivan and S. Willenbrock, *Higgs boson production in association with bottom quarks at next-to-leading order*, *Phys. Rev. D* **59** (1999) 094016 [[hep-ph/9811492](#)].
- [39] C. Balazs, H.-J. He and C. P. Yuan, *QCD corrections to scalar production via heavy quark fusion at hadron colliders*, *Phys. Rev. D* **60** (1999) 114001 [[hep-ph/9812263](#)].
- [40] F. Maltoni, Z. Sullivan and S. Willenbrock, *Higgs-Boson Production via Bottom-Quark Fusion*, *Phys. Rev. D* **67** (2003) 093005 [[hep-ph/0301033](#)].
- [41] R. V. Harlander and W. B. Kilgore, *Higgs boson production in bottom quark fusion at next-to-next-to leading order*, *Phys. Rev. D* **68** (2003) 013001 [[hep-ph/0304035](#)].
- [42] C. Duhr, F. Dulat and B. Mistlberger, *Higgs Boson Production in Bottom-Quark Fusion to Third Order in the Strong Coupling*, *Phys. Rev. Lett.* **125** (2020) 051804 [[1904.09990](#)].
- [43] S. Dittmaier, M. Krämer and M. Spira, *Higgs radiation off bottom quarks at the Tevatron and the CERN LHC*, *Phys. Rev. D* **70** (2004) 074010 [[hep-ph/0309204](#)].
- [44] S. Dawson, C. B. Jackson, L. Reina and D. Wackerroth, *Exclusive Higgs boson production with bottom quarks at hadron colliders*, *Phys. Rev. D* **69** (2004) 074027 [[hep-ph/0311067](#)].
- [45] M. Wiesemann, R. Frederix, S. Frixione, V. Hirschi, F. Maltoni and P. Torrielli, *Higgs production in association with bottom quarks*, *JHEP* **02** (2015) 132 [[1409.5301](#)].
- [46] M. A. G. Aivazis, J. C. Collins, F. I. Olness and W.-K. Tung, *Leptoproduction of heavy quarks. 2. A Unified QCD formulation of charged and neutral current processes from fixed target to collider energies*, *Phys. Rev. D* **50** (1994) 3102 [[hep-ph/9312319](#)].
- [47] M. Cacciari, M. Greco and P. Nason, *The P_T spectrum in heavy flavor hadroproduction*, *JHEP* **05** (1998) 007 [[hep-ph/9803400](#)].
- [48] S. Forte, E. Laenen, P. Nason and J. Rojo, *Heavy quarks in deep-inelastic scattering*, *Nucl. Phys. B* **834** (2010) 116 [[1001.2312](#)].
- [49] R. Harlander, M. Kramer and M. Schumacher, *Bottom-quark associated Higgs-boson production: reconciling the four- and five-flavour scheme approach*, **1112.3478**.
- [50] M. Bonvini, A. S. Papanastasiou and F. J. Tackmann, *Resummation and matching of b -quark mass effects in $b\bar{b}H$ production*, *JHEP* **11** (2015) 196 [[1508.03288](#)].
- [51] S. Forte, D. Napoletano and M. Ubiali, *Higgs production in bottom-quark fusion in a matched scheme*, *Phys. Lett. B* **751** (2015) 331 [[1508.01529](#)].
- [52] M. Bonvini, A. S. Papanastasiou and F. J. Tackmann, *Matched predictions for the $b\bar{b}H$ cross section at the 13 TeV LHC*, *JHEP* **10** (2016) 053 [[1605.01733](#)].
- [53] S. Forte, D. Napoletano and M. Ubiali, *Higgs production in bottom-quark fusion: matching beyond leading order*, *Phys. Lett. B* **763** (2016) 190 [[1607.00389](#)].
- [54] C. Duhr, F. Dulat, V. Hirschi and B. Mistlberger, *Higgs production in bottom quark fusion: matching the 4- and 5-flavour schemes to third order in the strong coupling*, *JHEP* **08** (2020) 017 [[2004.04752](#)].
- [55] A. H. Ajjath, A. Chakraborty, G. Das, P. Mukherjee and V. Ravindran, *Resummed prediction for Higgs boson production through $b\bar{b}$ annihilation at N^3LL* , *JHEP* **11** (2019) 006 [[1905.03771](#)].

- [56] G. Das, C. Dey, M. C. Kumar and K. Samanta, *Threshold enhanced cross sections for colorless productions*, *Phys. Rev. D* **107** (2023) 034038 [[2210.17534](#)].
- [57] A. H. Ajjath, P. Banerjee, A. Chakraborty, P. K. Dhani, P. Mukherjee, N. Rana et al., *NNLO $QCD \oplus QED$ corrections to Higgs production in bottom quark annihilation*, *Phys. Rev. D* **100** (2019) 114016 [[1906.09028](#)].
- [58] C. Anastasiou, K. Melnikov and F. Petriello, *Higgs boson production at hadron colliders: Differential cross sections through next-to-next-to-leading order*, *Phys. Rev. Lett.* **93** (2004) 262002 [[hep-ph/0409088](#)].
- [59] C. Anastasiou, K. Melnikov and F. Petriello, *Fully differential Higgs boson production and the di-photon signal through next-to-next-to-leading order*, *Nucl. Phys. B* **724** (2005) 197 [[hep-ph/0501130](#)].
- [60] S. Bühler, F. Herzog, A. Lazopoulos and R. Müller, *The fully differential hadronic production of a Higgs boson via bottom quark fusion at NNLO*, *JHEP* **07** (2012) 115 [[1204.4415](#)].
- [61] R. Mondini and C. Williams, *Bottom-induced contributions to Higgs plus jet at next-to-next-to-leading order*, *JHEP* **05** (2021) 045 [[2102.05487](#)].
- [62] V. Ravindran, J. Smith and W. L. van Neerven, *QCD threshold corrections to di-lepton and Higgs rapidity distributions beyond N^2 LO*, *Nucl. Phys. B* **767** (2007) 100 [[hep-ph/0608308](#)].
- [63] T. Ahmed, M. K. Mandal, N. Rana and V. Ravindran, *Rapidity Distributions in Drell-Yan and Higgs Productions at Threshold to Third Order in QCD*, *Phys. Rev. Lett.* **113** (2014) 212003 [[1404.6504](#)].
- [64] T. Ahmed, M. K. Mandal, N. Rana and V. Ravindran, *Higgs Rapidity Distribution in $b\bar{b}$ Annihilation at Threshold in N^3 LO QCD*, *JHEP* **02** (2015) 131 [[1411.5301](#)].
- [65] L. Cieri, X. Chen, T. Gehrmann, E. W. N. Glover and A. Huss, *Higgs boson production at the LHC using the q_T subtraction formalism at N^3 LO QCD*, *JHEP* **02** (2019) 096 [[1807.11501](#)].
- [66] X. Chen, T. Gehrmann, N. Glover, A. Huss, T.-Z. Yang and H. X. Zhu, *Dilepton Rapidity Distribution in Drell-Yan Production to Third Order in QCD*, *Phys. Rev. Lett.* **128** (2022) 052001 [[2107.09085](#)].
- [67] S. Catani and M. Grazzini, *An NNLO subtraction formalism in hadron collisions and its application to Higgs boson production at the LHC*, *Phys. Rev. Lett.* **98** (2007) 222002 [[hep-ph/0703012](#)].
- [68] S. Catani, L. Cieri, G. Ferrera, D. de Florian and M. Grazzini, *Vector boson production at hadron colliders: a fully exclusive QCD calculation at NNLO*, *Phys. Rev. Lett.* **103** (2009) 082001 [[0903.2120](#)].
- [69] S. Catani, G. Ferrera and M. Grazzini, *W Boson Production at Hadron Colliders: The Lepton Charge Asymmetry in NNLO QCD*, *JHEP* **05** (2010) 006 [[1002.3115](#)].
- [70] S. Catani, L. Cieri, D. de Florian, G. Ferrera and M. Grazzini, *Diphoton production at hadron colliders: a fully-differential QCD calculation at NNLO*, *Phys. Rev. Lett.* **108** (2012) 072001 [[1110.2375](#)].
- [71] P. Banerjee, G. Das, P. K. Dhani and V. Ravindran, *Threshold resummation of the rapidity*

- distribution for Higgs production at NNLO+NNLL, *Phys. Rev. D* **97** (2018) 054024 [[1708.05706](#)].
- [72] A. H. Ajjath, P. Mukherjee, V. Ravindran, A. Sankar and S. Tiwari, *Next-to-soft corrections for Drell-Yan and Higgs boson rapidity distributions beyond N^3 LO*, *Phys. Rev. D* **103** (2021) L111502 [[2010.00079](#)].
 - [73] S. Catani and L. Trentadue, *Resummation of the QCD Perturbative Series for Hard Processes*, *Nucl. Phys. B* **327** (1989) 323.
 - [74] D. Westmark and J. F. Owens, *Enhanced threshold resummation formalism for lepton pair production and its effects in the determination of parton distribution functions*, *Phys. Rev. D* **95** (2017) 056024 [[1701.06716](#)].
 - [75] P. Banerjee, G. Das, P. K. Dhani and V. Ravindran, *Threshold resummation of the rapidity distribution for Drell-Yan production at NNLO+NNLL*, *Phys. Rev. D* **98** (2018) 054018 [[1805.01186](#)].
 - [76] P. Banerjee, G. Das, P. K. Dhani and V. Ravindran, *Threshold resummation in the rapidity distribution for a colorless particle production at the LHC*, *PoS* **LL2018** (2018) 043 [[1807.04583](#)].
 - [77] T. Ahmed, A. A. H., P. Mukherjee, V. Ravindran and A. Sankar, *Rapidity distribution at soft-virtual and beyond for n -colorless particles to N^4 LO in QCD*, *Eur. Phys. J. C* **81** (2021) 943 [[2010.02980](#)].
 - [78] V. Ravindran and J. Smith, *Threshold corrections to rapidity distributions of Z and $W^\pm$ bosons beyond N^2 LO at hadron colliders*, *Phys. Rev. D* **76** (2007) 114004 [[0708.1689](#)].
 - [79] G. Lustermaans, J. K. L. Michel and F. J. Tackmann, *Generalized Threshold Factorization with Full Collinear Dynamics*, [1908.00985](#).
 - [80] M. Bonvini and G. Marinelli, *On the approaches to threshold resummation of rapidity distributions for the Drell-Yan process*, *Eur. Phys. J. C* **83** (2023) 931 [[2306.03568](#)].
 - [81] T. Ahmed, N. Rana and V. Ravindran, *Higgs boson production through $b\bar{b}$ annihilation at threshold in N^3 LO QCD*, *JHEP* **10** (2014) 139 [[1408.0787](#)].
 - [82] G. Das, *$Z, W^\pm$ rapidity distributions at NNLL and beyond*, [2303.16578](#).
 - [83] A. Buckley, J. Ferrando, S. Lloyd, K. Nordström, B. Page, M. Rüfenacht et al., *LHAPDF6: parton density access in the LHC precision era*, *Eur. Phys. J. C* **75** (2015) 132 [[1412.7420](#)].
 - [84] A. Kulesza, G. F. Sterman and W. Vogelsang, *Joint resummation in electroweak boson production*, *Phys. Rev. D* **66** (2002) 014011 [[hep-ph/0202251](#)].
 - [85] S. Catani, M. L. Mangano, P. Nason and L. Trentadue, *The Resummation of soft gluons in hadronic collisions*, *Nucl. Phys. B* **478** (1996) 273 [[hep-ph/9604351](#)].
 - [86] R. Boughezal, C. Focke, X. Liu and F. Petriello, *W -boson production in association with a jet at next-to-next-to-leading order in perturbative QCD*, *Phys. Rev. Lett.* **115** (2015) 062002 [[1504.02131](#)].
 - [87] J. Gaunt, M. Stahlhofen, F. J. Tackmann and J. R. Walsh, *N -jettiness Subtractions for NNLO QCD Calculations*, *JHEP* **09** (2015) 058 [[1505.04794](#)].
 - [88] J. M. Campbell and R. K. Ellis, *An Update on vector boson pair production at hadron colliders*, *Phys. Rev. D* **60** (1999) 113006 [[hep-ph/9905386](#)].

- [89] J. M. Campbell, R. K. Ellis and W. T. Giele, *A Multi-Threaded Version of MCFM*, *Eur. Phys. J. C* **75** (2015) 246 [[1503.06182](#)].
- [90] R. Boughezal, J. M. Campbell, R. K. Ellis, C. Focke, W. Giele, X. Liu et al., *Color singlet production at NNLO in MCFM*, *Eur. Phys. J. C* **77** (2017) 7 [[1605.08011](#)].
- [91] D. J. Gross and F. Wilczek, *Ultraviolet Behavior of Nonabelian Gauge Theories*, *Phys. Rev. Lett.* **30** (1973) 1343.
- [92] H. D. Politzer, *Reliable Perturbative Results for Strong Interactions?*, *Phys. Rev. Lett.* **30** (1973) 1346.
- [93] W. E. Caswell, *Asymptotic Behavior of Nonabelian Gauge Theories to Two Loop Order*, *Phys. Rev. Lett.* **33** (1974) 244.
- [94] D. R. T. Jones, *Two Loop Diagrams in Yang-Mills Theory*, *Nucl. Phys. B* **75** (1974) 531.
- [95] E. Egorian and O. V. Tarasov, *Two Loop Renormalization of the QCD in an Arbitrary Gauge*, *Teor. Mat. Fiz.* **41** (1979) 26.
- [96] O. V. Tarasov, A. A. Vladimirov and A. Y. Zharkov, *The Gell-Mann-Low Function of QCD in the Three Loop Approximation*, *Phys. Lett. B* **93** (1980) 429.
- [97] S. A. Larin and J. A. M. Vermaseren, *The Three loop QCD Beta function and anomalous dimensions*, *Phys. Lett. B* **303** (1993) 334 [[hep-ph/9302208](#)].
- [98] T. van Ritbergen, J. A. M. Vermaseren and S. A. Larin, *The Four loop beta function in quantum chromodynamics*, *Phys. Lett. B* **400** (1997) 379 [[hep-ph/9701390](#)].
- [99] M. Czakon, *The Four-loop QCD beta-function and anomalous dimensions*, *Nucl. Phys. B* **710** (2005) 485 [[hep-ph/0411261](#)].
- [100] P. A. Baikov, K. G. Chetyrkin and J. H. Kühn, *Five-Loop Running of the QCD coupling constant*, *Phys. Rev. Lett.* **118** (2017) 082002 [[1606.08659](#)].
- [101] F. Herzog, B. Ruijl, T. Ueda, J. A. M. Vermaseren and A. Vogt, *The five-loop beta function of Yang-Mills theory with fermions*, *JHEP* **02** (2017) 090 [[1701.01404](#)].
- [102] T. Luthe, A. Maier, P. Marquard and Y. Schroder, *The five-loop Beta function for a general gauge group and anomalous dimensions beyond Feynman gauge*, *JHEP* **10** (2017) 166 [[1709.07718](#)].
- [103] T. Hahn, *CUBA: A Library for multidimensional numerical integration*, *Comput. Phys. Commun.* **168** (2005) 78 [[hep-ph/0404043](#)].
- [104] T. Hahn, *Concurrent Cuba*, *J. Phys. Conf. Ser.* **608** (2015) 012066 [[1408.6373](#)].
- [105] G. Das, M. C. Kumar and K. Samanta, *Resummed inclusive cross-section in ADD model at N^3LL* , *JHEP* **10** (2020) 161 [[1912.13039](#)].
- [106] G. Das, M. C. Kumar and K. Samanta, *Resummed inclusive cross-section in Randall-Sundrum model at NNLO+NNLL*, *JHEP* **07** (2020) 040 [[2004.03938](#)].
- [107] G. Das, M. C. Kumar and K. Samanta, *Precision QCD phenomenology of exotic spin-2 search at the LHC*, *JHEP* **04** (2021) 111 [[2011.15121](#)].
- [108] W. Bizoń, X. Chen, A. Gehrmann-De Ridder, T. Gehrmann, N. Glover, A. Huss et al., *Fiducial distributions in Higgs and Drell-Yan production at $N^3LL+NNLO$* , *JHEP* **12** (2018) 132 [[1805.05916](#)].
- [109] J. A. M. Vermaseren, *New features of FORM*, [math-ph/0010025](#).

- [110] B. Ruijl, T. Ueda and J. Vermaseren, *FORM version 4.2*, [1707.06453](#).
- [111] J. M. Henn, G. P. Korchemsky and B. Mistlberger, *The full four-loop cusp anomalous dimension in $\mathcal{N} = 4$ super Yang-Mills and QCD*, *JHEP* **04** (2020) 018 [[1911.10174](#)].
- [112] T. Huber, A. von Manteuffel, E. Panzer, R. M. Schabinger and G. Yang, *The four-loop cusp anomalous dimension from the $N = 4$ Sudakov form factor*, *Phys. Lett. B* **807** (2020) 135543 [[1912.13459](#)].
- [113] A. von Manteuffel, E. Panzer and R. M. Schabinger, *Cusp and collinear anomalous dimensions in four-loop QCD from form factors*, *Phys. Rev. Lett.* **124** (2020) 162001 [[2002.04617](#)].
- [114] O. V. Tarasov, *Anomalous dimensions of quark masses in the three-loop approximation*, *Phys. Part. Nucl. Lett.* **17** (2020) 109 [[1910.12231](#)].
- [115] S. A. Larin, *The Renormalization of the axial anomaly in dimensional regularization*, *Phys. Lett. B* **303** (1993) 113 [[hep-ph/9302240](#)].
- [116] E. N. Alekseev, V. A. Matveev, K. S. Nirov and V. A. Rubakov, eds., *Particles and cosmology. Proceedings, International School, Baksan Valley, Russia, April 22-27, 1993*, 1994.
- [117] K. G. Chetyrkin, *Quark mass anomalous dimension to $\mathcal{O}(\alpha_S^4)$* , *Phys. Lett. B* **404** (1997) 161 [[hep-ph/9703278](#)].