Unifying fractors, gravitors and photons from a gauge theoretical approach

Rodrigo F. Sobreiro*

Instituto de Física, Universidade Federal Fluminense,
Campus da Praia Vermelha, Av. Litorânea s/n, 24210-346, Niterói, RJ, Brazil

We revisit the first principles gauge theoretical construction of relativistic gapless fracton theory recently developed by A. Blasi and N. Maggiore. The difference is that, instead of considering a symmetric tensor field, we consider a vector field with a gauge group index, *i.e.* the usual Einstein-Cartan variable used in the first order formalism of gravity. After discussing the most general quadratic action for this field, we explore the physical sectors contained in the model. Particularly, we show that the model contains not only linear gravity and fractons, but also ordinary Maxwell equations, suggesting an apparent electrically charged phase of, for instance, spin liquids and glassy dynamical systems. Moreover, by a suitable change of field variables, we recover the Blasi-Maggiore gauge model of fractons and linear gravity.

I. INTRODUCTION

In recent years, collective excitations called fractons have been drawing attention in solid state physics, see for instance [1-3]. These type of quasiparticles are supposed to exist in, for example, spin liquids [2, 3] and glassy dynamics [2–4]. Fractons have also been connected to elasticity theory [2, 3, 5] and gravity [2, 3, 6]. In a gauge theoretical description [3, 7–10], fractons carry fractonic charge and produce (and interact through) U(1) tensor fields which generalize the usual Maxwell electromagnetism [11, 12]. In such fractonic electrodynamics, there are two types [7]: one with scalar fractonic charges where not only charges but also dipole moment and one specific component of the quadrupole moment are conserved; and another with vector charges where charges and its angular momentum analogue are conserved. For instance, for vector charge theory, if $E^{ij} = E^{ji}$ stands for the tensor electric field and ρ^i for vector fractonic charge density, Gauss law reads¹

$$\partial_j E^{ij} = \rho^i \,, \tag{1}$$

whose natural covariant generalization [8] yields

$$\partial_{\alpha} F^{\mu\nu\alpha} = \tau^{\mu\nu} \,, \tag{2}$$

with $F^{\mu\nu\alpha} = F^{\nu\mu\alpha}$ being the fracton field strength (See Section IV) and $\tau^{\mu\nu} = \tau^{\nu\mu}$ the generalized fractonic current density. Indeed, it was shown in [8] that equation (2) accommodates a complete covariant generalization of vector fractonic electrodynamics.

Just like ordinary electrodynamics, fractonic electrodynamics can be formulated in terms of potentials and gauge symmetry [3, 7–10]. Moreover, a gravitational parallel between gravity and fractons was established in [6]. In fact, as demonstrated by Blasi and Maggiore [9], the

construction of a theory of fractons based on the gauge principle, produces an action composed by the usual fracton action and the Fierz-Pauli action for spin-2 massless gravitons [13, 14]. Therefore, the covariant Blasi-Maggiore (BM) model generalizes pure fractonic vector gauge theory to include gravitational modes on it. See also [10]. Later, Bertolini and Maggiore found their way to formulate the BM model in terms of the fracton field strength appearing in equation (2).

The BM model is constructed with a symmetric tensor field $h_{\mu\nu}$, transforming under gauge transformations of the form

$$\delta_f h_{\mu\nu} = \partial_\mu \partial_\nu \zeta \,\,\,\,(3)$$

with ζ being a local parameter. It turns out that transformation (3) is just a special case of the gauge transformation in linear gravity theories (linear diffeomorphisms) [15], namely

$$\delta_d h_{\mu\nu} = \partial_\mu \zeta_\nu + \partial_\nu \zeta_\mu , \qquad (4)$$

for $\zeta_{\mu} = \partial_{\mu} \zeta/2$. Therefore, because the Fierz-Pauli is invariant under (4) it is also invariant under (3). The main conclusion of their gauge approach is that gravitonic modes are inherent to fracton models due to the fact that transformations (3) are a special case of the linear diffeomorphism transformations (4). But the contrary is not true because the fractonic action is not invariant under transformations (4).

In the present paper we construct a gauge fractonic model using Einstein-Cartan (EC) linear variables [16–22]. For simplicity, and for the sake of comparison with the BM model [9], we neglect the independent sector of the spin-connection in this first work. The action we gain is indeed equivalent to the one in the BM model, as we prove by a simple change of field variables. One useful feature in working with EC variables in linear gravity is that we can work with two distinct sets of indices, both transforming under Lorentz transformations. But one set of indices refers to spacetime itself while another set refers to the local tangent space. Thence, by fixing a regime to our model where frames in tangent space do not

^{*} rodrigo_sobreiro@id.uff.br

¹ Latin indices from i to q run through $\{1,2,3\}$. We also define Greek indices to run through $\{0,1,2,3\}$ while Latin indices from a to h run through $\{\underline{0},\underline{1},\underline{2},\underline{3}\}$.

move with respect to each other (all observers have synchronized clocks), the model reveals a sector which is indistinguishable from ordinary electromagnetism. In this sense, the BM gauge theory of fractons embraces not only fractons and graviton-like modes, but also photon-like excitations. And this feature is made evident in terms of the EC variables.

This work is organized as follows: In Section II we construct the gauge model of fractons in EC variables. In Section III we discuss the regime of the model where fracton electrodynamics behave exactly as ordinary Maxwell theory. In Section IV it is shown that the model considered in this work coincides with BM model of fractons. Finally, In Section V we display our final considerations.

II. GAUGE THEORY OF A SPECIAL KIND

To begin with the construction of our model, we set the scenario to be the Minkowski spacetime with metric $\eta \equiv \text{diag}(+1,-1,-1,-1)$. We define the Minkowskian vierbein η_{μ}^{a} (and its inverse η_{a}^{μ}) to be the set of maps connecting local inertial frames x^a to world coordinates x^{μ} , namely $x^{a} = \eta_{\mu}^{a} x^{\mu}$ and $x^{\mu} = \eta_{a}^{\mu} x^{a}$. Because the whole spacetime is flat, all frames are therefore taken to be inertial. Geometrodynamically speaking, this means that no gravitational effects are in place. For simplicity, η_{μ}^{a} is thus assumed to carry constant components and all coordinate systems are taken as Cartesian ones. In the EC formalism of gravity [16–22], the vierbein e^a_μ locally maps general frames dx^{μ} to inertial ones dx^{a} . The inertial frames are usually taken as Cartesian, but they could be curvilinear as well. Hence, a more general approach in flat space could consider curvilinear yet inertial frames in both sides of the vierbein map. However, this is an unnecessary complication we avoid.

The fundamental field of the model is h_{μ}^{a} , usually the graviton field in linear gravity theories. Inhere, this is precisely the general fracton potential field. Together with h_{μ}^{a} , the field h_{a}^{μ} is also important for covariance and the following relations are valid:

$$\eta_{\mu}^{a} \eta_{b}^{\mu} = \delta_{b}^{a} ,
\eta_{\mu}^{a} \eta_{a}^{\nu} = \delta_{\mu}^{\nu} ,
h_{a}^{\mu} = \eta_{b}^{\mu} \eta_{a}^{\nu} h_{\nu}^{b} .$$
(5)

Naturally, Latin and Greek indices transform under global Lorentz transformations². We also define linear diffeomorphisms, a genuine local Abelian gauge transformation, in the form

$$\delta_d h_\mu^a = \frac{1}{2} \eta_\nu^a \left(\partial_\mu \zeta^\nu + \partial^\nu \zeta_\mu \right) ,$$

$$\delta_d h_a^\mu = \frac{1}{2} \eta_a^\nu \left(\partial^\mu \zeta_\nu + \partial_\nu \zeta^\mu \right) .$$
 (6)

In the EC formalism, besides the graviton, the spin-connection is also considered as a fundamental field and independent from h^a_μ . Nevertheless, this field is not important for our current goals and we do not consider it here.

Let us perform two exercises. The first one is to find the most general quadratic theory describing the graviton. This means to find the most general action, respecting the global Lorentz symmetries and the gauge symmetry (6), polynomially local on the fields and their derivatives and quadratic on h_{μ}^{a} . We also consider that the action depends mostly quadratically on the derivatives. Moreover, h_{μ}^{a} is taken as a massless field from the beginning. The most general possible action with arbitrary coefficients a_{I} , already respecting global Lorentz symmetries, is just

$$S^{h} = \int d^{4}x \left(a_{1}h \Box h + a_{2}h_{\mu}^{a} \Box h_{a}^{\mu} + a_{3}h\eta_{a}^{\mu}\partial_{\mu}\partial^{\nu}h_{\nu}^{a} + a_{4}h_{a}^{\mu}\partial_{\mu}\partial^{\nu}h_{\nu}^{a} \right) , \tag{7}$$

with $h = \eta_a^{\mu} h_{\mu}^a$ being the trace of the gauge potential and $\Box = \partial^{\mu} \partial_{\mu}$ the d'Alambert operator. Imposing, $\delta_d S^h = 0$, a straightforward calculation gives the constraints $a_2 = -a_1$, $a_3 = -2a_1$, and $a_4 = 2a_1$ over the coefficients. Thus,

$$S_{lin}^{h} = \frac{a}{2} \int d^{4}x \left(-h \Box h + h_{\mu}^{a} \Box h_{a}^{\mu} - 2h_{a}^{\mu} \partial_{\mu} \partial^{\nu} h_{\nu}^{a} + 2h \eta_{a}^{\mu} \partial_{\mu} \partial^{\nu} h_{\nu}^{a} \right) , \qquad (8)$$

where we have redefined $a_1 = -a/2$ for convenience. This is the usual Fierz-Pauli massless action [13, 14] in the EC formalism, *i.e.* the non-interacting theory for spin-2 massless gravitons.

Following [8, 10], the fracton theory can be obtained from a more restricted class of the gauge transformations (6) by setting $\zeta^{\alpha} = \partial^{\alpha} \zeta$. Let us call it fracton gauge symmetry and define δ_f by

$$\delta_f h_\mu^a = \eta_\nu^a \partial^\nu \partial_\mu \zeta ,
\delta_f h_\mu^a = \eta_\alpha^\nu \partial_\nu \partial^\mu \zeta .$$
(9)

Thence, if we now impose fracton symmetry to action (7), $\delta_f S^h = 0$, one straightforwardly obtains the constraints $a_3 = -2a_1$ and $a_2 = a_1 - a_4$. Hence, S^h reduces to

$$S^{h} = \int d^{4}x \left[b_{1} \left(-h \Box h + 2h\eta_{a}^{\mu} \partial_{\mu} \partial^{\nu} h_{\nu}^{a} - h_{a}^{\mu} \partial_{\mu} \partial^{\nu} h_{\nu}^{a} \right) + b_{2} \left(-h_{\mu}^{a} \Box h_{a}^{\mu} + h_{a}^{\mu} \partial_{\mu} \partial^{\nu} h_{\nu}^{a} \right) \right],$$

$$(10)$$

where we have redefined $a_1 \to -b_1$ and $a_2 \to -b_2$. Performing two more redefinitions, $b_1 \to a/2$ and $b_2 \to (b-a)/2$, we finally get

$$S^h = S_{lin}^h + S_{frac}^h , (11)$$

with

$$S_{frac}^{h} = \frac{b}{2} \int d^4x \left(-h_{\mu}^{a} \Box h_{a}^{\mu} + h_{a}^{\mu} \partial_{\mu} \partial^{\nu} h_{\nu}^{a} \right) , \qquad (12)$$

² All results attained in the present work are also achieved by considering the Poincaré group as well.

being the action of fractons in terms of the EC variables h_{μ}^{a} . Action (11) is just the EC version of the BM model [8–10] (See Sect. IV). From the gauge construction of the model, the parameters a and b are arbitrary constants (with dimension of mass squared). Nevertheless, a can be related with Newton's constant if action (11) refers to some kind of gravity model, by means of $a = 1/16\pi G$. In fact, as discussed in [10], pure linear gravity theory is achieved by setting b = 0 while pure fractonic action is obtained for a = 0. For any other non trivial values of these parameters, a mixed model of fractons and gravitons is at our disposal. Remarkably, in the case of a = 0, there is a regime where the field equations split into ordinary Maxwell equations for scalar fracton charges and SO(3)algebra-valued fractonic electrodynamics (See Sect. III).

It is useful to rewrite action (11) in terms of the gauge invariant field $H^a_{\mu\nu}$,

$$S^{h} = \frac{a}{2} \int d^{4}x H_{\mu} H^{\mu} + \frac{(b-a)}{4} \int d^{4}x H^{a}_{\mu\nu} H^{\mu\nu}_{a} , \quad (13)$$

with

$$H^{a}_{\mu\nu} = \partial_{\mu}h^{a}_{\nu} - \partial_{\nu}h^{a}_{\mu} ,$$

$$H_{\mu} = \eta^{\nu}_{a}H^{a}_{\mu\nu} .$$
 (14)

And, to complete the model, at least classically, we couple action (13) with an external source field τ_a^{μ} , so the total action of interest reads

$$S = S^h - \int d^4x \; h^a_\mu \tau^\mu_a \; . \tag{15}$$

The source τ_a^{μ} is regarded as a covariant current of fractonic charges. The corresponding field equations generated from the action (15) are easily computed,

$$\partial_{\nu} \left[(b - a) H_a^{\mu \nu} + a \left(\eta_a^{\mu} H^{\nu} - \eta_a^{\nu} H^{\mu} \right) \right] = \tau_a^{\mu} . \tag{16}$$

Taking the divergence of the field equations (16), one directly obtains the fracton charge conservation equation

$$\partial_{\mu}\tau_{a}^{\mu} = 0. \tag{17}$$

Equation (16) can be simplified a bit by taking its trace, yielding

$$\partial_{\mu}H^{\mu} = \frac{1}{(b+2a)}\tau. \tag{18}$$

Thus, (16) reduces to

$$\partial_{\nu} \left[(b-a) H_a^{\mu\nu} - a \eta_a^{\nu} H^{\mu} \right] = \theta_a^{\mu} , \qquad (19)$$

where

$$\theta_a^{\mu} = \tau_a^{\mu} - \frac{a}{(b+2a)} \eta_a^{\mu} \tau ,$$
 (20)

is an effective current whose trace is given by

$$\theta = \frac{(b-2a)}{(b+2a)}\tau \ . \tag{21}$$

Clearly, if b = 2a, the effective current is traceless. Although equation (19) is simpler than (16), the effective current is no longer a conserved quantity.

Besides the field equations (16), the Bianchi identities are also valid

$$\epsilon^{\mu\nu\alpha\beta}\partial_{\nu}H^{a}_{\alpha\beta} = 0. \tag{22}$$

The Bianchi identities complete the set of field equations for h_{μ}^{a} . Another way to look, inspired in electrodynamics, is to solve equations (16) and (22) for the gauge invariant field $H^a_{\mu\nu}$, the analogous field to the electromagnetic field $F_{\mu\nu} = \partial_{\mu}A_{\nu} - \partial_{\nu}A_{\mu}.$

To end this section, let us define

$$\mathcal{F}_{\mu\nu}^{a} = (b - a)H_{\mu\nu}^{a} + a(\eta_{\mu}^{a}H_{\nu} - \eta_{\nu}^{a}H_{\mu}), \qquad (23)$$

which is the tensor field appearing in equation (16). Thus.

$$\partial^{\nu} \mathcal{F}^{a}_{\mu\nu} = \tau^{a}_{\mu} \,, \tag{24}$$

is clearly the generalization of (1). First, because it is a covariant generalization. Second, due to the a and bparameters characterizing fracton and linear gravity sectors. Third, because the field and source carry a gauge index. In fact, the gauge index will reveal the photonic sector of the model in certain regime. Nevertheless, there is a difference: the field $H^a_{\mu\nu}$ is antisymmetric in spacetime indices while E_{ij} is symmetric. Indeed, our construction naturally led to an anti-symmetric representation of the gauge invariant fractoric fields. In Section IV, we will prove that this representation is equivalent to the symmetric representation of the BM model.

MAXWELL ELECTRODYNAMICS AND SO(3) ALGEBRA-VALUED **ELECTRODYNAMICS**

In this section, we turnoff the gravitational channel by setting the parameter a to zero and assume a pure covariant fracton electrodynamical theory with gauge index. The field equations (16) reduce to

$$\partial_{\nu} H_a^{\mu\nu} = \tau_a^{\mu} \ . \tag{25}$$

where the constant b has been absorbed in h_{μ}^{a} and τ_{a}^{μ} by

means of the rescaling $h^a_\mu \to b^{-1/2} h^a_\mu$ and $\tau^\mu_a \to b^{1/2} \tau^\mu_a$. Recalling that $H^a_{\mu\nu}$ is gauge invariant and thus observable, and that the Bianchi identity (22) are always valid, we have a set of equations very similar to Maxwell equations. Except, of course, for the gauge index. The gauge index, referring to the tangent space, is an internal index analogous to color indices in quantum chromodynamics [23], a genuine gauge charge index. However, inhere, by construction, the gauge index refers to a gauge symmetry identified with local spacetime isometries characterized by the Lorentz group of transformations connecting all local inertial frames. Therefore, the set of equations (25)

and (22) describe an SO(1,3) algebra-valued theory with gauge symmetry (9): The fields and charges are SO(1,3) algebra-valued objects and they transform under gauge transformations (9). These gauge transformations can be cast in the form

$$\delta_f h_\mu^a = \partial_\mu \zeta^a ,$$

$$\delta_f h_\mu^a = \partial^\mu \zeta_a ,$$
(26)

with $\zeta^a = \eta_{\nu}^a \partial^{\nu} \zeta$. Therefore, we have a $U(1)^4 = U(1) \times U(1) \times U(1) \times U(1)$ gauge theory for SO(1,3) algebra-valued fields.

At this point, we explore the fact that the gauge indices are tangent space indices: We impose a restrictive condition to the frames x^a : We fix the time axis and leave an SO(3) global gauge freedom for tangent indices. Physically, we are imposing that all frames x^a have synchronized clocks and they relate to each other only by spacial rotations, *i.e.* they do not move with respect to each other. The possibility of relativistic effects are contained in the frames x^{μ} and the action of the complete Lorentz group on they. Therefore, in this regime, the split between space and time in tangent indices is natural.

We consider first the index a projected in time direction, $a = \underline{0}$. Since we have fixed this action, these components do not transform under the residual SO(3) possible rotations. Therefore, in this regime we are allowed to omit all indices $a = \underline{0}$. Remarkably, in this case, we recover the equations of ordinary electrodynamics. In fact, identifying $h_{\mu}^{0} = A_{\mu}$ with the electrodynamical potential, we have $H_{\mu\nu}^{0} = \partial_{\mu}A_{\nu} - \partial_{\nu}A_{\mu} = F_{\mu\nu}$ being identified with the electromagnetic field. Moreover, we can identify $\tau_{\mu}^{0} = j_{\mu}$ with electric charge four-current $(j_{0} = \rho)$ is the scalar fractonic charge density and j_{i} is the fractonic vector current density.). Therefore, it is trivial to show that (25) and (22) are exactly reduced to Maxwell equations. Furthermore, electrodynamics is a realization of a U(1) gauge theory. Indeed, gauge transformations (9) (or (26)) reduce to U(1) gauge transformations,

$$\delta_e A_\mu = \partial_\mu \xi ,$$

$$\delta_e A^\mu = \partial^\mu \xi ,$$
 (27)

with $\xi = \eta_{\nu}^{0} \partial^{\nu} \zeta$ and $\delta_{f} \to \delta_{e}$. Hence, in this regime, this sector of fractonic charges manifest exactly like electric charges (including its conservation law, $\partial_{\mu} j^{\mu} = 0$ coming from (17)) and the fractonic fields in this sector, as well as the charges, have no apparent difference from ordinary electromagnetic fields.

For $a=\underline{i}$, we define $H^{0i}_{\underline{i}}=E^i_{\underline{i}},\ H^{ij}_{\underline{i}}=\epsilon^{ij}_{\ k}B^k_{\underline{i}},\ \tau^0_{\underline{i}}=\rho_{\underline{i}}$ and $\tau^i_{\underline{i}}=j^i_{\underline{i}}$, and equations (25) become

$$\vec{\nabla} \cdot \vec{E}_{\underline{i}} = \rho_{\underline{i}}.$$

$$\vec{\nabla} \times \vec{B}_{\underline{i}} - \frac{\partial \vec{E}_{\underline{i}}}{\partial t} = \vec{j}_{\underline{i}}.$$
(28)

Thus, we attained an SO(3) algebra-valued electrodynamics. Electric and magnetic fields are vector fields

but they carry a gauge index associated with the global SO(3) rotations in tangent space. The gauge transformations (9), on the other hand, become

$$\delta_p h_{\mu}^{\underline{i}} = \partial_{\mu} \xi^{\underline{i}},
\delta_p h_{\underline{i}}^{\mu} = \partial^{\mu} \xi_{\underline{i}},$$
(29)

with $\xi^{\underline{i}} = \eta^{\underline{i}}_{\nu} \partial^{\nu} \zeta$ and $\delta_f \to \delta_p$. Transformations (29) describe a $U(1)^3 = U(1) \times U(1) \times U(1)$ local gauge symmetry, classifying this sector of the model as a kind of triple electrodynamics whose conserved charges are SO(3) algebra-valued objects.

Let us go back to the full covariant equation (25) and the Bianchi identity (22). We have a generalized covariant fracton model where the four current τ_a^{μ} describes a generalization of fractonic vector charges. And, for a specific relativistic regime, the model splits in a sector indistinguishable from ordinary electrodynamics and a triple electrodynamics whose charges and fields are SO(3) algebra-valued objects. We conclude therefore that the full model (15) carries, not only fractonic and gravitonic modes, but also photonic modes. These new modes are indeed present in the general theory (15), but, in this case, they can be distinguished from ordinary electromagnetic physics by performing a Lorentz transformation in tangent space.

IV. FRACTON ELECTRODYNAMICS IN FULLY SPACETIME REPRESENTATION

We have constructed the model (15) based on the same gauge principles of the BM model [8–10], except for the basic field which is not a symmetric tensor but a vector field with internal charge index. We have attained a model of gauge invariant antisymmetric fractonic fields. Therefore, we still have to show how our model relates to the BM model, which is constructed in terms of symmetric tensor fields.

To connect our model with the BM model, we must perform a change of variables from h^a_μ to a fully space-time representation $h_{\mu\nu}$. The connection between EC variables h^a_μ and second order variables $h_{\mu\nu}$ is given by

$$h_{\mu\nu} = \frac{1}{2} \left(\eta_{a\mu} h_{\nu}^{a} + \eta_{a\nu} h_{\mu}^{a} \right) . \tag{30}$$

It is then straightforward to show that

$$F_{\mu\nu\alpha} = \eta_{a\mu} H^a_{\nu\alpha} + \eta_{a\nu} H^a_{\mu\alpha} = \partial_{\mu} h_{\alpha\nu} + \partial_{\nu} h_{\mu\alpha} - 2\partial_{\alpha} h_{\mu\nu} , \quad (31)$$

which is the fracton gauge invariant field strength defined in [8]. In fact, applying ∂^{α} in (31) and using the field equations (25) one readily finds equations (2), where the fracton current in spacetime representation reads

$$\tau_{\mu\nu} = \eta_{a\mu}\tau_{\nu}^{a} + \eta_{a\nu}\tau_{\mu}^{a} . \tag{32}$$

The field equation (2) is the pure fracton equation (with sources) of the BM model. The full equation in the BM

model actually reads

$$(b-a)\partial^{\alpha}F_{\mu\nu\alpha} + + a \left[\eta_{\mu\nu}\partial_{\alpha}F_{\beta}^{\beta\alpha} - \frac{1}{2} \left(\partial_{\mu}F_{\nu} + \partial_{\nu}F_{\mu} \right) \right] = \tau_{\mu\nu} , \quad (33)$$

with $F_{\mu} = F^{\nu}_{\nu\mu}$. It is a straightforward exercise to show that the substitution of (31) and (32) in the field equations (33) yields equations (16). An easier task is to show that action (15), in terms of the field strength (31), reads

$$S^{h}(F) = \frac{(b-a)}{12} \int d^{4}x F_{\mu\nu\alpha} F^{\mu\nu\alpha} + \frac{a}{8} \int d^{4}x F^{\nu}_{\nu\mu} F^{\alpha}_{\alpha\mu} ,$$
(34)

which is, up to normalization factors, the BM action found in [8]. This proves that the model described by action (15) is totally equivalent to the BM model [8].

V. CONCLUSIONS

In this paper we constructed a covariant model of fractons based on purely gauge principles and using EC variables in complete analogy to the BM gauge model of fractons [8–10]. The general action, quadratic on the gauge field, is composed by by pure fractoric sector and another sector identified as the Fierz-Pauli action for massless gravitons. We proved, by changing the EC variables to second order variables, that our model is totally equivalent to the BM model, as expected. In terms of gauge invariant fields fields, we have constructed a fractonic model in a representation of antisymmetric fields, in contrast to the BM model of symmetric fields. Nevertheless, we proved the equivalence between both models. In a nutshell, we constructed a $U(1)^4$ gauge theory of fractons with SO(1,4) algebra-valued antisymmetric fields and charges which is equivalent to the BM symmetric fractoric gauge theory of fractors.

Additionally, we showed that there is a relativistic regime where Maxwell electrodynamics emerges. In this regime, observers with synchronized clocks not moving with respect to each other could not distinguish this sector of fracton electrodynamics from ordinary Maxwell electromagnetism. This effect also works for non-relativistic observers, *i.e.* a relativistic regime where absolute time can be defined for local observers (*e.g.* Galilean frames [24]). Moreover, the SO(3) algebravalued vector charges and fields are also present in this regime, a kind of triple $U(1)^3$ electrodynamics.

It turns out that the relativistic regime favoring fractons to mimic electrodynamics is just the typical situation occurring in most solid state laboratories. Therefore, such effect could be probed somehow by observing, for instance, a kind of phase in a spin liquid looking like an electrically charged spin liquid. Such system would produce fake electromagnetic fields and interact with them. However, the system would not respond to ordinary electromagnetic fields normally because the charges are actually fracton charges and not electrical ones.

The results obtained can be improved by exploring further the model in EC representation. For instance, one can couple scalar fields instead of an external current, providing dynamics do the fracton charges. Another possibility is to couple fermions, since the EC formalism is a suitable scenario to couple gravity with spinors [20, 25].

ACKNOWLEDGEMENTS

This study was financed in part by The Coordenação de Aperfeiçoamento de Pessoal de Nível Superior - Brasil (CAPES) - Finance Code 001.

S. Vijay, J. Haah, and L. Fu, "Fracton Topological Order, Generalized Lattice Gauge Theory and Duality". Phys. Rev. B 94 no. 23, (2016) 235157.

^[2] R. M. Nandkishore and M. Hermele, "Fractons". Ann. Rev. Condensed Matter Phys. 10 (2019) 295–313.

^[3] M. Pretko, X. Chen, and Y. You, "Fracton Phases of Matter".

Int. J. Mod. Phys. A 35 no. 06, (2020) 2030003.

^[4] A. Prem, J. Haah, and R. Nandkishore, "Glassy quantum dynamics in translation invariant fracton models". *Phys. Rev. B* 95 no. 15, (2017) 155133.

^[5] M. Pretko and L. Radzihovsky, "Fracton-Elasticity Duality". Phys. Rev. Lett. 120 no. 19, (2018) 195301.

^[6] M. Pretko, "Emergent gravity of fractons: Mach's principle revisited". Phys. Rev. D 96 no. 2, (2017) 024051.

^[7] M. Pretko, "Generalized Electromagnetism of Subdimensional Particles: A Spin Liquid Story".

Phys. Rev. B 96 no. 3, (2017) 035119.

^[8] E. Bertolini and N. Maggiore, "Maxwell theory of fractons". Phys. Rev. D 106 no. 12, (2022) 125008.

^[9] A. Blasi and N. Maggiore, "The theory of symmetric tensor field: From fractons to gravitons and back". *Phys. Lett. B* 833 (2022) 137304.

^[10] E. Bertolini, A. Blasi, A. Damonte, and N. Maggiore, "Gauging fractons and linearized gravity". Symmetry 15 (2023) 945.

^[11] J. C. Maxwell, "A dynamical theory of the electromagnetic field". Phil. Trans. Roy. Soc. Lond. 155 (1865) 459–512.

^[12] J. D. Jackson, Classical Electrodynamics. Wiley, 1998.

^[13] M. Fierz and W. Pauli, "On relativistic wave equations for particles of arbitrary spin in an electromagnetic field". Proc. Roy. Soc. Lond. A 173 (1939) 211–232.

^[14] G. Gambuti and N. Maggiore, "Fierz–Pauli theory reloaded: from a theory of a symmetric tensor field to

- linearized massive gravity". Eur. Phys. J. C 81 no. 2, (2021) 171.
- [15] R. M. Wald, General Relativity. Chicago Univ. Pr., Chicago, USA, 1984.
- [16] R. Utiyama, "Invariant theoretical interpretation of interaction". Phys. Rev. 101 (1956) 1597–1607.
- [17] T. W. B. Kibble, "Lorentz invariance and the gravitational field". J. Math. Phys. 2 (1961) 212–221.
- [18] D. W. Sciama, "The Physical structure of general relativity". Rev. Mod. Phys. 36 (1964) 463–469.
 [Erratum: Rev.Mod.Phys. 36, 1103–1103 (1964)].
- [19] C. W. Misner, K. S. Thorne, and J. A. Wheeler, Gravitation. W. H. Freeman, San Francisco, 1973.
- [20] V. De Sabbata and M. Gasperini, INTRODUCTION TO GRAVITY. Singapore, Singapore: World Scientific 346p, 1985.

- [21] A. Mardones and J. Zanelli, "Lovelock-Cartan theory of gravity". Class. Quant. Grav. 8 (1991) 1545–1558.
- [22] J. Zanelli, "Lecture notes on Chern-Simons (super-)gravities. Second edition (February 2008)". in Proceedings, 7th Mexican Workshop on Particles and Fields (MWPF 1999): Merida, Mexico, November 10-17, 1999. 2005.
- [23] C. Itzykson and J. B. Zuber, Quantum Field Theory. International Series In Pure and Applied Physics. McGraw-Hill, New York, 1980. http://dx.doi.org/10.1063/1.2916419.
- [24] A. Guerrieri and R. F. Sobreiro, "Non-relativistic limit of gravity theories in the first order formalism". *JHEP* 03 (2021) 104.
- [25] F. W. Hehl, J. D. McCrea, E. W. Mielke, and Y. Ne'eman, "Metric affine gauge theory of gravity: Field equations, Noether identities, world spinors, and breaking of dilation invariance". *Phys. Rept.* 258 (1995) 1–171.