# **On** $\mathbb{P}^1$ -stabilization in unstable motivic homotopy theory

Aravind Asok\*

Tom Bachmann

Michael J. Hopkins

### Abstract

We analyze stabilization with respect to  $\mathbb{P}^1$  in the Morel–Voevodsky unstable motivic homotopy theory. We introduce a refined notion of cellularity in various motivic homotopy categories taking into account both the simplicial and Tate circles. Under suitable cellularity hypotheses, we refine the Whitehead theorem by showing that a map of nilpotent motivic spaces can be seen to be an equivalence if it so after taking (Voevodsky) motives. We then establish a version of the Freudenthal suspension theorem for  $\mathbb{P}^1$ -suspension, again under suitable cellularity hypotheses. As applications, we resolve Murthy's conjecture on splitting of corank 1 vector bundles on smooth affine algebras over algebraically closed fields having characteristic 0 and compute new unstable motivic homotopy of motivic spheres.

# Contents

1	Introduction	2
2	Preliminaries on unstable and stable motivic homotopy theory	5
	2.1 Motivic localization and the unstable connectivity theorem	 5
	2.2 Stable motivic homotopy theory and effectivity	
	2.3 Real-étale motivic homotopy theory	
3	Weak cellularity and nullity	20
	3.1 Weak cellularity and nullifications: basic properties	 21
	3.2 Stable weak $S^{p,q}$ -cellularity via <i>t</i> -structures	
	3.3 Examples	
4	A weakly-cellular Whitehead tower and consequences	35
	4.1 A weakly-cellular Whitehead tower	 35
	4.2 Unstable cellularity of loop spaces and fibers	
	4.3 Weak cellularity: comparing unstable and stable	
	4.4 Further examples	
5	Equivariant geometry of symmetric powers	42
	5.1 Motivic homotopy theory and group scheme actions	 43
	5.2 The motivic Dold-Thom theorem	
	5.3 Cellularity of the cofiber of the assembly map	 46
	5.4 Weak cellularity and quotients	
	5.5 Modifications in positive characteristic	
6	Weak cellular estimates of the fiber of the unit map	52
	6.1 The unit of stabilization and fiber sequences	 52
	6.2 The main result	
	6.3 Complements	

\*Aravind Asok was partially supported by National Science Foundation Awards DMS-1802060 and DMS-2101898

7	Applications	60
	7.1 Murthy's conjecture and complements	. 60
	7.2 Computations	. 62

# **1** Introduction

If n is an integer  $\geq 2$ , and X is an (n-1)-connected CW complex, then the classical Freudenthal suspension theorem asserts that the unit map  $X \to \Omega \Sigma X$  is 2n - 1-connected, i.e., induces an isomorphism on homotopy groups in degrees  $\leq 2n - 2$  and an epimorphism in degree 2n - 1. The suspension theorem gives rise to the notion of "stable" phenomena in topology, which according to Adams occur in "sufficiently large dimensions", in contrast to "unstable" phenomena, which occur in some definite dimensions. Typically, stable phenomena are more amenable to calculation than unstable phenomena, while unstable phenomena are more closely related to geometry; this distinction perhaps led Eilenberg to quip "we can distinguish two cases—the stable case and the interesting case" [Ada95, p. 125]. Granted control of the stabilization process, one may often "unwind" an unstable problem into a sequence of stable problems whose solutions are more computationally accessible.

In algebraic geometry, the influence of homotopy theory has followed a different course. Stable structures, e.g., cohomology theories of various sorts like algebraic K-theory have been studied for a long time. However, stable and unstable homotopy categories housing these structures have only comparatively recently aided in analysis of such theories. For concreteness, fix a field k and write  $Sm_k$  for the category of smooth k-varieties and Spc(k) for the Morel–Voevodsky unstable motivic homotopy category [MV99]. The stable motivic homotopy category, denoted SH(k), is built by stabilizing with respect to two circles: a "simplicial" circle  $S^1 =: S^{1,0}$  and a "Tate" circle  $\mathbb{G}_m = S^{1,1}$ ; equivalently, one may stabilize with respect to  $\mathbb{P}^1 \simeq S^{2,1} = S^1 \wedge \mathbb{G}_m$ . Stabilizing with respect to  $\mathbb{P}^1$  is a "natural" choice because known cohomology theories, e.g., algebraic K-theory, are representable in the resulting setup [Voe98]. The category SH(k)has been analyzed intensely since its construction, following analogies both structural and computational with classical stable homotopy theory.

To date, the link between unstable and stable motivic homotopy theory has been harder to quantify. Morel, in his foundational work [Mor12] analyzed stabilization with respect to  $S^1$ . His results imply that stabilization with respect to the simplicial circle behaves as one expects from the classical Freudenthal suspension theorem: if  $\mathscr{X}$  is an (n-1)-connected motivic space, then the map  $\mathscr{X} \to \Omega \Sigma \mathscr{X}$  is itself (2n-1)-connected in the sense above. However, stabilization with respect to  $S^1$  is not "geometrically" well-behaved. More importantly, it seems not as well connected with the "stable" phenomena suggested above and is therefore not so helpful from a computational standpoint.

All of the above points were understood shortly after Morel and Voevodsky introduced the motivic homotopy category. Morel's early computations (eventually published in [Mor12]) imply that a stabilization theorem for  $\mathbb{P}^1$ -suspension of form similar to  $S^1$ -suspension cannot hold without additional hypotheses. Moreover, Voevodsky wrote: "[a]n analog of the Fre[u]denthal suspension theorem must exist, but we have no idea what it is" ([Voe00a, p. 12]). That additional hypotheses are necessary is perhaps not surprising given the form of the suspension theorem in equivariant homotopy theory [tD87, Theorem II.2.10].

Thus, to date, the situation can be described as follows: unstable motivic homotopy theory does control geometric phenomena (e.g., vector bundles on smooth affine varieties), and stable motivic homotopy theory is calculable, but only in very special cases has one been able to achieve sufficient control over the stabilization process to link unstable and stable phenomena. Our goal in this paper is to bridge this gap by establishing an version of the Freudenthal suspension theorem for  $\mathbb{P}^1$ -suspension in unstable motivic homotopy theory.

In order to state the result, we introduce a notion of cellularity, extending that described by Dror

[DF96]. Given a pointed (compact) motivic space A, we construct a class of motivic spaces we call "weakly A-cellular". It contains  $A \wedge X_+$  for any smooth scheme X, is stable under (weak) equivalences, (homotopy) colimits and "(homotopy) cofiber extensions"; we write O(A) for the class of weakly A-cellular spaces. We will be particularly interested in the class of weakly  $S^{p,q}$ -cellular motivic spaces,  $p \ge q$ .

The class  $O(S^{n+1,0})$  is the class of *n*-connected motivic spaces, while  $O(S^{q,q})$  is an unstable analog of the notion of *q*-effective spectra studied by various authors. For applications to be described momentarily, it will suffice to observe that the punctured affine space  $\mathbb{A}^n \setminus 0 \in O(S^{4,2})$  if  $n \ge 3$ . With that terminology in hand, we may state our  $\mathbb{P}^1$ -stabilization theorem.

**Theorem 1** (See Theorem 6.3.1). Assume k is a field that has characteristic 0. If  $\mathscr{X} \in \text{Spc}(k)$  is a pointed motivic space and lies in  $O(S^{p,q})$  with  $p \ge 4, q \ge 2$ , then the fiber of the stabilization map

$$\mathscr{X} \longrightarrow \Omega^{2,1} \Sigma^{2,1} \mathscr{X}$$

lies in  $O(S^{a,2q})$  where a = min(2p - 1, p + 2q - 1).

*Remark* 2. In the body of the text, we will also establish a version of Theorem 1 when k is a field having characteristic exponent p > 0. Roughly speaking, the above statement holds after inverting p; see Theorem 6.3.1 for more details.

#### Applications

Theorem 1 has a host of consequences, some of which will be developed in forthcoming work. We content ourselves with listing here one concrete application to the theory of algebraic vector bundles and some computational consequences. Building on a long line of classical questions in the theory of projective modules over commutative rings, it is natural to search for "explicit" obstructions to splitting positive corank bundles on (smooth) affine varieties over fields (see [Mur99] or the introduction to [AF15] for some discussion of the history of this problem). For a precise problem in this direction, M.P. Murthy suggested that over an algebraically closed field k, the only obstruction to splitting off a trivial rank 1 summand from a rank d bundle E on a smooth affine variety of dimension d + 1,  $d \ge 1$ , was the vanishing of  $c_d(E) \in CH^d(X)$  [Mur99, p. 173]. That the preceding statement was true came to be known as Murthy's splitting conjecture [AF15, Conjecture 1].

**Theorem 3** (See Theorem 7.1.1). Assume k is an algebraically closed field having characteristic 0. If X is a smooth affine k-scheme of dimension d + 1,  $d \ge 1$ , and E is a rank d vector bundle on X, then E splits off a trivial rank 1 summand if and only if  $0 = c_d(E) \in CH^d(X)$ .

The theorem is immediate from definitions in the case d = 1 and as such is only interesting when  $d \ge 2$ . In that case, the connection between Theorem 1 and Theorem 3 was laid out in the introduction of [AF15]. In brief, the question of splitting a trivial rank 1 summand off a vector bundle can be phrased as an  $\mathbb{A}^1$ -homotopical lifting problem by appeal to the representability theorem for vector bundles [AHW17, Theorem 1]. In turn, the relevant lifting problem can be analyzed by techniques of obstruction theory.

In the case of Murthy's conjecture, the potentially non-vanishing obstructions rely on knowledge of the sheaves  $\pi_{d-1}(\mathbb{A}^d \setminus 0)$  and  $\pi_d(\mathbb{A}^d \setminus 0)$ ,  $d \ge 2$ . The sheaves  $\pi_{d-1}(\mathbb{A}^d \setminus 0)$  were computed by Morel and stabilize for  $d \ge 2$ . In [AF14b] and [AF15], the first author and J. Fasel computed the sheaves  $\pi_d(\mathbb{A}^d \setminus 0)$  for d = 2, 3, at least over fields having characteristic not equal to 2.

Building on some ideas of Morel, the first author and J. Fasel formulated a general conjecture for the form of  $\pi_d(\mathbb{A}^d \setminus 0)$  for  $d \ge 4$  in [AF13, Conjecture 7] from which Murthy's conjecture would follow. In fact, to establish Murthy's conjecture one only needs the computation of  $\pi_3(\mathbb{A}^3 \setminus 0)$  mentioned above and a

surjectivity statement that follows from Theorem 1, i.e., less than the complete computation of  $\pi_d(\mathbb{A}^d \setminus 0)$ . This is the path we follow to deduce Murthy's conjecture. On the other hand, Theorem 1 gives much more. Indeed, the entire structure of  $\pi_d(\mathbb{A}^d \setminus 0)$  for  $d \ge 4$  may be deduced from a corresponding  $\mathbb{P}^1$ -stable computation which appears in [RSØ19], at least over fields having characteristic 0 (see Theorem 7.2.1).

#### **Overview of the proof**

Freudenthal's original proof of the suspension theorem in the case of spheres was of a rather geometric nature [Fre38]. Blakers and Massey employed a more formal and conceptually clearer technique via homotopy excision [BM51], which is essentially the standard proof now. Our proof, by necessity, does not follow any of the classical proofs so we give here a brief outline for the argument.

First, there is the key geometric ingredient, which yields a cellular estimate regarding the cofiber of the assembly map in the motivic Eilenberg–MacLane spectrum  $H\mathbb{Z} \wedge \Sigma_{\mathbb{P}^1}^{\infty} X_+$  for X a smooth scheme. The key geometric input arises from Voevodsky's motivic Dold–Thom theorem which allows one to realize the above motivic Eilenberg–Mac Lane spaces as a colimit of symmetric powers. In that case, the the cofiber of the assembly map can be described very geometrically as a Thom complex of a suitable equivariant vector bundle and the cellular estimate can be obtained by stratifying with respect to stabilizer type; this kind of analysis goes back to the work of Nakaoka [Nak57] and was codified in [AD01]. In a motivic context, this kind of analysis also appears in Voevodsky's analysis of the zero slice of the sphere spectrum in characteristic 0 [Voe04]. In the context of this paper, the required result in characteristic 0 is Theorem 5.4.2 and a suitable version in positive characteristic can be found in Theorem 5.5.5.

Second, we reduce from  $H\mathbb{Z}$ -modules to  $\mathbb{P}^1$ -spectra. This uses a number of ingredients, but perhaps key is an adaptation of the main idea in Levine's argument for the convergence of Voeovodsky's slice tower [Lev13] as detailed in [BEØ21]. Levine's argument treats the case of fields of finite cohomological dimension, but in conjunction with some information about real étale homotopy theory, one can treat the case of arbitrary characteristic 0-fields. This reduction is handled in the proof of Theorem 6.2.1, but see Lemmas 6.2.7 and 6.2.6.

Third, we reduce from  $\mathbb{P}^1$ -spectra to spaces. This reduction uses a refinement of the classical Whitehead tower of a space taking cellularity into account. Building this refined tower and analyzing its properties is the content of Section 4, but see Amplification 4.1.13 for a precise statement. The input to this statement are results of Bachmann–Yakerson [BY20] and Bachmann [Bac21a] on conservativity of stabilization with respect to  $\Omega^{1,1}$  which we review here as Theorem 2.2.29.

All of these ideas are built on the notion of weak cellular classes that we develop in Section 3. This section contains the basic definitions and develops a host of natural properties, including relationship between cellularity and fiber and cofiber sequences etc.

#### Notation and conventions

Much of the notation to be used in this paper will be introduced in Sections 2 and 3; we try to present an over view here. Throughout, assume k is a field; further hypotheses will be imposed on k as necessary in the body of the paper: frequently k will be assumed perfect, and sometimes furthermore infinite. Write  $Sm_k$  for the category of smooth k-schemes and  $Sch_k$  for the category of finite-type separated k-schemes; we will also write  $Sch_k^{qp}$  for the category of quasi-projective k-schemes.

If  $S_k$  is one of the categories of schemes described above, then we write  $P(S_k)$  for the  $\infty$ -topos of presheaves of spaces on  $Sm_k$ . If  $\tau$  is a Grothendieck topology on  $S_k$ , then we write  $Shv_{\tau}(S_k) \subset P(S_k)$  for full subcategory of Nisnevich sheaves of spaces on  $S_k$ , which is once again an  $\infty$ -topos. There are corresponding pointed versions of all these constructions: we write  $P(S_k)_*$  for the  $\infty$ -topos of presheaves of pointed spaces and  $Shv_{\tau}(S_k)$  for the subcategory of  $\tau$ -sheaves of pointed spaces.

We write  $S^i$  for the simplicial *i*-circle, and we set  $S^{p,q} := S^{p-q} \wedge \mathbb{G}_m^{\wedge q}$ . We write  $\Sigma^i$  for the operation of smashing with  $S^i$ , and  $\Sigma^{p,q}$  for the operation of smashing with  $S^{p,q}$ . We write <u>Map</u> for the internal mapping space, i.e., given  $\mathscr{X}, \mathscr{Y} \in \operatorname{Spc}(k)$ , <u>Map</u>( $\mathscr{X}, \mathscr{Y}$ ) represents  $U \mapsto \operatorname{Hom}_{\operatorname{Shv}_{Nis}(\operatorname{Sm}_k)}(U \times \mathscr{X}, \mathscr{Y})$ . Likewise, if  $\mathscr{X}$  and  $\mathscr{Y}$  are pointed spaces, then we write <u>Map</u><sub>\*</sub>( $\mathscr{X}, \mathscr{Y}$ ) for the pointed internal mapping space. We write  $\Omega^i$  for the *i*-fold loops functor adjoint to  $\Sigma^i$  and, more generally,  $\Omega^{p,q}\mathscr{X}$  for Map<sub>\*</sub>( $S^{p,q}, \mathscr{X}$ ).

Suppose  $f : \mathscr{X} \to \mathscr{Y}$  is a morphism in an  $\infty$ -topos. By convention, we will also say that *any* morphism is -2-connected. For  $n \ge -1$  we will say that a morphism f is n-connected if it is an effective epimorphism and for every choice of local base-point the fibers of f are (n-1)-connected. In particular, a space  $\mathscr{X}$  is (n-1)-connected if the map  $\mathscr{X} \to *$  is n-connected.

Given a cofiber sequence

$$\mathscr{X} \longrightarrow \mathscr{Y} \longrightarrow \mathscr{Z}$$

we will refer to  $\mathscr{Z}$  as a cofiber extension of  $\mathscr{Z}$  by  $\mathscr{X}$ . We write

$$\begin{split} \Sigma^{\infty}_{S^1} : \operatorname{Spc}(k)_* &\leftrightarrows \operatorname{SH}^{S^1}(k) : \Omega^{\infty}_{S^1} \\ \Sigma^{\infty} : \operatorname{Spc}(k)_* &\leftrightarrows \operatorname{SH}(k) : \Omega^{\infty} \\ \sigma^{\infty} : \operatorname{SH}^{S^1}(k) &\leftrightarrows \operatorname{SH}(k) : \omega^{\infty} \end{split}$$

Further notation will be introduced below.

#### Acknowledgements

The authors would like to thank M. Levine for questions about the relation between structure of our proof of the Freudenthal suspension theorem with regards to the classical proof. We would also like to thank J. Fasel for discussion around Murthy's conjecture and J.Rognes for helpful comments.

# 2 Preliminaries on unstable and stable motivic homotopy theory

In this section, we recall the basic definitions and properties of unstable and stable motivic homotopy categories in various settings. Notation in the literature can vary quite a bit, but in an effort to keep our presentation as self-contained as possible, we have attempted to summarize everything we need here.

#### 2.1 Motivic localization and the unstable connectivity theorem

2.1.1 (Motivic spaces). A presheaf  $\mathscr{X} \in P(Sm_k)$  is  $\mathbb{A}^1$ -invariant if for every  $U \in Sm_k$  the projection  $U \times \mathbb{A}^1 \to U$  induces an equivalence  $\mathscr{X}(U) \to \mathscr{X}(U \times \mathbb{A}^1_k)$ . The  $\infty$ -category of motivic spaces Spc(k) is the full subcategory of  $Shv_{Nis}(Sm_k)$  spanned by  $\mathbb{A}^1$ -invariant Nisnevich sheaves of spaces. Since homotopy invariance is defined by a small set of conditions, the inclusion  $Spc(k) \subset Shv_{Nis}(Sm_k)$  is an accessible localization [Hoy17, §3.4]. The category Spc(k) is a presentable  $\infty$ -category and the inclusion  $Spc(k) \subset P(Sm_k)$  admits a left adjoint

 $L_{mot}: P(Sm_k) \longrightarrow Spc(k)$ 

that we call the motivic localization functor.

2.1.2 (Variants). There are two classes of variants of Spc(k) that will appear in the body of the text: modifications of the category of schemes we consider (e.g., to permit singularities), and alternative Grothendieck topologies  $\tau$  on such categories of schemes (typically finer than the Nisnevich topology). In practice, we will consider Voevodsky's cdh-topology, Scheiderer's real-étale topology or Kelly's  $\ell$ dh topology. The variants to be discussed will not reappear until Sections 2.3 and 5, and we will include more details and references there as necessary.

Write  $\operatorname{Sch}_k$  for the category of finite type, separated k-schemes. By an admissible subcategory  $\operatorname{S}_k$ of  $\operatorname{Sch}_k$ , we will mean one such that  $X \in \operatorname{S}_k \Longrightarrow \operatorname{A}_X^1 \in \operatorname{S}_k$ ,  $X \in \operatorname{S}_k \Longrightarrow U \in \operatorname{S}_k$  for  $f : U \to X$ any finite-type étale morphism, and  $\operatorname{S}_k$  is closed under finite products and coproducts. Examples include taking  $\operatorname{S}_k$  to be the subcategory  $\operatorname{Sch}_k^{pp}$  consisting of quasi-projective k-schemes or all of  $\operatorname{Sch}_k$ .

If  $\tau$  is a Grothendieck topology on an admissible subcategory  $S_k$  of  $Sch_k$ , then we write  $Spc_{\tau}(S_k)$  for the full subcategory of the  $\infty$ -topos  $Shv_{\tau}(S_k)$  spanned by  $\mathbb{A}^1$ -invariant sheaves of spaces and  $Spc_{\tau}(k)$  for the full subcategory of the  $\infty$ -topos  $Shv_{\tau}(Sm_k)$  spanned by  $\mathbb{A}^1$ -invariant sheaves of spaces. In these situations, we will write  $L_{\tau}$  for the localization functor to  $\tau$ -motivic spaces.

2.1.3 (Motivic localization). Write  $\Delta_k^*$  for the cosimplicial affine space with  $\Delta_k^n := \operatorname{Spec} k[x_0, \ldots, x_n] / \langle \sum_{i=0}^n x_i = 1 \rangle$ . One defines the singular functor by the formula:

$$\operatorname{Sing}^{\mathbb{A}^1} \mathscr{X} := \operatorname{colim}_{n \in \Delta^{\operatorname{op}}} \mathscr{X}(\Delta_k^n \times -).$$

This functor is left adjoint to the inclusion of  $\mathbb{A}^1$ -invariant presheaves of spaces into presheaves of spaces. Morel and Voevodsky construct  $L_{mot}$  explicitly as an infinite composition of  $\mathrm{Sing}^{\mathbb{A}^1}$  and Nisnevich sheafification [MV99, §2 Lemma 3.20].

2.1.4 (Properties of  $L_{mot}$ ). The simplex category is sifted [Lur09, Lemma 5.5.8.4], and one deduces that  $L_{mot}$  preserves finite products [Hoy14, C.6]. Moreover, the functor  $L_{mot}$  is locally cartesian in the sense that for any diagram  $\mathscr{X}_1 \to \mathscr{X}_0 \leftarrow \mathscr{X}_2 \in P(Sm_k)$  with  $\mathscr{X}_1, \mathscr{X}_0 \in Spc(k)$  the canonical map

$$L_{mot}(\mathscr{X}_1 \times_{\mathscr{X}_0} \mathscr{X}_2) \longrightarrow \mathscr{X}_1 \times_{\mathscr{X}_0} L_{mot}(\mathscr{X}_2)$$

is an equivalence. In particular, it follows that colimits in Spc(k) are universal (i.e., colimits are stable by pullback) [Hoy17, Proposition 3.15]. While the localization  $L_{mot}$  is known not to be left exact (i.e., Spc(k) is not an  $\infty$ -topos), universality of colimits in Spc(k) implies that Spc(k) is a semi-topos in the sense of [Lur09, Definition 6.2.3.1].

2.1.5 (Essentially smooth base change). We will call a morphism  $f : X \to Y$  of schemes *essentially* smooth if it can be written as a cofiltered limit  $\lim_{\alpha} X_{\alpha} \to Y$  of smooth, affine transition morphisms. If k is a perfect field and L/k is a field extension then  $f : \operatorname{Spec} L \to \operatorname{Spec} k$  is essentially smooth (see, e.g., [Hoy15, Lemma A.2]). In that case, the functors  $f^* : P(\operatorname{Sm}_k) \to P(\operatorname{Sm}_L)$  preserve  $\mathbb{A}^1$ - and Nisnevich local objects [Hoy15, Lemma A.4]. In this context, suppose we are given a finite diagram  $d : I \to \operatorname{Sm}_L$ which is a cofiltered limit of finite diagrams  $d_{\alpha} : I \to \operatorname{Sm}_{L_{\alpha}}$ . In that case, the evident map

$$\operatorname{colim}_{\alpha} \operatorname{Map}(\operatorname{colim} d_{\alpha}, f_{\alpha}^* \mathscr{X}) \longrightarrow \operatorname{Map}(\operatorname{colim} d, f^* \mathscr{X})$$

is an equivalence by [Hoy15, Lemma A.5]. We will typically appeal to such results by stating that some construction that we want to perform is "compatible with essentially smooth base change".

Morel's foundational work [Mor12] studied the interaction between  $L_{mot}$  and connectivity. Unlike the results above which hold over rather general base schemes, the following result requires strong hypotheses on k and is the main reason we assume k is a field (frequently perfect).

**Theorem 2.1.6** (Morel). Suppose  $n \ge -2$  is an integer and assume k is a field and  $\mathscr{X}$  is a space. Assume either  $n \le 0$  or  $\mathscr{X}$  is pulled back from a perfect subfield of k. If  $\mathscr{X}$  is n-connected, then  $L_{mot}\mathscr{X}$  is also n-connected.

*Proof.* The statement is trivial for n < 0, and for n = 0 follows essentially from the construction of the motivic localization functor [MV99, §2 Corollary 3.22]. Assuming k is perfect, for n > 0 it is [Mor12, Theorem 6.38] (though see [AWW17, Theorem 2.2.12] for a detailed proof). The assertion that the result holds if  $\mathscr{X}$  is pulled back from a perfect subfield then follows by compatibility with essentially smooth base change (see Paragraph 2.1.5).

Morel deduces the above results by analyzing the Postnikov tower of connected motivic local spaces. Key to deducing the above results are the notion of strictly and strongly  $\mathbb{A}^1$ -invariant sheaves of groups.

2.1.7 (Strong and strict  $\mathbb{A}^1$ -invariance). A Nisnevich sheaf of groups  $\mathbf{G}$  on  $\mathrm{Sm}_k$  is *strongly*  $\mathbb{A}^1$ -*invariant* if the cohomology presheaves  $H^i(-, \mathbf{G})$  are  $\mathbb{A}^1$ -invariant for i = 0, 1. A Nisnevich sheaf of abelian groups is *strictly*  $\mathbb{A}^1$ -*invariant* if the cohomology presheaves  $H^i(-, \mathbf{A})$  are  $\mathbb{A}^1$ -invariant for all  $i \ge 0$ .

*Remark* 2.1.8. Morel established that if k is a perfect field, then strongly  $\mathbb{A}^1$ -invariant sheaves of abelian groups are strictly  $\mathbb{A}^1$ -invariant [Mor12, Corollary 5.45], so in practice we will only speak of strong  $\mathbb{A}^1$ -invariance of Nisnevich sheaves of groups. The category of strictly  $\mathbb{A}^1$ -invariant sheaves is abelian (see Paragraph 2.2.15 for further discussion of this point).

2.1.9. A pointed connected space  $\mathscr{X} \in \text{Shv}_{Nis}(\text{Sm}_k)$  is  $\mathbb{A}^1$ -invariant if and only if the homotopy sheaves  $\pi_i(\mathscr{X})$  are strongly  $\mathbb{A}^1$ -invariant for all  $i \ge 1$  [AWW17, Lemma 2.2.11]. Taken together, the above results also imply that the layers of the Postnikov tower of any connected space  $\mathscr{X}$  are themselves  $\mathbb{A}^1$ -invariant.

2.1.10 (Contraction). Recall that if G is a pointed presheaf of groups on  $Sm_k$ , then its contraction  $G_{-1}$  is the presheaf defined by

$$1 \longrightarrow \mathbf{G}_{-1}(U) \longrightarrow \mathbf{G}(U \times \mathbb{G}_m) \longrightarrow \mathbf{G}(U),$$

where the second map is induced by  $id \times 1 : U \to U \times \mathbb{G}_m$  [Mor12, p. 33]. Morel showed that contraction is an endofunctor of the category of strongly  $\mathbb{A}^1$ -invariant sheaves of groups [Mor12, Lemma 2.32] that preserves exact sequences [Mor12, Lemma 7.33]. The key result about contractions that we need is: if  $\mathscr{X}$ is a pointed connected motivic space, then

$$\pi_i(\Omega^{1,1}\mathscr{X}) \simeq \pi_i(\mathscr{X})_{-1},$$

which is established in [Mor12, Theorem 6.13].

2.1.11 (Quotients of strongly  $\mathbb{A}^1$ -invariant sheaves).

**Lemma 2.1.12.** Let k be a perfect field and  $\mathscr{X} \in Shv_{Nis}(Sm_k)$ . Assume that  $\mathscr{X}$  is an H-space (grouplike monoid in the homotopy category) and that the sheaf of groups  $\pi_0 \mathscr{X}$  receives a surjection from a strongly  $\mathbb{A}^1$ -invariant sheaf. Then  $\pi_0 L_{mot} \mathscr{X}$  is strongly  $\mathbb{A}^1$ -invariant.

*Proof.* We know that  $L_{mot} \mathscr{X}$  is still an *H*-space (since  $L_{mot}$  preserves finite products, including the shearing equivalence  $\mathscr{X} \times \mathscr{X} \to \mathscr{X} \times \mathscr{X}$ ,  $(x, y) \mapsto (xy, y)$ ), and hence  $\pi_0 L_{mot} \mathscr{X}$  is  $\mathbb{A}^1$ -invariant [Cho14, Theorem 4.18]. We also know that  $\pi_0 \mathscr{X} \to \pi_0 L_{mot} \mathscr{X}$  is surjective [MV99, §2 Corollary 3.22]. Since  $\pi_0 \mathscr{X}$  receives a surjection from a strongly  $\mathbb{A}^1$ -invariant sheaf hence so does the  $\mathbb{A}^1$ -invariant sheaf  $\pi_0 L_{mot} \mathscr{X}$ , which is thus strongly  $\mathbb{A}^1$ -invariant [Cho14, Theorem 4.18].

Lemma 2.1.13. Let k be a perfect field. The forgetful functor

{strongly  $\mathbb{A}^1$ -invariant sheaves of groups}  $\rightarrow {\mathbb{A}^1$ -invariant sheaves of groups}

preserves sifted colimits.

*Proof.* By [Lur09, Corollary 5.5.8.17] it suffices to treat filtered colimits and geometric realizations. Since cohomology commutes with filtered colimits, this case is clear. Hence let  $\mathbf{G}_{\bullet}$  be a simplicial diagram of strongly  $\mathbb{A}^1$ -invariant sheaves of groups. Let  $|\mathbf{G}_{\bullet}|^{\text{Nis}}$  denote the colimit in Nisnevich sheaves. Then  $\pi_0|\mathbf{G}_{\bullet}|^{\text{Nis}}$  is a quotient of the strongly  $\mathbb{A}^1$ -invariant sheaf  $\pi_0\mathbf{G}_0$ . It follows via Lemma 2.1.12 that  $\mathbf{C} := \pi_0 L_{\text{mot}}|\mathbf{G}_{\bullet}|^{\text{Nis}}$  is (grouplike and) strongly  $\mathbb{A}^1$ -invariant. Thus  $\mathbf{C}$  is the common colimit of  $\mathbf{G}_{\bullet}$  in the two categories.

**Corollary 2.1.14.** Let k be a perfect field and  $\mathscr{X}_{\bullet} \in \text{Mon}(\text{Spc}(k))$  a sifted (respectively simplicial) diagram. Assume that each  $\pi_0(\mathscr{X}_{\bullet})$  (respectively just  $\pi_0(\mathscr{X}_0)$ ) is grouplike and strongly  $\mathbb{A}^1$ -invariant. Then  $\pi_0 \text{ colim } \mathscr{X}_{\bullet}$  is grouplike and strongly  $\mathbb{A}^1$ -invariant.

*Proof.* As before  $\pi_0 \operatorname{colim} X_{\bullet}$  is grouplike and  $\mathbb{A}^1$ -invariant by [Cho14, Theorem 4.18]. It follows that  $\pi_0 \operatorname{colim} \mathscr{X}_{\bullet}$  is the colimit of the diagram  $\pi_0 \mathscr{X}_{\bullet}$  in the category of  $\mathbb{A}^1$ -invariant sheaves of groups. In the sifted case, this colimit is strongly  $\mathbb{A}^1$ -invariant by Lemma 2.1.13. In the simplicial case we know that  $\pi_0 \mathscr{X}_0 \to \pi_0 \operatorname{colim} \mathscr{X}_{\bullet}$  is surjective, and so again the target is strongly  $\mathbb{A}^1$ -invariant by Lemma 2.1.12.  $\Box$ 

#### Nilpotence, solvability and functorial towers

2.1.15 (Lower central series). Assume  $\pi$  is a strongly  $\mathbb{A}^1$ -invariant sheaf of groups acting on a strongly  $\mathbb{A}^1$ -invariant sheaf of groups  $\mathbf{G}$ . In that case, there exists [ABH22, Construction 4.9] an initial strongly  $\mathbb{A}^1$ -invariant quotient of  $\mathbf{G}$  on which  $\pi$  acts trivially; we write  $\Gamma^2_{\pi}\mathbf{G} \subset \mathbf{G}$  for the (strongly  $\mathbb{A}^1$ -invariant) kernel of the quotient map. In the case where  $\mathbf{G}$  acts on itself by conjugation, we write  $[\mathbf{G}, \mathbf{G}]_{\mathbb{A}^1}$  for this subsheaf and refer to it as the  $\mathbb{A}^1$ -commutator subsheaf (a priori it could differ from commutator subgroup sheaf). Inductively, one then defines the  $\mathbb{A}^1$ -lower central series  $\Gamma^i_{\pi}\mathbf{G}$  for the action of  $\pi$  on  $\mathbf{G}$  in the usual way:  $\Gamma^1_{\pi}\mathbf{G} = \mathbf{G}$  and  $\Gamma^{i+1}_{\pi}\mathbf{G} = \Gamma^2_{\pi}\Gamma^i_{\pi}\mathbf{G}$ ; when  $\pi = \mathbf{G}$  acting by conjugation we simply drop  $\pi$  from the notation. An action of  $\pi$  on  $\mathbf{G}$  is called  $\mathbb{A}^1$ -nilpotent if and only if  $\Gamma^i_{\pi}\mathbf{G} = 1$  after finitely many steps, i.e., the  $\mathbb{A}^1$ -lower central series is finite.

2.1.16. One can also define the  $\mathbb{A}^1$ -derived series by iterating the  $\mathbb{A}^1$ -commutator construction. Given a strongly  $\mathbb{A}^1$ -invariant sheaf of groups  $\mathbf{G}$ , one sets  $\mathbf{G}_{\mathbb{A}^1}^{(0)} := \mathbf{G}$ , and defines inductively  $\mathbf{G}_{\mathbb{A}^1}^{(i+1)} = [\mathbf{G}_{\mathbb{A}^1}^{(i)}, \mathbf{G}_{\mathbb{A}^1}^{(i)}]_{\mathbb{A}^1}$ . We say that a strongly  $\mathbb{A}^1$ -invariant sheaf of groups is  $\mathbb{A}^1$ -solvable if  $\mathbf{G}_{\mathbb{A}^1}^{(i)}$  terminates after finitely many steps. By induction it follows that an  $\mathbb{A}^1$ -solvable group is a successive extension of strictly  $\mathbb{A}^1$ -invariant sheaves.

2.1.17 (Solvable motivic spaces). A pointed, connected motivic space  $\mathscr{X}$  will be called *solvable* if  $\pi_1(\mathscr{X})$  is  $\mathbb{A}^1$ -solvable.

**Proposition 2.1.18.** If  $\mathscr{X}$  is a solvable motivic space, then  $\mathscr{X}$  admits a functorial Whitehead tower. More precisely, there exist a weakly increasing sequence of integers  $1 \le n_1 \le n_2 \le \cdots$ , strictly  $\mathbb{A}^1$ -invariant sheaves  $\mathbf{A}_i(\mathscr{X})$ , and spaces  $\mathscr{X}\langle i \rangle$ ,  $i \ge 0$ , fitting into fiber sequences of the form

 $\mathscr{X}\langle i+1\rangle \longrightarrow \mathscr{X}\langle i\rangle \longrightarrow K(\mathbf{A}_i(\mathscr{X}), n_i)$ 

such that:

1.  $\mathscr{X}\langle 0\rangle = \mathscr{X},$ 

- 2.  $\mathscr{X}\langle i \rangle$  is  $n_i$ -connective, and
- 3.  $n_i \to \infty$  as  $i \to \infty$ .

*Proof.* Set  $\mathscr{X}\langle 0 \rangle = \mathscr{X}$ . We define  $\mathscr{X}\langle i \rangle$  inductively as follows. Since  $\mathscr{X}\langle i \rangle$  is  $n_i$ -connective, its first non-trivial homotopy sheaf appears in degree  $n_i$ . If  $n_i = 1$ , then we define  $\mathscr{X}\langle i+1 \rangle$  to be the fiber of the composite map:

$$\mathscr{X}\langle i\rangle \longrightarrow B\pi_1(\mathscr{X}\langle i\rangle) \longrightarrow B\pi_1(\mathscr{X})^{ab}_{\mathbb{A}^1}$$

If  $n_i > 1$ , then we define  $\mathscr{X}\langle i+1 \rangle$  to be the fiber of the corresponding k-invariant in the Postnikov tower. The assumption that  $\mathscr{X}$  is solvable guarantees that  $n_i$  is eventually greater than 1 and we conclude.

2.1.19 (Nilpotent motivic spaces). A pointed, connected motivic space  $\mathscr{X}$  is called *simple* if  $\pi_1(\mathscr{X})$  is abelian and the action of  $\pi_1(\mathscr{X})$  on higher homotopy sheaves is trivial. Any motivic *h*-space is simple. A motivic space is called *nilpotent* if  $\pi_1(\mathscr{X})$  is  $\mathbb{A}^1$ -nilpotent and acts  $\mathbb{A}^1$ -nilpotently on higher homotopy sheaves [AFH22, Definition 3.3.1]. Note that any nilpotent motivic space is solvable in the sense above. If  $\mathscr{X}$  is a nilpotent motivic space, then the usual Postnikov tower of  $\mathscr{X}$  admits a functorial principal refinement [ABH22, Theorem 5.1]. For later use, and to make the presentation more self-contained we summarize this in the following statement.

**Proposition 2.1.20.** If  $\mathscr{X}$  is a nilpotent motivic space, then the Postnikov tower admits a functorial principal refinement. More precisely, there exist a weakly increasing sequence of integers  $n_i, i \ge 2$ , strictly  $\mathbb{A}^1$ -invariant sheaves  $\mathbf{A}_i(\mathscr{X})$ , morphisms  $\mathscr{X} \to \mathscr{X}_i$ , and fiber sequences of the form

$$\mathscr{X}_{i+1} \longrightarrow \mathscr{X}_i \longrightarrow K(\mathbf{A}_i(\mathscr{X}), n_i)$$

such that:

1. the induced map  $\mathscr{X} \to \lim_i \mathscr{X}_i$  is an equivalence

2.  $n_i \to \infty$  as  $i \to \infty$ .

#### Localization at primes

If  $R \subset \mathbb{Q}$  is a subring. We will need to consider unstable *R*-localization of  $\operatorname{Spc}(k)$  as discussed in [AFH22, Definition 4.3.1]; we write  $\mathscr{X}_R$  or  $\mathscr{X} \otimes R$  for what we called  $\operatorname{L}_R \mathscr{X}$  in [AFH22]. In particular, we recall the following fact.

**Theorem 2.1.21** ([AFH22, Theorem 4.3.9]). Assume k is a field. If  $\mathscr{X} \in \text{Spc}(k)_*$  is a weakly nilpotent motivic space, then  $\mathscr{X}_R$  is connected and weakly nilpotent and the map

$$\pi_i(\mathscr{X})\otimes R\longrightarrow \pi_i(\mathscr{X}_R)$$

is an isomorphism for every  $i \ge 1$  (where for i = 1, the tensor product is interpreted as in [AFH22, Theorem 4.3.7])

#### Motivic localization, colimits and limits

2.1.22. Let  $\mathscr{X} \in \operatorname{Shv}_{\operatorname{Nis}}(\operatorname{Sm}_k)_*$ . Then  $\Omega \mathscr{X}$  has the structure of a group in the sense of homotopy theory. In fact the induced functor from pointed connected Nisnevich sheaves to groups is an equivalence [Lur09, Lemma 7.2.2.11] with inverse denoted  $\mathscr{B}$ . Furthermore given  $f : \mathscr{E} \to \mathscr{B}$ , fib(f) carries a canonical action by  $\Omega \mathscr{B}$ . If  $\mathscr{B}$  is connected, sending  $f : \mathscr{E} \to \mathscr{B}$  to fib(f) with its  $\Omega \mathscr{B}$ -action defines a functor from the category of spaces over  $\mathscr{B}$  to spaces with  $\Omega \mathscr{B}$ -action. In fact, this functor is an equivalence with inverse obtained by sending a space  $\mathscr{F}$  with action of  $\Omega \mathscr{B}$  to the map  $\mathscr{F}/\!\!/\Omega \mathscr{B} \to */\!\!/\Omega \mathscr{B}$  where  $\mathscr{F}/\!\!/\Omega \mathscr{B}$  denotes the (homotopy) quotient, a.k.a. bar construction, i.e., the geometric realization of

$$\mathscr{F} = \mathscr{F} \times \Omega \mathscr{B} = \cdots$$

These results carry to the motivic situation, with one small modification.

Lemma 2.1.23. Let k be a perfect field.

- 1. The functor  $\Omega$  :  $\operatorname{Spc}(k)_* \to \operatorname{Mon}(\operatorname{Spc}(k))$  induces an equivalence between pointed connected motivic spaces and grouplike monoids in motivic spaces with the additional condition that  $\pi_0$  be strongly  $\mathbb{A}^1$ -invariant; its inverse is the classifying space functor B.
- 2. Let  $\mathscr{B} \in \operatorname{Spc}(k)_*$  be connected. There is an equivalence of categories between  $\mathscr{E} \in \operatorname{Spc}(k)_{/\mathscr{B}}$  and the category of  $\mathscr{F} \in \operatorname{Spc}(k)$  with an action by  $\Omega \mathscr{B}$ . It sends  $\mathscr{E}$  to  $\operatorname{fib}(\mathscr{E} \to \mathscr{B})$  with the canonical action, and it sends  $\mathscr{F}$  to  $\mathscr{F}/\!\!/\Omega \mathscr{B}$ .

*Proof.* (1) is an immediate consequences of the analogous results in  $\operatorname{Shv}_{\operatorname{Nis}}(\operatorname{Sm}_k)$ , using the characterization of pointed, connected  $\mathbb{A}^1$ -local spaces in terms of homotopy sheaves (Paragraph 2.1.9). For (2), it is clear that the functor is fully faithful. To prove essential surjectivity it suffices to show that if  $\mathscr{F} \in \operatorname{Shv}_{\operatorname{Nis}}(\operatorname{Sm}_k)_*$  carries an action by  $\Omega \mathscr{B}$  with  $\mathscr{F}, \mathscr{B} \mathbb{A}^1$ -invariant, then also  $\mathscr{F}/\!\!/\Omega \mathscr{B}$  is  $\mathbb{A}^1$ -invariant. This follows (via [AWW17, Lemma 2.2.10]) by considering the fiber sequence

$$\mathscr{F} \to \mathscr{F} /\!\!/ \Omega \mathscr{B} \to \mathscr{B} \in \mathrm{Shv}_{\mathrm{Nis}}(\mathrm{Sm}_k),$$

in which by assumption  $\mathscr{B}$  is connected and  $\mathscr{F}, \mathscr{B}$  are both  $\mathbb{A}^1$ -invariant.

**Theorem 2.1.24.** Assume k is a field and suppose we have a cartesian diagram of pointed objects in  $Shv_{Nis}(Sm_k)$ :

$$\begin{array}{c} \mathscr{X}_{00} \xrightarrow{g_0} \mathscr{X}_{10} \\ \downarrow^{f_0} & \downarrow^{f_1} \\ \mathscr{X}_{01} \xrightarrow{g_1} \mathscr{X}_{11} \end{array}$$

that is pulled back from a perfect subfield. If  $\mathscr{X}_{11}$  is connected and  $\pi_0(L_{mot}\Omega\mathscr{X}_{11})$  is strongly  $\mathbb{A}^1$ -invariant (in particular, if  $\pi_1(\mathscr{X})$  is strongly  $\mathbb{A}^1$ -invariant, e.g., trivial), then the diagram

$$L_{mot} \mathscr{X}_{00} \xrightarrow{g_0} L_{mot} \mathscr{X}_{10}$$

$$\downarrow^{f_0} \qquad \qquad \downarrow^{f_1}$$

$$L_{mot} \mathscr{X}_{01} \xrightarrow{g_1} L_{mot} \mathscr{X}_{11}$$


is cartesian as well.

*Proof.* We can view  $\mathscr{X}_{00}, \mathscr{X}_{01}, \mathscr{X}_{10} \in \operatorname{Shv}_{\operatorname{Nis}}(\operatorname{Sm}_k)_{/\mathscr{X}_{11}}$ . Put  $\mathscr{G} = \Omega \mathscr{X}_{11}, \mathscr{F} := \operatorname{fib}(\mathscr{X}_{01} \to \mathscr{X}_{11}), \mathscr{H} := \operatorname{fib}(\mathscr{X}_{10} \to \mathscr{X}_{11})$ . Under the equivalence of  $\operatorname{Shv}_{\operatorname{Nis}}(\operatorname{Sm}_k)_{/\mathscr{X}_{11}}$  with spaces with an action by  $\mathscr{G}, \mathscr{X}_{01}, \mathscr{X}_{10}$  correspond respectively to  $\mathscr{F}, \mathscr{H}$ . Noting that  $\mathscr{X}_{00}$  is the product of  $\mathscr{X}_{01}, \mathscr{X}_{10} \in \operatorname{Shv}_{\operatorname{Nis}}(\operatorname{Sm}_k)_{/\mathscr{X}_{11}}$  and applying the inverse equivalence, we deduce that


$$\mathscr{X}_{00} \simeq (\mathscr{F} \times \mathscr{H}) /\!\!/ \mathscr{G}$$

The same logic applies to  $L_{mot} \mathscr{X}_{10} \times_{L_{mot} \mathscr{X}_{11}} L_{mot} \mathscr{X}_{01}$ . It thus suffices to prove that  $L_{mot}$  preserves the fiber sequences  $\mathscr{G} \to * \to \mathscr{X}_{11}, \mathscr{F} \to \mathscr{X}_{01} \to \mathscr{X}_{11}$  and  $\mathscr{H} \to \mathscr{X}_{10} \to \mathscr{X}_{11}$ . This is proved in [AWW17, Theorem 2.3.3].


**Proposition 2.1.25** (Realization fibrations). Assume k is a perfect field, and I is a sifted category. Assume  $\mathscr{X}^{ij}$ , i, j = 0, 1 are I-diagrams in Spc(k) fitting into a cartesian square of the form:



If  $\mathscr{X}^{11}$  is objectwise connected, then writing  $|\mathscr{X}^{ij}|_{\mathbf{I}} := \operatorname{colim}_{\mathbf{I}} \mathscr{X}^{ij}$  there is a cartesian square of motivic spaces of the form



*Proof.* Write  $|-|_{\mathbf{I}}^{\text{Nis}}$  for  $\operatorname{colim}_{\mathbf{I}} \mathscr{X}^{ij}$  computed in  $\operatorname{Shv}_{\operatorname{Nis}}(\operatorname{Sm}_k)$ . In that case, observe that the hypothesis that  $\mathscr{X}^{11}$  is objectwise connected implies that there is a cartesian square of the form



this result was mentioned by Rezk [Rez14, Theorem 4.4], see also [Lur17, Lemma 5.5.6.17]. The same argument shows that  $\Omega |\mathscr{X}^{11}|_{\mathbf{I}}^{\text{Nis}} \simeq |\Omega \mathscr{X}^{11}|_{\mathbf{I}}^{\text{Nis}}$ . Corollary 2.1.14 shows that  $\pi_0 L_{\text{mot}} |\Omega \mathscr{X}^{11}|_{\mathbf{I}}^{\text{Nis}}$  is strongly  $\mathbb{A}^1$ -invariant. The result thus follows by appeal to Theorem 2.1.24.

**Proposition 2.1.26.** If k is a perfect field, then the functor  $\Omega$  preserves sifted colimits of connected motivic spaces.

*Proof.* Assume I is a sifted category and suppose  $\mathscr{X} : \mathbf{I} \to \operatorname{Spc}(k)_*$  is a sifted diagram of motivic spaces. In that case, there is a cartesian square of the form



The result then follows immediately from Proposition 2.1.25 (using that sifted simplicial sets are weakly contractible [Lur09, Proposition 5.5.8.7]).  $\Box$ 

### 2.2 Stable motivic homotopy theory and effectivity

2.2.1 (Presentably symmetric monoidal categories). Write  $\Pr^L$  for the  $\infty$ -category whose objects are presentable  $\infty$ -categories and morphisms are colimit preserving functors [Lur09, Definition 5.5.3.1]. The category  $\Pr^L$  admits limits [Lur09, Proposition 5.5.3.13] and colimits [Lur09, Theorem 5.5.3.18]. The category  $\Pr^L$  can be equipped with a symmetric monoidal structure  $\Pr^{L,\otimes}$  [Lur17, Proposition 4.8.1.15]. By a *presentably symmetric monoidal category* we will mean a commutative algebra in  $\Pr^{L,\otimes}$  and we write  $\operatorname{CAlg}(\Pr^{L,\otimes})$  for the category of commutative algebra objects. Recall that a presentably symmetric monoidal  $\infty$ -category  $\mathscr{C}$  such that the tensor product  $\mathscr{C} \times \mathscr{C} \to \mathscr{C}$  preserves colimits in each variable separately [Lur09, Notation 4.8.1.2(iv)].

2.2.2 (Stable homotopy categories). Given any presentably symmetric monoidal  $\infty$ -category C and a set of objects X in C, we can invert the tensor product with the objects X. More precisely there exists a new presentably symmetric monoidal infinity category  $C[X^{-1}]$  with the following universal property in the infinity category  $CAlg(Pr^{L,\otimes})$ : given any functor  $C \to D$  such that f(x) is invertible for every  $x \in X$ , it factors uniquely though  $C[X^{-1}]$ ; we refer the reader to [Rob15, §2.1] for further details of this construction. 2.2.3 (S<sup>1</sup> and  $\mathbb{P}^1$ -spectra). The category of pointed motivic spaces  $\operatorname{Spc}(k)_*$  is a presentably symmetric monoidal infinity category via the smash product of pointed spaces. We define

$$SH^{S^{1}}(k) := Spc(k)_{*}[(S^{1})^{-1}]$$
$$SH(k) := Spc(k)_{[}(\mathbb{P}^{1})^{-1}].$$

These categories come equipped with canonical adjunctions

$$\operatorname{Spc}(k)_{*} \xrightarrow{\Sigma_{S^{1}}^{\infty}} \operatorname{SH}^{S^{1}}(k)$$
$$\operatorname{Spc}(k)_{*} \xrightarrow{\Sigma^{\infty}} \operatorname{SH}(k)$$

where  $\Sigma_{S^1}^{\infty}$  and  $\Sigma^{\infty}$  are symmetric monoidal. The categories  $\mathrm{SH}^{S^1}(k)$  and  $\mathrm{SH}(k)$  are stable  $\infty$ -categories and are generated under sifted colimits by  $S^1$  or  $\mathbb{P}^1$ -desuspensions of suspension spectra of smooth schemes. We refer the reader to [Hoy17, Proposition 6.4] for further details. For the moment, we observe only that The passage from  $S^1$ -spectra to  $\mathbb{P}^1$ -spectra also fits into an adjunction.

$$\operatorname{SH}^{S^1}(k) \xrightarrow[\omega^{\infty}]{\sigma^{\infty}} \operatorname{SH}(k)$$

having properties analogous to those mentioned above. (Indeed since  $\mathbb{P}^1\simeq S^1\wedge\mathbb{G}_m$  we have  $\mathrm{SH}(k)\simeq$  $\mathrm{SH}^{S^1}(k)[\mathbb{G}_m^{-1}].)$ 

2.2.4. If A is a strictly  $\mathbb{A}^1$ -invariant sheaf of groups, we will write HA for the Eilenberg-Mac Lane  $S^1$ -spectrum determined by **A**.

# **Proposition 2.2.5.** Assume k is a field.

- 1. The functor  $\Omega_{S^1}^{\infty} : \operatorname{SH}^{S^1}(k)_{\geq 0} \longrightarrow \operatorname{Spc}(k)_*$  preserves sifted colimits. 2. If k is furthermore perfect, then the functor  $\Omega^{1,1}$  on  $\operatorname{Spc}(k)_*$  preserve sifted colimits of 1-connected spaces.
- 3. If k is perfect, then the functor  $\Omega^{2,1}$  preserves sifted colimits of 1-connected spaces, while the functor  $\Omega^{2,1}\Sigma^{2,1}$  preserves sifted colimits of connected spaces.

*Proof.* For the first statement, we begin by observing that  $\mathbb{A}^1$ -invariant Nisnevich sheaves of spectra are closed under colimits in the larger category of Nisnevich sheaves of spectra. Therefore, it suffices to establish the result for Nisnevich sheaves of spectra. To establish the result for Nisnevich sheaves of spectra, it suffices to check stalkwise, in which case we may reduce to the case of ordinary spectra, which may be found in [Lur17, Proposition 1.4.3.9].

For the second statement, suppose I is a sifted category and  $\mathscr{X} : \mathbf{I} \to \operatorname{Spc}(k)_*$  is a sifted diagram of pointed 1-connected motivic spaces. Recall that  $\Omega^{1,1}$  preserves connectivity (see Paragraph 2.1.10). In that case, we can consider the *i*-connective covers  $\mathscr{X}(i)$ , which fit into principal fiber sequences of the form:

$$\mathscr{X}\langle i+1\rangle \longrightarrow \mathscr{X}\langle i\rangle \longrightarrow K(\boldsymbol{\pi}_i\mathscr{X},i)$$

where  $i \ge 2$ . Write  $|-|_{\mathbf{I}}$  for colim<sub>I</sub>. The above is an I-diagram of fiber sequences with 1-connected base, so it follows from Proposition 2.1.25 (which uses the assumption k is perfect because of the implicit appeal to the unstable connectivity theorem) that

$$|\mathscr{X}\langle i+1\rangle|_{\mathbf{I}} \longrightarrow |\mathscr{X}\langle i\rangle|_{\mathbf{I}} \longrightarrow |K(\pi_{i}\mathscr{X},i)|_{\mathbf{I}}$$

is a fiber sequence.

We first establish the result for  $|K(\pi_i \mathscr{X}, i)|_{\mathbf{I}}$  by comparison with the  $S^1$ -stable case. Indeed, each  $K(\pi_i \mathscr{X}, i)$  is  $\Omega_{S^1}^{\infty} \Sigma^i H \pi_i \mathscr{X}$  for  $i \ge 2$ , and in fact the entire diagram of Eilenberg–MacLane spaces lifts to a diagram of Eilenberg–MacLane spectra (e.g. use [Lur09, Proposition 1.4.3.9]). By the first point, the functor  $\Omega_{S^1}^{\infty}$  preserves sifted colimits. Granted that observation, observe that

$$\Omega^{1,1}|K(\boldsymbol{\pi}_{i}\mathscr{X},i)|_{\mathbf{I}} \simeq \Omega^{1,1}|\Omega_{S^{1}}^{\infty}\Sigma^{i}H\boldsymbol{\pi}_{i}\mathscr{X}|_{\mathbf{I}} \simeq \Omega^{1,1}\Omega_{S^{1}}^{\infty}|\Sigma^{i}H\boldsymbol{\pi}_{i}\mathscr{X}|_{\mathbf{I}} \simeq \Omega_{S^{1}}^{\infty}\Omega^{1,1}|\Sigma^{i}H\boldsymbol{\pi}_{i}\mathscr{X}|_{\mathbf{I}},$$

where the third equivalence follows from the fact established in the first point that  $\Omega_{S^1}^{\infty}$  preserves sifted colimits. Now,  $\Omega^{1,1}$  commutes with colimits of spectra and therefore, we conclude that

$$\Omega_{S^1}^{\infty}\Omega^{1,1}|\Sigma^i H\boldsymbol{\pi}_i \mathscr{X}|_{\mathbf{I}} \simeq \Omega_{S^1}^{\infty}|\Omega^{1,1}\Sigma^i H\boldsymbol{\pi}_i \mathscr{X}|_{\mathbf{I}} \simeq |\Omega^{1,1} K(\boldsymbol{\pi}_i \mathscr{X},i)|_{\mathbf{I}}.$$

where the last equivalence follows by repeating in the opposite order the equivalences described two displays above.

Now, since applying  $\Omega^{1,1}$  preserves fiber sequences and connectivity, another application of Proposition 2.1.25 implies that there is a fiber sequence of the form

$$|\Omega^{1,1}\mathscr{X}\langle i+1\rangle|_{\mathbf{I}} \longrightarrow |\Omega^{1,1}\mathscr{X}\langle i\rangle|_{\mathbf{I}} \longrightarrow |\Omega^{1,1}K(\boldsymbol{\pi}_{i}\mathscr{X},i)|_{\mathbf{I}}.$$

In that case, there is a commutative diagram of fiber sequences of the form

The right vertical map is an equivalence by the preceding discussion. Taking fibers horizontally and appealing to [ABH22, Proposition 3.1], we conclude that the map from the fiber  $\mathscr{F}\langle i+1\rangle$  of the left vertical map to the fiber  $\mathscr{F}\langle i\rangle$  of the middle vertical map is an equivalence. Note that  $\mathscr{F}\langle i\rangle$  is at least *i*-connective, so the equivalence above followed by induction shows that  $\mathscr{F}\langle i\rangle$  is  $\infty$ -connective and thus contractible, which is what we wanted to show.

Writing  $\Omega^{2,1} = \Omega \Omega^{1,1}$  and combining the conclusion of Point (2) with Proposition 2.1.26, we see that  $\Omega^{2,1}$  preserves sifted colimits of 1-connected spaces. Since  $\Sigma^{2,1}$  preserves colimits, and since  $\Sigma^{2,1}$  sends 0-connected spaces to 1-connected spaces by the unstable connectivity theorem 2.1.6, we also conclude that  $\Omega^{2,1}\Sigma^{2,1}(-)$  also preserves sifted colimits of 0-connected spaces.

*Remark* 2.2.6. Assume k is a field and Y is a pointed open subscheme of  $\mathbb{A}^1$ . The same style of proof can be used to establish that the internal mapping space  $\underline{Map}_*(Y, -)$  preserves sifted colimits. Indeed, by [Mor12, Remark 2.33], given a strongly  $\mathbb{A}^1$ -invariant sheaf G, we may define

$$\mathbf{G}^{(Y)}(U) := \ker(\mathbf{G}(Y \times U) \to \mathbf{G}(U))$$

This formula defines a sheaf of groups, and the same argument as in [Mor12, Lemma 2.32] shows that  $B\mathbf{G}^{(Y)}$  can be identified with the connected component of the identity map in  $\underline{\mathrm{Map}}_{*}(Y, B\mathbf{G})$  and thus  $\mathbf{G}^{(Y)}$  is again strongly  $\mathbb{A}^{1}$ -invariant. In that case, the evident analog of [Mor12, Lemma 6.14] holds using homotopy purity replacing  $\mathbb{G}_{m}$  by Y and thus the proof of [Mor12, Theorem 6.13] goes through with  $\mathbb{G}_{m}$  replaced by Y. In particular,  $\mathrm{Map}_{*}(Y, -)$  preserves connectivity so we can repeat the argument above.

2.2.7. If  $E \in \text{CAlg}(\text{SH}(k))$ , we write  $\text{Mod}_E(k)$  for the category of *E*-module spectra. Of particular interest will be the case where  $E = H\mathbb{Z}$ , Voevodsky's motivic Eilenberg–Mac Lane spectrum or  $H\mathbb{Z}[1/p]$ where *p* is the characteristic exponent of *k* (note: a minor conflict arises between this notation and that for the Eilenberg–Mac Lane  $S^1$ -spectrum attached to a strictly  $\mathbb{A}^1$ -invariant sheaf; i.e., Paragraph 2.2.4—we trust that this will not lead to confusion). By [RØ08, Theorem 1], we know that if *k* has characteristic 0, then  $\text{Mod}_{H\mathbb{Z}}(k)$  is equivalent to Voevodsky's (unbounded) derived category of motives  $\mathbf{DM}(k)$ . This result was extended to an equivalence of  $\text{Mod}_{H\mathbb{Z}[1/p]}(k)$  and Voevodsky's (unbounded) derived category of motives with  $\mathbb{Z}[1/p]$ -coefficients  $\mathbf{DM}(k, \mathbb{Z}[1/p])$  in [EK20b, Corollary 5.3].

2.2.8 (Essentially smooth base change revisited). The results of Paragraph 2.1.5 carry over to  $\text{SH}^{S^1}(k)$  or SH(k); see [Hoy15, Lemma A.7] for details. These results also carry over to  $\text{Mod}_{H\mathbb{Z}[1/p]}$  by the same result and the discussion of Paragraph 2.2.7.

2.2.9 (Effective spectra). We write  $\operatorname{SH}^{S^1}(k)(d)$  for the localizing subcategory generated by  $\Sigma^{d,d}\Sigma_{S^1}^{\infty}X_+$  for  $X \in \operatorname{Sm}_k$ ; we refer to this category as the subcategory of *d*-effective  $S^1$ -spectra. We also write  $\operatorname{SH}(k)^{\text{eff}}$  for the localizing subcategory of  $\operatorname{SH}(k)$  generated by spectra of the form  $\Sigma_{\mathbb{P}^1}^{\infty}X_+$  (i.e., no  $\mathbb{G}_m$ -desuspensions) and we write  $\operatorname{SH}(k)^{\text{eff}}(n)$  for the category  $\Sigma^{2n,n}\operatorname{SH}(k)^{\text{eff}}$ ; this category can be identified with  $\Sigma^{n,n}\operatorname{SH}(k)^{\text{eff}}$  because it is triangulated.

2.2.10 (Slices). The inclusion  $i_n : SH(k)^{\text{eff}}(n) \hookrightarrow SH(k)^{\text{eff}}$  admits a right adjoint  $r_n$  as a consequence of Neeman's Brown representability theorem [Nee01, Theorem 8.4.4]. We set  $f_n := i_n \circ r_n$ . There are natural transformations  $f_n \to f_m$  whenever m < n and a tower of the form

 $\cdots \longrightarrow f_n E \longrightarrow f_{n-1}E \longrightarrow \cdots \longrightarrow f_0E \longrightarrow f_{-1}E \longrightarrow \cdots \longrightarrow E.$ 

It is called the slice tower; the layers  $f_n E$  are *effective covers* of E and the *slices*  $s_i E$  are defined as the cofibers of  $f_{n+1}E \to f_n E$ .

Voevodsky showed [Voe04, Theorem 6.6] that  $s_0(\mathbb{1}_k) \simeq H\mathbb{Z}$  when k is a field having characteristic 0 and Levine generalized this result (by a completely different argument) to k an arbitrary field [Lev08, Theorem 10.5.1]. Now, the slice filtration is multiplicative, so the slices  $s_i E$  have the structure of a module over  $s_0\mathbb{1}$  [Pel11, Theorem 3.6.13(6)].

2.2.11 (Very effective spectra). Following Spitzweck–Østvær [SØ12], define  $SH(k)^{veff}$  to be the subcategory generated by taking homotopy colimits and extensions of objects of the form  $\Sigma^{n,0}\Sigma_{\mathbb{P}^1}^{\infty}X_+$ ,  $n \ge 0$ . More generally, if  $E \in CAlg(SH(k))$ , we write  $Mod_E(k)^{veff}$  for the subcategory of  $Mod_E(k)$  generated under colimits and extensions by  $E \wedge \Sigma_{\mathbb{P}^1}^{\infty}X_+$ ,  $X \in Sm_k$ .

#### Recollections on inverting primes, strong dualizability

Given a subring  $R \subset \mathbb{Q}$ , we can consider variants of all the stable categories above where with Rcoefficients, e.g.,  $\mathrm{SH}^{S^1}(k)_R$ ,  $\mathrm{SH}(k)_R$ , etc. Of particular interest will be the cases  $R = \mathbb{Q}$ ,  $R = \mathbb{Z}[\frac{1}{2}]$ and  $R = \mathbb{Z}[\frac{1}{n}]$  where p is the characteristic exponent of k.

**Proposition 2.2.12.** Assume k is a field with characteristic exponent p.

- 1. For any  $U \in \text{Sm}_k$ ,  $\Sigma^{\infty}U_+$  is strongly dualizable in  $\text{SH}(k)[\frac{1}{n}]$ .
- 2. The category  $\operatorname{SH}(k)[\frac{1}{p}]$  is generated under colimits by  $\Sigma^{\infty}X_{+}$  for  $X \in \operatorname{Sm}_{k}$  projective or, more generally, by strongly dualizable objects.

*Proof.* The case where k has characteristic 0 is [Rio05, Théoréme 1.4,2.2] appealing to resolution of singularities. By appeal to Gabber's refined alternations, Riou [LYZR19, B.1-2] extended this argument to the case where k was perfect. The assumption k perfect is removed in [EK20a, Theorem 3.2.1].

We will also need some comparisons between unstable and stable localizations. By checking stalkwise, the following result follows immediately from the corresponding classical statement (which can be proved by e.g. observing that  $\mathscr{X}$  below has vanishing *p*-completion).

**Proposition 2.2.13.** Assume k is a field and  $\mathscr{X}$  is a pointed 1-connected motivic space. If for all  $i, \pi_i(\mathscr{X})$  is uniquely p-divisible (for some  $p \in \mathbb{Z}$ ), then  $\sum_{S^1} \mathscr{X} \simeq \sum_{S^1} \mathscr{X}[\frac{1}{p}]$ .

**Proposition 2.2.14.** Assume k is a field and  $\mathscr{X}$  is a pointed 1-connected motivic space. If  $\Sigma_{S^1}^{\infty} \mathscr{X} \simeq \Sigma_{S^1}^{\infty} \mathscr{X}[\frac{1}{n}]$ , then  $\pi_i(\mathscr{X})$  is uniquely p-divisible for each  $i \geq 1$ .

*Proof.* Consider the  $\mathbb{Z}[\frac{1}{p}]$ -localization described in Theorem 2.1.21, which has the property that  $\pi_i(\mathscr{X}_{\mathbb{Z}[1/p]})$  tensors  $\pi_i(\mathscr{X})$  with  $\mathbb{Z}[1/p]$ . The map  $\Sigma_{S^1}^{\infty} \mathscr{X} \to \Sigma_{S^1}^{\infty}(\mathscr{X}_{\mathbb{Z}[1/p]})$  is a  $\mathbb{Z}[1/p]$ -equivalence of *p*-periodic spectra (by assumption for the first and by the preceding proposition for the second statement), whence an equivalence. The Whitehead theorem [ABH22, Theorem 5.3] implies that  $\mathscr{X} \to \mathscr{X}_{\mathbb{Z}[1/p]}$  is an equivalence as well.

#### **Generation of** *t***-structures**

We review here some facts about generation of *t*-structures, which seems to have been independently discovered by several authors including in unpublished work of Morel [Ayo07, §2.1.3 p. 280] and [ATJLSS03]. Lurie has also given a treatment of these ideas in the context of stable presentable  $\infty$ -categories.

2.2.15 (Homotopy *t*-structure on  $S^1$ -spectra). The category  $\operatorname{SH}^{S^1}(k)$  can be equipped with the homotopy *t*-structure:  $\operatorname{SH}^{S^1}(k)_{\leq 0}$  consists of those spectra that are coconnective (i.e.  $\pi_i = 0$  for i > 0), and  $\operatorname{SH}^{S^1}(k)_{\geq 0}$  consists of those spectra that are connective (i.e.  $\pi_i = 0$  for i < 0) [Mor05, Lemma 6.2.11]. Equivalently, the category  $\operatorname{SH}^{S^1}(k)_{\geq 0}$  is generated under colimits and extensions by  $\Sigma^{\infty}X_+$ ,  $X \in \operatorname{Sm}_k$ . The abelian category  $\operatorname{SH}^{S^1}(k)^{\heartsuit}$  can be identified with the category of strictly  $\mathbb{A}^1$ -invariant sheaves [Mor05, Lemma 6.2.13].

2.2.16 (Homotopy *t*-structure on SH(*k*) and homotopy modules). The category SH(*k*) can be equipped with a homotopy *t*-structure [Mor04, Theorem 5.2.3]. In this case, the non-negative part of the *t*-structure is generated under colimits and extensions by  $\Sigma^{p,q}\Sigma^{\infty}_{\mathbb{P}^1}X_+$  with  $p - q \ge 0$  (see [Hoy15, §2.1]). The category SH(*k*)<sup> $\heartsuit$ </sup> can be described in terms of *homotopy modules* via [Mor04, Theorem 5.2.6]. Recall that a homotopy module consists of a  $\mathbb{Z}$ -graded strictly  $\mathbb{A}^1$ -invariant sheaf  $\mathbf{M}_*$  together with isomorphisms  $\mu_n : \mathbf{M}_n \simeq (\mathbf{M}_{n+1})_{-1}$  (see 2.1.10).

2.2.17. The homotopy module  $\pi_0(\mathbf{1}_k)$  has been determined by Morel [Mor04, Theorem 6.4.1] and [Mor12, Theorem 6.40]: it is the graded unramified Milnor–Witt K-theory sheaf  $\mathbf{K}^{MW}_*$ . Our conventions for Milnor–Witt K-theory follow Morel's book [Mor12, 3.1]. For an extension K/k, the sections of  $\pi_0(\mathbf{1}_k)(K)$  are generated by the motivic Hopf map  $\eta$  of degree -1, and symbols [u], for  $u \in K^{\times}$ , satisfying relations. For a unit u, we write  $\langle u \rangle = 1 + \eta[u]$ , and the map  $u \to \langle u \rangle$  factors through an isomorphism  $GW(K) \to \pi_0(\mathbf{1}_k)(K)_0$ . We write  $\rho$  for the symbol [-1], h for  $1 + \langle -1 \rangle$  and  $\varepsilon$  for  $-\langle -1 \rangle$ .

2.2.18 (Homotopy modules ctd.). Every homotopy module has the structure of a module over  $\pi_0(\mathbb{1}_k)$ . The category of homotopy modules also admits several equivalent descriptions that are quite useful. Using motivic infinite loop space recognition machinery [EHK<sup>+</sup>21, Theorem 3.5.14], the degree 0 part of a homotopy module can be described in terms of strictly  $\mathbb{A}^1$ -invariant sheaves admitting "framed transfers" [BY20, Theorem 5.14]. Alternatively, there is a "field-theoretic" description a la Voevodsky's theory of sheaves with transfers, the so-called category of sheaves with Milnor–Witt transfers [BCD<sup>+</sup>20, Chapter 2]. Finally there is an approach by Morel using so-called "generalized transfers" [Fel20]. In all cases, the sheaves  $\mathbf{M}_*$  are determined by their sections over finitely generated extensions of the base field together

with suitable residue and transfer morphisms that are compatibe with action of  $\pi_0(\mathbb{1}_k)$ . Moreover, all the descriptions above are equivalent [AN19, Theorem 1.3] or [Fel21, Corollary 6.2.4].

2.2.19. If  $E \in CAlg(SH(k))$ , then  $Mod_E(k)$  admits a homotopy *t*-structure whose non-negative part is generated by  $E \wedge \Sigma_{\mathbb{P}^1}^{\infty} X_+$  for  $X \in Sm_k$ . The heart of this *t*-structure has been described for various *E*. Of particular interest to us is the case  $E = H\mathbb{Z}[1/p]$ , i.e., Voevodsky's motivic Eilenberg–Mac Lane spectrum, in which case  $Mod_{H\mathbb{Z}}(k)^{\heartsuit}$  coincides with Voevodsky's category of homotopy modules of  $\mathbb{Z}[1/p]$ -modules with transfers. Indeed, Voevodsky showed that the heart of a corresponding *t*-structure on **DM**(*k*) could be described in terms of strictly  $\mathbb{A}^1$ -invariant Nisnevich sheaves with transfers [Voe00b, §3.1 p. 205] and then one may transfer this equivalence to module spectra in conjunction with the discussion of Paragraph 2.2.7.

2.2.20. The category  $\operatorname{SH}(k)^{\operatorname{eff}}$  also has a *t*-structure. Indeed, we define the non-negative part to be  $\operatorname{SH}(k)^{\operatorname{eff}} \cap \operatorname{SH}(k)_{\geq 0}$ . By [Bac17, Proposition 4] it follows that this yields a *t*-structure on  $\operatorname{SH}(k)^{\operatorname{eff}}$  that is called the effective homotopy *t*-structure;  $E \in \operatorname{SH}(k)^{\operatorname{eff}}_{\leq 0}$  if and only if  $\pi_i(E)_0 = 0$  for i > 0. If follows from the discussion after [Bac17, Proposition 4] that the non-negative part of the homotopy *t*-structure on  $\operatorname{SH}(k)^{\operatorname{eff}}$  is generated under colimits and extensions by  $\Sigma^{n,0}\Sigma_{\mathbb{P}^1}^{\infty}X_+$  and thus coincides with  $\operatorname{SH}(k)^{\operatorname{veff}}$  (see Paragraph 2.2.11); but we recapitulate the argument in the next result.

**Proposition 2.2.21.** The categories  $\operatorname{SH}^{S^1}(k)_{\geq 0}$ ,  $\operatorname{SH}(k)^{\operatorname{veff}}$  and  $\operatorname{Mod}_E(k)^{\operatorname{veff}}$  for any  $E \in \operatorname{CAlg}(\operatorname{SH}(k)^{\operatorname{veff}})$  are generated under sifted colimits by the image of  $\operatorname{Sm}_k$ .

*Proof.* We first show that the categories are generated by the objects claimed under colimits. The categories  $\operatorname{SH}^{S^1}(k)_{\geq 0}$  and  $\operatorname{SH}(k)^{\operatorname{veff}}$  are the positive parts of presentably stable  $\infty$ -categories with *t*-structure by the discussion of Paragraph 2.2.15 and 2.2.16. In these stable  $\infty$ -categories, the  $\infty$ -connective objects are zero (this follows from hypercompleteness of the Nisnevich  $\infty$ -topos). Moreover, for every positive object, we can always find a family of maps from shifted suspension spectra of smooth schemes that induce an epimorphism after taking  $\pi_0$  (see "Details on killing cells" [Bac17, p. 1130]); the result follows from the discussion there, though see [Lur18, Corollary C.2.1.7] for a more abstract statement. The case of  $\operatorname{Mod}_E(k)^{\operatorname{veff}}$  is similar and reduces to the case of  $\operatorname{SH}(k)^{\operatorname{veff}}$  since the forgetful functor  $\operatorname{Mod}_E(k)^{\heartsuit} \to \operatorname{SH}(k)^{\heartsuit}$  is exact and conservative [Bac21b, Lemma 29 and its proof].

Now we reduce to sifted colimits. We treat  $\operatorname{SH}^{S^1}(k)_{\geq 0}$  as an example. Let  $\mathcal{C} \subset \operatorname{SH}^{S^1}(k)_{\geq 0}$  denote the subcategory generated under sifted colimits by  $\operatorname{Sm}_k$ . We claim  $\mathcal{C}$  is closed under binary sums. To see this, first let  $X \in \operatorname{Sm}_k$  and write  $\mathcal{C}_X \subset \mathcal{C}$  for the subcategory of those objects Y such that  $\Sigma^{\infty}_+ X \oplus Y \in \mathcal{C}$ . Then  $\mathcal{C}_X$  contains  $\Sigma^{\infty}_+ \operatorname{Sm}_k$  and is closed under sifted colimits (use that sifted categories are weakly contractible [Lur09, Proposition 5.5.8.7]), hence  $\mathcal{C}_X = \mathcal{C}$ . New let  $X \in \mathcal{C}$  be arbitrary. Then  $\mathcal{C}_X$  contains  $\Sigma^{\infty}_+ \operatorname{Sm}_k$  by what we just did, and is closed under sifted colimits as before; hence  $\mathcal{C}_X = \mathcal{C}$ . We have thus proved that  $\mathcal{C}$  is closed under finite sums and sifted colimits, whence all colimits [Lur09, Lemma 5.5.8.13].

#### **Decomposing** $SH(k)_{\mathbb{O}}$

2.2.22. Recall that there exists an endomorphism  $\varepsilon$  of the motivic sphere spectrum  $\mathbb{1}_k \in SH(k)$  such that  $\varepsilon^2 = id$ :  $\varepsilon$  corresponds to permutation of the two factors in  $\mathbb{G}_m \wedge \mathbb{G}_m$ . Morel computed that  $\varepsilon$  corresponds to the element  $-\langle -1 \rangle \in GW(k)$  under the identification of  $\pi_0(\mathbb{1})$  with  $\mathbf{K}_0^{MW}(k)$  from Paragraph 2.2.17. With that notation, the class of the hyperbolic form h is  $1 - \varepsilon$ , and the relation  $\eta h = 0$  in Milnor–Witt K-theory implies that  $\eta$  is killed by  $1 - \varepsilon$ . After inverting 2, we obtain a pair of orthogonal idempotents  $e_{\pm} := \frac{1}{2}(1 \mp \varepsilon)$  that decompose  $SH(k)_{\mathbb{Z}[1/2]}$  as a sum of eigen-subcategories. The image of  $e_+$ , i.e., the -1-eigenspace of  $\varepsilon$  is defined to be  $SH(k)_{\mathbb{Z}[1/2]}^+$ , while the image of  $e_-$  is defined to be  $SH(k)_{\mathbb{Z}[1/2]}^-$ , and there is a decomposition:

$$\operatorname{SH}(k)_{\mathbb{Z}[1/2]} \simeq \operatorname{SH}(k)^+_{\mathbb{Z}[1/2]} \times \operatorname{SH}(k)^-_{\mathbb{Z}[1/2]};$$

(this unfortunate terminological issue was caused by a change in conventions as regards  $\varepsilon$ .) If k has finite 2-cohomological dimension, then the --part vanishes [Lev13, Lemma 6.8]. After rationalizing, the +-part admits a description originally due to F. Morel.

# **Theorem 2.2.23** (Morel). If k is a field, then $\operatorname{SH}(k)^+_{\mathbb{O}} \simeq \operatorname{Mod}_{H\mathbb{Q}}(k)$ .

*Proof.* This result is obtained by composing a number of equivalences. First, the usual Dold–Kan adjunction defines an equivalence between  $\operatorname{SH}^{S^1}(k)_{\mathbb{Q}}$  and  $\operatorname{D}^{\mathbb{A}^1}(k)_{\mathbb{Q}}$  where the latter is Morel's  $\mathbb{A}^1$ -derived category [CD19, §5.3.35]. After inverting the Tate object, there is an induced equivalence  $\operatorname{SH}(k)_{\mathbb{Q}} \simeq D^{\mathbb{A}^1}(k)_{\mathbb{Q}}[\mathbb{G}_m^{-1}]$ . By [CD19, Theorem 16.2.13] we know that  $D^{\mathbb{A}^1}(k)_{\mathbb{Q}}[\mathbb{G}_m^{-1}]^+$  is the category of "Beilinson" motives [CD19, §14]. Finally, under the assumptions on k, we conclude that the category of "Beilinson" motives coincides with Voevodsky's rationalized derived category  $\operatorname{DM}(k, \mathbb{Q})$  [CD19, Theorem 16.1.14]. The equivalence in the statement then follows from the discussion of Paragraph 2.2.7.

# $S^1$ -spectra and monoids

Corollary 2.2.24. Let k be a perfect field. Taking infinite loops induces a fully faithful functor

$$\mathrm{SH}^{S^1}(k)_{\geq 0} \to \mathrm{CMon}(\mathrm{Spc}(k))$$

with image those grouplike commutative monoids such that  $\pi_0$  is strongly  $\mathbb{A}^1$ -invariant. This subcategory is closed under colimits in  $\mathrm{CMon}(\mathrm{Spc}(k))$ . In particular  $\mathrm{SH}^{S^1}(k)_{\geq 0} \to \mathrm{CMon}(\mathrm{Spc}(k))$  preserves colimits.

*Proof.* The fully faithfulness statement is [EHK<sup>+</sup>21, Proposition 3.1.13 and Corollary 3.1.15]; there the characterization of the image is also proved. For closure under colimits, it suffices to prove closure under finite coproducts, filtered colimits and geometric realizations [Lur09, Lemma 5.5.8.13]. We must show that such colimits in CMon(Spc(k)) preserve the subcategory of monoids with  $\pi_0$  grouplike and strongly  $\mathbb{A}^1$ -invariant. The case of finite coproducts (sums) is clear, and the same holds for filtered colimits. Geometric realizations are dealt with in Corollary 2.1.14.

**Corollary 2.2.25.** Let k be a perfect field. Denote the localization from Corollary 2.2.24 by

$$L: \operatorname{CMon}(\operatorname{Spc}(k)) \leftrightarrows \operatorname{SH}^{S^1}(k)_{\geq 0}: R.$$

*Consider a pushout square in* CMon(Spc(k))



Suppose that  $\pi_0(B), \pi_0(C)$  are grouplike and strongly  $\mathbb{A}^1$ -invariant. Then

$$D \simeq R(LC \amalg_{LA} LB).$$

*Proof.* Since R preserves colimits by Corollary 2.2.24, the assertion is just that  $\pi_0(D)$  is grouplike and strongly  $\mathbb{A}^1$ -invariant. Writing the pushout as a geometric realization, Corollary 2.1.14 reduces this to showing that  $\pi_0(C \oplus D)$  is grouplike and strongly  $\mathbb{A}^1$ -invariant, which is clear.

#### Conservativity of $\mathbb{G}_m$ -stabilization

2.2.26. Consider the  $\mathbb{G}_m$ -stabilization functor

$$\sigma^{\infty}: \mathrm{SH}^{S^1}(k) \longrightarrow \mathrm{SH}(k)$$

(see Paragraph 2.2.3). By construction, this functor sends the  $S^1$ -suspension spectrum of a smooth scheme X to its corresponding  $\mathbb{P}^1$ -suspension spectrum; it therefore sends d-effective  $S^1$ -spectra to d-effective  $\mathbb{P}^1$ -spectra and also sends  $\mathrm{SH}^{S^1}(k)_{\geq 0}$  to  $\mathrm{SH}(k)^{\mathrm{veff}}$  (see [BY20, §6.1]).

2.2.27. There is an induced functor

$$\sigma^{\infty}: \mathrm{SH}^{S^1}(k)^{\heartsuit} \longrightarrow \mathrm{SH}(k)^{\heartsuit}.$$

In order for a strictly  $\mathbb{A}^1$ -invariant sheaf  $\mathbf{A}$  to support the structure of a homotopy module (see Paragraphs 2.2.16 and 2.2.18), two things must happen. First, any element of  $\mathrm{SH}(k)^{\heartsuit}$  is a module over  $\pi_0(\mathbb{1}_k)$ . Second, we must be able to find compatible  $\mathbb{G}_m$ -deloopings, i.e., we must be able to find a sequence of strictly  $\mathbb{A}^1$ -invariant sheaves  $\mathbf{A}_i$ ,  $i \ge 0$  such that  $\mathbf{A}_0 = \mathbf{A}$  and  $(\mathbf{A}_{i+1})_{-1} = \mathbf{A}_i$ ; in particular, the sheaf  $\mathbf{A}$ must itself be a contraction. Moreover, the  $\mathbf{K}_{-*}^{MW}$ -module structure and the  $\mathbb{G}_m$ -deloopings must necessarily be compatible.

*Example* 2.2.28. The functor  $\sigma^{\infty}$  is not an equivalence in general. Indeed, if *C* is a smooth projective curve of genus g > 0 over a field *k*, then the zeroth  $\mathbb{A}^1$ -homology sheaf  $\mathbf{H}_0^{\mathbb{A}^1}(C)$  is known to be the free sheaf of abelian groups on *C* (see [AH11, Example 2.6]). A modification of the argument in [Lev10, §2] shows that  $\mathbf{H}_0^{\mathbb{A}^1}(C)$  does not underlie a homotopy module.

Morel asked whether any strictly  $\mathbb{A}^1$ -invariant sheaf that is a 1-fold contraction should underlie a homotopy module (see [Mor11, Remark 5.10]). Variants of this question were pursued by the second author, partly in collaboration with M. Yakerson. The following result summarizes the results we will use: loosely speaking,  $S^1$ -spectra that are "sufficiently effective" can be infinitely  $\mathbb{G}_m$ -delooped.

**Theorem 2.2.29.** Assume k is a perfect field. There exists a minimal integer  $\epsilon(k) \ge 0$  such that the following statements hold.

1. For  $E \in SH^{S^1}(k)_{\geq 0} \cap SH^{S^1}(k)(\epsilon(k))$ , the canonical map

$$\pi_0(E) \longrightarrow \pi_0(\omega^{\infty} \sigma^{\infty} E)$$

is an isomorphism.

2. The functor

$$\omega^{\infty} : \mathrm{SH}(k)^{\mathrm{eff}}(\epsilon(k))^{\heartsuit} \longrightarrow \mathrm{SH}^{S^{1}}(k)(\epsilon(k))$$

is an equivalence of categories. In fact  $\epsilon(k) \leq 2$ .

*Proof.* For any strictly  $\mathbb{A}^1$ -invariant sheaf, F. Morel constructed "canonical" or "Bass–Tate" transfers on  $\mathbb{M}_{-2}$  (see [Mor12, p. 99] or [Bac21a, §2.1.8]). If these "canonical transfers" give rise to framed transfers, then the result is the conclusion of [Bac21a, Corollary 4.9]. If *k* has characteristic 0, the condition that Morel's canonical transfers on 2-fold contractions give rise to framed transfers was established by [BY20, Theorem 5.19]; if *k* has positive characteristic, then the same statement follows from [Fel20, Theorem 6.1.5(iii,iv)]. The second statement is simply [Bac21a, Theorem 4.6] in light of the observations just made.

#### 2.3 Real-étale motivic homotopy theory

Here, we prove unstable analogs of some results from [Bac18a]. Much stronger results will be established in forthcoming work. We write rét for the real-étale topology of Scheiderer [Sch94, Definition 1.2.1]. Along the lines of Paragraph 2.1.2 we write  $\operatorname{Spc}_{rét}(k)$  for the  $\infty$ -category of real étale motivic spaces. We write  $L_{rét}$  for the corresponding localization functor to distinguish it from  $L_{mot}$ .

#### Unstable real-étale localization

**Proposition 2.3.1.** Let k be a perfect field and  $\mathscr{X} \in \text{Shv}_{Nis}(\text{Sm}_k)$  be connected. Assume that for  $i \leq n$ ,  $\pi_i \mathscr{X}$  is a strictly  $\mathbb{A}^1$ -invariant real étale sheaf. Then for  $i \leq n$ ,  $\pi_i L_{r\acute{e}t} \mathscr{X} \simeq \pi_i \mathscr{X}$ . Moreover all homotopy sheaves of  $L_{r\acute{e}t} \mathscr{X}$  are real étale sheaves.

In particular if  $\mathscr{X}$  is *n*-connective, so is  $L_{rét}\mathscr{X}$ .

*Proof.* Write  $L_{rét}^{naive}$  :  $Shv_{Nis}(Sm_k) \to Shv_{Nis}(Sm_k)$  for the real étale sheafification functor. We can obtain  $L_{rét}$  as a countably infinite composition of  $L_{mot}$  and  $L_{rét}^{naive}$ . It thus suffices to prove that if  $\mathscr{X} \in Shv_{Nis}(Sm_k)$  is connected such that  $\pi_i \mathscr{X}$  is both strongly  $\mathbb{A}^1$ -invariant and a real étale sheaf for  $i \leq n$ , then the maps  $\mathscr{X} \to L_{rét}^{naive} \mathscr{X}$  and  $\mathscr{X} \to L_{mot} \mathscr{X}$  both induce isomorphisms on  $\pi_i$  for  $i \leq n$ . Both functors preserve the fiber sequence  $\mathscr{X}_{>n} \to \mathscr{X} \to \mathscr{X}_{\leq n}$  as well as *n*-connected spaces (for  $L_{mot}$  this is by unstable connectivity). It hence suffices to show that  $L_{mot} \mathscr{X}_{\leq n} \simeq \mathscr{X}_{\leq n} \simeq L_{rét}^{naive} \mathscr{X}_{\leq n}$ ; this is clear for  $L_{mot}$ . For  $L_{rét}^{naive}$ , by an induction on the Postnikov tower, it suffices to prove that if **A** is a real étale sheaf of abelian groups, then  $K_{Nis}(\mathbf{A}, i)$  satisfies real étale descent. This follows from the fact that  $H_{rét}^*(-, \mathbf{A}) \simeq H_{Nis}^*(-, \mathbf{A})$ , which itself is a consequence of [Sch94, Proposition 19.2.1]. This also implies that the (Nisnevich) homotopy sheaves of  $L_{rét} \mathscr{X}$  are real étale sheaves. □

**Proposition 2.3.2.** Let k be a perfect field. For  $\mathscr{X} \in \operatorname{Spc}(k)_*$  simply connected, the following are equivalent: (1)  $\mathscr{X}$  satisfies real étale descent. (2)  $\rho : \mathscr{X} \to \mathscr{X} \wedge \mathbb{G}_m$  is an equivalence.

Moreover then  $\rho^* : \mathscr{X} \to \Omega_{\mathbb{G}_m} \mathscr{X}$  is an equivalence.

Proof. Put

$$\mathscr{X}[\rho^{-1}] = \operatorname{colim}(\mathscr{X} \xrightarrow{\rho} \mathscr{X} \wedge \mathbb{G}_m \xrightarrow{\rho} \mathscr{X} \wedge \mathbb{G}_m^{\wedge 2} \xrightarrow{\rho} \dots)$$

We first check that  $L_{rét} \mathscr{X} \to L_{rét} (\mathscr{X}[\rho^{-1}])$  is an equivalence. Since both spaces are simply connected by Proposition 2.3.1, it suffices to check the claim after mapping into a simply connected rét-local space. By Proposition 2.3.1, these admit a Postnikov tower with layers consisting of Eilenberg–Mac Lane spaces for real étale sheaves. This reduces to checking that  $\rho$  becomes an equivalence in  $L_{rét}SH^{S^1}(k)$ , which is proved in [?, Theorem 4.2].

The same argument also proves that if  $\mathscr{X}$  is rét-local, then  $\rho^*$  is an equivalence on  $\mathscr{X}$ : we reduce to the Postnikov layers, which come from  $L_{rét}SH^{S^1}(k)$ , where the claim holds.

Next we check that  $\rho : \mathscr{X}[\rho^{-1}] \to \mathscr{X}[\rho^{-1}] \land \mathbb{G}_m$  is an equivalence. Appeal to Theorem 3.2.12 reduces to the same statement in SH(k), which is clear.

Now let  $\mathscr{X}$  be rét-local (and simply connected) and consider the sequence of spaces

$$\mathscr{X} \to \mathscr{X}[\rho^{-1}] \to \mathcal{L}_{\text{rét}} \mathscr{X}[\rho^{-1}] \simeq \mathscr{X} \to \mathscr{X}[\rho^{-1}] \to \cdots$$

Considering the subsequence of spaces of the form  $\mathscr{X}[\rho^{-1}]$  shows that the colimit satisfies (2); on the other hand the subsequence of spaces of the form  $\mathscr{X}$  is along identity maps, and so the colimit is  $\mathscr{X}$ . Hence (1) implies (2).

Now suppose that  $\mathscr{X}$  satisfies (2). Suppose that  $\pi_i \mathscr{X}$  is a real étale sheaf, for  $i \leq n$  (e.g. n = 1). Denote by  $\mathscr{F}, \mathscr{C}$  the fiber and cofiber of  $\mathscr{X} \to \mathscr{X}_{\leq n}$ . It follows from Proposition 2.3.1 that  $\mathscr{X}_{\leq n}$  satisfies real étale descent, and hence satisfies (2) by what we already did. Consequently  $\mathscr{C}$  satisfies (2). We know that [ABH22, Proposition 3.5(2)]

$$\pi_{n+1}\mathscr{F}\simeq\pi_{n+2}\Sigma\mathscr{F}\simeq\pi_{n+2}\mathscr{C}.$$

On the other hand  $\pi_{n+2}\mathscr{C} \simeq \pi_{n+2}\Sigma^{\infty}\mathscr{C}$  by [Bac21a, Corollary 4.9], which is a real étale sheaf by [Bac18a, Proposition 41]. We have thus proved that all homotopy sheaves of  $\mathscr{X}$  are real étale sheaves, and so  $\mathscr{X} \simeq L_{\text{rét}}\mathscr{X}$  by Proposition 2.3.1.

### The rationalized - part

Morel initially proposed a description of the rationalized –-part complementing the description of Theorem 2.2.23 in terms of modules over the Witt sheaf. Ananyevskiy–Levine–Panin established Morel's description [ALP17, Theorem 4.2]. We will need the following alternative description of the rationalized –-part due to the second author.

**Theorem 2.3.3** ([Bac18a, Proposition 41]). There is an equivalence  $SH(k)_{\mathbb{Q}}^- \simeq D((Spec k)_{r\acute{e}t}, \mathbb{Q})$ , in particular, the minus part of a rationalized motivic spectrum has real-étale descent.

#### **Real realizations**

Let k be a field and  $\sigma$  be an ordering of k. Write  $k_{\sigma}$  for the induced real closure. Recall that  $SH(k_{\sigma})[\rho^{-1}] \simeq$ SH [Bac18a, Theorem 35].

Definition 2.3.4. We call the composite functor

$$r_{\sigma} : \mathrm{SH}(k) \to \mathrm{SH}(k_{\sigma})[\rho^{-1}] \simeq \mathrm{SH}$$

the real realization associated with  $\sigma$ .

# **3** Weak cellularity and nullity

The classical Postnikov tower (say in the category of simplicial sets) can be effected by localization with respect to the maps  $S^n \to *$  (or  $* \to S^n$ ), i.e., nullification [DF96, Example A.6]. More generally, one can build the Postnikov tower in any  $\infty$ -topos by means of nullifications [ABFJ20, Lemma 3.3.3].

In motivic homotopy theory, for  $p \ge q$  there is a bi-graded family of spheres  $S^{p,q} := S^{p-q} \wedge \mathbb{G}_m^{\wedge q}$ . Analogous to the classical situation we will consider nullifications of these spheres. While all maps  $S^i \to S^j$  are null-homotopic if i < j, this is no longer the case for  $\mathbb{G}_m^{\wedge i} \to \mathbb{G}_m^{\wedge j}$ . As such, one cannot expect that the nullifications with respect to bi-graded spheres will behave in exactly the same way as the classical Postnikov truncations, e.g., classically, the homotopy fiber of the map to the *n*-th Postnikov section is at least *n*-connected and it is not a priori clear that a corresponding property holds for maps to a nullification at  $S^{p,q}$  (this is one of the reasons why we eschew "connectivity" terminology); however see Theorem 4.1.9.

The relevant nullification functors may be defined stably and unstably. We will use the terminology weakly  $S^{p,q}$ -cellular for spaces that are killed by nullification. If  $\mathscr{X}$  is weakly  $S^{p,q}$ -cellular, then we establish a refinement of the Whitehead and Postnikov towers for  $\mathscr{X}$  in which all the layers are suitably cellular. We then use this to establish cellularity estimates for  $\Omega^{1,0}\mathscr{X}$  and  $\Omega^{1,1}\mathscr{X}$ . To do this, requires comparison of stable and unstable nullifications and relies on results of Levine and the conservativity of  $\mathbb{G}_m$ -stabilization results from Paragraphs 2.2.26–2.2.29.

#### 3.1 Weak cellularity and nullifications: basic properties

In this section, we study nullifications at a *pointed*, *compact*, *motivic space* A; in practice, A will be a motivic sphere  $S^{p,q}$ . We then introduce the class O(A) of weakly A-cellular spaces. By definition; these are spaces that admit no non-trivial maps to A-null spaces.

#### **Basic definitions and constructions**

3.1.1 (A-equivalences). Assume A is a pointed, compact, motivic space. Consider the set of morphisms  $\{A \times U \rightarrow U\}, U \in Sm_k$ ; we will call this set of morphisms in Spc(k) the generating A-equivalences. The left Bousfield localization at (the smallest strongly saturated class of Spc(k) generated by) the set of generating A-equivalences exists and is again a presentable  $\infty$ -category. We write

$$L_A: \operatorname{Spc}(k) \longrightarrow \operatorname{Spc}(k)$$

for the corresponding localization endofunctor.

3.1.2. If C is a presentable  $\infty$ -category equipped with a functor  $\operatorname{Sm}_k \to C$ , we use similar notation for the localization of  $\mathscr{C}$  generated by the image of the above maps. We have in mind  $C = \operatorname{Spc}(k)_*, \operatorname{SH}^{S^1}(k), \operatorname{SH}^{S^1}(k)_{>0}, \operatorname{SH}(k), \operatorname{SH}(k)^{\operatorname{veff}}$ .

As usual, the localization functors identify the left Bousfield localization of  $\operatorname{Spc}(k)$  with the subcategory consisting of local objects. The A-local spaces may be characterized in terms of the internal mapping objects: a space  $\mathscr{X}$  is A-local if and only if the map  $\mathscr{X} \simeq \operatorname{Map}(*, \mathscr{X}) \to \operatorname{Map}(A, \mathscr{X})$  is an equivalence. Likewise, A-equivalences can be detected by mapping into A-local spaces, i.e., a map  $f : \mathscr{X} \to \mathscr{Y}$  is an A-equivalence if and only if for every A-local space  $\mathscr{W}$ , the map  $\operatorname{Map}(\mathscr{Y}, \mathscr{W}) \to \operatorname{Map}(\mathscr{X}, \mathscr{W})$  is an equivalence. In the case of interest, we make the following definition.

**Definition 3.1.3.** Assume A is a pointed, compact motivic space. Suppose  $\mathscr{X}, \mathscr{Y}$  are motivic spaces and  $f : \mathscr{X} \to \mathscr{Y}$  is a map.

- 1. The space  $\mathscr{X}$  is A-null if  $\mathscr{X} \to L_A \mathscr{X}$  is an equivalence (i.e.,  $\mathscr{X}$  is A-local).
- 2. The space  $\mathscr{X}$  is weakly A-cellular and we write  $\mathscr{X} \in O(A)$ , if  $L_A \mathscr{X} = *$ ; equivalently the map  $\mathscr{W} \to \operatorname{Map}(\mathscr{X}, \mathscr{W})$  is an equivalence for every A-null space  $\mathscr{W}$ .
- 3. The map f is a A-equivalence if  $L_A(f)$  is an equivalence.

*Remark* 3.1.4. Of particular interest to us will be the case  $A = S^{p,q}$ , in which case we will write  $L^{p,q}$  for  $L_A$ . We will sometimes refer to  $S^{q,q}$ -null spaces as  $\mathbb{G}_m$ -q-truncated spaces, and spaces in  $O(S^{q,q})$  as (unstably) q-effective spaces, etc. We refer the reader to Section 3.3 for a number of examples that help to give a sense of how we will use these notions.

*Example* 3.1.5. In the case  $A = S_k^0$ , the generating A-equivalences are the maps  $S_k^0 \times X = X \sqcup X \to X$ . In any  $\infty$ -topos, the corresponding localization is -1-truncation [ABFJ20, Lemma 3.3.3]. Since any -1-truncated sheaf is  $\mathbb{A}^1$ -invariant, it follows that  $\mathbb{L}^{0,0} = \mathbb{L}_A = \tau_{<-1}^{\text{Nis}}$ .

We now establish basic properties of these localization functors; our treatment follows that of [DF96].

#### Weak cellularity, first properties

We record the compatiblity of localization with colimits in the following result.

**Lemma 3.1.6.** The localizations  $L_A$ , viewed as functors from motivic spaces to A-null motivic spaces preserve colimits. In particular, given a cofiber sequence,

$$\mathscr{X} \xrightarrow{f} \mathscr{Y} \xrightarrow{g} \mathscr{Z},$$

the following statements hold.

- 1. If  $\mathscr{X}$  and  $\mathscr{Z}$  are in O(A), then so is  $\mathscr{Y}$ .
- 2. If  $\mathscr{X} \in O(A)$ , then g is an A-equivalence, and
- 3. If f is a A-equivalence, then  $\mathscr{Z} \in O(A)$ .

*Proof.* All these statements follow from the fact that  $L_A$  is a homotopical left adjoint.

**Amplification 3.1.7.** If C is a presentable infinity category under  $Sm_k$  (see Paragraph 3.1.2), then the conclusions of Lemma 3.1.6 hold in C.

*Proof.* The left Bousfield localization  $L_A$  of C is still a homotopical left adjoint.

**Lemma 3.1.8.** The functor  $L_A$  preserves finite products.

*Proof.* We need only check that if f is a generating A-equivalence and  $X \in \text{Sm}_k$  then  $f \times \text{id}_X$  is an A-equivalence [Lur09, Proposition 2.2.1.9], which is clear.

We amplify this point. Write CMon(Spc(k)) for the category of commutative monoid objects on Spc(k). This category admits forgetful functors to  $Spc(k)_*$  and Spc(k). Applying 3.1.2 to the left adjoints, we transport  $L_A$  to these categories as well.

Amplification 3.1.9. The forgetful functors

$$\operatorname{CMon}(\operatorname{Spc}(k)) \longrightarrow \operatorname{Spc}(k)_* \longrightarrow \operatorname{Spc}(k)$$

*commute with*  $L_A$ .

*Proof.* By Lemma 3.1.8 we know that  $L_A$  is compatible with the symmetric monoidal structure on Spc(k). By definition the commutative monoid objects in Spc(k) are algebras over an  $\infty$ -operad, and the same holds for the pointed objects. The result then follows from [Lur17, Proposition 2.2.1.9] and [Lur17, Remark 7.3.2.13].

Before deducing some further consequences about A-null spaces from the above result, we amplify some of the consequences about A-localizations preserving symmetric monoidal structures.

**Amplification 3.1.10.** Suppose C is a presentably symmetric monoidal  $\infty$ -category under  $Sm_k$  such that

- 1. the functor  $Sm_k \rightarrow C$  is symmetric monoidal, and
- 2. the functor  $Sm_k \to C$  has dense image, i.e., the category C is generated under colimits by the image of the source.

*The localization functor*  $L_A$  *on* C *is monoidal.* 

*Proof.* The condition that the functor  $\operatorname{Sm}_k \to \mathcal{C}$  is symmetric monoidal guarantees that if  $\alpha$  is a generating equivalence, then  $\alpha \otimes \operatorname{id}_X$  is as well for any  $X \in \operatorname{Sm}_k$ . Combined with the density condition, this implies that  $\operatorname{L}_A : \mathscr{C} \to \operatorname{L}_A \mathscr{C}$  is symmetric monoidal [Lur09, Proposition 2.2.1.9].

**Corollary 3.1.11.** The conclusion of Amplification 3.1.10 holds for  $C = \text{Spc}(k)_*$ ,  $\text{SH}^{S^1}(k)_{\geq 0}$ ,  $\text{SH}(k)^{\text{veff}}$  or  $\text{Mod}_{H\mathbb{Z}}^{\text{veff}}$ .

*Proof.* Immediate by combining Amplification 3.1.10 with Proposition 2.2.21.

*Example* 3.1.12. If  $\mathscr{X} \in \text{Spc}(k)$ , then  $A \wedge \mathscr{X}_+$  is weakly A-cellular. Indeed by Corollary 3.1.11, it suffices to prove that A is A-cellular as a pointed space; this follows from Amplification 3.1.9.

Let us also mention the following.

**Lemma 3.1.13.** The functor  $L_A$  commutes with essentially smooth change of base field. In particular, if  $f : \operatorname{Spec}(k') \to \operatorname{Spec}(k)$  is an essentially smooth field extension, then  $f^*$  sends A-equivalences to  $f^*A$ -equivalences, O(A) to  $O(f^*A)$  and A-null spaces to  $f^*A$ -null spaces.

*Proof.* Since  $f^*$  preserves colimits and finite products, it sends A-equivalences to  $f^*A$ -equivalences. It hence remains to prove that it sends A-null spaces to  $f^*A$ -null spaces. Since A is compact, this follows immediately from essentially smooth base change (see Paragraph 2.1.5).

#### Null spaces and limits

Just as weakly cellular objects and weakly cellular maps interact well with colimits, local objects interact well with limits. We record the necessary facts in the following result (these facts are standard and amount to notational modifications of [DF96, 1.A.8])

#### Lemma 3.1.14. The following statements hold.

- 1. If  $\mathscr{X}$  is A-null, then for any space  $\mathscr{Y}$ , the space  $\operatorname{Map}(\mathscr{Y}, \mathscr{X})$  is again A-null.
- 2. Any limit of A-null spaces is again A-null.
- 3. If  $\mathscr{F} \to \mathscr{E} \to \mathscr{B}$  is a fiber sequence with  $\mathscr{F}$  and  $\mathscr{B}$  A-null spaces, and  $\mathscr{B}$  connected, then  $\mathscr{E}$  is A-null as well.
- 4. If  $\mathscr{X}$  is a pointed connected space, then  $\mathscr{X}$  is A-null if and only if Map  $(A, \mathscr{X}) = *$ .
- 5. If  $\mathscr{X}$  is a pointed, simply connected space, then  $\mathscr{X}$  is  $\Sigma A$ -null if and only if  $\Omega \mathscr{X}$  is A-null.

*Proof.* Point (1). To say that  $\mathscr{X}$  is A-null is equivalent to saying that the induced map

$$\mathscr{X} = \underline{\operatorname{Map}}(*, \mathscr{X}) \to \underline{\operatorname{Map}}(A, \mathscr{X})$$

is an equivalence. Mapping  $\mathscr{Y}$  into this equivalence and using the exponential law shows that

$$\underline{\mathrm{Map}}(\mathscr{Y},\mathscr{X}) \to \underline{\mathrm{Map}}(\mathscr{Y},\underline{\mathrm{Map}}(A,\mathscr{X})) \simeq \underline{\mathrm{Map}}(A,\underline{\mathrm{Map}}(\mathscr{Y},\mathscr{X}))$$

is an equivalence as well, which is what we wanted to show.

*Point* (2). Let I be a small diagram and let  $\mathscr{X} : \mathbf{I} \to \operatorname{Spc}(k)$ , be an I-diagram of motivic spaces. There is an equivalence

$$\operatorname{Map}(-, \lim_{\mathbf{I}} \mathscr{X}) \simeq \lim_{\mathbf{I}} \operatorname{Map}(-, \mathscr{X})$$

by the definition of mapping spaces. Now, for any  $i \in \mathbf{I}$ , the map  $\mathscr{X} \to \underline{\mathrm{Map}}(A, \mathscr{X})$  is an equivalence by assumption and we conclude that the induced map

$$\lim_{\mathbf{I}} \mathscr{X} \longrightarrow \lim_{\mathbf{I}} \underline{\operatorname{Map}}(A, \mathscr{X})$$


is an equivalence as well.

*Point (3).* Suppose  $\mathscr{F} \to \mathscr{E} \to \mathscr{B}$  is a fiber sequence with  $\mathscr{B}$  connected. In that case, we have the following morphism of fiber sequences

$$\begin{array}{c} \mathscr{F} & \longrightarrow \mathscr{E} & \longrightarrow \mathscr{B} \\ & \downarrow & & \downarrow \\ & & \downarrow & & \downarrow \\ \underline{\operatorname{Map}}(A, \mathscr{F}) & \longrightarrow \underline{\operatorname{Map}}(A, \mathscr{E}) & \longrightarrow \underline{\operatorname{Map}}(A, \mathscr{B}) \end{array}$$

where the outer two maps are equivalences and  $\mathscr{B}$  is connected. It follows that the middle map is an equivalence [Lur09, Lemma 6.2.3.16].

*Point (4).* We know that  $A \to *$  is an A-equivalence by Example 3.1.12. If  $\mathscr{X}$  is pointed and connected and A-null, it follows immediately that  $\underline{\operatorname{Map}}_*(A, \mathscr{X})$  is \*. Conversely, suppose  $\underline{\operatorname{Map}}_*(A, \mathscr{X}) = *$  and  $\mathscr{X}$  is connected. There is a fiber square of the form



linking pointed and free mapping spaces. Since  $\mathscr{X}$  is assumed connected,  $* \to \mathscr{X}$  is an effective epimorphism. Since  $\operatorname{Spc}(k) \subset \operatorname{Shv}_{\operatorname{Nis}}(\operatorname{Sm}_k)$  is closed under limits, [Lur09, Lemma 6.2.3.16] implies that the evaluation map ev in the above diagram is an equivalence. By 2-out-of-3, it follows that also  $\operatorname{Map}(A, \mathscr{X}) \to \operatorname{Map}(*, \mathscr{X})$  is an equivalence, as needed.

Point (5). This statement follows by adjunction of loops and suspension from the preceding point:  $\mathscr{X}$  is  $\Sigma A$ -null if and only if  $* = \operatorname{Map}_*(\Sigma A, \mathscr{X}) \simeq \operatorname{Map}_*(A, \Omega \mathscr{X})$ .

*Remark* 3.1.15. Let  $X \in \text{Sm}_k$  and  $\mathscr{X} \in \text{Spc}(k)_*$ . Note that  $\pi_0 \underline{\text{Map}}_*(A, \mathscr{X})(X) \simeq [A \wedge X_+, \mathscr{X}]_*$  and  $\pi_i(\underline{\text{Map}}_*(A, \mathscr{X})(X), 0) \simeq [\Sigma^i A \wedge X_+, \mathscr{X}]_*$ . In particular, by Lemma 3.1.14(4), if  $\mathscr{X}$  is connected then  $\mathscr{X}$  is *A*-null if and only if

$$[\Sigma^i A \wedge X_+, \mathscr{X}]_* = 0$$

for all  $i \ge 0$  and  $X \in \text{Sm}_k$ .

#### **Construction of the cellularization**

One may construct  $L_A$  explicitly by means of a small object argument; we summarize this observation in the next construction.

**Construction 3.1.16.** Let  $\mathscr{X} \in \operatorname{Spc}(k)_*$  be connected. Define spaces  $\mathscr{X}_n$  inductively as follows. Set  $\mathscr{X}_0 = \mathscr{X}$ . Assuming  $\mathscr{X}_n$  is defined, we define  $\mathscr{X}_{n+1}$  as the pushout of the diagram

$$* \longleftarrow \bigvee_{\{f: \Sigma^i A \land Y_+ \to \mathscr{X}_n\}} \Sigma^i A \land Y_+ \longrightarrow \mathscr{X}_n,$$

where  $Y \in \text{Sm}_k$  and *i* is an integer  $\geq 0$ . We then set  $\mathscr{X}_{\infty} = \text{colim}_n \mathscr{X}_n$ . The space  $\mathscr{X}_{\infty}$  is a model for  $L_A \mathscr{X}$ .

*Proof.* First note that each  $\mathscr{X}_i$  is connected (see Remark 3.1.17), and hence so is  $\mathscr{X}_\infty$ . Since  $\Sigma^i A \wedge Y_+ \stackrel{A}{\simeq} *$  we know that  $\mathscr{X} \to \mathscr{X}_\infty$  is an A-equivalence; hence it remains to verify that  $\mathscr{X}_\infty$  is A-null. By Remark 3.1.15, for this we need to check that for every  $Y \in \mathrm{Sm}_k$  and  $i \ge 0$ , every map  $\Sigma^i A \wedge Y_+ \to \mathscr{X}_\infty$  is null homotopic. Since the source is compact in  $\mathrm{Spc}(k)$ , this holds by construction.

*Remark* 3.1.17. Suppose that A is n-connective and  $\mathscr{X}$  is n-connected. One deduces from [Lur09, Corollary 6.5.1.17] (stability of n-connective morphisms under pushouts) and Theorem 2.1.6 (unstable connectivity) that  $\mathscr{X}_{\infty}$  is also n-connected (for n > 0, the reference requires k perfect).

An immediate consequence is the following.

**Lemma 3.1.18.** If  $\mathscr{X}$  is pointed and connected then  $\mathscr{X}_{\leq p-q-1} \simeq (L^{p,q}\mathscr{X})_{\leq p-q-1}$ . In particular if additionally  $\mathscr{X} \in O(S^{p,q})$  then  $\mathscr{X}$  is (p-q-1)-connected.

*Proof.* The space  $S^{p,q}$  is (p-q) - 1-connected. It follows that  $\mathscr{X}_n \to \mathscr{X}_{n+1}$  induces an equivalence on (p-q) - 1-truncations. The same is true for the colimit  $\mathscr{X} \to \mathscr{X}_{\infty} \simeq L^{p,q} \mathscr{X}$ .

We can deduce the following.

**Theorem 3.1.19.** If  $\mathscr{X}$  is pointed, connected and weakly A-cellular, and  $\mathscr{Y}$  is pointed, connected and weakly-B-cellular, t

hen  $\mathscr{X} \wedge \mathscr{Y}$  is weakly  $A \wedge B$ -cellular.

*Proof.* We first show that  $\mathscr{X} \wedge B \in O(A \wedge B)$ . To see this, note that  $\mathscr{X}_n \wedge B \to \mathscr{X}_{n+1} \wedge B$  is the pushout of a map of the form  $(-) \wedge A \wedge B \to *$ , and hence an  $A \wedge B$ -equivalence. Thus

$$\mathscr{X} \wedge B \stackrel{A \wedge B}{\simeq} \mathscr{X}_{\infty} \wedge B = *$$

as needed. Now note that  $\mathscr{X} \wedge \mathscr{Y}_n \to \mathscr{X} \wedge \mathscr{Y}_{n+1}$  is the pushout of a map  $(-) \wedge \mathscr{X} \wedge B \to *$ , and hence (by what we just did) also an  $A \wedge B$ -equivalence. Consequently

$$\mathscr{X}\wedge \mathscr{Y}\stackrel{A\wedge B}{\simeq} \mathscr{X}\wedge \mathscr{Y}_{\infty}=*,$$

concluding the proof.

### Characterizing $S^{p,q}$ -null and weakly $S^{p,q}$ -cellular spaces

The characterization of  $S^{p,q}$ -null spaces from Lemma 3.1.14(4) admits an interpretation in terms of homotopy sheaves using the notion of contraction (see Paragraph 2.1.10).

**Corollary 3.1.20.** If  $\mathscr{X}$  is a pointed connected space, then  $\mathscr{X}$  is  $S^{p,q}$ -null if and only if  $\pi_i(\mathscr{X})_{-q} = 0$  for all  $i \ge p - q$ .

*Proof.* Lemma 3.1.14(4) gives a necessary and sufficient condition for  $S^{p,q}$ -nullity of a pointed connected space in terms of contractibility of the (p,q)-fold loop space of  $\mathscr{X}$ . We may write

$$\Omega^{p,q}\mathscr{X}\simeq\Omega^{p-q}_{S^1}\Omega^{q,q}\mathscr{X},$$

and then the statement follows by appeal to the discussion of Paragraph 2.1.10 (in particular, [Mor12, Theorem 6.13]).  $\Box$ 

Lemma 3.1.21. Suppose *W* is a pointed connected space. The following statements are equivalent.

- 1. The space  $\mathscr{W}$  is  $S^{p,q}$ -null.
- 2. For every integer  $i \geq 1$ , the spaces  $K(\pi_i(\mathcal{W}), i)$  are  $S^{p,q}$ -null.
- 3. For every integer i, the Postnikov layers  $\tau_{\leq i} \mathcal{W}$  are  $S^{p,q}$ -null.

*Proof.* The implication (1)  $\Longrightarrow$  (2) is an immediate consequence of Corollary 3.1.20. The implication (2)  $\Longrightarrow$  (3) follows by induction using the fiber sequences  $K(\pi_i(\mathscr{X}), i) \to \tau_{\leq i} \mathscr{W} \to \tau_{\leq i-1} \mathscr{W}$  and Lemma 3.1.14(3). The implication (3)  $\Longrightarrow$  (1) follows from Lemma 3.1.14(2) and the fact that  $\mathscr{W} = \lim_i \tau_{\leq i} \mathscr{W}$ .

**Proposition 3.1.22.** Assume  $\mathscr{F} \to \mathscr{E} \xrightarrow{f} \mathscr{B}$  is a fiber sequence in  $\operatorname{Spc}(k)$  with  $\mathscr{B}$  connected.

- 1. The morphism f is a universal A-equivalence (that is, any base change of f in Spc(k) is an A-equivalence) if and only if  $\mathscr{F} \in O(A)$ .
- 2. If  $\mathscr{F}$  and  $\mathscr{B}$  lie in O(A), then  $\mathscr{E} \in O(A)$  as well.

*Proof.* The second statement is a special case of the first, so it remains to establish the first statement. If f is a universal A-equivalence, then  $\mathscr{F} \to *$  is as well by universality, so  $\mathscr{F} \in O(A)$  by definition. Conversely, suppose that  $\mathscr{F} \in O(A)$  and  $\mathscr{B}' \to \mathscr{B}$  is a morphism, with fiber denoted  $\mathscr{S}$ . Since  $\mathscr{B}$  is connected,  $* \to \mathscr{B}$  is an effective epimorphism and  $\mathscr{B} \simeq */\!/\Omega \mathscr{B}$ . Similarly  $\mathscr{X} \simeq \mathscr{F}/\!/\Omega \mathscr{B}$  and  $\mathscr{B}' \simeq \mathscr{S}/\!/\Omega \mathscr{B}$ . By universality of colimits, the base change of f along  $\mathscr{B}' \to \mathscr{B}$  is given by

$$(\mathscr{S} \times \mathscr{F}) / \!\!/ \Omega \mathscr{B} \to \mathscr{S} / \!\!/ \Omega \mathscr{B}$$

Since  $L_A$  preserves finite products (Lemma 3.1.8), this is a colimit of A-equivalences, and hence an A-equivalence.

**Corollary 3.1.23.** If  $f : \mathscr{X} \to \mathscr{Y}$  and  $g : \mathscr{Y} \to \mathscr{Z}$  are maps such that  $\operatorname{fib}(f), \operatorname{fib}(g) \in O(A)$  and  $\operatorname{fib}(g)$  is connected, then  $\operatorname{fib}(g \circ f)$  lies in O(A) as well.

Proof. By [ABH22, Proposition 3.1], there is a fiber sequence of the form

$$\operatorname{fib}(f) \longrightarrow \operatorname{fib}(g \circ f) \longrightarrow \operatorname{fib}(g).$$

We deduce that  $fib(g \circ f) \in O(A)$  by appeal to Proposition 3.1.22(2).

#### **Truncation and cellularity**

**Proposition 3.1.24.** Suppose  $\mathscr{X}$  is a pointed connected motivic space. Let  $q \ge 0$  and  $p \le d$  such that  $\Omega^{q,q}\mathscr{X}$  is d-connective. Then  $L^{p+q,q}\mathscr{X} \simeq L^{d+q,q}\mathscr{X}$ .

*Proof.* Since the map  $\mathscr{X} \to L^{d+q,q} \mathscr{X}$  is an  $S^{p+q,q}$ -equivalence (as  $d \ge p$ ), it suffices to prove that  $L^{d+q,q} \mathscr{X}$  is  $S^{p+q,q}$ -null. Since  $L^{d+q,q} \mathscr{X}$  is connected (Remark 3.1.17), for this it is enough to show that  $\pi_n(L^{d+q,q} \mathscr{X})_{-q} = 0$  for all  $n \ge p$  (Corollary 3.1.20). This is true for  $n \ge d$  by construction. By Lemma 3.1.18 we have  $\pi_n(L^{d+q,q} \mathscr{X}) \simeq \pi_n(\mathscr{X})$  for n < d, whence in this case the vanishing holds by assumption. This concludes the proof.

**Corollary 3.1.25.** Suppose  $\mathscr{X}$  is a pointed d-connected motivic space,  $d \geq 0$ . Then  $L^{p+q,q}\mathscr{X}$  is d-connected.

*Proof.* If  $p \ge d$  then  $(L^{p+q,q}\mathscr{X})_{\le d} \simeq \mathscr{X}_{\le d} = *$ , by Lemma 3.1.18. Otherwise  $L^{p+q,q}\mathscr{X} \simeq L^{d+q,q}\mathscr{X}$  by Proposition 3.1.24, and now the same argument works.

**Corollary 3.1.26.** Let  $\mathscr{X} \in O(S^{q,q})$  be pointed and connected. The following are equivalent:

1.  $\mathscr{X} \in O(S^{d,0}),$ 2.  $\Omega^{q,q} \mathscr{X} \in O(S^{d,0}),$ 3.  $\mathscr{X} \in O(S^{d+q,q}).$ 

*Proof.* It is clear that (1) implies (2) (see \$2.1.10), and (2) implies (3) by Proposition 3.1.24. Finally (3) implies (1) by Lemma 3.1.18.

The following is occasionally useful.

**Corollary 3.1.27.** Let  $f_i : \mathscr{X}_i \to \mathscr{Y}_i \in \operatorname{Spc}(k)_*$  be a cofiltered family of  $S^{p,q}$ -equivalences. Let  $f : \lim_i \mathscr{X}_i \to \lim_i \mathscr{Y}_i$  be the induced map. Assume that the connectivities of  $\mathscr{X} \to \mathscr{X}_i$  and  $\mathscr{Y} \to \mathscr{Y}_i$  tend to infinity with *i*, and  $\mathscr{X}, \mathscr{Y}$  connected. Then *f* is an  $S^{p,q}$ -equivalence.

*Proof.* Both  $\mathscr{X}$  and  $\mathscr{Y}$  being connected, the same is true for their  $S^{p,q}$ -nullifications (Corollary 3.1.25). Hence by Yoneda and Lemma 3.1.21, it will suffice to show the following: if  $\mathscr{W}$  is connected, *r*-truncated (for some *r*) and  $S^{p,q}$ -null, then  $\operatorname{Map}(\mathscr{X}, \mathscr{W}) \simeq \operatorname{Map}(\mathscr{Y}, \mathscr{W})$ . By assumption for  $i \gg 0$ , both  $\mathscr{X} \to \mathscr{X}_i$  and  $\mathscr{Y} \to \mathscr{Y}_i$  are *r*-connected. Hence

$$\operatorname{Map}(\mathscr{X}, \mathscr{W}) \simeq \operatorname{Map}(\mathscr{X}_i, \mathscr{W}) \simeq \operatorname{Map}(\operatorname{L}^{p,q} \mathscr{X}_i, \mathscr{W}) \simeq \operatorname{Map}(\mathscr{X}', \mathscr{W})$$

This concludes the proof.

#### Truncation, loop spaces and fiber sequences

**Lemma 3.1.28.** *Let k be a field.* 

- 1. Let  $\mathscr{G} \in \operatorname{Spc}(k)$  be a connected group. Then  $BL^{p,q}\mathscr{G} \simeq L^{p+1,q}B\mathscr{G}$ .
- 2. Let  $\mathscr{X} \in \operatorname{Spc}(k)_*$  be simply connected. Then  $\Omega L^{p+1,q} \mathscr{X} \simeq L^{p,q} \Omega \mathscr{X}$ .

*Proof.* (1) Note that  $L^{p+q,q}\mathscr{G}$  is again a connected group object because  $L^{p+q,q}$  preserves finite products by appeal to Lemma 3.1.8 and connectedness by appeal to Corollary 3.1.25. By construction  $L^{p+q,q}\mathscr{G}$ is  $S^{p+q,q}$ -null. It follows from Lemma 3.1.14(5) that since  $\mathscr{G} \equiv \Omega B\mathscr{G}$ ,  $BL^{p+q,q}\mathscr{G}$  is  $S^{p+1+q,q}$ -null. Hence the map  $B\mathscr{G} \to BL^{p+q,q}\mathscr{G}$  factors through a map  $L^{p+1+q,q}B\mathscr{G} \longrightarrow BL^{p+q,q}\mathscr{G}$ . We claim that this map is an equivalence. To see this, we construct an inverse, as follows. Consider the augmentation  $B\mathscr{G} \to L^{p+1+q,q}B\mathscr{G}$ . Since the target is  $S^{p+1+q,q}$ -null, it follows that  $\Omega L^{p+1+q,q}B\mathscr{G}$  is  $S^{p+q,q}$ -null by appeal to Lemma 3.1.14(5). Therefore, the map  $\mathscr{G} \to \Omega L^{p+1+q,q}B\mathscr{G}$  factors uniquely through a map  $L^{p+q,q}\mathscr{G} \to \Omega L^{p+1+q,q}B\mathscr{G}$ . Classifying this map provides the required inverse by universality.

(2) Applying (1) to  $\mathscr{G} = \Omega \mathscr{X}$  we get

$$\Omega L^{p+1,q} \mathscr{X} \simeq \Omega L^{p+1,q} B \Omega \mathscr{X} \simeq \Omega B L^{p,q} \Omega \mathscr{X} \simeq L^{p,q} \Omega \mathscr{X}.$$

Here we used Lemma 2.1.23, which applies since  $\Omega \mathscr{X}$ ,  $L^{p,q} \Omega \mathscr{X}$  are connected (Corollary 3.1.25).

Suppose  $\mathscr{F} \to \mathscr{E} \to \mathscr{B}$  is a fiber sequence of pointed motivic spaces. Lemma 3.1.14(2) implies that  $\operatorname{fib}(L^{p,q}\mathscr{E} \to L^{p,q}\mathscr{B})$  is  $S^{p,q}$ -null and therefore there is an induced map  $L^{p,q}\mathscr{F} \to \operatorname{fib}(L^{p,q}\mathscr{E} \to L^{p,q}\mathscr{B})$ . The following result gives some conditions under which this map is an equivalence. These conditions are not optimal, but will be sufficient for our future needs.

**Proposition 3.1.29.** Assume k is a field and suppose  $\mathscr{B}$  is a  $\max\{p, 1\}$ -connected motivic space,  $p \ge 0$ . If  $\mathscr{F} \to \mathscr{E} \to \mathscr{B}$  is a fiber sequence, then

$$\mathcal{L}^{p+q,q}\mathscr{F}\longrightarrow\mathcal{L}^{p+q,q}\mathscr{E}\longrightarrow\mathcal{L}^{p+q,q}\mathscr{B}$$

is also a fiber sequence.

*Proof.* We shall freely use that  $L^{p,q}$  preserves connectivity (Corollary 3.1.25).

Since  $\mathscr{B}$  is connected,  $\mathscr{E} \simeq \mathscr{F} /\!\!/ \Omega \mathscr{B}$ . Set  $\mathscr{E}' = L^{p+q,q} \mathscr{F} /\!\!/ L^{p+q,q} \Omega \mathscr{B}$ . Via Lemma 2.1.23 we obtain a fiber sequence

$$\mathcal{L}^{p+q,q}\mathscr{F}\longrightarrow \mathscr{E}'\longrightarrow B\mathcal{L}^{p+q,q}\Omega\mathscr{B}.$$

Since  $\mathscr{B}$  is (p+1)-connective, using Lemma 3.1.28 and Proposition 3.1.24 we identify the base as

$$BL^{p+q,q}\Omega\mathscr{B} \simeq B\Omega L^{p+q+1,q}\mathscr{B} \simeq L^{p+q+1,q}\mathscr{B} \simeq L^{p+q,q}\mathscr{B}.$$

It remains to show that  $\mathscr{E}' \simeq L^{p+q,q} \mathscr{E}$ . By construction we have an  $S^{p+q,q}$ -equivalence  $\mathscr{E} \to \mathscr{E}'$ , so it suffices to show that  $\mathscr{E}'$  is  $S^{p+q,q}$ -null. This holds by Lemma 3.1.14(3).

*Remark* 3.1.30. The functor  $L^{p,q}$  cannot preserve fiber sequences in general. Indeed, if A is a strictly  $\mathbb{A}^1$ -invariant sheaf of groups, then the fiber sequence

$$K(\mathbf{A}, p-1) \longrightarrow * \longrightarrow K(\mathbf{A}, p)$$

is not preserved by  $L^{p,0} = \tau_{< p}$ .

**Corollary 3.1.31.** If  $\mathscr{X}$  is a nilpotent motivic space, then  $L^{p+q,q}\mathscr{X}$  is again a nilpotent motivic space. More precisely, suppose given a principal refinement of  $\mathscr{X} \to \mathscr{X}_{\leq p}$ , consisting of fiber sequences

$$\mathscr{X}_{i+1} \longrightarrow \mathscr{X}_i \longrightarrow \mathscr{B}_i$$

with  $\mathscr{X}_0 = \mathscr{X}_{\leq p}$ ,  $\mathscr{B}_i = K(A_i, n_i)$  with  $n_i \geq p+1$ , and  $\lim_i \mathscr{X}_i \simeq \mathscr{X}$ . Then  $L^{p+q,q} \mathscr{X} \simeq \lim_i L^{p+q,q} \mathscr{X}_i$ , the induced sequences

$$\mathrm{L}^{p+q,q}\mathscr{X}_{i+1} \to \mathrm{L}^{p+q,q}\mathscr{X}_i \to \mathrm{L}^{p+q,q}\mathscr{B}_i$$

are fiber sequences, and  $L^{p+q,q}\mathscr{B}_i$  is an  $n_i$ -connective, grouplike commutative monoid.

*Proof.* The fiber sequences are preserved by  $L^{p+q,q}$  by Proposition 3.1.29, since  $\mathscr{B}_i$  is a *p*-connected. Since  $L^{p+q,q}$  preserves finite products it preserves grouplike commutative monoids, whence  $L^{p+q,q}\mathscr{B}_i$  is one. In particular it is nilpotent, and  $n_i$ -connective by Corollary 3.1.25. Also  $L^{p+q,q}\mathscr{X}_0 \simeq \mathscr{X}_0$  (via Corollary 3.1.20:  $\mathscr{X}_0$  is *p*-connected) is nilpotent, and hence by induction  $L^{p+q,q}\mathscr{X}_{i+1}$  is nilpotent. If we set  $\mathscr{X}' = \lim_i L^{p+q,q}\mathscr{X}_i$ , then  $\mathscr{X}'$  is nilpotent, since the connectivity of the  $L^{p+q,q}\mathscr{B}_i$  tends to infinity. It remains to prove that  $\mathscr{X}' \simeq L^{p+q,q}\mathscr{X}$ . Since  $\mathscr{X}'$  is  $S^{p+q,q}$ -null, for this it suffices to show that  $\mathscr{X} \to \mathscr{X}'$  is an  $S^{p+q,q}$ -equivalence. This follows from Corollary 3.1.27 (since the  $\mathscr{B}_i$  have connectivity tending to  $\infty$ , and the same is true for the  $L^{p,q}\mathscr{B}_i$  by Corollary 3.1.25, and hence so do  $\mathscr{X} \to \mathscr{X}_i$  and  $\mathscr{X}' \to L^{p,q}\mathscr{X}_i$ ).

#### **3.2** Stable weak $S^{p,q}$ -cellularity via *t*-structures

We now analyze weak  $S^{p,q}$ -cellularity in the context of stable homotopy categories (see Paragraph 3.1.2). Combining these observations with the discussion of conservativity of  $\mathbb{G}_m$ -stabilization (see 2.2.26–2.2.29), we will be able to deduce statements about weak  $S^{p,q}$ -cellularity *unstably*. Stably, the situation regarding weak  $S^{p,q}$ -cellularity is more well-behaved because the relevant localization functors can be realized using truncation with respect to suitable *t*-structures. (Our *t*-structures are homologically indexed, following topological conventions).

3.2.1. Set  $C = SH^{S^1}(k)$  or  $SH(k)^{\text{eff}}$ . The homotopy *t*-structures on each of these categories were described in Paragraphs 2.2.15 and 2.2.20. We write  $C_{\geq 0}$  for the positive part of the relevant *t*-structure. Given  $X \in Sm_k$ , we abuse notation and write  $X_+$  for the image of  $X \in Sm_k$  under the functor  $Sm_k \to C$ . We now introduce some new *t*-structures on each of these categories which we will then link to weak  $S^{p,q}$ -cellularity.

**Lemma 3.2.2.** The subcategory of C generated under colimits and extensions by objects of the form  $\Sigma^{p,q}X_+$  is the non-negative part of an accessible t-structure on C.

*Proof.* This is an immediate consequence of [Lur17, Proposition 1.4.4.11].

**Definition 3.2.3.** Write  $C_{\geq (p,q)}$  for the subcategory of C generated under colimits and extensions by objects of the form  $\Sigma^{p,q}X_+$  (the non-negative part of an accessible *t*-structure by Lemma 3.2.2);  $C_{\leq (p,q)}$  for the corresponding non-positive part and  $\tau_{\leq (p,q)}$  or  $\tau_{\geq (p,q)}$  for the corresponding truncation functors.

By Corollary 3.1.11, we know that the localization functor  $L^{p,q}$  on  $C_{\geq 0}$  is symmetric monoidal. The next result identifies  $L^{p,q}$  with truncation for the *t*-structures of Definition 3.2.3 and weakly  $S^{p,q}$ -cellular spaces with the non-negative part of a *t*-structure.

**Proposition 3.2.4.** Assume k is a field and C is as in Paragraph 3.2.1.

- 1. There is an identification  $\mathcal{C}_{>(p,q)} = O(S^{p,q})$ .
- 2. There is a canonical isomorphism of functors  $L^{p,q} \simeq \tau_{<(p,q)}$  on  $\mathcal{C}$ .

*Proof.* That  $C_{\geq(p,q)}$  consists of weakly  $S^{p,q}$ -cellular objects holds because  $C_{\geq(p,q)}$  is generated under colimits and cofiber extensions by  $\Sigma^{p,q}X_+$  and the latter is weakly  $S^{p,q}$ -cellular by Example 3.1.12, the fact that  $L^{p,q}$  is symmetric monoidal (Corollary 3.1.11) and preserves colimits (Amplification 3.1.7).

To conclude, it suffices to show that  $C_{<(p,q)}$  consists of  $S^{p,q}$ -null objects. Indeed, assuming that statement is true, suppose E is an arbitrary object of C. There is a cofiber sequence of the form

$$\tau_{\geq (p,q)}E \longrightarrow E \longrightarrow \tau_{<(p,q)}E.$$

Since  $\tau_{\geq (p,q)}E$  is weakly  $S^{p,q}$ -cellular, then since  $S^{p,q}$ -localization preserves colimits, Amplification 3.1.7 implies that  $E \to \tau_{<(p,q)}E$  is an  $S^{p,q}$ -equivalence, i.e., that the map

$$L^{p,q}E \longrightarrow L^{p,q}\tau_{<(p,q)}E$$

is an equivalence. On the other hand, since  $\tau_{<(p,q)}E$  is  $S^{p,q}$ -null by assumption, we conclude that the natural map  $\tau_{<(p,q)E} \to L^{p,q}\tau_{\le(p,q)}E$  is an equivalence as well. These two observations yield the natural transformation we seek.

To see that  $\mathcal{C}_{<(p,q)}$  consists of  $S^{p,q}$ -null objects, we proceed as follows. Generating  $S^{p,q}$ -equivalences are of the form  $S^{p,q}_+ \wedge X_+ \to X_+$ , and hence split as sums of maps of the form  $S^{p,q} \wedge X_+ \to 0$  and  $X_+ \xrightarrow{\mathrm{id}} X_+$ . Since any object is local with respect to the second class of maps,  $E \in \mathcal{C}_{\geq 0}$  being  $S^{p,q}$ -local is equivalent to requiring that  $\operatorname{Map}(S^{p,q} \wedge X_+, E) = 0$  for all  $X \in \operatorname{Sm}_k$ . This is precisely the definition of  $\mathcal{C}_{<(p,q)}$ .

*Remark* 3.2.5. Proposition 3.2.4 allows us to build a functorial weak  $S^{p,q}$ -cellularization of any spectrum. Moreover, it highlights the fundamental difference between  $S^{p,q}$ -localization in the unstable and stable situations. If E is any object, we have the cofiber sequence

$$\tau_{\geq (p,q)}E \longrightarrow E \longrightarrow \tau_{<(p,q)}E,$$

which is also a fiber sequence. Thus  $\operatorname{fib}(E \to L^{p,q}E)$  is itself weakly  $S^{p,q}$ -cellular. In contrast, working unstably, it is unclear that the homotopy fiber of the map  $\mathscr{X} \to L^{p,q}\mathscr{X}$  is itself weakly  $S^{p,q}$ -cellular. We now use the description of localization in terms of truncation with respect to a *t*-structure to establish a number of properties.

3.2.6. Up to shifts, the *t*-structures introduced above have been studied in several places including [Bac17, §4] and [BY20, §6]. We may identify the positive and negative parts in terms of homotopy sheaves. We summarize these observations in the following result.

Lemma 3.2.7. The following statements hold for C as in Paragraph 3.2.1.

- 1. The equality  $O(S^{p,q}) = C(q) \cap C_{\geq p}$  holds, i.e., the weakly  $S^{p,q}$ -cellular objects of C are the q-effective and p-connective objects.
- 2. If  $E \in C_{>0}$ , then the following statements are equivalent
  - (a)  $E \in \mathcal{C}_{<(p,q)}$ , *i.e.*, E is  $S^{p,q}$ -null,
  - (b)  $\pi_i(E)_{-q} = 0 \text{ for } i \ge p.$
- 3. If  $\alpha : E \to E' \in \mathcal{C}_{\geq 0}$  is such that  $\operatorname{fib}(\alpha) \in O(S^{p,q})$  and E' is connected, then  $\alpha$  is a  $S^{p,q}$ -equivalence.
- 4. If  $E \in O(S^{p,q})$  and  $E' \in O(S^{p',q'})$ , then  $E \wedge E' \in O(S^{p+p',q+q'})$ .

*Proof.* The first and the second statements follow from [BY20, Lemma 6.2(3)] and [Bac17, Proposition 4] after shifting. For the third statement, since E' is connected we conclude that there is a cofiber sequence in  $C_{>0}$  of the form

$$\operatorname{fib}(\alpha) \longrightarrow E \longrightarrow E',$$

and the result follows from Amplification 3.1.7. The final statement follows from the first two points.  $\Box$ 

*Remark* 3.2.8. Note that  $\tau_{\geq (n,n)} : C_{\geq 0} \to C_{\geq 0}$  coincides with the functor usually denoted  $f_n$ , i.e., the *n*-th slice cover (this follows from [BY20, Lemma 6.2(2)]).

One immediate consequence of the description of stable weak  $S^{p,q}$ -cellularity and  $S^{p,q}$ -nullity in terms of homotopy sheaves is the fact that cellularity is preserved by *R*-localization for  $R \subset \mathbb{Q}$  a subring. We record this consequence for ease of reference.

**Corollary 3.2.9.** Suppose  $R \subset \mathbb{Q}$  is a subring, and  $E \in C$  as in Paragraph 3.2.1. If E is  $S^{p,q}$ -null or weakly  $S^{p,q}$ -cellular, then so is  $E_R$ .

In the stable setting, the descriptions above allow us to establish weak cellular estimates for loop spaces. While the fact that  $\Omega$  changes weak cellular class in a controlled way is straightforward from the *t*-structure description, the weak cellular estimate for  $\Omega^{1,1}$  relies on Levine's analysis of the slice filtration [Lev08].

**Proposition 3.2.10.** Assume C is as in Paragraph 3.2.1. Suppose  $E \in C_{\geq 0}$  and  $p, q \geq 0$  are integers. For (2) and (3) below, assume that either k is infinite or  $C = SH(k)^{\text{eff}}$ .

- 1. If  $E \in O(S^{p,q})$ , then  $\Omega E \in O(S^{p-1,q})$ .
- 2. If  $E \in O(S^{p,q})$ , then  $\Omega^{1,1}E \in O(S^{p-1,q-1})$ .
- 3. If  $E \in O(S^{p,q})$ , then  $\Omega^{2,1}E \in O(S^{p-2,q-1})$ .

*Proof.* For the first statement, we want to show that  $E \in \Sigma C_{\geq (p,q)}$ , but this follows from Lemma 3.2.7(1). For the second statement we first treat  $C = SH(k)^{\text{eff}}$ . In that case, since  $\Omega^{1,1}\Sigma^{p,q}X_+ = \Sigma^{p-1,q-1}X_+$ , it follows that  $\Omega^{1,1}$  lowers the weight of the generators. Since  $\Omega^{1,1}$  preserves colimits, the result then follows by Proposition 2.2.21. When  $C = SH^{S^1}(k)_{\geq 0}$ , it suffices to observe that[Lev08, Theorem 7.4.2] implies that  $\Omega^{1,1}$  sends  $SH^{S^1}(k)(q+1)$  into  $SH^{S^1}(k)(q)$ . The third statement follows by combining the first two.

While the functors  $\sigma^{\infty}$  and  $\Sigma_{S^1}^{\infty}$  preserve weakly cellular objects since they preserve colimits and generators, the corresponding statements for the adjoint functors are not immediate. The following result is key to our comparison of stable and unstable cellularity in the sequel.

**Proposition 3.2.11.** Let k be a perfect field, and  $p \in \mathbb{Z}, q \ge 0$ . The functors

$$\omega^{\infty} : \mathrm{SH}(k)^{\mathrm{veff}} \to \mathrm{SH}^{S^1}(k)_{\geq 0}$$

and


$$\Omega_{S^1}^{\infty} : \mathrm{SH}^{S^1}(k)_{\geq 0} \to \mathrm{Spc}(k)$$

preserve weakly  $S^{p,q}$ -cellular objects,  $S^{p,q}$ -equivalences and  $S^{p,q}$ -null objects. (For the second functor we assume in addition that  $p - q \ge 0$ .)

*Proof.* It is clear that both functors preserve  $S^{p,q}$ -null objects. To conclude, it will be enough to show that the functors send  $S^{p,q}$ -nullifications to  $S^{p,q}$ -equivalences.

For the first functor, let  $E \in SH(k)^{\text{veff}}$ . We have the cofiber sequence  $\tau_{\geq (p,q)}E \to E \to \tau_{<(p,q)}E \simeq$  $L^{p,q}E$ . The second map is an  $S^{p,q}$ -nullifications, and so it will be enough to show that  $\omega^{\infty}$  preserves weakly  $S^{p,q}$ -cellular objects. It thus suffices to show that  $\omega^{\infty}$  sends  $SH(k)^{eff}(n)$  to  $SH^{S^1}(k)(n)$ ; this assertion is a consequence of [Lev08, Theorem 9.0.3] (note: this theorem implicitly assumes k is infinite and perfect because of its appeal to the techniques of proof used in that of [Lev08, Theorem 7.1.1.], however as mentioned in [Lev08, Remark 9.0.4], the assumption of infinitude of k is not necessary for spectra satisfying Axiom A3 [Lev08, p. 233]. That Axiom A3 always holds for spectra in the image of  $\omega^{\infty}$  is a consequence of [Lev06, Corollary 9.4.3]). <sup>1</sup>

For the second functor, let  $E \in SH^{S^1}(k)_{\geq 0}$ . Using a variant of Construction 3.1.16, we can form a (transfinite) sequence of pushout squares in  $SH^{S^1}(k)_{\geq 0}$ 



with  $i \ge 0$  and  $\operatorname{colim}_n E_n \simeq L^{p,q} E$ . Let us write  $L : \operatorname{CMon}(\operatorname{Spc}(k)) \leftrightarrows \operatorname{SH}^{S^1}(k)_{\ge 0} : R$  for the evident adjunction, and  $F : \operatorname{Spc}(k)_* \to \operatorname{CMon}(\operatorname{Spc}(k))$  for the evident left adjoint functor. Since R is fully faithful (Corollary 2.2.24), the above pushout square is obtained by applying L to a pushout square in  $\operatorname{CMon}(\operatorname{Spc}(k))$ 

$F\Sigma^i S^{p,q} \wedge X_+$	$\longrightarrow$	$RE_n$
		ļ
0	$\longrightarrow$	$E'_{n+1}.$

Corollary 2.2.25 tells us that  $E'_{n+1} \simeq RE_{n+1}$ , and so  $RE_n \to RE_{n+1}$  is an  $S^{p,q}$ -equivalence; hence so is  $\Omega_{S^1}^{\infty}E_n \to \Omega_{S^1}^{\infty}E_{n+1}$  (see Amplification 3.1.9). After all is done we conclude that  $\Omega_{S^1}^{\infty}E \to \Omega_{S^1}^{\infty}L^{p,q}E$  is an  $S^{p,q}$ -equivalence, as needed. 

#### Detecting weak-cellularity stably

We may also refine the Whitehead theorem for nilpotent motivic spaces [ABH22, Theorem 5.4] to a statement about stable detection of  $S^{p,q}$ -equivalences. In particular, the result below implies that for a given nilpotent space, weak cellularity can be detected stably.

**Theorem 3.2.12** (Weakly cellular Whitehead theorem). Let k be a perfect field of exponential characteristic e and  $f: \mathscr{X} \to \mathscr{Y}$  be a map of nilpotent motivic spaces. Suppose one of the following conditions holds.

- 1.  $\sum_{S^1}^{\infty} f$  has fiber in  $O(S^{p+q,q})$ 2.  $\sum_{S^1}^{\infty} f$  has cofiber in  $O(S^{2,2})$  and  $\sum^{\infty} f$  has fiber in  $O(S^{p+q,q})$

<sup>&</sup>lt;sup>1</sup>An alternative proof is found in [BE21, Theorem 5.1].

- 3.  $\sum_{S^1}^{\infty} f$  has cofiber in  $O(S^{2,2})$ ,  $\Sigma^{\infty} f$  has fiber in  $O(S^{p,0})$  and  $\Sigma^{\infty} f \wedge H\mathbb{Z}$  has cofiber in  $O(S^{q,q})$
- 4.  $\sum_{S^1}^{\infty} f$  has cofiber in  $O(S^{2,2})[1/e]$ , k has finite virtual 2-étale cohomological dimension,  $\sum^{\infty} f \wedge H\mathbb{Z}$
- has fiber in  $O(S^{p+q,q})$ , and for every ordering  $\sigma$  of k,  $r_{\sigma}(f) \wedge H\mathbb{Z}$  has fiber in  $O(S^p)$

Then f is an  $S^{p+q,q}$ -equivalence.

(Recall the real realization functor  $r_{\sigma}$  from Definition 2.3.4.)

*Proof.* (1) We want to show that  $L^{p+q,q}f$  is an equivalence. The spaces  $L^{p+q,q}\mathscr{X}$  and  $L^{p+q,q}\mathscr{Y}$  are again nilpotent by appeal to Corollary 3.1.31 (or simply Corollary 3.1.25 if  $p \ge 1$ ). By the Yoneda lemma, it suffices to show that for any nilpotent  $S^{p+q,q}$ -null space  $\mathscr{T}$ , the map  $Map(\mathscr{Y}, \mathscr{T}) \to Map(\mathscr{X}, \mathscr{T})$  is an equivalence. Since  $\mathscr{T}$  is nilpotent the map  $\mathscr{T} \to \mathscr{T}_{< p}$  admits a principal refinement [ABH22, Theorem 5.1]. In particular, there are a sequence of  $S^{p+q,q}$ -null spaces  $\mathscr{T}_i$  and fiber sequences of the form

$$\mathscr{T}_{i+1} \longrightarrow \mathscr{T}_i \longrightarrow K(\mathbf{A}_i, n_i)$$

where  $n_i \ge p+1$ ,  $\mathscr{T} = \lim_i \mathscr{T}_i$ ,  $\mathscr{T}_0 = \mathscr{T}_{<p}$  and  $\mathbf{A}_i$  is a subquotient of  $\pi_j(\mathscr{T})$  for some  $j \ge p$ . Since contraction is exact and  $\pi_j(\mathscr{T})_{-q} = 0$  also  $(\mathbf{A}_i)_{-q} = 0$ . It now suffices to show that  $\operatorname{Map}(\mathscr{V}, \mathscr{W}) \to$  $\operatorname{Map}(\mathscr{X}, \mathscr{W})$  is an isomorphism, for  $\mathscr{W} = K(\mathbf{A}_i, n_i)$  and  $\mathscr{W} = \mathscr{T}_0$ . In the former case we use that since  $(\mathbf{A}_i)_{-q} = 0$ ,  $\Sigma^{n_i} H \mathbf{A}_i \in \operatorname{SH}^{S^1}(k)_{\ge 0}$  is  $S^{p+q,q}$ -null. Since  $\Sigma_{S^1}^{\infty} f$  is an  $S^{p+q,q}$ -equivalence, we conclude that  $\operatorname{Map}(\mathscr{Y}, K(\mathbf{A}_i), n_i)) \to \operatorname{Map}(\mathscr{X}, K(\mathbf{A}_{i,i}))$  is an isomorphism for each  $i \ge 0$  as required. In the latter case we use since  $\Sigma_{S^1}^{\infty} f$  has fiber in  $O(S^{p+q,q})$  it is *p*-connected, and hence so is *f* [ABH22, Theorem 5.4]. Thus  $\mathscr{X}_{<p} \simeq \mathscr{Y}_{<p}$  and consequently

$$\operatorname{Map}(\mathscr{X}, \mathscr{T}_{< p}) \simeq \operatorname{Map}(\mathscr{X}_{< p}, \mathscr{T}_{< p}) \simeq \operatorname{Map}(\mathscr{Y}_{< p}, \mathscr{T}_{< p}) \simeq \operatorname{Map}(\mathscr{Y}, \mathscr{T}_{< p}).$$

(2) It will be enough to show: if  $E \in SH^{S^1}(k)(2)_{\geq 0}$  and  $\sigma^{\infty}E \in SH(k)(q)_{\geq p}$  then  $E \in SH^{S^1}(k)(q)_{\geq p}$ (apply this to fib $(\Sigma_{S^1}^{\infty}f)$  and then use (1)). Since  $SH^{S^1}(k)(q)^{\heartsuit} \simeq SH(k)(q)^{\heartsuit}$  (Theorem 2.2.29), it suffices to prove that  $E \in SH^{S^1}(k)(q)$  as soon as  $\sigma^{\infty}E \in SH(k)(q)$  (and  $E \in SH^{S^1}(k)(2)_{\geq 0}$ ). Replacing E by  $E/f_qE$ , we may assume that  $\Omega^{q,q}E = 0$ . Suppose that in addition  $E \in SH^{S^1}(k)_{\geq n}$  and consider the map  $\iota : E \to \omega^{\infty}\sigma^{\infty}E$ . Then  $\pi_n(\iota)$  is an isomorphism (Theorem 2.2.29). It follows that  $\pi_n(\omega^{\infty}\sigma^{\infty}E)_{-q} = 0$ , whence  $\pi_n(\omega^{\infty}\sigma^{\infty}E) = 0$  by [BY20, Lemma 6.2(iii)]. Thus  $E \in SH^{S^1}(k)_{\geq n+1}$ . Since n was arbitrary, E = 0 as needed.

(3) By Lemma 3.2.7 it is enough to show for  $E \in SH(k)^{veff}$  that if  $E \wedge H\mathbb{Z} \in SH(k)^{eff}(q)$  then  $E \in SH(k)^{eff}(q)$ . For this see [Bon20, Theorem 3.1.1(II)].

(4) Using (3) and the interaction between inverting  $\rho$  and real realization established in [Bac18a], this immediate from Proposition 3.2.13 below.

The following is a generalization of [Bac18b, Theorem 16, Lemma 19, Theorem 33].

**Proposition 3.2.13.** Let k be a perfect field of exponential characteristic e, and  $E \in SH(k)^{veff}[1/e]$ . Suppose that  $E[1/\rho] \in SH(k)_{\geq p}$  and  $E \wedge H\mathbb{Z} \in SH(k)_{\geq p}$ . If k has finite virtual 2-étale cohomological dimension, then  $E \in SH(k)_{>p}$ .

*Proof.* We may assume p = 1. We have  $\pi_0(E)_*[1/\rho] = 0$  and  $\pi_0(E)_*/\eta = 0$ ; we must prove that  $\pi_0(E)_* = 0$ . Let  $x \in \pi_0(E)_*(K)$  for some field K finitely generated over k. We have  $\rho^n x = 0$  for some n, and we can write  $x = \eta^n y$ . Now we have (see also §2.2.22)

$$0 = \eta^n \rho^n x = (\eta^2 \rho)^n y = (\eta(h-2))^n y = (-2\eta)^n y = \pm 2^n x.$$

In other words every element of  $\pi_0(E)_*$  is annihilated by a power of 2. We claim that there exists N = N(K) such that for d > N,  $\rho : \pi_0(E)_d(K) \to \pi_0(E)_{d+1}(K)$  is an isomorphism. This is proved at the

end. We deduce that if  $x \in \pi_0(E)_i(K)$  then  $\rho^e x = 0$ , where  $e = \max(N - i, 0)$ . Hence by the argument from before we see that  $2^e x = 0$ . Since e only depends on i (and K), it will be enough to show that every element of  $\pi_0(E)_i$  is divisible by 2. If i > N this is trivial. If  $i \le N$  the composite

$$\pi_0(E)_{N+1} \xrightarrow{\rho} \pi_0(E)_{N+2} \xrightarrow{\eta^{2+(N-i)}} \pi_0(E)_i$$

is surjective. But as before  $\eta^{2+(N-i)}\rho$  is divisible by 2.

It remains to prove the claim. Let  $M = \pi_0^{\text{eff}}(E) \in \text{SH}(k)^{\text{eff}}^{\heartsuit}$ . Then  $\pi_0(E)_* \simeq \pi_0(M)_*$ , by [Bac17, Proposition 5(1)]. It will suffice to show that for  $i \in \{0, 1\}$  and d > N = N(K) we have  $\pi_i(\operatorname{cof}(M \xrightarrow{\rho} M))_d(K) = 0$ . Since  $\pi_0(-)_0$  is an exact conservative functor on  $\operatorname{SH}(k)^{\text{eff}}^{\heartsuit}$  [Bac17, Proposition 5(3)],  $M \in \operatorname{SH}(k)^{\text{eff}}^{\heartsuit}$  is 2-primary torsion (i.e.  $M \simeq \operatorname{colim}_n M[2^n]$ ) and hence so are  $\pi_*(M)_*$ . It follows that we may apply [BEØ21, Lemma 5.23] to  $\operatorname{cof}(M \xrightarrow{\rho} M)$ . The result follows.

#### 3.3 Examples

Here, we collect some useful examples.

*Example* 3.3.1. If  $\mathscr{X} \in O(A)$  is a pointed space, then  $\mathscr{X}^n \in O(A)$  and  $\mathscr{X}^{\wedge n} \in O(A)$  as well. Indeed, the first statement is an immediate consequence of Lemma 3.1.14. The second statement follows by induction from the first and the observation that objects in O(A) are stable under colimits and extensions, i.e., Lemma 3.1.6. In more detail, repeated application of this observation shows that if  $\mathscr{X}, \mathscr{Y} \in O(A)$ , then so are  $\mathscr{X} \vee \mathscr{Y}$  and there is a cofiber sequence of the form

$$\mathscr{X} \lor \mathscr{Y} \longrightarrow \mathscr{X} \times \mathscr{Y} \longrightarrow \mathscr{X} \land \mathscr{Y}$$

which shows that  $\mathscr{X} \wedge \mathscr{Y} \in O(A)$  as well.

*Example* 3.3.2. It follows that if  $\mathscr{X} \in O(A)$  is in addition connected, then  $\Omega \Sigma \mathscr{X} \in O(A)$  as well. Indeed, by [AWW17, Theorems 2.1.1 and 2.4.2] we have the James model  $J(\mathscr{X})$  of  $\Omega \Sigma \mathscr{X}$ , which is a filtered colimit of spaces  $J_n(\mathscr{X})$ , where  $J_0(\mathscr{X}) = *$  and for  $i \ge 0$  there are cofiber sequences of the form

$$J_i(\mathscr{X}) \longrightarrow J_{i+1}(\mathscr{X}) \longrightarrow \mathscr{X}^{\wedge i+1}.$$

Induction using the fact that  $\mathscr{X}^{\wedge i+1} \in O(A)$  shows that  $J_i(\mathscr{X}) \in O(A)$  for all i, and thus  $J(\mathscr{X}) = \operatorname{colim}_i J_i(\mathscr{A}) \in O(A)$  as well.

*Example* 3.3.3 (Relative James constructions). If  $f : A \to X$  is a morphism of pointed spaces, then there is a James-style description of  $\operatorname{fib}(X \to \operatorname{cof}(f))$ , due initially to B. Gray. Indeed, one modifies  $J(\mathscr{X})$  to J(f) which is the sub-simplicial presheaf of J(X) consisting of words such that every letter beyond the first lies in A. In this case, the successive layers are of the  $X \wedge A^{\wedge n-1}$ . In particular, it follows that if A and X are weakly B-cellular, then  $\operatorname{fib}(X \to \operatorname{cof}(f))$  is also weakly B-cellular.

*Example* 3.3.4. If X is a smooth k-scheme and  $\pi : \mathscr{E} \to X$  is a rank r vector bundle on X, then  $\operatorname{Th}(\pi) \in O(S^{2r,r})$ . This statement is immediate if  $\pi$  is a trivial bundle by appeal to Example 3.1.12. In general, if we choose an open cover  $u : U \to X$  along which  $\pi$  trivializes, then there is an equivalence  $\operatorname{colim}_{n \in \Delta^{\operatorname{op}}} C(u) \to X$ . It follows that  $\operatorname{Th}(\pi)$  is realized as a colimit of Thom spaces of trivial bundles, and one concludes by appeal to Lemma 3.1.6.

*Example* 3.3.5. If X is an irreducible smooth k-scheme and  $Z \subset X$  is a smooth closed subscheme everywhere of codimension c, then homotopy purity yields a cofiber sequence of the form:

$$X \smallsetminus Z \longrightarrow X \longrightarrow \operatorname{Th}(\nu_{Z/X}),$$

where  $\nu_{Z/X}$  is the normal bundle of Z in X. If particular, if  $X \setminus Z \in O(S^{p,q})$  for  $p \ge 2c, q \ge c$ , then  $X \in O(S^{2c,c})$  as well.

*Example* 3.3.6 (Projective spaces). The space  $\mathbb{P}^n \in O(S^{2,1})$ . Indeed, there are cofiber sequences of the form  $\mathbb{P}^{n-1} \to \mathbb{P}^n \to \mathbb{P}^{1 \wedge n}$ . Example 3.3.1 shows that  $\mathbb{P}^{1 \wedge n} \in O(S^{2,1})$  and we conclude by appeal to Lemma 3.1.6. It follows also that  $\mathbb{P}^{\infty} \simeq K(\mathbb{Z}(1), 2) \in O(S^{2,1})$ .

Likewise, we may consider the Panin-Walter quaternionic projective spaces HP<sup>n</sup> [PW21, §3]. For every  $n \ge 1$  there are cofiber sequences of the form

$$\operatorname{HP}^n \longrightarrow \operatorname{HP}^{n+1} \longrightarrow \operatorname{Th}(\nu_n),$$

where  $\nu_n$  is a rank 2 bundle. Example 3.3.8 shows that  $\operatorname{Th}(\nu_n) \in O(S^{4,2})$  and  $\operatorname{HP}^1 \simeq S^{4,2}$  by [ADF17, Theorem 2]. By induction and appeal to Lemma 3.1.6 we conclude that  $\operatorname{HP}^n \in O(S^{4,2})$  and thus that  $\operatorname{HP}^{\infty} \simeq B \operatorname{SL}_2 \in O(S^{4,2})$  as well.

*Example* 3.3.7. Suppose X is a connected smooth projective variety equipped with a  $\mathbb{G}_m$ -action with isolated fixed points, then  $X \in O(S^{2,1})$ . In this case, the variety X admits a Bialynicki-Birula paving by affine spaces [ByB73] and can be written as an increasing union of smooth open subschemes with smooth closed complements at each stage. In this case, the result follows by induction appealing to Example 3.3.5 upon noticing that X has an open dense subscheme isomorphic to  $\mathbb{A}^n$ . It follows that if G is a split, simply connected, semi-simple k-group, then  $G/P \in O(S^{2,1})$  for any parabolic subgroup  $P \subset G$ . In particular, when  $G = GL_n$  we deduce that Grassmannians and generalized flag varieties lie in  $O(S^{2,1})$ . Since  $BGL_n \simeq \operatorname{colim}_n Gr_{r,N}$ , we see  $BGL_n \in O(S^{2,1})$  as well.

*Example* 3.3.8. The space  $MGL_n$  is the Thom space of the universal rank n vector bundle on  $BGL_n$ . Writing the former as a colimit of the universal rank n vector bundles over  $Gr_{n,N}$ , we conclude from Example 3.3.8 that  $MGL_n \in O(S^{2n,n})$ . Analogously,  $MSL_n$  is constructed as the Thom space of the universal rank n vector bundles over  $BSL_n$  and these may be written as a colimit of vector bundles over finite-dimensional approximations to  $BSL_n$ . Thus,  $MSL_n \in O(S^{2n,n})$ . Similarly,  $MSp_{2n} \in O(S^{4n,2n})$  since this may be realized as the Thom space of a universal rank 2n bundle on  $BSp_{2n}$ .

*Example* 3.3.9 (Motivic Eilenberg–Mac Lane spaces). Assume k is a field that has characteristic 0. Voevodsky showed in [Voe04] that if  $n \ge 2$ , then  $K(\mathbb{Z}(n), 2n) \in O(S^{2n,n})$ , generalizing the first part of Example 3.3.6; see Paragraph 5.2.5 for further explanations of the notation. Indeed, [Voe04, Lemma 6.5] shows that we may write  $K(\mathbb{Z}(n), 2n)$  as a filtered colimit of spaces  $K_{i,n}$  such that  $K_{0,n} = *$  and fitting into cofiber sequences of the form

$$K_{i,n} \longrightarrow K_{i+1,n} \longrightarrow \operatorname{Sym}^n(S^{2n,n}),$$

and [Voe04, Theorem 4.1] implies that  $\text{Sym}^n(S^{2n,n}) \in O(S^{2n,n})$  (this latter statement uses the assumption that k has characteristic 0). We will revisit and refine this observation in Section 5.

*Remark* 3.3.10. More generally, provided k is perfect, for any  $E \in SH(k)^{veff}$  we have  $E_n \in O(S^{2n,n})$ . Since  $E_n = \Omega^{\infty} \Sigma^{2n,n} E$ , this follows from Proposition 3.2.11.

**Proposition 3.3.11.** Suppose k is a field and  $n \ge 1$  is an integer.

- 1. The simply-connected split k-group schemes  $SL_{n+1}$ ,  $Sp_{2n}$ ,  $Spin_{n+2}$  and  $G_2$  lie in  $O(S^{3,2})$ .
- 2. The (Nisnevich) classifying spaces of any of the group schemes appearing in Point (1) lies in  $O(S^{4,2})$ .

*Proof.* Observe that  $SL_2 \simeq Sp_2 \simeq Spin_3$  and all are equivalent to  $S^{3,2}$ . Next, recall that there are fiber sequences of the form

$$SL_n \longrightarrow SL_{n+1} \longrightarrow S^{2n-1,n},$$
  

$$Spin_n \longrightarrow Spin_{n+1} \longrightarrow Q_n,$$
  

$$Sp_{2n} \longrightarrow Sp_{2n+2} \longrightarrow S^{4n+3,2n+2},$$
  

$$SL_3 \longrightarrow G_2 \longrightarrow Q_6.$$

The first and the fourth sequences are described in [AHW18, AHW20], the third in [AHW19], while the second is established in [Aso22]. Since  $Q_{2n} \simeq S^{2n,n}$  and  $Q_{2n-1} \simeq S^{2n-1,n}$  (see by [ADF17, Theorem 2]), the result follows immediately from Proposition 3.1.22(2). The second point follows from the first by appeal to Lemma 3.1.14(5) and the observation that  $\Omega B_{\text{Nis}} G \simeq G$ .

**Corollary 3.3.12.** The geometric classifying spaces  $B \operatorname{SL}_n$  and  $B \operatorname{Sp}_{2n} \in O(S^{4,2})$  for every  $n \ge 1$ .

*Example* 3.3.13. The classifying space  $B\mathbb{G}_m = K(\mathbb{Z}(1), 2)$  (see Example 5.2.7) is  $S^{4,2}$ -null. This observation has two natural generalizations: Voevodsky's motivic Eilenberg–Mac Lane space  $K(\mathbb{Z}(n), 2n)$  is  $S^{2n+2,n+1}$ -null. The first non-trivial stage of the simplicial Postnikov tower of  $K(\mathbb{Z}(n), 2n)$  is the Eilenberg–Mac Lane space  $K(\mathbb{K}_n^M, n)$ , which is again  $S^{2n+2,n+1}$ -null. All of these facts follow from standard facts about contractions; see, e.g., [HW19, Example 2.4].

# 4 A weakly-cellular Whitehead tower and consequences

In this section, we amplify some of the results of the previous section. We begin in Section 4.1 by constructing refinements of the classical Whitehead and Postnikov towers taking weak  $S^{p,q}$ -cellularity into account: the *k*-invariants in the refined towers can be taken to be  $\mathbb{P}^1$ -infinite loop spaces (see Amplification 4.1.13). We then deduce a number of consequences of this refinement including refined cellularity estimates on various kinds of loop spaces (Proposition 4.2.1), fibers, cofibers and we establish versions of the Whitehead theorem taking cellularity into account. We also show that weak and strong cellularity agree in favorable cases.

#### 4.1 A weakly-cellular Whitehead tower

If  $\mathscr{X}$  is a space that is weakly  $S^{p,q}$ -cellular for some (p,q), then there is no reason for the first non-trivial layer of the classical Postnikov tower of  $\mathscr{X}$  to again be weakly  $S^{p,q}$ -cellular as the following example shows.

*Example* 4.1.1. Consider the motivic Eilenberg–MacLane space  $K(\mathbb{Z}(n), 2n)$ . Note that  $K(\mathbb{Z}(n), 2n) \in O(S^{2n,n})$  if k has characteristic 0 by appeal to Example 3.3.9. The first non-trivial stage of the Postnikov tower and the first non-trivial k-invariant yield a sequence morphisms of the form

$$K(\mathbb{Z}(n),2n) \longrightarrow K(\mathbf{K}_n^M,n) \xrightarrow{k_n} K(\mathbf{H}_{\mathbb{Z}}^{n-1,n},n+1)$$

where  $\mathbf{H}_{\mathbb{Z}}^{n-1,n}$  is the sheaf associated with the presheaf  $U \mapsto H^{n-1,n}(U,\mathbb{Z})$ . Note that  $(\mathbf{H}_{\mathbb{Z}}^{n-1,n})_{-i} = \mathbf{H}_{\mathbb{Z}}^{n-1-i,n-i}$ . Since  $\mathbf{H}_{\mathbb{Z}}^{0,1} = 0$  by the computation of weight 1 motivic cohomology, it follows that  $K(\mathbf{H}_{\mathbb{Z}}^{n-1,n}, n+1)$  is  $S^{p+n,n}$ -null for any  $p \ge 0$  by appeal to Corollary 3.1.20. Observe that  $K(\mathbf{K}_{n}^{M}, n)$  cannot be weakly  $S^{p+n,n}$ -cellular as the k-invariant above shows that it admits a non-trivial map to an  $S^{p+n,n}$ -null space.

#### Weak cellularization of nilpotent spaces

Example 4.1.1 also demonstrates how cellularity can be "recovered" by adding cells in higher degrees.

**Definition 4.1.2.** For a nilpotent motivic space  $\mathscr{X}$  set  $\tau_{>(p,q)}\mathscr{X} = \operatorname{fib}(\mathscr{X} \to L^{p,q}\mathscr{X}).$ 

*Example* 4.1.3. Let  $\mathscr{X} = K(\mathbf{A}, n)$  with A strictly  $\mathbb{A}^1$ -invariant. Then  $\mathscr{X} = \Omega_{S^1}^{\infty} \Sigma H \mathbf{A}$  and hence by Proposition 3.2.11 we find

$$au_{\geq (p,q)} K(\mathbf{A}, n) \simeq \Omega^{\infty}_{S^1} \tau_{\geq (p,q)} \Sigma^n H \mathbf{A}.$$

In particular if  $p - q \le n$  then

$$\tau_{\geq (p,q)} K(\mathbf{A}, n) \simeq \Omega_{S^1}^{\infty} f_q \Sigma^n H \mathbf{A}$$

*Remark* 4.1.4. At this stage, the space  $\tau_{\geq (n+q,q)}K(\mathbf{A}, n)$  can be quite strange. For example, if  $\mathbf{A}$  is birationally invariant, then  $\mathbf{A}_{-1} = 0$  and so  $\tau_{\geq (n+q,q)}K(\mathbf{A}, n) = 0$  for q > 0.

*Example* 4.1.5. Note that

$$\Omega \tau_{\geq n+q,q} K(\mathbf{A}, n) \simeq \tau_{\geq n-1+q,q} K(\mathbf{A}, n-1) \in O(S^{n-1+q,q}).$$

**Lemma 4.1.6.** If  $\mathscr{Y} \in O(S^{p,q})$ , then  $\operatorname{Map}(\mathscr{Y}, \tau_{\geq (p,q)}\mathscr{X}) \simeq \operatorname{Map}(\mathscr{Y}, \mathscr{X})$ .

*Proof.* Indeed since  $Map(\mathscr{Y}, L^{p,q}\mathscr{X}) = *$ .

**Lemma 4.1.7.** If  $\Omega^{q,q} \mathscr{X}$  is d-connective for  $d \ge p$ , then  $\tau_{\ge (p+q,q)} \mathscr{X} \simeq \tau_{\ge (d+q,q)} \mathscr{X}$ .

Proof. Immediate from Proposition 3.1.24.

**Construction 4.1.8.** Suppose given a nilpotent space  $\mathscr{X}$ , p > 0, and a principal refinement of  $\mathscr{X} \to \mathscr{X}_{< p}$  as in Corollary 3.1.31. We call the sequences

$$\tau_{\geq (p+q,q)} \mathscr{X}_{i+1} \to \mathscr{X}_i \to \tau_{\geq (p+q,q)} K(\mathbf{A}_i, n_i)$$

a refined Postnikov tower of  $L^{p+q,q} \mathscr{X}$ . Note the following:

- $\tau_{>(p+q,q)} \mathscr{X}_0 = *$  (since  $\mathscr{X}_0 = \mathscr{X}_{< p}$  is  $S^{p+q,q}$ -local)
- $\tau_{\geq (p+q,q)} K(\mathbf{A}_i, n_i) \simeq \tau_{\geq (n_i+q,q)} K(\mathbf{A}_i, n_i)$  (by Lemma 4.1.7, since  $n_i \geq p$ )
- The sequences are fiber sequences (since  $n_i > p$ , so  $L^{p+q,q} \mathscr{X}_{i+1} \to L^{p+q,q} \mathscr{X}_i \to L^{p+q,q} K(\mathbf{A}_i, n_i)$  is a fiber sequence by Proposition 3.1.29).
- $\lim_{i} \tau_{\geq (p+q,q)} \mathscr{X}_{i} \simeq \tau_{\geq (p+q,q)} \mathscr{X}$  (since the same holds for  $L^{p+q,q}$ , by Corollary 3.1.31)

**Theorem 4.1.9.** Let  $\mathscr{X}$  be nilpotent. Then  $\tau_{\geq (p+q,q)}\mathscr{X} \in O(S^{p+q,q})$ .

*Proof.* We use the refined Postnikov tower of  $L^{p+q,q} \mathscr{X}$ . We first show by induction that each  $\tau_{\geq (p+q,q)} \mathscr{X}_i \in O(S^{p+q,q})$ . For i = 0 this is clear. The rotated fiber sequence

$$\Omega \tau_{\geq (p+q,q)} K(\mathbf{A}_i, n_i) \to \tau_{\geq (p+q,q)} \mathscr{X}_{i+1} \to \mathscr{X}_i$$

shows (via Example 4.1.5, Lemma 4.1.7 and Lemma 3.1.14) inductively that  $\tau_{\geq (p+q,q)} \mathscr{X}_{i+1} \in O(S^{p+q,q})$ .

It remains to prove that  $\tau_{\geq (p,q)} \mathscr{X} \in O(S^{p,q})$ . If  $\mathscr{Y}$  is  $S^{p+q,q}$ -null then so is  $\tau_{\leq n} \mathscr{Y}$  for any n by Lemma 3.1.21. Hence  $\operatorname{Map}(\tau_{\geq (p+q,q)} \mathscr{X}, \mathscr{Y}) \simeq \lim_{n} \operatorname{Map}(\tau_{\geq (p+q,q)} \mathscr{X}, \tau_{\leq n} \mathscr{Y})$ . However for i large enough,  $\operatorname{Map}(\tau_{\geq (p+q,q)} \mathscr{X}, \tau_{\leq n} \mathscr{Y}) \simeq \operatorname{Map}(\tau_{\geq (p+q,q)} \mathscr{X}_i, \tau_{\leq n} \mathscr{Y}) = 0$ , as needed.

**Theorem 4.1.10** (Refined Postnikov tower). Suppose that  $\mathscr{X} \in O(S^{q,q})$  is simple. We have  $\tau_{\geq (q,q)}\tau_{\leq i-1}\mathscr{X} \simeq L^{i+q,q}\mathscr{X}$ . Applying  $\tau_{\geq (q,q)}$  to the fiber sequences  $\mathscr{X}_{\leq i} \to \mathscr{X}_{\leq i} \to K(\pi_i \mathscr{X}, i+1)$  yields fiber sequences

$$\mathbf{L}^{i+1+q,q}\mathscr{X} \to \mathbf{L}^{i+q,q}\mathscr{X} \to \tau_{\geq (i+1+q,q)} K(\boldsymbol{\pi}_i \mathscr{X}, i+1).$$

We have  $L^{q,q} \mathscr{X} = *$  and  $\lim_{n} L^{q+n,q} \mathscr{X} = \mathscr{X}$ .

*Proof.* We apply Construction 4.1.8 with  $n_i = i$ ,  $\mathscr{X}_i = \mathscr{X}_{\leq i}$ . The fiber sequences

$$\tau_{\geq (j+q,q)} K(\pi_j \mathscr{X}, j) \to \tau_{\geq (1+q,q)} \mathscr{X}_{j+1} \to \tau_{\geq (1+q,q)} \mathscr{X}_j$$

show that for  $j \geq i$  the map  $\tau_{\geq(1+q,q)} \mathscr{X}_{j+1} \to \tau_{\geq(1+q,q)} \mathscr{X}_{j}$  is an  $S^{i+q,q}$ -equivalence (i.e.  $\tau_{\geq(j+q,q)} K(\pi_{j} \mathscr{X}, j) \in O(S^{i+q,q})$ ), whereas for j < i the space  $\tau_{\geq(1+q,q)} \mathscr{X}_{j+1}$  is  $S^{i+q,q}$ -null if  $\tau_{\geq(1+q,q)} \mathscr{X}_{j}$  is (i.e.  $\tau_{\geq(j+q,q)} K(\pi_{j} \mathscr{X}, j)$  is  $S^{i+q,q}$ -null). Thus  $\mathscr{X} \to \tau_{\geq(1+q,q)} \mathscr{X}_{i}$  is an  $S^{i+q,q}$ -equivalence to an  $S^{i+q,q}$ -null space, as needed.

Now we prove the remaining statements. The fiber sequence  $\mathscr{X}_{\leq i} \to \mathscr{X}_{< i} \to K(\pi_i \mathscr{X}, i+1)$  is preserved by  $L^{q,q}$  by Proposition 3.1.29, and hence so does  $\tau_{\geq (q,q)}$ . We have  $\tau_{\geq (q,q)}K(\pi_i \mathscr{X}, i+1) \simeq \tau_{\geq (i+1+q,q)}K(\pi_i \mathscr{X}, i+1)$  by Lemma 4.1.7. Since  $\mathscr{X} \in O(S^{q,q})$  we have  $L^{q,q} \mathscr{X} = *$ . Finally  $\lim_n L^{q+n,q} \mathscr{X} = \mathscr{X}$  by Lemma 3.1.18.

#### **Refined Whitehead towers**

Recall the definition of  $\mathbb{A}^1$ -solvable group and solvable motivic space from Paragraph 2.1.16 and 2.1.17. Recall also from Proposition 2.1.18 that a sovlable motivic space has a functorially defined Whitehead tower. We now show how to modify this tower in a fashion that preserves weak cellular classes.

**Construction 4.1.11.** Let  $\mathscr{X} \in \operatorname{Spc}(k)_*$  be a solvable motivic space and suppose further that  $\mathscr{X} \in O(S^{q,q})$ . Let  $n_i$ ,  $\mathbf{A}_i$  be the integers and sheaves from Proposition 2.1.18. We can inductively and uniquely build commutative diagrams of fiber sequences

such that the following statements hold:

1. 
$$\mathscr{X}\langle 0,q \rangle = \mathscr{X}$$
  
2.  $\mathscr{X}\langle i,q \rangle \in O(S^{n_i+q,q}),$   
3.  $\Omega^{q,q}\mathscr{X}\langle i,q \rangle \simeq \Omega^{q,q}\mathscr{X}\langle i \rangle.$ 

*Proof.* Suppose we have obtained  $\mathscr{X}\langle i,q\rangle$  and proved (2), (3) for it. (Note that this is trivial for i = 0.) Since  $\mathscr{X}\langle i,q\rangle \in O(S^{n_i+q,q})$ , the composite

$$\mathscr{X}\langle i,q\rangle \to \mathscr{X}\langle i\rangle \to K(\mathbf{A}_i,n_i)$$

factors uniquely through  $\tau_{\geq(n_i+q,q)}K(\mathbf{A}_i,n_i)$ . We thus uniquely obtain the space  $\mathscr{X}\langle i+1,q\rangle$ . The fiber sequence

$$\Omega \tau_{>(n_i+q,q)} K(\mathbf{A}_i, n_i) \to \mathscr{X} \langle i+1, q \rangle \to \mathscr{X} \langle i, q \rangle$$

shows (via Example 4.1.5 and Lemma 3.1.14) that  $\mathscr{X}\langle i+1,q\rangle \in O(S^{n_i-1+q,q})$ . By construction and induction we have  $\Omega^{q,q}\mathscr{X}\langle i+1,q\rangle \simeq \Omega^{q,q}\mathscr{X}\langle i+1\rangle$ , which is  $n_{i+1}$ -connective. We claim that also  $\mathscr{X}\langle i+1,q\rangle$  is connected; it will follow via Corollary 3.1.25 that  $\mathscr{X}\langle i+1,q\rangle \in O(S^{n_{i+1}+q,q})$ , as needed.

Proving that  $\mathscr{X}\langle i+1,q\rangle$  is connected is only an issue when  $n_i = 1$ . We must show that  $\pi_1 \mathscr{X}\langle i,q\rangle \rightarrow \pi_1 \tau_{>(1+q,q)} K(\mathbf{A}_i,1)$  is surjective. Consider the fiber sequence

$$F \to \Sigma_{S^1}^{\infty} \mathscr{X} \langle i, q \rangle \to f_q \Sigma H \mathbf{A}_i \simeq \tau_{\geq (1,q)} \Sigma H \mathbf{A}_i \in \mathrm{SH}^{S^1}(k).$$

Since  $\pi_1 \mathscr{X}\langle i,q\rangle \to \pi_1 \Sigma_{S^1}^{\infty} \mathscr{X}\langle i,q\rangle \simeq (\pi_1 \mathscr{X}\langle i,q\rangle)_{\mathbb{A}^1}^{ab}$  is surjective [ABH22, Lemma 4.1(2)], it suffices to prove that  $F \in \mathrm{SH}^{S^1}(k)_{\geq 1}$ . But also  $F \in \mathrm{SH}^{S^1}(k)(q)_{\geq 0}$ , and so it is enough to prove that

 $\pi_1(\Sigma_{S^1}^{\infty} \mathscr{X}\langle i,q\rangle)_{-q} \to \pi_1(f_q \Sigma H \mathbf{A}_i)_{-q} \text{ is surjective [BY20, Lemma 6.2(3)]. Since } \pi_1 \mathscr{X}\langle i,q\rangle \to \pi_1 \Sigma_{S^1}^{\infty} \mathscr{X}\langle i,q\rangle$ is surjective so is  $\pi_1(\mathscr{X}\langle i,q\rangle)_{-q} \to \pi_1(\Sigma_{S^1}^{\infty} \mathscr{X}\langle i,q\rangle)_{-q}$ . Furthermore  $\pi_1(f_q \Sigma H \mathbf{A}_i)_{-q} \simeq (\mathbf{A}_i)_{-q}$  and  $\pi_1(\mathscr{X}\langle i,q\rangle)_{-q} \simeq \pi_1(\mathscr{X}\langle i\rangle)_{-q}$ . It is thus enough to prove that  $\pi_1(\mathscr{X}\langle i\rangle) \to \mathbf{A}_i$  is surjective, which holds since  $\mathscr{X}\langle i+1\rangle$  is connected.  $\Box$ 

*Remark* 4.1.12. The construction shows that each  $\mathscr{X}\langle i,q\rangle$  is a solvable motivic space. Similarly if  $\mathscr{X}$  is a nilpotent motivic space, then an induction argument making repeated appeal to [AFH22, Corollary 3.3.7] together with the fact that the spaces  $\tau_{\geq (n_i+q,q)}K(\mathbf{A}_i,n_i)$  are nilpotent (being grouplike commutative monoids) implies that each  $\mathscr{X}\langle i,q\rangle$  is a nilpotent motivic space.

**Amplification 4.1.13.** Let k be a perfect field and  $\mathscr{X} \in O(S^{p,q})$  a solvable motivic space. Assume  $q \ge 2$ . If  $\mathscr{X}\langle i,q \rangle$  is the tower from Construction 4.1.11, then there exist fiber sequences

$$\mathscr{X}\langle i+1,q\rangle \longrightarrow \mathscr{X}\langle i,q\rangle \longrightarrow \Omega^{\infty}\Sigma^{n_i+q,q}\mathscr{B}_i$$

for some  $\mathscr{B}_i \in SH(k)^{eff \heartsuit}$ . The connectivity of  $\mathscr{X}\langle i,q \rangle$  as well as the sequence  $n_i$  tends to  $\infty$  with i.

*Proof.* The only thing that remains to be established is that  $\tau_{\geq (n_i+q,q)}K(\mathbf{A}_i, n_i) \simeq \Omega^{\infty}\Sigma^{n_i+q,q}\mathscr{B}_i$  for some  $\mathscr{B}_i$  as claimed. This fact follows from Theorem 2.2.29.

#### Digression: weak vs. strong cellularity

**Corollary 4.1.14.** Let  $f : \mathscr{X} \to \mathscr{Y}$ , with  $\mathscr{X}, Y \in O(S^{q,q})$  both  $\mathbb{A}^1$ -solvable. Then f is an equivalence if and only if the induced map  $\pi_i(\mathscr{X})_{-q} \to \pi_i(\mathscr{Y})_{-q}$  is an isomorphism for every  $i \geq 2$ , and also the induced maps

$$(\pi_1(\mathscr{X})^{(j-1)}_{\mathbb{A}^1}/\pi_1(\mathscr{X})^{(j)}_{\mathbb{A}^1})_{-q} \to (\pi_1(\mathscr{Y})^{(j-1)}_{\mathbb{A}^1}/\pi_1(\mathscr{Y})^{(j)}_{\mathbb{A}^1})_{-q}$$

are isomorphisms for every  $j \ge 1$ .

*Proof.* Let  $\mathscr{X}\langle \bullet, q \rangle, \mathscr{Y}\langle \bullet, q \rangle$  be functorial refined Whitehead towers for  $\mathscr{X}, \mathscr{Y}$ , with bases  $(\mathbf{A}_i, n_i)$  and  $(\mathbf{B}_i, n_i)$ . We can assume that  $n_i = 1$  for  $i \leq N + 1$  (a common bound for the derived lengths of  $\pi_1 \mathscr{X}, \pi_1 \mathscr{Y}$ ) and  $N_{N+i} = i$ . Also  $\mathbf{A}_{N+i} = \pi_i \mathscr{X}$  for i > 1, and similarly for  $\mathscr{Y}$ . Finally  $\mathbf{A}_i \simeq \pi_1(\mathscr{X})^{(i-1)}_{\mathbb{A}^1}/\pi_1(\mathscr{X})^{(i)}_{\mathbb{A}^1}$  for  $i \leq N + 1$ , and similarly for  $\mathscr{Y}$ . Now by construction, the map f is an equivalence if and only if so are the induced maps

$$\tau_{\geq (n_i+q,q)} K(\mathbf{A}_i, n_i) \to \tau_{\geq (n_i+q,q)} K(\mathbf{B}_i, n_i)$$

are equivalences. Using [BY20, Lemma 6.1(1)], this just means that  $(\mathbf{A}_i)_{-q} \xrightarrow{\simeq} (\mathbf{B}_i)_{-q}$ . This is the desired result.

**Proposition 4.1.15.** Let  $\mathscr{X} \in O(S^{p+q,q})$  with  $p \ge 2$ . Then  $\mathscr{X}$  is strongly  $S^{p+q,q}$ -cellular, that is,  $\mathscr{X}$  lies in the subcategory of  $\operatorname{Spc}(k)$  generated under colimits by  $S^{p+q,q} \wedge \operatorname{Spc}(k)_*$ .

*Proof.* We may assume  $\mathscr{X} \in \operatorname{Spc}(k)_*$ . We shall construct  $\mathscr{X}_1 \to \mathscr{X}_2 \to \cdots \to \mathscr{X}$  such that each  $\mathscr{X}_i$  can be built using colimits of the asserted form, and such that the induced map  $\pi_n(\Omega^{q,q}\mathscr{X}_i) \to \pi_n(\Omega^{q,q}\mathscr{X})$  is an isomorphism for  $n \leq i$ . Using Corollary 4.1.14 we deduce that  $\mathscr{X} \simeq \operatorname{colim}_i \mathscr{X}_i$ , concluding the proof.

It remains to construct the  $\mathscr{X}_i$ . Since  $\mathscr{X}$  is simply connected, we may put  $\mathscr{X}_1 = *$ ; in fact we may put  $\mathscr{X}_i = *$  for i < p. Now inductively suppose  $\mathscr{X}_{i-1}$  has been constructed; we shall construct  $\mathscr{X}_i$ . By construction we may assume  $i \ge p$ . First set

$$\mathscr{Y} = \mathscr{X}_{i-1} \lor \bigvee_{X_+ \land S^{i+q,q} \to \mathscr{X}} X_+ \land S^{i+q,q},$$

where the sum is over  $X \in \text{Sm}_k$  with a map as indicated. Then  $\pi_n(\mathscr{Y}) \simeq \pi_n(\mathscr{X}_{i-1})$  for n < i, and hence  $\pi_n(\Omega^{q,q}\mathscr{Y}) \simeq \pi_n(\Omega^{q,q}\mathscr{X}_{i-1})$  for the same n. Moreover by construction,  $\pi_i(\Omega^{q,q}\mathscr{Y}) \to \pi_i(\Omega^{q,q}\mathscr{X})$  is surjective. Let  $\mathscr{F} = \text{fib}(\mathscr{Y} \to \mathscr{X})$ . Set

$$\mathscr{K} = \bigvee_{X_+ \wedge S^{i+q,q} \to \mathscr{F}} X_+ \wedge S^{i+q,q}$$

and form the pushout

$$\begin{array}{cccc} \mathscr{K} & \longrightarrow & \mathscr{Y} \\ & & & \downarrow \\ & & & \downarrow \\ * & \longrightarrow & \mathscr{X}_i. \end{array}$$

We get

$$\pi_n(\mathscr{X}_i) \simeq \pi_n(\mathscr{Y}) \simeq \pi_n(\mathscr{X}_{i-1})$$

for n < i and

 $\pi_i(\mathscr{X}_i) \simeq \pi_i(\mathscr{Y})/\pi_i(K).$ 

It follows by induction (and §2.1.10) that  $\pi_n(\Omega^{q,q}\mathscr{X}_i) \simeq \pi_n(\Omega^{q,q}\mathscr{X})$ , for n < i. By construction,

$$\pi_i(\Omega^{q,q}\mathscr{Y})/\pi_i(\Omega^{q,q}\mathscr{F})\simeq\pi_i(\Omega^{q,q}\mathscr{X})$$

By construction,  $\pi_i(\Omega^{q,q}\mathscr{K}) \to \pi_i(\Omega^{q,q}\mathscr{F})$  is surjective. Using the relationship between  $\Omega^{q,q}$  and contraction, as exactness of this operation (see §2.1.10), we conclude that

$$\pi_i(\Omega^{q,q}\mathscr{X}_i) \simeq \pi_i(\mathscr{X}_i)_{-q} \simeq \pi_i(\mathscr{Y})_{-q} / \pi_i(\mathscr{K})_{-q} \simeq \pi_i(\Omega^{q,q}\mathscr{Y}) / \pi_i(\Omega^{q,q}\mathscr{F}) \simeq \pi_i(\Omega^{q,q}\mathscr{X}).$$

This concludes the proof.

## 4.2 Unstable cellularity of loop spaces and fibers

Using the weakly cellular Whitehead tower constructed above we can deduce consequences about weak cellularity of loop spaces.

**Proposition 4.2.1.** Assume  $\mathscr{X} \in O(S^{p,q})$  is a solvable motivic space.

- 1. The space  $\Omega \mathscr{X}$  lies in  $O(S^{p-1,q})$  (provided p > 0).
- 2. The space  $\Omega^{1,1} \mathscr{X}$  lies in  $O(S^{p-1,q-1})$  (provided q > 0).
- 3. The space  $\Omega^{2,1} \mathscr{X}$  lies in  $O(S^{p-2,q-1})$  (provided p > 1, q > 0).

*Proof.* The functors  $\Omega$ ,  $\Omega^{1,1}$  and  $\Omega^{2,1}$  preserve fiber sequences. Consider the weakly cellular Whitehead tower of Construction 4.1.11. Looping, we obtain fiber sequences of the form

$$\Omega^{\infty}_{S^1}\Omega^2\mathscr{B}_i \longrightarrow \Omega\mathscr{X}\langle i+1,q \rangle \longrightarrow \Omega\mathscr{X}\langle i,q \rangle.$$

By assumption  $\mathscr{B}_i \in O(S^{n_i+q,q})$ , so the space  $\Omega^2 \Omega_{S^1}^{\infty} \mathscr{B}_i$  lies in  $O(S^{n_i-2+q,q})$  by appeal to Proposition 3.2.10(1) and Proposition 3.2.11. Therefore, Proposition 3.1.22(1) implies that  $\Omega \mathscr{X}\langle i+1,q\rangle \rightarrow \Omega \mathscr{X}\langle i,q\rangle$  is an  $S^{p-1,q}$ -equivalence for *i* sufficiently large, say  $i \geq N$ . Since  $L^{p-1,q}$  preserves connectivity (Corollary 3.1.25), we find that the space  $L^{p-1,q}\Omega \mathscr{X}\langle N,q\rangle \simeq L^{p,q}\Omega \mathscr{X}\langle N,q\rangle$  is  $\infty$ -connected, and hence vanishes. Next consider the fiber sequences

$$\Omega \mathscr{X} \langle i+1, q \rangle \longrightarrow \Omega \mathscr{X} \langle i, q \rangle \longrightarrow \Omega_{S^1}^\infty \Omega \mathscr{B}_i.$$

By descending induction (starting with i = N - 1) we see using Proposition 3.1.22 that  $\Omega \mathscr{X} \langle i, q \rangle \in O(S^{p-1,q})$ , as needed.

The other statements are deduced in a completely analogous fashion, appealing to the other parts of Proposition 3.2.10. (Note in particular that (2) and (3) are trivially true if  $q \in \{0, 1\}$ , so we may assume that  $q \ge 2$  and hence use the version of the Whitehead tower with layers infinite  $\mathbb{P}^1$ -loop spaces, i.e., Amplification 4.1.13.)

Remark 4.2.2. Suppose given fiber sequences

$$\mathscr{X}_{i+1} \longrightarrow \mathscr{X}_i \longrightarrow \mathscr{B}_i$$

for  $i \ge 0$ , with all spaces connected,  $\mathscr{B}_i \in O(S^{p,q})$  for all  $i, \Omega \mathscr{B}_i \in O(S^{p,q})$  for i sufficiently large, and the connectivity of the  $\mathscr{X}_i$  tending to infinity with i. Then the argument in the proof of Proposition 4.2.1 shows that each  $\mathscr{X}_i \in O(S^{p,q})$ .

**Theorem 4.2.3.** If  $f : \mathscr{X} \to \mathscr{Y}$  is a map of pointed, connected spaces with  $\mathscr{X} \in O(S^{p-1,q}), \mathscr{Y} \in O(S^{p,q})$  and  $p-q \geq 2$ , then  $\operatorname{fib}(f) \in O(S^{p-1,q})$ .

*Proof.* Consider the rotated fiber sequence  $\Omega \mathscr{Y} \to \operatorname{fib}(f) \to \mathscr{X}$ . Since  $\Omega \mathscr{Y} \in O(S^{p-1,q})$  by Proposition 4.2.1, we deduce that  $L^{p-1,q}\operatorname{fib}(f) \simeq L^{p-1,q} \mathscr{X}$  (Proposition 3.1.22). This vanishes by assumption.

### 4.3 Weak cellularity: comparing unstable and stable

We can also refine the motivic Blakers–Massey theorem in a fashion that takes weak cellular estimates into account.

**Theorem 4.3.1.** Suppose  $\mathscr{F} \xrightarrow{\iota} \mathscr{E} \xrightarrow{f} \mathscr{B}$  is a fiber sequence of pointed connected motivic spaces.

- 1. If  $\mathscr{F} \in O(S^{p,q})$  and  $\mathscr{B} \in O(S^{p',q'})$  for  $p'-q' \ge 1$ , then  $\operatorname{fib}(\operatorname{cof}(\iota) \to \mathscr{B}) \in O(S^{p+p',q+q'})$ .
- 2. If  $\mathscr{F} \in O(S^{p,q})$  and  $\mathscr{B}$  is  $O(S^{p',q'})$  for  $p'-q' \ge 1$ , then the canonical map  $\Sigma \mathscr{F} \to \operatorname{cof}(f)$  is an  $S^{p+p',q+q'}$ -equivalence with cofiber in  $O(S^{p+p'+1,q+q'})$ .

*Proof.* For the first statement, recall that by [ABH22, Lemma 3.2(1)] we know that  $\operatorname{fbb}(\operatorname{cof}(\iota) \to \mathscr{B})$  is equivalent to  $\Sigma \mathscr{F} \land \Omega \mathscr{B}$ . Since  $\mathscr{B} \in O(S^{p',q'})$ ,  $p' - q' \ge 1$ , we conclude that  $\Omega \mathscr{B} \in O(S^{p'-1,q})$  by Proposition 4.2.1(1). On the other hand,  $\Sigma \mathscr{F} \in O(S^{p+1,q})$  by Theorem 3.1.19. It follows that  $\Sigma \mathscr{F} \land \Omega \mathscr{B} \in O(S^{p+p',q+q'})$  by another application of Theorem 3.1.19.

For the second statement, note that Proposition 3.1.22 in conjunction with the fiber sequence

$$\Sigma \mathscr{F} \land \Omega \mathscr{B} \longrightarrow \operatorname{cof}(\iota) \longrightarrow \mathscr{B}$$

together with the connectivity assertion of the first point implies that  $cof(\iota) \to \mathscr{B}$  is an  $S^{p+p',q+q'}$ -equivalence. Thus,  $cof(cof(\iota) \to \mathscr{B}) \in O(S^{p+p',q+q'})$  by Lemma 3.1.6.

Writing f as the composite  $\mathscr{E} \to cof(\iota) \to \mathscr{B}$  yields a cofiber sequence [ABH22, Lemma 3.2(2)] of the form

$$\Sigma \mathscr{F} \longrightarrow \operatorname{cof}(f) \longrightarrow \Sigma \operatorname{cof}(\operatorname{cof}(\iota) \to \mathscr{B})$$

Since  $\Sigma \mathscr{F}$  is 1-connected and [ABH22, Proposition 3.3] implies that  $\operatorname{cof}(f)$  is 1-connected as well, we conclude that  $\Sigma \mathscr{F} \to \operatorname{cof}(f)$  is an  $S^{p+p',q+q'}$ -equivalence by Theorem 3.2.12 (having  $S^1$ -stable fiber  $\operatorname{cof}(\operatorname{cof}(\iota) \to \mathscr{B}) \in O(S^{p+p',q+q'})$ ), as claimed.

**Corollary 4.3.2.** Suppose  $f : \mathscr{X} \to \mathscr{Y}$  is a map of nilpotent motivic spaces with  $\mathscr{Y}$  simply connected. 1. If fib $(f) \in O(S^{p,q})$ , then  $\operatorname{cof}(f) \in O(S^{p+1,q})$ . 2. If  $cof(f) \in O(S^{p+1,q}), \mathcal{Y} \in O(S^{3,1}), fib(f) \in O(S^{2,0}), then fib(f) \in O(S^{p,q}).$ 

In particular, if  $f : \mathscr{X} \to \mathscr{Y}$  is a map of 1-connected spaces,  $\mathscr{Y} \in O(S^{3,1})$  and  $\operatorname{fib}(f) \in O(S^{2,0})$ , then  $\operatorname{fib}(f) \in O(S^{p,q})$  if and only if  $\operatorname{cof}(f) \in O(S^{p+1,q})$ .

*Proof.* Since  $\operatorname{fib}(f) \in O(S^{p,q})$ ,  $\Sigma \operatorname{fib}(f) \in O(S^{p+1,q})$ . In that case, Theorem 4.3.1(2) then implies that  $\Sigma \operatorname{fib}(f) \to \operatorname{cof}(f)$  is an  $S^{p+2,q}$ -equivalence, so  $\operatorname{cof}(f) \in O(S^{p+1,q})$  as well.

For the second point, suppose  $\mathscr{F} \in O(S^{r,r'})$ ; we know this holds for r = 2, r' = 0 by assumption. Since  $\mathscr{B} \in O(S^{3,1})$  is simply connected, Theorem 4.3.1(2) we conclude that  $\Sigma \operatorname{fib}(f) \to \operatorname{cof}(f)$  is an  $S^{r+3,r'+1}$ -equivalence. Since  $\operatorname{cof}(f) \in O(S^{p+1,q})$  by assumption, we conclude that  $\Sigma \operatorname{fib}(f) \to \operatorname{cof}(f)$  is more highly connected. Since  $\mathscr{F}$  is simply connected by assumption, Theorem 3.2.12 implies  $\operatorname{fib}(f)$  has higher connectivity as well. By induction, the result follows.

We now refine F. Morel's Freudenthal suspension theorem [Mor12, Theorem 6.61] by providing a weak cellular estimate on the fiber of the unit map of the simplicial loops-suspension adjunction.

**Proposition 4.3.3.** If k is a field, and  $\mathscr{X} \in O(S^{p,q})$  with p-q > 1 (and  $q \ge 0$ ), then  $\operatorname{fib}(\mathscr{X} \to \Omega \Sigma \mathscr{X}) \in O(S^{2p-1,2q})$ .

*Proof.* The space  $\Omega \Sigma \mathscr{X} \in O(S^{p,q})$  by appeal to Example 3.3.2. Write *s* for the unit map  $\mathscr{X} \to \Omega \Sigma \mathscr{X}$ . Note that  $\Sigma \Omega \Sigma \mathscr{X}$  splits as  $\bigvee_{i\geq 0} \Sigma \mathscr{X}^{\wedge i}$  by the Hilton–Milnor splitting [WW19, §5] or [DH21, Corollary 2.11]. Thus,  $\Sigma \operatorname{cof}(s) \simeq \operatorname{cof}(\Sigma s) \simeq \bigvee_{i\geq 2} \Sigma \mathscr{X}^{\wedge i}$  and thus lies in  $O(S^{2p+1,2q})$  by appeal to Theorem 3.1.19.

Note that fib(s) is simply connected under the hypotheses, whence by Corollary 4.3.2 it will be enough to show that  $cof(s) \in O(S^{2p,2q})$ . Since cof(s) is also simply connected, that follows from what we just proved about  $\Sigma cof(s)$ .

#### Weak cellularity and *R*-localization

Assume  $R \subset \mathbb{Q}$  is a subring. Another consequence of the weakly cellular version of the Whitehead tower is the compatibility of *R*-localization and weak cellularity.

**Proposition 4.3.4.** Assume  $\mathscr{X} \in \operatorname{Spc}(k)_*$  is a nilpotent motivic space lying in  $O(S^{p,q})$  for some integers  $p \ge q \ge 0$ . If  $R \subset \mathbb{Q}$  is a subring, then  $\mathscr{X}_R$  lies in  $O(S^{p,q})$  as well.

*Proof.* Consider the refined Whitehead tower of  $\mathscr{X}$  from Construction 4.1.11, consisting of fiber sequences

$$\mathscr{X}\langle i+1,q\rangle \to \mathscr{X}\langle i,q\rangle \to \mathscr{B}_i = \tau_{>(n_i+q,q)}K(\mathbf{A}_i,n_i).$$

Since all involved spaces are nilpotent 4.1.12, these fiber sequences are preserved by *R*-localization [AFH22, Theorem 4.3.11]. Appealing to Remark 4.2.2, it remains to prove that  $(\mathscr{B}_i)_R \in O(S^{p,q})$  for all *i*, and  $(\Omega \mathscr{B}_i)_R \in O(S^{p,q})$  for *i* sufficiently large. This is true because  $\mathscr{B}_i = \Omega_{S^1}^{\infty} B_i$  for some  $B_i \in \mathrm{SH}^{S^1}(k)(q)_{\geq p-q}$ ,  $\Omega_{S^1}^{\infty}$  commutes with *R*-localization, and  $\mathrm{SH}^{S^1}(k)(q)_{\geq p-q}$  is stable under *R*-localization, being stable under colimits.

## 4.4 Further examples

*Example* 4.4.1. One consequence of Proposition 4.2.1 is the following weak cellular estimate, generalizing Example 3.3.2: if  $\mathscr{X}$  is a pointed, connected space that lies in  $O(S^{p,q})$ , then  $\Omega^{2,1}\Sigma^{2,1}\mathscr{X} \in O(S^{p,q})$ . Since  $\mathscr{X}$  is pointed and connected, we conclude that  $\Sigma^{2,1}\mathscr{X}$  is in particular 1-connected and thus nilpotent and furthermore lies in  $O(S^{p+2,q+1})$ . In particular, q + 1 > 0 and therefore, the cellularity estimate follows immediately from Proposition 4.2.1(3). More generally, it follows that  $\Omega^{2n,n}\Sigma^{2n,n}\mathscr{X} \in O(S^{p,q})$  under the above hypotheses as well, and also that  $Q(\mathscr{X}) \in O(S^{p,q})$ .

*Example* 4.4.2 (Motivic connective covers of BGL). The space  $\mathbb{Z} \times BGL$  is the  $\mathbb{P}^1$ -infinite loop space of the spectrum KGL. For any integer  $n \ge 1$ , we set

$$BGL\langle 2n,n\rangle := \Omega^{\infty}\tau_{>(2n,n)}\mathbf{KGL}.$$

In particular, observe that  $BGL(2n, n) \in O(S^{2n,n})$  by appeal to Proposition 3.2.11.

Recall from, e.g., [AF14a, §3.5] that Suslin described a morphism  $S^{2n-1,n} \to GL$  via an explicit matrix; this matrix is the clutching function of a morphism  $S^{2n,n} \to BGL$  by [ADF17, Theorem 4.3.4] that stabilizes to the unit map  $1 \to \text{KGL}$  [Sus82, Theorem 2.3]. Observe that the map  $S^{2n,n} \to BGL$  factors through  $BGL\langle 2n,n \rangle$ .

*Example* 4.4.3 (Motivic connective covers of BGL II). Continuing the discussion of Example 4.4.2, consider the composite map  $S^{2n,n} \to BGL \to K(\mathbb{Z}(n), 2n)$ , where the second map is given by the *n*-th Chern class. Such a map is uniquely determined by a morphism  $\pi_{n,0}(S^{2n,n}) = \mathbf{K}_n^{MW} \to \mathbf{K}_n^M$  (see the discussion of [ADF17, Proof of Theorem 4.2.3]) and thus is (n-1)! times a generator by [Sus82, Theorem 5.2].

The equivalence  $BGL_n \simeq Gr_n \simeq \operatorname{colim}_N Gr_{n,N}$  in conjunction with Example 3.3.7 shows that  $BGL_n$  and thus  $BGL \in O(S^{2,1})$ . We may identify  $BGL = BGL\langle 2, 1 \rangle$  and  $BSL = BGL\langle 4, 2 \rangle$ , the latter using the fact that BSL is the homotopy fiber of  $c_1 : BGL \to K(\mathbb{Z}(1), 2)$ .

Now, assume that we work over a field k that has characteristic 0 and consider the induced map  $c_2: BSL \to K(\mathbb{Z}(2), 4)$ . We claim that

$$BGL\langle 6,3\rangle \simeq \operatorname{fib}(c_2:BSL \to K(\mathbb{Z}(2),4)).$$

Indeed, the space  $K(\mathbb{Z}(2),4)$  is  $S^{6,3}$ -null by Example 3.3.13 and lies in  $O(S^{4,2})$  by Example 3.3.9. Putting these facts together with Proposition 3.1.29 we conclude that  $BGL\langle 6,3 \rangle \in O(S^{6,3})$  and the map  $BGL\langle 6,3 \rangle \to BSL$  is an  $S^{6,3}$ -equivalence. We view the space  $BGL\langle 6,3 \rangle$  as a motivic analog of the space  $BU\langle 6 \rangle$ .

*Example* 4.4.4. Assume G is a split semi-simple k-group scheme. In that case, we may consider the étale classifying space  $B_{\text{ét}}$  G. In general, this space fails to be  $\mathbb{A}^1$ -connected, but there is a canonical identification

$$\mathrm{H}^{1}_{\mathrm{\acute{e}t}}(\mathrm{G}) \longrightarrow \boldsymbol{\pi}_{0}(B_{\mathrm{\acute{e}t}}\mathbf{G}),$$

where the source is the Nisnevich sheafification of the presheaf  $U \mapsto H^1_{\acute{e}t}(U, G)$  [?]. The connected component of the base-point corresponding to the trivial torsor therefore coincides with  $B_{\text{Nis}}$  G. If G is a split semi-simple k-group scheme, then there is a unique simply connected convering group  $\tilde{G} \to G$ where the covering map is a central isogeny. By functoriality, there is an induced map  $B_{\text{Nis}}\tilde{G} \to B_{\text{Nis}}$  G. By Proposition 3.3.11, we conclude that  $B_{\text{Nis}}\tilde{G}$  lies in  $O(S^{4,2})$  if G has factors only of type A, B, C, D or  $G_2$ . In that case, we may realize  $B_{\text{Nis}}\tilde{G}$  as a connective cover of  $B_{\acute{e}t}$  G.

# 5 Equivariant geometry of symmetric powers

In this section, we analyze various facts about the geometry of symmetric powers. By means of Voevodsky's motivic Dold-Thom theorem [Voe10b, Theorem 3.7 and 3.11], the motivic Eilenberg–Mac Lane space  $K(\mathbb{Z}(n), 2n)$  admits a model in terms of symmetric powers of the motivic spheres  $S^{2n,n}$ . We will then analyze cellularity of symmetric powers.

## 5.1 Motivic homotopy theory and group scheme actions

Assume G is a finite group such that |G| is invertible in k.

5.1.1 (*G*-quasiprojective varieties). A *G*-scheme *X* is called *G*-quasiprojective if it admits an ample *G*-equivariant line bundle. Note that if *G* is finite, then any quasi-projective variety equipped with a *G*-action is *G*-quasiprojective: if  $\mathscr{L}$  is ample, then  $\det(\bigoplus_{a \in G} g^* \mathscr{L})$  is ample and *G*-equivariant.

5.1.2 (*G*-spaces). We extend some of the discussion of Paragraph 2.1.2 to spaces with *G*-action. Write  $\operatorname{Sch}_k^G$  for the category of *G*-quasiprojective schemes and  $\operatorname{Sm}_k^G$  for the subcategory of smooth *G*-quasiprojective varieties. The categories  $\operatorname{Sm}_k^G$  and  $\operatorname{Sch}_k^G$  can be equipped with the *G*-Nisnevich topology; see, e.g., [Hoy17, §3.3] for a definition. If  $\operatorname{S}_k^G$  is either of the categories  $\operatorname{Sm}_k^G$  or  $\operatorname{Sch}_k^G$ , we can consider the  $\infty$ -topos  $\operatorname{P}(\operatorname{S}_k^G)$  of presheaves of spaces. We write  $\operatorname{Shv}_{Nis}^G(\operatorname{Sm}_k)$  for the  $\infty$ -category of *G*-Nisnevich sheaves of spaces on  $\operatorname{Sm}_k^G$  and  $\operatorname{Shv}_{Nis}^G(\operatorname{Sch}_k)$  for the  $\infty$ -category of *G*-Nisnevich sheaves of spaces on  $\operatorname{Sch}_k^G$ .

5.1.3 (motivic *G*-spaces). We write  $\operatorname{Spc}^{G}(k)$  for the full subcategory of  $\operatorname{Shv}_{\operatorname{Nis}}^{G}(\operatorname{Sm}_{k})$  consisting of  $\mathbb{A}^{1}$ -invariant sheaves of spaces [Hoy17, Definition 3.12]. Abusing notation, we write  $\operatorname{L}_{mot}$  for the motivic localization functor in the *G*-equivariant context as well. Once again,  $\operatorname{Spc}^{G}(k)$  is a semi-topos [Hoy17, Proposition 3.15].

5.1.4 (Quotients). If X is a G-quasiprojective k-scheme, then a quotient of X by the action of G exists [SGA03, Expose V, Proposition 1.8] and is known to be a quasiprojective k-scheme. If X is furthermore assumed normal, then X/G is once again normal by the universal property of categorical quotients. The functor of taking the quotient is left adjoint to the functor  $\operatorname{Sch}_k \to \operatorname{Sch}_k^G$  sending a quasi-projective k-scheme X to X equipped with the trivial G-action. As such, the quotient construction preserves colimits. Likewise, the functor of adding a disjoint base-point is left adjoint to the forgetful functor from pointed schemes to schemes. It follows that the functor  $X_+ \to (X/G)_+$  is colimit preserving.

5.1.5 (Quotients and  $\mathbb{A}^1$ -equivalences). By left Kan extending the functor  $X_+ \longrightarrow (X/G)_+$  we obtain  $\operatorname{Quot}_G : \operatorname{P}(\operatorname{Sm}_k^G) \longrightarrow \operatorname{P}(\operatorname{Sch}_k)$  which induces a functor

$$\operatorname{Quot}_G: \operatorname{Spc}^G(k) \longrightarrow \operatorname{Spc}^{ft}_{Nis}(k)$$

(see [Voe10b, Proposition 2.10] or [Del09, Proposition 45 p. 403]).

### 5.2 The motivic Dold-Thom theorem

5.2.1 (Symmetric powers). Assume  $r \ge 0$  is an integer. If X is a smooth scheme, then we may consider  $X^{\times r}$  with the action of  $\Sigma_r$  by permutation of the factors. Passing to sifted cocompletions we obtain  $P_{\Sigma}(Sm_k) \to P_{\Sigma}(Sm_k^G)$ . This functor preserves Nisnevich and motivic equivalences [Voe10b, Theorem 2.5] and hence induces a functor  $Spc(k) \to Spc^G(k)$ . In particular, if  $\mathscr{X}$  is a pointed motivic space, we will write  $\mathscr{X}^{\times r}$  for motivic space with  $\Sigma_r$ -action obtained from the functor just described. We then set

$$\operatorname{Sym}^n \mathscr{X} := \operatorname{Quot}_{S_r}(\mathscr{X}^{\times r})$$

If  $\mathscr{X}$  is a pointed space, then the inclusion  $S_{n-1} \subset S_n$  and the inclusion  $\mathscr{X}^{\times n-1} \to \mathscr{X}^n$  (via the basepoint in the last factor) yield morphisms

$$\operatorname{Sym}^{n-1}\mathscr{X} \longrightarrow \operatorname{Sym}^n\mathscr{X}$$

We define

$$\operatorname{Sym}^{\infty} \mathscr{X} := \operatorname{colim}_n \operatorname{Sym}^n \mathscr{X}$$

which has naturally the structure of a (strictly) commutative monoid.

5.2.2 (Symmetric powers and spectra). For any motivic spaces  $\mathscr{X}$  and  $\mathscr{Y}$ , there is an  $S_r$ -equivariant map  $\mathscr{Y} \times \mathscr{X}^r \to (\mathscr{Y} \times \mathscr{X})^r$  induced by the diagonal. If  $\mathscr{Y}$  and  $\mathscr{X}$  are additionally pointed, the above maps induce morphisms  $\mathscr{Y} \wedge \operatorname{Sym}^r \mathscr{X} \longrightarrow \operatorname{Sym}^r (\mathscr{Y} \wedge \mathscr{X})$ . These morphisms are compatible with the stabilization morphisms  $\operatorname{Sym}^n \mathscr{X} \to \operatorname{Sym}^{n+1} \mathscr{X}$  described in Paragraph 5.2.1 in the evident sense and taking  $\mathscr{Y} = S^{2,1}$  we obtain morphisms

$$s_n: \Sigma^{2,1} \operatorname{Sym}^{\infty} \mathscr{X} \longrightarrow \operatorname{Sym}^{\infty} (\Sigma^{2,1} \mathscr{X}).$$

5.2.3 (Inverting the characteristic exponent). For an integer d we write

$$\operatorname{Sym}^{\infty}[1/d]\mathscr{X} := \operatorname{colim}(\operatorname{Sym}^{\infty}\mathscr{X} \xrightarrow{\times d} \operatorname{Sym}^{\infty}\mathscr{X} \xrightarrow{\times d} \cdots)$$

where  $\times d$  is multiplication defined by the monoid structure. In the sequel, we will take d to be the characteristic exponent of the base-field. This construction is also compatible with stabilization morphism of Paragraph 5.2.1 so there are induced morphisms

$$\Sigma^{2,1} \operatorname{Sym}^{\infty}[1/d] \mathscr{X} \longrightarrow \operatorname{Sym}^{\infty}[1/d](\Sigma^{2,1} \mathscr{X})$$

5.2.4. If  $S_k$  is an admissible category as in Paragraph 2.1.2, then we write  $\operatorname{Cor}(S_k)$  for the category whose objects are those of  $S_k$  but whose morphisms are finite k-correspondences. In the notation of [SV00, p. 36], the morphism between two objects of  $S_k$  in  $\operatorname{Cor}(S_k)$  is given by  $c_{equi}(X \times_k Y/X, 0)$ . This is a subgroup of the finite k-correspondences from X to Y, i.e., the free abelian group generated by closed subschemes  $Z \subset X \times_k Y$  such that  $Z \to X$  is finite and dominates an irreducible component of X. In general it is a proper subgroup [SV00, Example 3.4.7], but coincides with all finite correspondences when X is regular [SV00, Corollary 3.4.6]. The category  $\operatorname{Cor}(S_k)$  is additive with direct sum given by disjoint union and has a symmetric monoidal structure such that the graph functor  $\Gamma : S_k \to \operatorname{Cor}(S_k)$  is symmetric monoidal.

5.2.5 (Motivic Eilenberg–Mac Lane spaces). For U a normal scheme and X quasi-projective, we write  $\operatorname{Cor}_k(U, X)$  (resp.  $\operatorname{Cor}_k^{\operatorname{eff}}(U, X)$ ) for the presheaf of finite correspondences. We set

$$K(\mathbb{Z}(n), 2n) := a_{\text{Nis}} \mathbb{Z}_{tr}(\mathbb{A}^n) / \mathbb{Z}_{tr}(\mathbb{A}^n \setminus 0).$$

The construction just described makes sense more generally, e.g., we may define  $\mathbb{Z}_{tr}(\Sigma^{2n,n}X_+)$  for any smooth scheme X as the Nisnevich sheafification of a  $\mathbb{Z}_{tr}(X \times \mathbb{A}^n)/\mathbb{Z}_{tr}(X \times \mathbb{A}^n \setminus 0)$ .

5.2.6. Consider the model of  $S^{2n,n}$  given by  $\mathbb{A}^n \setminus 0$ , i.e., the Thom space of a trivialized rank *n* vector bundle over Spec *k*. There is a canonical map

$$S^{2n,n} \longrightarrow K(\mathbb{Z}(n), 2n)$$

which induces a morphism of commutative monoids

$$\operatorname{Sym}^{\infty} S^{2n,n} \longrightarrow K(\mathbb{Z}(n), 2n).$$

*Example* 5.2.7. The case n = 1 of the map from the preceding point is particularly simple and we do not have to leave smooth schemes to form the relevant quotient. In that case,  $S^{2,1} \simeq \mathbb{P}^1$  and we have  $\operatorname{Sym}^{\infty}\mathbb{P}^1 \simeq \mathbb{P}^{\infty}$ . There is an  $\mathbb{A}^1$ -equivalence  $\mathbb{P}^{\infty} \simeq B\mathbb{G}_m$  [MV99, §4 Proposition 4.8]. On the other hand, the description of weight 1 motivic cohomology shows that  $\mathbb{Z}_{tr}(\mathbb{A}^1)/\mathbb{Z}_{tr}(\mathbb{A}^1 \setminus 0) \simeq B\mathbb{G}_m$  [SV96, §2]. Putting these observations together, we conclude that the map

$$\operatorname{Sym}^{\infty}S^{2,1} \longrightarrow K(\mathbb{Z}(1),2)$$

is an equivalence.

5.2.8. Let E be a  $\mathbb{P}^1$ -prespectrum, that, is, a sequence of spaces  $E^0, E^1, \dots \in \operatorname{Spc}(k)_*$  together with bonding maps  $\Sigma^{2,1}E^i \to E^{i+1}$ . The assembly maps of Paragraph 5.2.2 supply bonding maps in a  $\mathbb{P}^1$ prespectrum  $\operatorname{Sym}^{\infty}[1/d]E$  with entries  $\operatorname{Sym}^{\infty}[1/d]E_i$ .

**Theorem 5.2.9** (motivic Dold-Thom theorem). Assume k is a perfect field with characteristic exponent c and  $n \ge 1$  is an integer.

1. The map

$$\operatorname{Sym}^{\infty}[1/c]S^{2n,n}|_{\operatorname{Sm}_k} \longrightarrow K(\mathbb{Z}(n),2n)[1/c]$$

is an isomorphism in Spc(k).

2. The assembly morphisms  $e_n : \Sigma^{2,1}K(\mathbb{Z}(n), 2n)[1/c] \to K(\mathbb{Z}(n+1), 2n+2)[1/c]$  are compatible with the equivalence of the first point, i.e., for every n the diagram

$$\begin{split} \Sigma^{2,1} \mathrm{Sym}^{\infty}[1/c] S^{2n,n} &\longrightarrow \Sigma^{2,1} K(\mathbb{Z}(n),2n)[1/c] \\ & \downarrow \\ \mathrm{Sym}^{\infty}[1/c] S^{2n+2,n+1} &\longrightarrow K(\mathbb{Z}(n+1),2n+2)[1/c]. \end{split}$$

commutes.

3. The morphisms above induce an isomorphism in SH(k):

 $\operatorname{Sym}^{\infty}[1/c]\mathbb{1}_k|_{\operatorname{Sm}_k} \longrightarrow H\mathbb{Z}[1/c].$ 

4. More generally for  $X \in \text{Sm}_k$  we have isomorphisms

 $\operatorname{Sym}^{\infty}[1/c]\Sigma^{2n,n}X_{+}|_{\operatorname{Sm}_{k}} \simeq \mathbb{Z}_{tr}(\Sigma^{2n,n}X_{+})[1/c] \in \operatorname{Spc}(k)$ 

and

$$\operatorname{Sym}^{\infty}[1/c]\Sigma^{\infty}X_{+}|_{\operatorname{Sm}_{k}}\longrightarrow H\mathbb{Z}[1/c]\wedge\Sigma^{\infty}X_{+}\in \operatorname{SH}(k)$$

*Proof.* The first statement is a consequence of [Voe10b, Theorem 3.7, Propositions 3.11,3.15] and the definition of the motivic EM space given above. Similarly for the first part of the fourth statement. The third point follows from the second and the discussion of Paragraphs 5.2.2 and 5.2.3. The second half of the fourth point follows similarly, using that  $H\mathbb{Z}[1/c] \wedge \Sigma^{\infty} X_{+}$  is equivalent to the prespectrum with entries  $(\mathbb{Z}_{tr}(X_{+})[1/c], \mathbb{Z}_{tr}(\Sigma^{2,1}X_{+})[1/c], \ldots)$ . (This latter claim follows from the fact that  $\mathbf{DM}(k, \mathbb{Z}[1/c]) \simeq Mod_{H\mathbb{Z}[1/c]}(k)$ , which is proved if c = 1 in [RØ08] and for c > 1 in [HKØ17, Theorem 5.8]).

For the second statement, recall that the assembly morphism in the motivic Eilenberg–Mac Lane spectrum is constructed as the composite

$$S^{2,1} \wedge K(\mathbb{Z}(n), 2n) \longrightarrow K(\mathbb{Z}(1), 2) \wedge K(\mathbb{Z}(n), 2n) \longrightarrow K(\mathbb{Z}(n+1), 2n+2);$$

where, at the level of cycles, the first morphism is induced by the map sending  $U \to S^{2,1}$  to its graph viewed as an element of  $\operatorname{Cor}_k(U, S^{2,1})$  and the second morphism is induced by external product of cycles: using the models we have, this arises from the commutative diagram:

$$\operatorname{Cor}_{k}(U, \mathbb{A}^{1} \setminus 0) \times \operatorname{Cor}_{k}(U, \mathbb{A}^{n} \setminus 0) \longrightarrow \operatorname{Cor}_{k}(U, \mathbb{A}^{n+1} \setminus 0)$$

$$\downarrow$$

$$\operatorname{Cor}_{k}(U, \mathbb{A}^{1}) \times \operatorname{Cor}_{k}(U, \mathbb{A}^{n}) \longrightarrow \operatorname{Cor}_{k}(U, \mathbb{A}^{n+1})$$

where the top map is the composite of the external product with the inclusion  $\mathbb{A}^1 \setminus 0 \times \mathbb{A}^n \setminus 0 \subset \mathbb{A}^{n+1} \setminus 0$ .

Now, the map  $\mathbb{P}^1 \wedge \operatorname{Sym}^d(S^{2n,n}) \longrightarrow \operatorname{Sym}^d(S^{2n+2,n+1})$  is induced by the diagonal map. On the other hand, the identification of Point (1) ultimately rests on [SV96, Theorem 6.8] (via [Voe10b, Proposition 3.5]). This identification tells us that for any finite type scheme X and any connected normal scheme U, the group  $\operatorname{Hom}(U, \sqcup_{d\geq 0}\operatorname{Sym}^d X)$  coincides after inverting c with the free commutative monoid on integral cycles on  $U \times X$  that are finite and surjective over U. The result then follows by comparing definitions: see [SV00, §3.7 p. 59] for external product of cycles.

## 5.3 Cellularity of the cofiber of the assembly map

Suppose  $X \in \text{Sm}_k$ . In this section, we establish the main geometric result, which amounts to a weak cellular estimate on the cofiber of the assembly maps in the spectrum  $H\mathbb{Z} \wedge \Sigma^{\infty} X_+$ . We begin by giving an "explicit" description of the cofiber of the assembly map as a colimit of spaces defined using representation theory of the symmetric group.

After Theorem 5.2.9, the discussion of Paragraph 5.2.2, then identifies the assembly map in  $H\mathbb{Z} \wedge \Sigma^{\infty}X_{+}$  as a colimit of maps of the form

(5.3.1) 
$$\Sigma^{2,1} \operatorname{Sym}^{r}(\Sigma^{2n,n} X_{+}) \longrightarrow \operatorname{Sym}^{r}(\Sigma^{2n+2,n+1} X_{+})$$

in the following terms. Write  $\rho_r$  for the standard representation of the symmetric group  $\Sigma_r$  on a based r-dimensional k-vector space by permutation of the basis vectors. Assuming r is coprime to the characteristic of k, the representation  $\rho_r$  decomposes as  $\mathbf{1} \oplus \bar{\rho}_r$  and we refer to  $\bar{\rho}_r$  as the reduced standard representation. View  $\rho_r^{\oplus n}$  as an  $\Sigma_r$ -equivariant vector bundle over Spec k. We write  $\mathbb{A}[\bar{\rho}_r]$  for the affine space attached to  $\bar{\rho}_r$ , i.e., the spectrum of the symmetric algebra of the dual, viewed as a  $\Sigma_r$ -scheme and  $p : \mathbb{A}[\bar{\rho}_r] \setminus 0 \to \text{Spec } k$  for the  $\Sigma_r$ -equivariant structure map.

**Proposition 5.3.1.** If  $X \in Sm_k$ , then there is a canonical identification

$$\operatorname{cof}(\Sigma^{2,1}\operatorname{Sym}^{r}\Sigma^{2n,n}X_{+} \to \operatorname{Sym}^{r}\Sigma^{2n+2,n+1}X_{+}) \simeq \Sigma^{2n+3,n+1}\operatorname{Quot}_{\Sigma_{r}}(\operatorname{Th}(p^{*}\bar{\rho}_{r}^{\oplus n}) \wedge (X^{\times r})_{+}).$$

*Proof.* The map of (5.3.1) is the map

$$\Sigma^{2,1}\operatorname{Quot}_{\Sigma_r}(\operatorname{Th}(\rho_r^{\oplus n}) \wedge (X^{\times r})_+) \longrightarrow \operatorname{Quot}_{\Sigma_r}(\operatorname{Th}(\rho_r^{\oplus n+1}) \wedge (X^{\times r})_+)$$

induced by the map  $\mathbf{1} \oplus \rho_r^{\oplus n} \to \rho_r \oplus \rho_r^{\oplus n}$  that includes  $\mathbf{1} \subset \rho_r$  in the first factor of the target. This map identifies with a map

$$\Sigma^{2n+2,n+1}\operatorname{Quot}_{\Sigma_r}(\operatorname{Th}(\bar{\rho}_r)^{\oplus n} \wedge (X^{\times r})_+) \longrightarrow \Sigma^{2n+2,n+1}\operatorname{Quot}_{\Sigma_r}(\operatorname{Th}(\bar{\rho}_r)^{\oplus n+1} \wedge (X^{\times r})_+)$$

which is itself obtained by suspension from the map

$$\operatorname{Quot}_{\Sigma_r}(\operatorname{Th}(\bar{\rho}_r)^{\oplus n} \wedge (X^{\times r})_+) \longrightarrow \operatorname{Quot}_{\Sigma_r}(\operatorname{Th}(\bar{\rho}_r)^{\oplus n+1} \wedge (X^{\times r})_+)$$

induced by the inclusion  $\bar{\rho}^{\oplus n} \to \bar{\rho}^{\oplus n+1}$ . In order to identify the cofiber of this map, we simply observe that  $\operatorname{Quot}_{\Sigma_n}$  is a colimit preserving functor and unwind the definitions.

Indeed, first consider the commutative square

$$\begin{split} \mathbb{A}[\bar{\rho}_r^{\oplus n}] \smallsetminus 0 & \longrightarrow \mathbb{A}[\bar{\rho}_r^{\oplus n}] \\ & \downarrow & \downarrow \\ \mathbb{A}[\bar{\rho}_r^{\oplus n+1}] \smallsetminus 0 & \longrightarrow \mathbb{A}[\bar{\rho}_r^{\oplus n}]. \end{split}$$

The map on horizontal cofibers is a map  $\operatorname{Th}(\bar{\rho}_r^{\oplus n}) \to \operatorname{Th}(\bar{\rho}_r^{\oplus n+1})$  that, after smashing with  $(X^{\times r})_+$  and applying  $\operatorname{Quot}_{\Sigma_r}$  is the map in which we are interested in. Observe that the left hand vertical map is  $\Sigma_r$ -equivariantly equivalent to the inclusion

$$(\mathbb{A}[\bar{\rho}_r^{\oplus n}] \smallsetminus 0) \times \mathbb{A}[\bar{\rho}_r] \hookrightarrow \mathbb{A}[\bar{\rho}_r^{\oplus n+1}] \smallsetminus 0,$$

with closed complement  $\Sigma_r$ -equivariantly isomorphic to  $\mathbb{A}[\bar{\rho}_r] \setminus 0$  and normal bundle  $p^* \bar{\rho}_r^{\oplus n}$ . Therefore,  $\Sigma_r$ -equivariant homotopy purity [Hoy17, Theorem 3.23] yields an identification of the cofiber of the left hand vertical map with  $\operatorname{Th}(p^* \bar{\rho}_r^{\oplus n})$ .

With those observations in mind, computing cofibers vertically, then yields a  $\Sigma_r$ -equivariant cofiber sequence of the form

$$\operatorname{Th}(p^*\bar{\rho}_r^{\oplus n}) \longrightarrow * \longrightarrow \operatorname{cof}(\operatorname{Th}(\bar{\rho}_r^{\oplus n}) \to \operatorname{Th}(\bar{\rho}_r^{\oplus n+1})).$$

Shifting the cofiber sequence to the right once, smashing with  $(X^{\times r})_+$  and applying  $\operatorname{Quot}_{\Sigma_r}$ , we therefore obtain an identification

$$\operatorname{cof}(\operatorname{Quot}_{\Sigma_r}(\operatorname{Th}(\bar{\rho}_r)^{\oplus n} \wedge (X^{\times r})_+) \to \operatorname{Quot}_{\Sigma_r}(\operatorname{Th}(\bar{\rho}_r)^{\oplus n+1}) \wedge (X^{\times r})_+) \simeq \Sigma \operatorname{Quot}_{\Sigma_r}(\operatorname{Th}(p^*\bar{\rho}_r^{\oplus n}) \wedge (X^{\times r})_+),$$

and the result follows.

*Example* 5.3.2. For concreteness, let us describe what happens when r = 2 and  $X = \operatorname{Spec} k$ . In that case,  $\bar{\rho}_2$  is the 1-dimensional sign representation of  $\Sigma_2$ . The action of  $\Sigma_2$  on  $\mathbb{G}_m = \mathbb{A}(\bar{\rho}_2) \setminus 0$  is the sign representation, which is free and has quotient  $\mathbb{G}_m$  via the squaring map. Thus, the cofiber in the lemma may be identified as:

$$\Sigma^{2n+3,n+1} \operatorname{Quot}_{\Sigma_2} \operatorname{Th}(p^* \bar{\rho}_2^{\oplus n}).$$

Since the action of  $\Sigma_2$  on  $\mathbb{G}_m$  is free, the bundle  $p^* \bar{\rho}_2^{\oplus n}$  is  $\Sigma_2$ -equivariantly trivial and we conclude the cofiber in question is

$$\mathbb{G}_{m+} \wedge S^{4n+3,2n+1},$$

which lies in  $O(S^{4n+3,2n+1})$ .

Remark 5.3.3. Intuitively speaking, since  $Quot_{\Sigma r}$  is a colimit preserving functor, we expect that it preserves weak cellularity classes as well. We thus expect that the space appearing in Proposition 5.3.1 lies in  $O(S^{4n+3,2n+1})$  as well. Unfortunately, the example above is essentially the only case in which the quotients that appear are smooth schemes. In dealing with singular schemes, we are forced to invoke resolution of singularities. Voeovdsky observed that there is a well-defined  $S^1$ -stable homotopy type for singular schemes, which we review in the next section. At the moment, we content ourselves with producing an equivariant version of a weak cellular estimate that we will eventually use.

We begin by making the following definition. In contrast with Section 3 we write down the cellular class directly instead of as the kernel of a localization since this is enough for our purposes. It seems likely that the more general theory can also be developed.

**Definition 5.3.4.** We will say that a pointed motivic *G*-space  $\mathscr{X}$  lies in  $O^G(S^{p,q})$  if it lies in the subcategory generated under colimits and cofiber extensions by objects of the form  $\Sigma^{p,q}X_+$  for  $X \in \text{Sm}_k^G$ .

Consider the  $\Sigma_r$ -action on  $\mathbb{A}[\bar{\rho}] \leq 0$ . The stabilizers of the action of  $\Sigma_r$  on  $\mathbb{A}[\bar{\rho}_r] \leq 0$  are always *proper* subgroups of  $\Sigma_r$ . We may stratify  $\mathbb{A}[\bar{\rho}] \leq 0$  or equivalently  $\mathbb{A}[\bar{\rho}]$  by stabilizer type; the resulting stratification is frequently called the Luna stratification.

**Proposition 5.3.5** (Luna stratification). Assume k is a perfect field, X is an irreducible smooth affine G-scheme and G a finite group whose order is invertible in k. There exists a decomposition of X into locally closed closed subschemes X(H) indexed by conjugacy classes of subgroups  $H \subset G$  such that the following properties hold:

- 1. all points of X(H) have stabilizer conjugate to H;
- 2. all X(H) are smooth;
- 3. there is a subgroup  $H_0$  (the generic stabilizer) and open dense subscheme  $X(H_0)$ ;
- 4. the closure of X(H) is a union of X(H') where H' runs through the conjugacy classes of subgroups of G containing the conjugacy class of H.

*Proof.* The existence of this stratification is "well-known" and we content ourselves with providing some references. Since |G| is invertible in k, the group scheme G is a linearly reductive étale group scheme. Since k is perfect, it suffices to prove these properties after base-change to an algebraic closure of k. The existence and properties of the stratification as stated are consequences of Luna's étale slice theorem which is established in characteristic 0 in [Lun73]. In positive characteristic the slice theorem holds for actions of linearly reductive finite groups by [BR85]. The existence and properties of the stratification above are described in [Lun73, III.2-4].

**Lemma 5.3.6.** Assume k is a perfect field, G is a finite group whose order is invertible in k, and X is a smooth G-quasi-projective k-scheme. If V is a G-equivariant vector bundle on X such that for every conjugacy class of subgroups  $H \subset G$ ,  $V|_{X(H)^H}$  splits H-equivariantly as  $\mathbf{1}_{X(H)^H}^{\oplus n} \oplus V'_H$ , then  $\operatorname{Th}(V) \in O^G(S^{2n,n})$ .

*Proof.* If W is a smooth G-scheme and V is a G-equivariant vector bundle that splits G-equivariantly as  $V \simeq \mathbf{1}_W \oplus V'$ , then  $\mathrm{Th}(V) \simeq \Sigma^{2n,n} \mathrm{Th}(V')$ .

Given a G-equivariant open immersion  $U \subset W$  with smooth closed complement Z and a G-equivariant vector bundle V on W, observe that the inclusion of  $V|_U \hookrightarrow V$  has closed complement  $V|_Z$  and the normal bundle of  $V|_Z \hookrightarrow V$  is identified G-equivariantly with  $V|_Z \oplus \nu_{Z/W}$ . In that case, by functoriality of the homotopy purity isomorphisms [Hoy17, Corollary 3.25], there are G-equivariant cofiber sequences of the form

$$\operatorname{Th}(V|_U) \longrightarrow \operatorname{Th}(V) \longrightarrow \operatorname{Th}(\nu_{Z/X} \oplus V|_Z).$$

We use these cofiber sequences together with induction to establish the result. If there exists a G-Nisnevich covering  $Z' \to Z$  such that  $V|_{Z'}$  splits off a summand G-equivariantly isomorphic to  $1^{\oplus n}$ , then  $\text{Th}(V|_Z)$  lies in  $O^G(S^{2n,n})$  since the latter is closed under formation of colimits.

If  $\operatorname{Th}(V|_U) \in O(S^{2n,n})$  and  $V|_{Z'}$  splits G-equivariantly as a sum of a trivial rank r vector bundle and a (not necessarily G-equivariantly trivial) complement V', then

$$\operatorname{Th}(\nu_{Z/X} \oplus V|_Z) \in O^G(S^{2n,n})$$

by the discussion of the preceding paragraph. We conclude that  $Th(V) \in O^G(S^{2r,r})$  since the latter is closed under cofiber extensions by definition.

Consider the Luna stratification from Proposition 5.3.5. Since there is a generic stabilizer type, the restriction of  $V|_{X(H_0)}$  splits off a trivial summand of rank n by assumption. The collection of orbit types is partially ordered, by the closure order (say  $H \le H'$  if  $H \subset H'$ ). In that case, the result follows from the preceding statements by induction. In more detail: consider the Nisnevich covering of X(H) by the  $X(H)^H \times^H G$ .

**Theorem 5.3.7.** If k has characteristic 0, and X is a smooth k-scheme, then

$$\operatorname{Th}(p^*\bar{\rho}_r^{\oplus n}) \wedge (X^{\times r})_+ \in O^{\Sigma_r}(S^{2n,n})$$

*Proof.* It suffices to establish that  $\operatorname{Th}(p^*\bar{\rho_r}^{\oplus n}) \in O^{\Sigma_r}(S^{2n,n})$ . Consider the stratification of  $\mathbb{A}[\bar{\rho_r}]$  by orbit type. The assumption that k has characteristic 0 guarantees that  $|\Sigma_r|$  is invertible in k for all r. By appeal to Lemma 5.3.6, to establish the result, it suffices to show that the restriction of  $p^*\bar{\rho_r}$  to the G-stable locally closed subscheme of  $\mathbb{A}[\bar{\rho_r}] \setminus 0$  described by each stabilizer type splits off a trivial bundle of rank 1.

Now if *H* is a subgroup of  $\Sigma_r$  occurring as the stabilizer of a point, then *H* is a partition subgroup, i.e., it is isomorphic to  $\prod_{i=1}^{d} \Sigma_{n_i}$  where  $\sum_i n_i = r$ . The stabilizers of the  $\Sigma_r$  action on  $\mathbb{A}[\bar{\rho}_r] \setminus 0$  are always *proper*. Indeed, it suffices to observe that the fixed point locus is empty, and this can be checked after base change to an algebraic closure of *k*, in which case it is clear.

Thus, suppose H is a proper partition subgroup of  $\Sigma_r$ . It suffices to prove that  $\operatorname{Res}_{H}^{\Sigma_r} \bar{\rho}_r$  splits off a trivial bundle. We have  $H = \Sigma_{n_1} \times \cdots \times \Sigma_{n_d}$  and  $\operatorname{Res}_{H}^{\Sigma_r}(\rho_r) \simeq \rho_{n_1} \oplus \cdots \oplus \rho_{n_d}$ . Since  $d \ge 2$  (H being a proper subgroup) we find that  $\operatorname{Res}_{H}^{\Sigma_r}(\rho_r)$  splits off at least two trivial representations, whence  $\operatorname{Res}_{H}^{\Sigma_r} \bar{\rho}_r$  splits off at least one.

#### 5.4 Weak cellularity and quotients

Assume that k has characteristic 0. By the discussion of [ABH22, §6.1] we have a commutative diagram of functors of the form

$$\begin{array}{c} \mathbf{P}(\mathbf{Sch}_{k}^{ft}) \xrightarrow{R} \mathbf{P}(\mathbf{Sm}_{k}) \\ \downarrow & \downarrow \\ \mathbf{SH}_{ft}^{S^{1}}(k) \xrightarrow{\pi^{*}} \mathbf{SH}^{S^{1}}(k) \end{array}$$

where  $\operatorname{Sch}_{k}^{ft}$  denotes the category of quasi-projective k-schemes and the top horizontal map is induced by the functor  $\operatorname{Sm}_{k} \to \operatorname{Sch}_{k}^{ft}$ . The category  $\operatorname{SH}_{ft}^{S^{1}}(k)$  is the  $S^{1}$ -stable homotopy category built from  $\operatorname{Sch}_{k}^{ft}$  equipped with the cdh-topology. The functor  $\pi^{*}$  is an equivalence by a result of Voevodsky under these hypotheses [Voe10c, Theorem 4.2]. [ABH22, Proposition 6.2] asserts that if  $\mathscr{X} \in \operatorname{P}(\operatorname{Sch}_{k}^{ft})_{*}$ , then  $\operatorname{L}_{mot} \Sigma_{S^{1}}^{\infty}(\mathscr{X}|_{\operatorname{Sm}_{k}}) \simeq \pi^{*} \operatorname{L}_{cdh} \Sigma_{S^{1}}^{\infty} \mathscr{X}$ .

We now discuss in more detail the interaction between quotients and cellularity. If X is a G-quasiprojective smooth k-scheme, then  $Quot_G(X)$  is a finite type k-scheme.

**Proposition 5.4.1.** Assume k has characteristic 0. If  $\mathscr{X} \in \operatorname{Spc}^{G}(k)_{*}$  lies in  $O^{G}(S^{p,q})$ , then  $\Sigma_{S^{1}}^{\infty}R\operatorname{Quot}_{G} \mathscr{X} \in O(S^{p,q})$  as well.

*Proof.* By construction of  $O^G(S^{p,q})$  and closure of  $O(S^{p,q})$  under colimits and cofiber extensions, it suffices to prove that if  $\mathscr{X} \in \operatorname{Spc}^G(k)$  then  $\Sigma_{S^1}^{\infty} R \operatorname{Quot}_G(\Sigma^{p,q} \mathscr{X}) \in O(S^{p,q})$ . By the above discussion this is the same as  $\pi^* \Sigma_{S^1}^{\infty} \operatorname{Quot}_G(\Sigma^{p,q} \mathscr{X})$ . Noting that  $\operatorname{Quot}_G : \operatorname{H}^G(k)_* \to \operatorname{H}^{ft}_{\operatorname{Nis}}(k)$  commutes with  $\Sigma^{p,q}$  and so does  $\pi^*$  (since the inverse does), this is the same as  $\Sigma^{p,q} \pi^* \Sigma_{S^1}^{\infty} \operatorname{Quot}_G(\mathscr{X})$ , which lies in  $O(S^{p,q})$ , as needed.

**Theorem 5.4.2.** Assume k has characteristic 0. If  $E \in Mod_{H\mathbb{Z}}^{veff}(k)$ , then if  $E_n := \Omega^{\infty} \Sigma^{2n,n} E$ ,

$$cof(\Sigma^{2,1}E_n \to E_{n+1}) \in O(S^{4n+3,2n+1}).$$

*Proof.* We know that  $\operatorname{Mod}_{H\mathbb{Z}}^{\operatorname{veff}}(k)$  is generated under sifted colimits by spectra of the form  $H\mathbb{Z} \wedge \Sigma^{\infty} X_+$  (see Proposition 2.2.21). The functor  $\Omega^{\infty}$  preserves these colimits. Since weakly cellular classes are closed under colimits (see Lemma 3.1.6), it suffices to prove the result for  $E = H\mathbb{Z} \wedge \Sigma^{\infty} X_+$ . In that case, Proposition 5.3.1 tells us that

$$\operatorname{cof}(\Sigma^{2,1}\operatorname{Sym}^{r}\Sigma^{2n,n}X_{+} \to \operatorname{Sym}^{r}\Sigma^{2n+2,n+1}X_{+}) \simeq \Sigma^{2n+3,n+1}\operatorname{Quot}_{\Sigma_{r}}(\operatorname{Th}(p^{*}\bar{\rho}_{r}^{\oplus n}) \wedge (X^{\times r})_{+}),$$

while Theorem 5.3.7 tells us that

$$\Sigma^{2n+3,n+1}(\operatorname{Th}(p^*\bar{\rho}_r^{\oplus n}) \wedge (X^{\times r})_+) \in O^{\Sigma_r}(S^{4n+3,2n+1}).$$

In that case, Proposition 5.4.1 tells us that weak cellularity is preserved by taking quotients, at least after passing to  $S^1$ -spectra, so we see that

$$\Sigma_{S^1}^{\infty} \operatorname{cof}(\Sigma^{2,1} E_n \to E_{n+1}) \in O(S^{4n+3,2n+1}).$$

Also since  $cof(\Sigma^{2,1}E_n \to E_{n+1})$  is a (3,1)-suspension it is simply connected, and hence Theorem 3.2.12(1) implies that  $cof(\Sigma^{2,1}E_n \to E_{n+1}) \in O(S^{4n+3,2n+1})$ .

## 5.5 Modifications in positive characteristic

Dealing with cellularity of quotients in positive characteristic is slightly more subtle for two reasons. First, at the moment there is not a well-defined  $S^1$ -stable homotopy for singular varieties even after inverting the characteristic of the base-field. If strong resolution of singularities were to hold in positive characteristic, then the argument above could be transferred verbatim to positive characteristic. Second, when analyzing symmetric products, we have to deal with non-tame group actions, since the characteristic exponent of the base field will divide the order of  $\Sigma_r$  for r sufficiently large. For these reasons, rather than attempting to work in  $S^1$ -spectra, we will work directly in Voevodsky's category of motives appealing to the results of [Kel17].

#### Motives of singular schemes in positive characteristic

Assume k is a field that has characteristic exponent p > 0 and  $\ell$  is a prime different from p. Write  $\mathbf{DM}_{ft}^{\ell dh}(k, \mathbb{Z}_{(\ell)})$  for the variant of Voevodsky's category of motives built out of finite type schemes and the  $\ell dh$ -topology. There is an evident functor

$$\operatorname{Mod}_{H\mathbb{Z}_{(\ell)}}(k) \simeq \mathbf{DM}(k, \mathbb{Z}_{(\ell)}) \to \mathbf{DM}_{ft}^{\ell dh}(k, \mathbb{Z}_{(\ell)})$$

whose right adjoint we denote by  $r_{H\mathbb{Z}_{(\ell)}}$ . In fact this is an adjoint equivalence [Kel17, Theorem 4.0.13].

**Theorem 5.5.1.** There is a commutative square of the form

*Proof.* This is [HKØ17, Theorem 3.1] in view of [HKØ17, Proposition 4.10] and [HKØ17, Corollary 4.27].  $\Box$ 

**Proposition 5.5.2.** Assume k has characteristic exponent p > 0 and  $\ell$  is a prime different from p. If  $\mathscr{X} \in \operatorname{Spc}^G(k)$  lies in  $O^G(S^{r,s})$ , then  $H\mathbb{Z}_{(\ell)} \wedge \Sigma^{\infty} R \operatorname{Quot}_G \mathscr{X} \in O(S^{r,s})$  as well.

*Proof.* The proof follows the same lines as that of Proposition 5.4.1. As before,  $O(S^{r,s})$  is stable under colimits and cofiber extensions. Therefore, by construction of  $O^G(S^{r,s})$ , without loss of generality, we may assume that  $\mathscr{X} = \Sigma^{r,s}Y_+$  for a smooth G-scheme Y. In that case,  $\operatorname{Quot}_G(\Sigma^{r,s}Y_+) = \Sigma^{r,s}(Y/G)_+$  by definition. As before the functor  $r_{H\mathbb{Z}_{(\ell)}}$  commutes with  $\Sigma^{r,s}$  (since its inverse does), and we conclude.

#### Transferring

Assume k is a field that has characteristic exponent p > 0. If G is a finite group, then G is linearly reductive if and only if |G| is coprime to k. In Theorem 5.3.7, we needed to consider symmetric groups  $\Sigma_r$  for arbitrary r, in particular r might be divisible by p and therefore the subgroups  $H \subset \Sigma_r$  we consider need not be linearly reductive.

We now discuss how to avoid this issue, at the expense of inverting the prime p, or localizing at a prime  $\ell$  different from p via a standard transfer argument. In practice, we will use the fact that if G is a finite group and H is an  $\ell$ -Sylow subgroup, then G/H has order coprime to  $\ell$ . If  $X \in \text{Sm}_k^G$ , then there is an evident map  $\text{Res}_H^G(X)/H \to X/G$  that is finite and flat morphism of degree |G/H| and thus coprime to  $\ell$ . This map induces a covariant transfer map of motives and pullback followed by transfer is multiplication by |G/H|. The following lemma is a precise version of this observation.

**Lemma 5.5.3.** Assume k is a field having characteristic exponent p and let  $\ell$  be a prime different from p. Let G be a finite group and  $H \subset G$  a finite subgroup of index coprime to  $\ell$  and suppose  $X \in \text{Sm}_k^G$ . The object  $\underline{M}(X/G)$  is a retract of  $\underline{M}(\text{Res}_H^G(X)/H)$ .

*Proof.* This is a consequence of [Voe10b, Proposition A.8(1)]: loosely speaking pullback followed by pushforward is multiplication by |G/H| which is invertible in  $\mathbb{Z}_{(\ell)}$ .

We now provide a variant of Theorem 5.3.7 in positive characteristic. If G is a finite group, then by definition, the  $\ell$ -Sylow subgroup  $Syl_{\ell}(G) \subset G$  has index coprime to  $\ell$  and has order invertible in k. As in Section 5.3 consider the standard representation  $\rho_r$  of  $\Sigma_r$ . Abusing notation, we write p for the  $Syl_{\ell,r}$ -equivariant structure map  $\mathbb{A}[\operatorname{Res}_{Syl_{\ell,r}}^{\Sigma_r}(\rho_r)] > 0.$ 

**Theorem 5.5.4.** Assume k has characteristic exponent p > 0, and  $\ell$  is a prime different from p and X is a smooth k-scheme. If  $Syl_{\ell,r} \subset \Sigma_r$  is an  $\ell$ -Sylow subgroup, then

$$\Sigma^{2n+3,n+1}(\operatorname{Th}(p^*(\operatorname{Res}_{Syl_{\ell,r}}^{\Sigma_r}\bar{\rho}_r)^{\oplus n})\wedge (X^{\times r})_+),$$

lies in  $O^{Syl_{\ell,r}}(S^{4n+3,2n+1})$ .

*Proof.* Consider the stratification of  $\mathbb{A}[\operatorname{Res}_{Syl_{\ell,r}}^{\Sigma_r}(\rho_r)] \leq 0$  by stabilizer type from Proposition 5.3.5 for the action of  $Syl_{\ell,r}$ . In this case, the stabilizers are intersections of the form  $Syl_{\ell,r} \cap H$  where H is a proper partition subgroup of  $\Sigma_r$ . Since the order of  $Syl_{\ell,r}$  is invertible in p, it follows that the orders of each of the groups  $Syl_{\ell,r} \cap H$  are also invertible in k, in particular, all these groups are linearly reductive.

By appeal to Lemma 5.3.6, it suffices to show that for group of the form  $Syl_{\ell,r} \cap H$  the representation  $\operatorname{Res}_{Syl_{\ell,r}\cap H}^{\Sigma_r} \bar{\rho_r}$  splits off a trivial summand. This follows in exactly the same way as at the end of Theorem 5.3.7.

Finally, we may establish a version of Theorem 5.4.2 in positive characteristic.

**Theorem 5.5.5.** Assume k is a field having characteristic exponent p > 1. Let n > 0 and  $E \in Mod_{H\mathbb{Z}[1/p]}^{\text{veff}}(k)$ . For  $E_n := \Omega^{\infty} \Sigma^{2n,n} E$  we have

$$cof(\Sigma^{2,1}E_n \to E_{n+1}) \in O(S^{4n+3,2n+1}).$$

*Proof.* As before we may reduce to  $E = H\mathbb{Z}[1/p] \wedge \Sigma^{\infty}X_{+}$  for some  $X \in Sm_{k}$ . By continuity, X is pulled back from a perfect subfield of finite cohomological dimension (the perfect closure of a finitely

generated extension of  $\mathbb{F}_p$ ). By essentially smooth base change (Lemma 3.1.13) we may thus assume that k has finite cohomological dimension. Since  $\Omega^{\infty} E_n \in O(S^{2n,n})$  by Proposition 3.2.11, we see that

$$Y := \operatorname{cof}(\Sigma^{2,1} E_n \to E_{n+1}) \in O(S^{2n+2,n+1}) \subset O(S^{4,2}).$$

Combining this with Proposition 2.2.13 we find that also

$$\Sigma_{S^1}^{\infty} Y \in O(S^{4,2})[1/p]$$

We may thus apply Theorem 3.2.12(4) to reduce to showing that  $\Sigma^{\infty}Y \wedge H\mathbb{Z} \in O(S^{4n+3,2n+1})$ . By standard arguments it suffices to prove the same for  $\Sigma^{\infty}Y \wedge H\mathbb{Z}_{(\ell)}$  for all  $\ell \neq p$ . This follows from Lemma 5.5.3 in conjunction with Proposition 5.5.2, Theorem 5.5.4 and Theorem 5.2.9.

## 6 Weak cellular estimates of the fiber of the unit map

In this section, we put everything together to deduce the Freudenthal theorem. For a pointed connected motivic space  $\mathscr{X}$ , write

$$s_{\mathscr{X}}: \mathscr{X} \longrightarrow \Omega^{2,1} \Sigma^{2,1} \mathscr{X};$$

for the stabilization map, i.e., the unit map of the loop-suspension adjunction. Our goal is to, assuming suitable hypotheses on the weak cellular class of  $\mathscr{X}$ , provide an estimate for the weak cellular class of  $\operatorname{fib}(s_{\mathscr{X}})$ . We proceed by a series of reductions.

## 6.1 The unit of stabilization and fiber sequences

In this section, we analyze how the weak cellular class of the fiber of the stabilization map behaves in fiber sequences. Proposition 6.1.1 provides fiber sequences linking the fiber of stabilization in the base, fiber and total space of a fiber sequence. Then Proposition 6.1.2 yields the main inductive tool we use to analyze the fiber of stabilization.

**Proposition 6.1.1** (Fibers). For a pointed connected motivic space  $\mathscr{X}$ , write

$$s_{\mathscr{X}}:\mathscr{X}\longrightarrow\Omega^{2,1}\Sigma^{2,1}\mathscr{X};$$

for the stabilization map, i.e., the unit map of the loop-suspension adjunction. Suppose we have a fiber sequence  $\mathscr{F} \to \mathscr{E} \to \mathscr{B}$  of pointed, connected motivic spaces. Set  $\mathscr{F}' := \operatorname{fib}(\Sigma^{2,1}\mathscr{E} \to \Sigma^{2,1}\mathscr{B})$ .

1. There is an induced factorization

$$\mathscr{F} \longrightarrow \Omega^{2,1} \Sigma^{2,1} \mathscr{F} \longrightarrow \Omega^{2,1} \mathscr{F}',$$

where the first morphism is the unit map.

2. There is an induced fiber sequence


$$\operatorname{fib}(\mathscr{F} \to \Omega^{2,1}\mathscr{F}') \longrightarrow \operatorname{fib}(s_{\mathscr{E}}) \longrightarrow \operatorname{fib}(s_{\mathscr{B}})$$

*3. There is an induced fiber sequence* 

$$\operatorname{fib}(s_{\mathscr{F}}) \longrightarrow \operatorname{fib}(\mathscr{F} \to \Omega^{2,1}\mathscr{F}') \longrightarrow \Omega^{2,1} \operatorname{fib}(\Sigma^{2,1}\mathscr{F} \to \mathscr{F}')$$

*Proof.* Since the composite map  $\mathscr{F} \to \mathscr{B}$  is null, so is the map  $\Sigma^{2,1}\mathscr{F} \to \Sigma^{2,1}\mathscr{B}$ . It follows that there is a map  $\Sigma^{2,1}\mathscr{F} \to \mathscr{F}'$ , which by looping yields the map  $\Omega^{2,1}\Sigma^{2,1}\mathscr{F} \to \Omega^{2,1}\mathscr{F}'$  and by adjunction yields the map  $\mathscr{F} \to \Omega^{2,1} \mathscr{F}'$  factoring as claimed.

Consider the commutative diagram:



where the vertical maps are unit maps. The fiber sequence of the second point arises by comparing the vertical and horizontal homotopy fibers in this diagram.


The third fiber sequence arises by taking the fiber of the composite via [ABH22, Proposition 3.1] for the factorization in the first point. 

**Proposition 6.1.2.** Suppose  $\mathscr{F} \xrightarrow{\iota} \mathscr{E} \xrightarrow{f} \mathscr{B}$  is a fiber sequence of pointed, motivic spaces, lying in  $O(S^{m,n}), m-n \ge 2, n \ge 1.$  Set  $\mathscr{F}' := \operatorname{fib}(\Sigma^{2,1}\mathscr{E} \to \Sigma^{2,1}\mathscr{B}).$ 

1. We have fib
$$(\Sigma^{2,1}\mathscr{F} \to \mathscr{F}') \in O(S^{2m+1,2n+1})$$
.

- 2. If  $\operatorname{fib}(s_{\mathscr{F}}) \in O(S^{2n-1,2m})$ , then  $\operatorname{fib}(s_{\mathscr{E}}) \in O(S^{2m-1,2n})$  if and only if  $\operatorname{fib}(s_{\mathscr{F}}) \in O(S^{2m-1,2n})$ . 3. If  $\operatorname{fib}(s_{\mathscr{F}}) \in O(S^{2m-1,2n})$ , then  $\operatorname{fib}(\operatorname{fib}(s_{\mathscr{F}}) \to \operatorname{fib}(s_{\mathscr{E}})) \in O(S^{2m-2,2n})$

*Proof.* Point 1. Application of Corollary 4.3.2 implies that it suffices to show that  $cof(\Sigma^{2,1}\mathscr{F} \to \mathscr{F}') \in$  $O(S^{2m+2,2n+1})$ , for which by Theorem 3.2.12 it is enough that  $\Sigma C := \Sigma^2 \operatorname{cof}(\Sigma^{2,1} \mathscr{F} \to \mathscr{F}') \in O(S^{2m+4,2n+1}).$ Consider the commutative square of cofiber sequences



The connectivities in the bottom row are obtained by Theorem 4.3.1(2) and imply what we need (using Corollary 3.1.26).

**Point 2.** By Proposition 6.1.1(3) there is a fiber sequence

$$\operatorname{fib}(s_{\mathscr{F}}) \longrightarrow \operatorname{fib}(\mathscr{F} \to \Omega^{2,1}\mathscr{F}') \longrightarrow \Omega^{2,1} \operatorname{fib}(\Sigma^{2,1}\mathscr{F} \to \mathscr{F}').$$

By the conclusion of the first point in conjunction with Propostion 4.2.1, we conclude that the last term lies in  $O(S^{2m-1,2n})$ . If  $fib(s_{\mathscr{F}}) \in O(S^{2m-1,2n})$ , then Proposition 3.1.22 tells us that the second map in the preceding display is an  $S^{2m-1,2n}$ -equivalence, and we deduce that  $\operatorname{fib}(\mathscr{F} \to \Omega^{2,1}\mathscr{F}') \in O(S^{2m-1,2n})$ as well.

Next, there is a fiber sequence

$$\operatorname{fib}(\mathscr{F} \to \Omega^{2,1}\mathscr{F}') \longrightarrow \operatorname{fib}(s_{\mathscr{E}}) \longrightarrow \operatorname{fib}(s_{\mathscr{B}}).$$

by Proposition 6.1.1(2). In that case, another application of Proposition 3.1.22 in conjunction with the estimate of the preceding paragraph implies that  $fib(s_{\mathscr{B}}) \in O(S^{2m-1,2n})$ , if and only if  $fib(s_{\mathscr{E}}) \in O(S^{2m-1,2n})$ .  $O(S^{2m-1,2n}).$ 

**Point 3.** Proposition 6.1.1 shows that the map  $fib(s_{\mathscr{F}}) \to fib(s_{\mathscr{E}})$  factors as

$$\operatorname{fib}(s_{\mathscr{F}}) \longrightarrow \operatorname{fib}(\mathscr{F} \to \Omega^{2,1}\mathscr{F}') \longrightarrow \operatorname{fib}(s_{\mathscr{E}})$$

The proposition also shows that the fiber of the first map is  $\Omega^{3,1} \operatorname{fib}(\Sigma^{2,1} \mathscr{F} \to \mathscr{F}')$ , while the fiber of the second map is  $\Omega \operatorname{fib}(s_{\mathscr{B}})$ . Assuming  $\operatorname{fib}(s_{\mathscr{B}}) \in O(S^{2m-1,2n})$ , we conclude that  $\Omega \operatorname{fib}(s_{\mathscr{B}}) \in O(S^{2m-2,2n})$ . The first point yields  $\Omega^{3,1} \operatorname{fib}(\Sigma^{2,1} \mathscr{F} \to \mathscr{F}') \in O(S^{2m-2,2n})$ . In that case Corollary 3.1.23 implies that  $\operatorname{fib}(\operatorname{fib}(s_{\mathscr{F}}) \to \operatorname{fib}(s_{\mathscr{E}})) \in O(S^{2m-2,2n})$  as well.

#### 6.2 The main result

In this section, we combine all the tools developed so far to analyze  $\mathbb{P}^1$ -suspension. If E is any  $\mathbb{P}^1$ -spectrum, then set

$$\underline{E}^{a,b} := \Omega^{\infty} \Sigma^{a,b} E$$

We may now state the main result.

**Theorem 6.2.1** (The  $\mathbb{P}^1$ -stabilization theorem). Assume k is a perfect field with characteristic exponent p,  $\mathscr{X}$  is a motivic space and  $s_{\mathscr{X}} : \mathscr{X} \to \Omega^{2,1} \Sigma^{2,1} \mathscr{X}$  is the unit of the loop suspension adjunction. Assume m and n are integers with  $m \leq 2n, m-n \geq 2$ , and  $n \geq 2$ . If  $\mathscr{X} \in O(S^{m,n})$  and  $\pi_*(\mathscr{X})$  is uniquely p-divisible, then  $\operatorname{fib}(s_{\mathscr{X}}) \in O(S^{2m-1,2n})$ .

Before delving into the proof, we outline the strategy. We quickly reduce to the case m = 2n and by appeal to the weakly cellular Whitehead tower, and the inductive machinery described in the previous section, we reduce the analysis of the weak cellular class of the fiber of the unit map  $s_{\mathscr{X}}$  to the case of  $\mathbb{P}^1$ -infinite loop spaces; essentially all the effort is focused on treating this latter case.

Intuitively, we analyze the fiber of the stabilization map for  $\mathbb{P}^1$ -infinite loop spaces by "fracturing" with respect to behavior with respect to the motivic Hopf map  $\eta$  (recall the conventions of Paragraph 2.2.17). In particular, we would like to separately analyze  $\eta$ -complete and  $\eta$ -local  $\mathbb{P}^1$ -infinite loop spaces. In practice, the geometric analysis of the cofiber of the assembly map for Eilenberg–Mac Lane spaces will give us control over the  $\eta$ -complete situation.

There is an equality of the form  $1 - \eta \rho = 1 + \eta [-1] = \langle -1 \rangle$  and therefore,  $1 + \varepsilon = \eta \rho$  in  $\pi_{0,0}(\mathbb{1}_k)$ . From this observation, the  $\eta$ -local situation can be accessed, if we additionally invert 2, by studying the  $\rho$ -local situation, which in turn can be analyzed by appeal to the real-étale topology. To build a fracture square, we then need to study  $\eta$  and 2-torsion, which we do appeal to a modification of Levine's slice convergence argument. There are a number of ways to perform this kind of "fracture" in practice: here we analyze the torsion and rational parts separately, but other configurations are also possible.

*Proof.* We proceed by a string of reductions.

Step 1. (Reduction to the Chow diagonal) We may assume that m = 2n. Indeed, if  $\mathscr{X} \in O(S^{m,n})$ , then  $\Sigma^{2n-m}\mathscr{X} \in O(S^{2n,n})$ . Since  $\mathscr{X}$  is 1-connected by assumption, so is  $\Sigma^{2n-m}\mathscr{X}$ . If  $\pi_*(\mathscr{X})$  is uniquely *p*-divisible, then  $\Sigma_{S^1}^{\infty}\mathscr{X}$  is as well, and thus so is  $\Sigma_{S^1}^{\infty}\Sigma^{2n-m}\mathscr{X}$ . Therefore, Proposition 2.2.14 implies that  $\pi_*(\Sigma^{2n-m}\mathscr{X})$  is uniquely *p*-divisible as well. In that case, assuming  $\operatorname{fib}(s_{\Sigma^{2n-m}}\mathscr{X}) \in O(S^{4n-1,2n})$ , appealing inductively to Lemma 6.2.2, we conclude that  $\operatorname{fib}(s_{\mathscr{X}}) \in O(S^{2m-1,2n})$  as required.

Step 2. (Reduction to  $\mathbb{P}^1$ -infinite loop spaces) To establish the result for spaces, it suffices to establish the result for  $\mathbb{P}^1$ -infinite loop spaces; this reduction is the content of Lemma 6.2.3. In more detail, under the hypotheses, to show that  $\operatorname{fib}(s_{\mathscr{X}}) \in O(S^{4n-1,2n})$ , it suffices to establish that for E is  $\operatorname{SH}(k)^{\operatorname{veff}}[\frac{1}{p}]$  we have  $\operatorname{fib}(s_{\underline{E}^{2n,n}}) \in O(S^{4n-1,2n})$ . Moreover, Proposition 6.1.2(2) implies that the collection of  $E \in \operatorname{SH}(k)^{\operatorname{veff}}[1/p]$  satisfying this estimate on the weak-cellular class of the fiber of stabilization is stable under extensions and cofibers; it is also stable under filtered colimits. Thus, it even suffices to analyze  $\operatorname{fib}(s_{\underline{E}^{2n,n}})$  with  $E = \Sigma^{\infty}_+ X[1/p]$  and  $X \in \operatorname{Sm}_k$  (see Proposition 2.2.21).

**Step 3.** (Reduction to the case of finite vcd) We may assume that k has finite vcd. Indeed, since  $X \in Sm_k$  has finite-presentation, it is necessarily pulled back from the perfect closure of some finitely generated extension of the prime field. Since cellularity estimates are stable under essentially smooth base-change (Lemma 3.1.13), we conclude.

**Step 4.** (Reduction to slices and effective covers) It suffices to establish the weak cellular estimate on the fiber of the stabilization map for the effective cover  $f_N E$  for N sufficiently large and  $s_i E$  for arbitrary *i*; see Step 5 for the latter, and proceed to Step 7 for the former. Indeed, there are fiber sequences of the form

$$f_{i+1}E \longrightarrow f_iE \longrightarrow s_iE,$$

and thus corresponding fiber sequences after applying  $\Omega^{\infty}$ . In that case, the assumption that  $\operatorname{fib}(s_{\Omega^{\infty}f_NE}) \in O(S^{4n-1,2n})$  for N large and  $\operatorname{fib}(s_{\Omega^{\infty}s_iE})$  for all  $i \geq 0$  implies that  $\operatorname{fib}(s_{\Omega^{\infty}f_iE}) \in O(S^{4n-1,2n})$  by repeated appeal to Proposition 6.1.2(2). In that case, the required weakly cellular estimate follows by descending induction.

**Step 5.** (Reducing from slices to very effective motives) If  $E \in SH(k)^{veff}$ , then  $s_i E$  remains very effective and admits the structure of an  $H\mathbb{Z}[1/p]$ -module (see Paragraph 2.2.10 for more details and references).

Step 6. (Treating effective motives) The weakly cellular estimate on the fiber of the stabilization map holds for  $E \in \operatorname{Mod}_{H\mathbb{Z}[1/p]}^{\operatorname{veff}}(k)$ . Indeed, in that case, the relevant weak cellular estimate is contained in Lemma 6.2.5.

Step 7. (Treating the rational plus part of effective covers) To establish the result for  $f_N E$  as above, we proceed in two steps, first, we treat the rationalization  $f_N E_{\mathbb{Q}}$ ; this breaks up into  $\pm$  pieces under the decomposition of  $\mathrm{SH}(k)_{\mathbb{Q}}$  into eigenspaces of  $\varepsilon$  (see the conventions of Paragraph 2.2.22). The +-part is described in terms of Voevodsky motives as detailed in Theorem 2.2.23. Lemma 6.2.4 implies that  $(f_N E_{\mathbb{Q}})^+ = 0$ , so we conclude that  $f_N E_{\mathbb{Q}} \simeq (f_N E_{\mathbb{Q}})^-$ .

**Step 8.** (Reducing the rational minus part of effective covers to the real-étale case) The - part of the decomposition of  $SH(k)_{\mathbb{Q}}$  was identified in terms of the derived category of real-étale sheaves of  $\mathbb{Q}$ -vector spaces (Theorem 2.3.3). In particular, we conclude  $(f_N E_{\mathbb{Q}})^-$  has real-étale descent.

Step 9. (Separating out the fields.) Let us set  $\mathscr{X}_i = \Omega^{\infty} \Sigma^{2n,n} f_i E$ . We want to prove that  $\operatorname{fib}(s_{\mathscr{X}_i}) \in O(S^{4n-1,2n})$  for some *i*. Taking i > 2n ensures that  $\operatorname{fib}(s_{\mathscr{X}_i}) \in O(S^{2n+1,2n})$  and hence by Corollary 3.1.26 it will be enough to show that  $\pi_r(\operatorname{fib}(s_{\mathscr{X}_i})) = 0$  for  $r \leq 2n-2$ . Since  $s_i(E)$  has already been dealt with (step 5), we have  $\operatorname{fib}(s_{\Omega^{\infty}\Sigma^{2n,n}s_iE}) \in O(S^{4n-1,2n})$ . Now Proposition 6.1.2(3) shows that  $\pi_r(\operatorname{fib}(s_{\mathscr{X}_i+1})) \to \pi_r(\operatorname{fib}(s_{\mathscr{X}_i}))$  is surjective for  $r \leq 2n-2$ . Since homotopy sheaves are unramified, it hence suffices to prove: for every finitely generated extension K/k there exists  $i \gg 0$  with

 $\pi_r(\operatorname{fib}(s_{\mathscr{X}_i}))(K) = 0 \text{ for } r \leq 2n-2.$ 

Step 10. Consider the commutative diagram

$$\begin{array}{cccc} \mathscr{X}_i & \longrightarrow & L_{\mathrm{r\acute{e}t}} \mathscr{X}_i \\ & & \downarrow \\ \Omega^{2,1} \Sigma^{2,1} \mathscr{X}_i & \longrightarrow & \Omega^{2,1} L_{\mathrm{r\acute{e}t}} \Sigma^{2,1} \mathscr{X}_i. \end{array}$$


Since  $(\mathscr{X}_i)_{\mathbb{Q}}$  satisfies real étale descent (step 8), the same is true for  $(\Sigma^{2,1}\mathscr{X}_i)_{\mathbb{Q}}$  (use that being a real étale sheaf is equivalent to being  $\rho$ -periodic by Proposition 2.3.2), and so Lemma 6.2.7 shows that on sections over K, the horizontal maps are arbitrarily connective (for  $i \gg 0$ ). It follows that

$$\operatorname{fib}(s_{\mathscr{X}_i}) \to \operatorname{fib}(L_{\operatorname{r\acute{e}t}} \mathscr{X}_i \to \Omega^{2,1} L_{\operatorname{r\acute{e}t}} \Sigma^{2,1} \mathscr{X}_i)$$

is arbitrarily connective on sections over K (for  $i \gg 0$ ). We conclude since the right hand space lies in  $O(S^{4n-1,2n})$  by Lemma 6.2.6.

**Lemma 6.2.2.** Suppose  $\mathscr{X} \in O(S^{m,n})$  with m < 2n and  $m - n \ge 2$  (and  $n \ge 0$ ). If  $\operatorname{fib}(s_{\Sigma}\mathscr{X}) \in O(S^{2m+1,2n})$ , then  $\operatorname{fib}(s_{\mathscr{X}}) \in O(S^{2m-1,2n})$ .

Proof. Consider the diagram



where the bottom horizontal map is  $\Omega^{2,1}$  applied to the map  $\Sigma^{2,1} \mathscr{X} \to \Omega \Sigma^{3,1} \mathscr{X}$ . Both composites are equivalent to the evident unit map, so the diagram commutes.

Since  $\mathscr{X} \in O(S^{m,n})$ , we conclude that  $\Sigma^{2,1} \mathscr{X} \in O(S^{m+2,n+1})$ . Proposition 4.3.3 tells us the fiber of top horizontal map lies in  $O(S^{2m-1,2n})$  and that

$$\operatorname{fib}(\Sigma^{2,1}\mathscr{X}\longrightarrow \Omega\Sigma^{3,1}\mathscr{X})\in O(S^{2m+3,2n+2}).$$

Then, Proposition 4.2.1 tells us that  $\Omega^{2,1} \text{fib}(\Sigma^{2,1} \mathscr{X} \to \Omega \Sigma \mathscr{X}) \in O(S^{2m+1,2n+1})$ . On the other hand, the hypothesis is that  $\text{fib}(s_{\Sigma \mathscr{X}}) \in O(S^{2m+1,2n})$  and another application of Proposition 4.2.1 tells us that  $\Omega \text{fib}(s_{\Sigma \mathscr{X}}) \in O(S^{2m,2n})$ .

The fiber sequence attached to a composite of morphisms [ABH22, Proposition 3.1] yields:

$$\operatorname{fib}(\mathscr{X} \to \Omega \Sigma \mathscr{X}) \longrightarrow \operatorname{fib}(\mathscr{X} \to \Omega^{3,1} \Sigma^{3,1} \mathscr{X}) \longrightarrow \Omega \operatorname{fib}(s_{\Sigma \mathscr{X}}).$$

The weak connectivity estimates above in conjunction with Proposition 3.1.22 tell us that  $\operatorname{fib}(\mathscr{X} \to \Omega^{3,1}\Sigma^{3,1}\mathscr{X}) \in O(S^{2m-1,2n})$  as well.

Another application of [ABH22, Proposition 3.1] yields a fiber sequence of the form

$$\operatorname{fib}(s_{\mathscr{X}}) \longrightarrow \operatorname{fib}(\mathscr{X} \to \Omega^{3,1}\Sigma^{3,1}\mathscr{X}) \longrightarrow \Omega^{2,1}\operatorname{fib}(\Sigma^{2,1}\mathscr{X} \to \Omega\Sigma^{3,1}\mathscr{X}),$$

which rotates to a fiber sequence of the form

$$\Omega^{3,1}\mathrm{fib}(\Sigma^{2,1}\mathscr{X}\to\Omega\Sigma^{3,1}\mathscr{X})\longrightarrow\mathrm{fib}(s_\mathscr{X})\longrightarrow\mathrm{fib}(\mathscr{X}\to\Omega^{3,1}\Sigma^{3,1}\mathscr{X})$$

We have  $\Omega^{3,1} \text{fib}(\Sigma^{2,1} \mathscr{X} \to \Omega \Sigma^{3,1} \mathscr{X}) \in O(S^{2m,2n+1})$  by appeal to Proposition 4.2.1. Another application of Proposition 3.1.22 implies that  $\text{fib}(s_{\mathscr{X}}) \in O(S^{2m-1,2n})$  as required.

**Lemma 6.2.3.** Suppose  $\mathscr{X} \in O(S^{2n,n})$ ,  $n \geq 2$ . The space  $\operatorname{fib}(s_{\mathscr{X}}) \in O(S^{4n-1,2n})$  as soon as for  $i \geq 0$  the spaces  $\operatorname{fib}(s_{\tau > a,a}K(\pi_i \mathscr{X}, i)) \in O(S^{4n-1,2n})$ .

Proof. Consider the Postnikov tower

$$\tau_{\geq a,a} K(\boldsymbol{\pi}_i \mathscr{X}, i) \to \mathcal{L}^{2n+i+1,n} \mathscr{X} \to \mathcal{L}^{2n+i,n} \mathscr{X}$$

of Theorem 4.1.10. By induction using Proposition 6.1.2(2) we find that  $\operatorname{fib}(s_{L^{2n+i,n}\mathscr{X}}) \in O(S^{4n-1,2n})$ for every *i*. Consider the fiber sequence  $\mathscr{F}_i \to \mathscr{X} \to L^{2n+i,n}\mathscr{X}$ . In the language of Proposition 6.1.1, both  $\mathscr{F}_i$  and  $\mathscr{F}'_i$  are *i*-connective (use Lemma 3.1.18), and hence  $\operatorname{fib}(s_{\mathscr{X}}) \to \operatorname{fib}(s_{L^{2n+i,n}\mathscr{X}})$  is (i-1)connective. The result now follows via Corollary 3.1.27.

# **Lemma 6.2.4.** If $X \in \text{Sm}_k$ , then $(f_N \Sigma^{\infty} X_+)^+_{\mathbb{O}} = 0$ for N sufficiently large depending on X.

*Proof.* This argument is a modification of one due to Levine [Lev13, Proposition 6.9(2)]. Since we are working with rational coefficients, and strongly dualizable objects generate SH(k) (see Proposition 2.2.12), we know that  $\Sigma^{\infty}X_{+}$  can be built out finitely many  $\Sigma^{\infty}Y_{+}$  with Y smooth and projective. We will show that we may take N to be the maximum of the dimensions of Y and we reduce to demonstrating the vanishing above for X smooth and projective.

Let  $d = \dim X$ . For any  $E \in SH(k)$  the map  $f_N E \to E$  is universal among maps from spectra in  $\Sigma^{2n,n}SH(k)^{\text{eff}}$  to E. Since  $\Sigma^{2n,n}SH(k)^{\text{eff}}$  is the localizing subcategory generated by  $\Sigma^{q,q}\Sigma^{\infty}U_+$  for  $q \ge n$ , to show  $f_N E = 0$  for N > d, it suffices to show that  $\operatorname{Hom}_{SH(k)_{\mathbb{Q}}}(\Sigma^{*+N,N}\Sigma^{\infty}U_+, E) = 0$  for all N > d.

By the equivalence of  $\text{SH}(k)^+_{\mathbb{Q}}$  with  $\text{Mod}_{H\mathbb{Q}}(k)$  (see Theorem 2.2.23), in order to demonstrate  $f_N E^+$  vanishes we need only establish the vanishing of

$$\operatorname{Hom}_{\operatorname{Mod}_{H\mathbb{Q}}(k)}(\mathbf{M}_{\mathbb{Q}}(U)[*+N](N), \mathbf{M}_{\mathbb{Q}}(X)).$$

Since X is smooth and projective of dimension d, Atiyah duality yields identifications of the form:

$$\operatorname{Hom}_{\operatorname{Mod}_{H\mathbb{Q}}(k)}(\mathbf{M}_{\mathbb{Q}}(U)[*+N](N), \mathbf{M}_{\mathbb{Q}}(X)) \simeq \operatorname{Hom}_{\operatorname{Mod}_{H\mathbb{Q}}(k)}(\mathbf{M}_{\mathbb{Q}}(U \times X), \mathbb{Q}(d-N)[2d-N-*])$$
$$\simeq H^{2d-N-*, d-N}(U \times X, \mathbb{O})$$

For any smooth k-scheme, motivic cohomology vanishes in negative weight (this follows either from the identification of motivic cohomology with higher Chow groups [Voe02, Corollary 2], or from the definition of motivic complexes in conjunction with the cancellation theorem [Voe10a, Corollary 4.10]), so the preceding motivic cohomology group vanishes when N > d as required.

**Lemma 6.2.5.** If  $E \in Mod^{veff}(H\mathbb{Z})[1/p]$  and  $n \ge 0$ , then

$$fib(s_{E^{2n,n}}) \in O(S^{4n-1,2n}).$$

(If k has positive characteristic, assume  $n \ge 1$ .)

*Proof.* Observe that there is a factorization of the identity map

$$\underline{E}^{2n,n} \stackrel{s_{\underline{E}^{2n,n}}}{\longrightarrow} \Omega^{2,1} \underline{\Sigma}^{2,1} \underline{\underline{E}}^{2n,n} \stackrel{\Omega^{2,1}a_n}{\longrightarrow} \Omega^{2,1} \underline{\underline{E}}^{2n+2,n+1} \simeq \underline{\underline{E}}^{2n,n}$$

where the first map is the unit map and the map  $a_n$  is the assembly map for the spectrum. By appeal to [ABH22, Proposition 3.1] we thus obtain a fiber sequence of the form

$$\operatorname{fib}(s_{E^{2n,n}}) \longrightarrow * \longrightarrow \Omega^{2,1} \operatorname{fib}(a_n),$$

i.e., an equivalence  $\Omega^{3,1} \operatorname{fib}(a_n) \simeq \operatorname{fib}(s_{E^{2n,n}}).$ 

If k has characteristic 0, then Theorem 5.4.2 yields a cellularity estimate for the cofiber of  $a_n$ . If k has positive characteristic, Theorem 5.5.5 achieves the same. In either case, Corollary 4.3.2(2) allows us to turn this into a weak cellular estimate for fib $(a_n)$ ; taken together, we conclude that fib $(a_n) \in O(S^{4n+2,2n+1})$ . Proposition 4.2.1 then allows us to give a cellularity estimate for the loop space  $\Omega^{3,1}(-)$ : writing  $\Omega^{3,1}$ fib $(a_n) = \Omega \Omega^{2,1}$ fib $(a_n)$ , we conclude that  $\Omega^{2,1}$ fib $(a_n) \in O(S^{4n-1,2n})$  as required.

**Lemma 6.2.6.** If  $\mathscr{X} \in O(S^{2n,n})$ ,  $n \geq 2$ , then  $\operatorname{fib}(\operatorname{L}_{r\acute{e}t}\mathscr{X} \to \Omega^{2,1}\operatorname{L}_{r\acute{e}t}\Sigma^{2,1}\mathscr{X}) \in O(S^{4n-1,2n})$ .

*Proof.* Since  $\mathscr{X} \in O(S^{2n,n}), n \geq 2$ , it follows that  $\mathscr{X}$  is 1-connected. In that case, Proposition 2.3.1 implies that  $L_{\text{rét}}\mathscr{X}$  is 1-connected as well. Moreover, Proposition 2.3.2 implies that being a real-étale sheaf is equivalent to being  $\rho$ -periodic. In that case, it follows that  $L_{\text{rét}}\Sigma^{2,1}\mathscr{X} \simeq \Sigma^{2,1}L_{\text{rét}}\mathscr{X} \simeq \Sigma L_{\text{rét}}\mathscr{X}$ . Likewise,  $\Omega^{2,1}\Sigma L_{\text{rét}} \simeq \Omega\Sigma L_{\text{rét}}\mathscr{X}$  (use again Proposition 2.3.2). Moreover both spaces lie in  $O(S^{q,q})$  for any q, being  $\rho$ -periodic. The result follows by appeal to Proposition 4.3.3.

**Lemma 6.2.7.** Assume k is a field of finite virtual cohomological dimension and d and b are positive integers. Suppose  $\mathscr{X} \in O(S^{2,0})$ . For any finitely generated field extension K/k of transcendence degree  $\leq d$  and any integer r, there exists an integer N (depending on r, d, b and K, but not  $\mathscr{X}$ ) such that if  $\mathscr{X} \in O(S^{N,N})$  and  $\mathscr{X}_{\mathbb{O}}$  satisfies real étale descent, then

$$\pi_a(\mathscr{X})_{-b}(K) \longrightarrow \pi_a(\operatorname{L}_{r\acute{e}t}\mathscr{X})_{-b}(K)$$

is an isomorphism for  $a \leq r$ .

*Proof.* We claim that there exists a function N(v) such that for any field K with  $vcd_2(K) \leq v$  and N > N(v) the maps

$$\pi_a(\mathscr{X})_{-b}(K) \to (a_{\text{rét}}\pi_a(\mathscr{X}))_{-b}(K)$$

are isomorphisms, and also that  $a_{r\acute{e}t}\pi_a(\mathscr{X})$  is strictly  $\mathbb{A}^1$ -invariant. Assuming this, from Proposition 2.3.1 we deduce that  $\pi_a L_{r\acute{e}t}\mathscr{X} \simeq a_{r\acute{e}t}\pi_a(\mathscr{X})$ , and hence the first part of the claim concludes the proof.

For a nilpotent space  $\mathscr{Y}$ , put  $\mathscr{Y}_{tors} = \operatorname{fib}(\mathscr{Y} \to \mathscr{Y}_{\mathbb{Q}})$ . Since  $\mathscr{X}_{\mathbb{Q}}$  is rét-local, its homotopy sheaves are real étale sheaves (Proposition 2.3.1), and hence it suffices to prove the claim for  $\mathscr{X}_{tors}$ . Consider the Postnikov tower

$$\tau_{\geq q,q} K(\boldsymbol{\pi}_i \mathscr{X}, i) \to \mathcal{L}^{n+i+1,n} \mathscr{X} \to \mathcal{L}^{n+i,n} \mathscr{X}$$

of Theorem 4.1.10. This induces a tower

$$(\tau_{\geq q,q}K(\boldsymbol{\pi}_{i}\mathscr{X},i))_{tors} \to (\mathcal{L}^{n+i+1,n}\mathscr{X})_{tors} \to (\mathcal{L}^{n+i,n}\mathscr{X})_{tors}$$

By induction, it will suffice to prove the claim for  $(\tau_{\geq q,q}K(\pi_i \mathscr{X}, i))_{tors}$ . Hence we may assume that  $\mathscr{X} = \Omega^{\infty} E$ , where  $E \in SH(k)^{eff}(N)_{\geq 0}$  is torsion.

Combining the fist part of the claim with rigidity of E [EHIK21, Theorem 3.4.4] in the case v = 0 (the  $vcd_2$  of real closed fields) we see that also

$$a_{\text{rét}} \boldsymbol{\pi}_a(\Omega^\infty E) \simeq \boldsymbol{\pi}_a(\Omega^\infty(E[\rho^{-1}]))$$

(check on stalks), which is strictly  $\mathbb{A}^1$ -invariant. Hence it suffices to prove the first part of the claim, for which it suffices to prove that the homotopy groups of  $(E/\rho)(K)$  vanish in a certain range. This follows from [BEØ21, Lemma 5.23].

## 6.3 Complements

The statement of Theorem 6.2.1 had a restriction  $m \le 2n$  on the weak cellular class of the space under consideration. We now remove this assumption.

**Theorem 6.3.1.** Assume k is a perfect field with characteristic exponent p. If  $\mathscr{X} \in O(S^{m,n})$  with  $m \ge 4, n \ge 2$ , and  $\pi_*(\mathscr{X})$  is uniquely p-divisible, then

$$\operatorname{fib}(s_{\mathscr{X}}) \in O(S^{a,2n}).$$

where  $a = \min(2m - 1, m + 2n - 1)$  (which agree when m = 2n).

*Proof.* When  $m \leq 2n$ , this is precisely Theorem 6.2.1. We thus treat the case where m > 2n. We describe an inductive machine to reduce to the case m = 2n. By assumption  $\mathscr{X}$  is 1-connected, and write  $\mathscr{G} := \Omega \mathscr{X}$ . In that case, recall that  $\mathscr{X} \simeq B\mathscr{G}$  (see Paragraph 2.1.22), i.e.,  $\mathscr{X}$  can be written as a colimit over  $\Delta^{\mathrm{op}}$  with terms isomorphic to finite products  $\mathscr{G}$ . If  $\mathscr{G} \in O(S^{m,n})$  then the same thing is true for any finite product  $\mathscr{G}^{\times r}$  by appeal to Lemma 3.1.8.

Assume fib $(s_{\mathscr{G}\times n}) \in O(S^{a,b})$ . The functor  $\Omega^{2,1}\Sigma^{2,1}(-)$  preserves sifted colimits by appeal to Proposition 2.2.5(3). Since sifted colimits are stable under pullbacks by Proposition 2.1.25 we conclude that there is a fiber sequence of the form

$$\operatorname{colim}_{\Delta^{\operatorname{op}}} \operatorname{fib}(B_{\bullet}\mathscr{G} \longrightarrow \Omega^{2,1}\Sigma^{2,1}B_{\bullet}\mathscr{G}) \longrightarrow \mathscr{X} \longrightarrow \Omega^{2,1}\Sigma^{2,1}\mathscr{X}.$$

We claim that  $\operatorname{colim}_{\Delta^{\operatorname{op}}} \operatorname{fib}(B_{\bullet}\mathscr{G} \longrightarrow \Omega^{2,1}\Sigma^{2,1}B_{\bullet}\mathscr{G}) \in O(S^{a+1,b})$ . It is clear that the colimit lies in  $O(S^{a,b})$ , since each term does. It hence remains to prove that the colimit is (a - b)-connected (Corollary 3.1.26). This holds since each term in the geometric realization is (a - b)-connective and the zeroth term is contractible.<sup>2</sup>

If m > 2n, it follows that  $\Omega^{m-2n} \mathscr{X} \in O(S^{2n,n})$ . In that case, Theorem 6.2.1 implies that  $\operatorname{fib}(s_{\Omega^{m-2n}} \mathscr{X}) \in O(S^{4n-1,2n})$ . Therefore by m - 2n-fold application of the comparison result above, we conclude that

$$\operatorname{fib}(s_{\mathscr{X}}) \in O(S^{4n-1+(m-2n),2n}),$$

which is exactly what we needed to show.

*Remark* 6.3.2. Let us make some remarks on the hypotheses of the main theorem. Assume  $\mathscr{X} \in O(S^{m,n})$ . Without some hypothesis on n, Theorem 6.2.1 is false. Indeed, consider the map  $S^n \to S^{n+1,1}$  for any integer  $n \ge 2$ . In that case, [Mor12, Corollary 6.43] implies that the map  $\pi_n(S^n) \to \pi_n(S^{n+1,1})$  is the canonical map  $\mathbb{Z} \to \mathbf{GW}$ . In particular, this map fails to be an isomorphism. The assumption that  $n \ge 2$  arises in exactly one place: via the implicit appeal to Theorem 2.2.29. As a consequence, it is natural to ask whether the hypothesis on n can be weakened.

Question 6.3.3. Is the conclusion of Theorem 6.3.1 true for  $\mathscr{X} \in O(S^{m,n})$  with  $m \ge 2, n \ge 1$ ? Theorem 6.3.4. Let  $\mathscr{X} \in O(S^{p,q})$  with  $p \ge q \ge 2$ ,  $p - q \ge 2$ . In that case, for any integer  $i \ge 1$ fib $(\mathscr{X} \to \Omega^{2i,i} \Sigma^{2i,i} \mathscr{X}) \in O(S^{a,2q})$ ,

where  $a = \min(2p - 1, p + 2q - 1)$ , and likewise,

$$\operatorname{fib}(\mathscr{X} \to Q\mathscr{X}) \in O(S^{a,2q})$$

as well.

<sup>&</sup>lt;sup>2</sup>Since geometric realizations of connected spaces commute with loops, this reduces to the claim that if  $X_{\bullet}$  is a simplicial set with  $X_0 = *$  then  $\pi_0|X_{\bullet}| = *$ , which is clear.

*Proof.* The case i = 1 is Theorem 6.3.1, so assume i > 0 and consider the map  $s_{\Sigma^{2i-2,i-1}\mathscr{X}}$ . Since  $\mathscr{X} \in O(S^{p,q})$ , it follows that  $\Sigma^{2i-2,i-1}\mathscr{X} \in O(S^{p+2i-2,q+i-1})$ . In that case, Theorem 6.3.1 implies that  $\operatorname{fib}(s_{\Sigma^{2i-2,i-1}\mathscr{X}}) \in O(S^{a'(i),2(q+i-1)})$  where  $a'(i) = \min(2(p+2i-2)-1, p+2i-2+2(q+i-1))$ . Applying  $\Omega^{2i-2,i-1}$ , we see that

$$\Omega^{2i-2,i-1} \operatorname{fib}(s_{\Sigma^{2i-2,i-1}} \mathscr{X}) \in O(S^{a(i),b})$$

where a(i) = min(2p + 2i - 2 - 1, p + 2i - 2 + 2q + i - 1). The result follows by appeal to [ABH22, Proposition 3.1]. The second statement follows from the first.

# 7 Applications

In this section, we deduce various concrete consequences of Theorem 6.2.1. In particular, we establish Theorem 3 from the introduction.

## 7.1 Murthy's conjecture and complements

The strategy of argument is laid out in [AF15] and it is observed in [AF13] that a suitable version of the Freudenthal suspension theorem would suffice to establish Murthy's conjecture; see [AF21, §4.3-4.4] for further details of this approach. The discussion of [AF21] reduces Murthy's conjecture to the triviality of a certain sheaf [AF21, Conjecture 4.3.2]; we describe in the next section how to establish this vanishing statement. However, to establish Murthy's conjecture we actually need significantly less, as we now describe.

**Theorem 7.1.1** (Murthy's conjecture in char. 0). Assume k is an algebraically closed field having characteristic 0 and X is a smooth affine d + 1-fold,  $d \ge 1$ . If E is a rank d vector bundle on X, then E splits off a trivial rank 1 summand if and only if  $0 = c_d(E) \in CH^d(X)$ .

*Proof.* By [AHW17, Theorem 1], the vector bundle E corresponds to an element of  $[X, BGL_d]$ . Moreover, by [AHW18, Theorem 2.2.4], there is a fiber sequence

$$S^{2d-1,d} \longrightarrow BGL_{d-1} \rightarrow BGL_{d}$$

where we have used the identification  $GL_d/GL_{d-1} \simeq Q_{2d-1} \simeq S^{2d-1,d}$ . We proceed by analyzing the Moore–Postnikov factorization of the stabilization map  $BGL_{d-1} \rightarrow BGL_d$  (see [AF15, Theorem 6.1.1] for a convenient summary).

Recall [AF15, Proposition 6.3.1]: since X has dimension d, a lift of an element of  $[X, BGL_d]_{\mathbb{A}^1}$  along the map  $BGL_{d-1} \rightarrow BGL_d$  exists if and only if a primary and secondary obstruction vanish. Moreover, the primary obstruction is the twisted Euler class

$$e(E) \in H^d(X, \pi_{d-1,0}(S^{2d-1,d})(\det E)),$$

but under the hypothesis that k is algebraically closed the primary obstruction vanishes if and only if  $c_d(E)$  vanishes. By hypothesis, it suffices to show that the secondary obstruction vanishes, and by [AF15, Corollary 6.3.3] this obstruction lies in

$$o_2(E) \in H^{d+1}(X, \pi_{d,0}(S^{2d-1,d})(\det E)).$$

We will deduce this vanishing from Theorem 6.2.1 via analysis of the Gersten complex of the homotopy sheaf  $\pi_{d,0}(S^{2d-1,d})$ .

In order to set things up, we recall some facts about the sheaf  $\pi_{d,0}(S^{2d-1,d})$ . First, an explicit construction via "Suslin matrices" yields maps:

$$S^{2n-1,n} \longrightarrow \Omega^{-2n+1,-n}(\mathbb{Z} \times B_{\acute{e}t} \mathcal{O})$$

(where we make sense of the negative loopings by geometric Bott periodicity via the work of [ST15]). At the level of homotopy sheaves, the induced maps

$$\pi_{d+j,j}(S^{2d-1,d}) \longrightarrow \mathbf{GW}_{d+1-j}^{d-j}$$

are epimorphisms for  $j \ge d - 3$  [AF17, Theorem 5].

The cohomology of strictly  $\mathbb{A}^1$ -invariant sheaves may be computed by appeal to Gersten resolutions. In more detail, if **A** is a strictly  $\mathbb{A}^1$ -invariant sheaf on  $\mathscr{X}$  and *L* is a line bundle on *X*, then  $H^{d+1}(X, \mathbf{A}(L))$  can be computed by means of a Gersten resolution and is the cokernel of a map

$$\bigoplus_{x \in X^{(d)}} \mathbf{A}_{-d}(\kappa_x; L_x) \longrightarrow \bigoplus_{x \in X^{(d+1)}} \mathbf{A}_{-d-1}(\kappa_x; L_x).$$

The fact that the morphism  $\pi_{d+j,j}(S^{2d-1,d}) \to \mathbf{GW}_{d+1-j}^{d-j}$  described in the preceding paragraph is an epimorphism for  $j \ge d-3$  implies that the induced map on cohomology

$$H^{d+1}(X, \pi_{d,0}(S^{2d-1,d})(\det E)) \longrightarrow H^{d+1}(X, \mathbf{GW}^d_{d+1}(\det E))$$

is itself an epimorphism. Under the assumptions on k, the latter group vanishes by appeal to [AF15, Theorem 3.7.1]. In other words, the image of the secondary obstruction  $o_2(E)$  in the target is zero. Set

$$\mathbf{A}(d) := \ker(\boldsymbol{\pi}_{d,0}(S^{2d-1,d}) \longrightarrow \mathbf{GW}_{d+1}^d).$$

To establish the result it suffices to show that  $H^{d+1}(X, \mathbf{A}(d)(\det E))$  vanishes. For this vanishing, we appeal to Theorem 6.2.1.

Indeed, Theorem 6.2.1 tells us that the map  $S^{2d-1,d} \to \Omega^{2,1}S^{2d+1,d+1}$  has fiber lying in  $O(S^{4d-3,2d})$ . In particular, if  $d \ge 3$  and  $j \ge 0$ , then there are epimorphisms

$$\pi_{d+j,j}(S^{2d-1,d}) \to \pi_{d+j,j}(\Omega^{2,1}S^{2d+1,d+1}).$$

On the other hand, by [AF15, Theorem 4.4.1] we may identify A(3) with the sheaf  $F_5$  which fits into an exact sequence of the form:

$$\mathbf{I}^6 \longrightarrow \mathbf{F}_5 \longrightarrow \mathbf{K}_5^M/24 \longrightarrow 0.$$

The contractions of these sheaves are well-known (see, e.g., [AF15, Example 2.3.2]). Indeed, we deduce that  $\mathbf{A}(3)_{-4}$  is an extension of  $\mathbf{K}_1^M/24$  by  $\mathbf{I}^2$ . The discussion above implies that  $\mathbf{A}(d)_{-d-1}$  is a quotient of  $(\mathbf{F}_5)_{-3}$ . In particular, for  $x \in X^{(d+1)}$ , since  $\kappa_x$  is algebraically closed, we see that  $\mathbf{I}^2(\kappa_x) = \mathbf{K}_1^M/24(\kappa_x) = 0$ . The required vanishing of

$$H^{d+1}(X, \mathbf{A}(d)(\det \xi)),$$

then follows from the description of the Gersten resolution above and the fact that the terms  $\mathbf{A}(d)_{-d-1}(\kappa_x; (\det \xi)_x)$  are isomorphic as abstract groups to  $\mathbf{A}(d)_{-d-1}(\kappa_x)$ .

*Remark* 7.1.2. The vanishing of  $H^{d+1}(X, \pi_{d,0}(S^{2d-1,d})(\det E))$  established in the theorem above also implies Suslin's cancellation conjecture in corank 1 [AF21, Conjecute 4.5.1] over algebraically closed fields having characteristic 0 by appeal to [Du22, Theorem 1.2]. However, in [Fas21], J. Fasel establishes Suslin's cancellation conjecture in corank 1 for smooth affine *d*-folds over an algebraically closed field *k* in which (d-1)! is invertible without establishing the above vanishing.

## 7.2 Computations

We now describe computations of unstable homotopy sheaves of motivic spheres. Recall that the unit map  $1 \rightarrow KO$  desuspends to morphisms

$$S^{2n-i,n} \longrightarrow \Omega^{-2n+i,n}(\mathbb{Z} \times B_{\acute{e}t} O)$$

by [ST15, Theorems 1 and 3]; the loop spaces admit explicit geometric descriptions. It was observed in [AF17, Theorem 5] that the maps with source  $S^{2n-1,n}$  are given by explicit "Suslin matrices", and in [ADF17, Theorem 4.3.6] that those with source  $S^{2n,n}$  are realized by explicit vector bundles.

On the other hand, consider the motivic Hopf map  $\nu : S^{7,4} \to S^{4,2}$ , obtained by performing the Hopf construction on the multiplication map  $SL_2 \times SL_2 \to SL_2$ . This map is also part of the motivic *J*-homomorphism. The motivic Hopf map induces morphisms  $\nu_d : S^{2d+2,d+2} \to S^{2d-1,d}$ . The induced map on homotopy sheaves factors through a map

$$(\nu_d)_*: \mathbf{K}^M_{d+2-j}/24 \longrightarrow \pi^{\mathbb{A}^1}_{d+j,j}(S^{2d-1,d});$$

this is [AFW20, Theorem 3.18(2)], which builds on [AWW17, Theorem 5.2.5]. With those observations in mind, the following result, establishes [AF21, Conjecture 4.3.2], which is a stronger form of [AWW17, Conjecture 5.2.9].

**Theorem 7.2.1.** Assume k is a field having characteristic 0. For any integer  $d \ge 4$  there is a short exact sequence of the form

$$0 \longrightarrow \mathbf{K}_{d+2-j}^{M}/24 \xrightarrow{(\nu_d)_*} \pi_{d+j,j}^{\mathbb{A}^1}(S^{2d-1,d}) \longrightarrow \mathbf{GW}_{d+1-j}^{d-j},$$

which is an isomorphism if  $j \ge d - 3$ .

*Proof.* Combine Theorem 6.2.1 with [RSØ19, Theorem 5.5].

We include here a number of further unstable computations of homotopy sheaves.

**Theorem 7.2.2.** If k is a perfect field, then the following statements hold.

*1.* For any  $d \ge 0$  there are isomorphisms

$$\pi_{4+r,r}(S^{3+d,d}) \xrightarrow{\sim} \pi_{d+r,r}(S^{2d-1,d}).$$

2. For any  $d \ge 1$  there is a short exact sequence

$$0 \longrightarrow \operatorname{coker}(\pi_{6+r,r}(S^{4+d,d}) \longrightarrow \mathbf{K}_{2d-r}^{MW}/h^{\frac{(1+(-1)^{d+1})}{2}}) \longrightarrow \pi_{3+r,r}(S^{2+d,d}) \longrightarrow \pi_{4+r,r}(S^{3+d,d}) \longrightarrow 0$$

3. If k has characteristic 0, then for any  $d \ge 2$ , there is a short exact sequence of the form

$$0 \longrightarrow \mathbf{K}_{d+2-r}^{M}/24 \longrightarrow \pi_{4+r,r}(S^{3+d,d}) \longrightarrow \mathbf{GW}_{d+1-r}^{d-r};$$

where the right vertical map is surjective for  $r \ge d - 3$ .

*Proof.* The first point is an immediate consequence of the simplicial EHP exact sequence [AWW17, Theorem 3.2.1]. The second point follows from the simplicial EHP sequence in conjunction with the computation of the  $d_1$ -differential [AWW17, Theorem 4.4.1]: when d is even, the differential is 0, while if d is odd, it is multiplication by h. The final point follows from the first point in conjunction with Theorem 7.2.1.

*Remark* 7.2.3. In a range, depending on r, the cokernel in Point (2) of the previous statement coincides with  $\mathbf{K}_{2d-r}^{MW}/h^{\frac{(1+(-1)^{d+1})}{2}}$  because the source groups are zero. Lacking applications, we do not make this precise.

# References

- [ABFJ20] M. Anel, G. Biedermann, E. Finster, and A. Joyal. A generalized Blakers-Massey theorem. J. Topol., 13(4):1521– 1553, 2020. 20, 21
- [ABH22] A. Asok, T. Bachmann, and M.J. Hopkins. On the Whitehead theorem for nilpotent motivic spaces. 2022. Available at https://arxiv.org/abs/2210.05933. 8, 9, 13, 15, 20, 26, 31, 32, 37, 40, 49, 53, 56, 57, 60
- [AD01] G. Z. Arone and W. G. Dwyer. Partition complexes, Tits buildings and symmetric products. Proc. London Math. Soc. (3), 82(1):229–256, 2001. 4
- [Ada95] J. F. Adams. Stable homotopy and generalised homology. Chicago Lectures in Mathematics. University of Chicago Press, Chicago, IL, 1995. Reprint of the 1974 original. 2
- [ADF17] A. Asok, B. Doran, and J. Fasel. Smooth models of motivic spheres and the clutching construction. Int. Math. Res. Not. IMRN, (6):1890–1925, 2017. 34, 35, 42, 62
- [AF13] A. Asok and J. Fasel. Toward a meta-stable range in  $\mathbb{A}^1$ -homotopy theory of punctured affine spaces. *Oberwolfach Rep.*, 10(2):1892–1895, 2013. 3, 60
- [AF14a] A. Asok and J. Fasel. Algebraic vector bundles on spheres. J. Topol., 7(3):894–926, 2014. 42
- [AF14b] A. Asok and J. Fasel. A cohomological classification of vector bundles on smooth affine threefolds. *Duke Math. J.*, 163(14):2561–2601, 2014. **3**
- [AF15] A. Asok and J. Fasel. Splitting vector bundles outside the stable range and A<sup>1</sup>-homotopy sheaves of punctured affine spaces. J. Amer. Math. Soc., 28(4):1031–1062, 2015. 3, 60, 61
- [AF17] A. Asok and J. Fasel. An explicit KO-degree map and applications. J. Topol., 10(1):268–300, 2017. 61, 62
- [AF21] A. Asok and J. Fasel. Vector bundles on algebraic varieties, 2021. Available at https://arxiv.org/abs/2111.03107\right). 60, 61, 62
- [AFH22] A. Asok, J. Fasel, and M. J. Hopkins. Localization and nilpotent spaces in A<sup>1</sup>-homotopy theory. *Compos. Math.*, 158(3):654–720, 2022. 9, 38, 41
- [AFW20] A. Asok, J. Fasel, and B. Williams. Motivic spheres and the image of the Suslin-Hurewicz map. *Invent. Math.*, 219(1):39–73, 2020. 62
- [AH11] A. Asok and C. Haesemeyer. Stable  $\mathbb{A}^1$ -homotopy and *R*-equivalence. J. Pure Appl. Algebra, 215(10):2469–2472, 2011. 18
- [AHW17] A. Asok, M. Hoyois, and M. Wendt. Affine representability results in  $\mathbb{A}^1$ -homotopy theory, I: vector bundles. *Duke Math. J.*, 166(10):1923–1953, 2017. 3, 60
- [AHW18] A. Asok, M. Hoyois, and M. Wendt. Affine representability results in A<sup>1</sup>-homotopy theory, II: Principal bundles and homogeneous spaces. *Geom. Topol.*, 22(2):1181–1225, 2018. **35**, 60
- [AHW19] A. Asok, M. Hoyois, and M. Wendt. Generically split octonion algebras and A<sup>1</sup>-homotopy theory. Algebra Number Theory, 13(3):695–747, 2019. 35
- [AHW20] A. Asok, M. Hoyois, and M. Wendt. Affine representability results in A<sup>1</sup>-homotopy theory III: finite fields and complements. *Algebr. Geom.*, 7(5):634–644, 2020. 35
- [ALP17] A. Ananyevskiy, M. Levine, and I. Panin. Witt sheaves and the  $\eta$ -inverted sphere spectrum. J. Topol., 10(2):370– 385, 2017. 20
- [AN19] A. Ananyevskiy and A. Neshitov. Framed and MW-transfers for homotopy modules. *Selecta Math.* (*N.S.*), 25(2):Paper No. 26, 41, 2019. 16
- [Aso22] A. Asok. Affine representability of quadrics revisited. J. Algebra, 608:37–51, 2022. 35
- [ATJLSS03] L. Alonso Tarrío, A. Jeremías López, and M. J. Souto Salorio. Construction of t-structures and equivalences of derived categories. Trans. Amer. Math. Soc., 355(6):2523–2543, 2003. 15
- [AWW17] A. Asok, K. Wickelgren, and T.B. Williams. The simplicial suspension sequence in A<sup>1</sup>-homotopy. *Geom. Topol.*, 21(4):2093–2160, 2017. 7, 10, 33, 62
- [Ayo07] J. Ayoub. Les six opérations de Grothendieck et le formalisme des cycles évanescents dans le monde motivique. I. Astérisque, (314):x+466 pp. (2008), 2007. 15
- [Bac17] T. Bachmann. The generalized slices of Hermitian K-theory. J. Topol., 10(4):1124–1144, 2017. 16, 29, 30, 33
- [Bac18a] T. Bachmann. Motivic and real étale stable homotopy theory. Compos. Math., 154(5):883–917, 2018. 19, 20, 32

[Bac18b]	Tom Bachmann. On the conservativity of the functor assigning to a motivic spectrum its motive. <i>Duke Math. J.</i> , 167(8):1525–1571, 06 2018. arXiv:1506.07375. 32
[Bac21a]	T. Bachmann. The zeroth $\mathbb{P}^1$ -stable homotopy sheaf of a motivic space. <i>Journal of the Institute of Mathematics of Jussieu</i> , page 1–25, 2021. Available at https://doi.org/10.1017/S1474748021000396. 4, 18, 20
[Bac21b]	Tom Bachmann. Motivic tambara functors. <i>Mathematische Zeitschrift</i> , 297:1825–1852, 2021. arXiv:1807.02981. 16
[BCD <sup>+</sup> 20]	T. Bachmann, B. Calmès, F. Déglise, J. Fasel, and PA. Østvær. Milnor-Witt Motives, 2020. 15
[BE21]	Tom Bachmann and Elden Elmanto. Voevodsky's slice conjectures via hilbert schemes. <i>Algebraic Geometry</i> , 8(5):626–636, 2021. arXiv:1912.01595. 31
[BEØ21]	T. Bachmann, E. Elmanto, and PA. Østvær. On étale motivic spectra and Voevoedsky's convergence conjecture. 2021. Available at https://arxiv.org/abs/2003.04006. 4, 33, 58
[BM51]	A. L. Blakers and W. S. Massey. The homotopy groups of a triad. I. Ann. of Math. (2), 53:161-205, 1951. 4
[Bon20]	Mikhail V. Bondarko. On infinite effectivity of motivic spectra and the vanishing of their motives. <i>Doc. Math.</i> , 25:811–840, 2020. 32
[BR85]	P. Bardsley and R. W. Richardson. Étale slices for algebraic transformation groups in characteristic <i>p. Proc. London Math. Soc.</i> (3), 51(2):295–317, 1985. 48
[BY20]	T. Bachmann and M. Yakerson. Towards conservativity of $\mathbb{G}_m$ -stabilization. Geom. Topol., 24(4):1969–2034, 2020. 4, 15, 18, 29, 30, 32, 38
[ByB73]	A. Biał ynicki Birula. Some theorems on actions of algebraic groups. Ann. of Math. (2), 98:480-497, 1973. 34
[CD19]	DC. Cisinski and F. Déglise. <i>Triangulated categories of mixed motives</i> . Springer Monographs in Mathematics. Springer, Cham, [2019] ©2019. 17
[Cho14]	U. Choudhury. Connectivity of motivic H-spaces. Algebr. Geom. Topol., 14(1):37-55, 2014. 7, 8
[Del09]	P. Deligne. Voevodsky's lectures on motivic cohomology 2000/2001. In <i>Algebraic topology</i> , volume 4 of <i>Abel Symp.</i> , pages 355–409. Springer, Berlin, 2009. 43
[DF96]	E. Dror-Farjoun. <i>Cellular spaces, null spaces and homotopy localization</i> , volume 1622 of <i>Lecture Notes in Mathematics</i> . Springer-Verlag, Berlin, 1996. 3, 20, 21, 23
[DH21]	S. Devalapurkar and P. Haine. On the James and Hilton-Milnor Splittings, and the metastable EHP sequence. <i>Doc. Math.</i> , 26:1423–1464, 2021. 41
[Du22]	P. Du. Enumerating non-stable vector bundles. Int. Math. Res. Not. IMRN, (19):14797-14864, 2022. 61
[EHIK21]	E. Elmanto, M. Hoyois, R. Iwasa, and S. Kelly. Cdh descent, cdarc descent, and Milnor excision. <i>Math. Ann.</i> , 379(3-4):1011–1045, 2021. 58
[EHK <sup>+</sup> 21]	E. Elmanto, M. Hoyois, A. A. Khan, V. Sosnilo, and M. Yakerson. Motivic infinite loop spaces. <i>Camb. J. Math.</i> , 9(2):431–549, 2021. 15, 17
[EK20a]	E. Elmanto and A. A. Khan. Perfection in motivic homotopy theory. <i>Proc. Lond. Math. Soc. (3)</i> , 120(1):28–38, 2020. 14
[EK20b]	E. Elmanto and H. Kolderup. On modules over motivic ring spectra. Ann. K-Theory, 5(2):327-355, 2020. 14
[Fas21]	Jean Fasel. Suslin's cancellation conjecture in the smooth case, 2021. Available at: https://arxiv.org/abs/2111.13088.61
[Fel20]	N. Feld. Milnor-Witt cycle modules. J. Pure Appl. Algebra, 224(7):106298, 44, 2020. 15, 18
[Fel21]	N. Feld. Morel homotopy modules and Milnor-Witt cycle modules. Doc. Math., 26:617-659, 2021. 16
[Fre38]	H. Freudenthal. Über die Klassen der Sphärenabbildungen I. Große Dimensionen. <i>Compositio Math.</i> , 5:299–314, 1938. 4
[HKØ17]	M. Hoyois, S. Kelly, and PA. Østvær. The motivic Steenrod algebra in positive characteristic. <i>J. Eur. Math. Soc.</i> ( <i>JEMS</i> ), 19(12):3813–3849, 2017. 45, 50
[Hoy14]	M. Hoyois. A quadratic refinement of the Grothendieck-Lefschetz-Verdier trace formula. <i>Algebr. Geom. Topol.</i> , 14(6):3603–3658, 2014. 6
[Hoy15]	M. Hoyois. From algebraic cobordism to motivic cohomology. J. Reine Angew. Math., 702:173–226, 2015. 6, 14, 15

- [Hoy17] M. Hoyois. The six operations in equivariant motivic homotopy theory. *Adv. Math.*, 305:197–279, 2017. 5, 6, 12, 43, 47, 48
- [HW19] C. Haesemeyer and C. A. Weibel. The norm residue theorem in motivic cohomology, volume 200 of Annals of Mathematics Studies. Princeton University Press, Princeton, NJ, 2019. 35
- [Kel17] S. Kelly. Voevodsky motives and *l*dh descent. *Astérisque*, 391, 2017. 50
- [Lev06] M. Levine. Chow's moving lemma and the homotopy coniveau tower. *K-Theory*, 37(1-2):129–209, 2006. 31
- [Lev08] M. Levine. The homotopy coniveau tower. J. Topol., 1(1):217–267, 2008. 14, 30, 31
- [Lev10] M. Levine. Slices and transfers. *Doc. Math.*, (Extra vol.: Andrei A. Suslin sixtieth birthday):393–443, 2010. 18
- [Lev13] M. Levine. Convergence of Voevodsky's slice tower. Doc. Math., 18:907–941, 2013. 4, 17, 57
- [Lun73] D. Luna. Slices étales. In Sur les groupes algébriques, Bull. Soc. Math. France, Mém. 33, pages 81–105. Soc. Math. France, Paris, 1973. 48
- [Lur09] J. Lurie. *Higher topos theory*, volume 170 of *Annals of Mathematics Studies*. Princeton University Press, Princeton, NJ, 2009. 6, 8, 9, 11, 13, 16, 17, 22, 24
- [Lur17] J. Lurie. Higher Algebra. 2017. Available at https://www.math.ias.edu/~lurie/papers/HA.pdf. 11, 12, 22, 28
- [Lur18] J. Lurie. Spectral Algebraic Geometry. 2018. Available at https://www.math.ias.edu/~lurie/papers/HA.pdf. 16
- [LYZR19] M. Levine, Y. Yang, G. Zhao, and J. Riou. Algebraic elliptic cohomology theory and flops I. Math. Ann., 375(3-4):1823–1855, 2019. 14
- [Mor04] F. Morel. An introduction to  $\mathbb{A}^1$ -homotopy theory. In *Contemporary developments in algebraic K-theory*, ICTP Lect. Notes, XV, pages 357–441. Abdus Salam Int. Cent. Theoret. Phys., Trieste, 2004. 15
- [Mor05] F. Morel. The stable  $\mathbb{A}^1$ -connectivity theorems. *K*-Theory, 35(1-2):1–68, 2005. 15
- [Mor11] F. Morel. On the Friedlander-Milnor conjecture for groups of small rank. In *Current developments in mathematics,* 2010, pages 45–93. Int. Press, Somerville, MA, 2011. 18
- [Mor12] F. Morel. A<sup>1</sup>-algebraic topology over a field, volume 2052 of Lecture Notes in Mathematics. Springer, Heidelberg, 2012. 2, 6, 7, 13, 15, 18, 25, 41, 59
- [Mur99] M. P. Murthy. A survey of obstruction theory for projective modules of top rank. In Algebra, K-theory, groups, and education (New York, 1997), volume 243 of Contemp. Math., pages 153–174. Amer. Math. Soc., Providence, RI, 1999. 3
- [MV99] F. Morel and V. Voevodsky. A<sup>1</sup>-homotopy theory of schemes. *Publications Mathématiques de l'Institut des Hautes Études Scientifiques*, 90(1):45–143, 1999. 2, 6, 7, 44
- [Nak57] M. Nakaoka. Cohomology of symmetric products. J. Inst. Polytech. Osaka City Univ. Ser. A, 8:121–145, 1957. 4
- [Nee01] A. Neeman. *Triangulated categories*, volume 148 of *Annals of Mathematics Studies*. Princeton University Press, Princeton, NJ, 2001. 14
- [Pel11] P. Pelaez. Multiplicative properties of the slice filtration. *Astérisque*, (335):xvi+289, 2011. 14
- [PW21] I. Panin and C. Walter. Quaternionic Grassmannians and Borel classes in algebraic geometry. Algebra i Analiz, 33(1):136–193, 2021. 34
- [Rez14] C. Rezk. When are homotopy colimits compatible with homotopy base change?, 2014. Available at https://faculty.math.illinois.edu/~rezk/i-hate-the-pi-star-kan-condition.pdf. 11
- [Rio05] J. Riou. Dualité de Spanier-Whitehead en géométrie algébrique. C. R. Math. Acad. Sci. Paris, 340(6):431–436, 2005. 14
- [RØ08] O. Röndigs and P.-A. Østvær. Modules over motivic cohomology. Adv. Math., 219(2):689–727, 2008. 14, 45
- [Rob15] M. Robalo. *K*-theory and the bridge from motives to noncommutative motives. *Adv. Math.*, 269:399–550, 2015.
- [RSØ19] O. Röndigs, M. Spitzweck, and P.-A. Østvær. The first stable homotopy groups of motivic spheres. Ann. of Math.
   (2), 189(1):1–74, 2019. 4, 62
- [Sch94] C. Scheiderer. *Real and étale cohomology*, volume 1588 of *Lecture Notes in Mathematics*. Springer-Verlag, Berlin, 1994. 19

[SGA03]	Revêtements étales et groupe fondamental (SGA 1), volume 3 of Documents Mathématiques (Paris) [Mathematical Documents (Paris)]. Société Mathématique de France, Paris, 2003. Séminaire de géométrie algébrique du Bois Marie 1960–61. [Algebraic Geometry Seminar of Bois Marie 1960-61], Directed by A. Grothendieck, With two papers by M. Raynaud, Updated and annotated reprint of the 1971 original [Lecture Notes in Math., 224, Springer, Berlin; MR0354651 (50 #7129)]. 43
[SØ12]	M. Spitzweck and PA. Østvær. Motivic twisted K-theory. Algebr. Geom. Topol., 12(1):565-599, 2012. 14
[ST15]	M. Schlichting and G. S. Tripathi. Geometric models for higher Grothendieck-Witt groups in $\mathbb{A}^1$ -homotopy theory. <i>Math. Ann.</i> , 362(3-4):1143–1167, 2015. 61, 62
[Sus82]	A. A. Suslin. Mennicke symbols and their applications in the K-theory of fields. In Algebraic K-theory, Part I (Oberwolfach, 1980), volume 966 of Lecture Notes in Math., pages 334–356. Springer, Berlin-New York, 1982. 42
[SV96]	A. Suslin and V. Voevodsky. Singular homology of abstract algebraic varieties. <i>Invent. Math.</i> , 123(1):61–94, 1996. 44, 46
[SV00]	A. Suslin and V. Voevodsky. Relative cycles and Chow sheaves. In Cycles, transfers, and motivic homology theories, volume 143 of Ann. of Math. Stud., pages 10–86. Princeton Univ. Press, Princeton, NJ, 2000. 44, 46
[tD87]	T. tom Dieck. <i>Transformation groups</i> , volume 8 of <i>De Gruyter Studies in Mathematics</i> . Walter de Gruyter & Co., Berlin, 1987. 2
[Voe98]	V. Voevodsky. A <sup>1</sup> -homotopy theory. In <i>Proceedings of the International Congress of Mathematicians, Vol. I</i> ( <i>Berlin, 1998</i> ), number Extra Vol. I, pages 579–604, 1998. 2
[Voe00a]	V. Voeovodsky, 2000. Available at https://www.math.ias.edu/Voevodsky/files/files-annotated/Dropbox/Pul 2
[Voe00b]	V. Voevodsky. Triangulated categories of motives over a field. In <i>Cycles, transfers, and motivic homology theories</i> , volume 143 of <i>Ann. of Math. Stud.</i> , pages 188–238. Princeton Univ. Press, Princeton, NJ, 2000. 16
[Voe02]	V. Voevodsky. Motivic cohomology groups are isomorphic to higher Chow groups in any characteristic. <i>Int. Math. Res. Not.</i> , (7):351–355, 2002. 57
[Voe04]	V. Voevodsky. On the zero slice of the sphere spectrum. <i>Tr. Mat. Inst. Steklova</i> , 246(Algebr. Geom. Metody, Svyazi i Prilozh.):106–115, 2004. 4, 14, 34
[Voe10a]	V. Voevodsky. Cancellation theorem. <i>Doc. Math.</i> , (Extra vol.: Andrei A. Suslin sixtieth birthday):671–685, 2010. 57
[Voe10b]	V. Voevodsky. Motivic Eilenberg-Maclane spaces. Publ. Math. Inst. Hautes Études Sci., (112):1–99, 2010. 42, 43, 45, 46, 51
[Voe10c]	V. Voevodsky. Unstable motivic homotopy categories in Nisnevich and cdh-topologies. J. Pure Appl. Algebra, 214(8):1399–1406, 2010. 49
[WW19]	K. Wickelgren and T.B. Williams. The simplicial EHP sequence in $\mathbb{A}^1$ -algebraic topology. <i>Geom. Topol.</i> , 23(4):1691–1777, 2019. 41

A. Asok, Department of Mathematics, University of Southern California, 3620 S. Vermont Ave., Los Angeles, CA 90089-2532, United States; E-mail address: asok@usc.edu

T. Bachmann, Department of Mathematics, University of Oslo, Postboks 1053 Blindern, 0316 Oslo, Norway; E-mail address: tom.bachmann@zoho.com

M.J. Hopkins, Department of Mathematics, Harvard University, One Oxford Street, Cambridge, MA 02138, United States *E-mail address:* mjh@math.harvard.edu no