

Signals of strong parity violation in deep inelastic scattering

Alessandro Bacchetta,^{1,2,*} Matteo Cerutti,^{1,2,†} Ludovico Manna,^{1,3,‡} Marco Radici,^{2,§} and Xiaochao Zheng^{4,¶}

¹*Dipartimento di Fisica, Università di Pavia, via Bassi 6, I-27100 Pavia, Italy*

²*INFN Sezione di Pavia, via Bassi 6, I-27100 Pavia, Italy*

³*Dipartimento di Scienza della Terra e dell'Ambiente,
Università di Pavia, via Ferrata 7, I-27100 Pavia, Italy*

⁴*University of Virginia, Charlottesville, VA 22904, USA*

(Dated: Wednesday 17th January, 2024, 03:35)

We include strong parity-violating contributions to inclusive deep inelastic scattering (DIS) of longitudinally polarized leptons off an unpolarized target. At variance with standard results, we obtain nonvanishing parity-violating structure functions in the case of pure photon exchange. The addition of these strong parity-violating contributions improves the description of existing experimental data on DIS parity-violating asymmetries. We find the size of these contributions small but exhibiting a deviation from zero of about 1.5σ . The associated p -value is 0.063, indicating that the probability of making an error by rejecting the hypothesis of no parity-violating contributions is 6.3%, which is small but not negligible. Further improvement on the limit of the strong parity-violation can be expected from the future SoLID program at Jefferson Lab.

I. INTRODUCTION

Would the internal structure of the proton be identical in a mirrored world? According to the Standard Model (SM), it should remain the same, because Quantum ChromoDynamics (QCD) is invariant under parity (P) transformations. However, there is no first principle to guarantee parity invariance in strong interactions, and in this article we consider the impact of its violation (i.e., “strong P violation”) on the internal structure of the nucleons, and we show that a virtual photon probing an unpolarized proton may see more left-handed than right-handed quarks.

The observation of strong P violation would have far-fetched consequences beyond the specific problem of studying the internal structure of the nucleons. The violation of charge-parity (CP) symmetry is the most plausible explanation of the matter-antimatter imbalance in the Universe, but CP violation in the SM electroweak sector alone is not sufficient to justify such an asymmetry. An alternative solution is to go beyond the SM and introduce CP violation in the QCD Lagrangian. This can be done, e.g., with the addition of the so-called “ θ -term,” which is however constrained to be extremely small by experimental evidence. Other higher-dimensional CP-violating operators have been taken into consideration in the context of SM Effective Field Theory (SMEFT) framework, which is based on the assumption that effective low-energy modifications can arise from new physics at significantly heavier scales (see, e.g., Refs. [1–3]). All these extensions should give observable effects. The most studied one is the generation of permanent electric dipole moments in various particles (see, e.g., [4] and references therein). In the last thirty years, no experiment has been able to observe non-zero strong CP violation.

In this article, we show that strong P violation could be observable also in the details of the internal structure of the nucleon. The relaxation of P invariance leads to the introduction of new terms in the hadronic tensor for the Deep-Inelastic Scattering (DIS) process, generated by new Parton Distribution Functions (PDFs). The new P-odd terms can be either C-even (CP-odd) or C-odd (CP-even).

We focus our interest mainly on DIS of a longitudinally polarized lepton beam off an unpolarized proton or nuclear target and we briefly discuss the case of DIS of an unpolarized lepton beam off a longitudinally polarized target. In the first case, a new contribution in the structure function F_3 appears in the pure photon-exchange channel. Such

*Electronic address: alessandro.bacchetta@unipv.it
ORCID: 0000-0002-8824-8355

†Electronic address: matteo.cerutti@pv.infn.it
ORCID: 0000-0001-7238-5657

‡Electronic address: ludovico.manna01@ateneopv.it
ORCID: 0009-0003-7952-157X

§Electronic address: marco.radici@pv.infn.it
ORCID: 0000-0002-4542-9797

¶Electronic address: xiaochao@jlab.org
ORCID: 0000-0001-7300-2929

contribution would be non-zero only with the presence of a new strong P-violating (PV) PDF. In order to estimate the size of such parton density, we perform a fit to available experimental data from HERA, SLAC, and Jefferson Lab (JLab) that can be sensitive to this kind of PV effects.

Our study could open the door for new analyses of experimental data from DIS processes with different lepton beam polarizations and charges. In particular, we perform impact studies of future measurements described in the Solenoidal Large Intensity Device (SoLID) program [5] at JLab 12 GeV, and the future Electron Ion Collider (EIC) [6, 7].

II. FORMALISM

The expression of the cross section for neutral-current inclusive DIS with an initial electron or positron off an unpolarized target reads

$$\frac{d^2\sigma}{dx_B dy} = \frac{2\pi\alpha^2}{x_B y Q^2} \left[\left(Y_+ + R^2 y^2 / 2 \right) (F_{2,UU} + \lambda F_{2,LU}) - y^2 (F_{L,UU} + \lambda F_{L,LU}) - Y_- (x_B F_{3,UU} + \lambda x_B F_{3,LU}) \right], \quad (1)$$

where α is the fine structure constant, y is the inelasticity, x_B is the Bjorken variable, Q^2 is the negative of the 4-momentum transfer squared of the scattering, $R = 2Mx_B/Q$, and $Y_{\pm} = 1 \pm (1-y)^2$. For convenience, we explicitly separated the terms proportional to λ , the helicity of the electron or positron. More details about the derivation of this expression are given in Appendix A.

The structure functions involved in Eq. (1) can be expressed as

$$F_{2,UU}(x_B, Q^2) = F_2^{(\gamma)} - g_V^e \eta_{\gamma Z} F_2^{(\gamma Z)} + (g_V^e{}^2 + g_A^e{}^2) \eta_Z F_2^{(Z)}, \quad (2)$$

$$F_{2,LU}(x_B, Q^2) = g_A^e \eta_{\gamma Z} F_2^{(\gamma Z)} - 2g_V^e g_A^e \eta_Z F_2^{(Z)}, \quad (3)$$

$$F_{3,UU}(x_B, Q^2) = g_A^e \eta_{\gamma Z} F_3^{(\gamma Z)} - 2g_V^e g_A^e \eta_Z F_3^{(Z)}, \quad (4)$$

$$F_{3,LU}(x_B, Q^2) = F_3^{(\gamma)} - g_V^e \eta_{\gamma Z} F_3^{(\gamma Z)} + (g_V^e{}^2 + g_A^e{}^2) \eta_Z F_3^{(Z)}, \quad (5)$$

where

$$\eta_{\gamma Z} = \left(\frac{G_F M_Z^2}{2\sqrt{2}\pi\alpha} \right) \left(\frac{Q^2}{Q^2 + M_Z^2} \right), \quad \eta_Z = \eta_{\gamma Z}^2, \quad (6)$$

with G_F the Fermi constant and M_Z the Z^0 mass. The $g_{V,A}^e$ are electron's neutral weak couplings to the Z^0 . For positron scattering, one would change the sign of terms containing g_A^e , affecting Eqs. (3),(4). The two structure functions $F_{L,UU}$ and $F_{L,LU}$ have the same decomposition as the corresponding F_2 ones, Eqs. (2) and (3).

The above results correspond to the standard literature (see, e.g., Ref. [8] and the PDG review [9]). The only difference is that in the SM there is no contribution to the F_3 structure function from pure- γ exchange: $F_3^{(\gamma)}$ is the new ingredient in our analysis.

The observable that is affected by this additional term is the parity-violating asymmetry

$$\begin{aligned} A_{PV} &\equiv \frac{d\sigma(\lambda=1) - d\sigma(\lambda=-1)}{d\sigma(\lambda=1) + d\sigma(\lambda=-1)} \\ &= \frac{\left(Y_+ + \frac{R^2 y^2}{2} \right) F_{2,LU} - y^2 F_{L,LU} - Y_- x_B F_{3,LU}}{\left(Y_+ + \frac{R^2 y^2}{2} \right) F_{2,UU} - y^2 F_{L,UU} - Y_- x_B F_{3,UU}}. \end{aligned} \quad (7)$$

The structure functions can be written in terms of PDFs, which stem from the decomposition of the quark correlator. In particular, if we consider the correlator for unpolarized nucleons and if we include strong PV terms, the general expression at leading twist is ¹

$$\Phi^q(x, Q^2) = \left\{ f_1^q(x, Q^2) + g_1^{PVq}(x, Q^2) \gamma_5 \right\} \frac{\not{n}_+}{2}, \quad (8)$$

¹ A similar decomposition for the transverse-momentum-dependent correlator was studied in Ref. [10]. Our function g_1^{PV} corresponds to the integral of the function u_1 in that reference.

where x is the light-cone momentum fraction carried by quarks (neglecting target mass corrections, $x \approx x_B$). The second term is ignored in the SM. It contains the PDF g_1^{PV} , which describes the difference in the probability to find right-handed vs. left-handed quarks inside an unpolarized proton. It is P-odd and CP-even. Its behavior under QCD evolution is the same as the helicity distribution g_1 . The integral of this function is connected to the anapole moment of the proton and nuclei (see, e.g., Refs. [11–14]), which could contribute to PV effects in electron-proton elastic scattering experiments [15, 16].

Neglecting strong P violation and corrections in the strong coupling constant α_S , but including target mass corrections (see, e.g., Ref. [17] for a recent review), the structure function F_3 can be written in terms of the PDF f_1 [9], evaluated at the Nachtmann variable [17]

$$x_N = \frac{2x_B}{1 + \sqrt{1 + R^2}}. \quad (9)$$

For convenience, in the following we avoid explicitly writing the arguments of the PDFs. The detailed expression of the structure functions F_3 for each channel is

$$F_3^{(\gamma)}(x_B, Q^2) = 0, \quad (10)$$

$$F_3^{(\gamma Z)}(x_B, Q^2) = \frac{1}{\sqrt{1 + R^2}} \sum_q 2e_q g_A^q f_1^{(q-\bar{q})}, \quad (11)$$

$$F_3^{(Z)}(x_B, Q^2) = \frac{1}{\sqrt{1 + R^2}} \sum_q 2g_V^q g_A^q f_1^{(q-\bar{q})}, \quad (12)$$

where e_q is the quark charge, $g_{V,A}^q$ are quark's neutral weak couplings to the Z^0 , and R has been defined below Eq. (1). Note that we use the simplified notation $f_1^{(q\pm\bar{q})} = f_1^q \pm f_1^{\bar{q}}$ and similarly for the g_1 's below. Including strong P violation, the standard results are modified by the following additional contributions:

$$\Delta F_3^{(\gamma)}(x_B, Q^2) = -\frac{1}{\sqrt{1 + R^2}} \sum_q e_q^2 g_1^{\text{PV}(q+\bar{q})}, \quad (13)$$

$$\Delta F_3^{(\gamma Z)}(x_B, Q^2) = -\frac{1}{\sqrt{1 + R^2}} \sum_q 2e_q g_V^q g_1^{\text{PV}(q+\bar{q})}, \quad (14)$$

$$\Delta F_3^{(Z)}(x_B, Q^2) = -\frac{1}{\sqrt{1 + R^2}} \sum_q (g_V^{q2} + g_A^{q2}) g_1^{\text{PV}(q+\bar{q})}. \quad (15)$$

The detailed SM expression of the structure function F_2 for each channel is

$$F_2^{(\gamma)}(x_B, Q^2) = \sum_q e_q^2 f_1^{(q+\bar{q})}, \quad (16)$$

$$F_2^{(\gamma Z)}(x_B, Q^2) = \sum_q 2e_q g_V^q f_1^{(q+\bar{q})}, \quad (17)$$

$$F_2^{(Z)}(x_B, Q^2) = \sum_q (g_V^{q2} + g_A^{q2}) f_1^{(q+\bar{q})}. \quad (18)$$

Note that also the structure function F_2 gets modified by the inclusion of strong P-violation contributions:

$$\Delta F_2^{(\gamma)}(x_B, Q^2) = 0, \quad (19)$$

$$\Delta F_2^{(\gamma Z)}(x_B, Q^2) = -\sum_q 2e_q g_A^q x_B g_1^{\text{PV}(q-\bar{q})}, \quad (20)$$

$$\Delta F_2^{(Z)}(x_B, Q^2) = -\sum_q 2g_V^q g_A^q x_B g_1^{\text{PV}(q-\bar{q})}. \quad (21)$$

For completeness, we discuss the case of a polarized nucleon. The correlator becomes ²

$$\Phi^q(x, Q^2) = \left\{ f_1^q(x, Q^2) + g_1^{\text{PV}q}(x, Q^2)\gamma_5 + S_L \left(g_1^q(x, Q^2)\gamma_5 + f_{1L}^{\text{PV}q}(x, Q^2) \right) - \not{s}_T \left(h_1^q(x, Q^2)\gamma_5 - e_{1T}^{\text{PV}q}(x, Q^2) \right) \right\} \frac{\not{p}_+}{2}. \quad (22)$$

The fourth term, f_{1L}^{PV} , is P-odd, CP-odd and should be connected to the electric dipole moment of the proton.

In the case of DIS off longitudinally polarized protons, the g_5 structure function can be introduced [9]. In this case, the inclusion of strong PV terms will generate a difference from the weak parity violation standard results, with the additional term:

$$\Delta g_5(x_B, Q^2) \approx \Delta g_5^{(\gamma)}(x_B, Q^2) = \frac{1}{2} \sum_q e_q^2 f_{1L}^{\text{PV}(q-\bar{q})}. \quad (23)$$

III. COMPATIBILITY OF EXPERIMENTAL DATA WITH STRONG PARITY VIOLATION

There are currently no models that generate strong P-violating PDFs. Modifications of QCD through the inclusion of a θ -term would lead to extremely small effects, since the value of the θ parameter is constrained by measurements of the neutron electric dipole moment. Other higher-dimensional P-violating SMEFT operators could possibly generate nonzero P-violating PDFs. A model with a topologically non-trivial QCD background has been used to generate transverse-momentum-dependent fragmentation functions [18]. The inclusion of electroweak corrections in the evolution of PDFs can also produce PV PDFs. However, given that QED corrections are below 1% [19, 20], PV contributions would be smaller by a factor Q^2/M_Z^2 , thus negligible at low energies.

In order to obtain a first estimate of the size of the newly introduced PV PDFs, we assume they are proportional to their parity-even counterparts, i.e., $g_1^{\text{PV}} = a g_1$, with a being a very small number. This leads to

$$\Delta F_3^{(\gamma)}(x_B, Q^2) = -\frac{a}{\sqrt{1+R^2}} \sum_q e_q^2 g_1^{(q+\bar{q})}. \quad (24)$$

The total contributions to be added to the standard expression of F_3 are

$$\begin{aligned} \Delta F_{3,UU}(x_B, Q^2) &= -\frac{a}{\sqrt{1+R^2}} \left(g_A^e \eta_{\gamma Z} \sum_q 2e_q g_V^q g_1^{(q+\bar{q})} - 2g_V^e g_A^e \eta_Z \sum_q (g_V^{q^2} + g_A^{q^2}) g_1^{(q+\bar{q})} \right), \\ \Delta F_{3,LU}(x_B, Q^2) &= -\frac{a}{\sqrt{1+R^2}} \left(\sum_q e_q^2 g_1^{(q+\bar{q})} - g_V^e \eta_{\gamma Z} \sum_q 2e_q g_V^q g_1^{(q+\bar{q})} + (g_V^{e^2} + g_A^{e^2}) \eta_Z \sum_q (g_V^{q^2} + g_A^{q^2}) g_1^{(q+\bar{q})} \right). \end{aligned} \quad (25)$$

Moreover, the total contributions to be added to F_2 are

$$\begin{aligned} \Delta F_{2,UU}(x_B, Q^2) &= g_V^e \eta_{\gamma Z} a \sum_q 2e_q g_A^q x_B g_1^{(q-\bar{q})} - (g_V^{e^2} + g_A^{e^2}) \eta_Z a \sum_q 2g_V^q g_A^q x_B g_1^{(q-\bar{q})}, \\ \Delta F_{2,LU}(x_B, Q^2) &= -g_A^e \eta_{\gamma Z} a \sum_q 2e_q g_A^q x_B g_1^{(q-\bar{q})} + 2g_V^e g_A^e \eta_Z a \sum_q 2g_V^q g_A^q x_B g_1^{(q-\bar{q})}. \end{aligned} \quad (26)$$

To estimate the possible size of the new PDFs, we proceed in the following way: we *assume* the validity of the electroweak sector of the Standard Model and we attribute any discrepancy between low-energy measurements and predictions *entirely* to the newly introduced PDFs. In other words, we estimate how large the PV PDFs should be to be compatible with low-energy measurements.

We fit the theoretical predictions at NLO for the electron and positron PV asymmetries of Eq. (7) to the DIS experimental data from HERA [21] with proton beams (274 experimental points), and from JLab 6 GeV PVDIS [22, 23] (2 points) and the SLAC E122 experiment [24] (11 points) with a deuterium fixed target ³. In Tab.I, we list details of

² Our functions f_{1L}^{PV} and e_{1T}^{PV} correspond to the integral of the functions v_{1L} and w_{1T} in Ref. [10], respectively.

³ In first approximation, the deuterium target is described as a incoherent sum of free nucleons (one proton and one neutron)

Experiment	N_{dat}	Observable	Hadron	\sqrt{s} [GeV]	Q^2 [GeV ²]	y	Ref.
HERA	136	A_{PV} for e^+	proton	319	120 - 30000	0.033 - 0.9	[21]
HERA	138	A_{PV} for e^-	proton	319	120 - 30000	0.033 - 0.9	[21]
JLab PVDIS	2	A_{PV} for e^-	deuterium	4.77	1.085; 1.901	0.20; 0.28	[22]
SLAC E122	11	A_{PV} for e^-	deuterium	5.5 - 6.5	0.92 - 1.96	0.15 - 0.36	[24]
Total	287						

TABLE I: Breakdown of the data sets considered in this analysis. For each data set, the table includes information on: the number of data points (N_{dat}), the measured observable, the initial-state hadron, the center-of-mass energy \sqrt{s} , the covered range(s) in Q^2 , the inelasticity y , and the published reference. The total number of data points amounts to 287.

the data sets included in our analysis. In all, we analyze 287 experimental data points, 136 for positron asymmetry and 151 for electron asymmetry. In a future study, it could be interesting to estimate the effect of the new PV PDF g_1^{PV} also in Drell-Yan processes.

The value of the energy scale Q^2 is very small for E122 and JLab PVDIS data sets ($Q^2 \simeq 1 - 2$ GeV²). For them, the inclusion of target mass corrections (see Eq. (1) and Eqs. (11)-(18)) has a significant effect, while it does not modify the results for HERA data. Moreover, to be consistent with Refs. [22, 23] we introduce electroweak radiative corrections according to Ref. [25]. These corrections could be included also at higher Q^2 (using, e.g., the Djangoh event generator [26]) but they are small compared to the experimental uncertainties.

In our analysis, we choose the NNPDF4.0 [27] and the NNPDFpol1.1 [28] sets of unpolarized and polarized PDFs, respectively. We include the full set of PDF replicas to account for their uncertainty, which is interpreted as a source of systematic theoretical error to be added to the experimental systematic error. We checked that this theoretical error is much smaller than the experimental errors (of order 1% or less). We use PDF sets at NLO accuracy and compute the structure functions F_2 , F_L and F_3 at $\mathcal{O}(\alpha_s)$.

The error analysis is performed with the so-called bootstrap method, which consists in fitting an ensemble of Monte Carlo (MC) replicas of the experimental data. We generate 100 replicas of the experimental data and we relate each one of them to a single replica of unpolarized and polarized PDFs. In this way, we obtain a distribution of 100 values for the fit parameter a .

The resulting quality of the fit is shown in Table II, where the χ^2 per number of data points N_{dat} are provided for each of the considered experimental data sets, including one standard deviation from the full ensemble of replicas. We indicate also the values of χ^2/N_{dat} obtained with the SM predictions.

		SM predictions	Our analysis
Data set	N_{dat}	χ^2/N_{dat}	χ^2/N_{dat}
HERA e^+ (p)	136	1.12 ± 0.01	1.12 ± 0.01
HERA e^- (p)	138	0.98 ± 0.01	0.98 ± 0.01
JLab PVDIS e^- (d)	2	0.67 ± 0.12	0.42 ± 0.40
SLAC E122 e^- (d)	11	0.97 ± 0.01	0.94 ± 0.02
Total	287	1.042 ± 0.001	1.037 ± 0.004

TABLE II: Breakdown of the values of χ^2 per number of data points N_{dat} for all data sets considered in our analysis (hadron targets in brackets). The values of χ^2 and uncertainties refer to the mean value and one standard deviation from the ensemble of replicas of the experimental data.

The mean value of the global χ^2/N_{dat} is slightly smaller than the SM result, the description is thus improved by including our model for the PV PDF. However, it must be noted that the values of χ^2 are all close to or smaller than 1 because of large experimental errors affecting many data points. New data with higher precision would help to better assess the impact of PV contributions.

The resulting value for the fit parameter a is

$$a = \hat{a} \pm \Delta a = (-1.01 \pm 0.66) \times 10^{-4}, \quad (27)$$

where \hat{a} and Δa are the mean value and one standard deviation from the ensemble of 100 values, respectively. Our results indicate that the null hypothesis can be rejected with a p -value = 0.063: the probability of making an error by rejecting the hypothesis of no parity-violating contributions is 6.3%. It is not negligible but small. In other words,

current experimental data indicate that strong P violation is statistically possible and, actually, is slightly favoured with respect to the SM description. The negative value of a indicates also that in the proton there could be more left-handed quarks than right-handed ones. We checked that this result does not depend on the choice of the PDF by repeating the fit with different sets and obtaining no significant differences.




FIG. 1: Comparison between theoretical predictions and HERA data for e^+p (open red points) and e^-p (solid black points) as a function of Q^2 at $x = 0.13$. Upper panel: helicity asymmetry of Eq. (7); central panel: relative difference with respect to SM predictions for e^+ asymmetry; lower panel: same for e^- asymmetry. The (barely visible) colored uncertainty bands correspond to the 68% C.L.

In Fig. 1, the upper panel shows the comparison between theoretical predictions for A_{PV} of Eq. (7) (colored bands) and HERA data for inclusive DIS between a proton beam and electron e^- (solid black points) or positron e^+ (open red points) beams, as a function of Q^2 at the given $x = 0.13$. The central (lower) panel shows the relative difference of data and results of our fit with respect to SM predictions for the e^+ (e^-) asymmetry. The (barely visible) colored uncertainty bands correspond to the 68% confidence level (C.L.), obtained by excluding the largest and smallest 16% of the MC replicas. The narrow width of the bands reflects the small theoretical uncertainty of the PDFs used in this work.

As can be seen from Fig. 1 and Tab. II, the HERA data are nicely described in our framework although the χ^2 for the e^+ asymmetry is slightly worse than the e^- case because of the behaviour of more precise data points at the lowest Q^2 bins (see central panel). This may leave room for improvements of our model in a future work. In any case, both values of χ^2 for e^+ and e^- asymmetries are the same as in the SM framework (see Tab. II), indicating that PV contributions do not impact the description of HERA data: in fact, the fit is driven by the other experimental data sets with much smaller errors, described below.

In Fig. 2, the upper panel shows the comparison between theoretical predictions for electron A_{PV} of Eq. (7) (colored bands) and E122 data (open points) and JLab PVDIS data (solid points) as a function of Q^2 . The central and the lower panel show the relative difference of data and results of our fit with respect to SM predictions for the E122 and JLab PVDIS asymmetries, respectively. Similar to Fig. 1, the colored uncertainty bands correspond to the 68% C.L. Both experimental data sets are nicely described in our framework. This is confirmed in Tab. II by the systematic improvement of the quality of the fit with respect to the SM framework. The result is particularly relevant for the JLab PVDIS data, due to their very small uncertainty.




FIG. 2: Comparison between theoretical predictions and data from JLab 6 GeV PVDIS (solid points) and SLAC E122 (open points) experiments as a function of Q^2 . Upper panel: electron e^- asymmetry A_{PV} of Eq. (7). Central panel: relative difference with SM predictions for the E122 asymmetry. Lower panel: same for the PVDIS asymmetry. Uncertainty bands correspond to the 68% C.L.

IV. IMPACT STUDY OF FUTURE EXPERIMENTS

As already remarked when commenting our results in Tab. II, further insights into the existence of PV contribution in the structure of the proton (or light nuclei) could be provided by new high-precision experimental data, such as the ones expected from the upcoming SoLID program [5] at JLab 12 GeV (and possibly future upgrades of JLab [29]). SoLID is a versatile spectrometer designed specifically to achieve high precision measurements on a variety of topics. The nominal running conditions for SoLID's PVDIS measurement [30] include a 50 μ A, 11 GeV longitudinally polarized electron beam of 85% polarization, incident on a 40-cm liquid deuterium or liquid hydrogen target. The pseudodata used in our study are based on PVDIS measurement with either 120 days of beam (at 100% efficiency) on the deuterium target, or 90 days on the hydrogen target. The relative systematic uncertainty $\delta A_{PV}(\text{syst})/A_{PV}^{\text{SM}}$ consists of an uncorrelated 0.28% contribution and a 0.45% contribution correlated across all kinematic bins.

In order to estimate the reduction of the uncertainties on the g_1^{PV} PDF, we generate pseudodata by calculating the theoretical predictions for the A_{PV} asymmetry and include the experimental uncertainties based on the running conditions given above. We consider the SoLID pseudodata on either deuteron or proton target and we impose a (conservative) cut $x_B < 0.5$, to avoid the kinematic region where nuclear corrections for deuteron target and higher-twist contributions become relevant. The impact of the SoLID pseudodata on the fitted parameter a is reported in Tab. III. We obtain at least a factor 3 reduction of the uncertainty with a slightly larger impact when using the proton target. If the central value of the parameter a were to be confirmed by future measurements, the result in Tab. III would represent a deviation of at least 5σ from the SM result of $a = 0$. Figure 3 shows the impact of SoLID pseudodata on g_1^{PV} of the up quark normalized to its central value. We note that the uncertainty bands decrease in the region $0.2 < x_B < 0.5$, which is consistent with the region covered by the SoLID pseudodata. The reduction outside this range may be due to the low flexibility of our assumption. A more refined model will be needed when SoLID data become available.

These results indicate that SoLID PVDIS data will potentially put a drastic limit on the strong PV effects. Furthermore, they show that the SoLID PVDIS proton measurements will be valuable not only at high x_B in providing

Fit	a (10^{-4})	δa (10^{-4})
Baseline	1.01	0.66
Baseline + SoLID (d)	1.01	0.21
Baseline + SoLID (p)	1.01	0.15

TABLE III: Values of parameter a and its uncertainty δa related to baseline fit and the impact studies of SoLID pseudodata on deuteron (d) or proton (p) targets.




FIG. 3: Estimate of the impact of the SoLID pseudodata on the error bands of the u quark g_1^{PV} in the proton in x space at $Q = 2$ GeV, based on the baseline analysis presented in this paper. Purple bands: current 68% C.L. uncertainties on g_1^{PV} from the baseline fit. Yellow bands: 68% C.L. uncertainties after the inclusion of SoLID pseudodata on deuteron target. Red bands: 68% C.L. uncertainties after the inclusion of SoLID pseudodata on proton target.

a model-independent measurement of d/u PDF ratio [5], but also at medium x_B by offering a new way to explore physics beyond SM.

We also note that more measurements on the PVDIS asymmetries are expected from the future EIC at higher energies [6, 31]. We considered the EIC energy configurations 10×275 GeV and 10×137 GeV for electron-proton and electron-deuteron collisions, respectively, each corresponds to the maximum annual integrated luminosity of 100 fb^{-1} [7, 31]. However, we found that adding these EIC pseudodata brings little impact to the limit on the strong PV effects, due to the large projected uncertainties.

V. CONCLUSIONS

In this paper, we explored the impact of violating QCD parity invariance (strong P violation) on nucleon structure, focusing on inclusive DIS with longitudinally polarized leptons and unpolarized targets. This leads to the introduction of a new P-odd and CP-even PDF, denoted as g_1^{PV} , that describes the difference in the probability to find right-handed vs. left-handed quarks inside an unpolarized proton. This function generates a new contribution to the structure function F_3 from pure photon exchange.

To estimate the size of g_1^{PV} , we perform a fit to relevant experimental data from HERA, SLAC, and JLab. As a preliminary model, we assume that the PV PDFs are proportional to their parity-even counterparts and we fit the proportionality constant, a . Our analysis shows that including strong PV contributions improves the description of the data. We obtain the value $a = (-1.01 \pm 0.66) \times 10^{-4}$, which indicates that there could be more left-handed quarks than right-handed ones in an unpolarized proton. Furthermore, we carried out impact studies for the future EIC and the SoLID PVDIS measurement of JLab's 12 GeV program. While in the former case we don't find any significant impact, in the latter case we find that anticipated SoLID deuteron (proton) measurement will potentially provide a factor 3 (4) reduction in the uncertainty of a . Should the central value of the parameter a be confirmed by SoLID

measurements, it would represent at least a 5σ deviation from the SM result of no strong PV effects. We emphasize that detecting strong P violation could have implications beyond nucleon structure, potentially shedding light on the matter-antimatter asymmetry in the universe.

Acknowledgments

We gratefully acknowledge discussion with H. Spiesberger and P. Souder. The work of X. Zheng is supported by the U.S. Department of Energy, Office of Science, Office of Nuclear Physics under contract number DE-SC0014434.

Appendix A: Derivation of the cross section

For completeness, in this Appendix we provide the conventions used and the analytical derivation of the cross section in Eq. (1).

For a fast-moving lepton with initial (final) 4-momentum k ($k' = k - q$) and helicity λ , the leptonic tensor for the three channels (γ exchange, $\gamma - Z$ interference and Z exchange) can be written as

$$\begin{aligned} L_{\mu\nu}^{(j)} &= C^{(j)} L_{\mu\nu}^{(\gamma)}, \\ L_{\mu\nu}^{(\gamma)} &= 2 [k_\mu k'_\nu + k'_\mu k_\nu - (k \cdot k') g_{\mu\nu} - i \lambda \varepsilon_{\mu\nu\alpha\beta} k^\alpha k'^\beta], \end{aligned} \quad (\text{A1})$$

where the index j runs over the channels $i = \gamma, \gamma Z, Z$, and $C^{(\gamma)} = 1$, $C^{(\gamma Z)} = -(g_V^e - \lambda g_A^e)$, $C^{(Z)} = (g_V^e - \lambda g_A^e)^2$ [8]. For an anti-lepton, the same formula holds but with the sign of g_A^e flipped.

For an unpolarized hadron, the hadronic tensor is given by [9]

$$W^{\mu\nu} = \left(-g^{\mu\nu} + \frac{q^\mu q^\nu}{q^2} \right) F_1 + \frac{\tilde{P}^\mu \tilde{P}^\nu}{P \cdot q} F_2 + i \frac{\varepsilon^{\mu\nu\rho\sigma}}{2(P \cdot q)} P_\rho q_\sigma F_3, \quad (\text{A2})$$

where $\tilde{P}^\mu = P^\mu - q^\mu (P \cdot q)/q^2$.

The cross section becomes

$$\frac{d^2\sigma}{dx_B dy} = \frac{2\pi y \alpha^2}{Q^4} \sum_{j=\gamma, \gamma Z, Z} \eta^{(j)} L_{\mu\nu}^{(j)} W^{\mu\nu} = \frac{2\pi y \alpha^2}{Q^4} \sum_{j=\gamma, \gamma Z, Z} \eta^{(j)} C^{(j)} L_{\mu\nu}^{(\gamma)} W^{\mu\nu}, \quad (\text{A3})$$

where

$$\eta^\gamma = 1; \quad \eta^{\gamma Z} = \left(\frac{G_F M_Z^2}{2\sqrt{2}\pi\alpha} \right) \left(\frac{Q^2}{Q^2 + M_Z^2} \right); \quad \eta^Z = (\eta^{\gamma Z})^2, \quad (\text{A4})$$

with α the fine structure constant, M_Z the mass of the Z^0 boson, and G_F the Fermi coupling constant.

By neglecting the lepton mass, the contraction of the leptonic tensor $L_{\mu\nu}^{(\gamma)}$ of Eq. (A1) with the hadronic tensor $W^{\mu\nu}$ of Eq. (A2) gives

$$L_{\mu\nu}^{(\gamma)} W^{\mu\nu} = 2 \left[Q^2 F_1^{(\gamma)} + \frac{Q^2}{x_B y^2} \left(1 - y - \frac{1}{4} R^2 y^2 \right) F_2^{(\gamma)} - \lambda \frac{Q^2}{2} \frac{2-y}{y} F_3^{(\gamma)} \right], \quad (\text{A5})$$

where $R = 2M x_B/Q$. By including such target mass corrections, we make the replacement $2x_B F_1 = (1 + R^2)F_2 - F_L$ in the above contraction and we get

$$L_{\mu\nu}^{(\gamma)} W^{\mu\nu} = \frac{Q^2}{x_B y^2} \left[\left(Y_+ + \frac{1}{2} R^2 y^2 \right) F_2^{(\gamma)} - y^2 F_L^{(\gamma)} - \lambda Y_- x_B F_3^{(\gamma)} \right], \quad (\text{A6})$$

with $Y_\pm = 1 \pm (1 - y)^2$.

If we insert the above result into Eq. (A3) and we use the definitions of Eqs. (2)-(5), we finally get

$$\begin{aligned} \frac{d^2\sigma}{dx_B dy} &= \frac{2\pi\alpha^2}{x_B y Q^2} \left[\left(Y_+ + \frac{R^2 y^2}{2} \right) (F_{2,UU} + \lambda F_{2,LU}) - y^2 (F_{L,UU} + \lambda F_{L,LU}) \right. \\ &\quad \left. - Y_- (x_B F_{3,UU} + \lambda x_B F_{3,LU}) \right]. \end{aligned} \quad (\text{A7})$$

-
- [1] I. Brivio and M. Trott, Phys. Rept. **793**, 1 (2019).
 - [2] R. Boughezal, F. Petriello, and D. Wiegand, Phys. Rev. D **104**, 016005 (2021).
 - [3] A. S. Kronfeld et al. (USQCD) (2022), arXiv:2207.07641 [hep-lat].
 - [4] N. Blinov, N. Craig, M. J. Dolan, J. de Vries, P. Draper, I. G. Garcia, B. Lillard, and J. Shelton, in *Snowmass 2021* (2022), arXiv:2203.07218 [hep-ph].
 - [5] J. Arrington et al. (Jefferson Lab SoLID), J. Phys. G **50**, 110501 (2023).
 - [6] R. Abdul Khalek et al., Nucl. Phys. A **1026**, 122447 (2022).
 - [7] R. Boughezal, A. Emmert, T. Kutz, S. Mantry, M. Nycz, F. Petriello, K. Şimşek, D. Wiegand, and X. Zheng, Phys. Rev. D **106**, 016006 (2022).
 - [8] P. Anderson, D. Higinbotham, S. Mantry, and X. Zheng, in *30th International Workshop on Deep-Inelastic Scattering and Related Subjects* (2023), arXiv:2306.00097 [hep-ph].
 - [9] M. Tanabashi et al. (Particle Data Group), Phys. Rev. D **98**, 030001 (2018).
 - [10] W. Yang, Int. J. Mod. Phys. A **34**, 1950145 (2019).
 - [11] V. V. Flambaum, I. B. Khriplovich, and O. P. Sushkov, Phys. Lett. B **146**, 367 (1984).
 - [12] C. S. Wood, S. C. Bennett, D. Cho, B. P. Masterson, J. L. Roberts, C. E. Tanner, and C. E. Wieman, Science **275**, 1759 (1997).
 - [13] S.-L. Zhu, S. J. Puglia, B. R. Holstein, and M. J. Ramsey-Musolf, Phys. Rev. D **62**, 033008 (2000).
 - [14] R. D. Young, J. Roche, R. D. Carlini, and A. W. Thomas, Phys. Rev. Lett. **97**, 102002 (2006).
 - [15] R. Hasty et al. (SAMPLE), Science **290**, 2117 (2000).
 - [16] D. Androić et al. (Qweak), Nature **557**, 207 (2018).
 - [17] R. Ruiz et al. (2023), arXiv:2301.07715 [hep-ph].
 - [18] Z.-B. Kang and D. E. Kharzeev, Phys. Rev. Lett. **106**, 042001 (2011).
 - [19] R. D. Ball, V. Bertone, S. Carrazza, L. Del Debbio, S. Forte, A. Guffanti, N. P. Hartland, and J. Rojo (NNPDF), Nucl. Phys. B **877**, 290 (2013).
 - [20] A. Manohar, P. Nason, G. P. Salam, and G. Zanderighi, Phys. Rev. Lett. **117**, 242002 (2016).
 - [21] V. Andreev et al. (H1), Eur. Phys. J. C **78**, 777 (2018).
 - [22] D. Wang et al. (PVDIS), Nature **506**, 67 (2014).
 - [23] D. Wang et al., Phys. Rev. C **91**, 045506 (2015).
 - [24] C. Y. Prescott et al., Phys. Lett. B **84**, 524 (1979).
 - [25] J. Erler and S. Su, Prog. Part. Nucl. Phys. **71**, 119 (2013).
 - [26] E. C. Aschenauer, T. Burton, T. Martini, H. Spiesberger, and M. Stratmann, Phys. Rev. D **88**, 114025 (2013).
 - [27] R. D. Ball et al. (NNPDF), Eur. Phys. J. C **82**, 428 (2022).
 - [28] E. R. Nocera, R. D. Ball, S. Forte, G. Ridolfi, and J. Rojo (NNPDF), Nucl. Phys. B **887**, 276 (2014).
 - [29] A. Accardi et al. (2023), arXiv:2306.09360 [nucl-ex].
 - [30] P. A. Souder (contact), P. E. Reimer, X. Zheng, et al., *Precision Measurement of Parity-violation in Deep Inelastic Scattering Over a Broad Kinematic Range*, Jefferson Lab Experiment E12-10-007 (2010 with 2022 update).
 - [31] R. Abdul Khalek et al. (2022), arXiv:2203.13199 [hep-ph].