

From characteristic functions to multivariate distribution functions and European option prices by the damped COS method

Gero Junike*, Hauke Stier†

April 4, 2024

Abstract

We provide a unified framework for the computation of the distribution function and the computation of prices of financial options from the characteristic function of some density by the COS method. The classical COS method is numerically very efficient in one-dimension but cannot deal very well with certain financial options in general dimensions. Therefore, we introduce the damped COS method which can handle a large class of integrands very efficiently. We prove the convergence of the (damped) COS method and study its order of convergence. The (damped) COS method converges exponentially if the characteristic function decays exponentially. To apply the (damped) COS method, one has to specify two parameters: a truncation range for the multivariate density and the number of terms to approximate the truncated density by a cosine series. We provide an explicit formula for the truncation range and an implicit formula for the number of terms. Numerical experiments up to five dimensions confirm the theoretical results.

Keywords: Fourier-transform, numerical integration, inversion theorem, COS method, CDF, rainbow options

Mathematics Subject Classification 65D30 · 65T40 · 60E10 62P05

1 Introduction

We aim to solve the following integral numerically:

$$\int_{\mathbb{R}^d} w(\mathbf{x})g(\mathbf{x})d\mathbf{x}. \quad (1)$$

The function g is usually a density and the function w is called *function of interest*. Integrals as in (1) appear in a wide range of applications: The integral is equal to the cumulative distribution function (CDF) of the density g if w is an indicator function. CDFs appear in many scientific disciplines. In a financial context, the function w might also describe some *rainbow option*, which depends on several assets, and w is also called *payoff function*. The function g is then the density of the logarithmic returns of the assets and the integral describes the price of the rainbow option.

In many cases, the precise structure of g is unknown but the Fourier transform \hat{g} is often given in closed-form. For example: while the joint density of the sum of two independent random variables can only be expressed as a convolution and is usually not given explicitly, the joint characteristic function is much simpler to obtain, it is just the product of the marginal characteristic functions. Moreover, the characteristic function of a Lévy process at a particular time-point is usually given explicitly thanks to the Lévy-Khinchin formula.

The integral in (1) can be solved numerically using different techniques such as (quasi) Monte Carlo simulation, numerical quadrature and Fourier inversion, see Ruijter and Oosterlee (2012); Eberlein et al. (2010) and references therein. Special Fourier-inversion methods exist in case of a CDF, e.g., the Gil-Pelaez formula, see Gil-Pelaez (1951) and extensions, e.g., Schorr (1975); Abate and Whitt (1992); Waller et al. (1995); Hughett (1998) in one-dimension and Shephard (1991a,b) in d dimensions.

*Corresponding author. Carl von Ossietzky Universität, Institut für Mathematik, 26129 Oldenburg, Germany, ORCID: 0000-0001-8686-2661, E-mail: gero.junike@uol.de

†Carl von Ossietzky Universität, Institut für Mathematik, 26129 Oldenburg, Germany.

The COS method, see Fang and Oosterlee (2009a) for $d = 1$ and Ruijter and Oosterlee (2012) for $d > 1$, is a Fourier inversion technique. The COS method has been applied extensively in computational finance and economics, see Fang and Oosterlee (2009b, 2011); Grzelak and Oosterlee (2011); Ruijter and Oosterlee (2012); Zhang and Oosterlee (2013); Leitao et al. (2018); Liu et al. (2019a,b); Oosterlee and Grzelak (2019); Bardgett et al. (2019). Other one-dimensional Fourier pricing techniques can be found in Carr and Madan (1999); Lord et al. (2008); Ortiz-Gracia and Oosterlee (2013, 2016). The COS method compares favorably to other Fourier inversion techniques, see Fang and Oosterlee (2009a).

The main idea of the COS method is to truncate the integration range in (1) to some finite hypercube and to approximate the density g on the finite truncation range by a classical Fourier-cosine expansion. There is a clever trick to approximate the cosine coefficients for g in a very fast and robust way using \hat{g} . The COS method is particularly fast when the Fourier-cosine coefficients of the function of interest w are given analytically, too. For instance, in multivariate dimensions, the Fourier-cosine coefficients of a CDF can be obtained analytically. However, for many rainbow options the Fourier-cosine coefficients are not given in closed-form (e.g. arithmetic basket options). Ruijter and Oosterlee (2012) propose in this case to obtain the Fourier coefficients of the function of interest numerically by a discrete cosine transform but this significantly slows the COS method down. In this article, we introduce the *damped COS method*, which is able to avoid the expensive application of the discrete cosine transform if the Fourier transform of the function of interest is given in closed-form. The Fourier transform \hat{w} is known for many rainbow options, see Hurd and Zhou (2010); Eberlein et al. (2010).

In moderate dimensions, the COS method is a fast, robust and straightforward to implement alternative to the d -dimensional Gil-Pelaez formula, see Shephard (1991a,b) or the multivariate Lewis formula, see Eberlein et al. (2010), in particular if g is smooth and has semi-heavy tails.

This article makes the following main contributions: We prove the convergence of the multidimensional (damped) COS method, we analyze the order of convergence of the (damped) COS method and provide explicit and implicit formulas for the truncation range and the number of terms, respectively. Unlike Ruijter and Oosterlee (2012), who analyze the classical COS method, we include in our analysis numerical uncertainty on the characteristic function \hat{g} and on the Fourier-cosine coefficients of the function of interest. This helps to understand how approximations on \hat{g} and the Fourier-cosine coefficients of the function of interest affect the total error of the COS method.

This article is structured as follows: In Section 2, we fix some notation. In Section 3 we introduce the multidimensional (damped) COS method, prove its convergence, analyze the order of convergence and provide explicit and implicit formulas for the truncation range and the number of terms. In Section 4 we discuss some examples for g and \hat{g} . In Section 5 we discuss some functions of interest, i.e., examples for w . In Section 6 we provide numerical experiments and compare the COS method to a Monte Carlo simulation. Section 7 concludes.

2 Notation

Let $d \in \mathbb{N}$. Let \mathcal{L}^1 and \mathcal{L}^2 denote the sets of integrable and square integrable functions from \mathbb{R}^d to \mathbb{R} , and by $\langle \cdot, \cdot \rangle$ and $\|\cdot\|_2$ we denote the scalar product and the (semi)norm on \mathcal{L}^2 , respectively. The supremum norm of a function $g : \mathbb{R}^d \rightarrow \mathbb{C}$ is defined by $\|g\|_\infty := \sup_{\mathbf{x} \in \mathbb{R}^d} |g(\mathbf{x})|$. By $\Re\{z\}$ and $\Im\{z\}$ we denote the real part and imaginary part of a complex number $z \in \mathbb{C}$. The complex unit is denoted by i . By Γ , we denote the Gamma function. The Euclidean norm and the maximum norm on \mathbb{R}^d is denoted by $|\cdot|$ and by $|\cdot|_\infty$ respectively. For $\mathbf{x}, \mathbf{y} \in \mathbb{R}^d$ we define

$$\mathbf{x} \geq \mathbf{y} :\Leftrightarrow x_1 \geq y_1, \dots, x_d \geq y_d$$

and treat “ \leq ”, “ $<$ ”, “ $>$ ”, “ $=$ ” and “ \neq ” similarly. We set $\mathbb{R}_+^d := \{\mathbf{x} \in \mathbb{R}^d, \mathbf{x} > \mathbf{0}\}$. For $\mathbf{a}, \mathbf{b} \in \mathbb{R}^d$ with $\mathbf{a} \leq \mathbf{b}$, two complex vectors $\mathbf{z}, \mathbf{y} \in \mathbb{C}^d$ and $\lambda \in \mathbb{C}$ we define $\mathbf{z} + \mathbf{y} := (z_1 + y_1, \dots, z_d + y_d) \in \mathbb{C}^d$ and treat $\mathbf{z}\mathbf{y}$ and $\frac{\mathbf{z}}{\mathbf{y}}$

similarly. We further define

$$\begin{aligned}
\mathbf{z} \cdot \mathbf{y} &:= z_1 y_1 + \dots + z_d y_d \in \mathbb{C} \\
\lambda \mathbf{z} &:= (\lambda z_1, \dots, \lambda z_d) \in \mathbb{C}^d \\
[\mathbf{a}, \mathbf{b}] &:= [a_1, b_1] \times \dots \times [a_d, b_d] \subset \mathbb{R}^d \\
(-\infty, \mathbf{b}] &:= (-\infty, b_1] \times \dots \times (-\infty, b_d] \subset \mathbb{R}^d \\
\exp(\mathbf{x}) &:= (\exp(x_1), \dots, \exp(x_d)), \quad \mathbf{x} \in \mathbb{R}^d \\
\log(\mathbf{x}) &:= (\log(x_1), \dots, \log(x_d)), \quad \mathbf{x} \in \mathbb{R}_+^d.
\end{aligned}$$

For a subset $A \subset \mathbb{R}^d$, we define the indicator function $1_A(\mathbf{x})$ by one if $\mathbf{x} \in A$ and zero otherwise. Let $\mathbb{N}_0 = \mathbb{N} \cup \{0\}$. For $\mathbf{N} = (N_1, \dots, N_d) \in \mathbb{N}_0^d$ and a sequence $(a_{\mathbf{k}})_{\mathbf{k} \in \mathbb{N}_0^d} \subset \mathbb{C}$, we define

$$\sum'_{\mathbf{0} \leq \mathbf{k} \leq \mathbf{N}} a_{\mathbf{k}} := \sum_{\mathbf{0} \leq \mathbf{k} \leq \mathbf{N}} \frac{1}{2^{\Lambda(\mathbf{k})}} a_{\mathbf{k}},$$

where $\Lambda(\mathbf{k})$ is the number of components of the vector \mathbf{k} which are equal to zero, i.e., $\Lambda(\mathbf{k}) := \sum_{h=1}^d 1_{\{0\}}(k_h)$. For an integrable function $g : \mathbb{R}^d \rightarrow \mathbb{C}$ define its *Fourier transform* by

$$\widehat{g}(\mathbf{u}) := \int_{\mathbb{R}^d} g(\mathbf{x}) e^{i\mathbf{u} \cdot \mathbf{x}} d\mathbf{x}, \quad \mathbf{u} \in \mathbb{R}^d. \quad (2)$$

This definition of the Fourier transform also appear in Bauer (1996, Def. 22.6) and Eberlein et al. (2010). Provided the integral in (2) exists, the domain of \widehat{g} may also be extended to parts of the complex plane. If $g \geq 0$ and $\int g(\mathbf{x}) d\mathbf{x} = 1$ then g is called *density*, \widehat{g} is called *characteristic function* and the map $\mathbf{y} \mapsto \int_{(-\infty, \mathbf{y}]} g(\mathbf{x}) d\mathbf{x}$ is called *cumulative distribution function* (CDF).

3 Damped COS method

Typically, the function of interest, w , is only locally integrable but $w \notin \mathcal{L}^1$. We provide two examples: The integral in (1) is equal to the CDF of g evaluated at $\mathbf{y} \in \mathbb{R}^d$ if $w(\mathbf{x}) = 1_{(-\infty, \mathbf{y}]}(\mathbf{x})$ for $\mathbf{x} \in \mathbb{R}^d$. A rainbow option like an arithmetic basket put option with strike $K > 0$ is defined by $w(\mathbf{x}) = \max(K - \sum_{h=1}^d e^{x_h}, 0)$, $\mathbf{x} \in \mathbb{R}^d$. To introduce the *damped COS method*, we will consider a damped function of interest which is assumed to be integrable. The idea introducing a damping factor and to consider modified (integrable) functions of interest dates back at least to Carr and Madan (1999). Note that for many models and many rainbow options, both \widehat{g} and \widehat{w} are given in closed-form, see, e.g., Ruijter and Oosterlee (2012); Eberlein et al. (2010) and references therein.

For a scaling factor $\lambda > 0$, shift parameter $\boldsymbol{\mu} \in \mathbb{R}^d$ and a damping factor $\boldsymbol{\alpha} \in \mathbb{R}^d$, define the *damped density* by

$$f(\mathbf{x}) = \lambda e^{\boldsymbol{\alpha} \cdot (\mathbf{x} + \boldsymbol{\mu})} g(\mathbf{x} + \boldsymbol{\mu}), \quad \mathbf{x} \in \mathbb{R}^d \quad (3)$$

and the *damped function of interest* by

$$v(\mathbf{x}) = \frac{1}{\lambda} e^{-\boldsymbol{\alpha} \cdot (\mathbf{x} + \boldsymbol{\mu})} w(\mathbf{x} + \boldsymbol{\mu}), \quad \mathbf{x} \in \mathbb{R}^d. \quad (4)$$

By definition, it follows that

$$\int_{\mathbb{R}^d} w(\mathbf{x}) g(\mathbf{x}) d\mathbf{x} = \int_{\mathbb{R}^d} v(\mathbf{x}) f(\mathbf{x}) d\mathbf{x}. \quad (5)$$

Thanks to Proposition 3.1, f is a density centered around zero if we choose λ and $\boldsymbol{\mu}$ carefully and \widehat{f} is given in closed-form if \widehat{g} is given in closed-form.

Proposition 3.1. Let $g \in \mathcal{L}_1$ and $\alpha \in \mathbb{R}_+^d$. Assume that g is a density and that $\mathbf{x} \mapsto |\mathbf{x}|e^{\alpha \cdot \mathbf{x}}g(\mathbf{x})$ is integrable. Let $\lambda = (\widehat{g}(-i\alpha))^{-1}$ then $\lambda \in (0, \infty)$. Choose $\mu \in \mathbb{R}^d$ by

$$\mu_h = -\lambda i \frac{\partial}{\partial u_h} \widehat{g}(\mathbf{u} - i\alpha) \Big|_{\mathbf{u}=\mathbf{0}}, \quad h = 1, \dots, d. \quad (6)$$

Define $f(\mathbf{x}) = \lambda e^{\alpha \cdot (\mathbf{x} + \mu)} g(\mathbf{x} + \mu)$, $\mathbf{x} \in \mathbb{R}^d$. Then f is a density with characteristic function

$$\widehat{f}(\mathbf{u}) = \lambda e^{-i\mathbf{u} \cdot \mu} \widehat{g}(\mathbf{u} - i\alpha), \quad \mathbf{u} \in \mathbb{R}^d. \quad (7)$$

Further, the moments of f of first order are zero, i.e., $\int_{\mathbb{R}^d} f(\mathbf{x}) x_h d\mathbf{x} = 0$, $h = 1, \dots, d$.

Proof. Use $\int |\mathbf{x}|e^{\alpha \cdot \mathbf{x}}g(\mathbf{x})d\mathbf{x} < \infty$ and split the integration range into $\mathbb{R}^d \setminus B_1$ and B_1 , where B_1 is the unit ball, to see that $\mathbf{x} \mapsto e^{\alpha \cdot \mathbf{x}}g(\mathbf{x})$ is integrable. Since $\lambda = (\int_{\mathbb{R}^d} e^{\alpha \cdot \mathbf{x}}g(\mathbf{x})d\mathbf{x})^{-1}$ and g is a density we have $\lambda \in (0, \infty)$. By the definition of λ , f is a density. Since $f \in \mathcal{L}^1$, \widehat{f} exists. A direct analysis shows (7). By Bauer (1996, Thm 25.2), the partial derivatives in Equation (6) exist and it holds that $\mu_h = \lambda \int_{\mathbb{R}^d} e^{\alpha \cdot \mathbf{x}}g(\mathbf{x})x_h d\mathbf{x}$. Finally, we have that

$$\int_{\mathbb{R}^d} f(\mathbf{x}) x_h d\mathbf{x} = \int_{\mathbb{R}^d} \lambda e^{\alpha \cdot (\mathbf{x} + \mu)} g(\mathbf{x} + \mu) x_h d\mathbf{x} \lambda \int_{\mathbb{R}^d} e^{\alpha \cdot \mathbf{x}}g(\mathbf{x})x_h d\mathbf{x} - \mu_h \lambda \int_{\mathbb{R}^d} e^{\alpha \cdot \mathbf{x}}g(\mathbf{x})d\mathbf{x} = 0.$$

□

For some models \widehat{g} need to be approximated numerically, e.g., in Duffie et al. (2003), \widehat{g} is the solution to some ordinary differential equation, which itself need to be solved numerically before applying the COS method. From now on, we assume that $f \in \mathcal{L}_1$ and \widehat{f} is given explicitly or can be approximated numerically efficiently by some function ϑ and that v is (at least) locally integrable. At several places, we assume $v \in \mathcal{L}_1$, which usually can be achieved by setting $\alpha \neq \mathbf{0}$. We describe the COS method in detail to approximate the right hand-side of Equation (5). Let $\mathbf{M} \in \mathbb{R}_+^d$ large enough, so that

$$\int_{\mathbb{R}^d} v(\mathbf{x})f(\mathbf{x})d\mathbf{x} \approx \int_{[-\mathbf{M}, \mathbf{M}]} v(\mathbf{x})f(\mathbf{x})d\mathbf{x}. \quad (8)$$

Let $\mathbf{L} \geq \mathbf{M}$. If f is centered around zero, we truncate f and on $[-\mathbf{L}, \mathbf{L}]$ and approximate the truncated damped density by a Fourier-series. We intuitively have that

$$f \approx f1_{[-\mathbf{L}, \mathbf{L}]} \approx \sum'_{\mathbf{0} \leq \mathbf{k} \leq \mathbf{N}} a_{\mathbf{k}} e_{\mathbf{k}} 1_{[-\mathbf{L}, \mathbf{L}]} \approx \sum'_{\mathbf{0} \leq \mathbf{k} \leq \mathbf{N}} c_{\mathbf{k}} e_{\mathbf{k}} 1_{[-\mathbf{L}, \mathbf{L}]} \approx \sum'_{\mathbf{0} \leq \mathbf{k} \leq \mathbf{N}} \tilde{c}_{\mathbf{k}} e_{\mathbf{k}} 1_{[-\mathbf{L}, \mathbf{L}]}, \quad (9)$$

where we define the basis functions

$$e_{\mathbf{k}}(\mathbf{x}) = \prod_{h=1}^d \cos\left(k_h \pi \frac{x_h + L_h}{2L_h}\right), \quad \mathbf{x} \in \mathbb{R}^d, \quad \mathbf{k} \in \mathbb{N}_0^d,$$

and the classical Fourier-cosine coefficients of $f1_{[-\mathbf{L}, \mathbf{L}]}$ are given for $\mathbf{k} \in \mathbb{N}_0^d$ by

$$\begin{aligned} a_{\mathbf{k}} &= \frac{1}{\prod_{h=1}^d L_h} \int_{[-\mathbf{L}, \mathbf{L}]} f(\mathbf{x}) e_{\mathbf{k}}(\mathbf{x}) d\mathbf{x} \\ &\approx \frac{1}{\prod_{h=1}^d L_h} \int_{\mathbb{R}^d} f(\mathbf{x}) e_{\mathbf{k}}(\mathbf{x}) d\mathbf{x} \end{aligned} \quad (10)$$

$$= \frac{1}{2^{d-1} \prod_{h=1}^d L_h} \sum_{\mathbf{s}=(1, \pm 1, \dots, \pm 1) \in \mathbb{R}^d} \Re \left\{ \widehat{f} \left(\frac{\pi}{2} \frac{\mathbf{s} \mathbf{k}}{\mathbf{L}} \right) \exp \left(i \frac{\pi}{2} \mathbf{s} \cdot \mathbf{k} \right) \right\} =: c_{\mathbf{k}} \quad (11)$$

$$\approx \frac{1}{2^{d-1} \prod_{h=1}^d L_h} \sum_{\mathbf{s}=(1, \pm 1, \dots, \pm 1) \in \mathbb{R}^d} \Re \left\{ \vartheta \left(\frac{\pi}{2} \frac{\mathbf{s} \mathbf{k}}{\mathbf{L}} \right) \exp \left(i \frac{\pi}{2} \mathbf{s} \cdot \mathbf{k} \right) \right\} =: \tilde{c}_{\mathbf{k}}. \quad (12)$$

Sometimes it is necessary to choose $\mathbf{L} > \mathbf{M}$ to ensure that $c_{\mathbf{k}}$ is close enough to $a_{\mathbf{k}}$. The key insight of the COS method is the fact that the integral at the right-hand side of Equation (10) can be solved explicitly¹. If \hat{f} need to be approximated by some function ϑ , we use $\tilde{c}_{\mathbf{k}}$ instead of $c_{\mathbf{k}}$.

The idea of the multidimensional COS method is to approximate f as in (9) and hence the right-hand side of Equation (8) by

$$\int_{[-M, M]} v(\mathbf{x}) f(\mathbf{x}) d\mathbf{x} \approx \int_{[-M, M]} v(\mathbf{x}) \sum'_{\mathbf{0} \leq \mathbf{k} \leq \mathbf{N}} \tilde{c}_{\mathbf{k}} e_{\mathbf{k}}(\mathbf{x}) d\mathbf{x} \quad (13)$$

$$= \sum'_{\mathbf{0} \leq \mathbf{k} \leq \mathbf{N}} \tilde{c}_{\mathbf{k}} \underbrace{\int_{[-M, M]} v(\mathbf{x}) e_{\mathbf{k}}(\mathbf{x}) d\mathbf{x}}_{=: v_{\mathbf{k}}} \quad (14)$$

$$\approx \sum'_{\mathbf{0} \leq \mathbf{k} \leq \mathbf{N}} \tilde{c}_{\mathbf{k}} \underbrace{\int_{\mathbb{R}^d} v(\mathbf{x}) e_{\mathbf{k}}(\mathbf{x}) d\mathbf{x}}_{=: \tilde{v}_{\mathbf{k}}} \quad (15)$$

Classical COS method: If $\boldsymbol{\alpha} = \mathbf{0}$, we speak of the classical COS method. In important cases, the coefficients $v_{\mathbf{k}}$ can be obtained explicitly, i.e., the integral at right hand-side of Equation (14) can be solved analytically. Examples in finance include in one dimension plain vanilla put and call options and digital options, see Fang and Oosterlee (2009a) and in two dimensions geometric basket options, call-on-maximum options or put-on-minimum options, see Ruijter and Oosterlee (2012). In general dimensions, $v_{\mathbf{k}}$ of a CDF are given in closed-form, see Example 5.1. In the case that integral in Equation (14) cannot be solved directly (e.g. for arithmetic basket options), Ruijter and Oosterlee (2012) propose to solve the integral in Equation (14) numerically to obtain $v_{\mathbf{k}}$, e.g., by the discrete cosine transform or some quadrature rule. However, solving the integral in (14) numerically for each \mathbf{k} is expensive and slows down the COS method significantly.

Damped COS method: If $\boldsymbol{\alpha} \neq \mathbf{0}$, we speak of the damped COS method. Assume that v is integrable, which usually can be achieved by setting $\boldsymbol{\alpha} \neq \mathbf{0}$. Then we propose to approximate $v_{\mathbf{k}}$ by $\tilde{v}_{\mathbf{k}}$. This works if \mathbf{M} is large enough. Similar to the solution presented in Equation (11), the coefficients $\tilde{v}_{\mathbf{k}}$ are given analytically:

$$\tilde{v}_{\mathbf{k}} = \frac{1}{2^{d-1}} \sum_{\mathbf{s}=(1, \pm 1, \dots, \pm 1) \in \mathbb{R}^d} \Re \left\{ \hat{v} \left(\frac{\pi}{2} \frac{\mathbf{s} \mathbf{k}}{\mathbf{L}} \right) \exp \left(i \frac{\pi}{2} \mathbf{s} \cdot \mathbf{k} \right) \right\}. \quad (16)$$

In finance, the function \hat{v} is known explicitly for many rainbow options, e.g., arithmetic basket options, spread options and put and call options on the minimum or maximum of d assets.

In the remainder of the article, we will prove under which conditions the integral (1) can be approximated by the (damped) COS method.

Remark 3.2. In the special case that \hat{f} only takes real values, the computational cost of the COS method can be reduced by (about) the factor a half, since $c_{\mathbf{k}} = 0$ if $\sum_{h=1}^d k_h$ is odd.

In order to prove the convergence of the COS method in Theorem 3.5 and Corollary 3.6, we need the concept of *COS-admissibility*, which is introduced in Definition 3.3 and extends Junike and Pankrashkin (2022, Def. 1) to the multidimensional setting.

Definition 3.3. Let $\mathbf{L} = (L_1, \dots, L_d) \in \mathbb{R}_+^d$. A function $f \in \mathcal{L}^1$ is called *COS-admissible*, if

$$B_f(\mathbf{L}) := \sum'_{\mathbf{k} \in \mathbb{N}_0^d} \frac{1}{\prod_{h=1}^d L_h} \left| \int_{\mathbb{R}^d \setminus [-L, L]} f(\mathbf{x}) e_{\mathbf{k}}(\mathbf{x}) d\mathbf{x} \right|^2 \rightarrow 0, \quad \min_{h=1, \dots, d} L_h \rightarrow \infty.$$

By Proposition 3.4, it follows that bounded densities with existing moments are COS-admissible, which indicates that the class of d -dimensional, COS-admissible densities is large.

¹We use $\prod_{h=1}^d \cos \theta_h = \frac{1}{2^{d-1}} \sum_{\mathbf{s}=(1, \pm 1, \dots, \pm 1) \in \mathbb{R}^d} \cos(\mathbf{s} \cdot \boldsymbol{\theta})$, $\boldsymbol{\theta} \in \mathbb{R}^d$, which follows by mathematical induction, the fact that cosine is an even function and the trigonometric identities stated in Equations (4.3.17) and (4.3.31) in Abramowitz and Stegun (1972, Eqs. (4.3.17, 4.3.31)).

Proposition 3.4. Assume $f \in \mathcal{L}^1 \cap \mathcal{L}^2$ with

$$\int_{\mathbb{R}^d} |\mathbf{x}|^{2d} |f(\mathbf{x})|^2 d\mathbf{x} < \infty. \quad (17)$$

Then f is COS-admissible. Let $\mathbf{L} = (L_1, \dots, L_d) \in \mathbb{R}_+^d$ then it holds that

$$B_f(\mathbf{L}) \leq \Xi \int_{\mathbb{R}^d \setminus [-\mathbf{L}, \mathbf{L}]} \prod_{h=1}^d \max \{x_h^2 L_h^{-2}, 1\} |f(\mathbf{x})|^2 d\mathbf{x} \quad (18)$$

$$\leq \frac{\Xi}{d \min_{h=1, \dots, d} L_h^{2d}} \int_{\mathbb{R}^d \setminus [-\mathbf{L}, \mathbf{L}]} |\mathbf{x}|^{2d} |f(\mathbf{x})|^2 d\mathbf{x} + \Xi \int_{\mathbb{R}^d \setminus [-\mathbf{L}, \mathbf{L}]} |f(\mathbf{x})|^2 d\mathbf{x}, \quad (19)$$

where $\Xi = \frac{\pi^2}{3} \sum_{h=1}^d \left(\frac{\pi^2}{3} + 1 \right)^{h-1}$.

Proof. Let $\mathbf{L} \in \mathbb{R}_+^d$ and $\mathbf{j} \in \mathbb{Z}^d$. It follows by Parseval's identity

$$\begin{aligned} \int_{[2\mathbf{j}\mathbf{L}-\mathbf{L}, 2\mathbf{j}\mathbf{L}+\mathbf{L}]} |f(\mathbf{x})|^2 d\mathbf{x} &= \sum'_{\mathbf{k} \in \mathbb{N}_0^d} \frac{1}{\prod_{h=1}^d L_h} \left| \int_{[2\mathbf{j}\mathbf{L}-\mathbf{L}, 2\mathbf{j}\mathbf{L}+\mathbf{L}]} f(\mathbf{x}) \prod_{h=1}^d \underbrace{\cos \left(k_h \pi \frac{x_h - (2j_h L_h - L_h)}{2L_h} \right)}_{=(-1)^{j_h k_h} \cos \left(k_h \pi \frac{x_h + L_h}{2L_h} \right)} d\mathbf{x} \right|^2 \\ &= \sum'_{\mathbf{k} \in \mathbb{N}_0^d} \frac{1}{\prod_{h=1}^d L_h} \left| \int_{[2\mathbf{j}\mathbf{L}-\mathbf{L}, 2\mathbf{j}\mathbf{L}+\mathbf{L}]} f(\mathbf{x}) e_{\mathbf{k}}(\mathbf{x}) d\mathbf{x} \right|^2. \end{aligned} \quad (20)$$

By the Cauchy-Schwarz inequality, we obtain with $g(\mathbf{j}) := \prod_{h=1}^d \max\{|j_h|, 1\}$,

$$\begin{aligned} \left| \int_{\mathbb{R}^d \setminus [-\mathbf{L}, \mathbf{L}]} f(\mathbf{x}) e_{\mathbf{k}}(\mathbf{x}) d\mathbf{x} \right|^2 &= \left| \sum_{\mathbf{j} \in \mathbb{Z}^d \setminus \{\mathbf{0}\}} \frac{g(\mathbf{j})}{g(\mathbf{j})} \int_{[2\mathbf{j}\mathbf{L}-\mathbf{L}, 2\mathbf{j}\mathbf{L}+\mathbf{L}]} f(\mathbf{x}) e_{\mathbf{k}}(\mathbf{x}) d\mathbf{x} \right|^2 \\ &\leq \underbrace{\left(\sum_{\mathbf{j} \in \mathbb{Z}^d \setminus \{\mathbf{0}\}} \frac{1}{(g(\mathbf{j}))^2} \right)}_{=\Xi} \sum_{\mathbf{j} \in \mathbb{Z}^d \setminus \{\mathbf{0}\}} (g(\mathbf{j}))^2 \left| \int_{[2\mathbf{j}\mathbf{L}-\mathbf{L}, 2\mathbf{j}\mathbf{L}+\mathbf{L}]} f(\mathbf{x}) e_{\mathbf{k}}(\mathbf{x}) d\mathbf{x} \right|^2. \end{aligned} \quad (21)$$

The fact that $\Xi = \frac{\pi^2}{3} \sum_{h=1}^d \left(\frac{\pi^2}{3} + 1 \right)^{h-1}$ can be shown by mathematical induction over d . Then it follows that

$$\begin{aligned} B_f(\mathbf{L}) &\stackrel{(21)}{\leq} \Xi \sum_{\mathbf{j} \in \mathbb{Z}^d \setminus \{\mathbf{0}\}} (g(\mathbf{j}))^2 \sum'_{\mathbf{k} \in \mathbb{N}_0^d} \frac{1}{\prod_{h=1}^d L_h} \left| \int_{[2\mathbf{j}\mathbf{L}-\mathbf{L}, 2\mathbf{j}\mathbf{L}+\mathbf{L}]} f(\mathbf{x}) e_{\mathbf{k}}(\mathbf{x}) d\mathbf{x} \right|^2 \\ &\stackrel{(20)}{=} \Xi \sum_{\mathbf{j} \in \mathbb{Z}^d \setminus \{\mathbf{0}\}} (g(\mathbf{j}))^2 \int_{[2\mathbf{j}\mathbf{L}-\mathbf{L}, 2\mathbf{j}\mathbf{L}+\mathbf{L}]} |f(\mathbf{x})|^2 d\mathbf{x}. \end{aligned}$$

For $\mathbf{j} \in \mathbb{Z}^d$ and $\mathbf{x} \in [2\mathbf{j}\mathbf{L}-\mathbf{L}, 2\mathbf{j}\mathbf{L}+\mathbf{L}]$, one has $|j_h| \leq \frac{|x_h|}{L_h}$, $h = 1, \dots, d$. It follows that $(g(\mathbf{j}))^2 \leq \prod_{h=1}^d \max \{x_h^2 L_h^{-2}, 1\}$, which implies Inequality (18). By Young's inequality, it holds that

$$\prod_{h=1}^d (\max \{x_h^{2d} L_h^{-2d}, 1\})^{\frac{1}{d}} \leq \frac{1}{d} \sum_{h=1}^d \max \{x_h^{2d} L_h^{-2d}, 1\} \leq \frac{|\mathbf{x}|^{2d}}{d \min_{h=1, \dots, d} L_h^{2d}} + 1.$$

In the last inequality, we used $\max\{a, b\} \leq a + b$ for any $a, b \geq 0$ and $\sum_{h=1}^d x_h^{2d} \leq |\mathbf{x}|^{2d}$, which follows from the monotonicity of the p -norm. Hence, Inequality (19) holds. Assumption (17) and $f \in \mathcal{L}^2$ imply $B_f(\mathbf{L}) \rightarrow 0$, $\min_{h=1, \dots, d} L_h \rightarrow \infty$. \square

The following theorem shows that multivariate densities can be approximated by a cosine expansion. The theorem also includes numerical uncertainty on the Fourier transform \hat{f} .

Theorem 3.5. *Assume that $f \in \mathcal{L}^1 \cap \mathcal{L}^2$ is COS-admissible. Let $\vartheta : \mathbb{R}^d \rightarrow \mathbb{C}$ and define $\tilde{c}_{\mathbf{k}}$ as in Equation (12). For any $\varepsilon > 0$ there is a $\mathbf{L} \in \mathbb{R}_+^d$, a $\mathbf{N} \in \mathbb{N}^d$ and a $\gamma > 0$ so that $\|\hat{f} - \vartheta\|_\infty < \gamma$ implies*

$$\left\| f - \sum'_{\mathbf{0} \leq \mathbf{k} \leq \mathbf{N}} \tilde{c}_{\mathbf{k}} e_{\mathbf{k}} 1_{[-\mathbf{L}, \mathbf{L}]} \right\|_2 < \varepsilon.$$

Note that \mathbf{N} depends on \mathbf{L} and γ depends on both \mathbf{L} and \mathbf{N} .

Proof. Define $e_{\mathbf{k}}^{\mathbf{L}} = e_{\mathbf{k}} 1_{[-\mathbf{L}, \mathbf{L}]}$. It holds for $\mathbf{l}, \mathbf{k} \in \mathbb{N}_0^d$ that

$$\langle e_{\mathbf{k}}^{\mathbf{L}}, e_{\mathbf{l}}^{\mathbf{L}} \rangle = \begin{cases} 2^{\Lambda(\mathbf{k})} \prod_{h=1}^d L_h & , \mathbf{k} = \mathbf{l}, \\ 0 & , \text{otherwise} \end{cases} \quad (22)$$

where Λ is defined Section 2. For any $\mathbf{L} \in \mathbb{R}_+^d$ and $\mathbf{N} \in \mathbb{N}^d$, it holds that

$$\begin{aligned} \left\| f - \sum'_{\mathbf{0} \leq \mathbf{k} \leq \mathbf{N}} \tilde{c}_{\mathbf{k}} e_{\mathbf{k}}^{\mathbf{L}} \right\|_2 &\leq \underbrace{\|f - f 1_{[-\mathbf{L}, \mathbf{L}]} \|_2}_{=: A_1(\mathbf{L})} + \underbrace{\left\| f 1_{[-\mathbf{L}, \mathbf{L}]} - \sum'_{\mathbf{0} \leq \mathbf{k} \leq \mathbf{N}} a_{\mathbf{k}} e_{\mathbf{k}}^{\mathbf{L}} \right\|_2}_{=: A_2(\mathbf{L}, \mathbf{N})} \\ &\quad + \underbrace{\left\| \sum'_{\mathbf{0} \leq \mathbf{k} \leq \mathbf{N}} (a_{\mathbf{k}} - c_{\mathbf{k}}) e_{\mathbf{k}}^{\mathbf{L}} \right\|_2}_{=: A_3(\mathbf{L}, \mathbf{N})} + \underbrace{\left\| \sum'_{\mathbf{0} \leq \mathbf{k} \leq \mathbf{N}} (c_{\mathbf{k}} - \tilde{c}_{\mathbf{k}}) e_{\mathbf{k}}^{\mathbf{L}} \right\|_2}_{=: A_4(\mathbf{L}, \mathbf{N})}. \end{aligned}$$

Further,

$$A_3(\mathbf{L}, \mathbf{N})^2 = \sum_{\mathbf{0} \leq \mathbf{k} \leq \mathbf{N}} \sum_{\mathbf{0} \leq \mathbf{l} \leq \mathbf{N}} \frac{1}{2^{\Lambda(\mathbf{k}) + \Lambda(\mathbf{l})}} (a_{\mathbf{k}} - c_{\mathbf{k}})(a_{\mathbf{l}} - c_{\mathbf{l}}) \langle e_{\mathbf{k}}^{\mathbf{L}}, e_{\mathbf{l}}^{\mathbf{L}} \rangle \leq \sum'_{\mathbf{k} \in \mathbb{N}_0^d} \prod_{h=1}^d \{L_h\} |a_{\mathbf{k}} - c_{\mathbf{k}}|^2 = B_f(\mathbf{L}),$$

see Definition 3.3. For $\varepsilon > 0$, choose $\mathbf{L} \in \mathbb{R}_+^d$ such that $A_1(\mathbf{L}) < \frac{\varepsilon}{4}$ and $B_f(\mathbf{L}) < (\frac{\varepsilon}{4})^2$. Hence, $A_3(\mathbf{L}, \mathbf{N}) < \frac{\varepsilon}{4}$. Then choose $\mathbf{N} \in \mathbb{N}^d$ such that $A_2(\mathbf{L}, \mathbf{N}) < \frac{\varepsilon}{4}$. Such \mathbf{N} exists by classically Fourier analysis. By the definition of $c_{\mathbf{k}}$ and $\tilde{c}_{\mathbf{k}}$, see Equation (11), it follows that

$$|c_{\mathbf{k}} - \tilde{c}_{\mathbf{k}}| \leq \frac{1}{2^{d-1} \prod_{h=1}^d L_h} \sum_{\mathbf{s}=(1, \pm 1, \dots, \pm 1) \in \mathbb{R}^d} \left| \hat{f}\left(\frac{\pi}{2} \frac{\mathbf{s} \mathbf{k}}{\mathbf{L}}\right) - \vartheta\left(\frac{\pi}{2} \frac{\mathbf{s} \mathbf{k}}{\mathbf{L}}\right) \right| \leq \frac{\|\hat{f} - \vartheta\|_\infty}{\prod_{h=1}^d L_h}.$$

Similarly to the analysis of A_3 , we have

$$A_4(\mathbf{L}, \mathbf{N})^2 \leq \sum_{\mathbf{0} \leq \mathbf{k} \leq \mathbf{N}} \prod_{h=1}^d \{L_h\} |c_{\mathbf{k}} - \tilde{c}_{\mathbf{k}}|^2 \leq \frac{\|\hat{f} - \vartheta\|_\infty^2}{\prod_{h=1}^d L_h} \prod_{h=1}^d \{N_h + 1\}. \quad (23)$$

Choose $\gamma = \frac{\varepsilon}{4} \sqrt{\prod_{h=1}^d L_h} \left(\prod_{h=1}^d \{N_h + 1\} \right)^{-\frac{1}{2}}$. Then $\|\hat{f} - \vartheta\|_\infty < \gamma$ implies $A_4(\mathbf{L}, \mathbf{N}) < \frac{\varepsilon}{4}$, which concludes the proof. \square

Corollary 3.6 provides sufficient conditions in order to ensure that the COS method approximates the price of a rainbow option within a predefined error tolerance $\varepsilon > 0$ including numerical uncertainty on \hat{f} and numerical uncertainty on the cosine coefficients of the function of interest v : either because the $v_{\mathbf{k}}$ are approximated by solving the integral in Equation (14) numerically or because $v_{\mathbf{k}}$ are approximated by $\tilde{v}_{\mathbf{k}}$ defined in Equation (15).

Corollary 3.6. (Convergence of the COS method). Let $f \in \mathcal{L}^1 \cap \mathcal{L}^2$ be COS-admissible and $v : \mathbb{R}^d \rightarrow \mathbb{R}$ be locally in \mathcal{L}^2 , that is, $v1_{[-\mathbf{M}, \mathbf{M}]} \in \mathcal{L}^2$ for any $\mathbf{M} \in \mathbb{R}_+^d$. Assume $vf \in \mathcal{L}^1$, then the integral of the product of f and v can be approximated by a finite sum as follows: Let $\varepsilon > 0$. Let $\mathbf{M} \in \mathbb{R}_+^d$ and $\xi > 0$ such that

$$\int_{\mathbb{R}^d \setminus [-\mathbf{M}, \mathbf{M}]} |v(\mathbf{x})f(\mathbf{x})| d\mathbf{x} \leq \frac{\varepsilon}{3}, \quad \|v1_{[-\mathbf{M}, \mathbf{M}]} \|_2 \leq \xi. \quad (24)$$

Let $\mathbf{L} \geq \mathbf{M}$ such that

$$\|f - f1_{[-\mathbf{L}, \mathbf{L}]} \|_2 \leq \frac{\varepsilon}{12\xi} \quad (25)$$

and

$$B_f(\mathbf{L}) \leq \left(\frac{\varepsilon}{12\xi} \right)^2. \quad (26)$$

Choose $\mathbf{N} \in \mathbb{N}^d$ large enough, so that

$$\left\| f1_{[-\mathbf{L}, \mathbf{L}]} - \sum'_{\mathbf{0} \leq \mathbf{k} \leq \mathbf{N}} a_{\mathbf{k}} e_{\mathbf{k}} 1_{[-\mathbf{L}, \mathbf{L}]} \right\|_2 \leq \frac{\varepsilon}{12\xi}. \quad (27)$$

For some $\vartheta : \mathbb{R}^d \rightarrow \mathbb{C}$ assume

$$\|\widehat{f} - \vartheta\|_{\infty} \leq \frac{\varepsilon}{12\xi} \frac{\sqrt{\prod_{h=1}^d L_h}}{\sqrt{\prod_{h=1}^d \{N_h + 1\}}}. \quad (28)$$

Let $\eta > 0$ such that $\sum'_{\mathbf{0} \leq \mathbf{k} \leq \mathbf{N}} |\tilde{c}_{\mathbf{k}}|^2 \leq \eta$. Let $(\tilde{v}_{\mathbf{k}})_{\mathbf{k} \in \mathbb{N}_0^d}$ such that

$$\sum'_{\mathbf{0} \leq \mathbf{k} \leq \mathbf{N}} |\tilde{v}_{\mathbf{k}} - v_{\mathbf{k}}|^2 \leq \frac{\varepsilon^2}{9\eta}. \quad (29)$$

Then it follows that

$$\left| \int_{\mathbb{R}^d} v(\mathbf{x})f(\mathbf{x})d\mathbf{x} - \sum'_{\mathbf{0} \leq \mathbf{k} \leq \mathbf{N}} \tilde{c}_{\mathbf{k}} \tilde{v}_{\mathbf{k}} \right| \leq \varepsilon. \quad (30)$$

Proof. Define $e_{\mathbf{k}}^L = e_{\mathbf{k}} 1_{[-\mathbf{L}, \mathbf{L}]}$. Let $A_1(\mathbf{L})$, $A_2(\mathbf{L}, \mathbf{N})$ and $A_4(\mathbf{L}, \mathbf{N})$ be as in the proof of Theorem 3.5. By Inequalities (23, 28) it follows that $A_4(\mathbf{L}, \mathbf{N}) \leq \frac{\varepsilon}{12\xi}$. Due to $v_{\mathbf{k}} = \langle v1_{[-\mathbf{M}, \mathbf{M}]}, e_{\mathbf{k}}^L \rangle$ and applying Theorem 3.5 and the Cauchy-Schwarz inequality, we have that

$$\begin{aligned} & \left| \int_{\mathbb{R}^d} v(\mathbf{x})f(\mathbf{x})d\mathbf{x} - \sum'_{\mathbf{0} \leq \mathbf{k} \leq \mathbf{N}} \tilde{c}_{\mathbf{k}} \tilde{v}_{\mathbf{k}} \right| \\ &= \left| \int_{\mathbb{R}^d \setminus [-\mathbf{M}, \mathbf{M}]} v(\mathbf{x})f(\mathbf{x})d\mathbf{x} + \langle v1_{[-\mathbf{M}, \mathbf{M}]}, f \rangle - \sum'_{\mathbf{0} \leq \mathbf{k} \leq \mathbf{N}} \tilde{c}_{\mathbf{k}} \langle v1_{[-\mathbf{M}, \mathbf{M}]}, e_{\mathbf{k}}^L \rangle - \sum'_{\mathbf{0} \leq \mathbf{k} \leq \mathbf{N}} \tilde{c}_{\mathbf{k}} (\tilde{v}_{\mathbf{k}} - v_{\mathbf{k}}) \right| \\ &\leq \underbrace{\int_{\mathbb{R}^d \setminus [-\mathbf{M}, \mathbf{M}]} |v(\mathbf{x})f(\mathbf{x})| d\mathbf{x}}_{=: D_1(\mathbf{M})} + \left| \langle v1_{[-\mathbf{M}, \mathbf{M}]}, f - \sum'_{\mathbf{0} \leq \mathbf{k} \leq \mathbf{N}} \tilde{c}_{\mathbf{k}} e_{\mathbf{k}}^L \rangle \right| + \underbrace{\sqrt{\sum'_{\mathbf{0} \leq \mathbf{k} \leq \mathbf{N}} |\tilde{v}_{\mathbf{k}} - v_{\mathbf{k}}|^2 \sum'_{\mathbf{0} \leq \mathbf{k} \leq \mathbf{N}} |\tilde{c}_{\mathbf{k}}|^2}}_{=: D_2(\mathbf{N}, \mathbf{L}, \mathbf{M})} \\ &< \frac{\varepsilon}{3} + \|v1_{[-\mathbf{M}, \mathbf{M}]} \|_2 \left\| f - \sum'_{\mathbf{0} \leq \mathbf{k} \leq \mathbf{N}} \tilde{c}_{\mathbf{k}} e_{\mathbf{k}}^L \right\|_2 + \frac{\varepsilon}{3} \\ &< \frac{\varepsilon}{3} + \xi \left(A_1(\mathbf{L}) + A_2(\mathbf{L}, \mathbf{N}) + \sqrt{B_f(\mathbf{L})} + A_4(\mathbf{L}, \mathbf{N}) \right) + \frac{\varepsilon}{3} \\ &\leq \frac{\varepsilon}{3} + \xi \left(\frac{\varepsilon}{12\xi} + \frac{\varepsilon}{12\xi} + \frac{\varepsilon}{12\xi} + \frac{\varepsilon}{12\xi} \right) + \frac{\varepsilon}{3} = \varepsilon. \end{aligned}$$

□

In Junike and Pankrashkin (2022) and Junike (2024), it is assumed that f has semi-heavy tails, i.e., f decays exponentially or faster. Here, we make the same assumption in multivariate dimensions to be able to estimate \mathbf{M} , \mathbf{L} and \mathbf{N} .

Definition 3.7. A function $f : \mathbb{R}^d \rightarrow \mathbb{R}$ decays exponentially if there are $C_1, C_2, m > 0$ such that for $|\mathbf{x}| > m$ it holds that $|f(\mathbf{x})| \leq C_1 e^{-C_2 |\mathbf{x}|}$.

Lemma 3.8. Let $f \in \mathcal{L}^1 \cap \mathcal{L}^2$. Let $\mathbf{M}, \mathbf{L} \in \mathbb{R}_+^d$ with $\mathbf{M} \leq \mathbf{L}$ then it holds that

$$\|f1_{[-\mathbf{L}, \mathbf{L}]} - \sum'_{\mathbf{0} \leq \mathbf{k} \leq \mathbf{N}} a_{\mathbf{k}} e_{\mathbf{k}} 1_{[-\mathbf{L}, \mathbf{L}]} \|_2^2 \leq \int_{\mathbb{R}^d} |f(\mathbf{x})|^2 d\mathbf{x} - \prod_{h=1}^d L_h \sum'_{\mathbf{0} \leq \mathbf{k} \leq \mathbf{N}} |c_{\mathbf{k}}|^2 + G(\mathbf{L}),$$

where

$$G(\mathbf{L}) := B_f(\mathbf{L}) + 2\sqrt{B_f(\mathbf{L}) \int_{\mathbb{R}^d} |f(\mathbf{x})|^2 d\mathbf{x}}. \quad (31)$$

Proof. Let

$$\phi_{\mathbf{k}} := \frac{1}{\prod_{h=1}^d L_h} \int_{\mathbb{R}^d \setminus [-\mathbf{L}, \mathbf{L}]} f(\mathbf{x}) e_{\mathbf{k}}(\mathbf{x}) d\mathbf{x}, \quad \mathbf{k} \in \mathbb{N}_0^d.$$

It holds that $c_{\mathbf{k}} = a_{\mathbf{k}} + \phi_{\mathbf{k}}$. It follows by the Cauchy-Schwarz inequality that

$$\begin{aligned} \prod_{h=1}^d L_h \sum'_{\mathbf{0} \leq \mathbf{k} \leq \mathbf{N}} |c_{\mathbf{k}}|^2 &= \prod_{h=1}^d L_h \left(\sum'_{\mathbf{0} \leq \mathbf{k} \leq \mathbf{N}} |a_{\mathbf{k}}|^2 + \sum'_{\mathbf{0} \leq \mathbf{k} \leq \mathbf{N}} |\phi_{\mathbf{k}}|^2 + 2 \sum'_{\mathbf{0} \leq \mathbf{k} \leq \mathbf{N}} |\phi_{\mathbf{k}}| |a_{\mathbf{k}}| \right) \\ &\leq \prod_{h=1}^d L_h \sum'_{\mathbf{0} \leq \mathbf{k} \leq \mathbf{N}} |a_{\mathbf{k}}|^2 + B_f(\mathbf{L}) + 2 \prod_{h=1}^d L_h \sqrt{\sum'_{\mathbf{0} \leq \mathbf{k} \leq \mathbf{N}} |\phi_{\mathbf{k}}|^2 \sum'_{\mathbf{0} \leq \mathbf{k} \leq \mathbf{N}} |a_{\mathbf{k}}|^2} \\ &\stackrel{(20)}{\leq} \prod_{h=1}^d L_h \sum'_{\mathbf{0} \leq \mathbf{k} \leq \mathbf{N}} |a_{\mathbf{k}}|^2 + \underbrace{B_f(\mathbf{L}) + 2\sqrt{B_f(\mathbf{L}) \int_{\mathbb{R}^d} |f(\mathbf{x})|^2 d\mathbf{x}}}_{=G(\mathbf{L})} \end{aligned} \quad (32)$$

$$\stackrel{(20)}{\leq} \int_{\mathbb{R}^d} |f(\mathbf{x})|^2 d\mathbf{x} + G(\mathbf{L}). \quad (33)$$

Hence,

$$\begin{aligned} \|f1_{[-\mathbf{L}, \mathbf{L}]} - \sum'_{\mathbf{0} \leq \mathbf{k} \leq \mathbf{N}} a_{\mathbf{k}} e_{\mathbf{k}} 1_{[-\mathbf{L}, \mathbf{L}]} \|_2^2 &\stackrel{(22)}{\leq} \prod_{h=1}^d L_h \sum'_{\substack{\mathbf{k}_1 > N_1 \text{ or } \dots \text{ or } \mathbf{k}_d > N_d}} |a_{\mathbf{k}}|^2 \\ &= \prod_{h=1}^d L_h \sum'_{\mathbf{k} \in \mathbb{N}_0^d} |a_{\mathbf{k}}|^2 - \prod_{h=1}^d L_h \sum'_{\mathbf{0} \leq \mathbf{k} \leq \mathbf{N}} |a_{\mathbf{k}}|^2 \\ &\stackrel{(20,32)}{\leq} \int_{\mathbb{R}^d} |f(\mathbf{x})|^2 d\mathbf{x} - \prod_{h=1}^d L_h \sum'_{\mathbf{0} \leq \mathbf{k} \leq \mathbf{N}} |c_{\mathbf{k}}|^2 + G(\mathbf{L}). \end{aligned}$$

□

Theorem 3.9. (Classical COS method: Find \mathbf{M} and \mathbf{L}). Let $f \in \mathcal{L}^1 \cap \mathcal{L}^2$ be a nonnegative function satisfying Inequality (17). Let $v : \mathbb{R}^d \rightarrow \mathbb{R}$ be bounded with $\|v\|_{\infty} \in (0, \infty)$. Let $n \geq 2$ be some even number and assume the moments of f of n^{th} -order exist, i.e.,

$$m_h(n) := \int_{\mathbb{R}^d} x_h^n f(\mathbf{x}) d\mathbf{x} = i^{-n} \frac{\partial^n}{\partial u_h^n} \widehat{f}(\mathbf{u}) \Big|_{\mathbf{u}=\mathbf{0}} \in (0, \infty), \quad h = 1, \dots, d. \quad (34)$$

Assume that f decays exponentially. Let $\varepsilon > 0$ be small enough. Define

$$M_h := \left(\frac{3d \|v\|_\infty m_h(n)}{\varepsilon} \right)^{\frac{1}{n}}, \quad h = 1, \dots, d, \quad (35)$$

and $\mathbf{L} = \mathbf{M} = (M_1, \dots, M_d) \in \mathbb{R}_+^d$. There is a $\mathbf{N} \in \mathbb{N}_0^d$ so that

$$\left| \int_{\mathbb{R}^d} v(\mathbf{x}) f(\mathbf{x}) d\mathbf{x} - \sum'_{\mathbf{0} \leq \mathbf{k} \leq \mathbf{N}} c_{\mathbf{k}} v_{\mathbf{k}} \right| \leq \varepsilon. \quad (36)$$

Corollary 3.10. (Damped COS method: Find \mathbf{M} and \mathbf{L}). Assume that all assumptions in Theorem 3.9 hold and that $v \in \mathcal{L}^1 \cap \mathcal{L}^2$, v satisfies Inequality (17) and v decays exponentially. Define $\tilde{v}_{\mathbf{k}}$ as in (15). Then Inequality (36) still holds, if we replace $v_{\mathbf{k}}$ by $\tilde{v}_{\mathbf{k}}$.

Corollary 3.11. (Find \mathbf{N}). Assume that all assumptions in Corollary 3.10 hold. The number of terms can be chosen by any $\mathbf{N} \in \mathbb{N}_0^d$ such that

$$\left| (2\pi)^{-d} \int_{\mathbb{R}^d} |\widehat{f}(\mathbf{u})|^2 d\mathbf{u} - \prod_{h=1}^d L_h \sum'_{\mathbf{0} \leq \mathbf{k} \leq \mathbf{N}} |c_{\mathbf{k}}|^2 \right| \leq \frac{\varepsilon^2}{162 \|v\|_2^2}. \quad (37)$$

Proof. We first prove Theorem 3.9: Equation (34) follows by Bauer (1996, Thm 25.2). For $h \in \{1, \dots, d\}$ let $\pi_h : \mathbb{R}^d \rightarrow \mathbb{R}$, $\mathbf{x} \mapsto x_h$. Let λ^d be the Lebesgue measure on \mathbb{R}^d and define the finite and positive measure $\mu := f \lambda^d$. By Markov's inequality, see Bauer (1992, Lemma 20.1), it follows that

$$\int_{\mathbb{R}^d \setminus [-\mathbf{M}, \mathbf{M}]} |v(\mathbf{x}) f(\mathbf{x})| d\mathbf{x} \leq \|v\|_\infty \sum_{h=1}^d \mu(\{\mathbf{x} \in \mathbb{R}^d : |\pi_h(\mathbf{x})| \geq M_h\}) \leq \|v\|_\infty \sum_{h=1}^d \frac{m_h(n)}{M_h^n} = \frac{\varepsilon}{3}.$$

The last inequality follows by the definition of \mathbf{M} . Define $\xi := \|v\|_\infty \sqrt{2^d \prod_{h=1}^d M_h}$. It holds that $\|v 1_{[-\mathbf{M}, \mathbf{M}]}\|_2 \leq \xi$. Hence, the inequalities in (24) are satisfied. Next, we use the following auxiliary result: Let $s \geq 0$, $a > 0$ and $n \in \mathbb{N}_0$ and $d \in \mathbb{N}$. Then it holds by mathematical induction over n and Amann and Escher (2009, Theorem 8.11) that

$$\int_{\{\mathbf{x} \in \mathbb{R}^d : |\mathbf{x}| > s\}} e^{-a|\mathbf{x}|} |\mathbf{x}|^n d\mathbf{x} = \frac{d\pi^{\frac{d}{2}}}{\Gamma(1 + \frac{d}{2})} e^{-as} \frac{(n+d-1)!}{a^{n+d}} \sum_{k=0}^{n+d-1} \frac{(as)^k}{k!}. \quad (38)$$

For ε small enough, \mathbf{L} is large enough. Using that f decays exponentially and applying Equation (38), we obtain with $\ell := \min_{h=1, \dots, d} L_h$ that

$$\|f - f 1_{[-\mathbf{L}, \mathbf{L}]}\|_2 \leq C_1 \sqrt{\int_{\{\mathbf{x} \in \mathbb{R}^d : |\mathbf{x}| > \ell\}} e^{-2C_2|\mathbf{x}|} d\mathbf{x}} \leq \frac{\varepsilon}{12\xi}. \quad (39)$$

The last inequality holds true if ε is small enough because, thanks to Inequality (38), the term in the middle of (39) decreases exponentially in ε , while the term at the right-hand side of (39) goes to zero like $\varepsilon^{1+\frac{d}{2n}}$ for $\varepsilon \searrow 0$. Hence, Inequality (25) holds. By Inequality (19) it holds that $B_f(\mathbf{L}) \leq \varepsilon^2 (12\xi)^{-2}$ if ε is small enough because $B_f(\mathbf{L})$ decreases exponentially in ε : use Inequality (39) and observe that the term $\int_{\mathbb{R}^d \setminus [-\mathbf{L}, \mathbf{L}]} |\mathbf{x}|^{2d} |f(\mathbf{x})|^2 d\mathbf{x}$ converges exponentially thanks to Inequality (38). Hence, Inequality (26) holds. By classical Fourier analysis, there is a $\mathbf{N} \in \mathbb{N}_0^d$ such that Inequality (27) is satisfied. By assumption we have $\vartheta = \widehat{f}$ and $v_{\mathbf{k}} = \tilde{v}_{\mathbf{k}}$. Inequalities (28) and (29) hold trivially. Apply Corollary 3.6 to finish the proof of Theorem 3.9.

We prove Corollary 3.10: We have to show that Inequality (29) holds to proof Corollary 3.10. Let $G(\mathbf{L})$ be as in Equality (31). Observe $G(\mathbf{L}) \rightarrow 0$, $\min_h L_h \rightarrow \infty$ because f is COS-admissible by Proposition 3.4.

There is $\mathbf{P} \in \mathbb{R}_+^d$ and a $\gamma > 0$ such that $G(\mathbf{L}) \leq \gamma$ for all $\mathbf{L} \geq \mathbf{P}$. By Inequality (33), it follows for all $\mathbf{N} \in \mathbb{N}^d$ and all $\mathbf{L} \geq \mathbf{P}$ that

$$\sum'_{\mathbf{0} \leq \mathbf{k} \leq \mathbf{N}} |c_{\mathbf{k}}|^2 \leq \frac{\int_{\mathbb{R}^d} |f(\mathbf{x})|^2 d\mathbf{x} + \gamma}{\prod_{h=1}^d P_h} =: \eta < \infty. \quad (40)$$

It follows by Proposition 3.4 for all $\mathbf{N} \in \mathbb{N}^d$ that

$$\sum'_{\mathbf{0} \leq \mathbf{k} \leq \mathbf{N}} |\tilde{v}_{\mathbf{k}} - v_{\mathbf{k}}|^2 \leq \sum'_{\mathbf{k} \in \mathbb{N}_0^d} \left| \int_{\mathbb{R}^d \setminus [-\mathbf{M}, \mathbf{M}]} v(\mathbf{x}) e_{\mathbf{k}}(\mathbf{x}) d\mathbf{x} \right|^2 \leq \prod_{h=1}^d M_h B_v(\mathbf{M}) \leq \frac{\varepsilon^2}{9\eta} \quad (41)$$

the last inequality holds true if ε is small enough because the term in the middle of (41) decreases exponentially in ε since v decays exponentially, while the right-hand side of (41) goes to zero like ε^2 for $\varepsilon \searrow 0$.

We prove Corollary 3.11: For ε small enough, \mathbf{M} and \mathbf{L} are large enough. Apply the Plancherel theorem and Corollary 3.10 to f and $v := f$ to see that there is a $\mathbf{N} \in \mathbb{N}^d$ such that Inequality (37) holds. Let $G(\mathbf{L})$ be defined as in Equation (31). Note that $\|v1_{[-\mathbf{M}, \mathbf{M}]\|_2 \leq \|v\|_2$ for any $\mathbf{M} \in \mathbb{R}_+^d$. By Lemma 3.8 and the Plancherel theorem, it follows that

$$\begin{aligned} \|f1_{[-\mathbf{L}, \mathbf{L}]} - \sum'_{\mathbf{0} \leq \mathbf{k} \leq \mathbf{N}} a_{\mathbf{k}} e_{\mathbf{k}} 1_{[-\mathbf{L}, \mathbf{L}]\|_2^2 &\leq \left| (2\pi)^{-d} \int_{\mathbb{R}^d} |\hat{f}(\mathbf{u})|^2 d\mathbf{u} - \prod_{h=1}^d L_h \sum'_{\mathbf{0} \leq \mathbf{k} \leq \mathbf{N}} |c_{\mathbf{k}}|^2 \right| + G(\mathbf{L}) \\ &\leq \frac{\varepsilon^2}{162 \|v\|_2^2} + \frac{\varepsilon^2}{162 \|v\|_2^2} = \left(\frac{\varepsilon}{9 \|v\|_2} \right)^2. \end{aligned} \quad (42)$$

The last inequality holds because for $\varepsilon > 0$ small enough, \mathbf{L} is large enough so that $G(\mathbf{L}) \leq \frac{\varepsilon^2}{162 \|v\|_2^2}$. Note that we may replace the term $\frac{\varepsilon^2}{12\xi}$ in Inequalities (25, 26, 27) in Corollary 3.6 by $\frac{\varepsilon}{9 \|v\|_2}$ since $\hat{f} = \vartheta$ and $\|v\|_2 < \infty$. Apply Inequality (42) to conclude. \square

Remark 3.12. Provided the expression $(2\pi)^{-d} \int_{\mathbb{R}^d} |\hat{f}(\mathbf{u})|^2 d\mathbf{u}$ can be obtained precisely, Inequality (37) makes it possible to define a stopping criterion for \mathbf{N} . In particular, Inequality (37) enables us to determine \mathbf{N} while computing the coefficients $c_{\mathbf{k}}$: incrementally increase \mathbf{N} and compute $|c_{\mathbf{k}}|$ and $|c_{\mathbf{k}}|^2$ simultaneously. Stop when Inequality (37) is met. However, since right-hand side of Equation (37) converges to zero like $O(\varepsilon^2)$, rounding-off errors makes it difficult to find \mathbf{N} by Inequality (37) for very small ε . Using arbitrary-precision arithmetic instead of fixed-precision arithmetic should overcome this drawback.

The next theorem implies that the COS method converges exponentially if \hat{f} decays exponentially, i.e., if Inequality (43) holds for all $p > 0$. The cases (i) and (ii) in Theorem 3.13 treat the classical and the damped COS method, respectively. The bound for the order of convergence of the damped COS method is slightly sharper.

Theorem 3.13. (Order of convergence). Assume $f \in \mathcal{L}^1 \cap \mathcal{L}^2$ satisfies Inequality (17) and decays exponentially. Assume v is bounded. Let $\gamma > 0$ and $\beta \in (0, 1)$. For $n \in \mathbb{N}$, let $\mathbf{N} = (n, \dots, n) \in \mathbb{N}^d$ and $\mathbf{M} = \mathbf{L} = (\gamma n^\beta, \dots, \gamma n^\beta)$. Define $\tilde{v}_{\mathbf{k}}$ as in Equation (14) or Equation (15). Assume for some $p > \frac{d}{2}$ that

$$|\hat{f}(\mathbf{u})| \leq O(|\mathbf{u}|_\infty^{-p}), \quad |\mathbf{u}|_\infty \rightarrow \infty. \quad (43)$$

(i) Define $v_{\mathbf{k}}$ as in Equation (14). Then it holds that

$$\left| \int_{\mathbb{R}^d} v(\mathbf{x}) f(\mathbf{x}) d\mathbf{x} - \sum'_{\mathbf{0} \leq \mathbf{k} \leq \mathbf{N}} c_{\mathbf{k}} v_{\mathbf{k}} \right| \leq O\left(n^{-(1-\beta)p + \frac{d}{2}}\right), \quad n \rightarrow \infty.$$

(ii) Assume that $v \in \mathcal{L}^1 \cap \mathcal{L}^2$, v satisfies Inequality (17) and v decays exponentially. Define $\tilde{v}_{\mathbf{k}}$ as in Equation (15). Then it holds that

$$\left| \int_{\mathbb{R}^d} v(\mathbf{x}) f(\mathbf{x}) d\mathbf{x} - \sum'_{\mathbf{0} \leq \mathbf{k} \leq \mathbf{N}} c_{\mathbf{k}} \tilde{v}_{\mathbf{k}} \right| \leq O\left(n^{-(1-\beta)(p - \frac{d}{2})}\right), \quad n \rightarrow \infty.$$

Proof. Let $A_1(\mathbf{L})$, $A_2(\mathbf{L}, \mathbf{N})$, $D_1(\mathbf{M})$ and $D_2(\mathbf{N}, \mathbf{L}, \mathbf{M})$ be as in the proof of Corollary 3.6. Since $v_{\mathbf{k}} = \langle v1_{[-\mathbf{M}, \mathbf{M}]}, e_{\mathbf{k}}^{\mathbf{L}} \rangle$ and similarly to the proof of Corollary 3.6 we have that

$$\begin{aligned} & \left| \int_{\mathbb{R}^d} v(\mathbf{x}) f(\mathbf{x}) d\mathbf{x} - \sum'_{\mathbf{0} \leq \mathbf{k} \leq \mathbf{N}} c_{\mathbf{k}} \tilde{v}_{\mathbf{k}} \right| \\ & \leq D_1(\mathbf{M}) + \|v1_{[-\mathbf{M}, \mathbf{M}]\|_2 \left(A_1(\mathbf{L}) + A_2(\mathbf{L}, \mathbf{N}) + \sqrt{B_f(\mathbf{L})} \right) + D_2(\mathbf{N}, \mathbf{L}, \mathbf{M}). \end{aligned} \quad (44)$$

We will analyze the order of convergence of each term at the right-hand side of Inequality (44): Since v is bounded and f decays exponentially, $D_1(\mathbf{M})$, $A_1(\mathbf{L})$ and $\sqrt{B_f(\mathbf{L})}$ decay exponentially, i.e., can be bounded by $O(\exp(-C_3 n^\beta))$, $n \rightarrow \infty$ for some C_3 , see proof of Theorem 3.9. By Inequality (40), the term $\sum' |c_{\mathbf{k}}|^2$ is bounded. If $w_{\mathbf{k}} = v_{\mathbf{k}}$ then $D_2(\mathbf{N}, \mathbf{L}, \mathbf{M}) = 0$. If $w_{\mathbf{k}} = \tilde{v}_{\mathbf{k}}$ then $D_2(\mathbf{N}, \mathbf{L}, \mathbf{M})$ decays exponentially, see proof of Corollary 3.10. Last, we treat $A_2(\mathbf{L}, \mathbf{N})$. Let $j \in \{1, \dots, d\}$. Let n be large enough. Let $\mathbf{k} \in \mathbb{N}_0^d$ such that $k_j > n$. By Equation (11) and Inequality (43), there is a constant $a_1 > 0$ so that

$$|c_{\mathbf{k}}|^2 \stackrel{(11)}{\leq} \left(\frac{1}{2^{d-1} \prod_{h=1}^d L_h} \sum_{\mathbf{s}=(1, \pm 1, \dots, \pm 1) \in \mathbb{R}^d} \left| \hat{f}\left(\frac{\pi \mathbf{s} \mathbf{k}}{2 \mathbf{L}}\right) \right| \right)^2 \stackrel{(43)}{\leq} a_1 n^{2\beta(p-d)} |\mathbf{k}|_\infty^{-2p}.$$

By mathematical induction over d and the applying the integral test of convergence, one can show that

$$\sum_{\mathbf{k} \in \mathbb{N}_0^d, k_j > n} |\mathbf{k}|_\infty^{-2p} \leq \frac{2^{d-1}}{(2p-d)n^{2p-d}}. \quad (45)$$

It follows by Inequality (45) for some $a_2 > 0$ that

$$\prod_{h=1}^d L_h \sum_{\mathbf{k} \in \mathbb{N}_0^d, k_j > n} |c_{\mathbf{k}}|^2 \leq a_2 n^{-(1-\beta)(2p-d)}. \quad (46)$$

Let $G(\mathbf{L})$ be defined as in Equality (31). By Equality (22), the Cauchy-Schwarz (CS) inequality and Inequality (33), we obtain

$$\begin{aligned} A_2(\mathbf{L}, \mathbf{N})^2 & \stackrel{(22)}{\leq} \prod_{h=1}^d L_h \sum'_{k_1 > N_1 \text{ or } \dots \text{ or } k_d > N_d} |a_{\mathbf{k}} + c_{\mathbf{k}} - c_{\mathbf{k}}|^2 \\ & \stackrel{(CS)}{\leq} \prod_{h=1}^d L_h \sum'_{k_1 > N_1 \text{ or } \dots \text{ or } k_d > N_d} |c_{\mathbf{k}}|^2 + \prod_{h=1}^d L_h \sum'_{\mathbf{k} \in \mathbb{N}_0^d} |a_{\mathbf{k}} - c_{\mathbf{k}}|^2 \\ & \quad + 2 \sqrt{\prod_{h=1}^d L_h \sum'_{\mathbf{k} \in \mathbb{N}_0^d} |c_{\mathbf{k}}|^2 \prod_{h=1}^d L_h \sum'_{\mathbf{k} \in \mathbb{N}_0^d} |a_{\mathbf{k}} - c_{\mathbf{k}}|^2} \\ & \stackrel{(33)}{\leq} \sum_{j=1}^d \left(\prod_{h=1}^d L_h \sum_{\mathbf{k} \in \mathbb{N}_0^d, k_j > n} |c_{\mathbf{k}}|^2 \right) + B_f(\mathbf{L}) + 2 \sqrt{\left(\int_{\mathbb{R}^d} |f(\mathbf{x})|^2 d\mathbf{x} + G(\mathbf{L}) \right) B_f(\mathbf{L})} \\ & \stackrel{(46)}{\leq} O\left(n^{-(1-\beta)(2p-d)}\right), \quad n \rightarrow \infty, \end{aligned}$$

since $B_f(\mathbf{L})$ and $G(\mathbf{L})$, converge exponentially to zero. Since v is bounded, we have that $\|v1_{[-\mathbf{M}, \mathbf{M}]\|_2 \leq O\left(n^{\frac{d\beta}{2}}\right)$, $n \rightarrow \infty$. Noting that $\frac{-(1-\beta)(2p-d)+d\beta}{2} = -(1-\beta)p + \frac{d}{2}$, shows (i). It holds $\|v1_{[-\mathbf{M}, \mathbf{M}]\|_2 \leq \|v\|_2$ if $v \in \mathcal{L}_2$, which implies (ii). \square

4 Characteristic functions

In this section, in Examples 4.1 and 4.2, we recall the normal and the Variance Gamma distribution from the literature. Remarks 4.3, 4.4 and 4.5 provide a financial context.

Example 4.1. (Normal distribution). Let \mathbf{X} be a multivariate normal random variable with location $\boldsymbol{\eta} \in \mathbb{R}^d$ and covariance matrix $\Sigma \in \mathbb{R}^d \times \mathbb{R}^d$. The random variable \mathbf{X} has characteristic function $\widehat{g}(\mathbf{u}) = \exp(i\boldsymbol{\eta} \cdot \mathbf{u} - \frac{1}{2}\mathbf{u} \cdot \Sigma \mathbf{u})$, which can be extended to \mathbb{C}^d , i.e., $\widehat{g}(\mathbf{u} - i\boldsymbol{\alpha})$ exists for all $\boldsymbol{\alpha} \in \mathbb{R}^d$. By Proposition 3.1, we set $\lambda = \exp(-\boldsymbol{\eta} \cdot \boldsymbol{\alpha} - \frac{1}{2}\boldsymbol{\alpha} \cdot \Sigma \boldsymbol{\alpha})$ and $\boldsymbol{\mu} = \boldsymbol{\eta} + \Sigma \boldsymbol{\alpha}$. The characteristic function of the damped density f , defined in Equation (3), is given by $\widehat{f}(\mathbf{u}) = \exp(-\frac{1}{2}\mathbf{u} \cdot \Sigma \mathbf{u})$. A straightforward computation shows that

$$(2\pi)^{-d} \int_{\mathbb{R}^d} |\widehat{f}(\mathbf{u})|^2 d\mathbf{u} = \frac{2^{-d}}{\sqrt{\pi^d \det(\Sigma)}}.$$

Example 4.2. (Variance Gamma distribution). Let \mathbf{Z} be a d -dimensional, standard normal random variable. Let G be a Gamma distributed random variable, independent of \mathbf{Z} , with shape $a > 0$ and scale $s > 0$. Let $\boldsymbol{\eta}, \boldsymbol{\theta} \in \mathbb{R}^d$ and $\boldsymbol{\sigma} \in \mathbb{R}_+^d$. Consider $\mathbf{X} = \boldsymbol{\eta} + \boldsymbol{\theta}G + \sqrt{G}\boldsymbol{\sigma}\mathbf{Z}$. The distribution of \mathbf{X} is denoted by $\text{VG}(a, s, \boldsymbol{\eta}, \boldsymbol{\theta}, \boldsymbol{\sigma})$. Define $\Sigma \in \mathbb{R}^d \times \mathbb{R}^d$ such that $\Sigma_{ii} = \sigma_i^2$ and $\Sigma_{ij} = 0$ for $i \neq j$. Then \mathbf{X} has characteristic function

$$\widehat{g}(\mathbf{u}) = \exp(i\boldsymbol{\eta} \cdot \mathbf{u}) \left(1 - is\boldsymbol{\theta} \cdot \mathbf{u} + \frac{1}{2}s\mathbf{u} \cdot \Sigma \mathbf{u}\right)^{-a},$$

see Luciano and Schoutens (2006). The (extended) Fourier transform $\widehat{g}(\mathbf{u} - i\boldsymbol{\alpha})$ exists for all $\boldsymbol{\alpha} \in \mathbb{R}^d$ with $\zeta := 1 - s\boldsymbol{\theta} \cdot \boldsymbol{\alpha} - \frac{1}{2}s\boldsymbol{\alpha} \cdot \Sigma \boldsymbol{\alpha} > 0$, see Bayer et al. (2022). By Proposition 3.1, we set $\lambda = \exp(-\boldsymbol{\eta} \cdot \boldsymbol{\alpha})\zeta^a$ and $\boldsymbol{\mu} = \boldsymbol{\eta} + as\zeta^{-1}(\boldsymbol{\theta} + \Sigma \boldsymbol{\alpha})$. The characteristic function of the damped density f , defined in Equation (3), is given by

$$\widehat{f}(\mathbf{u}) = \exp\left(-i\frac{as}{\zeta}(\boldsymbol{\theta} + \Sigma \boldsymbol{\alpha}) \cdot \mathbf{u}\right) \left(1 - i\frac{s}{\zeta}(\boldsymbol{\theta} + \Sigma \boldsymbol{\alpha}) \cdot \mathbf{u} + \frac{1}{2}\frac{s}{\zeta}\mathbf{u} \cdot \Sigma \mathbf{u}\right)^{-a}.$$

Apply the Courant–Fischer–Weyl min-max principle to see that $|\widehat{f}(\mathbf{u})| \leq O(|\mathbf{u}|_\infty^{-2a})$ for $|\mathbf{u}|_\infty \rightarrow \infty$.

Remark 4.3. In a financial context, we model d stock prices over time by a d -dimensional positive semi-martingale $(\mathbf{S}(t))_{t \geq 0}$ on a filtered probability space $(\Omega, \mathcal{F}, P, (\mathcal{F}_t)_{t \geq 0})$. The filtration $(\mathcal{F}_t)_{t \geq 0}$ satisfies the usual conditions with $\mathcal{F}_0 = \{\Omega, \emptyset\}$. The logarithmic returns are defined by $\mathbf{X}(t) := \log(\mathbf{S}(t))$, $t \geq 0$. There is a bank account paying continuous compound interest $r \in \mathbb{R}$. There is a European rainbow option $w : \mathbb{R}^d \rightarrow \mathbb{R}$ with maturity $T > 0$ and payoff $w(\mathbf{X}(T))$ at time T . We denote by g the (risk-neutral) density of $\log(\mathbf{S}(T))$. The time-0 price of the European option with payoff w is then given by $e^{-rT} \int_{\mathbb{R}^d} w(\mathbf{x})g(\mathbf{x})d\mathbf{x}$. This integral can be approximated by the (damped) COS method.

Remark 4.4. Let $\Sigma \in \mathbb{R}^d \times \mathbb{R}^d$ be a symmetric positive definite matrix. For the Black-Scholes (BS) model, see Karatzas and Shreve (1998), the logarithmic returns $\mathbf{X}(T)$ are normally distributed with location $\boldsymbol{\eta} := \log(\mathbf{S}(0)) + (\mathbf{r} - \frac{1}{2}\text{diag}(\Sigma))T$ and covariance matrix $T\Sigma$, where $\mathbf{r} = (r, \dots, r) \in \mathbb{R}^d$ and $\text{diag}(\Sigma) \in \mathbb{R}^d$ denotes the diagonal of Σ .

Remark 4.5. Let $\nu > 0$, $\boldsymbol{\sigma} \in \mathbb{R}_+^d$ and $\boldsymbol{\theta} \in \mathbb{R}^d$. In the multivariate Variance Gamma (VG) model, see Luciano and Schoutens (2006), the logarithmic returns $\mathbf{X}(T)$ follow a $\text{VG}(\frac{T}{\nu}, \nu, \boldsymbol{\eta}, \boldsymbol{\theta}, \boldsymbol{\sigma})$ distribution, where

$$\eta_h := \log(S_h(0)) + \left(r + \frac{1}{\nu} \log\left(1 - \frac{1}{2}\sigma_h^2\nu - \theta_h\nu\right)\right)T, \quad h = 1, \dots, d.$$

5 Functions of interest

Example 5.1. (CDF). Let $w(\mathbf{x}) = 1_{(-\infty, \mathbf{y}]}(\mathbf{x})$, $\mathbf{x} \in \mathbb{R}^d$ for some $\mathbf{y} \in \mathbb{R}^d$. The integral in (1) is equal to the CDF of the density g evaluated at \mathbf{y} . The coefficients $v_{\mathbf{k}}$, defined in Equation (14), can be obtained in

closed-form if $\alpha = \mathbf{0}$. Let $\mathbf{M}, \mathbf{L} \in \mathbb{R}_+^d$ as in Section 3. Let $\gamma_h := \min(y_h - \mu_h, M_h)$, $h = 1, \dots, d$. It holds for $\mathbf{k} \in \mathbb{N}_0^d$ that $v_{\mathbf{k}} = 0$ if $\gamma_h < -M_h$ for any h and otherwise

$$v_{\mathbf{k}} = \lambda^{-1} \prod_{\substack{h=1 \\ k_h=0}}^d \{\gamma_h + M_h\} \prod_{\substack{h=1 \\ k_h>0}}^d \left\{ \frac{2L_h}{\pi k_h} \left(\sin\left(k_h \pi \frac{\gamma_h + L_h}{2L_h}\right) - \sin\left(k_h \pi \frac{-M_h + L_h}{2L_h}\right) \right) \right\}.$$

Next, we assume for some $\alpha \in \mathbb{R}^d$ that the map $\mathbf{x} \mapsto w(\mathbf{x})e^{-\alpha \cdot \mathbf{x}}$ is integrable. The Fourier-transform of w then exists at all points $\mathbf{u} + i\alpha \in \mathbb{C}^d$, $\mathbf{u} \in \mathbb{R}^d$. Let $\lambda > 0$ and $\mu \in \mathbb{R}^d$. Let v be as in Equation (4). The Fourier-transform of v is given by $\hat{v}(\mathbf{u}) = \lambda^{-1} e^{-i\mathbf{u} \cdot \mu} \hat{w}(\mathbf{u} + i\alpha)$. Hence, a closed-form expression for \hat{w} is sufficient to obtain a closed-form expression for \hat{v} . We can then directly obtain $\hat{v}_{\mathbf{k}}$, defined in Equation (15) via Equation (16). For many functions of interest, \hat{w} is known in closed-form in d dimensions. We provide some examples from finance, where integral in (1) is then interpreted as a price: Digital cash-or-nothing put options and arithmetic basket options are discussed in Examples 5.1 and 5.3, respectively. For put and call options on the maximum or minimum of d assets, see Eberlein et al. (2010); for spread options, see Hurd and Zhou (2010). Note that a digital cash-or-nothing put option with strike \mathbf{K} is equal to the CDF of g evaluated at $\log(\mathbf{K})$. We discuss Example 5.2 to test the damped COS method for $\alpha \neq \mathbf{0}$.

Example 5.2. (Digital cash-or-nothing put option). The payoff function of a cash-or-nothing put option is defined by $w(\mathbf{x}) = 1_{[0, \mathbf{K}]}(e^{\mathbf{x}})$, $\mathbf{x} \in \mathbb{R}^d$ for some strikes $\mathbf{K} \in \mathbb{R}_+^d$. The option pays 1\$ at maturity if $\mathbf{S}(T) \leq \mathbf{K}$ and nothing otherwise. The integral $\int w(\mathbf{x})g(\mathbf{x})d\mathbf{x}$ is equal to $G(\log(\mathbf{K}))$, where G is the CDF of g . A simple calculation shows that the Fourier-transform of w exists for $\mathbf{z} \in \mathbb{C}^d$ such that $\Im\{z_h\} < 0$ and is given by $\hat{w}(\mathbf{z}) = \prod_{h=1}^d \frac{K_h^{iz_h}}{iz_h}$. For $\lambda > 0$ and $\mu \in \mathbb{R}^d$, let v be as in Equation (4). It holds for $\alpha < \mathbf{0}$ that $\|v\|_{\infty} \leq \lambda^{-1} e^{-\alpha \cdot \log(\mathbf{K})}$ and

$$\|v\|_2^2 = \lambda^{-2} \prod_{h=1}^d \frac{\exp(-2\alpha_h(\log(K_h)))}{-2\alpha_h}.$$

Example 5.3. (Arithmetic basket put option). The payoff function of an arithmetic basket put option is defined by $w(\mathbf{x}) = \max(K - \sum_{h=1}^d e^{x_h}, 0)$, $\mathbf{x} \in \mathbb{R}^d$ for some strike $K > 0$. The Fourier-transform of w exists for $\mathbf{z} \in \mathbb{C}^d$ such that $\Im\{z_h\} < 0$ and is given by

$$\hat{w}(\mathbf{z}) = \int_{\mathbb{R}^d} e^{i\mathbf{z} \cdot \mathbf{x}} w(\mathbf{x}) d\mathbf{x} = \frac{K^{(1+i\sum_{h=1}^d z_h)} \prod_{h=1}^d \Gamma(iz_h)}{\Gamma\left(i\sum_{h=1}^d z_h + 2\right)}. \quad (47)$$

Equation (47) follows by an elementary substitution² from Olver et al. (2010, Eq. (5.14.1)) and is also mentioned in a similar form in Hubalek and Kallsen (2003). If $\alpha < \mathbf{0}$ it holds $\|v\|_{\infty} \leq \lambda^{-1} K^{1-\sum_{h=1}^d \alpha_h}$ and, using Olver et al. (2010, Eq. (5.14.1)), it holds that

$$\|v\|_2^2 \leq \frac{K^{2-2\sum_{h=1}^d \alpha_h}}{\lambda^2} \frac{\prod_{h=1}^d \Gamma(-2\alpha_h)}{\Gamma(1 + \sum_{h=1}^d (-2\alpha_h))}.$$

6 Numerical experiments

We provide several numerical experiments to solve the integral in (1) by the COS method. Reference values are obtained by Eberlein et al. (2010, Theorem 3.2) with damping factor $\mathbf{R} = (-4, \dots, -4)$, who express the integral in (1) by another integral involving the Fourier-transforms \hat{g} and \hat{w} . To obtain reference values and to solve the integral in Corollary 3.11, we use the command *cubintegrate* with the method *cuhre* from the R-package *cubature* with relative tolerance 10^{-11} . We confirm all reference values using the COS method with $\mathbf{N} = (2000, \dots, 2000)$ and a truncation range obtained from Equation (35) with $\varepsilon = 10^{-10}$ using $n = 8$

²We thank Friedrich Hubalek from TU Wien for pointing this out to us.

moments. For the normal distribution, reference values are also given in closed-form in the uncorrelated case for a CDF and the digital cash-or-nothing put option. All experiments are performed on a laptop with Intel i7-11850H processor and 32 GB RAM. The COS method and Monte Carlo simulations are implemented in C++ using for-loops without parallelization. The memory requirements are minimal.

We first investigate the influence of the damping factor α on the accuracy of the COS method to obtain the price of a cash-or-nothing put option with strike $\mathbf{K} \in \mathbb{R}_+^d$ in the BS model, which is just the CDF of a normal distribution evaluated at $\log(\mathbf{K})$, hence reference values can be obtained in close-form. Figure 1 shows the behavior of the COS method for different damping factors in dimensions $d \in \{2, 3, 4\}$. If α is too close to zero, almost no damping takes place and the difference between $v_{\mathbf{k}}$ and $\tilde{v}_{\mathbf{k}}$ is large, which implies a relative high error of the COS method. If $|\alpha|$ is too big, $\|v\|_\infty$ and $\|v\|_2$ become very large and the truncation error increases. However, we observe in the example that a wide range of damping factor α work well in various dimensions. Further, fixing the number of terms N and the truncation range L , the accuracy of the classical COS method with $\alpha = 0$ and the damped COS method with $\alpha \neq 0$ is very similar for some damping factors.

We illustrate the order of convergence of the COS method for an arithmetic basket put option in the VG model. We compare three different maturities. In Figure 1 we can see that the theoretical bound from Theorem 3.13 for the order of convergence is sharp and close the empirical order of convergence.




Figure 1: *Left*: Error of the price of a cash-or-nothing put option in the BS model for different damping factors with $\mathbf{K} = \mathbf{S}(0) = (100, \dots, 100)$, $T = 1$, $r = 0$ and $\Sigma_{ii} = \sigma^2$, $\Sigma_{ij} = 0$, $i \neq j$, where $\sigma = 0.2$. Further, $\mathbf{M} = \mathbf{L} = (20\sigma, \dots, 20\sigma)$ and $\mathbf{N} = (70, \dots, 70)$. Reference values are obtained by the closed-form solution. *Right*: Logarithmic error of the price by the COS method for the VG model over the logarithmic number of terms for an arithmetic basket put option and $d = 2$. We choose $\mathbf{N} = (n, n)$ and $\mathbf{M} = \mathbf{L} = (\frac{1}{2}n^\beta, \frac{1}{2}n^\beta)$ with $\beta = \frac{1}{2}$. We set $\mathbf{S}(0) = (100, 100)$, $K = 200$, $\sigma = (0.2, 0.2)$, $\theta = (-0.03, -0.03)$, $\nu = 0.1$, $r = 0$ and $\alpha = (-4, -4)$. The theoretical bound from Theorem 3.13, i.e., a line with slope $-(1-\beta)(p-\frac{d}{2})$, is shown in gray. For the VG model, we have $p = \frac{2T}{\nu}$. Reference values are obtained by Eberlein et al. (2010).

6.1 Other methods

We compare the COS method with a Monte Carlo (MC) simulation to obtain the price of a cash-or-nothing put option in the BS model, which is equal to the CDF of the normal distribution evaluated at $\log(\mathbf{K})$, hence reference values can be obtained in close-form. The computational complexity of a MC simulation with $U \in \mathbb{N}$ runs scales like $O(Ud)$. We estimate U by the central limit theorem and a statistical error of 0.99. The COS method consist of d -nested sums. According to Equation (11), the computational complexity of the COS method scales like $O\left(\prod_{h=1}^d \{N_h\} 2^{d-1}\right)$. A MC simulation converges relative slowly but hardly

depends on the dimension. On the other hand, the complexity of the COS method grows exponentially in the dimension, however, the COS method also converges exponentially for the BS model. The choice between MC and the COS method depends both on the dimension and the error tolerance ε : the higher d the faster MC compared to the COS method but the smaller ε , the better performs the COS method. In Table 1 and Figure 2, we observe that the COS method is faster than MC for $d \leq 3$ and $\varepsilon \leq 10^{-2}$. For $d = 4$ ($d = 5$), the COS method outperforms MC for $\varepsilon \leq 10^{-3}$ ($\varepsilon \leq 10^{-5}$), otherwise a MC simulation is faster. If $\varepsilon = 10^{-9}$ and $d = 4$, the COS method needs 220 number of terms in each dimensions to stay below the error tolerance and the CPU time is about one hour. We estimate that a MC simulation would need more than 20,000 years.

d	N	L	U	τ_{COS}	τ_{MC}	Reference value
1	30	2.0	16481995016	8.9e-6	5.1e+3	0.539827
2	30	2.4	13700525367	3.7e-4	1.6e+4	0.291414
3	40	3.0	8795611829	4.9e-2	1.2e+4	0.157313
4	50	3.6	5156004587	1.1e+1	8.1e+3	0.084922
5	50	4.2	2902219256	1.4e+3	7.8e+3	0.045843

Table 1: CPU time of the COS method (τ_{COS}) and CPU time of a MC simulation (τ_{MC}) for the BS model to price a cash-or-nothing put option. We set $\varepsilon = 10^{-5}$, $\alpha = (-7, \dots, -7)$, $\Sigma_{ii} = \sigma^2$, $\Sigma_{ij} = 0$, $i \neq j$, where $\sigma = 0.2$, and $S(0) = K = (100, \dots, 100)$. We set $N_1 = \dots = N_d = N$. We obtain the truncation range $L = (L, \dots, L)$ from Inequality (35) using $n = 8$ moments. The reference value can be obtained in closed-form. CPU time is measured in seconds.




Figure 2: Ratio of the CPU time of the COS method (τ_{COS}) and the CPU time of a MC simulation (τ_{MC}) for the BS model to price a cash-or-nothing put option. Parameters are as in Table 1.

6.2 On the choice of N

In this section, we consider an arithmetic basket put option in the BS and the VG models. The methodology can also be applied to other rainbow options or a CDF. We compare different strategies to choose the number of terms N . For $d = 1$ we also consider the bound for N from Junike (2024), which can be obtained as follows:

If the (damped) density f is $J + 1$ times differentiable with bounded derivatives, the number of terms can be chosen by

$$N \geq \left(\frac{2^{s+\frac{5}{2}} \|f^{(s+1)}\|_{\infty} L^{s+2}}{s\pi^{s+1}} \frac{12Ke^{-rT}}{\varepsilon} \right)^{\frac{1}{s}}, \quad (48)$$

where $s \in \{1, \dots, J\}$. The term $\|f^{(s+1)}\|_{\infty}$ can be bounded by

$$\|f^{(s+1)}\|_{\infty} \leq \frac{1}{2\pi} \int_{\mathbb{R}} |u|^{s+1} |\varphi(u)| du. \quad (49)$$

For the BS model, we choose $s = 40$. According to Küchler and Tappe (2008), the density of the VG model is $J + 1$ times continuously differentiable if J is equal to the largest natural number which is less than $\frac{2T}{\nu} - 2$. For the VG model, we use $s = J$.

In Table 2 one can see that Corollary 3.11 provides a sharper bound for \mathbf{N} than Junike (2024). This is particularly noticeable for the VG model, which is less smooth than the BS model. However, the formula in Junike (2024) is more stable, compare with Remark 3.12. The number of terms obtained by Corollary 3.11 is at most three times larger than the minimal number of terms necessary to stay below the error tolerance.

	d	Model	N	CPU time COS (num. int.)	Parameters	\mathbf{L}	Ref. value
Minimal N	1	BS	25	0.02	$\sigma = 0.2$	1.8	7.965567
Cor. 3.11	1	BS	28	0.03			
Junike (2024)	1	BS	34	0.03			
Minimal N	1	VG	95	0.12	$\sigma = 0.1213,$ $\theta = -0.1436,$ $\nu = 0.1686$	5.5	5.195700
Cor. 3.11	1	VG	264	0.28 (1.95)			
Junike (2024)	1	VG	729	0.96 (12.23)			
Minimal \mathbf{N}	2	BS	65	5.35	$\Sigma_{11} = 0.2^2, \Sigma_{22} = 0.4^2$ $\Sigma_{12} = \Sigma_{21} = \frac{1}{2}\sqrt{\Sigma_{11}\Sigma_{22}}$	(5.7, 11.5)	21.010354
Cor. 3.11	2	BS	116	17.02			
Minimal \mathbf{N}	2	VG	55	8.55	$\sigma = (0.2, 0.25), \nu = 0.1$ $\theta = (-0.03, -0.05),$	(5.8, 7.5)	12.670179
Cor. 3.11	2	VG	154	68.20 (1092.2)			

Table 2: Comparison of different strategies to choose \mathbf{N} for an arithmetic basket put option in the BS model and the VG model. We use the error tolerance $\varepsilon = 10^{-3}$. We set $\mathbf{N} = (N, \dots, N)$, $\mathbf{S}(0) = (100, \dots, 100)$, $K = 100d$, $\boldsymbol{\alpha} = (-4, \dots, -4)$, $T = 1$ and $r = 0$. We obtain the truncation range $\mathbf{L} = (L, \dots, L)$ from Inequality (35) using $n = 8$ moments. Reference values are obtained by Eberlein et al. (2010). We average over ten runs to obtain the CPU time, which is measured in milliseconds.

7 Conclusions

In this article we introduced and discussed the damped COS method, which is a numerical tool to solve certain multidimensional integrals numerically, e.g., to obtain a CDF from a characteristic function or to price rainbow options in a financial context. The (damped) COS method requires several parameters: In particular, one has to specify a truncation range \mathbf{L} for the density f , a truncation range \mathbf{M} for the integral and the number of terms \mathbf{N} of cosine functions to approximate the truncated density. Corollary 3.6 provides sufficient conditions on \mathbf{M} , \mathbf{L} and \mathbf{N} to ensure the convergence of the COS method within a given error tolerance $\varepsilon > 0$. Theorem 3.9 and Corollary 3.11 provide formulas for the truncation ranges \mathbf{M} and \mathbf{L} and the number of terms \mathbf{N} , respectively. Theorem 3.13 provides an upper bound of the order of convergence of the COS method. Numerical experiments indicate that the bound is sharp. In particular, the (damped) COS method converges exponentially if the Fourier transform \hat{f} decays exponentially.

References

Joseph Abate and Ward Whitt. The fourier-series method for inverting transforms of probability distributions. *Queueing systems*, 10:5–87, 1992.

- M. Abramowitz and I. A. Stegun. *Handbook of mathematical functions with formulas, graphs, and mathematical tables*, volume 55. US Government printing office, 1972.
- H. Amann and J. Escher. *Analysis III*. Birkhäuser Verlag AG, Springer, 2009.
- C. Bardgett, E. Gourier, and M. Leippold. Inferring volatility dynamics and risk premia from the S&P 500 and VIX markets. *Journal of Financial Economics*, 131(3):593–618, 2019.
- Heinz Bauer. *Maß-und Integrationstheorie*. Walter de Gruyter, 1992.
- Heinz Bauer. *Probability Theory*, volume 23. Walter de Gruyter, 1996.
- Christian Bayer, Chiheb Ben Hammouda, Antonis Papapantoleon, Michael Samet, and Raúl Tempone. Optimal damping with hierarchical adaptive quadrature for efficient fourier pricing of multi-asset options in Levy models. *arXiv preprint arXiv:2203.08196*, 2022.
- P. Carr and D. Madan. Option valuation using the fast Fourier transform. *Journal of Computational Finance*, 2(4):61–73, 1999.
- Darrell Duffie, Damir Filipović, and Walter Schachermayer. Affine processes and applications in finance. *The Annals of Applied Probability*, 13(3):984–1053, 2003.
- Ernst Eberlein, Kathrin Glau, and Antonis Papapantoleon. Analysis of fourier transform valuation formulas and applications. *Applied Mathematical Finance*, 17(3):211–240, 2010.
- F. Fang and C. W. Oosterlee. A novel pricing method for European options based on Fourier-cosine series expansions. *SIAM Journal on Scientific Computing*, 31(2):826–848, 2009a.
- F. Fang and C. W. Oosterlee. Pricing early-exercise and discrete barrier options by Fourier-cosine series expansions. *Numerische Mathematik*, 114(1):27, 2009b.
- F. Fang and C. W. Oosterlee. A Fourier-based valuation method for Bermudan and barrier options under Heston’s model. *SIAM Journal on Financial Mathematics*, 2(1):439–463, 2011.
- J Gil-Pelaez. Note on the inversion theorem. *Biometrika*, 38(3-4):481–482, 1951.
- L. A. Grzelak and C. W. Oosterlee. On the Heston model with stochastic interest rates. *SIAM Journal on Financial Mathematics*, 2(1):255–286, 2011.
- Friedrich Hubalek and Jan Kallsen. Variance-optimal hedging and markowitz-efficient portfolios for multivariate processes with stationary independent increments with and without constraints. *TU München*, 2003.
- Paul Hughett. Error bounds for numerical inversion of a probability characteristic function. *SIAM journal on numerical analysis*, 35(4):1368–1392, 1998.
- Thomas R Hurd and Zhuowei Zhou. A fourier transform method for spread option pricing. *SIAM Journal on Financial Mathematics*, 1(1):142–157, 2010.
- G. Junike and K. Pankrashkin. Precise option pricing by the COS method—How to choose the truncation range. *Applied Mathematics and Computation*, 421:126935, 2022.
- Gero Junike. On the number of terms in the cos method for european option pricing. *Numerische Mathematik*, 2024.
- I. Karatzas and S. E. Shreve. *Methods of Mathematical finance*, volume 39. Springer, 1998.
- U. Küchler and S. Tappe. On the shapes of bilateral Gamma densities. *Statistics & Probability Letters*, 78(15):2478–2484, 2008.
- Á. Leitaó, C. W. Oosterlee, L. Ortiz-Gracia, and S. M. Bohte. On the data-driven COS method. *Applied Mathematics and Computation*, 317:68–84, 2018.

- S. Liu, A. Borovykh, L. A. Grzelak, and C. W. Oosterlee. A neural network-based framework for financial model calibration. *Journal of Mathematics in Industry*, 9:1–28, 2019a.
- S. Liu, C. W. Oosterlee, and S. M. Bohte. Pricing options and computing implied volatilities using neural networks. *Risks*, 7(1):16, 2019b.
- R. Lord, F. Fang, F. Bervoets, and C. W. Oosterlee. A fast and accurate FFT-based method for pricing early-exercise options under Lévy processes. *SIAM Journal on Scientific Computing*, 30(4):1678–1705, 2008.
- Elisa Luciano and Wim Schoutens. A multivariate jump-driven financial asset model. *Quantitative finance*, 6(5):385–402, 2006.
- Frank W. J. Olver, Daniel W. Lozier, Ronald F. Boisvert, and Charles W. Clark. *NIST handbook of mathematical functions*. Cambridge university press, 2010.
- C. W. Oosterlee and L. A. Grzelak. *Mathematical modeling and computation in finance: with exercises and Python and MATLAB computer codes*. World Scientific, 2019.
- L. Ortiz-Gracia and C. W. Oosterlee. Robust pricing of European options with wavelets and the characteristic function. *SIAM Journal on Scientific Computing*, 35(5):B1055–B1084, 2013.
- L. Ortiz-Gracia and C. W. Oosterlee. A highly efficient Shannon wavelet inverse Fourier technique for pricing European options. *SIAM Journal on Scientific Computing*, 38(1):B118–B143, 2016.
- M. J. Ruijter and C. W. Oosterlee. Two-dimensional Fourier cosine series expansion method for pricing financial options. *SIAM Journal on Scientific Computing*, 34(5):B642–B671, 2012.
- Benno Schorr. Numerical inversion of a class of characteristic functions. *BIT Numerical Mathematics*, 15(1):94–102, 1975.
- Neil G Shephard. From characteristic function to distribution function: a simple framework for the theory. *Econometric theory*, 7(4):519–529, 1991a.
- Neil G Shephard. Numerical integration rules for multivariate inversions. *Journal of Statistical Computation and Simulation*, 39(1-2):37–46, 1991b.
- Lance A Waller, Bruce W Turnbull, and J Michael Hardin. Obtaining distribution functions by numerical inversion of characteristic functions with applications. *The American Statistician*, 49(4):346–350, 1995.
- B. Zhang and C. W. Oosterlee. Efficient pricing of European-style Asian options under exponential Lévy processes based on Fourier cosine expansions. *SIAM Journal on Financial Mathematics*, 4(1):399–426, 2013.