

ONE-STEP SMOOTHING SPLINES INSTRUMENTAL REGRESSION

JAD BEYHUM[†], ELIA LAPENTA[‡] AND PASCAL LAVERGNE^{*}

ABSTRACT. We extend nonparametric regression smoothing splines to a context where there is endogeneity and instrumental variables are available. Unlike popular existing estimators, the resulting estimator is one-step and relies on a unique regularization parameter. We derive uniform rates of the convergence for the estimator and its first derivative. We also address the issue of imposing monotonicity in estimation and extend the approach to a partly linear model. Simulations confirm the good performances of our estimator compared to two-step procedures. Our method yields economically sensible results when used to estimate Engel curves.

1. INTRODUCTION

We consider the prototypical model

$$(1.1) \quad Y = g_0(Z) + \varepsilon \quad E[\varepsilon|W] = 0,$$

where $Y \in \mathbb{R}$ is the dependent variable, $Z \in \mathbb{R}$ is the endogenous continuous explanatory variable, and $W \in \mathbb{R}^p$ are the instrumental variables (IVs). The goal is to estimate nonparametrically g_0 , the causal effect of the variable Z on Y , using W to account for endogeneity. If we assumed linear relationships, we could use the two-stage least squares estimator: in a first stage, one obtains the linear projection of Z given W , then in a second stage one linearly regresses Y onto the previously estimated linear projection. Considering a nonparametric function g_0 allows estimating the causal relationship of Y and Z in a more flexible manner.

Date: April 2024.

Jad Beyhum undertook most of this work while employed by CREST, ENSAI (Rennes). He is grateful for support from the Research Fund KU Leuven under grant STG/23/014. Elia Lapenta acknowledges funding from the French National Research Agency (ANR) under grant ANR-23-CE26-0008-01. Pascal Lavergne acknowledges funding from the French National Research Agency (ANR) under the Investments for the Future program (Investissements d'Avenir, grant ANR-17-EURE-0010).

The authors thank Xavier D'Haultfoeuille, Ben Deaner, Frank Windmeijer, Daniel Wilhelm, Aureo de Paula, Kaspar Wuthrich, Christoph Breunig, Joachim Freyberger, Elie Tamer, seminar and conference participants at the University of Bonn, the 2023 Encounters in Econometric Theory at Nuffield College, Oxford University, the 2023 Bristol Econometrics Study Group conference, and the 34th (EC)² conference in Manchester.

Existing nonparametric estimators of g_0 typically rely on two steps. [Newey and Powell \(2003\)](#) develop a nonparametric equivalent to the two-stage least squares estimator: they use linear-in-parameter series expansions of $E[Y|W]$ and $E[g(Z)|W]$ in a generalized method of moments framework, see also [Ai and Chen \(2003\)](#), [Hall and Horowitz \(2005\)](#), [Blundell et al. \(2007\)](#), [Johannes et al. \(2011\)](#), [Horowitz \(2014\)](#) for other series-based methods. Alternatively, [Florens \(2003\)](#), [Hall and Horowitz \(2005\)](#), [Darolles et al. \(2011\)](#), and [Gagliardini and Scaillet \(2012\)](#) rely on kernel methods to estimate the unknown conditional expectations. As is well-known, backing up a nonparametric estimate of g_0 is an ill-posed inverse problem. Hence, one needs some kind of regularization, such as hard thresholding, see [Horowitz \(2011\)](#), [Chen and Pouzo \(2012\)](#), Tikhonov or ridge-type regularization, see [Newey and Powell \(2003\)](#), [Darolles et al. \(2011\)](#), [Florens et al. \(2011\)](#), [Gagliardini and Scaillet \(2012\)](#), [Singh et al. \(2019\)](#), or a Landweber-type iterative method, see [Johannes et al. \(2013\)](#), [Dunker et al. \(2014\)](#). A general exposition of some of these methods is given by [Carrasco et al. \(2007\)](#). A recent machine learning literature considers solving a saddle point problem that is dual to a generalized method of moments criterion. Here one first maximizes an objective function with respect to a function of the instruments W , then one minimizes with respect to a function of Z to obtain g_0 , see, e.g., [Bennett et al. \(2019\)](#). If the set of functions upon which one optimizes is large, then one has in addition to introduce some penalization in the optimization problem, see [Dikkala et al. \(2020\)](#), [Liao et al. \(2020\)](#). [Muandet et al. \(2020\)](#) consider a related but different saddle point problem.

We here develop a smoothing splines instrumental regression estimator for g_0 that fully avoids nonparametric first stage estimation. Our estimator has several characteristics that should make it appealing for empirical work. First, our approach is particularly attractive because it is one-step. Two-step procedures typically lead to theoretical and practical issues: one may need to estimate in a first step an object that may be more complex than the final object of interest; and first-stage estimation typically affects the second-stage small sample and asymptotic properties. Second, a key benefit of the one-step nature of our estimator is that it depends upon one regularization parameter only. In existing two-step methods, each stage relies on a particular choice of a smoothing or regularization parameter, whose fine-tuning may be difficult in practice, while affecting the final results. In some methods, a third parameter is introduced to deal with the ill-posed nature of the inverse problem. To choose our unique regularization parameter, we devise a practical cross-validation method that yields good performances in simulations. Third, by contrast to previous approaches based on series or kernel methods, our estimator is a natural generalization of the popular smoothing splines estimator, see [Wahba \(1990\)](#), [Green and Silverman \(1993\)](#). The appeal of splines lies in their simplicity together with their excellent

approximations of smooth functions, see [Schumaker \(2007\)](#). Splines-based methods have been extensively studied, see, e.g., [Hall and Opsomer \(2005\)](#), [Li and Ruppert \(2008\)](#), [Claeskens et al. \(2009\)](#), [Schwarz and Krivobokova \(2016\)](#), and have been found to have excellent performances in practice. Fourth, due to its spline nature, our estimator is computationally simple, and a closed-form expression is easily obtained for the spline coefficients. Fifth, as an additional advantage, one obtains straightforward estimators of derivatives.

We also propose some extensions to our method. First, we show how to impose monotonicity constraints by relying on a method proposed by [Hall and Huang \(2001\)](#). The constrained estimator is simple to implement in practice. Second, to illustrate the versatility of our method, we extend our results to a partly linear model. When used for estimating Engel curves, our smoothing splines estimator and its monotone constrained version yield comparable results, that are reasonable from an economic viewpoint.

The paper is organized as follows. In [Section 2](#), we detail the formulation of our estimator. We exhibit a global quantity that accounts for all the information contained in [Model \(1.1\)](#), and that is minimized by the true function g_0 . We then consider an empirical equivalent, and we set up a minimization problem penalized by a roughness measure of the function to regularize the solution. We show that our estimator extends smoothing splines to the instrumental variables context, and we give a closed form formula for its computation. The asymptotic properties of our estimator are analyzed in [Section 3](#), where uniform rates of convergences are derived for the function itself and its derivative. [Section 4](#) deals with estimation under monotonicity constraints. In [Section 5](#), we report selected simulation results, where our estimator exhibits excellent finite sample performance compared to some existing two-step methods, and we illustrate our method for Engel curves estimation. Finally, we extend our estimator to the partly linear model in [Section 6](#). Concluding remarks are given in [Section 7](#). Details and supplementary results of simulations, as well as the proof of subsidiary results are included in the online appendix.

2. OUR ESTIMATOR

2.1. General formulation. We assume that g_0 belong to some space of functions $\mathcal{G}$ on which identification holds, that is,

$$(2.2) \quad E[Y - g(Z)|W] = 0 \quad \text{a.s.} \Rightarrow g = g_0 \quad \text{a.s.}$$

For a discussion of this condition called *completeness*, see e.g. [D'Haultfoeuille \(2011\)](#) and [Freyberger \(2017\)](#). When Z is continuous, as we assume here, W should typically have at least one continuous component for completeness to hold. Some of the instruments, however, could be discrete, and this will not affect further our exposition and reasoning.

Instead of dealing directly with (2.2), as done by most previous work, we consider an equivalent formulation that does not require estimating a conditional expectation given the instruments W . By the results of Bierens (1982),

$$(2.3) \quad E[Y - g(Z)|W] = 0 \quad \text{a.s.} \Leftrightarrow E[(Y - g(Z)) \exp(\mathbf{i}W^\top t)] = 0 \quad \forall t \in \mathbb{R}^p.$$

Consider now

$$(2.4) \quad g_0 = \arg \min_{g \in \mathcal{G}} M(g), \quad M(g) = \int |E[(Y - g(Z)) \exp(\mathbf{i}W^\top t)]|^2 d\mu(t),$$

where μ is a symmetric probability measure. Then it is straightforward to see that $M(g) \geq 0$ for all $g \in \mathcal{G}$, and that under (2.2)

$$M(g) = 0 \Leftrightarrow g = g_0 \quad \text{a.s.}$$

With a random sample $\{(Y_i, Z_i, W_i), i = 1, \dots, n\}$ at hand, a natural estimator of $M(g)$ is

$$(2.5) \quad \begin{aligned} M_n(g) &= \int \left| \frac{1}{n} \sum_{i=1}^n (Y_i - g(Z_i)) \exp(\mathbf{i}W_i^\top t) \right|^2 d\mu(t) \\ &= \frac{1}{n^2} \sum_{1 \leq i, j \leq n} (Y_i - g(Z_i)) (Y_j - g(Z_j)) \omega(W_i - W_j), \end{aligned}$$

where

$$\omega(z) = \int_{\mathbb{R}^p} \exp(it'z) d\mu(t) = \int_{\mathbb{R}^p} \cos(t'z) d\mu(t),$$

due to the symmetry of μ : ω is (up to a constant) the Fourier transform of the density of μ . The above formulation as a V-statistic will be used in practice for computational purposes. This statistic accounts for an infinity of moment conditions, as stated in (2.3). It is different from generalized method of moments criteria used in previous work, that account for an increasing but finite number of moment conditions.

The condition for μ to have support $\mathbb{R}^p$ translates into the restriction that ω should have a strictly positive Fourier transform almost everywhere. Examples include products of triangular, normal, logistic, see Johnson et al. (1995, Section 23.3), Student, including Cauchy, see Dreier and Kotz (2002), or Laplace densities. To achieve scale invariance, we recommend, as in Bierens (1982), to scale the exogenous instruments by a measure of dispersion, such as their empirical standard deviation. Note that the function ω is *not* similar to a typical kernel used in nonparametric estimation, as there is no smoothing parameter entering ω , which is thus a fixed function that does not vary with the sample size. Hence, our estimation procedure introduces no smoothing on the instruments.

If W has bounded support, results from Bierens (1982) yield that the equivalence (2.3) holds when t is restricted to lie in a (arbitrary) neighborhood of 0 in $\mathbb{R}^p$. Hence, μ can be taken as any symmetric probability measure that contains 0 in the interior of its support.

As noted by [Bierens \(1982\)](#), there is no loss of generality assuming a bounded support, as his equivalence result equally applies to a one-to-one transformation of W , which can be chosen with bounded image.

Minimizing $M_n(g)$ would lead to interpolation. We regularize the problem by assuming some smoothness for the function g . We assume that Z has compact support, say $[0, 1]$ without loss of generality, and that $\mathcal{G}$ is the space of differentiable functions on $[0, 1]$ with absolutely continuous first derivative. That is, if $g \in \mathcal{G}$, there is an integrable function g'' such that $\int_0^z g''(t) dt = g'(z) - g'(0)$. We then estimate g_0 as a minimizer of a penalized version of $M_n(g)$ on $\mathcal{G}$. Specifically,

$$(2.6) \quad \hat{g} \in \arg \min_{g \in \mathcal{G}} M_n(g) + \lambda \int_0^1 |g''(z)|^2 dz,$$

where $\lambda > 0$ is a regularization parameter.

A recent approach we became aware of when preparing this paper is the “kernel maximum moment loss” approach proposed by [Zhang et al. \(2023\)](#). While it does not smooth on the instruments, it assumes that the regression of interest belongs to a Reproducing Kernel Hilbert Space (RKHS), and solves a minimization problem by penalizing by the norm on such a space. The estimator thus depends on the chosen RKHS. Differently, we assume that the regression of interest belongs to a space of smooth functions, and we penalize by the integral of the squared second derivative of the regression, which is a very intuitive measure of roughness, but not a RKHS norm.

2.2. Closed-form solution. We here show that (2.6) has a unique solution, a natural cubic spline, that we characterize in Proposition (2.1) below. We begin with some definitions. For $a < Z_1 < \dots < Z_n < b$, a function g on $[a, b]$ is a cubic spline if two conditions are satisfied: on each of the intervals $(a, Z_1), \dots, (Z_n, b)$, g is a cubic polynomial; the polynomial pieces fit together at each Z_i in such a way that g and its first and second derivatives are continuous. The points Z_i are called knots. A cubic spline on $[a, b]$ is said to be a *natural cubic spline* if its second and third derivatives are zero at a and b . Without loss of generality, we consider hereafter that $[a, b] = [0, 1]$. Given any values (g_i, Z_i) , $i = 1, \dots, n$, there is a unique interpolating natural cubic spline, that is, a natural cubic spline g with knots Z_i such that $g(Z_i) = g_i$, $i = 1, \dots, n$. For details, see e.g. [Green and Silverman \(1993\)](#). A key result for our analysis is the following.

Theorem 2.1 ([Green and Silverman \(1993, Th. 2.3\)](#)). *Suppose $n \geq 2$, $0 \leq Z_1 < \dots < Z_n \leq 1$, and let g be the interpolating natural cubic spline with values g_i at knots Z_i , $i = 1, \dots, n$. Let $\tilde{g}$ be any function in $\mathcal{G}$ for which $\tilde{g}(Z_i) = g_i$, $i = 1, \dots, n$. Then*

$$(2.7) \quad \int_0^1 |\tilde{g}''(t)|^2 dt \geq \int_0^1 |g''(t)|^2 dt.$$

with equality only if $\tilde{g} = g$.

This result allows us to restrict our attention to natural cubic splines, when studying the potential minimizers of

$$(2.8) \quad S_n(g) = M_n(g) + \lambda \int_0^1 |g''(z)|^2 dz.$$

Indeed, suppose $\tilde{g}$ is any function in $\mathcal{G}$ that is not a natural cubic spline with knots at Z_i . Let g be the natural cubic spline interpolant to the values $\tilde{g}(Z_i)$. Then $M_n(g) = M_n(\tilde{g})$. Because of the above optimality property of the natural cubic spline interpolant, (2.7) holds with strict inequality, and thus $S_n(\tilde{g}) > S_n(g)$. This means that, unless $\tilde{g}$ itself is a natural cubic spline with knots at Z_i , we can find a natural cubic spline with knots at Z_i that attains a smaller value of $S_n(g)$. It follows at once that a minimizer g of $S_n(g)$, if it exists, must be a natural cubic spline. It is key to notice that we have not forced g to be a natural cubic spline. This arises as a mathematical consequence of the choice of the roughness penalty. Now, as detailed below, we only need to minimize $S_n(g)$ over a finite-dimensional class of functions.

Assuming the Z_i 's are all different, which happens with probability one for a continuous Z , a natural cubic spline with knots at Z_i can be written as

$$(2.9) \quad g(z) = a_0 + a_1 z + \frac{1}{12} \sum_{i=1}^n \delta_i |z - Z_i|^3, \quad \sum_{i=1}^n \delta_i = \sum_{i=1}^n \delta_i Z_i = 0,$$

see [Green and Silverman \(1993, Section 7.3\)](#). The function g is uniquely defined by the coefficients a_0 , a_1 , and $\delta_i, i = 1, \dots, n$, or equivalently by its value at the knots, see [Proposition 2.1](#)'s proof for details.

It will be useful for what follows to use matrix notations. Let

$$\mathbf{Z} = \begin{pmatrix} 1 & Z_1 \\ \vdots & \vdots \\ 1 & Z_n \end{pmatrix},$$

$\mathbf{E} = [\frac{1}{12}|Z_i - Z_j|^3, i, j = 1, \dots, n]$, and $\mathbf{g} = (g(Z_1), \dots, g(Z_n))^T$. Then $\mathbf{g} = \mathbf{Z}\mathbf{a} + \mathbf{E}\boldsymbol{\delta}$ with constraints $\mathbf{Z}^T\boldsymbol{\delta} = 0$. Also, one can check that

$$\int g''(z)^2 dz = \boldsymbol{\delta}^T \mathbf{E} \boldsymbol{\delta},$$

see [Green and Silverman \(1993, Section 7.3\)](#). Let $\mathbf{Y}$ be the vector $(Y_1, \dots, Y_n)^T$, then

$$(2.10) \quad M_n(g) + \lambda \int (g''(z))^2 dz = (\mathbf{Y} - \mathbf{Z}\mathbf{a} - \mathbf{E}\boldsymbol{\delta})^T \boldsymbol{\Omega} (\mathbf{Y} - \mathbf{Z}\mathbf{a} - \mathbf{E}\boldsymbol{\delta}) + \lambda \boldsymbol{\delta}^T \mathbf{E} \boldsymbol{\delta},$$

where $\boldsymbol{\Omega}$ is the matrix with generic element $n^{-2}\omega(W_i - W_j)$. Hence, we want to minimize a quadratic function in parameters under the constraints $\mathbf{Z}^T\boldsymbol{\delta} = 0$. This yields a

unique solution under the usual requirements. The following proposition gives a precise characterization.

Proposition 2.1. *For any $\lambda > 0$, if all Z_i 's are different and all W_i 's are different, the solution to (2.6) exists, is unique, and is a natural cubic spline $\hat{g}$ characterized by*

$$(2.11) \quad \begin{bmatrix} \tilde{\mathbf{E}} & \mathbf{Z} \\ \mathbf{Z}^T & \mathbf{0} \end{bmatrix} \begin{pmatrix} \hat{\boldsymbol{\delta}} \\ \hat{\mathbf{a}} \end{pmatrix} = \begin{pmatrix} \mathbf{Y} \\ \mathbf{0} \end{pmatrix}, \quad \tilde{\mathbf{E}} = \mathbf{E} + \lambda \mathbf{\Omega}^{-1}.$$

Moreover, the values at the knots are

$$\hat{\mathbf{g}} = \left[\mathbf{P} + \mathbf{E} \tilde{\mathbf{E}}^{-1} (\mathbf{I} - \mathbf{P}) \right] \mathbf{Y},$$

where $\mathbf{P} = \mathbf{Z} \left(\mathbf{Z}^T \tilde{\mathbf{E}}^{-1} \mathbf{Z} \right)^{-1} \mathbf{Z}^T \tilde{\mathbf{E}}^{-1}$.

Our estimator is obtained directly by solving the linear system of equations (2.11). It does not necessitate estimation of other nonparametric quantities, and relies on only one regularization parameter λ . It also directly provides an estimator of the first derivative of g as

$$(2.12) \quad \tilde{g}'(z) = \hat{a}_1 + \frac{1}{4} \sum_{i=1}^n \hat{\delta}_i \text{sign}(z - Z_i)(z - Z_i)^2, \quad \text{sign}(u) = \mathbf{1}(u \geq 0) - \mathbf{1}(u < 0).$$

There are alternative ways to (2.9) for expressing a natural cubic spline. We focus on this formulation as it does not rely on a particular support of Z , nor on the fact that the knots Z_i are arranged in increasing order. In particular, the closed-form expression in Proposition 2.1 is valid regardless of the support of Z and therefore it can be used without first transforming Z into $(0, 1)$. We also found this formulation to be convenient for practical implementation. For large samples, where the above formula may not be computationally efficient, one can adapt to our context the Reinsch algorithm, see [Green and Silverman \(1993\)](#).

There is only one penalty parameter λ to choose to implement our estimator. We propose to choose it via cross-validation, where the quantity to be minimized is $M_n(g)$. For instance, we can split the data at random into two equally sized folds, and let each fold play the role of the training set in turn. Specifically, for each λ , we compute $\hat{g}_{k,\lambda}$ for each fold $k = 1, 2$, and we create the cross-validated vector of fitted values $\tilde{\mathbf{g}}_\lambda$, with typical element $\hat{g}_{k,\lambda}(Z_i)$, where k is the fold that does not include Z_i . We then choose the value of λ that minimizes $M_n(\tilde{\mathbf{g}}_\lambda)$ (note that $M_n(g) = M_n(\mathbf{g})$ depends only upon the values at the knots). This two-fold cross-validation method can easily be extended to (possibly repeated) k -fold cross-validation. In empirical implementations, we used the above two-fold cross-validation method with good results.

3. ASYMPTOTIC ANALYSIS

The formal study of our estimator is based on a reformulation of $M(g)$ in (2.4). Let $\mathcal{D}^2$ be the set of twice weakly differentiable functions. Consider

$$\mathcal{G} = \left\{ g : [0, 1] \rightarrow \mathbb{R}, g \in \mathcal{D}^2 : \int_0^1 |g''(t)|^2 dt < \infty \right\}, \quad \mathcal{H} = \{h \in \mathcal{G} : h(0) = h'(0) = 0\},$$

and the inner product $\langle h_1, h_2 \rangle_{\mathcal{H}} = \int_0^1 h_1''(z) h_2''(z) dz$ on $\mathcal{H}$. Then each $g \in \mathcal{G}$ can be uniquely written as $g(z) = (1, z)\beta + h(z)$, where $\beta = (g(0), g'(0)) \in \mathbb{R}^2$, $h(z) = g(z) - g(0) - g'(0)z$, $h \in \mathcal{H}$. Denote by L_μ^2 the space of complex functions l from $\mathbb{R}^q$ onto $\mathbb{C}$ such that

$$\|l\|_\mu^2 = \int |l(t)|^2 d\mu(t) < \infty.$$

Consider the operators $A : \mathcal{H} \mapsto L_\mu^2$ and $B : \mathbb{R}^2 \mapsto L_\mu^2$ such that

$$(3.13) \quad A h = E[h(Z) \exp(\mathbf{i}W^T \cdot)] \quad \text{and} \quad B \beta = E[(1, Z)\beta \exp(\mathbf{i}W^T \cdot)].$$

The minimization problem (2.4) identifying g_0 can be expressed as

$$(3.14) \quad \min_{\beta, h} \|E[Y \exp(\mathbf{i}W^T \cdot)] - B \beta - A h\|_\mu^2$$

for $(\beta, h) \in \mathbb{R}^2 \times \mathcal{H}$. The above quantity reaches its minimum zero at (β_0, h_0) , with $g_0(z) = (1, z)\beta_0 + h_0(z)$. A key advantage of this formulation for theoretical analysis is that using orthogonal projections, we can profile (3.14) to first determine h_0 , then β_0 as a function of h_0 . In our proofs, we will also consider the penalized empirical counterpart of (3.14) and use a similar profiling method to obtain $(\hat{\beta}, \hat{h})$, and then $\hat{g}(z) = (1, z)\hat{\beta} + \hat{h}(z)$.

The following assumption ensures that $E[Y \exp(\mathbf{i}W^T \cdot)] \in L_\mu^2$, and that A and B are onto L_μ^2 .

Assumption 3.1. (a) $E[Y^2] < \infty$; (b) Z has a bounded density f_Z on $[0, 1]$; (c) μ is a symmetric probability measure with support $\mathbb{R}^p$; (d) $\int \int E[\exp(\mathbf{i}W^T t) f_Z(z)]^2 \mu(t) dz dt < \infty$; (e) W has at least one continuous component.

Our assumptions on the support of Z is without much loss of generality, since we can always use a one-to-one transformation that maps Z into $[0, 1]$. We then formalize the completeness assumption, under which the problem (3.14) admits a unique solution (β_0, h_0) .

Assumption 3.2. g_0 belongs to $\mathcal{G}$ and the mapping $g \in \mathcal{G} \mapsto E[g(Z)|W = \cdot]$ is injective.

We introduce now a *source condition*, which is common in the literature on inverse problems. While it is not needed to establish the consistency of $\hat{g}$ and its first derivative, it is necessary to obtain convergence rates.

Assumption 3.3. Let M be the orthogonal projection onto the orthogonal of the span of B , and let $T = MA$. Let $(\sigma_j, \varphi_j, \psi_j)_j$ be the singular system of T , where $(\varphi_j)_j$ is a sequence of orthonormal elements in $\mathcal{H}$, $(\psi_j)_j$ is a sequence of orthonormal elements in L^2_μ , and $(\sigma_j)_j$ is a sequence of strictly positive values in $\mathbb{R}$. Then there exists $\gamma > 0$ such that

$$\sum_j \sigma_j^{-2\gamma} |\langle h_0, \varphi_j \rangle_{\mathcal{H}}|^2 < \infty.$$

Theorem 3.2. Under Assumptions 3.1, 3.2, if $\lambda \rightarrow 0$ and $n\lambda \rightarrow \infty$, then

$$\sup_{z \in [0,1]} |\widehat{g}(z) - g_0(z)| = o_p(1) \quad \text{and} \quad \sup_{z \in [0,1]} |\widehat{g}'(z) - g'_0(z)| = o_p(1).$$

If moreover Assumption 3.3 holds, then

$$\sup_{z \in [0,1]} |\widehat{g}(z) - g_0(z)| \quad \text{and} \quad \sup_{z \in [0,1]} |\widehat{g}'(z) - g'_0(z)| \quad \text{are both } O_p \left(\frac{1}{\sqrt{n\lambda}} + \lambda^{\frac{\gamma \wedge 2}{2}} \right).$$

We obtain consistency of our estimator and its derivative under mild assumptions, that only involve a standard condition on the regularization parameter λ . Also, since we avoid first stage estimation, we do not need any smoothness assumption on the distribution of Z given W . In two-step estimation methods that smooth over the instruments, one has to ensure that first-step estimation is consistent, and one typically needs conditions that relate the different smoothing parameters, see e.g. Ai and Chen (2003), Chen and Pouzo (2012). In some instances, consistency may further necessitate regularization parameters, see Chen and Pouzo (2012), and a source condition, see Gagliardini and Scaillet (2012). A general discussion can be found in Carrasco et al. (2007).

Turning now to our consistency rates, we do not claim that these are sharp or optimal. However, by contrast to previous results in this literature, they depend upon only one smoothing parameter. From the proofs, the rate $1/\sqrt{n\lambda}$ corresponds to a standard deviation term, while the second rate $\lambda^{\frac{\gamma \wedge 2}{2}}$ corresponds to a bias term. If λ is chosen to balance these two rates, we obtain the convergence rate $n^{-\frac{\gamma \wedge 2}{2(1+\gamma \wedge 2)}}$. For $\gamma = 2$ or 1 , this respectively yields $n^{-1/3}$ and $n^{-1/4}$. It is unclear how to compare the above rates to existing results on optimal convergence rates, see Hall and Horowitz (2005), Chen and Christensen (2018), as these authors make assumptions on conditional expectation operators such as $E\{g(Z)|W\}$, while Assumption 3.3 concerns an unconditional expectation operator. Our assumption, however, assumes that the problem is mildly ill-posed, while some previous work also considers the case of a severely ill-posed inverse problems.

4. ESTIMATION UNDER MONOTONICITY

In some instances, we may expect the function of interest g_0 to be monotonic. If g_0 is the Engel curve that relates the proportion of consumer expenditure on a good as a function

of total expenditure, we typically expect this function to be increasing for a “normal” good and decreasing for an “inferior” good. Accounting for monotonicity in estimation is expected to improve accuracy in small and moderate samples, see [Chetverikov and Wilhelm \(2017\)](#).

To implement such a monotonicity constraint into estimation, we note that since our smoothing splines estimator is linear, the derivative estimator (2.12) is linear as well. Let express it in matrix form. Since $\mathbf{g} = \mathbf{Z}\mathbf{a} + \mathbf{E}\boldsymbol{\delta}$, we can write $\mathbf{g}' = \mathbf{O}\mathbf{a} + \mathbf{D}\boldsymbol{\delta}$, where $\mathbf{g}' = (g'(Z_1), \dots, g'(Z_n))^T$, $\mathbf{D} = [\frac{1}{4} \text{sign}(Z_i - Z_j) |Z_i - Z_j|^2, i, j = 1, \dots, n]$, and

$$\mathbf{O} = \begin{pmatrix} 0 & 1 \\ \vdots & \vdots \\ 0 & 1 \end{pmatrix}.$$

From Proposition 2.1,

$$(4.15) \quad \begin{pmatrix} \hat{\boldsymbol{\delta}} \\ \hat{\mathbf{a}} \end{pmatrix} = \mathbf{S} \begin{pmatrix} \mathbf{Y} \\ \mathbf{0} \end{pmatrix}, \quad \mathbf{S} = \begin{bmatrix} \tilde{\mathbf{E}} & \mathbf{Z} \\ \mathbf{Z}^T & \mathbf{0} \end{bmatrix}^{-1}.$$

Hence,

$$\mathbf{g}' = (\mathbf{D}, \mathbf{O}) \mathbf{S} \begin{pmatrix} \mathbf{Y} \\ \mathbf{0} \end{pmatrix}.$$

We rely on a method proposed by [Hall and Huang \(2001\)](#), that is based on the same linear estimator but reweights the observations Y_i to impose monotonicity at observations points. It adjusts the unconstrained estimator by tilting the empirical distribution to make the least possible change, in the sense of a distance measure, subject to imposing the constraint of monotonicity at observation points. Specifically, if g_0 is assumed to be monotonically increasing, we consider the constrained optimization program

$$(4.16) \quad \min_{p_1, \dots, p_n} n - \sum_{i=1}^n (np_i)^{1/2}$$

subject to $\sum_{i=1}^n p_i = 1, \quad p_i \geq 0 \text{ for all } i = 1, \dots, n, \quad (\mathbf{D}, \mathbf{O}) \mathbf{S} \begin{pmatrix} \mathbf{p} \circ \mathbf{Y} \\ \mathbf{0} \end{pmatrix} \geq 0,$

where $\mathbf{p} \circ \mathbf{Y} = (p_1 Y_1, \dots, p_n Y_n)^T$ is the Hadamard product between vectors. If g_0 was assumed to be monotonically decreasing, we would modify the last inequalities. [Hall and Huang \(2001\)](#) considered more general optimization problems based on a family of Cressie-Read divergences, but we focus on the above program for convenience. It is strictly convex, so it admits a unique solution $\mathbf{p}^*$, and it is computationally fast to solve. The final estimator $\hat{g}^*$ is the natural cubic spline with coefficients $\mathbf{a}^*$ and $\boldsymbol{\delta}^*$ defined as in (4.15), with $\mathbf{p}^* \circ \mathbf{Y}$ in place of $\mathbf{Y}$.

We now state the asymptotic properties of our constrained smoothing splines estimator.

Assumption 4.4. *There exists $\eta > 0$ such that $g'_0(z) \geq \eta$ for all $z \in [0, 1]$.*

Theorem 4.3. *Under Assumptions 3.1, 3.2, and 4.4 if $\lambda \rightarrow 0$ and $n\lambda \rightarrow \infty$, then*

$$\sup_{z \in [0,1]} |\hat{g}^*(z) - g_0(z)| = o_p(1) \quad \text{and} \quad \sup_{z \in [0,1]} |\hat{g}^{*'}(z) - g'_0(z)| = o_p(1).$$

If moreover Assumption 3.3 holds, then

$$\sup_{z \in [0,1]} |\hat{g}^*(z) - g_0(z)| \quad \text{and} \quad \sup_{z \in [0,1]} |\hat{g}^{*'}(z) - g'_0(z)| \quad \text{are both } O_p\left(\frac{1}{\sqrt{n\lambda}} + \lambda^{\frac{\gamma \wedge 2}{2}}\right).$$

The above result directly follows from Theorem 3.2. Indeed, as $\hat{g}'$ is uniformly consistent, the constraint in the optimization problem (4.16) becomes asymptotically irrelevant from Assumption 4.4. Accordingly, $\hat{g}^* = \hat{g}$ with probability approaching one, and our results readily follow. While the monotonicity constraints become asymptotically irrelevant, they can matter in finite samples, as shown by Chetverikov and Wilhelm (2017) and illustrated by our empirical results.

5. NUMERICAL RESULTS

5.1. Small sample behavior. We used a DGP in line with Equation (1.1), where

$$\varepsilon = \frac{aV + \eta}{\sqrt{1 + a^2}}, \quad a = \sqrt{\frac{\rho_{\varepsilon V}^2}{1 - \rho_{\varepsilon V}^2}},$$

$$Z = \frac{\beta W + V}{\sqrt{1 + \beta^2}}, \quad \beta = \sqrt{\frac{\rho_{WZ}^2}{1 - \rho_{WZ}^2}},$$

and (W, V, η) are independent standard Gaussian. This yields standard Gaussian marginal distributions for ε and Z whatever the values of the parameters. The correlation $\rho_{\varepsilon V}$ measures the level of endogeneity of Z . The correlation ρ_{WZ} measures instead the strength of the instrument W .

We implemented our smoothing splines estimator with ω equal to the density of a Laplace distribution with mean zero and variance 1. The choice of the penalty parameter λ was based on two-fold cross-validation, as previously detailed. We considered λ within the grid $\{p/(1-p), p = 10^{-5} + k * (0.7 - 10^{-5})/399, k = 0, \dots, 399\}$.

We compared our estimator to two existing methods, for which a data-driven procedure has been proposed for the choice of smoothing or regularization parameters. We considered first the kernel-based Tikhonov estimator of Darolles et al. (2011), hereafter referred as Tikhonov. We also considered the series estimator of Horowitz (2014) based on a basis of Legendre polynomials. The implementation details of both methods are given in the online supplement, together with supplementary results.

We first considered two functional forms for g_0 , each normalized to have unit variance: a quadratic function $g_{0,1}(z) = z^2/\sqrt{2}$, and a non-polynomial function $g_{0,2}(z) = \sqrt{3\sqrt{3}}z \exp(-z^2/2)$. We ran 2000 Monte Carlo simulations with sample sizes $n = 200$ and 400. We consider three couples of values for $(\rho_{\varepsilon V}, \rho_{WZ})$: (a) (0.5, 0.9), a setting with low endogeneity and a strong instrument, (b) (0.8, 0.9), corresponding to high endogeneity and a strong instrument, (c) (0.8, 0.7), a more complex setting with high endogeneity level but a weaker instrument. To evaluate the gains of imposing monotonicity, we then considered a third function $g_{0,3}(z) = (\sqrt{(10/3)} \log(|z-1|+1) \text{sign}(z-1) - 0.6z + 2z^3)/8$. The regularization parameter λ was chosen before the monotonicizing step, and we used the R package CVXR to solve (4.16), see Fu et al. (2020).

Table 1 reports our results. The Tikhonov estimator is severely biased in all cases, while our estimator is almost unbiased. The series estimator mostly lies in between, but with large differences depending on the setup. In terms of variance, Tikhonov does better than smoothing splines, that itself does better than series. Smoothing splines performs best in terms of MSE in almost all cases. Exceptions are cases corresponding to the second function with $n = 400$ and strong instruments, where the series estimator is close to unbiased. Overall, the severity of endogeneity does not affect much the estimators' performances. However, the strength of instruments has important effects. Our smoothing splines estimator appears to be more robust than its competitors to a decrease in the strength of the instrument. Finally, imposing monotonicity does not affect much bias, but yields a substantial decrease in variance. Depending on the particular setup, it can be more than halved.

5.2. Empirical application. We applied the smoothing spline estimator to the estimation of Engel curves, which relate the proportion of spending on a given good as a function of total expenditures. We used the “Engel95” data, from the R package np, see Hayfield and Racine (2008). This dataset is a random sample from the 1995 British Family Expenditure Survey and contains data for 1655 households of married couples for which the head-of-household is employed and between 25 and 55 years old. We focused on the subsample of 628 households with no kids. We report results for two Engel curves, pertaining to the expenditure shares on leisure and fuel. Economic theory suggests that the Engel curve for leisure is increasing and the one for fuel is decreasing. Following Blundell et al. (2007), we instrumented the *logarithm of total household's expenditure*, which is likely endogenous, by the *logarithm of total earnings before tax*. We consider the four estimators used in our simulations, and implementation details remain the same.

The estimated nonparametric functions are reported in Figure 1. The Tikhonov estimate exhibits a non-monotonic and quite irregular behavior, while the series estimate is mainly monotonic and very regular. Since our smoothing splines estimates are monotonic,

TABLE 1. Simulation results

		$g_{0,1}$			$g_{0,2}$			$g_{0,3}$			
n		Sm.	Tikh.	Ser.	Sm.	Tikh.	Ser.	Cons.	Sm.	Tikh.	Ser.
$\rho_{ZW} = 0.9, \rho_{\varepsilon V} = 0.5$											
200	Bias ²	0.000	0.148	0.092	0.001	0.065	0.005	0.000	0.000	0.088	0.104
200	Var	0.069	0.030	0.060	0.074	0.028	0.073	0.041	0.076	0.022	0.116
200	MSE	0.069	0.177	0.152	0.075	0.092	0.078	0.044	0.076	0.111	0.219
400	Bias ²	0.000	0.098	0.094	0.001	0.044	0.005	0.000	0.000	0.054	0.140
400	Var	0.052	0.020	0.031	0.053	0.018	0.025	0.026	0.056	0.015	0.044
400	MSE	0.052	0.118	0.125	0.054	0.063	0.030	0.027	0.056	0.069	0.184
$\rho_{ZW} = 0.9, \rho_{\varepsilon V} = 0.8$											
200	Bias ²	0.001	0.148	0.092	0.001	0.064	0.005	0.000	0.000	0.093	0.107
200	Var	0.066	0.028	0.060	0.072	0.026	0.074	0.039	0.072	0.019	0.107
200	MSE	0.067	0.176	0.152	0.072	0.090	0.079	0.042	0.073	0.112	0.214
400	Bias ²	0.000	0.098	0.094	0.000	0.043	0.005	0.000	0.000	0.055	0.141
400	Var	0.049	0.019	0.030	0.051	0.018	0.026	0.025	0.054	0.013	0.042
400	MSE	0.050	0.117	0.125	0.052	0.061	0.030	0.026	0.054	0.068	0.183
$\rho_{ZW} = 0.7, \rho_{\varepsilon V} = 0.8$											
200	Bias ²	0.009	0.307	0.133	0.004	0.158	0.158	0.012	0.012	0.184	0.189
200	Var	0.091	0.032	0.130	0.120	0.028	0.053	0.056	0.101	0.019	0.050
200	MSE	0.099	0.338	0.262	0.124	0.186	0.211	0.080	0.113	0.203	0.238
400	Bias ²	0.003	0.218	0.139	0.004	0.110	0.157	0.002	0.002	0.123	0.190
400	Var	0.069	0.024	0.053	0.087	0.021	0.024	0.041	0.086	0.014	0.021
400	MSE	0.073	0.242	0.192	0.090	0.131	0.180	0.051	0.089	0.138	0.212

Note: Average over a grid of 100 equidistant points on $[-2, 2]$ and 2000 Monte Carlo replications of the squared bias (Bias²), the variance (Var), and the Mean Squared Error (MSE) for the constrained smoothing splines estimator (Cons.), the smoothing splines estimator (Sm.), the Tikhonov estimator (Tikh.), and the series estimator (Ser.).

but at the boundaries of the data, our constrained and unconstrained estimates are very close. Both are in line with the findings of [Blundell et al. \(2007\)](#).

6. EXTENSION TO A PARTLY LINEAR MODEL

We have proposed a generalization of regression smoothing splines to the context where there is endogeneity and instrumental variables are available. While we detail our estimator and its properties in the simple univariate context, a multivariate extension could be considered. However, including more covariates in a fully nonparametric way would submit us to the curse of dimensionality typical of functional estimation. Hence, we focus




FIGURE 1. Estimated Engel curves.

here on a partly linear model, as considered by e.g. [Heckman \(1986\)](#), [Robinson \(1988\)](#), [Blundell et al. \(2007\)](#), [Chen and Pouzo \(2009\)](#), and [Florens et al. \(2012\)](#). This model provides a simple and economical way to include additional controls. We thus consider

$$(6.17) \quad Y = X^T \gamma_0 + g_0(Z) + \varepsilon \quad E[\varepsilon|W] = 0,$$

where $X \in \mathbb{R}^q$ is a vector of exogenous covariates, whose components are thus included in W , while, as earlier, $Z \in \mathbb{R}$ is the endogenous variable. The following condition ensures identification of (γ_0, g_0) in (6.17).

Assumption 6.5. (γ_0, g_0) belongs to $\mathbb{R}^q \times \mathcal{G}$ and the mapping $(\gamma_0, g_0) \in \mathbb{R}^q \times \mathcal{G} \mapsto E[X^T \gamma + g(Z)|W]$ is injective.

Our identification assumption is similar to ones imposed in other work, see, e.g., [Chen and Pouzo \(2012\)](#) or [Florens et al. \(2012\)](#). First, it excludes collinearity between the components of X . Second, it requires that the distribution of Z given W must be complete so that the mapping $g \mapsto E[g(Z)|W]$ is injective. Third, it rules out the presence of an intercept in X , since an intercept can always be absorbed by the nonparametric function g_0 . Fourth, it requires that no function g is such that $E[g(Z)|W]$ is a linear function of the variables in X . Under Assumption 6.5,

$$(\gamma_0, g_0) = \arg \min_{(\gamma, g) \in \mathbb{R}^q \times \mathcal{G}} \int |E[(Y - X^T \gamma - g(Z)) \exp(\mathbf{i}W^T t)]|^2 d\mu(t).$$

We proceed as in the benchmark model and estimate (γ_0, g_0) by minimizing the empirical counterpart of the above penalized by a roughness measure of the nonparametric component, that is

$$(6.18) \quad (\hat{\gamma}, \hat{g}) \in \arg \min_{(\gamma, g) \in \mathbb{R}^q \times \mathcal{G}} M_n^{PL}(\gamma, g) + \lambda \int |g''(z)|^2 dz,$$

where $M_n^{PL}(\gamma, g) = \int \left| \frac{1}{n} \sum_{i=1}^n (Y_i - X_i^T \gamma - g(Z_i)) \exp(\mathbf{i} W^T t) \right|^2 d\mu(t).$

The estimators $(\hat{\gamma}, \hat{g})$ can be computed in the same way as in the benchmark model. To show this, let $\mathbf{L}$ be the $n \times (q+2)$ matrix whose row i is $(1, Z_i, X_i^T)$. The following is a direct extension of Proposition 2.1 to the partly linear model.

Proposition 6.2. *For any $\lambda > 0$, if all Z_i 's and all W_i 's are different, and $\mathbf{L}$ is full-column rank, the solution to (6.18) exists and is unique. The estimator $\hat{g}$ is a natural cubic spline. The coefficients $\hat{\mathbf{a}} = (\hat{a}_0, \hat{a}_1, \hat{\gamma}^T)^T$ and $\hat{\boldsymbol{\delta}}$ are characterized by*

$$(6.19) \quad \begin{bmatrix} \tilde{\mathbf{E}} & \mathbf{L} \\ \mathbf{L}^T & \mathbf{0} \end{bmatrix} \begin{pmatrix} \hat{\boldsymbol{\delta}} \\ \hat{\mathbf{a}} \end{pmatrix} = \begin{pmatrix} \mathbf{Y} \\ \mathbf{0} \end{pmatrix}, \quad \tilde{\mathbf{E}} = \mathbf{E} + \lambda \boldsymbol{\Omega}^{-1}.$$

We now focus on the consistency and the convergence rates of the regression function. Let us define the operator $\mathbf{D} : \mathbb{R}^{q+2} \mapsto L_\mu^2$ such that

$$(6.20) \quad \mathbf{D}(\gamma, \beta) = E[(X^T \gamma + (1, Z)\beta) \exp(\mathbf{i} W^T \cdot)].$$

Assumption 6.6. *Assumption 3.1 holds and $E\|X\|^2 < \infty$.*

Assumption 6.7. *Assumption 3.3 holds with $\mathbf{D}$ replacing $\mathbf{B}$ and $\mathbf{M}$ being the orthogonal projection operator onto the orthogonal of the span of $\mathbf{D}$.*

The following is a direct extension of Theorem 3.2.

Theorem 6.4. *Under Assumptions 6.5 and 6.6, if $\lambda \rightarrow 0$ and $n\lambda \rightarrow \infty$, then*

$$\sup_{z \in [0,1]} |\hat{g}(z) - g_0(z)| = o_p(1) \quad \text{and} \quad \sup_{z \in [0,1]} |\hat{g}'(z) - g_0'(z)| = o_p(1).$$

If moreover Assumption 6.7 holds, then

$$\sup_{z \in [0,1]} |\hat{g}(z) - g_0(z)| \quad \text{and} \quad \sup_{z \in [0,1]} |\hat{g}'(z) - g_0'(z)| \quad \text{are both } O_p \left(\frac{1}{\sqrt{n\lambda}} + \lambda^{\frac{\gamma \wedge 2}{2}} \right).$$

7. CONCLUDING REMARKS

Further extensions to our method could be considered, such as a nonparametrically additive model, see [Linton and Nielsen \(1995\)](#). A theory of inference on the nonparametric function of interest, or some functional of it, is to be developed, see, e.g., [Chen and Christensen \(2018\)](#) for results on series-based estimators. This would likely necessitate extending recent work on splines asymptotics, see [Claeskens et al. \(2009\)](#), [Schwarz and Krivobokova \(2016\)](#). These issues are left for further research.

REFERENCES

- Ai, C. and Chen, X. (2003). Efficient estimation of models with conditional moment restrictions containing unknown functions. *Econometrica*, 71:1795–843.
- Bennett, A., Kallus, N., and Schnabel, T. (2019). Deep generalized method of moments for instrumental variable analysis. In Wallach, H., Larochelle, H., Beygelzimer, A., Alché-Buc, F. d., Fox, E., and Garnett, R., editors, *Advances in Neural Information Processing Systems*, volume 32, pages 3564–74. Curran Associates, Inc.
- Bierens, H. J. (1982). Consistent model specification tests. *J Econometrics*, 20:105–34.
- Blundell, R., Chen, X., and Kristensen, D. (2007). Semi-nonparametric IV estimation of shape-invariant engel curves. *Econometrica*, 75:1613–69.
- Busby, R. C., Schochetman, I., and Smith, H. A. (1972). Integral operators and the compactness of induced representations. *Transactions of the American Mathematical Society*, 164:461–77.
- Carrasco, M., Florens, J.-P., and Renault, E. (2007). Linear inverse problems in structural econometrics estimation based on spectral decomposition and regularization. In Heckman, J. J. and Leamer, E. E., editors, *Handbook of Econometrics*, volume 6, chapter 77, pages 5633–751. Amsterdam: Elsevier.
- Centorrino, S., Fève, F., and Florens, J.-P. (2017). Additive nonparametric instrumental regressions: A guide to implementation. *Journal of Econometric Methods*, 6:1–25.
- Chen, X. and Christensen, T. M. (2018). Optimal sup-norm rates and uniform inference on nonlinear functionals of nonparametric IV regression. *Quantitative Economics*, 9:39–84.
- Chen, X. and Pouzo, D. (2009). Efficient estimation of semiparametric conditional moment models with possibly nonsmooth residuals. *Journal of Econometrics*, 152:46–60.
- Chen, X. and Pouzo, D. (2012). Estimation of nonparametric conditional moment models with possibly nonsmooth generalized residuals. *Econometrica*, 80:277–321.
- Chetverikov, D. and Wilhelm, D. (2017). Nonparametric instrumental variable estimation under monotonicity. *Econometrica*, 85:1303–320.

- Claeskens, G., Krivobokova, T., and Opsomer, J. D. (2009). Asymptotic properties of penalized spline estimators. *Biometrika*, 96:529–44.
- Darolles, S., Fan, Y., Florens, J.-P., and Renault, E. (2011). Nonparametric instrumental regression. *Econometrica*, 79:1541–65.
- Dikkala, N., Lewis, G., Mackey, L., and Syrgkanis, V. (2020). Minimax estimation of conditional moment models. In Larochelle, H., Ranzato, M., Hadsell, R., Balcan, M. F., and Lin, H., editors, *Advances in Neural Information Processing Systems*, volume 33, pages 12248–12262. Curran Associates, Inc.
- Dreier, I. and Kotz, S. (2002). A note on the characteristic function of the t-distribution. *Statistics & Probability Letters*, 57:221 – 24.
- Dunker, F., Florens, J.-P., Hohage, T., Johannes, J., and Mammen, E. (2014). Iterative estimation of solutions to noisy nonlinear operator equations in nonparametric instrumental regression. *Journal of Econometrics*, 178:444–55.
- D’Haultfoeuille, X. (2011). On the completeness condition in nonparametric instrumental problems. *Econometric Theory*, 27:460–71.
- Florens, J.-P. (2003). Inverse problems and structural econometrics: The example of instrumental variables. In Dewatripont, M., Hansen, L. P., and Turnovsky, S. J., editors, *Advances in Economics and Econometrics: Theory and Applications, Eighth World Congress*, volume 2 of *Econometric Society Monographs*, pages 284–311. Cambridge: Cambridge University Press.
- Florens, J.-P., Johannes, J., and Van Bellegem, S. (2011). Identification and estimation by penalization in nonparametric instrumental regression. *Econometric Theory*, 27:472–96.
- Florens, J.-P., Johannes, J., and Van Bellegem, S. (2012). Instrumental regression in partially linear models. *The Econometrics Journal*, 15:304–24.
- Freyberger, J. (2017). On completeness and consistency in nonparametric instrumental variable models. *Econometrica*, 85:1629–644.
- Fu, A., Narasimhan, B., and Boyd, S. (2020). CVXR: An R package for disciplined convex optimization. *Journal of Statistical Software*, 94:1–34.
- Gagliardini, P. and Scaillet, O. (2012). Tikhonov regularization for nonparametric instrumental variable estimators. *Journal of Econometrics*, 167:61–75.
- Green, P. J. and Silverman, B. W. (1993). *Nonparametric regression and generalized linear models: a roughness penalty approach*. Boca Raton, FL: CRC Press.
- Hall, P. and Horowitz, J. L. (2005). Nonparametric methods for inference in the presence of instrumental variables. *The Annals of Statistics*, 33:2904–29.
- Hall, P. and Huang, L.-S. (2001). Nonparametric kernel regression subject to monotonicity constraints. *The Annals of Statistics*, 29:624–47.

- Hall, P. and Opsomer, J. D. (2005). Theory for penalised spline regression. *Biometrika*, 92:105–18.
- Hayfield, T. and Racine, J. S. (2008). Nonparametric econometrics: The np package. *Journal of Statistical Software*, 27(5):1–32.
- Heckman, N. E. (1986). Spline smoothing in a partly linear model. *Journal of the Royal Statistical Society. Series B (Methodological)*, 48:244–48.
- Horowitz, J. L. (2011). Applied nonparametric instrumental variables estimation. *Econometrica*, 79:347–94.
- Horowitz, J. L. (2014). Adaptive nonparametric instrumental variables estimation: Empirical choice of the regularization parameter. *Journal of Econometrics*, 180:158–73.
- Johannes, J., Van Bellegem, S., and Vanhems, A. (2011). Convergence rates for ill-posed inverse problems with an unknown operator. *Econometric Theory*, 27:522–45.
- Johannes, J., Van Bellegem, S., and Vanhems, A. (2013). Iterative regularisation in nonparametric instrumental regression. *Journal of Statistical Planning and Inference*, 143:24–39.
- Johnson, N., Kotz, S., and Balakrishnan, N. (1995). *Continuous univariate distributions*, volume 2 of *Wiley series in probability and mathematical statistics: Applied probability and statistics*. Hoboken, NJ: John Wiley & Sons.
- Kress, R. (1999). *Linear integral equations*. New York, NY: Springer.
- Kreyszig, E. (1978). *Introductory functional analysis with applications*. Hoboken, NJ: John Wiley & Sons.
- Li, Y. and Ruppert, D. (2008). On the asymptotics of penalized splines. *Biometrika*, 95:415–36.
- Liao, L., Chen, Y.-L., Yang, Z., Dai, B., Kolar, M., and Wang, Z. (2020). Provably efficient neural estimation of structural equation models: An adversarial approach. In *Advances in Neural Information Processing Systems*, volume 33, pages 8947–58.
- Linton, O. and Nielsen, J. P. (1995). A kernel method of estimating structured nonparametric regression based on marginal integration. *Biometrika*, 82:93–100.
- Luenberger, D. G. and Ye, Y. (2008). *Linear and nonlinear programming*, volume 116 of *International Series in Operations Research & Management Science*. Springer US, New York, NY.
- Muandet, K., Mehrjou, A., Lee, S. K., and Raj, A. (2020). Dual instrumental variable regression. *Advances in Neural Information Processing Systems*, 33:2710–21.
- Newey, W. K. and Powell, J. L. (2003). Instrumental variable estimation of nonparametric models. *Econometrica*, 71:1565–578.
- Robinson, P. M. (1988). Root-n consistent semiparametric regression. *Econometrica*, 56:931–54.

- Schumaker, L. (2007). *Spline functions: Basic theory*. Cambridge: Cambridge University Press, 3 edition.
- Schwarz, K. and Krivobokova, T. (2016). A unified framework for spline estimators. *Biometrika*, 103:121–31.
- Singh, R., Sahani, M., and Gretton, A. (2019). Kernel instrumental variable regression. In Wallach, H., Larochelle, H., Beygelzimer, A., Alché-Buc, F. d., Fox, E., and Garnett, R., editors, *Advances in Neural Information Processing Systems*, volume 32. Curran Associates, Inc.
- Wahba, G. (1990). *Spline models for observational data*. Philadelphia, PA, Society for Industrial and Applied Mathematics.
- Zhang, R., Imaizumi, M., Schölkopf, B., and Muandet, K. (2023). Instrumental variable regression via kernel maximum moment loss. *Journal of Causal Inference*, 11:20220073.

APPENDIX A. MONTE CARLO: ADDITIONAL INFORMATION

Implementation of the Tikhonov estimator. The Tikhonov optimization problem at the population level is

$$\arg \min_g \|E[Y|W] - E[g(Z)|W]\|_{L^2_W}^2 + \lambda \|g\|_{L^2_Z}^2.$$

Solving the analog sample level problem involves estimating not only the conditional expectation operator given W , but also the adjoint conditional expectation operator given Z . Both are estimated by kernel smoothing. We used Gaussian kernels of order 2, while the bandwidths were set using Silverman’s rule of thumb, i.e., equal to $n^{-1/5}$ times the empirical standard deviation of the variable on which smoothing is performed (either Z or W). To select the regularization parameter, we use the pseudo-cross-validation procedure of [Centorrino et al. \(2017\)](#), searching for the minimum of the criterion on a grid between 0 and 1.

Implementation of the series estimator. We also considered a series estimator based on a basis of Legendre polynomials. The main idea is to consider the equality

$$E[Y|W] f_W(W) = \int g(Z) f_{(Z,W)}(Z, W) dZ.$$

The right-hand side of the equation, the function g , and the joint density $f_{(Z,W)}$ are each approximated by a series expansion, respectively on J , K , and $J \times K$ terms. We used a method proposed by [Horowitz \(2014\)](#), who considered the case $J = K$ and derived an adaptive procedure to select J . Since this method is designed for variables belonging to $[0, 1]$, we transformed observations of Z and W by their respective empirical cumulative distribution functions (cdf). (In unreported simulations, we found that using the true cdfs instead did not affect our results much.) This implies in particular that even if the




FIGURE 2. True regression function and average estimators for $n = 200$, $\rho_{\epsilon V} = 0.8$, and $\rho_{ZW} = 0.7$.

relation between Z and W is linear, the first-stage equation is not linear anymore in the transformed variables.

Supplementary results. Figures 2a, 2b and 2c graph the pointwise average of each estimator for $n = 200$ and $(\rho_{\epsilon V}, \rho_{WZ}) = (0.8, 0.7)$. Here, the series estimator is much

steeper than the true quadratic curve $g_{0,1}$, while it fails to fit the sign changes in the first derivative for $g_{0,2}$. In all cases, the Tikhonov estimator is much smoother than the true curves, while the smoothing splines estimator is almost unbiased. From our figures, the degree of smoothing appears to be quite different among the averaged estimators. However, there is no clearly accepted way to measure degrees of freedom in nonparametric instrumental variable regression. The issue is particularly intricate for our competitors. The Tikhonov estimator depends on two bandwidths parameters as well as a regularization parameter, and the influence of each choice on the final estimator is far from clear. The same comment applies to the series estimator, which relies on three estimated nonparametric components. The method proposed by Horowitz (2014) simplifies the matter by making each dependent upon a single parameter J , but the effect of this choice on the final estimator remains to be investigated. By contrast, our smoothing spline estimator depends upon a single regularization parameter.

APPENDIX B. PROOF OF PROPOSITION 2.1

(a) *Unicity.* We begin by studying Ω . Let $\mathbf{b} = (b_1, \dots, b_n)^T \in \mathbb{R}^n$, then

$$\mathbf{b}^T \Omega \mathbf{b} = \int \left| \frac{1}{n} \sum_{i=1}^n b_i \exp(\mathbf{i} W_i^T t) \right|^2 \mu(dt) \geq 0.$$

Hence $\mathbf{b}^T \Omega \mathbf{b} = 0$ iff $(1/n) \sum_{i=1}^n b_i \exp(\mathbf{i} W_i^T t) = 0$ for all $t \in \mathbb{R}^q$. Define the random vector $(\tilde{b}, \tilde{\mathbf{W}})$ that equals (b_i, W_i) with probability $1/n$, and $\tilde{E}$ the corresponding expectation. Then, $\tilde{E}[\tilde{b} \exp(\mathbf{i} \tilde{\mathbf{W}}^T t)] = 0$ for all $t \in \mathbb{R}^q$. From Bierens (1982), this implies that $\tilde{E}[\tilde{b} | \tilde{\mathbf{W}} = \mathbf{W}_i] = 0$. Since $\tilde{E}[\tilde{b} | \tilde{\mathbf{W}} = \mathbf{W}_i] = b_i$ if all W_i s are different, $b_i = 0$ for all $i = 1, \dots, n$. Hence, Ω is positive definite.

From Green and Silverman (1993, Chapter 2), a natural cubic spline is uniquely defined by the vector of its values at the knots $\mathbf{g}$, and we can write

$$\int g''(z)^2 dz = \mathbf{g}^T \mathbf{K} \mathbf{g},$$

for a positive semi-definite matrix $\mathbf{K}$. Hence, our minimization problem writes

$$\min_{\mathbf{g}} (\mathbf{Y} - \mathbf{g})^T \Omega (\mathbf{Y} - \mathbf{g}) + \lambda \mathbf{g}^T \mathbf{K} \mathbf{g} = \min_{\mathbf{g}} \mathbf{g}^T (\Omega + \lambda \mathbf{K}) \mathbf{g} + 2 \mathbf{g}^T \Omega \mathbf{Y} + \mathbf{Y}^T \Omega \mathbf{Y}.$$

Since $\Omega + \lambda \mathbf{K}$ is positive-definite for any $\lambda > 0$, the problem is convex and has a unique minimum.

(b) *Solution.* A natural cubic spline g can also be uniquely written as

$$(A.1) \quad g(z) = a_0 + a_1 z + \frac{1}{12} \sum_{i=1}^n \delta_i |z - Z_i|^3, \quad \sum_{i=1}^n \delta_i = \sum_{i=1}^n \delta_i Z_i = 0,$$

whenever the Z_i 's are all different. One can thus write $\mathbf{g} = \mathbf{E} \boldsymbol{\delta} + \mathbf{Z} \mathbf{a}$, with $\mathbf{Z}^T \boldsymbol{\delta} = 0$.

To show that the above formulation is unique, let us check that $\mathbf{E}\boldsymbol{\delta} + \mathbf{Z}\mathbf{a} = \mathbf{0}$ with $\mathbf{Z}^T\boldsymbol{\delta} = \mathbf{0}$ implies the nullity of all coefficients. Our premises yield $\boldsymbol{\delta}^T(\mathbf{E}\boldsymbol{\delta} + \mathbf{Z}\mathbf{a}) = \boldsymbol{\delta}^T\mathbf{E}\boldsymbol{\delta} = 0$. From [Green and Silverman \(1993, Section 7.3\)](#),

$$(A.2) \quad \int g''(z)^2 dz = \boldsymbol{\delta}^T \mathbf{E} \boldsymbol{\delta} \geq 0.$$

Hence, since g'' is continuous, it should be that g'' is identically zero. Since the increments of the third derivative at the knots are $g'''(Z_i^+) - g'''(Z_i^-) = \delta_i$, $i = 1, \dots, n$, this implies that $\boldsymbol{\delta} = \mathbf{0}$. Finally, $\mathbf{Z}\mathbf{a} = \mathbf{0}$ implies $\mathbf{a} = \mathbf{0}$, as $\mathbf{Z}$ is full rank.

From the uniqueness obtained in Part (a),

$$(A.3) \quad (\mathbf{Y} - \mathbf{Z}\mathbf{a} - \mathbf{E}\boldsymbol{\delta})^T \boldsymbol{\Omega} (\mathbf{Y} - \mathbf{Z}\mathbf{a} - \mathbf{E}\boldsymbol{\delta}) + \lambda \boldsymbol{\delta}^T \mathbf{E} \boldsymbol{\delta}$$

admits a unique global minimum under the constraint $\mathbf{Z}^T\boldsymbol{\delta} = \mathbf{0}$. To characterize such a minimum, consider the Lagrangian

$$(\mathbf{Y} - \mathbf{Z}\mathbf{a} - \mathbf{E}\boldsymbol{\delta})^T \boldsymbol{\Omega} (\mathbf{Y} - \mathbf{Z}\mathbf{a} - \mathbf{E}\boldsymbol{\delta}) + \lambda \boldsymbol{\delta}^T \mathbf{E} \boldsymbol{\delta} + \mathbf{l}^T \mathbf{Z}^T \boldsymbol{\delta},$$

where $\mathbf{l} \in \mathbb{R}^2$ is the vector of Lagrange multipliers, and the associated first-order conditions with respect to $(\boldsymbol{\delta}^T, \mathbf{a}^T)^T$

$$\begin{bmatrix} \mathbf{E}^T \boldsymbol{\Omega} & \mathbf{0} \\ \mathbf{Z}^T \boldsymbol{\Omega} & -\lambda \mathbf{I} \end{bmatrix} \begin{bmatrix} \tilde{\mathbf{E}} & \mathbf{Z} \\ \mathbf{Z}^T & \mathbf{0} \end{bmatrix} \begin{pmatrix} \boldsymbol{\delta} \\ \mathbf{a} \end{pmatrix} - \begin{bmatrix} \mathbf{E}^T \boldsymbol{\Omega} & \mathbf{0} \\ \mathbf{Z}^T \boldsymbol{\Omega} & -\lambda \mathbf{I} \end{bmatrix} \begin{pmatrix} \mathbf{Y} \\ \mathbf{0} \end{pmatrix} - \begin{pmatrix} \mathbf{Z} \mathbf{l} \\ \mathbf{0} \end{pmatrix} = \begin{pmatrix} \mathbf{0} \\ \mathbf{0} \end{pmatrix}.$$

One solution to these first-order conditions is given by $\mathbf{l} = (0, 0)^T$ and $(\hat{\boldsymbol{\delta}}^T, \hat{\mathbf{a}}^T)^T$ satisfying

$$(A.4) \quad \begin{bmatrix} \tilde{\mathbf{E}} & \mathbf{Z} \\ \mathbf{Z}^T & \mathbf{0} \end{bmatrix} \begin{pmatrix} \hat{\boldsymbol{\delta}} \\ \hat{\mathbf{a}} \end{pmatrix} = \begin{pmatrix} \mathbf{Y} \\ \mathbf{0} \end{pmatrix}, \quad \tilde{\mathbf{E}} = \mathbf{E} + \lambda \boldsymbol{\Omega}^{-1},$$

and this solution satisfies the constraint $\mathbf{Z}^T \hat{\boldsymbol{\delta}} = \mathbf{0}$. This solution is a strict local minimum if the second-order sufficient conditions hold ([Luenberger and Ye, 2008](#), page 334), that is for any non-zero $(\boldsymbol{\delta}^T, \mathbf{a}^T)^T$ satisfying $\mathbf{Z}^T \boldsymbol{\delta} = \mathbf{0}$,

$$(A.5) \quad \begin{pmatrix} \boldsymbol{\delta} \\ \mathbf{a} \end{pmatrix}^T \begin{bmatrix} \mathbf{E}^T \boldsymbol{\Omega} \mathbf{E} + \lambda \mathbf{E} & \mathbf{E}^T \boldsymbol{\Omega} \mathbf{Z} \\ \mathbf{Z}^T \boldsymbol{\Omega} \mathbf{E} & \mathbf{Z}^T \boldsymbol{\Omega} \mathbf{Z} \end{bmatrix} \begin{pmatrix} \boldsymbol{\delta} \\ \mathbf{a} \end{pmatrix} = (\mathbf{E}\boldsymbol{\delta} + \mathbf{Z}\mathbf{a})^T \boldsymbol{\Omega} (\mathbf{E}\boldsymbol{\delta} + \mathbf{Z}\mathbf{a}) + \lambda \boldsymbol{\delta}^T \mathbf{E} \boldsymbol{\delta} > 0.$$

As $\boldsymbol{\Omega}$ is positive definite, and using (A.2), (A.5) is non-negative, and is zero iff $\mathbf{E}\boldsymbol{\delta} + \mathbf{Z}\mathbf{a} = \mathbf{0}$. But this would imply $(\boldsymbol{\delta}^T, \mathbf{a}^T)^T = \mathbf{0}$ as shown above.

The right-hand side matrix in (A.4) is full rank as $\mathbf{Z}$ is full rank and $\lambda > 0$. Indeed, assume

$$\begin{bmatrix} \mathbf{E} + \lambda \boldsymbol{\Omega}^{-1} & \mathbf{Z} \\ \mathbf{Z}^T & \mathbf{0} \end{bmatrix} \begin{pmatrix} \boldsymbol{\delta} \\ \mathbf{a} \end{pmatrix} = \begin{pmatrix} \mathbf{0} \\ \mathbf{0} \end{pmatrix},$$

this implies $\mathbf{Z}^T \boldsymbol{\delta} = \mathbf{0}$ and $\mathbf{0} = \boldsymbol{\delta}^T [(\mathbf{E} + \lambda \boldsymbol{\Omega}^{-1}) \boldsymbol{\delta} + \mathbf{Z} \mathbf{a}] = \boldsymbol{\delta}^T \mathbf{E} \boldsymbol{\delta} + \lambda \boldsymbol{\delta}^T \boldsymbol{\Omega}^{-1} \boldsymbol{\delta}$. From (A.2) and the positive-definiteness of $\boldsymbol{\Omega}$, this yields $\boldsymbol{\delta} = \mathbf{0}$, and in turn $\mathbf{a} = \mathbf{0}$.

To obtain the values at the knots $\hat{\mathbf{g}}$, note that the inverse of the matrix in (A.4) is

$$(A.6) \quad \begin{bmatrix} \tilde{\mathbf{E}}^{-1} (\mathbf{I} - \mathbf{P}) & \tilde{\mathbf{E}}^{-1} \mathbf{Z} (\mathbf{Z}^T \tilde{\mathbf{E}}^{-1} \mathbf{Z})^{-1} \\ (\mathbf{Z}^T \tilde{\mathbf{E}}^{-1} \mathbf{Z})^{-1} \mathbf{Z}^T \tilde{\mathbf{E}}^{-1} & -(\mathbf{Z}^T \tilde{\mathbf{E}}^{-1} \mathbf{Z})^{-1} \end{bmatrix},$$

where $\mathbf{P}$ is the oblique projection on the span of $\mathbf{Z}$ along the direction spanned by vectors $\mathbf{h}$ such that $\mathbf{Z}^T \tilde{\mathbf{E}}^{-1} \mathbf{h} = 0$. Hence,

$$\begin{pmatrix} \mathbf{E} \hat{\boldsymbol{\delta}} \\ \mathbf{Z} \hat{\mathbf{a}} \end{pmatrix} = \begin{bmatrix} \mathbf{E} \tilde{\mathbf{E}}^{-1} (\mathbf{I} - \mathbf{P}) \mathbf{Y} \\ \mathbf{P} \mathbf{Y} \end{bmatrix}.$$

Use $\hat{\mathbf{g}} = \mathbf{Z} \hat{\mathbf{a}} + \mathbf{E} \hat{\boldsymbol{\delta}}$ to obtain the desired result. $\square$

APPENDIX C. PROOF OF THEOREM 3.2

We start by introducing some useful notations and results. Let $\mathcal{X}$ and $\mathcal{Y}$ be Hilbert spaces with corresponding inner products $\langle \cdot, \cdot \rangle_{\mathcal{X}}$ and $\langle \cdot, \cdot \rangle_{\mathcal{Y}}$, and consider a linear operator $D : \mathcal{X} \mapsto \mathcal{Y}$. The norm of D is $\|D\|_{op} = \sup_{f \in \mathcal{X}, \|f\|_{\mathcal{X}}=1} \|Df\|_{\mathcal{Y}}$. When $\|D\|_{op} < \infty$, D is said to be bounded (or continuous), see Kress (1999, Chapter 2). Let D^* be the adjoint of D , defined as $D^* : \mathcal{Y} \mapsto \mathcal{X}$ such that $\langle Df, \psi \rangle_{\mathcal{Y}} = \langle f, D^* \psi \rangle_{\mathcal{X}}$ for any $(f, \psi) \in \mathcal{X} \times \mathcal{Y}$. When D is bounded, D^* always exists and $\|D\|_{op} = \|D^*\|_{op}$, see Kress (1999, Theorem 4.9). In what follows, we will repeatedly use the following properties: (a) $\|Df\|_{\mathcal{Y}} \leq \|D\|_{op} \|f\|_{\mathcal{X}}$ for any $f \in \mathcal{X}$, and (b) if C is another linear operator, then $\|CD\|_{op} \leq \|C\|_{op} \|D\|_{op}$, whenever the composition CD is well defined.

We divide the proof into several steps. In Step 1, we analyze the minimization problem at the population level. In Step 2, we analyze the problem at the sample level. In Step 3, we bound the norm of $\hat{h} - h_0$. In Step 4 and 5, we combine the results to first establish uniform consistency of $\hat{g}$ and its first derivative, second to obtain uniform rates of convergence. The proof relies on Lemmas S3.1 and S3.2, which are stated in the online supplement.

Step 1. From Assumption 3.2 and Bierens (1982),

$$g = 0 \Leftrightarrow E[g(Z)|W] = 0 \Leftrightarrow E[g(Z) \exp(\mathbf{i}W^T t)] = 0 \quad \forall t \in \mathbb{R}^p.$$

Hence, the null space of the linear mapping $g \mapsto E[g(Z) \exp(\mathbf{i}W^T \cdot)]$ only contains the zero element, and such a mapping is injective (one-to-one). This implies that $Ah = E[h(Z) \exp(\mathbf{i}W^T \cdot)]$ and $B\beta = E[(1, Z)\beta \exp(\mathbf{i}W^T \cdot)]$ are also injective.

Each $g \in \mathcal{G}$ can be uniquely written as $g(z) = (1, z)\beta + h(z)$, where $\beta = (g(0), g'(0))$, $h(z) = g(z) - g(0) - g'(0)z$, $h \in \mathcal{H}$. Hence, the intersection of the ranges of the operators A and B is the null function, since $Ah = B\beta$ iff $(1, z)\beta - h(z) = 0$.

Consider the problem

$$(A.7) \quad \min_{\beta, h} \|r - B\beta - Ah\|_\mu^2, \quad r = E[Y \exp(iW^T \cdot)],$$

where $\|\cdot\|_\mu$ is the L_μ^2 norm. If $g_0(z) = (1, z)\beta_0 + h_0(z)$, then (β_0, h_0) is the unique solution. We now obtain an explicit expression of (β_0, h_0) solving (A.7). Let P be the orthogonal projection operator of functions in L_μ^2 onto $\mathcal{R}(B)$ the range of B. Since B is defined on $\mathbb{R}^2$, its range $\mathcal{R}(B)$ is a linear finite dimensional space. As linear finite dimensional spaces are complete, see Kreyszig (1978, Theorem 2.4-2), $\mathcal{R}(B)$ is also linear and complete. By Kress (1999, Theorem 1.26), projection operators onto linear and complete spaces are well-defined, and so is P.

We now show that P writes as $B(B^*B)^{-1}B^*$, where B^* is the adjoint of B. As previously noted, B is injective and its null space is $\mathcal{N}(B) = \{0\}$. Then $\mathcal{N}(B^*B) = \mathcal{N}(B) = \{0\}$, B^*B is injective, and $(B^*B)^{-1}$ exists. As linear operators mapping $\mathbb{R}^2$ into $\mathbb{R}^2$ are uniquely characterized by second order matrices, see Kreyszig (1978, Section 2.9), B^*B is a second order matrix, as well as its inverse. Hence, the operator $B(B^*B)^{-1}B^* : L_\mu^2 \mapsto L_\mu^2$ is well-defined. For any $f \in L_\mu^2$ and $\beta \in \mathbb{R}^2$,

$$\langle f - B(B^*B)^{-1}B^*f, B\beta \rangle_\mu = \langle B^*f - B^*f, \beta \rangle = 0.$$

Hence, $f - B(B^*B)^{-1}B^*f \perp \mathcal{R}(B)$, and $B(B^*B)^{-1}B^*f$ indeed represents the projection of f onto $\mathcal{R}(B)$, see Kress (1999). Therefore, $P = B(B^*B)^{-1}B^*$.

Let $M = I - P$ be the orthogonal projection onto the orthogonal complement of $\mathcal{R}(B)$. Then

$$r = B\beta_0 + Ah_0 \Rightarrow Mr = MAh_0 = Th_0.$$

The operator $T = MA$ is injective, since the intersection of the ranges of A and B is the null function and A is injective. This yields

$$h_0 = T^{-1}Mr, \quad \beta_0 = (B^*B)^{-1}B^*(r - Ah_0).$$

Consider now the penalized problem

$$(A.8) \quad \min_{(\beta, h) \in \mathbb{R}^2 \times \mathcal{H}} \|r - Ah - B\beta\|_\mu^2 + \lambda \|h\|_{\mathcal{H}}^2.$$

Let us profile with respect to β . For any fixed h ,

$$\min_{\beta \in \mathbb{R}^2} \|r - Ah - B\beta\|_\mu^2 = \|r - Ah - P(r - Ah)\|_\mu^2 = \|Mr - Th\|_\mu^2.$$

We thus need to solve

$$\min_{h \in \mathcal{H}} \|Mr - Th\|_\mu^2 + \lambda \|h\|_{\mathcal{H}}^2.$$

From Lemma S3.2(a), T is compact, and thus bounded. A direct application of [Kress \(1999, Theorem 16.1\)](#) ensures that the unique solution h_λ satisfies $(T^*T + \lambda I)h_\lambda = T^*Mr$. Now, for any h ,

$$\lambda \|h\|_{\mathcal{H}}^2 \leq \lambda \|h\|_{\mathcal{H}}^2 + \|Th\|_{\mu}^2 = \lambda \langle h, h \rangle_{\mathcal{H}} + \langle h, T^*Th \rangle_{\mathcal{H}} = \langle h, (T^*T + \lambda I)h \rangle.$$

Hence, $(T^*T + \lambda I)$ is strictly coercive and has a bounded inverse by the Lax-Milgram Theorem, see [Kress \(1999, Theorem 13.26\)](#). Therefore,

$$(A.9) \quad h_\lambda = (T^*T + \lambda I)^{-1}T^*Mr.$$

Step 2. We study the minimization problem at the sample level and we obtain sample counterparts of the population objects of Step 1. Recall that $\hat{g}$ solves

$$(A.10) \quad \min_{g \in \mathcal{G}} \int \left| \frac{1}{n} \sum_{i=1}^n [Y_i - g((Z_i))] \exp(\mathbf{i}W_i^T t) \right|^2 \mu(dt) + \lambda \int_0^1 |g''(z)|^2 dz.$$

By Proposition 2.1, under Assumption 3.1, the solution $\hat{g}$ is unique with probability 1, and since each $g \in \mathcal{G}$ writes uniquely as $g(z) = (1, z)\beta + h(z)$, there is a unique $(\hat{\beta}, \hat{h})$ such that $\hat{g}(z) = (1, z)\hat{\beta} + \hat{h}(z)$. Define

$$(A.11) \quad \hat{A} : \mathcal{H} \mapsto L_{\mu}^2, \quad \hat{A}h = \frac{1}{n} \sum_{i=1}^n h(Z_i) \exp(\mathbf{i}W_i^T \cdot),$$

$$(A.12) \quad \hat{B} : \mathbb{R}^2 \mapsto L_{\mu}^2, \quad \hat{B}\beta = \frac{1}{n} \sum_{i=1}^n (1, Z_i)\beta \exp(\mathbf{i}W_i^T \cdot),$$

and $\hat{r} = (1/n) \sum_{i=1}^n Y_i \exp(\mathbf{i}W_i^T \cdot)$. The optimization problem (A.10) is equivalent to

$$(A.13) \quad \min_{(\beta, h) \in \mathbb{R}^2 \times \mathcal{H}} \|\hat{r} - \hat{A}h - \hat{B}\beta\|_{\mu}^2 + \lambda \|h\|_{\mathcal{H}}^2.$$

We will profile with respect to β , and to do so requires dealing with the orthogonal projection onto the range of $\hat{B}$. Let us proceed as in Step 1. First,

$$\|\hat{B}\beta\|_{\mu}^2 = 0 \Leftrightarrow \int \left| \frac{1}{n} \sum_{i=1}^n (1, Z_i)\beta \exp(\mathbf{i}W_i^T t) \right|^2 \mu(dt) = 0 \Leftrightarrow \beta^T \mathbf{Z}^T \mathbf{\Omega} \mathbf{Z} \beta = 0.$$

From Assumption 3.1(e), W has at least one continuous component, so that all W_i 's are different with probability 1, and thus $\mathbf{\Omega} > 0$ with probability 1 from the proof of Proposition 2.1. Hence, $\|\hat{B}\beta\|_{\mu}^2 = 0$ iff $\mathbf{Z}\beta = 0$. As $\mathbf{Z}$ has full column rank with probability 1 from Assumption 3.1, $\|\hat{B}\beta\|_{\mu}^2 = 0$ iff $\beta = 0$, and $\hat{B}$ is injective. Let $\hat{P}$ be the orthogonal projection onto the range of $\hat{B}$, which is well defined and can be expressed as $\hat{P} = \hat{B}(\hat{B}^* \hat{B})^{-1} \hat{B}^*$. Then,

$$\min_{\beta \in \mathbb{R}^2} \|\hat{r} - \hat{A}h - \hat{B}\beta\|_{\mu}^2 = \|\hat{r} - \hat{A}h - \hat{P}(\hat{r} - \hat{A}h)\|_{\mu} = \|\hat{M}\hat{r} - \hat{T}h\|_{\mu}^2,$$

where $\widehat{M} = I - \widehat{P}$ and $\widehat{T} = \widehat{M}\widehat{A}$. We thus need to solve

$$(A.14) \quad \min_{h \in \mathcal{H}} \|\widehat{M}\widehat{r} - \widehat{T}h\|_{\mu}^2 + \lambda \|h\|_{\mathcal{H}}^2.$$

From Lemma S3.2(e), $\widehat{T}$ is compact, and thus bounded. Thus, using a similar reasoning as in Step 1, the unique solution is

$$(A.15) \quad \widehat{h} = (\widehat{T}^* \widehat{T} + \lambda I)^{-1} \widehat{T}^* \widehat{M}\widehat{r},$$

which in turn yields

$$(A.16) \quad \widehat{\beta} = (\widehat{B}^* \widehat{B})^{-1} \widehat{B}^* (\widehat{r} - \widehat{A}\widehat{h}).$$

Step 3. We now prove that

$$(A.17) \quad \|\widehat{h} - h_0\|_{\mathcal{H}} = O_p \left(\frac{1}{\sqrt{n\lambda}} + \|h_{\lambda} - h_0\|_{\mathcal{H}} \right).$$

We consider the decomposition $\widehat{h} - h_0 = S_1 + S_2 + S_3 + S_4 + h_{\lambda} - h_0$, where

$$\begin{aligned} S_1 &= (T^* T + \lambda I)^{-1} T^* (\widehat{M}\widehat{r} - \widehat{T}h_0), \quad S_2 = (T^* T + \lambda I)^{-1} (\widehat{T}^* - T^*) (\widehat{M}\widehat{r} - \widehat{T}h_0), \\ S_3 &= \left[(\widehat{T}^* \widehat{T} + \lambda I)^{-1} - (T^* T + \lambda I)^{-1} \right] \widehat{T}^* (\widehat{M}\widehat{r} - \widehat{T}h_0), \quad S_4 = (\widehat{T}^* \widehat{T} + \lambda I)^{-1} \widehat{T}^* \widehat{T}h_0 - h_{\lambda}. \end{aligned}$$

We have

$$\|S_1\|_{\mathcal{H}} \leq \|(T^* T + \lambda I)^{-1} T^*\|_{op} \|\widehat{M}\widehat{r} - \widehat{T}h_0\|_{\mu} = O_p \left(\frac{1}{\sqrt{n\lambda}} \right).$$

Indeed, T is a compact operator from Lemma S3.2(a), $\|(T^* T + \lambda I)^{-1} T^*\|_{op} \leq c'/\sqrt{\lambda}$ from Lemma S3.1(b), and $\|\widehat{M}\widehat{r} - \widehat{T}h_0\|_{\mu} = O_p(1/\sqrt{n})$ from Lemma S3.2(h). Next,

$$\|S_2\|_{\mathcal{H}} \leq \|(T^* T + \lambda I)^{-1}\|_{op} \|\widehat{T}^* - T^*\|_{op} \|\widehat{M}\widehat{r} - \widehat{T}h_0\|_{\mu} = O_p \left(\frac{1}{n\lambda} \right),$$

as $\|(T^* T + \lambda I)^{-1}\|_{op} \leq c/\lambda$ from Lemma S3.1(a), $\|\widehat{T} - T\|_{op} = O_p(1/\sqrt{n})$ and $\|\widehat{M}\widehat{r} - \widehat{T}h_0\|_{\mu} = O_p(1/\sqrt{n})$ from Lemma S3.2(f) and (h). Next,

$$\begin{aligned} \|S_3\|_{\mathcal{H}} &\leq \|(\widehat{T}^* \widehat{T} + \lambda I)^{-1} \widehat{T}^* - (T^* T + \lambda I)^{-1} T^*\|_{op} \|\widehat{M}\widehat{r} - \widehat{T}h_0\|_{\mu} \\ &\leq \|(\widehat{T}^* \widehat{T} + \lambda I)^{-1} \widehat{T}^* - (T^* T + \lambda I)^{-1} T^* - (T^* T + \lambda I)^{-1} (\widehat{T}^* - T^*)\|_{op} \|\widehat{M}\widehat{r} - \widehat{T}h_0\|_{\mu} \\ &\leq \|(\widehat{T}^* \widehat{T} + \lambda I)^{-1} \widehat{T}^* - (T^* T + \lambda I)^{-1} T^*\|_{op} \|\widehat{M}\widehat{r} - \widehat{T}h_0\|_{\mu} \\ &\quad + \|(T^* T + \lambda I)^{-1}\|_{op} \|\widehat{T}^* - T^*\|_{op} \|\widehat{M}\widehat{r} - \widehat{T}h_0\|_{\mu} = O_p \left(\frac{1}{\sqrt{n\lambda}} \right). \end{aligned}$$

Indeed, $\widehat{T}$ and T are compact operators from Lemma S3.2(a) and (e), so $\|(\widehat{T}^* \widehat{T} + \lambda I)^{-1} \widehat{T}^* - (T^* T + \lambda I)^{-1} T^*\|_{op} \leq 2c'/\sqrt{\lambda}$ and $\|(T^* T + \lambda I)^{-1}\|_{op} \leq c/\lambda$ by Lemma S3.1. Moreover,

$\|\hat{T}^* - T^*\|_{op} = O_p(1/\sqrt{n})$ from Lemma S3.2(f). Finally,

$$\begin{aligned}
\|S_4\|_{\mathcal{H}} &= \|(\hat{T}^* \hat{T} + \lambda I)^{-1} \hat{T}^* \hat{T} h_0 - (T^* T + \lambda I)^{-1} T^* T h_0\|_{\mathcal{H}} \\
&= \|(\hat{T}^* \hat{T} + \lambda I)^{-1} (\hat{T}^* \hat{T} + \lambda I - \lambda I) h_0 - (T^* T + \lambda I)^{-1} (T^* T + \lambda I - \lambda I) h_0\|_{\mathcal{H}} \\
&= \|\lambda[(T^* T + \lambda I)^{-1} - (\hat{T}^* \hat{T} + \lambda I)^{-1}] h_0\|_{\mathcal{H}} \\
&= \|\lambda(\hat{T}^* \hat{T} + \lambda I)^{-1} [\hat{T}^* \hat{T} - T^* T] (T^* T + \lambda I)^{-1} h_0\|_{\mathcal{H}} \\
&= \|\lambda(\hat{T}^* \hat{T} + \lambda I)^{-1} [\hat{T}^* (\hat{T} - T) + (\hat{T}^* - T^*) T] (T^* T + \lambda I)^{-1} h_0\|_{\mathcal{H}} \\
&\leq \|(\hat{T}^* \hat{T} + \lambda I)^{-1} \hat{T}^*\|_{op} \|\hat{T} - T\|_{op} \|\lambda(T^* T + \lambda I)^{-1}\|_{op} \|h_0\|_{\mathcal{H}} \\
&\quad + \|\lambda(\hat{T}^* \hat{T} + \lambda I)^{-1}\|_{op} \|\hat{T}^* - T^*\|_{op} \|T(T^* T + \lambda I)^{-1}\|_{op} \|h_0\|_{\mathcal{H}} = O_p\left(\frac{1}{\sqrt{n\lambda}}\right).
\end{aligned}$$

Here we use that $\|\hat{T} - T\|_{op} = \|\hat{T}^* - T^*\|_{op} = O_p(1/\sqrt{n})$ from Lemma S3.2(f), and that $\|\lambda(T^* T + \lambda I)^{-1}\|_{op} \leq c$, $\|\lambda(\hat{T}^* \hat{T} + \lambda I)^{-1}\|_{op} \leq c$, and $\|(\hat{T}^* \hat{T} + \lambda I)^{-1} \hat{T}^*\|_{op} \leq c'/\sqrt{\lambda}$ from Lemma S3.1. Gathering results gives (A.17).

Step 4. We here show convergence of our estimators. Since T is injective from Step 1 and compact from Lemma S3.2(a), $\|(T^* T + \lambda I)^{-1} T^* T h - h\|_{\mathcal{H}} = o(1)$ for all h whenever $\lambda \rightarrow 0$, see Kress (1999, Definition 15.5 and Theorem 15.23). Hence $\|h_{\lambda} - h_0\|_{\mathcal{H}} = o(1)$. This and (A.17) yields $\|\hat{h} - h_0\|_{\mathcal{H}} = o_p(1)$ if in addition $n\lambda \rightarrow \infty$.

We now show that $\|\hat{\beta} - \beta_0\| = O_p\left(1/\sqrt{n} + \|\hat{h} - h_0\|_{\mathcal{H}}\right)$. From (A.16),

$$\begin{aligned}
\hat{\beta} - \beta_0 &= [(\hat{B}^* \hat{B})^{-1} - (B^* B)^{-1}] \hat{B}^* (\hat{r} - \hat{A} \hat{h}) + (B^* B)^{-1} [\hat{B}^* - B^*] (\hat{r} - \hat{A} \hat{h}) \\
&\quad + (B^* B)^{-1} B^* (\hat{r} - r) + (B^* B)^{-1} B^* (\hat{A} - A) \hat{h} - (B^* B)^{-1} B^* A (\hat{h} - h_0) \\
\Rightarrow \|\hat{\beta} - \beta_0\| &\leq \|(\hat{B}^* \hat{B})^{-1} - (B^* B)^{-1}\|_{op} \|\hat{B}^*\|_{op} \left(\|\hat{r}\|_{\mu} + \|\hat{A}\|_{op} \|\hat{h}\|_{\mathcal{H}}\right) \\
&\quad + \|(B^* B)^{-1}\|_{op} \|\hat{B}^* - B^*\|_{op} \left(\|\hat{r}\|_{\mu} + \|\hat{A}\|_{op} \|\hat{h}\|_{\mathcal{H}}\right) \\
&\quad + \|(B^* B)^{-1}\|_{op} \|B^*\|_{op} \left(\|\hat{r} - r\|_{\mu} + \|\hat{A} - A\|_{op} \|\hat{h}\|_{\mathcal{H}} + \|A\|_{op} \|\hat{h} - h_0\|_{\mathcal{H}}\right).
\end{aligned}$$

Lemma S3.1 ensures that $\|(\hat{B}^* \hat{B})^{-1} - (B^* B)^{-1}\|_{op}$, $\|\hat{B}^* - B^*\|_{op} = \|\hat{B} - B\|_{op}$, $\|\hat{r} - r\|_{\mu}$, and $\|\hat{A} - A\|_{op}$ all are $O_p(n^{-1/2})$. We have $\|B^*\|_{op} = \|B\|_{op} < \infty$, as B is a linear operator with finite dimensional domain, see Kreyszig (1978, Theorem 2.7-8), and $\|B^* B\|_{op} = \|B\|_{op}^2$. Similarly, $\|(B^* B)^{-1}\|_{op} < \infty$ as B is injective. From Lemma S3.2(a), A is compact and hence bounded, and from Lemma S3.2(d) $\|\hat{A}\|_{op} = O_p(1)$. From a similar reasoning, $\|\hat{B}^*\|_{op} = O_p(1)$. Also $\|\hat{h} - h_0\|_{\mathcal{H}} = o_p(1)$ implies $\|\hat{h}\|_{\mathcal{H}} = O_p(1)$. Combine these results to obtain that

$$\|\hat{\beta} - \beta_0\| = O_p\left(n^{-1/2} + \|\hat{h} - h_0\|_{\mathcal{H}}\right) = o_p(1).$$

Since $\widehat{g}(z) = (1, z)\widehat{\beta} + \widehat{h}(z)$, to show uniform consistency of $\widehat{g}$ and $\widehat{g}'$, it now suffices to show that $\sup_{z \in [0,1]} |\widehat{h}(z) - h_0(z)|$ and $\sup_{z \in [0,1]} |\widehat{h}'(z) - h'_0(z)|$ are bounded by $\|\widehat{h} - h_0\|_{\mathcal{H}}$. As for any $h \in \mathcal{H}$, $h'(z) = \int_0^z h''(t)dt$,

$$\sup_{z \in [0,1]} |\widehat{h}'(z) - h'_0(z)| \leq \sup_{z \in [0,1]} \int_0^z |\widehat{h}''(t) - h''_0(t)|dt \leq \int_0^1 |\widehat{h}''(t) - h''_0(t)|dt \leq \|\widehat{h} - h_0\|_{\mathcal{H}},$$

from Cauchy-Schwartz inequality. Since $h(z) = \int_0^z h'(t)dt$, a similar reasoning yields

$$\sup_{z \in [0,1]} |\widehat{h}(z) - h_0(z)| \leq \sup_{z \in [0,1]} \int_0^z |\widehat{h}'(t) - h'_0(t)|dt \leq \sup_{z \in [0,1]} |\widehat{h}'(z) - h'_0(z)|.$$

Step 5. We now obtain uniform convergence rates. Assumption 3.3 allows applying Proposition 3.11 in Carrasco et al. (2007) to the operator T and yields $\|h_\lambda - h_0\|_{\mathcal{H}} = O\left(\lambda^{\frac{\gamma \wedge 2}{2}}\right)$. Combining with the results of Step 3 gives

$$\|\widehat{h} - h_0\|_{\mathcal{H}} = O_p\left(\frac{1}{\sqrt{n\lambda}} + \lambda^{\frac{\gamma \wedge 2}{2}}\right) \text{ and } \|\widehat{\beta} - \beta_0\| = O_p\left(\frac{1}{\sqrt{n\lambda}} + \lambda^{\frac{\gamma \wedge 2}{2}}\right).$$

Use the same arguments as in Step 4 to obtain uniform convergence rates. $\square$

APPENDIX D. PROOF OF THEOREM 4.3

We first prove that Assumption 6.5 implies

- (a) D and A are injective,
- (b) $\mathcal{R}(A) \cap \mathcal{R}(D) = \{0\}$,
- (c) L is full column rank with probability approaching one,

where $\mathcal{R}(A)$ denotes the range of A , with A defined in (3.13), and $\mathcal{R}(D)$ denotes the range of D , with D is defined in (6.20).

The proof of condition (a) uses arguments contained in the proof of Step 1 of Theorem 3.2 in the main text. For completeness, we also provide it here. Given $\beta \in \mathbb{R}^2$ and $h \in \mathcal{H}$ (with $\mathcal{H}$ defined in Section 3), for $g(z) = (1, z)\beta + h(z)$ we have that $g \in \mathcal{G}$. So,

$$\begin{aligned} (\beta, \gamma) = 0, h = 0 & \Leftrightarrow E[X^T \gamma + (1, Z)\beta + h(Z)|W] = 0 \\ & \Leftrightarrow E[(X^T \gamma + (1, Z)\beta + h(Z)) \exp(iW^T t)] = 0, \quad \forall t \in \mathbb{R}^p \\ & \Leftrightarrow D(\gamma, \beta) + A h = 0, \end{aligned}$$

where the first equivalence follows from Assumption 6.5, the second equivalence from Bierens (1982), and the last equivalence from the definition of D and A . Since $A 0 = 0$, by the above display $D(\gamma, \beta) = 0$ implies $(\gamma, \beta) = 0$. So, D is injective. Similarly, the above display also ensures that A is injective. Thus, condition (a) is proved. To prove condition (b), consider an element belonging to $\mathcal{R}(A) \cap \mathcal{R}(D)$, say $A h = D(\gamma, \beta)$. Then,

$D(\gamma, \beta) + A(-h) = 0$ and by the above display, we get that $(\gamma, \beta) = 0$ and $h = 0$. So, condition (b) is proved. Let us finally show condition (c). Recall that $\mathbf{L}$ is the $n \times (q+2)$ matrix whose t th row is $(1, Z_i, X_i^T)$. Take (γ, β) such that $(1, Z)\beta + X^T\gamma = 0$. Then, $D(\gamma, \beta) = 0$, and by the injectivity of D , we obtain that $\gamma = 0$ and $\beta = 0$. Thus, $(1, Z, X^T)$ are linearly independent. This implies that $E[(1, Z, X^T)^T (1, Z, X^T)]$ is full rank. Since $(1/n)\mathbf{L}^T \mathbf{L} = \sum_{i=1}^n (1, Z_i, X_i^T)^T (1, Z_i, X_i^T) = E[(1, Z, X^T)^T (1, Z, X^T)] + o_p(1)$, we get that $\mathbf{L}^T \mathbf{L}$ is full rank with probability approaching one. So $\mathbf{L}$ is full column rank with probability approaching one and condition (c) is proved.

Given conditions (a), (b), and (c), the proof of Theorem 6.4 proceeds along the same arguments as the proof of Theorem 3.2 in the main text. $\square$

APPENDIX E. AUXILIARY LEMMAS

The following lemma is from Florens et al. (2011, Lemma A.1).

Lemma 5.1. *Consider two Hilbert spaces $\mathcal{X}$ and $\mathcal{Y}$ and a linear compact operator $K : \mathcal{X} \mapsto \mathcal{Y}$. Then there are universal constants c and c' such that (a) $\|\lambda(\lambda I + K^* K)^{-1}\|_{op} \leq c$; (b) $\|(\lambda I + K^* K)^{-1} K^*\|_{op} \leq \frac{c'}{\sqrt{\lambda}}$.*

Lemma 5.2. *Under Assumptions 3.2 and 3.1, the following holds: (a) The operators A and T are compact; (b) $\|\hat{B} - B\|_{op} = O_p(n^{-1/2})$; (c) $\|(\hat{B}^* \hat{B})^{-1} - (B^* B)^{-1}\|_{op} = O_p(n^{-1/2})$ and $\|\hat{M} - M\|_{op} = O_p(n^{-1/2})$; (d) $\|\hat{A} - A\|_{op} = O_p(n^{-1/2})$; (e) The operator $\hat{T}$ is compact; (f) $\|\hat{T} - T\|_{op} = O_p(n^{-1/2})$; (g) $\|\hat{r} - r\|_\mu = O_p(n^{-1/2})$; (h) $\|\hat{M}\hat{r} - \hat{T}h_0\|_\mu = O_p(n^{-1/2})$.*

Proof of Lemma 5.2: (a). Let us show A is compact by compact embedding. Define $\tilde{A}$ as the extension of A to $L^2([0, 1])$, where $L^2([0, 1])$ is the space of real-valued squared-integrable functions on $[0, 1]$, i.e. $\tilde{A}h = E[h(Z)e^{iW^T}]$ for any $h \in L^2([0, 1])$. For all $h \in \mathcal{H}$, we have $h(z) = \int_0^z \int_0^x h''(t) dt dx$, so that

$$\begin{aligned} \|h\|_{L^2[0,1]}^2 &= \int_0^1 |h(z)|^2 dz \leq \sup_{z \in [0,1]} |h(z)| \leq \sup_{z \in [0,1]} \left| \int_0^z h'(t) dt \right| \leq \sup_{t \in [0,1]} |h'(t)| \\ &\leq \sup_{t \in [0,1]} \left| \int_0^t h''(u) du \right| \leq \int_0^1 |h''(u)| du \leq \|h\|_{\mathcal{H}}^2. \end{aligned}$$

Therefore, every bounded set on $(\mathcal{H}, \|\cdot\|_{\mathcal{H}})$ is also a bounded set on $(L^2([0, 1]), \|\cdot\|)$. Hence, compactness of $\tilde{A}$ implies compactness of A . Now for any $h \in L^2[0, 1]$,

$$(\tilde{A}h)(t) = E[h(Z) E[\exp(iW^T t)|Z]] = \int h(z) E[\exp(iW^T t)|Z = z] f_Z(z) dz,$$

where

$$\int |E[\exp(iW^T t)|Z = z]|^2 \mu(t) f_Z(z) dt dz \leq 1,$$

as $|\exp(\mathbf{i}\cdot)| \leq 1$. Since $\tilde{A}$ is an integral operator whose kernel is Hilbert-Schmidt, i.e. squared integrable, we can apply [Busby et al. \(1972, Proposition 2.1\)](#) to conclude that $\tilde{A}$ is compact.

Let us now show that T is compact. The range of B , $\mathcal{R}(B)$, is finite dimensional, linear, and closed. P is the orthogonal projection onto $\mathcal{R}(B)$, and is thus bounded by [Kress \(1999, Theorem 13.3\)](#). Hence, $M = I - P$ is bounded as well. Since $T = MA$ is the composition of a bounded and a compact operator, it is compact by [Kress \(1999, Theorems 2.14 and 2.16\)](#).

(b). For $\beta \in \mathbb{R}^2$, we have

$$\|(\hat{B} - B)\beta\|_\mu^2 = \int |(E_n - E)[\exp(\mathbf{i}W^\top t)(1, Z)]\beta|^2 \mu(dt),$$

where E_n denotes the empirical expectation. By the Cauchy-Schwarz inequality,

$$E\|\hat{B} - B\|_{op}^2 \leq E \left[\int (|(E_n - E)[\exp(\mathbf{i}W^\top t)]|^2 + |(E_n - E)[Z \exp(\mathbf{i}W^\top t)]|^2) \mu(dt) \right].$$

Since data are i.i.d.,

$$\begin{aligned} E[|(E_n - E)[Z \exp(\mathbf{i}W^\top t)]|^2] &= E \left[\left| n^{-1} \sum_{i=1}^n Z_i \exp(\mathbf{i}W_i^\top t) - E[Z \exp(\mathbf{i}W^\top t)] \right|^2 \right] \\ &= \text{Var} \left(\frac{1}{n} \sum_{i=1}^n Z_i \exp(\mathbf{i}W_i^\top t) \right) \\ &= n^{-1} (E[|Z \exp(\mathbf{i}W^\top t)|^2] - |E[Z \exp(\mathbf{i}W^\top t)]|^2) = O(n^{-1}), \end{aligned}$$

as $|Z \exp(\mathbf{i}W^\top t)| \leq 1$ for all $t \in \mathcal{T}$. Similarly, $E[|(E_n - E)[\exp(\mathbf{i}W^\top t)]|^2] = O(n^{-1})$. This implies $E\|\hat{B} - B\|_{op}^2 = O(n^{-1})$, and by Markov's inequality, $\|\hat{B} - B\|_{op}^2 = O_p(n^{-1})$.

(c). From [Kreyszig \(1978, Theorem 2.7-8\)](#), as B is a linear operator with a finite dimensional domain, it is bounded, and $\|B\|_{op} < \infty$. Also $\|B^*\|_{op} = \|B\|_{op}$ and $\|B^*B\|_{op} = \|B\|_{op}^2$. The operator B^*B maps $\mathbb{R}^2$ into $\mathbb{R}^2$, and is thus a matrix. From (b), $\|\hat{B}\|_{op}$ and $\|\hat{B}^*\|_{op}$ are $o_p(1)$, and

$$\begin{aligned} \|B^*B - \hat{B}^*\hat{B}\|_{op} &= \|(B^* - \hat{B}^*)B + \hat{B}^*(B - \hat{B})\|_{op} \\ &\leq \|B^* - \hat{B}^*\|_{op} \|B\|_{op} + \|\hat{B}^*\|_{op} \|B - \hat{B}\|_{op} = o_p(n^{-1/2}). \end{aligned}$$

Since B is injective, B^*B is invertible, $(B^*B)^{-1}$ exists and is bounded. By the continuous mapping theorem, $\|(\hat{B}^*\hat{B})^{-1} - (B^*B)^{-1}\|_{op} = o_p(1)$. Hence $\|(\hat{B}^*\hat{B})^{-1}\|_{op} \leq \|(\hat{B}^*\hat{B})^{-1} - (B^*B)^{-1}\|_{op} + \|(B^*B)^{-1}\|_{op} = o_p(1)$. Moreover,

$$\begin{aligned} \|(\hat{B}^*\hat{B})^{-1} - (B^*B)^{-1}\|_{op} &= \|(\hat{B}^*\hat{B})^{-1}(B^*B - \hat{B}^*\hat{B})(B^*B)^{-1}\|_{op} \\ &\leq \|(\hat{B}^*\hat{B})^{-1}\|_{op} \|B^*B - \hat{B}^*\hat{B}\|_{op} \|(B^*B)^{-1}\|_{op} = o_p(n^{-1/2}). \end{aligned}$$

For the difference between $M = I - B(B^* B)^{-1} B^*$ and $\widehat{M} = I - \widehat{B}(\widehat{B}^* \widehat{B})^{-1} \widehat{B}^*$, we have

$$\begin{aligned} \|\widehat{M} - M\|_{op} &= \|(\widehat{B} - B)(\widehat{B}^* \widehat{B})^{-1} \widehat{B}^* + B[(\widehat{B}^* \widehat{B})^{-1} - (B^* B)^{-1}] \widehat{B}^* + B(B^* B)^{-1} [\widehat{B}^* - B^*]\|_{op} \\ &\leq \|\widehat{B} - B\|_{op} \|(\widehat{B}^* \widehat{B})^{-1}\|_{op} \|\widehat{B}^*\|_{op} + \|B\|_{op} \|(\widehat{B}^* \widehat{B})^{-1} - (B^* B)^{-1}\| \|\widehat{B}^*\|_{op} \\ &\quad + \|B\|_{op} \|(B^* B)^{-1}\|_{op} \|\widehat{B}^* - B^*\|_{op} = o_p(n^{-1/2}). \end{aligned}$$

(d). Recall that for any $h \in \mathcal{H}$, $h(z) = \int_0^z \int_0^x h''(u) du dx$. Thus,

$$\begin{aligned} (\widehat{A}h)(t) &= \frac{1}{n} \sum_{i=1}^n h(Z_i) \exp(\mathbf{i}W_i^T t) = \frac{1}{n} \sum_{i=1}^n \int_0^{Z_i} \int_0^x h''(u) du dx \exp(\mathbf{i}W_i^T t) \\ &= \int_{[0,1]^2} h''(u) \left[\frac{1}{n} \sum_{i=1}^n \mathbf{1}(0 < u < x) \mathbf{1}(0 < x < Z_i) \exp(\mathbf{i}W_i^T t) \right] du dx \\ &= \int_0^1 h''(u) \left[\int_0^1 \frac{1}{n} \sum_{i=1}^n \mathbf{1}(u < x < Z_i) \exp(\mathbf{i}W_i^T t) dx \right] du \\ (A.18) \quad &= \int_0^1 h''(u) \widehat{k}(u, t) du, \end{aligned}$$

where $\widehat{k}(u, t)$ is defined implicitly above. Exchanging the empirical measure with the population probability and using the same steps as above yield

$$(A h)(t) = \int_0^1 h''(u) \left[\int_0^1 E\{\mathbf{1}(u < x < Z) \exp(\mathbf{i}W^T t)\} dx \right] = \int_0^1 h''(u) k(u, t) du.$$

where $k(u, t) = E\widehat{k}(u, t)$ is defined implicitly above. Next,

$$\begin{aligned} \|\widehat{A} - A\|_{op}^2 &= \sup_{h \in \mathcal{H} \mid \|h\|_{\mathcal{H}}=1} \|\widehat{A}h - Ah\|_{\mu}^2 = \sup_{h \in \mathcal{H} \mid \|h\|_{\mathcal{H}}=1} \int \left| (\widehat{A}h)(t) - (Ah)(t) \right|^2 \mu(dt) \\ &= \sup_{h \in \mathcal{H} \mid \|h\|_{\mathcal{H}}=1} \int \left| h''(u) \left[\widehat{k}(u, t) - k(u, t) \right] \right|^2 du \mu(dt) \\ &\leq \sup_{h \in \mathcal{H} \mid \|h\|_{\mathcal{H}}=1} \int \left(\int_0^1 |h''(u)|^2 du \int_0^1 \left| \widehat{k}(u, t) - k(u, t) \right|^2 du \right) \mu(dt) \\ &= \sup_{h \in \mathcal{H} \mid \|h\|_{\mathcal{H}}=1} \|h\|_{\mathcal{H}} \int \left(\int_0^1 \left| \widehat{k}(u, t) - k(u, t) \right|^2 du \right) \mu(dt) \\ &= \int_{[0,1] \times \mathbb{R}^q} \left| \widehat{k}(u, t) - k(u, t) \right|^2 du \otimes \mu(dt) \\ \Rightarrow E\|\widehat{A} - A\|_{op}^2 &\leq \int_{[0,1] \times \mathbb{R}^q} E \left| \widehat{k}(u, t) - k(u, t) \right|^2 du \otimes \mu(dt). \end{aligned}$$

Now,

$$\begin{aligned}
E \left| \widehat{k}(u, t) - k(u, t) \right|^2 &= E \left| \int_0^1 (E_n - E) \mathbf{1}(u < x < Z) \exp(\mathbf{i} W^T t) dx \right|^2 \\
&\leq \int_0^1 E \left| (E_n - E) \mathbf{1}(u < x < Z) \exp(\mathbf{i} W^T t) \right|^2 dx \\
&= \int_0^1 \text{Var} \left(\frac{1}{n} \sum_{i=1}^n \mathbf{1}(u < x < Z_i) \exp(\mathbf{i} W_i^T t) \right) dx \\
&= \frac{1}{n} \int_0^1 \text{Var} \left(\mathbf{1}(u < x < Z) \exp(\mathbf{i} W^T t) \right) dx = O(n^{-1}).
\end{aligned}$$

Use Markov's inequality to obtain the desired result.

(e). By reasoning as in the proof of (a), compactness of $\widehat{T} = \widehat{M}\widehat{A}$ follows if $\widehat{M}$ is bounded and $\widehat{A}$ is compact. The first claim is shown following similar arguments as in (a). To obtain compactness of $\widehat{A}$, we will use Theorem 8.1-4 in Kreyszig (1978) stating that a bounded operator with a finite dimensional range is compact. As

$$\widehat{A}h = \sum_{i=1}^n h(Z_i) \frac{1}{n} \exp(W_i^T \cdot) \in \text{Span} \left(\frac{1}{n} \exp(W_1^T \cdot), \dots, \frac{1}{n} \exp(W_n^T \cdot) \right),$$

the range of $\widehat{A}$ is finite dimensional for all n . Moreover, using (A.18)

$$\|\widehat{A}h\|_\mu^2 = \int \left| \int_0^1 h''(u) \widehat{k}(u, t) du \right|^2 \mu(dt) \leq \|h\|_{\mathcal{H}}^2 \sup_{u,t} |\widehat{k}(u, t)|^2 du \leq \|h\|_{\mathcal{H}}^2,$$

as $|\widehat{k}(u, t)| \leq 1$. Hence, $\|\widehat{A}\|_{op} \leq 1$, and $\widehat{A}$ is compact.

(f). Since $\widehat{T} - T = (\widehat{M} - M)\widehat{A} + M(\widehat{A} - A)$, the result follows from (c), (d), and the fact that M is bounded.

(g). The proof is analogous to the proof of (b).

(h). Write $\widehat{M}\widehat{r} - M r = (\widehat{M} - M)\widehat{r} + M(\widehat{r} - r)$, and use $\|\widehat{M} - M\|_{op} = O_p(n^{-1/2})$, $\|\widehat{r} - r\|_\mu = O_p(n^{-1/2})$, and $\|M\|_{op} < \infty$ from previous items to obtain $\|\widehat{M}\widehat{r} - M r\|_\mu = O_p(n^{-1/2})$. Use (f) above to get $\|(\widehat{M}\widehat{r} - \widehat{T}h_0) - (M r - T h_0)\|_\mu = O_p(n^{-1/2})$, and note that $M r - T h_0 = 0$.

□

[†]DEPARTMENT OF ECONOMICS, KU LEUVEN, NAAMSESTRAAT 69, 3000 LEUVEN, BELGIUM.
Email address: jad.beyhum@gmail.com

[‡]ENSAE-CREST, 5 AVENUE HENRY LE CHATELIER 91120 PALAISEAU, FRANCE.
Email address: elia.lapenta@ensae.fr

*TOULOUSE SCHOOL OF ECONOMICS, UNIVERSITÉ TOULOUSE CAPITOLE, 1 ESPLANADE DE L'UNIVERSITÉ, 31000 TOULOUSE, FRANCE.

Email address: pascal.lavergne@ut-capitole.fr