

# Dynamical invariant based shortcut to equilibration in open quantum systems

Mohamed Boubakour,<sup>1,\*</sup> Shimpei Endo,<sup>2,3</sup> Thomás Fogarty,<sup>1</sup> and Thomas Busch<sup>1</sup>

<sup>1</sup>*Quantum Systems Unit, Okinawa Institute of Science and Technology Graduate University, Okinawa 904-0495, Japan*

<sup>2</sup>*Department of Engineering Science, The University of Electro-Communications, Tokyo 182-8585, Japan*

<sup>3</sup>*Department of Physics, Tohoku University, Sendai 980-8578, Japan*

(Dated: April 17, 2024)

We propose using the dynamical invariants, also known as the Lewis–Riesenfeld invariants, to speed-up the equilibration of a driven open quantum system. This allows us to reverse engineer the time-dependent master equation that describes the dynamics of the open quantum system and systematically derive a protocol that realizes a shortcut to equilibration. The method does not require additional constraints on the timescale of the dynamics beside the Born–Markov approximation and can be generically applied to boost single particle quantum engines significantly. We demonstrate it with the damped harmonic oscillator, and show that our protocol can achieve a high-fidelity control in shorter timescales than simple non-optimized protocols. We find that the system is heated during the dynamics to speed-up the equilibration, which can be considered as an analogue of the Mpemba effect in quantum control.

## I. INTRODUCTION

Understanding and controlling open quantum systems is a major challenge for the exploration of quantum phenomena in the presence of dissipative effects, the deterministic preparation of quantum states, and the development of quantum devices [1–4]. A particular interesting and relevant question is how to accelerate the equilibration of open quantum systems, which, for example, has applications in enhancing the performance of quantum heat engines [5–8]. Recently this question has attracted some attention and various techniques based on approaches such as optimal control [9–12], linear response theory [13], techniques inspired by shortcuts to adiabaticity [14, 15] and reverse engineering [16–18] have been developed. Some of these have also been successfully applied to design optimized quantum Carnot cycles [19] or an optimized quantum Otto cycle [20, 21], both of which show that a potential quantum advantage can be achieved.

The challenge of accelerating the equilibration of an open quantum system comes with the challenge of describing its dynamics in a practical way for control. Even in the well-established framework of using master equations to describe open quantum systems [22], their derivation can be a very difficult task for an underlying general time-dependent Hamiltonian. For example, a shortcut to equilibration in Refs. [16, 18] was realized by deriving a non-adiabatic time-dependent master equation in the inertial limit [23, 24], which assumes small variations of the adiabatic parameter of the system. While this allows to obtain the Lindblad operators explicitly, the resulting driving protocol can be restricted.

In this work, we propose a shortcut to equilibration realized by using the dynamical invariant (DI) to describe the dynamics of driven open quantum systems.

Also known as the Lewis–Riesenfeld invariant, it was originally introduced by the latter to solve the time-dependent dynamics of closed quantum systems [25] and recently has been applied to shortcuts to adiabaticity in open quantum systems to assist in minimizing dissipation [26]. However, they can also be used to derive a time-dependent master equation in a comprehensive way, without restrictions on the driving protocol, and which also gives a clear picture of the influence of the driving protocol on the dissipative effects [27–29]. Using the master equation, we present a general formulation of the shortcut between the equilibrium states of the initial and final Hamiltonian through reverse engineering.

To show the power of this approach, we apply this technique to the damped harmonic oscillator. This is a well-known model that can describe atomic and optomechanical systems, and is commonly used to explore the operation of quantum heat engines [30–32]. Moreover, dissipative harmonic oscillator systems have recently attracted additional attention as an interesting framework to describe Bose polaron systems [33, 34]. These systems are promising platforms to explore quantum thermodynamics phenomena like heat transport between mesoscopic quantum gases [35] or thermometry [36]. We find that our protocol outperforms non-optimized protocols, showing much higher fidelity and shorter time to reach the target equilibrium state. Notably, the effective temperature of the system in the middle of the dynamics needs to increase to achieve faster equilibration.

## II. DYNAMICAL INVARIANT BASED TIME-DEPENDENT MASTER EQUATION

A driven, open quantum system is in general described by a Hamiltonian of the form

$$H(t) = H_S(t) + H_B + H_I, \quad (1)$$

where  $H_S(t)$  is the time-dependent Hamiltonian of the system of interest and  $H_B$  is the Hamiltonian of the en-

\* mohamed.boubakour@oist.jp

vironment. In our case, the environment corresponds to a thermal bath at a given temperature  $T$  and thus its state is given by  $\rho_B = Z_B^{-1} \exp\left(-\frac{H_B}{k_B T}\right)$ . The final term describes the interaction between the system and the environment,  $H_I = \sum_k A_k \otimes B_k$ , where the  $A_k$  act on the system and the  $B_k$  act on the bath. To derive a solvable time-dependent master equation, we use the Born approximation that assumes that the coupling strength is sufficiently weak to neglect correlations between the system and the bath during the dynamics. The state of the system and the bath can then be written as a product of the reduced states,  $\rho(t) \approx \rho_S(t) \otimes \rho_B$ . We also use the Markov approximation i.e. the correlations inside the bath decay much faster than any other timescale of the system. This allows one to derive a master equation that is local in time. After tracing out the bath, we obtain a Redfield master equation in the interaction picture of the form [22, 37]

$$\begin{aligned} \frac{d\tilde{\rho}_S(t)}{dt} = & -\frac{1}{\hbar^2} \sum_{k,l} \int_0^\infty B_{kl}(\tau) \left[ \tilde{A}_k(t), \tilde{A}_l(t-\tau) \tilde{\rho}_S(t) \right] \\ & - B_{lk}(-\tau) \left[ \tilde{A}_k(t), \tilde{\rho}_S(t) \tilde{A}_l(t-\tau) \right] d\tau, \end{aligned} \quad (2)$$

where  $B_{kl}(\tau) = \text{Tr}_B \left( \tilde{B}_k(\tau) B_l \rho_B \right)$  is the two-point correlation function of the bath and the tilde indicates operators in the interaction picture. In the presence of a time-dependent system Hamiltonian, the evaluation of  $\tilde{A}_k(t)$  can be challenging since there is no general procedure that allows to calculate the time evolution operator of the system  $U_S(t) = T_{\leftarrow} \exp\left(-\frac{i}{\hbar} \int_0^t H_S(\tau) d\tau\right)$  (where  $T_{\leftarrow}$  is the time-ordering operator).

However, this problem can be solved by using the DI. Let us consider the closed dynamics of the system  $i\hbar \partial_t |\psi(t)\rangle = H_S(t) |\psi(t)\rangle$ . A DI of the Hamiltonian  $H_S(t)$  is a Hermitian operator  $I$  obeying [25]

$$\frac{dI(t)}{dt} = \frac{\partial I(t)}{\partial t} + \frac{1}{i\hbar} [I(t), H_S(t)] = 0. \quad (3)$$

This means that the solution of the closed dynamics can be written as a linear combination of the instantaneous eigenstates of the DI  $|\psi(t)\rangle = \sum_n c_n e^{i\alpha_n(t)} |\phi_n(t)\rangle$  where the dynamical phases are given by  $\alpha_n(t) = \frac{1}{\hbar} \int_0^t \langle \phi_n(\tau) | i\hbar \frac{\partial}{\partial \tau} - H(\tau) | \phi_n(\tau) \rangle d\tau$ . The time-evolution operator of the system can therefore also be obtained from the eigenstates of the DI as

$$U_S(t) = \sum_n e^{i\alpha_n(t)} |\phi_n(t)\rangle \langle \phi_n(0)|, \quad (4)$$

which in turn allows one to calculate the operators acting on the system in the interaction picture as [27]

$$\tilde{A}_k(t) = \sum_{m,n} e^{i(\alpha_n(t) - \alpha_m(t))} \langle \phi_m(t) | A_k | \phi_n(t) \rangle F_{mn}, \quad (5)$$

with  $F_{mn} = |\phi_m(0)\rangle \langle \phi_n(0)|$ . The operators can therefore be written as products of time-dependent scalar functions that contain the information on the driving protocol, and time-independent operators  $F_{mn}$ . Those operators are jump operators constructed with the DI eigenstate suggesting that the dissipative part of the open dynamics will involve transitions of the system between those states.

Focusing on the case of the time-dependent harmonic oscillator, the Hamiltonian of the system is given by  $H_S(t) = \frac{p^2}{2m} + \frac{1}{2} m \omega(t)^2 x^2$ . A DI of this Hamiltonian is [25, 38]

$$I(t) = \frac{(b(t)p - m\dot{b}(t)x)^2}{2m} + \frac{1}{2} m \omega_0^2 \left( \frac{x}{b(t)} \right)^2, \quad (6)$$

where  $\omega_0 = \omega(0)$  and  $b(t)$  corresponds to a dimensionless scaling function that satisfies the Ermakov equation

$$\ddot{b}(t) + \omega^2(t)b(t) = \frac{\omega_0^2}{b^3(t)}. \quad (7)$$

We notice that the invariant has the structure of a harmonic oscillator with a constant frequency  $\omega_0$ , a position  $x/b(t)$  and momentum  $b(t)p - m\dot{b}(t)x$ . Thus the eigenstates can be obtained by using the standard ladder operators.

### III. SHORTCUT TO EQUILIBRATION FOR THE DAMPED HARMONIC OSCILLATOR

#### A. Time-dependent master equation

In what follows, we design a shortcut to equilibration (STE) protocol for the time-dependent damped harmonic oscillator (DHO). The Hamiltonian of the bath is given by  $H_B = \sum_n \hbar \omega_n (b_n^\dagger b_n + \frac{1}{2})$  and the interaction is described in the rotating wave approximation by  $H_I = \sum_n g_n (a^\dagger b_n + a b_n^\dagger)$ , where the  $g_n$  are constant coupling strengths between the particle and the  $n$ -th mode of the bath. By using the DI of Eq. (6), we describe the dynamics of the driven system by deriving the following Lindblad master equation in the interaction picture (see Appendix A for details)

$$\begin{aligned} \frac{d\tilde{\rho}_S(t)}{dt} = & -\frac{i}{\hbar} \left[ \tilde{H}_{LS}(t), \tilde{\rho}_S(t) \right] \\ & + \frac{|D(t)|^2}{2\hbar^2} \gamma_+(\tilde{\omega}(t)) \left( a_I \tilde{\rho}_S(t) a_I^\dagger - \frac{1}{2} \{ a_I^\dagger a_I, \tilde{\rho}_S(t) \} \right) \\ & + \frac{|D(t)|^2}{2\hbar^2} \gamma_-(\tilde{\omega}(t)) \left( a_I^\dagger \tilde{\rho}_S(t) a_I - \frac{1}{2} \{ a_I a_I^\dagger, \tilde{\rho}_S(t) \} \right), \end{aligned} \quad (8)$$

where  $\tilde{H}_{LS}(t)$  is the time-dependent Lamb shift in the interaction picture and  $D(t) = b(t) + 1/b(t) + i\dot{b}(t)/\omega_0$ . The time-dependent decay rates characterizing the emission and absorption are given by  $\gamma_{\pm}(\tilde{\omega}(t)) =$

$\pi J(\tilde{\omega}(t)) (1 + n(\tilde{\omega}(t)))$  and  $\gamma(\tilde{\omega}(t)) = \gamma_+(\tilde{\omega}(t))e^{-\frac{\hbar\tilde{\omega}(t)}{k_B T}}$ , with  $J(\omega) = \sum_n g_n^2 \delta(\omega - \omega_n)$  being the bath spectral density function and  $n(\omega) = (e^{\frac{\hbar\omega}{k_B T}} - 1)^{-1}$  the Planck distribution. The time-dependent decay rates  $\gamma$  and  $\gamma_+$  are characterized by an effective Bohr frequency  $\tilde{\omega}(t) = \omega_0/b(t)^2$ . Therefore the scaling function  $b(t)$  sets the frequency at which the absorption and emission of excitations from the particle occur during the dynamics. The Lindblad operators  $a_I$  and  $a_I^\dagger$  correspond respectively to the annihilation and creation operator of the invariant at  $t = 0$ . However, if one considers a driving protocol with a continuous start from the initial Hamiltonian  $[H_S(0), I(0)] = 0$ , we recover the creation and annihilation operators of the particle,  $a_I = a$  ( $a_I^\dagger = a^\dagger$ ). The range of validity of the master equation imposes constraints on the driving protocol  $\omega(t)$ . The Markov approximation implies  $\omega(t)^{-1} \gg \tau_B$  where  $\tau_B$  is the characteristic decay time of the bath two-point correlation functions. We also used a secular approximation that implies  $\int_0^t \tilde{\omega}(\tau) d\tau \gg \tilde{\omega}(t)\tau_B$  (more details are given in Appendix A).

### B. Formulation of the shortcut protocol and reverse engineering

We design a shortcut protocol for an isothermal stroke where the trap frequency of the system is driven from  $\omega_0$  to  $\omega_f$  while coupled to a bath at temperature  $T$ . In principle the system needs to be driven slowly in order to remain at equilibrium with the bath at all times during the process. In order to speed up this process our STE protocol will consist of mimicking the stroke at finite time by connecting the initial and final desired equilibrium states. Initially the particle is at equilibrium with the frequency  $\omega(0) = \omega_0$  i.e.  $\rho_S(0) = Z_0^{-1} e^{-H_S(0)/k_B T}$ . The final desired state is an equilibrium state of the particle with the bath at the trapping frequency  $\omega(t_f) = \omega_f$  i.e.  $\rho_S(t_f) = Z_f^{-1} e^{-H_S(t_f)/k_B T}$ , where  $t_f$  is the duration of the protocol. We focus on the compression stroke  $\omega_f > \omega_0$  but the expansion can also be done in the same way.

We know that the state of the particle will be Gaussian during the dynamics since the Hamiltonian is quadratic [39]. Therefore it can be fully determined by the expectation values  $\langle a^\dagger a \rangle(t) = \text{Tr}(a^\dagger a \rho_S(t))$  and  $\langle a^2 \rangle(t) = \text{Tr}(a^2 \rho_S(t))$  characterizing the excitation and the squeezing of the particle. The equations describing their evolution during the driving protocol are obtained in the interaction picture from the master equation (8) as

$$\frac{d\langle \tilde{a}^\dagger \tilde{a} \rangle}{dt} = \frac{\pi}{2\hbar^2} |D(t)|^2 J(\tilde{\omega}(t)) (n(\tilde{\omega}(t)) - \langle \tilde{a}^\dagger \tilde{a} \rangle), \quad (9)$$

$$\frac{d\langle \tilde{a}^2 \rangle}{dt} = -\frac{\pi}{2\hbar^2} |D(t)|^2 J(\tilde{\omega}(t)) \langle \tilde{a}^2 \rangle. \quad (10)$$

Since the particle is initially in a Gibbs state with  $\langle \tilde{a}^2 \rangle(0) = 0$ , it follows from Eq. (10) that  $\langle \tilde{a}^2 \rangle = 0$




Figure 1. (a) Fidelity between the final state and the target state as a function of  $t_f$ . (b) Profile of the trap frequency for the STE protocol as a function of time for different protocol durations. The black dashed line shows the reference ramp. (c) Coherence generated during the dynamics of the STE as a function of time. (d) Effective temperature of the particle during the STE protocol as a function of time. The final compression is  $\omega_f = 3\omega_0$  and the bath temperature is  $T = \hbar\omega_0/k_B$ . The spectral density function is an Ohmic distribution with an abrupt cut-off  $J(\omega) = \gamma\omega\Theta(\Lambda - \omega)$  with  $\gamma = \hbar^2/500$  and  $\Lambda = 100\omega_0$ , and the number of particles in the bath is  $N = 600$ .

at any time during the dynamics. Thus, the protocol is described by the differential equation (9) alone. Furthermore, the initial and target state are equilibrium states which implies  $\langle \tilde{a}^\dagger \tilde{a} \rangle(0) = (e^{\frac{\hbar\omega_0}{k_B T}} - 1)^{-1}$ ,  $\langle \tilde{a}^\dagger \tilde{a} \rangle(t_f) = (e^{\frac{\hbar\omega_f}{k_B T}} - 1)^{-1}$  and  $\frac{d\langle \tilde{a}^\dagger \tilde{a} \rangle(0)}{dt} = \frac{d\langle \tilde{a}^\dagger \tilde{a} \rangle(t_f)}{dt} = 0$ . In addition to these boundary conditions, we impose  $\frac{d^2\langle \tilde{a}^\dagger \tilde{a} \rangle(0)}{dt^2} = \frac{d^2\langle \tilde{a}^\dagger \tilde{a} \rangle(t_f)}{dt^2} = 0$  to ensure a smooth evolution of the system between the initial state and the target state.

The protocol can now be found by reverse-engineering Eq. (9) to obtain the scaling function  $b(t)$ , which in turn allows one to obtain the trap frequency from the Ermakov equation (7). For that, we assume that the evolution of  $\langle \tilde{a}^\dagger \tilde{a} \rangle$  is given by a 5-th order polynomial function that ensures that the boundary conditions obtained above are satisfied. The boundary conditions on  $\langle \tilde{a}^\dagger \tilde{a} \rangle$  combined with Eq. (9) also imply  $b(0) = 1$ ,  $b(t_f) = \sqrt{\omega_0/\omega_f}$  and  $\dot{b}(0) = \dot{b}(t_f) = 0$ . We also obtain additional boundary conditions from the Ermakov equation  $\ddot{b}(0) = \ddot{b}(t_f) = 0$ . We can thus consider a 6-th order polynomial ansatz for the scaling function  $b(t) = \sum_{n=0}^6 a_n (t/t_f)^n$ , in which the first 6 coefficients allow to satisfy the above boundary conditions. The 6-th order term can then ensure that the scaling function connects the initial state and the target state through Eq. (9). The coefficient  $a_6$  is simply found by maximizing the fidelity between the target state and

the state of the particle at the end of the protocol. Using this ansatz for the scaling function allows to avoid the divergence issues that one can encounter when solving Eq. (9), where at  $t = 0$  and  $t = t_f$  the average excitation  $\langle a^\dagger a \rangle$  and the Planck distribution  $n(\tilde{\omega}(t))$  converge to the same values.

### C. Properties of the shortcut

To quantify the performance of the shortcut, we calculate the fidelity between the target state  $\rho_T$  and the state of the particle at the end of the protocol

$$F(\rho_S(t_f), \rho_T) = \text{Tr} \left( \sqrt{\sqrt{\rho_T} \rho_S(t_f) \sqrt{\rho_T}} \right)^2. \quad (11)$$

Since the states of the particle and the bath remain Gaussian, we can use an efficient numerical method to solve the dynamics of the total Hamiltonian (1) during the protocol. In fact, this can be done by time-evolving the covariance matrix of the system plus the bath, which corresponds to a  $(2N+2) \times (2N+2)$  matrix (where  $N$  is the number of particles in the bath), with the Heisenberg equations of motion [40]. This allows us to see when the validity of the master equation actually breaks down for a given set of bath parameters (number of particles, spectral density function) and when the shortcut therefore does not work anymore. To demonstrate the benefit of our shortcut, we compare it with two simple protocols: the sudden quench  $\omega_q(t > 0) = \omega_f$  and a reference ramp described by a polynomial function  $\omega_r(t) = \omega_0 + 10\Delta\omega(t/t_f)^3 - 15\Delta\omega(t/t_f)^4 + 6\Delta\omega(t/t_f)^5$  with  $\Delta\omega = \omega_f - \omega_0$ . For the DHO, the quench protocol is characterized by an asymptotic exponential convergence of the fidelity to one as  $F \approx 1 - e^{-t_f/\tau_q}$  where  $\tau_q$  is a characteristic time related to the decay rates [16].

The fidelity as a function of  $t_f$  is shown in Fig. 1(a). The STE outperforms the sudden quench and the reference ramp that also shows an asymptotic exponential behavior but with a longer characteristic time than the sudden quench. However, one can see that for short durations, the fidelity for the STE protocol collapses because at shorter times the description provided by the master equation deviates from the exact dynamics. The fidelity reaches approximately the value 0.999 around  $t_f \approx 16/\omega_0$  and then keeps increasing to one.

In order to obtain physical insights on the STE protocol the profile of the trap frequency  $\omega(t)$  is shown in Fig. 1(b) for different protocol durations  $t_f$  and compared to the reference ramp (black dashed line). We see that the profiles are quite different: while for the reference ramp the frequency increases monotonically toward  $\omega_f$ , in the STE protocol the particle is driven to large trap frequency values at intermediate times before decreasing to reach the final frequency at  $t = t_f$ . Notably, for faster protocols the particle is driven to higher frequency values. Inversely, for larger  $t_f$ , the amplitude of the trap

frequency decreases and we observe that the STE protocol gets closer to the reference ramp when  $t_f$  approaches the adiabatic limit.

Such a trap frequency profile necessarily implies non-equilibrium features in the dynamics. To quantify them, we calculate the coherence in the system during the STE, which has been suggested to play a key role in the control of open quantum systems [19]. We define it as the change of entropy between the diagonal part of the state and the full density matrix in the instantaneous eigenenergy basis [41]

$$C(t) = S(\rho_{diag}(t)) - S(\rho_S(t)), \quad (12)$$

where  $S(\rho) = -\text{Tr}(\rho \log(\rho))$  is the von Neumann entropy. The coherence is shown in Fig. 1(c). One can see that its profile reflects the results we have shown before: coherence is generated in the system when the trap frequency is changing. It vanishes when the maximum trap frequency is reached, and increases again when the trap frequency decreases toward the final value. There is no coherence remaining in the system at the end of the protocol since the particle reaches the target Gibbs state. Moreover the amount generated during the dynamics increases for faster protocols. This also allows to explain why the quench protocol works better than the reference ramp. Indeed, coherence causes transitions of the particle between its eigenstates and can be used as a catalysis that helps to accelerate the thermalization of an open quantum system. A controlled manipulation of coherence allows to reach the new equilibrium state and this is what the STE achieves. Even though the non-optimal protocols do not reach the target state, the sudden quench will always perform better than a smooth ramp that generates much less non-adiabatic excitations. Notably, this is in contrast with shortcuts to adiabaticity protocols which require the suppression of such excitations to reach the desired target state [42].

Finally we characterize the dynamics of the STE by calculating the effective temperature of the system. During the dynamics, the state of the particle can be written in the interaction picture as  $\tilde{\rho}_S(t) = Z(t)^{-1} \sum_n e^{-\epsilon(t)n} |\phi_n(0)\rangle \langle \phi_n(0)|$  with

$$\langle \tilde{a}^\dagger \tilde{a} \rangle(t) = (e^{\epsilon(t)} - 1)^{-1}, \quad (13)$$

and the effective partition function is given by  $Z(t) = (1 - e^{-\epsilon(t)})^{-1}$ . Thus, back to the Schrödinger picture the Hamiltonian is always diagonal in the instantaneous eigenbasis of the DI  $\rho_S(t) = Z(t)^{-1} \sum_n e^{-\epsilon(t)n} |\phi_n(t)\rangle \langle \phi_n(t)|$ , which shows that similarly to the closed dynamics, the eigenstates of the DI give us the states that the system will explore for a given protocol. We then simply define the effective temperature of the system as

$$T_{\text{eff}}(t) = \frac{\hbar\omega(t)}{k_B\epsilon(t)}. \quad (14)$$

Defining a temperature for non-equilibrium states can be useful in many cases [43, 44] and is defined in our case

from the effective Planck distribution Eq. (13). Therefore besides characterizing how far from equilibrium the system is during the protocol, it is also related to both the energy and entropy of the particle. A high effective temperature is therefore associated with a state of high energy and mixedness. The effective temperature of the particle during the STE is shown in Fig. 1(d). It deviates significantly from the bath temperature before returning to it at the end of the protocol. More interestingly, the particle is driven to states that are effectively hotter, and the faster the shortcut is, the hotter the state of the particle is. While the shortcut is designed for an isothermal compression, which corresponds to a cooling process, the strategy adopted by the STE actually consists of warming up the particle in order to cool it down faster. This is reminiscent of the Mpemba effect [45], an empirical phenomenon where a hot liquid can freeze faster than a cold liquid. Recently, the Mpemba effect has been discussed and predicted for a quantum dot coupled to two reservoirs [46]. Here, we observe a similar feature to the thermal Mpemba effect in the context of driven open quantum systems, with the same behavior being previously observed in the effective temperature for an STE of a two-level system [18].

#### IV. CONCLUSION

In the present work, we have presented results that pave the way for improving the control of driven open quantum systems and the performance of quantum heat engines. We have shown that dynamical invariants can be a powerful tool for describing and accelerating the equilibration for these systems, and that besides the Born-Markov approximation, it lacks additional restrictions on the timescale of the dynamics. Our work also brings new physical interpretations of the dynamical invariant. Indeed the scaling function  $b(t)$ , that fully characterizes the invariant, sets both the driving protocol and the decay rates. This allows to derive protocols that modify both the unitary part and the dissipative part of the dynamics. Therefore an experimental implementation is feasible as one only needs to control the trapping potential unlike the method proposed in [15] where one also

needs to change the decay rates in a specific way. The STE protocol is characterized by a manipulation of the coherence that drives the particle to hotter states when designed for an isothermal compression. Our observation resonates with the thermal Mpemba effect and a rigorous formulation of this phenomenon in the context of driven open quantum systems would be an interesting direction to take in the future.

It is also an interesting avenue to design shortcuts and address the question whether the geometric bound [47–49] can be reached by minimizing the dissipated work. While we have considered the isothermal stroke in this work, one can consider different strokes, or start from non-equilibrium states. It would be also possible to design shortcuts for applications other than ones thermodynamics. For example, our method could be used to design fast and robust protocols for quantum gates [3, 50]. Also recently, a similar approach has been used to quickly generate entangled states in a double two-level system [51]. Finally, an important extension would be to go beyond the single particle problem and optimize the equilibration of interacting many-body states. While not an easy task, two possible directions are within reach: two particles with short-range interactions [52] and the hardcore Tonks–Girardeau limit [53]. Beside having well-known analytical results, both systems have showed enhanced performances compared to non-interacting quantum engines [54–56], paving the way for fully-optimized many-body quantum heat engines.

#### ACKNOWLEDGMENTS

This work was supported by the Okinawa Institute of Science and Technology Graduate University, and used the computing resources of the Scientific Computing and Data Analysis section at OIST. The authors also acknowledge support from a SHINKA grant from OIST and Tohoku University. SE is supported by JSPS KAKENHI Grant Numbers JP21H00116 and JP22K03492. TF acknowledges support from JSPS KAKENHI Grant Number JP23K03290. TF and TB are also supported by JST Grant Number JPMJPF2221.

- 
- [1] C. Brif, R. Chakrabarti, and H. Rabitz, Control of quantum phenomena: past, present and future, *New J. Phys.* **12**, 075008 (2010).
  - [2] C. P. Koch, Controlling open quantum systems: tools, achievements, and limitations, *Journal of Physics: Condensed Matter* **28**, 213001 (2016).
  - [3] S. Kallush, R. Dann, and R. Kosloff, Controlling the uncontrollable: Quantum control of open-system dynamics, *Science Advances* **8**, eadd0828 (2022), <https://www.science.org/doi/pdf/10.1126/sciadv.add0828>.
  - [4] Koch, Christiane P., Boscain, Ugo, Calarco, Tommaso, Dirr, Gunther, Philipp, Stefan, Glaser, Stefan J., Kosloff, Ronnie, Montangero, Simone, Schulte-Herbrüggen, Thomas, Sugny, Dominique, and Wilhelm, Frank K., Quantum optimal control in quantum technologies. strategic report on current status, visions and goals for research in Europe, *EPJ Quantum Technol.* **9**, 19 (2022).
  - [5] R. Kosloff and A. Levy, Quantum Heat Engines and Refrigerators: Continuous Devices, *Annual Review of Physical Chemistry* **65**, 365 (2014).

- [6] S. Vinjanampathy and J. Anders, Quantum thermodynamics, *Contemporary Physics* **57**, 545 (2016).
- [7] N. M. Myers, O. Abah, and S. Deffner, Quantum thermodynamic devices: From theoretical proposals to experimental reality, *AVS Quantum Science* **4**, 027101 (2022), [https://pubs.aip.org/avs/aqs/article-pdf/doi/10.1116/5.0083192/16494008/027101\\_1\\_online.pdf](https://pubs.aip.org/avs/aqs/article-pdf/doi/10.1116/5.0083192/16494008/027101_1_online.pdf).
- [8] G. Kurizki and A. G. Kofman, *Thermodynamics and Control of Open Quantum Systems* (Cambridge University Press, 2022).
- [9] V. Mukherjee, A. Carlini, A. Mari, T. Caneva, S. Montangero, T. Calarco, R. Fazio, and V. Giovannetti, Speeding up and slowing down the relaxation of a qubit by optimal control, *Phys. Rev. A* **88**, 062326 (2013).
- [10] N. Suri, F. C. Binder, B. Muralidharan, and S. Vinjanampathy, Speeding up thermalisation via open quantum systems variational optimisation, *Eur. Phys. J. Spec. Top.* **227**, 203 (2018).
- [11] V. Cavina, A. Mari, A. Carlini, and V. Giovannetti, Optimal thermodynamic control in open quantum systems, *Phys. Rev. A* **98**, 012139 (2018).
- [12] R. Xu, Reinforcement learning approach to shortcuts between thermodynamic states with minimum entropy production, *Phys. Rev. E* **105**, 054123 (2022).
- [13] N. Pancotti, M. Scandi, M. T. Mitchison, and M. Perarnau-Llobet, Speed-ups to isothermality: Enhanced quantum thermal machines through control of the system-bath coupling, *Phys. Rev. X* **10**, 031015 (2020).
- [14] T. Villazon, A. Polkovnikov, and A. Chandran, Swift heat transfer by fast-forward driving in open quantum systems, *Phys. Rev. A* **100**, 012126 (2019).
- [15] S. Alipour, A. Chenu, A. T. Rezakhani, and A. del Campo, Shortcuts to Adiabaticity in Driven Open Quantum Systems: Balanced Gain and Loss and Non-Markovian Evolution, *Quantum* **4**, 336 (2020).
- [16] R. Dann, A. Tobalina, and R. Kosloff, Shortcut to equilibration of an open quantum system, *Phys. Rev. Lett.* **122**, 250402 (2019).
- [17] L. Dupays, I. L. Egusquiza, A. del Campo, and A. Chenu, Superadiabatic thermalization of a quantum oscillator by engineered dephasing, *Phys. Rev. Res.* **2**, 033178 (2020).
- [18] R. Dann, A. Tobalina, and R. Kosloff, Fast route to equilibration, *Phys. Rev. A* **101**, 052102 (2020).
- [19] R. Dann and R. Kosloff, Quantum signatures in the quantum Carnot cycle, *New J. Phys.* **22**, 013055 (2020).
- [20] A. Das and V. Mukherjee, Quantum-enhanced finite-time Otto cycle, *Phys. Rev. Research* **2**, 033083 (2020).
- [21] A. Pedram, S. C. Kadioğlu, A. Kabakçioğlu, and O. E. Müstecaplıoğlu, A quantum otto engine with shortcuts to thermalization and adiabaticity, *New Journal of Physics* **25**, 113014 (2023).
- [22] H.-P. Breuer and F. Petruccione, *The theory of Open Quantum Systems* (Oxford University Press, 2002).
- [23] R. Dann, A. Levy, and R. Kosloff, Time-dependent markovian quantum master equation, *Phys. Rev. A* **98**, 052129 (2018).
- [24] R. Dann and R. Kosloff, Inertial theorem: Overcoming the quantum adiabatic limit, *Phys. Rev. Res.* **3**, 013064 (2021).
- [25] H. R. Lewis and W. B. Riesenfeld, An exact quantum theory of the time-dependent harmonic oscillator and of a charged particle in a time-dependent electromagnetic field, *J. Math. Phys.* **10**, 1458 (1969), <https://doi.org/10.1063/1.1664991>.
- [26] L. M. Cangemi, H. Espinós, R. Puebla, E. Torrontegui, and A. Levy, Control of open quantum systems via dynamical invariants (2023), [arXiv:2311.13164 \[quant-ph\]](https://arxiv.org/abs/2311.13164).
- [27] S. L. Wu, X. L. Huang, and X. X. Yi, Driven Markovian master equation based on the Lewis-Riesenfeld-invariant theory, *Phys. Rev. A* **106**, 052217 (2022).
- [28] S. L. Wu, X. Y. Zhang, and X. X. Yi, Dynamical invariants of open quantum systems, *Phys. Rev. A* **92**, 062122 (2015).
- [29] D.-W. Luo, P. V. Pyshkin, C.-H. Lam, T. Yu, H.-Q. Lin, J. Q. You, and L.-A. Wu, Dynamical invariants in a non-markovian quantum-state-diffusion equation, *Phys. Rev. A* **92**, 062127 (2015).
- [30] J. Roßnagel, S. T. Dawkins, K. N. Tolazzi, O. Abah, E. Lutz, F. Schmidt-Kaler, and K. Singer, A single-atom heat engine, *Science* **352**, 325 (2016).
- [31] V. Blickle and C. Bechinger, Realization of a micrometre-sized stochastic heat engine, *Nature Physics* **8**, 143 (2012).
- [32] R. Kosloff and Y. Rezek, The Quantum Harmonic Otto Cycle, *Entropy* **19**, 136 (2017).
- [33] A. Lampo, S. H. Lim, M. Á. García-March, and M. Lewenstein, Bose polaron as an instance of quantum Brownian motion, *Quantum* **1**, 30 (2017).
- [34] A. Lampo, M. A. García March, and M. Lewenstein, eds., *Quantum Brownian Motion Revisited* (Springer Cham, 2019).
- [35] C. Charalambous, M. A. Garcia-March, M. Mehboudi, and M. Lewenstein, Heat current control in trapped Bose-Einstein condensates, *New J. Phys.* **21**, 083037 (2019).
- [36] M. Mehboudi, A. Lampo, C. Charalambous, L. A. Correa, M. A. García-March, and M. Lewenstein, Using polarons for sub-nk quantum nondemolition thermometry in a Bose-Einstein condensate, *Phys. Rev. Lett.* **122**, 030403 (2019).
- [37] A. REDFIELD, The theory of relaxation processes\*  
\*this work was started while the author was at harvard university, and was then partially supported by joint services contract n5ori-76, project order i., in *Advances in Magnetic Resonance*, Advances in Magnetic and Optical Resonance, Vol. 1, edited by J. S. Waugh (Academic Press, 1965) pp. 1-32.
- [38] X. Chen, A. Ruschhaupt, S. Schmidt, A. del Campo, D. Guéry-Odelin, and J. G. Muga, Fast Optimal Frictionless Atom Cooling in Harmonic Traps: Shortcut to Adiabaticity, *Phys. Rev. Lett.* **104**, 063002 (2010).
- [39] J. Eisert and M. B. Plenio, Introduction to the basics of entanglement theory in continuous-variable systems, *International Journal of Quantum Information* **01**, 479 (2003), <https://doi.org/10.1142/S0219749903000371>.
- [40] Ángel Rivas, A. D. K. Plato, S. F. Huelga, and M. B. Plenio, Markovian master equations: a critical study, *New J. Phys.* **12**, 113032 (2010).
- [41] T. Baumgratz, M. Cramer, and M. B. Plenio, Quantifying coherence, *Phys. Rev. Lett.* **113**, 140401 (2014).
- [42] D. Guéry-Odelin, A. Ruschhaupt, A. Kiely, E. Torrontegui, S. Martínez-Garaot, and J. G. Muga, Shortcuts to adiabaticity: Concepts, methods, and applications, *Rev. Mod. Phys.* **91**, 045001 (2019).
- [43] J. Casas-Vázquez and D. Jou, Temperature in non-equilibrium states: a review of open problems and cur-

rent proposals, *Reports on Progress in Physics* **66**, 1937 (2003).

- [44] A. Puglisi, A. Sarracino, and A. Vulpiani, Temperature in and out of equilibrium: A review of concepts, tools and attempts, *Physics Reports* **709-710**, 1 (2017), temperature in and out of equilibrium: a review of concepts, tools and attempts.
- [45] E. B. Mpemba and D. G. Osborne, Cool?, *Physics Education* **4**, 172 (1969).
- [46] A. K. Chatterjee, S. Takada, and H. Hayakawa, Quantum Mpemba effect in a quantum dot with reservoirs, *Phys. Rev. Lett.* **131**, 080402 (2023).
- [47] P. Salamon and R. S. Berry, Thermodynamic length and dissipated availability, *Phys. Rev. Lett.* **51**, 1127 (1983).
- [48] M. Scandi and M. Perarnau-Llobet, Thermodynamic length in open quantum systems, *Quantum* **3**, 197 (2019).
- [49] G. Li, J.-F. Chen, C. P. Sun, and H. Dong, Geodesic path for the minimal energy cost in shortcuts to isothermality, *Phys. Rev. Lett.* **128**, 230603 (2022).
- [50] D. Turyansky, O. Ovdatt, R. Dann, Z. Aqua, R. Kosloff, B. Dayan, and A. Pick, Inertial geometric quantum logic gates (2023), [arXiv:2303.13674 \[quant-ph\]](https://arxiv.org/abs/2303.13674).
- [51] W. Ma, X. L. Huang, and S. L. Wu, Dynamics of a driven open double two-level system and its entanglement generation, *Phys. Rev. A* **107**, 032409 (2023).
- [52] T. Busch, B.-G. Englert, K. Rzażewski, and M. Wilkens, Two Cold Atoms in a Harmonic Trap, *Foundations of Physics* **28**, 549 (1998).
- [53] M. Girardeau, Relationship between Systems of Impenetrable Bosons and Fermions in One Dimension, *J. Math. Phys.* **1**, 516 (1960).
- [54] J. Jaramillo, M. Beau, and A. del Campo, Quantum supremacy of many-particle thermal machines, *New J. Phys.* **18**, 075019 (2016).
- [55] T. Fogarty and T. Busch, A many-body heat engine at criticality, *Quantum Science and Technology* **6**, 015003 (2020).
- [56] M. Boubakour, T. Fogarty, and T. Busch, Interaction-enhanced quantum heat engine, *Phys. Rev. Res.* **5**, 013088 (2023).

### Appendix A: Time-dependent master equation of the damped harmonic oscillator

We start from the Redfield equation given by Eq. (2) applied to the damped harmonic oscillator

$$\begin{aligned} \frac{d\tilde{\rho}_S(t)}{dt} = & -\frac{1}{\hbar^2} \int_0^\infty B_{12}(\tau) [\tilde{a}^\dagger(t), \tilde{a}(t-\tau)\tilde{\rho}_S(t)] \\ & - B_{21}(-\tau) [\tilde{a}^\dagger(t), \tilde{\rho}_S(t)\tilde{a}(t-\tau)] \\ & + B_{21}(\tau) [\tilde{a}(t), \tilde{a}^\dagger(t-\tau)\tilde{\rho}_S(t)] \\ & - B_{12}(-\tau) [\tilde{a}(t), \tilde{\rho}_S(t)\tilde{a}^\dagger(t-\tau)] d\tau, \end{aligned} \quad (\text{A1})$$

where the bath two-point correlation functions are given by

$$\begin{aligned} B_{12}(\tau) &= \sum_n g_n^2 \text{Tr}_B(\tilde{b}_n(\tau)b_n^\dagger\rho_B) = \sum_n e^{-i\omega_n\tau} g_n^2 (1+n(\omega_n)) \\ &= \int_0^\infty e^{-i\omega\tau} J(\omega)(1+n(\omega))d\omega, \\ B_{21}(\tau) &= \sum_n g_n^2 \text{Tr}_B(\tilde{b}_n^\dagger(\tau)b_n\rho_B) = \sum_n e^{i\omega_n\tau} g_n^2 n(\omega_n) \\ &= \int_0^\infty e^{i\omega\tau} J(\omega)n(\omega)d\omega. \end{aligned} \quad (\text{A2})$$

Now we need to evaluate the ladder operators of the particle in the interaction picture. Since they are conjugate, we can just focus on the annihilation operator, which we can write with the position and momentum operators in the interaction picture as

$$\tilde{a}(t) = \sqrt{\frac{m\omega_0}{2\hbar}} \left( \tilde{x}(t) + i\frac{\tilde{p}(t)}{m\omega_0} \right). \quad (\text{A3})$$

We use the invariant (6) to evaluate the operator in the interaction picture. Since the invariant is a harmonic oscillator with a position  $x/b(t)$  and momentum  $\Pi = b(t)p - m\dot{b}(t)x$ , we can express the position and momentum of the particle in terms of the instantaneous ladder operators of the invariant that we denote  $a_{I_t}$  and  $a_{I_t}^\dagger$

$$\begin{aligned} x &= b(t)\sqrt{\frac{\hbar}{2m\omega_0}}(a_{I_t} + a_{I_t}^\dagger), \\ p &= \frac{\Pi}{b(t)} + m\dot{b}(t)\frac{x}{b(t)} = \sqrt{\frac{\hbar m\omega_0}{2}} \left( C(t)a_{I_t} + C^*(t)a_{I_t}^\dagger \right), \end{aligned} \quad (\text{A4})$$

with the complex function

$$C(t) = \frac{\dot{b}(t)}{\omega_0} - \frac{i}{b(t)}. \quad (\text{A5})$$

We deduce that the annihilation operator of the harmonic oscillator in the Schrödinger picture is related to the ladder operators of the invariant through the following Bogoliubov transformation

$$a = \frac{1}{2} \left( D_1(t)a_{I_t} + D_2^*(t)a_{I_t}^\dagger \right), \quad (\text{A6})$$

where

$$D_{1,2}(t) = b(t) \pm \frac{1}{b(t)} \pm i\frac{\dot{b}(t)}{\omega_0}. \quad (\text{A7})$$

We now need to calculate the ladder operators of the invariant in the interaction picture. This is easily done by using the time-evolution operator written with the

eigenstates of the invariant (Eq. (4)) and using the expression of the dynamical phase for the harmonic oscillator  $\alpha_n(t) = -\omega_0(n+1/2) \int_0^t 1/b(\tau)^2 d\tau$

$$\begin{aligned} \tilde{a}_{I_t}(t) &= \sum_{n,m} e^{i(\alpha_m(t)-\alpha_n(t))} |\phi_n(0)\rangle \langle \phi_n(t)| a_{I_t} |\phi_m(t)\rangle \langle \phi_m(0)| \\ &= \sum_{n,m} e^{i(\alpha_m(t)-\alpha_n(t))} \sqrt{m} \delta_{n,m-1} |\phi_n(0)\rangle \langle \phi_m(0)| \\ &= e^{-i\varphi(t)} \sum_n \sqrt{n+1} |\phi_n(0)\rangle \langle \phi_{n+1}(0)| = e^{-i\varphi(t)} a_{I_0}, \end{aligned} \quad (\text{A8})$$

where the phase  $\varphi$  is given by

$$\begin{aligned} [\tilde{a}^\dagger(t), \tilde{a}(t-\tau)\tilde{\rho}_S(t)] &= \frac{1}{4} \left( e^{-i(\varphi(t)-\varphi(t-\tau))} D_2(t) D_2^*(t-\tau) [a_I, a_I^\dagger \tilde{\rho}_S(t)] + e^{-i(\varphi(t)+\varphi(t-\tau))} D_2(t) D_1(t-\tau) [a_I, a_I \tilde{\rho}_S(t)] \right. \\ &\quad \left. + e^{i(\varphi(t)+\varphi(t-\tau))} D_1^*(t) D_2^*(t-\tau) [a_I^\dagger, a_I^\dagger \tilde{\rho}_S(t)] + e^{i(\varphi(t)-\varphi(t-\tau))} D_1^*(t) D_1(t-\tau) [a_I^\dagger, a_I \tilde{\rho}_S(t)] \right). \end{aligned} \quad (\text{A11})$$

The integral in the Redfield equation is dominated by the bath two-point correlation function that rapidly decays with a characteristic time  $\tau_B$ . The decay time is given by the cut-off of the bath  $\tau_B \sim \Lambda^{-1}$ . Based on the Markov approximation, the decay time must be much smaller than the typical timescale of the system given by  $\omega(t)^{-1}$  i.e  $\tau_B \ll \omega(t)^{-1}$ . We can thus use the first order approximation of the phase in the integral  $\varphi(t-\tau) \approx \varphi(t) - \tilde{\omega}(t)\tau$ . We also make the zero-th order approximation  $D_i(t-\tau) \approx D_i(t)$  meaning that the variations of the scaling function  $b(t)$  and its derivative are negligible in the time window  $[0, \tau_B]$ . Formally, it implies  $\tau_B \ll \left| \frac{D_i(t)}{\dot{D}_i(t)} \right|$ . This approximation can be reformulated as  $\tau_B \ll \tau_D$  where we introduce a driving timescale  $\tau_D = \min_{i,t} \left| \frac{D_i(t)}{\dot{D}_i(t)} \right|$  [23]. Let us remark that this approx-

$$\int_0^\infty B_{12}(\tau) [\tilde{a}^\dagger(t), \tilde{a}(t-\tau)\tilde{\rho}_S(t)] d\tau \approx \frac{|D_2(t)|^2}{4} \int_0^\infty B_{12}(\tau) e^{-i\tilde{\omega}(t)\tau} d\tau [a_I, a_I^\dagger \tilde{\rho}_S(t)] + \frac{|D_1(t)|^2}{4} \int_0^\infty B_{12}(\tau) e^{i\tilde{\omega}(t)\tau} d\tau [a_I^\dagger, a_I \tilde{\rho}_S(t)]. \quad (\text{A12})$$

The integrals can be calculated by using the well-known result  $\int_0^\infty e^{i\omega\tau} d\tau = \pi\delta(\omega) + iP(1/\omega)$  where  $P$  denotes the principal value. After combining the different terms and few lines of algebra, we obtain the time-dependent master equation for the damped harmonic oscillator given by Eq. (8), where we replaced  $D_1(t)$  by

$$\varphi(t) = \int_0^t \frac{\omega_0}{b(\tau)^2} d\tau = \int_0^t \tilde{\omega}(\tau) d\tau. \quad (\text{A9})$$

For the next we will use  $a_I$  ( $a_I^\dagger$ ) instead of  $a_{I_0}$  ( $a_{I_0}^\dagger$ ) to denote the ladder operators of the invariant at  $t=0$ . Now we can obtain an explicit expression of the annihilation operator of the particle in the interaction picture

$$\tilde{a}(t) = \frac{1}{2} \left( D_1(t) e^{-i\varphi(t)} a_I + D_2^*(t) e^{i\varphi(t)} a_I^\dagger \right). \quad (\text{A10})$$

We can insert Eq. (A10) in the Redfield equation (A1). We only explicitly write it down for the first commutator in the right-hand side of Eq.(A1) since the same treatment can be straightforwardly done for the other terms. After expanding the commutator, we obtain

imation is not necessary to derive the master equation, however it allows to simplify the reverse-engineering for the shortcut.

The last approximation we will use is the secular approximation. We neglect the non-secular terms, to derive a master equation in Lindblad form and ensure that the state of the system remains physical. This means that the non-secular contributions contain fast oscillating terms that average to zero. This implies  $\varphi(t) + \varphi(t-\tau) \gg \varphi(t) - \varphi(t-\tau)$ . By using a first order expansion and the Markov approximation, we obtain  $\varphi(t) \gg \tilde{\omega}(t)\tau_B$  i.e  $\int_0^t \tilde{\omega}(\tau) d\tau \gg \tilde{\omega}(t)\tau_B$ .

Taking account of the different approximations in Eq. (A11), we obtain

$D(t)$ . The time-dependent Lamb shift is given by

$$\begin{aligned} \tilde{H}_{LS}(t) &= \frac{\hbar}{4} \left( |D_1(t)|^2 P \int_0^\infty \frac{J(\omega)}{\tilde{\omega}(t) - \omega} d\omega \right. \\ &\quad \left. - |D_2(t)|^2 P \int_0^\infty \frac{J(\omega)}{\tilde{\omega}(t) + \omega} d\omega \right). \end{aligned} \quad (\text{A13})$$