

Global existence and uniqueness of strong solutions to the 2D nonhomogeneous primitive equations with density-dependent viscosity

Quansen Jiu,^{a,*} Lin Ma^{a,†} and Fengchao Wang^{b,‡}

^a School of Mathematical Sciences, Capital Normal University, Beijing 100048, P. R. China.

^b College of Mathematics and Physics, Beijing University of Chemical Technology, Beijing 100029, P.R. China.

Abstract: This paper is concerned with an initial-boundary value problem of the two-dimensional inhomogeneous primitive equations with density-dependent viscosity. The global well-posedness of strong solutions is established, provided the initial horizontal velocity is suitably small, that is, $\|\nabla u_0\|_{L^2} \leq \eta_0$ for suitably small $\eta_0 > 0$. The initial data may contain vacuum. The proof is based on the local well-posedness and the blow-up criterion proved in [15], which states that if T^* is the maximal existence time of the local strong solutions (ρ, u, w, P) and $T^* < \infty$, then

$$\sup_{0 \leq t < T^*} (\|\nabla \rho(t)\|_{L^\infty} + \|\nabla^2 \rho(t)\|_{L^2} + \|\nabla u(t)\|_{L^2}) = \infty.$$

To complete the proof, it is required to make an estimate on a key term $\|\nabla u_t\|_{L_t^1 L_\Omega^2}$. We prove that it is bounded and could be as small as desired under certain smallness conditions, by making use of the regularity result of hydrostatic Stokes equations and some careful time weighted estimates.

Key Words: Nonhomogeneous primitive equations; Global well-posedness; Density-dependent viscosity

Mathematics Subject Classification: 35Q35, 35D35, 35B30, 35Q86, 76D03, 86A10

1 Introduction

In this paper, we consider the global existence and uniqueness of strong solutions to the

*E-mail: jiuqs@mail.cnu.edu.cn.

†E-mail: malin.cnu@foxmail.com.

‡E-mail: wfcwymmm@163.com.

nonhomogeneous primitive equations in R^2 , which read as

$$\begin{cases} \rho_t + \partial_x(\rho u) + \partial_y(\rho w) = 0, \\ \partial_t(\rho u) + \partial_x(\rho u^2) + \partial_y(\rho uw) + \partial_x P - \partial_x(\mu(\rho)\partial_x u) - \partial_y(\mu(\rho)\partial_y u) = 0, \\ \partial_x u + \partial_y w = 0, \\ \partial_y P = 0, \end{cases} \quad (1.1)$$

where the density ρ , the horizontal velocity u , the vertical velocity w and the pressure P are the unknown functions. The viscosity coefficient $\mu = \mu(\rho)$ is a function of the density ρ and assumed to satisfy

$$\mu(\rho) \in C^2[0, \infty) \quad \text{and} \quad \mu(\rho) \geq \underline{\mu} > 0 \quad \text{on} \quad [0, \infty). \quad (1.2)$$

The primitive equations are based on the so called hydrostatic approximation, and they form a fundamental block in models for planetary oceanic and atmospheric dynamics (see, for example, Pedlosky [29] and Vallis [34]). As far as we know, it was essentially introduced by Richardson in [30], to study the complicated atmospheric phenomena and for predicting the weather and possible climate changes. Until 1990s, the mathematical studies of primitive equations were initiated by Lions, Temam and Wang in [21]-[23], and they established the global existence of weak solutions to the system for the initial data $a \in L^2$; The uniqueness of weak solutions for the two-dimensional case was later proved by Bresch, Kazhikjov and Lemoine [1] for initial data a with $\partial_z a \in L^2$ and by Kukavica, Pei, Rusin and Ziane [18] for the initial data which is only continuous on the space variables. However, the uniqueness of these weak solutions for the three-dimensional case is still unclear. Concerning the strong solutions, the local existence results to the 2D primitive equations were proved by Guill-Gonzalez, Masmoudi and Rodruiz-Bellido in [12], while the global existence and uniqueness of strong solutions for the 2D primitive equations was achieved by Bresch, Kazhikhov and Lemoine in [1] and Temam and Ziane in [32]. The global existence of strong solutions for the 3D incompressible primitive equations has been known since the breakthrough work by Cao and Titi [6], see also Kobelkov [17] and Kukavica and Ziane [19]-[20]. In the partial viscosities and partial diffusivity case, one can see the works by Cao and Titi in [7] and by Cao, Li and Titi in [2]-[5]. However, the above-mentioned results are phrased within the L^2 -setting. Based on the maximal regularity technique, Hieber and kashiwabara [13] and Giga, Gries, Hieber, Hussein and kashiwabara [10]-[11] gave the existing results in the L^p type spaces.

In recent years, the mathematical analysis of compressible primitive equations has been attracting the attention of many mathematicians. Gatapov and Kazhikhov firstly studied in [9] and they proved the global existence of weak solutions to the 2D compressible primitive equations. The uniqueness of such weak solutions was proved by Jiu, Li and Wang in [14]. Ersoy, Ngom and Sy [8] derived the 3D compressible primitive equations and obtained the stability of weak solutions to the 3D compressible primitive equations with degenerate viscosity (see also Tang and Gao [33] for 2D case). Recently, Wang, Dou and Jiu [36] established the global existence of weak solutions to 3D compressible primitive equations with gravity field. The similar result is referred to Liu and Titi [26]. However, the uniqueness of these weak solutions for 3D case is still an open problem. Moreover, Liu and Titi [25] established local well-posedness results in the three-dimensional case. Meanwhile, they also considered the zero Mach number limit of compressible primitive equations in [24] and [27]. From the point of view of physics, the viscosity may depend on the density. Wang [35] established the local existence and uniqueness of strong solutions to the 2D compressible primitive equations with density-dependent viscosity.

Very recently, Wang and Jiu [38] proved the global existence and uniqueness of strong solutions to the 2D compressible primitive equations with density-dependent viscosity.

In this paper, we consider the nonhomogeneous primitive equations (1.1) in the domain $\Omega = M \times (0, 1)$, with $M = (0, L)$, where $L > 0$. We complement the system (1.1) with boundary conditions and initial conditions as follows.

$$\begin{aligned} \rho, u, w, P & \text{ are periodic in the direction } x, \\ u|_{y=0} &= u|_{y=1} = 0, \\ w|_{y=0} &= w|_{y=1} = 0, \end{aligned} \tag{1.3}$$

and

$$\begin{aligned} u|_{t=0} &= u_0(x, y), \\ w|_{t=0} &= w_0(x, y), \\ \rho|_{t=0} &= \rho_0(x, y). \end{aligned} \tag{1.4}$$

Throughout this paper, we define the following function spaces:

$$\begin{aligned} V &= \{v \in H^1(\Omega); \partial_y v = 0\}, \\ H_{0,per}^1(\Omega) &= \{v \in H^1(\Omega); \partial_x \int_0^1 v(x, y) dy = 0, v|_{y=0,1} = 0, v(0, y) = v(L, y)\}. \end{aligned}$$

The nonhomogeneous primitive equations, as the intermediate equations between incompressible primitive equations and compressible primitive equations, were firstly studied by Jiu and Wang in [16], and they proposed the following compatibility condition:

$$\mu \Delta u_0 - \partial_x P_0 = \rho_0^{\frac{1}{2}} g(x, z) \text{ in } \Omega \text{ and } \partial_x \int_0^1 u_0 dz = 0, \quad \text{for some } (P_0, g) \in V \times L^2$$

and established the local existence and uniqueness of strong solutions to the 2D nonhomogeneous primitive equations with large data. Furthermore, Wang, Jiu and Xu [37] obtained an global existence result on strong solutions to the 2D nonhomogeneous primitive equations for initial data with small $H^{\frac{1}{2}}$ -norm. Recently, for the case that the viscosity coefficient depends on the density, Jiu, Ma and Wang [15] established the local-in-time well-posedness of strong solutions to the 2D nonhomogeneous primitive equations (1.1), which is listed as follows.

Theorem 1.1 *Suppose that the initial data (ρ_0, u_0) satisfies the regularity conditions*

$$0 \leq \rho_0 \in W^{2,2}(\Omega), \quad \nabla \rho_0 \in L^\infty, \quad u_0 \in H_{0,per}^1(\Omega) \cap H^2(\Omega) \tag{1.5}$$

and the compatibility condition

$$\begin{aligned} -\rho_0 u_0 \partial_x u_0 + \rho_0 \int_0^y \partial_x u_0(s) ds \partial_y u_0 - \partial_x P_0 + \partial_x(\mu(\rho_0) \partial_x u_0) + \partial_y(\mu(\rho_0) \partial_y u_0) &:= \rho_0 V_1, \\ \partial_x \int_0^1 u_0(x, y) dy &= 0, \end{aligned} \tag{1.6}$$

for some $(P_0, V_1) \in V \times L^2$. Then there exists a small time $T_ \in (0, T)$, such that the initial boundary value problem (1.1)-(1.4) has a unique strong solution (ρ, u, w, P) satisfying*

$$\begin{aligned} \rho &\geq 0, \quad \rho \in C([0, T_*]; W^{1,\infty}) \cap C([0, T_*]; W^{2,2}), \quad \rho_t \in C([0, T_*]; L^q), \quad \sqrt{\rho} u_t \in L^\infty(0, T_*; L^2), \\ P &\in C([0, T_*]; H^1) \cap L^2(0, T_*; H^2), \quad \nabla u \in C([0, T_*]; H^1) \cap L^2(0, T_*; H^2), \\ u_t &\in L^2(0, T_*; H_{0,per}^1), \quad w \in C([0, T_*]; H^1) \cap L^2(0, T_*; H^2), \end{aligned} \tag{1.7}$$

where $1 \leq q < \infty$. Furthermore, if T^* is the maximal existence time of the local strong solutions (ρ, u, w, P) , then

$$\sup_{t \in [0, T^*]} (\|\nabla \rho(t)\|_{L^\infty} + \|\nabla^2 \rho(t)\|_{L^2} + \|\nabla u(t)\|_{L^2}) = \infty. \quad (1.8)$$

However, up to now, the global existence and uniqueness of strong solutions to the 2D nonhomogeneous primitive equations is open. The main purpose of this paper is to establish the global well-posedness of strong solutions to the problem (1.1)-(1.4) with small initial data in some sense. Our main result can be stated as

Theorem 1.2 *Suppose that the initial data (ρ_0, u_0, P_0) satisfies the conditions in Theorem 1.1. Then there exists a small positive constant η_0 , depending on Ω , $\sup_\Omega \rho_0 = \bar{\rho}$, $\|\mu(\rho)\|_{C^2}$, $\|\nabla \rho_0\|_{L^\infty}$ and $\|\nabla^2 \rho_0\|_{L^2}$, such that if*

$$\|\nabla u_0\|_{L^2} \leq \eta_0,$$

then the initial boundary value problem (1.1)-(1.4) admits a unique global strong solution (ρ, u, w, P) with

$$\begin{aligned} \rho &\geq 0, \quad \rho \in C(0, \infty; W^{1,\infty}) \cap C(0, \infty; W^{2,2}), \quad \rho_t \in C(0, \infty; L^q), \quad \sqrt{\rho} u_t \in L^\infty(0, \infty; L^2), \\ P &\in C(0, \infty; H^1) \cap L^2(0, \infty; H^2), \quad \nabla u \in C(0, \infty; H^1) \cap L^2(0, \infty; H^2) \\ u_t &\in L^2(0, \infty; H_{0,per}^1), \quad w \in C(0, \infty; H^1) \cap L^2(0, \infty; H^2). \end{aligned} \quad (1.9)$$

Moreover, the following decay property holds

$$\|\sqrt{\rho} u_t\|_{L^2}^2 \leq C(\Omega, \|\mu\|_{C^1}, \bar{\rho}, \|\nabla \rho_0\|_{L^\infty}, \|\nabla^2 \rho_0\|_{L^2}) \|\nabla u_0\|_{L^2}^2 t^{-2}, \quad (1.10)$$

and

$$\|\nabla u\|_{L^2}^2 \leq C(\Omega, \|\mu\|_{C^1}) (1 + \frac{\bar{\rho}}{\|\rho_0\|_{L^p}}) e^{-\sigma t}, \quad (1.11)$$

where $\sigma = \frac{\mu}{\|\rho_0\|_{L^p}}$ for any $p > 1$.

Based on the result in [15], we apply the continuity method to establish the global well-posedness of strong solutions to (1.1)-(1.4). It was shown in [15] that if T^* is the maximal existence time of the local strong solutions (ρ, u, w, P) and $T^* < T$, then

$$\sup_{0 \leq t < T^*} (\|\nabla \rho(t)\|_{L^\infty} + \|\nabla^2 \rho(t)\|_{L^2} + \|\nabla u(t)\|_{L^2}) = \infty,$$

which implies that the proof of global existence relies on the estimates on the gradient of horizontal velocity and the one-order and two-order derivative of the density. We start with the a priori hypothesis $\|\nabla \rho\|_{L^\infty} + \|\nabla^2 \rho\|_{L^2} \leq 8(\|\nabla \rho_0\|_{L^\infty} + \|\nabla^2 \rho_0\|_{L^2})$ and $\|\nabla u\|_{L^2}^2 \leq 4\frac{\bar{\mu}}{\mu} \|\nabla u_0\|_{L^2}^2$, where $\bar{\mu} = \sup_{[0, \bar{\rho}]} \mu(\rho)$. Then we prove that $\|\nabla \rho\|_{L^\infty} + \|\nabla^2 \rho\|_{L^2}$ is less than $4(\|\nabla \rho_0\|_{L^\infty} + \|\nabla^2 \rho_0\|_{L^2})$ and $\|\nabla u\|_{L^2}^2$ is less than $(2\ln 2 + 1)\frac{\bar{\mu}}{\mu} \|\nabla u_0\|_{L^2}^2$ on $[0, T]$ with the assumption $\|\nabla u_0\|_{L^2}$ is suitably small. In fact, by making use of the regularity result of hydrostatic Stokes equations (see Lemma 3.3), it is easy to obtain the desired bound of $\|\nabla u\|_{L^2}$ under the assumption that $\|\nabla u_0\|_{L^2}$ is suitably small (see Lemma 3.2). Then, the next key step is to show that $\|\nabla \rho\|_{L^\infty} + \|\nabla^2 \rho\|_{L^2}$ is less than $4(\|\nabla \rho_0\|_{L^\infty} + \|\nabla^2 \rho_0\|_{L^2})$. To do this, it suffices to deal with a trouble term $\|\nabla u_t\|_{L_t^1 L_\Omega^2}$ which appears when estimating the term $\|u\|_{L_t^1 H_\Omega^3}$ (see Lemma 3.8). Indeed, based on the regularity result of hydrostatic Stokes equations and some careful time weighted estimates, we find

that the norm of $\|\nabla u_t\|_{L_t^1 L_\Omega^2}$ is bounded and could be as small as desired under certain smallness conditions. More precisely, we split it into two terms

$$\int_0^{T_*} \|\nabla u_t\|_{L^2} dt \leq \left(\int_0^{T^0} \|\nabla u_t\|_{L^2}^2 dt \right)^{\frac{1}{2}} \sqrt{T^0} + \left(\int_{T^0}^{T_*} t^2 \|\nabla u_t\|_{L^2}^2 dt \right)^{\frac{1}{2}} \left(\int_{T^0}^{T_*} t^{-2} dt \right)^{\frac{1}{2}}, \quad (1.12)$$

where T^0 is a small enough number independent on the norm of $\|\nabla u_0\|_{L^2}$. We remark that the local time T_* might depend on $\|\nabla u_0\|_{L^2}$ (see [15]). We can consider the local existence of the strong solutions under the additional assumption of $\|\nabla u_0\|_{L^2} \leq 1$ such that the local time T_* we obtain is independent on the norm of $\|\nabla u_0\|_{L^2}$ (see Lemma 2.4). Therefore, we can choose some small T_0 independent on the norm of $\|\nabla u_0\|_{L^2}$ such that the first term of (1.12) is small enough. Meanwhile, the second term of (1.12) can be controlled by the time weighted estimates (see (3.26) for more details). Then, it follows that $\int_0^T \|u\|_{H^3} dt \leq \ln 2$ under the assumption that $\|\nabla u_0\|_{L^2}$ is suitably small. As a result, the control of $\|\nabla \rho\|_{L^\infty} + \|\nabla^2 \rho\|_{L^2}$ and $\|\nabla u\|_{L^2}$ lead to uniform estimates for other higher order quantities, which guarantees the extension of local strong solutions.

The rest of this paper is organized as follows. In Section 2, we recall some lemmas and give an auxiliary lemma which is used later. In Section 3, we give an outline of the proof of the main Theorem.

The following notations will be used:

- (1) $\|f\|_{L_y^2} = \left(\int_0^1 |f|^2 dy \right)^{\frac{1}{2}},$
- (2) $\|f\|_{L_y^2 L_x^\infty} = \left(\int_0^1 \|f\|_{L_x^\infty}^2 dy \right)^{\frac{1}{2}}, \|f\|_{L_y^\infty L_x^\infty} = \text{esssup}_{(x,y) \in \Omega} |f|,$
- (3) For $1 < p < \infty$, $\|f\|_{L^p} = \|f\|_{L_y^p L_x^p} = \|f\|_{L_x^p L_y^p} = \left(\int_0^1 \int_0^L |f|^p dx dy \right)^{\frac{1}{p}},$
- (4) $\int_\Omega = \int_\Omega dx dy.$

2 Preliminaries

In this section, we recall some basic facts and elementary inequalities which will be used later. First of all, we present the well-known Gagliardo-Nirenberg interpolation inequality.

Lemma 2.1 (see [28])(Gagliardo-Nirenberg interpolation inequality). *For a function $u : \Omega \rightarrow \mathbb{R}$ defined on a bounded Lipschitz domain $\Omega \subset \mathbb{R}^n, \forall 1 \leq q, r \leq \infty$, and a natural number m . Suppose also that a real number θ and a natural number j are such that*

$$\frac{1}{p} = \frac{j}{n} + \left(\frac{1}{r} - \frac{m}{n} \right) \theta + \frac{1-\theta}{q},$$

and

$$\frac{j}{m} \leq \theta \leq 1.$$

Then, we have

$$\|D^j u\|_{L^p} \leq C_1 \|D^m u\|_{L^r}^\theta \|u\|_{L^q}^{1-\theta} + C_2 \|u\|_{L^s},$$

where s is a positive constant. The constants C_1 and C_2 depend on the domain Ω and m, n, j, r, q, θ .

The derivations of high-order estimates on the horizontal velocity u rely on the following regularity results for the density-dependent hydrostatic Stokes equations.

Lemma 2.2 Assume that $\rho \in W^{2,2}$, $\nabla \rho \in L^\infty$ and $0 \leq \rho \leq \bar{\rho}$, let $(u, P) \in H_{0,per}^1 \times L^2$ be the unique weak solution to the problem

$$-\partial_x(\mu(\rho)\partial_x u) - \partial_y(\mu(\rho)\partial_y u) + \partial_x P = f, \quad \partial_x \int_0^1 u(x, y) dy = 0, \quad \int_\Omega P dx dy = 0. \quad (2.1)$$

Then we have the following regularity results:

(1) If $f \in L^2$, then $(u, P) \in H^2 \times H^1$ and

$$\|u\|_{H^2} + \|P\|_{H^1} \leq C(1 + \|\nabla \rho\|_{L^\infty}) \|f\|_{L^2}.$$

(2) If $f \in L^r$, then $(u, P) \in W^{2,r} \times W^{1,r}$ and

$$\|u\|_{W^{2,r}} + \|P\|_{W^{1,r}} \leq C(1 + \|\nabla \rho\|_{L^\infty})^2 \|f\|_{L^r}, \quad (2.2)$$

where $r > 2$.

(3) If $f \in H^1$, then $(u, P) \in H^3 \times H^2$ and

$$\|u\|_{H^3} + \|P\|_{H^2} \leq \tilde{C}(1 + \|\nabla \rho\|_{L^\infty} + \|\nabla^2 \rho\|_{L^2})^3 \|f\|_{H^1}, \quad (2.3)$$

the constant \tilde{C} depend also on $\|\partial^2 \mu / \partial x^2\|_C$.

Lemma 2.3 Assume that $u \in L^1([0, T], H^3(\Omega))$, let ρ be the unique solution to the problem

$$\partial_t \rho + \partial_x \rho u - \partial_y \rho \int_0^y \partial_x u(s) ds, \quad \rho|_{t=0} = \rho_0.$$

If $\nabla \rho_0 \in L^\infty(\Omega)$ and $\rho_0 \in W^{2,2}(\Omega)$, then we have the following regularity results:

$$\|\nabla \rho(t)\|_{L^\infty(\Omega)} \leq \|\nabla \rho_0\|_{L^\infty(\Omega)} \exp\left(\int_0^t \|u(s)\|_{H^3(\Omega)} ds\right), \quad (2.4)$$

and

$$\|\nabla^2 \rho(t)\|_{L^2(\Omega)} \leq (\|\nabla \rho_0\|_{L^\infty(\Omega)} + \|\nabla^2 \rho_0\|_{L^2(\Omega)}) \exp \int_0^t \|u(s)\|_{H^3(\Omega)} ds. \quad (2.5)$$

The proof of (2.4) and (2.5) is referred to [15] and we omit it here.

As aforementioned, the global existence and uniqueness of strong solutions will be proved by combining the global a priori estimates in section 3 with the following local-in-time existence results which can follow as a corollary of Theorem 1.1 in [15].

Lemma 2.4 Suppose that the initial data (ρ_0, u_0, P_0) satisfies the regularity conditions in Theorem 1.1 and the additional condition $\|\nabla u_0\|_{L^2}^2 \leq 1$. Then there exists a small time $T_* \in (0, T)$ such that the initial boundary value problem (1.1)-(1.4) has a unique strong solution (ρ, u, w, P) satisfying

$$\begin{aligned} \rho &\in C([0, T_*]; W^{1,\infty}) \cap C([0, T_*]; W^{2,2}), \quad \rho_t \in C([0, T_*]; L^q), \quad \sqrt{\rho} u_t \in L^\infty(0, T_*; L^2), \\ P &\in C([0, T_*]; H^1) \cap L^2(0, T_*; H^2), \quad \nabla u \in C([0, T_*]; H^1) \cap L^2(0, T_*; H^2), \\ u_t &\in L^2(0, T_*; H_{0,per}^1), \quad w \in C([0, T_*]; H^1) \cap L^2(0, T_*; H^2), \end{aligned}$$

where $1 \leq q < \infty$ and T_* is independent on the norm of $\|\nabla u_0\|_{L^2}$.

3 Proof of Theorem 1.2

This section is devoted to the proof of Theorem 1.2. Indeed, we follow the idea in [15] and first prove the estimates for the case of a positive initial density and obtain the uniform bounds which are independent of the lower bounds of the initial density. Then the vacuum case can be proved by applying standard regularizing techniques and compactness arguments.

3.1 A priori estimates

In this subsection, we establish several a priori estimates for strong solutions to the system (1.1)-(1.4), which will play a key role in extending local strong solutions to the global ones. For simplicity, throughout this subsection, a constant C will denote some positive one which may be dependent on Ω and $\|\mu\|_{C_1}$, but is independent of ρ_0 and u_0 , and \tilde{C} depends on C and $\|\partial^2 \mu / \partial x^2\|_C$. We remark that the value of the constant C will be different from line to line.

We begin with the boundedness of density and the basic energy estimates, which can be easily derived from the system (1.1).

Lemma 3.1 *Assume that (ρ, u, w, P) is the unique strong solution to the system (1.1)-(1.4) on $\Omega \times (0, T)$ with the initial data satisfying the assumptions in Theorem 1.1. Then, for every $(x, y, t) \in \Omega \times (0, T)$, it holds that*

$$\|\rho\|_{L^\infty} = \|\rho_0\|_{L^\infty},$$

and

$$\int_{\Omega} \rho \frac{1}{2} u^2(t) + \int_0^t \int_{\Omega} \mu(\rho) |\nabla u|^2 ds \leq \frac{1}{2} \bar{\rho} \|u_0\|_{L^2}^2, \quad (3.1)$$

which yields

$$\frac{1}{2} \int_{\Omega} \rho u^2(t) + \underline{\mu} \int_0^t \int_{\Omega} |\nabla u|^2 ds \leq \frac{1}{2} \bar{\rho} \|u_0\|_{L^2}^2. \quad (3.2)$$

The next lemma is concerned with the L^2 -norm of the gradient of horizontal velocity. Before state and prove it, we denote

$$M_0 := \|\nabla \rho_0\|_{L^\infty} + \|\nabla^2 \rho_0\|_{L^2}.$$

Lemma 3.2 *Assume that (ρ, u, w, P) is the unique strong solution to the system (1.1)-(1.4) on $\Omega \times (0, T)$ with the initial data satisfying the assumptions in Theorem 1.1, satisfying*

$$\sup_{t \in [0, T]} \|\nabla \rho(t)\|_{L^\infty} + \sup_{t \in [0, T]} \|\nabla^2 \rho(t)\|_{L^2} \leq 8M_0, \quad (3.3)$$

and

$$\sup_{t \in [0, T]} \|\nabla u(t)\|_{L^2}^2 \leq 4 \frac{\bar{\mu}}{\underline{\mu}} \|\nabla u_0\|_{L^2}^2 \leq 1. \quad (3.4)$$

There exists a positive number γ_0 , depending on Ω , $\sup_{\Omega} \rho_0 = \bar{\rho}$, $\|\mu\|_{C^2}$, $\|\nabla \rho_0\|_{L^\infty}$ and $\|\nabla^2 \rho_0\|_{L^2}$, such that, if

$$\|\nabla u_0\|_{L^2} \leq \gamma_0, \quad (3.5)$$

then

$$\frac{2}{\underline{\mu}} \int_0^T \int_{\Omega} \rho u_t^2 dt + \sup_{t \in [0, T]} \int_{\Omega} |\nabla u|^2 \leq (2\ln 2 + 1) \frac{\bar{\mu}}{\underline{\mu}} \|\nabla u_0\|_{L^2}^2. \quad (3.6)$$

Before proving Lemma 3.2, we need the following lemma.

Lemma 3.3 *Assume that (ρ, u, w, P) is the unique strong solution to the system (1.1)-(1.4) on $\Omega \times (0, T)$ with the initial data satisfying the assumptions in Theorem 1.1, and satisfies*

$$\sup_{t \in [0, T]} \|\nabla \rho(t)\|_{L^\infty} + \sup_{t \in [0, T]} \|\nabla^2 \rho(t)\|_{L^2} \leq 8M_0.$$

Then it holds that

$$\|\nabla u\|_{H^1} \leq C(\bar{\rho}^{1/2} M \|\sqrt{\rho} u_t\|_{L^2} + \bar{\rho} M \|\nabla u\|_{L^2}^2 + \bar{\rho}^{\frac{3}{2}} M^{\frac{3}{2}} \|\nabla u\|_{L^2}^{\frac{5}{2}} + \bar{\rho}^2 M^2 \|\nabla u\|_{L^2}^3), \quad (3.7)$$

where $M = 1 + 8M_0$.

Proof. The horizontal momentum equations $(1.1)_2$ can be rewritten as follows,

$$-\partial_x(\mu(\rho)\partial_x u) - \partial_y(\mu(\rho)\partial_y u) + \partial_x P = -\rho\partial_t u - \rho u\partial_x u + \rho \int_0^y \partial_x u(s) ds \partial_y u. \quad (3.8)$$

It follows from Lemma 2.1 and Lemma 2.2 that

$$\begin{aligned} \|\nabla u\|_{H^1} &\leq CM(\|\rho u_t\|_{L^2} + \bar{\rho}\|u\|_{L^6}\|\partial_x u\|_{L^3} + \bar{\rho}\|\int_0^y \partial_x u(s) ds\|_{L_y^2 L_x^\infty} \|\partial_y u\|_{L_x^2 L_y^\infty}) \\ &\leq CM[\bar{\rho}^{1/2} \|\sqrt{\rho} u_t\|_{L^2} + \bar{\rho}\|\nabla u\|_{L^2}(C_1\|\nabla^2 u\|_{L^2}^{\frac{1}{3}}\|\nabla u\|_{L^2}^{\frac{2}{3}} + C_2\|\nabla u\|_{L^2}) \\ &\quad + \bar{\rho}\|\nabla u\|_{L^2}(C_1\|\nabla^2 u\|_{L^2}^{\frac{1}{2}}\|\nabla u\|_{L^2}^{\frac{1}{2}} + C_2\|\nabla u\|_{L^2})] \\ &\leq C(\bar{\rho}^{1/2} M \|\sqrt{\rho} u_t\|_{L^2} + \bar{\rho} M \|\nabla u\|_{L^2}^2 + \bar{\rho} M \|\nabla^2 u\|_{L^2}^{\frac{1}{3}} \|\nabla u\|_{L^2}^{\frac{5}{3}} \\ &\quad + M \bar{\rho} \|\nabla^2 u\|_{L^2}^{\frac{1}{2}} \|\nabla u\|_{L^2}^{\frac{3}{2}}). \end{aligned} \quad (3.9)$$

Applying Young's inequality, we deduce that

$$\|\nabla u\|_{H^1} \leq C(\bar{\rho}^{1/2} M \|\sqrt{\rho} u_t\|_{L^2} + \bar{\rho} M \|\nabla u\|_{L^2}^2 + \bar{\rho}^{\frac{3}{2}} M^{\frac{3}{2}} \|\nabla u\|_{L^2}^{\frac{5}{2}} + \bar{\rho}^2 M^2 \|\nabla u\|_{L^2}^3). \quad (3.10)$$

□

Now we give the proof of Lemma 3.2.

Proof of Lemma 3.2. Multiplying the horizontal momentum equations $(1.1)_2$ by u_t and integrating by part over Ω yields

$$\begin{aligned} \int_{\Omega} \rho u_t^2 + \frac{1}{2} \frac{d}{dt} \int_{\Omega} \mu(\rho) |\nabla u|^2 &= - \int_{\Omega} \rho u \partial_x u u_t + \int_{\Omega} (\rho \int_0^y \partial_x u(s) ds \partial_y u u_t) + \frac{1}{2} \int_{\Omega} \mu_t(\rho(t)) |\nabla u|^2 \\ &= - \int_{\Omega} \rho u \partial_x u u_t + \int_{\Omega} (\rho \int_0^y \partial_x u(s) ds \partial_y u u_t) \\ &\quad + \frac{1}{2} \int_{\Omega} (\int_0^y \partial_x u(s) ds \partial_y \mu(\rho) - u(x) \partial_x \mu(\rho)) |\nabla u|^2. \end{aligned} \quad (3.11)$$

Next, we will estimate the terms on the right-hand side in (3.11) one by one. For the first term, applying Young inequality and Lemma 3.3, we have

$$\begin{aligned}
\int_{\Omega} \rho u_t u \partial_x u &\leq \|\sqrt{\rho} u_t\|_{L^2} \|\sqrt{\rho}\|_{L^\infty} \|u\|_{L^6} \|\partial_x u\|_{L^3} \\
&\leq \bar{\rho}^{\frac{1}{2}} \|\sqrt{\rho} u_t\|_{L^2} \|\nabla u\|_{L^2} (C_1 \|\nabla^2 u\|_{L^2}^{\frac{1}{3}} \|\nabla u\|_{L^2}^{\frac{2}{3}} + C_2 \|\nabla u\|_{L^2}) \\
&\leq \epsilon \|\sqrt{\rho} u_t\|_{L^2}^2 + \bar{\rho} \|\nabla u\|_{L^2}^2 (C_1 \|\nabla^2 u\|_{L^2}^{\frac{2}{3}} \|\nabla u\|_{L^2}^{\frac{4}{3}} + C_2 \|\nabla u\|_{L^2}^2) \\
&\leq \epsilon \|\sqrt{\rho} u_t\|_{L^2}^2 + C(\bar{\rho} \|\nabla u\|_{L^2}^{\frac{10}{3}} \|\nabla^2 u\|_{L^2}^{\frac{2}{3}} + \bar{\rho} \|\nabla u\|_{L^2}^4) \\
&\leq \epsilon \|\sqrt{\rho} u_t\|_{L^2}^2 + C(\bar{\rho}^{\frac{4}{3}} M^{\frac{2}{3}} \|\sqrt{\rho} u_t\|_{L^2}^{\frac{2}{3}} \|\nabla u\|_{L^2}^{\frac{10}{3}} + M^{\frac{2}{3}} \bar{\rho}^{\frac{5}{3}} \|\nabla u\|_{L^2}^{\frac{14}{3}} \\
&\quad + M \bar{\rho}^2 \|\nabla u\|_{L^2}^5 + M^{\frac{4}{3}} \bar{\rho}^{\frac{7}{3}} \|\nabla u\|_{L^2}^{\frac{16}{3}} + \bar{\rho} \|\nabla u\|_{L^2}^4) \\
&\leq \epsilon \|\sqrt{\rho} u_t\|_{L^2}^2 + C(\bar{\rho}^2 M \|\nabla u\|_{L^2}^5 + M^{\frac{2}{3}} \bar{\rho}^{\frac{5}{3}} \|\nabla u\|_{L^2}^{\frac{14}{3}} \\
&\quad + M^{\frac{4}{3}} \bar{\rho}^{\frac{7}{3}} \|\nabla u\|_{L^2}^{\frac{16}{3}} + \bar{\rho} \|\nabla u\|_{L^2}^4).
\end{aligned} \tag{3.12}$$

Similarly, the second term can be bounded as

$$\begin{aligned}
\int_{\Omega} (\rho \int_0^y \partial_x u(s) ds \partial_y u u_t) &\leq \epsilon \|\sqrt{\rho} u_t\|_{L^2}^2 + \bar{\rho} \|\nabla u\|_{L^2}^2 (C_1 \|\nabla^2 u\|_{L^2} \|\nabla u\|_{L^2} + C_2 \|\nabla u\|_{L^2}^2) \\
&\leq \epsilon \|\sqrt{\rho} u_t\|_{L^2}^2 + C(\bar{\rho} \|\nabla u\|_{L^2}^3 \|\nabla^2 u\|_{L^2} + \bar{\rho} \|\nabla u\|_{L^2}^4) \\
&\leq \epsilon \|\sqrt{\rho} u_t\|_{L^2}^2 + C(\bar{\rho}^{3/2} M \|\sqrt{\rho} u_t\|_{L^2} \|\nabla u\|_{L^2}^3 + \bar{\rho}^2 M \|\nabla u\|_{L^2}^5 \\
&\quad + \bar{\rho}^{\frac{5}{2}} M^{\frac{3}{2}} \|\nabla u\|_{L^2}^{\frac{11}{2}} + \bar{\rho}^3 M^2 \|\nabla u\|_{L^2}^6 + \bar{\rho} \|\nabla u\|_{L^2}^4) \\
&\leq \epsilon \|\sqrt{\rho} u_t\|_{L^2}^2 + C(\bar{\rho}^2 M \|\nabla u\|_{L^2}^5 + \bar{\rho}^{\frac{5}{2}} M^{\frac{3}{2}} \|\nabla u\|_{L^2}^{\frac{11}{2}} \\
&\quad + \bar{\rho}^3 M^2 \|\nabla u\|_{L^2}^6 + \bar{\rho} \|\nabla u\|_{L^2}^4).
\end{aligned} \tag{3.13}$$

For the last term, by direct estimates, we get

$$\begin{aligned}
&\int_{\Omega} [\int_0^y \partial_x u(s) ds \partial_y \mu(\rho) |\nabla u| |\nabla u| - u(x) \partial_x \mu(\rho) |\nabla u| |\nabla u|] \\
&\leq \|\int_0^y \partial_x u(s) ds\|_{L^\infty} \|\partial_y \mu(\rho)\|_{L^\infty} \|\nabla u\|_{L^2} \|\nabla u\|_{L^2} - \|u\|_{L^\infty} \|\partial_x \mu(\rho)\|_{L^\infty} \|\nabla u\|_{L^2} \|\nabla u\|_{L^2} \\
&\leq \|u\|_{H^2} \|\partial_y \mu(\rho)\|_{L^\infty} \|\nabla u\|_{L^2}^2 \\
&\leq C(\bar{\rho}^{1/2} M \|\sqrt{\rho} u_t\|_{L^2} \|\nabla u\|_{L^2}^2 + \bar{\rho} M \|\nabla u\|_{L^2}^4 + \bar{\rho}^{\frac{3}{2}} M^{\frac{3}{2}} \|\nabla u\|_{L^2}^{\frac{9}{2}} + \bar{\rho}^2 M^2 \|\nabla u\|_{L^2}^5) M_0 \\
&\leq \epsilon \|\sqrt{\rho} u_t\|_{L^2}^2 + C(\bar{\rho} M^2 M_0^2 \|\nabla u\|_{L^2}^4 + \bar{\rho} M M_0 \|\nabla u\|_{L^2}^4 + \bar{\rho}^{\frac{3}{2}} M^{\frac{3}{2}} M_0 \|\nabla u\|_{L^2}^{\frac{9}{2}} + \bar{\rho}^2 M^2 M_0 \|\nabla u\|_{L^2}^5).
\end{aligned} \tag{3.14}$$

Substituting (3.12)-(3.14) into (3.11), we obtain

$$\int_{\Omega} \rho u_t^2 + \frac{1}{2} \frac{d}{dt} \int_{\Omega} \mu(\rho) |\nabla u|^2 \leq C \mathcal{Q} \|\nabla u\|_{L^2}^2, \tag{3.15}$$

where

$$\begin{aligned}
\mathcal{Q} &:= \bar{\rho} \|\nabla u\|_{L^2}^2 + \bar{\rho} M^2 M_0^2 \|\nabla u\|_{L^2}^2 + \bar{\rho} M M_0 \|\nabla u\|_{L^2}^2 + M^{\frac{2}{3}} \bar{\rho}^{\frac{5}{3}} \|\nabla u\|_{L^2}^{\frac{8}{3}} \\
&\quad + M^{\frac{4}{3}} \bar{\rho}^{\frac{7}{3}} \|\nabla u\|_{L^2}^{\frac{10}{3}} + \bar{\rho}^2 M \|\nabla u\|_{L^2}^3 + \bar{\rho}^2 M^2 M_0 \|\nabla u\|_{L^2}^3 + \bar{\rho}^{\frac{5}{2}} M^{\frac{3}{2}} \|\nabla u\|_{L^2}^{\frac{7}{2}} \\
&\quad + \bar{\rho}^3 M^2 \|\nabla u\|_{L^2}^4 + \bar{\rho}^{\frac{3}{2}} M^{\frac{3}{2}} M_0 \|\nabla u\|_{L^2}^{\frac{5}{2}}.
\end{aligned} \tag{3.16}$$

Integrating with respect to time on $[0, t]$ and using the fact that $\mu \geq \underline{\mu} > 0$ gives

$$\begin{aligned} \frac{2}{\underline{\mu}} \int_0^t \int_{\Omega} \rho u_t^2 ds + \int_{\Omega} |\nabla u|^2 &\leq \int_0^t \frac{CQ}{\underline{\mu}} \|\nabla u(s)\|_{L^2}^2 ds + \int_{\Omega} \frac{\mu(\rho_0)}{\underline{\mu}} |\nabla u_0|^2 \\ &\leq \int_0^t \frac{CQ}{\underline{\mu}} (\|\nabla u(s)\|_{L^2}^2 + \frac{2}{\underline{\mu}} \int_0^s \int_{\Omega} \rho u_t^2(\tau) d\tau) ds + \int_{\Omega} \frac{\mu(\rho_0)}{\underline{\mu}} |\nabla u_0|^2 \end{aligned} \quad (3.17)$$

Applying Gronwall inequality, we deduce that

$$\begin{aligned} \frac{2}{\underline{\mu}} \int_0^T \int_{\Omega} \rho u_t^2 dt + \sup_{t \in [0, T]} \int_{\Omega} |\nabla u|^2 \\ \leq \frac{1}{\underline{\mu}} \int_{\Omega} \mu(\rho_0) |\nabla u_0|^2 + \frac{1}{\underline{\mu}} \int_{\Omega} \mu(\rho_0) |\nabla u_0|^2 \cdot \int_0^T \frac{CQ}{\underline{\mu}} dt \cdot \exp\left(\int_0^T \frac{CQ}{\underline{\mu}} dt\right). \end{aligned} \quad (3.18)$$

According to Lemmas 3.1, (3.4) and Poincaré inequality, one has

$$\begin{aligned} \int_0^T \frac{CQ}{\underline{\mu}} dt \leq C \left\{ \bar{\rho} + \bar{\rho} M^2 M_0^2 + \bar{\rho} M M_0 + M^{\frac{2}{3}} \bar{\rho}^{\frac{5}{3}} + M^{\frac{4}{3}} \bar{\rho}^{\frac{7}{3}} + \bar{\rho}^2 M + \bar{\rho}^2 M^2 M_0 \right. \\ \left. + \bar{\rho}^{\frac{5}{2}} M^{\frac{3}{2}} + \bar{\rho}^3 M^2 + \bar{\rho}^{\frac{3}{2}} M^{\frac{3}{2}} M_0 \right\} \frac{\bar{\rho}}{\underline{\mu}^2} \|\nabla u_0\|_{L^2}^2. \end{aligned} \quad (3.19)$$

Now we choose γ_0 small enough such that

$$C(\bar{\rho} + \bar{\rho} M^4 + \bar{\rho} M^2 + M^{\frac{2}{3}} \bar{\rho}^{\frac{5}{3}} + M^{\frac{4}{3}} \bar{\rho}^{\frac{7}{3}} + \bar{\rho}^2 M + \bar{\rho}^2 M^3 + \bar{\rho}^{\frac{5}{2}} M^{\frac{3}{2}} + \bar{\rho}^3 M^2 + \bar{\rho}^{\frac{3}{2}} M^{\frac{5}{2}}) \frac{\bar{\rho}}{\underline{\mu}^2} \gamma_0^2 \leq \ln 2.$$

It follows from (3.18), (3.19) and (3.5) that

$$\begin{aligned} \frac{2}{\underline{\mu}} \int_0^T \int_{\Omega} \rho u_t^2 dt + \sup_{t \in [0, T]} \int_{\Omega} |\nabla u|^2 &\leq \frac{\bar{\mu}}{\underline{\mu}} \|\nabla u_0\|_{L^2}^2 + \frac{\bar{\mu}}{\underline{\mu}} \|\nabla u_0\|^2 \cdot \ln 2 \cdot 2 \\ &= (2 \ln 2 + 1) \frac{\bar{\mu}}{\underline{\mu}} \|\nabla u_0\|_{L^2}^2. \end{aligned} \quad (3.20)$$

□

Recall that $\|\nabla \rho\|_{L^\infty} + \|\nabla^2 \rho(t)\|_{L^2(\Omega)} \leq (\|\nabla \rho_0\|_{L^\infty(\Omega)} + \|\nabla^2 \rho_0\|_{L^2(\Omega)}) \exp \int_0^t \|u(s)\|_{H^3(\Omega)} ds$. To prove that $\|\nabla \rho(t)\|_{L^\infty} + \|\nabla^2 \rho(t)\|_{L^2} \leq 4M_0$, we still need the H^3 -regularity of the velocity u . In fact, we need the following several lemmas to deal with $\|u\|_{H^3}$. First, we establish some time weighted estimates.

Lemma 3.4 *Assume that (ρ, u, w, P) is the unique strong solution to the system (1.1)-(1.4) on $\Omega \times (0, T)$ with the initial data satisfying the assumptions in Theorem 1.1 and (3.3)-(3.5). Then it holds that*

$$\frac{2}{\underline{\mu}} \int_0^T \left(t \int_{\Omega} \rho u_t^2 \right) dt + \sup_{t \in [0, T]} \left(t \int_{\Omega} |\nabla u|^2 \right) \leq (2 \ln 2 + 1) \frac{\bar{\mu} \bar{\rho}}{2 \underline{\mu}^2} \|\nabla u_0\|_{L^2}^2. \quad (3.21)$$

Proof. Multiplying (3.15) by t and adding the term $\frac{1}{2} \int_{\Omega} \mu(\rho) |\nabla u|^2$, we have

$$t \int_{\Omega} \rho u_t^2 + \frac{1}{2} \frac{d}{dt} \left(t \int_{\Omega} \mu(\rho) |\nabla u|^2 \right) \leq t C Q \|\nabla u\|_{L^2}^2 + \frac{1}{2} \int_{\Omega} \mu(\rho) |\nabla u|^2. \quad (3.22)$$

Integrating with respect to time on $[0, t]$ and using the fact that $\mu \geq \underline{\mu} > 0$, we get

$$\begin{aligned} & \frac{2}{\underline{\mu}} \int_0^t (s \int_{\Omega} \rho u_t^2) ds + \sup_{t \in [0, T]} (t \int_{\Omega} |\nabla u|^2) \\ & \leq \int_0^t (\frac{CQ}{\underline{\mu}} s \|\nabla u\|_{L^2}^2) ds + \frac{1}{\underline{\mu}} \int_0^t (\int_{\Omega} \mu(\rho) |\nabla u|^2) ds \\ & \leq \int_0^t \frac{CQ}{\underline{\mu}} (s \|\nabla u(s)\|_{L^2}^2 + \frac{2}{\underline{\mu}} \int_0^s \tau \int_{\Omega} \rho u_t^2(\tau) d\tau) ds + \frac{1}{\underline{\mu}} \int_0^t (\int_{\Omega} \mu(\rho) |\nabla u|^2) ds. \end{aligned} \quad (3.23)$$

By Gronwall inequality, assumption (3.5), Poincaré inequality and Lemma 3.1, we deduce

$$\begin{aligned} & \frac{2}{\underline{\mu}} \int_0^T (t \int_{\Omega} \rho u_t^2) dt + \sup_{t \in [0, T]} (t \int_{\Omega} |\nabla u|^2) \\ & \leq \frac{1}{\underline{\mu}} \int_0^T (\int_{\Omega} \mu(\rho) |\nabla u|^2) dt + \frac{1}{\underline{\mu}} \int_0^T (\int_{\Omega} \mu(\rho) |\nabla u|^2) dt \cdot \int_0^T \frac{CQ}{\underline{\mu}} dt \cdot \exp(\int_0^T \frac{CQ}{\underline{\mu}} dt) \\ & \leq (2 \ln 2 + 1) \frac{\bar{\mu} \bar{\rho}}{2 \underline{\mu}^2} \|\nabla u_0\|_{L^2}^2. \end{aligned} \quad (3.24)$$

□

Lemma 3.5 Assume that (ρ, u, w, P) is the unique strong solution to the system (1.1)-(1.4) on $\Omega \times (0, T)$ with the initial data satisfying the assumptions in Theorem 1.1 and (3.3)-(3.5). Then it holds that

$$\sup_{t \in [0, T]} t \int_{\Omega} \rho u_t^2 + 2 \underline{\mu} \int_0^T (t \int_{\Omega} |\nabla u_t|^2) dt \leq C \Theta_1 \|\nabla u_0\|_{L^2}^2 \cdot \exp[C \Theta_2], \quad (3.25)$$

and

$$\sup_{t \in [0, T]} t^2 \int_{\Omega} \rho u_t^2 + 2 \underline{\mu} \int_0^T (t^2 \int_{\Omega} |\nabla u_t|^2) dt \leq C \bar{\rho} \Theta_1 \|\nabla u_0\|_{L^2}^2 \cdot \exp[C \Theta_2], \quad (3.26)$$

where $\Theta_1 = 1 + M^4 \bar{\rho}^4 + \bar{\rho}^5 M^5 + M^6 \bar{\rho}^6$ and $\Theta_2 = C(M^4 \bar{\rho}^2 + \bar{\rho}^2 + \bar{\rho}^4)$.

Proof. Taking t-derivative of the horizontal momentum equations (1.1)₂, we deduce that

$$\begin{aligned} & \rho u_{tt} + \rho u \partial_x u_t - \rho \int_0^y \partial_x u(s) ds \partial_y u_t - \partial_x(\mu(\rho) \partial_x u_t) - \partial_y(\mu(\rho) \partial_y u_t) + \partial_x P_t \\ & = -\rho_t u_t - (\rho u)_t \partial_x u + (\rho \int_0^y \partial_x u(s) ds)_t \partial_y u + \partial_x(\mu(\rho)_t \partial_x u) + \partial_y(\mu(\rho)_t \partial_y u). \end{aligned} \quad (3.27)$$

Multiplying (3.27) by tu_t and integrating (by parts) over Ω yields

$$\begin{aligned} & \frac{t}{2} \frac{d}{dt} \int_{\Omega} \rho u_t^2 + t \int_{\Omega} \mu(\rho) |\nabla u_t|^2 \\ & = -\frac{t}{2} \int_{\Omega} \rho_t u_t^2 - t \int_{\Omega} (\rho u)_t \partial_x u u_t + t \int_{\Omega} (\rho \int_0^y \partial_x u(s) ds)_t \partial_y u u_t - \frac{t}{2} \int_{\Omega} (\mu(\rho))_t \frac{d|\nabla u|^2}{dt}. \end{aligned} \quad (3.28)$$

Noting the fact that $0 < \underline{\mu} \leq \mu$, one has

$$\begin{aligned} & \frac{t}{2 \underline{\mu}} \frac{d}{dt} \int_{\Omega} \rho u_t^2 + t \int_{\Omega} |\nabla u_t|^2 = -\frac{t}{2 \underline{\mu}} \int_{\Omega} \rho_t u_t^2 - \frac{t}{\underline{\mu}} \int_{\Omega} (\rho u)_t \partial_x u u_t \\ & + \frac{t}{\underline{\mu}} \int_{\Omega} (\rho \int_0^y \partial_x u(s) ds)_t \partial_y u u_t - \frac{t}{2 \underline{\mu}} \int_{\Omega} (\mu(\rho))_t \frac{d|\nabla u|^2}{dt} = \sum_{i=1}^4 I_i. \end{aligned} \quad (3.29)$$

Using (1.1)₁ and Poincaré inequality, we estimate the term I_1 as

$$\begin{aligned}
I_1 &= -\frac{t}{2\underline{\mu}} \int_{\Omega} \rho_t u_t^2 \\
&= \frac{t}{2\underline{\mu}} \int_{\Omega} [\partial_x(\rho u) - \partial_y(\rho \int_0^y \partial_x u(s) ds)] u_t u_t \\
&= -\frac{t}{\underline{\mu}} \int_{\Omega} \rho u \partial_x u_t u_t + \frac{t}{\underline{\mu}} \int_{\Omega} \rho \int_0^y \partial_x u(s) ds \partial_y u_t u_t \\
&\leq \frac{t}{\underline{\mu}} \|\rho u_t\|_{L_y^2 L_x^\infty} \|\partial_x u_t\|_{L^2} \|\nabla u\|_{L^2} + \frac{t}{\underline{\mu}} \|\rho u_t\|_{L_y^2 L_x^\infty} \left\| \int_0^y \partial_x u(s) ds \right\|_{L_x^2 L_y^\infty} \|\partial_y u_t\|_{L^2} \\
&\leq C \frac{t}{\underline{\mu}} (\|\partial_x \rho u_t + \rho \partial_x u_t\|_{L^2}^{\frac{1}{2}} \|\rho u_t\|_{L^2}^{\frac{1}{2}} + \|\rho u_t\|_{L^2}) \|\nabla u_t\|_{L^2} \|\nabla u\|_{L^2} \\
&\leq C \frac{t}{\underline{\mu}} [(M_0 \|u_t\|_{L^2}^{\frac{1}{2}} + \bar{\rho}^{\frac{1}{2}} \|\partial_x u_t\|_{L^2}^{\frac{1}{2}}) \|\rho u_t\|_{L^2}^{\frac{1}{2}} + \|\rho u_t\|_{L^2}] \|\nabla u_t\|_{L^2} \|\nabla u\|_{L^2} \\
&\leq C \left(\frac{t}{\underline{\mu}} \bar{\rho}^{\frac{1}{4}} M_0 \|\nabla u_t\|_{L^2}^{\frac{3}{2}} \|\nabla u\|_{L^2} \|\sqrt{\rho} u_t\|_{L^2}^{\frac{1}{2}} + \frac{t}{\underline{\mu}} \bar{\rho}^{\frac{3}{4}} \|\nabla u_t\|_{L^2}^{\frac{3}{2}} \|\nabla u\|_{L^2} \|\sqrt{\rho} u_t\|_{L^2}^{\frac{1}{2}} \right. \\
&\quad \left. + \frac{t}{\underline{\mu}} \bar{\rho}^{\frac{1}{2}} \|\nabla u_t\|_{L^2} \|\nabla u\|_{L^2} \|\sqrt{\rho} u_t\|_{L^2} \right).
\end{aligned} \tag{3.30}$$

Applying Young's inequality, we deduce that

$$\begin{aligned}
I_1 &\leq \epsilon t \|\nabla u_t\|^2 + Ct \bar{\rho} M_0^4 \|\sqrt{\rho} u_t\|^2 \|\nabla u\|^4 + Ct \bar{\rho}^3 \|\sqrt{\rho} u_t\|^2 \|\nabla u\|^4 \\
&\quad + Ct \bar{\rho} \|\sqrt{\rho} u_t\|^2 \|\nabla u\|^2,
\end{aligned} \tag{3.31}$$

where ϵ is some small number.

To estimate I_2 , taking the mass equation (1.1)₁ into consideration, we get

$$\begin{aligned}
I_2 &= -\frac{t}{\underline{\mu}} \int_{\Omega} (\rho u)_t \partial_x u u_t = -\frac{t}{\underline{\mu}} \int_{\Omega} \rho_t u \partial_x u u_t - \frac{t}{\underline{\mu}} \int_{\Omega} \rho u_t \partial_x u u_t \\
&= \frac{t}{\underline{\mu}} \int_{\Omega} [\partial_x \rho u - \partial_y \rho \int_0^y \partial_x u(s) ds] u \partial_x u u_t - \frac{t}{\underline{\mu}} \int_{\Omega} \rho u_t \partial_x u u_t = I_{21} + I_{22}.
\end{aligned}$$

It follows from Sobolev embedding inequality, Gagliardo-Nirenberg inequality, and Lemma 3.3 that

$$\begin{aligned}
I_{21} &= \frac{t}{\underline{\mu}} \int_{\Omega} [\partial_x \rho u u \partial_x u u_t - \partial_y \rho \int_0^y \partial_x u(s) ds u \partial_x u u_t] \\
&\leq \frac{t}{\underline{\mu}} \|\partial_x \rho\|_{L^\infty} \|u\|_{L^\infty} \|u\|_{L_x^2 L_y^\infty} \|\partial_x u\|_{L^2} \|u_t\|_{L_y^2 L_x^\infty} \\
&\quad + \frac{t}{\underline{\mu}} \|\partial_y \rho\|_{L^\infty} \|u\|_{L^\infty} \left\| \int_0^y \partial_x u(s) ds \right\|_{L_x^2 L_y^\infty} \|\partial_x u\|_{L^2} \|u_t\|_{L_y^2 L_x^\infty} \\
&\leq \frac{t}{\underline{\mu}} M_0 \|u\|_{H^2} \|\nabla u\|_{L^2}^2 \|\nabla u_t\|_{L^2} \\
&\leq C \frac{t}{\underline{\mu}} M_0 (\bar{\rho}^{1/2} M \|\sqrt{\rho} u_t\|_{L^2} + \bar{\rho} M \|\nabla u\|_{L^2}^2 + \bar{\rho}^{\frac{3}{2}} M^{\frac{3}{2}} \|\nabla u\|_{L^2}^{\frac{5}{2}} + \bar{\rho}^2 M^2 \|\nabla u\|_{L^2}^3) \|\nabla u\|_{L^2}^2 \|\nabla u_t\|_{L^2} \\
&\leq \epsilon t \|\nabla u_t\|^2 + Ct (M_0^2 \bar{\rho} M^2 \|\sqrt{\rho} u_t\|_{L^2}^2 \|\nabla u\|^4 + M_0^2 M^2 \bar{\rho}^2 \|\nabla u\|_{L^2}^8 + M_0^2 M^3 \bar{\rho}^3 \|\nabla u\|_{L^2}^9 \\
&\quad + M_0^2 M^4 \bar{\rho}^4 \|\nabla u\|_{L^2}^{10}),
\end{aligned} \tag{3.32}$$

where ϵ is some small number. Similarly, it holds that

$$\begin{aligned}
I_{22} &= -\frac{t}{\underline{\mu}} \int_{\Omega} \rho u_t \partial_x u u_t \\
&\leq \frac{t}{\underline{\mu}} \|\rho u_t\|_{L_y^2 L_x^\infty} \|\nabla u\|_{L^2} \|\nabla u_t\|_{L^2} \\
&\leq \epsilon t \|\nabla u_t\|^2 + Ct \bar{\rho} M_0^4 \|\sqrt{\rho} u_t\|^2 \|\nabla u\|^4 + Ct \bar{\rho}^3 \|\sqrt{\rho} u_t\|^2 \|\nabla u\|^4 + Ct \bar{\rho} \|\sqrt{\rho} u_t\|^2 \|\nabla u\|^2,
\end{aligned} \tag{3.33}$$

where ϵ is some small number.

To bound the term I_3 , we rewrite it as

$$\begin{aligned}
I_3 &= \frac{t}{\underline{\mu}} \int_{\Omega} (\rho \int_0^y \partial_x u(s) ds)_t \partial_y u u_t \\
&= \frac{t}{\underline{\mu}} \int_{\Omega} [\rho_t \int_0^y \partial_x u(s) ds \partial_y u u_t] + \frac{t}{\underline{\mu}} \int_{\Omega} [\rho \int_0^y \partial_x u_t(s) ds \partial_y u u_t] \\
&= -\frac{t}{\underline{\mu}} \int_{\Omega} [(\partial_x \rho u - \partial_y \rho \int_0^y \partial_x u(s) ds) \int_0^y \partial_x u(s) ds \partial_y u u_t] + \frac{t}{\underline{\mu}} \int_{\Omega} [\rho \int_0^y \partial_x u_t(s) ds \partial_y u u_t] \\
&= I_{31} + I_{32}.
\end{aligned}$$

Owing to Young's inequality and Sobolev embedding inequality, we deduce that

$$\begin{aligned}
I_{31} &\leq \frac{t}{\underline{\mu}} \|\partial_x \rho\|_{L^\infty} \|u\|_{L^\infty} \left\| \int_0^y \partial_x u(s) ds \right\|_{L_y^\infty L_x^2} \|\partial_y u\|_{L^2} \|u_t\|_{L_x^\infty L_y^2} \\
&\quad + \frac{t}{\underline{\mu}} \|\partial_y \rho\|_{L^\infty} \left\| \int_0^y \partial_x u(s) ds \right\|_{L^\infty} \left\| \int_0^y \partial_x u(s) ds \right\|_{L_y^\infty L_x^2} \|\partial_y u\|_{L^2} \|u_t\|_{L_x^\infty L_y^2} \\
&\leq \frac{t}{\underline{\mu}} M_0 \|u\|_{H^2} \|\nabla u\|_{L^2}^2 \|\nabla u_t\|_{L^2} \\
&\leq \epsilon t \|\nabla u_t\|_{L^2}^2 + Ct (M_0^2 \bar{\rho} M^2 \|\sqrt{\rho} u_t\|_{L^2}^2 \|\nabla u\|_{L^2}^4 + M_0^2 M^2 \bar{\rho}^2 \|\nabla u\|_{L^2}^8 \\
&\quad + M_0^2 M^3 \bar{\rho}^3 \|\nabla u\|_{L^2}^9 + M_0^2 M^4 \bar{\rho}^4 \|\nabla u\|_{L^2}^{10}),
\end{aligned} \tag{3.34}$$

where ϵ is some small number. A similar argument to I_{32} yields

$$\begin{aligned}
I_{32} &= \frac{t}{\underline{\mu}} \int_{\Omega} [\rho \int_0^y \partial_x u_t(s) ds \partial_y u u_t] \\
&\leq \frac{t}{\underline{\mu}} \|\rho u_t\|_{L_y^2 L_x^\infty} \left\| \int_0^y \partial_x u_t(s) ds \right\|_{L_x^2 L_y^\infty} \|\partial_y u\|_{L^2} \\
&\leq \frac{t}{\underline{\mu}} \|\rho u_t\|_{L_y^2 L_x^\infty} \|\nabla u\|_{L^2} \|\nabla u_t\|_{L^2} \\
&\leq \epsilon t \|\nabla u_t\|^2 + Ct \bar{\rho} M_0^4 \|\sqrt{\rho} u_t\|^2 \|\nabla u\|^4 + Ct \bar{\rho}^3 \|\sqrt{\rho} u_t\|^2 \|\nabla u\|^4 + Ct \bar{\rho} \|\sqrt{\rho} u_t\|^2 \|\nabla u\|^2,
\end{aligned} \tag{3.35}$$

where ϵ is some small number.

Concerning the last term I_4 , by direct estimates, one has

$$\begin{aligned}
I_4 &= -\frac{t}{\underline{\mu}} \int_{\Omega} \mu(\rho)_t \nabla u \nabla u_t \\
&= \frac{t}{\underline{\mu}} \int_{\Omega} [\partial_x \mu(\rho) u - \partial_y \mu(\rho) \int_0^y \partial_x u(s) ds] \nabla u \nabla u_t \\
&\leq \frac{t}{\underline{\mu}} \|\nabla \mu(\rho)\|_{L^\infty} \|u\|_{H^2} \|\nabla u\|_{L^2} \|\nabla u_t\|_{L^2} \\
&\leq C \frac{t}{\underline{\mu}} M_0 (\bar{\rho}^{1/2} M \|\sqrt{\rho} u_t\|_{L^2} + \bar{\rho} M \|\nabla u\|_{L^2}^2 + \bar{\rho}^{\frac{3}{2}} M^{\frac{3}{2}} \|\nabla u\|_{L^2}^{\frac{5}{2}} + \bar{\rho}^2 M^2 \|\nabla u\|_{L^2}^3) \|\nabla u\|_{L^2} \|\nabla u_t\|_{L^2} \\
&\leq \epsilon t \|\nabla u_t\|^2 + Ct (M_0^2 \bar{\rho} M^2 \|\sqrt{\rho} u_t\|_{L^2}^2 \|\nabla u\|_{L^2}^2 + M_0^2 M^2 \bar{\rho}^2 \|\nabla u\|_{L^2}^6 + M_0^2 M^3 \bar{\rho}^3 \|\nabla u\|_{L^2}^7 \\
&\quad + M_0^2 M^4 \bar{\rho}^4 \|\nabla u\|_{L^2}^8),
\end{aligned} \tag{3.36}$$

where ϵ is some small number.

Substituting all the above estimates (3.31)-(3.36) into (3.29) and choosing ϵ small enough, we deduce that

$$\begin{aligned}
&\frac{t}{2\underline{\mu}} \frac{d}{dt} \int_{\Omega} \rho u_t^2 + t \int_{\Omega} |\nabla u_t|^2 \\
&\leq Ct (\bar{\rho} M_0^4 \|\sqrt{\rho} u_t\|_{L^2}^2 \|\nabla u\|_{L^2}^4 + \bar{\rho}^3 \|\sqrt{\rho} u_t\|_{L^2}^2 \|\nabla u\|_{L^2}^4 + \bar{\rho} \|\sqrt{\rho} u_t\|_{L^2}^2 \|\nabla u\|_{L^2}^2 \\
&\quad + M_0^2 \bar{\rho} M^2 \|\sqrt{\rho} u_t\|_{L^2}^2 \|\nabla u\|_{L^2}^2 + M_0^2 M^2 \bar{\rho}^2 \|\nabla u\|_{L^2}^6 + M_0^2 M^3 \bar{\rho}^3 \|\nabla u\|_{L^2}^7 + M_0^2 M^4 \bar{\rho}^4 \|\nabla u\|_{L^2}^8 \\
&\quad + M_0^2 \bar{\rho} M^2 \|\sqrt{\rho} u_t\|_{L^2}^2 \|\nabla u\|_{L^2}^4 + M_0^2 M^2 \bar{\rho}^2 \|\nabla u\|_{L^2}^8 + M_0^2 M^3 \bar{\rho}^3 \|\nabla u\|_{L^2}^9 + M_0^2 M^4 \bar{\rho}^4 \|\nabla u\|_{L^2}^{10}).
\end{aligned} \tag{3.37}$$

Adding the term $\frac{1}{2\underline{\mu}} \int_{\Omega} \rho u_t^2$ into (3.37), one has

$$\begin{aligned}
&\frac{1}{2\underline{\mu}} \frac{d}{dt} (t \int_{\Omega} \rho u_t^2) + t \int_{\Omega} |\nabla u_t|^2 \\
&\leq Ct (\|\sqrt{\rho} u_t\|_{L^2}^2 (\bar{\rho} M_0^4 \|\nabla u\|_{L^2}^4 + \bar{\rho}^3 \|\nabla u\|_{L^2}^4 + \bar{\rho} \|\nabla u\|_{L^2}^2 + M_0^2 \bar{\rho} M^2 \|\nabla u\|_{L^2}^2 + M_0^2 \bar{\rho} M^2 \|\nabla u\|_{L^2}^4) \\
&\quad + Ct (M_0^2 M^2 \bar{\rho}^2 \|\nabla u\|_{L^2}^6 + M_0^2 M^3 \bar{\rho}^3 \|\nabla u\|_{L^2}^7 + M_0^2 M^4 \bar{\rho}^4 \|\nabla u\|_{L^2}^8 + M_0^2 M^2 \bar{\rho}^2 \|\nabla u\|_{L^2}^8 \\
&\quad + M_0^2 M^3 \bar{\rho}^3 \|\nabla u\|_{L^2}^9 + M_0^2 M^4 \bar{\rho}^4 \|\nabla u\|_{L^2}^{10}) + \frac{1}{2\underline{\mu}} \|\sqrt{\rho} u_t\|_{L^2}^2.
\end{aligned} \tag{3.38}$$

Integrating with respect to time on $[0, t]$ and using the fact that $\mu \geq \underline{\mu} > 0$, we get

$$\begin{aligned}
&t \int_{\Omega} \rho u_t^2 + 2\underline{\mu} \int_0^t s \int_{\Omega} |\nabla u_t|^2 ds \\
&\leq \int_0^t Cs \|\sqrt{\rho} u_t\|_{L^2}^2 (\bar{\rho} M_0^4 \|\nabla u\|_{L^2}^4 + \bar{\rho}^3 \|\nabla u\|_{L^2}^4 + \bar{\rho} \|\nabla u\|_{L^2}^2 + M_0^2 \bar{\rho} M^2 \|\nabla u\|_{L^2}^2 \\
&\quad + M_0^2 \bar{\rho} M^2 \|\nabla u\|_{L^2}^4) ds + \int_0^t Cs (M_0^2 M^2 \bar{\rho}^2 \|\nabla u\|_{L^2}^6 + M_0^2 M^3 \bar{\rho}^3 \|\nabla u\|_{L^2}^7 + M_0^2 M^4 \bar{\rho}^4 \|\nabla u\|_{L^2}^8 \\
&\quad + M_0^2 M^2 \bar{\rho}^2 \|\nabla u\|_{L^2}^8 + M_0^2 M^3 \bar{\rho}^3 \|\nabla u\|_{L^2}^9 + M_0^2 M^4 \bar{\rho}^4 \|\nabla u\|_{L^2}^{10}) ds + \int_0^t \|\sqrt{\rho} u_t\|_{L^2}^2 ds \\
&\leq \int_0^t C \mathcal{Q}_2(s) \|\sqrt{\rho} u_t\|_{L^2}^2 + 2\underline{\mu} \int_0^s \tau \int_{\Omega} |\nabla u_t|^2 d\tau ds + t \|\nabla u\|_{L^2}^2 \int_0^t C \mathcal{Q}_1 ds + \int_0^t \|\sqrt{\rho} u_t\|_{L^2}^2 ds,
\end{aligned}$$

where

$$\begin{aligned}\mathcal{Q}_1 = & M_0^2 M^2 \bar{\rho}^2 \|\nabla u\|_{L^2}^4 + M_0^2 M^3 \bar{\rho}^3 \|\nabla u\|_{L^2}^5 + M_0^2 M^4 \bar{\rho}^4 \|\nabla u\|_{L^2}^6 \\ & + M_0^2 M^2 \bar{\rho}^2 \|\nabla u\|_{L^2}^6 + M_0^2 M^3 \bar{\rho}^3 \|\nabla u\|_{L^2}^7 + M_0^2 M^4 \bar{\rho}^4 \|\nabla u\|_{L^2}^8,\end{aligned}$$

and

$$\mathcal{Q}_2 = \bar{\rho} M_0^4 \|\nabla u\|^4 + \bar{\rho}^3 \|\nabla u\|^4 + \bar{\rho} \|\nabla u\|^2 + M_0^2 \bar{\rho} M^2 \|\nabla u\|^2 + M_0^2 \bar{\rho} M^2 \|\nabla u\|^4.$$

Gronwall's inequality yields

$$\begin{aligned}& \sup_{t \in [0, T]} t \int_{\Omega} \rho u_t^2 + 2\bar{\mu} \int_0^T (t \int_{\Omega} |\nabla u_t|^2) dt \\ & \leq \int_0^T \|\sqrt{\rho} u_t\|_{L^2}^2 dt + \sup_{t \in [0, T]} t \|\nabla u\|_{L^2}^2 \int_0^T C \mathcal{Q}_1 dt \\ & + \int_0^T (t \|\nabla u\|_{L^2}^2 \int_0^t C \mathcal{Q}_1 ds + \int_0^t \|\sqrt{\rho} u_t\|_{L^2}^2 ds) \cdot C \mathcal{Q}_2 \cdot \exp\left(\int_0^t C \mathcal{Q}_2 ds\right) dt \\ & \leq \left(\sup_{t \in [0, T]} t \|\nabla u\|_{L^2}^2 \int_0^T C \mathcal{Q}_1 dt + \int_0^T \|\sqrt{\rho} u_t\|_{L^2}^2 dt\right) \cdot \int_0^T C \mathcal{Q}_2 dt \cdot \exp\left(\int_0^T C \mathcal{Q}_2 dt\right), \\ & \leq \left(\sup_{t \in [0, T]} t \|\nabla u\|_{L^2}^2 \int_0^T C \mathcal{Q}_1 dt + \int_0^T \|\sqrt{\rho} u_t\|_{L^2}^2 dt\right) \cdot \exp\left(\int_0^T C \mathcal{Q}_2 dt\right)\end{aligned}\tag{3.39}$$

By using Lemma 3.1 and (3.4), one obtains

$$\begin{aligned}\int_0^T C \mathcal{Q}_1 dt & \leq \int_0^T C (M_0^2 M^2 \bar{\rho}^2 \|\nabla u\|_{L^2}^4 + M_0^2 M^3 \bar{\rho}^3 \|\nabla u\|_{L^2}^5 + M_0^2 M^4 \bar{\rho}^4 \|\nabla u\|_{L^2}^6 \\ & + M_0^2 M^2 \bar{\rho}^2 \|\nabla u\|_{L^2}^6 + M_0^2 M^3 \bar{\rho}^3 \|\nabla u\|_{L^2}^7 + M_0^2 M^4 \bar{\rho}^4 \|\nabla u\|_{L^2}^8) dt \\ & \leq C(M^4 \bar{\rho}^3 + M^5 \bar{\rho}^4 + M^6 \bar{\rho}^5) \|u_0\|_{L^2}^2,\end{aligned}\tag{3.40}$$

and

$$\begin{aligned}\int_0^T C \mathcal{Q}_2 dt & \leq \int_0^T C (\bar{\rho} M_0^4 \|\nabla u\|^4 + \bar{\rho}^3 \|\nabla u\|^4 + \bar{\rho} \|\nabla u\|^2 + M_0^2 \bar{\rho} M^2 \|\nabla u\|^2 + M_0^2 \bar{\rho} M^2 \|\nabla u\|^4) dt \\ & \leq C(M^4 \bar{\rho}^2 + \bar{\rho}^2 + \bar{\rho}^4) \|u_0\|_{L^2}^2.\end{aligned}\tag{3.41}$$

Taking (3.21) and (3.6) into account and using Poincaré inequality, we obtain

$$\begin{aligned}& \sup_{t \in [0, T]} t \int_{\Omega} \rho u_t^2 + 2\bar{\mu} \int_0^T (t \int_{\Omega} |\nabla u_t|^2) dt \\ & \leq [C \bar{\rho} (M^4 \bar{\rho}^3 + M^5 \bar{\rho}^4 + M^6 \bar{\rho}^5) \|\nabla u_0\|_{L^2}^2 \\ & + (2 \ln 2 + 1) \frac{\bar{\mu}}{2} \|\nabla u_0\|_{L^2}^2] \cdot \exp [C (M^4 \bar{\rho}^2 + \bar{\rho}^2 + \bar{\rho}^4) \|u_0\|_{L^2}^2] \\ & \leq C (M^4 \bar{\rho}^4 + M^5 \bar{\rho}^5 + M^6 \bar{\rho}^6 + 1) \|\nabla u_0\|_{L^2}^2 \cdot \exp [C (M^4 \bar{\rho}^2 + \bar{\rho}^2 + \bar{\rho}^4)].\end{aligned}\tag{3.42}$$

Next we turn to the t^2 -weighted estimates. To do this, multiplying (3.38) by t , one has

$$\begin{aligned}
& \frac{t}{2\underline{\mu}} \frac{d}{dt} \left(t \int_{\Omega} \rho u_t^2 \right) + t^2 \int_{\Omega} |\nabla u_t|^2 \\
& \leq Ct^2 \|\sqrt{\rho} u_t\|_{L^2}^2 (\bar{\rho} M_0^4 \|\nabla u\|_{L^2}^4 + \bar{\rho}^3 \|\nabla u\|_{L^2}^4 + \bar{\rho} \|\nabla u\|_{L^2}^2 + M_0^2 \bar{\rho} M^2 \|\nabla u\|_{L^2}^2 + M_0^2 \bar{\rho} M^2 \|\nabla u\|_{L^2}^4) \\
& \quad + Ct^2 (M_0^2 M^2 \bar{\rho}^2 \|\nabla u\|_{L^2}^6 + M_0^2 M^3 \bar{\rho}^3 \|\nabla u\|_{L^2}^7 + M_0^2 M^4 \bar{\rho}^4 \|\nabla u\|_{L^2}^8 + M_0^2 M^2 \bar{\rho}^2 \|\nabla u\|_{L^2}^8 \\
& \quad + M_0^2 M^3 \bar{\rho}^3 \|\nabla u\|_{L^2}^9 + M_0^2 M^4 \bar{\rho}^4 \|\nabla u\|_{L^2}^{10}) + \frac{t}{2\underline{\mu}} \|\sqrt{\rho} u_t\|^2.
\end{aligned} \tag{3.43}$$

Similar to the method of (3.38), adding the term $\frac{t}{2\underline{\mu}} \int_{\Omega} \rho u_t^2$ into (3.43), we deduce that

$$\begin{aligned}
& \frac{1}{2\underline{\mu}} \frac{d}{dt} [t^2 \int_{\Omega} \rho u_t^2] + t^2 \int_{\Omega} |\nabla u_t|^2 \\
& \leq Ct^2 \|\sqrt{\rho} u_t\|_{L^2}^2 (\bar{\rho} M_0^4 \|\nabla u\|_{L^2}^4 + \bar{\rho}^3 \|\nabla u\|_{L^2}^4 + \bar{\rho} \|\nabla u\|_{L^2}^2 + M_0^2 \bar{\rho} M^2 \|\nabla u\|_{L^2}^2 + M_0^2 \bar{\rho} M^2 \|\nabla u\|_{L^2}^4) \\
& \quad + Ct^2 \|\nabla u\|_{L^2}^4 (M_0^2 M^2 \bar{\rho}^2 \|\nabla u\|_{L^2}^2 + M_0^2 M^3 \bar{\rho}^3 \|\nabla u\|_{L^2}^3 + M_0^2 M^4 \bar{\rho}^4 \|\nabla u\|_{L^2}^4 \\
& \quad + M_0^2 M^2 \bar{\rho}^2 \|\nabla u\|_{L^2}^4 + M_0^2 M^3 \bar{\rho}^3 \|\nabla u\|_{L^2}^5 + M_0^2 M^4 \bar{\rho}^4 \|\nabla u\|_{L^2}^6) + \frac{t}{\underline{\mu}} \|\sqrt{\rho} u_t\|_{L^2}^2.
\end{aligned} \tag{3.44}$$

Integrating with respect to time on $[0, t]$ and using the fact that $\mu \geq \underline{\mu} > 0$, we get

$$\begin{aligned}
& t^2 \int_{\Omega} \rho u_t^2 + 2\underline{\mu} \int_0^t s^2 \int_{\Omega} |\nabla u_t|^2 ds \\
& \leq \int_0^t C s^2 \|\sqrt{\rho} u_t\|_{L^2}^2 \mathcal{Q}_2 ds + \int_0^t C s^2 \|\nabla u\|_{L^2}^4 \mathcal{Q}_3 ds + \int_0^t 2s \|\sqrt{\rho} u_t\|_{L^2}^2 ds \\
& \leq \int_0^t C \mathcal{Q}_2 (s^2 \|\sqrt{\rho} u_t\|_{L^2}^2 + 2\underline{\mu} \int_0^s \tau^2 \int_{\Omega} |\nabla u_t|^2 d\tau) ds + \int_0^t C s^2 \|\nabla u\|_{L^2}^4 \mathcal{Q}_3 ds + \int_0^t 2s \|\sqrt{\rho} u_t\|_{L^2}^2 ds,
\end{aligned}$$

where

$$\begin{aligned}
\mathcal{Q}_3 = & M_0^2 M^2 \bar{\rho}^2 \|\nabla u\|_{L^2}^2 + M_0^2 M^3 \bar{\rho}^3 \|\nabla u\|_{L^2}^3 + M_0^2 M^4 \bar{\rho}^4 \|\nabla u\|_{L^2}^4 \\
& + M_0^2 M^2 \bar{\rho}^2 \|\nabla u\|_{L^2}^4 + M_0^2 M^3 \bar{\rho}^3 \|\nabla u\|_{L^2}^5 + M_0^2 M^4 \bar{\rho}^4 \|\nabla u\|_{L^2}^6.
\end{aligned}$$

Applying Gronwall's inequality, Lemma 3.1, (3.4) and Lemma 3.4, we obtain that

$$\begin{aligned}
& \sup_{t \in [0, T]} t^2 \int_{\Omega} \rho u_t^2 + 2\underline{\mu} \int_0^T (t^2 \int_{\Omega} |\nabla u_t|^2) dt \\
& \leq \sup_{t \in [0, T]} t^2 \|\nabla u\|_{L^2}^4 \int_0^T C \mathcal{Q}_3 dt + 2 \int_0^T t \|\sqrt{\rho} u_t\|_{L^2}^2 dt \\
& \quad + \left(\sup_{t \in [0, T]} t^2 \|\nabla u\|_{L^2}^4 \int_0^T C \mathcal{Q}_3 dt + \int_0^T t \|\sqrt{\rho} u_t\|_{L^2}^2 dt \right) \cdot \int_0^T C \mathcal{Q}_2 dt \cdot \exp \left(\int_0^T C \mathcal{Q}_2 dt \right) \\
& \leq \left(\sup_{t \in [0, T]} t^2 \|\nabla u\|_{L^2}^4 \int_0^T C \mathcal{Q}_3 dt + \int_0^T t \|\sqrt{\rho} u_t\|_{L^2}^2 dt \right) \cdot \exp \left(\int_0^T C \mathcal{Q}_2 dt \right) \\
& \leq C (M^4 \bar{\rho}^5 + M^5 \bar{\rho}^6 + M^6 \bar{\rho}^7 + \bar{\rho}) \|\nabla u_0\|_{L^2}^2 \cdot \exp [C (M^4 \bar{\rho}^2 + \bar{\rho}^2 + \bar{\rho}^4)],
\end{aligned} \tag{3.45}$$

where we use

$$\int_0^T C \mathcal{Q}_3 dt \leq C (M^4 \bar{\rho}^3 + M^5 \bar{\rho}^4 + M^6 \bar{\rho}^5) \|\nabla u_0\|_{L^2}^2.$$

Thus, we complete the proof of this lemma. \square

We also have the following decay estimates.

Lemma 3.6 *Assume that (ρ, u, w, P) is the unique strong solution to the system (1.1)-(1.4) on $\Omega \times (0, T)$ with the initial data satisfying the assumptions in Theorem 1.1 and (3.3)-(3.5). Then it holds that*

$$\sup_{t \in [0, T]} e^{\sigma t} \int_{\Omega} \rho u^2(t) + \underline{\mu} \int_0^T (e^{\sigma t} \int_{\Omega} |\nabla u|^2) dt \leq \bar{\rho} \|u_0\|_{L^2}^2, \quad (3.46)$$

and

$$\int_0^T (e^{\sigma t} \frac{2}{\underline{\mu}} \int_{\Omega} \rho u_t^2) dt + \sup_{t \in [0, T]} e^{\sigma t} \int_{\Omega} |\nabla u|^2 \leq C(1 + \frac{\bar{\rho}}{\|\rho_0\|_{L^p}}) \|u_0\|_{L^2}^2, \quad (3.47)$$

where $\sigma = \frac{\underline{\mu}}{C\|\rho_0\|_{L^p}}$ for any $p > 1$.

Proof. Multiplying the horizontal momentum equations $(1.1)_2$ by u and integrating by parts over Ω , we obtain

$$\frac{1}{2} \frac{d}{dt} \|\sqrt{\rho} u\|_{L^2}^2 + \int_{\Omega} \mu(\rho) |\nabla u|^2 = 0. \quad (3.48)$$

Moreover, noticing that

$$\begin{aligned} \|\sqrt{\rho} u\|_{L^2}^2 &= \int_{\Omega} \rho u^2 \leq \left(\int_{\Omega} \rho^p \right)^{\frac{1}{p}} \left(\int_{\Omega} u^{2q} \right)^{\frac{1}{q}} \leq \|\rho\|_{L^p} \|u\|_{L^{2q}}^2 \leq C \|\rho\|_{L^p} \|\nabla u\|_{L^2}^2 \\ &\leq C \|\rho_0\|_{L^p} \|\nabla u\|_{L^2}^2 \leq \sigma^{-1} \int_{\Omega} \mu(\rho) |\nabla u|^2, \end{aligned} \quad (3.49)$$

where $\sigma = \frac{\underline{\mu}}{C\|\rho_0\|_{L^p}}$ for $p > 1$.

Multiplying both sides of (3.48) and (3.49) by the term $e^{\sigma t}$, respectively, and collecting them, we have

$$\frac{d}{dt} [e^{\sigma t} \|\sqrt{\rho} u\|_{L^2}^2] + e^{\sigma t} \int_{\Omega} \mu(\rho) |\nabla u|^2 \leq 0.$$

It follows that

$$\sup_{t \in [0, T]} e^{\sigma t} \|\sqrt{\rho} u\|_{L^2}^2 + \underline{\mu} \int_0^T (e^{\sigma t} \int_{\Omega} |\nabla u|^2) dt \leq \|\sqrt{\rho_0} u_0\|_{L^2}^2 \leq \bar{\rho} \|u_0\|_{L^2}^2. \quad (3.50)$$

Recall that

$$2 \int_{\Omega} \rho u_t^2 + \frac{d}{dt} \int_{\Omega} \mu(\rho) |\nabla u|^2 \leq C \mathcal{Q} \|\nabla u\|_{L^2}^2. \quad (3.51)$$

Multiplying both sides of (3.51) by the term $e^{\sigma t}$ and adding the term $e^{\sigma t} \sigma \int_{\Omega} \mu(\rho) |\nabla u|^2$ to both sides, we get

$$\frac{d}{dt} (e^{\sigma t} \int_{\Omega} \mu(\rho) |\nabla u|^2) + e^{\sigma t} 2 \int_{\Omega} \rho u_t^2 \leq e^{\sigma t} \|\nabla u\|_{L^2}^2 C \mathcal{Q} + e^{\sigma t} \sigma \int_{\Omega} \mu(\rho) |\nabla u|^2. \quad (3.52)$$

Integrating with respect to time on $[0, t]$ and using the fact that $\mu \geq \underline{\mu} > 0$, we get

$$\begin{aligned}
& e^{\sigma t} \int_{\Omega} |\nabla u|^2 + \int_0^t \left(\frac{2}{\underline{\mu}} e^{\sigma s} \int_{\Omega} \rho u_t^2 \right) ds \\
& \leq \int_0^t \frac{CQ}{\underline{\mu}} e^{\sigma s} \|\nabla u\|_{L^2}^2 ds + \int_0^t e^{\sigma s} \frac{\sigma}{\underline{\mu}} \int_{\Omega} \mu(\rho) |\nabla u|^2 ds + \int_{\Omega} \frac{\mu(\rho_0)}{\underline{\mu}} |\nabla u_0|^2 \\
& \leq \int_0^t \frac{CQ}{\underline{\mu}} (e^{\sigma s} \|\nabla u\|_{L^2}^2 + \int_0^s \left(\frac{2}{\underline{\mu}} e^{\sigma \tau} \int_{\Omega} \rho u_t^2 d\tau \right) ds \\
& \quad + \int_0^t e^{\sigma s} \frac{\sigma}{\underline{\mu}} \int_{\Omega} \mu(\rho) |\nabla u|^2 ds + \int_{\Omega} \frac{\mu(\rho_0)}{\underline{\mu}} |\nabla u_0|^2
\end{aligned}$$

By Gronwall inequality in integral form

$$\begin{aligned}
& \sup_{t \in [0, T]} e^{\sigma t} \int_{\Omega} |\nabla u|^2 + \int_0^T \left(e^{\sigma t} \frac{2}{\underline{\mu}} \int_{\Omega} \rho u_t^2 \right) dt \\
& \leq \int_0^T \left(e^{\sigma t} \frac{\sigma}{\underline{\mu}} \int_{\Omega} \mu(\rho) |\nabla u|^2 \right) dt + \int_{\Omega} \mu(\rho_0) \frac{|\nabla u_0|^2}{\underline{\mu}} \\
& \quad + \left[\int_0^T \left(e^{\sigma t} \frac{\sigma}{\underline{\mu}} \int_{\Omega} \mu(\rho) |\nabla u|^2 \right) + \int_{\Omega} \mu(\rho_0) \frac{|\nabla u_0|^2}{\underline{\mu}} dt \right] \cdot \int_0^T \frac{CQ}{\underline{\mu}} dt \cdot \exp\left(\int_0^T \frac{CQ}{\underline{\mu}} dt\right) \\
& \leq \left(1 + \frac{\bar{\rho}}{\|\rho_0\|_{L^p}}\right) (1 + 2\ln 2) \frac{\bar{\mu}}{\underline{\mu}} \|\nabla u_0\|_{L^2}^2.
\end{aligned} \tag{3.53}$$

Hence, it follows that

$$\int_0^T \left(e^{\sigma t} \frac{2}{\underline{\mu}} \int_{\Omega} \rho u_t^2 \right) dt + \sup_{t \in [0, T]} e^{\sigma t} \int_{\Omega} |\nabla u|^2 \leq C \left(1 + \frac{\bar{\rho}}{\|\rho_0\|_{L^p}}\right) \|\nabla u_0\|_{L^2}^2. \tag{3.54}$$

This completes the proof of Lemma 3.6. \square

Before stating the H^3 -regularity of the velocity u , we present an estimate for $\int_0^T \|\nabla u_t\|_{L^2}^2 ds$ as follows.

Lemma 3.7 *Assume that (ρ, u, w, P) is the unique strong solution to the system (1.1)-(1.4) on $\Omega \times [0, T]$, and the initial data satisfies the assumptions in Theorem 1.1 and (3.3)-(3.5). Then, for every $t \in [0, T]$, it holds that*

$$\|\sqrt{\rho} u_t\|_{L^2}^2 + \underline{\mu} \int_0^t \|\nabla u_t(s)\|_{L^2}^2 ds \leq (\bar{\rho} \|V_1\|_{L^2}^2 + CM_0^2 \Theta_3 \|\nabla u_0\|_{L^2}^2) \exp(C \bar{\rho} \Theta_3 \|\nabla u_0\|_{L^2}^2), \tag{3.55}$$

where $\Theta_3 = \bar{\rho} M^2 + \bar{\rho}^3 M^2 + \bar{\rho}^4 M^3 + \bar{\rho}^5 M^4$.

Proof. Multiplying (3.27) by u_t and integrating by parts over Ω , we have

$$\begin{aligned}
& \frac{1}{2} \frac{d}{dt} \int_{\Omega} \rho u_t^2 + \int_{\Omega} \mu(\rho) |\nabla u_t|^2 \\
& = - \int_{\Omega} \frac{\rho_t}{2} u_t^2 - \int_{\Omega} (\rho u)_t \partial_x u u_t + \int_{\Omega} \left(\rho \int_0^y \partial_x u(s) ds \right)_t \partial_y u u_t - \frac{1}{2} \int_{\Omega} \mu_t(\rho) \frac{d|\nabla u|^2}{dt} \\
& = - \int_{\Omega} \frac{\rho_t}{2} u_t^2 - \int_{\Omega} \rho_t u \partial_x u u_t - \int_{\Omega} \rho u_t \partial_x u u_t + \int_{\Omega} \left[\rho_t \int_0^y \partial_x u(s) ds \partial_y u u_t \right. \\
& \quad \left. + \int_{\Omega} \left[\rho \int_0^y \partial_x u_t(s) ds \partial_y u u_t \right] - \int_{\Omega} \mu_t(\rho) \nabla u \nabla u_t \right] = \sum_{i=1}^6 J_i.
\end{aligned} \tag{3.56}$$

By direct estimates and Young inequality, we have

$$\begin{aligned}
J_1 &= \frac{1}{2} \int_{\Omega} (u \partial_x \rho - \int_0^y \partial_x u(s) ds \partial_y \rho) u_t u_t \\
&\leq \bar{\rho}^{\frac{1}{2}} \|u\|_{H^2} \|\sqrt{\rho} u_t\|_{L^2} \|\nabla u_t\|_{L^2} \\
&\leq C \bar{\rho} \|u\|_{H^2}^2 \|\sqrt{\rho} u_t\|_{L^2}^2 + \frac{\mu}{2} \|\nabla u_t\|_{L^2}^2.
\end{aligned} \tag{3.57}$$

It follows from Hölder's inequality and Gagliardo-Nirenberg inequality that

$$\begin{aligned}
J_2 &= - \int_{\Omega} \rho_t u \partial_x u u_t = \|u\|_{H^2} \|\partial_x \rho\|_{L^\infty} \|\partial_x u\|_{L^2} \|u_t\|_{L_x^\infty L_y^2} \|u\|_{L_x^2 L_y^\infty} \\
&\quad + \|u\|_{H^2} \|\partial_x \rho\|_{L^\infty} \|\partial_x u\|_{L^2} \|u_t\|_{L_x^\infty L_y^2} \left\| \int_0^y \partial_x u(s) ds \right\|_{L_x^2 L_y^\infty} \\
&\leq C \|\nabla \rho\|_{L^\infty}^2 \|u\|_{H^2}^2 \|\nabla u\|_{L^2}^4 + \frac{\mu}{2} \|\nabla u_t\|_{L^2}^2.
\end{aligned} \tag{3.58}$$

Similarly,

$$\begin{aligned}
J_3 &= - \int_{\Omega} \rho u_t \partial_x u u_t \\
&\leq \sqrt{\bar{\rho}} \|\sqrt{\rho} u_t\|_{L^2} \|u_t\|_{L_x^\infty L_y^2} \|\partial_x u\|_{L_x^2 L_y^\infty} \\
&\leq C \bar{\rho} \|\sqrt{\rho} u_t\|_{L^2}^2 \|u\|_{H^2}^2 + \frac{\mu}{2} \|\nabla u_t\|_{L^2}^2.
\end{aligned} \tag{3.59}$$

Applying anisotropic Sobolev embedding theorem and Young inequality, one has

$$\begin{aligned}
J_4 &= \int_{\Omega} [(u \partial_x \rho - \int_0^y \partial_x u(s) ds \partial_y \rho) \int_0^y \partial_x u(s) ds \partial_y u u_t] \\
&\leq \|u\|_{L^\infty} \|\partial_x \rho\|_{L^\infty} \left\| \int_0^y \partial_x u(s) ds \right\|_{L_x^2 L_y^\infty} \|\partial_y u\|_{L^2} \|u_t\|_{L_x^\infty L_y^2} \\
&\quad + \left\| \int_0^y \partial_x u(s) ds \right\|_{L^\infty} \|\partial_y \rho\|_{L^\infty} \left\| \int_0^y \partial_x u(s) ds \right\|_{L_x^2 L_y^\infty} \|\partial_y u\|_{L^2} \|u_t\|_{L_x^\infty L_y^2} \\
&\leq C \|\nabla \rho\|_{L^\infty}^2 \|u\|_{H^2}^2 \|\nabla u\|_{L^2}^4 + \frac{\mu}{2} \|\nabla u_t\|_{L^2}^2,
\end{aligned} \tag{3.60}$$

$$\begin{aligned}
J_5 &= \int_{\Omega} [\rho \int_0^y \partial_x u_t(s) ds \partial_y u u_t] \\
&\leq \left\| \int_0^y \partial_x u_t(s) ds \right\|_{L_x^2 L_y^\infty} \|\partial_y u\|_{L_y^2 L_x^\infty} \|\rho u_t\|_{L^2} \\
&\leq C \bar{\rho} \|\sqrt{\rho} u_t\|_{L^2}^2 \|u\|_{H^2}^2 + \frac{\mu}{2} \|\nabla u_t\|_{L^2}^2,
\end{aligned} \tag{3.61}$$

and

$$J_6 = \int_{\Omega} \mu_t(\rho) \nabla u \nabla u_t \leq C \|\nabla \rho\|_{L^\infty}^2 \|u\|_{H^2}^2 \|\nabla u\|_{L^2}^4 + \frac{\mu}{2} \|\nabla u_t\|_{L^2}^2. \tag{3.62}$$

Substituting the above estimates J_1 - J_6 into (3.57) we obtain

$$\frac{d}{dt} \int_{\Omega} \rho u_t^2 + \underline{\mu} \int_{\Omega} |\nabla u_t|^2 \leq C(\bar{\rho} \|u\|_{H^2}^2 \|\sqrt{\rho} u_t\|_{L^2}^2 + M_0^2 \|\nabla u\|_{L^2}^4 \|u\|_{H^2}^2). \tag{3.63}$$

Then, integrating it over $(0, t) \subset (0, T)$ and using Lemma 3.1-3.3, one has

$$\begin{aligned}
& \|\sqrt{\rho}u_t\|_{L^2}^2 + \underline{\mu} \int_0^t \int_{\Omega} |\nabla u_t|^2 ds \\
& \leq [\|\sqrt{\rho}u_t(0)\|_{L^2}^2 + C \int_0^T (M_0^2 \|\nabla u\|_{L^2}^4 \|u\|_{H^2}^2) ds] \exp(C\bar{\rho} \int_0^T \|u\|_{H^2}^2 ds) \\
& \leq [\bar{\rho} \|V_1\|_{L^2}^2 + CM_0^2 \Theta_3 \|\nabla u_0\|_{L^2}^2] \exp(C\bar{\rho} \Theta_3 \|\nabla u_0\|_{L^2}^2),
\end{aligned} \tag{3.64}$$

where $\Theta_3 = \bar{\rho}M^2 + \bar{\rho}^3M^2 + \bar{\rho}^4M^3 + \bar{\rho}^5M^4$. Then, we complete the proof of this lemma. \square

Remark 3.1 When we deal with the term of $\|\sqrt{\rho}u_t(0)\|_{L^2}$, we first need to regularize the initial data (ρ_0, u_0, P_0) . That is, for each $\delta \in (0, 1)$, that we choose $\rho_0^\delta \in C^{2+\alpha}$ such that $0 < \delta \leq \rho_0^\delta \leq \rho_0 + 1$, $\rho_0^\delta \rightarrow \rho_0$ in $W^{2,2}$ as $\delta \rightarrow 0$, and denote by $(u_0^\delta, P_0^\delta) \in H^1(\Omega) \times L^2(\Omega)$ a solution to the problem

$$-\rho_0^\delta u_0^\delta \partial_x u_0^\delta + \rho_0^\delta \int_0^y \partial_x u_0^\delta(s) ds \partial_y u_0^\delta - \partial_x P_0^\delta + \partial_x(\mu(\rho_0^\delta) \partial_x u_0^\delta) + \partial_y(\mu(\rho_0^\delta) \partial_y u_0^\delta) := \rho_0^\delta V_1. \tag{3.65}$$

In fact, with such regularized initial data and the compatibility condition (3.65), we can get corresponding solutions $(\rho^\delta, u^\delta, P^\delta)$ satisfying the estimates in Lemma 3.1-3.9 and these estimates are independent of δ . Then we use the standard compactness arguments to get the desired result.

With the help of Lemmas 3.4-3.7, we can now estimate $\|u\|_{L^1(0,T;H^3(\Omega))}$.

Lemma 3.8 Assume that (ρ, u, w, P) is the unique strong solution to the system (1.1)-(1.4) on $\Omega \times (0, T)$ with the initial data satisfying the assumptions in Theorem 1.1 and (3.3)-(3.5). Then it holds that

$$\begin{aligned}
\int_0^T \|u\|_{H^3} dt & \leq \tilde{C} \Theta_5 (\bar{\rho}C + M_0^2 \Theta_3 \|\nabla u_0\|_{L^2}^2) \exp(C\bar{\rho} \Theta_3 \|\nabla u_0\|_{L^2}^2) \sqrt{T_0} \\
& + \tilde{C} \Theta_5 (\bar{\rho} \|\nabla u_0\|_{L^2}^2 \cdot \Theta_1 \cdot \exp(\Theta_2))^{\frac{1}{2}} \frac{1}{\sqrt{T_0}} + \tilde{C} M^5 \bar{\rho}^2 \|\nabla u_0\|_{L^2}^2 + \tilde{C} \Theta_6 \bar{\rho} \|\nabla u_0\|_{L^2}^2,
\end{aligned} \tag{3.66}$$

where $\Theta_5 = M^5 \bar{\rho} + M^4$, $\Theta_6 = M^5 \bar{\rho} + M^5 \bar{\rho}^3 + M^{\frac{11}{2}} \bar{\rho}^{\frac{3}{2}} + M^6 \bar{\rho}^2 + M^6 \bar{\rho}^4 + M^7 \bar{\rho}^5$, and $T_0 < T$ which is to be determined later.

Proof. It follows from the regularity theory for the density-dependent hydrostatic Stokes system (see Lemma 2.2) that

$$\|u\|_{H^3} \leq \tilde{C} M^3 \|\rho u_t + \rho u \partial_x u + \rho \int_0^y \partial_x u(s) ds \partial_y u\|_{H^1} \triangleq L_1 + L_2, \tag{3.67}$$

where

$$L_1 = \tilde{C} M^3 (\|\sqrt{\rho} \sqrt{\rho} u_t\|_{L^2} + \|\rho u \partial_x u\|_{L^2} + \|\rho \int_0^y \partial_x u(s) ds \partial_y u\|_{L^2}),$$

and

$$\begin{aligned}
L_2 & = \tilde{C} M^3 (\|\nabla \rho u_t\|_{L^2} + \|\rho \nabla u_t\|_{L^2} + \|\nabla \rho u \partial_x u\|_{L^2} + \|\rho \nabla u \partial_x u\|_{L^2} + \|\rho u \nabla \partial_x u\|_{L^2} \\
& + \|\nabla \rho \int_0^y \partial_x u(s) ds \partial_y u\|_{L^2} + \|\rho \partial_x u \partial_y u\|_{L^2} + \|\rho \int_0^y \partial_{xx} u(s) ds \partial_y u\|_{L^2} \\
& + \|\rho \int_0^y \partial_x u(s) ds \nabla \partial_y u\|_{L^2}).
\end{aligned} \tag{3.68}$$

Similar to the proof of (3.9) in Lemma 3.3, we obtain

$$\begin{aligned}
L_1 &\leq \tilde{C}M^3(\bar{\rho}^{1/2}\|\sqrt{\rho}u_t\|_{L^2} + \bar{\rho}\|\nabla u\|_{L^2}^2 + \bar{\rho}\|\nabla u\|_{L^2}^{\frac{5}{2}}\|\nabla^2 u\|_{L^2}^{\frac{1}{2}} + \bar{\rho}\|\nabla u\|_{L^2}^{\frac{3}{2}}\|\nabla^2 u\|_{L^2}^{\frac{1}{2}}) \\
&\leq \frac{1}{2}\|\nabla u\|_{H^2} + \tilde{C}M^3\bar{\rho}^{1/2}\|\sqrt{\rho}u_t\|_{L^2} + \tilde{C}\bar{\rho}M^3\|\nabla u\|_{L^2}^2 + \tilde{C}\bar{\rho}^{\frac{3}{2}}M^{\frac{9}{2}}\|\nabla u\|_{L^2}^{\frac{5}{2}} + \tilde{C}\bar{\rho}^2M^6\|\nabla u\|_{L^2}^3.
\end{aligned} \tag{3.69}$$

We rewrite L_2 as follow

$$\begin{aligned}
L_2 &\leq \tilde{C}M^3\|\nabla\rho u_t\|_{L^2} + \tilde{C}M^3\|\rho\nabla u_t\|_{L^2} + \tilde{C}M^3\|\nabla\rho u\partial_x u\|_{L^2} + \tilde{C}M^3\|\rho\nabla u\nabla u\|_{L^2} \\
&\quad + \tilde{C}M^3\|\rho u\nabla^2 u\|_{L^2} + \tilde{C}M^3\|\nabla\rho\int_0^y\partial_x u(s)ds\partial_y u\|_{L^2} + \tilde{C}M^3\|\rho\int_0^y\partial_{xx}u(s)ds\partial_y u\|_{L^2} \\
&\quad + \tilde{C}M^3\|\rho\int_0^y\partial_x u(s)ds\nabla\partial_y u\|_{L^2}) = \sum_{i=1}^8 J_{2i}.
\end{aligned} \tag{3.70}$$

By direct estimates, one has

$$L_{21} + L_{22} \leq \tilde{C}M^4\|\nabla u_t\|_{L^2} + \tilde{C}M^3\bar{\rho}\|\nabla u_t\|_{L^2}. \tag{3.71}$$

Applying Hölder's inequality, (3.10) and Gagliardo-Nirenberg inequality, we deduce that

$$\begin{aligned}
L_{23} &\leq \tilde{C}M^3M_0\|u\|_{H^2}\|\partial_x u\|_{L^2} \\
&\leq \tilde{C}M^3M_0\|\nabla u\|_{L^2}(\bar{\rho}^{1/2}M\|\sqrt{\rho}u_t\|_{L^2} + \bar{\rho}M\|\nabla u\|_{L^2}^2 + \bar{\rho}^{\frac{3}{2}}M^{\frac{3}{2}}\|\nabla u\|_{L^2}^{\frac{5}{2}} + \bar{\rho}^2M^2\|\nabla u\|_{L^2}^3) \\
&\leq \tilde{C}M^5\bar{\rho}^{\frac{1}{2}}\|\sqrt{\rho}u_t\|_{L^2}\|\nabla u\|_{L^2} + \tilde{C}M^5\bar{\rho}\|\nabla u\|_{L^2}^3 + \tilde{C}M^{\frac{11}{2}}\bar{\rho}^{\frac{3}{2}}\|\nabla u\|_{L^2}^{\frac{7}{2}} + \tilde{C}M^6\bar{\rho}^2\|\nabla u\|_{L^2}^4,
\end{aligned} \tag{3.72}$$

$$\begin{aligned}
L_{24} &\leq \tilde{C}M^3\bar{\rho}\|\nabla u\|_{L_y^2L_x^\infty}\|\nabla u\|_{L_x^2L_y^\infty} \leq \tilde{C}M^3\bar{\rho}\|u\|_{H^2}^2 \\
&\leq \tilde{C}M^3\bar{\rho}(\bar{\rho}^{1/2}M\|\sqrt{\rho}u_t\|_{L^2} + \bar{\rho}M\|\nabla u\|_{L^2}^2 + \bar{\rho}^{\frac{3}{2}}M^{\frac{3}{2}}\|\nabla u\|_{L^2}^{\frac{5}{2}} + \bar{\rho}^2M^2\|\nabla u\|_{L^2}^3)^2 \\
&\leq \tilde{C}M^3\bar{\rho}(\bar{\rho}M^2\|\sqrt{\rho}u_t\|_{L^2}^2 + \bar{\rho}^2M^2\|\nabla u\|_{L^2}^4 + \bar{\rho}^3M^3\|\nabla u\|_{L^2}^5 + \bar{\rho}^4M^4\|\nabla u\|_{L^2}^6) \\
&\leq \tilde{C}M^5\bar{\rho}^2\|\sqrt{\rho}u_t\|_{L^2}^2 + \tilde{C}M^5\bar{\rho}^3\|\nabla u\|_{L^2}^4 + \tilde{C}M^6\bar{\rho}^4\|\nabla u\|_{L^2}^5 + \tilde{C}M^7\bar{\rho}^5\|\nabla u\|_{L^2}^6,
\end{aligned} \tag{3.73}$$

and

$$L_{25} \leq \tilde{C}M^3\bar{\rho}\|u\|_{H^2}^2. \tag{3.74}$$

By Sobolev and Poincaré inequalities, We estimate L_{26} as follows

$$L_{26} \leq \tilde{C}M^3\|\nabla\rho\|_{L^\infty}\int_0^y\partial_x u(s)ds\|\partial_y u\|_{L_x^\infty L_y^2} \leq \tilde{C}M^3M_0\|\nabla u\|_{L^2}\|u\|_{H^2}. \tag{3.75}$$

Similarly, one has

$$L_{27} + L_{28} \leq \tilde{C}M^3\bar{\rho}\|u\|_{H^2}^2. \tag{3.76}$$

Substituting the above estimates L_{21} - L_{28} into (3.70), we have

$$\begin{aligned}
L_2 &\leq \tilde{C}M^4\|\nabla u_t\|_{L^2} + \tilde{C}M^3\bar{\rho}\|\nabla u_t\|_{L^2} + \tilde{C}M^5\bar{\rho}^2\|\sqrt{\rho}u_t\|_{L^2}^2 \\
&\quad + \tilde{C}M^5\bar{\rho}^{\frac{1}{2}}\|\sqrt{\rho}u_t\|_{L^2}\|\nabla u\|_{L^2} + \tilde{C}M^5\bar{\rho}\|\nabla u\|_{L^2}^3 + \tilde{C}M^{\frac{11}{2}}\bar{\rho}^{\frac{3}{2}}\|\nabla u\|_{L^2}^{\frac{7}{2}} \\
&\quad + \tilde{C}M^6\bar{\rho}^2\|\nabla u\|_{L^2}^4 + \tilde{C}M^5\bar{\rho}^3\|\nabla u\|_{L^2}^4 + \tilde{C}M^6\bar{\rho}^4\|\nabla u\|_{L^2}^5 + \tilde{C}M^7\bar{\rho}^5\|\nabla u\|_{L^2}^6.
\end{aligned} \tag{3.77}$$

Combining (3.69) and (3.77), one has

$$\begin{aligned}
\int_0^T \|u\|_{H^3} dt &\leq \int_0^T [\tilde{C}M^3\bar{\rho}^{1/2}\|\sqrt{\rho}u_t\|_{L^2} + \tilde{C}M^4\|\nabla u_t\|_{L^2} + \tilde{C}M^3\bar{\rho}\|\nabla u_t\|_{L^2} \\
&\quad + \tilde{C}M^5\bar{\rho}^{1/2}\|\sqrt{\rho}u_t\|_{L^2}\|\nabla u\|_{L^2} + \tilde{C}M^5\bar{\rho}^2\|\sqrt{\rho}u_t\|_{L^2}^2 + \tilde{C}\bar{\rho}M^3\|\nabla u\|_{L^2}^2 \\
&\quad + \tilde{C}\bar{\rho}^{3/2}M^{9/2}\|\nabla u\|_{L^2}^{5/2} + \tilde{C}\bar{\rho}^2M^6\|\nabla u\|_{L^2}^3 + \tilde{C}M^5\bar{\rho}\|\nabla u\|_{L^2}^3 + \tilde{C}M^{11/2}\bar{\rho}^{3/2}\|\nabla u\|_{L^2}^{7/2} \\
&\quad + \tilde{C}M^6\bar{\rho}^2\|\nabla u\|_{L^2}^4 + \tilde{C}M^5\bar{\rho}^3\|\nabla u\|_{L^2}^4 + \tilde{C}M^6\bar{\rho}^4\|\nabla u\|_{L^2}^5 + \tilde{C}M^7\bar{\rho}^5\|\nabla u\|_{L^2}^6] dt \\
&\leq \tilde{C}\Theta_5 \int_0^T \|\nabla u_t\|_{L^2} dt + \tilde{C}M^5(\bar{\rho}^2 + 1) \int_0^T \|\sqrt{\rho}u_t\|_{L^2}^2 dt + \tilde{C}\Theta_6 \int_0^T \|\nabla u\|_{L^2}^2 dt,
\end{aligned} \tag{3.78}$$

where $\Theta_5 = M^3\bar{\rho} + M^4 + M^3\bar{\rho}^{1/2}$ and $\Theta_6 = M^5\bar{\rho} + M^5\bar{\rho}^3 + M^{11/2}\bar{\rho}^{3/2} + M^6\bar{\rho}^2 + M^6\bar{\rho}^4 + M^7\bar{\rho}^5$. Then, in order to obtain the estimate of $\|u\|_{L^1(0,T;H^3(\Omega))}$, we have to deal with the term of

$$\int_0^T \|\nabla u_t(t)\|_{L^2} dt.$$

Applying Hölder's inequality, (3.26) and (3.55), one has

$$\begin{aligned}
\int_0^T \|\nabla u_t\|_{L^2} dt &\leq \left(\int_0^{T_0} \|\nabla u_t\|_{L^2}^2 dt\right)^{1/2} \sqrt{T_0} + \left(\int_{T_0}^T t^2 \|\nabla u_t\|_{L^2}^2 dt\right)^{1/2} \left(\int_{T_0}^T t^{-2} dt\right)^{1/2} \\
&\leq C[\bar{\rho} + M_0^2\Theta_3\|\nabla u_0\|_{L^2}^2] \exp(C\bar{\rho}\Theta_3\|\nabla u_0\|_{L^2}^2) \sqrt{T_0} \\
&\quad + C[\bar{\rho}\|\nabla u_0\|_{L^2}^2 \cdot \Theta_1 \cdot \exp(\Theta_2)]^{1/2} \frac{1}{\sqrt{T_0}},
\end{aligned} \tag{3.79}$$

where $T_0 < T$ is a small number which is determined later.

Thus, substituting (3.79) into (3.78) and using Poincaré inequalities, we obtain that

$$\begin{aligned}
\int_0^T \|u\|_{H^3} dt &\leq \tilde{C}\Theta_5[\bar{\rho} + M_0^2\Theta_3\|\nabla u_0\|_{L^2}^2] \exp(C\bar{\rho}\Theta_3\|\nabla u_0\|_{L^2}^2) \sqrt{T_0} \\
&\quad + \tilde{C}\Theta_5[\bar{\rho}\|\nabla u_0\|_{L^2}^2 \cdot \Theta_1 \cdot \exp(\Theta_2)]^{1/2} \frac{1}{\sqrt{T_0}} \\
&\quad + \tilde{C}M^5(\bar{\rho}^2 + 1)\|\nabla u_0\|_{L^2}^2 + \tilde{C}\Theta_6\bar{\rho}\|\nabla u_0\|_{L^2}^2.
\end{aligned} \tag{3.80}$$

Here the proof of this lemma is completed. \square

In view of (3.80), we are now in a position to prove that $\|\nabla \rho(t)\|_{L^\infty} + \|\nabla^2 \rho(t)\|_{L^2} \leq 4M_0$, provided the initial data $\|\nabla u_0\|_{L^2}$ and the time T_0 are suitably small in some sense.

Lemma 3.9 *Assume that (ρ, u, w, P) is the unique strong solution to the system (1.1)-(1.4) on $\Omega \times [0, T]$, and the initial data satisfies the assumptions in Theorem 1.1 and (3.3)-(3.5). Moreover, suppose that there exists a positive number ϵ_0 , depending on Ω , $\sup_\Omega \rho_0 = \bar{\rho}$, $\|\mu\|_{C^2}$, $\|\nabla \rho_0\|_{L^\infty}$, $\|\nabla^2 \rho_0\|_{L^2}$, such that*

$$\|\nabla u_0\|_{L^2} \leq \epsilon_0.$$

Then it holds that

$$\sup_{t \in [0, T]} \|\nabla \rho(t)\|_{L^\infty} + \sup_{t \in [0, T]} \|\nabla^2 \rho(t)\|_{L^2} \leq 4M_0. \tag{3.81}$$

Proof. Recall that

$$\|\nabla \rho\|_{L^\infty} \leq \|\nabla \rho_0\|_{L^\infty} \exp\left(\int_0^t \|u(s)\|_{H^3} ds\right), \quad (3.82)$$

and

$$\|\nabla^2 \rho\|_{L^2} \leq (\|\nabla \rho_0\|_{L^\infty} + \|\nabla^2 \rho_0\|_{L^2}) \exp\left(\int_0^t \|u(s)\|_{H^3} ds\right). \quad (3.83)$$

Now, we can choose some small positive constant ϵ_0 and then time $T_0 < T$, satisfying

$$\begin{aligned} & \tilde{C}\Theta_5[\bar{\rho} + M_0^2\Theta_3\epsilon_0^2] \exp(C\bar{\rho}\Theta_3\epsilon_0^2) \sqrt{T_0} \\ & + \tilde{C}\Theta_5[\bar{\rho}\epsilon_0^2 \cdot \Theta_1 \cdot \exp(\Theta_2)]^{\frac{1}{2}} \frac{1}{\sqrt{T_0}} + \tilde{C}M^5\bar{\rho}^2\epsilon_0^2 + \tilde{C}\Theta_6\bar{\rho}\epsilon_0^2 \leq \ln 2 \end{aligned} \quad (3.84)$$

Thus, if $\|\nabla u_0\|_{L^2} \leq \epsilon_0$, it follows from Lemma 3.8 that

$$\|\nabla \rho\|_{L^\infty} + \|\nabla^2 \rho\|_{L^2} \leq 2(2\|\rho_0\|_{L^\infty} + \|\nabla^2 \rho_0\|_{L^2}) \leq 4(\|\nabla \rho_0\|_{L^\infty} + \|\nabla^2 \rho_0\|_{L^2}) = 4M_0.$$

Here we complete the proof of Lemma 3.9. \square

3.2 Proof of Theorem 1.2

With the above uniform estimates in hand, we are ready to prove Theorem 1.2.

Proof of Theorem 1.2. According to Lemma 2.4, there exists a $T_* > 0$ such that the primitive equations have a unique local strong solution (ρ, u, w, P) on $[0, T_*]$. Next, we apply the above a priori estimates in Section 3.1 to extend the local solution to a global one.

Since $\|\nabla \rho_0\|_{L^\infty} + \|\nabla^2 \rho_0\|_{L^2} = M_0 < 8M_0$, and thanks to the continuity of $\|\nabla \rho(t)\|_{L^\infty} + \|\nabla^2 \rho(t)\|_{L^2}$ and $\|\nabla u(t)\|_{L^2}$, there exists a $T_1 \in (0, T_*)$ such that

$$\sup_{0 \leq t \leq T_1} \|\nabla \rho(t)\|_{L^\infty} + \|\nabla^2 \rho(t)\|_{L^2} \leq 8M_0,$$

and

$$\sup_{0 \leq t \leq T_1} \|\nabla u(t)\|_{L^2}^2 \leq 4 \frac{\bar{\mu}}{\underline{\mu}} \|\nabla u_0\|_{L^2}^2.$$

Let

$$T^* = \sup \left\{ T \mid (\rho, u, w, P) \text{ is a strong solution to (1.1) on } [0, T] \right\},$$

and

$$T_1^* = \sup \left\{ T \mid \begin{aligned} & (\rho, u, w, P) \text{ is a strong solution to (1.1) on } [0, T], \sup_{0 \leq t \leq T} (\|\nabla \rho(t)\|_{L^\infty} \\ & + \|\nabla^2 \rho(t)\|_{L^2}) \leq 8M_0, \text{ and } \sup_{0 \leq t \leq T} \|\nabla u(t)\|_{L^2}^2 \leq (2\ln 2 + 1) \frac{\bar{\mu}}{\underline{\mu}} \|\nabla u_0\|_{L^2}^2 \end{aligned} \right\}.$$

Then $T_1^* \geq T_1 > 0$. Recalling Lemma 3.2 and Lemma 3.9, it is easy to verify that

$$T^* = T_1^*,$$

provided that $\|\nabla u_0\|_{L^2} \leq \eta_0 = \min\{\gamma_0, \epsilon_0\}$ as assumed.

We claim that $T^* = \infty$. Otherwise assume that $T^* < \infty$. By virtue of Lemma 3.2 and Lemma 3.9, for every $t \in [0, T^*)$, it holds that

$$\|\nabla \rho(t)\|_{L^\infty} + \|\nabla^2 \rho\|_{L^2} \leq 4M_0 \text{ and } \|\nabla u(t)\|_{L^2}^2 \leq (2ln2 + 1) \frac{\bar{\mu}}{\underline{\mu}} \|\nabla u_0\|_{L^2}^2,$$

which contradicts the blow-up criterion (1.8) in Theorem 1.1. Hence we complete the proof of Theorem 1.2.

Acknowledgements. Q. Jiu is partially supported by National Natural Sciences Foundation of China (No. 11931010). F. Wang is partially supported by the National Natural Science Foundation of China (No.12201028), Chinese Postdoctoral Science Foundation (No.2022M720383)

References

- [1] D. Bresch, A.V. Kazhikhov, J. Lemoine, On the two-dimensional hydrostatic Navier-Stokes equations, *SIAM J. Math. Anal.* 36(3) (2004) 796-814.
- [2] C. Cao, J. Li, E. S. Titi, Global well-posedness of strong solutions to the 3D primitive equations with horizontal eddy diffusivity. *J Differential Equations*, 2014, 257(11): 4108-4132.
- [3] C. Cao, J. Li, E. S. Titi, Local and global well-posedness of strong solutions to the 3D primitive equations with vertical eddy diffusivity. *Arch Ration Mech Anal*, 2014, 214(1): 35-76.
- [4] C. Cao, J. Li, E. S. Titi, Global well-posedness of the three-dimensional primitive equations with only horizontal viscosity and diffusion. *Comm Pure Appl Math*, 2016, 69(8): 1492-1531.
- [5] C. Cao, J. Li, E. S. Titi, Strong solutions to the 3D primitive equations with only horizontal dissipation: near H^1 initial data. *J Funct Anal*, 2017, 272(11): 4606-4641.
- [6] C. Cao, E. S. Titi, Global well-posedness of the three-dimensional viscous primitive equations of large scale ocean and atmosphere dynamics. *Ann Math*, 2007, 166: 245-267.
- [7] C. Cao, E. S. Titi, Global well-posedness of the 3D primitive equations with partial vertical turbulence mixing heat diffusion. *Comm Math Phys*, 2012, 310: 537-568.
- [8] M. Ersoy, T. Ngom, M. Sy, Compressible primitive equations : formal derivation and stability of weak solutions , *Nonlinearity*, 24(2011), 79-96.
- [9] B. V. Gatapov, A. V. Kazhikhov, Existence of a global solution of a model problem of atmospheric dynamics, *Siberian Math. J.*, 46(2005), 805-812.
- [10] Y. Giga, M. Gries, M. Hieber, A. Hussein, T. Kashiwabara, The primitive equations in the scaling-invariant space $L^\infty(L^1)$. *J. Evol. Equ.* 2021, no. 4, 4145-4169.
- [11] Y. Giga, M. Gries, M. Hieber, A. Hussein, T. Kashiwabara, Analyticity of solutions to the primitive equations. *Math. Nachr.* 2020, no. 2, 284-304.

- [12] F. Guill-Gonzalez, N. Masmoudi, and M. A. Rodriguez-Bellido, Anisotropic estimates and strong solutions of the primitive equations, *Differ. Integral Equations* 14(11), 1381-1408(2001).
- [13] M. Hieber, T. Kashiwabara, Global strong well-posedness of the three dimensional primitive equations in L^p -spaces. *Arch. Ration. Mech. Anal.* 2016, no. 3, 1077-1115.
- [14] Q. Jiu, M. Li, F. Wang, Uniqueness of the global weak solutions to 2D compressible primitive equations. *J. Math. Anal. Appl.* 461(2), 1653-1671(2018).
- [15] Q.S. Jiu, L. Ma, F. C. Wang, Local Well-Posedness of Strong Solutions to the 2D density-dependent incompressible primitive equations, 2023, preprint.
- [16] Q. Jiu, F. Wang, Local existence and uniqueness of strong solutions to the two dimensional nonhomogeneous incompressible primitive equations. *Acta Math. Sci. Ser. B (Engl. Ed.)* 2020, no. 5, 1316-1334.
- [17] G. M. Kobelkov, Existence of a solution in the large for ocean dynamics equations, *J.Math.Fluid Mech.*, 2007, 588-610.
- [18] I. Kukavica, P.Yuan, Walter, R, M. Ziane, Primitive equations with continuous initial data. *Nonlinearity* 27 (2014), no. 6, 1135-1155.
- [19] I. Kukavica, M. Ziane, On the regularity of the primitive equations of the ocean, *Nonlinearity*, 2007, 2739-2753.
- [20] I. Kukavica, M. Ziane, Uniform gradient bounds for the primitive equations of the ocean, *Differ. Integral Equ.*, 2008, 837-849.
- [21] J.L.Lions, R.Temam, S.Wang, New formulations of the primitive equations of atmosphere and applications, *Nonlinearity*, 1992, 237-288.
- [22] J.L.Lions, R.Temam, S.Wang, On the equation of the large scale ocean, *Nonlinearity*, 1992, 1007-1053.
- [23] J.L.Lions, R.Temam, S.Wang, Mathematical theory of the coupled atmosphere-ocean model, *J.Math.Pures Appl.*, 1995, 105-163.
- [24] X. Liu, E. S. Titi, Zero Mach number limit of the compressible Primitive Equations Part I : well-prepared initial data. *Arch. Ration. Mech. Anal.* 238, 705-747(2020).
- [25] X. Liu, E. S. Titi, Local well-posedness of strong solutions to the three-dimensional compressible primitive equations. *Arch. Ration. Mech. Anal.* 241 (2021), no. 2, 729-764.
- [26] X.Liu, E. S. Titi, Global existence of weak solutions to the compressible Primitive Equations of atmospheric dynamics with degenerate viscosities, *SIAMJ. Math. Anal.*, 51(2019), 1913-1964.
- [27] X.Liu, E. S. Titi, Zero Mach number limit of the compressible primitive equations: Ill-prepared initial data. *J. Differential Equations* 356 (2023).
- [28] L. Nirenberg, On elliptic partial differential equations, *Ann. Scuola Norm. Sup. Pisa* (3) 1959, 115-162 .

- [29] J. Pedlosky, *Geophysical Fluid Dynamics*. 2nd ed. New-York: Springer-Verlag, 1987.
- [30] L. F. Richardson, *Weather Prediction by Numerical Press* (Cambridge University Press, Cambridge, 1922) [Reprinted by Dover, New York(1965)].
- [31] J. Simon, Nonhomogeneous viscous incompressible fluids: existence of velocity, density, and pressure, *SIAM J. Math. Anal.* 1990, 1093-1117.
- [32] R. Temam and M. Ziane, Some mathematical problems in geophysical fluid dynamics, *Handbook Math. Fluid Dyn.* 3, 535-658 (2005).
- [33] T.Tang, H.J.Gao, On the stability of weak solution for compressible primitive equations, *Acta Appl. Math.*, 140(2015), 133-145.
- [34] G.K. Vallis, *Atmospheric and Oceanic Fluid Dynamics*, Cambridge Univ. Press, 2006.
- [35] F.C, Wang, Local existence and uniqueness of strong solutions to the two dimensional compressible primitive equations with density-dependent viscosity. *J. Math. Fluid Mech.* 23 (2021), no. 3, Paper No. 78, 21 pp.
- [36] F. C. Wang, C. S. Dou, Q. S. Jiu, Global existence of weak solutions to 3D compressible primitive equations with degenerate viscosity. *J. Math. Phys.* 61 (2020).
- [37] F.C, Wang, Q. Jiu, X. Xu, Global well-posedness of strong solutions to the 2D nonhomogeneous incompressible primitive equations with vacuum, *J. Differential Equations* 2021, 624-644.
- [38] F.C, Wang, Q. Jiu, Global well-posedness of strong solutions to the 2D compressible primitive equations with density-dependent viscosity. 2023, preprint.