

STRONG SURVIVAL AND EXTINCTION FOR BRANCHING RANDOM WALKS VIA A NEW ORDER FOR GENERATING FUNCTIONS

DANIELA BERTACCHI AND FABIO ZUCCA

ABSTRACT. We consider general discrete-time branching random walks on a countable set X . According to these processes, a particle at $x \in X$ generates a random number of children and places them at (some of) the sites of X , not necessarily independently nor with the same law at different starting vertices x . We introduce a new type of stochastic ordering of branching random walks, generalizing the germ order introduced by Hutchcroft in [13], which relies on the generating function of the process. We prove that given two branching random walks with law μ and ν respectively, with $\mu \geq \nu$, then in every set where there is extinction according to μ , there is extinction also according to ν . Moreover, in every set where there is strong local survival according to ν , there is strong local survival also according to μ , provided that the supremum of the global extinction probabilities, for the ν -process, taken over all starting points x , is strictly smaller than 1. New conditions for survival and strong survival for inhomogeneous branching random walks are provided. We also extend a result of Moyal [16] which claims that, under some conditions, the global extinction probability for a branching random walk is the only fixed point of its generating function, whose supremum over all starting coordinates, may be equal to 1.

Keywords: branching random walk, generating function, fixed point, extinction probability vectors, germ order, pgf order, strong survival.

AMS subject classification: 60J80.

1. INTRODUCTION

The branching random walk (or briefly *BRW*) on an at most countable set X is a process which describes the evolution of a population breeding and dying on X . The elements of X can be seen as the locations where individuals of the population may live.

A general BRW is defined once we fix the reproduction law $\mu = \{\mu_x\}_{x \in X}$ (see Section 2.1 for details). All particles at site x breed and place children according to μ_x , which incorporates not only information about how many the children are, but also about where they are sent to live. In this sense particles do not walk, but there is a random walk of the population as a whole.

The *branching process* can be seen as a particular case of the BRW, where X is reduced to a singleton and the only information needed is the reproduction law ρ defined on $\mathbb{N}$. If locations are seen as types, the branching process has only one type, while the BRW is also known as *multi-type branching process*. A natural way to define a BRW on X is to couple a family of branching processes, given by reproduction laws $\{\rho_x\}_{x \in X}$, and a random walk with transition matrix P on X . Each individual at x has a ρ_x -distributed number of offspring, which are independently dispersed according to the random walk. We call this kind of process BRW *with independent diffusion*. We remark that for general BRW, the dispersal of the progeny may not be independent nor based on a random walk (for instance we may place two children at a given vertex with probability p and one child at each of a couple of other vertices with probability $1 - p$).

We are interested in the long-term behaviour of the process in fixed subsets of X . In the long run, for any $A \subseteq X$, a BRW starting with one individual at $x \in X$ can go extinct in A (no individuals alive in A from a certain time on) or survive in A (infinitely many visits to A). If the probability of extinction in A is equal to 1, we say that there is *extinction* in A , and we say that there is *survival* in A otherwise. There is *global survival* when there is survival in X and we have *strong survival* in A when, conditioned on global survival, there is survival in A .

Clearly, the probability of extinction in A depends on the starting vertex x . Then, letting x vary in X , we get an extinction probability vector that we denote by $\mathbf{q}(A)$. If we allow A to vary among the subsets of X , we have the family of all extinction probability vectors of the BRW.

For the branching process, it is well known that the long-term behaviour and the extinction probability are linked with the generating function of ρ : $G(z) := \sum_n \rho(n)z^n$, $z \in [0, 1]$. Provided that the process is nontrivial, that is $\rho(1) < 1$, this generating function has at most two fixed points: the extinction probability and 1. In the case of a general BRW it is possible to define a (multi-dimensional) generating function which plays a similar role, but as soon as X is not finite, the situation gets far more complex: there might be infinitely many fixed points and infinitely many extinction probability vectors (see Section 2.3); moreover there can be fixed points that are not extinction probability vectors. It is still true, however, that the vector $\mathbf{1}$ is always a fixed point of the generating function and the *global extinction* probability vector (that is, the probability of extinction in the whole space X) is always the smallest fixed point.

The fact that the generating function of the process entails all the information on its behaviour is exploited in the main result of the present paper. In [13] the author focussed on BRWs with independent diffusion and reproduction law ρ equal for all sites and introduced a new stochastic ordering. This order is named *germ order* and is based on a comparison between the one-dimensional generating functions of the reproduction laws. The author was able to compare BRWs which are defined by the same underlying random walk P on X and differ only by the reproduction law, which is constant along X .

We define the germ order for general BRWs and extend [13, Theorem 1.3], by proving the following (a more precise statement is given by Theorem 4.3).

Theorem 1.1. *Let μ and ν be the law of two BRWs on a countable space X and let $\mu \geq_{\text{germ}} \nu$.*

- (1) *In any set where there is μ -extinction, there is ν -extinction.*
- (2) *If the supremum of the global ν -extinction probabilities, over all starting coordinates, is smaller than 1, then in any set where there is ν -strong survival, there is μ -strong survival.*

The request in the second part of the statement may appear technical at first glance, but it is worth remarking that under very mild conditions, among all fixed points, only the global extinction probability vector may satisfy this condition. Indeed we extend a result of [16], which states that, under certain conditions, the global extinction probability vector is the only fixed point which may have coordinates bounded from above by some $\delta < 1$. We are able to prove that under no conditions at all, the global extinction probability vector is the only extinction probability vector with this property. Moreover, if a mild condition is satisfied (see Theorem 3.1), it is also the only fixed point with supremum different from 1. This result allows us to simplify the original proof of [13, Theorem 1.3] and extend it to the case of general BRWs.

The paper is organized as follows. In Section 2 we introduce the process formally and we give the basic definitions. Section 2.3 is devoted to the main object of study: the generating function G of the BRW, which is already known to be useful since extinction probabilities are (some of) its fixed points. In particular, Proposition 2.4 shows that fixed points of G can be found by iterating the function itself on suitable starting vectors. As shown in Section 2.4, fixed points can be used to construct special martingales which are useful tools for proving the main result of Section 3: Theorem 3.1. This theorem aims to show that, given a generic (not necessarily irreducible) BRW, if there exists $\delta < 1$ such that the probability of extinction in A , starting from any $x \in X$, is smaller than δ , then $\mathbf{q}(A) = \mathbf{q}(X)$. The same can be said for any fixed point $\mathbf{z}$: if all its coordinates are small than δ for some $\delta < 1$, then $\mathbf{z} = \mathbf{q}(X)$, provided that the BRW satisfies a mild sufficient condition. We show that without this condition, there are examples where the property does not hold for fixed points (see Examples 3.2 and 3.3). Section 4 is devoted to the relation between survival (resp. strong survival) for two BRWs satisfying a non-trivial order (called *germ order*). We recall the usual stochastic order for BRWs and introduce the pgf and the germ orders, which are based on the behavior of the two generating functions. These two definitions extend to general BRWs the

corresponding ones which were given in [13] for BRWs which can be projected onto a branching process. The germ order is weaker than the pgf order which is weaker than the usual stochastic order (the latter is the classic order used for coupling BRWs). The main result of the section, Theorem 4.3, deals with the germ order, while Theorem 4.4 deals with the pgf order. For two BRWs with independent diffusion we find a condition equivalent to germ-order (see Proposition 4.7). The results of this Section generalize the results in [13]. As explained in details at the end of the section, by using Theorem 4.3 and Proposition 4.7, we are able to prove new and powerful conditions for survival and strong survival for inhomogeneous BRWs. All the proofs, along with technical lemmas can be found in Section 5. The final Appendix is devoted to the construction of $\mathbb{R}^X$ as a Polish space which is essential for coupling processes stochastically ordered in the classical way.

2. BASIC DEFINITIONS AND PROPERTIES

2.1. The process. Given an at most countable set X , a discrete-time BRW is a process $\{\eta_n\}_{n \in \mathbb{N}}$, where $\eta_n(x)$ is the number of particles alive at $x \in X$ at time n . The dynamics is described as follows: consider the (countable) measurable space $(S_X, 2^{S_X})$ where $S_X := \{f : X \rightarrow \mathbb{N} : \sum_y f(y) < +\infty\}$ and let $\mu = \{\mu_x\}_{x \in X}$ be a family of probability measures on $(S_X, 2^{S_X})$. A particle of generation n , at site $x \in X$, lives one unit of time; after that, a function $f \in S_X$ is chosen at random according to the law μ_x . This function describes the number of children and their positions, that is, the original particle is replaced by $f(y)$ particles at y , for all $y \in X$. The choice of f is independent for all breeding particles.

An explicit construction is the following: given a family $\{f_{i,n,x}\}_{i,n \in \mathbb{N}, x \in X}$ of independent S_X -valued random variable such that, for every $x \in X$, $\{f_{i,n,x}\}_{i,n \in \mathbb{N}}$ have the common law μ_x , then the discrete-time BRW $\{\eta_n\}_{n \in \mathbb{N}}$ is defined iteratively as follow

$$\eta_{n+1}(x) = \sum_{y \in X} \sum_{i=1}^{\eta_n(y)} f_{i,n,y}(x) = \sum_{y \in X} \sum_{j=0}^{\infty} \mathbb{1}_{\{\eta_n(y)=j\}} \sum_{i=1}^j f_{i,n,y}(x) \quad (2.1)$$

starting from an initial condition η_0 . The actual canonical construction can be carried on, by using Kolmogorov's Theorem, in such a way that the probability space and the process $\{\eta_n\}_{n \in \mathbb{N}}$ are fixed, while the probability measure depends on the starting configuration and the family μ . When the initial configuration is η , then the corresponding probability measure is denoted by $\mathbb{P}_\mu^\eta$ and the expectation by $\mathbb{E}_\mu^\eta$. In the particular case when the initial state is one particle at x , namely $\eta = \delta_x$ a.s., then we write $\mathbb{P}_\mu^x$ and $\mathbb{E}_\mu^x$. When μ is fixed, we avoid the subscript μ in the above notations. Similarly, when a result holds for every initial condition η (or when the initial condition is fixed) we avoid the superscripts η and x . We denote the BRW by (X, μ) ; if needed, the initial value will be indicated each time. Clearly, (X, μ) is a Markov chain with absorbing state $\mathbf{0}$, the configuration with no particles at all sites. We denote by $\{\mathcal{F}_n\}_{n \in \mathbb{N}}$ the filtration associated to the process, namely, $\mathcal{F}_n := \sigma(f_{i,j,x} : i, j \in \mathbb{N}, j < n, x \in X)$. Note that $\mathcal{F}_0$ is the trivial σ -algebra. By using (2.1) it is easy to see that the BRW is *adapted* to $\{\mathcal{F}_n\}_{n \in \mathbb{N}}$, that is, η_n is $\mathcal{F}_n$ -measurable for every $n \in \mathbb{N}$.

The total number of children associated to f is represented by the function $\mathcal{H} : S_X \rightarrow \mathbb{N}$ defined by $\mathcal{H}(f) := \sum_{y \in X} f(y)$; the associated law $\rho_x(\cdot) := \mu_x(\mathcal{H}^{-1}(\cdot))$ is the law of the random number of children of a particle living at x . We denote by $m_{xy} := \sum_{f \in S_X} f(y) \mu_x(f)$ the expected number of children that a particle living at x sends to y . It is easy to show that $\sum_{y \in X} m_{xy} = \bar{\rho}_x$ where $\bar{\rho}_x$ is the expected value of the law ρ_x .

In particular, if ρ_x does not depend on $x \in X$, we say that the BRW can be *projected on a branching process* (see [6] for details). If a BRW can be projected onto a BRW defined on a finite set, then it is called $\mathcal{F}$ -BRW (see [8, Section 2.3], Remark 4.6 or [2, 19] for the details on the properties of this projection map). The case of the projection on a branching process, is a particular case of $\mathcal{F}$ -BRW, where the finite set reduces to a singleton. Other examples are the so called *quasi-transitive BRWs* (see [7, Section 2.4, p. 408] for the formal definition), where the action of the

group of automorphisms of the BRW (namely, bijective maps preserving the reproduction laws) has a finite number j of orbits: the finite set onto which we project has cardinality j . When there is just one orbit, then the BRW is called *transitive* (which is thus a particular case of BRW projected on a branching process). We note that in general, an $\mathcal{F}$ -BRW does not need to be transitive nor quasi-transitive.

It is important to note that, for a generic BRW, the locations of the offsprings are not (necessarily) chosen independently, but they are assigned by the function $f \in S_X$. We denote by P the *diffusion matrix* with entries $p(x, y) = m_{xy}/\bar{\rho}_x$. When the children are dispersed independently, they are placed according to P and the process is called *BRW with independent diffusion*: in this case

$$\mu_x(f) = \rho_x \left(\sum_{y \in X} f(y) \right) \frac{(\sum_{y \in X} f(y))!}{\prod_{y \in X} f(y)!} \prod_{y \in X} p(x, y)^{f(y)}, \quad \forall f \in S_X. \quad (2.2)$$

To a generic discrete-time BRW we associate a graph (X, E_μ) where $(x, y) \in E_\mu$ if and only if $m_{xy} > 0$. We say that there is a path from x to y of length n , and we write $x \xrightarrow{n} y$, if it is possible to find a finite sequence $\{x_i\}_{i=0}^n$ (where $n \in \mathbb{N}$) such that $x_0 = x$, $x_n = y$ and $(x_i, x_{i+1}) \in E_\mu$ for all $i = 0, \dots, n-1$. Clearly $x \xrightarrow{0} x$ for all $x \in X$; if there exists $n \in \mathbb{N}$ such that $x \xrightarrow{n} y$, then we write $x \rightarrow y$. Whenever $x \rightarrow y$ and $y \rightarrow x$ we write $x \rightleftharpoons y$. If the graph (X, E_μ) is *connected*, then we say that the BRW is *irreducible*.

In order to avoid trivial situations where particles have one offspring almost surely, we assume henceforth the following.

Assumption 2.1. *For all $x \in X$ there is a vertex $y \rightleftharpoons x$ such that $\mu_y(f: \sum_{w: w \rightleftharpoons y} f(w) = 1) < 1$.*

2.2. Survival and extinction.

Definition 2.2. *We call survival in $A \subseteq X$ the event*

$$\mathcal{S}(A) := \left\{ \limsup_{n \rightarrow +\infty} \sum_{y \in A} \eta_n(y) > 0 \right\},$$

and we denote by $\mathcal{E}(A) = \mathcal{S}(A)^c$ the event that we call extinction in A . We define the extinction probability vector $\mathbf{q}(A)$ as $\mathbf{q}(x, A) := \mathbb{P}^x(\mathcal{E}(A))$ for $x \in X$.

It is important to note that, in the canonical construction, the events $\{\mathcal{E}(A), \mathcal{S}(A)\}_{A \subseteq X}$ and the corresponding random variables $\{\mathbb{1}_{\mathcal{E}}(A), \mathbb{1}_{\mathcal{S}}(A)\}_{A \subseteq X}$ are fixed and do not depend on $\boldsymbol{\mu}$ and the initial configuration η . The dependence on $\boldsymbol{\mu}$ and η is in the probability measure $\mathbb{P}_{\boldsymbol{\mu}}^x$.

Definition 2.3.

- (1) *The process survives in $A \subseteq X$, starting from $x \in X$, if*

$$\mathbf{q}(x, A) < 1;$$

otherwise the process goes extinct in A (or dies out in A).

- (2) *The process survives globally, starting from x , if it survives in X .*
(3) *There is strong survival in $A \subseteq X$, starting from $x \in X$, if $\mathbf{q}(x, A) = \mathbf{q}(x, X) < 1$*

In the rest of the paper we use the notation $\mathbf{q}(x, y)$ instead of $\mathbf{q}(x, \{y\})$ for all $x, y \in X$. When there is no survival with positive probability, we say that there is extinction and the fact that extinction occurs almost surely will be tacitly understood. It is worth noting that, in the irreducible case, for every $A \subseteq X$, the inequality $\mathbf{q}(x, A) < 1$ holds for some $x \in X$ if and only if it holds for every $x \in X$ (although it may be $\mathbf{q}(x, A) \neq \mathbf{q}(y, A)$ for some $x \neq y$). For details and results on survival and extinction see for instance [6, 19].

2.3. Infinite-dimensional generating function. To the family $\{\mu_x\}_{x \in X}$, we associate the following generating function $G_\mu : [0, 1]^X \rightarrow [0, 1]^X$,

$$G_\mu(\mathbf{z}|x) := \sum_{f \in S_X} \mu_x(f) \prod_{y \in X} \mathbf{z}(y)^{f(y)} = \mathbb{E}^x \left[\prod_{y \in X} \mathbf{z}(y)^{\eta_1(y)} \right], \quad (2.3)$$

where $G_\mu(\mathbf{z}|x)$ is the x coordinate of $G_\mu(\mathbf{z})$. The family $\{\mu_x\}_{x \in X}$ is uniquely determined by G (see for instance [7, Section 2.3] or [8, Section 2.2] and Lemma 5.1). Henceforth, when possible, we write G instead of G_μ . The function G is continuous with respect to the *pointwise convergence topology* of $[0, 1]^X$ and nondecreasing with respect to the usual partial order of $[0, 1]^X$ (see [5, Sections 2 and 3] for further details). It is easy to show that for all $\mathbf{z} \leq \mathbf{v}$, $t \mapsto G(\mathbf{z} + t(\mathbf{v} - \mathbf{z}))$ is a convex function and, in some cases, it is a strictly convex function (see [7, Lemma 5.1]); nevertheless, in general, the function G is convex (see [8, Section 3.1]). The generating function of the total number of children satisfies $\phi_x(t) := \sum_{n \in \mathbb{N}} \rho_x(n) t^n = G(t\mathbf{1}|x)$ for all $x \in X$ and $t \in [0, 1]$.

As in the case of a branching process, extinction probabilities are fixed points of G . The smallest fixed point is $\mathbf{q}(X)$: more generally, given a solution of $G(\mathbf{z}) \leq \mathbf{z}$, then $\mathbf{z} \geq \mathbf{q}(X)$. Consider now the closed sets $F_G := \{\mathbf{z} \in [0, 1]^X : G(\mathbf{z}) = \mathbf{z}\}$, $U_G := \{\mathbf{z} \in [0, 1]^X : G(\mathbf{z}) \leq \mathbf{z}\}$ and $L_G := \{\mathbf{z} \in [0, 1]^X : G(\mathbf{z}) \geq \mathbf{z}\}$; clearly $F_G = U_G \cap L_G$. Moreover, by the monotonicity property, $G(U_G) \subseteq U_G$ and $G(L_G) \subseteq L_G$. The iteration of G produces sequences converging to fixed points.

Proposition 2.4. *Fix $\mathbf{z}_0 \in [0, 1]^X$ and define, iteratively, $\mathbf{z}_{n+1} := G(\mathbf{z}_n)$ for all $n \in \mathbb{N}$. Suppose that $\mathbf{z}_n \rightarrow \mathbf{z}$ as $n \rightarrow +\infty$ for some $\mathbf{z} \in [0, 1]^X$. Then $\mathbf{z} \in F_G$. Moreover, fix $\mathbf{w} \in [0, 1]^X$.*

- (1) *If $\mathbf{w} \in U_G$ then $\mathbf{w} \geq \mathbf{z}_0$ implies $\mathbf{w} \geq \mathbf{z}$ (the converse holds for $\mathbf{z}_0 \in L_G$).*
- (2) *If $\mathbf{w} \in L_G$ then $\mathbf{w} \leq \mathbf{z}_0$ implies $\mathbf{w} \leq \mathbf{z}$ (the converse holds for $\mathbf{z}_0 \in U_G$).*

The proof is straightforward (see for instance [5]). The sequence $\{\mathbf{z}_n\}_{n \in \mathbb{N}}$ defined in the previous proposition converges if $\mathbf{z}_0 \in L_G$ (resp. $\mathbf{z}_0 \in U_G$): in that case $\mathbf{z}_n \uparrow \mathbf{z}$ (resp. $\mathbf{z}_n \downarrow \mathbf{z}$) for some $\mathbf{z} \in F_G$.

We note that $\mathbf{q}(X)$ is not only the smallest fixed point of G , but also of any of its iterates $G^{(n)}$, where $G^{(1)} := G$ and $G^{(n+1)} := G \circ G^{(n)}$ for every $n \geq 1$. Indeed it is known (see for instance [19]) that $\mathbf{q}(X) = \lim_{i \rightarrow +\infty} G^{(i)}(\mathbf{0}) = \lim_{i \rightarrow +\infty} G^{(i \cdot n)}(\mathbf{0})$ for every $n \geq 1$. By Proposition 2.4, since $\mathbf{0} \in L_G$ is the smallest point of $[0, 1]^X$, the above sequence converges to the smallest fixed point of $G^{(n)}$ for all $n \geq 1$.

Let us briefly address the question of the cardinality of the set of fixed points F_G and its subset $\text{ext}(G) := \{\mathbf{z} \in [0, 1]^X : \mathbf{z} = \mathbf{q}(A), A \subseteq X\}$, that is, the set of extinction probability vectors; the question is relevant in the case of irreducible processes, otherwise it is very easy to find examples where these sets are finite or infinite. It is clear that the cardinality of both sets is at most $\mathfrak{c} := 2^{\aleph_0}$ (where $\aleph_0$ is the cardinality of $\mathbb{N}$). Let us denote the cardinality of a set by $\#$. An example can be found in [8] where $\#F_G = \mathfrak{c}$ while $\#\text{ext}(G) = 2$; whence there are fixed points which are not extinction probability vectors. In [7, Example 4.2] there is an irreducible BRW where $\#\text{ext}(G) \geq 3$, in [10] there is an example where $\#\text{ext}(G) \geq 4$ and in [9] there is an example where $\#\text{ext}(G) = \mathfrak{c}$. The question on the cardinality of $\text{ext}(G)$ was completely solved in [1] where it has been shown that, for every choice of $N \in \mathbb{N} \cup \{\aleph_0, \mathfrak{c}\}$ there exists an irreducible BRW where the cardinality of $\text{ext}(G)$ is N .

We note that if the BRW has independent diffusion (see equation (2.2)), then

$$G(\mathbf{z}|x) = \phi_x(P\mathbf{z}(x)), \quad \forall x \in X, \mathbf{z} \in [0, 1]^X, \quad (2.4)$$

where $\phi_x(t) := \sum_{i \in \mathbb{N}} \rho_x(i) t^i$ is the generating function of the total number of children of a particle at x and $P\mathbf{z}(x) = \sum_{y \in X} p(x, y) \mathbf{z}(y)$. Indeed, by using the definition of G and equation (2.2)

$$\begin{aligned} G(\mathbf{z}|x) &= \sum_{f \in S_X} \rho_x \left(\sum_{y \in X} f(y) \right) \frac{(\sum_{y \in X} f(y))!}{\prod_{y \in X} f(y)!} \prod_{y \in X} p(x, y)^{f(y)} \prod_{y \in X} \mathbf{z}(y)^{f(y)} \\ &= \sum_{n \in \mathbb{N}} \rho_x(n) \sum_{\substack{f \in S_X \\ \sum_{y \in X} f(y) = n}} \frac{n!}{\prod_{y \in X} f(y)!} \prod_{y \in X} (p(x, y) \mathbf{z}(y))^{f(y)} \\ &= \sum_{n \in \mathbb{N}} \rho_x(n) \left(\sum_{y \in X} p(x, y) \mathbf{z}(y) \right)^n = \phi_x(P\mathbf{z}(x)). \end{aligned}$$

Clearly $\phi_x(t) = \phi_x(t\mathbf{1}(x)) = G(t\mathbf{1}|x)$.

2.4. A useful martingale. Given $\mathbf{z} \in [0, 1]^X$ and $\mathbf{w} \in [0, +\infty)^X$, we define $\mathbf{z}^{\mathbf{w}} \in [0, 1]$ as

$$\mathbf{z}^{\mathbf{w}} := \prod_{x \in X} \mathbf{z}(x)^{\mathbf{w}(x)}.$$

Note that this infinite product always converges, being the limit of a nonincreasing sequence (for any choice of ordering of the elements in X).

The first result gives an explicit expression of the conditional expectation of the above product in terms of the generating function of the process.

Lemma 2.5. *For every $\mathbf{z} \in [0, 1]^X$, $m \geq 0$, $k \geq 1$ and for every initial condition η , we have*

$$\mathbb{E}^\eta [\mathbf{z}^{\eta_{m+k}} | \mathcal{F}_m] = (G^{(k)}(\mathbf{z}))^{\eta_m}, \quad \mathbb{P}^\eta\text{-a.s.}$$

The previous lemma and Doob's Martingale Theorem imply the following.

Proposition 2.6. *For every give initial state η , if $\mathbf{z} \in L_G$ (resp. $\mathbf{z} \in U_G$), then $\mathbb{E}^\eta[\mathbf{z}^{\eta_{n+1}} | \mathcal{F}_n] \geq \mathbf{z}^{\eta_n}$ (resp. $\mathbb{E}^\eta[\mathbf{z}^{\eta_{n+1}} | \mathcal{F}_n] \leq \mathbf{z}^{\eta_n}$) for all $n \geq 0$.*

In particular if $\mathbf{z} \in F_G$, then $\{\mathbf{z}^{\eta_n}\}_{n \in \mathbb{N}}$ is a martingale and there exists a $[0, 1]$ -valued, $\mathcal{F}_\infty$ -measurable random variable $W_{\mathbf{z}}$ such that,

$$\mathbf{z}^{\eta_n} \rightarrow W_{\mathbf{z}}, \quad \mathbb{P}^\eta\text{-a.s. and in } L^p(\mathbb{P}^\eta) \quad \forall p \geq 1.$$

Moreover $\mathbb{E}^\eta[W_{\mathbf{z}} | \mathcal{F}_n] = \mathbf{z}^{\eta_n}$ $\mathbb{P}^\eta$ -a.s.

Note that, for every $\mathbf{z} \in [0, 1]^X$, we have that $\mathbf{z}^{\eta_n} \rightarrow 1$ on $\mathcal{E}(X)$; whence if $\mathbf{z} \in F_G$, then $W_{\mathbf{z}} = 1$, $\mathbb{P}^\eta$ -a.s. on $\mathcal{E}(X)$. Moreover, by the martingale property, $\mathbb{E}^\eta[W_{\mathbf{z}}] = \mathbf{z}^\eta$ and $\mathbb{E}^x[W_{\mathbf{z}}] = \mathbf{z}(x)$. This martingale plays a crucial role in the proof of Theorem 3.1.

Corollary 2.7. *If $A \subseteq X$, then $\mathbf{q}(A)^{\eta_n} \rightarrow \mathbb{1}_{\mathcal{E}(A)}$ $\mathbb{P}^\eta$ -a.s. and in $L^p(\mathbb{P}^\eta)$ for all $p \geq 1$.*

The following corollary gives monotonicity of the limit $W_{\mathbf{z}}$ with respect to $\mathbf{z}$.

Corollary 2.8. *If $\mathbf{z}$ and $\mathbf{v}$ are two fixed points, then the following are equivalent*

- (1) $\mathbf{z} \geq \mathbf{v}$
- (2) $\mathbb{P}^\eta(W_{\mathbf{z}} \geq W_{\mathbf{v}}) = 1$ for every initial condition $\eta \in S_X$
- (3) $\mathbb{P}^x(W_{\mathbf{z}} \geq W_{\mathbf{v}}) = 1$ for every $x \in X$.

3. UPPER BOUNDS RESULTS FOR EXTINCTION PROBABILITIES AND FIXED POINTS

By using the martingales of Section 2.4, we can remove the assumption of irreducibility from [16, Lemma 3.3], a result which says that, under a mild condition, if the coordinates of $\mathbf{v} \in F_G$ are bounded away from 1, then $\mathbf{v} = \mathbf{q}(X)$. Note that Theorem 3.1 (1) says that no assumptions are needed to prove that the same property holds for all $\mathbf{v}$ extinction probability vectors. Theorem 3.1 plays a key role in Section 4.

Theorem 3.1. *Let (X, μ) be a generic BRW (not necessarily irreducible).*

- (1) *If $A \subseteq X$ such that $\mathbf{q}(A) \neq \mathbf{q}(X)$, then $\sup_{x \in X} \mathbf{q}(x, A) = 1$.*
- (2) *If $\inf_{x \in X} \mathbf{q}(x, X) > 0$, then for all $\mathbf{z} \in F_G$, $\mathbf{z} \neq \mathbf{q}(X)$ we have that $\sup_{x \in X} \mathbf{z}(x) = 1$.*

The assumption $\inf_{x \in X} \mathbf{q}(x, X) > 0$, which is needed in the second part of the previous proposition, cannot be removed without replacing it by other assumptions. Indeed, without this assumption, there are examples of BRWs with an uncountable number of fixed points $\mathbf{z}$ (clearly different from $\mathbf{q}(X)$) such that $\sup_{x \in X} \mathbf{z}(x) < 1$. Example 3.2 shows a reducible case, while an irreducible one can be found in Example 3.3.

Example 3.2. *Let $X = \mathbb{N}$ and $\{p_n\}_{n \in \mathbb{N}}$ such that $p_n \in (0, 1)$ for all $n \in \mathbb{N}$ and $\sum_{i=0}^n (1-p_i) < +\infty$; this implies that $\prod_{i=0}^{\infty} p_i \in (0, 1)$ and $\prod_{i=n}^{\infty} p_i \uparrow 1$ as $n \rightarrow +\infty$. Consider a BRW where a particle at n has 1 child at $n+1$ with probability p_n and no children with probability $1-p_n$. Clearly $\eta_n = \delta_n$ or $\eta_n = \mathbf{0}$.*

A straightforward computation shows that $G(\mathbf{z}|n) = 1 - p_n + p_n \mathbf{z}(n+1)$. Moreover it is easy to show that $\mathbf{q}(n, X) = 1 - \prod_{i=n}^{\infty} p_i$ whence $\inf_{n \in \mathbb{N}} \mathbf{q}(n, X) = 0$. More generally, $\mathbf{q}(A) = \mathbf{q}(X)$ if A is infinite and $\mathbf{q}(A) = \mathbf{1}$ if A is finite.

Given $z_0 \in (1 - \prod_{i=0}^{\infty} p_i, 1) = (\mathbf{q}(0, X), 1)$, then the recursive relation $z_{n+1} := 1 - (1 - z_n)/p_n$ uniquely defines a strictly decreasing and strictly positive sequence such that $z_n > 1 - \prod_{i=n}^{\infty} p_i$. Indeed, by rewriting the recursive equality, $1 - z_{n+1} = (1 - z_n)/p_n > 1 - z_n$ for all $n \in \mathbb{N}$. The inequality $z_n > 1 - \prod_{i=n}^{\infty} p_i$ can be proven easily by induction on n . Note that $\mathbf{z}(n) := z_n$ for all $n \in \mathbb{N}$ defines a fixed point of G . Moreover $\sup_{n \in \mathbb{N}} \mathbf{z}(n) = \mathbf{z}(0) = z_0 < 1$.

We observe, that every fixed point $\mathbf{w}$ can be constructed by iteration $\mathbf{w}(n+1) := 1 - (1 - \mathbf{w}(n))/p_n$ for all $n \in \mathbb{N}$ starting from $\mathbf{w}(0) \in [\mathbf{q}(0, X), 1]$. Indeed the 0-th coordinate of a fixed point belongs to the interval $[\mathbf{q}(0, X), 1]$ and the iteration equality is equivalent to $G(\mathbf{w}|n) = \mathbf{w}(n)$. Thus, in this case, for every fixed point $\mathbf{w}$ (different from $\mathbf{1}$) we have $\sup_{n \in \mathbb{N}} \mathbf{w}(n) < 1$.

Example 3.3. *Let $X = \mathbb{N}$ and $\{p_n\}_{n \in \mathbb{N}}$ as in Example 3.2. Moreover let $\{r_n\}$ be a sequence such that $1 - p_n - r_n > 0$. Consider a BRW where a particle at $n \geq 1$ has 1 child at $n+1$ with probability p_n , 1 child at $n-1$ with probability r_n and no children with probability $1 - p_n - r_n$. Suppose that $r_0 = 0$, whence a particle at 0 has 1 child at 1 with probability p_0 and no children with probability $1 - p_0$. A straightforward computation shows that*

$$G(\mathbf{z}|n) = \begin{cases} 1 - p_n - r_n + p_n \mathbf{z}(n+1) + r_n \mathbf{z}(n-1) & n \geq 1 \\ 1 - p_0 + p_0 \mathbf{z}(1) & n = 0. \end{cases}$$

Clearly the generating function is smaller than the generating function of Example 3.2, since $G(\mathbf{z}|n) = 1 - p_n + p_n \mathbf{z}(n+1) - r_n(1 - \mathbf{z}(n-1)) \leq 1 - p_n + p_n \mathbf{z}(n+1)$; whence $\mathbf{q}(n, X) \leq 1 - \prod_{i=n}^{\infty} p_i$; again, $\inf_{n \in \mathbb{N}} \mathbf{q}(n, X) = 0$.

In order to prove that there are fixed points, different from $\mathbf{q}(X)$, with all coordinates smaller than δ (for some $\delta < 1$), it suffices to find at least two distinct fixed points with this property.

Given $z_0 \in (1 - \prod_{i=0}^{\infty} p_i, 1) \subset (\mathbf{q}(0, X), 1)$, the recursive relation

$$z_{n+1} := \begin{cases} 1 - (1 - z_0)/p_0 & n = 0 \\ 1 + (1 - z_{n-1})r_n/p_n - (1 - z_n)/p_n & n \geq 1 \end{cases}$$

uniquely defines a strictly decreasing and strictly positive sequence such that $z_n > 1 - \prod_{i=n}^{\infty} p_i$. more precisely, we prove that $z_0 \geq z_{n-1} > z_n > 1 - \prod_{i=n}^{\infty} p_i$ by induction on n . The inequality $1 - \prod_{i=1}^{\infty} p_i < z_1 < z_0$ is trivial. Suppose that $1 - \prod_{i=n}^{\infty} p_i < z_n < z_{n-1} \leq z_0$, that is $\prod_{i=n}^{\infty} p_i > 1 - z_n > 1 - z_{n-1} \geq 1 - z_0$. Note that, $1 - z_{n+1} = ((1 - z_n) - (1 - z_{n-1})r_n)/p_n > ((1 - z_n) - (1 - z_n)r_n)/p_n > (1 - z_n)(1 - r_n)/p_n > 1 - z_n \geq 1 - z_0$ since, by hypothesis, $1 - p_n - r_n > 0$, that is, $(1 - r_n)/p_n > 1$. On the other hand, since $1 - z_{n-1} > 1 - z_0 > 0$, we have $1 - z_{n+1} = ((1 - z_n) - (1 - z_{n-1})r_n)/p_n < (1 - z_n)/p_n < p_n^{-1} \prod_{i=n}^{\infty} p_i = \prod_{i=n+1}^{\infty} p_i$. Then $\mathbf{z}(n) := z_n$ for all $n \in \mathbb{N}$ defines a fixed point of G with $\sup_{n \in \mathbb{N}} \mathbf{z}(n) = \mathbf{z}(0) = z_0 < 1$.

Moreover, as in Example 3.2, all fixed points $\mathbf{w}$ (different from $\mathbf{1}$) satisfy $\sup_{n \in \mathbb{N}} \mathbf{w}(n) < 1$.

One may wonder when $\inf_{x \in X} \mathbf{q}(x, X) > 0$; the following remark gives a sufficient condition.

Remark 3.4. If $\inf_{x \in X} \mu_x(\mathbf{0}) > 0$, then $\inf_{x \in X} \mathbf{z}(x) > 0$ for every fixed point $\mathbf{z}$ (including $\mathbf{q}(A)$ for every $A \subseteq X$). Indeed $\mathbf{z}(x) = G(\mathbf{z}|x) \geq \mu_x(\mathbf{0})$.

Note that the existence of a nonempty subset A satisfying $\inf_{x \in X} \mathbf{q}(x, A) > 0$ implies the existence of $y \in X$ such that $\inf_{x \in X} \mathbf{q}(x, y) > 0$.

It is worth noting that the existence of a positive lower bound for an extinction probability vector is a sufficient condition for the asymptotic explosion of the population. A precise statement is given by the following lemma.

Lemma 3.5. Let $A \subseteq X$. If $\inf_{x \in X} \mathbf{q}(x, A) > 0$, then $\mathbb{P}^\eta(\{\sum_{x \in X} \eta_n(x) \rightarrow +\infty\} \cap \mathcal{S}(A)) = \mathbb{P}^\eta(\mathcal{S}(A))$.

4. GERM ORDER: EXTINCTION AND STRONG SURVIVAL

Here we discuss extinction and strong survival for BRWs under different types of stochastic dominations. We are generalizing the results in [13] by considering generic BRWs instead of BRWs projected on a branching process (see Section 2.1 for the definition).

Define $L(A) := \sum_{x \in A, n \in \mathbb{N}} \eta_n(x)$ the total number of visits in A ; clearly, $\mathbf{q}(x, A) = \mathbb{P}^x(L(A) < +\infty)$ for all $x \in X$. Moreover let $L_n(A) := \sum_{x \in A, i \leq n} \eta_i(x)$ be the number of visits in A before time n ; clearly $L_n(A) \uparrow L(A)$ as $n \rightarrow +\infty$.

Definition 4.1. Let $\mu := \{\mu_x\}_{x \in X}$ and $\nu := \{\nu_x\}_{x \in X}$ be two families of measures on $\mathbb{R}^X$ with support on S_X . Let G_μ and G_ν be the generating functions.

- (1) $\mu \succeq \nu$ if and only if $\mu_x \succeq \nu_x$ for all $x \in X$, that is, if and only if for every $\mathbb{R}^X$ -valued nondecreasing measurable function F we have $\int F d\mu_x \geq \int F d\nu_x$ for all $x \in X$ such that the integrals are well defined.
- (2) $\mu \geq_{\text{pgf}} \nu$ if and only if $G_\mu(\mathbf{z}) \leq G_\nu(\mathbf{z})$ for all $\mathbf{z} \in [0, 1]^X$.
- (3) $\mu \geq_{\text{germ}} \nu$ if and only if there exists $\delta \in [0, 1]$ $G_\mu(\mathbf{z}) \leq G_\nu(\mathbf{z})$ for all $\mathbf{z} \in [\delta, 1]^X$.

We observe that $\mu \succeq \nu \Rightarrow \mu \geq_{\text{pgf}} \nu \Rightarrow \mu \geq_{\text{germ}} \nu$ but the reverse implications do not hold. We recall that for real-valued measures (that is, when X is a singleton), $\mu \succeq \nu$ is equivalent to the existence of two random variables η, ζ with laws μ and ν respectively, such that $\eta \geq \zeta$ a.s. (this construction is usually referred as an ordered coupling). This result can be extended to measures on partially ordered, compact metric spaces ([15, Theorem 2.4]) and to measures on partially ordered Polish spaces (see for instance [14, Theorem 1]).

It is not difficult to show that $\mathbb{R}^X$, with a suitable finite metric, is a partially ordered Polish space, as we show in the Appendix.

In this paper we deal with the weaker order $\geq_{\text{germ}}$. The following result shows that $\geq_{\text{germ}}$ is a partial order.

Proposition 4.2. The binary relation $\geq_{\text{germ}}$ is a partial order.

We note that if X is a singleton, by [17, Theorem 10.18], 1 is an isolated solution of $G_\mu(z) = G_\nu(z)$, whence in this case $\geq_{\text{germ}}$ defines a total order; as a consequence, $\geq_{\text{germ}}$ is a total order on the class of BRWs projected on a branching process (which is the class studied in [13]). As soon as $\#X \geq 2$ and the total offspring distributions $\{\rho_x\}_{x \in X}$ are not constant with respect to x , then $\geq_{\text{germ}}$ is not a total order (see the discussion after Proposition 4.7). Note that the germ order defined here is a generalization of the germ order defined in [13] (see Proposition 4.7).

The main result of this section is the following; this result generalizes [13, Theorem 1.3]. Although our proof uses similar arguments, we stress that Theorem 3.1 allows us to simplify part of the proof of Theorem 4.3 compared to [13, Theorem 1.3] obtaining a more general result at the same time.

Theorem 4.3. Let $\mu \geq_{\text{germ}} \nu$ and $A \subseteq X$.

- (1) If $x \in X$, then $\mathbf{q}^\mu(x, A) = 1$ implies $\mathbf{q}^\nu(x, A) = 1$.
- (2) If $\sup_{x \in X} \mathbf{q}^\nu(x, X) < 1$, then $\mathbf{q}^\nu(x, A) = \mathbf{q}^\nu(x, X)$ for all $x \in X$ implies $\mathbf{q}^\mu(x, A) = \mathbf{q}^\mu(x, X)$ for all $x \in X$.

Roughly speaking, extinction in A for (X, μ) implies extinction in A for (X, ν) . Moreover strong survival in A for (X, ν) implies strong survival in A for (X, μ) (the same statement for survival follows from Theorem 4.3 (1)).

Clearly, the germ order is not the only condition which allows to deduce strong survival for (X, μ) given the same behavior for (X, ν) . For instance if μ_x and ν_x agree outside a set A , then strong survival in A for (X, μ) is equivalent to strong survival for (X, ν) (see [8, Theorem 4.2] or [9, Theorem 2.4]).

As a warm-up in Section 5 we start by proving the same result under the stronger assumption $\mu \geq_{\text{pgf}} \nu$. Under this assumption, one can easily prove that $\mathbf{q}^\mu(X) \leq \mathbf{q}^\nu(X)$; indeed $G_\mu(\mathbf{q}^\nu(X)) \leq G_\nu(\mathbf{q}^\nu(X)) = \mathbf{q}^\nu(X)$. The following result generalizes [13, Corollary 2.2]. As in the previous case, Theorem 3.1 simplifies part of the proof of Theorem 4.4 compared to [13, Corollary 2.2].

Theorem 4.4. Let $\mu \geq_{\text{pgf}} \nu$ and $A \subseteq X$.

- (1) If $x \in X$, then $\mathbf{q}^\mu(x, A) = 1$ implies $\mathbf{q}^\nu(x, A) = 1$.
- (2) If $\sup_{x \in X} \mathbf{q}^\nu(x, X) < 1$, then $\mathbf{q}^\nu(x, A) = \mathbf{q}^\nu(x, X)$ for all $x \in X$ implies $\mathbf{q}^\mu(x, A) = \mathbf{q}^\mu(x, X)$ for all $x \in X$.

Remark 4.5. One may wonder when condition $\sup_{x \in X} \mathbf{q}(x, X) < 1$ is satisfied. We note that it holds if and only if there exist $\mathbf{v} \in [0, 1]$ and $\delta \in [0, 1]$ such that $G^{(n)}(\mathbf{v}) \leq \mathbf{v} \leq \delta \mathbf{1}$ for some $n \geq 1$ (apply Proposition 2.4). In particular if

$$G^{(n)}(\delta \mathbf{1}) \leq \delta \mathbf{1} \text{ for some } n \geq 1 \text{ and } \delta \in [0, 1], \quad (4.5)$$

then $\sup_{x \in X} \mathbf{q}(x, X) \leq \delta$. An easy computation shows that $G(\delta \mathbf{1}|x) = \sum_{n \in \mathbb{N}} \rho_x(n) \delta^n$ where ρ_x is the law of the number of children of a particle at x (see the definition in Section 2.1). Whence if the family of laws $\{\rho_x : x \in X\}$ is finite and they are all supercritical, then equation (4.5) holds. Indeed, in this case, for each x there exist $\delta_x \in [0, 1]$ such that $\sum_{n \in \mathbb{N}} \rho_x(n) \delta_x^n \leq \delta_x$ (choose $\delta_x = \delta_y$ if $\rho_x = \rho_y$) thus $G(\delta \mathbf{1}) \leq \delta \mathbf{1}$ where $\delta = \max_{x \in X} \delta_x$. However, condition (4.5) may be satisfied even when ρ_x is subcritical for some $x \in X$.

Remark 4.6. Another setting where it is easy to verify that $\sup_{x \in X} \mathbf{q}(x, X) < 1$ is the case of $\mathcal{F}$ -BRWs, which are BRWs with finitely many "neighborhood types" (a generalization of quasi-transitive BRWs, see for instance [8, Section 2.3] for the precise definition and properties).

For these BRWs, $\mathbf{q}(\cdot, X)$ assumes only a finite number of values. Indeed, (X, μ) is an $\mathcal{F}$ -BRW if it can be projected onto a BRW (Y, ν) where Y is finite; more precisely, there exist a surjective map $g : X \rightarrow Y$ such that $G_\mu(\mathbf{z} \circ g) = G_\nu(\mathbf{z}) \circ g$ for all $\mathbf{z} \in [0, 1]^Y$ (see [6, Section 3.1] for explicit computations). In [8, Section 2.3] it has been shown that $\mathbf{q}^\mu(X) = \mathbf{q}^\nu(X) \circ g$ whence, if (Y, ν) is supercritical and irreducible, then $\sup_{x \in X} \mathbf{q}^\mu(x, X) = \sup_{x \in X} \mathbf{q}^\nu(g(x), Y) = \max_{y \in Y} \mathbf{q}^\nu(y, Y) < 1$. A characterization of $\mathcal{F}$ -BRWs with independent diffusion is given in [3, Proposition 4.8].

The following result characterizes germ domination for BRWs with independent diffusion in terms of germ order of its one-dimensional laws of the total number of offspring, which also proves to be necessary in the case of general BRWs. Recall that the (one-dimensional) generating function ϕ_x of the total number of offspring can be retrieved from the generating function G of the process. Indeed $\phi_x(t) = G_\mu(t \mathbf{1}|x)$.

Proposition 4.7. Suppose that (X, μ) and (X, ν) are BRWs and denote by ϕ_x^μ and ϕ_x^ν the generating functions of the total number of offspring of the particles at x , according to the laws μ and ν respectively, for all $x \in X$. Consider the following for any fixed $\delta < 1$:

- (1) $G_\mu(\mathbf{z}) \leq G_\nu(\mathbf{z})$ for all $\mathbf{z} \in [\delta, 1]^X$;

(2) $\phi_x^\mu(t) \leq \phi_x^\nu(t)$ for all $t \in [\delta, 1]$ and all $x \in X$.

Then (1) $\Rightarrow$ (2). Moreover if (X, μ) and (X, ν) are BRWs with independent diffusion with the same matrix P (see equations (2.2) and (2.4)), then (2) $\Rightarrow$ (1).

We note that, in general, germ domination of all breeding laws does not imply germ ordering, even when X is finite. Take for instance $X = \{x, y\}$ and

$$G_\mu(x, y) := \begin{pmatrix} \frac{5}{6}xy + \frac{1}{6} \\ \frac{1+2x}{3} \end{pmatrix} \quad G_\nu(x, y) := \begin{pmatrix} \frac{4}{5}\left(\frac{5x+y}{6}\right)^2 + \frac{1}{5} \\ \frac{1+(5x+y)/6}{3} \end{pmatrix}.$$

Roughly speaking, according to μ every particle in x has 2 children (1 in x and 1 in y) with probability $5/6$ and no children with probability $1/6$, while every particle in y has 1 child in x with probability $2/3$ and no children with probability $1/3$. According to ν , on the other hand, each particle in x has 2 children with probability $4/5$ and no children with probability $1/5$ while each particle in y has 1 child with probability $2/3$ and no children with probability $1/3$; in this case each newborn is placed independently at random in x with probability $5/6$ or in y with probability $1/6$. Clearly (2) holds for $\delta = 0$ indeed, for all $t \in [0, 1)$ we have

$$G_\mu(t, t) = \begin{pmatrix} \frac{5}{6}t^2 + \frac{1}{6} \\ \frac{1+2t}{3} \end{pmatrix} < \begin{pmatrix} \frac{4}{5}t^2 + \frac{1}{5} \\ \frac{1+2t}{3} \end{pmatrix} = G_\nu(t, t)$$

nevertheless

$$G_\mu(x, 1) = \frac{5t}{6} + \frac{1}{6} > \frac{(5t+1)^2}{45} + \frac{1}{5} = G_\nu(t, 1|x)$$

for all $t \in (1/10, 1)$; thus G_μ and G_ν are incomparable. Moreover, if there exist $x, y \in X$ such that $\phi_x^\mu >_{\text{germ}} \phi_x^\nu$ and $\phi_y^\nu >_{\text{germ}} \phi_y^\mu$, then, according to Proposition 4.7, G_μ and G_ν are incomparable.

Conditions for survival in A or in X for general BRWs are usually difficult to find (see for instance [19, Theorem 4.1] and [7, Theorems 3.1 and 3.2]). Theorem 4.3 (1) and Proposition 4.7 together provide a powerful tool to prove survival for BRWs with independent diffusion. Indeed suppose that (X, ν) is a BRW with independent diffusion and survives in A . Then, any other BRW (X, μ) with independent diffusion, with the same matrix P , such that condition (2) of Proposition 4.7 holds, survives in A , no matter how inhomogeneous the offspring distributions of (X, μ) are. This applies for instance to the case of global survival ($A = X$). If (X, ν) is an $\mathcal{F}$ -BRW, then it survives globally if and only if the Perron-Frobenius eigenvalue of a finite matrix is strictly larger than 1 (see [19, Theorem 4.3], [7, Theorem 3.1] and [7, Section 2.4]). An $\mathcal{F}$ -BRW with independent diffusion is completely described by [3, Proposition 4.8]. Thus, we may be able to identify when (X, ν) survives globally and, by Theorem 4.3 (1), claim that (X, μ) survives globally as well, even if (X, μ) can be fairly inhomogeneous.

Another application of Proposition 4.7 is the following: suppose that (X, ν) is an irreducible and quasi-transitive BRW with independent diffusion (see for instance [7, Section 2.4]). Consider another BRW with independent diffusion (X, μ) such that condition (2) of Proposition 4.7 holds. If there exists $x \in X$ such that $\mathbf{q}^\nu(x, x) < 1$, then for every nonempty set $A \subseteq X$ we have $\mathbf{q}^\mu(w, X) = \mathbf{q}^\mu(w, A) < 1$ for all $w \in X$. Indeed, if $y \in A \subseteq X$, according to [7, Corollary 3.2], $\mathbf{q}^\nu(x, x) < 1$ implies $\mathbf{q}^\nu(w, X) \leq \mathbf{q}^\nu(x, A) \leq \mathbf{q}^\nu(w, y) = \mathbf{q}^\nu(w, X) < 1$ for all $w \in X$. Moreover, since a quasi-transitive BRW is an $\mathcal{F}$ -BRW, by Remark 4.6 we have $\sup_{w \in X} \mathbf{q}^\nu(w, X) < 1$. Proposition 4.7 and Theorem 4.3 yields the claim.

The following example shows that if we have two BRWs with independent diffusion and the offspring distribution has a certain expression, then the pgf ordering is equivalent to the coordinate-wise ordering of the first moment matrices.

Example 4.8. If $\boldsymbol{\mu}$ satisfies equation (2.2), then $G(\mathbf{z}|x) = \sum_{n \in \mathbb{N}} \rho_x(n) (P\mathbf{z}(x))^n$ (see Section 2.3). If, in particular, $\rho_x(n) = \frac{1}{1+\bar{\rho}_x} (\frac{\bar{\rho}_x}{1+\bar{\rho}_x})^n$ (as in the discrete-time counterpart of a continuous-time BRW), then the previous expression becomes $G(\mathbf{z}|x) = (1 + \bar{\rho}_x P(\mathbf{1} - \mathbf{z})(x))^{-1}$ or, in a more compact way,

$$G(\mathbf{z}) = \frac{1}{1 + M(\mathbf{1} - \mathbf{z})} \quad (4.6)$$

where M is the first-moment matrix and $M\mathbf{v}(x) = \bar{\rho}_x P\mathbf{v}(x)$. Suppose that $\boldsymbol{\mu}$ and $\boldsymbol{\nu}$ satisfy equation (2.2); let $M_{\boldsymbol{\mu}}$ and $M_{\boldsymbol{\nu}}$ be the first moment matrices of $\boldsymbol{\mu}$ and $\boldsymbol{\nu}$ respectively. By using equation (4.6), $M_{\boldsymbol{\mu}} \geq M_{\boldsymbol{\nu}}$ (with the usual natural partial order) if and only if $\boldsymbol{\mu} \geq_{pgf} \boldsymbol{\nu}$. Therefore, Theorem 4.4(1) applies.

In order to apply Theorem 4.4(2) we need $\sup_{x \in X} \mathbf{q}^{\boldsymbol{\nu}}(x, X) < 1$. According to Remark 4.5, a sufficient condition for $\sup_{x \in X} \mathbf{q}^{\boldsymbol{\nu}}(x, X) < 1$ is the existence of $\delta < 1$ such that $G_{\boldsymbol{\nu}}(\delta \mathbb{1}|x) \leq \delta$ for all $x \in X$: if G is as in equation (4.6) this condition is equivalent to $\inf_{x \in X} \bar{\rho}_x < +\infty$.

We close this section with an application of Theorem 4.3 to survival in a sequence of subsets.

Corollary 4.9. Let $\boldsymbol{\mu} \geq_{germ} \boldsymbol{\nu}$ and consider a sequence $\{A_n\}_{n \in \mathbb{N}}$ of subsets of X .

- (1) If $x \in X$ and $\mathbb{P}_{\boldsymbol{\nu}}^x(\limsup_{n \rightarrow +\infty} \{\sum_{y \in A_n} \eta_n(y) > 0\}) > 0$, then $\mathbb{P}_{\boldsymbol{\mu}}^x(\limsup_{n \rightarrow +\infty} \{\sum_{y \in A_n} \eta_n(y) > 0\}) > 0$.
- (2) If $\sup_{x \in X} \mathbf{q}^{\boldsymbol{\nu}}(x, X) < 1$ and $\mathbb{P}_{\boldsymbol{\nu}}^x(\liminf_{n \rightarrow +\infty} \{\sum_{y \in A_n} \eta_n(y) = 0\}) = \mathbf{q}^{\boldsymbol{\nu}}(x, X)$ for all $x \in X$, then $\mathbb{P}_{\boldsymbol{\mu}}^x(\liminf_{n \rightarrow +\infty} \{\sum_{y \in A_n} \eta_n(y) = 0\}) = \mathbf{q}^{\boldsymbol{\mu}}(x, X)$ for all $x \in X$.
- (3) If $\sup_{x \in X} \mathbf{q}^{\boldsymbol{\nu}}(x, X) < 1$ and $\mathbb{P}_{\boldsymbol{\nu}}^x(\limsup_{n \rightarrow +\infty} \{\sum_{y \in A_n} \eta_n(y) > 0\} | \mathcal{S}(X)) = 1$ for all $x \in X$, then $\mathbb{P}_{\boldsymbol{\mu}}^x(\limsup_{n \rightarrow +\infty} \{\sum_{y \in A_n} \eta_n(y) > 0\} | \mathcal{S}(X)) = 1 \ \forall x \in X$.

Example 4.10. As an application of Corollary 4.9 consider a metric d on X ; for instance, d could be the natural metric induced by a connected graph structure on X . Fix $x_0 \in X$ and define the maximal displacement as $M_n := \mathbb{1}_{\mathcal{S}(X)} \cdot \max\{d(x_0, y) : y \in X, \eta_n(y) > 0\}$. Then, given $\alpha > 0$ and $f : \mathbb{N} \mapsto (0, +\infty)$,

$$\limsup_{n \rightarrow +\infty} M_n / f(n) \leq \alpha, \mathbb{P}_{\boldsymbol{\mu}}^{x_0} \text{-a.s.} \implies \limsup_{n \rightarrow +\infty} M_n / f(n) \leq \alpha, \mathbb{P}_{\boldsymbol{\nu}}^{x_0} \text{-a.s.}$$

The details can be found in Section 5.

We observe that, in principle, the main results of this section can be extended to BRWs in varying environment; these are BRWs where $\boldsymbol{\mu} = \{\mu_{x,n}\}_{x \in X, n \in \mathbb{N}}$ and the reproduction law of a particle at x at time n is $\mu_{x,n}$. Such processes admit a space-time counterpart (as in the proof of Lemma 5.4, see also [4]) which is a BRW in a fixed environment. Such an extension, however, goes beyond the purpose of this paper.

5. PROOFS

Since Lemma 2.5 and Proposition 2.6 hold for every initial condition η , in order to avoid a cumbersome notation, in the proofs we use $\mathbb{P}$ and $\mathbb{E}$ instead of $\mathbb{P}^\eta$ and $\mathbb{E}^\eta$.

Proof of Lemma 2.5. Let $k = 1$. We write the explicit expression of η_{m+1} as a function of η_m and identify $\eta_m(\omega)$ with a function $h \in S_X$. Then

$$\begin{aligned} \mathbb{E}[\mathbf{z}^{\eta_{m+1}} | \mathcal{F}_m] &= \mathbb{E}\left[\prod_{x \in X} \mathbf{z}(x)^{\sum_{y \in X} \sum_{i=1}^{\eta_m(y)} f_{i,m,y}(x)} | \mathcal{F}_m\right] \\ &= \sum_{h \in S_X} \mathbb{1}(\eta_m = h) \mathbb{E}\left[\prod_{x \in X} \mathbf{z}(x)^{\sum_{y \in X} \sum_{i=1}^{h(y)} f_{i,m,y}(x)} | \mathcal{F}_m\right], \mathbb{P}\text{-a.s.} \end{aligned} \quad (5.7)$$

where in the last equality we used the fact that η_m is $\mathcal{F}_m$ -measurable. Using independence of $f_{i,m,y}$ and $\mathcal{F}_m$, we get

$$\mathbb{E}\left[\prod_{x \in X} \mathbf{z}(x)^{\sum_{y \in X} \sum_{i=1}^{h(y)} f_{i,m,y}(x)} \middle| \mathcal{F}_m\right] = \mathbb{E}\left[\prod_{x \in X} \prod_{y \in X} \prod_{i=1}^{h(y)} \mathbf{z}(x)^{f_{i,m,y}(x)}\right], \quad \mathbb{P}\text{-a.s.}$$

Now, since $\{f_{i,m,y}(x)\}_{i,m \in \mathbb{N}, y \in X}$ is a family of independent random variables, this expectation can be written as (by definition of G)

$$\prod_{y \in X} \prod_{i=1}^{h(y)} \mathbb{E}\left[\prod_{x \in X} \mathbf{z}(x)^{f_{i,m,y}(x)}\right] = \prod_{y \in X} \prod_{i=1}^{h(y)} G(\mathbf{z}|y).$$

Thus (5.7) becomes

$$\begin{aligned} \mathbb{E}[\mathbf{z}^{\eta_{m+1}} | \mathcal{F}_m] &= \sum_{h \in S_X} \mathbb{1}(\eta_m = h) \prod_{y \in X} \prod_{i=1}^{h(y)} G(\mathbf{z}|y) = \sum_{h \in S_X} \mathbb{1}(\eta_m = h) \prod_{y \in X} G(\mathbf{z}|y)^{h(y)} \\ &= \sum_{h \in S_X} \mathbb{1}(\eta_m = h) \prod_{y \in X} G(\mathbf{z}|y)^{\eta_m(y)} = G(\mathbf{z})^{\eta_m}, \quad \mathbb{P}\text{-a.s.} \end{aligned} \quad (5.8)$$

which proves the claim for $k = 1$.

The claim is proven by induction on k . Indeed

$$\begin{aligned} \mathbb{E}[\mathbf{z}^{\eta_{m+k}} | \mathcal{F}_m] &= \mathbb{E}\left[\mathbb{E}[\mathbf{z}^{\eta_{m+k}} | \mathcal{F}_{m+k-1}] | \mathcal{F}_m\right] \\ &= \mathbb{E}\left[G(\mathbf{z})^{\eta_{m+k-1}} | \mathcal{F}_m\right] \\ &= \left(G^{(k-1)}(G(\mathbf{z}))\right)^{\eta_m} = (G^{(k)}(\mathbf{z}))^{\eta_m}, \quad \mathbb{P}\text{-a.s.} \end{aligned}$$

where in the last line we used the induction hypothesis and the definition of $G^{(k)}$. $\square$

Proof of Proposition 2.6. The two inequalities come from Lemma 2.5 and they hold for every initial state of the process; in particular if $\mathbf{z} \in F_G$, then $\{\mathbf{z}^{\eta_n}\}_{n \in \mathbb{N}}$ is a martingale.

Note that $\{\mathbf{z}^{\eta_n}\}_{n \in \mathbb{N}}$ is uniformly bounded by the constant function 1, whence it is a uniformly integrable family. Thus [18, Theorem 14.1] (or [11, Theorem 9.4.5]) implies that, for all $p \geq 1$

$$\mathbf{z}^{\eta_n} \rightarrow W_{\mathbf{z}}, \quad \mathbb{P}\text{-a.s. and in } L^p(\mathbb{P})$$

where the $L^p(\mathbb{P})$ convergence comes from the a.s. convergence and the Bounded Convergence Theorem. The $L^1(\mathbb{P})$ -convergence and the fact that $\mathbb{E}[\mathbf{z}^{\eta_m} | \mathcal{F}_n] = \mathbf{z}^{\eta_n}$ for all $m \geq n$ implies $\mathbb{E}[W_{\mathbf{z}} | \mathcal{F}_n] = \mathbf{z}^{\eta_n}$. $\square$

Proof of Corollary 2.7. Note that $\mathbb{E}^\eta[\mathbb{1}_{\mathcal{E}(A)} | \mathcal{F}_n] = \mathbf{q}(A)^{\eta_n}$. Indeed, it is enough to note that, for every sequence $\{f_i\}_{i=1}^n$ in S_X , the Markov property implies

$$\mathbb{P}^\eta(\mathcal{E}(A) | \eta_1 = f_1, \dots, \eta_n = f_n) = \mathbb{P}^x(\mathcal{E}(A) | \eta_n = f_n) = \mathbf{q}(A)^{f_n}.$$

By Proposition 2.6, $\{\mathbf{q}(A)^{\eta_n}\}_{n \in \mathbb{N}}$ is a martingale and, by [18, Theorem 14.2] (or [11, Theorem 9.4.8]), since $\mathbb{1}_{\mathcal{E}(A)} \in L^p(\mathbb{P}^x)$, then $\mathbb{E}^\eta[\mathbb{1}_{\mathcal{E}(A)} | \mathcal{F}_n] \rightarrow \mathbb{E}^\eta[\mathbb{1}_{\mathcal{E}(A)} | \mathcal{F}_\infty] = \mathbb{1}_{\mathcal{E}(A)}$, $\mathbb{P}^\eta$ -a.s. and in $L^p(\mathbb{P}^\eta)$ for all $p \geq 1$. $\square$

Proof of Corollary 2.8. The equivalence between (2) and (3) follows from the identity $\mathbb{P}^\eta = *_{x \in X} *_{i=1}^{\eta(x)} \mathbb{P}^x$ where $*$ is the usual convolution product of measures. Let us see the details.

(2) $\Rightarrow$ (3). There is nothing to prove.

(3) $\Rightarrow$ (2). Consider, on a suitable probability space, a family $\{\{\eta_n^{i,x}\}_n\}_{i \in \mathbb{N}, x \in X}$ of independent BRWs such that $\eta_0^{i,x} = \delta_x$. By the superimposition property (or by equation (2.1)) we have that $\eta_n := \sum_{x \in X} \sum_{i=1}^{\eta_n^{(x)}} \eta_n^{i,x}$ is a BRW with initial condition η . Whence the law of $\{\eta_n^{i,x}\}_n$ is $\mathbb{P}^x$ and

the law of $\{\eta_n\}_n$ is $\mathbb{P}^\eta$. Clearly $\mathbf{z}^{\eta_n} = \prod_{x \in X} \prod_{i=1}^{\eta(x)} \mathbf{z}^{\eta_n^{i,x}}$. Since $\mathbf{z}^{\eta_n^{i,x}} \rightarrow W_{\mathbf{z}}^{i,x}$ a.s., then $\mathbf{z}^{\eta_n} \rightarrow \prod_{x \in X} \prod_{i=1}^{\eta(x)} W_{\mathbf{z}}^{i,x}$ a.s. A similar argument holds for $\mathbf{v}$ instead of $\mathbf{z}$. By hypothesis $W_{\mathbf{z}}^{i,x} \geq W_{\mathbf{v}}^{i,x}$ a.s. whence $\prod_{x \in X} \prod_{i=1}^{\eta(x)} W_{\mathbf{z}}^{i,x} \geq \prod_{x \in X} \prod_{i=1}^{\eta(x)} W_{\mathbf{v}}^{i,x}$ a.s.

(1) $\Rightarrow$ (2). Suppose that $\mathbf{z} \geq \mathbf{v}$, then $\mathbf{z}^{\eta_n} \geq \mathbf{v}^{\eta_n}$ $\mathbb{P}^\eta$ -a.s. thus by taking the limit as $n \rightarrow +\infty$, Proposition 2.6 yields $\mathbb{P}^\eta(W_{\mathbf{z}} \geq W_{\mathbf{v}}) = 1$.

(2) $\Rightarrow$ (1). Finally, suppose that $\mathbb{P}^x(W_{\mathbf{z}} \geq W_{\mathbf{v}}) = 1$ for every $x \in X$. Then $\mathbf{z}(x) = \mathbb{E}^x[W_{\mathbf{z}}] \geq \mathbb{E}^x[W_{\mathbf{v}}] = \mathbf{v}(x)$ for all $x \in X$. $\square$

In order to prove Theorem 3.1(2), we need Lemma 3.5, thus we proceed with its proof first.

Proof of Lemma 3.5. Let $\inf_{x \in X} \mathbf{q}(x, A) =: \alpha > 0$. If $\alpha = 1$, then there is nothing to prove, since $\mathbb{P}^\eta(\mathcal{S}(A)) = 0$. If $\alpha < 1$, from Corollary 2.7 we have that, $\mathbb{P}^\eta$ -a.s. on $\mathcal{S}(A)$,

$$0 = \lim_{n \rightarrow +\infty} \mathbf{q}(A)^{\eta_n} \geq \lim_{n \rightarrow +\infty} \alpha^{\sum_{x \in X} \eta_n(x)}.$$

Thus $\mathbb{P}^\eta(\{\lim_{n \rightarrow +\infty} \alpha^{\sum_{x \in X} \eta_n(x)} = 0\} \cap \mathcal{S}(A)) = \mathbb{P}^\eta(\mathcal{S}(A))$, which implies the claim. $\square$

Proof of Theorem 3.1.

(1) The statement is [1, Corollary 4.2].

(2) Assume now that $\inf_{x \in X} \mathbf{q}(x, X) > 0$. By hypothesis, $\mathbf{z}(x) \geq \mathbf{q}(x, X)$ for all $x \in X$ and there exists x_0 such that $\mathbf{z}(x_0) > \mathbf{q}(x_0, X)$. Suppose by contradiction that $\mathbf{z}(x) \leq 1 - \varepsilon$ for all $x \in X$, for some $\varepsilon > 0$. Let $W_{\mathbf{z}} := \lim_{n \rightarrow +\infty} \mathbf{z}^{\eta_n}$ (which exists by Lemma 2.5). On $\mathcal{E}(X)$ we have $W_{\mathbf{z}} = 1$ (see discussion after Proposition 2.6). By Lemma 3.5, on $\mathcal{S}(X)$, $W_{\mathbf{z}} = \lim_{n \rightarrow +\infty} \mathbf{z}^{\eta_n} \leq \lim_{n \rightarrow +\infty} (1 - \varepsilon)^{\sum_{x \in X} \eta_n(x)} = 0$, $\mathbb{P}^{x_0}$ -a.s. Whence $W_{\mathbf{z}} = \mathbb{1}_{\mathcal{E}(X)}$, $\mathbb{P}^{x_0}$ -a.s. Thus

$$\mathbf{q}(x_0, X) < \mathbf{z}(x_0) = \mathbb{E}^{x_0}[W_{\mathbf{z}}] = \mathbb{E}^{x_0}[\mathbb{1}_{\mathcal{E}(X)}] = \mathbf{q}(x_0, X)$$

which is a contradiction. $\square$

We prove now that the binary relation $\geq_{\text{germ}}$ is a partial order on the space of all generating functions defined on $[0, 1]^X$. To this aim we need a lemma.

Lemma 5.1. *Let $G(\mathbf{z})$ be a holomorphic function defined on D^n where D is the closed unit ball in $\mathbb{C}$. Suppose that G vanishes on $[\delta, 1]^n$ for some $0 \leq \delta < 1$. Then G vanishes on D^n .*

Proof. We first prove the statement for $n = 2$. Let $\mathbf{z} = (\mathbf{z}(1), \mathbf{z}(2))$. Fix $\mathbf{z}(1) \in [\delta, 1]$ (meaning that the imaginary part of $\mathbf{z}(1)$ is 0 and the real part belongs to the interval) and let $G_{\mathbf{z}(1)}(w) := G(\mathbf{z}(1), w)$ for all $w \in D$. By hypothesis, $G_{\mathbf{z}(1)}$ is a holomorphic function of one complex variable which vanishes on the real interval $[\delta, 1]$. Since this interval has at least one limit point in D , then it is well-known that $G_{\mathbf{z}(1)}(w) = 0$ for all $w \in D$ (see for instance [17, Theorem 10.18]). This proves that G vanishes on $[\delta, 1] \times D$.

Now fix $s \in D$ and define $G_s(\xi) := G(\xi, s)$ for all $\xi \in D$. This is again a holomorphic function of one complex variable which vanishes on the real interval $[\delta, 1]$. By the same argument as before, $G_s(\xi) = 0$ for all $\xi \in D$. This means that $G(\mathbf{z})$ vanishes in D^2 . The statement for a general $n > 2$ follows easily by induction. $\square$

Proof of Proposition 4.2. The relation $\geq_{\text{germ}}$ is clearly reflexive and transitive. Let us prove it is antisymmetric. Suppose that $\boldsymbol{\mu} \geq_{\text{germ}} \boldsymbol{\nu}$ and $\boldsymbol{\nu} \geq_{\text{germ}} \boldsymbol{\mu}$, that is, there exists $\delta < 1$ such that for all $\mathbf{z} \in [\delta, 1]^X$, then $G_{\boldsymbol{\mu}}(\mathbf{z}) = G_{\boldsymbol{\nu}}(\mathbf{z})$; we prove that $\boldsymbol{\mu} = \boldsymbol{\nu}$ (this is equivalent to $G_{\boldsymbol{\mu}} = G_{\boldsymbol{\nu}}$ as discussed in Section 2.3). It is enough to prove that $\mu_x = \nu_x$ (or equivalently that $G_{\boldsymbol{\mu}}(\cdot|x) = G_{\boldsymbol{\nu}}(\cdot|x)$) for every fixed $x \in X$.

To this aim, note that equation (2.3) defines a continuous function G on D^X where $D := \{z \in \mathbb{C} : |z| \leq 1\}$ is the closed disk of radius 1 in the complex plane. Whence when X is finite, for

every fixed $x \in X$, the generating function $G(\cdot|x)$ can be seen as a holomorphic function of several variables. In this case the result follow from Lemma 5.1; indeed since $G_{\mu}(\cdot|x) - G_{\nu}(\cdot|x)$ vanishes on $[\delta, 1]^X$, then, by Lemma 5.1, it vanishes on D^X whence $\mu_x(f) - \nu_x(f) = 0$ for every $f \in S_X$.

Now let X be infinite; given a subset $Y \subseteq X$ define $V(Y) := \{\mathbf{z} \in [0, 1]^X : \mathbf{z}(x) = 1, \forall x \in X \setminus Y\}$ and let $\pi : V(Y) \mapsto [0, 1]^Y$ be the bijective map defined as $\pi(\mathbf{z}) := \mathbf{z}|_Y$ (the restriction of $\mathbf{z}$ to Y). Given $f \in S_X$ define $\langle f \rangle_Y := \{g \in S_X : g|_Y = f|_Y\}$ the set of functions extending the restriction of f to Y ; moreover define $S_X(Y) := \{f \in S_X : \{f > 0\} \subseteq Y\}$ the set of finitely supported functions on X whose support is in Y . Clearly, since $x \mapsto f(x)\mathbb{1}(x \in Y)$ is a map in $S_X(Y)$ and $\langle f \rangle_Y = \langle f(\cdot)\mathbb{1}(\cdot \in Y) \rangle_Y$, then the map $f \mapsto \langle f \rangle_Y$ is a bijection from $S_X(Y)$ onto $\{\langle f \rangle_f : f \in S_X\}$. Roughly speaking, $\langle f \rangle_Y$ are equivalence classes containing exactly one function $g \in S_X(Y)$ and since every $g \in S_X(Y)$ belongs to a class, there is a one to one correspondence between $S_X(Y)$ and $\{\langle f \rangle_f : f \in S_X\}$.

We observe now that $G_{\mu}(\cdot|x)|_{V(Y)}$ and $G_{\nu}(\cdot|x)|_{V(Y)}$ can be seen as functions defined on $[0, 1]^Y$, indeed $G_{\mu}(\pi^{-1}(\cdot)|x)|_{V(Y)} = G_{\mu}(\pi^{-1}(\cdot)|x)$ and π^{-1} is a bijection from $[0, 1]^Y$ onto $V(Y)$ (and the same holds for ν). More precisely

$$G_{\mu}(\pi^{-1}(\mathbf{z})|x) = \sum_{f \in S_X(Y)} \mu_x(\langle f \rangle_Y) \prod_{y \in Y} \mathbf{z}(y)^{f(y)},$$

and an analogous expression holds for G_{ν} . Suppose that Y is finite; since $G_{\mu}(\pi^{-1}(\cdot)|x) = G_{\nu}(\pi^{-1}(\cdot)|x)$ on $[\delta, 1]^Y$ the same equality holds on $V(Y)$ (by Lemma 5.1). This implies easily that $\mu_x(\langle f \rangle_Y) = \nu_x(\langle f \rangle_Y)$ for every $f \in S_X(Y)$ or, equivalently, for every $f \in S_X$. Consider now a fixed sequence of finite subsets of X , say $\{Y_n\}_{n \in \mathbb{N}}$, such that $Y_n \subseteq Y_{n+1}$ and $\bigcup_{n \in \mathbb{N}} Y_n = X$. Then, for all $f \in S_X$ we have $\langle f \rangle_{Y_{n+1}} \subseteq \langle f \rangle_{Y_n}$ and $\bigcap_{n \in \mathbb{N}} \langle f \rangle_{Y_n} = \{f\}$, therefore

$$\mu_x(f) = \lim_{n \rightarrow +\infty} \mu_x(\langle f \rangle_{Y_n}) = \lim_{n \rightarrow +\infty} \nu_x(\langle f \rangle_{Y_n}) = \nu_x(f).$$

□

Before proving Theorem 4.3, as a warm-up we prove Theorem 4.4; to this aim we need a preparatory lemma.

Lemma 5.2. *Let $\mu \geq_{pgf} \nu$ and $A \subseteq X$. Then $\mathbb{E}_{\mu}^x[\exp(-tL(A))] \leq \mathbb{E}_{\nu}^x[\exp(-tL(A))]$ for all $t \geq 0$, $x \in X$.*

Proof. We prove by induction on n that $\mathbb{E}_{\mu}^x[\exp(-tL_n(A))] \leq \mathbb{E}_{\nu}^x[\exp(-tL_n(A))]$ for all $t \geq 0$, $x \in X$. the claim follows from the Bounded Convergence Theorem.

If $n = 0$, then, for all $x \in X$ and $t \in [0, +\infty)$, $\mathbb{E}_{\mu}^x[\exp(-tL_0(A))] = \exp(-t)\mathbb{1}_A(x) + \mathbb{1}_{A^c}(x) = \mathbb{E}_{\nu}^x[\exp(-tL_0(A))]$.

Let $n \geq 0$ and suppose that $\mathbb{E}_{\mu}^x[\exp(-tL_n(A))] \leq \mathbb{E}_{\nu}^x[\exp(-tL_n(A))]$ for all $t \geq 0$, $x \in X$. We have

$$\begin{aligned} \mathbb{E}_{\mu}^x[\exp(-tL_{n+1}(A))] &= (\exp(-t)\mathbb{1}_A(x) + \mathbb{1}_{A^c}(x)) \sum_{f \in S_X} \mu_x(f) \prod_{y \in X} \mathbb{E}_{\mu}^y[\exp(-tL_n(A))]^{f(y)} \\ &= (\exp(-t)\mathbb{1}_A(x) + \mathbb{1}_{A^c}(x)) G_{\mu}(\mathbb{E}_{\mu}^{(\cdot)}[\exp(-tL_n(A))]|x) \\ &\stackrel{(\text{induction})}{\leq} (\exp(-t)\mathbb{1}_A(x) + \mathbb{1}_{A^c}(x)) G_{\mu}(\mathbb{E}_{\nu}^{(\cdot)}[\exp(-tL_n(A))]|x) \\ &\stackrel{(\text{pgf order})}{\leq} (\exp(-t)\mathbb{1}_A(x) + \mathbb{1}_{A^c}(x)) G_{\nu}(\mathbb{E}_{\nu}^{(\cdot)}[\exp(-tL_n(A))]|x) \\ &= \mathbb{E}_{\nu}^x[\exp(-tL_{n+1}(A))]. \end{aligned}$$

□

Proof of Theorem 4.4.

(1) By the Bounded Convergence Theorem and Lemma 5.2, for all $x \in X$ and $A \subseteq X$ we have

$$\begin{aligned} \mathbf{q}^\mu(x, A) &= \mathbb{P}_\mu^x(L(A) < +\infty) = \lim_{t \rightarrow 0^+} \mathbb{E}_\mu^x[\exp(-tL(A))] \\ &\leq \lim_{t \rightarrow 0^+} \mathbb{E}_\nu^x[\exp(-tL(A))] = \mathbb{P}_\nu^x(L(A) < +\infty) = \mathbf{q}^\nu(x, A). \end{aligned}$$

(2) We know from (1) and from the hypotheses that,

$$\mathbf{q}^\mu(A) \leq \mathbf{q}^\nu(A) = \mathbf{q}^\nu(X).$$

Then $\sup_{x \in X} \mathbf{q}^\mu(x, A) < 1$ which, according to Theorem 3.1, implies $\mathbf{q}^\mu(A) = \mathbf{q}^\mu(X)$. $\square$

We can prove now Theorem 4.3. We need two preparatory lemmas. The first one is the analogous of [13, Lemma 2.3] and the proof is on the same line. As usual, $\vee$ and $\wedge$ denote the maximum and the minimum respectively.

Lemma 5.3. *Let $\mu \geq_{\text{germ}} \nu$ and $A \subseteq X$. Then there exists $\delta < 1$ such that for all $t \in (\delta, 1]$ and all $x \in X$,*

$$\mathbb{E}_\nu^x[t^{\mathbb{1}(L(A) > 0)}] \geq t \vee \mathbb{E}_\mu^x[t^{L(A)}].$$

Proof. From Definition 4.1, for every $\mathbf{z} \in [\delta, 1]^X$ (that is, for every $\mathbf{z} \in [0, 1]^X$ such that $\delta \mathbf{1} \leq \mathbf{z} \leq \mathbf{1}$) we have $G_\mu(\mathbf{z}) \leq G_\nu(\mathbf{z})$. If $t = 1$ there is nothing to prove. Let us fix $t \in (\delta, 1)$. Clearly $G_\mu(\mathbf{z}) \leq G_\nu(\mathbf{z})$ for all $\mathbf{z} \in [t, 1]^X$.

The strategy of the proof is to find $\mathbf{v}_\infty, \mathbf{w}_\infty \in [t, 1]^X$ such that $\mathbb{E}_\nu^x[t^{\mathbb{1}(L(A) > 0)}] \geq \mathbf{v}_\infty(x) \geq \mathbf{w}_\infty(x) \geq t \vee \mathbb{E}_\mu^x[t^{L(A)}]$ for all $x \in X$. To this aim define $I_\mu, I_\nu : [t, 1]^X \mapsto [t, 1]^X$ as follows

$$\begin{aligned} I_\mu \mathbf{z}(x) &:= (t \vee t^{\mathbb{1}(x \in A)} G_\mu(\mathbf{z}|x)) \wedge \mathbf{z}(x) \\ &= t \vee (t^{\mathbb{1}(x \in A)} G_\mu(\mathbf{z}|x) \wedge \mathbf{z}(x)) = \begin{cases} t & x \in A \\ t \vee (G_\mu(\mathbf{z}|x) \wedge \mathbf{z}(x)) & x \notin A \end{cases} \end{aligned} \quad (5.9)$$

and I_ν is defined analogously by using G_ν instead of G_μ . It is easy to show that I_μ, I_ν are nondecreasing, continuous functions on $[t, 1]^X$. Moreover, for all $\mathbf{z} \in [t, 1]^X$ we have $t\mathbf{1} \leq I_\mu \mathbf{z} \leq I_\nu \mathbf{z} \leq \mathbf{z}$. Define recursively

$$\begin{cases} \mathbf{v}_0(x) = \mathbf{w}_0(x) := t^{\mathbb{1}(x \in A)}, & \forall x \in X, \\ \mathbf{v}_{n+1} := I_\nu \mathbf{v}_n, & \forall n \in \mathbb{N}, \\ \mathbf{w}_{n+1} := I_\mu \mathbf{w}_n, & \forall n \in \mathbb{N}, \end{cases}$$

whence $\{\mathbf{w}_n\}_{n \in \mathbb{N}}$ and $\{\mathbf{v}_n\}_{n \in \mathbb{N}}$ are nonincreasing sequences in $[t, 1]^X$ such that $t\mathbf{1} \leq \mathbf{w}_n \leq \mathbf{v}_n \leq \mathbf{z}$, therefore $\mathbf{v}_n \downarrow \mathbf{v}_\infty$, $\mathbf{w}_n \downarrow \mathbf{w}_\infty$ and $t\mathbf{1} \leq \mathbf{w}_\infty \leq \mathbf{v}_\infty \leq \mathbf{z}$. By the same arguments of Proposition 2.4, we have $I_\nu \mathbf{v}_\infty = \mathbf{v}_\infty$ and $I_\mu \mathbf{w}_\infty = \mathbf{w}_\infty$. We prove now, by induction on $n \in \mathbb{N}$, that $\mathbf{w}_n(x) \geq t \vee \mathbb{E}_\mu^x[t^{L_n(A)}]$ for all $n \in \mathbb{N}$ which, in turn, implies $\mathbf{w}_\infty(x) \geq t \vee \mathbb{E}_\mu^x[t^{L(A)}]$. If $n = 0$, then $\mathbf{w}_0(x) = t^{\mathbb{1}(x \in A)} \geq t \vee \mathbb{E}_\mu^x[t^{L_0(A)}]$ since $\mathbb{1}(x \in A) = L_0(A)$. Suppose that the inequality holds for $n \in \mathbb{N}$, then, by using that the BRW is a stationary Markov process and that the set of descendants of different particles belonging to a fixed generation are independent, we have for all $x \in X$

$$\begin{aligned} \mathbb{E}_\mu^x[t^{L_{n+1}(A)}] &= \mathbb{E}_\mu^x[\mathbb{E}_\mu^x[t^{L_{n+1}(A)} | \mathcal{F}_1]] = t^{\mathbb{1}(x \in A)} \sum_{f \in S_X} \mu_x(f) \prod_{y \in X} \mathbb{E}_\mu^y[t^{L_n(A)}]^{f(y)} \\ &= t^{\mathbb{1}(x \in A)} G_\mu(\mathbb{E}_\mu^{(\cdot)}[t^{L_n(A)}] | x) \leq t^{\mathbb{1}(x \in A)} G_\mu(\mathbf{w}_n | x) \end{aligned}$$

(where $\mathbb{E}_\mu^{(\cdot)}[t^{L_n(A)}]$ represents the vector $y \mapsto \mathbb{E}_\mu^y[t^{L_n(A)}]$). Note that in the last inequality we used the induction hypothesis and the fact that G_μ is nondecreasing. Clearly $\mathbb{E}_\mu^x[t^{L_{n+1}(A)}] \leq \mathbb{E}_\mu^x[t^{L_n(A)}] \leq \mathbf{w}_n(x)$, thus

$$t \vee \mathbb{E}_\mu^x[t^{L_{n+1}(A)}] \leq t \vee (\mathbf{w}_n(x) \wedge t^{\mathbb{1}(x \in A)} G_\mu(\mathbf{w}_n | x)) = I_\mu \mathbf{w}_n = \mathbf{w}_{n+1}.$$

Now we prove that $\mathbb{E}_{\nu}^x[t^{\mathbb{1}(L(A)>0)}] \geq \mathbf{v}_{\infty}(x)$ for all $x \in X$. Let us define $D := \{x \in X : \mathbf{v}_{\infty}(x) = t\}$; clearly, since $t \leq \mathbf{v}_{\infty}(x) \leq t^{\mathbb{1}(x \in A)}$ for all $x \in X$, then $D \supseteq A$. Define recursively

$$\begin{cases} \mathbf{h}_0(x) := t^{\mathbb{1}(x \in D)}, & \forall x \in X \\ \mathbf{h}_{n+1} := I_{\nu} \mathbf{h}_n & \forall n \in \mathbb{N}. \end{cases}$$

The sequence $\{\mathbf{h}_n\}_{n \in \mathbb{N}}$ is nondecreasing therefore $\mathbf{h}_n \downarrow \mathbf{h}_{\infty}$ for some $\mathbf{h}_{\infty} \in [t, 1]^X$. Moreover, since $I_{\nu} \mathbf{v}_{\infty} = \mathbf{v}_{\infty} \leq \mathbf{h}_0$, then $t \leq \mathbf{v}_{\infty}(x) \leq \mathbf{h}_{\infty}(x) \leq t^{\mathbb{1}(x \in D)}$; thus $\mathbf{h}_n(x) = t$ for all $x \in D$. On the other hand, if $x \notin D$, then, by definition of D , $t < \mathbf{v}_{\infty}(x) \leq \mathbf{h}_n(x)$ for all $n \in \mathbb{N}$ and $G_{\nu}(\mathbf{h}_n|x) \geq G_{\nu}(\mathbf{v}_{\infty}|x) = \mathbf{v}_{\infty}(x) > t$ for all $n \in \mathbb{N}$. Therefore, by using equation (5.9),

$$\mathbf{h}_{n+1}(x) = \begin{cases} t & x \in D \\ G_{\nu}(\mathbf{h}_n|x) \wedge \mathbf{h}_n(x) & x \notin D. \end{cases}$$

Define $E_n(D)$ as the number of particles in D by time n with no ancestors in D and let $E(D) := \lim_{n \rightarrow +\infty} E_n(D)$ (note that $E_{n+1}(D) \geq E_n(D)$). If, for instance, $x \in D$, then $E_n(D) = 1$ for all $n \in \mathbb{N}$. We want to prove that $\mathbf{h}_n(x) = \mathbb{E}_{\nu}^x[t^{E_n(D)}]$ for all $x \in X$ which, according to the Bounded Convergence Theorem, implies $\mathbf{h}_{\infty}(x) = \mathbb{E}_{\nu}^x[t^{E(D)}]$ for all $x \in X$. To this aim note that $L(A) > 0$ implies $E(D) \geq 1$, therefore $\mathbb{E}_{\nu}^x[t^{E(D)}] \leq \mathbb{E}_{\nu}^x[t^{\mathbb{1}(L(A)>0)}]$ for all $x \in X$. Define $\tilde{\mathbf{h}}_n(x) := \mathbb{E}_{\nu}^x[t^{E_n(D)}]$ for all $x \in X$. By using again the fact that the BRW is a stationary Markov process and that the progenies of different particles are independent, we see that the (nonincreasing) sequence $\{\tilde{\mathbf{h}}_n(x)\}_{n \in \mathbb{N}}$ satisfies the following recursive equation for all $x \in X$

$$\begin{aligned} \tilde{\mathbf{h}}_{n+1}(x) &= \mathbb{E}_{\nu}^x[t^{E_{n+1}(D)}] = \begin{cases} t & x \in D \\ \mathbb{E}_{\nu}^x[\mathbb{E}_{\nu}^x[t^{E_{n+1}(D)}|\mathcal{F}_1]] = (\spadesuit) & x \notin D \end{cases} \\ (\spadesuit) &= \sum_{f \in S_X} \nu_x(f) \prod_{y \in X} \mathbb{E}_{\nu}^y[t^{E_n(D)}]^{f(y)} = G_{\nu}(\tilde{\mathbf{h}}_n|x) = \tilde{\mathbf{h}}_n(x) \wedge G_{\nu}(\tilde{\mathbf{h}}_n|x) \end{aligned}$$

where, in the last equality, we used the fact that, by definition, $\tilde{\mathbf{h}}_{n+1}(x) \leq \tilde{\mathbf{h}}_n(x)$ for all $x \in X$, which implies $G_{\nu}(\tilde{\mathbf{h}}_n|x) = \tilde{\mathbf{h}}_{n+1}(x) \leq \tilde{\mathbf{h}}_n(x)$ for all $x \notin D$. We observe that $\tilde{\mathbf{h}}_0 = \mathbf{h}_0$ since $\mathbb{E}_{\nu}^x[E_0(D)] = \mathbb{1}(x \in D)$ for all $x \in X$; moreover the sequences $\{\tilde{\mathbf{h}}_n(x)\}_{n \in \mathbb{N}}$ and $\{\mathbf{h}_n(x)\}_{n \in \mathbb{N}}$ satisfy the same recursive equation, hence $\tilde{\mathbf{h}}_n = \mathbf{h}_n$ for all $n \in \mathbb{N}$. This yields

$$\mathbb{E}_{\nu}^x[t^{\mathbb{1}(L(A)>0)}] \geq \mathbb{E}_{\nu}^x[t^{E(D)}] = \lim_{n \rightarrow +\infty} \tilde{\mathbf{h}}_n(x) = \lim_{n \rightarrow +\infty} \mathbf{h}_n(x) = \mathbf{h}_{\infty} \geq \mathbf{v}_{\infty}.$$

□

Lemma 5.4. *Let $\mu \geq_{\text{germ}} \nu$ and $A \subseteq X$. Then there exists $\delta < 1$ such that for all $t \in (\delta, 1)$ and all $x \in X$,*

$$\mathbf{q}^{\nu}(x, A) \geq \frac{\mathbf{q}^{\mu}(x, A) \vee t - t}{1 - t}.$$

Proof. In order to prove this lemma (by using Lemma 5.3) we define an auxiliary space-time version of the BRW (as in [13, Lemma 2.3]). More precisely, given a BRW $\{\eta_n\}_{n \in \mathbb{N}}$ on X we denote by $\{\eta_n^{st}\}_{n \in \mathbb{N}}$ a BRW on $X \times \mathbb{N}$ that we call *space-time version* of the original process and which is defined by $\eta_n^{st}(x, m) := \eta_n(x) \delta(n, m)$ (where $\delta(n, m) = 1$ if $n = m$ and 0 otherwise). Roughly speaking, the particles in x at time n in the original BRW, are now placed in (x, n) at time n in the st-BRW. The space-time version of μ , say μ^{st} is defined as follows, $\forall g \in S_{X \times \mathbb{N}}$ and $\forall (x, n) \in X \times \mathbb{N}$,

$$\mu_{(x, n)}^{st}(g) = \begin{cases} \mu_x(f) & \text{if } g = f \otimes \delta_{n+1} \\ 0 & \text{otherwise} \end{cases}$$

where $(f \otimes \delta_i)(y, j) := f(y) \delta(i, j)$ for all $(y, j) \in X \times \mathbb{N}$.

Elementary computations show that for all $\mathbf{z} \in [0, 1]^{X \times \mathbb{N}}$, $G_{\boldsymbol{\mu}^{st}}(\mathbf{z}|(x, n)) = G_{\boldsymbol{\mu}}(\mathbf{z}(\cdot, n+1)|x)$ and $G_{\boldsymbol{\nu}^{st}}(\mathbf{z}|(x, n)) = G_{\boldsymbol{\nu}}(\mathbf{z}(\cdot, n+1)|x)$. If $\boldsymbol{\mu} \geq_{\text{germ}} \boldsymbol{\nu}$, then $\boldsymbol{\mu}^{st} \geq_{\text{germ}} \boldsymbol{\nu}^{st}$. Indeed if $\mathbf{z} \in [\delta, 1]^{X \times \mathbb{N}}$ (where $\delta < 1$), then $\mathbf{z}(\cdot, n) \in [\delta, 1]^X$ for all $n \in \mathbb{N}$ whence

$$G_{\boldsymbol{\mu}^{st}}(\mathbf{z}|(x, n)) = G_{\boldsymbol{\mu}}(\mathbf{z}(\cdot, n+1)|x) \leq G_{\boldsymbol{\nu}}(\mathbf{z}(\cdot, n+1)|x) = G_{\boldsymbol{\nu}^{st}}(\mathbf{z}|(x, n))$$

for all $(x, n) \in X \times \mathbb{N}$.

Moreover $A \subseteq X$ is visited infinitely often by $(X, \boldsymbol{\mu})$ (resp. $(X, \boldsymbol{\nu})$) if and only if $A \times \mathbb{N}$ is visited infinitely often by $(X \times \mathbb{N}, \boldsymbol{\mu}^{st})$ (resp. $(X \times \mathbb{N}, \boldsymbol{\nu}^{st})$). In particular $\mathbf{q}^{\boldsymbol{\mu}}(x, A) = \mathbf{q}^{\boldsymbol{\mu}^{st}}((x, n), A \times \mathbb{N})$ and $\mathbf{q}^{\boldsymbol{\nu}}(x, A) = \mathbf{q}^{\boldsymbol{\nu}^{st}}((x, n), A \times \mathbb{N})$ for all $(x, n) \in X \times \mathbb{N}$, $A \subseteq X$. Thus, it suffices to prove the lemma for the space-time version of the BRW.

To avoid a cumbersome notation, for the rest of the proof we write $\boldsymbol{\mu}$ and $\boldsymbol{\nu}$ instead of $\boldsymbol{\mu}^{st}$ and $\boldsymbol{\nu}^{st}$ respectively. Moreover we use $\mathbb{P}_{\boldsymbol{\mu}}^{x,n}$ and $\mathbb{P}_{\boldsymbol{\nu}}^{x,n}$ to denote the laws of the space-time processes starting from (x, n) . Given $A \subseteq X \times \mathbb{N}$, we define $A_k := A \cap (X \times [k, +\infty))$. We observe that

$$L(A) = +\infty \iff L(A_k) > 0, \forall k \in \mathbb{N} \iff L(A_k) > 0, \text{ for infinitely many } k \in \mathbb{N}$$

since $\{L(A_{k+1}) > 0\} \subseteq \{L(A_k) > 0\}$ and at every fixed time the number of particles is finite. Whence $\{L(A) = +\infty\} = \bigcap_{k \in \mathbb{N}} \{L(A_k) > 0\}$ and $\{L(A) < +\infty\} = \liminf_{k \in \mathbb{N}} \{L(A_k) = 0\}$. This implies $\mathbb{1}(L(A_k) > 0) \downarrow \mathbb{1}(L(A) = +\infty)$. Note that $L(A) = +\infty$ implies $L(A_k) = +\infty$ for all $k \in \mathbb{N}$ while $L(A) < +\infty$ implies $L(A_k) = 0$ eventually as $k \rightarrow +\infty$.

We apply Lemma 5.3 to A_k and, for every fixed $(x, n) \in X \times \mathbb{N}$, we obtain

$$\mathbb{E}_{\boldsymbol{\nu}}^{x,n}[t^{\mathbb{1}(L(A_k) > 0)}] \geq t \wedge \mathbb{E}_{\boldsymbol{\mu}}^{x,n}[t^{L(A_k)}]. \quad (5.10)$$

According to the Monotone Convergence Theorem

$$\lim_{k \rightarrow +\infty} \mathbb{E}_{\boldsymbol{\nu}}^{x,n}[t^{\mathbb{1}(L(A_k) > 0)}] = \mathbb{E}_{\boldsymbol{\nu}}^{x,n}[t^{\mathbb{1}(L(A) = +\infty)}] = t(1 - \mathbf{q}^{\boldsymbol{\nu}}((x, n), A)) + \mathbf{q}^{\boldsymbol{\nu}}((x, n), A). \quad (5.11)$$

According to the Bounded Convergence Theorem, if $t < 1$

$$\lim_{k \rightarrow +\infty} \mathbb{E}_{\boldsymbol{\mu}}^{x,n}[t^{L(A_k)}] = \mathbb{E}_{\boldsymbol{\mu}}^{x,n}[\mathbb{1}(L(A) < +\infty)] = \mathbf{q}^{\boldsymbol{\mu}}((x, n), A). \quad (5.12)$$

By using equations (5.10), (5.11) and (5.12) we obtain

$$t(1 - \mathbf{q}^{\boldsymbol{\nu}}((x, n), A)) + \mathbf{q}^{\boldsymbol{\nu}}((x, n), A) \geq t \wedge \mathbf{q}^{\boldsymbol{\mu}}((x, n), A)$$

which yields the result. $\square$

Proof of Theorem 4.3.

- (1) Fix $x \in X$ and suppose that $\mathbf{q}^{\boldsymbol{\mu}}(x, A) = 1$. Then by Lemma 5.4, if we choose $t \in (\delta, 1)$ we have

$$\mathbf{q}^{\boldsymbol{\nu}}(x, A) \geq \frac{\mathbf{q}^{\boldsymbol{\mu}}(x, A) \vee t - t}{1 - t} = \frac{1 - t}{1 - t} = 1.$$

- (2) Suppose that $\mathbf{q}^{\boldsymbol{\nu}}(A) = \mathbf{q}^{\boldsymbol{\nu}}(X)$ and that $\sup_{x \in X} \mathbf{q}^{\boldsymbol{\nu}}(x, X) < 1$. Then, by Lemma 5.4,

$$1 > \sup_{x \in X} \mathbf{q}^{\boldsymbol{\nu}}(x, X) = \sup_{x \in X} \mathbf{q}^{\boldsymbol{\nu}}(x, A) \geq \frac{\sup_{x \in X} \mathbf{q}^{\boldsymbol{\mu}}(x, A) \vee t - t}{1 - t} \geq \frac{\sup_{x \in X} \mathbf{q}^{\boldsymbol{\mu}}(x, A) - t}{1 - t}$$

which is equivalent to $\sup_{x \in X} \mathbf{q}^{\boldsymbol{\mu}}(x, A) < 1$. According to Theorem 3.1 the last inequality implies $\mathbf{q}^{\boldsymbol{\mu}}(A) = \mathbf{q}^{\boldsymbol{\mu}}(X)$. $\square$

Proof of Proposition 4.7. (1) $\Rightarrow$ (2). Using the hypothesis and the expression for ϕ_x , we get that for every $t \in [\delta, 1]$ and for all $x \in X$,

$$\phi_x^{\boldsymbol{\mu}}(t) = G_{\boldsymbol{\mu}}(t\mathbf{1}|x) \leq G_{\boldsymbol{\nu}}(t\mathbf{1}|x) = \phi_x^{\boldsymbol{\nu}}(t).$$

- (2) $\Rightarrow$ (1). Recall that, for BRWs with independent diffusion, the generating functions are $G_{\boldsymbol{\mu}}(\mathbf{z}|x) = \phi_x^{\boldsymbol{\mu}}(P\mathbf{z}(x))$ and $G_{\boldsymbol{\nu}}(\mathbf{z}|x) = \phi_x^{\boldsymbol{\nu}}(P\mathbf{z}(x))$. We observe that the map $\mathbf{z} \mapsto P\mathbf{z}$ is nondecreasing and

continuous from $[0, 1]^X$ into itself; in particular, if $\mathbf{z} \in [\delta, 1]^X$ for some $\delta < 1$, then $P\mathbf{z} \in [\delta, 1]^X$. Indeed $Pt\mathbf{1} = t\mathbf{1}$ therefore $\delta\mathbf{1} = P\delta\mathbf{1} \leq P\mathbf{z} \leq P\mathbf{1} = \mathbf{1}$. Take $\mathbf{z} \in [\delta, 1]^X$; then for all $x \in X$

$$G_\mu(\mathbf{z}|x) = \phi_x^\mu(P\mathbf{z}(x)) \leq \phi_x^\nu(P\mathbf{z}(x)) = G_\nu(\mathbf{z}|x)$$

where we used the inequality $\phi_x^\mu(t) \leq \phi_x^\nu(t)$ for $t = P\mathbf{z}(x) \in [\delta, 1]$ (due to the monotonicity of P). $\square$

Proof of Corollary 4.9. Consider, as in the proof of Lemma 5.4, the space-time version $\{\eta_n^{st}\}_{n \in \mathbb{N}}$ of the process. Clearly

$$\limsup_{n \rightarrow +\infty} \left\{ \sum_{y \in A_n} \eta_n(y) > 0 \right\} = \mathcal{S}^{st} \left(\bigcup_{n \in \mathbb{N}} (A_n \times \{n\}) \right)$$

where $\mathcal{S}^{st}(\cdot)$ is the survival event of the space-time process. Recall that, for all $A \subseteq X$, $\mathbf{q}^\mu(x, A) = \mathbf{q}^{\mu^{st}}((x, n), A \times \mathbb{N})$ and $\mathbf{q}^\nu(x, A) = \mathbf{q}^{\nu^{st}}((x, n), A \times \mathbb{N})$ for all $(x, n) \in X \times \mathbb{N}$. (1) and (2) follows from Theorem 4.3 applied to the space-time process. (3) follow from (2) by noting that $\limsup_{n \rightarrow +\infty} \{\sum_{y \in A_n} \eta_n(y) > 0\} \subseteq \mathcal{S}(X)$. $\square$

Details on Example 4.10. We note that

$$\mathbb{P}_\mu^{x_0}(\limsup_{n \rightarrow +\infty} M_n/f(n) \leq \alpha) = 1 \iff \mathbb{P}_\mu^{x_0} \left(\limsup_{n \rightarrow +\infty} \left\{ \sum_{y: d(x_0, y) \geq (\alpha + \varepsilon)f(n)} \eta_n(y) > 0 \right\} \right) = 0, \forall \varepsilon > 0$$

and a similar equality holds for ν . The result follows by applying Corollary 4.9 to $A_n^\varepsilon := \{y \in X: d(x_0, y) \geq (\alpha + \varepsilon)f(n)\}$. $\square$

6. APPENDIX: PRODUCT OF METRIC SPACES

In this appendix we show how the product of metric spaces can be endowed with a finite metric which generates the pointwise convergence topology. We also address separability and completeness. We note that $\mathbb{R}^X$ can be endowed with a finite metric which turns it into a Polish space.

Lemma 6.1. *Consider a metric space (Y, d) and a function $f \in L^1([0, +\infty))$ such that f is non increasing a.e. and $\int_0^\varepsilon f(t)dt > 0$ for all $\varepsilon > 0$. Then $d_1(x, y) := \int_0^{d(x, y)} f(t)dt$ for all $x, y \in Y$ defines a finite metric which generates the same topology.*

Proof. Note that f is a.s. nonnegative and $\int_0^a f(t)dt = 0$ if and only if $a = 0$. Whence $d_1(x, y) \geq 0$ for all $x, y \in Y$ and the equality holds if and only if $d(x, y) = 0$, that is, $x = y$. As for the triangle inequality

$$\begin{aligned} d_1(x, z) + d_1(z, y) &= \int_0^{d(x, z)} f(t)dt + \int_0^{d(z, y)} f(t)dt \\ &\geq \int_0^{d(x, z)} f(t)dt + \int_0^{d(z, y)} f(t + d(x, z))dt \\ &= \int_0^{d(x, z)} f(t)dt + \int_{d(x, z)}^{d(x, z) + d(z, y)} f(t)dt \\ &= \int_0^{d(x, z) + d(z, y)} f(t)dt \geq \int_0^{d(x, y)} f(t)dt = d_1(x, y). \end{aligned}$$

Finally $d_1(x, y) \leq \|f\|_1 := \int_0^\infty f(t)dt < +\infty$ for all $x, y \in Y$.

Let us prove that the topology is the same. On the one hand $B(x, r) = B_1(x, \int_0^r f(t)dt)$ for all $r > 0$. On the other hand, $\varepsilon \mapsto \int_0^\varepsilon f(t)dt$ is right continuous in 0, whence for every $r > 0$ there exists $\varepsilon > 0$ such that $0 < \int_0^\varepsilon f(t)dt =: r_1 < r$, that is, $B(x, r) \supseteq B_1(x, r_1)$. $\square$

An example is given by $f := \mathbb{1}_{[0,M]}$ which gives $d_1(x, y) = \min(d(x, y), M)$, where $M > 0$. Since the topology is the same, if the original metric space is separable (resp. complete) the same hold for the new one. The advantage of a finite metric is clear in the following lemma.

We suppose that $\{(Y_n, d_n)\}_{n \in J}$ is a countable (finite or infinite) sequence of finite metric spaces where $\sup_{x, y \in Y_n} d_n(x, y) = M_n < +\infty$.

Proposition 6.2. *Let $\{\alpha_n\}_{n \in J}$ a sequence of positive real numbers such that $\sum_{n \in J} \alpha_n M_n < +\infty$. Consider the product space $\prod_{n \in J} Y_n$ endowed with the product topology (the pointwise convergence topology). Then*

$$d(\mathbf{z}, \mathbf{v}) := \sum_{n \in J} \alpha_n d_n(\mathbf{z}(n), \mathbf{v}(n)) \quad (6.13)$$

is a finite metric on $\prod_{n \in J} Y_n$ which generates the pointwise convergence topology.

Proof. The defining properties of a metric for d follow easily from the corresponding properties for every d_n .

We denote by $\mathbf{y}$ an element of the product space and $\mathbf{y}(i)$ is called the i th coordinate. Recall that the product topology of $\prod_{n \in J} Y_n$ is the smallest topology containing the basic open sets $\langle E_n \rangle_{n \in S} := \{\mathbf{y} \in \prod_{n \in J} Y_n : \mathbf{y}(i) \in E_i, \forall i \in S\}$, where $S \subseteq J$ is finite and E_i is an open subset of Y_i (for every $i \in S$).

Suppose that $A \subseteq \prod_{n \in J} Y_n$ is an open set and $\mathbf{y} \in A$. Then, by definition of product topology, there exist a finite $S \subseteq J$ and a collection of open sets $\{E_i\}_{i \in S}$ such that $\langle E_i \rangle_{i \in S} \subseteq A$. Since $\mathbf{y}(i) \in E_i$ and E_i is open, then for every $i \in S$, there exists $r_i > 0$ such that $\mathbf{y}(i) \in B_n(\mathbf{y}(i), r_i) \subseteq E_i$. Define $\beta := \min\{\alpha_i r_i : i \in S\}$; it is easy to show that $B(\mathbf{y}, \beta) \subseteq \langle E_i \rangle_{i \in S}$. Indeed, if $\mathbf{z} \in B(\mathbf{y}, \beta)$, then $d(\mathbf{y}, \mathbf{z}) \leq \beta$ which implies $d_i(\mathbf{y}(i), \mathbf{z}(i)) \leq \alpha^{-1} \beta \leq r_i$ for all $i \in S$. Whence, $\mathbf{z} \in \langle E_i \rangle_{i \in S}$.

Conversely, consider $B(\mathbf{y}, r)$. We show that there exist a finite $S \subseteq J$ and a collection of open sets $\{E_i\}_{i \in S}$ such that $\langle E_i \rangle_{i \in S} \subseteq B(\mathbf{y}, r)$. Since $\sum_{n \in J} \alpha_n M_n < +\infty$, there exists a finite $S \subseteq J$ such that $\sum_{n \in J \setminus S} \alpha_n M_n < r/2$. Define $r_n := r/(2\alpha_n \#S)$ for every $n \in S$ where $\#S < +\infty$ is the cardinality of S . If $\mathbf{z}$ is such that $d_n(\mathbf{y}(n), \mathbf{z}(n)) \leq r_n$ for all $n \in S$. Then

$$\begin{aligned} d(\mathbf{y}, \mathbf{z}) &= \sum_{n \in S} d_n(\mathbf{y}(n), \mathbf{z}(n)) \alpha_n + \sum_{n \in J \setminus S} d_n(\mathbf{y}(n), \mathbf{z}(n)) \alpha_n \\ &\leq \sum_{n \in S} r_n \alpha_n + \sum_{n \in J \setminus S} M_n \alpha_n < r/2 + r/2 = r \end{aligned}$$

Whence, if $E_n := B_n(\mathbf{y}(n), r_n)$ for all $n \in S$, then $\langle E_i \rangle_{i \in S} \subseteq B(\mathbf{y}, r)$. □

The following lemma is elementary but we include it for the sake of completeness. It generalizes to metric spaces a well-known result on *total convergence* in normed space.

Lemma 6.3. *Let (Y, d) be a metric space. The space is complete if and only if every sequence $\{y_i\}_{i \in \mathbb{N}}$ such that $\sum_{i \in \mathbb{N}} d(y_i, y_{i+1}) < +\infty$ converges.*

Proof. Suppose that (Y, d) is complete. By using the triangle inequality, $d(y_n, y_m) \leq \sum_{i=n}^{m-1} d(y_i, y_{i+1})$ for all $n < m$, if $\sum_{i \in \mathbb{N}} d(y_i, y_{i+1}) < +\infty$, then $\{y_i\}_{i \in \mathbb{N}}$ is a Cauchy sequence, whence it is convergent.

Conversely, suppose that every sequence $\{y_i\}_{i \in \mathbb{N}}$ such that $\sum_{i \in \mathbb{N}} d(y_i, y_{i+1}) < +\infty$ converges. Let $\{y_i\}_{i \in \mathbb{N}}$ be a Cauchy sequence. Define $n_i := \min\{n \in \mathbb{N} : d(y_j, y_m) \leq 1/2^{i+1}, \forall j, m \geq n\}$. By construction $\sum_{i \in \mathbb{N}} d(y_{n_i}, y_{n_{i+1}}) \leq \sum_{i \in \mathbb{N}} 1/2^{i+1} = 1 < +\infty$, whence the subsequence $\{y_{n_i}\}_{i \in \mathbb{N}}$ converges to some $z \in Y$. Let $\varepsilon > 0$ and i_ε such that $1/2^{i_\varepsilon} \leq \varepsilon$. By continuity, $d(y_{n_{i_\varepsilon}}, z) \leq 1/2^{i_\varepsilon+1}$ and $d(y_{n_{i_\varepsilon}}, y_n) \leq 1/2^{i_\varepsilon+1}$ for every $n \geq n_{i_\varepsilon}$. Thus, for all $n \geq n_{i_\varepsilon}$, $d(y_n, z) \leq d(y_{n_{i_\varepsilon}}, z) + d(y_{n_{i_\varepsilon}}, y_n) \leq 1/2^{i_\varepsilon} \leq \varepsilon$ and this proves that the space is complete. □

Remark 6.4. It is known, see for instance [12], that if every Y_i is separable and the cardinality of J is at most $2^{\aleph_0}$, then $\prod_{n \in J} Y_n$ is separable. The converse is trivial.

Moreover, by using Lemma 6.3 it is easy to show that every finite metric space (Y_i, d_i) is complete if and only if $\prod_{n \in J} Y_n$ is complete with the distance (6.13). Indeed, suppose that every finite metric space (Y_i, d_i) is complete. Since $d(\mathbf{y}, \mathbf{z})/\alpha_i \geq d_i(\mathbf{y}(i), \mathbf{z}(i))$ for every $i \in J$, if $\sum_{n \in \mathbb{N}} d(\mathbf{y}_n, \mathbf{y}_{n+1}) < +\infty$, then $\sum_{n \in \mathbb{N}} d_i(\mathbf{y}_n(i), \mathbf{y}_{n+1}(i)) < +\infty$ for every $i \in J$; thus $d_i(\mathbf{y}_n(i), \mathbf{z}(i)) \rightarrow 0$ as $n \rightarrow +\infty$ for some $\mathbf{z}(i) \in Y_i$. Since the topology generated by d is the pointwise convergence topology (or by direct computation by using the Bounded Convergence Theorem) we have $d(\mathbf{y}_n, \mathbf{z}) \rightarrow 0$ as $n \rightarrow +\infty$ where $\mathbf{z}(i) := z(i)$ for all $i \in J$; whence $(\prod_{n \in J} Y_n, d)$ is complete. Conversely suppose that $(\prod_{n \in J} Y_n, d)$ is complete and fix $j \in J$. Fix also $\mathbf{z} \in \prod_{n \in J} Y_n$ and suppose that $\sum_{n \in \mathbb{N}} d_j(y_i, y_{i+1}) < +\infty$ where $\{y_i\}_{i \in \mathbb{N}}$ is a sequence in Y_j . For every fixed $i \in \mathbb{N}$, define $\mathbf{y}_i$ as $\mathbf{y}_i(n) := \mathbf{z}(n)$ for all $n \neq j$ and $\mathbf{y}_i(j) := y_i$. Then $\sum_{i \in \mathbb{N}} d(\mathbf{y}_i, \mathbf{y}_{i+1}) = \alpha_j \sum_{i \in \mathbb{N}} d_j(y_i, y_{i+1}) < +\infty$ whence $d(\mathbf{y}_i, \mathbf{w}) \rightarrow 0$ as $i \rightarrow +\infty$ for some $\mathbf{w} \in \prod_{n \in J} Y_n$ which implies $d_j(y_i, \mathbf{w}(j)) = d_j(\mathbf{y}_i(j), \mathbf{w}(j)) \rightarrow 0$ as $i \rightarrow +\infty$. This proves that (Y_j, d_j) is complete.

Thus, every finite metric space (Y_i, d_i) is Polish if and only if $\prod_{n \in J} Y_n$ is a Polish metric space with the distance defined by equation (6.13). This applies for instance to $\mathbb{R}^X$ endowed with the distance

$$d(\mathbf{z}, \mathbf{v}) := \sum_{n \in J} \frac{\min(|\mathbf{z}(n) - \mathbf{v}(n)|, 1)}{2^n}$$

where $\{x_i : i \in J\}$ is a (finite or infinite) enumeration of X and $J := \{1, \dots, \#X\}$. Whence $\mathbb{R}^X$ is a Polish metric space and the metric d generates the pointwise convergence topology.

Since $[0, 1]^X$ and $\mathbb{N}^X$ are closed subsets of $\mathbb{R}^X$, they are Polish metric spaces as well. In particular every measure μ_x , supported on $S_X \subseteq \mathbb{N}^X$, can be seen as a measure defined on $\mathbb{N}^X$ or $\mathbb{R}^X$. We note that $\mathbb{R}^X$ is a partially ordered Polish metric space, meaning that the set $\{(\mathbf{z}, \mathbf{v}) \in \mathbb{R}^X \times \mathbb{R}^X : \mathbf{z} \leq \mathbf{v}\}$ is a closed subset of $\mathbb{R}^X \times \mathbb{R}^X$.

ACKNOWLEDGEMENTS

The authors are grateful to Elisabetta Candellero for useful discussions. They acknowledge support by INdAM-GNAMP.

REFERENCES

- [1] D. Bertacchi, P. Braunsteins, S. Hautphenne, F. Zucca, Extinction probabilities in branching processes with countably many types: a general framework, *ALEA*, **19**, (2022), 311–338.
- [2] D. Bertacchi, F. Zucca, Critical behaviours and critical values of branching random walks on multigraphs, *J. Appl. Probab.* **45** (2008), 481–497.
- [3] D. Bertacchi, E. Candellero, F. Zucca, Martin boundaries and asymptotic behavior of branching random walks, (2023), <https://doi.org/10.48550/arXiv.2308.03711>.
- [4] D. Bertacchi, P. M. Rodriguez, F. Zucca, Galton-Watson processes in varying environment and accessibility percolation, *Braz. J. Probab. Stat.* **34** n. 3 (2020), 613–628.
- [5] D. Bertacchi, F. Zucca, Characterization of the critical values of branching random walks on weighted graphs through infinite-type branching processes, *J. Stat. Phys.* **134** n. 1 (2009), 53–65.
- [6] D. Bertacchi, F. Zucca, Recent results on branching random walks, *Statistical Mechanics and Random Walks: Principles, Processes and Applications*, Nova Science Publishers (2012), 289–340.
- [7] D. Bertacchi, F. Zucca, Strong local survival of branching random walks is not monotone, *Adv. Appl. Probab.* **46** n.2 (2014), 400–421.
- [8] D. Bertacchi, F. Zucca, A generating function approach to branching random walks, *Braz. J. Probab. Stat.* **31** n.2 (2017), 229–253.
- [9] D. Bertacchi and F. Zucca, Branching random walks with uncountably many extinction probability vectors, *Braz. J. Probab. Stat.* **34**(2), 426–438, 2020.
- [10] P. Braunsteins, S. Hautphenne, The probabilities of extinction in a branching random walk on a strip, *J. Appl. Probab.* **57** . 3 (2020), 811–831.
- [11] D.L. Chung, *A course in probability theory*, 3rd. edition, Academic Press, 2001.
- [12] E. Hewitt, A remark on density characters, *Bull. Amer. Math. Soc.* **52** n. 8 (1946), 641–643.

- [13] T. Hutchcroft, Transience and recurrence of sets for branching random walk via non-standard stochastic orders, *Ann. Inst. H. Poincaré Probab. Statist.*, **58** n. 2 (2022), 1041–1051.
- [14] T. Kamae, U. Krengel and G. L. O’Brien, Stochastic Inequalities on Partially Ordered Spaces, *Ann. Appl. Probab.* **5** n. 6 (1977), 899–912.
- [15] T. Liggett, *Interacting particle systems*. Springer-Verlag, Berlin, 2005.
- [16] J. E. Moyal, Multiplicative population chains, *Proceedings of the Royal Society of London. Series A, Mathematical and Physical Sciences*, **266** (1962), 518–526.
- [17] W. Rudin, *Real and Complex Analysis*, 3rd Edition, McGraw-Hill, New York, 1987.
- [18] D. Williams, *Probability with martingales*, Cambridge Univ. Press, 1991.
- [19] F. Zucca, Survival, extinction and approximation of discrete-time branching random walks, *J. Stat. Phys.*, **142** n.4 (2011), 726–753.

D. BERTACCHI, DIPARTIMENTO DI MATEMATICA E APPLICAZIONI, UNIVERSITÀ DI MILANO–BICOCCA, VIA COZZI 53, 20125 MILANO, ITALY.

Email address: `daniela.bertacchi@unimib.it`

F. ZUCCA, DIPARTIMENTO DI MATEMATICA, POLITECNICO DI MILANO, PIAZZA LEONARDO DA VINCI 32, 20133 MILANO, ITALY.

Email address: `fabio.zucca@polimi.it`