

On fresh sets in iterations of Prikry type forcing notions

Moti Gitik* and Eyal Kaplan

March 5, 2024

Abstract

We examine the existence (and mostly non-existence) of fresh sets in commonly used iterations of Prikry type forcing notions. Results of [4] are generalized. As an application, a question of a referee of [9] is answered. In addition stationary sets preservation is addressed.

1 Introduction

We continue here the series of papers [4], [5],[8], [6] on iterations of Prikry type forcing notions and concentrate on existence or non-existence of fresh subsets.

Our framework throughout the paper will be as follows:

1. GCH,
2. κ is an inaccessible cardinal,
3. $\langle P_\alpha, \mathcal{Q}_\beta \mid \alpha \leq \kappa, \beta < \kappa \rangle$ be either Magidor (full) or non-stationary or Easton support iterations of Prikry type forcing notions.
4. For every $p \in P_\kappa$ and $\gamma < \kappa$, let $p(\gamma)$ denotes the γ -th coordinate of p , i.e. $p = \langle \mathop{\mathcal{P}}\nolimits(p(\gamma) \mid \gamma < \kappa) \rangle$. $\text{supp}(p) = \{\gamma < \kappa \mid p \restriction \gamma \Vdash \mathop{\mathcal{P}}\nolimits(p(\gamma)) \neq 0_{\mathcal{Q}_\gamma}\}$.
5. For every $\beta < \kappa$, $\langle \mathcal{Q}_\beta, \leq_{\mathcal{Q}_\beta}^* \rangle$ is forced to be $|\beta|$ -closed.
Note that for a singular β this implies $|\beta|^+$ -closure.
6. For every $\beta < \kappa$, $|\mathcal{Q}_\beta| < \kappa$.

*We are grateful to participants of TAU Set Theory seminar and in particular to Menachem Magidor and to Omer Ben-Neria for their comments and remarks. The work of the first author was partially supported by ISF grant No. 882/22.

7. If the Easton support is used, then for every $p \in P_\kappa$ and every inaccessible $\alpha \leq \kappa$, $\alpha > |\text{supp}(p) \cap \alpha|$, provided that for every $\beta < \alpha$, $|P_\beta| < \alpha$.
8. If the non-stationary support is used, then for every $p \in P_\kappa$ and every inaccessible $\alpha \leq \kappa$, $\text{supp}(p) \cap \alpha$ is non-stationary in α , provided that for every $\beta < \alpha$, $|P_\beta| < \alpha$.

Definition 1.1 (J. Hamkins) A set A of ordinals is called a *fresh set* if all its initial segments are in the ground model, but A is not there.

Hamkins [7] proved that if a forcing has a gap at κ ,¹ then no fresh subsets of κ are added. We proved in [4] results on non-existence of fresh sets in iterations of certain Prikry type forcing notions P_κ , as above. Here we would like to generalize this results.

In addition, results regarding preservation of stationary sets when forcing with each of the supports.

The paper is organized as follows. In Section 2 we deal with preservations of stationarity. Later sections address fresh subsets. Section 3 deal with non-stationary support iterations. Sections 4 and 5 address Easton and full supports respectively.

2 On preservation of stationarity in Prikry type extensions

Let κ be an inaccessible cardinal and let S be a stationary subset of κ . Suppose that $\langle P_\alpha, \mathcal{Q}_\beta \mid \alpha \leq \kappa, \beta < \kappa \rangle$ is an Easton or a full (Magidor) or a non-stationary support iteration of Prikry type forcing notions as in the introduction. Let G_κ be a generic subset of P_κ .

We would like to address here a question *whether S remains stationary in $V[G_\kappa]$* .

- Remark 2.1**
1. Note that we can assume that κ is a regular cardinal in V , since otherwise it is possible to replace it by $\text{cof}(\kappa)$.
 2. Our main interest will be in situations where κ is an inaccessible cardinal in $V[G]$.
 3. If κ is a Mahlo cardinal, then Easton support iterations satisfy κ -c.c. and so preserve stationary subsets of κ .

¹ P has a gap at κ , if $P = R * \mathcal{Q}$ such that $|R| < \kappa$ and $\mathcal{Q}$ is $\kappa + 1$ -strategically closed.

4. If a full support iteration of Prikry forcings is used, then the set of former measurables which changed their cofinality will be non-stationary, as witnessed by the regressive function which maps each one of them to the first element in its Prikry sequence.

Theorem 2.2 *Assume GCH. Let κ be an inaccessible cardinal. Let $P = P_\kappa$ be as in the introduction. Let $G \subseteq P_\kappa$ be generic over V . Then κ is inaccessible in $V[G]$, and–*

1. *If $S \subseteq \kappa$ is stationary in V and consists of singulars, then S remains stationary in $V[G]$.*
2. *If the Easton or nonstationary support is used, then P_κ preserves stationary subsets of κ .*
3. *If the full support is used, $S \subseteq \kappa$ is stationary in V , and–*
 - (a) *For every $\alpha \in S$, $\langle Q_\alpha, \leq^* \rangle$ is $|\alpha|^+$ -complete.*
 - (b) *For every $\alpha < \kappa$, Q_α has the property that for every $p, q, r \in Q_\alpha$, if $p, q \geq^* r$, then there is $t \in Q_\alpha$ such that $t \geq^* p, q$.*

Then S remains stationary in $V[G]$.

The inaccessibility of κ in $V[G]$ is proved in 2.6. Point 1 of the Theorem is proved in 2.7. Point 2 of the Theorem is proved in Theorems 2.9 and 2.11. Point 3 of the Theorem is proved in 2.10.

Let us start with the following lemma which is a weak form of a strong Prikry condition:

Lemma 2.3 *Let $D \subseteq P_\kappa$ be a dense open and let $p \in P_\kappa$. Then there are $\alpha < \kappa$ and $q \geq^* p$ such that for every $r \in P_\alpha$, $r \geq_{P_\alpha} q \restriction \alpha$ there is $r' \geq_{P_\alpha} r$ such that $r' \frown q \restriction \alpha \in D$.*

Proof. Let D and $p = \langle p_\gamma \mid \gamma < \kappa \rangle$ be as in the statement of the lemma.

We prove the lemma for the full support iteration. The arguments for the Easton and for the non-stationary support iterations are very similar, only coordinates in supports should be considered.

Suppose that the conclusion of the lemma fails.

We define by recursion, for each $\beta < \kappa$, a condition–

$$p(\beta) = \langle p_\gamma^* \mid \gamma < \beta \rangle \frown \langle p_\gamma \mid \beta \leq \gamma < \kappa \rangle$$

so that $p(\beta) \restriction \beta \Vdash \neg \sigma_\beta$, where

$$\sigma_\beta \equiv \exists t \in P_\kappa \setminus \beta, t \geq^* p \restriction \beta \exists r \in G_\beta \quad r \restriction t \in D.$$

For the first stage, we have—

$$\sigma_1 \equiv \exists t \in P_\kappa \setminus 1, t \geq^* p \restriction 1 \exists r \in G_1 \quad r \restriction t \in D.$$

Q_0 satisfies the Prikry condition, so there is $p_0^* \geq^* p_0$ which decides σ_1 . If $p_0^* \Vdash \sigma_1$, then $p_0^* \restriction \underline{t}$ will be as desired. So, assume that $p_0^* \Vdash \neg \sigma_1$, and continue.

The successor step is similar to the first stage above.

For limit steps, suppose that β is a limit ordinal. Let us show that—

$$p(\beta) = \langle \underline{p}_\gamma^* \mid \gamma < \beta \rangle \restriction \langle \underline{p}_\gamma \mid \beta \leq \gamma < \kappa \rangle$$

is as desired, i.e. $p(\beta) \restriction \beta \Vdash \neg \sigma_\beta$. Suppose otherwise, then there is $r = \langle \underline{r}_\gamma \mid \gamma < \beta \rangle \in P_\beta$ such that $r \geq p(\beta) \restriction \beta$ and $r \Vdash \sigma_\beta$. Extend it, if necessary, so that for some $\underline{t}$

$$r \restriction \underline{t} \geq^* p \restriction \beta \text{ and } r \restriction \underline{t} \in D.$$

By the definition of order on $\mathcal{P}_\beta$, there is $\beta^* < \beta$ such that for every γ , $\beta^* \leq \gamma < \beta$,

$$r \restriction \gamma \restriction \underline{r}_\gamma \geq_\gamma^* \underline{p}_\gamma^*.$$

Consider a P_{β^*} -name

$$\underline{t}' = \langle \underline{r}_\gamma \mid \beta^* \leq \gamma < \beta \rangle \restriction \underline{t}.$$

Then

$$r \restriction \beta^* \restriction \underline{t}' \geq^* p \restriction \beta^* \text{ and } r \restriction \beta^* \restriction \underline{t}' \in D.$$

But $r \restriction \beta^* \geq p(\beta^*) \restriction \beta^* \Vdash \neg \sigma_{\beta^*}$. Contradiction.

This completes the construction.

Consider $p(\alpha) = \langle \underline{p}_\gamma^* \mid \gamma < \kappa \rangle$. Pick some $r \geq p(\kappa)$ in D . Now we obtain a contradiction as in the limit stage above.

□

Assuming that κ is measurable in the ground model, a stronger version of Lemma 2.3 can be proved:

Lemma 2.4 *Assume that κ is measurable in V and U is a normal measure on κ . Let $D \subseteq P_\kappa$ be a dense open and let $p \in P_\kappa$. Then there are $\alpha < \kappa$ and $q \geq^* p$, such that $q \restriction \alpha = p \restriction \alpha$, and for every $r \in P_\alpha, r \geq_{P_\alpha} p \restriction \alpha$ there is $r' \geq_{P_\alpha} r$ such that $r' \restriction q \restriction \alpha \in D$. Moreover, for every $X \in U$, α, q above can be chosen such that $\alpha \in X$.*

Proof. Let σ_β be as in lemma 2.3. If there exists $\beta \in X$ such that $p \restriction \beta \Vdash \sigma_\beta$, we are done.

Assume otherwise. For every $\beta \in X$ there exists $r_\beta \geq p \restriction \beta$ such that $r_\beta \Vdash \neg \sigma_\beta$. For each such limit β , there exists $\beta' < \beta$ such that $r_\beta \restriction \beta' \Vdash r_\beta \restriction \beta' \geq^* p \restriction [\beta', \beta)$. The function $\beta \mapsto \beta'$ is regressive, and thus there exist a set $A \in U$ and $\beta^* < \kappa$ such that for every $\beta \in A$, $r_\beta \restriction \beta^* \Vdash r_\beta \restriction \beta^* \geq^* p \restriction [\beta^*, \beta)$. Since $|P_{\beta^*}| < \kappa$, we can shrink $A \in U$ further, and assume that there exists $r^* \in P_{\beta^*}$, such that, for every $\beta \in A$, $r_\beta \restriction \beta^* = r^*$.

Then r^* has the following property: for every $\beta \in A$ there exists $s(\beta) \geq^* p \restriction [\beta^*, \beta)$ such that $r^* \restriction s(\beta) \Vdash \neg \sigma_\beta$. Now apply ineffability: we can find $A^* \subseteq A$, $A^* \in U$, and a P_{β^*} -name for a condition $s^* = [\beta \mapsto s(\beta)]_U \in P \restriction \beta^*$, such that $r^* \Vdash s^* \geq^* p \restriction \beta^*$, and, for every $\beta \in A^*$,

$$r^* \restriction s^* \restriction [\beta^*, \beta) = r^* \restriction s(\beta) \Vdash \neg \sigma_\beta.$$

Finally, pick some $q \geq r^* \restriction s^*$, $q \in D$. Let $\beta \in A^* \setminus \beta^* + 1$ be such that—

$$q \restriction \beta \Vdash q \restriction \beta \geq^* s^* \restriction \beta$$

and in particular,

$$q \restriction \beta \Vdash q \restriction \beta \geq^* p \restriction \beta, \text{ and } q \restriction \beta \restriction q \restriction \beta \in D$$

and thus $q \restriction \beta \Vdash \sigma_\beta$; however, since $\beta \in A^*$,

$$q \restriction \beta \geq r^* \restriction s^* \restriction [\beta^*, \beta) \Vdash \neg \sigma_\beta$$

which is a contradiction.

□

Lemma 2.5 *Let $p \in P_\kappa$ be a condition, and assume that ζ a P_κ -name for an ordinal. Then there exists $q \geq^* p$ and a set of ordinals $A \in V$ with $|A| < \kappa$ such that $q \Vdash \zeta \in A$.*

Proof. Apply lemma 2.3 on the dense open set D of conditions which decide the value of ζ . Then there exists $q \geq^* p$ and $\alpha < \kappa$ such that, for every $r \geq q \restriction \alpha$ there exists $r' \geq r$ such that $r' \restriction q \restriction \alpha \parallel \zeta$. Let $A \in V$ be the set of all possible values of ζ as decided by some extension of q . We argue that $|A| < \kappa$.

Assume that $q' \geq q$ decides the value of ζ . Denote $r = q' \restriction \alpha$. Then there exists $r' \geq r$ such that $r' \restriction q \restriction \alpha \parallel \zeta$. So both conditions q' , $r' \restriction q \restriction \alpha$ decide the value of ζ ; but those conditions are compatible, since $r' \geq p' \restriction \alpha$, and $q' \restriction \alpha \geq q \restriction \alpha$. This shows that every element of A can be realized as the decided value of ζ by a condition of the form $r \restriction q \restriction \alpha$ for some $r \in P_\alpha$. But the cardinality of the set of such conditions is strictly below κ , since $|P_\alpha| < \kappa$, by the assumption on the cardinality of the forcings Q_β for $\beta < \kappa$.

□

Corollary 2.6 *Let $G_\kappa \subseteq P_\kappa$. Then κ remains inaccessible in $V[G_\kappa]$.*

Proof. We concentrate on the proof that κ remains a regular cardinal after forcing with P_κ , since it's routine to verify that it remains strong limit.

Assume that $\check{f}$ is a P_κ -name for a function from some ordinal $\tau < \kappa$ to κ . Let $p \in P_\kappa$ be a condition which forces this. We argue that there exists $p^* \geq p$ and some $\mu^* < \kappa$, such that $p^* \Vdash \text{rng}(\check{f}) \subseteq \mu^*$.

Let $G_{\tau+1} \subseteq P_{\tau+1}$ be an arbitrary generic extension containing $p \restriction \tau + 1$. We prove that, in $V[G_{\tau+1}]$, there exist $q \geq^* p \restriction \tau + 1$ and $\mu < \kappa$ such that $q \Vdash \text{rng}(\check{f}) \subseteq \mu$. Once we prove that, we are done: let $\check{q}, \check{\mu}$ be $P_{\tau+1}$ -names which are forced by $p \restriction \tau + 1$ to have the above properties. Let $\mu^* < \kappa$ be an upper bound on the set of possible values of $\check{\mu}$, as forced by extensions of $p \restriction \tau + 1$. Since $|P_{\tau+1}| < \kappa$, this set is bounded in κ , and thus there exists such an upper bound below κ . Then $p^* = p \restriction \tau + 1 \check{\cap} \check{q} \Vdash \text{rng}(\check{f}) \subseteq \mu^*$, as desired.

Work in $V[G_{\tau+1}]$. Apply lemma 2.5 over and over to construct a $\leq^*$ -increasing sequence of conditions $\langle p_\xi : \xi \leq \tau \rangle$ in $P_\kappa \setminus \tau + 1$, such that, for each $\xi < \tau$ there exists some $\mu_\xi < \kappa$ such that $p_\xi \Vdash \check{f}(\xi) < \mu_\xi$. Note that in limit steps (including the last step) we may take upper bound, since the direct extension order of $P_\kappa \setminus \tau + 1$ is more than τ -closed. Finally, $q = p_\tau$ forces that the image of $\check{f}$ is bounded by $\mu = \bigcup_{\xi < \tau} \mu_\xi < \kappa$.

□

Theorem 2.7 *Let $S \subseteq \kappa$ be a stationary set consisting of singulars. Let $G_\kappa \subseteq P_\kappa$. Then S remains stationary in $V[G_\kappa]$.*

Proof. Let $C \subseteq \kappa$ be a club in $V[G]$. Let $p \in G$ be condition which forces this.

Work in V . Pick an elementary submodel $M \preceq H_\chi$ such that:

1. $|M| = \delta < \kappa$,
2. $M \cap \kappa = \delta$,
3. $\delta \in S$ (in particular, δ is singular),
4. ${}^{\text{cof}(\delta)}M \subseteq M$,
5. $\kappa, P_\kappa, S, \check{C}, p \in M$.

Pick a cofinal sequence in δ , $\langle \delta_i \mid i < \text{cof}(\delta) \rangle$.

Apply lemma 2.5. Construct (in V) a $\leq^*$ -increasing sequence of conditions in the forcing $P_\kappa \setminus \text{cof}(\delta)$, $\langle p_\xi : \xi \leq \text{cof}(\delta) \rangle$, such that each condition p_ξ belongs to M .

We first construct $p_0 \geq^* p$ in M , such that $p_0 \restriction \text{cof}(\delta) + 1 = p \restriction \text{cof}(\delta) + 1$, and, using Lemma 2.5,

$$p_0 \restriction \text{cof}(\delta) + 1 \Vdash \exists \alpha < \kappa, p_0 \restriction \text{cof}(\delta) + 1 \Vdash \min(\mathcal{C} \setminus \delta_0) < \alpha.$$

Let α_0 be the least upper bound of the set of all possible values for the $P_{\text{cof}(\delta)+1}$ -name α . Then $p_0 \Vdash \min(\mathcal{C} \setminus \delta_0) < \alpha_0$. Note that by elementarity, $\alpha_0 < \delta$.

Assuming that $i < \text{cof}(\delta)$ and p_i, α_i have been defined and both are in M , and let us define p_{i+1}, α_{i+1} . Let $p_{i+1} \geq^* p_i$ in M be such that $p_{i+1} \restriction \text{cof}(\delta) + 1 = p \restriction \text{cof}(\delta) + 1$, and there exists $\alpha_{i+1} < \kappa$, $p_{i+1} \Vdash \min(\mathcal{C} \setminus \max\{\delta_{i+1}, \alpha_i\}) < \alpha_{i+1}$. Take α_{i+1} to be minimal with this property. Then $\alpha_{i+1} < \delta$.

For the limit step, assume that $j < \text{cof}(\delta)$ is limit and $\langle p_i : i < j \rangle, \langle \alpha_i : i < j \rangle$ have been constructed. Use the fact that M is closed under $< \text{cof}(\delta)$ -sequences to find an upper bound $q \in M$ of $\langle p_i : i < j \rangle$, such that $q \restriction \text{cof}(\delta) + 1 = p \restriction \text{cof}(\delta) + 1$. We used here the fact that $P_\kappa \setminus \text{cof}(\delta) + 1$ is $\text{cof}(\delta)$ -closed. Finally, let $p_j \geq^* q$ be chosen in M such that $p_j \restriction \text{cof}(\delta) + 1 = p \restriction \text{cof}(\delta) + 1$ and, for some $\alpha_j < \kappa$, $p_j \Vdash \min(\mathcal{C} \setminus \max\{\delta_j, \sup\{\alpha_i : i < j\}\}) < \alpha_j$. We used here again the fact that M is closed under $< \text{cof}(\delta)$ -sequences and thus $\langle \alpha_i : i < j \rangle \in N$. Let α_j be the minimal with the above property. Then $\alpha_j \in M$.

This concludes the inductive construction. In the final limit step, take, in V , an upper bound q^* for all the conditions $\langle p_i : i < \text{cof}(\delta) \rangle$. Then $q^* \Vdash \delta = \bigcup_{i < \text{cof}(\delta)} \alpha_i \in \mathcal{C}$, since C is forced by p to be closed. Thus $\delta \in C \cap S$, as desired.

□

Remark 2.8 Basically the same argument works for κ replaced by κ^+ .

Now let us try to extend the theorem to S 's which consists of regular cardinals. We deal with three supports separately.

First point out the following:

Theorem 2.9 *Suppose that the Easton support is used in P_κ . Assume that $S \subseteq \kappa$ is stationary. Then S remains stationary in $V[G]$.*

Proof. If κ is a Mahlo cardinal, the Easton support iteration is κ -c.c. and thus preserves stationary subsets of κ . Thus, we can assume that κ is not Mahlo. In this case, there exists a club $C \subseteq \kappa$ of singular cardinals. Thus, by shrinking S we can assume that it consists of

singulars. Then, Theorem 2.7 applies.

□

Turn to the full support.

Theorem 2.10 *Suppose that P_κ is the full support iteration.*

Assume that for every $\beta < \kappa$, for every $p, q, r \in \mathcal{Q}_\alpha$, if $p, q \geq^ r$, then there is $t \in \mathcal{Q}_\alpha$ such that $t \geq^* p, q$.*

Let $S \subseteq \kappa$ be a stationary such that, for every $\alpha \in S$, $\mathcal{Q}_\alpha$ is $|\alpha|^+$ -complete.

Then S remains stationary in $V[G]$.

Proof. Suppose otherwise. Pick some $p \in P_\kappa$ and a name $\mathcal{C}$ such that

$$p \Vdash \mathcal{C} \text{ is a club in } \kappa \text{ and } \mathcal{C} \cap S = \emptyset.$$

Pick now $M \preceq H_\chi$ and $\delta \in S$, as in Theorem 2.7.

It is enough to find a condition $r \geq p$ which forces “ $\mathcal{C}$ is unbounded in δ ” in order to derive a contradiction.

Suppose that there is no such r . Let $G_\delta \subseteq P_\delta$ be a generic with $p \restriction \delta \in G_\delta$. Then, in $V[G_\delta]$,

$$p \restriction \delta \Vdash \mathcal{C}_{G_\delta} \cap \delta \text{ is bounded in } \delta.$$

We have $\delta \in S$, so, by the assumption of the theorem, $\mathcal{Q}_\delta$ is δ^+ -complete, and hence, P_κ/G_δ is δ^+ -complete. Then there are $p' \in P_\kappa/G_\delta, p' \geq^* p \restriction \delta$ and $\rho < \delta$ such that

$$p' \Vdash \mathcal{C}_{G_\delta} \cap \delta \subseteq \rho.$$

Pick some $t \in G_\delta, t \geq p \restriction \delta$ such that

$$t \restriction p' \Vdash \mathcal{C}_{G_\delta} \cap \delta \subseteq \rho.$$

Since $t \geq p \restriction \delta$, there exists $\gamma < \delta$ such that $t = t \restriction \gamma \smallfrown t \restriction \gamma \smallfrown t \restriction \gamma$ and $t \restriction \gamma \geq^* p \restriction \gamma$. In particular, $t \restriction \gamma \Vdash t \restriction \gamma \smallfrown p' \geq^* p \restriction \gamma$.

Now work in M above the condition p . Note that γ, ρ above are below δ and thus are in M .

Also $p, \mathcal{C} \in M$, and recall that

$$p \Vdash \mathcal{C} \text{ is a club in } \kappa$$

Let $\zeta \in M$ be a P_κ -name such that $p \Vdash \zeta = \min(\mathcal{C} \setminus \check{\rho} + 1)$. By Lemma 2.5, we can find (in M) a P_γ -name for a condition $q_1 \geq^* p \restriction \gamma$ such that $p \restriction \gamma \Vdash \exists \mu < \kappa \ q_1 \Vdash \zeta < \mu$. Let $\mu \in M$ be a P_γ -name such that $p \restriction \gamma \Vdash q_1 \Vdash \zeta < \mu$. In M , let μ^* be the supremum of all

possible values of μ , as forced by extensions of $p \restriction \gamma$. Then $\mu^* < \delta$ (since $\mu^* \in M \cap \kappa$). By elementarity, $p \restriction \gamma \widehat{\smile} q_1 \Vdash \zeta < \mu^*$ holds in V as well.

Finally, let $p^* \in P_\kappa$ be a condition such that $q^* \restriction \gamma = t \restriction \gamma$ and $p^* \setminus \gamma$ direct extends both the conditions $t \setminus \gamma \widehat{\smile} p'$, q_1 . Note that, since $t \setminus \gamma \widehat{\smile} p'$ and q_1 direct extend $p \setminus \gamma$, the extra assumption of the theorem allows us to construct such p^* .

Then on the one hand, p^* forces that $\mathcal{C} \cap \delta \subseteq \rho$; on the other hand, it forces that $\min(\mathcal{C} \setminus \rho + 1) \leq \mu^* < \delta$. A contradiction.

□

Turn now to the non-stationary support.

Theorem 2.11 *Assume that the non-stationary support is used. Let $S \subseteq \kappa$ be stationary. Then S is stationary in $V[G_\kappa]$.*

Proof. Let $\mathcal{C}$ be a P_κ -name and $p \in P_\kappa$, $p \Vdash \mathcal{C}$ is a club disjoint to S .

We construct:

1. $\leq^*$ -increasing sequence of conditions $\langle p_i : i < \kappa \rangle$,
2. decreasing sequence of clubs $\langle C_i : i < \kappa \rangle$, each is disjoint to the support of p_i ,
3. increasing continuous sequences of ordinals $\langle \nu_i : i < \kappa \rangle$, $\langle \alpha_i : i < \kappa \rangle$.

The sequence of conditions will be a fusion sequence, in the sense that, for every $i < j$, $p_i \leq^* p_j$ and $p_j \restriction \nu_i = p_i \restriction \nu_i$.

Let $p_0 = p$ and $\alpha_0 = 0$. Pick a club C_0 disjoint to $\text{supp}(p_0)$. Let $\nu_0 = \min(C_0)$.

Let $G_{\nu_0} \subseteq P_{\nu_0}$ be a generic with $p_0 \restriction \nu_0 \in G_{\nu_0}$. Apply Lemma 2.5, inside $V[G_{\nu_0}]$, and find $p'_1 \in P_\kappa/G_{\nu_0}$, $p'_1 \geq^* p_0 \setminus \nu_0$ and $A'_1 \subseteq \kappa$, $|A'_1| < \kappa$ such that $p'_1 \Vdash \min(\mathcal{C}) \in A'_1$.

Now back to V , we have $|P_{\nu_0}| < \kappa$, hence there is A_1 , $|A_1| < \kappa$ such that

$$p_1 = p_0 \restriction \nu_0 \widehat{\smile} p'_1 \Vdash \min(\mathcal{C}) \in A_1.$$

Set $\alpha_1 = \sup(A_1)$.

Next, we pick a club $C_1 \subseteq C_0$ disjoint to $\text{supp}(p_1)$. Let $\nu_1 = \min(C_1) \setminus \max(\nu_0, \alpha_1)$.

Let $G_{\nu_1} \subseteq P_{\nu_1}$ be a generic with $p_1 \restriction \nu_1 \in G_{\nu_1}$. Apply Lemma 2.5, inside $V[G_{\nu_1}]$, and find $p'_2 \in P_\kappa/G_{\nu_1}$, $p'_2 \geq^* p_1 \setminus \nu_1$ and $A'_2 \subseteq \kappa$, $|A'_2| < \kappa$ such that $p'_2 \Vdash \min(\mathcal{C} \setminus \nu_1 + 1) \in A'_1$.

Now back to V , we have $|P_{\nu_1}| < \kappa$, hence there is A_2 , $|A_2| < \kappa$ such that

$$p_2 = p_1 \restriction \nu_1 \widehat{\smile} p'_2 \Vdash \min(\mathcal{C} \setminus \nu_1 + 1) \in A_2.$$

Set $\alpha_2 = \sup(A_2)$.

We do the same at each successor stage $i < \kappa$ of the construction.

Suppose now that $i < \kappa$ is limit. Set $\alpha_i = \nu_i = \bigcup_{j < i} \nu_j = \bigcup_{j < i} \alpha_j$ and let p_i be the coordinateswise union of $p_j, j < i$. Let $C_i = \bigcap_{j < i} C_j$. Pick $\nu_{i+1} = \min(C_i) \setminus \nu_i + 1$. Continue as above.

Finally, let p_κ be the coordinateswise union of $p_i, i < \kappa$. It is in P_κ since for every $i < j$, $p_i \leq^* p_j$ and $p_j \upharpoonright \nu_i = p_i \upharpoonright \nu_i$ and $\Delta_{i < \kappa} C_i$ is disjoint to its support.

Now,

$$p_\kappa \Vdash \{ \alpha_i \mid i \text{ limit} \} \subseteq \mathcal{C}.$$

But $\{ \alpha_i \mid i \text{ limit} \}$ is club in V , and so, $S \cap \{ \alpha_i \mid i \text{ limit} \} \neq \emptyset$. Contradiction.

□

A similar argument can be using elementary submodel shows preservation of stationary subsets of κ^+ with non-stationary support iterations.

Note $|P_\kappa| = \kappa$, in the Easton support case, and so, κ^+ is preserved. By [1], κ^+ is preserved in non-stationary support iterations. Let us consider the full support iterations. Note that the Magidor iteration of Prikry forcing notions [10] satisfies $\kappa^+ - \text{c.c.}$. A bit more general:

Proposition 2.12 *Suppose that for every $\beta < \kappa$, for every $s, t, r \in \mathcal{Q}_\beta$, if $s \leq_{\mathcal{Q}_\beta}^* t, r$, then there is $e \in \mathcal{Q}_\beta, e \geq_{\mathcal{Q}_\beta}^* t, r$. Then P_κ satisfies $\kappa^+ - \text{c.c.}$*

In general it turns out that κ^+ may be collapsed with full support iteration.

Proposition 2.13 *Suppose that κ is a measurable cardinal. Let P_κ be the full support iteration of $\text{Col}(\alpha, \alpha^+) = \{ f \mid f \in {}^\xi \alpha^+, \xi < \alpha \}$, for every regular $\alpha < \kappa$. Then κ^+ is collapsed in V^{P_κ} .*

Proof. Let U be a normal measure over κ .

We start with the following claim:

Claim 1 Let $p = \langle p(\beta) \mid \beta < \kappa \rangle \in P_\kappa$. Then there are $A^* \in U$, $\tau^* < \kappa$ and $p^* \geq p$ such that $p^* \upharpoonright \beta \Vdash \text{dom}(\check{p}^*(\beta)) = \tau^*$, for every $\beta \in A^*$.

Proof. For every regular $\beta < \kappa$ there are $s_\beta \in P_\beta, s_\beta \geq p \upharpoonright \beta$ and $\tau_\beta^0 < \beta$ such that $s_\beta \Vdash \text{dom}(\check{p}(\beta)) = \tau_\beta^0$.

Find $A'_0 \in U$ and $p_0 \in P_\kappa$ such that $p_0 \upharpoonright \beta = s_\beta$, for every $\beta \in A'_0$. For example, take $p_0 = [\beta \mapsto s_\beta]_U$.

Then we consider a regressive function $\beta \mapsto \tau_\beta^0$ on A'_0 . Find $A_0 \subseteq A'_0, A_0 \in U$ and $\tau^0 < \kappa$

such that $\tau_\beta^0 = \tau^0$, for every $\beta \in A_0$.

Repeat the process with p_0 replacing p and find $A_1 \subseteq A_0, A_1 \in U, p_1$ and τ^1 such that $p_1 \restriction \beta \Vdash \text{dom}(p_0(\beta)) = \tau^1$.

Continue by induction. Let $A^* = \bigcap_{n < \omega} A_n$ and p^* be the coordinatewise union of p_n 's. Set $\tau^* = \bigcup_{n < \omega} \tau^n$.

Then $p^* \restriction \beta \Vdash \text{dom}(p^*(\beta)) = \tau^*$, for every $\beta \in A^*$, will be as desired.

□ of the claim.

For every $\tau < \kappa$, define a maximal antichain A_τ in P_κ . Proceed as follows.

Let us pick functions $\langle h_\gamma \mid \gamma < \kappa^+ \rangle$ such that $\text{dom}(h_\gamma) \in U$, for every $\alpha \in \text{dom}(h_\gamma), h_\gamma(\alpha) < \alpha^+$ and $[h_\gamma]_U = \gamma$, for example κ^+ -canonical functions will do the job.

Fix $\tau < \kappa$. Let A_τ be a maximal antichain in P_κ of cardinality κ^+ which consists of $p \in P_\kappa$ such that:

(*) if for some $B \in U$ and $\gamma < \kappa^+$, the condition $t_{B\gamma} = \langle t_{B\gamma}(\alpha) \mid \alpha < \kappa \rangle$ is compatible with p , then, for some $B' \in U, p \geq t_{B'\gamma}$,

where, for $E \in U, t_{E\gamma}(\alpha) = 0_\alpha$, unless $\alpha \in E \cap \text{dom}(h_\gamma) \setminus \tau + 1$, and if $\alpha \in E \cap \text{dom}(h_\gamma) \setminus \tau + 1$, then $t_{E\gamma}(\alpha) = \{(\tau, h_\gamma(\alpha))\}$, i.e. the value of the generic function for α at τ is $h_\gamma(\alpha)$.

Let $\langle p_i^\tau \mid i < \kappa^+ \rangle$ be an enumeration of A_τ .

Let $G \subseteq P_\kappa$ be a generic. Define $F : \kappa \rightarrow (\kappa^+)^V$ by setting $F(\tau) = i$ iff $p_i^\tau \in G$.

We claim that such F is onto.

Suppose otherwise. Pick some $p \in G$ and $\eta < \kappa^+$ such that $p \Vdash \text{rng}(F) \subseteq \eta$.

Apply Claim 1. Let $A^* \in U, \tau^* < \kappa$ and $p^* \geq p$ be as in the conclusion of the claim.

Now for every $\gamma < \kappa^+$, we can extend p^* to a condition p^γ by adding a pair $(\tau^*, h_\gamma(\alpha))$ to $p^*(\alpha)$, for every $\alpha \in A^* \cap \text{dom}(h_\gamma) \setminus \gamma + 1$. Note that if $\gamma \neq \gamma'$, then $p^\gamma, p^{\gamma'}$ are incompatible. So, the set $\{p^\gamma \mid \gamma < \kappa^+\}$ consists of κ^+ -many incompatible conditions.

Then, each of p^γ 's must be compatible with a member of A_{τ^*} with index below η . Hence, there is $i^* < \eta$ such that $p_{i^*}^{\tau^*}$ is compatible with κ^+ -many p^γ 's. Pick two of them $\gamma \neq \gamma'$.

By (*), then $p_{i^*}^{\tau^*} \geq t_{B\gamma}, t_{B'\gamma'}$, for some $B, B' \in U$. However, $\gamma \neq \gamma'$ implies that there is $\alpha \in B \cap B'$, such that $h_\gamma(\alpha) \neq h_{\gamma'}(\alpha)$, and this is impossible due to the compatibility.

Contradiction.

□.

We conjecture that the measurability assumption can be much weaken. However, the following positive result can be proved:

Proposition 2.14 *Suppose that there is a club $C \subseteq \kappa$ such that for every $\alpha \in C, \langle Q_\alpha, \leq_{Q_\alpha}^* \rangle$ is forced to be α^+ -closed. Then P_κ preserves κ^+ .*

In particular, if κ is not a Mahlo cardinal, then P_κ preserves κ^+ .

Proof. Let $p \in P_\kappa$ and $\underset{\sim}{f}$ be a name such that

$$p \Vdash \underset{\sim}{f} : \kappa \rightarrow \kappa^+.$$

Fix a club C such that for every $\alpha \in C$, $\langle Q_\alpha, \leq^*_{Q_\alpha} \rangle$ is forced to be α^+ -closed. Assume also that for every $\alpha \in C$, for every $\beta < \alpha$, $|P_\beta| < \alpha$. Let $\langle \alpha_i \mid i < \kappa \rangle$ be an increasing continuous enumeration of C . Apply Lemma 2.5 and find $p_0 \geq^* p$, $p_0 \restriction \alpha_0 = p \restriction \alpha_0$ and $\eta_0 < \kappa^+$ such that $p_0 \Vdash \underset{\sim}{f}(0) < \eta_0$.

Continue by induction and define a $\leq^*$ -increasing sequence $\langle p_i \mid i < \kappa \rangle$ and sequence $\langle \eta_i \mid i < \kappa \rangle$ of ordinals below κ^+ such that

1. $p_i \Vdash \underset{\sim}{f}(i) < \eta_i$,
2. for every $i < j$, $p_j \restriction \alpha_i = p_i \restriction \alpha_i$,

There is no problem at limit stages i , since $\langle P_\kappa \setminus \alpha_i, \leq^* \rangle$ is α_i^+ -closed since $\alpha_i \in C$.

The second item insures that there is $p^* \in P_\kappa$ such that $p^* \geq^* p_i$, for every $i < \kappa$. Then

$$p^* \Vdash \text{rng}(\underset{\sim}{f}) \subseteq \bigcup_{i < \kappa} \eta_i.$$

□

3 Non-stationary support iterations

We assume GCH throughout as before. Let κ be an inaccessible cardinal.

Let $\langle P_\alpha, Q_\beta : \alpha \leq \kappa, \beta < \kappa \rangle$ be a non-stationary support iteration of Prikry type forcings, with the properties stated in the introduction.

Let I be a stationary subset of κ which consists of singular cardinals μ such that for every $\gamma < \mu$, $|P_\gamma| < \mu$.

Assume that-

If $\alpha \in I$, then $\Vdash_{P_\alpha} \langle P_\kappa \setminus \alpha, \leq^* \rangle$ is α^{++} -closed.

Note that for a singular cardinal α , we have $\Vdash_{P_\alpha} \langle P_\kappa \setminus \alpha, \leq^* \rangle$ is α^+ -closed. We will need a slightly more closure. A typical situation is where κ is Mahlo, Q_α is trivial at every accessible α and each forcing Q_α has cardinality below the least inaccessible above α .

Theorem 3.1 P_κ does not add fresh unbounded subsets to κ .

Proof. Recall the following fusion lemma for the non-stationary support iteration of Prikry forcings²:

Lemma 3.2 Let $p \in P_\kappa$. For every $\beta < \kappa$, let $F(\beta)$ be a P_β -name for a $\leq^*$ -dense open subset of $P \setminus \beta$ above $p \restriction \beta$, and assume that this is forced by $p \restriction \beta$. Then there exist $p^* \geq^* p$ and a club $C \subseteq \kappa$ such that for every singular $\beta \in C$, $p^* \restriction \beta \Vdash p^* \restriction \beta \in F(\beta)$.

Let $G \subseteq P_\kappa$ be generic over V , and assume for contradiction that there exists a function $f \in 2^\kappa$ which is the characteristic function of a fresh subset of κ . Let $\tilde{f}$ be a P_κ -name for it, and assume that this is forced by some condition in G . For simplicity, assume that this is the weakest condition.

Let $\zeta \in \kappa \cap I$ be the least ordinal for which a new subset is added in the extension from V to $V[G]$. Such ζ exists, since the forcings Q_α (for $\alpha < \kappa$) have cardinality below κ , and at least one of them is non-trivial.

Note that $\zeta \in \kappa \cap I$, $\langle P_\kappa \restriction \zeta, \leq^* \rangle$ is ζ^{++} -closed. Pick a condition $q \in P_\zeta$ which forces that a new subset is added to ζ . For simplicity, assume that the weakest condition in P_ζ forces this (else, work above a condition in P_κ whose restriction to P_ζ equals q).

We divide into two cases:

Case 1.³ There exists $\mu \in (\zeta, \kappa) \cap I$ and a condition $p^* \in P_\mu$ which forces that the following property holds:

$$\begin{aligned} & \exists p \in G_\mu \exists s \in P \setminus \mu \forall r \geq^* s \exists \xi < \kappa \exists r_0, r_1 \geq^* r, \\ & V \models (p \restriction r_0 \parallel \tilde{f} \restriction \xi, p \restriction r_1 \parallel \tilde{f} \restriction \xi), \text{ and the decisions are different.} \end{aligned}$$

(here, G_μ denotes the canonical name for the generic set for P_μ). By extending p^* , we can decide the value of p in the statement above, and thus assume that $p^* \geq p$. Let $\tilde{s}$ be a P_μ -name for s from the above property, and assume that this is forced by p^* .

Let us apply the same methods as in the main lemma in [7]. We construct, in V , a binary tree of conditions, $\langle \langle p^*, \tilde{s}_\sigma \rangle : \sigma \in {}^{\mu>}2 \rangle$ and a tree of functions $\langle b_\sigma : \sigma \in {}^{\mu>}2 \rangle$ such that $\tilde{s}_\emptyset = \tilde{s}$, and for every $\sigma \in {}^{\mu>}2$:

$$1. \forall i < 2, \langle p^*, \tilde{s}_{\sigma \smallfrown \langle i \rangle} \rangle \parallel \tilde{f} \restriction lh(b_{\sigma \smallfrown \langle i \rangle}) = b_{\sigma \smallfrown \langle i \rangle}.$$

²The fusion property for non-stationary support iterations is due to Friedman and Magidor [2]. A version suitable for iterations of Prikry type forcings appeared in [1]. The proof is basically given in lemma 3.3 in [4].

³It basically repeats those of 4.11, [4].

2. $b_{\sigma \smallfrown \langle 0 \rangle} \perp b_{\sigma \smallfrown \langle 1 \rangle}$.
3. If $\text{lh}(\sigma)$ is limit, then p^* forces that $\mathcal{S}_\sigma$ is an upper bound, with respect to the direct extension order, of $\langle \mathcal{S}_{\sigma \restriction \xi} : \xi < \text{lh}(\sigma) \rangle$.
4. b_σ is an end extension of $b_{\sigma \restriction \xi}$ for every $\xi < \text{lh}(\sigma)$.

Now assume that $g \subseteq P_\mu$ is generic over V with $p^* \in g$. In $V[g]$, let $h \in 2^{<\mu}$ be the characteristic function of a new subset of μ (such a new subset exists because μ is above ζ). h defines a branch through the binary tree, $\langle \langle p^*, \mathcal{S}_{h \restriction \xi} \rangle : \xi < \mu \rangle$. Since $\langle \mathcal{S}_{h \restriction \xi} : \xi < \mu \rangle$ form the sequence of moves of the first player, when the second plays according to the strategy τ , there exists an upper bound $s^* \in P \setminus \mu$, which extends all the conditions in the sequence. Thus, there exists an upper bound for the branch, of the form $\langle p^*, \mathcal{S}^* \rangle$. It forces that—

$$b = \bigcup_{\xi < \mu} b_{h \restriction \xi}$$

is an initial segment of f . We argue that this must be a strict initial segment of f . Indeed, otherwise, $(\text{cf}(\kappa))^{V[G_\mu]} \leq \mu$. But, since κ is inaccessible, $G_\mu \subseteq P_\mu$ is a forcing whose cardinality is strictly below κ , so it preserves cofinalities greater of equal to κ .

Therefore, b is a strict initial segment of f , and thus $b \in V$. Therefore, h can be defined, in V , using the binary tree and the set b . This is a contradiction to the choice of h .

Case 2. For every $\mu \in (\zeta, \kappa) \cap I$, every condition in P_μ forces that—

$$\begin{aligned} & \forall p \in G_\mu \ \forall s \in P_\kappa \setminus \mu \exists r \geq^* s \ \forall \xi < \kappa \ \forall r_0, r_1 \geq^* r, \\ & V \models \text{ If } p \smallfrown r_0 \parallel \mathcal{f} \restriction \xi \text{ and } p \smallfrown r_1 \parallel \mathcal{f} \restriction \xi \text{ then the decisions are the same.} \end{aligned}$$

Define for every $\mu < \kappa$

$$\begin{aligned} e(\mu) = \{ & r \in P_\kappa \setminus \mu : \forall p \in G_\mu \ \forall \xi < \kappa \ \forall r_0, r_1 \geq^* r, \\ & V \models \text{ If } p \smallfrown r_0 \parallel \mathcal{f} \restriction \xi \text{ and } p \smallfrown r_1 \parallel \mathcal{f} \restriction \xi \text{ then the decisions are the same.} \} \end{aligned}$$

Claim 2 Suppose that $\mu \in (\zeta, \kappa) \cap I$. Then $e(\mu)$ is a dense open in $\langle P_\kappa \setminus \mu, \leq^* \rangle$.⁴

Proof. Just note that $|G_\mu| \leq \mu^+$ and $\langle P_\kappa \setminus \mu, \leq^* \rangle$ is μ^{++} -closed.

□ of the claim.

⁴It is the only place in the proof where μ^{++} -completeness is used.

Given $\mu \in (\zeta, \kappa) \cap I$ as above, the following set is also forced to be $\leq^*$ -dense open in $P \setminus \mu$:

$$d(\mu) = \{r \in P \setminus \mu : \exists g \in 2^\mu, r \Vdash \check{f} \restriction \mu = g\}.$$

The $\leq^*$ -density of $d(\mu) \subseteq P \setminus \mu$ follows as well from the fact that the direct extension order of $P \setminus \mu$ is more than μ -closed.

We can now apply the standard fusion argument 3.2. There exists $p \in P_\kappa$ and a club $C \subseteq \kappa$ such that $\min(C) > \zeta$, and, for every $\mu \in C \cap I$,

$$p \restriction \mu \Vdash p \setminus \mu \in d(\mu) \cap e(\mu).$$

For each $\mu \in C \cap I$, there exists a condition in G of the form $q_\mu \hat{\smallfrown} p \setminus \mu$, where $q_\mu \in P_\mu$, which decides the value of $\check{f} \restriction \mu$. The reason is that $p \restriction \mu$ forces that $p \setminus \mu$ is in $d(\mu)$, and thus the value of $\check{f} \restriction \mu$ is decided by the forcing P_μ .

For each such μ , q_μ is an extension of $p \restriction \mu$, and thus there exists a finite set $b_\mu \subseteq \mu$ such that at every $\delta \in \mu \setminus b_\mu$, $q_\mu(\delta)$ direct extends $p(\delta)$ (as forced by $q_\mu \restriction \delta$).

The function $\mu \mapsto \max b_\mu$ is a regressive function in $V[G]$, and its domain is the set $C \cap I$. $C \cap I$ is stationary in V since I is assumed to be stationary. By theorem 2.2, it is also stationary in $V[G]$. Since κ is still regular in $V[G]$, we can find an unbounded subset $S \subseteq \kappa$ and an ordinal $\mu^* < \kappa$ such that for every $\mu \in S$, $b_\mu \subseteq \mu^*$. By increasing μ^* , we can assume that it belongs to $C \cap I$.

Now, shrink S further to stabilize the function $\mu \mapsto q_\mu \restriction \mu^*$. This is possible since S is unbounded in κ , and $q_\mu \restriction \mu^*$ is a condition in P_{μ^*} which has a small cardinality (and, again, κ is inaccessible in $V[G]$).

So we can assume that there exists a condition $q^* \in P_{\mu^*}$, such that for every $\mu \in S$, there exists some direct extension $r_\mu \in P \setminus \mu^*$ of $p \setminus \mu^*$, such that—

$$q^* \hat{\smallfrown} r_\mu \Vdash \check{f} \restriction \mu$$

and $q^* \hat{\smallfrown} r_\mu \in G$.

Recall that $\mu^* \in C \cap I$ and the condition p obtained by fusion above also satisfies that $p \restriction \mu^* \Vdash p \setminus \mu^* \in e(\mu^*)$. Since $q^* \geq p \restriction \mu^*$, the condition q^* forces that, for every $\xi < \kappa$, any pair of direct extensions of $p \setminus \mu^*$ which decide $\check{f} \restriction \xi$, decide this initial segment the same way.

It follows that $q^* \hat{\smallfrown} p \setminus \mu^*$ decides $\check{f}$ entirely, and forces it to be the following function of V :

$$h = \bigcup_{\mu \in \kappa \setminus \mu^*} \{g \in 2^\mu : \text{there exists a } P_{\mu^*}\text{-name for an extension } \mathcal{I} \geq^* p \setminus \mu^*, \\ \text{such that } q^* \restriction r \Vdash \mathcal{I} \restriction \mu = g\}$$

which is a contradiction. $\square$

Remark 3.3 Lemma 3.2 implies that $V[G] \models 2^\kappa = \kappa^+$. Indeed, assume that A is a subset of κ in $V[G]$. Let $\mathcal{A}$ be a P_κ -name for it. For every singular $\beta < \kappa$, define in V^{P_β} the set–

$$F(\beta) = \{q \in P_\kappa \setminus \beta : \exists A_\beta \subseteq \beta, q \Vdash \mathcal{A} \cap \beta = A_\beta\}.$$

Note that β is singular and thus $F(\beta)$ is $\leq^*$ -dense open. Let $p^* \in G$ and $C \subseteq \kappa$ be such that for every singular $\beta \in C$, $p^* \restriction \beta \Vdash p^* \setminus \beta \in F(\beta)$. Then there exists a P_β -name $\mathcal{A}_\beta$ for a subset of β , such that $p^* \Vdash \mathcal{A} \cap \beta = (\mathcal{A}_\beta)_{\mathcal{I} \restriction P_\beta}$. Then $A = (\mathcal{A})_G$ can be computed in $V[G]$ from the sequence $\langle \mathcal{A}_\beta : \beta \in C \rangle$. By using canonical names for bounded subsets of κ , and by GCH in V , there are at most κ^+ such sequences. So there are at most κ^+ –many subset of κ in $V[G]$.

The situation with higher cardinals was clarified in [4]. The following was shown basically in [4], 4.11:

Lemma 3.4 *P_κ does not add fresh unbounded subsets to κ^+ , or to any cardinal λ of V with $\text{cf}(\lambda) > \kappa$.*

Proof. The proof is a variation of the proof of theorem 3.1, and it basically appears in [4]. Assume that $\mathcal{f}$ is a P_κ -name for the characteristic function of a fresh unbounded subset of λ . Divide into cases as in the proof of theorem 3.1. Case 1 remains the same. Case 2 is simplified, since the sets $d(\mu)$ is no longer required. Indeed, in the notations of the proof of theorem 3.1, assume that $p \in P_\kappa$ is a condition and $C \subseteq \kappa$ is a club, such that for every $\mu \in C \cap I$,

$$p \restriction \mu \Vdash p \setminus \mu \in e(\mu)$$

Now, work in $V[G]$. For each $\xi < \lambda$, let $q_\xi \in P_\kappa$ be an extension of p which decides $\mathcal{f} \restriction \xi$. Let $b_\xi \subseteq \kappa$ be a finite set such that, for every $\alpha \in b_\xi$, $q_\xi \restriction \alpha \Vdash q_\xi(\alpha) \geq^* p(\alpha)$. Let $\mu_\xi < \kappa$ be an upper bound on b_ξ . Since $\text{cf}(\lambda) > \kappa$ in V , the same holds true in $V[G]$ as well (by the same proof as in corollary 2.6). Thus, there exists $\mu^* \in C \cap I$ such that, for an unbounded

$S \subseteq \lambda$, $\mu_\xi < \mu^*$. By shrinking S , we can assume that, for some $q^* \in P_{\mu^*}$, $q_\xi \restriction \mu^* = q^*$. Then q^* satisfies that, for every $\xi \in S$, there exists a direct extension $r_\xi \in P \restriction \mu^*$ of $p \restriction \mu^*$, such that $q^* \restriction r_\xi \parallel \check{f} \restriction \xi$. Now, as in the proof of case 2 in theorem 3.1, the condition $q^* \restriction p \restriction \mu^*$ forces $\check{f}$ to be the following function of V :

$$h = \bigcup_{\xi < \kappa^+} \{g \in 2^\mu : \text{there exists a } P_{\mu^*}\text{-name for an extension } \check{r} \geq^* p \restriction \mu^*, \\ \text{such that } q^* \restriction r \Vdash \check{f} \restriction \xi = g\}$$

which is a contradiction.

□

We finish with an application for iterations of Prikry type forcings with the nonstationary support. The referee of [9] asked if such an iteration, below a cardinal κ , can add new measurable cardinals below κ . We show that the answer is negative⁵.

Theorem 3.5 *Assume GCH and let κ be an inaccessible cardinal.*

Let P_κ be a non-stationary support iteration of Prikry-type forcings satisfying the conditions from the beginning of the section. Assume:

() For every Mahlo cardinal $\alpha < \kappa$ which is not measurable in V , $\langle P_\kappa \restriction \alpha, \leq^* \rangle$ is α^+ -closed.*

Let λ be a cardinal such that $\forall \tau < \lambda (|P_\tau| < \lambda)$. Let $G \subseteq P_\kappa$ be generic over V . Then If λ is measurable in $V[G]$, it was already measurable in V .

Furthermore, if the assumption () is strengthened to–*

*(**) For every Mahlo cardinal $\alpha < \kappa$, $\langle P_\kappa \restriction \alpha, \leq^* \rangle$ is α^+ -closed.*

then λ is measurable in $V[G]$ if and only if it is measurable in V .

Proof. Assume first that (**) holds and λ is a measurable cardinal in V . If $\lambda > \kappa$, the Levy-Solovay Theorem [11] shows that λ is measurable in $V[G]$. If $\lambda \leq \kappa$, standard arguments show that λ remains measurable in $V[G \restriction P_\lambda]$ (see [5]). If $\lambda < \kappa$, the forcing $\langle P_\kappa \restriction \lambda, \leq^* \rangle$ is λ^+ -closed, so $P_\kappa \restriction \lambda$ does not add subsets to λ and λ remains measurable in $V[G]$. Thus, let us concentrate on the other direction assuming (*).

We first recall Lemma 2.1 in [4]: given a forcing notion which does not add new fresh unbounded subsets to cardinals of V in the interval $[\kappa, (2^\kappa)^V] = [\kappa, \kappa^+]$, every κ -complete ultrafilter in the generic extension extends a κ -complete ultrafilter from V . Assume now

⁵The answer for the same question in the full or Easton support is known to be negative, see [3].

that α is measurable in $V[G]$, and let $W \in V[G]$ be a nontrivial κ -complete ultrafilter on α . If $\alpha \geq \kappa$, then by the results in this section, P_κ does not add fresh unbounded subsets to α , α^+ , and thus $W \cap V \in V$ by Lemma 2.1 in [4]. Thus, assume that $\alpha < \kappa$, and assume that α is not measurable in V . So, α is a Mahlo cardinal in V . If the forcing $\langle P_\kappa \setminus \alpha, \leq^* \rangle$ is α^{++} -closed, then $W \in V[G \restriction P_\alpha]$.

However we assumed only that $\langle P_\kappa \setminus \alpha, \leq^* \rangle$ is α^+ -closed. In this case W need not be in $V[G \restriction P_\alpha]$.

Proceed then as follows. Work in $V[G \restriction P_\alpha]$. Let $\langle A_i \mid i < \alpha^+ \rangle$ be an enumeration of all subsets of α (such enumeration exists by applying remark 3.3 on P_α). Define a $\leq^*$ -increasing sequence of conditions $\langle p_i \mid i < \alpha \rangle$ in $V[G \restriction P_\alpha]$ such that for every $i < \alpha$, $p_i \Vdash A_i \in \mathcal{W}$. Set $W' = \{A_i \mid i < \alpha, p_i \Vdash A_i \in \mathcal{W}\}$. Then W' will be an α -complete ultrafilter over α in $V[G \restriction P_\alpha]$.

Apply now 3.1 and 3.4 to P_α . It follows that no fresh subsets are added to α, α^+ . Now, by Lemma 2.1 from [4], $W' \cap V \in V$ is a nontrivial α -complete ultrafilter over α in V . A contradiction.

□

Remark 3.6 1. The closure assumptions made on $\langle P_\kappa \setminus \alpha, \leq^* \rangle$ are needed.

For example, start with $V = L[U]$, where U is a normal ultrafilter over α . Iterate $Cohen(\beta), \beta < \alpha$. Let V be this model. Then α is not a measurable in V . Force with $Cohen(\alpha)$ over V . Then α will be a measurable in the extension. Here we take Q_β to be trivial for every $\beta < \alpha$ and $Q_\alpha = Cohen(\alpha)$.

2. Also, the assumption $\forall \tau < \lambda (|P_\tau| < \lambda)$ is necessary. Just use the previous example (with $\lambda = \alpha$). $Q_0 = Cohen(\alpha)$ resurrects measurability of α .
3. Note that a measurable cardinal in V need not be such in $V[G]$ without assuming $(*)$. Just use the Prikry forcing or the iteration of such forcings.

4 On fresh sets in the Easton support iterations

We start with an easier case of the Easton support iterations with κ being a Mahlo cardinal. κ -c.c. of the forcing will be used to show the following:

Theorem 4.1 *Suppose that κ is a Mahlo cardinal and P_κ is an Easton support iteration. Then no fresh sets are added to κ .*

Proof. Suppose otherwise. Work in V . Let $\mathcal{A}$ be a name of such subset.

Define a tree T of possibilities as follows. Fix an increasing enumeration $\langle \kappa_\xi \mid \xi < \kappa \rangle$ of all inaccessible cardinals below κ .

For every $\xi < \kappa$, let

$$Lev_\xi(T) = \{x \subseteq \kappa_\xi \mid \exists p \in P_\kappa \quad p \Vdash \mathcal{A} \cap \kappa_\xi = x\}.$$

Let $x \in Lev_\alpha(T), y \in Lev_\beta(T)$. Set $x >_T y$ iff $\alpha > \beta$ and $x \cap \kappa_\beta = y$.

Then $\langle T, <_T \rangle$ is a κ -tree, since κ is an inaccessible.

Lemma 4.2 $\langle T, <_T \rangle$ has a κ -branch.

Proof. Let $\langle x_\gamma \mid \gamma < \kappa \rangle$ be an enumeration of T .

There is a club $C \subseteq \kappa$ such that for every $\gamma, \delta \in C$ the following hold:

1. $\kappa_\gamma = \gamma$,
2. the level of $x_\gamma \geq \gamma$,
3. if $\gamma < \delta$, then the level of $x_\gamma < \delta$.

For every $\gamma \in C$, pick $p_\gamma \in P_\kappa$ such that $p_\gamma \Vdash \mathcal{A} \cap \kappa_{\xi_\gamma} = x_\gamma$, where $\xi_\gamma \geq \gamma$ denotes the level of x_γ .

Now, κ is a Mahlo cardinal and an Easton support was used, hence there is a stationary $S \subseteq C$ such that for every $\gamma, \delta \in S$, p_γ and p_δ are compatible.

Take any two $\gamma < \delta$ in S . Then $x_\delta \cap \xi_\gamma = x_\gamma$ due to the compatibility of p_γ and p_δ .

So, $\{x_\gamma \mid \gamma \in S\}$ is a κ -branch.

□

Let now $b = \{x_i \mid i < \kappa\}$ be a maximal κ -branch in T . For every $i < \kappa$ fix $p_i \in P_\kappa$ which witnesses that $x_i \in T$.

By the assumption made, $\bigcup_{i < \kappa} x_i \neq A$, since this union is in V .

Then for every $i < \kappa$, there is $i' \geq i$ such that $x_{i'}$ is a splitting point of T .

Denote by $y_{i'}$ an immediate successor of $x_{i'}$ which is not in b . Let $q_{i'}$ be a condition which witnesses that $y_{i'} \in T$.

Let $C \subseteq \kappa$ be a club such that for every $i_1, i_2 \in C, i_1 < i_2$, we have $i'_1 < i_2$.

The next lemma provides the desired contradiction, since P_κ satisfies κ -c.c.

Lemma 4.3 The conditions $\{q_{i'} \mid i \in C\}$ are pairwise incompatible.

Proof. Let $i_1 < i_2$ be in C . Then

$$q_{i'_1} \Vdash \underset{\sim}{A} \cap \kappa_{i'_1} \neq x_{i'_1+1}.$$

However,

$$q_{i'_2} \Vdash \underset{\sim}{A} \cap \kappa_{i'_1} = x_{i'_1+1},$$

since $y_{i'_2} >_T x_{i_2} >_T x_{i'_1+1}$. This is possible onle when $q_{i'_1}$ and $q_{i'_2}$ are incompatible.

□

Let us give an example of the Easton support iteration P_κ which adds a fresh subset, however we give up here the assumption that $|\underset{\sim}{Q}_\beta| < \kappa$.

Let $\langle \kappa_\beta \mid \beta < \kappa \rangle$ be an increasing sequence of measurable cardinals above an inaccessible κ .

Let $\langle P_\alpha, \underset{\sim}{Q}_\beta \mid \alpha \leq \kappa, \beta < \kappa \rangle$ be an Easton support iterations of the Prikry forcings, i.e. for each $\beta < \kappa$, $\underset{\sim}{Q}_\beta$ is the Prikry forcing with a normal ultrafilter over κ_β .

Let G_κ be a generic subset of P_κ . For every $\beta < \kappa$, let b_β be the Prikry sequence added by G_κ to κ_β .

Lemma 4.4 *The set*

$$A = \{\alpha < \kappa \mid \text{the first element of the sequence } b_{\kappa_\alpha} \text{ is } 0\}$$

is a fresh subset of } \kappa.

Proof. Every initial segment of A is in V due to the support used. On the other hand $A \notin V$, since every condition in the forcing P_κ should be bounded in κ , and so it can be extended to one which forces $b_{\kappa_\alpha}(0) = 0$ or to one forcing $b_{\kappa_\alpha}(0) \neq 0$.

□

Let us turn now to a general case, i.e. we assume only that κ is an inaccessible.

Our aim will be to prove the following :

Theorem 4.5 *Let κ be an inaccessible cardinal and $\langle P_\alpha, \underset{\sim}{Q}_\beta \mid \alpha \leq \kappa, \beta < \kappa \rangle$ be the Easton support iteration of Prikry type forcing notions. Let I be a stationary subset of κ which consists of singular cardinals μ such that for every $\gamma < \mu$, $|P_\gamma| < \mu$. Suppose that for every $\alpha \in I$, $\Vdash_{P_\alpha} \langle P_\kappa \setminus \alpha, \leq^* \rangle$ is α^{++} -closed.*

Let $G_\kappa \subseteq P_\kappa$ be a generic. Then, in $V[G_\kappa]$, there is no fresh subsets of κ .

Remark 4.6 Similar results were proved in [4] for cardinals above κ . The proof there is based on the fact that $|P_\kappa| = \kappa$ and it is much easier.

Proof. Let G_κ be a generic subset of P_κ .

We would like to show that there is no fresh subset of κ in $V[G_\kappa]$.

Suppose otherwise. Work in V . Let $\mathcal{A}$ be a name of such subset and let $\mathcal{f}$ be a name of the characteristic function of A . Fix some $p \in G_\kappa$ which forces this.

Let $\zeta < \kappa$ be an ordinal for which a new subset is added in the extension from V to $V[G]$. Such ζ exists, since the forcings Q_α (for $\alpha < \kappa$) have cardinality below κ , and at least one of them is non-trivial. By increasing if necessary, we can assume that ζ is a singular cardinal. Then $\langle P_\kappa \setminus \zeta, \leq^* \rangle$ is more than ζ -closed. Thus, there exists a condition $q \in G_\kappa \restriction P_\zeta$ which forces that a new subset is added to ζ . For simplicity, assume that $p \restriction \zeta$ forces this.

Given a condition $r \in P_\kappa$, let us denote by $r(\gamma)$ its γ -th coordinate, i.e. $r = \langle r(\gamma) \mid \gamma < \kappa \rangle$.

We divide into two cases as in Theorem 3.1.

Case 1. There exists $\mu \in (\zeta, \kappa) \cap I$ and a condition $p^* \in P_\mu$ which forces that the following property holds:

$$\begin{aligned} \exists p \in G_\mu \exists s \in P \setminus \mu \forall r \geq^* s \exists \xi < \kappa \exists r_0, r_1 \geq^* r, \\ V \models (p \restriction r_0 \parallel \mathcal{f} \restriction \xi, p \restriction r_1 \parallel \mathcal{f} \restriction \xi), \text{ and the decisions are different.} \end{aligned}$$

Case 2. For every $\mu \in (\zeta, \kappa) \cap I$, every condition in P_μ forces that—

$$\begin{aligned} \forall p \in G_\mu \forall s \in P_\kappa \setminus \mu \exists r \geq^* s \forall \xi < \kappa \forall r_0, r_1 \geq^* r, \\ V \models \text{If } p \restriction r_0 \parallel \mathcal{f} \restriction \xi \text{ and } p \restriction r_1 \parallel \mathcal{f} \restriction \xi \text{ then the decisions are the same.} \end{aligned}$$

The treatment of the first case is exactly as in Theorem 3.1. Let us deal with the second case.

As in 3.1, we define for every $\mu < \kappa$

$$\begin{aligned} e(\mu) = \{r \in P_\kappa \setminus \mu : \forall p \in G_\mu \forall \xi < \kappa \forall r_0, r_1 \geq^* r, \\ V \models \text{If } p \restriction r_0 \parallel \mathcal{f} \restriction \xi \text{ and } p \restriction r_1 \parallel \mathcal{f} \restriction \xi \text{ then the decisions are the same.}\} \end{aligned}$$

By Claim 1 of 3.1, it is $\leq^*$ -dense open subset of $P_\kappa \setminus \mu$, for every $\mu \in (\zeta, \kappa) \cap I$. Again, here is the only place where μ^{++} -closure of the direct order on $P_\kappa \setminus \mu$ is used.

Given a generic $G_\kappa \subseteq P_\kappa$, $p \in G_\kappa$, define in $V[G_\kappa]$,

$$S = \{\xi < \kappa \mid \xi \text{ is limit ordinal and } \exists t \geq^* p, t \in G_\kappa \text{ such that } t \restriction \xi \Vdash_{P_\xi} t \restriction \xi \parallel_{P_\kappa \setminus \xi} \mathcal{A} \cap \xi\}.$$

For every $\xi \in S$, fix some $t^\xi \in G_\kappa$ such that $t^\xi \geq^* p$ and $t^\xi \restriction \xi \Vdash_{P_\xi} t^\xi \restriction \xi \parallel_{P_\kappa \setminus \xi} \mathcal{A} \cap \xi$. So, there is $p^\xi \in G \restriction P_\xi$, $p^\xi \geq t^\xi \restriction \xi$, $(p^\xi)^\frown t^\xi \restriction \xi \Vdash \mathcal{A} \cap \xi = a_\xi$, for some $a_\xi \in V$. Then there is a finite $b^\xi \subseteq \xi$ such that $p^\xi \restriction b^\xi \geq^* p \restriction (\xi \setminus b^\xi)$.

Suppose for a moment that S is stationary in $V[G_\kappa]$.

Then we can find a stationary subset S' of S and a finite b such that for every $\xi \in S'$, $b^\xi = b$. Now we can freeze $p^\xi \upharpoonright \max(b)$. Denote $\max(b)$ by μ^* .

Let $\mu \in I$ be a cardinal above μ^* .

Consider the set $e(\mu)$ defined above. It is $\leq^*$ -dense open subset of $P_\kappa \setminus \mu$, in $V[G_\mu]$ above $p \setminus \mu$. In particular there is $r \in P_\kappa \setminus \mu$, $r \geq^* p \setminus \mu$ such that for every $p' \in G_\mu$, for every $\xi < \kappa$ and for every $r_0, r_1 \geq^* r$

$$V \models \text{If } p' \frown r_0 \Vdash_{\mathcal{A}} A \cap \xi \text{ and } p' \frown r_1 \Vdash_{\mathcal{A}} A \cap \xi \text{ then the decisions are the same.}$$

Recall that $t^\xi \geq^* p$ and $p^\xi \setminus \mu^* \geq^* p \upharpoonright (\mu^*, \xi)$, for every $\xi \in S'$. If we were able to conclude from this that $p^\xi \setminus \mu \geq^* r$, then it will imply that $A \in V$. However it need not be the case since the support of r may be bigger than those of $p \setminus \mu$ and incompatibility may occur on coordinates outside of $\text{supp}(p)$.

Let us argue that it is possible to overcome this obstacle.

Work in V . Set $p_0 = p$. Use Lemma 2.5 to find $q \geq^* p$ and $\mu_0 \geq \sup(\text{supp}(p))$, $\mu_0 \in I$ such that $q \Vdash_{\mu_0} \mu^* \leq \mu_0$.

Then, we use the density of $e(\mu_0)$ to find $r_0 \geq^* q \setminus \mu_0$ such that

$$q \upharpoonright \mu_0 \Vdash_{P_{\mu_0}} r_0 \in e(\mu_0).$$

Set $p_1 = q \upharpoonright \mu_0 \frown r_0$.

Next, we run the argument above with $p = p_0$ replaced by p_1 . Again, using Lemma 2.5 find $q_1 \geq^* p_1$ and $\mu_1 \geq \sup(\text{supp}(p_1))$, $\mu_0 + 1, \mu_1 \in I$ such that $q_1 \Vdash_{\mu_1} \mu^* \leq \mu_1$, where μ^* is now defined using p_1 instead of p .

Then, use the density of $e(\mu_1)$ to find $r_1 \geq^* q_1 \setminus \mu_1$ such that

$$q_1 \upharpoonright \mu_1 \Vdash_{P_{\mu_1}} r_1 \in e(\mu_1).$$

Set $p_2 = q \upharpoonright \mu_1 \frown r_1$.

Continue by induction and define p_n, q_n, r_n, μ_n , for every $n < \omega$.

Finally set $\mu_\omega = \bigcup_{n < \omega} \mu_n$ and $p_\omega = \bigcup_{n < \omega} p_n = q_\omega = \bigcup_{n < \omega} q_n$. Then, for every $n < \omega$,

$$p_\omega \upharpoonright \mu_n \Vdash_{P_{\mu_n}} p_\omega \setminus \mu_n \in e(\mu_n),$$

since $p_\omega \setminus \mu_n \geq^* r_n \in e(\mu_n)$ and $e(\mu_n)$ is dense open. Also, $\sup(\text{supp}(p_\omega)) = \mu_\omega$.

Pick now a generic $G \subseteq P_\kappa$ with $p_\omega \in G$. Let S_{p_ω} be defined as S above only with p_ω replacing p . Assuming its stationarity, define $q^\xi \geq^* p_\omega$, $p^\xi \in G \upharpoonright P_\xi$ for $\xi \in S'$ exactly as above. Then

there will be a stationary $S' \subseteq S$ and $n^* < \omega$ such that $p^\xi \restriction (\mu_{n^*}, \xi) \geq^* p_\omega \restriction (\mu_{n^*}, \xi)$, since $\sup(\text{supp}(p_\omega)) = \mu_\omega$ and a non-direct extension is used at finitely many places only.

Shrink S' further to S'' and stabilise the value of the function $\xi \mapsto p^\xi \restriction \mu_{n^*}^+$.

Finally, we use that $p_\omega \restriction \mu_{n^*} \in e(\mu_{n^*})$.

Hence, the following lemma will complete the proof. We prove it for the initial p , but the same argument works for S_{p_ω} or any S_x with $x \geq p$.

Lemma 4.7 *S is stationary in $V[G_\kappa]$.*

Proof. The argument will be similar to those of 2.2.

Suppose otherwise. Let $C \subseteq \kappa$ be a club disjoint from S . Assume that $p \in G_\kappa$ forces this, otherwise replace it by a stronger condition doing this.

Work in V . Pick an elementary submodel $M \preceq H_\chi$ such that

1. $|M| = \delta < \kappa$,
2. $M \cap \kappa = \delta$,
3. $\text{cof}(\delta) < \delta$,
4. ${}^{\text{cof}(\delta)}M \subseteq M$,
5. $\kappa, P_\kappa, \mathcal{Q}, p \in M$.

Pick a cofinal in δ sequence $\langle \delta_i \mid i < \text{cof}(\delta) \rangle$ consisting of singulars and with $\delta_0 > \text{cof}(\delta)$.

Let $G_{\text{cof}(\delta)+1} \subseteq P_{\text{cof}(\delta)+1}$ be a generic with $p \restriction \text{cof}(\delta) + 1 \in G_{\text{cof}(\delta)+1}$.

Work in $V[G_{\text{cof}(\delta)+1}]$.

Consider

$$D^0 = \{r \geq p \mid \exists c < \kappa \quad r \Vdash c = \min(\mathcal{Q} \setminus \delta_0)\}.$$

Clearly, D^0 is a dense open and it belongs to M .

Apply Lemma 2.3. Then there will be $\alpha_0 < \kappa$ and $q_0 \geq^* p, q \restriction \text{cof}(\delta) + 1 = p \restriction \text{cof}(\delta) + 1$, in M such that for every $r \in P_{\alpha_0}, r \geq_{P_{\alpha_0}} q_0 \restriction \alpha_0$ there is $r' \geq_{P_{\alpha_0}} r$ such that $r' \restriction q_0 \setminus \alpha_0 \in D_0$. So, for every such r' there is $c(r') < \kappa$ such that

$$r' \restriction q_0 \setminus \alpha_0 \Vdash c(r') = \min(\mathcal{Q} \setminus \delta_0).$$

Note that $P_{\alpha_0} \subseteq M$. Hence all $c(r')$'s are in M . Also $|P_{\alpha_0}| < \kappa$. Hence their sup is below κ , and then, by elementarity, in M . Denote it by c_0^* .

Define, for every $\tau < \kappa$,

$$D_\tau = \{r \in P \mid r \geq^* p \text{ and } \exists \xi \in \kappa \setminus \tau \quad r \restriction \xi \Vdash_{P_\xi} (r \restriction \xi) \parallel_{P_{>\xi}} \dot{A} \cap \xi\}.$$

Claim 3 D_τ is $\leq^*$ -dense open above p .

Proof. Set $\tau_0 = \tau + 1$. Consider

$$D(\tau_0) = \{r \in P_\kappa \mid r \perp p \text{ or } (r \geq p \text{ and } r \parallel \dot{A} \cap \tau_0)\}.$$

By Lemma 2.3, there are α_0 and $q \geq^* p$ such that

$$q \restriction \alpha_0 \Vdash \exists b \in \dot{G}_{\alpha_0} \quad b \restriction q \setminus \alpha_0 \parallel \dot{A} \cap \tau_0.$$

If $\alpha_0 \leq \tau_0$ then we are done. Suppose that $\alpha_0 > \tau_0$. Consider $D(\alpha_0)$ and again, using Lemma 2.3, pick α_1 and $q_1 \geq^* q$ such that

$$q_1 \restriction \alpha_1 \Vdash \exists b \in \dot{G}_{\alpha_1} \quad b \restriction q_1 \setminus \alpha_1 \parallel \dot{A} \cap \alpha_0.$$

If $\alpha_1 \leq \alpha_0$, then $q_1 \in D(\alpha_0)$ and we are done. If $\alpha_1 > \alpha_0$, then continue and define in the same fashion α_2, q_2 etc.

Suppose that the process continues infinitely many steps. Then we will have

$$\alpha_0 < \alpha_1 < \dots < \alpha_i < \dots \text{ and } q_0 \leq^* q_1 \leq^* \dots \leq^* q_i \leq^* \dots, i < \omega.$$

Let $\alpha^* = \bigcup_{i < \omega} \alpha_i$ and $q^* \geq^* q_i$, for every $i < \omega$. Let $G \subseteq P_\kappa$ be a generic with $q^* \in G$. For every $i < \omega$, let $G_{\alpha_i} = G \restriction P_{\alpha_i}$ and $G^* = G \restriction P_{\alpha^*}$.

Now, for every $i < \omega$, there are $r_{i+1} \in G_{\alpha_{i+1}}$ and $a_i \in V$ such that

$$r_{i+1} \restriction q_{i+1} \setminus \alpha_{i+1} \Vdash \dot{A} \cap \alpha_i = a_i.$$

Then,

$$r_{i+1} \restriction q^* \setminus \alpha_{i+1} \Vdash \dot{A} \cap \alpha_i = a_i.$$

Set $a = \bigcup_{i < \omega} a_i$. Then, in $V[G_{\alpha^*}]$, $q^* \restriction \alpha^* \Vdash \dot{A} \cap \alpha^* = a$, since of r_i 's is in G_{α^*} . So, there is $r \in G_{\alpha^*}$ such that

$$r \restriction q^* \setminus \alpha^* \Vdash \dot{A} \cap \alpha^* = a.$$

Hence,

$$r \Vdash_{P_{\alpha^*}} (q^* \setminus \alpha^* \parallel \dot{A} \cap \alpha^*).$$

The only requirement on G_{α^*} is that $q^* \restriction \alpha^* \in G_{\alpha^*}$. Hence,

$$q^* \restriction \alpha^* \Vdash q^* \setminus \alpha^* \parallel \dot{A} \cap \alpha^*.$$

So, $q^* \in D_\tau$.

□ of the claim.

Consider $D_{c_0^*}$.

It is in M , as well. So, inside M , we can pick $\xi_0 \geq c_0^*, \xi_0 \in I$ and $t_0 \geq^* p$ such that

$$t_0 \restriction \xi_0 \Vdash_{P_{\xi_0}} t_0 \setminus \xi_0 \parallel_{P_{\kappa \setminus \xi_0}} \dot{A} \cap \xi_0 \}.$$

We continue the same process only with c_0 replaced by $\max(\xi_0, \delta_1)$ and p by q_0 .

At limit stages of the construction and at the stage $\text{cof}(\delta)$ itself we would like to put $\langle q_i \mid i < \text{cof}(\delta) \rangle$ into a single condition q^* . This is possible since $\delta_0 > \text{cof}(\delta)$, and so, we have enough completeness to proceed.

We have $p \restriction \text{cof}(\delta) + 1 \restriction q^* \Vdash \delta \in \dot{C}$, since C is forced to be closed by p .

Let $G \subseteq P_\kappa$ be a generic with $q^* \in G$. Set $G_\delta = G \restriction P_\delta$. Then for every $i < \text{cof}(\delta)$, $q_i \restriction \delta \in G_\delta$. Hence, there are $r_i \in G_{\xi_i}, r_i \geq q_i \restriction \xi_i$ and $a_i \in V$ such that

$$r_i \restriction q_i \setminus \xi_i \Vdash \dot{A} \cap \xi_i = a_i.$$

Set $a = \bigcup_{i < \text{cof}(\delta)} a_i$. Then, in $V[G]$, $A \cap \delta = a$, since each $r_i \restriction q_i \setminus \xi_i$ is in G .

Remember that the only requirement on a generic set G was that q^* belongs to it. Hence, back in V ,

$$q^* \restriction \delta \Vdash_{P_\delta} q^* \setminus \delta \parallel_{P_{>\delta}} \dot{A} \cap \delta.$$

So, $\delta \in S$ and as it was shown above $\delta \in C$ as well. Contradiction.

□

5 No fresh subsets of κ in the full support

Our aim will be to prove the following :

Theorem 5.1 *Let κ be an inaccessible cardinal and $\langle P_\alpha, \mathcal{Q}_\beta \mid \alpha \leq \kappa, \beta < \kappa \rangle$ be the full iteration of Prikry type forcing notions. Suppose that for every $\beta < \kappa$, for every $x, y, z \in \mathcal{Q}_\beta$, if $z \leq_{\mathcal{Q}_\beta}^* x, y$ and x, y are compatible according to $\leq_{\mathcal{Q}_\beta}$, then they are compatible according to $\leq_{\mathcal{Q}_\beta}^*$, i.e. there is $e \in \mathcal{Q}_\beta, e \geq_{\mathcal{Q}_\beta}^* x, y$.⁶*

Let $G_\kappa \subseteq P_\kappa$ be a generic. Then, in $V[G_\kappa]$, there is no fresh subsets of κ .

Remark 5.2 More restrictive results were proved in [4] for cardinals above κ . The present proof can be easily modified for higher cardinals.

Proof. Let G_κ be a generic subset of P_κ .

We would like to show that there is no fresh subset of κ in $V[G_\kappa]$.

Suppose otherwise. Work in V . Let $\mathcal{A}$ be a name of such subset and let $\mathcal{f}$ be a name of the characteristic function of A . Fix some $p \in G_\kappa$ which forces this.

Let $\zeta < \kappa$ be an ordinal for which a new subset is added in the extension from V to $V[G]$. Such ζ exists, since the forcings Q_α (for $\alpha < \kappa$) have cardinality below κ , and at least one of them is non-trivial. By increasing if necessary, we can assume that ζ is a singular cardinal. Then $\langle P_\kappa \setminus \zeta, \leq^* \rangle$ is more than ζ -closed. Thus, there exists a condition $q \in G_\kappa \restriction P_\zeta$ which forces that a new subset is added to ζ . For simplicity, assume that $p \restriction \zeta$ forces this.

Let I be a subset of κ which consists of singular cardinals τ such that for every $\rho < \tau, |P_\rho| < \tau$.

As before, we divide into two cases:

Case 1. There exists $\mu \in (\zeta, \kappa) \cap I$ and a condition $p^* \in P_\mu, p^* \geq p \restriction \mu$ which forces that the following property holds:

$$\begin{aligned} \exists p' \in G_\mu \exists s \in P_\kappa \setminus \mu, s \geq p \setminus \mu \forall r \geq^* s, r(\mu) = s(\mu), r(\mu^+) = s(\mu^+) \exists \xi < \kappa \exists r_0, r_1 \geq^* r, \\ r_0(\mu) = r_1(\mu) = r(\mu), r_0(\mu^+) = r_1(\mu^+) = r(\mu^+), \end{aligned}$$

$$V \models (p' \restriction r_0 \parallel \mathcal{f} \restriction \xi, p' \restriction r_1 \parallel \mathcal{f} \restriction \xi), \text{ and the decisions are different,}$$

where G_μ denotes the canonical name for the generic set for P_μ .

By extending p^* , if necessary, we can decide the value of p' in the statement above, and thus assume that $p^* \geq p'$. Let $\mathcal{s}$ be a P_μ -name for s from the above property, and assume that this is forced by p^* .

Note that here we do not assume μ^{++} -completeness of the direct order. Additional requirements are included in order to compensate this. Still, the treatment of this case repeats completely Case 1 of 3.1.

⁶Note that if $\leq = \leq^*$, then this holds trivially. Prikry, Magidor, Radin forcings, their supercompact versions, etc., have this property. Actually, any reasonable forcing of this type has this property.

Case 2. For every $\mu \in (\zeta, \kappa) \cap I$, every condition in P_μ stronger than $p \restriction \mu$ forces (and so, $p \restriction \mu$ forces) that–

$$\begin{aligned} \forall p' \in G_\mu \ \forall s \in P \setminus \mu, s \geq p \setminus \mu \exists r \geq^* s, r(\mu) = s(\mu), r(\mu^+) = s(\mu^+) \ \forall \xi < \kappa \ \forall r_0, r_1 \geq^* r, \\ r_0(\mu) = r_1(\mu) = r(\mu), r_0(\mu^+) = r_1(\mu^+) = r(\mu^+), \\ V \models \text{ If } p' \restriction r_0 \parallel \dot{A} \cap \xi \text{ and } p' \restriction r_1 \parallel \dot{A} \cap \xi \text{ then the decisions are the same.} \end{aligned}$$

For every $\mu \in (\zeta, \kappa) \cap I$, define (in $V[G_\mu]$)

$$\begin{aligned} e'(\mu) = \{r \in P_\kappa \setminus \mu : \forall p' \in G_\mu \ \forall \xi < \kappa \ \forall r_0, r_1 \geq^* r, \\ r_0(\mu) = r_1(\mu) = r(\mu), r_0(\mu^+) = r_1(\mu^+) = r(\mu^+), \\ V \models \text{ If } p' \restriction r_0 \parallel \dot{f} \restriction \xi \text{ and } p' \restriction r_1 \parallel \dot{f} \restriction \xi \text{ then the decisions are the same.}\} \end{aligned}$$

Claim 4 For every $\mu \in (\zeta, \kappa) \cap I$,

$e'(\mu)$ is $\leq^*$ -dense in $P \setminus \mu$ (above $p \setminus \mu$).⁷

Proof. Note that $|G_\mu| \leq \mu^+$, the forcing $\langle P_\kappa \setminus \mu^{++} \rangle$ is μ^{++} -closed and the coordinates μ, μ^+ do not change. Hence, for every given $r' \in P_\kappa \setminus \mu$, we can construct a $\leq^*$ -increasing sequence of a length $|G_\mu|$ of conditions stronger than r' , which takes care of each $p' \in G_\mu$. Then its upper bound will be in $e(\mu)$.

□ of the claim.

Given a generic $G_\kappa \subseteq P_\kappa, p \in G_\kappa$, define in $V[G_\kappa]$,

$$S = \{\xi < \kappa \mid \xi \text{ is limit ordinal and } \exists t \geq^* p, t \in G_\kappa \text{ such that } t \restriction \xi \Vdash_{P_\xi} t \setminus \xi \parallel_{P_\kappa \setminus \xi} \dot{A} \cap \xi\}.$$

For every $\xi \in S$, fix some $t^\xi \in G_\kappa$ such that $t^\xi \geq^* p$ and $t^\xi \restriction \xi \Vdash_{P_\xi} t^\xi \setminus \xi \parallel_{P_\kappa \setminus \xi} \dot{A} \cap \xi$. So, there is $p^\xi \in G \restriction P_\xi, p^\xi \geq t^\xi \restriction \xi, (p^\xi) \restriction t^\xi \setminus \xi \Vdash \dot{A} \cap \xi = a_\xi$, for some $a_\xi \in V$. Then there is a finite $b^\xi \subseteq \xi$ such that $p^\xi \setminus b^\xi \geq^* p \restriction (\xi \setminus b^\xi)$.

The argument of Lemma 4.7 applies without changes in the present situation, and shows that S is stationary (in $V[G_\kappa]$).

Then we can find a stationary subset S' of S and a finite b such that for every $\xi \in S'$, $b^\xi = b$.

Now we can freeze $p^\xi \restriction \max(b)^{++}$.

If the following holds,

⁷It need not be open.

($\aleph$): For every $\beta < \kappa$, for every $s, t, r \in Q_\beta$, if $s \leq_{Q_\beta}^* t, r$, then there is $e \in Q_\beta, e \geq_{Q_\beta}^* t, r$.⁸

then it is easy to conclude that $A \in V$, which is impossible.

Deal with the general situation. Let $\mu \in I$ be a cardinal $\geq \max(b)$. Freeze $p^\xi \restriction \mu^{++} + 1$.

Then for every $\xi, \rho \in S'$,

$$p^\xi(\mu) = p^\rho(\mu) \text{ and } p^\xi(\mu^+) = p^\rho(\mu^+).$$

There is $s \in G_\kappa, s \geq p, s \restriction \mu^{++} \geq p^\xi \restriction \mu^{++}, \xi \in S'$, such that

$s \Vdash \forall \xi \in S' \exists t^\xi \geq^* p, t^\xi \in G_\kappa \quad \exists p^\xi \in G_\kappa \restriction P_\xi \quad b_\xi = b$ which has the properties above.

Consider the set $e'(\mu)$ defined above. By the claim, it is $\leq^*$ -dense subset of $P_\kappa \setminus \mu$, in $V[G_\mu]$ above $p \restriction \mu$. In particular there is $r \in P_\kappa \setminus \mu, r \geq^* s \restriction \mu$ such that for every $p' \in G_\mu$, for every $\xi < \kappa$ and for every $r_0, r_1 \geq^* r$

$$V \models \text{If } p' \restriction r_0 \restriction A \cap \xi \text{ and } p' \restriction r_1 \restriction A \cap \xi \text{ then the decisions are the same.}$$

Pick now a generic $G' \subseteq P_\kappa$ with $G_\mu = G' \restriction P_\mu$ and $s \restriction \mu \cap r \in G'$.

We assumed the following,

($\beth$): For every $\beta < \kappa$, for every $x, y, z \in Q_\beta$, if $z \leq_{Q_\beta}^* x, y$ and x, y are compatible according to $\leq_{Q_\beta}$, then they are compatible according to $\leq_{Q_\beta}^*$, i.e. there is $e \in Q_\beta, e \geq_{Q_\beta}^* x, y$.

Now, P_κ is a full support iteration. So, ($\beth$) implies -

($\beth$): For every $\beta < \kappa$, for every $u, v, w \in P_\kappa$, if $w \leq_{P_\kappa}^* u, v$ and u, v are compatible according to $\leq_{P_\kappa}$, then they are compatible according to $\leq_{P_\kappa}^*$.⁹

It follows, in this case, that $A \in V$. Contradiction.

□

⁸For example if P_κ is the Magidor iteration of Prikry forcings. Also note that we do not to split into Cases 1,2 in this type of situation.

⁹It need not be the case for Easton or non-stationary support, since then the support of w may be strictly smaller than those of u, v , and u, v may disagree on a common coordinate outside the support of w .

References

- [1] Omer Ben-Neria and Spencer Unger, Homogeneous changes in cofinality with applications to HOD, *Journal Math. logic*, vol 17 (2017) no.2
- [2] Sy Freidman and Menachem Magidor, The JSL, vol 74, (2009), pp. 1069-1080.
- [3] Moti Gitik. Prikry-type forcings. In *Handbook of set theory*, pages 1351–1447. Springer, 2010.
- [4] Moti Gitik and Eyal Kaplan. On restrictions of ultrafilters from generic extensions to ground models. *The Journal of Symbolic Logic*, pages 1–31, 2021.
- [5] Moti Gitik and Eyal Kaplan. Non-stationary support iterations of prikry forcings and restrictions of ultrapower embeddings to the ground model. *Annals of Pure and Applied Logic*, 174(1):103164, 2023.
- [6] Moti Gitik and Eyal Kaplan. On Easton support iteration of Prikry-type forcing notions. *arXiv preprint arXiv:2301.12421*, 2023.
- [7] Joel David Hamkins. Gap forcing. *Israel Journal of Mathematics*, 125(1):237–252, 2001.
- [8] Eyal Kaplan. The Magidor iteration and restrictions of ultrapowers to the ground model. *arXiv preprint arXiv:2202.04980*, 2022.
- [9] Eyal Kaplan, PhD Thesis, Tel Aviv University, 2023.
- [10] Menachem Magidor, How large is the first strongly compact cardinal? Or study of identity crises, *Ann. Math. Logic*, vol. 10 (1976), pp. 33-57.
- [11] Azriel Levy and Robert M. Solovay. Measurable cardinals and the continuum hypothesis. *Israel Journal of Mathematics* 5:234-248, 1967.