ON THE LOGARITHMIC SLICE FILTRATION

FEDERICO BINDA, DOOSUNG PARK, AND PAUL ARNE ØSTVÆR

ABSTRACT. We consider slice filtrations in logarithmic motivic homotopy theory. Our main results establish conjectured compatibilities with the Beilinson, BMS, and HKR filtrations on (topological, log) Hochschild homology and related invariants. In the case of perfect fields admitting resolution of singularities, we show that the slice filtration realizes the BMS filtration on the *p*-completed topological cyclic homology. Furthermore, the motivic trace map is compatible with the slice and BMS filtrations, yielding a natural morphism from the motivic slice spectral sequence to the BMS spectral sequence. Finally, we consider the Kummer étale hypersheafification of logarithmic K-theory and show that its very effective slices compute Lichtenbaum étale motivic cohomology.

1. INTRODUCTION

Logarithmic motives provide a framework for studying non- \mathbb{A}^1 -invariant cohomology theories in arithmetic geometry [11], [12], [13], [14], [15], [16], [17], [38], [47]. This paper introduces logarithmic analogs of slice filtrations in motivic homotopy theory defined by Voevodsky [49] and Spitzweck-Østvær [48]. The slice perspective produces explicit calculations, see [29] for a survey, and remains a powerful tool in homotopy theory following Hill-Hopkins-Ravenel's solution of the Kervaire invariant one problem [27]. Examples of applications of motivic slices include a proof of the Milnor conjecture on quadratic forms in [43] and calculations of universal motivic invariants in [45], [46]. One of the main purposes of this paper is to compare the logarithmic slice filtration with the Bhatt-Morrow-Scholze filtration on topological Hochschild homology and refinements [9].

Suppose that k is a perfect field of characteristic p and A is a smooth k-algebra. Bhatt, Morrow, and Scholze [9, §1.4] expected that their filtrations on topological Hochschild homology $\text{THH}(A; \mathbb{Z}_p)$ (and the related theories $\text{TC}^-(A; \mathbb{Z}_p)$, $\text{TP}(A; \mathbb{Z}_p)$, and $\text{TC}(A; \mathbb{Z}_p)$) afford a precise relation with filtrations on algebraic K-theory K(A), see [24], [31], via the trace maps

 $K(A) \to THH(A; \mathbb{Z}_p), TC^{-}(A; \mathbb{Z}_p), TP(A; \mathbb{Z}_p), TC(A; \mathbb{Z}_p).$

In the literature, these filtrations are often called "motivic filtrations". Mathew verified this expectation in [36, p. 4] by showing that the filtrations are compatible using Postnikov towers in (pro-)Nisnevich and (pro-)étale topologies. Moreover, Mathew raised a deeper question about the existence of a common construction, realizing both filtrations.

This paper aims to explore the motivic properties of these filtrations using logarithmic motivic homotopy theory developed in [17] and provide a first positive answer in this direction. To explain this, let us recall the main idea behind Voevodsky's slice filtration in \mathbb{A}^1 -homotopy theory. In algebraic topology, a standard method to understand a spectrum E is to study its Postnikov tower, i.e., how it is built out of topological Eilenberg-MacLane spectra. Sending E to its n - 1-connected cover $E^{(n)}$ defines a functor from SH to the full subcategory $\Sigma^n SH_{\geq 0}$ of n - 1-connected spectra, that is right adjoint to the obvious inclusion. Replacing the category $\Sigma^n SH_{\geq 0}$ with the category $\Sigma^{2n,n}SH(S)^{\text{eff}}$ measuring \mathbb{G}_m -effectivity, Voevodsky defined a motivic analog of the Postnikov tower, known as the slice tower

$$f_{n+1}E \to f_nE \to f_{n-1}E \to \dots \to E$$

Date: April 10, 2024.

for any motivic spectrum E. When $E = \mathbf{KGL}$ is the spectrum representing algebraic K-theory, the induced spectral sequence is the motivic Atiyah-Hirzebruch spectral sequence of Levine-Voevodsky [32], [50].

Since (topological) Hochschild homology and its cousins are not \mathbb{A}^1 -homotopy invariant (in fact, the \mathbb{A}^1 -localization of THH is trivial, see, e.g., [21]), there is no motivic spectrum in SH(S) representing them and thus Voevodsky's slice machinery is inapplicable. In [17], we developed an extension of SH(S) using the language of log schemes and proved that there are indeed \mathbb{P}^1 -spectra representing THH and variants. The HKR filtration, the Beilinson filtration, and the Bhatt-Morrow-Scholze filtration on HH, HC⁻, THH, TC⁻ and so on are all compatible with the bounding maps defining the motivic spectra **HH**, **HC**⁻, **THH**, **TC**⁻ and relatives, giving rise to filtered \mathbb{P}^1 -motivic spectra. See Propositions 4.9 and 5.8.

In this work, we consider an analog of Voevodsky's slice filtration in logSH(S) and in the linearized variant logDA(S). We define the effective category logSH(S)^{eff} as the localizing subcategory of logSH(S) generated by \mathbb{P}^1 -suspensions of $(X, \partial X) \in \text{SmlSm}/S$, where $X \in$ Sm/S and ∂X is a strict normal crossing divisor on X. Similarly, we define the coeffective category logSH(S)^{coeff} as its right orthogonal in the obvious sense. By (de)suspension, we obtain the *n*-effective and the *m*-coeffective categories for every $n, m \in \mathbb{Z}$. Following Spitzweck-Østvær [48], we will consider the very effective version of the slice filtration in logSH(S). Rather than measuring \mathbb{G}_m -effectivity, the very effective slice filtration measures \mathbb{P}^1 -effectivity and has strong convergence properties. Our main results are the following. Below, when Fil $_{\bullet}\mathcal{F}$ is a filtration on \mathcal{F} , we set Fil $^n\mathcal{F} := \text{cofib}(\text{Fil}_{n+1}\mathcal{F} \to \mathcal{F})$ for $n \in \mathbb{Z}$.

Theorem 1.1 (See Theorem 4.14). Let R be any ring, and let $S \to \text{Spec}(R)$ be a smooth R-scheme. For $n \in \mathbb{Z}$, the spectra

$$\operatorname{Fil}_{\operatorname{HKB}}^{n}\mathbf{HH}(-/R), \operatorname{Fil}_{\operatorname{B}}^{n}\mathbf{HC}^{-}(-/R), \operatorname{Fil}_{\operatorname{B}}^{n}\mathbf{HP}(-/R)$$

are n-co-very effective, that is, they belong to the subcategory $\log DA(S)_{\leq n}^{\text{veff}}$ of Construction 2.3. In particular, there are canonical morphisms

$$\mathbf{\tilde{f}}_n \mathbf{H} \mathbf{H}(-/R) \simeq \Sigma^{2n,n} \mathbf{\tilde{f}}_0 \mathbf{H} \mathbf{H}(-/R) \to \mathrm{Fil}_n^{\mathrm{HKR}} \mathbf{H} \mathbf{H}(-/R),$$

and similarly for \mathbf{HC}^- and \mathbf{HP} , where $\tilde{\mathbf{f}}_n$ denotes the nth very effective slice functor in $\log \mathrm{DA}(S)$.

Theorem 1.2 (See Theorem 5.16). Let S be the spectrum of a p-adic quasi-syntomic ring R. For $n \in \mathbb{Z}$, the filtered spectra

$$\operatorname{Fil}_{\operatorname{BMS}}^{n}\operatorname{\mathbf{THH}}(-;\mathbb{Z}_{p}), \operatorname{Fil}_{\operatorname{BMS}}^{n}\operatorname{\mathbf{TC}}^{-}(-;\mathbb{Z}_{p}), \operatorname{Fil}_{\operatorname{BMS}}^{n}\operatorname{\mathbf{TP}}(-;\mathbb{Z}_{p}), \operatorname{Fil}_{\operatorname{BMS}}^{n}\operatorname{\mathbf{TC}}(-;\mathbb{Z}_{p})$$

are in logDA(S, \mathbb{Z}_p)^{veff} $\leq n$. In particular, there are canonical morphisms

$$\tilde{\mathrm{f}}_{n}\mathbf{THH}(-;\mathbb{Z}_{p})\simeq\Sigma^{2n,n}\tilde{\mathrm{f}}_{0}\mathbf{THH}(-;\mathbb{Z}_{p})\rightarrow\mathrm{Fil}_{n}^{\mathrm{BMS}}\mathbf{THH}(-;\mathbb{Z}_{p}),$$

and similarly for TC^- , TP and TC.

Assume now that S = Spec(k) is a field of positive characteristic p, admitting resolution of singularities. Since topological cyclic homology and variants are, in any case, étale sheaves, we can consider an étale variant of the very effective slice filtration in the subcategory spanned by *p*-completed objects in logDA(k). Our result is then the following.

Theorem 1.3 (See Theorem 7.5). Let k be a perfect field admitting resolution of singularities. Then the induced morphism

$$\tilde{\mathbf{f}}_i^{\text{k\acute{e}t}}\mathbf{TC}(-;\mathbb{Z}_p) \to \mathrm{Fil}_i^{\mathrm{BMS}}\mathbf{TC}(-;\mathbb{Z}_p)$$

is an equivalence in the ∞ -category of p-complete Kummer étale motives $\log DA^{\wedge}_{k\acute{e}t}(k)^{\wedge}_{p}$ for every integer i. Here, we denote by $\tilde{f}^{k\acute{e}t}_{i}$ the ith very effective cover in $\log DA^{\wedge}_{k\acute{e}t}(k)^{\wedge}_{p}$. In other words, despite having very different origins, we can identify the slice and the BMS filtration. We see this as an answer to Mathew's question for $TC(-;\mathbb{Z}_p)$, i.e., the slice filtrations realize the two "motivic filtrations" on K and $TC(-;\mathbb{Z}_p)$. We remark that the proof of the above theorem crucially relies on Geisser-Levine [26], together with a motivic version of the prismatic-crystalline comparison [11].

It a natural question to compare the slice towers in $\log SH(S)$ and SH(S). When the base S is the spectrum of a field that admits resolution of singularities, we can combine the above results with the trace methods using the motivic (logarithmic) cyclotomic trace constructed in [17, Corollary 8.6.2], and obtain the following.

Theorem 1.4. Let k be a perfect field of characteristic p > 0 admitting resolution of singularities.

(1) The log motivic trace maps in $\log SH(k)$

$$\begin{split} \mathbf{KGL} &\to \mathbf{THH}(-;\mathbb{Z}_p), \quad \mathbf{KGL} \to \mathbf{TC}^-(-;\mathbb{Z}_p), \\ \mathbf{KGL} \to \mathbf{TP}(-;\mathbb{Z}_p), \quad \mathbf{KGL} \to \mathbf{TC}(-;\mathbb{Z}_p) \end{split}$$

are compatible with the slice filtrations on the left-hand sides and the BMS filtrations on the right-hand sides.

(2) The graded pieces of the log motivic trace maps are identified with

$$\begin{split} \mathbf{M}\mathbb{Z}(i)[2i] &\to \mathrm{Fil}_{\mathrm{conj}}^{i} \mathbf{\Omega}_{/k}[2i], \\ \mathbf{M}\mathbb{Z}(i)[2i] &\to \mathrm{Fil}_{i}^{\mathrm{N}} \mathbf{W} \mathbf{\Omega}_{/k}[2i], \\ \mathbf{M}\mathbb{Z}(i)[2i] &\to \mathbf{W} \mathbf{\Omega}_{/k}[2i], \\ \mathbf{M}\mathbb{Z}(i)[2i] &\to \mathbf{W} \mathbf{\Omega}_{/k,\log}^{i}[i]. \end{split}$$

(3) The graded pieces of the log trace maps yield natural morphisms

$$R\Gamma_{\rm mot}(X - \partial X, \mathbb{Z}(i)) \to R\Gamma_{\rm s\acute{e}t}(X, \tau^{\leq i}\Omega_{/k}),$$

$$R\Gamma_{\rm mot}(X - \partial X, \mathbb{Z}(i)) \to R\Gamma_{\rm s\acute{e}t}(X, {\rm Fil}_{i}^{\rm N}W\Omega_{/k}),$$

$$R\Gamma_{\rm mot}(X - \partial X, \mathbb{Z}(i)) \to R\Gamma_{\rm s\acute{e}t}(X, W\Omega_{/k}),$$

$$R\Gamma_{\rm mot}(X - \partial X, \mathbb{Z}(i)) \to R\Gamma_{\rm spro\acute{e}t}(X, W\Omega_{/k, \log}^{i}[-i]),$$

where the left-hand side denotes Suslin-Friedlander motivic cohomology with integral coefficients.

In light of Levine's comparison between the homotopy conveau tower and the slice tower [31, Theorem 9.0.3], we obtain in particular that for $X \in \text{SmlSm}/k$, the log trace maps

$$\begin{aligned} \mathrm{K}(X - \partial X) &\to \mathrm{THH}(X; \mathbb{Z}_p), \quad \mathrm{K}(X - \partial X) \to \mathrm{TC}^-(X; \mathbb{Z}_p), \\ \mathrm{K}(X - \partial X) &\to \mathrm{TP}(X; \mathbb{Z}_p), \quad \mathrm{K}(X - \partial X) \to \mathrm{TC}(X; \mathbb{Z}_p) \end{aligned}$$

are compatible with the homotopy coniveau filtrations on the left-hand sides and the BMS filtrations on the right-hand sides. As an immediate corollary, we obtain a natural morphism from the motivic Atiyah-Hirzebruch spectral sequence [31, §11.3]

$$E_2^{pq} = H^{i-j}_{\text{mot}}(X - \partial X, \mathbb{Z}(-j)) \Rightarrow \pi_{-i-j} \mathcal{K}(X - \partial X)$$

to the log version of the BMS spectral sequence [9, Theorem 1.12(5)], (see [13, Theorem 1.3] for the log case)

$$E_2^{ij} = H_{\text{syn}}^{i-j}(X, \mathbb{Z}_p(-j)) \Rightarrow \pi_{-i-j} \text{TC}(X; \mathbb{Z}_p)$$

for $X \in \text{SmlSm}/k$ and in particular for $X \in \text{Sm}/k$. In particular, we get directly that the morphism

$$H^i_{\text{mot}}(X - \partial X, \mathbb{Z}(j)) \to H^i_{\text{syn}}(X - \partial X, \mathbb{Z}_p(j))$$

considered in [9, Corollary 8.21], [5, Theorem 6.15] (using Geisser-Levine to identify p-adic motivic cohomology) factors canonically through the log syntomic cohomology $H^i_{\text{syn}}(X, \mathbb{Z}_p(j))$. Note that the input of [26] is not necessary to obtain the map: it is just a consequence of the compatibility between the slice filtration and the BMS filtration, together with [31, Theorem 6.4.2]. For p-adic motivic cohomology with \mathbb{Q}_p -coefficients, a similar refinement has been considered by Ertl and Niziol in [23] (see Remark 7.11). The comparison between motivic cohomology and syntomic cohomology (the non-log case) is also considered in the recent work of Elmanto-Morrow [22].

The non-topological counterpart (for classical cyclic, periodic and Hochschild homology) also holds.

Theorem 1.5. Let k be a perfect field of admitting resolution of singularities (e.g., char k = 0).

(1) The log motivic trace maps in $\log SH(k)$

$$\mathbf{KGL} \to \mathbf{HH}(-/k), \quad \mathbf{KGL} \to \mathbf{HC}^{-}(-/k), \quad \mathbf{KGL} \to \mathbf{HP}(-/k)$$

are compatible with the slice filtrations on the left-hand sides and the HKR filtration on **HH** and Beilinson filtrations on \mathbf{HC}^- and \mathbf{HP} .

(2) The graded pieces of the log motivic trace maps are identified with

$$\begin{split} \mathbf{M}\mathbb{Z}(n)[2i] &\to \mathbf{\Omega}_{/k}^{i}[2i],\\ \mathbf{M}\mathbb{Z}(i)[2i] &\to \mathbf{\Omega}_{/k}^{\geq i}[2i],\\ \mathbf{M}\mathbb{Z}(i)[2i] &\to \mathbf{\Omega}_{/k}[2i]. \end{split}$$

(3) The graded pieces of the log trace maps yield natural morphisms

$$R\Gamma_{\rm mot}(X - \partial X, \mathbb{Z}(i)) \to R\Gamma_{\rm sNis}(X, \mathbf{\Omega}_{/k}^{i}),$$

$$R\Gamma_{\rm mot}(X - \partial X, \mathbb{Z}(i)) \to R\Gamma_{\rm sNis}(X, \mathbf{\Omega}_{/k}^{\geq i}),$$

$$R\Gamma_{\rm mot}(X - \partial X, \mathbb{Z}(i)) \to R\Gamma_{\rm sNis}(X, \mathbf{\Omega}_{/k}).$$

We remark that the compatibility between the slice filtration and the BMS filtration along the trace map has been established using completely different methods (see again [22], and the upcoming work of Bachmann-Elmanto-Morrow). Our results give a refinement of such statements (at least over a field), since the trace map further factors through the log invariants, in a way that is compatible with the motivic filtrations on the source and on the target.

Finally, in the last section of the paper we consider the slice filtration in the (hypercomplete) Kummer étale version $\log SH^{\wedge}_{k\acute{e}t}(k)$ of $\log SH(k)$.

Theorem 1.6 (See Theorem 7.7). Let k be a perfect field admitting resolution of singularities, and assume that the étale cohomological dimension of k is finite.

There are natural equivalences

 $\mathbf{s}_{i}^{\mathrm{k\acute{e}t}} L_{\mathrm{k\acute{e}t}} \mathbf{KGL} \simeq \tilde{\mathbf{s}}_{i}^{\mathrm{k\acute{e}t}} L_{\mathrm{k\acute{e}t}} \mathbf{KGL} \simeq L_{\mathrm{k\acute{e}t}} \mathbf{s}_{i} \mathbf{KGL} \simeq L_{\mathrm{k\acute{e}t}} \tilde{\mathbf{s}}_{i} \mathbf{KGL} \simeq \Sigma^{2i,i} L_{\mathrm{k\acute{e}t}} \mathbf{MZ}$

in logSH^{\wedge}_{két}(k). Moreover, $L_{két}$ **M**Z represents Lichtenbaum étale motivic cohomology [37, Lecture 10] when restricted to Sm/k.

For $X \in \text{SmlSm}/k$ note moreover that there is a natural equivalence

$$\operatorname{nap}_{\operatorname{logSH}_{\operatorname{het}}^{\wedge}(k)}(\Sigma^{\infty}X_{+}, L_{\operatorname{k\acute{e}t}}\mathbf{KGL}) \simeq L_{\operatorname{k\acute{e}t}}\mathbf{K}(X),$$

where the right hand-side is the Kummer étale hypersheafification of the functor $X \mapsto K(X - \partial X)$ for $X \in \text{Sm}/k$. In particular, for $X \in \text{Sm}/k$ is just étale K-theory.

Remark 1.7 (On resolution of singularities). For the K-theoretic applications we have assumed to be working over a field k admitting resolution of singularities. This is used in the explicit computation of the right adjoint ω^* from Voevodsky's motives to log motives, that on a smooth k-scheme X satisfies $\omega^*(X) = (\overline{X}, \partial X)$ where \overline{X} is a smooth proper compactification of X, with normal crossing boundary ∂X . This simple formula allows us, for example, to identify $\omega^* \mathbf{KGL}$ as well as $\omega^* \mathbf{MZ}$, and to show that ω^* commutes with taking (very effective) slices. Using that, we can directly invoke the known results in $\mathrm{SH}(k)$. We believe, however, that the compatibility between ω^* and f_n is general, so the assumption on resolution of singularities should be superfluous. One would expect that the analog of Levine's computation holds for the log K-theory spectrum **logKGL** defined

1.8. Outline of the proofs. Let us discuss the main idea of the proof of Theorems 1.1 and 1.2. In [17], we constructed motivic spectra in $\log SH(S)$ representing logarithmic Hochschild, periodic, and cyclic homology. Such spectra are Bott periodic spectra, in the sense that the strict Nisnevich sheaf HH(-) of spectra on lSch/S satisfies $\Omega_{\mathbb{P}^1}HH \simeq HH$ (and similarly for HP(-) and HC⁻(-)). The compatibility of the \mathbb{P}^1 -bundle formula with the HKR and the Beilinson filtration allows us to show (see Proposition 4.11) that we can promote the construction and obtain filtered motivic \mathbb{P}^1 -spectra, satisfying the equivalence

in $[17, \S6.5]$. We plan to address this question in future work.

$$\operatorname{Fil}_{i}^{\operatorname{HKR}}\operatorname{HH}(-/R)(1)[2] \simeq \operatorname{Fil}_{i+1}^{\operatorname{HKR}}\operatorname{HH}(-/R)$$

and similarly for HP and HC⁻ with the Beilinson filtration, for any ring R. At this point, the co-effectivity of $\operatorname{Fil}^n_{\mathrm{HKR}}$ HH with respect to the *t*-structure given by the very effective slice tower is a direct consequence of the vanishing of the sheaf cohomology $H^{2j}(X, \Omega^j_{X/R})$ for negative j.

The topological counterpart, that is, for THH and variant, is proven by a conceptually similar method. More precisely, one shows using the \mathbb{P}^1 bundle formula that the BMS filtration on THH gives an equivalence

$$\operatorname{Fil}_{i}^{\operatorname{BMS}}\operatorname{THH}(-;\mathbb{Z}_{p})(1)[2] \simeq \operatorname{Fil}_{i+1}^{\operatorname{BMS}}\operatorname{THH}(-;\mathbb{Z}_{p}),$$

and similarly for TP, TC^- and TC, see Proposition 5.11. The co-effectivity statement can then be deduced by passing to the associated graded, reducing to vanishing in (log) prismatic cohomology. This is shown by log quasi-syntomic descent. Overall, the coeffectivity results immediately imply the existence of the natural morphisms between the very effective slice tower and the BMS, HKR, and Beilinson filtrations.

Let us now consider the natural functor $\omega^* \colon \mathrm{SH}(S) \to \mathrm{logSH}(S)$, right adjoint to the \mathbb{A}^1 -localization functor from $\mathrm{logSH}(S)$ to $\mathrm{SH}(S)$ for S a scheme. When $S = \mathrm{Spec}(k)$ is a perfect field that admits resolution of singularities, it is easy to show that ω^* commutes with the (very effective) slice filtration on both sides. Using the results in [17, §7.7] and the main result of Levine in [31, Theorems 6.4.2, 9.0.2], we can then identify the very effective slices of $\omega^* \mathbf{KGL}$ with motivic cohomology in $\mathrm{logSH}(k)$. Combining this with Theorems 1.1 and 1.2 we obtain Theorems 1.4 and 1.5.

Finally, in §7, we consider the very effective slices of Kummer étale K-theory. This allows us to consider a spectrum representing Lichtenbaum étale motivic cohomology in logSH_{két}(k): if k admits resolution of singularities, this is computed by the zeroth very effective slice of the Kummer étale sheafification of **KGL**. See Theorem 7.7. Using Geisser-Levine as input, together with the identification $\mathbf{MZ}_p^{\text{syn}}(i) \simeq L_{\text{sproét}} \mathbf{W\Omega}_{/k,\log}^i[-i]$ as motivic spectra from [11], we can further show that in the Kummer étale (or in the strict étale) *p*-complete category, we have $\operatorname{Fil}_i^{\text{BMS}} \mathbf{TC}(-; \mathbb{Z}_p) \in (\log \text{DA}_{\text{két}}^{\wedge}(k)_p^{\wedge})_{\geq i}^{\text{veff}}$. Since we have already proved in general that $\operatorname{Fil}_i^{\text{BMS}} \mathbf{TC}(-; \mathbb{Z}_p)$ is also *i*-co-very effective, we conclude.

Acknowledgements. We thank Marc Levine and Alberto Merici for inspiring discussions and Matthew Morrow for helpful comments on a preliminary version of the paper. F.B. is partially supported by the PRIN 2022 "The arithmetic of motives and L-functions" at MUR (Italy). D.P. is partially supported by the research training group GRK 2240 "Algebro-Geometric Methods in Algebra, Arithmetic and Topology." P.A.Ø. was partially supported by The European Commission – Horizon-MSCA-PF-2022 "Motivic integral *p*-adic cohomologies", the RCN Project no. 312472 "Equations in Motivic Homotopy Theory", and a Guest Professorship awarded by The Radboud Excellence Initiative.

2. SLICES FOR LOGARITHMIC MOTIVES

Throughout this section, let S be a quasi-compact quasi-separated scheme, and let τ be any topology on lSch/S generated by a family of morphisms in lSm/S. Recall that the category SmlSm/S is the full subcategory of lSm/S consisting of $X \in \text{lSm/S}$ such that $\underline{X} \in \text{Sm/S}$. By [17, Remark 2.3.19], [16, Lemma A.5.10], it can be equivalently described as the category of pairs $X = (\underline{X}, \partial X)$ such that $\underline{X} \in \text{Sm/S}, X - \partial X$ is an open subscheme of \underline{X} , and the closed complement ∂X is a strict normal crossing divisor on \underline{X} . We see $(\underline{X}, \partial X)$ as a log scheme with the compactifying log structure given by the embedding $X - \partial X \subset \underline{X}$.

We work with the Nisnevich sheaf of stable presentable symmetric monoidal ∞ -categories

$$\operatorname{Sch}^{\operatorname{op}} \to \operatorname{CAlg}(\operatorname{Pr}_{\operatorname{Mod}_{\Lambda}}^{\operatorname{L}}), \quad S \mapsto \operatorname{logSH}_{\tau}(S, \Lambda)^{\otimes}$$

given by

$$\mathrm{logSH}_{\tau}(S,\Lambda)^{\otimes} := \mathrm{Sp}_{\mathbb{P}^1}(\mathrm{Sh}_{\tau}(\mathrm{SmlSm}/S,\mathrm{Mod}_{\Lambda})[(\mathbb{P}^{\bullet},\mathbb{P}^{\bullet-1})^{-1}])^{\otimes}$$

introduced in [17], where Λ is an \mathbb{E}_{∞} -ring, and Mod_{\Lambda} denotes the ∞ -category of Λ -module objects of Sp. It admits a symmetric monoidal structure and is presentable and stable by [35, Theorems 4.5.2.1(1), 7.1.2.1, Corollary 7.1.1.5]. To simplify the notation, we will often omit the superscript \otimes . If Λ is the sphere spectrum \mathbb{S} , then we omit Λ in logSH_{τ}(S, Λ). If Λ is the Eilenberg-MacLane spectrum HR for a commutative ring R, then we set

$$\log DA_{\tau}(S) := \log SH_{\tau}(S, H\mathbb{Z})$$
 and $\log DA_{\tau}(S, R) := \log SH_{\tau}(S, HR)$

If τ is the strict Nisnevich topology, then we omit the subscript τ .

We will be using the following conventions for filtrations. Let \mathcal{F} be an object of an ∞ -category \mathcal{C} . The ∞ -category of filtered objects of \mathcal{C} is $\operatorname{Fun}(\mathbb{Z}^{op}, \mathcal{C})$, where \mathbb{Z}^{op} is the category consisting of the integers such that $\operatorname{Hom}_{\mathbb{Z}^{op}}(i, j)$ is * if $i \geq j$ and \emptyset if i < j. A filtered object $\mathcal{F}(-)$ of \mathcal{C} is complete if $\lim_{i} \mathcal{F}(i) \simeq 0$. The underlying object of $\mathcal{F}(-)$ is $\mathcal{F}(-\infty) := \operatorname{colim}_i \mathcal{F}(i)$. The filtration $\mathcal{F}(-)$ on $\mathcal{F}(-\infty)$ is exhaustive. We write

$$\begin{aligned} \operatorname{Fil}_{i}\mathcal{F} &:= \mathcal{F}(i),\\ \operatorname{Fil}^{i}\mathcal{F} &:= \operatorname{cofib}(\operatorname{Fil}_{i+1}\mathcal{F} \to \mathcal{F}),\\ \operatorname{gr}_{i}\mathcal{F} &:= \operatorname{cofib}(\operatorname{Fil}_{i+1}\mathcal{F} \to \operatorname{Fil}_{i}\mathcal{F}). \end{aligned}$$

Observe that we have $\operatorname{gr}_i \mathcal{F} \simeq \operatorname{fib}(\operatorname{Fil}^i \mathcal{F} \to \operatorname{Fil}^{i-1} \mathcal{F}).$

The slice filtration on SH(S) is due to Voevodsky [49, §2]. We have its direct analog on logSH(S) as follows.

Construction 2.1. Let $\log SH_{\tau}(S, \Lambda)^{eff} \subset \log SH_{\tau}(S, \Lambda)$ be the category of *effective spectra*, that is, the full stable ∞ -subcategory of $\log SH_{\tau}(S, \Lambda)$ generated under colimits by $\Sigma^{n,0}\Sigma_{\mathbb{P}^1}^{\infty}X_+$ for $X \in SmlSm/S$ and integer n (equivalently, it is the localizing subcategory as a triangulated generated by $\Sigma_{\mathbb{P}^1}^{\infty}X_+$ for varying X, which means the smallest full triangulated subcategory containing those objects and closed under direct sums).

For every integer $i \in \mathbb{Z}$, let $\Sigma^i_{\mathbb{P}^1} \log \operatorname{SH}_{\tau}(S, \Lambda)^{\operatorname{eff}} \subset \log \operatorname{SH}_{\tau}(S, \Lambda)$ be the full subcategory of $\operatorname{logSH}_{\tau}(S, \Lambda)$ generated under colimits by $\Sigma^i_{\mathbb{P}^1} E$ for $E \in \operatorname{logSH}_{\tau}(S, \Lambda)^{\operatorname{eff}}$. These categories form an exhaustive filtration of $\operatorname{logSH}_{\tau}(S, \Lambda)$ by subcategories

$$\cdots \subset \Sigma^{i+1}_{\mathbb{P}^1} \mathrm{logSH}_{\tau}(S, \Lambda)^{\mathrm{eff}} \subset \Sigma^{i}_{\mathbb{P}^1} \mathrm{logSH}_{\tau}(S, \Lambda)^{\mathrm{eff}} \subset \cdots \subset \mathrm{logSH}_{\tau}(S, \Lambda)^{\mathrm{eff}}$$

To simplify the notation, we will write $\log \operatorname{SH}_{\tau}(S, \Lambda)^{\operatorname{eff}}_{\geq i}$ for $\Sigma^{i}_{\mathbb{P}^{1}} \log \operatorname{SH}_{\tau}(S, \Lambda)^{\operatorname{eff}}$. Let $\log \operatorname{SH}_{\tau}(S, \Lambda)^{\operatorname{eff}}_{\leq i}$ be the category of *i-coeffective spectra*, that is the full subcategory of $\log \operatorname{SH}_{\tau}(S, \Lambda)$ spanned by those objects \mathcal{G} such that $\operatorname{Hom}_{\log \operatorname{SH}_{\tau}(S, \Lambda)}(\mathcal{F}, \mathcal{G}) \simeq 0$ for every $\mathcal{F} \in \log \operatorname{SH}_{\tau}(S, \Lambda)^{\operatorname{eff}}_{\geq i}$.

The inclusion $\iota_i \colon \Sigma^i_{\mathbb{P}^1} \log \operatorname{SH}_{\tau}(S, \Lambda)^{\operatorname{eff}} \subset \log \operatorname{SH}_{\tau}(S, \Lambda)$ preserves colimits and hence admits a right adjoint \mathbf{r}_i . Let $\mathbf{f}_i^{\tau} \to \operatorname{id}$ be the counit of this adjunction pair, and let \mathbf{f}_{τ}^{i-1} be the cofiber of this natural transformation. Observe that the essential image of \mathbf{f}_{τ}^i is in $\Sigma^i_{\mathbb{P}^1} \log \operatorname{SH}_{\tau}(S, \Lambda)^{\operatorname{eff}}$. We set $\mathbf{s}_i^{\tau} := \operatorname{cofib}(\mathbf{f}_{i+1}^{\tau} \to \mathbf{f}_i^{\tau})$.

We also remark that the inclusion ι_i preserves compact objects. Thus, its right adjoint \mathbf{r}_i preserves filtered colimits [34, Proposition 5.5.7.2], and hence the composite $\mathbf{f}_i = \mathbf{r}_i \iota_i$ preserves filtered colimits too.

We omit τ in f_i^{τ} and f_{τ}^i when $\tau =$ sNis.

Remark 2.2. We note that $\log SH(S)^{\text{eff}}$ is a stable symmetric monoidal ∞ -subcategory of $\log SH(S)$.

We also consider the log version of the very effective slice filtration following Spitzweck-Østvær [48, §5].

Construction 2.3. Let $\log \operatorname{SH}_{\tau}(S, \Lambda)^{\operatorname{veff}}$ be the smallest full subcategory of $\log \operatorname{SH}_{\tau}(S, \Lambda)$ containing $\Sigma_{\mathbb{P}^1}^{\infty} X_+$, closed under colimits and closed under extensions in the sense that if $X \to Y \to Z$ is a cofiber sequence in $\operatorname{logSH}_{\tau}(S, \Lambda)$ with X and Z in $\operatorname{logSH}_{\tau}(S, \Lambda)^{\operatorname{veff}}$, then also $Y \in \operatorname{logSH}_{\tau}(S, \Lambda)^{\operatorname{veff}}$. We note that $\operatorname{logSH}_{\tau}(S, \Lambda)^{\operatorname{veff}}$ is a symmetric monoidal ∞ -subcategory of $\operatorname{logSH}_{\tau}(S, \Lambda)$, but unlike $\operatorname{logSH}_{\tau}(S, \Lambda)^{\operatorname{eff}}$ it is not stable.

We can consider the colimit-preserving inclusions (we omit τ and Λ for simplicity)

$$\Sigma_{\mathbb{P}^1}^{i+1} \mathrm{logSH}(S)^{\mathrm{veff}} \subset \Sigma_{\mathbb{P}^1}^{i} \mathrm{logSH}(S)^{\mathrm{veff}} \subset \Sigma_{\mathbb{P}^1}^{i-1} \mathrm{logSH}(S)^{\mathrm{veff}} \subset \cdots \subset \mathrm{logSH}(S)$$

giving rise via their right adjoints to functorial filtrations

$$\tilde{\mathbf{f}}_{i+1}^{\tau} E \to \tilde{\mathbf{f}}_{i}^{\tau} E \to \tilde{\mathbf{f}}_{i-1}^{\tau} E \to \dots \to E$$

where each $\tilde{\mathbf{f}}_i^{\tau}$ is the right adjoint to the inclusion $\Sigma_{\mathbb{P}^1}^i \log \operatorname{SH}_{\tau}(S, \Lambda) \subset \log \operatorname{SH}_{\tau}(S, \Lambda)$. The very effective slice functors $\tilde{\mathbf{s}}_i^{\tau} \colon \log \operatorname{SH}_{\tau}(S, \Lambda) \to \log \operatorname{SH}_{\tau}(S, \Lambda)$ are then defined by the cofiber sequence

$$\tilde{\mathbf{f}}_{i+1}^{\tau} \to \tilde{\mathbf{f}}_{i}^{\tau} \to \tilde{\mathbf{s}}_{i}^{\tau}$$

Note that one can effectively "compute" the *i*th slice as $\tilde{s}_i^{\tau} E \simeq \Sigma_{\mathbb{P}^1}^i \tilde{s}_0^{\tau} E$ for every motivic spectrum E that satisfies Bott periodicity $\Sigma_{\mathbb{P}^1} E \simeq E$.

We omit τ in f_i^{τ} and f_{τ}^i when $\tau =$ sNis.

To simplify the notation, we will write $\log \operatorname{SH}_{\tau}(S, \Lambda)_{\geq i}^{\operatorname{veff}}$ for $\Sigma_{\mathbb{P}^1}^i \log \operatorname{SH}_{\tau}(S, \Lambda)^{\operatorname{veff}}$. We define for every integer *i* the category $\log \operatorname{SH}_{\tau}(S, \Lambda)_{\leq 0}^{\operatorname{veff}}$ to be the category of *co-very effective* spectra, that is the full subcategory of $\log \operatorname{SH}_{\tau}(S, \Lambda)$ that is spanned by those objects \mathcal{G} such that $\operatorname{Hom}_{\log \operatorname{SH}_{\tau}(S,\Lambda)}(\mathcal{F},\mathcal{G}) \simeq 0$ for every $\mathcal{F} \in \operatorname{logSH}_{\tau}(S,\Lambda)_{\geq 1}^{\operatorname{veff}}$. We similarly write $\operatorname{logSH}_{\tau}(S,\Lambda)_{\leq i}^{\operatorname{veff}}$ for the right orthogonal subcategory to $\operatorname{logSH}_{\tau}(S,\Lambda)_{>i+1}^{\operatorname{veff}}$.

Remark 2.4. For every integer *i*, observe that we have the obvious inclusions

$$\log \operatorname{SH}(S)_{\geq i}^{\operatorname{veff}} \subset \log \operatorname{SH}(S)_{\geq i}^{\operatorname{eff}}$$
 and $\log \operatorname{SH}(S)_{\leq i}^{\operatorname{veff}} \subset \log \operatorname{SH}(S)_{\leq i}^{\operatorname{veff}}$.

Remark 2.5. Assume that S = Spec(k) is a field and that $\tau = d\text{Nis}$. As in \mathbb{A}^1 -motivic homotopy theory, $\log \text{SH}_{\tau}(k, \Lambda)^{\text{veff}}$ is the connective part of *t*-structures on $\log \text{SH}_{\tau}(S, \Lambda)$ and on $\log \text{SH}_{\tau}(S, \Lambda)^{\text{eff}}$ by [35, Proposition 1.4.4.11]. Note that the analog of Morel's connectivity theorem in the log setting [14, Theorem 3.2], together with semi-purity [14, Theorem 4.4], gives that this is exactly the non-negative part of the standard homotopy *t*-structure (see [39, Theorem 5.2.3]).

Remark 2.6. A similar construction can be performed in other motivic settings. For example, one could consider in the category of \mathbb{P}^1 -spectra MS_S introduced by Annala-Iwasa [2] the subcategory MS_S^{veff} generated by $\Sigma_{\mathbb{P}^1}^{\infty}X_+$ for $X \in \text{Sm}/S$ and closed under colimits and extensions. Again by [35, Proposition 1.4.4.11], this is the connective part of a *t*-structure on MS_S (that we can call the homotopy *t*-structure, if the expected connection to homotopy sheaves holds). We can consider the colimit-preserving inclusions

$$\Sigma_{\mathbb{P}^1}^{i+1}\mathrm{MS}_S^{\mathrm{veff}} \subset \Sigma_{\mathbb{P}^1}^i\mathrm{MS}_S^{\mathrm{veff}} \subset \cdots \subset \mathrm{MS}_S$$

giving rise via their right adjoints to functorial filtrations

$$\mathbf{f}_{i+1}E \to \mathbf{f}_iE \to \dots \to E$$

for any spectrum E.

Recall from [17, Construction 4.0.18] that there is a symmetric monoidal functor

 $\lambda_{\sharp} \colon \mathrm{MS}_S \to \mathrm{logSH}(S)$

induced by the functor $\lambda \colon \text{Sm}/S \to \text{Sm}/S$ given by $\lambda(X) := X$ (seen as log scheme with trivial log structure). The following is immediate from the definitions.

Proposition 2.7. Let S be any quasi-compact quasi-separated scheme. Then for every integer n we have the inclusion $\lambda_{\sharp} \Sigma_{\mathbb{P}^1}^n \mathrm{MS}_S^{\mathrm{veff}} \subset \Sigma_{\mathbb{P}^1}^n \mathrm{logSH}(S)^{\mathrm{veff}}$.

We record some elementary properties of the slice filtration.

Lemma 2.8. Let τ' be a topology on SmlSm/S finer than τ . Then for every integer *i*, the τ' -localization functor

$$L_{\tau'} \colon \mathrm{logSH}_{\tau}(S, \Lambda) \to \mathrm{logSH}_{\tau'}(S, \Lambda)$$

sends $\log SH_{\tau}(S, \Lambda)_{>i}^{eff}$ into $\log SH_{\tau'}(S, \Lambda)_{>i}^{eff}$, and its right adjoint

 $\iota \colon \mathrm{logSH}_{\tau'}(S,\Lambda) \hookrightarrow \mathrm{logSH}_{\tau}(S,\Lambda)$

sends $\log SH_{\tau'}(S, \Lambda)_{\leq i}^{\text{eff}}$ into $\log SH_{\tau}(S, \Lambda)_{\leq i}^{\text{eff}}$. A similar result holds for the very effective version, too.

Proof. We only give proof for the effective version. The statement about the left adjoint is a consequence of the fact that $L_{\tau'}$ preserves colimits and sends $\Sigma^{2n,n}\Sigma^{\infty}X_+$ to $\Sigma^{2n,n}\Sigma^{\infty}X_+$ for every integer n. Use the natural isomorphism

 $\operatorname{Hom}_{\operatorname{logSH}_{\tau}(S,\Lambda)}(\mathcal{F},\iota\mathcal{G}) \cong \operatorname{Hom}_{\operatorname{logSH}_{\tau}(S,\Lambda)}(L_{\tau'}\mathcal{F},\mathcal{G})$

for $\mathcal{F} \in \log SH_{\tau}(S, \Lambda)_{\geq i+1}^{\text{eff}}$ and $\mathcal{G} \in \log SH_{\tau'}(S, \Lambda)_{\leq i}^{\text{eff}}$ to deduce the statement about the right adjoint.

Lemma 2.9. Let $\Lambda \to \Lambda'$ be a map of \mathbb{E}_{∞} -rings. Then the base-change functor

$$-\otimes_{\Lambda} \Lambda' \colon \mathrm{logSH}_{\tau}(S,\Lambda) \to \mathrm{logSH}_{\tau}(S,\Lambda')$$

sends $\log SH_{\tau}(S, \Lambda)_{\geq i}^{\text{eff}}$ into $\log SH_{\tau}(S, \Lambda')_{\geq i}^{\text{eff}}$, and the restriction functor

 $\log \operatorname{SH}_{\tau}(S, \Lambda') \to \log \operatorname{SH}_{\tau}(S, \Lambda)$

sends $\log SH_{\tau}(S, \Lambda')_{\leq i}^{\text{eff}}$ to $\log SH_{\tau}(S, \Lambda)_{\leq i}^{\text{eff}}$. A similar result holds for the very effective version, too.

Proof. This is a consequence of the fact that $-\otimes_{\Lambda}\Lambda'$ preserves colimits and sends $\Sigma^{2n,n}\Sigma^{\infty}X_+$ to $\Sigma^{2n,n}\Sigma^{\infty}X_+$ for every $X \in \text{SmlSm}/S$ and integer n.

In particular, the restriction functor $\log DA(S) \rightarrow \log SH(S)$ sends $\log DA(S)_{\leq i}^{\text{eff}}$ to $\log SH(S)_{\leq i}^{\text{eff}}$. The following easy Lemma will be useful for the main theorem later in the text.

Lemma 2.10. If $\mathcal{F} \in \log SH_{\tau}(S, \Lambda)$ admits a filtration such that $Fil^{i}\mathcal{F} \in \log SH_{\tau}(S, \Lambda)_{\leq i}^{eff}$ for every integer *i*, then, there exists a unique natural morphism of filtered objects

$$f_{\bullet}^{\tau}\mathcal{F} \to Fil_{\bullet}\mathcal{F}$$

whose morphism of underlying objects is id: $\mathcal{F} \to \mathcal{F}$. Similarly, if $\mathcal{F} \in \log SH_{\tau}(S, \Lambda)$ admits a filtration such that $Fil^i \mathcal{F} \in \log SH_{\tau}(S, \Lambda)_{\leq i}^{\text{veff}}$ for every *i*, then there exists a unique natural morphism of filtered objects

$$\tilde{f}_{\bullet}^{\tau}\mathcal{F} \to \mathrm{Fil}_{\bullet}\mathcal{F}$$

whose morphism of underlying objects is $id: \mathcal{F} \to \mathcal{F}$.

Proof. We only give proof for the effective version. Consider the naturally induced maps

$$f^i_{\tau}\mathcal{F} \xrightarrow{\simeq} f^i_{\tau}(\mathrm{Fil}^i\mathcal{F}) \to \mathrm{Fil}^i_{\tau}\mathcal{F},$$

where the first map is obtained by the assumption $f_{\tau}^{i} \mathcal{F} \in \log SH_{\tau}(S, \Lambda)_{\leq i}^{\text{eff}}$ for every integer i. By taking fib $(\mathcal{F} \to (-))$, we get the desired morphism $f_{\bullet}^{\tau} \mathcal{F} \to Fil_{\bullet} \mathcal{F}$. For its uniqueness, consider the induced exact sequence

$$\operatorname{Hom}_{\operatorname{logSH}_{\tau}(S,\Lambda)}(\Sigma^{1,0}\mathrm{f}_{i+1}^{\tau}\mathcal{F},\operatorname{Fil}^{i}\mathcal{F}) \to \operatorname{Hom}_{\operatorname{logSH}_{\tau}(S,\Lambda)}(\mathrm{f}_{\tau}^{i}\mathcal{F},\operatorname{Fil}^{i}\mathcal{F}) \\ \to \operatorname{Hom}_{\operatorname{logSH}_{-}(S,\Lambda)}(\mathcal{F},\operatorname{Fil}^{i}\mathcal{F}) \to \operatorname{Hom}_{\operatorname{logSH}_{-}(S,\Lambda)}(\mathrm{f}_{i+1}^{\tau}\mathcal{F},\operatorname{Fil}^{i}\mathcal{F}).$$

The first and fourth terms are 0 since $\Sigma^{1,0}$ sends $\log \operatorname{SH}_{\tau}(S, \Lambda)^{\operatorname{eff}}_{\geq i+1}$ into itself. This establishes the uniqueness of $f^{\bullet}\mathcal{F} \to \operatorname{Fil}^{\bullet}\mathcal{F}$ and hence the uniqueness of $f^{\bullet}\mathcal{F} \to \operatorname{Fil}_{\bullet}\mathcal{F}$. \Box

3. SLICE FILTRATION ON K-THEORY

One of the fundamental results in \mathbb{A}^1 -homotopy theory is the equivalence $s_i \mathbf{KGL} \simeq \Sigma^{2i,i} \mathbf{MZ}$ in $\mathbf{SH}(k)$, where k is a perfect field, due to Voevodsky [51] in characteristic zero and Levine [31] in general (this is now known more generally, e.g., over Dedekind schemes by [7], [45, Theorem 2.19]. The effective and the very effective slices of \mathbf{KGL} coincide, see [1, Lemma 2.4], so that, in particular, the motivic Eilenberg-MacLane spectrum \mathbf{MZ} is very effective.

In this section, we promote this to $\log SH(k)$, assuming resolution of singularities.

Proposition 3.1. Let S be a quasi-compact quasi-separated scheme. Then, for every integer i, we have the inclusion

$$\omega^* \mathrm{SH}(S)_{\leq i}^{\mathrm{eff}} \subset \mathrm{logSH}(S)_{\leq i}^{\mathrm{eff}}$$

A similar result holds for the very effective version, too.

Proof. By adjunction, it suffices to show $\omega_{\sharp} \log \operatorname{SH}(S)_{\geq i}^{\operatorname{eff}} \subset \operatorname{SH}(S)_{\geq i}^{\operatorname{eff}}$ and $\omega_{\sharp} \log \operatorname{SH}(S)_{\geq i}^{\operatorname{veff}} \subset \operatorname{SH}(S)_{\geq i}^{\operatorname{veff}}$. This holds since ω_{\sharp} preserves colimits and $\omega_{\sharp} \Sigma_{\mathbb{P}^1}^n \Sigma^{\infty} Y_+ \simeq \Sigma_{\mathbb{P}^1}^n \Sigma^{\infty} (Y - \partial Y)_+$ for $Y \in \operatorname{SmlSm}/S$ and integer n.

Proposition 3.2. Let k be a perfect field admitting resolution of singularities. Then, for every integer i, we have the inclusion

$$\omega^* \mathrm{SH}(k)^{\mathrm{eff}}_{>i} \subset \mathrm{logSH}(k)^{\mathrm{eff}}_{>i}.$$

A similar result holds for the very effective version, too.

Proof. For $X \in \text{Sm}/k$, there exists proper $Y \in \text{Sm}\text{Sm}/k$ such that $Y - \partial Y \cong X$ by resolution of singularities. Due to [42, Theorem 1.1(2)], we have an equivalence $\omega^* \Sigma_{\mathbb{P}^1}^n \Sigma^\infty X_+ \simeq \Sigma_{\mathbb{P}^1}^n \Sigma^\infty Y_+$ for every integer *n*. To conclude, observe that ω^* preserves colimits by [42, Theorem 1.1(4)].

Proposition 3.3. Let k be a perfect field admitting resolution of singularities. For $\mathcal{F} \in$ SH(k) and integer i, there is a natural equivalence

$$\omega^* \mathbf{f}_i \mathcal{F} \simeq \mathbf{f}_i \omega^* \mathcal{F}$$

in $\log SH(k)$. A similar result holds for the very effective version, too.

Proof. We have the natural fiber sequence

$$f_i \omega^* f_i \mathcal{F} \to f_i \omega^* \mathcal{F} \to f_i \omega^* f^{i-1} \mathcal{F}$$

in logSH(k). Proposition 3.2 implies $f_i \omega^* f_i \mathcal{F} \simeq \omega^* f_i$, and Proposition 3.1 implies $f_i \omega^* f^{i-1} \mathcal{F} \simeq 0$. From these, we obtain the desired equivalence.

Theorem 3.4. Let k be a perfect field admitting resolution of singularities. Then the slice filtration and the very effective slice filtration on $\mathbf{KGL} \in \log SH(k)$ agree and are complete and exhaustive. Furthermore, there are natural equivalences

$$s_i KGL \simeq \widetilde{s}_i KGL \simeq \Sigma^{2i,i} M\mathbb{Z}$$

in $\log SH(k)$ for every integer i.

Proof. We have equivalences

$$s_i \mathbf{KGL} \simeq^{(1)} s_i \omega^* \mathbf{KGL} \simeq^{(2)} \omega^* s_i \mathbf{KGL} \simeq^{(3)} \omega^* \Sigma^{2i,i} \mathbf{MZ} \simeq^{(4)} \Sigma^{2i,i} \mathbf{MZ}$$

in $\log SH(k)$, where (1) is due to [17, Definition 6.2.6], (2) is due to Proposition 3.3, (3) is due to [31, Theorems 6.4.2, 9.0.3], and (4) is due to [17, Proposition 7.7.6].

We also have $f_i \mathbf{KGL} \simeq \omega^* f_i \mathbf{KGL}$. To show that the slice filtration on $\mathbf{KGL} \in \log SH(k)$ is complete and exhaustive, it suffices to show that the slice filtration on $\mathbf{KGL} \in SH(k)$ is complete and exhaustive since ω^* preserves colimits and limits. The fact that \mathbf{KGL} is slice complete can be found, e.g., in [44, Lemma 3.11].

For the very effective slice of $\mathbf{KGL} \in \log \mathrm{SH}(k)$, use [1, Lemma 2.4] and argue similarly as above.

4. HKR filtration on logarithmic Hochschild homology

Throughout this section, we fix (R, P) a pre-log ring, and S a quasi-compact quasiseparated scheme over Spec(R). In practice, the reader can ignore P in most statements and assume that R is always considered with a trivial log structure.

Our goal is to construct the HKR filtration on $\mathbf{HH}(-/R)$ and Beilinson filtrations on $\mathbf{HC}^{-}(-/R)$ and $\mathbf{HP}(-/R)$ in logDA(S), which are the \mathbb{P}^{1} -stabilized versions of the corresponding filtrations on $\mathrm{HH}(-/R)$, $\mathrm{HC}^{-}(-/R)$, and $\mathrm{HP}(-/R)$, built, in the log setting, in our previous works [12] and [13]. We also explore their fundamental properties.

Definition 4.1. For a map of pre-log rings $(R, P) \to (A, M)$ and integer *i*, we set

$$\mathbb{L}\Omega^{i}_{(A,M)/(R,P)} := \wedge^{i}_{A}\mathbb{L}_{(A,M)/(R,P)}.$$

where $\mathbb{L}_{(A,M)/(R,P)}$ is Gabber's logarithmic cotangent complex [41, §8], and $\wedge_A^i(-)$ is the *i*th derived exterior power. The *logarithmic derived de Rham cohomology* of $(R, P) \rightarrow (A, M)$ is defined as the total complex

$$\mathbb{L}\Omega_{(A,M)/(R,P)} := \operatorname{Tot}(\mathbb{L}\Omega^0_{(A,M)/(R,P)} \to \mathbb{L}\Omega^1_{(A,M)/(R,P)} \to \cdots).$$

This admits the Hodge filtration given by

$$\mathbb{L}\Omega^{\geq i}_{(A,M)/(R,P)} := \operatorname{Tot}(0 \to \dots \to 0 \to \mathbb{L}\Omega^{i}_{(A,M)/(R,P)} \to \mathbb{L}\Omega^{i+1}_{(A,M)/(R,P)} \to \dots)$$

where $\mathbb{L}\Omega^{i}_{(A,M)/(R,P)}$ is in cohomological degree i for $i \geq 0$ and $\mathbb{L}\Omega^{\geq i}_{(A,M)/(R,P)} := \mathbb{L}\Omega_{(A,M)/(R,P)}$ for i < 0. Let $\widehat{\mathbb{L}\Omega}_{(A,M)/(R,P)}$ be the completion of $\mathbb{L}\Omega_{(A,M)/(R,P)}$ with respect to the Hodge filtration, and let $\widehat{\mathbb{L}\Omega}^{\geq n}_{X/(R,P)}$ be the *n*th term of the induced Hodge filtration of $\widehat{\mathbb{L}\Omega}_{(A,M)/(R,P)}$.

Remark 4.2. By Zariski descent, we can also define the sheaf of complexes $\mathbb{L}\Omega^{i}_{X/(R,P)}$, $\mathbb{L}\Omega^{\geq i}_{X/(R,P)}$, $\widehat{\mathbb{L}\Omega}_{X/(R,P)}$, $\widehat{\mathbb{L}\Omega}_{X/(R,P)}$, and $\widehat{\mathbb{L}\Omega}^{\geq i}_{X/(R,P)}$ on X_{Zar} for every log scheme X over (R, P).

We now list some facts that we will use in the paper.

- (1) For any map of pre-log rings $(R, P) \to (A, M)$, the logarithmic Hochschild homology HH((A, M)/(R, P)) of [12, Definition 5.3] admits a separated, descending filtration with graded pieces $\wedge_A^i \mathbb{L}_{(A,M)/(R,P)}[i]$. This is obtained by Kan extension of the Postnikov filtration in the polynomial case. We shall refer to this as the log HKR filtration, see [12, Theorem 1.1].
- (2) The negative cyclic homology and the periodic homology $\text{HC}^{-}((A, M)/(R, P))$ and HP((A, M)/(R, P)) admit the complete exhaustive *Beilinson filtrations*

$$\operatorname{Fil}_{\geq \bullet}^{\operatorname{B}}\operatorname{HC}^{-}((A, M)/(R, P)), \quad \operatorname{Fil}_{\geq \bullet}^{\operatorname{B}}\operatorname{HP}((A, M)/(R, P))$$

with graded pieces $\widehat{\mathbb{L}\Omega}_{(A,M)/(R,P)}^{\geq n}[2n]$ and $\widehat{\mathbb{L}\Omega}_{(A,M)/(R,P)}[2n]$. The non-log case is due to Antieau [4, Theorem 1.1], while the log case is [13, Theorem 1.1]. If $(A, M) \in \operatorname{Poly}_{(R,P)}$, then

$$\operatorname{Fil}_{i}^{\mathrm{B}}\operatorname{HC}^{-}((A, M)/(R, P)) := \tau_{\geq 2i}^{\mathrm{B}} \tau_{\geq \bullet} \operatorname{HC}^{-}((A, M)/(R, P)),$$

$$\operatorname{Fil}_{i}^{\mathrm{B}}\operatorname{HP}((A, M)/(R, P)) := \tau_{\geq 2i}^{\mathrm{B}} \tau_{\geq \bullet} \operatorname{HP}((A, M)/(R, P)),$$

where $\tau_{\geq i}^{\text{B}}$ denotes the truncation functor for the Beilinson *t*-structure on the filtered derived category DF(A) [4, Definition 2.2]. We obtain the filtrations for general (A, M) by left Kan extension and regarding HC⁻ and HP as bicomplete bifiltered complexes, see [4, Remark 4.4], or [13, 2.2].

Construction 4.3. Let X be a log scheme. The inclusion functor from the category of line bundles over \underline{X} to the category of vector bundles over \underline{X} yields the canonical morphism

$$c_1^{\mathrm{K}} \colon R\Gamma_{\mathrm{Zar}}(\underline{X}, \mathbb{G}_m)[1] \to \mathrm{K}(\underline{X}).$$

The first Chern class for TC is the composite morphism

$$c_1^{\mathrm{TC}} \colon R\Gamma_{\mathrm{Zar}}(\underline{X}, \mathbb{G}_m)[1] \xrightarrow{c_1^{\mathrm{K}}} \mathrm{K}(\underline{X}) \xrightarrow{\mathrm{Tr}} \mathrm{TC}(\underline{X}) \xrightarrow{p^*} \mathrm{TC}(X),$$

where $p: X \to \underline{X}$ is the morphism removing the log structure. We similarly obtain the morphisms c_1^{THH} , $c_1^{\text{TC}^-}$, c_1^{TP} , c_1^{HH} , $c_1^{\text{HC}^-}$, and c_1^{HP} to THH(X), $\text{TC}^-(X)$, TP(X), HH(X), $\text{HC}^-(X)$, and HP(X).

Remark 4.4. We will set the following notation for convenience when dealing with projective bundle formulas below: For a morphism of commutative ring spectra $f: \mathcal{F} \to \mathcal{G}$ with $c \in \pi_0(\mathcal{G})$, we will often write fc (or simply c if f is clear from the context) for the composite morphism of spectra

(4.1)
$$fc\colon \mathcal{F} \xrightarrow{\simeq} \mathcal{F} \otimes_{\mathbb{S}} \mathbb{S} \xrightarrow{f \otimes c} \mathcal{G} \otimes_{\mathbb{S}} \mathcal{G} \xrightarrow{\mu} \mathcal{G},$$

where μ is the multiplication. We also consider the unit $1 \in \pi_0(\mathcal{G})$.

Recall the projective bundle formula for $K(\underline{X} \times \mathbb{P}^1)$: The morphism of spectra

$$(1, c_1^{\mathcal{K}}(\mathcal{O}(1))) \colon \mathcal{K}(\underline{X}) \oplus \mathcal{K}(\underline{X}) \to \mathcal{K}(\underline{X} \times \mathbb{P}^1)$$

is an equivalence, where $\mathcal{O}(1) \in \operatorname{Pic}(\underline{X} \times \mathbb{P}^1) \cong \pi_0(R\Gamma_{\operatorname{Zar}}(\underline{X}, \mathbb{G}_m)[1])$, and $c_1^{\mathrm{K}}(\mathcal{O}(1)) \in \pi_0(K(\underline{X} \times \mathbb{P}^1))$ (implicitly we are writing 1 for the pull-back map along $\pi : \underline{X} \times \mathbb{P}^1 \to \underline{X}$). Due to [19, Theorem 1.5], we similarly have the projective bundle formula for $\operatorname{TC}(\underline{X} \times \mathbb{P}^1)$: The morphism of spectra

(4.2)
$$(1, c_1^{\mathrm{TC}}(\mathcal{O}(1))) \colon \mathrm{TC}(\underline{X}) \oplus \mathrm{TC}(\underline{X}) \to \mathrm{TC}(\underline{X} \times \mathbb{P}^1)$$

is an equivalence. Analogous results hold for THH, TC⁻, TP, HH, HC⁻, and HP.

Proposition 4.5. There is an S^1 -equivariant equivalence of filtered complexes

(4.3)
$$\mathbb{Z} \oplus (\mathbb{Z}[1])\{-1\} \simeq \mathrm{HH}(\mathbb{P}^1/\mathbb{Z}),$$

where the filtrations on \mathbb{Z} and $\mathbb{Z}[1]$ are the Postnikov filtrations. Here, the notation $(\mathbb{Z}[1])\{-1\}$ means

$$\operatorname{Fil}_{i}((\mathbb{Z}[1])\{-1\}) = \begin{cases} \mathbb{Z} & \text{if } i \leq 1, \\ 0 & \text{otherwise} \end{cases}$$

Proof. Every S^1 -action on $\mathbb{Z} \oplus \mathbb{Z}$ is trivial, so it suffices to show the claim after forgetting the S^1 -actions. We have the equivalences of filtered complexes

$$\mathbb{Z}[x] \oplus \mathbb{Z}[x] dx[1] \simeq \operatorname{HH}(\mathbb{Z}[x]),$$
$$\mathbb{Z}[x^{-1}] \oplus \mathbb{Z}[x^{-1}] d(x^{-1})[1] \simeq \operatorname{HH}(\mathbb{Z}[x^{-1}]),$$
$$\mathbb{Z}[x, x^{-1}] \oplus \mathbb{Z}[x, x^{-1}] dx[1] \simeq \operatorname{HH}(\mathbb{Z}[x, x^{-1}]),$$

where the filtrations are the Postnikov filtrations on both sides. Consider the standard cover on \mathbb{P}^1 , and use the equivalences of complexes

$$\mathbb{Z} \simeq \operatorname{fib}(\mathbb{Z}[x] \oplus \mathbb{Z}[x^{-1}] \to \mathbb{Z}[x, x^{-1}])$$
$$\mathbb{Z}[-1] \simeq \operatorname{fib}(\mathbb{Z}[x]dx \oplus \mathbb{Z}[x^{-1}]d(x^{-1}) \to \mathbb{Z}[x, x^{-1}]dx)$$

to obtain the desired equivalence.

Proposition 4.6. Let X be a log scheme. Then the morphism of spectra

(4.4)
$$(1, c_1^{\mathrm{TC}}(\mathcal{O}(1))) \colon \mathrm{TC}(X) \oplus \mathrm{TC}(X) \to \mathrm{TC}(X \times \mathbb{P}^1)$$

is an equivalence, and this is obtained by applying $TC(X) \otimes_{\mathbb{Z}} - to$ (4.3). We have similar results for THH, TC^- , TP, HH, HC^- , and HP.

Proof. We focus on TC since the proofs are similar. By base-change along $TC(\underline{X}) \rightarrow TC(X)$ from (4.2) we get (4.4). Hence we reduce to the case when X has a trivial log structure.

By [19, Theorem 9.1 and p. 1102], we have the commutative diagram of spectra

$$\begin{array}{ccc} \mathrm{K}(X)[1] & \xrightarrow{\tau_{\mathrm{K}}} & \mathrm{K}(X \times \mathbb{G}_m) & \xrightarrow{\partial} & \mathrm{K}(X \times \mathbb{P}^1)[1] \\ & & & & & \downarrow \mathrm{Tr} & & & \downarrow \mathrm{Tr} \\ & & & & \mathrm{TC}(X)[1] & \xrightarrow{\tau_{\mathrm{TC}}} & \mathrm{TC}(X \times \mathbb{G}_m) & \xrightarrow{\partial} & \mathrm{TC}(X \times \mathbb{P}^1)[1], \end{array}$$

where $\tau_{\rm K}$ and $\tau_{\rm TC}$ are obtained by the Bass functor structures on K and TC, and the boundary morphisms ∂ are obtained by the standard cover of \mathbb{P}^1 . The composite $\partial \circ \tau_K$ sends $1 \in \pi_0 {\rm K}(X)$ to $c_1^{\rm K}(\mathcal{O}(1)) \in \pi_0 {\rm K}(X \times \mathbb{P}^1)$, so the composite $\partial \circ \tau_{\rm TC}$ sends $1 \in \pi_0 {\rm TC}(X)$ to $c_1^{\rm TC}(\mathcal{O}(1)) \in \pi_0 {\rm TC}(X \times \mathbb{P}^1)$. It follows that the morphism

$$(1, \partial \circ \tau_{\mathrm{TC}}) \colon \mathrm{TC}(X) \oplus \mathrm{TC}(X) \to \mathrm{TC}(X \times \mathbb{P}^1)$$

agrees with $(1, c_1^{\mathrm{TC}}(\mathcal{O}(1)))$.

By [19, pp. 1102–1103], we see that τ_{THH} : $\text{THH}(X)[1] \to \text{THH}(X \times \mathbb{G}_m)$ is obtained by applying $\text{THH}(X) \otimes_{\mathbb{Z}} - \text{to } \tau_{\text{HH}} : \text{HH}(\mathbb{Z})[1] \to \text{HH}(\mathbb{Z}[x, x^{-1}])$ that sends $1 \in \pi_0 \text{HH}(\mathbb{Z})$ to $dx/x \in \Omega_{\mathbb{Z}[x, x^{-1}]/\mathbb{Z}} \cong \pi_1 \text{HH}(\mathbb{Z}[x, x^{-1}])$. Furthermore, τ_{TC} is obtained by applying TC to τ_{THH} . Compare this with (4.3) to conclude.

Proposition 4.7. Let (R, P) be a pre-log ring. For a quasi-compact quasi-separated log scheme X over (R, P) and $i \in \mathbb{Z} \cup \{-\infty\}$, the projective bundle formula for $HH(X \times$

 $\mathbb{P}^1/(R,P)$) restricts to natural equivalences of complexes

$$\begin{aligned} \operatorname{Fil}_{i}^{\operatorname{HKR}}\operatorname{HH}(X \times \mathbb{P}^{1}/(R, P)) &\simeq \operatorname{Fil}_{i}^{\operatorname{HKR}}\operatorname{HH}(X/(R, P)) \oplus \operatorname{Fil}_{i-1}^{\operatorname{HKR}}\operatorname{HH}(X/(R, P)), \\ \operatorname{Fil}_{i}^{\operatorname{B}}\operatorname{HC}^{-}(X \times \mathbb{P}^{1}/(R, P)) &\simeq \operatorname{Fil}_{i}^{\operatorname{B}}\operatorname{HC}^{-}(X/(R, P)) \oplus \operatorname{Fil}_{i-1}^{\operatorname{B}}\operatorname{HC}^{-}(X/(R, P)), \\ \operatorname{Fil}_{i}^{\operatorname{B}}\operatorname{HP}(X \times \mathbb{P}^{1}/(R, P)) &\simeq \operatorname{Fil}_{i}^{\operatorname{B}}\operatorname{HP}(X/(R, P)) \oplus \operatorname{Fil}_{i-1}^{\operatorname{B}}\operatorname{HP}(X/(R, P)). \end{aligned}$$

Proof. We refer to [13, §2.2] for the category $\operatorname{Poly}_{(R,P)}$ of polynomial (R, P)-algebras. By Zariski descent and left Kan extension, we reduce to the case when $X := \operatorname{Spec}(A, M)$ with $(A, M) \in \operatorname{Poly}_{(R,P)}$. In this case, the HKR filtration on $\operatorname{HH}(X/(R, P))$ is the Postnikov filtration. Hence we have the S^1 -equivariant equivalence of filtered complexes

$$\operatorname{HH}(X \times \mathbb{P}^1/(R, P)) \simeq \operatorname{HH}(X/(R, P)) \otimes \operatorname{HH}(\mathbb{P}^1/\mathbb{Z}).$$

Together with Proposition 4.5, we obtain the S^1 -equivariant decomposition of filtered complexes

(4.5)
$$\operatorname{HH}(X \times \mathbb{P}^1/(R, P)) \simeq \operatorname{HH}(X/(R, P)) \oplus (\operatorname{HH}(X/(R, P))[1])\{-1\},$$

where the filtrations on HH(X/(R, P)) and HH(X/(R, P))[1] are the Postnikov filtrations. This implies the claim for HH.

By applying $(-)^{hS^1}$ on both sides of (4.5), we obtain an equivalence of complexes induced by the homotopy fixed point spectra sequence:

$$\operatorname{Fil}_{\bullet}^{\operatorname{HKR}}\operatorname{HC}^{-}(X \times \mathbb{P}^{1}/(R, P)) \simeq \operatorname{Fil}_{\bullet}^{\operatorname{HKR}}\operatorname{HC}^{-}(X/(R, P)) \oplus \operatorname{Fil}_{\bullet-1}^{\operatorname{HKR}}\operatorname{HC}^{-}(X/(R, P)).$$

Take $\tau^{\rm B}_{\geq 2i}$ on both sides to show the claim for HC⁻. The claim for HP can be proven similarly.

Construction 4.8. The local-to-global spectral sequence $E_2^{st} = H^s(X, \Omega_X^t) \Rightarrow \pi_{t-s} \operatorname{HH}(X)$ for $X = \mathbb{P}^1$, together with the decomposition of $\operatorname{HH}(\mathbb{P}^1/\mathbb{Z})$ of Proposition 4.5, induces by restriction of $c_1^{\operatorname{HH}}(\mathcal{O}(1)) \in \pi_0 \operatorname{HH}(\mathbb{P}^1)$ a class we obtain

$$c_1^{\operatorname{Hod}}(\mathcal{O}(1)) \in H^1_{\operatorname{Zar}}(\mathbb{P}^1, \Omega^1_{\mathbb{P}^1/\mathbb{Z}}),$$

that agrees with the classical first Chern class in Hodge cohomology. After pulling back, we obtain

$$c_1^{\operatorname{Hod}}(\mathcal{O}(1)) \in H^1_{\operatorname{Zar}}(X \times \mathbb{P}^1, \mathbb{L}\Omega^1_{X/(R,P)})$$

for a log scheme X over a pre-log ring (R, P). By descent and reduction to the polynomial case, we have the following \mathbb{P}^1 -bundle formulas: the natural morphisms of complexes

$$\begin{aligned} &R\Gamma_{\mathrm{Zar}}(X, \mathbb{L}\Omega^{i}_{X/(R,P)}) \oplus R\Gamma_{\mathrm{Zar}}(X, \mathbb{L}\Omega^{i-1}_{X/(R,P)})[-1] \to R\Gamma_{\mathrm{Zar}}(X \times \mathbb{P}^{1}, \mathbb{L}\Omega^{i}_{X \times \mathbb{P}^{1}/(R,P)}), \\ &R\Gamma_{\mathrm{Zar}}(X, \widehat{\mathbb{L}\Omega}^{\geq i}_{X/(R,P)}) \oplus R\Gamma_{\mathrm{Zar}}(X, \widehat{\mathbb{L}\Omega}^{\geq i-1}_{X/(R,P)})[-1] \to R\Gamma_{\mathrm{Zar}}(X \times \mathbb{P}^{1}, \widehat{\mathbb{L}\Omega}^{\geq i}_{X \times \mathbb{P}^{1}/(R,P)}), \\ &R\Gamma_{\mathrm{Zar}}(X, \widehat{\mathbb{L}\Omega}_{X/(R,P)}) \oplus R\Gamma_{\mathrm{Zar}}(X, \widehat{\mathbb{L}\Omega}_{X/(R,P)})[-1] \to R\Gamma_{\mathrm{Zar}}(X \times \mathbb{P}^{1}, \widehat{\mathbb{L}\Omega}_{X \times \mathbb{P}^{1}/(R,P)}), \end{aligned}$$

induced by 1 and $c_1^{\text{Hod}}(\mathcal{O}(1))$ are equivalences (see proof of [11, Proposition 2.20]).

If X is quasi-compact and quasi-separated, these morphisms are identified with the natural morphisms

$$\begin{aligned} \operatorname{gr}_{i}^{\operatorname{HKR}} \operatorname{HH}(X/(R,P)) &\oplus \operatorname{gr}_{i-1}^{\operatorname{HKR}} \operatorname{HH}(X/(R,P)) \simeq \operatorname{gr}_{i}^{\operatorname{HKR}} \operatorname{HH}(X \times \mathbb{P}^{1}/(R,P)), \\ \operatorname{gr}_{i}^{\operatorname{B}} \operatorname{HC}^{-}(X/(R,P)) &\oplus \operatorname{gr}_{i-1}^{\operatorname{B}} \operatorname{HC}^{-}(X/(R,P)) \simeq \operatorname{gr}_{i}^{\operatorname{B}} \operatorname{HC}^{-}(X \times \mathbb{P}^{1}/(R,P)), \\ \operatorname{gr}_{i}^{\operatorname{B}} \operatorname{HP}(X/(R,P)) &\oplus \operatorname{gr}_{i-1}^{\operatorname{B}} \operatorname{HP}(X/(R,P)) \simeq \operatorname{gr}_{i}^{\operatorname{B}} \operatorname{HP}(X \times \mathbb{P}^{1}/(R,P)) \end{aligned}$$

obtained by Proposition 4.7. Indeed, this can be shown for HH by comparing (4.1) with the composite map

$$R\Gamma_{\operatorname{Zar}}(X, \mathbb{L}\Omega^{i-1}_{X/(R,P)}) \otimes_{\mathbb{Z}} \mathbb{Z}[-1]$$

$$\rightarrow R\Gamma_{\operatorname{Zar}}(X \times \mathbb{P}^{1}, \mathbb{L}\Omega^{i-1}_{X \times \mathbb{P}^{1}/(R,P)}) \otimes_{\mathbb{Z}} R\Gamma_{\operatorname{Zar}}(X \times \mathbb{P}^{1}, \mathbb{L}\Omega^{1}_{X \times \mathbb{P}^{1}/(R,P)})$$

$$\rightarrow R\Gamma_{\operatorname{Zar}}(X \times \mathbb{P}^{1}, \mathbb{L}\Omega^{i}_{X \times \mathbb{P}^{1}/(R,P)}).$$

A similar argument proves the claim for HC⁻ and HP.

Proposition 4.9. Let (R, P) be a pre-log ring. Then the presheaves of complexes

 $\mathrm{Fil}_i^{\mathrm{HKR}}\mathrm{HH}(-/(R,P)), \ \mathrm{Fil}_i^{\mathrm{B}}\mathrm{HC}^-(-/(R,P)), \ and \quad \mathrm{Fil}_i^{\mathrm{B}}\mathrm{HP}(-/(R,P))$

on log schemes over (R, P) are $(\mathbb{P}^n, \mathbb{P}^{n-1})$ -invariant for n > 0 and $i \in \mathbb{Z} \cup \{-\infty\}$. Moreover, the presheaves of complexes $\mathbb{L}\Omega^i_{/(R,P)}$, $\widehat{\mathbb{L}\Omega}^{\geq i}_{/(R,P)}$, and $\widehat{\mathbb{L}\Omega}_{/(R,P)}$ on the category of log schemes over (R, P) given by

$$X \mapsto R\Gamma_{\operatorname{Zar}}(X, \mathbb{L}\Omega^{i}_{X/(R,P)}), \ R\Gamma_{\operatorname{Zar}}(X, \widehat{\mathbb{L}\Omega}^{\leq i}_{X/(R,P)}), \ R\Gamma_{\operatorname{Zar}}(X, \widehat{\mathbb{L}\Omega}_{X/(R,P)})$$

are $(\mathbb{P}^n, \mathbb{P}^{n-1})$ -invariant for n > 0 and $i \in \mathbb{Z}$.

Proof. This is basically discussed in [12, §7, 8], forgetting the filtrations. Let X be a log scheme over (R, P). Using the completeness and exhaustiveness of the filtrations, it suffices to show that the natural map of complex

$$\operatorname{gr}_{i}^{\operatorname{HKR}}\operatorname{HH}(X/(R,P)) \to \operatorname{gr}_{i}^{\operatorname{HKR}}\operatorname{HH}(X \times (\mathbb{P}^{n}, \mathbb{P}^{n-1})/(R,P))$$

and the similar maps for $\operatorname{gr}_{i}^{\operatorname{B}}\operatorname{HC}^{-}$ and $\operatorname{gr}_{i}^{\operatorname{B}}\operatorname{HP}$ are equivalences. Furthermore, we reduce to the case when $X = \operatorname{Spec}(A, M)$ with $(A, M) \in \operatorname{Poly}_{(R,P)}$ by left Kan extension. Then the claim is a consequence of the $(\mathbb{P}^{n}, \mathbb{P}^{n-1})$ -invariance of Hodge cohomology, see e.g. the proof of [16, Proposition 9.2.1].

Definition 4.10. For $X \in \text{SmlSm}/S$ and $i \in \mathbb{Z} \cup \{-\infty\}$, Proposition 4.7 yields a natural equivalence of complexes

(4.6)
$$\operatorname{Fil}_{i}^{\operatorname{HKR}}\operatorname{HH}(X/R) \simeq \Omega_{\mathbb{P}^{1}}\operatorname{Fil}_{i+1}^{\operatorname{HKR}}\operatorname{THH}(X/R).$$

Using this as bonding maps, Proposition 4.9 enables us to define the complete exhaustive filtration

$$\operatorname{Fil}_{i}^{\operatorname{HKR}}\mathbf{HH}(-/R) := (\operatorname{Fil}_{i}^{\operatorname{HKR}}\operatorname{HH}(-/R), \operatorname{Fil}_{i+1}^{\operatorname{HKR}}\operatorname{HH}(-/R), \ldots)$$

on $\mathbf{HH}(-/R) \simeq \operatorname{Fil}_{-\infty}^{\operatorname{HKR}} \mathbf{HH}(-/R) \in \operatorname{logDA}(S)$, where $\mathbf{HH}(-/R)$ is the \mathbb{P}^1 -spectrum representing (log) Hochschild homology constructed in [12, §8]. We similarly define the complete exhaustive filtrations

$$\operatorname{Fil}_{i}^{\mathrm{B}}\mathbf{H}\mathbf{C}^{-}(-/R)$$
 and $\operatorname{Fil}_{i}^{\mathrm{B}}\mathbf{H}\mathbf{P}(-/R)$.

Proposition 4.11. For $i \in \mathbb{Z} \cup \{-\infty\}$, we have a natural equivalence of complexes

$$\operatorname{Fil}_{i}^{\operatorname{HKR}}\mathbf{HH}(-/R)(1)[2] \simeq \operatorname{Fil}_{i+1}^{\operatorname{HKR}}\mathbf{HH}(-/R).$$

We have similar equivalences for $\operatorname{Fil}_{i}^{\operatorname{B}}\operatorname{HC}^{-}$ and $\operatorname{Fil}_{i}^{\operatorname{B}}\operatorname{HP}$.

Proof. For $\operatorname{Fil}_{i}^{\operatorname{HKR}}$ **HH**, this is a direct consequence of (4.6). The proofs for the other cases are similar.

Definition 4.12. For $i \in \mathbb{Z}$, Proposition 4.9 enables us to define the \mathbb{P}^1 -spectra

$$\mathbb{L}\mathbf{\Omega}^{i}_{/R} := (\mathbb{L}\Omega^{i}_{/R}, \mathbb{L}\Omega^{i+1}_{/R}[1], \ldots),$$
$$\widehat{\mathbb{L}\mathbf{\Omega}}^{\geq i}_{/R} := (\widehat{\mathbb{L}\Omega}^{\geq i}_{/R}, \widehat{\mathbb{L}\Omega}^{\geq i+1}_{/R}[1], \ldots),$$
$$\widehat{\mathbb{L}\mathbf{\Omega}}_{/R} := (\widehat{\mathbb{L}\Omega}_{/R}, \widehat{\mathbb{L}\Omega}_{/R}[1], \ldots)$$

in logDA(S) whose bonding maps are given by the projective bundle formulas in Construction 4.8. If $S \to \operatorname{Spec}(R)$ is smooth, then we omit \mathbb{L} in $\mathbb{L}\Omega$.

Proposition 4.13. For $i \in \mathbb{Z}$, we have natural equivalences

$$\begin{split} \mathrm{gr}_{i}^{\mathrm{HKR}}\mathbf{H}\mathbf{H}(-/R) &\simeq \mathbb{L}\mathbf{\Omega}_{/R}^{i}[i],\\ \mathrm{gr}_{i}^{\mathrm{B}}\mathbf{H}\mathbf{C}^{-}(-/R) &\simeq \widehat{\mathbb{L}\mathbf{\Omega}_{/R}^{\geq i}}[i],\\ \mathrm{gr}_{i}^{\mathrm{B}}\mathbf{H}\mathbf{P}(-/R) &\simeq \widehat{\mathbb{L}\mathbf{\Omega}_{/R}}[i] \end{split}$$

in $\log DA(S)$.

Proof. The bonding maps defining the \mathbb{P}^1 -spectra on both sides are compatible, as observed in Construction 4.8.

We are ready to prove our main result on HH and variants.

Theorem 4.14. Assume that $S \to \operatorname{Spec}(R)$ is smooth. For $i \in \mathbb{Z}$, we have

$$\operatorname{Fil}_{\operatorname{HKR}}^{i}\operatorname{HH}(-/R), \operatorname{Fil}_{\operatorname{B}}^{i}\operatorname{HC}^{-}(-/R), \operatorname{Fil}_{\operatorname{B}}^{i}\operatorname{HP}(-/R) \in \operatorname{logDA}(S)_{< i}^{\operatorname{veff}}$$

In particular, there are canonical morphisms

$$\begin{split} \tilde{\mathbf{f}}_{n}\mathbf{H}\mathbf{H}(-/R) &\simeq \Sigma^{2n,n} \tilde{\mathbf{f}}_{0}\mathbf{H}\mathbf{H}(-/R) \to \mathrm{Fil}_{n}^{\mathrm{HKR}}\mathbf{H}\mathbf{H}(-/R), \\ \tilde{\mathbf{f}}_{n}\mathbf{H}\mathbf{C}^{-}(-/R) &\simeq \Sigma^{2n,n} \tilde{\mathbf{f}}_{0}\mathbf{H}\mathbf{C}^{-}(-/R) \to \mathrm{Fil}_{n}^{\mathrm{B}}\mathbf{H}\mathbf{C}^{-}(-/R), \\ \tilde{\mathbf{f}}_{n}\mathbf{H}\mathbf{P}(-/R) &\simeq \Sigma^{2n,n} \tilde{\mathbf{f}}_{0}\mathbf{H}\mathbf{P}(-/R) \to \mathrm{Fil}_{n}^{\mathrm{B}}\mathbf{H}\mathbf{P}(-/R), \end{split}$$

Proof. By Proposition 4.11, we reduce to the case when i = -1. Since $\Sigma^{\infty} X_{+}$ is compact in logDA(S), it suffices to show the vanishing

$$\operatorname{Hom}_{\log \mathrm{DA}(S)}(\Sigma^{\infty}X_{+}, \operatorname{gr}_{-j}^{\mathrm{HKR}}\mathbf{HH}(-/R)) \simeq 0$$

for every integer j > 0 and similar vanishings for $\operatorname{gr}_{-j}^{\mathrm{B}}\mathbf{H}\mathbf{C}^{-}$ and $\operatorname{gr}_{-j}^{\mathrm{B}}\mathbf{H}\mathbf{P}$. By the smoothness assumption and Proposition 4.13, it suffices to show the vanishings

$$H_{\text{Zar}}^{-2j}(X, \Omega_{X/R}^{-j}) = 0 \text{ and } H_{\text{Zar}}^{-2j}(X, \Omega_{X/R}^{\ge 0}) = 0$$

for j > 0. This is clear.

Remark 4.15. Since $H_{Zar}^{2j}(X, \Omega_{X/R}^{\geq 0})$ does not vanish for $j \geq 0$ and nontrivial X, the proof of Theorem 4.14 shows that $\operatorname{Fil}_{\mathrm{B}}^{i}\mathbf{HC}^{-}(-/R)$ and $\operatorname{Fil}_{\mathrm{B}}^{i}\mathbf{HP}(-/R)$ are not in $\operatorname{logDA}(S)_{\leq i}^{\operatorname{eff}}$. On the other hand, we have $\operatorname{Fil}_{\mathrm{HKR}}^{i}\mathbf{HH}(-/R) \in \operatorname{logDA}_{\leq i}^{\operatorname{eff}}(S)$.

Proof of Theorem 1.5. (1) is a consequence of Lemma 2.10 and Theorems 3.4 and 4.14. (2) is a consequence of (1) and Proposition 4.13, and (3) is a consequence of (2). \Box

Question 4.16. Despite having a very different origin, Theorem 4.14 shows an intimate connection between the (very effective) slice filtration and the HKR filtration (and the Beilinson filtrations). We leave it as an open question whether the (very effective) slice and HKR filtrations agree on $\mathbf{HH}(-/(R, P))$, and whether the slice and Beilinson filtrations agree on $\mathbf{HC}^{-}(-/(R, P))$ and $\mathbf{HP}(-/(R, P))$.

5. BMS filtration on logarithmic topological Hochschild homology

Let us fix a prime p. We refer the reader to [13] for the definitions and the main properties of the category lQSyn of log quasi-syntomic rings, the category lQRSPerfd of log quasi-regular semiperfectoid rings, and the log quasi-syntomic topology on lQSyn.

Definition 5.1. Let \mathfrak{X} be a quasi-coherent bounded *p*-adic formal log scheme. We say that \mathfrak{X} is *log quasi-syntomic* if strict étale locally it is isomorphic to $\text{Spf}(A, M)^a$ for $(A, M) \in \text{lQSyn}$. Let FlQSyn denote the category of quasi-compact quasi-separated log quasi-syntomic formal log schemes.

Our goal is to define the BMS filtrations on the motivic \mathbb{P}^1 -spectra representing (log) THH, TC⁻, TP, and TC, defined in [17, Section 8] and explore their fundamental properties.

Proposition 5.2. The presheaves $\text{THH}(-;\mathbb{Z}_p)$, $\text{THH}(-;\mathbb{Z}_p)_{hS^1}$, $\text{TC}(-;\mathbb{Z}_p)^-$, and $\text{TP}(-;\mathbb{Z}_p)$ on lQSyn^{op} are log quasi-syntomic sheaves.

Proof. Argue as in [36, Theorem 5.5], but use the *p*-completed version of [13, Theorem 2.3] as [9, Remark 4.9]. See [13, Proposition 3.10] for the non *p*-completed version. \Box

In particular, for every $\mathfrak{X} \in \operatorname{FlQSyn}$, we can consider $\operatorname{THH}(\mathfrak{X};\mathbb{Z}_p)$ and its cousins.

Remark 5.3. Let A be a commutative ring. Then we have an equivalence $\text{THH}(A_p^{\wedge}; \mathbb{Z}_p) \simeq \text{THH}(A; \mathbb{Z}_p)$. Thus, the quasi-syntomic sheaf $\text{THH}(-; \mathbb{Z}_p)$ restricted to FlSm/R in [11, Definition 2.2] commutes with the p-completion functor from lSm/R for any p-complete ring R. For this reason, we will implicitly extend the \mathbb{P}^1 -spectrum **THH** in $\log \text{SH}(R)$ of [17, §8] (denoted **logTHH** in loc. cit.) to a \mathbb{P}^1 -spectrum, indicated with the same letters, in $\log \text{FSH}(R)$. For the same reason, we will implicitly consider the sheaf $\text{THH}(-;\mathbb{Z}_p)$ and variants as defined on (log) schemes or on p-adic formal (log) schemes without explicitly mentioning it.

Let us quickly review how we can extend the BMS filtrations in [9] to pre-log rings, see [13]. For $(A, M) \in lQRSPerfd$, the BMS filtrations on

 $\operatorname{THH}((A, M); \mathbb{Z}_p), \operatorname{TC}^{-}((A, M); \mathbb{Z}_p), \operatorname{TP}((A, M); \mathbb{Z}_p), \operatorname{TC}((A, M); \mathbb{Z}_p)$

are the double-speed Postnikov filtrations, that is, $\operatorname{Fil}^{i}_{BMS} := \tau_{\leq 2i}$. Equivalently, we have $\operatorname{Fil}^{BMS}_{i} := \tau_{\geq 2i-1}$. By log quasi-syntomic descent, we obtain the *BMS filtrations* for $(A, M) \in \operatorname{IQSyn}$ too. Observe that we have a natural equivalence of spectra

(5.1)
$$\operatorname{gr}_{i}^{\operatorname{BMS}}\operatorname{TC}((A, M); \mathbb{Z}_{p}) \simeq \operatorname{fib}(\operatorname{gr}_{i}^{\operatorname{BMS}}\operatorname{TC}^{-}((A, M); \mathbb{Z}_{p}) \xrightarrow{\varphi - \operatorname{can}} \operatorname{gr}_{i}^{\operatorname{BMS}}\operatorname{TP}((A, M); \mathbb{Z}_{p})).$$

For $\mathfrak{X} \in FlQSyn$, we also obtain the BMS filtrations on

 $\operatorname{THH}(\mathfrak{X};\mathbb{Z}_p), \operatorname{TC}^{-}(\mathfrak{X};\mathbb{Z}_p), \operatorname{TP}(\mathfrak{X};\mathbb{Z}_p), \operatorname{TC}(\mathfrak{X};\mathbb{Z}_p)$

by Zariski descent. Note that the graded pieces satisfy quasi-syntomic descent as discussed in [11, §3].

For a *p*-adic formal scheme \mathfrak{X} over $\operatorname{Spf}(R)$ for a quasi-syntomic ring R, let $\mathfrak{X} \times \mathbb{P}^n$ denote the *p*-adic formal scheme $\mathfrak{X} \times (\mathbb{P}^n_{\operatorname{Spf}(R)})_p^{\wedge}$ to simplify notation.

Proposition 5.4. The presheaves

$$\operatorname{Fil}_{i}^{\operatorname{BMS}}\operatorname{THH}(-;\mathbb{Z}_{p}), \ \operatorname{Fil}_{i}^{\operatorname{BMS}}\operatorname{TC}^{-}(-;\mathbb{Z}_{p}), \ and \ \operatorname{Fil}_{i}^{\operatorname{BMS}}\operatorname{TP}(-;\mathbb{Z}_{p})$$

on FlQSyn are $(\mathbb{P}^n, \mathbb{P}^{n-1})$ -invariant for $n \in \mathbb{Z}_{>0}$ and $i \in \mathbb{Z} \cup \{-\infty\}$.

Proof. Using the completeness and exhaustiveness of the filtrations, it suffices to show that the natural map of complex

$$\operatorname{gr}_{i}^{\operatorname{BMS}}\operatorname{THH}(\mathfrak{X};\mathbb{Z}_{p}) \to \operatorname{gr}_{i}^{\operatorname{BMS}}\operatorname{THH}(\mathfrak{X}\times(\mathbb{P}^{n},\mathbb{P}^{n-1});\mathbb{Z}_{p})$$

and the similar maps for TC⁻, TP, and TC are equivalences. This is a consequence of [13, Theorem 1.3(3),(4)] and [11, Corollary 3.11].

Proposition 5.5. For $(A, M) \in lQSyn$ and integer i < 0, we have the vanishing

$$\operatorname{Fil}_{\operatorname{BMS}}^{i}\operatorname{TC}((A, M); \mathbb{Z}_p) \simeq 0.$$

Proof. By (log) quasi-syntomic descent, we reduce to the case $(A, M) \in \text{lQRSPerfd}$. To conclude, observe that we have $\pi_i \text{TC}((A, M); \mathbb{Z}_p) \simeq 0$ for i < -1 using [40, Theorem II.4.10] as in the proof of [9, Proposition 7.16].

Construction 5.6. Let A be a commutative ring. Apply the étale sheafification and p-completion functors to the first Chern class $c_1^{\text{TC}} \colon R\Gamma_{\text{Zar}}(A, \mathbb{G}_m)[1] \to \text{TC}(A)$ to obtain the natural morphism

$$\widehat{c}_1^{\mathrm{TC}} \colon R\Gamma_{\mathrm{\acute{e}t}}(A, \mathbb{G}_m)_p^{\wedge}[1] \to \mathrm{TC}(A; \mathbb{Z}_p)$$

Apply $\pi_2(-)[2]$ to this to obtain the morphism

$$T_p(A^{\times})[1] \to (\pi_2 \mathrm{TC}(A; \mathbb{Z}_p))[2].$$

The étale sheafification of the left-hand side is $R\Gamma_{\text{\acute{e}t}}(A, \mathbb{G}_m)_p^{\wedge}[1]$, and the quasi-syntomic sheafification of the right-hand side is $R\Gamma_{\text{syn}}(A, \mathbb{Z}_p(1))[2]$ since

$$R\Gamma_{\mathrm{syn}}(A, \mathbb{Z}_p(1))[2] \simeq (\pi_2 \mathrm{TC}(A; \mathbb{Z}_p))[2]$$

if A is quasi-regular semiperfectoid. Hence we obtain the induced natural morphism

$$\widehat{c}_1^{\mathrm{syn}} \colon R\Gamma_{\mathrm{\acute{e}t}}(A, \mathbb{G}_m)_p^{\wedge}[1] \to R\Gamma_{\mathrm{syn}}(A, \mathbb{Z}_p(1))[2]$$

for every quasi-syntomic ring A. This agrees with \hat{c}_1^{syn} in [8, Theorem 7.5.6] up to shift for quasi-regular semiperfectoid $\mathbb{Z}_p^{\text{cycl}} = \mathbb{Z}_p[\mu_{p^{\infty}}]_p^{\wedge}$ -algebra by comparing [3, Theorem 6.7] and [8, Notation 7.5.1], for quasi-regular semiperfectoid ring by quasi-syntomic descent for the cover $A \to A \widehat{\otimes}_{\mathbb{Z}_p} \mathbb{Z}_p^{\text{cycl}}$, and for quasi-syntomic ring by quasi-syntomic descent again.

For a quasi-regular semiperfectoid ring A such that A is w-local in the sense of [10, Definition 1.6] (in particular any Zariski cover of Spec A is split) and A^{\times} is p-divisible (such rings form a base for the quasi-syntomic topology in light of the proof of [9, Proposition 7.17]), we see that $T_p(A^{\times})[2] \simeq R\Gamma_{\text{ét}}(A, \mathbb{G}_m)_p^{\wedge}[1]$ is concentrated in degree 2. Hence for such a ring A, we have a natural equivalence $\tau_{[1,2]} \hat{c}_1^{\text{TC}} \simeq \hat{c}_1^{\text{syn}}$. In particular, we obtain the natural commutative diagram

$$\tau_{[1,2]} \operatorname{TC}(A; \mathbb{Z}_p) \longleftrightarrow \tau_{\geq 1} \operatorname{TC}(A; \mathbb{Z}_p) \longrightarrow \operatorname{TC}(A; \mathbb{Z}_p).$$

For $\mathfrak{X} \in FlQSyn$, we obtain the natural commutative diagram

by quasi-syntomic descent.

For any *p*-adically complete ring A, write $A\langle x \rangle$ for the *p*-adic completion of the polynomial ring A[x].

Lemma 5.7. For $(A, M) \in \text{IQRSPerfd}$, the BMS filtrations on $\text{THH}((A\langle x \rangle, M); \mathbb{Z}_p)$ and $\text{THH}((A\langle x, x^{-1} \rangle, M); \mathbb{Z}_p)$ are the double-speed Postnikov filtrations. Similar results hold for TC^- and TP too.

Proof. We focus on the case of $(A\langle x \rangle, M)$ since the proofs are similar. By definition of quasi-regular semiperfectoid ring, there exists a perfectoid ring R with a map $R \to A$. By base-change, we have equivalences of complexes

$$\mathbb{L}_{(A[x],M)/R} \simeq \mathbb{L}_{R[x]/R} \otimes_{R}^{\mathbb{L}} A \oplus \mathbb{L}_{(A,M)/R} \otimes_{R}^{\mathbb{L}} R[x] \simeq A[x] \oplus \mathbb{L}_{(A,M)/R} \otimes_{R}^{\mathbb{L}} R[x].$$

After taking p-completions, we get an equivalence of complexes

$$\mathbb{L}_{(A\langle x\rangle,M)/R} \simeq A\langle x\rangle \oplus \mathbb{L}_{(A,M)/R}\widehat{\otimes}_R^{\mathbb{L}} R\langle x\rangle.$$

Since $\mathbb{L}_{(A,M)/R}$ has p-complete Tor amplitude in degree -1 by e.g. [13, Lemma 4.15] and $\wedge_{A\langle x\rangle}^{j}A\langle x\rangle \simeq 0$ for $j \geq 2, \ \wedge_{A\langle x\rangle}^{i}\mathbb{L}_{(A\langle x\rangle,M)/R}$ has *p*-complete Tor amplitude in [-i,-i+1]for $i \geq 0$. By [13, Proposition 7.3] for THH and [9, Theorem 7.2(3),(4), Proposition 7.8] for TC⁻ and TP, we see that the *i*th graded pieces of THH($(A\langle x \rangle, M); \mathbb{Z}_p)$, $TC^{-}((A\langle x\rangle, M); \mathbb{Z}_p)$, and $TP((A\langle x\rangle, M); \mathbb{Z}_p)$ are in $D^{[-2i, -2i+1]}(A\langle x\rangle)$. Using this, we deduce the claim.

The following proposition is crucial for us.

Proposition 5.8. For $\mathfrak{X} \in \operatorname{FlQSyn}_R$ and $i \in \mathbb{Z} \cup \{-\infty\}$, the projective bundle formula for $\text{THH}(\mathfrak{X} \times \mathbb{P}^1; \mathbb{Z}_p)$ restricts to an equivalence

$$\operatorname{Fil}_{i}^{\operatorname{BMS}}\operatorname{THH}(\mathfrak{X} \times \mathbb{P}^{1}; \mathbb{Z}_{p}) \simeq \operatorname{Fil}_{i}^{\operatorname{BMS}}\operatorname{THH}(\mathfrak{X}; \mathbb{Z}_{p}) \oplus \operatorname{Fil}_{i-1}^{\operatorname{BMS}}\operatorname{THH}(\mathfrak{X}; \mathbb{Z}_{p})$$

We have similar equivalences for TC⁻, TP, and TC too.

Proof. By log quasi-syntomic descent, we reduce to the case when $\mathfrak{X} = \mathrm{Spf}(A, M)$ and $(A, M) \in$ lQRSPerfd. By Remark 5.3 and Lemma 5.7, the standard cover on \mathbb{P}^1 yields the cartesian square of S^1 -equivariant filtered complexes

such that the filtrations on the three corners other than $\operatorname{THH}((\mathbb{P}^1_{\operatorname{Spf}(A,M)})_p^{\wedge};\mathbb{Z}_p)$ are the double speed Postnikov filtrations. Next, we note that we have an S^1 -equivariant equivalence

(5.3)
$$\operatorname{THH}((A\langle x\rangle, M); \mathbb{Z}_p) \simeq \operatorname{THH}((A, M); \mathbb{Z}_p) \otimes_{\mathbb{Z}} \operatorname{HH}(\mathbb{Z}[x])$$

and similarly for $A\langle x^{-1}\rangle$ and $A\langle x, x^{-1}\rangle$ (note that the right hand-side is already pcomplete). Hence, the descriptions of $HH(\mathbb{Z}[x])$, $HH(\mathbb{Z}[x^{-1}])$, and $HH(\mathbb{Z}[x, x^{-1}])$ in Proposition 4.5 tell that the double-speed Postnikov filtrations on these agree with the usual Postnikov filtrations: in other words, the equivalence (5.3) is an equivalence of S^1 -equivariant filtered complexes. Hence we obtain the S^1 -equivariant equivalence of filtered complexes

$$\operatorname{THH}((\mathbb{P}^{1}_{\operatorname{Spf}(A,M)})_{p}^{\wedge};\mathbb{Z}_{p}) \simeq \operatorname{THH}(X;\mathbb{Z}_{p}) \otimes_{\mathbb{Z}} \operatorname{HH}(\mathbb{P}^{1})$$

where the filtration on $HH(\mathbb{P}^1)$ is given by Proposition 4.5. Hence we obtain the S^1 equivariant decomposition

(5.4)
$$\operatorname{THH}(\mathfrak{X} \times \mathbb{P}^{1}; \mathbb{Z}_{p}) \simeq \operatorname{THH}(\mathfrak{X}; \mathbb{Z}_{p}) \oplus (\operatorname{THH}(\mathfrak{X}; \mathbb{Z}_{p})[1])\{-1\},$$

D 10

D) (0

where the filtration on $\text{THH}(X;\mathbb{Z}_p)$ is the double-speed Postnikov filtration, and

$$\operatorname{Fil}_{i}(\operatorname{THH}(\mathfrak{X};\mathbb{Z}_{p})[1])\{-1\}) := \operatorname{Fil}_{i-1}\operatorname{THH}(\mathfrak{X};\mathbb{Z}_{p}).$$

From this, we obtain the desired equivalence for THH. To obtain the desired equivalences for TC⁻ and TP, apply $(-)^{hS^1}$ and $(-)^{tS^1}$ to (5.4), and use Lemma 5.7. For TC, use (5.1).

Construction 5.9. For $\mathfrak{X} \in \mathrm{FlQSyn}_B$, Proposition 5.8 yields the natural equivalences of complexes

$$\begin{aligned} &\operatorname{gr}_{i}^{\operatorname{BMS}}\operatorname{THH}(\mathfrak{X};\mathbb{Z}_{p})\oplus\operatorname{gr}_{i-1}^{\operatorname{BMS}}\operatorname{THH}(\mathfrak{X};\mathbb{Z}_{p})\simeq\operatorname{gr}_{i}^{\operatorname{BMS}}\operatorname{THH}(\mathfrak{X}\times\mathbb{P}^{1};\mathbb{Z}_{p}),\\ &\operatorname{gr}_{i}^{\operatorname{BMS}}\operatorname{TC}^{-}(\mathfrak{X};\mathbb{Z}_{p})\oplus\operatorname{gr}_{i-1}^{\operatorname{BMS}}\operatorname{TC}^{-}(\mathfrak{X};\mathbb{Z}_{p})\simeq\operatorname{gr}_{i}^{\operatorname{BMS}}\operatorname{TC}^{-}(\mathfrak{X}\times\mathbb{P}^{1};\mathbb{Z}_{p}),\\ &\operatorname{gr}_{i}^{\operatorname{BMS}}\operatorname{TP}(\mathfrak{X};\mathbb{Z}_{p})\oplus\operatorname{gr}_{i-1}^{\operatorname{BMS}}\operatorname{TP}(\mathfrak{X};\mathbb{Z}_{p})\simeq\operatorname{gr}_{i}^{\operatorname{BMS}}\operatorname{TP}(\mathfrak{X}\times\mathbb{P}^{1};\mathbb{Z}_{p}),\\ &\operatorname{gr}_{i}^{\operatorname{BMS}}\operatorname{TC}(\mathfrak{X};\mathbb{Z}_{p})\oplus\operatorname{gr}_{i-1}^{\operatorname{BMS}}\operatorname{TC}(\mathfrak{X};\mathbb{Z}_{p})\simeq\operatorname{gr}_{i}^{\operatorname{BMS}}\operatorname{TC}(\mathfrak{X}\times\mathbb{P}^{1};\mathbb{Z}_{p}).\end{aligned}$$

As in Construction 4.8 these are identified with the projective bundle formulas for $\mathfrak{X} \times \mathbb{P}^1$

$$\begin{aligned} \operatorname{gr}_{i}^{\mathrm{N}} R\Gamma_{\widehat{\mathbb{A}}}(\mathfrak{X})\{i\} \oplus \operatorname{gr}_{i-1}^{\mathrm{N}} R\Gamma_{\widehat{\mathbb{A}}}(\mathfrak{X})\{i-1\}[-2] &\xrightarrow{\simeq} \operatorname{gr}_{i}^{\mathrm{N}} R\Gamma_{\widehat{\mathbb{A}}}(\mathfrak{X} \times \mathbb{P}^{1})\{i\} \\ \operatorname{Fil}_{i}^{\mathrm{N}} R\Gamma_{\widehat{\mathbb{A}}}(\mathfrak{X})\{i\} \oplus \operatorname{Fil}_{i-1}^{\mathrm{N}} R\Gamma_{\widehat{\mathbb{A}}}(\mathfrak{X})\{i-1\}[-2] &\xrightarrow{\simeq} \operatorname{Fil}_{i}^{\mathrm{N}} R\Gamma_{\widehat{\mathbb{A}}}(\mathfrak{X} \times \mathbb{P}^{1})\{i\} \\ R\Gamma_{\widehat{\mathbb{A}}}(\mathfrak{X})\{i\} \oplus R\Gamma_{\widehat{\mathbb{A}}}(\mathfrak{X})\{i-1\}[-2] &\xrightarrow{\simeq} R\Gamma_{\widehat{\mathbb{A}}}(\mathfrak{X} \times \mathbb{P}^{1})\{i\} \\ R\Gamma_{\operatorname{syn}}(\mathfrak{X}, \mathbb{Z}_{p}(i)) \oplus R\Gamma_{\operatorname{syn}}(\mathfrak{X}, \mathbb{Z}_{p}(i-1))[-2] &\xrightarrow{\simeq} R\Gamma_{\operatorname{syn}}(\mathfrak{X} \times \mathbb{P}^{1}, \mathbb{Z}_{p}(i)) \end{aligned}$$

induced by $(1, \hat{c}_1^{\text{syn}}(\mathcal{O}(1)))$. These are special cases of [8, Lemma 9.1.4(4)–(6)] when \mathfrak{X} has trivial log structure since the two constructions of \hat{c}_1^{syn} discussed in Construction 5.6 agree.

We now fix the spectrum S of a quasi-syntomic ring R.

Definition 5.10. For $X \in \text{SmlSm}/S$ and $i \in \mathbb{Z} \cup \{-\infty\}$, Proposition 5.8 yields a natural equivalence of complexes

(5.5)
$$\operatorname{Fil}_{i}^{\mathrm{BMS}}\mathrm{THH}(X;\mathbb{Z}_{p}) \simeq \Omega_{\mathbb{P}^{1}}\mathrm{Fil}_{i+1}^{\mathrm{BMS}}\mathrm{THH}(X;\mathbb{Z}_{p}).$$

Here, we are implicitly composing with the *p*-completion functor. Note that every $X \in \text{SmlSm}/S$ automatically satisfies the property that $X_p^{\wedge} \in \text{FlQSyn}$. Using this as bonding maps, Proposition 5.4 enables us to define the complete exhaustive filtration

$$\operatorname{Fil}_{i}^{\operatorname{BMS}}\mathbf{THH}(-;\mathbb{Z}_{p}) := (\operatorname{Fil}_{i}^{\operatorname{BMS}}\operatorname{THH}(-;\mathbb{Z}_{p}), \operatorname{Fil}_{i+1}^{\operatorname{BMS}}\operatorname{THH}(-;\mathbb{Z}_{p}), \ldots)$$

on the \mathbb{P}^1 -spectrum $\mathbf{THH}(-;\mathbb{Z}_p) \simeq \operatorname{Fil}_{-\infty}^{\operatorname{BMS}} \mathbf{THH}(-;\mathbb{Z}_p) \in \operatorname{logDA}(S,\mathbb{Z}_p)$. We similarly define the complete exhaustive filtrations

$$\operatorname{Fil}_{i}^{\operatorname{BMS}}\mathbf{TC}^{-}(-;\mathbb{Z}_{p}), \operatorname{Fil}_{i}^{\operatorname{BMS}}\mathbf{TP}(-;\mathbb{Z}_{p}), \operatorname{Fil}_{i}^{\operatorname{BMS}}\mathbf{TC}(-;\mathbb{Z}_{p})$$

Proposition 5.11. For $i \in \mathbb{Z} \cup \{-\infty\}$, we have a natural equivalence in logDA (S, \mathbb{Z}_p)

$$\operatorname{Fil}_{i}^{\operatorname{BMS}}\mathbf{THH}(-;\mathbb{Z}_{p})(1)[2] \simeq \operatorname{Fil}_{i+1}^{\operatorname{BMS}}\mathbf{THH}(-;\mathbb{Z}_{p}).$$

We have similar equivalences for TC^- , TP, and TC too.

Proof. For **THH**, this is a direct consequence of (5.5). The proofs of the other cases are similar.

Definition 5.12. For integers d and i, we define

$$\operatorname{Fil}_{i}^{N}\widehat{\Delta}\{d\} := (\operatorname{Fil}_{i}^{N}\widehat{\Delta}\{d\}, \operatorname{Fil}_{i+1}^{N}\widehat{\Delta}\{d+1\}[2], \ldots),$$
$$\widehat{\Delta}\{d\} := (\widehat{\Delta}\{d\}, \widehat{\Delta}\{d+1\}[2], \ldots),$$
$$\mathbf{M}\mathbb{Z}_{p}^{\operatorname{syn}}(d) := \operatorname{fib}(\varphi_{p} - \operatorname{can}: \operatorname{Fil}_{d}^{N}\widehat{\Delta}\{d\} \to \widehat{\Delta}\{d\}),$$

where the bonding maps for $\operatorname{Fil}_{n}^{N}\widehat{\Delta}\{d\}$, $\widehat{\Delta}\{d\}$, and $\mathbf{M}\mathbb{Z}_{p}^{\operatorname{syn}}(d)$ are obtained by the projective bundle formulas in Construction 5.9, and the morphisms φ_{p} and can are the pointwise cyclotomic trace and canonical morphisms. We will omit $\{d\}$ and (d) if d = 0. In [11, Theorems 3.15, 3.16], (formal variants of) $\widehat{\Delta}$, $\operatorname{Fil}_{0}^{N}\widehat{\Delta}$, and $\mathbf{M}\mathbb{Z}_{p}^{\operatorname{syn}}$ are denoted by $\mathbf{E}^{\widehat{\Delta}}$, $\mathbf{E}^{\operatorname{Fil}\widehat{\widehat{\Delta}}}$, and $\mathbf{E}^{\operatorname{Fsyn}}$.

Proposition 5.13. For $i \in \mathbb{Z}$, we have natural equivalences

$$gr_i^{BMS}\mathbf{THH}(-;\mathbb{Z}_p) \simeq gr_i^{N}\widehat{\boldsymbol{\Delta}}\{i\}[2i],$$

$$gr_i^{BMS}\mathbf{TC}^{-}(-;\mathbb{Z}_p) \simeq Fil_i^{N}\widehat{\boldsymbol{\Delta}}\{i\}[2i],$$

$$gr_i^{BMS}\mathbf{TP}(-;\mathbb{Z}_p) \simeq \widehat{\boldsymbol{\Delta}}\{i\}[2i],$$

$$gr_i^{BMS}\mathbf{TC}(-;\mathbb{Z}_p) \simeq \mathbf{M}\mathbb{Z}_p^{\text{syn}}(i)[2i]$$

in logDA(S, \mathbb{Z}_p).

Proof. By [13, Theorem 1.3(3),(4)], it remains to show that the bonding maps for the left-hand sides can be identified with the bonding maps for the right-hand sides. This is a consequence of Construction 5.9.

The Tate twist is related to the Breuil-Kisin twist in the following sense:

Proposition 5.14. For $d \in \mathbb{Z}$ and $i \in \mathbb{Z} \cup \{-\infty\}$, we have a natural equivalence

$$\operatorname{Fil}_{i}^{\operatorname{N}}\widehat{\mathbf{\Delta}}(d) \simeq \operatorname{Fil}_{i+d}^{\operatorname{N}}\widehat{\mathbf{\Delta}}\{d\}$$

in logDA(S, \mathbb{Z}_p).

Proof. We have

$$\operatorname{Fil}_{i}^{\mathrm{N}}\widehat{\boldsymbol{\Delta}} := (\operatorname{Fil}_{i}^{\mathrm{N}}\widehat{\mathbb{A}}, \operatorname{Fil}_{i+1}^{\mathrm{N}}\widehat{\mathbb{A}}\{1\}[2], \ldots)$$

Tensoring with (d)[2d] in logDA(S) means shifting d terms, so we have

$$\operatorname{Fil}_{i}^{N}\widehat{\Delta}(d)[2d] \simeq (\operatorname{Fil}_{i+d}^{N}\widehat{\mathbb{A}}\{d\}[2d], \operatorname{Fil}_{i+d+1}^{N}\widehat{\mathbb{A}}\{d+1\}[2d+1], \ldots).$$

We obtain the desired equivalence after applying [-2d].

If $i = -\infty$, then observe that Proposition 5.14 yields an equivalence

$$\widehat{\boldsymbol{\Delta}}(d) \simeq \widehat{\boldsymbol{\Delta}}\{d\}$$

in logDA (S, \mathbb{Z}_p) for $d \in \mathbb{Z}$.

The notation $\mathbf{M}\mathbb{Z}_p^{\text{syn}}(i)$ is compatible with the Tate twist in the following sense:

Proposition 5.15. For $d, i \in \mathbb{Z}$, we have a natural equivalence

$$(\mathbf{M}\mathbb{Z}_p^{\mathrm{syn}}(i))(d) \simeq \mathbf{M}\mathbb{Z}_p^{\mathrm{syn}}(i+d)$$

in logDA(S, \mathbb{Z}_p).

Proof. This is a direct consequence of Proposition 5.14.

Theorem 5.16. For $i \in \mathbb{Z}$,

 $\operatorname{Fil}^{i}_{\operatorname{BMS}}\mathbf{THH}(-;\mathbb{Z}_{p}), \operatorname{Fil}^{i}_{\operatorname{BMS}}\mathbf{TC}^{-}(-;\mathbb{Z}_{p}), \operatorname{Fil}^{i}_{\operatorname{BMS}}\mathbf{TP}(-;\mathbb{Z}_{p}), \operatorname{Fil}^{i}_{\operatorname{BMS}}\mathbf{TC}(-;\mathbb{Z}_{p})$ are in logDA $(S,\mathbb{Z}_{p})_{\leq i}^{\operatorname{veff}}$. In particular, there are canonical morphisms

$$\tilde{\mathbf{f}}_n \mathrm{THH}(-; \mathbb{Z}_p) \simeq \Sigma^{2n, n} \tilde{\mathbf{f}}_0 \mathrm{THH}(-; \mathbb{Z}_p) \to \mathrm{Fil}_n^{\mathrm{BMS}} \mathrm{THH}(-; \mathbb{Z}_p),$$

and similarly for \mathbf{TC}^- , \mathbf{TP} and \mathbf{TC} .

Proof. By Proposition 5.11, we reduce to the case when i = -1. It suffices to show the vanishing

$$\operatorname{Hom}_{\operatorname{logDA}(S,\mathbb{Z}_p)}(\Sigma^{\infty}X_+,\operatorname{Fil}_{\operatorname{BMS}}^{-1}\mathbf{THH}(-;\mathbb{Z}_p))=0$$

and similar vanishings for \mathbf{TC}^- , \mathbf{TP} , and \mathbf{TC} . Since $\operatorname{Fil}_{BMS}^{-1}\operatorname{THH}(-;\mathbb{Z}_p) = 0$ by construction and $\operatorname{Fil}_{BMS}^{-1}\operatorname{TC}(-;\mathbb{Z}_p) = 0$ by Proposition 5.5, the claim holds for **THH** and **TC**. Using the compactness of $\Sigma^{\infty}X_+$ in $\operatorname{logDA}(S,\mathbb{Z}_p)$, it suffices to show the vanishing

$$\operatorname{Hom}_{\operatorname{logDA}(S,\mathbb{Z}_p)}(\Sigma^{\infty}X_+, \operatorname{gr}_{-j}^{\operatorname{BMS}}\mathbf{TC}^-(-;\mathbb{Z}_p)) = 0$$

for every integer j > 0 and similar vanishing for **TP**. By Proposition 5.13, it suffices to show the vanishings

$$H_{\mathrm{sNis}}^{-2j}(X,\widehat{\mathbb{A}}\{-j\}) = 0.$$

This follows from $\widehat{\mathbb{A}}_{(A,M)}\{-j\} \in \mathbb{D}^{\geq 0}(A)$ for $(A, M) \in lQSyn$, which can be deduced from the case of $(A, M) \in lQRSPerfd$ by log quasi-syntomic descent. \Box

Question 5.17. As for the HKR filtration, we leave it as an open question whether the slice and BMS filtrations agree on $\mathbf{THH}(-;\mathbb{Z}_p)$, $\mathbf{TC}^-(-;\mathbb{Z}_p)$, $\mathbf{TP}(-;\mathbb{Z}_p)$, and $\mathbf{TC}(-;\mathbb{Z}_p)$. Since T is the analog of S^2 , the double-speed convergence of the very effective slice filtration on **KGL** gives at least an analogy with the double-speed Postnikov filtration defining the BMS filtration.

6. The characteristic p case

Throughout this section, k is a perfect field of characteristic p. Our goal is to identify the graded pieces of the filtered spectra $\mathbf{THH}(-;\mathbb{Z}_p)$, $\mathbf{TC}^-(-;\mathbb{Z}_p)$, $\mathbf{TP}(-;\mathbb{Z}_p)$, and $\mathbf{TC}(-;\mathbb{Z}_p)$ in this case. We will use [9] for the identifications at the level of S^1 -spectra, and then we will use the projective bundle formulas in [8] to identify the bonding maps.

Construction 6.1. For a log smooth saturated pre-log k-algebra (A, M), recall from [11, §4.3] that the chain complex $W\Omega_{(A,M)/k}$ comes equipped with the Nygaard filtration $\operatorname{Fil}_{N}^{i}W\Omega_{(A,M)/k}$ given by the subcomplex

$$p^{i-1}VW(A) \to p^{i-2}VW\Omega^{1}_{(A,M)/k} \to \cdots \to pVW\Omega^{i-2}_{(A,M)/k}$$
$$\to VW\Omega^{i-1}_{(A,M)/k} \to W\Omega^{i}_{(A,M)/k} \to W\Omega^{i+1}_{(A,M)/k} \to \cdots$$

For $X \in lSm/k$, the log Crystalline cohomology $R\Gamma_{crys}(X) := R\Gamma_{Zar}(X, W\Omega_{X/k})$ is then equipped with the Nygaard filtration $\operatorname{Fil}_{i}^{N}R\Gamma_{crys}$ by Zariski descent. Furthermore, we obtain a natural equivalence of complexes

(6.1)
$$\operatorname{gr}_{i}^{\mathrm{N}}R\Gamma_{\mathrm{crys}}(X) \simeq R\Gamma_{\mathrm{Zar}}(X, \tau^{\leq i}\Omega_{X/k})$$

from [11, (4.22.1), (4.22.2)].

By [11, Theorem 4.30], we have a natural equivalence of filtered \mathbb{E}_{∞} -rings

$$R\Gamma_{\widehat{\mathbb{A}}}(X) \simeq R\Gamma_{\mathrm{crys}}(X).$$

Construction 6.2. For $X \in lSm/k$, recall from [8, Construction 7.3.1] the morphism of complexes

$$c_1^{\operatorname{crys}} \colon R\Gamma_{\operatorname{\acute{e}t}}(\underline{X}, \mathbb{G}_m)[1] \to \operatorname{Fil}_1^N R\Gamma_{\operatorname{crys}}(\underline{X})[2].$$

Compose this with the canonical morphism $\operatorname{Fil}_1^N R\Gamma_{\operatorname{crys}}(\underline{X})[2] \to \operatorname{Fil}_1^N R\Gamma_{\operatorname{crys}}(X)[2]$ to obtain

$$c_1^{\operatorname{crys}} \colon R\Gamma_{\operatorname{\acute{e}t}}(\underline{X}, \mathbb{G}_m)[1] \to \operatorname{Fil}_1^N R\Gamma_{\operatorname{crys}}(X)[2].$$

As a consequence of [8, Proposition 7.5.5], we see that the triangle

(6.2)
$$\begin{array}{c} R\Gamma_{\text{\acute{e}t}}(\underline{X}, \mathbb{G}_m)[1] \\ & \overbrace{\mathrm{Fil}_1^N R\Gamma_{\widehat{\mathbb{A}}}}^{c_1^{\widehat{\mathbb{A}}}}(X)[2] \xrightarrow{\simeq} \mathrm{Fil}_1^N R\Gamma_{\mathrm{crys}}(X)[2] \end{array}$$

commutes.

Construction 6.3. For $X \in ISm/k$, recall from [30, Theorem 4.12] that we have the inverse Cartier operator, which is a natural isomorphism of graded \mathcal{O}_X -algebras

$$C_{X/k}^{-1} \colon \Omega_{X/k}^* \xrightarrow{\cong} \mathcal{H}^*(\Omega_{X/k})$$

sending $d \log x$ to $d \log x$ for every local section x of \mathcal{M}_X . Using this, we have the composite morphism

$$c_1^{\operatorname{Hod}} \colon R\Gamma_{\operatorname{Zar}}(\underline{X}, \mathbb{G}_m)[1] \xrightarrow{c_1^{\operatorname{Hod}}} R\Gamma_{\operatorname{Zar}}(X, \Omega^1_{X/k})[1] \xrightarrow{C_{X/k}^{-1}} R\Gamma_{\operatorname{Zar}}(X, \mathcal{H}^1(\Omega_{X/k}))[1],$$

and we have the projective bundle formulas for \mathbb{P}^1

$$R\Gamma_{\operatorname{Zar}}(X,\mathcal{H}^{i}(\Omega_{X/k})) \oplus R\Gamma_{\operatorname{Zar}}(X,\mathcal{H}^{i-1}(\Omega_{X/k}))[-1] \xrightarrow{\simeq} R\Gamma_{\operatorname{Zar}}(X\times\mathbb{P}^{1},\mathcal{H}^{i}(\Omega_{X/k})),$$
$$R\Gamma_{\operatorname{Zar}}(X,\tau^{\leq i}\Omega_{X/k}) \oplus R\Gamma_{\operatorname{Zar}}(X,\tau^{\leq i-1}\Omega_{X/k})[-1] \xrightarrow{\simeq} R\Gamma_{\operatorname{Zar}}(X\times\mathbb{P}^{1},\tau^{\leq i}\Omega_{X/k})$$

induced by $(1, c_1^{\text{Hod}}(\mathcal{O}(1)))$. Using (6.1), the last equivalence can be identified with

$$\operatorname{gr}_{i}^{\mathrm{N}} R\Gamma_{\operatorname{crys}}(X) \oplus \operatorname{gr}_{i-1}^{\mathrm{N}} R\Gamma_{\operatorname{crys}}(X)[-1] \xrightarrow{\simeq} \operatorname{gr}_{i}^{\mathrm{N}} R\Gamma_{\operatorname{crys}}(X \times \mathbb{P}^{1}).$$

Proposition 6.4. For $X \in lSm/k$ and integer *i*, the natural square

commutes.

Proof. It suffices to show that the image of $c_1^{\operatorname{crys}}(\mathcal{O}(1)) \in H^2(\operatorname{Fil}_1^N R\Gamma_{\operatorname{crys}}(X \times \mathbb{P}^1))$ in $H^2(\operatorname{gr}_1^N R\Gamma_{\operatorname{crys}}(X \times \mathbb{P}^1))$ agrees with $c_1^{\operatorname{Hod}}(\mathcal{O}(1))$ since these induce the projective bundle formulas for \mathbb{P}^1 . For this, we reduce to the case when $X = \operatorname{Spec}(k)$ since the general case can be shown by considering the pullbacks of the first Chern classes.

We set $A := k[x, x^{-1}]$ for simplicity of notation. Since $\mathcal{O}(1) \in R\Gamma_{Zar}(\mathbb{P}^1_k, \mathbb{G}_m)$ is the image of $x^{-1} \in A^*$ under the boundary map $A^* \cong R\Gamma_{Zar}(\mathbb{G}_{m,k}, \mathbb{G}_m) \to R\Gamma_{Zar}(\mathbb{P}^1_k, \mathbb{G}_m)$, it suffices to show that the image of $c_1^{\operatorname{crys}}(x^{-1}) \in H^1(\operatorname{Fil}_1^N W\Omega_{A/k})$ in $H^1(\operatorname{gr}_1^N W\Omega_{A/k})$ agrees with $c_1^{\operatorname{Hod}}(x^{-1})$.

Consider the canonical morphism $W\Omega_{A/k} \to \Omega_{A/k}$, which induces $\operatorname{Fil}_1^N W\Omega_{A/k} \to \operatorname{Fil}_1^{\operatorname{Hod}}\Omega_{A/k}$ and hence $\gamma^{\mathrm{dR}} \colon \operatorname{gr}_1^N W\Omega_{A/k} \to \operatorname{gr}_1^{\operatorname{Hod}}\Omega_{A/k}$. The equivalence $\operatorname{gr}_1^N W\Omega_{A/k} \simeq \tau^{\leq 1}\Omega_{A/k}$ is explicitly given by the isomorphism of chain complexes

$$VW(A)/pVW(A) \xrightarrow{F/p} A$$

$$\downarrow^{d} \qquad \qquad \downarrow^{d}$$

$$W\Omega^{1}_{A/k}/VW\Omega^{1}_{A/k} \xrightarrow{F} Z\Omega^{1}_{A/k},$$

where the lower horizontal map is due to [28, (I.3.11.3)]. Together with [28, Proposition I.3.3], we see that γ^{dR} is given by the composite morphism $\tau^{\leq 1}\Omega_{A/k} \to H^1(\Omega_{A/k})[-1] \xrightarrow{C_{X/k}} \Omega^1_{A/k}$. In particular, $H^1(\gamma^{dR})$ is an isomorphism. Hence to compare $c_1^{\text{crys}}(x^{-1})$ and $c_1^{\text{Hod}}(x^{-1})$ in $H^1(\text{gr}_1^{\text{NW}}\Omega_{A/k})$, it suffices to compare their images in $H^1(\text{gr}_1^{\text{Hod}}\Omega_{A/k})$.

By [8, Example 5.2.4, Theorem 7.6.2], the image of $c_1^{\text{crys}}(x^{-1})$ in $H^1(\text{gr}_1^{\text{Hod}}\Omega_{A/k})$ is identified with $c_1^{\text{Hod}}(x^{-1}) = d\log x \in \Omega^1_{A/k}$. To conclude, observe that $H^1(\gamma^{dR}) = C_{X/k} \colon H^1(\Omega_{A/k}) \to \Omega_{A/k}$ sends $c_1^{\text{Hod}}(x^{-1}) = d\log x$ to $d\log x$.

Definition 6.5. The *strict pro-étale topology* on the category of log schemes is the topology generated by the families $\{X_i \to X\}_{i \in I}$ such that $\{\underline{X}_i \to \underline{X}\}_{i \in I}$ is a pro-étale covering Let sproét be the shorthand for this topology. We denote by $L_{\text{sproét}}(-)$ the strict pro-étale sheafification functor.

Recall the following construction from, e.g., [33].

Definition 6.6. Let X be a fine log scheme over k and let dlog: $\mathcal{M}_X^{\mathrm{gp}} \to \Omega_{X/k}^1$ be the natural map. For every integer $r \geq 1$, write dlog[·]: $(\mathcal{M}_X^{\mathrm{gp}})^{\otimes i} \to W_r \Omega_{X/k}^i$ for the map of strict étale sheaves locally given by $m_1 \otimes \ldots \otimes m_i \to \mathrm{dlog}[m_1] \wedge \ldots \wedge \mathrm{dlog}[m_i]$, where [m] denotes the lift of m as in [33, 2.5]. We denote by $W_r \Omega_{X/k,\log}^i \subseteq W_r \Omega_{X/k}^i$ the strict étale subsheaf generated by the image of dlog[·], and by $W \Omega_{X/k,\log}^i$ the limit $\lim_r W_r \Omega_{X/k,\log}^i$ as a strict pro-étale sheaf.

Proposition 6.7. For $X \in \text{SmlSm}/k$ and integer *i*, the sequence of chain complexes of strict pro-étale sheaves

$$0 \to W\Omega^{i}_{X/k,\log}[-i] \to \operatorname{Fil}_{i}^{\operatorname{N}}W\Omega^{\bullet}_{X/k} \xrightarrow{\varphi/p^{i}-1} W\Omega^{\bullet}_{X/k} \to 0$$

is exact in each degree. Furthermore, we have a natural equivalence of complexes of strict pro-étale sheaves $W\Omega^i_{X/k,\log} \simeq R \lim_m W_m \Omega^i_{X/k,\log}$ (that is, $W\Omega^i_{X/k,\log}$ is also the derived inverse limit).

Proof. This is formally identical to [9, Proposition 8.4], using [33, Corollary 2.14] instead of [28, Theorem I.5.7.2].

Remark 6.8. Recall the following fact. For any fs log scheme X and any quasi-coherent sheaf \mathcal{F} on \underline{X} , we have $R\Gamma_{\text{sét}}(X,\mathcal{F}) \simeq R\Gamma_{\text{két}}(X,\mathcal{F})$ (this is due to Kato, see e.g. [16, Proposition 9.2.3]), and this agrees with $R\Gamma_{\text{Zar}}(X,\mathcal{F})$ and $R\Gamma_{\text{sNis}}(X,\mathcal{F})$ by [16, (9.1.1)]. Moreover, we have an equivalence

$$R\Gamma(X, L_{\text{lét}}\mathcal{F}) \simeq \operatorname{colim}_{Y \in X_{\text{div}}} R\Gamma_{\text{két}}(Y, \mathcal{F})$$

by [16, Theorem 5.1.2]. Hence if \mathcal{G} is a complex of bounded below $(\mathbb{P}^n, \mathbb{P}^{n-1})$ -invariant coherent sheaves on SmlSm/k, then they are automatically log étale sheaves.

For $X \in \mathrm{ISm}/k$ and integer $i \geq 0$, we set $B\Omega^{i}_{X/k} := \mathrm{im}(\Omega^{i-1}_{X/k} \xrightarrow{d} \Omega^{i}_{X/k})$ and $Z\Omega^{i}_{X/k} := \mathrm{ker}(\Omega^{i}_{X/k} \xrightarrow{d} \Omega^{i+1}_{X/k})$. The following result is a consequence of [38, Theorem 4.4] and Remark 6.8 above, using, for example, the fact that the above presheaves restricted to smooth k-schemes are reciprocity sheaves (see [18, §11]).

Proposition 6.9. The presheaves of complexes

$$\Omega^{i}_{/k}, \ B\Omega^{i}_{/k}, \ Z\Omega^{i}_{/k}, \ \tau^{\leq i}\Omega_{/k}, \ W_{m}\Omega_{/k}, \ W\Omega_{/k}$$

on SmlSm/k given by

$$X \mapsto R\Gamma_{\operatorname{Zar}}(X, \Omega^{i}_{X/k}), \ R\Gamma_{\operatorname{Zar}}(X, B\Omega^{i}_{X/k}), \ R\Gamma_{\operatorname{Zar}}(X, Z\Omega^{i}_{X/k}), R\Gamma_{\operatorname{Zar}}(X, \tau^{\leq i}\Omega_{X/k}), \ R\Gamma_{\operatorname{Zar}}(X, W_m\Omega_{X/k}), \ R\Gamma_{\operatorname{Zar}}(X, W\Omega_{X/k})$$

are $(\mathbb{P}^n, \mathbb{P}^{n-1})$ -invariant and log étale hypersheaves for all integers m, n > 0 and i.

We note that $R\Gamma_{Zar}(X, W\Omega_{X/k})$ represents Hyodo-Kato cohomology with the trivial log structure on k, see [33, Corollary 1.23].

Proposition 6.10. The presheaf of complexes

$$L_{\text{sét}} W_m \Omega^i_{/k, \log}, \ L_{\text{sproét}} W \Omega^i_{/k, \log}$$

on SmlSm/k given by

$$X \mapsto R\Gamma_{\text{sét}}(X, W_m \Omega^i_{X/k, \log}), \ R\Gamma_{\text{sproét}}(X, W \Omega^i_{X/k, \log})$$

are $(\mathbb{P}^n, \mathbb{P}^{n-1})$ -invariant and log étale hypersheaves for all integers m, n > 0 and i.

Proof. Consider the exact sequence of strict étale sheaves

$$0 \to \Omega^i_{X/k, \log} \to \Omega^i_{X/k} \to \Omega^i_{X/k}/B\Omega^i_{X/k} \to 0$$

obtained by [33, Proposition 2.13]. By [33, (2.12)], Remark 6.8 and induction on m, we see that $L_{\text{sét}}W_m\Omega^i_{/k,\log}$ is $(\mathbb{P}^n, \mathbb{P}^{n-1})$ -invariant and a log étale hypersheaf. Together with Proposition 6.7, we deduce that $L_{\text{spro\acute{e}t}}W\Omega^i_{/k,\log}$ is $(\mathbb{P}^n, \mathbb{P}^{n-1})$ -invariant and a log étale hypersheaf.

Construction 6.11. For $X \in lSm/k$ and integer *i*, we have a natural equivalence of complexes

(6.3)
$$R\Gamma_{\rm syn}(X, \mathbb{Z}_p(i)) \simeq R\Gamma_{\rm spro\acute{e}t}(X, W\Omega^i_{X/k, \log}).$$

by [11, Theorem 4.30] and Proposition 6.7, and see [11, Theorem 4.27] for the φ -compatibility. Together with [8, Theorem 7.3.5], we see that $c_1^{\text{crys}} \colon R\Gamma_{\text{\acute{e}t}}(X, \mathbb{G}_m)[1] \to R\Gamma_{\text{crys}}(X)[2]$ naturally factors through a morphism of complexes

$$c_1^{W\Omega_{\log}} \colon R\Gamma_{\text{\'et}}(X, \mathbb{G}_m)[1] \to R\Gamma_{\text{spro\acute{e}t}}(X, W\Omega^1_{X/k, \log})[1].$$

As a consequence of [8, Proposition 7.5.5], we see that the triangle

(6.4)
$$R\Gamma_{\acute{e}t}(\underline{X}, \mathbb{G}_m)[1] \xrightarrow{c_1^{W\Omega_{\log}}} R\Gamma_{syn}(X)[2] \xrightarrow{\simeq} R\Gamma_{syn\acute{e}t}(X, W\Omega^1_{X/k, \log})[2]$$

commutes.

Construction 6.12. For $X \in lSm/k$, recall from [8, Construction 7.3.1] the morphism of complexes

$$c_1^{\text{syn}} \colon R\Gamma_{\text{\'et}}(\underline{X}, \mathbb{G}_m)[1] \to \operatorname{Fil}_1^{\mathrm{N}} R\Gamma_{\operatorname{crys}}(\underline{X})[2].$$

Compose this with the canonical morphism $\operatorname{Fil}_1^N R\Gamma_{\operatorname{crys}}(\underline{X})[2] \to \operatorname{Fil}_1^N R\Gamma_{\operatorname{crys}}(X)[2]$ to obtain

$$c_1^{\operatorname{crys}} \colon R\Gamma_{\operatorname{\acute{e}t}}(\underline{X}, \mathbb{G}_m)[1] \to \operatorname{Fil}_1^{\operatorname{N}} R\Gamma_{\operatorname{crys}}(X)[2]$$

Definition 6.13. By Proposition 6.9, we can define the objects

$$\operatorname{Fil}_{i}^{\operatorname{conj}} \mathbf{\Omega}_{/k} := (\tau_{i} \Omega_{/k}, (\tau_{\geq i+1} \Omega_{/k})[1], \ldots),$$

$$\operatorname{Fil}_{i}^{N} \mathbf{W} \mathbf{\Omega}_{/k} := (\operatorname{Fil}_{i}^{N} W \Omega_{/k}, (\operatorname{Fil}_{i+1}^{N} W \Omega_{/k})[1], \ldots),$$

$$L_{\operatorname{spro\acute{e}t}} \mathbf{W} \mathbf{\Omega}_{/k, \log}^{i} := (L_{\operatorname{spro\acute{e}t}} W \Omega_{/k, \log}^{i}, L_{\operatorname{spro\acute{e}t}} W \Omega_{/k, \log}^{i+1}[1], \ldots)$$

in $\log DA(k)$, where the bonding maps are given by the following composite morphisms:

$$\tau_{i}\Omega_{/k} \xrightarrow{c_{1}^{\text{Hod}}} \tau_{i}(\Omega_{\mathbb{P}^{1}}\Omega_{/k}[1]) \xrightarrow{\simeq} \Omega_{\mathbb{P}^{1}}(\tau_{\geq i+1}\Omega_{/k})[1],$$

$$\operatorname{Fil}_{i}^{N}W\Omega_{/k} \xrightarrow{c_{1}^{\operatorname{crys}}} \operatorname{Fil}_{i}^{N}(\Omega_{\mathbb{P}^{1}}W\Omega_{/k}[1]) \xrightarrow{\simeq} \Omega_{\mathbb{P}^{1}}(\operatorname{Fil}_{i+1}^{N}W\Omega_{/k})[1],$$

$$L_{\operatorname{spro\acute{e}t}}W\Omega_{/k,\log}^{i} \xrightarrow{c_{1}^{W\Omega_{\log}}} L_{\operatorname{spro\acute{e}t}}\Omega_{\mathbb{P}^{1}}W\Omega_{/k,\log}^{i+1}[1] \xrightarrow{\simeq} \Omega_{\mathbb{P}^{1}}L_{\operatorname{spro\acute{e}t}}W\Omega_{/k,\log}^{i+1}[1].$$

Proposition 6.14. For $i \in \mathbb{Z}$, we have natural equivalences

$$\begin{aligned} \operatorname{gr}_{i}^{\mathrm{N}}\widehat{\boldsymbol{\Delta}} &\simeq \operatorname{Fil}_{\operatorname{conj}}^{i}\boldsymbol{\Omega}_{/k}, \\ \operatorname{Fil}_{i}^{\mathrm{N}}\widehat{\boldsymbol{\Delta}} &\simeq \operatorname{Fil}_{i}^{\mathrm{N}}\mathbf{W}\boldsymbol{\Omega}_{/k}, \\ \widehat{\boldsymbol{\Delta}} &\simeq \mathbf{W}\boldsymbol{\Omega}_{/k}, \\ \mathbf{M}\mathbb{Z}_{p}^{\operatorname{syn}}(i) &\simeq L_{\operatorname{spro\acute{e}t}}\mathbf{W}\boldsymbol{\Omega}_{/k,\log}^{i}[-i] \end{aligned}$$

in $\log DA(k)$.

Proof. By [11, Theorem 4.30, Proposition 5.1] and (6.3), it suffices to show that the bonding maps on both sides are compatible. This follows from Proposition 6.4 and the commutativity of (6.2) and (6.4). \Box

Proof of Theorem 1.4. (1) is a consequence of Lemma 2.10 and Theorems 3.4 and 5.16. (2) is a consequence of (1) and Propositions 5.13 and 6.14, and (3) is a consequence of (2). \Box

7. Very effective slices of Kummer étale K-theory

Let k be a perfect field. In this section, we construct a morphism in $\log SH^{\wedge}_{k\acute{e}t}(k)$ (the hypercomplete version of $\log SH_{k\acute{e}t}(k)$) representing a morphism from the Lichtenbaum étale cohomology to the syntomic cohomology induced by the cyclotomic trace. We also discuss the very effective slices of Kummer étale K-theory.

By Proposition 6.10, the functors $L_{\text{spro\acute{e}t}}W\Omega^{i}_{-/k,\log}$ are log étale hypersheaves, and represent *p*-adic log syntomic cohomology. Hence we obtain the morphism $L_{\text{k\acute{e}t}}M\mathbb{Z} \to M\mathbb{Z}_{p}^{\text{syn}}$ in $\log \mathrm{SH}^{\wedge}_{\text{k\acute{e}t}}(k)$ by adjunction from the morphism $\mathbb{M}\mathbb{Z} \to \mathbb{M}\mathbb{Z}_{p}^{\text{syn}}$ in $\log \mathrm{SH}(k)$ induced by the motivic cyclotomic trace map. We will show that $L_{\text{k\acute{e}t}}\mathbb{M}\mathbb{Z}$ represents the Lichtenbaum étale motivic cohomology.

For every integer i and a commutative ring R, let R(i) be the Nisnevich sheaf of complexes on Sm/k given by the motivic cohomology

$$X \mapsto R\Gamma_{\mathrm{mot}}(X, R(i)).$$

Consider again the functor

$$\omega^* \colon \operatorname{Sh}_{Nis}(\operatorname{Sm}/S, \operatorname{Sp}) \to \operatorname{Sh}_{sNis}(\operatorname{SmlSm}/S, \operatorname{Sp})$$

such that $\omega^* \mathcal{F}(X) := \mathcal{F}(X - \partial X)$ for $\mathcal{F} \in \mathrm{Sh}_{Nis}(\mathrm{Sm}/k, \mathrm{Sp})$ and $X \in \mathrm{SmlSm}/k$.

Proposition 7.1. The presheaf of complexes $L_{k\acute{e}t}\omega^*\mathbb{Z}(i)$ on SmlSm/k is $(\mathbb{P}^n, \mathbb{P}^{n-1})$ -invariant for every integer n > 0 and i.

Proof. For every complex \mathcal{F} , we have the arithmetic fracture square

$$\begin{array}{c} \mathcal{F} \longrightarrow \prod_{\ell} \mathcal{F}_{\ell}^{\wedge} \\ \downarrow \qquad \qquad \downarrow \\ \mathcal{F}_{\mathbb{Q}} \longrightarrow \prod_{\ell} (\mathcal{F}_{\ell}^{\wedge})_{\mathbb{Q}} \end{array}$$

that is cartesian, where ℓ runs over primes, $(-)^{\wedge}_{\ell}$ is the ℓ -adic completion, and $(-)_{\mathbb{Q}}$ is the rationalization. Use this square for

$$\mathcal{F} := \operatorname{fib}(L_{\operatorname{k\acute{e}t}}\omega^*\mathbb{Z}(i)(X) \to L_{\operatorname{k\acute{e}t}}\omega^*\mathbb{Z}(i)(X \times (\mathbb{P}^n, \mathbb{P}^{n-1})))$$

to reduce to showing that the presheaves of complexes $L_{k\acute{e}t}\omega^*\mathbb{Q}(i)$ and $(L_{k\acute{e}t}\omega^*\mathbb{Z}(i))^{\wedge}_{\ell}$ are $(\mathbb{P}^n, \mathbb{P}^{n-1})$ -invariant.

If $X_{\bullet} \to X$ is a Kummer étale hypercover, then $X_{\bullet} - \partial X_{\bullet} \to X - \partial X$ is an étale hypercover. Since [37, Theorem 14.30] implies that $X \in \mathrm{Sm}/k \to R\Gamma_{\mathrm{mot}}(X, \mathbb{Q}(i))$ is an étale hypersheaf, we see that $X \in \mathrm{SmlSm}/k \mapsto R\Gamma_{\mathrm{mot}}(X - \partial X, \mathbb{Q}(i))$ is a Kummer étale hypersheaf. It follows that we have an equivalence of presheaves of complexes with rational coefficients $L_{\mathrm{k\acute{e}t}}\omega^*\mathbb{Q}(i) \simeq L_{\mathrm{sNis}}\omega^*\mathbb{Q}(i)$. The latter is $(\mathbb{P}^n, \mathbb{P}^{n-1})$ -invariant by [16, Proposition 8.1.12]. Hence it remains to show that $(L_{\mathrm{k\acute{e}t}}\omega^*\mathbb{Z}(i))^{\wedge}_{\ell}$ is $(\mathbb{P}^n, \mathbb{P}^{n-1})$ -invariant. Since $(L_{\mathrm{k\acute{e}t}}\omega^*\mathbb{Z}(i))^{\wedge}_{\ell} \simeq \lim_{d} L_{\mathrm{k\acute{e}t}}\omega^*\mathbb{Z}/\ell^d(i)$, it suffices to show that $L_{\mathrm{k\acute{e}t}}\omega^*\mathbb{Z}/\ell^d(i)$ is $(\mathbb{P}^n, \mathbb{P}^{n-1})$ invariant. By induction and using the exact sequence of abelian groups

$$0 \to \mathbb{Z}/\ell \to \mathbb{Z}/\ell^n \to \mathbb{Z}/\ell^{n-1} \to 0,$$

we reduce to showing that $L_{k\acute{e}t}\omega^*\mathbb{Z}/\ell(i)$ is $(\mathbb{P}^n,\mathbb{P}^{n-1})$ -invariant.

Assume $\ell \neq p$. We have an equivalence

$$L_{
m k\acute{e}t}\omega^*\mathbb{Z}/\ell(i)\simeq L_{
m k\acute{e}t}\omega^*\mu_{
ho}^{\otimes i}$$

by [37, Theorem 10.2] since the Kummer étale topology is finer than the strict étale topology. Hence $L_{k\acute{e}t}\omega^*\mathbb{Z}/\ell(i)(X)$ is equivalent to the Kummer étale cohomology $R\Gamma_{k\acute{e}t}(X,\mathbb{Z}/\ell(i))$. This is \Box -invariant and a log étale hypersheaf by [17, Theorem 9.1.5, Proposition 9.1.6], so this is $(\mathbb{P}^n, \mathbb{P}^{n-1})$ -invariant by [17, §3.5]. Assume $\ell = p$. then we have an equivalence of Nisnevich sheaves $\mathbb{Z}/p^n(i) \simeq W_n \Omega^i_{/k,\log}[-i]$ on Sm/k by Geisser-Levine [26, Theorem 8.5]. Hence we have an equivalence of Kummer étale sheaves

(7.1)
$$L_{\text{k\acute{e}t}}\omega^*\mathbb{Z}/p^n(i) \simeq L_{\text{k\acute{e}t}}W_n\Omega^i_{/k,\log}[-i]$$

on SmlSm/k since $\omega^* W_n \Omega^i_{/k,\log} \simeq W_n \Omega^i_{/k,\log}$ by the definition of logarithmic forms. This is $(\mathbb{P}^n, \mathbb{P}^{n-1})$ -invariant by Proposition 6.10.

Definition 7.2. For $X \in \text{SmlSm}/k$, the Kummer étale motivic cohomology $R\Gamma_L(X, \mathbb{Z}(i))$ is the Kummer étale hypersheafification of $X \mapsto R\Gamma_{\text{mot}}(X, \mathbb{Z}(i)) := R\Gamma_{\text{mot}}(X - \partial X, \mathbb{Z}(i))$. For every integer *i*, Proposition 7.1 yields a natural equivalence

(7.2)
$$\operatorname{map}_{\log \operatorname{SH}_{\mathrm{k\acute{e}t}}^{\wedge}(k)}(\Sigma^{\infty}X_{+},\Sigma^{0,i}L_{\mathrm{k\acute{e}t}}\mathbf{M}\mathbb{Z}) \simeq R\Gamma_{\mathrm{L}}(X,\mathbb{Z}(i)),$$

where $\log \mathrm{SH}^{\wedge}_{\mathrm{k\acute{e}t}}(k)$ is the Kummer étale hyper-localization of $\log \mathrm{SH}(k)$, and $L_{\mathrm{k\acute{e}t}}$: $\log \mathrm{SH}(k) \rightarrow \log \mathrm{SH}^{\wedge}_{\mathrm{k\acute{e}t}}(k)$ is the localization functor. The Kummer étale K-theory $L_{\mathrm{k\acute{e}t}}\mathrm{K}(X)$ is the Kummer étale hypersheafification of $X \mapsto \mathrm{K}(X) := \mathrm{K}(X - \partial X)$.

Assume that X has a trivial log structure. Then the small Kummer étale site $X_{k\acute{e}t}$ and small étale site $X_{\acute{e}t}$ agree. Hence $R\Gamma_L(X, \mathbb{Z}(i))$ agrees with the Lichtenbaum étale motivic cohomology, and $L_{k\acute{e}t}K(X)$ agrees with the étale K-theory of X.

We also consider the localization functor $L_{\text{sét}} : \log \text{SH}(k) \to \log \text{SH}^{\wedge}_{\text{sét}}(k)$.

Definition 7.3. For a topology τ on SmlSm/k, let logDA_{τ}(k)^{\wedge}_p be the full subcategory of logDA_{τ}(k) spanned by *p*-complete objects. We define the slice filtration on logDA_{τ}(k)^{\wedge}_p as we did on logDA_{τ}(k).

Proposition 7.4. For every integer i, there are equivalences in $\log DA(k)_p^{\wedge}$

$$(L_{\text{s\acute{e}t}}\mathbf{M}\mathbb{Z}(i))_p^{\wedge} \simeq (L_{\text{k\acute{e}t}}\mathbf{M}\mathbb{Z}(i))_p^{\wedge} \simeq \mathbf{M}\mathbb{Z}_p^{\text{syn}}(i).$$

Proof. We have an equivalence in $\log DA(k)$

$$\mathbf{M}\mathbb{Z}_p^{\mathrm{syn}}(i) \simeq \lim_m (L_{\mathrm{k\acute{e}t}} \mathbf{W}_m \mathbf{\Omega}^i_{/k, \log}[-i])$$

by Proposition 6.7. The right-hand side is equivalent to $(L_{\text{k\acute{e}t}}\mathbf{M}\mathbb{Z}(i))_p^{\wedge}$ by (7.1). To show $(L_{\text{s\acute{e}t}}\mathbf{M}\mathbb{Z}(i))_p^{\wedge} \simeq (L_{\text{k\acute{e}t}}\mathbf{M}\mathbb{Z}(i))_p^{\wedge}$, use Proposition 6.10 and (7.1).

Theorem 7.5. Let k be a perfect field admitting resolution of singularities. Then the induced morphism

$$\tilde{\mathrm{f}}_{i}^{\mathrm{k\acute{e}t}}\mathbf{TC}(-;\mathbb{Z}_{p}) \to \mathrm{Fil}_{i}^{\mathrm{BMS}}\mathbf{TC}(-;\mathbb{Z}_{p})$$

is an equivalence in $\log DA^{\wedge}_{k\acute{e}t}(k)^{\wedge}_p$ for every integer *i*. Here, we denote by $\tilde{f}^{k\acute{e}t}_i$ the *i*th very effective cover in $\log DA^{\wedge}_{k\acute{e}t}(k)^{\wedge}_p$. We have a similar result for $\tilde{f}^{s\acute{e}t}_i$ and $\log DA^{\wedge}_{s\acute{e}t}(k)^{\wedge}_p$ too.

Proof. We focus on the proof for the Kummer étale case since the proof for the strict étale case proceeds verbatim. By Lemma 2.8 and Theorem 3.4, we have $L_{k\acute{e}t} \mathbf{M}\mathbb{Z} \in \log \mathrm{DA}^{\wedge}_{k\acute{e}t}(k)^{\mathrm{veff}}_{\geq i}$. Since the *p*-completion functor $\log \mathrm{DA}^{\wedge}_{k\acute{e}t}(k) \to \log \mathrm{DA}^{\wedge}_{k\acute{e}t}(k)^{\wedge}_{p}$ is a left adjoint, we can show that this preserves (very) effectiveness arguing as in Lemma 2.8. Hence we have $\mathrm{gr}_{i}^{\mathrm{BMS}}\mathbf{TC}(-;\mathbb{Z}_{p}) \in (\log \mathrm{DA}^{\wedge}_{k\acute{e}t}(k)^{\wedge}_{p})^{\mathrm{veff}}_{\geq i}$ using Proposition 7.4. For $j \geq i$, we have

$$\operatorname{fib}(\operatorname{Fil}_{j}^{\operatorname{BMS}}\mathbf{TC}(-;\mathbb{Z}_{p}) \to \operatorname{Fil}_{i-1}^{\operatorname{BMS}}\mathbf{TC}(-;\mathbb{Z}_{p})) \in (\operatorname{logDA}_{\operatorname{k\acute{e}t}}^{\wedge}(k)_{p}^{\wedge})_{\geq i}^{\operatorname{veff}}$$

by induction on j. Take colimits on j to have $\operatorname{Fil}_{i}^{\operatorname{BMS}}\mathbf{TC}(-;\mathbb{Z}_{p}) \in (\operatorname{logDA}_{\operatorname{k\acute{e}t}}^{\wedge}(k)_{p}^{\wedge})_{\geq i}^{\operatorname{veff}}$. Theorem 5.16 finishes the proof.

Remark 7.6. (1) In the proof of Theorem 7.5, we have used in particular the fact that $(L_{\text{k\acute{e}t}}\mathbf{M}\mathbb{Z}(i))_p^{\wedge}$ is very effective in the category $\log \mathrm{DA}^{\wedge}_{\mathrm{k\acute{e}t}}(k)_p^{\wedge}$. Since the inclusion $\log \mathrm{DA}^{\wedge}_{\mathrm{k\acute{e}t}}(k)_p^{\wedge} \subset \log \mathrm{DA}^{\wedge}_{\mathrm{k\acute{e}t}}(k)$ does not obviously preserve the very effective subcategory, we are not claiming that the same holds when we see $(L_{\mathrm{k\acute{e}t}}\mathbf{M}\mathbb{Z}(i))_p^{\wedge}$ as object in $\log \mathrm{DA}^{\wedge}_{\mathrm{k\acute{e}t}}(k)$.

(2) One could reasonably ask if the canonical morphism constructed in Theorem 5.16 becomes an equivalence for k a field (say, assuming resolution of singularities), after applying the canonical functor from logDA (k, \mathbb{Z}_p) to logDA $_{k\acute{e}t}^{\wedge}(k)_p^{\wedge}$, and if this equivalence coincides with the one provided by Theorem 7.5. Equivalently, one could ask if the p-completion of the Kummer étale very effective slice filtration for $TC(-;\mathbb{Z}_p)$ coincides with the Kummer étale very effective slice filtration computed in logDA $_{k\acute{e}t}^{\wedge}(k)_p^{\wedge}$. We expect this to be the case.

Theorem 7.7. Let k be a perfect field admitting resolution of singularities, and assume that the étale cohomological dimension of k is finite.

(1) The filtrations

$$f_{\bullet}^{k\acute{e}t}L_{k\acute{e}t}\mathbf{KGL}, \quad \tilde{f}_{\bullet}^{k\acute{e}t}L_{k\acute{e}t}\mathbf{KGL}, \quad and \quad L_{k\acute{e}t}\tilde{f}_{\bullet}\mathbf{KGL}$$

on $L_{k\acute{e}t} \mathbf{KGL} \in \log \mathrm{SH}^{\wedge}_{k\acute{e}t}(k)$ agree and are complete and exhaustive. (2) There are natural equivalences

$$s_i^{k\acute{e}t} L_{k\acute{e}t} \mathbf{KGL} \simeq \tilde{s}_i^{k\acute{e}t} L_{k\acute{e}t} \mathbf{KGL} \simeq L_{k\acute{e}t} s_i \mathbf{KGL} \simeq L_{k\acute{e}t} \tilde{s}_i \mathbf{KGL} \simeq \Sigma^{2i,i} L_{k\acute{e}t} \mathbf{MZ}$$

in logSH[^]_{k\acute{e}t}(k).

(3) For $X \in \text{SmlSm}/k$, there is a natural equivalence

$$\operatorname{map}_{\operatorname{logSH}_{\operatorname{k\acute{e}t}}^{\wedge}(k)}(\Sigma^{\infty}X_{+}, L_{\operatorname{k\acute{e}t}}\mathbf{KGL}) \simeq L_{\operatorname{k\acute{e}t}}\mathbf{K}(X).$$

(4) For $X \in \text{SmlSm}/k$, there are natural equivalences

$$\max_{\substack{\log \operatorname{SH}_{\operatorname{k\acute{e}t}}^{\wedge}(k,\mathbb{S}/p^n)}} (\Sigma^{\infty} X_+, L_{\operatorname{k\acute{e}t}} \operatorname{\mathbf{KGL}} \otimes_{\mathbb{S}} \mathbb{S}/p^n) \simeq L_{\operatorname{k\acute{e}t}} \operatorname{K}(X) \otimes_{\mathbb{S}} \mathbb{S}/p^n, \\ \max_{\substack{\log \operatorname{SH}_{\operatorname{k\acute{e}t}}^{\wedge}(k)_p^{\wedge}}} (\Sigma^{\infty} X_+, (L_{\operatorname{k\acute{e}t}} \operatorname{\mathbf{KGL}})_p^{\wedge}) \simeq (L_{\operatorname{k\acute{e}t}} \operatorname{K}(X))_p^{\wedge}$$

without the assumption on the étale cohomological dimension.

Proof. As pointed out by Spitzweck-Østvær, the connectivity of $f_n E$ of any motivic spectrum increases with n (see [48, Proposition 5.11]). Hence the very effective slice filtration is automatically complete.

By Proposition 7.1, we have

$$\operatorname{hap}_{\operatorname{logSH}_{\operatorname{k\acute{e}t}}^{\wedge}(k)}(\Sigma^{\infty}X_{+},\Sigma^{0,i}L_{\operatorname{k\acute{e}t}}\mathbf{M}\mathbb{Z})\simeq 0$$

for $X \in \text{SmlSm}/k$ and i < 0. By [16, Corollary 5.1.7] and [17, Proposition 2.7.3], $\Sigma^{\infty}X_{+}$ is compact. Since $L_{\text{k\acute{e}t}}$ preserve colimits, Proposition 3.4 implies that the filtration $L_{\text{k\acute{e}t}} \tilde{\mathbf{f}}_{\bullet} \mathbf{KGL}$ is exhaustive. It follows that we have

$$\operatorname{map}_{\operatorname{logSH}_{\operatorname{k\acute{e}t}}^{\wedge}(k)}(\Sigma^{\infty}X_{+}, L_{\operatorname{k\acute{e}t}}\tilde{\mathrm{f}}^{-1}\mathbf{KGL}) \simeq 0.$$

This implies $L_{\text{k\acute{e}t}}\tilde{f}^{-1}\mathbf{KGL} \in \log \mathrm{SH}^{\wedge}_{\mathrm{k\acute{e}t}}(k)^{\mathrm{eff}}_{\leq -1}$. On the other hand, Lemma 2.8 implies $L_{\mathrm{k\acute{e}t}}\tilde{f}_{0}\mathbf{KGL} \in \log \mathrm{SH}^{\wedge}_{\mathrm{k\acute{e}t}}(k)^{\mathrm{veff}}_{\geq 0}$. From these, we have natural equivalences

$$f_0 L_{k\acute{e}t} \mathbf{KGL} \simeq f_0 L_{k\acute{e}t} \mathbf{KGL} \simeq L_{k\acute{e}t} f_0 \mathbf{KGL}.$$

Apply $\Sigma^{2i,i}$ to this and use Proposition 3.4 to finish the proof of (1).

(2) is an immediate consequence of (1).

Consider the localization functor

$$L_{\text{k\acute{e}t}} \colon \operatorname{Sp}_{\mathbb{P}^1}(\operatorname{Sh}_{sNis}(\operatorname{SmlSm}/k, \operatorname{Sp})) \to \operatorname{Sp}_{\mathbb{P}^1}(\operatorname{Sh}_{\text{k\acute{e}t}}(\operatorname{SmlSm}/k, \operatorname{Sp}))$$

without the further $(\mathbb{P}^n, \mathbb{P}^{n-1})$ -localizations. To show (3), we only need to show that $L_{\text{k\acute{e}t}}\mathbf{KGL}$ is $(\mathbb{P}^n, \mathbb{P}^{n-1})$ -invariant for every integer n. The above argument shows that the filtration $L_{\text{k\acute{e}t}}\tilde{F}_{\geq \bullet}\mathbf{KGL}$ on $L_{\text{k\acute{e}t}}\mathbf{KGL}$ is complete and exhaustive since this argument does not require $(\mathbb{P}^n, \mathbb{P}^{n-1})$ -localizations. Hence it suffices to show that the graded pieces $L_{\text{k\acute{e}t}}\tilde{s}_i\mathbf{KGL}$ are $(\mathbb{P}^n, \mathbb{P}^{n-1})$ -invariant. This is a consequence of Theorem 3.4 and Proposition 7.1.

Argue similarly as above and use \lim_n for (4), but note that the étale \mathbb{Z}/p^n -cohomological dimension (=étale \mathbb{S}/p^n -cohomological dimension) of k is always finite by [6, Théorème X.5.1].

Remark 7.8. Even though $\mathbf{KGL} \in \log \mathrm{SH}(k)$ is \mathbb{A}^1 -invariant, $L_{\mathrm{k\acute{e}t}}\mathbf{KGL} \in \log \mathrm{SH}^{\wedge}_{\mathrm{k\acute{e}t}}(k)$ is not \mathbb{A}^1 -invariant in the above case of k with char k > 0 since $R\Gamma_{\mathrm{L}}$ is not \mathbb{A}^1 -invariant as observed in [37, Lecture 10].

Construction 7.9. Let k be a perfect field admitting resolution of singularities. By Proposition 6.10 and (6.3), log syntomic cohomology on SmlSm/k is a Kummer étale hypersheaf. It follows that the log motivic cyclotomic trace map

$$\mathrm{Tr}\colon\mathbf{KGL}\to\mathbf{TC}$$

in $\log SH(k)$ yields the map

$$L_{\text{két}} \text{Tr} \colon L_{\text{két}} \mathbf{KGL} \to \mathbf{TC}$$

in $\log SH^{\wedge}_{k\acute{e}t}(k)$.

If we apply map_{logSH}^{\wedge}_{két}(k)($\Sigma^{\infty}X_{+}, -$) to $L_{két}$ Tr, then we get

 $L_{k\acute{e}t} \mathbf{K}(X) \to \mathbf{TC}(X)$

by Theorem 7.7(3) when the étale cohomological dimension of k is finite, which is the Kummer étale hypersheafification of the log cyclotomic trace map $K(X) \to TC(X)$. Similarly, if we apply map_{logSH^A_{két}(k)^A_p($\Sigma^{\infty}X_{+}, -)$ to $(L_{két}Tr)^{A}_{p}$, then we get}

$$(L_{\text{két}} \mathcal{K}(X))_p^{\wedge} \to \mathcal{TC}(X; \mathbb{Z}_p)$$

by Theorem 7.7(4).

As an application of Kummer étale K-theory, we show the following:

Proposition 7.10. Let k be a perfect field admitting resolution of singularities. For $X \in \text{SmlSm}/k$, there is a natural equivalence

$$(L_{\mathrm{k\acute{e}t}}\mathrm{K}(X))_p^{\wedge} \simeq \mathrm{TC}(X;\mathbb{Z}_p).$$

Proof. Assume first that $X \in \text{Sm}/k$. Then the equivalence $(L_{\text{ét}} K(X))_p^{\wedge} \simeq \text{TC}(X; \mathbb{Z}_p)$ is due to Geisser and Hesselholt [25, Theorem 4.2.6]. In general, the desired equivalence for $X \in \text{SmlSm}/k$ follows from the Gysin sequence [17, Theorem 3.2.21] by induction on the number of smooth irreducible components r of ∂X . More precisely, if Z is a smooth divisor on proper $X \in \text{Sm}/k$, then we have the fiber sequence in logSH(k)

(7.3)
$$\Sigma^{\infty}(X,Z)_{+} \to \Sigma^{\infty}X_{+} \to \operatorname{Th}(N_{Z}X),$$

where $\operatorname{Th}(N_Z X)$ is the motivic Thom space defined in [17, Definition 3.2.7]. Note that by applying the natural functor $\omega_{\sharp} \colon \log \operatorname{SH}(k) \to \operatorname{SH}(k)$, left adjoint to ω^* , the above sequence reduces to the standard localization sequence in $\operatorname{SH}(k)$. In particular, applying the log K-theory spectrum **KGL**, we obtain the localization sequence in algebraic K-theory in light of [17, Theorem 6.5.7]. On the other hand, if we apply the log TC spectrum **TC** to (7.3) we obtain by definition the Gysin or residue sequence in logarithmic topological cyclic homology. By Construction 7.9, we obtain a commutative diagram of spectra

whose horizontal sequences are fiber sequences. The general case follows similarly by induction on r.

Remark 7.11. Assume that the boundary ∂X of $X \in \text{SmlSm}/k$ has r irreducible components. For integers $m \ge 1$ and $n \le d - r$, the induced natural map

$$\pi_n(R\Gamma_{\mathrm{mot}}(X,\mathbb{Z}/p^m(d))) \to \pi_n(R\Gamma_{\mathrm{L}}(X,\mathbb{Z}/p^m(d)))$$

is an isomorphism. Indeed, if r = 0, then this is a consequence of [26, Theorem 8.5]. If r > 0, then proceed by induction on r, and use the Gysin sequence [17, Theorem 3.2.21] and the five lemma. See [23, Corollary 1.1] for a related result with \mathbb{Q}_p -coefficients and the inequality $n \leq d$.

Remark 7.12. The fact that p-adic étale K-theory is identified with topological cyclic homology holds in larger generality (namely, without smoothness assumption) in light of [20, Theorem C]. However, this does not immediately imply a generalization of Proposition 7.10.

References

- A. ANANYEVSKIY, O. RÖNDIGS, AND P. A. ØSTVÆR, On very effective hermitian K-theory, Math. Z., 294 (2020), pp. 1021–1034.
- [2] T. ANNALA AND R. IWASA, *Motivic spectra and universality of K-theory*. ArXiv preprint: http://arxiv.org/abs/2204.03434, (2023).
- [3] J. ANSCHÜTZ AND A.-C. LE BRAS, The p-completed cyclotomic trace in degree 2, Ann. K-Theory, 5 (2020), pp. 539–580.
- [4] B. ANTIEAU, Periodic cyclic homology and derived de Rham cohomology, Ann. K-Theory, 4 (2019), pp. 505–519.
- [5] B. ANTIEAU, A. MATHEW, M. MORROW, AND T. NIKOLAUS, On the Beilinson fiber square, Duke Math. J., 171 (2022), pp. 3707–3806.
- [6] M. ARTIN, A. GROTHENDIECK, AND J. L. VERDIER, Théorie des topos et cohomologie étale des schémas, vol. 269, 270, 305 of Lecture Notes in Mathematics, Springer-Verlag, 1972–1973. Séminaire de Géométrie Algébrique du Bois-Marie 1963—64.
- [7] T. BACHMANN, The very effective covers of KO and KGL over Dedekind schemes, J. Eur. Math. Soc. (JEMS), (2023). to appear.
- [8] B. BHATT AND J. LURIE, Absolute prismatic cohomology. ArXiv preprint: https://arxiv.org/abs/2201.06120, 2022.
- B. BHATT, M. MORROW, AND P. SCHOLZE, Topological Hochschild homology and integral p-adic Hodge theory, Publ. Math. Inst. Hautes Études Sci., 129 (2019), pp. 199–310.
- [10] B. BHATT AND P. SCHOLZE, The pro-étale topology for schemes, Astérisque, (2015), pp. 99–201.
- [11] F. BINDA, T. LUNDEMO, A. MERICI, AND D. PARK, Logarithmic prismatic cohomology, motivic sheaves, and comparison theorems. ArXiv preprint: https://arxiv.org/abs/2312.13129, 2023.
- [12] F. BINDA, T. LUNDEMO, D. PARK, AND P. A. ØSTVÆR, A Hochschild-Kostant-Rosenberg theorem and residue sequences for logarithmic Hochschild homology, Advances in Math., 435 (2023), p. 109354.
- [13] —, Logarithmic prismatic cohomology via logarithmic THH, Int. Math. Res. Not, 2023 (2023), pp. 19641–19696.
- [14] F. BINDA AND A. MERICI, Connectivity and purity for logarithmic motives, J. Inst. Math. Jussieu, 22 (2023), pp. 335–381.
- [15] F. BINDA, D. PARK, AND P. A. ØSTVÆR, Motives and homotopy theory in logarithmic geometry, C. R., Math., Acad. Sci. Paris, 360 (2022), pp. 717–727.
- [16] , Triangulated categories of logarithmic motives over a field, vol. 433 of Astérisque, Paris: Société Mathématique de France (SMF), 2022.
- [17] —, Logarithmic motivic homotopy theory. ArXiv preprint: https://arxiv.org/abs/2303.02729, 2023.
- [18] F. BINDA, K. RÜLLING, AND S. SAITO, On the cohomology of reciprocity sheaves, Forum Math. Sigma, 10 (2022), pp. Paper No. e72, 111.
- [19] A. J. BLUMBERG AND M. A. MANDELL, Localization theorems in topological Hochschild homology and topological cyclic homology, Geom. Topol., 16 (2012), pp. 1053–1120.
- [20] D. CLAUSEN, A. MATHEW, AND M. MORROW, K-theory and topological cyclic homology of henselian pairs, J. Amer. Math. Soc., 34 (2021), pp. 411–473.
- [21] E. ELMANTO, THH and TC are (very) far from being homotopy functors, J. Pure Appl. Algebra, 225 (2021), p. 12. Id/No 106640.
- [22] E. ELMANTO AND M. MORROW, *Motivic cohomology of equicharacteristic schemes*. ArXiv preprint: https://arxiv.org/abs/2309.08463v1, 2023.

- [23] V. ERTL AND W. NIZIOŁ, Syntomic cohomology and p-adic motivic cohomology, Algebr. Geom., 6 (2019), pp. 100–131.
- [24] E. M. FRIEDLANDER AND A. SUSLIN, The spectral sequence relating algebraic K-theory to motivic cohomology, Ann. Sci. École Norm. Sup. (4), 35 (2002), pp. 773–875.
- [25] T. GEISSER AND L. HESSELHOLT, Topological cyclic homology of schemes, in Algebraic K-theory (Seattle, WA, 1997), vol. 67 of Proc. Sympos. Pure Math., Amer. Math. Soc., Providence, RI, 1999, pp. 41–87.
- [26] T. GEISSER AND M. LEVINE, The k-theory of fields in characteristic p, Invent. Math., 139 (2000), pp. 459–493.
- [27] M. A. HILL, M. J. HOPKINS, AND D. C. RAVENEL, On the nonexistence of elements of Kervaire invariant one, Ann. Math. (2), 184 (2016), pp. 1–262.
- [28] L. ILLUSIE, Complexe de de Rham-Witt et cohomologie cristalline, Ann. Sci. École Norm. Sup. (4), 12 (1979), pp. 501–661.
- [29] D. C. ISAKSEN AND P. A. ØSTVÆR, *Motivic stable homotopy groups*, in Handbook of homotopy theory, Boca Raton, FL: CRC Press, 2020, pp. 757–791.
- [30] K. KATO, Logarithmic structures of Fontaine-Illusie, in Algebraic analysis, geometry, and number theory (Baltimore, MD, 1988), Johns Hopkins Univ. Press, Baltimore, MD, 1989, pp. 191–224.
- [31] M. LEVINE, The homotopy conveau tower, J. Topol., 1 (2008), pp. 217–267.
- [32] —, The homotopy coniveau tower, J. Topol., 1 (2008), pp. 217–267.
- [33] P. LORENZON, Logarithmic Hodge-Witt forms and Hyodo-Kato cohomology, Journal of Algebra, 249 (2002), pp. 247 – 265.
- [34] J. LURIE, Higher topos theory, vol. 170, Princeton, NJ: Princeton University Press, 2009.
- [35] _____, Higher algebra. Available at https://www.math.ias.edu/~lurie/papers/HA.pdf, 2017.
- [36] A. MATHEW, Some recent advances in topological Hochschild homology, Bull. Lond. Math. Soc., 54 (2022), pp. 1–44.
- [37] C. MAZZA, V. VOEVODSKY, AND C. WEIBEL, Lecture notes on motivic cohomology, vol. 2 of Clay Mathematics Monographs, American Mathematical Society, Providence, RI; Clay Mathematics Institute, Cambridge, MA, 2006.
- [38] A. MERICI, A motivic integral p-adic cohomology. ArXiv preprint: https://arxiv.org/abs/2211.14303, 2023.
- [39] F. MOREL, An introduction to A¹-homotopy theory, in Contemporary developments in algebraic Ktheory, vol. XV of ICTP Lect. Notes, Abdus Salam Int. Cent. Theoret. Phys., Trieste, 2004, pp. 357– 441.
- [40] T. NIKOLAUS AND P. SCHOLZE, On topological cyclic homology, Acta Math., 221 (2018), pp. 203–409.
- [41] M. C. OLSSON, The logarithmic cotangent complex, Math. Ann., 333 (2005), pp. 859–931.
- [42] D. PARK, On the log motivic stable homotopy groups, Homology, Homotopy Appl., to appear (2023). ArXiv preprint: https://arxiv.org/abs/2308.07683.
- [43] O. RÖNDIGS AND P. A. ØSTVÆR, Slices of Hermitian K-theory and Milnor's conjecture on quadratic forms, Geom. Topol., 20 (2016), pp. 1157–1212.
- [44] O. RÖNDIGS, M. SPITZWECK, AND P. A. ØSTVÆR, The motivic Hopf map solves the homotopy limit problem for K-theory, Doc. Math., 23 (2018), pp. 1405–1424.
- [45] O. RÖNDIGS, M. SPITZWECK, AND P. A. ØSTVÆR, The first stable homotopy groups of motivic spheres, Ann. Math. (2), 189 (2019), pp. 1–74.
- [46] O. RÖNDIGS, M. SPITZWECK, AND P. A. ØSTVÆR, The second stable homotopy groups of motivic spheres, 2021; to appear in Duke Math. J.
- [47] S. SAITO, Reciprocity sheaves and logarithmic motives. Preprint, (2020).
- [48] M. SPITZWECK AND P. A. ØSTVÆR, Motivic twisted K-theory, Algebr. Geom. Topol., 12 (2012), pp. 565–599.
- [49] V. VOEVODSKY, Open problems in the motivic stable homotopy theory. I, in Motives, polylogarithms and Hodge theory, Part I (Irvine, CA, 1998), vol. 3, I of Int. Press Lect. Ser., Int. Press, Somerville, MA, 2002, pp. 3–34.
- [50] —, A possible new approach to the motivic spectral sequence for algebraic K-theory, in Recent progress in homotopy theory. Proceedings of a conference, Baltimore, MD, USA, March 17–27, 2000, Providence, RI: American Mathematical Society (AMS), 2002, pp. 371–379.
- [51] V. VOEVODSKY, On the zero slice of the sphere spectrum, Tr. Mat. Inst. Steklova, 246 (2004), pp. 106– 115.

DEPARTMENT OF MATHEMATICS "F. ENRIQUES", UNIVERSITY OF MILAN, ITALY *Email address*, F. Binda: federico.binda@unimi.it

DEPARTMENT OF MATHEMATICS AND INFORMATICS, UNIVERSITY OF WUPPERTAL, GERMANY *Email address*: dpark@uni-wuppertal.de

DEPARTMENT OF MATHEMATICS "F. ENRIQUES", UNIVERSITY OF MILAN, ITALY & DEPARTMENT OF MATHEMATICS, UNIVERSITY OF OSLO, NORWAY

Email address: paul.oestvaer@unimi.it & paularne@math.uio.no