

NONLINEAR SCHRÖDINGER EQUATION IN TERMS OF ELLIPTIC AND HYPERELLIPTIC σ FUNCTIONS

SHIGEKI MATSUTANI

ABSTRACT. It is known that the elliptic function solutions of the nonlinear Schrödinger equation are reduced to the algebraic differential relation in terms of the Weierstrass sigma function, $\left[-i\frac{\partial}{\partial t} + \alpha\frac{\partial}{\partial u}\right]\Psi - \frac{1}{2}\frac{\partial^2}{\partial u^2}\Psi + (\Psi^*\Psi)\Psi = \frac{1}{2}(2\beta + \alpha^2 - 3\wp(v))\Psi$, where $\Psi(u; v, t) := e^{\alpha u + i\beta t + c} \frac{e^{-\zeta(v)u} \sigma(u+v)}{\sigma(u)\sigma(v)}$, its dual $\Psi^*(u; v, t)$, and certain complex numbers α, β and c . In this paper, we generalize the algebraic differential relation to those of genera two and three in terms of the hyperelliptic sigma functions.

CONTENTS

1. Introduction	1
2. Genus one case	3
3. Hyperelliptic curve X and sigma functions	5
3.1. Geometrical setting of hyperelliptic curves	5
3.2. Sigma function and its derivatives	7
4. The addition formulae	9
5. Baker function for X	13
6. The algebraic differential relation of genera two and three	14
6.1. Genus two	14
6.2. Genus three	15
Acknowledgments:	15
References	16

1. INTRODUCTION

Our purpose of this work is to investigate the algebraic solutions of the nonlinear Schrödinger (NLS) equation in terms of elliptic and hyperelliptic σ functions.

Date: March 15, 2024.

2020 Mathematics Subject Classification. Primary 14H70, 14H05; Secondary 14H42, 14H45 .

Key words and phrases. Nonlinear Schrödinger equation, hyperelliptic sigma functions .

As mentioned in Section 2, the elliptic solutions of the NLS equation are reduced to the algebraic differential relation [Ka],

$$(1.1) \quad \left[-i \frac{\partial}{\partial t} + \alpha \frac{\partial}{\partial u} \right] \Psi - \frac{1}{2} \frac{\partial^2}{\partial u^2} \Psi + (\Psi^* \Psi) \Psi = \frac{1}{2} (2\beta + \alpha^2 - 3\wp(v)) \Psi,$$

where

$$\Psi(u; v, t) := e^{\alpha u + i\beta t + c} \frac{e^{-\zeta(v)u} \sigma(u+v)}{\sigma(u)\sigma(v)}, \quad \Psi^*(u) := -e^{-\alpha u - i\beta t - c} \frac{e^{-\zeta(v)u} \sigma(u-v)}{\sigma(u)\sigma(v)}$$

for the elliptic curve $E : \widehat{y}^2 = 4(x - e_1)(x - e_2)(x - e_3)$, $\alpha, \beta, c \in \mathbb{C}$, and the Weierstrass sigma functions, σ , and zeta function, ζ , of E . After we fix the point v and e_i so that Ψ^* is the complex conjugate of Ψ and (α, u) belongs to $i\mathbb{R} \times \mathbb{R}$ or $\mathbb{R} \times i\mathbb{R}$, we find the elliptic solutions of the NLS equation. We refer these conditions as the reality conditions of the NLS equation. Determination of the conditions depends on the situations which we require, and is not simple because we tune the moduli parameter e_i as in [Ka].

The construction of the algebraic solutions of the NLS equation was found by Previato 1985 [P] and is written in the book by Belokolos, Bobenko, Enolskii, Its and Matveev [BBEIM] precisely. On the other hand reevaluation of the theory of the hyperelliptic sigma function, which was discovered by Klein [Kl] and studied by Baker [B], has been performed by several authors [BEL1, BEL2, EEL, Ma6]. We found the hyperelliptic solutions of the KdV equation [BEL1, Ma1], MKdV equation [Ma1], Toda lattice equation [KMP], sine-Gordon equation [Ma3] in terms of the sigma functions and meromorphic functions of the symmetric product of algebraic curves. However, hyperelliptic solutions of the NLS equation in terms of the hyperelliptic sigma functions have not yet been obtained. When we obtain them, we can find the algebraic relations among the invariances of the solutions and numerically evaluate the time development for a certain class of curves as in [Ma1, Ma5]; the evaluation is, sometimes, much easier than the approach by means of the theta functions.

In this paper, we leave aside the reality conditions for now and present a generalization of the algebraic differential relation (1.1) of genera two and three, $g = 2, 3$, in terms of the hyperelliptic sigma functions associated with the hyperelliptic solutions of the NLS equation. To obtain this, we express the Baker function in terms of the sigma function reported in [BEL1], though our Baker function differs from that in [BEL1]. Using the Baker equation, we demonstrate this generalization. We note that the Baker equation, or the generalized Lamé equation, of the Baker function contains the potential \wp_{gg} , while the addition theorem of the sigma functions gives the meromorphic functions of the Jacobian J_X , which does not coincide with \wp_{gg} itself except for $g = 1$. Thus, this discrepancy prevents us from obtaining a generalization of (1.1). Even for genera two and three, we find a generalization of the algebraic differential relation (1.1) in this paper. In other words, the purpose of this paper is to show how to overcome the discrepancy so that we can generalize (1.1) associated with the hyperelliptic solution of the NLS equation.

We note that in the NLS equation, we have crucial problems related to the reality conditions depending on the “nonfocusing” and “focusing” NLS equations. These correspond to the sign of the nonlinear terms in the NLS equation [P] and to the choice of (α, u) which belongs to $i\mathbb{R} \times \mathbb{R}$ or $\mathbb{R} \times i\mathbb{R}$. However, we will not address these problems at this stage, because we plan to consider them in the next step after considering the algebraic properties of the generalized algebraic differential relation.

The remainder of this paper is organized as follows. Section 2 is devoted to the genus one case or the derivation of (1.1). Section 3 presents a preliminary of the hyperelliptic sigma function. Section 4 provides the addition formula related to the NLS equation. In Section 5, we demonstrate the Baker functions in terms of hyperelliptic sigma functions. In Section 6, we generalize (1.1) for the genera two and three cases using the Baker function and the addition formulae in Theorems 6.2 and 6.3.

2. GENUS ONE CASE

In this Section, we demonstrate how one gets an elliptic solution of the NLS equation (1.1) [Ka]. Let E be an elliptic curve given by

$$(2.1) \quad E : \frac{1}{4}\widehat{y}^2 = y^2 = x^3 + \lambda_2 x^2 + \lambda_1 x + \lambda_0 = (x - e_1)(x - e_2)(x - e_3),$$

where the e 's are complex numbers, $\lambda_2 = -(e_1 + e_2 + e_3)$ and $\widehat{y} = 2y$. In Weierstrass's convention, $\lambda_2 = 0$, but we leave it nonzero.

The Weierstrass elliptic σ function associated with the curve E is connected with the Weierstrass \wp and ζ functions by

$$(2.2) \quad \wp(u) = -\frac{d^2}{du^2} \log \sigma(u), \quad \zeta(u) = \frac{d}{du} \log \sigma(u).$$

Here the coordinate u in the Jacobian J_E associated with E is given by

$$(2.3) \quad u = \int_{\infty}^{(x,y)} \frac{dx}{2y},$$

with $x(u) = \wp(u)$, $\widehat{y}(u) = \wp'(u)$ and ∞ is the infinite point of E . The Jacobian is given by $J_E = \mathbb{C}/(2\mathbb{Z}\omega' + 2\mathbb{Z}\omega'')$ using the double period $(2\omega', 2\omega'')$ for $\omega' = \int_{\infty}^{(e_1,0)} \frac{dx}{2y}$ and $\omega'' = \int_{\infty}^{(e_3,0)} \frac{dx}{2y}$.

The key to obtain a \wp function solution of the NLS equation is the addition formula,

$$(2.4) \quad \wp(u) - \wp(v) = -\frac{\sigma(u+v)\sigma(u-v)}{[\sigma(v)\sigma(u)]^2}.$$

Let $(\wp(u), \wp'(u)) = (x_1, 2y_1)$ and $(\wp(v), \wp'(v)) = (x_2, 2y_2)$. Using $\frac{d}{du} = 2y_1 \frac{d}{dx_1}$ and $\frac{d}{dv} = 2y_2 \frac{d}{dx_2}$, we find that (2.4) leads to the following relations:

$$(2.5) \quad \zeta(u+v) + \zeta(u-v) - 2\zeta(u) = -\frac{2y_1}{x_2 - x_1},$$

$$(2.6) \quad \zeta(u+v) - \zeta(u) - \zeta(v) = \frac{y_2 - y_1}{x_2 - x_1},$$

$$(2.7) \quad \wp(u+v) + \wp(u-v) = \frac{2f_\Omega(x_1, x_2)}{(x_2 - x_1)^2},$$

$$(2.8) \quad \wp(u-v) = \frac{f_\Omega(x_1, x_2) + 2y_1y_2}{(x_2 - x_1)^2},$$

$$(2.9) \quad \wp(u+v) + \wp(u) + \wp(v) = \left[\frac{y_2 - y_1}{x_2 - x_1} \right]^2 - \lambda_2.$$

where

$$(2.10) \quad f_\Omega(x_1, x_2) = x_1x_2(x_1 + x_2 + 2\lambda_2) + \lambda_1(x_1 + x_2) + \lambda_0.$$

By using (2.5)-(2.9) and by defining $\psi(u; v) = \frac{e^{-\zeta(v)u}\sigma(u+v)}{\sigma(u)\sigma(v)}$, we have the relation,

$$(2.11) \quad \left[\frac{d^2}{du^2} - 2\wp(u) \right] \psi(u; v) = (\wp(v) + \lambda_2)\psi(u; v),$$

which is known as the Lamé equation. ψ is known as the Baker function, or the Baker-Akhiezer function [BEL1, Ma6].

We also have $\frac{d\psi(u; v)}{du} = (\zeta(u+v) - \zeta(u) - \zeta(v))\psi(u; v)$, or

$$(2.12) \quad \left[\frac{d}{du} - (\zeta(u+v) - \zeta(u)) \right] \psi(u; v) = \zeta(v)\psi(u; v).$$

Further, using the Miura relation, $2x + \lambda_2 + \tilde{x} = \mu^2 + \mu_u$ for $\mu := \frac{d}{du} \log \psi(u; v) = \frac{y - \tilde{y}}{x - \tilde{x}}$, we obtain the static MKdV equation,

$$(2.13) \quad 2(\lambda_2 + 3\tilde{x})\mu_u + 6\mu^2\mu_u + \mu_{uuu} = 0.$$

We introduce

$$\begin{aligned}\Psi(u; v, t) &:= e^{\alpha u + i\beta t + c} \psi(u; v) = \frac{e^{\alpha u + i\beta t + c - \zeta(v)u} \sigma(u + v)}{\sigma(u)\sigma(v)}, \\ \Psi^*(u; v, t) &:= e^{-\alpha u - i\beta t - c} \psi(u; -v) = -\frac{e^{-\alpha u - i\beta t - c + \zeta(v)u} \sigma(u - v)}{\sigma(u)\sigma(v)},\end{aligned}$$

for $\alpha, \beta, c \in \mathbb{C}$. Here we use $\sigma(-v) = -\sigma(v)$ and $\zeta(-v) = -\zeta(v)$. We have

$$\left[-i \frac{\partial}{\partial t} + \alpha \frac{\partial}{\partial u} \right] \Psi(u) - \frac{1}{2} \frac{\partial^2}{\partial u^2} \Psi(u) + \wp(u) \Psi(u) = \frac{1}{2} (2\beta + \alpha^2 - \wp(v) - \lambda_2) \Psi(u).$$

Then the addition theorem (2.4), $\Psi^* \Psi = \wp(u) - \wp(v)$, leads the relation,

$$(2.14) \quad \left[-i \frac{\partial}{\partial t} + \alpha \frac{\partial}{\partial u} \right] \Psi - \frac{1}{2} \frac{\partial^2}{\partial u^2} \Psi + (\Psi^* \Psi) \Psi = \frac{1}{2} (2\beta + \alpha^2 - 3\wp(v) - \lambda_2) \Psi,$$

which is the same as (1.1) when $\lambda_2 = 0$.

3. HYPERELLIPTIC CURVE X AND SIGMA FUNCTIONS

In this Section, we provide a background information on the hyperelliptic θ functions and the σ functions as a generalization of the Weierstrass elliptic σ function.

3.1. Geometrical setting of hyperelliptic curves. Let X be a hyperelliptic curve defined by

$$X : y^2 = f(x) := x^{2g+1} + \lambda_{2g} x^{2g} + \cdots + \lambda_0$$

together with a smooth point ∞ at infinity. Here λ 's are complex numbers. The affine ring related to X is denoted as $R_X := \mathbb{C}[x, y]/(y^2 - f(x))$. We fix the basis of the holomorphic one-form

$$\nu_i^I = \frac{x^{i-1} dx}{2y} \quad (i = 1, \dots, g),$$

and the homology basis for the curve X ,

$$H_1(X, \mathbb{Z}) = \bigoplus_{j=1}^g \mathbb{Z} \alpha_j \oplus \bigoplus_{j=1}^g \mathbb{Z} \beta_j,$$

so that their intersections are given by $[\alpha_i, \alpha_j] = 0$, $[\beta_i, \beta_j] = 0$ and $[\alpha_i, \beta_j] = -[\beta_i, \alpha_j] \delta_{ij}$.

We take the half-period matrix $\omega = \begin{bmatrix} \omega' \\ \omega'' \end{bmatrix}$ of X with respect to the given basis where

$$\omega' = \frac{1}{2} \left[\oint_{\alpha_j} \nu_i^I \right], \quad \omega'' = \frac{1}{2} \left[\oint_{\beta_j} \nu_i^I \right].$$

The period matrices $2\omega'$ and $2\omega''$ form the lattice Γ_X in \mathbb{C}^g as a \mathbb{Z} -module. The Jacobian variety of X is denoted by J_X , $\kappa_X : \mathbb{C}^g \rightarrow J_X = \mathbb{C}^g / \Gamma_X$. The abelian covering of X generated by the path space of X with the basepoint $\infty \in X$ is denoted by \tilde{X} , $\kappa_X : \tilde{X} \rightarrow X$

$\kappa_X(\gamma_{P,\infty}) = P$, where $\gamma_{P,\infty}$ is a path from ∞ to $P \in X$. For a non-negative integer k , we define the Abelian integral \tilde{w} from k -th symmetric product $S^k \tilde{X}$ of \tilde{X} to \mathbb{C}^g and the Abel-Jacobi map w from $S^k X$ by

$$\tilde{w} : S^k \tilde{X} \rightarrow \mathbb{C}, \quad \tilde{w}(\gamma_1, \dots, \gamma_k) = \sum_{i=1}^k \int_{\gamma_i} \begin{bmatrix} \nu_1^I \\ \vdots \\ \nu_g^I \end{bmatrix},$$

$$w : S^k X \rightarrow J_X, \quad w((x_1, y_1), \dots, (x_k, y_k)) = \sum_{i=1}^k \int_{\infty}^{(x_i, y_i)} \begin{bmatrix} \nu_1^I \\ \vdots \\ \nu_g^I \end{bmatrix} \bmod \Lambda.$$

The image of w is denoted by $W_X^k = w(S^k X)$, i.e., $J_X = W_X^g$. Furthermore we introduce an injection $\iota_X : X \rightarrow \tilde{X}$ and will fix it. We find that $w = \kappa_J \circ \tilde{w} \circ \iota_X$.

Definition 3.1. Let $(x_i, y_i)_{i=1, \dots, g} \in S^g X$.

(1) We define the polynomials associated with $F(x) = (x - x_1) \cdots (x - x_g)$ by

$$(3.1) \quad \pi_i(x) := \frac{F(x)}{x - x_i} = \chi_{i,g-1}x^{g-1} + \chi_{i,g-2}x^{g-2} + \cdots + \chi_{i,1} + \chi_{i,0},$$

(2) We define $g \times g$ matrices as follows.

$$\mathcal{X}_g := \begin{pmatrix} \chi_{1,0} & \chi_{1,1} & \cdots & \chi_{1,g-1} \\ \chi_{2,0} & \chi_{2,1} & \cdots & \chi_{2,g-1} \\ \vdots & \vdots & \ddots & \vdots \\ \chi_{g,0} & \chi_{g,1} & \cdots & \chi_{g,g-1} \end{pmatrix}, \quad \mathcal{Y}_g := \begin{pmatrix} y_1 & & & \\ & y_2 & & \\ & & \ddots & \\ & & & y_g \end{pmatrix} = \frac{1}{2} \mathcal{H}_g,$$

$$\mathcal{F}'_g := \begin{pmatrix} F'(x_1) & & & \\ & F'(x_2) & & \\ & & \ddots & \\ & & & F'(x_g) \end{pmatrix}, \quad \mathcal{V}_g := \begin{pmatrix} 1 & 1 & \cdots & 1 \\ x_1 & x_2 & \cdots & x_g \\ x_1^2 & x_2^2 & \cdots & x_g^2 \\ \vdots & \vdots & \ddots & \vdots \\ x_1^{g-1} & x_2^{g-1} & \cdots & x_g^{g-1} \end{pmatrix},$$

$$\mathcal{E}_g := \begin{pmatrix} 1 & & & & \\ \varepsilon_{g-1} & 1 & & & 0 \\ \varepsilon_{g-2} & \varepsilon_{g-1} & 1 & & \\ \vdots & \vdots & \vdots & \ddots & \\ \varepsilon_1 & \varepsilon_2 & \cdots & \varepsilon_{g-1} & 1 \end{pmatrix}, \quad \mathcal{K}_g := \begin{pmatrix} x_1^{g-1} & x_1^{g-2} & \cdots & 1 \\ x_2^{g-1} & x_2^{g-2} & \cdots & 1 \\ \vdots & \vdots & \ddots & \vdots \\ x_g^{g-1} & x_g^{g-2} & \cdots & 1 \end{pmatrix},$$

where $F(x) = x^g + \varepsilon_{g-1}x^{g-1} + \cdots + \varepsilon_1x + \varepsilon_0$ and $F'(x) := dF(x)/dx$.

Lemma 3.2. Let $u \in \tilde{w}(\iota_X((x_1, y_1), \dots, (x_g, y_g)))$.

(1) The inverse matrix of \mathcal{X}_g is given as $\mathcal{X}_g^{-1} = \mathcal{F}_g'^{-1} \mathcal{V}_g$.

(2) For $\partial_{u_i} := \partial/\partial u_i$, $\partial_{x_i} := \partial/\partial x_i$, and $\partial_{u_i}^{(r)} := \partial/\partial u_i^{(r)}$, we have

$$\begin{bmatrix} \partial_{u_1} \\ \partial_{u_2} \\ \vdots \\ \partial_{u_g} \end{bmatrix} = 2\mathcal{Y}_g \mathcal{F}_g'^{-1} \mathcal{X}_g \begin{bmatrix} \partial_{x_1} \\ \partial_{x_2} \\ \vdots \\ \partial_{x_g} \end{bmatrix}, \quad \frac{\partial x_i}{\partial u_r} = \frac{2y_i}{F'(x_i)} \chi_{i,r-1}, \quad \frac{\partial y_i}{\partial u_r} = \frac{f'(x_i)}{F'(x_i)} \chi_{i,r-1},$$

$$(3.2) \quad \frac{\partial}{\partial u_g} = \sum_{i=1}^g \frac{2y_i}{F'(x_i)} \frac{\partial}{\partial x_i}, \quad \frac{\partial}{\partial u_{g-1}} = \sum_{i=1}^g \frac{2y_i \chi_{i,g-1}}{F'(x_i)} \frac{\partial}{\partial x_i}.$$

3.2. Sigma function and its derivatives. We define differentials of the second kind,

$$\nu_j^\Pi = \frac{1}{2y} \sum_{k=j}^{2g-j} (k+1-j) \lambda_{k+1+j} x^k dx, \quad (j = 1, \dots, g)$$

and complete hyperelliptic integrals of the second kind

$$\eta' = \frac{1}{2} \left[\oint_{\alpha_j} \nu_i^\Pi \right], \quad \eta'' = \frac{1}{2} \left[\oint_{\beta_j} \nu_i^\Pi \right].$$

For this basis of the $2g$ -dimensional space of meromorphic differentials, the half-periods $\omega', \omega'', \eta', \eta''$ satisfy the generalized Legendre relation

$$(3.3) \quad \mathfrak{M} \begin{pmatrix} 0 & -1_g \\ 1_g & 0 \end{pmatrix} \mathfrak{M}^T = \frac{i\pi}{2} \begin{pmatrix} 0 & -1_g \\ 1_g & 0 \end{pmatrix},$$

where $\mathfrak{M} = \begin{pmatrix} \omega' & \omega'' \\ \eta' & \eta'' \end{pmatrix}$. Let $\tau = \omega'^{-1} \omega''$. The theta function on \mathbb{C}^g with modulus τ and characteristics $\tau a + b$ for $a, b \in \mathbb{C}^g$ is given by

$$\theta \begin{bmatrix} a \\ b \end{bmatrix} (z; \tau) = \sum_{n \in \mathbb{Z}^g} \exp \left[2\pi i \left\{ \frac{1}{2} {}^t(n+a)\tau(n+a) + {}^t(n+a)(z+b) \right\} \right].$$

The σ -function ([B], p.336, [BEL1]), an analytic function on the space \mathbb{C}^g and a theta series having modular invariance of a given weight with respect to \mathfrak{M} , is given by the formula

$$\sigma(u) = \gamma_0 \exp \left\{ -\frac{1}{2} {}^t u \eta' \omega'^{-1} u \right\} \theta \begin{bmatrix} \delta'' \\ \delta' \end{bmatrix} \left(\frac{1}{2} \omega'^{-1} u; \mathbb{T} \right),$$

where δ' and δ'' are half-integer characteristics giving the vector of Riemann constants with basepoint at ∞ and γ_0 is a certain non-zero constant. The σ -function vanishes only on $\kappa_J^{-1}(W_X^{g-1})$ (see for example [B], p. 252). The hyperelliptic \wp and ζ functions are defined by

$$\wp_{ij} = -\frac{\partial^2}{\partial u_i \partial u_j} \log \sigma(u), \quad \zeta_i = \frac{\partial}{\partial u_i} \log \sigma(u).$$

As the Jacobi inversion formula, we have the relation,

$$F(x) = x^g - \sum_{i=1}^g \wp_{gi}(u) x^{i-1}.$$

Let $\{\phi_i(x, y)\}$ be an ordered set of $\mathbb{C} \cup \{\infty\}$ -valued functions over X defined by

$$(3.4) \quad \phi_i(x, y) = \begin{cases} x^i & \text{for } i \leq g, \\ x^{\lfloor (i-g)/2 \rfloor + g} & \text{for } i > g \text{ and } i - g \text{ even,} \\ x^{\lfloor (i-g)/2 \rfloor} y & \text{for } i > g \text{ and } i - g \text{ odd.} \end{cases}$$

Here we note that $\{\phi_i(x, y)\}$ is a set of the bases of R_X as the \mathbb{C} vector space.

Following [O], we introduce a multi-index \mathfrak{h}^n . For n with $1 \leq n < g$, we let \mathfrak{h}^n be the set of positive integers i such that $n + 1 \leq i \leq g$ with $i \equiv n + 1 \pmod{2}$. Namely,

$$\mathfrak{h}^n = \begin{cases} n + 1, n + 3, \dots, g - 1 & \text{for } g - n \equiv 0 \pmod{2} \\ n, n + 2, \dots, g & \text{for } g - n \equiv 1 \pmod{2} \end{cases}$$

and partial derivative over the multi-index \mathfrak{h}^n

$$\sigma_{\mathfrak{h}^n} = \left(\prod_{i \in \mathfrak{h}^n} \frac{\partial}{\partial u_i} \right) \sigma(u),$$

For $n \geq g$, we define \mathfrak{h}^n as empty and $\sigma_{\mathfrak{h}^n}$ as σ itself.

Table 1

genus	$\sigma_{\mathfrak{h}^1}$	$\sigma_{\mathfrak{h}^2}$	$\sigma_{\mathfrak{h}^3}$	$\sigma_{\mathfrak{h}^4}$	$\sigma_{\mathfrak{h}^5}$	$\sigma_{\mathfrak{h}^6}$	$\sigma_{\mathfrak{h}^7}$	$\sigma_{\mathfrak{h}^8}$	\dots
1	σ	σ	σ	σ	σ	σ	σ	σ	\dots
2	σ_2	σ	σ	σ	σ	σ	σ	σ	\dots
3	σ_2	σ_3	σ	σ	σ	σ	σ	σ	\dots
4	σ_{24}	σ_3	σ_4	σ	σ	σ	σ	σ	\dots
5	σ_{24}	σ_{35}	σ_4	σ_5	σ	σ	σ	σ	\dots
6	σ_{246}	σ_{35}	σ_{46}	σ_5	σ_6	σ	σ	σ	\dots
7	σ_{246}	σ_{357}	σ_{46}	σ_{57}	σ_6	σ_7	σ	σ	\dots
8	σ_{2468}	σ_{357}	σ_{468}	σ_{57}	σ_{68}	σ_7	σ_8	σ	\dots
\vdots	\vdots	\vdots	\vdots	\vdots	\vdots	\vdots	\vdots	\vdots	\ddots

For $u \in \mathbb{C}^g$, we denote by u' and u'' the unique vectors in \mathbb{R}^g such that

$$u = 2 {}^t \omega' u' + 2 {}^t \omega'' u''.$$

We define

$$\begin{aligned} L(u, v) &= {}^t u (2 {}^t \eta' v' + 2 {}^t \eta'' v''), \\ \chi(\ell) &= \exp \left\{ 2\pi i \left({}^t \ell' \delta'' - {}^t \ell'' \delta' + \frac{1}{2} {}^t \ell' \ell'' \right) \right\} \quad (\in \{1, -1\}) \end{aligned}$$

for $u, v \in \mathbb{C}^g$ and for $\ell (= 2 {}^t\omega' \ell' + 2 {}^t\omega'' \ell'') \in \Lambda$. Then $\sigma_{\mathfrak{h}^n}(u)$ for $u \in \kappa_J^{-1}(W_X^n)$ satisfies the translational relation ([O], Lemma 7.3):

$$(3.5) \quad \sigma_{\mathfrak{h}^n}(u + \ell) = \chi(\ell) \sigma_{\mathfrak{h}^n}(u) \exp L(u + \frac{1}{2}\ell, \ell) \quad \text{for } u \in \kappa_J^{-1}(W_X^n).$$

Then we also have the translational formulas of \wp and ζ for $u \in \mathbb{C}^g$,

$$(3.6) \quad \wp_{ij}(u + \ell) = \wp_{ij}(u), \quad \zeta_i(u + \ell) = \zeta_i(u) + 2 \sum_{j=1}^g (\eta'_{ij} \ell'_j + \eta''_{ij} \ell''_j).$$

Further for $n \leq g$, we note that $\sigma_{\mathfrak{h}^n}(-u) = (-1)^{ng + \frac{1}{2}n(n-1)} \sigma_{\mathfrak{h}^n}(u)$ for $u \in \kappa_J^{-1}(W_X^n)$, especially,

$$(3.7) \quad \begin{cases} \sigma_{\mathfrak{b}}(-u) = -\sigma_{\mathfrak{b}}(u) & \text{for } u \in \kappa_J^{-1}(W_X^2) \\ \sigma_{\#}(-u) = (-1)^g \sigma_{\#}(u) & \text{for } u \in \kappa_J^{-1}(W_X^1) \end{cases}$$

by Proposition 7.5 in [O].

4. THE ADDITION FORMULAE

In this Section, we give the addition formulae of the hyperelliptic σ functions which are the generalization of the addition formula (2.4). Those then provide the key to construct the hyperelliptic version of the algebraic differential relation (1.1) and (2.14).

Let us introduce the Frobenius-Stickelberger determinant:

Definition 4.1. For a positive integer $n \geq 1$ and $(x_1, y_1), \dots, (x_n, y_n)$ in X , we define the Frobenius-Stickelberger determinant [KMP, Ma6],

$$\Psi_n((x_1, y_1), \dots, (x_n, y_n)) := \begin{vmatrix} 1 & \phi_1(x_1, y_1) & \cdots & \phi_{n-2}(x_1, y_1) & \phi_{n-1}(x_1, y_1) \\ 1 & \phi_1(x_2, y_2) & \cdots & \phi_{n-2}(x_2, y_2) & \phi_{n-1}(x_2, y_2) \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ 1 & \phi_1(x_{n-1}, y_{n-1}) & \cdots & \phi_{n-2}(x_{n-1}, y_{n-1}) & \phi_{n-1}(x_{n-1}, y_{n-1}) \\ 1 & \phi_1(x_n, y_n) & \cdots & \phi_{n-2}(x_n, y_n) & \phi_{n-1}(x_n, y_n) \end{vmatrix},$$

where $\phi_i(x_j, y_j)$'s are the monomials defined in (3.4).

We have the addition formula for the hyperelliptic σ functions as the generalization of (2.4). (Theorem 5.1 in [EEMOP]):

Theorem 4.2. Assume that (m, n) is a pair of positive integers. Let (x_i, y_i) ($i = 1, \dots, m$), (x'_j, y'_j) ($j = 1, \dots, n$) in X and $u \in \kappa_J^{-1}(W_X^m)$, $v \in \kappa_J^{-1}(W_X^n)$ be points such

that $u = \tilde{w}(\iota_X(x_1, y_1), \dots, \iota_X(x_m, y_m))$ and $v = \tilde{w}(\iota_X(x'_1, y'_1), \dots, \iota_X(x'_n, y'_n))$. Then the following relation holds:

$$\begin{aligned}
& \frac{\sigma_{\mathbb{H}^{m+n}}(u+v)\sigma_{\mathbb{H}^{m+n}}(u-v)}{\sigma_{\mathbb{H}^m}(u)^2\sigma_{\mathbb{H}^n}(v)^2} \\
&= \delta(g, m, n) \frac{\prod_{i=0}^1 \Psi_{m+n}((x_1, y_1), \dots, (x_m, y_m), (x'_1, (-1)^i y'_1), \dots, (x'_n, (-1)^i y'_n))}{(\Psi_m((x_1, y_1), \dots, (x_m, y_m))\Psi_n((x'_1, y'_1), \dots, (x'_n, y'_n)))^2} \\
(4.1) \quad & \times \prod_{i=1}^m \prod_{j=1}^n \frac{1}{\Psi_2((x_i, y_i), (x'_j, y'_j))},
\end{aligned}$$

where $\delta(g, m, n) = (-1)^{gm + \frac{1}{2}n(n-1) + mn}$.

Theorem 4.2 with $m = g$ and $n = 1$ leads to the following Corollary, which Belokolos, Enolskii, and Salerno found [BES, Theorem 3.2];

Corollary 4.3. *Let $(x_i, y_i) \in X$ ($i = 1, \dots, g$), $(x, y) := (x'_1, y'_1) \in X$, $u \in \mathbb{C}^g$ and $v \in \kappa_J^{-1}(W_X^1)$ such that $u = w((x_1, y_1), \dots, (x_g, y_g))$ modulo Γ_X and $v = w((x, y))$ modulo Γ_X . Then we have*

$$\begin{aligned}
(4.2) \quad \frac{\sigma(u+v)\sigma(u-v)}{\sigma(u)^2\sigma_{\mathbb{H}^1}(v)^2} &= x^g - \sum_{i=1}^g \wp_{gj}(u)x^{i-1} \\
&= F(x) = (x - x_1)(x - x_2) \cdots (x - x_g).
\end{aligned}$$

Proof: Direct computation shows it. ■

Considering the difference in the differential operators in Jacobian J_X , we generalize (2.5)-(2.9). Let $(x_i, y_i) \in X$ ($i = 1, \dots, g$) and $(x, y) \in X$ as follows.

Lemma 4.4. *For $u = \tilde{w}(\iota_X(x_1, y_1), \dots, \iota_X(x_g, y_g))$ and $v = \tilde{w}(\iota_X(x, y))$ in the case of $k = g$ of Corollary 4.3, the following hold.*

$$\begin{aligned}
(1) \quad & \sum_{i=1}^g e^{i-1} (\zeta_i(u+v) + \zeta_i(u-v) - 2\zeta_i(u)) = - \sum_{\ell=1}^g \frac{2y_\ell F(e)}{(e - x_\ell)(x - x_\ell)F'(x_\ell)}, \\
(2) \quad & \zeta_i(u+v) + \zeta_i(u-v) - 2\zeta_i(u) = - \sum_{\ell=1}^g \frac{2y_\ell \chi_{\ell, i-1}}{(x - x_\ell)F'(x_\ell)}, \\
(3) \quad & \sum_{i=1}^g x_r^{i-1} (\zeta_i(u+v) + \zeta_i(u-v) - 2\zeta_i(u)) = - \frac{2y_r}{(x - x_r)}, \\
(4) \quad & \sum_{i=1}^g x^{i-1} (\zeta_i(u+v) - \zeta_i(u-v) - 2\zeta_i^{\mathbb{H}^1}(v)) = \frac{2yF'(x)}{F(x)} = \sum_{\ell=1}^g \frac{2y}{(x - x_\ell)}, \\
(5) \quad & \sum_{i,j}^g x^{i-1} x_r^{j-1} (\wp_{ij}(u+v) - \wp_{ij}(u-v)) = - \frac{4y_r y}{(x - x_r)^2}.
\end{aligned}$$

where e is a generic parameter. Further we let $\zeta_i^{\natural j}(v) := \left[\frac{\partial}{\partial \tilde{u}_i} \log \sigma_{\natural j}(\tilde{u}) \right]_{u=v}$ for $\tilde{u} \in \mathbb{C}^g$.

Proof: Lemma 3.2 and (3.1) show 1. Noting $\partial/\partial x = \sum_i (2y/x^{i-1})\partial/\partial v_i$, we have 3, 4, and 5. ■

Proposition 4.5. Let $u = \tilde{w}(\iota_X(x_1, y_1), \dots, \iota_X(x_g, y_g))$, and $v = \tilde{w}(\iota_X(x, y))$.

$$(4.3) \quad \wp_{gg}(u+v) + \wp_{gg}(u-v) + 2\wp_{gg}(u) = 2 \left[\sum_r \frac{y_r}{(x-x_r)F'(x_r)} \right]^2 + 2 \frac{f(x)}{F(x)^2} - 2\lambda_{2g} - 2x.$$

Proof: We consider the differential of Lemma 4.4 (ii) with respect to u . We interpret the differential (3.2) with respect to u into differentials with respect to x_r , then we have

$$\begin{aligned} \wp_{gg}(u+v) + \wp_{gg}(u-v) - 2\wp_{gg}(u) &= \sum_\ell \frac{2y_\ell}{F'(x_\ell)} \frac{\partial}{\partial x_\ell} \sum_r \frac{2y_r}{(x-x_r)F'(x_r)} \\ &= 2 \left[\sum_r \frac{y_r}{(x-x_r)F'(x_r)} \right]^2 + \sum_r \frac{2}{F'(x_r)} \frac{\partial}{\partial x} \left[\frac{f(x)}{(x-x_r)F'(x)} \right] \Big|_{x=x_r} \\ &= 2 \left[\sum_r \frac{y_r}{(x-x_r)F'(x_r)} \right]^2 + 2 \left[\frac{f(x)}{F(x)^2} - \lambda_{2g} - x - 2\wp_{gg}(u) \right]. \end{aligned}$$

The second equality is asserted by Lemma 4.6. Lemma 4.7 shows the final step. ■

Lemma 4.6. $\frac{\partial}{\partial x_k} F'(x_k) = \frac{1}{2} \left[\frac{\partial^2}{\partial x^2} F(x) \right]_{x=x_k}$.

Lemma 4.7. For $(\tilde{x}, \tilde{y}) \in X$ and $F(x)$, we have

$$\sum_{k=1}^g \frac{1}{F'(x_k)} \left[\frac{\partial}{\partial x} \left(\frac{f(x)}{(x-\tilde{x})F'(x)} \right) \right]_{x=x_k} = -\frac{f(\tilde{x})}{F(\tilde{x})^2} + \lambda_{2g} + b_r + 2\wp_{gg}.$$

Proof: The relation is proved by the integral.

- (1) Let us consider the polygon expansion of the curve X by D_X associated with the standard homology basis. Let its boundary be denoted by ∂D_X .

$$(4.4) \quad \oint_{\partial D_X} \frac{f(x)}{(x-\tilde{x})F(x)^2} dx = 0.$$

(2) The divisor of the integrand in (4.4) is

$$(4.5) \quad \left(\frac{f(x)}{(x - b_r)F(x)^2} dx \right) = 3 \sum_{i=1, b_i}^{2g+1} (b_i, 0) - (\tilde{x}, \tilde{y}) - (\tilde{x}, -\tilde{y}) \\ - 2 \sum_{i=1}^g (x_i, y_i) - 2 \sum_{i=1}^g (x_i, -y_i) - 3\infty.$$

(3) By considering the local parameter t ($x = 1/t^2$) at ∞ , we have

$$\text{res}_\infty \frac{f(x)}{(x - \tilde{x})F(x)^2} dx = -2(\lambda_{2g} + \tilde{x} + 2\wp_{gg}).$$

(4) By considering the local parameter t ($t = x - \tilde{x}$) at $(\tilde{x}, \pm\tilde{y})$, we obtain

$$\text{res}_{(\tilde{x}, \pm\tilde{y})} \frac{f(x)}{(x - \tilde{x})F(x)^2} dx = \frac{f(\tilde{x})}{F(\tilde{x})^2}.$$

(5) By considering the local parameter t ($t = x - x_k$) at $(x_k, \pm y_k)$, we have

$$\text{res}_{(x_k, \pm y_k)} \frac{f(x)}{(x - \tilde{x})F(x)^2} dx = \frac{1}{F'(x_k)} \left[\frac{\partial}{\partial x} \left(\frac{f(x)}{(x - \tilde{x})F'(x)} \right) \right]_{x=x_k}.$$

Using them we prove the relation. ■

Here we introduce $\tilde{\zeta}_i(P) := - \int_\infty^P \nu_i^\Pi$.

Lemma 4.8. *Let $u = \tilde{w}(\iota_X(x_1, y_1), \dots, \iota_X(x_g, y_g))$, $v = \tilde{w}(\iota_X(x, y))$. Then we have*

$$(4.6) \quad [\zeta_g(u + v) - \zeta_g(u - v) - 2\tilde{\zeta}_g(v)] = \frac{2y}{F(x)}.$$

Proof: The left hand side is the meromorphic function on $J_X \times W_X^1$ because of the translational formula (3.6). The right hand side is a meromorphic function on $S^g X \times X$ and thus, on $J_X \times W_X^1$ by the Abel-Jacobi theorem. Thus we show the equality of both sides. We compare the divisors. As a function of v , both sides vanish at every finite branch point $(x, y) = (b_r, 0)$ from the translational formula (3.6) and the definition of $\tilde{\zeta}_g$. At $v = 0$ or $P = \infty$, both sides diverge in order of one. At $v = u$ or $v = -u$, corresponding to $x = x_i$, the order of the pole is one. We note that both sides are odd functions with respect to v . For v , we expand both sides at $v = 0$,

$$\zeta_g(u + v) - \zeta_g(u - v) = -2 \sum_j \wp_{gj} v_j - \frac{2}{3!} \sum_{jkl} \wp_{gjk\ell} v_j v_k v_\ell + \text{the order of } v \geq 3$$

The local parameter $t = x^{g+1}/y$ shows

$$\zeta_g(u + v) - \zeta_g(u - v) - 2\tilde{\zeta}_g(v) = 2(x_1 + \dots + x_g)t + \frac{2}{t} + \text{the order of } t > 0.$$

$$\frac{2\tilde{y}}{F(\tilde{x})} = \frac{2}{t} (1 + (x_1 + \dots + x_g)t^2 + \text{the order of } t > 0).$$

The leading terms of both sides match. Hence we prove it. ■

Remark 4.9. Lemma 4.8 may be generalized as follows.

$$(4.7) \quad \begin{aligned} [\zeta_{g-1}(u+v) - \zeta_{g-1}(u-v) + 2\tilde{\zeta}_{g-1}(v)] &= \frac{2y}{F(x)}(x - \wp_{gg}), \\ [\zeta_{g-2}(u+v) - \zeta_{g-2}(u-v) + 2\tilde{\zeta}_{g-2}(v)] &= \frac{2y}{F(x)}(x^2 - \wp_{gg}x + \wp_{g,g-1}). \end{aligned}$$

Using above relations, we have the following lemma:

Lemma 4.10. For $u = \tilde{w}(\iota_X(x_1, y_1), \dots, \iota_X(x_g, y_g))$ and $v = \tilde{w}(\iota_X(x, y))$

$$\begin{aligned} \zeta_g(u+v) - \zeta_g(u) - \tilde{\zeta}_g(v) &= \frac{y}{F(x)} - \sum_{r=1}^g \frac{y_r}{(x - x_r)F'(x_r)}, \\ \wp_{gg}(u+v) - \wp_{gg}(u-v) &= - \sum_{r=1}^g \frac{4yy_r}{(x - x_r)F'(x_r)F(x)}, \\ \wp_{gg}(u+v) + \wp_{gg}(u) + x &= \left[\sum_r \frac{y_r}{(x - x_r)F'(x_r)} - \frac{y}{F(x)} \right]^2 - \lambda_{2g}. \end{aligned}$$

5. BAKER FUNCTION FOR X

Let us generalize (2.11) and $\psi(u; v) = \frac{e^{-\zeta(v)u}\sigma(u+v)}{\sigma(u)\sigma(v)}$ of genus one to general g .

Let us define

$$(5.1) \quad \psi(u; v) = \frac{e^{-\sum_i \tilde{\zeta}_i(P)u_i}\sigma(u+v)}{\sigma(u)\sigma_{\mathfrak{h}_1}(v)}$$

Here $\tilde{\zeta}_i(P) := - \int_{\infty}^P \nu_i^{\Pi}$ for $u \in \mathbb{C}^g$ and $P \in X$, and $v = \tilde{w}(\iota_X P)$. We define it as a meromorphic function of $J_X \times W_X^1$ while Buchstaber, Enolskii, and Leykin defined it as $J_X \times J_X$ [BEL1]. Thus ours is different from the one in [BEL1].

Proposition 4.5, Lemmas 4.8 and 4.10 show the generalization of (2.11).

Proposition 5.1. For $u = \tilde{w}(\iota_X(x_1, y_1), \dots, \iota_X(x_g, y_g))$ and $v = \tilde{w}(\iota_X(\tilde{x}, \tilde{y}))$, we have

$$(5.2) \quad \left[\frac{\partial^2}{\partial u_g^2} - 2\wp_{gg}(u) \right] \psi(u; v) = (\tilde{x} + \lambda_{2g})\psi(u; v),$$

$$(5.3) \quad \left[\frac{\partial}{\partial u_g} - (\zeta_g(u+v) - \zeta_g(u)) \right] \psi(u; v) = \tilde{\zeta}_g(v)\psi(u; v).$$

Proof: The same as the case of $g = 1$. ■

6. THE ALGEBRAIC DIFFERENTIAL RELATION OF GENERA TWO AND THREE

We extend the algebraic differential relation (1.1) of genus one to genera two and three.

6.1. Genus two. Let us consider only the genus $g = 2$ case as follows: We assume $(P_1, P_2, \tilde{P}) \in S^2X \times X$, $(P_i = (x_i, y_i)$, and $\tilde{P} = (\tilde{x}, \tilde{y})$ such that $u = \tilde{w}(\iota_X(P_1, P_2))$ and $v = \tilde{w}(\iota_X(\tilde{P}))$.

Corollary 4.3 of $g = 2$ shows

$$(6.1) \quad \frac{\sigma(u+v)\sigma(u-v)}{\sigma(u)^2\sigma_{\mathfrak{h}_1}(v)^2} = \tilde{x}^2 - \wp_{22}\tilde{x} - \wp_{21}.$$

We immediately obtain the following relation from (6.1), though we do not use it in this paper,

Lemma 6.1. *Let us consider the case $\tilde{P}^{(\pm)} = (\pm\tilde{x}, \tilde{y}^{(\pm)})$ and $v^{(\pm)} := w(P^{(\pm)})$. Then we have*

$$(6.2) \quad \frac{\sigma(u+v^{(+)})\sigma(u-v^{(+)})}{\sigma(u)^2\sigma_{\mathfrak{h}_1}(v^{(+)})^2} - \frac{\sigma(u+v^{(-)})\sigma(u-v^{(-)})}{\sigma(u)^2\sigma_{\mathfrak{h}_1}(v^{(-)})^2} = -2\wp_{22}\tilde{x}.$$

Now we state our first main theorem in this paper, in which we obtain the generalization of (2.14) or (1.1).

Theorem 6.2. *For $(P_1, P_2, \tilde{P}) \in S^2X \times X$ such that 1) $x_1x_2 = -\tilde{x}^2$, or 2) $x_1x_2 = 0$, and $\alpha \in \mathbb{C}^2$ and $c, \beta \in \mathbb{C}$, we let*

$$\Psi(u; v, t) := \frac{1}{\sqrt{\tilde{x}}} e^{t\alpha u + i\beta t + c} \psi(u; v), \quad \Psi^*(u; v, t) := \frac{1}{\sqrt{\tilde{x}}} e^{-t\alpha u - i\beta t - c} \psi(u; -v).$$

The following formulae ($g = 2$) hold for the conditions 1) and 2),

$$\begin{aligned} 1) : \quad & \left[-i \frac{\partial}{\partial t} - \alpha_g \frac{\partial}{\partial u_g} \right] \Psi - \frac{1}{2} \frac{\partial^2}{\partial u_g^2} \Psi + (\Psi^* \Psi) \Psi = \frac{1}{2} (2\beta + \alpha_g^2 - \tilde{x} - \lambda_2) \Psi, \\ 2) : \quad & \left[-i \frac{\partial}{\partial t} - \alpha_g \frac{\partial}{\partial u_g} \right] \Psi - \frac{1}{2} \frac{\partial^2}{\partial u_g^2} \Psi + (\Psi^* \Psi) \Psi = \frac{1}{2} (2\beta + \alpha_g^2 - 3\tilde{x} - \lambda_2) \Psi. \end{aligned}$$

Proof: From (5.2), we have

$$\left[-i \frac{\partial}{\partial t} - \alpha_g \frac{\partial}{\partial u_g} \right] \Psi - \frac{1}{2} \frac{\partial^2}{\partial u_g^2} \Psi + \wp_{gg} \Psi = \frac{1}{2} (2\beta + \alpha_g^2 - \tilde{x} - \lambda_2) \Psi.$$

Under the restriction 1), $(\Psi^* \Psi)$ is reduced to $(1/\tilde{x})$ times $\tilde{x}\wp_{22}(u)$. whereas under the restriction 2), $(\Psi^* \Psi)$ is equal to $-(\tilde{x} - \wp_{22}(u))$. ■

6.2. Genus three. Let us consider the genus $g = 3$ case as follows. We assume $(P_1, P_2, P_3, \tilde{P}) \in S^3X \times X$, $(P_i = (x_i, y_i))$, and $\tilde{P} = (\tilde{x}, \tilde{y})$ such that $u = \tilde{w}(\iota_X(P_1, P_2, P_3))$ and $v = \tilde{w}(\iota_X(\tilde{P}))$.

Corollary 4.3 of $g = 3$ shows

$$(6.3) \quad \frac{\sigma(u+v)\sigma(u-v)}{\sigma(u)^2\sigma_{\mathfrak{h}_1}(v)^2} = \tilde{x}^3 - \wp_{33}\tilde{x}^2 - \wp_{32}\tilde{x} - \wp_{31}.$$

Now we state our second main theorem in this paper, in which we obtain the generalization of (2.14) or (1.1).

Theorem 6.3. *For $(P_1, P_2, P_3, \tilde{P}) \in S^3X \times X$ such that $\tilde{x}(x_1x_2 + x_2x_3 + x_3x_1) - x_1x_2x_3 = 0$, $\alpha \in \mathbb{C}^g$ and $c, \beta \in \mathbb{C}$, we let*

$$\Psi(u; v, t) := \frac{1}{\tilde{x}} e^{t\alpha u + i\beta t + c} \psi(u; v), \quad \Psi^*(u; v, t) := \frac{1}{\tilde{x}} e^{-t\alpha u - i\beta t - c} \psi(u; -v).$$

The following formula ($g = 3$) holds.

$$\left[-i \frac{\partial}{\partial t} - \alpha_g \frac{\partial}{\partial u_g} \right] \Psi - \frac{1}{2} \frac{\partial^2}{\partial u_g^2} \Psi + (\Psi^* \Psi) \Psi = \frac{1}{2} (2\beta + \alpha_g^2 - 3\tilde{x} - \lambda_4) \Psi.$$

Proof: From (5.2), we have

$$\left[-i \frac{\partial}{\partial t} - \alpha_g \frac{\partial}{\partial u_g} \right] \Psi - \frac{1}{2} \frac{\partial^2}{\partial u_g^2} \Psi + \wp_{gg} \Psi = \frac{1}{2} (2\beta + \alpha_g^2 - \tilde{x} - \lambda_4) \Psi.$$

Under the restriction, $(\Psi^* \Psi)$ is reduced to $(1/\tilde{x})^2$ times $-(\tilde{x}^3 - \tilde{x}^2 \wp_{33}(u))$. ■

Remark 6.4. As mentioned in Introduction the Baker equation (5.2) contains the potential \wp_{gg} , while the terms \wp in $F(\tilde{x})$ in the addition theorem (4.2) do not coincide with \wp_{gg} except for $g = 1$. Thus, this discrepancy prevents us from obtaining the generalization of (1.1) directly for a long time.

Even for genera two and three, we found the generalization by restricting the configurations of points in S^gX . In other words, the restriction may be effective for the hyperelliptic solution of the NLS equation in terms of σ function, allowing us to easily find a generalization to the general genus.

However we should consider how the restriction effects on the Jacobian J_X . Further we note that in the NLS equation, we have crucial problems related to the reality conditions depending on the “nonfocusing” and “focusing” NLS equations [P] or the choice of (α, u) which belongs to $i\mathbb{R} \times \mathbb{R}$ or $\mathbb{R} \times i\mathbb{R}$. Since these problems are basically complicated as in [Ma4], we will consider them as the next steps.

Acknowledgments: The author is grateful to Professor Emma Previato for suggestions on this study at a very early stage and to Dr. Kayo Kinjo for discussing the algebraic solution of the NLS equation based on [BBEIM]. The author has been supported by the Grant-in-Aid for Scientific Research (C) of the Japan Society for the Promotion of Science Grant, No.21K03289.

REFERENCES

- [B] H.F. Baker, Abelian Functions, Abel's theorem and the allied theory of theta functions, Cambridge University Press, Cambridge, 1897.
- [BES] E. D. Belokolos, V.Z. Enolskii, and M. Salerno, *Wannier functions for quasi-periodic finite-gap potentials*, Theor. Math. Phys., **144** (2005) 1081-1099.
- [BBEIM] E. D. Belokolos, A. I. Bobenko, V. Z. Enolskii, A. R. Its, and V. B. Matveev, *Algebro-geometric approach to nonlinear integrable equations*, Springer-Verlag, New York, 1994.
- [BEL1] V. M. Buchstaber, V. Z. Enolskii, and D. V. Leykin, *Kleinian Functions, Hyperelliptic Jacobians and Applications*, Reviews in Mathematics and Mathematical Physics (London), eds. Novikov, S. P. and Krichever, I. M. Gordon and Breach, India, 1997 1-125.
- [BEL2] V. M. Buchstaber, V. Z. Enolskii, and D. V. Leykin, *σ functions: old and new results*, in Integrable systems and algebraic geometry vol.2, edited by R. Donagi and T. Shaska, London Math Soc. Lect. Note Series **459** (2020) 175–214.
- [EEL] J. C. Eilbeck, V. Z. Enolskii and D. V. Leykin, , *On the Kleinian construction of Abelian functions of canonical algebraic curves*, Proceedings of the 1998 SIDE III conference (2000).
- [EEMOP] J.C. Eilbeck, V.Z. Enol'skii, S. Matsutani, Y. Ônishi, and E. Previato, *Addition formulae over the Jacobian pre-image of hyperelliptic Wirtinger varieties*, Journal für die reine und angewandte Mathematik (Crelles Journal), (2008) **2008** No. 619 37-48
- [Ka] A.M. Kamchatnov New approach to periodic solutions of integrable equations and nonlinear theory of modulational instability Physics Reports 286 (1997) 199-270.
- [Kl] F. Klein, *Ueber hyperelliptische sigmafunctionen*, Math. Ann., **27** (1886) 431–464.
- [KMP] Y. Kodama, S. Matsutani, and E. Previato, *Quasi-periodic and periodic solutions of the Toda lattice via the hyperelliptic sigma function*, Annales de l'institut Fourier, (2012) **63** (2013) 655–688
- [Ma1] S. Matsutani, *Hyperelliptic solution of KdV and KP equations: re-evaluation of Baker's study on hyperelliptic sigma functions*, J. Phys. A: Math. Gen., **34** (2001) 4721-4732.
- [Ma2] S. Matsutani, *Hyperelliptic solutions of modified Kortweg-de Vries equation of genus g: essentials of Miura transformation*, J. Phys. A: Math. & Gen., **35** (2002) 4321–4333,
- [Ma3] S. Matsutani, *Relations of al functions over subvarieties in a hyperelliptic Jacobian*, CUBO A Math. J., **7** (2005) 75–85.
- [Ma4] S. Matsutani, *On real hyperelliptic solutions of focusing modified KdV equation*, arXiv:2309.04904,
- [Ma5] S. Matsutani, *A graphical representation of hyperelliptic KdV solutions*, arXiv:2310.14656 , (2023)
- [Ma6] S. Matsutani, *The Weierstrass sigma function in higher genus and applications to integrable equations*, Springer-Verlag, to appear, .
- [MP] S. Matsutani, E. Previato, *An algebro-geometric model for the shape of supercoiled DNA*, Physica D **430** (2022) 133073
- [O] Y. Ônishi, *Determinant expressions for hyperelliptic functions*, Proc. Edinburgh Math. Soc., **48** (2005) 705-742.
- [P] E. Previato, *Hyperelliptic quasi-periodic and soliton solutions of the nonlinear Schrödinger equation*, Duke Math. J., **52** (1985) 329–377.