

A Plausible Path Towards Unification of Interactions via Gauge Fields Consistent with the Equivalence Principle-II

James Lindesay*

Computational Physics Laboratory
Howard University, Washington, D.C. 20059

Abstract

An extension of the Lorentz group that includes generators Γ^μ carrying a space-time index has been previously demonstrated to *explicitly* construct the Minkowski metric *within* the internal group space as a consequence of the non-vanishing commutation relations between those generators. Fields that transform under representations of this extended group can fundamentally incorporate microscopic causality as a crucial property of physical fields. The first part of this exploration focused on the fundamental representation fermions (which satisfy the Dirac equation), and explored additional internal symmetries associated with those fermions. Any interactions that could result from those symmetries were demonstrated to necessarily be consistent with gravitational equivalence under curvilinear extensions of the (abelian) space-time translations.

The first boson representation of this algebra is the focus of this second paper. In particular, the equations of motion for a group of massive vector bosons are degenerate with that of massless vector bosons, allowing them to unitarily mix to form physical states with differing masses and dynamics. Thus, this representation exhibits a potential for enhancing insights into the standard modeling of electro-weak mixing of bosons. The various spinors that represent these bosons exhibit kinematic factors, and those factors are related during unitary mixing to generate the resultant physical states. For this reason, analytic and kinematic coincidences associated with known electro-weak masses will be explored for insights into possible predictive relationships between their masses and those of this first causal boson representation. To conclude, a plausible model will be constructed using examined coincidences for critique and insights into the potential viability of the approach.

1 Introduction

The relativistic formulation of electron dynamics by Dirac[1] incorporates the charge-conjugate spinor components necessary to insure microscopic causality[2]

*E-mail address, jlindesay@howard.edu

into an equation that is linear in the energy-momentum operator. Some properties of fundamental fermions that require $\hat{\Gamma}^\mu \hat{P}_\mu$ to be a Lorentz scalar operation satisfying

$$\mathbf{\Gamma}^\mu \cdot \frac{\hbar}{i} \frac{\partial}{\partial x^\mu} \hat{\Psi}_\gamma^{(\Gamma)}(\vec{x}) = -\gamma mc \hat{\Psi}_\gamma^{(\Gamma)}(\vec{x}), \quad (1.1)$$

have been explored in part I of this presentation[3]. In this formulation, the masses (which are always non-negative) label the standard states, and the factor γ is the eigenvalue of the hermitian operator $\hat{\Gamma}^0$. Dirac fields transform under the $\Gamma = \frac{1}{2}$ finite dimensional representation of this extended group, and the Dirac matrices γ^μ are given by $\gamma^\mu = 2\mathbf{\Gamma}^\mu$.

In this paper, spinors transforming under the $\Gamma = 1$ finite dimensional representation of this extended group will be developed. In particular, the eigenvalues of the hermitian 10×10 matrix $\mathbf{\Gamma}^0$ take on integral values $-1 \leq \gamma \leq +1$, and eigenvalues of angular momentum satisfy $0 \leq J \leq \Gamma$ and $-J \leq \gamma \leq J$, providing associative relationships between vector and scalar components. A direct examination of (1.1) for this representation expresses a degeneracy in the equation for $\gamma = 0$ massive components, along with massless fields. For this reason, a focus will be placed upon the potential of these fields towards enhancing standard modeling of electro-weak bosons. Plausible kinematic coincidences relating various electro-weak parameters using presently known phenomenology will be explored, in hopes of motivating the development of a compelling model describing general transformations of physical states. Some exemplary coincidences that depend upon un-discovered dark particle states and have been presented elsewhere[4] will be expanded and critiqued in light of more recent experimental measurements. To end the discussion, plausible models that depends *only* on known electro-weak parameters will be developed as further exemplars of the potential usefulness of this approach.

2 A Brief Extended Lorentz Group Primer

To begin, a brief summary of the general properties of the extended Lorentz group will be developed. A more detailed exposition can be found in [3] or [5].

2.1 Group algebra

The group algebra inclusive of operators Γ^μ requires the non-vanishing commutators to satisfy

$$\begin{aligned} [\Gamma^0, \Gamma^k] &= \frac{i}{\hbar} K_k, & [\Gamma^0, J_k] &= 0, & [\Gamma^0, K_k] &= -i\hbar \Gamma^k, \\ [\Gamma^j, \Gamma^k] &= -\frac{i}{\hbar} \epsilon_{jkm} J_m, & [\Gamma^j, J_k] &= i\hbar \epsilon_{jkm} \Gamma^m, & [\Gamma^j, K_k] &= -i\hbar \delta_{jk} \Gamma^0. \end{aligned} \quad (2.1)$$

Operators J_k and Γ^0 are hermitian observables, and K_k and Γ^k are anti-hermitian. For general algebras satisfying $[\hat{G}_r, \hat{G}_s] = -i \sum_m (c_s)_r^m \hat{G}_m$, the structure constants can be used to develop a group metric $\eta_{ab} \equiv \sum_{s,r} (c_a)_r^s (c_b)_s^r$, $\eta^{ab} \equiv ((\eta)^{-1})_{ab}$ that can construct the Casimir operator $\hat{C}_G \equiv \sum_{r,s} \eta^{G_r G_s} \hat{G}_r \hat{G}_s$ whose eigenvalues label the irreducible representations. The Casimir operator for this extended Lorentz group is $C_\Gamma = \frac{1}{6} [(\underline{J} \cdot \underline{J} - \underline{K} \cdot \underline{K})/\hbar^2 + \Gamma^0 \Gamma^0 - \underline{\Gamma} \cdot \underline{\Gamma}]$, and finite dimensional representations have dimension $N_\Gamma = \frac{1}{3}(\Gamma+1)(2\Gamma+1)(2\Gamma+3) = 10$ for the $\Gamma = 1$ representation being discussed. Furthermore, a metric for operators carrying the space time indexes under Lorentz sub-group transformations takes the form $\eta^{\Gamma^\mu \Gamma^\nu} = -\frac{1}{6} \eta_{\mu\nu}$ in terms of the Minkowski metric. *General* spinor forms that satisfy (2.1) are developed in reference [5].

2.2 Kinematics of spinors

Dimensionless spinors $\mathbf{u}_\gamma^{(\Gamma)}$ generally satisfy the equation

$$\Gamma^\mu p_\mu \mathbf{u}_\gamma^{(\Gamma)}(\underline{\beta}, s, s_z) = -\gamma m_{(st)} c \mathbf{u}_\gamma^{(\Gamma)}(\underline{\beta}, s, s_z), \quad (2.2)$$

where $\underline{\beta} = \underline{p}/\epsilon$ defines the relative Lorentz frame of the state for massive particles. The standard state is typically identified with $\underline{\beta} = 0$. For massless particles, the standard state is traditionally specified in terms of z -moving helicity states with contra-variant 4-momentum components $\vec{p}_{(st)} = (1, 0, 0, 1)$. More generally, standard state massless 4-momentum can be associate with $\vec{p}_{(st)} = \mu_{(st)} c(1, 0, 0, 1)$, where $\mu_{(st)}$ must be a definable kinematic scale uniquely associated with the creation of that state.

The definition of the Dirac adjoint spinor $\bar{\mathbf{u}} \equiv \mathbf{u}^\dagger \mathbf{g}_\Gamma$ to be identified with Dirac products between spinors requires the development of a N_Γ dimensional matrix that satisfies $\mathbf{G}_s^\dagger = \mathbf{g}_\Gamma \mathbf{G}_s \mathbf{g}_\Gamma$ for all finite dimensional forms of the generators of the extended group. This matrix turns out to be purely diagonal with elements $\mathbf{g}_\Gamma = (-1)^{\Gamma-\gamma}$, and only corresponds to $\mathbf{g}_{\frac{1}{2}} = \gamma^0$ for Dirac fermions.

3 Causal $\Gamma = 1$ Boson States

For massive particles, the momentum-space forms of the $\Gamma = 1$ spinors depend on the Lorentz boost $\underline{\beta}$ from the standard states, and the $m \rightarrow 0$ form of z -moving spinors are pure numbers, just as was the case for the $\Gamma = \frac{1}{2}$ representation spinors. *Unlike* the $\Gamma = \frac{1}{2}$ fields, the form (1.1) includes a degenerate set of $\gamma = 0$ distinct states that can be mixed while continuing to satisfy the field equation. Unitary mixing of these degenerate states establishes relationships between the kinematic spinors in terms of invariant mixing angles and mass ratios.

Of particular interest, standard model electro-weak bosons include electrically charged massive $W^\pm$ bosons and a massive neutral scalar H boson, as well as the

massive electrically neutral vector Z boson and photon A that result from the unitary mixing of the neutral W^3 and B bosons[6, 7]. In a prior publication[4], kinematic conditions resulting from the unitary mixing of the degenerate $\Gamma = 1$ bosonic causal spinors were demonstrated to potentially model electro-weak bosons consistent with experimental results. The matrix representations of the generators are 10-dimensional, inherently relating scalar and vector components. In what follows, plausible relationships and kinematic coincidences involving electro-weak and electromagnetically dark bosons will be further critiqued and explored. Henceforth, natural units with $\hbar \rightarrow 1, c \rightarrow 1$ will be utilized, and Einstein's summation convention over repeated *super/subscripts* will be assumed.

3.1 General form and properties of bosonic spinors

The $\Gamma = 1$ momentum-space spinor fields Φ of the fields described in (1.1) satisfy (2.2) with $\Gamma = 1$. The spin values s are integers satisfying $0 \leq s \leq 1$, and the eigenvalues of Γ^0 satisfying $-s \leq \gamma \leq +s$ represent an additional set of discrete integer valued quantum numbers. The Γ^β are 10×10 matrices[8] that will be ordered with the first component being the single scalar component with $\gamma = 0$, the next three components representing the s_z eigenstates with $\gamma = +1$, the following three components representing the s_z eigenstates with $\gamma = 0$, and the final three components representing the s_z eigenstates with $\gamma = -1$. The representation is irreducible due to the mixing of components in the anti-hermitian matrices Γ^j and $\mathbf{K}_j$.

All Dirac orthonormalized z-moving spinors $\mathbf{u}(\beta_z, s, s_z)$ satisfying $\mathbf{u}_a^\dagger \mathbf{g}_{\Gamma=1} \mathbf{u}_b \equiv \bar{\mathbf{u}}_a \mathbf{u}_b = (-1)^{1-\gamma_b} \delta_{ab}$ have components that become singular when $m \rightarrow 0$, where $\mathbf{g}_{\Gamma=1}$ is a diagonal matrix with elements $(-1)^{1-\gamma}$ for all spin states. Other than the subset of $\gamma = 0$ spinors, the *hermitian* normalized spinors are not orthogonal. However, only the hermitian orthonormalized $\gamma = 0$ spinors are non-singular for massless systems. Massive standard state (i.e. at rest) spinors are always eigenstates of Γ^0 satisfying $\Gamma^0 \mathbf{u}_a = \gamma_a \mathbf{u}_a|_{\beta \rightarrow 0}$, where $a = 1$ for the $\gamma = 0$ scalar spinor, for the $\gamma = +1$ vectors $a \in \{2, 3, 4\}$, for the degenerate $\gamma = 0$ vectors $a \in \{5, 6, 7\}$, and for the $\gamma = -1$ (anti-particle) vectors $a \in \{8, 9, 10\}$.

The massless spinors are *never* eigenstates of Γ^0 . However, they *do* satisfy $p_\mu \Gamma^\mu \mathbf{u}_a|_{m=0} = 0$. Generally, for massless spinors the forms $\mathcal{A}_{\gamma, s_z}^{(s)\beta}(\vec{p}) \equiv \Gamma^\beta \Phi_{\gamma, s_z}^{(1,s)}(\vec{p})$ satisfying (2.2) are momentum space spinor forms transverse to the 4-momentum $p_\beta \mathcal{A}_{\gamma, s_z}^{(s)\beta}(\vec{p}) = 0$. The configuration space form of these fields thus satisfy the Lorentz gauge.

Only the components $\mathcal{A}_{0,0}^{(s=0)\beta}(\vec{p})$ generate *new* spinor forms that are also eigenstates of $\mathbf{S}_z \frac{\mathcal{A}_{0,0}^{(s=0)x} \pm i \mathcal{A}_{0,0}^{(s=0)y}}{\sqrt{2}} = \pm 1 \frac{\mathcal{A}_{0,0}^{(s=0)x} \pm i \mathcal{A}_{0,0}^{(s=0)y}}{\sqrt{2}}$, $\mathbf{S}^2 \frac{\mathcal{A}_{0,0}^{(s=0)x} \pm i \mathcal{A}_{0,0}^{(s=0)y}}{\sqrt{2}} = 2 \frac{\mathcal{A}_{0,0}^{(s=0)x} \pm i \mathcal{A}_{0,0}^{(s=0)y}}{\sqrt{2}}$, and can also orthogonally mix the original spinors $\Phi_{\gamma, s_z}^{(1,s)}(\vec{p})$. The forms of $\mathcal{A}_{0,0}^{(s=1)\beta}(\vec{p})$ are *not* spinor eigenstates of angular momentum $\mathbf{S}_z, \mathbf{S}^2$. Thus, the configuration

space form of the $\mathcal{A}_{0,0}^{(0)\beta}$ fields can represent covariant gauge potentials that directly generate the field strengths $F_{\mu\nu} \equiv \partial_\mu \mathcal{A}_{0,0\nu}^{(0)} - \partial_\nu \mathcal{A}_{0,0\mu}^{(0)}$ that express common representations of spatial polarizations.

3.2 Unitary kinematic mixing of degenerate components

The mixing of previously degenerate eigenstates to develop dynamically alternative representations is a phenomenon that sometimes occurs in perturbative atomic physics (e.g. Stark effect). In this subsection, relationships for unitary kinematic mixing of degenerate ($\Gamma = 1, \gamma = 0$) states will be briefly developed. The focus will be on mixing degenerate spinors moving parallel to the z-axis expressed in terms of a kinematic parameter $\zeta(\beta_z)$ which is a function only of the Lorentz transformation from the standard state β_z . The *hermitian* normalized scalar spinor has the form

$$\mathbf{U}_X(\zeta_X, s=0, s_z=0) = \begin{pmatrix} \cos \zeta_X \\ 0 \\ -\frac{\sin \zeta_X}{\sqrt{2}} \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ \frac{\sin \zeta_X}{\sqrt{2}} \\ 0 \end{pmatrix} \xrightarrow{m_X \rightarrow 0} \begin{pmatrix} \frac{1}{\sqrt{2}} \\ 0 \\ -\frac{1}{2} \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ \frac{1}{2} \\ 0 \end{pmatrix}, \quad (3.1)$$

where the dimensionless *kinematic angle* for particle X is defined by

$$\begin{aligned} \zeta_X &\equiv \sin^{-1} \left(\frac{p_X}{\sqrt{m_X^2 + 2p_X^2}} \right) = \sin^{-1} \sqrt{\frac{\epsilon_X^2 - m_X^2}{2\epsilon_X^2 - m_X^2}} \\ &\rightarrow p_X = m_X \frac{\sin \zeta_X}{\sqrt{\cos 2\zeta_X}}, \quad \epsilon_X = m_X \frac{\cos \zeta_X}{\sqrt{\cos 2\zeta_X}}, \end{aligned} \quad (3.2)$$

with the Lorentz transformation from the standard state parameterized by $\beta_z = \tan \zeta_X$. It is noteworthy that systems with identical $\frac{p}{\epsilon}$ are co-moving, and thus have identical spinor forms. Massive spinors have kinematic angles in the range $-\frac{\pi}{4} < \zeta_X < \frac{\pi}{4}$, and massless spinors have $\zeta_X = \pm \frac{\pi}{4}$. The connection between $\gamma = 0$ Dirac orthonormal spinors $\mathbf{u}_X$ discussed previously and hermitian normalized spinors $\mathbf{U}_X$ is given by $\mathbf{u}_X = \frac{\mathbf{U}_X}{\sqrt{\cos 2\zeta_X}}$, which clearly demonstrates a breakdown of Dirac-type normalization for *massless* bosons.

Consider the general mixing of orthogonal states $X + V \Rightarrow R + Y$. Unitary mixing of the spinor components requires

$$\begin{aligned} \cos \theta_{XV} \mathbf{U}_X(\zeta_X, s, s_z) + \sin \theta_{XV} \mathbf{U}_V(\zeta_V, s, s_z) &= \mathbf{U}_R(\zeta_R, s, s_z), \\ -\sin \theta_{XV} \mathbf{U}_X(\zeta'_X, s, s_z) + \cos \theta_{XV} \mathbf{U}_V(\zeta'_V, s, s_z) &= \mathbf{U}_Y(\zeta_Y, s, s_z), \end{aligned} \quad (3.3)$$

where R and Y types interchange for $\theta_{XV} \rightarrow -\theta_{XV}$. For instance, if $\zeta_X > 0$ the spinor $\mathbf{U}_R(\zeta_R, s, s_z)$ has the form $\mathbf{U}_X(\zeta_X - \theta_{XV}, s, s_z)$, and the spinor $\mathbf{U}_Y(\zeta_Y, s, s_z)$ has the form $\mathbf{U}_V(\zeta_X - \theta_{XV}, s, s_z)$. Orthogonality requires that if X is mass/energy-like, then V must be tachyon/momentum-like, guaranteed by kinematic angles satisfying $\zeta_V = \zeta_X - \frac{\pi}{2}$.

Expressing the tachyonic form for system V via $m_V \rightarrow i\mu_V$, orthogonal mixing relates the kinematic angles as follows:

$$p_V \rightarrow \mp \frac{\mu_V}{m_X} \sqrt{m_X^2 + p_X^2} \quad , \quad \sqrt{m_V^2 + p_V^2} \rightarrow \pm \frac{\mu_V}{m_X} p_X \quad , \quad \zeta_V = \zeta_X \mp \frac{\pi}{2}. \quad (3.4)$$

The resulting spinors through trigonometric re-combination have kinematic angle $\zeta_R = \zeta_X \mp \theta_{XV}$. Furthermore, the spinor $\mathbf{u}_R$ will have the same kinematic form as $\mathbf{u}_X$, and $\mathbf{u}_Y$ will share the orthogonal form of $\mathbf{u}_V$. For massless resultant spinors, $\mathbf{u}_R(\zeta = \pm \frac{\pi}{4}) = \mp \mathbf{u}_Y(\zeta = \mp \frac{\pi}{4})$. In most of what follows, only the upper signs will be utilized unless the sign is explicitly demonstrated. The resultant energy-momentum of the $\mathbf{R}$ or $\mathbf{Y}$ spinor is required to satisfy $\vec{P}_X + \vec{P}_V = \vec{P}_{R,Y}$ as appropriate for the resultant mass. Given this expression, one can derive the form of the kinematic angle associated with mixing a massive X with an orthogonal V resulting in a product with invariant energy M , given by

$$\zeta X_{\mu_V}^M = \pm \sin^{-1} \left(\frac{\sqrt{\sqrt{M^4 + 2M^2(\mu_V^2 - m_X^2)} + (\mu_V^2 + m_X^2)^2} - 2m_X\mu_V}{\sqrt{2}(M^4 + 2M^2(\mu_V^2 - m_X^2) + (\mu_V^2 + m_X^2)^2)^{\frac{1}{4}}} \right), \quad (3.5)$$

where the sign follows that of the momentum p_X . This form vanishes when $M^2 = m_X^2 - \mu_V^2$. For ‘bootstrap’ self mixing, the kinematic angle takes the value $\zeta X_{\mu_X}^{m_X} = \sin^{-1} \sqrt{\frac{1}{2} - \frac{1}{\sqrt{5}}}$. Generally, the form $\vec{p} \cdot \vec{x} = m \frac{z \sin \zeta - ct |\cos \zeta|}{\sqrt{\cos(2\zeta)}}$ that appears in the 4-momentum conserving exponentials of quantum fields differ *only* in the masses of co-moving systems. It should be emphasized that all massive co-moving states will share the same kinematic angle.

Consistency of (3.5) with mixing to a *massless* state $M_Y = 0 \rightarrow \zeta_Y = \pm \frac{\pi}{4}$ requires that $\theta_{XV} = \tan^{-1} \frac{\mu_V}{m_X}$ and $|\zeta X_{\mu_V}^0| = \frac{\pi}{4} - \theta_{XV} = |\zeta X_{\mu_V}^{M_{Y_{xv}}}|$, where an additional invariant energy $M_{Y_{xv}}$ satisfies this relationship if $m_X > \mu_V$. The energy of this resultant state and an invariant representation label $M_{Y_{xv}}$ satisfy

$$\epsilon_{Y_{xv}} = \frac{m_X^2 + \mu_V^2}{2\sqrt{m_X\mu_V}} \quad \text{and} \quad M_{Y_{xv}} = \sqrt{2}\sqrt{m_X^2 - \mu_V^2}. \quad (3.6)$$

Note that generally, $\tan^{-1} \frac{\mu_V}{m_X} + \tan^{-1} \frac{m_X}{\mu_V} = \frac{\pi}{2}$, which relates the mixing angles resulting from interchange of X and V . The kinematic angle of the spinor $\mathbf{R}$ resulting from this mixing thus satisfies $\zeta_R = \zeta X_{\mu_V}^M - \theta_{XV} = \zeta X_{\mu_V}^M - \tan^{-1} \frac{\mu_V}{m_X}$. Furthermore, orthogonal *self*-mixing to $M_{XX} = 0$ states with $\zeta X_{\mu_X}^0 = 0$ (i.e. X at rest) generates massless states of momenta $\pm m_X$.

3.3 Modeling of electro-weak bosons

3.3.1 Standard electro-weak modeling

In the standard model of electro-weak interactions, the Lagrangian density

$$\mathcal{L}_{EW} = \left((\partial_\mu - ig_W W_\mu^{(a)} \hat{\tau}_a - ig_B B_\mu \hat{Y}) \phi \right)^\dagger \eta^{\mu\nu} (\partial_\nu - ig_W W_\nu^{(b)} \hat{\tau}_b - ig_B B_\nu \hat{Y}) \phi - V(\phi^\dagger \phi) \quad (3.7)$$

manifests local gauge invariance under $SU[2] \times U[1]$ transformations, where the $W_\nu^{(b)}$ potentials are associated with the $SU[2]$ generators $\hat{\tau}_b$, and the B_ν potentials are associated with a $U[1]$ hypercharge generator $\hat{Y}$. For a potential that takes the form $V(\phi^\dagger \phi) \cong m_\phi^2 \phi^\dagger \phi + \lambda(\phi^\dagger \phi)^2$, there is a minimum when $\langle \phi \rangle = 0$, and *this* ground state maintains the symmetry of the Lagrangian..

However, if the mass term becomes ‘tachyonic’ $m_\phi \rightarrow \pm i\mu_\phi$, then the potential is minimum for *any* particular value of the field that satisfies $\langle \phi^\dagger \phi \rangle \cong \frac{\mu_\phi^2}{2\lambda}$, so that the system settles into a particular ground state that breaks the symmetry of the Lagrangian. Choosing the $\hat{\phi}^0$ component to manifest the broken symmetry, longitudinal excitations about this new minimum $\hat{\phi}^0 \equiv \langle \phi \rangle + \hat{h}$ then define a massive Higgs scalar field $\hat{h}$, with real mass $m_H = \sqrt{2}\mu_\phi$ and vanishing ground state expectation value. Furthermore, transverse excitations along the symmetry of the Lagrangian result in massless (Goldstone) modes.

Substitution of the form of $\hat{\phi}^0$ into (3.7) yields quadratic terms generating a mass for the gauge bosons $W_\mu^{(\pm)}$ of $m_{W^{(\pm)}} = \left| \frac{g_W \langle \phi \rangle}{\sqrt{2}} \right|$, which is independent of the units of charge utilized. Furthermore, the remaining gauge bosons $W_\mu^{(3)}$ and B_μ are mixed to generate a massive charge-neutral Z boson and a massless photon that appropriately couples to electric charges of the $W_\mu^{(\pm)}$ bosons using

$$Z_\mu = W_\mu^{(3)} \cos \vartheta_w + B_\mu \sin \vartheta_w, \quad A_\mu = -W_\mu^{(3)} \sin \vartheta_w + B_\mu \cos \vartheta_w. \quad (3.8)$$

The charge coupling to the $W_\mu^{(\pm)}$ that results upon substitution of (3.8) into (3.7), and the quadratic term in the field Z_μ , allow the following identifications:

$$e = g_W \sin \vartheta_w = g_B \cos \vartheta_w, \quad \cos \vartheta_w = \frac{m_W}{m_Z}. \quad (3.9)$$

3.3.2 Unitary mixings involving $H + \phi$ spinors

An algebra of transformations whose boson representation generates spinors with both scalar and vector components presents an intriguing characteristic of possible relevance to the standard electro-weak bosons. To begin an exploration towards identifying electro-weak bosons as causal spinor fields, examine the orthogonal mixing of the degenerate components of a spinor inclusive of the scalar H with the tachyonic μ_ϕ that induces the electro-weak symmetry breaking using (3.5).

The mixing of m_H with $\mu_\phi = \frac{m_H}{\sqrt{2}}$ exhibits several unique properties. First, the unitary mixing angle satisfies $\theta_{H\phi} = \tan^{-1} \frac{m_\phi}{m_H} = \tan^{-1} \frac{1}{\sqrt{2}}$, defining a massless state with energy $\epsilon_{Yh\varphi} = \frac{3m_H}{2^{7/4}}$ generated by kinematic angle $|\zeta H_{m_\phi}^{M=0}| = \frac{\pi}{4} - \theta_{H\phi}$. Furthermore *this* combination, uniquely defines kinematic angles $|\zeta H_{\mu_\phi}^{m_H}| = |\zeta H_{\mu_\phi}^{M=0}| = \sin^{-1} \sqrt{\frac{1}{6}(3 - 2\sqrt{2})}$, which implies opposite momenta $p_H(\zeta H_{\mu_\phi}^{m_H}) = -p_H(\zeta H_{\mu_\phi}^0)$. This allows the self-generating mix $H + \phi \rightarrow \tilde{H}$ to an R -like $\tilde{H}$ with a mass $m_{\tilde{H}} = m_H$ of momentum p_H , along with $H' + \phi \rightarrow \tilde{G}$ to a *massless* R -like $\tilde{G}$ at the same invariant $M = m_H$ of momentum $-p_H$, where $\tilde{G}$ can be associated with a massless mode generated by an H' with opposite kinematic angle $-\zeta H_{\mu_\phi}^{m_H}$. In addition, the mix $H + \phi \rightarrow Y$ at $M = 0$ generates a massless $\mathbf{Y}$ spinor with the same energy as the $\tilde{G}$, but orthogonal to it. Similarly, $H + \phi \rightarrow Y$ at $M = m_H$ generates a massive Y -like spinor with the same kinematics as the $\tilde{G}$. This is apparently a type of bootstrap consistency unique amongst particles whose mass is generated by this type of symmetry breaking. The energy-momentum of the massless modes satisfy $\vec{P} = (\frac{3m_H}{2^{7/4}}, 0, 0, \mp \frac{3m_H}{2^{7/4}})$ from (3.6).

It is perhaps noteworthy that $H + \phi$ mixing to result in an R -like ϕ occurs with the H at rest $\zeta H_{\mu_\phi}^{m_\phi} = 0$. Analogous relationships are satisfied for mixing of the mass scale $g_W \langle \phi \rangle$ with a tachyonic $\mu_W = \left| \frac{g_W \langle \phi \rangle}{\sqrt{2}} \right|$, as well as for the mixing of $g_B \langle \phi \rangle$ with the tachyonic mass scale $\mu_B = \left| \frac{g_B \langle \phi \rangle}{\sqrt{2}} \right| = m_W \tan \theta_W$, to be discussed next. Finally, the orthogonal mixing of two $\mathbf{H}$ spinors generates *both* orthogonal massless modes with the kinematic angle $\zeta H_{\mu_H}^{M=0} = 0$ at rest and resultant massless modes of 4-momentum $\vec{p} = (m_H, 0, 0, \pm m_H)$, since $\theta_{HH} = \frac{\pi}{4}$.

3.3.3 Mixing of W+B spinors

For $W + B$ mixing, the unique case of invariant mass $M = \frac{m_W}{\cos \theta_{WB}} \equiv m_Z$ *analytically* generates an opposite helicity state for which

$$\zeta W_{\frac{m_W}{\cos \theta_{WB}} \tan \theta_{WB}} \equiv \zeta W_{\mu_B}^{m_Z} = \frac{\theta_{WB}}{2} = -(\zeta W_{\mu_B}^{m_Z} - \theta_{WB}) = -\zeta_{Zwb}, \quad (3.10)$$

i.e. the standard state frame of the $\mathbf{W}_0$ spinor is equal and opposite that of the resultant $\mathbf{Z}$ spinor. This means that the spinor structure of any co-moving *massive* states with equal and opposite momenta can be donated to the $\mathbf{W}_0$ and $\mathbf{Z}$ spinors while preserving Lorentz invariance, even if those states are initially massless. It also defines the mass scale of the B as $\mu_B = m_W \tan \theta_{WB} = \sqrt{m_Z^2 - m_W^2}$.

3.3.4 Analytic coincidences involving electro-weak mass scales

Several analytic coincidences beyond those already discussed can be ascertained. For instance, the ratios of the mass scales $m_H + \mu_\phi$, $g_W \langle \phi \rangle + \mu_W$, and $g_B \langle \phi \rangle + \mu_B$ are

identical, implying relationships defined by the fundamental symmetry breaking of the form

$$|\zeta\langle g\phi\rangle_{\mu_W}^{\langle g\phi\rangle}| = |\zeta\langle \tilde{g}\phi\rangle_{\mu_B}^{\langle \tilde{g}\phi\rangle}| = |\zeta H_{\mu_\phi}^{m_H}| = \frac{\pi}{4} - \theta_{H\phi} = |\zeta H_{\mu_\phi}^0| = |\zeta\langle \tilde{g}\phi\rangle_{\mu_B}^0| = |\zeta\langle g\phi\rangle_{\mu_W}^0|, \quad (3.11)$$

where for brevity $\langle g\phi\rangle \equiv g_W\langle\phi\rangle$ and $\langle \tilde{g}\phi\rangle \equiv g_B\langle\phi\rangle$. Also, due to relationships between energy scales of kinematic angles like (3.5), several mixes from rest generate relevant states, e.g.

$$\zeta H_{\mu_\phi}^{m_\phi} = 0, \quad \zeta\langle g\phi\rangle_{\mu_W}^{m_W} = 0, \quad \text{and} \quad \zeta\langle g\phi\rangle_{\langle \tilde{g}\phi\rangle}^{M_{Awb}} = 0. \quad (3.12)$$

Examples of co-moving states mixing to massless resultants include

$$\zeta W_{\mu_H}^0 = \zeta H_{\mu_W}^0 \quad \text{and} \quad \zeta\langle g\phi\rangle_{\langle \tilde{g}\phi\rangle}^0 = \zeta W_{\mu_B}^0. \quad (3.13)$$

Furthermore, the exchange $W \leftrightarrow B$ relates the kinematic angles via

$$\zeta W_{\mu_B}^{m_Z} + \zeta B_{\mu_W}^{m_Z} = \frac{\pi}{4} \quad \text{and} \quad \zeta B_{\mu_W}^0 + \theta_{WB} = \frac{\pi}{4}. \quad (3.14)$$

Finally, as $M_{Yh\phi} = m_H$ was the case for $H + \phi$ mixing, $M_{Y\langle g\phi\rangle w} = \langle g\phi\rangle$ is identically true for $\langle g\phi\rangle + W$ mixing.

3.3.5 Kinematic coincidences involving electro-weak mass scales

Kinematic coincidences associated with unitary mixing of degenerate boson representations of causal states will next be explored as a means to motivate the development of a compelling model incorporating the known properties of electro-weak bosons. To this end, various kinematic invariants and energies related to $W, B, Z, H, \phi, \langle g_W\phi\rangle, \langle g_B\phi\rangle$, and the various mixing angles will be explored.

Beyond masses, additional scales were included for examination. For instance, the invariant energy that satisfies $|\zeta W_{\mu_B}^{M_{Awb}}| + \theta_{WB} = \frac{\pi}{4}$ perhaps generating a *massless* state labeled by $M_{Awb} \simeq 96\text{GeV}$ in $W + B$ mixing, the energy of the photon $\epsilon_{Awb} \simeq 70.7\text{GeV}$ using (3.6), as well as the analogous energies (e.g. $\epsilon_{Yh\phi} \simeq 111.7\text{GeV}$) for other mixings, were examined. Also of interest was the observation that the kinematic angle describing the self-mixing bootstrap $\zeta m_m^m = \sin^{-1} \sqrt{\frac{1}{2} - \frac{1}{\sqrt{5}}}$ is the product of $W + B$ mixing at the bootstrap invariant energy $M_{bs} \simeq 245.8\text{GeV}$ satisfying $\zeta W_{\mu_B}^{M_{bs}} - \theta_{WB} \equiv \zeta m_m^m$. Furthermore, a unique invariant energy $M_* \simeq 246.4\text{GeV}$ relating a previously examined kinematic symmetry involving W and Z threshold energies in the H rest frame[4] given by $\zeta_Z(\epsilon = \frac{m_H^2}{m_W}) = \zeta_W(\epsilon = \frac{m_H^2}{m_Z}) \equiv \zeta W_{\mu_B}^{M_*}$, was included.

Previously published predictions of the masses and ratios will here be updated using the 2022 Particle Data Group (PDG) tables[9], as well as compared to the more recent 2022 re-analysis of the Collider Detector at Fermilab (CDF) W boson

mass data[10]. The reported mass values here utilized are as follows: $m_Z \simeq 91.1876(21)$, $m_H \simeq 125.25(17)$, $m_W^{PDG} \simeq 80.377(12)$, and $m_W^{CDF} \simeq 80.4335(94)$. For future comparisons, the difference between the reported PDG and CDF values of the W mass is $\Delta m_W \equiv m_W^{CDF} - m_W^{PDG} \simeq 4.71 \sigma_W^{PDF} \simeq 6.01 \sigma_W^{CDF}$.

The phenomenological coincidences associated directly with various kinematic angles of mixing follow below in no particular order. In addition to attempting to find meaningful coincidences with regards to model building, an additional purpose of such searches is a determination of just how difficult it is to find coincidences consistent with present observations:

$$KC1: \zeta H_{\mu\phi}^{m_H} - \theta_{H\phi} \simeq -\zeta W_{\mu H}^{m_H} \rightarrow \begin{aligned} m_W &\simeq (m_W - 0.077\sigma_W)_{PDG} \text{ or} \\ m_H &\simeq (m_H + 0.008\sigma_H)_{PDG}, \end{aligned} \quad (3.15)$$

$$\frac{m_H}{m_W} \simeq \frac{7+\sqrt{113}}{8\sqrt{2}},$$

i.e., $H + \phi$ mixing at m_H generates anti-moving kinematics with $W + H$ mixing at m_H ;

$$KC2: \zeta W_{\mu H}^{\langle g_B \phi \rangle} \simeq \frac{\theta_{HW}}{2} = \frac{\pi}{4} - \frac{\theta_{WH}}{2} \rightarrow \begin{aligned} m_W &\simeq (m_W + 1.62\sigma_W)_{PDG} \text{ or} \\ m_H &\simeq (m_H - 0.367\sigma_H)_{PDG}, \end{aligned} \quad (3.16)$$

with an interpretation yet to be ascertained;

$$KC3: \begin{aligned} \zeta_H(p_W(\zeta W_{\mu H}^{m_Z})) &\simeq \zeta W_{\mu B}^{m_Z} \rightarrow m_W \simeq (m_W + 0.801\sigma_W)_{PDG} \text{ or} \\ \text{or } p_W(\zeta W_{\mu H}^{m_Z}) &\simeq p_H(\frac{\theta_{WB}}{2}) \rightarrow m_H \simeq (m_H - 1.05\sigma_H)_{PDG}, \\ \frac{m_W}{m_Z} &\simeq \frac{2m_H^3}{2m_H^3 - 2m_H m_W^2 + m_W \sqrt{m_H^4 + (m_Z^2 - m_W^2)^2 + 2m_H^2(m_Z^2 + m_W^2)}} \end{aligned} \quad (3.17)$$

i.e., the kinematic angle of an **H** with the momentum of the **W** which mixes with an orthogonal **H** to generate a **Z** is co-moving with the **W** in $W + B$ mixing to generate a **Z**, perhaps suggestive of the B as an off μ -shell version of an H ;

$$KC4: M_{bs} \simeq M_* \rightarrow \begin{aligned} m_W &\simeq (m_W - 1.1\sigma_W)_{PDG} \text{ or} \\ m_H &\simeq (m_H - 0.751\sigma_H)_{PDG}, \end{aligned} \quad (3.18)$$

i.e., the mass of the bootstrap generated by $W + B$ mixing is same as the mass generated by the $W + B$ energy invariant symmetric under $m_W \leftrightarrow m_Z$ exchange;

$$KC5: \zeta Z_{\mu\phi}^{M_*} \simeq \zeta_W(\epsilon_W = m_H) \rightarrow \begin{aligned} m_W &\simeq (m_W + 0.219\sigma_W)_{CDF} \text{ or} \\ m_H &\simeq (m_H - 0.031\sigma_H)_{CDF}, \end{aligned} \quad (3.19)$$

i.e., the invariant energy M_* for $W + B$ mixing from (3.18) is the same as that for $Z + \phi$ mixing (the only single coincidence presented that is most consistent with the CDF results);

$$KC6: \zeta H_{m_W}^{M_{Awb}} \simeq 0 \rightarrow \begin{aligned} m_W &\simeq (m_W + 1.62\sigma_W)_{PDG} \text{ or} \\ m_H &\simeq (m_H - 0.367\sigma_H)_{PDG}, \end{aligned} \quad (3.20)$$

i.e., the H is at rest in $H + W$ mixing at the invariant energy M_{Awb} for (possible) mixing of $W + B$ to a same helicity massless state;

$$\begin{aligned}
& \text{KC7: } \zeta H_{\mu_B}^{m_H} - \theta_{HB} \simeq -\frac{\theta_{WB}}{2} \rightarrow \begin{array}{l} m_W \simeq (m_W - 4.86\sigma_W)_{PDG} \text{ or} \\ m_H \simeq (m_H + 0.37\sigma_H)_{\text{using } m_W^{PDG}} \text{ or} \\ m_H \simeq (m_H + 0.72\sigma_H)_{\text{using } m_W^{CDF}}, \end{array} \\
& \frac{m_H}{m_Z} \simeq \frac{1}{2} \left(\cos \theta_{WB} + (\cos \theta_{WB} (1 - \sin \theta_{WB})^2)^{1/3} + (\cos \theta_{WB} (1 + \sin \theta_{WB})^2)^{1/3} \right), \quad (3.21)
\end{aligned}$$

i.e., the spinor generated by $H + B$ mixing to generate a new $\mathbf{H}$ is co-moving with the $\mathbf{Z}$ generated from $W + B$ mixing, and anti-moving with that $\mathbf{W}$;

$$\begin{aligned}
& \text{KC11: } \zeta W_{<g_B\phi>}^{<g_W\phi>} - \theta_{W<g_B\phi>} \simeq -\frac{\theta_{WB}}{2} \rightarrow \begin{array}{l} m_W \simeq (m_W \simeq +1.3\sigma_W)_{PDG} \text{ or} \\ m_W \simeq (m_W - 4.3\sigma_H)_{CDF}, \end{array} \\
& \frac{m_W}{m_Z} \simeq \sqrt{\frac{2}{7} \left(8 + 5\sqrt{2} - \sqrt{79 + 52\sqrt{2}} \right)}, \quad (3.22)
\end{aligned}$$

a relationship involving a $\mathbf{W}$ mixing with energy scales in the Lagrangian.

Any consistent combination of two of the previous kinematic coincidences results in a determination of two mass ratios. Since the Z mass is presently best measured, the closest predictions of m_W and m_H resulting from a requirement of two coincidences follow:

$$\begin{aligned}
(3.15)+(3.17) & \rightarrow m_W \simeq (m_W + 0.881\sigma_W)_{PDG} \text{ and } m_H \simeq (m_H + 0.105\sigma_H)_{PDG}, \\
(3.15)+(3.18) & \rightarrow m_W \simeq (m_W - 0.957\sigma_W)_{PDG} \text{ and } m_H \simeq (m_H - 0.097\sigma_H)_{PDG}, \\
(3.16)+(3.17) & \rightarrow m_W \simeq (m_W + 0.631\sigma_W)_{PDG} \text{ and } m_H \simeq (m_H - 0.224\sigma_H)_{PDG}, \\
(3.16)+(3.18) & \rightarrow m_W \simeq (m_W - 0.422\sigma_W)_{CDF} \text{ and } m_H \simeq (m_H - 0.462\sigma_H)_{CDF}, \\
(3.17)+(3.18) & \rightarrow m_W \simeq (m_W + 0.152\sigma_W)_{PDG} \text{ and } m_H \simeq (m_H - 0.854\sigma_H)_{PDG}, \\
(3.17)+(3.19) & \rightarrow m_W \simeq (m_W + 0.159\sigma_W)_{PDG} \text{ and } m_H \simeq (m_H - 0.846\sigma_H)_{PDG}, \\
(3.17)+(3.20) & \rightarrow m_W \simeq (m_W + 0.631\sigma_W)_{PDG} \text{ and } m_H \simeq (m_H - 0.224\sigma_H)_{PDG}, \\
(3.17)+(3.21) & \rightarrow m_W \simeq (m_W + 1.14\sigma_W)_{PDG} \text{ and } m_H \simeq (m_H + 0.452\sigma_H)_{PDG}, \\
(3.18)+(3.19) & \rightarrow m_W \simeq (m_W + 0.144\sigma_W)_{PDG} \text{ and } m_H \simeq (m_H - 0.848\sigma_H)_{PDG}.
\end{aligned}$$

The coincidence (3.21) has been purposefully included as a cautionary example of an individual coincidence several standard deviations from *presently* measured results providing better predictions when combined with another coincidence.

3.3.6 Kinematic coincidences involving photon normalization

For clarity, this section will utilize gaussian units for all charges $e_G \equiv \sqrt{\alpha} = g_W \sin \theta_{WB} = g_B \cos \theta_{WB}$. Typical normalizations for the photon field utilize either forms of the type $e_G \sqrt{\frac{\epsilon_s}{\epsilon_A}}$ (see e.g. [2, 5, 11, 12]) or $e_G \frac{\epsilon_s}{\epsilon_A}$ (see e.g. [13, 14]), where ϵ_A is the energy of the photon, and ϵ_s is its standard state energy, customarily assigned to unity ($\epsilon_s \rightarrow 1$). The reported value of the fine structure constant here

utilized is $\frac{1}{\alpha_{PDG}} \simeq 137.03599907$. Two coincidences presented only constrain m_H :

$$\alpha H1 : \frac{1}{\sqrt{2}}(g_W \sin \theta_{W\phi}) \sqrt{\epsilon_{Yh\phi}} \simeq 1 \rightarrow m_H \simeq (m_H - 0.79\sigma_H)_{PDG}, \quad (3.23)$$

which defines the energy of the massless resultant of $H + \phi$ mixing as the standard state energy of electromagnetic quanta, in terms of the coupling defined for $W + \phi$ mixing, and

$$\alpha H2 : \frac{\sqrt{4\pi}e_G \cos \theta_{H\phi}}{\sin \theta_{Z\phi}} \sqrt{\epsilon_s} \simeq \frac{1}{2(2\pi)^{3/2}} \sqrt{m_H} \rightarrow m_H \simeq (m_H - 0.841\sigma_H)_{PDG}, \quad (3.24)$$

which suggests that a relationship between couplings involving $W + B$, $H + \phi$, and $Z + \phi$ mixing using the customary value of the photon standard state energy *inherits* a normalization choice for a massive H . This also relates the photon normalization to that of a massless mode generated *from* that H .

Three additional coincidences will constrain either m_W or m_H :

$$\alpha W1 : \frac{1}{\sqrt{\pi}} g_W \sqrt{M_{Awb}} \simeq 1 \rightarrow m_W \simeq (m_W + 0.356\sigma_W)_{PDG}, \quad (3.25)$$

$$\alpha W2 : \frac{1}{2\sqrt{4\pi}} (g_W \sin \theta_{H\phi}) M_{Awb} \simeq 1 \rightarrow m_W \simeq (m_W + 0.407\sigma_W)_{PDG}, \quad (3.26)$$

which both assign a standard state energy of the invariant M_{Awb} for electromagnetic quanta, and

$$\alpha W3 : \sqrt{\frac{\pi}{2}} (g_W \cos \theta_{W\phi}) \sqrt{\mu_B} \simeq 1 \rightarrow \begin{array}{l} m_W \simeq (m_W + 0.617\sigma_W)_{PDG} \text{ or} \\ m_H \simeq (m_H - 0.283\sigma_H)_{PDG}, \end{array} \quad (3.27)$$

which assigns a standard state energy of μ_B for electromagnetic quanta resulting from $W + \phi$ mixing.

The following coincidence determines only m_W :

$$\alpha W13 : \frac{4\pi g_B \sin \theta_{H\phi}}{\sqrt{2}} \simeq \sqrt{\frac{\langle g_B \phi \rangle}{\epsilon_{Awb}}} \rightarrow m_W \simeq (m_W + 0.55\sigma_W)_{PDG}, \quad (3.28)$$

suggestive of mixing of $H + \phi$ to generate a massless mode with energy ϵ_{Awb} . Two additional coincidences involve generation of a massless $H + \phi$ mode with charge quantization $\frac{e_G}{3}$:

$$\alpha W41 : \frac{\sqrt{4\pi} g_B \sin \theta_{W\phi}}{3\sqrt{2}} \sqrt{\frac{m_H}{\epsilon_{Yh\phi}}} \simeq \frac{1}{(2\pi)^{2/3}} \rightarrow \begin{array}{l} m_W \simeq (m_W - 0.04\sigma_W)_{PDG} \text{ or} \\ m_H \simeq (m_H + 0.01\sigma_H)_{PDG}, \end{array} \quad (3.29)$$

suggestive of a state labeled by m_H off mass shell towards m_W generating a massless mode, and

$$\alpha A45 : \frac{4\pi g_W \sin \theta_{H\phi}}{3} \simeq \sqrt{\frac{1}{2}} \sqrt{\frac{\mu_B}{\epsilon_{Yh\phi}}} \rightarrow \begin{array}{l} m_W \simeq (m_W - 0.004\sigma_W)_{CDF} \text{ or} \\ m_H \simeq (m_H - 0.004\sigma_H), \end{array} \quad (3.30)$$

suggestive of a state labeled by B generating a massless $H + \phi$ mode. The plausible physical relevance of these various coincidences are not considered to be equivalent.

Exemplars pairing a kinematic coincidence with a photon normalization include

$$\begin{aligned} (3.15)+(3.25) &\rightarrow m_W \simeq (m_W + 0.356\sigma_W)_{PDG} \text{ and } m_H \simeq (m_H + 0.048\sigma_H)_{PDG}; \\ (3.17)+(3.23) &\rightarrow m_W \simeq (m_W + 0.159\sigma_W)_{PDG} \text{ and } m_H \simeq (m_H - 0.845\sigma_H)_{PDG}; \\ (3.17)+(3.25) &\rightarrow m_W \simeq (m_W + 0.356\sigma_W)_{PDG} \text{ and } m_H \simeq (m_H - 0.587\sigma_H)_{PDG}. \end{aligned}$$

Alternatively, coincidences involving the reported renormalized value for the fine structure constant $\frac{1}{\alpha_R(m_Z)} \simeq 127.951[9, 15]$ can be explored. The form of the running fine structure constant at this fixed point is taken to be

$$\alpha_Z(\mu) = \frac{\alpha_R(m_Z)}{1 - \frac{2\alpha_R(m_Z)}{3\pi} g(\mu) \log \frac{\mu^2}{m_Z^2}}, \quad (3.31)$$

where $g(\mu)$ sums over the squares of charges (in units of e_G) below charged pair productions at momentum transfer μ . A few example coincidences follow:

$$\alpha_Z 2 : \frac{1}{\sqrt{2}} \frac{\sqrt{\alpha_Z(m_W)}}{3 \sin \theta_{WB}} \simeq \sqrt{4\pi} \frac{\epsilon_s}{m_W} \rightarrow m_W \simeq (m_W - 0.57\sigma_W)_{PDG}; \quad (3.32)$$

$$\alpha_Z r 2 : \frac{4\pi \sqrt{\alpha_Z(< g_B \phi >)}}{\sin \theta_{WB}} \sin \theta_{H\phi} \sqrt{\frac{< g_B \phi >}{\epsilon_{Yh\phi}}} \simeq 1 \rightarrow \begin{array}{l} m_W \simeq (m_W + 0.19\sigma_W)_{PDG} \\ \text{or } m_H \simeq (m_H - 0.07\sigma_H); \end{array} \quad (3.33)$$

$$\alpha_Z r 3 : \frac{1}{2(2\pi)^{3/2}} \frac{\sqrt{4\pi \alpha_Z(\epsilon_{Yhb})}}{\cos \theta_{WB}} \sin \theta_{W\phi} \simeq \frac{\epsilon_s}{\epsilon_{Yhb}} \rightarrow \begin{array}{l} m_W \simeq (m_W + 1.31\sigma_W)_{PDG} \\ \text{or } m_H \simeq (m_H - 0.04\sigma_H); \end{array} \quad (3.34)$$

$$\alpha_Z r 25 : \frac{\sqrt{4\pi \alpha_Z(M_{Awb})}}{3 \sin \theta_{WB}} \cos \theta_{H\phi} \frac{\epsilon_{Yhb}}{M_{Awb}} \simeq \frac{1}{(2\pi)^{3/2}} \rightarrow \begin{array}{l} m_W \simeq (m_W + 0.09\sigma_W)_{CDF} \\ \text{or } m_H \simeq (m_H + 0.01\sigma_H). \end{array} \quad (3.35)$$

Next, selected predictions involving possible combinations of kinematic coincidences (3.15)-(3.21) with photon normalization coincidences (3.23)-(3.27) will be presented (recall $\frac{1}{\alpha_{PDG}} \simeq 137.03599907$):

- (3.17)+(3.19)+(3.23) $\rightarrow m_Z \simeq (m_Z + 0.095\sigma_Z)_{PDG}$, $m_W \simeq (m_W + 0.173\sigma_W)_{PDG}$ and $m_H \simeq (m_H - 0.844\sigma_H)_{PDG}$, or alternatively $\frac{1}{\alpha} \simeq 137.0357$, $m_W \simeq (m_W + 0.159\sigma_W)_{PDG}$ and $m_H \simeq (m_H - 0.846\sigma_H)_{PDG}$;

- (3.17)+(3.19)+(3.24) $\rightarrow m_Z \simeq (m_Z + 0.634\sigma_Z)_{PDG}$, $m_W \simeq (m_W + 0.257\sigma_W)_{PDG}$ and $m_H \simeq (m_H - 0.835\sigma_H)_{PDG}$, or alternatively $\frac{1}{\alpha} \simeq 137.038$, $m_W \simeq (m_W + 0.159\sigma_W)_{PDG}$ and $m_H \simeq (m_H - 0.846\sigma_H)_{PDG}$;
- (3.21)+(3.22)+(3.29) $\rightarrow \frac{1}{\alpha} \simeq 137.0329$, $m_W \simeq (m_W + 1.33\sigma_W)_{PDG}$ and $m_H \simeq (m_H + 0.47\sigma_H)$.

A few observations are noteworthy. First, the originally reported values of the kinematic angles $\zeta W \simeq \frac{\theta_{WB}}{2} \simeq -\zeta Z$ for $W + B \rightarrow Z$ mixing were just *extremely* close numerical coincidences[4]. Subsequently, the analytic result (3.10) was found to *require* this to be the case, providing substantial motivational support for this approach. Furthermore, there are some cautionary observations that should be noted. Previous coincidences involving (3.17) and (3.19) using 2018 Particle Data Group (PDG) masses were very close, while they are only somewhat close using the more recent 2022 PDG data. Finally, one might be tempted to utilize two compatible photon normalization coincidences with two massive kinematic coincidences to attempt to calculate *all* mass and coupling values. For instance, if the four equations (3.17)+(3.19)+(3.23)+(3.24) were all physically relevant, one could *calculate* all electroweak parameters as satisfying $\frac{1}{\alpha} \simeq 137.0368$ (vs. 137.03599907), $m_Z \simeq (m_Z + 0.365\sigma_Z)_{PDG}$, $m_W \simeq (m_W + 0.215\sigma_W)_{PDG}$ and $m_H \simeq (m_H - 0.84\sigma_H)_{PDG}$. However, such a set of mathematically consistent relationships would make a disturbing inference about the ‘arbitrariness’ of the choice of the standard state photon energy ϵ_s in the photon normalization terms, somehow mathematically fixing *choice* of units of measurement.

From the representative set of kinematic coincidences presented, it seems clear that PDG values provide far more kinematic coincidences than CDF values. It should be noted that there are other kinematic coincidences with a broad range of *subjective* plausibility that have been examined, but excluded for brevity. Some are consistent with the overall predictions described in this section, while others are incompatible.

3.4 Modeling using only anti-moving frames

As has been mentioned, degenerate neutral $\Gamma = 1$ causal bosons $W + B$ that can mix to form a massless state also *necessarily* generates a massive $m_Z = \frac{m_W}{\cos \theta_{WB}}$ that moves in the opposing Lorentz frame $\underline{\beta}_Z = -\underline{\beta}_W$. Since the primary purpose of this paper is to demonstrate a plausible pathway towards enhancing predictability and understanding of standard model phenomenology, a final effort will be made to demonstrably model electro-weak phenomenology consistent with kinematic coincidences involving kinematic angles $\zeta = \pm \frac{\theta_{WB}}{2}$ resulting from degenerate boson mixing. The focus will remain only on the kinematics of *known* particles, exclusive of any undiscovered potential massive dark particles.

The phenomenological coincidences involving anti-moving frames that have been presented include KC3 (3.17), KC7 (3.21), and KC11 (3.22). However, since KC11 involves energy scales in the Lagrangian (not mass scales of particle states), the focus will be on particle state mixing to generate new particles. For convenience, the coincidences are repeated below, and plausible kinematic diagrams for unitary mixing are exhibited in Figure 1:

$$KC3 : p_W(\zeta W_{\mu_H}^{m_Z}) \simeq p_H(\zeta W_{\mu_B}^{m_Z}), \quad KC7 : \zeta H_{\mu_B}^{m_H} - \theta_{HB} \simeq -\frac{\theta_{WB}}{2}. \quad (3.36)$$

The modeling should produce all known particle states in a kinematically consis-




Figure 1: Unitary diagrammatic representations of KC3 (left) and KC7 (right) generated by anti-moving kinematics.

tent manner. For KC3, an **H** and anti-**H** colliding via exchange of an **H** scatter into precisely the needed reference frame $\zeta_H = \frac{\theta_{WB}}{2}$ (leaving the scalar **H** component unaffected), then go off-shell as intermediate **W**s with the same momenta. The neutral component of an intermediate **W** has the momentum needed to mix with an orthogonal vector **H** to generate a **Z**, while the charged components **W**[±] which are not degenerate remain unaffected. Thus the diagram generates all observed states. For KC7, an **H** and anti-**H** colliding via exchange of a vector **B** scatter into precisely the needed reference frames $\zeta_H = \pm \frac{\theta_{WB}}{2}$ (leaving the scalar **H** component unaffected), then go off-shell as an intermediate **W** with the correct kinematic parameter needed to mix with a **B** to generate a **Z**. The charged components **W**[±] which are not degenerate remain unaffected, likewise generating all observed states. This diagram is suggestive of the intermediate **B**s as off mass-shell tachyonic **H**s that can unitarily mix to generate **Z** and **A** bosons.

4 Discussion and Conclusions

The model described in this paper (examining bosons) and its companion part I paper (which examined fundamental fermions) has been demonstrated to mirror

causal spinor fields that exhibit properties consistent with observed leptons and bosons. Just as is the case for Dirac fermions, the $\Gamma = 1$ boson representation spinors have the kinematically dependent forms that allow the equation of motion to be linear in the energy-momentum *operators*, simplifying cluster decomposability. However, unlike the fermion representation, there are a set of degenerate (massive and massless) spinor fields corresponding to a vanishing eigenvalue for the equation of motion, inclusive of both vector and scalar components. This allows the degenerate set of spinors to be unitarily mixed to generate physical states that continue to satisfy the defining equation of motion, suggestive of a meaningful application of the formulation to describe electro-weak bosons.

Unitary mixing of orthogonal spinors consistent with one resultant state possibly being massless defines analytic forms relating the kinematic angles of the involved spinors in terms of the invariant mass of that resultant. In particular, the unique relationship of an $\mathbf{H}$ spinor with an orthogonal Φ spinor with (tachyonic) mass $\mu_\phi = \frac{m_H}{\sqrt{2}}$ to generate a resultant spinor of mass m_H , might *also* generate an orthogonal massless (Goldstone-like) mode utilizing the same invariant (extended Poincare) group label m_H . Furthermore, mixing $W + B$ at invariant energy $m_Z = \frac{m_W}{\cos \theta_{WB}}$ analytically *requires* that the kinematic frame of the $\mathbf{W}$ must be equal but opposite that of the resultant $\mathbf{Z}$, with a value precisely half the unitary mixing angle $\frac{\theta_{WB}}{2}$. These (and other) analytic relationships compelled the author to search for meaningful kinematic coincidences in a manner analogous to the examination of planetary motions by Kepler and others in hopes of discovering a set of rules that can lead to the development of acceptable supplementations to standard model canon.

Several representative (but by no means comprehensive) analytic relationships and kinematic coincidences that fall within a (subjectively) chosen range of plausibility and consistency with measured electro-weak parameters have been presented for demonstrative purposes. The selection includes exemplars that illustrate how the formulation might provide insight into how the Higgs sector provides kinematic structure to the *massive* W and Z spinor fields, as well as substantive support for massless modes. Furthermore, it should be noted that the formulation has the potential to provide insight into states that are electromagnetically dark, yet have physical and cosmological significance.

To conclude, an attempt has been made to develop exemplary model candidates consistent with what was initially an extremely close experimental kinematic coincidence involving $W + B \rightarrow Z$ mixing that has now been demonstrated to be an analytic necessity. The exemplars predict kinematic parameters consistent with present data. The approach recognizes that the development of a *compelling* model ultimately embraces the most elegant combination of analytic relationships with plausible kinematic coincidences that remain consistent with future measurements, as well as predictive of yet to be discovered phenomena.

References

- [1] P.A.M. Dirac, Proc. Roy. Soc. (London), A117, 610 (1928); *ibid*, A118, 351 (1928).
- [2] S. Weinberg, *The Quantum Theory of Fields*, Cambridge University Press (1995), ISBN 0-521-55001-7.
- [3] J. Lindesay, “A Plausible Path Towards Unification of Interactions via Gauge Fields Consistent with the Equivalence Principle-I”, arXiv:2403.07930v1 (2024), 15 pages.
- [4] J. Lindesay, “Unitary kinematic mixing of electro-weak bosons, *Int.J. Theor. Phys.*, (2022), doi:10.1007/s10773-022-05050-9
- [5] J. Lindesay, *Foundations of Quantum Gravity*, Cambridge University Press (2013), ISBN 978-1-107-00840-3, see ch. 4 and 9.
- [6] S. Weinberg, *Phys. Rev. Lett.* **19**, 1264 (1967).
- [7] M. Peskin, *Concepts of Elementary Particle Physics*, Oxford University Press, Oxford (2019), ISBN 978-0-19-881219-7.
- [8] See Appendix D2 in [5]
- [9] R.L. Workman et.al. (Particle Data Group), *Prog. Theor. Exp. Phys.* 2022, 083C01(2022)
- [10] CDF Collaboration, “High-precision measurement of the W boson mass with the CDF II detector”, *Science*, Vol 376, Issue 6589, pp. 170-176 (2022). DOI: 10.1126/science.abk1781
- [11] Michio Kaku, *Quantum Field Theory, A Modern Introduction*, Oxford University Press, New York (1993), ISBN 0-19-507652-4. See eg. Equations (3.99) and (3.100).
- [12] J.D. Bjorken and S.D. Drell, *Relativistic Quantum Mechanics*, McGraw-Hill, New York (1964).
- [13] E. Fradkin, *Quantum Field Theory*, Princeton University Press (2021), ISBN 9780691 I 49080 (print).
- [14] M. Srednicki, *Quantum Field Theory*, Cambridge University Press, Cambridge (2007).

- [15] M.D. Schwartz, *Quantum Field Theory and the Standard Model*, Cambridge University Press (2014) ISBN 978-1-107-03473-0.