




Continuous Gravitational Waves: A New Window to Look for Heavy Non-annihilating Dark Matter

Sulagna Bhattacharya ^{1, a}, Andrew L. Miller ^{2, 3, b} and Anupam Ray ^{4, 5, c}

¹*Tata Institute of Fundamental Research, Homi Bhabha Road, Mumbai 400005, India*

²*Nikhef – National Institute for Subatomic Physics,
Science Park 105, 1098 XG Amsterdam, The Netherlands*

³*Institute for Gravitational and Subatomic Physics (GRASP),
Utrecht University, Princetonplein 1, 3584 CC Utrecht, The Netherlands*

⁴*Department of Physics, University of California Berkeley, Berkeley, California 94720, USA*

⁵*School of Physics and Astronomy, University of Minnesota, Minneapolis, MN 55455, USA*

(Dated: March 22, 2024)

Sun-like stars can transmute into comparable mass black holes by steadily accumulating heavy non-annihilating dark matter particles over the course of their lives. If such stars form in binary systems, they could give rise to quasi-monochromatic, persistent gravitational waves, commonly known as continuous gravitational waves, as they inspiral towards one another. We demonstrate that next-generation space-based detectors, e.g., Laser Interferometer Space Antenna (LISA) and Big Bang Observer (BBO), can provide novel constraints on dark matter parameters (dark matter mass and its interaction cross-section with the nucleons) by probing gravitational waves from transmuted Sun-like stars that are in close binaries. Our projected constraints depend on several astrophysical uncertainties, nevertheless, are competitive with the existing constraints obtained from cosmological measurements as well as terrestrial direct searches, demonstrating a notable science-case for these space-based gravitational wave detectors as probes of particle dark matter.

I. INTRODUCTION

LIGO, Virgo and KAGRA [1–3] have observed $\mathcal{O}(100)$ binary black hole (BH) and neutron star mergers [4–7] in their various observing runs over the past few years. However, another class of gravitational waves (GWs), commonly known as, “continuous gravitational waves”, have so far eluded detection. Continuous waves are characterized as quasi-monochromatic (a sinusoid with a tiny frequency drift over time) as well as quasi-infinite (a signal whose duration greatly exceeds the observation time), and can be emitted from an array of sources, e.g., asymmetrically rotating isolated neutron stars [8–12], millisecond pulsars at the galactic center [13, 14], annihilating ultralight dark matter (DM) clouds around rotating BHs [15–18], inspiraling planetary-mass primordial black holes [19–25], and even ultralight particle DM that directly couples to the interferometers [26–34].

In this work, we point out the exciting possibility that continuous gravitational wave measurements can also probe strongly-interacting heavy non-annihilating DM. Such a DM model is hard to probe in the terrestrial detectors because of their tiny fluxes (DM flux in the terrestrial detector scales inversely with its mass) and remains to be scrutinized thoroughly (see, e.g., Ref. [35] for a bird’s-eye view of existing exclusions on such DM model). Here, we demonstrate that next-generation space based GW detectors, such as Laser Interferometer Space Antenna

(LISA) and Big Bang Observer (BBO) [36, 37], are excellent testing grounds for strongly-interacting heavy non-annihilating DM. Our proposal can be summarized as follows: accumulation of strongly-interacting heavy non-annihilating DM inside binary stellar objects (symmetric Sun-like binaries, to be more specific) can transmute them into low mass (comparable with the progenitor masses) black hole binaries. These low mass BH binaries, commonly known as transmuted black hole (TBH) binaries, if sufficiently close, can emit quasi-monochromatic continuous GWs in their inspiral phase, and can be observed in the next generation GW detectors, such as LISA and BBO. We theoretically estimate the occurrence rate-density of such TBH binaries whose progenitors are closely-spaced symmetric Sun-like binaries, and search these binaries in the future space-based GW detectors. Assuming a null detection of these TBH binaries with a year of observation time, we first calculate the upper limits on the occurrence rate-density, and by translating these upper limits on the occurrence rate-density, we provide novel (projected) constraints on heavy non-annihilating dark matter interactions. The constraints derived in this work are subject to several astrophysical uncertainties but are complementary with the existing constraints from terrestrial direct searches and cosmological measurements.

The rest of the work is organized as follows. In Sec. II, we briefly review the formation of low mass transmuted black holes. In Sec. III, we theoretically estimate the occurrence rate density of such transmuted black holes whose progenitors are closely-spaced symmetric Sun-like binaries. In Sec. IV, we estimate the upper limit on the occurrence rate density by assuming a null detection of these low mass TBH binaries in future space-based GW

^a sulagna@theory.tifr.res.in

^b andrew.miller@nikhef.nl

^c anupam.ray@berkeley.edu

detectors, such as LISA & BBO. In Sec. V, we use the upper limits on the occurrence rate density to derive the exclusion limits on the dark matter parameters (mass and its interaction cross-section with the nucleons). Finally, we culminate the paper with conclusions in Sec. VI.

II. FORMATION OF LOW MASS TRANSMUTED BLACK HOLES

In this section, we briefly review the transmutation process of binary stellar objects (for ease, we refer to binary stellar objects as stellar objects hereafter). Non-annihilating DM particles from the Galactic halo that transit through a stellar object can be captured due to their collisions with stellar nuclei [38–40]. For sufficiently large DM-nucleon scattering cross-sections, the capture process becomes very effective. This is simply because, in this regime, DM particles typically scatter many times while transiting through the stellar object, leading to a larger energy loss probability and almost all of the transiting DM particles get captured. For heavy DM, i.e., if the DM mass m_χ is much heavier than the nuclei mass m_A ($m_\chi \gg m_A$), which is of primary interest here, captured DM particles sink towards the stellar core and settle into a small core-region. This leads to a huge number density of these captured DM particles within the stellar core, which eventually collapses, followed by a small BH formation in the stellar core. This nascent BH, if not sufficiently light, can quickly devour the progenitor, transforming it into comparable mass BHs, which we refer as *low-mass transmuted BHs*. In the following, we systematically go over the various stages of DM-induced transmutation of a stellar object (more specifically, Sun-like stars which are the progenitors in our analysis).

Capture: To begin with, we first define the maximal capture rate as geometric capture rate (C_{geo}), and it represents the total number of DM particles that can pass through a stellar object. For a specific velocity distribution of the incoming DM particles $f(u)$, the maximal capture rate is [40]

$$C_{\text{geo}} = \frac{\rho_\chi}{m_\chi} \pi R^2 \int \frac{f(u) du}{u} (u^2 + v_{\text{esc}}^2), \quad (1)$$

where v_{esc} is the escape velocity of the planetary systems, R denotes the size of the stellar object, and $\rho_\chi = 0.4 \text{ GeV/cm}^3$ is the Galactic DM density in the Solar neighborhood. Specifically, for a Maxwell-Boltzmann velocity distribution, C_{geo} becomes

$$C_{\text{geo}} = \frac{\rho_\chi}{m_\chi} \pi R^2 \sqrt{\frac{8}{3\pi}} \bar{v} \left(1 + \frac{3v_{\text{esc}}^2}{2\bar{v}^2} \right), \quad (2)$$

where $\bar{v} = 270 \text{ km/s}$ denotes the average velocity of the DM particles in the Galactic halo. For Sun-like stars, the maximal capture rate is

$$C_{\text{geo}}|_{\text{Sun-like}} = 1.3 \times 10^{24} \text{ s}^{-1} \left(\frac{10^6 \text{ GeV}}{m_\chi} \right). \quad (3)$$




FIG. 1. The capture fraction ($f_{\text{cap}} = C/C_{\text{geo}}$) is shown with the DM-nucleon scattering cross-section ($\sigma_{\chi n}$) for DM masses of 10^5 GeV and 10^7 GeV . f_{cap} becomes very small for low $\sigma_{\chi n}$, and approaches unity for sufficiently large $\sigma_{\chi n}$. For heavier DM, relatively large $\sigma_{\chi n}$ is needed in order to reach $f_{\text{cap}} = 1$ as heavier DM are harder to stop due to their larger kinetic energy.

Depending on the DM-nucleon scattering cross-section and DM mass, a certain fraction (f_{cap}) of the DM particles that transit get trapped, implying a capture rate $C = f_{\text{cap}} C_{\text{geo}}$ [41–45]. For heavy DM ($m_\chi \gg m_A$) and large DM-nucleon scattering cross-sections, which is of primary interest here, f_{cap} can even reach unity [41]. Of course, for low DM-nucleon scattering cross-sections, capture becomes inefficient, and f_{cap} becomes extremely small (as shown in Fig. 1). We closely follow Ref. [46] to compute the capture fraction for our analysis, however, we have verified that using f_{cap} from Ref. [47] yields a slightly weaker result. We also note that enhancement of DM capture rate due to close binary is negligible in this analysis [48].

Spatial distribution inside the stellar volume: After accumulation, DM particles thermalize with the stellar nuclei via successive collisions. The timescale for thermalization depends strongly on the DM-nucleon scattering cross-section: for large $\sigma_{\chi n}$, it occurs quickly, i.e., over a time much shorter than the stellar object lifetime [49–55]. The thermalized DM particles then become spatially distributed in a way that depends crucially on their mass [56, 57]. For example, heavy DM tends to shrink towards the stellar-core and stabilize into a tiny radius around the core, commonly known as the thermalization radius. Quantitatively, for Sun-like stars, DM particles of mass 10^6 GeV thermalize within a radius of $r_{\text{th}} \sim 91 \text{ km}$ ($1.3 \times 10^{-4} R_\odot$), which reduces further as $m_\chi^{-1/2}$ for heavier DM.

Dark collapse and Black Hole formation: The number of captured DM particles grows linearly with time. As a consequence, stellar objects with cosmologically long lifetimes ($\sim \text{Gyr}$) can accumulate an enormous number of DM particles. Quantitatively, for a DM mass of 10^6 GeV , and sufficiently high DM-nuclei scattering cross-section (say 10^{-28} cm^2), $\mathcal{O}(10^{41})$ DM particles accumulate inside a Sun-like star. When they concentrate within a $\sim 91 \text{ km}$ radius, they give rise to a core-density of $\sim 6 \times 10^{25} \text{ GeV cm}^{-3}$, around 26 orders of magnitude higher than the local Galactic DM density. This huge core-density undergoes gravitational collapse once it exceeds the critical threshold value, and eventually leads to a micro-BH formation within the stellar core. The necessary and sufficient criterion for such a BH formation has been extensively studied in the literature [49–51, 53, 54, 58–77], and is essentially determined by the self-gravitating criterion and quantum degeneracy pressure. More specifically, BH formation occurs if

$$N_{\chi}^{\text{BH}}|_{t_{\text{age}}} = C \times t_{\text{age}} \geq \max [N_{\chi}^{\text{self}}, N_{\chi}^{\text{Cha}}], \quad (4)$$

where $N_{\chi}^{\text{BH}}|_{t_{\text{age}}}$ denotes the total number of accumulated DM particles in the stellar object throughout its lifetime (t_{age}), N_{χ}^{self} represents the self-gravitating criterion, and N_{χ}^{Cha} represents the quantum degeneracy pressure criterion. Below, we briefly describe these two criteria.

Inside the thermalization volume, if the DM density overcomes the baryonic density, then the DM particles self-gravitate inside the core. This self-gravitating criterion (N_{χ}^{self}) is same for bosonic/fermionic DM, and it only depends on the core-density ρ_{core} as well as core-temperature T_{core} of the stellar objects [49] and the dependencies can be found as,

$$N_{\chi}^{\text{self}} \sim 2.8 \times 10^{41} \left(\frac{\rho_{\text{core}}}{154 \text{ g/cm}^3} \right) \left(\frac{T_{\text{core}}}{1.54 \times 10^7 \text{ K}} \right)^{3/2} \times \left(\frac{10^6 \text{ GeV}}{m_{\chi}} \right)^{5/2}. \quad (5)$$

The quantum degeneracy pressure criterion (N_{χ}^{Cha}) represents the maximum number of DM particles beyond which BH formation occurs. It solely depends on the spin of the DM particles, and for bosonic (fermionic) DM, it stems from the Heisenberg Uncertainty principle (Pauli exclusion principle) [49]. Quantitatively,

$$N_{\chi}^{\text{Cha}}|_{\text{boson}} = 9.5 \times 10^{25} \left(\frac{10^6 \text{ GeV}}{m_{\chi}} \right)^2, \quad (6)$$

$$N_{\chi}^{\text{Cha}}|_{\text{fermion}} = 1.8 \times 10^{39} \left(\frac{10^6 \text{ GeV}}{m_{\chi}} \right)^3. \quad (7)$$

From these numerical estimates, it is evident that, in our parameter range of interest, the transmutation criterion for Sun-like stars (which are the progenitors in our analysis) is determined by the self-gravitating criterion. This

simply leads to the fact that the results derived from our analysis are applicable for both bosonic as well as fermionic DM. The initial mass of the micro-BH formed at the stellar core M_{BH} can also be estimated as

$$M_{\text{BH}} = m_{\chi} \times \max [N_{\chi}^{\text{self}}, N_{\chi}^{\text{Cha}}] = m_{\chi} \times N_{\chi}^{\text{self}} \quad (8) \\ \sim 2.5 \times 10^{-10} M_{\odot} \left(\frac{10^6 \text{ GeV}}{m_{\chi}} \right)^{3/2},$$

where we take the Solar core-density as 154 g/cm^3 and Solar core-temperature as $1.54 \times 10^7 \text{ K}$ [54].

Growth and Evaporation of newly-formed Black Holes: The newly produced micro-BH at the center of the stellar core can accrete from the surrounding material and swallow the progenitor on a very short timescale (as compared to the stellar object's lifetime). It will also evaporate by emitting particles via Hawking radiation. For the time-evolution of the nascent BH, we can conservatively¹ consider the baryonic matter accretion by the micro-BH [49, 50] (first term) and the BH evaporation (second term)

$$\frac{dM_{\text{BH}}}{dt} = \frac{4\pi\rho_{\text{core}}G^2M_{\text{BH}}^2}{c_s^3} - \frac{P(M_{\text{BH}})}{G^2M_{\text{BH}}^2}, \quad (9)$$

where M_{BH} denotes the mass of the newly formed BH and $c_s = \sqrt{T_{\text{core}}/m_n}$ denotes the sound speed in the core of the stellar object, T_{core} is the core-temperature and m_n is the nucleon mass. ρ_{core} denotes the core-density of the stellar object, G is the Gravitational constant, and $P(M_{\text{BH}})$ denotes the Page factor [78, 79]. It is important to note that the Page factor properly accounts for gray-body corrections of the Hawking evaporation spectrum, as well as the number of Standard Model (SM) species emissions from an evaporating BH (BHs heavier than 10^{17} g only emit mass-less particles, such as photons and neutrinos, whereas, lighter BHs emit massive SM particles too). In the classical black-body radiation limit, the Page factor evaluates to $1/(15360\pi)$ and is commonly used in the literature. Considering the gray-body corrections and by accounting for the number of SM species emitted, the Page factor ranges from $1/(1135\pi)$ to $1/(74\pi)$ [79, 80].

Since the accretion term scales as M_{BH}^2 , and the evaporation term scales as $1/M_{\text{BH}}^2$, for low BH masses, evaporation dominates over the accretion process. Therefore, for sufficiently light micro-BHs, successful transmutation of the hosts do not occur. This sets a cut-off on the DM mass that can be probed via transmutation as the mass of the micro-BH decreases with heavier DM (Eq. 8). For Sun-like systems, this cut-off mass is around $m_{\chi} \sim 10^{10} \text{ GeV}$ [75].

Drift time and maximal possible scattering cross-section: Transmutation of stellar objects does not occur at very large DM-nucleon scattering cross-sections.

¹ This is conservative because we do not account for the accretion of newly incoming DM particles by the micro-BH.

This is simply because, at very large DM-nucleon cross-sections, DM particles lose a significant amount of energy in the outer shells of the stellar object, and therefore, may not reach the stellar core (or take a significantly longer time to reach the core). We estimate the drift time, i.e., the time required by the DM particles to reach the stellar core by using the stellar density, temperature, and compositional profiles [59, 81]

$$t_{\text{drift}} = \frac{1}{Gm_\chi} \sum_j \sigma_{\chi j} \int_0^R \frac{n_j(r) \sqrt{3A_j T(r)}}{\int_0^r d^3r' \rho_j(r')} dr, \quad (10)$$

where $\sigma_{\chi j}$ denotes the DM-nuclei scattering cross-section, and is related to the DM-nucleon scattering cross-section via $\sigma_{\chi j} = \sigma_{\chi n} A_j^2 (\mu_{\chi A_j} / \mu_{\chi n})^2$. A_j denotes the mass number of the j -th nuclei, and $\mu_{\chi n}$ is the reduced mass of the DM-nucleon system. We set the ceilings of our results by demanding that $t_{\text{drift}} \leq 1$ Gyr. Quantitatively, for Sun-like systems (for both bosonic and fermionic DM), it corresponds to [54, 75]

$$\sigma_{\chi n} \leq 10^{-18} \text{ cm}^2 \left(\frac{m_\chi}{10^6 \text{ GeV}} \right). \quad (11)$$

III. OCCURRENCE RATE-DENSITY OF LOW MASS TRANSMUTED BLACK HOLES

Accumulation of strongly-interacting heavy non-annihilating DM particles inside stellar objects can lead to the formation of comparable mass TBHs. These low-mass TBHs, while in binaries, can emit GWs as they inspiral towards each other, and could be detected in the next-generation space-based GW detectors, such as LISA. As a concrete example, binaries with component masses of $1M_\odot$ and separation of $4R_\odot$ emit GWs at a frequency of 3.5×10^{-5} Hz, potentially detectable by LISA (as shown in Fig. 2). Of course, the initial separation of these progenitors binaries can vary, which would then lead to GW emission at a different frequency. For wide binaries (binaries with larger separation), the GW frequency decreases and eventually falls outside of the LISA sensitivity band. Quantitatively, for symmetric Sun-like binaries (which are the progenitors in our analysis), if the orbital separation exceeds $9.5R_\odot$, the GW frequency becomes $< 10^{-5}$ Hz and falls outside the LISA sensitivity (as shown in Fig. 2). This implies that in our analysis we only consider GW emission from Sun-like binaries with orbital separations between $4R_\odot - 9.5R_\odot$. Note, we choose Sun-like systems as our progenitors because (1) they have much larger sizes as compared to the planets ensuring maximum DM accumulation (Eq. 1), and (2) stars form binaries much more easily than planets do. In the following, we theoretically estimate the occurrence rate-density of such TBH binaries whose progenitors are closely-spaced symmetric Sun-like binaries. First, we define the key quantity for this estimation: transmutation time (τ_{trans}) which dictates the total time required in

order to have a successful transmutation of the progenitors. τ_{trans} is essentially a sum of two timescales, where the first timescale represents the time required to form a micro-BH inside the stellar core, and the second timescale represents the time required to swallow the progenitor by the newly formed micro-BH. In our parameter range of interest, the first timescale, which depends on the DM mass as well as the DM-nucleon scattering cross-section, always exceeds the second timescale, which only depends on the DM mass. Of course, transmutation time (τ_{trans}) also depends on the ambient DM density which we take as 0.4 GeV/cm^3 as our region of interest resides in the Solar neighborhood.

We consider a closely-spaced, sun-like binary system, which is formed at time t_f . For successful transmutation, the time required for transmutation has to be shorter than the available time, implying,

$$\tau_{\text{trans}} \leq (t_0 - t_f), \quad (12)$$

where $t_0 = 13.79$ Gyr denotes the current age of the Universe. Clearly, only a fraction of these binaries that satisfy the above criterion will undergo a transmutation. Therefore, the occurrence rate density of such TBH binaries (whose progenitors are closely-spaced symmetric Sun-like binaries) can be written as

$$R_{\text{TBH}} \propto \frac{1}{t_{\text{obs}}} \int_{t_*}^{t_0} dt_f \lambda \frac{d\rho^*}{dt} [t_f] \times \Theta [t_0 - t_f - \tau_{\text{trans}}(m_\chi, \sigma_{\chi n})], \quad (13)$$

where $t_{\text{obs}} = 1$ year denotes the observation time and $t_* = 4.9 \times 10^8 \text{ year}$ ($z_* = 10$) [82] denotes the earliest binary formation time. λ denotes the fraction of stellar mass in binaries and $\frac{d\rho^*}{dt} [t_f]$ denotes the cosmic star formation rate density which we take as Madau and Dickinson star formation rate density. For the normalization of Eq.(13), we use the total number of progenitors (closely-spaced symmetric Sun-like binaries) within our volume of interest and assume a uniform spatial distribution of the progenitors in our Galaxy.

In the following, we provide a brief estimate of the normalization criterion that has been used in our analysis. Our Galaxy contains almost 200×10^9 stars, and of which about 20% are Sun-like stars, implying 40×10^9 Sun-like stars in our Galaxy [83]. Among them, a certain fraction can form a closely-spaced symmetric binary, which we denote as α . This implies the total number of progenitors (closely-spaced symmetric Sun-like binaries) in our Galaxy is $40 \times 10^9 \alpha$, and by assuming a spatial distribution of the progenitors in our Galaxy, this leads to the normalization condition for R_{TBH} is $\frac{40 \times 10^9 \alpha}{\frac{4}{3} \pi (15 \text{ kpc})^3}$ for an observation time of 1 year. Here, we note that the closely-spaced symmetric binary fraction α is rare and it has large astrophysical uncertainties (see e.g., Ref. [84] and references therein). Quantitatively, for Sun-like ($M \approx 0.5 - 1.6M_\odot$) binaries with solar metallicity, the close binary fraction (orbital separation $\lesssim 100R_\odot$) is reported to be smaller than 0.05 [84]. Therefore, our




FIG. 2. GW frequency (f_{GW}) is shown (solid red line) as a function of the orbital separation for a symmetric Sun-like binary. We probe binaries with orbital separations ranging from $4R_{\odot}$ to $9.5R_{\odot}$, which corresponds to the frequency range of $(10^{-5}$ to $3.5 \times 10^{-5})$ Hz. Above an orbital separation of $9.5R_{\odot}$, the frequency falls outside the LISA band.

choices of α , which are used in this analysis, are well-justified. As these estimates are subject to astrophysical uncertainties, we take α as a free parameter and vary it over a wide range of values ($\alpha \sim 10^{-3} - 10^{-11}$), which are consistent with the literature.

IV. CONTINUOUS-WAVE PROBES OF LOW MASS TRANSMUTED BLACK HOLES

In the previous section, we estimated the occurrence rate density of TBH binary systems that originated from closely-spaced sun-like stars. We now must calculate the expected sensitivity of space-based GW detectors towards these TBH systems that slowly inspiral towards one another, specifically the expected rate density constraints on such systems, and compare that to Eq. (13) for different choices of the fraction of stars that form in binaries α .

The slow inspiral of two objects orbiting around each other can be described as a continuous wave if they are either far enough away from each other or at low enough frequencies. When these conditions are met, the time/frequency “chirp” characteristic of detected binary inspirals can be approximated as a slow linear increase of the GW frequency over time.

Quantitatively, consider the rate of change of GW frequency over time, \dot{f}_{GW} , of an inspiraling system suffi-

ciently far from merger [85]

$$\begin{aligned} \dot{f}_{\text{GW}} &= \frac{96}{5} \pi^{8/3} \left(\frac{G\mathcal{M}}{c^3} \right)^{5/3} f_{\text{GW}}^{11/3} \\ &\simeq 10^{-23} \text{ Hz/s} \left(\frac{\mathcal{M}}{0.87M_{\odot}} \right)^{5/3} \left(\frac{f_{\text{GW}}}{3.5 \times 10^{-5} \text{ Hz}} \right)^{11/3} \end{aligned} \quad (14)$$

where $\mathcal{M} \equiv \frac{(m_1 m_2)^{3/5}}{(m_1 + m_2)^{1/5}}$ is the chirp mass, m_1 and m_2 are the component masses, and f_{GW} is the GW frequency. Its integral is

$$f_{\text{GW}}(t) = f_0 \left[1 - \frac{8}{3} \frac{\dot{f}_{\text{GW}}}{f_0} (t - t_{\text{ref}}) \right]^{-3/8}, \quad (15)$$

where f_0 is the GW frequency at a reference time t_{ref} . From Eq. (14), we can see that, for a system with $m_1 = m_2 = 1M_{\odot}$, and at low-enough frequencies ($f_0 \sim 10^{-5}$ Hz), the second term in Eq. (15) will be much less than 1, and can be binomially expanded to

$$f_{\text{GW}} = f_0 + \dot{f}_{\text{GW}}(t - t_{\text{ref}}), \quad (16)$$

which describes quasi-monochromatic continuous waves. For a particular detector, we can compute the minimum detectable signal amplitude h_0 in a matched-filtering search for a quasi-monochromatic GW with a signal-to-noise ratio $\rho = 1$ (corresponding to the best possible sensitivity that we could have) [85, 86]. Matched filtering correlates a model waveform with the data, and in the case of a purely sinusoidal signal, it is simply a fast Fourier transform of the data

$$h_{0,\text{min}} = 2 \sqrt{\frac{S_n(f_{\text{GW}})}{t_{\text{obs}}}}, \quad (17)$$

where $S_n(f)$ is the noise power spectral density as a function of frequency f , and t_{obs} is the observation time. Here, we use the sensitivity curves for LISA and BBO given in [87].

We can also write an equation for the distance reach to binaries at different frequencies [85]

$$\begin{aligned} d(f_{\text{GW}}) &= \frac{4}{h_0} \left(\frac{G\mathcal{M}}{c^2} \right)^{5/3} \left(\frac{\pi f_{\text{GW}}}{c} \right)^{2/3} \\ &= 0.27 \text{ pc} \left(\frac{3.66 \times 10^{-19}}{h_0} \right) \left(\frac{\mathcal{M}}{0.87M_{\odot}} \right)^{5/3} \\ &\quad \times \left(\frac{f_{\text{GW}}}{3.5 \times 10^{-5} \text{ Hz}} \right)^{2/3} \end{aligned} \quad (18)$$

From the distance reach, and following the procedure in [21], we can calculate the number of binaries that we would detect in a given observing run by multiplying the space-time volume $\langle VT \rangle$ (which, for nearby binaries, is the volume of a sphere times the duration of the sources) by the expected formation rate density \mathcal{R} [21]

$$N_{\text{bin}}(f_{\text{GW}}) \simeq \langle VT \rangle \mathcal{R} = \frac{4}{3} \pi [d(f_{\text{GW}})]^3 \mathcal{R} T(f_{\text{GW}}), \quad (19)$$

T is the time over which we integrate the binary systems' frequency evolutions: $T = \max(t_{\text{obs}}, \Delta T)$, and ΔT is how long the binary system spends in the frequency range $[f, f + \delta f]$, which can be obtained by inverting Eq. (15)

$$\Delta T = \frac{5}{256} \pi^{-8/3} \left(\frac{c^3}{G\mathcal{M}} \right)^{5/3} \left[f_{\text{GW}}^{-8/3} - (f_{\text{GW}} + \delta f)^{-8/3} \right]. \quad (20)$$

When ΔT exceeds t_{obs} , the number of detectable sources is dominated by those between the frequencies $(f_{\text{GW}}, f_{\text{GW}} + \delta f)$, including those that began emitting gravitational waves well before the observation run. Here, $\delta f = 1/t_{\text{obs}}$ indicates that the frequency of the GW signal does not vary by more than one frequency bin during t_{obs} . Essentially, when $\Delta T > t_{\text{obs}}$, we allow for the enhancement of the signal during the observation time due to the fact that many possible systems inspiraling forever would be overlapping and thus covering the same frequency range. Here, we also impose that ΔT cannot exceed the time at which the first stars begin to form binaries, i.e., $\Delta T \lesssim t_0/10$.

After calculating $N_{\text{bin}}(f_{\text{GW}})$, we sum the number of binaries emitting gravitational waves at each frequency

$$N_{\text{bin}}^{\text{tot}} = \sum_i N_{\text{bin}}(f_{\text{GW},i}) < 1 \quad (21)$$

where we have required $N_{\text{bin}}^{\text{tot}} < 1$ to be in the case in which we do not observe these systems in the future, in order to get the strongest constraint on \mathcal{R} . Then, we solve for \mathcal{R} , which provides an upper limit on the formation rate densities of such binary planet systems as a function of their chirp masses

$$\mathcal{R} = \frac{3}{4\pi} \left(\sum_i T(f_{\text{GW},i}) d(f_{\text{GW},i})^3 \right)^{-1}. \quad (22)$$

In order to perform the above computation, we must impose a minimum separation between the two objects, and this condition leads to the maximum GW frequency at which we could detect these systems. On the other hand, the minimum frequency arises from the detector's sensitivity band. Thus, we need to restrict the orbital radius to be

$$R_{\text{orb}} > 2(R_1 + R_2), \quad (23)$$

where R_1 and R_2 are the radii of the two planets, $R_{\text{orb}} = \left(\frac{Gm}{\pi^2 f_{\text{GW}}^2} \right)^{1/3}$ and $m = m_1 + m_2$.

While we can in theory calculate the rate densities for any given m_1 and m_2 , here we restrict ourselves to solar-mass binaries, i.e. $R_1 = R_2 = R_{\odot}$ and $m_1 = m_2 = M_{\odot}$. To perform this computation, we sum over the luminosity distance reaches as a function of frequency only at frequencies for which Eq. (23) is satisfied. In Fig. 2, we show the orbital separation of the binary as a function of the GW frequency, for an equal solar-mass binary. The

maximum frequency for which the condition in Eq. (23) is satisfied, is drawn as a horizontal line (dashed red).

Applying the procedure described above for an equal solar-mass binary, and assuming that we do not find any such systems in future space-based detectors, such as LISA & BBO, we obtain a projection for the upper limit on the occurrence density of such systems

$$\mathcal{R} \leq \begin{cases} 19 \text{ kpc}^{-3}\text{yr}^{-1}, & \text{LISA,} \\ 8.22 \times 10^{-6} \text{ kpc}^{-3}\text{yr}^{-1}, & \text{BBO.} \end{cases} \quad (24)$$

If we, instead, assume $t_{\text{obs}} = 5$ years, a possible lifetime for space-based detectors, our rate density upper limits would become $8.74 \text{ kpc}^{-3}\text{yr}^{-1}$ and $3.65 \times 10^{-6} \text{ kpc}^{-3}\text{yr}^{-1}$ for LISA and BBO, respectively, which are stronger than those in Eq. (24). Note that our estimates of the rate densities depend both on the distance reach and ΔT , as $\Delta T \gg t_{\text{obs}}$ at all frequencies considered here. From Eq. (17) and Eq. (18), we can see that the distance reach scales with $\sqrt{t_{\text{obs}}}$, while ΔT depends on t_{obs} via the frequency range through which we consider the source to sweep. The interplay between these two factors – longer observation times improve d , but also shrink ΔT , since the frequency spread is smaller – results in a nontrivial change of the rate density upper limits when changing t_{obs} .

This is our main result in Sec. IV, and with these rate density upper limits, we now compute the projected constraints on strongly-interacting heavy non-annihilating particle DM parameters in Sec. V.

V. PROJECTED CONSTRAINTS ON DARK MATTER PARAMETERS

For *conservative* constraints on particle DM parameters (DM mass and its interaction strength with nucleons), we compare the theoretical occurrence rate density of TBH binaries (whose progenitors are symmetric closely-spaced Sun-like binaries) (Eq. 13) to the corresponding upper limits obtained from continuous GW searches (Eq. 24). More specifically, our projected constraints are simply derived from $R_{\text{TBH}} \leq 19 \text{ kpc}^{-3}\text{yr}^{-1}$ (for LISA) and $R_{\text{TBH}} \leq 8.22 \times 10^{-6} \text{ kpc}^{-3}\text{yr}^{-1}$ (for BBO).

In Fig. 3, we show the projected exclusion limits, assuming non-observation of continuous gravitational waves from stellar binaries, on $\{m_{\chi}, \sigma_{\chi n}\}$ obtained from future space-based detectors like LISA (left) and BBO (right) for strongly-interacting heavy non-annihilating DM particles. In the left (right) panel, red (yellow) shaded regions are the projected exclusions from our analysis, whereas, the gray shaded regions are the existing exclusions from a variety of searches. For our analysis, we assume an observation time of 1 year, and with a larger observation time the constraints get even stronger. Note that, the constraints in Fig. 3 applies for both bosonic as well as fermionic DM. This is simply because for Sun-




FIG. 3. Projected constraints on DM mass and its interaction cross-section with the nucleons from continuous GW searches with the future space-based detectors, such as LISA (left) and BBO (right). The constraints are derived by assuming a null detection of the transmuted black binaries (whose progenitors are symmetric Sun-like binaries) with a year of observation time and apply for both bosonic/fermionic DM particles. In the (left) right panel, (red) yellow-shaded regions denote the constraints derived in this work, whereas, the gray-shaded regions denote the existing constraints (see text). Our constraints crucially depend on α , closely-spaced symmetric binary fraction, and we show our constraints for $\alpha = 10^{-3}$ ($\alpha = 10^{-9}$) and $\alpha = 10^{-5}$ ($\alpha = 10^{-11}$) for LISA (BBO), consistent with the literature (see e.g., Ref. [84] and references therein). We note that for $\alpha \leq 10^{-6}$ ($\alpha \leq 10^{-12}$), we do not obtain any exclusion on the DM parameters for LISA (BBO).

like stars, the transmutation criterion is essentially determined by N_{χ}^{self} , and this is independent of the spin of the DM particles. Our constraints also crucially depend on the closely-space symmetric binary fraction α , which has large astrophysical uncertainties [84]. Here, we show our constraints for some reasonable choices of α ($\alpha = 10^{-3}$, 10^{-5} for LISA & $\alpha = 10^{-9}$, 10^{-11} for BBO), which are quite consistent with the literature [84]. So, it is evident that even if the closely-spaced symmetric binary fraction (α) is rare, the constraints obtained from continuous GW searches are already competitive with the existing exclusion limits, demonstrating that future space-based GW detectors are ideal laboratories to probe heavy non-annihilating dark matter.

In the following we briefly describe the existing exclusion limits which are shown in gray shaded regions. Cosmological constraints, labelled as, “CMB” and “MW Satellites” denote exclusions obtained from Planck measurements of temperature and polarization anisotropy of the cosmic microwave background [88, 89] and observations of Milky Way satellite galaxies [90, 91], respectively. Large panels of etched plastic, placed aboard the Skylab Space Station, also provide significant exclusion limits on DM-nucleon interactions, labelled as “Skylab” [92–94]. Constraints labelled as “Terrestrial detectors” represents a bird’s-eye view of the existing constraints from underground, surface, and high altitude detectors, and is taken from [35, 95, 96]. Other astrophysical constraints, such as, disk stability [92], interstellar gas cooling [97], as well as Galactic Center gas-cloud heating [98, 99];

terrestrial constraints, such as MAJORANA demonstrator at the Sanford underground research facility [100], DEAP-3600 detector at SNOLAB [101], a shallow-depth experiment performed at the University of Chicago [102], and Rocket-based X-ray Quantum Calorimetry (XQC) experiment [103] do not cover any additional parameter space, and hence, are not shown for clarity. Exclusion limits obtained from the mere existence of the Sun, and other Solar-system planets cover a similar parameter space [54, 75], and are also not shown for clarity. Finally, constraints obtained from cosmic ray silicon detector satellite (IMP7/8), and balloon-borne experiment (IMAX) are shown in gray (thin) dashed line as they are not based on detailed analyses in peer-reviewed papers [94].

In this analysis, we consider Sun-like systems as our progenitors because in the strongly-interacting regime, the Sun captures a lot more DM particles as compared to other planetary bodies, and more importantly, the binary formation for stars is much more favorable than planetary bodies (for planets, α is significantly small). The exclusion limits in Fig. 3 can be understood qualitatively from the following. For lighter DM, transmutation criterion is not attainable as the total number of captured DM particles inside the stellar core ($\sim 1/m_{\chi}$) is not sufficient for transmutation ($\sim 1/m_{\chi}^{5/2}$). This sets the sharp vertical cut-offs in lighter DM mass (around $m_{\chi} \sim 10^6$ GeV). For heavier DM, transmutation is more favourable, but, the mass of micro-BH decreases with an increase in DM mass (Eq. 8). As a result, the micro-BH

takes a substantially longer time to consume the host (much longer than the age of the stellar object), and Hawking radiation becomes dominant over the accretion processes. The combination of these two effects prevents successful transmutation and provides the vertical cut-offs at the heavier DM masses (around $m_\chi \sim 10^{10}$ GeV). For sufficiently low DM-nucleon scattering cross-sections, the capture fraction (f_{cap}) decreases, and eventually the total number of captured DM particles inside the stellar core is not sufficient for BH formation. Because of this inefficient capture with low DM-nucleon scattering cross-sections, we can not probe arbitrarily low DM-nucleon scattering cross-sections, leading to the lower boundary of our exclusion regions. Very large DM-nucleon scattering cross-sections are also not probed via transmutations as the drift time of the DM particles becomes substantially longer. In Fig. 3, we show the ceilings of our results by demanding that $t_{\text{drift}} \leq 1$ Gyr, which corresponds to $\sigma_{\chi n} \leq 10^{-18} \text{ cm}^2$ for $m_\chi = 10^6$ GeV and linearly increases with heavier DM mass. We also note that for $\alpha \leq 10^{-6}$ ($\alpha \leq 10^{-12}$), we do not obtain any exclusion on the DM parameters for LISA (BBO).

VI. SUMMARY & CONCLUSIONS

In this work, we propose a novel way to probe strongly interacting heavy non-annihilating DM interactions with ordinary baryonic matter. Such DM model is hard to probe in the conventional direct detection experiments because of their tiny fluxes, and demands new techniques. We show that next-generation space-based GW detectors, such as LISA, are ideal testing room for such DM model. We propose gradual accumulation of heavy, non-annihilating DM in the ordinary stars (Sun-like) can lead to comparable mass black holes, and these low mass BHs, while in binaries, can emit quasi mono-chromatic GWs as they inspiral towards each other. For closely-spaced symmetric Sun-like binaries (which are the progenitors in our analysis) the frequency of these GWs is $\mathcal{O}(10^{-5})$ Hz, detectable by future space-based GW detectors, such as LISA. So, we search these closely-spaced Solar-mass transmuted binaries in LISA, and estimate model-independent upper limits on their rate densities by assuming a null detection of these binaries with a year of observation time. Finally, we translate these limits to put significant (projected) exclusions on heavy non-annihilating DM interactions, demonstrating the potential of space-based

GW detectors as probes of heavy non-annihilating DM. The novel exclusion limits obtained in the analysis apply to bosonic as well as fermion dark matter particles and cover a mass window from 10^6 GeV to 10^{10} GeV. The lower mass cut-off arises from the fact that the transmutation criterion is harder to achieve for light DM particles, whereas, the higher mass cut-off stems from the fact that the newly produced micro-BH is inefficient for causing a successful transmutation. The exclusion limits also depend on the closely-spaced symmetric binary fraction of the progenitors (α), which is rare and currently uncertain. However, for reasonable choices of the closely-spaced symmetric binary fraction of the progenitors, we find that the exclusion limits derived in this analysis are competitive with the existing constraints obtained from cosmological as well as direct searches. Note that, while we have focused on symmetric Sun-like stars in this analysis, our results can easily be generalized for binaries with any component masses, by correctly accounting for the closely-spaced binary fraction, which is even rarer for planetary bodies. Therefore, our work opens up a new window to probe heavy non-annihilating DM interactions by using continuous GW techniques. We must also mention that our results are sensitive to whatever the true low-frequency cut-off of the future space-based detectors will be. If future space-based detectors cannot reach such a low frequency, it would be difficult to probe Solar-mass binaries in this way, and instead, we would have to consider much heavier systems, for which we need to estimate the close binary fraction correctly. Such a task depends heavily on uncertain astrophysics and can be looked at in future work. Additionally, other future space-based detectors, such as Taiji and TianQin [104, 105], may also be useful in probing this kind of DM.

VII. ACKNOWLEDGMENTS

We sincerely thank Basudeb Dasgupta, Ranjan Laha, & Nirmal Raj for helpful discussions and important inputs on the manuscript. S.B. acknowledges ICGC-2023 conference at IIT Guwahati for its kind hospitality where part of this research was completed. A.L.M acknowledges the support for this research by the Netherlands Organisation for Scientific Research (NWO). A.R. acknowledges support from the National Science Foundation (Grant No. PHY-2020275) and to the Heising-Simons Foundation (Grant No. 2017-228).

-
- [1] LSC collaboration, J. Aasi et al., *Advanced LIGO, Classical Quantum Gravity* **32** (2015) 074001 [1411.4547].
 - [2] VIRGO collaboration, F. Acernese et al., *Advanced Virgo: a second-generation interferometric*

- gravitational wave detector, Classical Quantum Gravity* **32** (2015) 024001 [1408.3978].
- [3] KAGRA collaboration, T. Akutsu et al., *Overview of KAGRA: Detector design and construction history, PTEP* **2021** (2021) 05A101 [2005.05574].

- [4] LIGO SCIENTIFIC COLLABORATION, VIRGO, KAGRA collaboration, R. Abbott et al., *GWTC-3: Compact Binary Coalescences Observed by LIGO and Virgo During the Second Part of the Third Observing Run*, [2111.03606](#).
- [5] T. Venumadhav, B. Zackay, J. Roulet, L. Dai and M. Zaldarriaga, *New binary black hole mergers in the second observing run of Advanced LIGO and Advanced Virgo*, *Phys. Rev. D* **101** (2020) 083030 [[1904.07214](#)].
- [6] B. Zackay, L. Dai, T. Venumadhav, J. Roulet and M. Zaldarriaga, *Detecting gravitational waves with disparate detector responses: Two new binary black hole mergers*, *Phys. Rev. D* **104** (2021) 063030 [[1910.09528](#)].
- [7] S. Olsen, T. Venumadhav, J. Mushkin, J. Roulet, B. Zackay and M. Zaldarriaga, *New binary black hole mergers in the LIGO-Virgo O3a data*, *Phys. Rev. D* **106** (2022) 043009 [[2201.02252](#)].
- [8] K. Riles, *Recent searches for continuous gravitational waves*, *Mod. Phys. Lett. A* **32** (2017) 1730035 [[1712.05897](#)].
- [9] R. Tenorio, D. Keitel and A. M. Sintes, *Search Methods for Continuous Gravitational-Wave Signals from Unknown Sources in the Advanced-Detector Era*, *Universe* **7** (2021) 474 [[2111.12575](#)].
- [10] O. J. Piccinni, *Status and Perspectives of CW Searches*, *Galaxies* **10** (2022) 72 [[2202.01088](#)].
- [11] K. Riles, *Searches for continuous-wave gravitational radiation*, *Living Rev. Rel.* **26** (2023) 3 [[2206.06447](#)].
- [12] LIGO SCIENTIFIC COLLABORATION, VIRGO, KAGRA collaboration, A. L. Miller, *Recent results from continuous gravitational wave searches using data from LIGO/Virgo/KAGRA's third observing run, in 57th Rencontres de Moriond on Gravitation*, 5, 2023, [2305.15185](#).
- [13] KAGRA, LIGO SCIENTIFIC, VIRGO collaboration, R. Abbott et al., *Search for continuous gravitational wave emission from the Milky Way center in O3 LIGO-Virgo data*, *Phys. Rev. D* **106** (2022) 042003 [[2204.04523](#)].
- [14] A. L. Miller and Y. Zhao, *Probing the Pulsar Explanation of the Galactic-Center GeV Excess Using Continuous Gravitational-Wave Searches*, *Phys. Rev. Lett.* **131** (2023) 081401 [[2301.10239](#)].
- [15] S. D'Antonio et al., *Semicoherent analysis method to search for continuous gravitational waves emitted by ultralight boson clouds around spinning black holes*, *Phys. Rev. D* **98** (2018) 103017 [[1809.07202](#)].
- [16] M. Isi, L. Sun, R. Brito and A. Melatos, *Directed searches for gravitational waves from ultralight bosons*, *Phys. Rev. D* **99** (2019) 084042 [[1810.03812](#)].
- [17] L. Sun, R. Brito and M. Isi, *Search for ultralight bosons in Cygnus X-1 with Advanced LIGO*, *Phys. Rev. D* **101** (2020) 063020 [[1909.11267](#)].
- [18] LIGO SCIENTIFIC COLLABORATION, VIRGO, KAGRA collaboration, R. Abbott et al., *All-sky search for gravitational wave emission from scalar boson clouds around spinning black holes in LIGO O3 data*, *Phys. Rev. D* **105** (2022) 102001 [[2111.15507](#)].
- [19] A. Miller et al., *Method to search for long duration gravitational wave transients from isolated neutron stars using the generalized frequency-Hough transform*, *Phys. Rev. D* **98** (2018) 102004 [[1810.09784](#)].
- [20] A. L. Miller, S. Clesse, F. De Lillo, G. Bruno, A. Depasse and A. Tanasijczuk, *Probing planetary-mass primordial black holes with continuous gravitational waves*, *Phys. Dark Univ.* **32** (2021) 100836 [[2012.12983](#)].
- [21] A. L. Miller, N. Aggarwal, S. Clesse and F. De Lillo, *Constraints on planetary and asteroid-mass primordial black holes from continuous gravitational-wave searches*, *Phys. Rev. D* **105** (2022) 062008 [[2110.06188](#)].
- [22] H.-K. Guo and A. Miller, *Searching for Mini Extreme Mass Ratio Inspirals with Gravitational-Wave Detectors*, [2205.10359](#).
- [23] M. Andrés-Carcasona, O. J. Piccinni, M. Martínez and L.-M. Mir, *BSD-COBI: New search pipeline to target inspiraling light dark compact objects.*, *PoS EPS-HEP2023* (2024) 067.
- [24] G. Alestas, G. Morras, T. S. Yamamoto, J. Garcia-Bellido, S. Kuroyanagi and S. Nesseris, *Applying the Viterbi Algorithm to Planetary-Mass Black Hole Searches*, [2401.02314](#).
- [25] A. L. Miller, N. Aggarwal, S. Clesse, F. De Lillo, S. Sachdev, P. Astone et al., *Gravitational wave constraints on planetary-mass primordial black holes using LIGO O3a data*, [2402.19468](#).
- [26] A. Pierce, K. Riles and Y. Zhao, *Searching for Dark Photon Dark Matter with Gravitational Wave Detectors*, *Phys. Rev. Lett.* **121** (2018) 061102 [[1801.10161](#)].
- [27] H.-K. Guo, K. Riles, F.-W. Yang and Y. Zhao, *Searching for Dark Photon Dark Matter in LIGO O1 Data*, *Commun. Phys.* **2** (2019) 155 [[1905.04316](#)].
- [28] A. L. Miller et al., *Probing new light gauge bosons with gravitational-wave interferometers using an adapted semicoherent method*, *Phys. Rev. D* **103** (2021) 103002 [[2010.01925](#)].
- [29] Y. Michimura, T. Fujita, S. Morisaki, H. Nakatsuka and I. Obata, *Ultralight vector dark matter search with auxiliary length channels of gravitational wave detectors*, *Phys. Rev. D* **102** (2020) 102001 [[2008.02482](#)].
- [30] LIGO SCIENTIFIC COLLABORATION, VIRGO, KAGRA collaboration, R. Abbott et al., *Constraints on dark photon dark matter using data from LIGO's and Virgo's third observing run*, *Phys. Rev. D* **105** (2022) 063030 [[2105.13085](#)].
- [31] S. M. Vermeulen et al., *Direct limits for scalar field dark matter from a gravitational-wave detector*, *Nature* **600** (2021) 424.
- [32] LIGO SCIENTIFIC COLLABORATION, VIRGO, KAGRA collaboration, A. L. Miller, F. Badaracco and C. Palomba, *Distinguishing between dark-matter interactions with gravitational-wave detectors*, *Phys. Rev. D* **105** (2022) 103035 [[2204.03814](#)].
- [33] A. L. Miller and L. Mendes, *First search for ultralight dark matter with a space-based gravitational-wave antenna: LISA Pathfinder*, *Phys. Rev. D* **107** (2023) 063015 [[2301.08736](#)].
- [34] Y. Manita, H. Takeda, K. Aoki, T. Fujita and S. Mukohyama, *Exploring spin of ultralight dark matter with gravitational wave detectors*, [2310.10646](#).
- [35] D. Carney et al., *Snowmass2021 Cosmic Frontier White Paper: Ultraheavy particle dark matter*, [2203.06508](#).

- [36] P. Amaro-Seoane, H. Audley, S. Babak, J. Baker, E. Barausse, P. Bender et al., *Laser Interferometer Space Antenna*, *arXiv e-prints* (2017) [arXiv:1702.00786](#) [[1702.00786](#)].
- [37] J. Crowder and N. J. Cornish, *Beyond LISA: Exploring future gravitational wave missions*, *Phys. Rev. D* **72** (2005) 083005 [[gr-qc/0506015](#)].
- [38] W. H. Press and D. N. Spergel, *Capture by the sun of a galactic population of weakly interacting, massive particles*, *Astrophysical Journal* **296** (1985) 679.
- [39] A. Gould, *WIMP Distribution in and Evaporation From the Sun*, *Astrophys. J.* **321** (1987) 560.
- [40] A. Gould, *Resonant Enhancements in WIMP Capture by the Earth*, *Astrophys. J.* **321** (1987) 571.
- [41] J. Bramante, J. Kumar, G. Mohlabeng, N. Raj and N. Song, *Light Dark Matter Accumulating in Terrestrial Planets: Nuclear Scattering*, **2210.01812**.
- [42] D. A. Neufeld, G. R. Farrar and C. F. McKee, *Dark Matter that Interacts with Baryons: Density Distribution within the Earth and New Constraints on the Interaction Cross-section*, *Astrophys. J.* **866** (2018) 111 [[1805.08794](#)].
- [43] D. McKeen, D. E. Morrissey, M. Pospelov, H. Ramani and A. Ray, *Dark Matter Annihilation inside Large-Volume Neutrino Detectors*, *Phys. Rev. Lett.* **131** (2023) 011005 [[2303.03416](#)].
- [44] M. Pospelov and A. Ray, *Neutrinos from Earth-bound dark matter annihilation*, *JCAP* **01** (2024) 029 [[2309.10032](#)].
- [45] Y. Ema, M. Pospelov and A. Ray, *Probing Earth-Bound Dark Matter with Nuclear Reactors*, **2402.03431**.
- [46] R. Garani and S. Palomares-Ruiz, *Dark matter in the Sun: scattering off electrons vs nucleons*, *JCAP* **05** (2017) 007 [[1702.02768](#)].
- [47] R. K. Leane and J. Smirnov, *Dark matter capture in celestial objects: treatment across kinematic and interaction regimes*, *JCAP* **12** (2023) 040 [[2309.00669](#)].
- [48] L. Brayeur and P. Tinyakov, *Enhancement of dark matter capture by neutron stars in binary systems*, *Phys. Rev. Lett.* **109** (2012) 061301.
- [49] S. D. McDermott, H.-B. Yu and K. M. Zurek, *Constraints on Scalar Asymmetric Dark Matter from Black Hole Formation in Neutron Stars*, *Phys. Rev. D* **85** (2012) 023519 [[1103.5472](#)].
- [50] C. Kouvaris and P. Tinyakov, *Constraining Asymmetric Dark Matter through observations of compact stars*, *Phys. Rev. D* **83** (2011) 083512 [[1012.2039](#)].
- [51] C. Kouvaris and P. Tinyakov, *Excluding Light Asymmetric Bosonic Dark Matter*, *Phys. Rev. Lett.* **107** (2011) 091301 [[1104.0382](#)].
- [52] B. Bertoni, A. E. Nelson and S. Reddy, *Dark Matter Thermalization in Neutron Stars*, *Phys. Rev. D* **88** (2013) 123505 [[1309.1721](#)].
- [53] R. Garani, Y. Genolini and T. Hambye, *New Analysis of Neutron Star Constraints on Asymmetric Dark Matter*, *JCAP* **05** (2019) 035 [[1812.08773](#)].
- [54] J. F. Acevedo, J. Bramante, A. Goodman, J. Kopp and T. Opferkuch, *Dark Matter, Destroyer of Worlds: Neutrino, Thermal, and Existential Signatures from Black Holes in the Sun and Earth*, *JCAP* **04** (2021) 026 [[2012.09176](#)].
- [55] R. Garani, A. Gupta and N. Raj, *Observing the thermalization of dark matter in neutron stars*, *Phys. Rev. D* **103** (2021) 043019 [[2009.10728](#)].
- [56] A. Gould and G. Raffelt, *Thermal Conduction of Massive Particles*, *Astrophys. J.* **352** (1990) 654.
- [57] R. K. Leane and J. Smirnov, *Floating Dark Matter in Celestial Bodies*, **2209.09834**.
- [58] I. Goldman and S. Nussinov, *Weakly Interacting Massive Particles and Neutron Stars*, *Phys. Rev. D* **40** (1989) 3221.
- [59] A. Gould, B. T. Draine, R. W. Romani and S. Nussinov, *Neutron Stars: Graveyard of Charged Dark Matter*, *Phys. Lett. B* **238** (1990) 337.
- [60] G. Bertone and M. Fairbairn, *Compact Stars as Dark Matter Probes*, *Phys. Rev. D* **77** (2008) 043515 [[0709.1485](#)].
- [61] A. de Lavallaz and M. Fairbairn, *Neutron Stars as Dark Matter Probes*, *Phys. Rev. D* **81** (2010) 123521 [[1004.0629](#)].
- [62] N. F. Bell, A. Melatos and K. Petraki, *Realistic neutron star constraints on bosonic asymmetric dark matter*, *Phys. Rev. D* **87** (2013) 123507 [[1301.6811](#)].
- [63] T. Güver, A. E. Erkoca, M. Hall Reno and I. Sarcevic, *On the capture of dark matter by neutron stars*, *JCAP* **05** (2014) 013 [[1201.2400](#)].
- [64] J. Bramante, K. Fukushima and J. Kumar, *Constraints on bosonic dark matter from observation of old neutron stars*, *Phys. Rev. D* **87** (2013) 055012 [[1301.0036](#)].
- [65] J. Bramante, K. Fukushima, J. Kumar and E. Stopnitzky, *Bounds on self-interacting fermion dark matter from observations of old neutron stars*, *Phys. Rev. D* **89** (2014) 015010 [[1310.3509](#)].
- [66] C. Kouvaris and P. Tinyakov, *Growth of Black Holes in the interior of Rotating Neutron Stars*, *Phys. Rev. D* **90** (2014) 043512 [[1312.3764](#)].
- [67] J. Bramante and T. Linden, *Detecting Dark Matter with Imploding Pulsars in the Galactic Center*, *Phys. Rev. Lett.* **113** (2014) 191301 [[1405.1031](#)].
- [68] C. Kouvaris, P. Tinyakov and M. H. G. Tytgat, *NonPrimordial Solar Mass Black Holes*, *Phys. Rev. Lett.* **121** (2018) 221102 [[1804.06740](#)].
- [69] B. Dasgupta, A. Gupta and A. Ray, *Dark matter capture in celestial objects: light mediators, self-interactions, and complementarity with direct detection*, *JCAP* **10** (2020) 023 [[2006.10773](#)].
- [70] G.-L. Lin and Y.-H. Lin, *Analysis on the black hole formations inside old neutron stars by isospin-violating dark matter with self-interaction*, *JCAP* **08** (2020) 022 [[2004.05312](#)].
- [71] B. Dasgupta, R. Laha and A. Ray, *Low Mass Black Holes from Dark Core Collapse*, *Phys. Rev. Lett.* **126** (2021) 141105 [[2009.01825](#)].
- [72] R. Garani, D. Levkov and P. Tinyakov, *Solar mass black holes from neutron stars and bosonic dark matter*, *Phys. Rev. D* **105** (2022) 063019 [[2112.09716](#)].
- [73] H. Steigerwald, V. Marra and S. Profumo, *Revisiting constraints on asymmetric dark matter from collapse in white dwarf stars*, *Phys. Rev. D* **105** (2022) 083507 [[2203.09054](#)].
- [74] D. Singh, A. Gupta, E. Berti, S. Reddy and B. S. Sathyaprakash, *Constraining properties of asymmetric dark matter candidates from gravitational-wave observations*, **2210.15739**.

- [75] A. Ray, *Celestial objects as strongly-interacting nonannihilating dark matter detectors*, *Phys. Rev. D* **107** (2023) 083012 [2301.03625].
- [76] S. Bhattacharya, B. Dasgupta, R. Laha and A. Ray, *Can LIGO Detect Nonannihilating Dark Matter?*, *Phys. Rev. Lett.* **131** (2023) 091401 [2302.07898].
- [77] J. Bramante and N. Raj, *Dark matter in compact stars*, *Phys. Rept.* **1052** (2024) 1 [2307.14435].
- [78] D. N. Page, *Particle Emission Rates from a Black Hole: Massless Particles from an Uncharged, Nonrotating Hole*, *Phys. Rev. D* **13** (1976) 198.
- [79] J. H. MacGibbon, *Quark and gluon jet emission from primordial black holes. 2. The Lifetime emission*, *Phys. Rev. D* **44** (1991) 376.
- [80] A. Arbey and J. Auffinger, *Physics Beyond the Standard Model with BlackHawk v2.0*, *Eur. Phys. J. C* **81** (2021) 910 [2108.02737].
- [81] J. Bramante, A. Buchanan, A. Goodman and E. Lodhi, *Terrestrial and Martian Heat Flow Limits on Dark Matter*, *Phys. Rev. D* **101** (2020) 043001 [1909.11683].
- [82] S. R. Taylor and J. R. Gair, *Cosmology with the lights off: standard sirens in the Einstein Telescope era*, *Phys. Rev. D* **86** (2012) 023502 [1204.6739].
- [83] S. B. Popov and M. E. Prokhorov, *Population synthesis in astrophysics*, *Phys. Usp.* **50** (2007) 1123.
- [84] M. Moe, K. M. Kratter and C. Badenes, *The Close Binary Fraction of Solar-type Stars Is Strongly Anticorrelated with Metallicity*, *Astrophysical Journal* **875** (2019) 61 [1808.02116].
- [85] M. Maggiore, *Gravitational Waves: Volume 1: Theory and Experiments*, vol. 1. Oxford University Press, 2008.
- [86] P. Astone, A. Colla, S. D'Antonio, S. Frasca and C. Palomba, *Method for all-sky searches of continuous gravitational wave signals using the Frequency-Hough transform*, *Physical Review D* **90** (2014) 042002.
- [87] C. J. Moore, R. H. Cole and C. P. L. Berry, *Gravitational-wave sensitivity curves*, *Class. Quant. Grav.* **32** (2015) 015014 [1408.0740].
- [88] V. Gluscevic and K. K. Boddy, *Constraints on Scattering of keV–TeV Dark Matter with Protons in the Early Universe*, *Phys. Rev. Lett.* **121** (2018) 081301 [1712.07133].
- [89] K. K. Boddy, V. Gluscevic, V. Poulin, E. D. Kovetz, M. Kamionkowski and R. Barkana, *Critical assessment of CMB limits on dark matter-baryon scattering: New treatment of the relative bulk velocity*, *Phys. Rev. D* **98** (2018) 123506 [1808.00001].
- [90] DES collaboration, E. O. Nadler et al., *Milky Way Satellite Census. III. Constraints on Dark Matter Properties from Observations of Milky Way Satellite Galaxies*, *Phys. Rev. Lett.* **126** (2021) 091101 [2008.00022].
- [91] E. O. Nadler, V. Gluscevic, K. K. Boddy and R. H. Wechsler, *Constraints on Dark Matter Microphysics from the Milky Way Satellite Population*, *Astrophys. J. Lett.* **878** (2019) 32 [1904.10000].
- [92] G. D. Starkman, A. Gould, R. Esmailzadeh and S. Dimopoulos, *Opening the Window on Strongly Interacting Dark Matter*, *Phys. Rev. D* **41** (1990) 3594.
- [93] A. Bhoonah, J. Bramante, B. Courtman and N. Song, *Etched plastic searches for dark matter*, *Phys. Rev. D* **103** (2021) 103001 [2012.13406].
- [94] B. D. Wandelt, R. Dave, G. R. Farrar, P. C. McGuire, D. N. Spergel and P. J. Steinhardt, *Selfinteracting dark matter*, in *4th International Symposium on Sources and Detection of Dark Matter in the Universe (DM 2000)*, pp. 263–274, 6, 2000, [astro-ph/0006344](#).
- [95] B. J. Kavanagh, *Earth scattering of superheavy dark matter: Updated constraints from detectors old and new*, *Phys. Rev. D* **97** (2018) 123013 [1712.04901].
- [96] M. C. Digman, C. V. Cappiello, J. F. Beacom, C. M. Hirata and A. H. G. Peter, *Not as big as a barn: Upper bounds on dark matter-nucleus cross sections*, *Phys. Rev. D* **100** (2019) 063013 [1907.10618].
- [97] R. S. Chivukula, A. G. Cohen, S. Dimopoulos and T. P. Walker, *Bounds on Halo Particle Interactions From Interstellar Calorimetry*, *Phys. Rev. Lett.* **65** (1990) 957.
- [98] A. Bhoonah, J. Bramante, F. Elahi and S. Schon, *Calorimetric Dark Matter Detection With Galactic Center Gas Clouds*, *Phys. Rev. Lett.* **121** (2018) 131101 [1806.06857].
- [99] A. Bhoonah, J. Bramante, S. Schon and N. Song, *Detecting composite dark matter with long-range and contact interactions in gas clouds*, *Phys. Rev. D* **103** (2021) 123026 [2010.07240].
- [100] M. Clark, A. Depoian, B. Elshimy, A. Kopec, R. F. Lang, S. Li et al., *Direct Detection Limits on Heavy Dark Matter*, *Phys. Rev. D* **102** (2020) 123026 [2009.07909].
- [101] (DEAP COLLABORATION)†, DEAP collaboration, P. Adhikari et al., *First Direct Detection Constraints on Planck-Scale Mass Dark Matter with Multiple-Scatter Signatures Using the DEAP-3600 Detector*, *Phys. Rev. Lett.* **128** (2022) 011801 [2108.09405].
- [102] C. V. Cappiello, J. I. Collar and J. F. Beacom, *New experimental constraints in a new landscape for composite dark matter*, *Phys. Rev. D* **103** (2021) 023019 [2008.10646].
- [103] A. L. Erickcek, P. J. Steinhardt, D. McCammon and P. C. McGuire, *Constraints on the Interactions between Dark Matter and Baryons from the X-ray Quantum Calorimetry Experiment*, *Phys. Rev. D* **76** (2007) 042007 [0704.0794].
- [104] W.-R. Hu and Y.-L. Wu, *The Taiji Program in Space for gravitational wave physics and the nature of gravity*, *Natl. Sci. Rev.* **4** (2017) 685.
- [105] TIANQIN collaboration, J. Luo et al., *TianQin: a space-borne gravitational wave detector*, *Class. Quant. Grav.* **33** (2016) 035010 [1512.02076].