

QUANTUM RESONANCES AND SCATTERING POLES OF CLASSICAL RANK ONE LOCALLY SYMMETRIC SPACES

BENJAMIN DELARUE AND JOACHIM HILGERT

ABSTRACT. For negatively curved symmetric spaces it is known from [HHP19] that the poles of the scattering matrices defined via the standard intertwining operators for the spherical principal representations of the isometry group are either given as poles of the intertwining operators or as quantum resonances, i.e. poles of the meromorphically continued resolvents of the Laplace-Beltrami operator. We extend this result to classical locally symmetric spaces of negative curvature with convex-cocompact fundamental group using results of Bunke and Olbrich. The method of proof forces us to exclude the spectral parameters corresponding to singular Poisson transforms.

1. INTRODUCTION

Let G be a non-compact connected classical real simple Lie group of real rank 1. We assume for simplicity that the center of G is finite and fix a maximal compact subgroup K of G . The quotient $X := G/K$ is a non-compact Riemannian symmetric space of rank 1.

Suppose that $\Gamma \subset G$ is a discrete torsion-free subgroup. We denote the limit set of Γ in the geodesic boundary ∂X of X by Λ and set $\Omega := \partial X \setminus \Lambda$. Further, we assume that Γ is a convex-cocompact subgroup, by which we mean that the quotient $\Gamma \backslash (X \cup \Omega)$ is a compact manifold with boundary. Ω is then called the *domain of discontinuity* of Γ in ∂X and $\Gamma \backslash \Omega$ is called the *boundary at infinity* of $\Gamma \backslash X$. We will always assume that Γ is not cocompact, which means that the boundary of $\Gamma \backslash (X \cup \Omega)$ is non-empty.

Let $\Delta_X : C_c^\infty(X) \rightarrow C_c^\infty(X)$ denote the positive Laplacian of X which is a G -invariant differential operator. Furthermore, there is the positive Laplacian $\Delta_{\Gamma \backslash X} : C_c^\infty(\Gamma \backslash X) \rightarrow C_c^\infty(\Gamma \backslash X)$ of $\Gamma \backslash X$. Since X and $\Gamma \backslash X$ are complete Riemannian manifolds, Δ_X and $\Delta_{\Gamma \backslash X}$ are essentially self-adjoint operators in $L^2(X)$ and $L^2(\Gamma \backslash X)$, respectively; we denote their unique self-adjoint extensions again by Δ_X and $\Delta_{\Gamma \backslash X}$.

The *quantum resonances* are the poles of the meromorphically continued modified L^2 -resolvents

$$\begin{aligned} R_\zeta &:= (\Delta_X - \|\rho\|^2 - \zeta^2)^{-1} : L^2(X) \rightarrow L^2(X), \\ R_\zeta^\Gamma &:= (\Delta_{\Gamma \backslash X} - \|\rho\|^2 - \zeta^2)^{-1} : L^2(\Gamma \backslash X) \rightarrow L^2(\Gamma \backslash X), \end{aligned} \tag{1.1}$$

which are holomorphic operator families, respectively, on the *physical half-plane* $\mathcal{H}_{\text{phys}} := \{\zeta \in \mathbb{C} \mid \text{Im } \zeta > 0\}$. Here ρ denotes the usual half-sum of positive restricted roots and $\|\cdot\|$ is the norm defined by the Killing form, see Section 2 for the setup of notation.

The *scattering poles* mentioned in the title are the poles of the meromorphic families S_λ and S_λ^Γ of scattering matrices which are constructed from the standard intertwining

operators for the spherical principal series representations of G . For the definition of the scattering matrices S_λ^Γ on $\Gamma \backslash X$ we build on the constructions from [BO00], which for technical reasons exclude the case $G = F_4^{-20}$. This is the only reason why we restrict our attention to the classical rank one spaces.

The relation between quantum resonances and scattering poles for the global rank one symmetric spaces was described explicitly in [HHP19]. The quantum resonances are given as the zeros of the function $c(i\zeta)c(-i\zeta)$ with $\text{Im } \zeta > 0$, where c is a variant of the Harish-Chandra c -function. The zeros of c are known (see e.g. [HHP19, Rem. 5.1]). [HHP19, Thm. 7.1]. The main result of that paper says that the scattering poles contained in $\mathcal{H}_{\text{phys}}$ are precisely the quantum resonances, whereas the scattering poles which are not quantum resonances are the poles of the family of standard intertwining operators.

In [Hil24, Problem 6.5] it was suggested that one can use results of Bunke and Olbrich to extend the relation between quantum resonances and scattering poles to convex-cocompact locally symmetric spaces of rank one. In fact, in [BO12] Bunke and Olbrich studied the meromorphic continuation of the resolvent kernels as well as the scattering poles for the classical locally symmetric spaces with negative curvature and convex-cocompact fundamental group. Their analysis is based on the meromorphic continuation of a family of operators extending Γ -invariant distributions on Ω to Γ -invariant distributions on ∂X . Our point of departure is [BO12, Lemma 11], where they show that the difference between the resolvent kernels on X and the lift of the resolvent kernels on $\Gamma \backslash X$ has boundary values on $\partial X \times \partial X$ which depend meromorphically on the spectral parameters. A careful analysis of the proof of this lemma yields a meromorphic function relating the the boundary values of the lifted resolvent kernels of $\Gamma \backslash X$ and the lifted kernels of the scattering matrices. This allows us to generalize the statement about the scattering poles being quantum resonances except for the exceptional spectral parameters for which the Poisson transform is singular so that no boundary value map is defined. More precisely, we can define a physical half-plane $\mathcal{H}_{\text{phys}}^\Gamma$ depending on Γ such that our main result can be formulated as follows (see Theorem 5.4).

Theorem. *Outside the set $-\frac{i}{2} \|\alpha_0\| \mathbb{N}$, the scattering poles which are not in the physical plane $\mathcal{H}_{\text{phys}}^\Gamma$ are precisely the quantum resonances.*

Here α_0 is the unique reduced root in a chosen positive system, see Section 2.

Acknowledgements. BD has received funding from the Deutsche Forschungsgemeinschaft (German Research Foundation, DFG) through the Priority Program (SPP) 2026 “Geometry at Infinity”. JH acknowledges support by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) via the grant SFB-TRR 358/1 2023 — 491392403.

2. SETUP

Let $\mathfrak{g} = \mathfrak{k} \oplus \mathfrak{p}$ be the Cartan decomposition of the Lie algebra of G , where \mathfrak{k} is the Lie algebra of K , and denote by $\langle \cdot, \cdot \rangle$ the inner product on \mathfrak{g} defined by the Killing form and the Cartan involution θ , and by $\|\cdot\|$ the associated norm. Let $\mathfrak{a} \subset \mathfrak{p}$ be a maximal abelian subspace, $\Sigma \subset \mathfrak{a}^*$ the root system of $(\mathfrak{g}, \mathfrak{a})$, and fix a choice $\Sigma^+ \subset \Sigma$ of positive

roots, corresponding to a choice of positive Weyl chamber $\mathfrak{a}^+ \subset \mathfrak{a}$. Denote by $\alpha_0 \in \Sigma^+$ the unique reduced root. Since $\dim \mathfrak{a} = 1$, there is precisely one element $H_0 \in \mathfrak{a}^+$ with $\|H_0\| = 1$; it satisfies $\alpha_0(H_0) = \|\alpha_0\|$. Let $\mathfrak{n} := \bigoplus_{\alpha \in \Sigma^+} \mathfrak{g}_\alpha$ be the nilpotent Lie algebra spanned by the root spaces \mathfrak{g}_α and $\rho := \frac{1}{2} \sum_{\alpha \in \Sigma^+} m_\alpha \alpha \in \mathfrak{a}^*$ the half-sum of the positive roots, weighted according to their multiplicities $m_\alpha := \dim \mathfrak{g}_\alpha$.

Let $A, N \subset G$ be the analytic subgroups with Lie algebras \mathfrak{a} and \mathfrak{n} , respectively, and let $M := Z_K(A)$ be the centralizer of A in K . Then the Iwasawa decomposition of G is given by $G = KAN$ and $P := MAN$ is a minimal parabolic subgroup of G . We denote by $\exp : \mathfrak{g} \rightarrow G$ the exponential map and by $\kappa : G \rightarrow K$, $H : G \rightarrow \mathfrak{a}$ the smooth maps such that $g \in \kappa(g) \exp(H(g))N$ for all $g \in G$. For $Y \in \mathfrak{a}$, $a = \exp(Y) \in A$, $\lambda \in \mathfrak{a}_\mathbb{C}^* := \mathfrak{a}^* \otimes_\mathbb{R} \mathbb{C}$, we write $a^\lambda := e^{\lambda(Y)} \in \mathbb{C}$. We fix the Haar measure dk on K of volume 1.

3. RESOLVENT KERNELS

It is known from [HC76] and [BO00] that the continuous spectra of Δ_X as well as $\Delta_{\Gamma \backslash X}$ are given by the interval $[\|\rho\|^2, \infty)$, with $\Delta_{\Gamma \backslash X}$ having at most finitely many eigenvalues outside this interval. $\|\rho\|^2$ is actually the minimum of the spectrum of Δ_X , but in order to describe the minimum of the spectrum of $\Delta_{\Gamma \backslash X}$ we have to introduce the *critical exponent*

$$\begin{aligned} \delta_\Gamma &:= \inf \left\{ s > 0 \mid \sum_{\gamma \in \Gamma} e^{-sd_X(o, \gamma o)} < +\infty \right\} \\ &= \inf \left\{ s > 0 \mid \forall x, y \in X : \sum_{\gamma \in \Gamma} e^{-sd_X(x, \gamma y)} < +\infty \right\}, \end{aligned} \tag{3.1}$$

where $d_X : X \times X \rightarrow \mathbb{R}$ is the Riemannian distance and $o := K \in G/K = X$ is the canonical base point. The critical exponent lies in the interval $[0, 2\|\rho\|]$ (see [Web08, Lemma 3.1]).

By [Cor90, Thm. 4.2] (see also [AZ22, Thm. 1] as well as [HC76] and [BO00]) the minimum of the spectrum of $\Delta_{\Gamma \backslash X}$ is $\|\rho\|^2$ if $0 \leq \delta_\Gamma \leq \|\rho\|$, whereas in the case $\|\rho\| < \delta_\Gamma \leq 2\|\rho\|$ the latter minimum is given by $\|\rho\|^2 - (\delta_\Gamma - \|\rho\|)^2$. Thus, while the family $\zeta \mapsto R_\zeta$ is holomorphic on the physical halfplane $\mathcal{H}_{\text{phys}}$, the family $\zeta \mapsto R_\zeta^\Gamma$ is holomorphic only on the halfplane

$$\mathcal{H}_{\text{phys}}^\Gamma := \begin{cases} \{\zeta \in \mathbb{C} \mid \text{Im } \zeta > 0\} & \text{for } 0 \leq \delta_\Gamma \leq \|\rho\| \\ \{\zeta \in \mathbb{C} \mid \text{Im } \zeta > \delta_\Gamma - \|\rho\|\} & \text{for } \|\rho\| < \delta_\Gamma \leq 2\|\rho\|. \end{cases}$$

Denote by $r_\zeta \in \mathcal{D}'(X \times X)$ and $r_\zeta^\Gamma \in \mathcal{D}'((\Gamma \backslash X) \times (\Gamma \backslash X))$ the Schwartz kernels of R_ζ and R_ζ^Γ , respectively, and by $\tilde{r}_\zeta^\Gamma \in \mathcal{D}'(X \times X)^{\Gamma \times \Gamma}$ the distribution corresponding to r_ζ^Γ when identifying $\mathcal{D}'((\Gamma \backslash X) \times (\Gamma \backslash X)) = \mathcal{D}'(X \times X)^{\Gamma \times \Gamma}$. Note here that the identification uses the pullback via the canonical projection $\text{pr}_\Gamma : X \rightarrow \Gamma \backslash X$, which is a submersion (cf. [Hör90, Thm. 6.12]).

When ζ lies sufficiently deep inside the physical half-plane, \tilde{r}_ζ^Γ is nothing but the “ Γ -summation” (or “ Γ -average”) of r_ζ . To make this precise we formulate the following lemma.

Lemma 3.1 (C.f. [BO12, Lemma 9]). *For all $\zeta \in \mathcal{H}_{\text{phys}}^\Gamma$ satisfying $\text{Im } \zeta > \sqrt{(\text{Re } \zeta)^2 + c_\Gamma}$, one has*

$$\tilde{r}_\zeta^\Gamma = \sum_{\gamma \in \Gamma} (e, \gamma)^* r_\zeta, \quad (3.2)$$

where

$$c_\Gamma := \delta_\Gamma^2 + \max(0, \delta_\Gamma - \|\rho\|)^2 \in [\delta_\Gamma^2, 2\delta_\Gamma^2]. \quad (3.3)$$

and $(\gamma_1, \gamma_2)^*$ denotes the pullback action of an element $(\gamma_1, \gamma_2) \in \Gamma \times \Gamma$. In particular, the right-hand side of (3.2) is a well-defined distribution on $X \times X$.

Proof. We argue as in [BO12, proof of Lemma 9], adding (and repeating) some details. Let $\text{diag}_X \subset X \times X$ be the diagonal and define

$$D_\Gamma := (\{e\} \times \Gamma) \cdot \text{diag}_X = (\Gamma \times \{e\}) \cdot \text{diag}_X \subset X \times X.$$

The proper discontinuity of the Γ -action on X implies that D_Γ is a closed subset of $X \times X$ (c.f. [Kap23, Thm. 11 (6)]). As pointed out in [BO12, p. 137], the distribution r_ζ is smooth outside the diagonal, which implies that on $X \times X \setminus D_\Gamma$ all the distributions $(e, \gamma)^* r_\zeta$, $\gamma \in \Gamma$, are smooth. Now, let $f \in C_c^\infty(X \times X)$ and let $\Gamma = \{\gamma_1, \gamma_2, \dots\}$ be an enumeration of the group Γ . Then we have for each $N \in \mathbb{N}$

$$\begin{aligned} \sum_{n=1}^N |((e, \gamma_n)^* r_\zeta)(f)| &= \sum_{\substack{n \leq N: \\ (\text{supp } f) \cap (e, \gamma_n^{-1}) \cdot (\text{supp } f) = \emptyset}} |((e, \gamma_n)^* r_\zeta)(f)| \\ &\quad + \sum_{\substack{n \leq N: \\ (\text{supp } f) \cap (e, \gamma_n^{-1}) \cdot (\text{supp } f) \neq \emptyset}} |((e, \gamma_n)^* r_\zeta)(f)| \\ &\leq \sum_{\substack{n \leq N: \\ (\text{supp } f) \cap (e, \gamma_n^{-1}) \cdot (\text{supp } f) = \emptyset}} |((e, \gamma_n)^* r_\zeta)(f)| + \sum_{\substack{\gamma \in \Gamma: \\ (\text{supp } f) \cap (e, \gamma) \cdot (\text{supp } f) \neq \emptyset}} |((e, \gamma)^* r_\zeta)(f)|. \end{aligned}$$

Here the last term is a finite sum because the $(\{e\} \times \Gamma)$ -action on $X \times X$ is properly discontinuous. On the other hand, for each $\gamma \in \Gamma$ such that $(\text{supp } f) \cap (e, \gamma^{-1}) \cdot (\text{supp } f) = \emptyset$, we can write

$$\begin{aligned} |((e, \gamma)^* r_\zeta)(f)| &= |((e, \gamma)^* r_\zeta)(f|_{X \times X \setminus D_\Gamma})| \\ &= \left| \int_{X \times X \setminus D_\Gamma} ((e, \gamma)^* r_\zeta)(x, y) \overline{f(x, y)} \, dx dy \right| \end{aligned}$$

and then apply [BO12, Lemma 8], as well as the expression for the minimum of the spectrum of $\Delta_{\Gamma \backslash X}$ stated above, to estimate for every compact set $W \subset \mathcal{H}_{\text{phys}}^\Gamma$:

$$|((e, \gamma)^* r_\zeta)(f)| \leq C \|f\|_\infty e^{-a \sqrt{-\max(0, \delta_\Gamma - \|\rho\|)^2 - \text{Re } (\zeta^2)}} \quad \forall \zeta \in W$$

with a constant $C > 0$ depending on W and $\text{supp } f$, and the exponent

$$a := \inf_{(x, y) \in \text{supp } f} d_X(x, \gamma y) > 0.$$

Now, in view of the definition of the critical exponent δ_Γ , the condition for convergence as $N \rightarrow \infty$ becomes

$$\sqrt{-\max(0, \delta_\Gamma - \|\rho\|)^2 - \operatorname{Re}(\zeta^2)} > \delta_\Gamma,$$

which is equivalent to

$$(\operatorname{Im} \zeta)^2 > (\operatorname{Re} \zeta)^2 + \underbrace{\delta_\Gamma^2 + \max(0, \delta_\Gamma - \|\rho\|)^2}_{=c_\Gamma}.$$

The convergence estimate depends on f only via $\operatorname{supp} f$ and $\|f\|_\infty$, so that we obtain distributional convergence of the right-hand side of (3.2) for all $\zeta \in \mathcal{H}_{\text{phys}}$ with $\operatorname{Im} \zeta > \sqrt{(\operatorname{Re} \zeta)^2 + c_\Gamma}$.

It remains to show that the right-hand side of (3.2) equals \tilde{r}_ζ^Γ , which can be done exactly as in the end of the proof of [BO12, Lemma 9]. \square

Remark 3.2. In (3.2) one could equivalently take $\sum_{\gamma \in \Gamma} (\gamma, e)^* r_\zeta$: since Δ_X is G -equivariant, so is each of its resolvents R_ζ for $\zeta \in \mathcal{H}_{\text{phys}}$. Thus r_ζ satisfies $(g, g^{-1}) \cdot r_\zeta = r_\zeta$ for all $g \in G$.

By [BO12, Lemma 6], r_ζ and r_ζ^Γ (equivalently \tilde{r}_ζ^Γ) extend to \mathbb{C} as meromorphic families of distributions. The meromorphic families of operators $C_c^\infty(X) \rightarrow \mathcal{D}'(X)$ defined by the distribution kernels r_ζ and \tilde{r}_ζ^Γ will be denoted by R_ζ and \tilde{R}_ζ^Γ , respectively (where \tilde{R}_ζ^Γ maps actually into $\mathcal{D}'(X)^\Gamma$ thanks to the $\Gamma \times \Gamma$ -invariance of \tilde{r}_ζ^Γ), while we denote the meromorphic family of operators $C_c^\infty(\Gamma \backslash X) \rightarrow \mathcal{D}'(\Gamma \backslash X)$ with distribution kernels r_ζ^Γ by R_ζ^Γ . This is compatible with the previous notation in the sense that R_ζ and R_ζ^Γ are meromorphic extensions of the operator families from (1.1) with domains shrunk to compactly supported smooth functions and codomains enlarged to distributions.

4. SCATTERING MATRICES

In this section we keep the setup from Section 2 and follow [BO00] in the definition of $\lambda \mapsto S_\lambda^\Gamma$ as a meromorphic family of the scattering matrices for $\Gamma \backslash X$. It is our goal to relate the poles of this family, which we call *scattering poles*, to the resonances of $\Gamma \backslash X$, i.e. the poles of the meromorphically continued family $\zeta \mapsto R_\zeta^\Gamma$ of resolvents.

4.1. Spherical principal series representations. In order to introduce the scattering matrices we have to review some facts about spherical principal series representations. We warn the reader that there is no fixed standard notation in the literature. As we will have to use spectral information from different sources, we will have to keep track of the different conventions (see Remark 4.2 below).

For $\lambda \in \mathfrak{a}_\mathbb{C}^*$ we define the one-dimensional group representation

$$\sigma_\lambda : P = MAN \rightarrow \mathbb{C} \setminus \{0\}, \quad man \mapsto a^{\rho+\lambda}, \quad (4.1)$$

and the associated homogeneous line bundle

$$V(\sigma_\lambda) := G \times_P \mathbb{C} \rightarrow G/P, \quad [g, z] \mapsto gP,$$

where $[g, z]$ is the P -orbit of $(g, z) \in G \times \mathbb{C}$ with respect to the right P -action given by $(g, z) \cdot p = (gp, \sigma_\lambda(p^{-1})z)$. Note that G acts on the total space of $V(\sigma_\lambda)$ from the left. The space $C^\infty(G/P; V(\sigma_\lambda))$ of all smooth sections $s : G/P \rightarrow V(\sigma_\lambda)$ can be identified with

$$H^{\lambda, \infty} := \{f \in C^\infty(G) \mid \forall g \in G, p \in P : f(gp) = \sigma_\lambda(p)^{-1}f(g)\} \quad (4.2)$$

so that $f \in H^{\lambda, \infty}$ corresponds to the section s of $V(\sigma_\lambda)$ with $s(gP) = [g, f(g)]$.

The left-regular representation of G on $H^{\lambda, \infty}$ will be denoted by $\pi^{\lambda, \infty}$:

$$(\pi^{\lambda, \infty}(g)f)(g') := f(g^{-1}g'), \quad f \in H^{\lambda, \infty}. \quad (4.3)$$

It is called (spherical, smooth) principal series representation with parameter λ .

Due to the Iwasawa decomposition the restriction map

$$\begin{aligned} H^{\lambda, \infty} &\rightarrow C^\infty(K)^M := \{f \in C^\infty(K) \mid \forall k \in K, m \in M : f(km) = f(k)\} \\ f &\mapsto f|_K \end{aligned} \quad (4.4)$$

is an isomorphism of vector spaces. This allows us to consider $H^{\lambda, \infty}$ and $C^\infty(G/P; V(\sigma_\lambda))$ as λ -independent vector spaces. It also shows that $V(\sigma_\lambda)$ is in fact a trivial line bundle but we will soon pass to its Γ -quotient, which need no longer be trivial.

We equip the vector space $C^\infty(G/P; V(\sigma_\lambda)) \cong H^{\lambda, \infty}$ with the Fréchet topology such that (4.4) is an isomorphism of topological vector spaces, where $C^\infty(K)^M$ is equipped with the subspace topology inherited from the standard Fréchet topology on $C^\infty(K)$. This way, we can consider $H^{\lambda, \infty}$ and $C^\infty(G/P; V(\sigma_\lambda))$ as λ -independent Fréchet spaces.

By restricting the line bundle $V(\sigma_\lambda)$ to $\Omega \subset G/P = \partial X$ and passing to the Γ -quotient, we obtain the locally homogeneous vector bundle

$$V_{\Gamma \backslash \Omega}(\sigma_\lambda) := \Gamma \backslash V(\sigma_\lambda)|_\Omega$$

over the smooth manifold $\Gamma \backslash \Omega$. The vector space $C^\infty(\Gamma \backslash \Omega; V_{\Gamma \backslash \Omega}(\sigma_\lambda))$ corresponds canonically to the space of Γ -invariant smooth sections of $V(\sigma_\lambda)|_\Omega$, which in turn corresponds canonically to the closed subspace of left- Γ -invariant functions in $H^{\lambda, \infty}$. Thus, the space $C^\infty(\Gamma \backslash \Omega; V_{\Gamma \backslash \Omega}(\sigma_\lambda))$ inherits a Fréchet topology from $H^{\lambda, \infty}$. In particular, we can consider $C^\infty(\Gamma \backslash \Omega; V_{\Gamma \backslash \Omega}(\sigma_\lambda))$ as a λ -independent Fréchet space, too.

Definition 4.1. We call the topological vector space

$$C^{-\infty}(G/P; V(\sigma_\lambda)) := C^\infty(G/P; V(\sigma_{-\lambda}))',$$

where “prime” denotes the topological dual, the space of *distributional sections* of $V(\sigma_\lambda)$.

We equip $C^{-\infty}(G/P; V(\sigma_\lambda))$ with the dual G -representation of $\pi^{-\lambda, \infty}$, which we denote by $\pi^{\lambda, -\infty}$, and write the subspace of Γ -invariant elements as ${}^\Gamma C^{-\infty}(G/P; V(\sigma_\lambda))$. The map

$$\begin{aligned} C^\infty(G/P; V(\sigma_\lambda)) &\cong H^{\lambda, \infty} \rightarrow C^{-\infty}(G/P; V(\sigma_\lambda)), \\ f &\mapsto \langle f|_K, \cdot \rangle_{L^2(K)} \end{aligned} \quad (4.5)$$

is well-defined, injective (since the restriction (4.4) is injective), and continuous. Moreover, the injection (4.5) is G -equivariant; for this, the sign change of λ in Definition 4.1 is essential. By virtue of this equivariant injection we can view the principal series representation

$(\pi^{\lambda,\infty}, C^\infty(G/P; V(\sigma_\lambda)))$ as a subrepresentation of $(\pi^{\lambda,-\infty}, C^{-\infty}(G/P; V(\sigma_\lambda)))$. We call the latter *distributional principal series representation* with parameter λ .

4.2. Comparing conventions. We will have to keep track of the conventions for meromorphic continuations with respect to complex parameters used in the various sources we rely on. To simplify this task we include a couple of remarks containing the pertinent comparisons.

Remark 4.2. Our definitions (4.1), (4.2) differ from the one in [BO00, Section 3], [Olb94, § 2.2] and [HHP19, § 2] by the sign of λ . We chose this sign convention to be consistent with [Kna86, VII, §1]. This is also consistent with the sign convention in [Vog08, Chap. 3]. In all these references the G -action is the left regular action. On the other hand, [Wal92, § 10.1] and [KS71] use the right regular version of the principal series representations. Note that the map $f \mapsto \tilde{f}$ with $\tilde{f}(g) := f(g^{-1})$ intertwines left and right regular actions. Up to this intertwiner the sign conventions in [Wal92, § 10.1] and [KS71] agree with the sign conventions in [Kna86, VII, §1]. Finally, we point out that our topologies on $C^\infty(G/P; V(\sigma_\lambda))$ and $C^\infty(\Gamma \backslash \Omega; V_{\Gamma \backslash \Omega}(\sigma_\lambda))$ are equivalent to those of [BO00, p. 89].

Remark 4.3 (Compatibility with [BO12, Section 6] and [HHP19]). Our choice of physical half-plane is compatible with [HHP19] when identifying $\mathfrak{a}_{\mathbb{C}}^* = \mathfrak{a}^* \otimes_{\mathbb{R}} \mathbb{C}$ with \mathbb{C} by mapping α_0 to 1 (c.f. the end of Remark 5.1 iii), taking into account the sign change mentioned in Remark 4.2. For compatibility with [BO12, Section 6.1], the bilinear form b in [BO12, Section 6.1] must be chosen such that $b|_{\mathfrak{a} \times \mathfrak{a}} = \langle \cdot, \cdot \rangle|_{\mathfrak{a} \times \mathfrak{a}}$, where $\langle \cdot, \cdot \rangle$ is the Killing form of \mathfrak{g} , which ensures that the norms on $\mathfrak{a}_{\mathbb{C}}^*$ considered in [BO12, Section 6.1] and [HHP19] agree (recall Remark 5.1 iii). Now, let $\tilde{\mathbb{C}}^\pm$ be the branched covers of \mathbb{C} on which the functions

$$\tilde{\mathbb{C}}^\pm \ni z \mapsto \sqrt{\pm(\|\rho\|^2 - z)} \in \mathbb{C} \quad (4.6)$$

are holomorphic, respectively. As Riemann surfaces, both $\tilde{\mathbb{C}}^+$ and $\tilde{\mathbb{C}}^-$ are isomorphic to the Riemann surface $\tilde{\mathbb{C}}$ of the complex square root. This allows us to compare points in $\tilde{\mathbb{C}}^+$ and $\tilde{\mathbb{C}}^-$ by identifying these spaces with $\tilde{\mathbb{C}}$. To achieve compatibility with [BO12, Section 6] and [HHP19] one finds that for both $\tilde{\mathbb{C}}^+$ and $\tilde{\mathbb{C}}^-$ one needs to choose $(\|\rho\|^2, \infty)$ as the branch cut. For $\tilde{\mathbb{C}}^+$ this corresponds to the “principal square root” branch cut of $\tilde{\mathbb{C}}$ along $(-\infty, 0]$, while for $\tilde{\mathbb{C}}^-$ it corresponds to the “opposite” branch cut of $\tilde{\mathbb{C}}$ along $[0, \infty)$. We define the *physical sheet* $\tilde{\mathbb{C}}_{\text{phys}}^- \subset \tilde{\mathbb{C}}^-$ over $\mathbb{C} \setminus (\|\rho\|^2, \infty)$ as the sheet which is mapped bi-holomorphically onto $\mathcal{H}_{\text{phys}}$ by the function with the $-$ sign in (4.6)

$$\tilde{\mathbb{C}}_{\text{phys}}^- \ni z \mapsto \sqrt{z - \|\rho\|^2} =: \zeta =: \zeta(z) \in \mathcal{H}_{\text{phys}},$$

providing us with the “physical parameter” $\zeta \in \mathcal{H}_{\text{phys}}$.

Bunke and Olbrich, on the other hand, work with the parameter

$$\lambda := \lambda(z) := \sqrt{\|\rho\|^2 - z}$$

obtained by considering the $+$ sign in (4.6). For the conversion from λ to ζ and vice-versa, there are the two possibilities $\lambda = \pm i\zeta$, which amount to choosing $\{\mp \operatorname{Re} \lambda > 0\} \subset \mathbb{C}$ as the physical λ -half-plane corresponding to the physical ζ -half-plane $\mathcal{H}_{\text{phys}}$. In [BO12, p. 135] Bunke and Olbrich work with the right half-plane but their argument works just as well with the left half-plane. Indeed, there is no preferred choice of the physical sheet in [BO12, Section 6], so we can use their results regardless of which sign we choose in the relation between the parameters λ and ζ . The following choice will turn out to be convenient:

$$\begin{aligned} \lambda &:= \lambda(\zeta) := -i\zeta & \forall \zeta \in \mathcal{H}_{\text{phys}}, \\ \zeta &= \zeta(\lambda) = i\lambda, & \forall \lambda \in \mathbb{C}, \operatorname{Re} \lambda > 0. \end{aligned} \quad (4.7)$$

4.3. Scattering matrices. We build on the descriptions of scattering matrices given in [BO00] and [HHP19].

Then, according to [BO00, pp. 79–80], using their isometric identification $\mathbb{C} \ni \lambda \leftrightarrow \lambda \frac{\rho}{\|\rho\|} \in \mathfrak{a}_{\mathbb{C}}^*$, we have the three meromorphic families on \mathbb{C} of maps

$$\operatorname{ext}_{\lambda} : C^{-\infty}(\Gamma \backslash \Omega, V_{\Gamma \backslash \Omega}(\sigma_{\lambda})) \rightarrow {}^{\Gamma}C^{-\infty}(\partial X, V(\sigma_{\lambda})) \quad (4.8)$$

$$\operatorname{res}_{\lambda} : {}^{\Gamma}C^{-\infty}(\partial X, V(\sigma_{\lambda})) \mapsto C^{-\infty}(\Gamma \backslash \Omega, V_{\Gamma \backslash \Omega}(\sigma_{\lambda})) \quad (4.9)$$

$$\hat{J}_{\lambda} : C^{-\infty}(\partial X, V(\sigma_{\lambda})) \rightarrow C^{-\infty}(\partial X, V(\sigma_{\lambda})), \quad (4.10)$$

where the latter is the “unnormalized” Knapp-Stein intertwiner described in detail in [KS71, Thm. 3] (in which the identification $\mathbb{C} \ni \lambda \leftrightarrow \lambda \rho \in \mathfrak{a}_{\mathbb{C}}^*$ is used, so that our λ corresponds to $\frac{\lambda}{\|\rho\|}$ in [KS71, Thm. 3]). Recall from Remark 4.2 that the P -representation σ_{λ} defined in (4.1) is the same as in [KS71] but corresponds to $\sigma_{-\lambda}$ in [BO00]. The Harish-Chandra c -function is the meromorphic function on $\mathfrak{a}_{\mathbb{C}}^*$ satisfying

$$c(\lambda \rho) := \int_{\bar{N}} e^{-(\lambda+1)\rho(H(\bar{n}))} d\bar{n}, \quad \forall \lambda \in \mathbb{C}, \operatorname{Re} \lambda > 0.$$

In view of the chosen identification of $\mathfrak{a}_{\mathbb{C}}^*$ with \mathbb{C} and the sign difference in the conventions for the principal series parameter used here and in [BO00], respectively, it is convenient to consider the meromorphic function $c_{\mathbb{C}}$ on \mathbb{C} defined by

$$c_{\mathbb{C}}(\lambda) := c\left(\frac{-\lambda}{\|\rho\|}\rho\right).$$

Then we define the normalized Knapp-Stein intertwiner

$$J_{\lambda} := \frac{1}{c_{\mathbb{C}}(\lambda)} \hat{J}_{\lambda}.$$

Note that our J_{λ} corresponds to $J_{1,-\lambda}$ in [BO00, p. 101].

Recall the critical exponent δ_{Γ} from (3.1) and note that if $\delta_{\Gamma}^{\text{BO}}$ denotes the critical exponent from [BO00, Def. 2.2], then $\delta_{\Gamma} = \delta_{\Gamma}^{\text{BO}} + \|\rho\|$.

Definition 4.4 (Scattering matrices, [BO00, Def. 5.6]). Let $\operatorname{Re}(\lambda) < \|\rho\| - \delta_{\Gamma}$. Then $S_{\lambda}^{\Gamma} := \operatorname{res}_{\lambda} \circ J_{\lambda} \circ \operatorname{ext}_{\lambda}$ is called the λ -scattering matrix for $\Gamma \backslash X$.

Note that our S_λ^Γ corresponds to $S_{-\lambda}$ in [BO00, Def. 5.6].

From [BO00, Lemma 5.7] we know that for $\operatorname{Re}(\lambda) < \|\rho\| - \delta_\Gamma$ the family

$$\lambda \mapsto S_\lambda^\Gamma : C^{\pm\infty}(\Gamma \backslash \Omega, V_{\Gamma \backslash \Omega}(\sigma_\lambda)) \rightarrow C^{\pm\infty}(\Gamma \backslash \Omega, V_{\Gamma \backslash \Omega}(\sigma_{-\lambda}))$$

is meromorphic. More precisely, we have the following result.

Proposition 4.5 (Scattering matrices, [BO00, Thm. 5.10]). *Suppose that $G \neq F_4^{-20}$. Then the families*

$$\lambda \mapsto S_\lambda^\Gamma : C^{\pm\infty}(\Gamma \backslash \Omega, V_{\Gamma \backslash \Omega}(\sigma_\lambda)) \rightarrow C^{\pm\infty}(\Gamma \backslash \Omega, V_{\Gamma \backslash \Omega}(\sigma_{-\lambda}))$$

and

$$\lambda \mapsto \operatorname{ext}_\lambda : C^{-\infty}(\Gamma \backslash \Omega, V_{\Gamma \backslash \Omega}(\sigma_\lambda)) \rightarrow {}^\Gamma C^{-\infty}(\partial X, V(\sigma_\lambda)),$$

originally defined for $\operatorname{Re}(\lambda) < \|\rho\| - \delta_\Gamma$, have meromorphic continuations to all of \mathbb{C} . The family ext has at most finite-dimensional singularities. Moreover, the following identities hold:

$$\operatorname{ext}_\lambda = J_{-\lambda} \circ \operatorname{ext}_\lambda \circ S_\lambda^\Gamma, \quad (4.11)$$

$$S_{-\lambda}^\Gamma \circ S_\lambda^\Gamma = \operatorname{id}. \quad (4.12)$$

It is expected that the necessity of hypothesis $G \neq F_4^{-20}$ in Proposition 4.5 is an artefact of its proof. Note that the use of this result is the only reason why we had to restrict our attention to classical groups. We will address this issue elsewhere.

Definition 4.6 (Scattering matrices for the globally symmetric space, c.f. [HHP19, §7]). For $\zeta \in \mathcal{H}_{\text{phys}}$ we define

$$S_\zeta := J_{i\zeta}, \quad (4.13)$$

the ζ -scattering matrix for X .

Note that our S_ζ corresponds to $S_{-\frac{\zeta\alpha_0}{\|\alpha_0\|}}$ in [HHP19, §7]. This takes into account that our principal series representation parameter λ corresponds to $-\lambda$ in [HHP19], while our resolvent R_ζ corresponds to $R_{-\frac{\zeta\alpha_0}{\|\alpha_0\|}} = R_{\frac{\zeta\alpha_0}{\|\alpha_0\|}}$ in [HHP19].

5. RELATING SCATTERING MATRICES AND RESOLVENT KERNELS

The key to the comparison between scattering matrices and resolvent kernels for *global* symmetric spaces given in [HHP19] was the possibility to take boundary values of Laplace eigenfunctions on X . The boundary value maps are given as the inverses of Poisson transforms. Moreover, the proof of [BO12, Lemma 11] suggests a way how to use the knowledge about the relation between resolvent poles and scattering poles of rank one global symmetric spaces in order to determine the corresponding relation for rank one *locally* symmetric spaces. We will follow this strategy and start by reviewing a few facts about Poisson transforms and boundary value maps.

5.1. Poisson transforms and boundary values. We define the *Poisson transform* with parameter $\lambda \in \mathfrak{a}_{\mathbb{C}}^*$ by

$$\begin{aligned}\mathcal{P}_\lambda : C^{-\infty}(G/P; V(\sigma_\lambda)) &\rightarrow C^\infty(G/K), \\ \mathcal{P}_\lambda(u)(g) &:= (\pi^{\lambda, -\infty}(g^{-1})u)(1_{- \lambda}),\end{aligned}$$

where $1_{- \lambda} \in C^\infty(G/P; V(\sigma_{-\lambda})) \cong H^{-\lambda, \infty}$ corresponds to the constant function on K with value 1 via the isomorphism (4.4). If $f \in H^{\lambda, \infty} \cong C^\infty(G/P; V(\sigma_\lambda))$ is identified with an element of $C^{-\infty}(G/P; V(\sigma_\lambda))$ via the injection (4.5), then

$$\mathcal{P}_\lambda(f)(g) = \int_K f(gk) dk = \int_K e^{(\lambda - \rho)(H(g^{-1}k))} f(k) dk. \quad (5.1)$$

Remark 5.1 (Comparing conventions). Note that due to our sign convention, see Remark 4.2, our Poisson transform \mathcal{P}_λ corresponds to $P_{-\lambda}^T$ in [BO12, § 4.2] and [BO00, Def. 4.8], respectively to $P_{-\lambda}$ in [HHP19, (2)].

For generic parameters λ the Poisson transformations are essentially inverted by *boundary value* maps. In [HHP19, (29)] this reads

$$\beta_{\rho - \lambda} P_\lambda = c(\lambda) \text{Id}.$$

We need to relate the boundary value maps used in [BO12, § 6.3] to the Poisson transforms used in [BO12, § 4.2] in order to relate them to the boundary value maps β_λ used in [HHP19, Lemma 6.1]. In [BO12, § 6.3] the authors are not very precise about the construction of boundary values they are using. They simply refer to [KO77] and [OS80], but they give the set of leading exponents. We assume they are using the conventions of [Olb94], which are compatible with [BO00, BO12, HHP19]. Note that [Olb94, Thm. 4.17] (see also [Olb94, Thm. 3.9] and [Kna86, Lemma 9.28]) relates the corresponding boundary value maps with the Poisson transform in the following formula

$$\beta_{1, \lambda} \circ P_\lambda^\gamma = c_\gamma(\lambda) \text{Id}.$$

The definitions of the c -functions in [HHP19] and [Olb94] are compatible. In our situation σ and γ are trivial, which means that the boundary value map $\beta_{1, \lambda}$ from [Olb94] equals the boundary value map $\beta_{\rho - \lambda}$ from [HHP19].

In [BO12, § 6.3] Bunke and Olbrich invoke [KO77] to guarantee the existence of boundary values for eigenfunctions of the Laplacians on the two factors of $(X \times X) \setminus D_\Gamma$, see [BO12, (48)]. It is defined for parameters determined by the leading exponents of Casimir eigenfunctions (see [BO12, Lemmas 10 & 11] and recall that in our situation σ and τ are trivial).

In principle the construction of the boundary value maps is built on the study of asymptotic behavior of eigenfunctions. The construction in [KO77] depends on a specific and very technical theory, the algebraic analysis initiated by Sato. If the boundary value map were applied to moderately growing eigenfunctions on all of $X \times X$ we could use [vdBS87] to describe the boundary value maps. This description shows that the direct product situation leads to a decoupling of the asymptotics which leads to a decoupling of the boundary

value map. Taking into account the notation and the various conventions from Remarks 5.1 and 4.3, the boundary value map from [BO12, § 6.3] suitably restricted is given by

$$\beta_\nu = \beta_{\nu(\zeta)} := \beta_{\rho-i\frac{\zeta\alpha_0}{\|\alpha_0\|}} \otimes \beta_{\rho-i\frac{\zeta\alpha_0}{\|\alpha_0\|}}, \quad \zeta \in \mathcal{H}_{\text{phys}}, \quad (5.2)$$

with the boundary value map of [HHP19] on the right-hand side. and $\nu \in (\mathfrak{a}_{\mathbb{C}}^*)^2 = \mathbb{C}^2$ denoting the leading exponents of the eigensections one takes boundary values of.

We restrict our attention to the non-vanishing boundary value maps. For these the parameter $\nu \in \mathbb{C}^2$ is given by

$$\nu := \nu(z) := (-\|\rho\| - \lambda(z), -\|\rho\| - \lambda(z)),$$

where $\lambda(z) := \sqrt{\|\rho\|^2 - z} \in \{\text{Re} > 0\} \subset \mathbb{C}$ (c.f. Remark 4.3), $z \in \mathbb{C}$ is a Casimir eigenvalue, and we use the convention from [BO12, § 6.1] that the complex vector space $\mathfrak{a}_{\mathbb{C}}^*$ is identified with \mathbb{C} by complex linear extension of the unique order-preserving isometric isomorphism $\mathfrak{a}^* \cong \mathbb{R}$. This means that $\rho \in \mathfrak{a}^*$ is identified with the number $\|\rho\| \in \mathbb{R}$ (which accordingly is written simply as ρ in [BO12, § 6]), where the norm is with respect to the inner product on $\mathfrak{a}^* \cong \mathfrak{a}$ given by restriction of the chosen bilinear form b on \mathfrak{g} defining the Riemannian metric on $X = G/K$. In particular, the number $\lambda(z) \in \mathbb{C}$ corresponds to the element

$$\lambda(z) = \sqrt{\|\rho\|^2 - z} \frac{\rho}{\|\rho\|} = \sqrt{1 - \frac{z}{\|\rho\|^2}} \rho \in \mathfrak{a}_{\mathbb{C}}^*. \quad (5.3)$$

The number $-\|\rho\| - \lambda(z)$ is a leading exponent of a Casimir eigenfunction for the eigenvalue z . The boundary value map maps tensor products of Casimir eigenfunctions for the eigenvalue z to tensor products of elements of $H^{1,\lambda(z)}$ (see [BO12, Lemmas 10 & 11] for all this and take the change of signs described in Remark 4.2 into account). Thus β_ν coincides with $\beta_{1,-\lambda(z)} \otimes \beta_{1,-\lambda(z)}$, where $\beta_{1,\lambda}$ is as in [Ol94], so that in view of (ii) we have

$$\beta_\nu = \beta_{\rho+\lambda(z)} \otimes \beta_{\rho+\lambda(z)} \quad (5.4)$$

with the boundary value maps from [HHP19] on the right hand side and the parameters considered as elements of $\mathfrak{a}_{\mathbb{C}}^*$.

To express the relation (5.4) using parameters in \mathbb{C} , one needs to take into account that in [HHP19] the identification of $\mathfrak{a}_{\mathbb{C}}^*$ with \mathbb{C} differs from that in [BO12, § 6]: In [HHP19] the element $\rho \in \mathfrak{a}^*$ gets identified with the number $\frac{\|\rho\|}{\|\alpha_0\|} \in \mathbb{R}$, where $\alpha_0 \in \Sigma^+ \subset \mathfrak{a}^*$ is the unique reduced positive root and the norm is with respect to the Killing form. Let us choose the bilinear form b in [BO12, § 6.1] such that on $\mathfrak{a} \times \mathfrak{a}$ it agrees with the Killing form, so that the norms on $\mathfrak{a}_{\mathbb{C}}^*$ considered in [BO12, § 6.1] and [HHP19] agree. Furthermore, let us denote for any element $\lambda \in \mathfrak{a}_{\mathbb{C}}^*$ the corresponding complex numbers in [BO12, § 6] and [HHP19] by $\lambda_{\text{BO}} \in \mathbb{C}$ and $\lambda_{\text{HHP}} \in \mathbb{C}$, respectively. Then

$$\lambda_{\text{BO}} = \|\alpha_0\| \lambda_{\text{HHP}}. \quad (5.5)$$

In particular, for the element $\lambda(z) \in \mathfrak{a}_{\mathbb{C}}^*$ from (5.3), which satisfies $\lambda(z)_{\text{BO}} = \sqrt{\|\rho\|^2 - z} = \sqrt{\rho_{\text{BO}}^2 - z}$, we find

$$\lambda(z)_{\text{HHP}} = \frac{1}{\|\alpha_0\|} \sqrt{\|\rho\|^2 - z} = \sqrt{\left(\frac{\|\rho\|}{\|\alpha_0\|}\right)^2 - \frac{z}{\|\alpha_0\|^2}} = \sqrt{\rho_{\text{HHP}}^2 - \frac{z}{\|\alpha_0\|^2}}. \quad (5.6)$$

Lemma 5.2. *We have the identity of meromorphic functions*

$$s_{\zeta}|_{\partial X \times \partial X \setminus \text{diag}_{\partial X}} = -2i \|\alpha_0\| \zeta \beta_{\nu(\zeta)}(r_{\zeta}|_{X \times X \setminus \text{diag}_X}). \quad (5.7)$$

Proof. Let $f, g \in C^\infty(\partial X)$ have disjoint supports. Then applying [HHP19, last eq. on p. 20] with the parameter $-\frac{\zeta\alpha_0}{\|\alpha_0\|}$ gives

$$\begin{aligned} \langle s_{\zeta}, f \otimes g \rangle &= \langle S_{\zeta} f, g \rangle \\ &= -2i \|\alpha_0\| \zeta \langle \beta_{\rho-i\frac{\zeta\alpha_0}{\|\alpha_0\|}} (\beta_{\rho-i\frac{\zeta\alpha_0}{\|\alpha_0\|}} R_{\zeta})' f, g \rangle \\ &= -2i \|\alpha_0\| \zeta \langle R_{\zeta} \beta'_{\rho-i\frac{\zeta\alpha_0}{\|\alpha_0\|}} f, \beta'_{\rho-i\frac{\zeta\alpha_0}{\|\alpha_0\|}} g \rangle \\ &= -2i \|\alpha_0\| \zeta \langle r_{\zeta}|_{X \times X \setminus \text{diag}_X}, (\beta'_{\rho-i\frac{\zeta\alpha_0}{\|\alpha_0\|}} \otimes \beta'_{\rho-i\frac{\zeta\alpha_0}{\|\alpha_0\|}})(f \otimes g) \rangle \\ &= -2i \|\alpha_0\| \zeta \langle r_{\zeta}|_{X \times X \setminus \text{diag}_X}, \beta'_{\nu(\zeta)}(f \otimes g) \rangle \\ &= -2i \|\alpha_0\| \zeta \langle \beta_{\nu(\zeta)}(r_{\zeta}|_{X \times X \setminus \text{diag}_X}), f \otimes g \rangle. \end{aligned}$$

The claim follows by density. \square

5.2. Local-global comparisons. Recall the Schwartz kernels r_{ζ} and $\tilde{r}_{\zeta}^{\Gamma}$ of R_{ζ} and $\tilde{R}_{\zeta}^{\Gamma}$, respectively from Section 3. Further, let s_{ζ} and $\tilde{s}_{\zeta}^{\Gamma}$ be the Schwartz kernels of S_{ζ} and the Γ -invariant lifts of the Schwartz kernel of $S_{i\zeta\|\alpha_0\|}^{\Gamma}$, respectively. We consider all of these as meromorphically extended to \mathbb{C} .

The proof of [BO12, Lemma 11] establishes the existence of a scalar-valued meromorphic function $\mathbb{C} \ni \zeta \mapsto a_{\zeta}$ satisfying the following formulas.

$$\beta_{\nu(\zeta)}(r_{\zeta}|_{X \times X \setminus \text{diag}_X}) = a_{\zeta} s_{\zeta}|_{\partial X \times \partial X \setminus \text{diag}_{\partial X}} \quad (5.8)$$

$$\beta_{\nu(\zeta)}((\tilde{r}_{\zeta}^{\Gamma} - r_{\zeta})|_{X \times X \setminus (D_{\Gamma} \setminus \text{diag}_X)}) = a_{\zeta} (\tilde{s}_{\zeta}^{\Gamma} - s_{\zeta})|_{\Omega \times \Omega \setminus (D_{\Gamma}^{\Omega} \setminus \text{diag}_{\Omega})}, \quad (5.9)$$

where

$$D_{\Gamma}^{\Omega} := (\{e\} \times \Gamma) \cdot \text{diag}_{\Omega} = (\Gamma \times \{e\}) \cdot \text{diag}_{\Omega} \subset \Omega \times \Omega.$$

In other words, we also have

$$\beta_{\nu(\zeta)}(\tilde{r}_{\zeta}^{\Gamma}|_{X \times X \setminus D_{\Gamma}}) = a_{\zeta} \tilde{s}_{\zeta}^{\Gamma}|_{\Omega \times \Omega \setminus D_{\Gamma}^{\Omega}}.$$

Actually, [BO12, p. 140] does only yield Equation (5.9) on a neighborhood of $\Omega \times \Omega \setminus (D_{\Gamma}^{\Omega} \setminus \text{diag}_{\Omega})$ in the closure of $X \times X$ (intersected with $X \times X \setminus \text{diag}_X$). But this suffices as the boundary values only depend on the values on a neighborhood of the boundary.

Lemma 5.2 shows that we know the function a_ζ explicitly on $(\partial X \times \partial X) \setminus \text{diag}_{\partial X}$:

$$a_\zeta = \frac{i}{2 \|\alpha_0\| \zeta}. \quad (5.10)$$

We thus arrive at the result

$$-2i \|\alpha_0\| \zeta \beta_{\nu(\zeta)}(\tilde{r}_\zeta^\Gamma|_{X \times X \setminus D_\Gamma}) = \tilde{s}_\zeta^\Gamma|_{\Omega \times \Omega \setminus D_\Gamma^\Omega}. \quad (5.11)$$

The following lemma will help us to remove D_Γ and D_Γ^Ω from Equation (5.11).

Lemma 5.3. *Let M be a smooth Riemannian manifold and*

$$P : C^\infty(M) \rightarrow C^\infty(M)$$

an elliptic differential operator of order > 0 (i.e., non-constant). Suppose that the resolvent

$$(P - z)^{-1} : L^2(M) \rightarrow L^2(M)$$

is defined for all z in an open subset $U \subset \mathbb{C}$ and extends to a connected open subset $V \supset U$ of \mathbb{C} as a meromorphic family of operators

$$R(z) : C_c^\infty(M) \rightarrow \mathcal{D}'(M).$$

Let $r(z) \in \mathcal{D}'(M \times M)$ be the meromorphic family of distributions given by the Schwartz kernel of $R(z)$. Then a complex number $z_0 \in V$ is a pole of $r(z)$ if, and only if, it is a pole of the meromorphic family

$$r(z)|_{M \times M \setminus \text{diag}_M} \in \mathcal{D}'(M \times M \setminus \text{diag}_M)$$

given by the restriction of $r(z)$ to the complement of the diagonal in $M \times M$.

Proof. Denote by $P_{C_c^\infty} : C_c^\infty(M) \rightarrow C_c^\infty(M)$ and $P_{\mathcal{D}'} : \mathcal{D}'(M) \rightarrow \mathcal{D}'(M)$ the operators obtained from P by restriction and duality, respectively. Then for all z in the set U , where $R(z)$ is $(P - z)^{-1}$ equipped with a smaller domain and a larger codomain, we have

$$(P_{\mathcal{D}'} - z \text{Id}_{\mathcal{D}'(M)}) \circ R(z) = I, \quad (5.12)$$

$$R(z) \circ (P_{C_c^\infty} - z \text{Id}_{C_c^\infty(M)}) = I, \quad (5.13)$$

where $I : C_c^\infty(M) \rightarrow \mathcal{D}'(M)$ is the inclusion. Now (5.12) and (5.13) are identities of holomorphic operator families on U which continue meromorphically to the connected open set $V \supset U$, so (5.12) and (5.13) also hold as identities of meromorphic operator families on V .

Let $z_0 \in V$. If z_0 is a pole of $r(z)|_{M \times M \setminus \text{diag}_M}$, then it clearly is also a pole of $r(z)$. Conversely, assume that z_0 is a pole of $r(z)$. This means that for all $z \neq z_0$ in a neighborhood U_0 of z_0 in V we have a unique presentation of $r(z)$ as a Laurent series

$$r(z) = \sum_{k \geq -k_0} (z - z_0)^k a_k \quad (5.14)$$

with distributions $a_k \in \mathcal{D}'(M \times M)$ and $k_0 \in \mathbb{N} = \{1, 2, \dots\}$ with $a_{-k_0} \neq 0$. Now suppose, in order to provoke a contradiction, that z_0 is not a pole of $r(z)|_{M \times M \setminus \text{diag}_M}$. This means that for all z in a neighborhood U_1 of z_0 in V we have a unique presentation

$$r(z)|_{M \times M \setminus \text{diag}_M} = \sum_{k \geq 0} (z - z_0)^k b_k \quad (5.15)$$

with distributions $b_k \in \mathcal{D}'(M \times M \setminus \text{diag}_M)$. By restricting $r(z)$ to $M \times M \setminus \text{diag}_M$ for $z \in U_0 \cap U_1$, the uniqueness of the presentation (5.15) implies that

$$a_k|_{M \times M \setminus \text{diag}_M} = 0 \quad \forall k < 0. \quad (5.16)$$

For each $k \geq -k_0$, let $A_k := C_c^\infty(M) \rightarrow \mathcal{D}'(M)$ be the operator with Schwartz kernel a_k . Then (5.14) implies that we have a convergent series

$$R(z) = \sum_{k \geq -k_0} (z - z_0)^k A_k \quad (5.17)$$

for all $z \in U_0$, and (5.16) implies that A_k is a local operator for all $k < 0$: If $f \in C_c^\infty(M)$ and $\chi \in C_c^\infty(M \setminus \text{supp } f)$, then (5.16) gives

$$A_k(f)(\chi) = a_k(f \otimes \chi) = 0$$

since $\text{supp}(f \otimes \chi) \subset M \times M \setminus \text{diag}_M$. Thus $\text{supp } A_k(f) \subset \text{supp } f$.

By (5.17) we can extract the operator A_{-k_0} from the meromorphic family $R(z)$ as

$$A_{-k_0} = \lim_{\substack{z \rightarrow z_0 \\ z \neq z_0}} (z - z_0)^{k_0} R(z).$$

On the other hand, by (5.12), we have

$$\begin{aligned} (P_{\mathcal{D}'} - z \text{Id}_{\mathcal{D}'(M)}) \circ (z - z_0)^{k_0} R(z) &= (z - z_0)^{k_0} (P_{\mathcal{D}'} - z \text{Id}_{\mathcal{D}'(M)}) \circ R(z) \\ &= (z - z_0)^{k_0} I \\ &\xrightarrow{z \rightarrow z_0} 0. \end{aligned}$$

Using the equicontinuity of the operator family $P_{\mathcal{D}'} - z \text{Id}_{\mathcal{D}'(M)}$ at $z = z_0$ we deduce that

$$(P_{\mathcal{D}'} - z_0 \text{Id}_{\mathcal{D}'(M)}) \circ A_{-k_0} = 0, \quad (5.18)$$

i.e.,

$$\text{im}(A_{-k_0}) \subset \ker(P_{\mathcal{D}'} - z_0 \text{Id}_{\mathcal{D}'(M)}). \quad (5.19)$$

Now P is elliptic, so (5.19) gives $\text{im}(A_{-k_0}) \subset C^\infty(M)$ by elliptic regularity. We already know that A_{-k_0} is a local operator, so by Peetre's Theorem we conclude that A_{-k_0} is a differential operator. In particular, $\text{im}(A_{-k_0}) \subset C_c^\infty(M)$. Knowing this, (5.18) implies that

$$(P_{C_c^\infty} - z_0 \text{Id}_{C_c^\infty(M)}) \circ A_{-k_0} = 0. \quad (5.20)$$

Now, around each $x \in M$ we can choose a neighborhood W of x in M such that the differential operator A_{-k_0} has a constant order N on W . Let $\sigma_P, \sigma_{A_{-k_0}} \in C^\infty(T^*M)$ be the principal symbols of P and A_{-k_0} , respectively. By the ellipticity of P we have $\sigma_P(x, \xi) \neq 0$

for all $\xi \neq 0$. Since P has order > 0 it follows that $\sigma_P(x, \xi) - z_0 \neq 0$ for all ξ with $\|\xi\|$ sufficiently large. From the composition theorem and (5.20) we get

$$(\sigma_P(x, \xi) - z_0)\sigma_{A_{-k_0}} = 0,$$

which implies that $\sigma_{A_{-k_0}}(x, \xi) = 0$ for all ξ with $\|\xi\|$ sufficiently large and consequently $\sigma_{A_{-k_0}}(x, \xi) = 0$ for all ξ because it is a polynomial. Thus the order N component of A_{-k_0} vanishes at x , which implies that $A_{-k_0}|_W = 0$. Since $x \in M$ was arbitrary, we conclude that $A_{-k_0} = 0$ and hence $a_{-k_0} = 0$, a contradiction. \square

We can now prove our main result:

Theorem 5.4. *Outside the set $-\frac{i}{2}\|\alpha_0\|\mathbb{N}$, the scattering poles which are not in the physical plane $\mathcal{H}_{\text{phys}}^\Gamma$ are precisely the quantum resonances.*

Proof. Let ζ_0 be a quantum resonance of $\Gamma \backslash X$, i.e. a pole of \tilde{r}_ζ^Γ . Then ζ_0 is *not* in the physical halfplane $\mathcal{H}_{\text{phys}}^\Gamma$ and Lemma 5.3 implies that ζ_0 is also a pole of $\tilde{r}_\zeta^\Gamma|_{X \times X \backslash D_\Gamma}$.

Case 1 $\zeta_0 \notin -\frac{i}{2}\|\alpha_0\|\mathbb{N}$ and $c(i\zeta_0 \frac{\alpha_0}{\|\alpha_0\|}) \neq 0$: In this case Equation (5.2) and the result [HHP19, Thm. 6.4] imply that $\zeta \mapsto \beta_{\nu(\zeta)}$ is a holomorphic family of injective operators in a neighborhood of ζ_0 . Thus by Equation (5.11) ζ_0 is a pole of $\tilde{s}_\zeta^\Gamma|_{\Omega \times \Omega \backslash D_\Gamma^\Omega}$. Hence ζ_0 is also a pole of \tilde{s}_ζ^Γ , i.e. a scattering pole of $\Gamma \backslash X$.

Case 2 $\zeta_0 \notin -\frac{i}{2}\|\alpha_0\|\mathbb{N}$ and $c(i\zeta_0 \frac{\alpha_0}{\|\alpha_0\|}) = 0$: In view of [HHP19, Re. 5.1] $c(i\zeta_0 \frac{\alpha_0}{\|\alpha_0\|}) = 0$ implies $\text{Im } \zeta_0 > \|\rho\|$. If $\delta_\Gamma < 2\|\rho\|$, then $i\rho \in \mathcal{H}_{\text{phys}}^\Gamma$, hence also $\zeta_0 \in \mathcal{H}_{\text{phys}}^\Gamma$. Thus, this case cannot occur, i.e. $\delta_\Gamma = 2\|\rho\|$. But then 0 is in the L^2 -spectrum of $\Delta_{\Gamma \backslash X}$. Since the continuous L^2 -spectrum of $\Delta_{\Gamma \backslash X}$ is $[\|\rho\|^2, \infty[$ the point 0 has to be an eigenvalue. But $\Delta_{\Gamma \backslash X}f = 0$ for $f \in L^2(\Gamma \backslash X)$ is equivalent to $df = 0$ and $d^*f = 0$, so f has to be constant. But this implies that $\Gamma \backslash X$ has finite volume, which contradicts our hypothesis the Γ is convex-cocompact, but not cocompact.

Conversely, assume that ζ_0 is a scattering pole which is not in the physical halfplane $\mathcal{H}_{\text{phys}}^\Gamma$ and satisfies $\zeta_0 \notin -\frac{i}{2}\|\alpha_0\|\mathbb{N}$. Then, by the argument given in Case 2, $c(i\zeta_0 \frac{\alpha_0}{\|\alpha_0\|}) \neq 0$ and again Equation (5.2) and [HHP19, Thm. 6.4] imply that $\zeta \mapsto \beta_{\nu(\zeta)}$ is a holomorphic family of injective operators in a neighborhood of ζ_0 .

For any $\lambda \in \mathfrak{a}_\mathbb{C}^*$ and $f \in C^{-\infty}(\Gamma \backslash \Omega, V_{\Gamma \backslash \Omega}(\sigma_\lambda))$ [BO12, p. 134] yields an Eisenstein series, which we denote by $\mathcal{P}_\lambda^\Gamma f$ since it takes the role of the Poisson transform in the case of locally symmetric spaces. Then $\mathcal{P}_\lambda^\Gamma f \in C^\infty(\Gamma \backslash X)$ is a $\Delta_{\Gamma \backslash X}$ -eigenfunction, which has f as boundary value. Applying this to the product $\Gamma \backslash \Omega \times \Gamma \backslash \Omega$ and scattering kernel \tilde{s}_ζ^Γ with the appropriate λ determined by ζ we find a kernel function $r_\zeta^{\Gamma, \text{Eis}}$ on $\Gamma \backslash X \times \Gamma \backslash X$ which has the same boundary values as r_ζ^Γ . But then, since $\zeta \mapsto \beta_{\nu(\zeta)}$ in a neighborhood of ζ_0 is a holomorphic family of injective maps, we have $r_\zeta^{\Gamma, \text{Eis}} = r_\zeta^\Gamma$ and ζ_0 is a pole of r_ζ^Γ , i.e. it is also a quantum resonance. \square

REFERENCES

- [AZ22] J.-Ph. Anker and H.-W. Zhang, *Bottom of the L^2 spectrum of the Laplacian on locally symmetric spaces*, Geom. Dedicata **216** (2022), no. 1, Paper No. 3, 12.
- [BO00] U. Bunke and M. Olbrich, *The spectrum of Kleinian manifolds*, J. Funct. Anal. **172** (2000), no. 1, 76–164.
- [BO12] ———, *Towards the trace formula for convex-cocompact groups*, Contributions in analytic and algebraic number theory, Springer Proc. Math., vol. 9, Springer, New York, 2012, pp. 97–148.
- [Cor90] K. Corlette, *Hausdorff dimensions of limit sets. I*, Invent. Math. **102** (1990), no. 3, 521–541.
- [HC76] Harish-Chandra, *Harmonic analysis on real reductive groups. III. The Maass-Selberg relations and the Plancherel formula*, Ann. of Math. (2) **104** (1976), no. 1, 117–201.
- [HHP19] S. Hansen, J. Hilgert, and A. Parthasarathy, *Resonances and scattering poles in symmetric spaces of rank one*, Int. Math. Res. Not. IMRN (2019), no. 20, 6362–6389.
- [Hil24] J. Hilgert, *Quantum-classical correspondences for locally symmetric spaces*, Symmetry in Geometry and Analysis — Festschrift for Toshiyuki Kobayashi, Birkhäuser, 2024.
- [Hör90] L. Hörmander, *The analysis of linear partial differential operators. I*, second ed., Grundlehren der mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences] **256**, Springer-Verlag, Berlin, 1990, Distribution theory and Fourier analysis.
- [Kap23] M. Kapovich, *A note on properly discontinuous actions*, São Paulo Journal of Mathematical Sciences (2023).
- [Kna86] A. W. Knap, *Representation theory of semisimple groups*, Princeton Mathematical Series, vol. 36, Princeton University Press, Princeton, NJ, 1986, An overview based on examples.
- [KO77] M. Kashiwara and T. Ōshima, *Systems of differential equations with regular singularities and their boundary value problems*, Ann. of Math. (2) **106** (1977), no. 1, 145–200.
- [KS71] A. W. Knap and E. M. Stein, *Intertwining operators for semisimple groups*, Ann. of Math. (2) **93** (1971), 489–578.
- [Olb94] M. Olbrich, *Die Poisson-Transformation für homogene Vektorbündel*, Ph.D. thesis, HU Berlin, 1994.
- [OS80] T. Ōshima and J. Sekiguchi, *Eigenspaces of invariant differential operators on an affine symmetric space*, Invent. Math. **57** (1980), no. 1, 1–81.
- [vdBS87] E. P. van den Ban and H. Schlichtkrull, *Asymptotic expansions and boundary values of eigenfunctions on Riemannian symmetric spaces*, J. Reine Angew. Math. **380** (1987), 108–165.
- [Vog08] D. A. Vogan, *Unitary representations and complex analysis*, Representation theory and complex analysis, Lecture Notes in Math. **1931**, Springer, Berlin, 2008, pp. 259–344.
- [Wal92] N. R. Wallach, *Real reductive groups II*, Pure and Applied Mathematics, vol. 132, Academic Press, Inc., Boston, MA, 1992.
- [Web08] A. Weber, *Heat kernel bounds, Poincaré series, and L^2 spectrum for locally symmetric spaces*, Bull. Aust. Math. Soc. **78** (2008), no. 1, 73–86.

Email address: bdelarue@math.upb.de

UNIVERSITÄT PADERBORN, WARBURGER STR. 100, 33098 PADERBORN, GERMANY

Email address: hilgert@math.upb.de

UNIVERSITÄT PADERBORN, WARBURGER STR. 100, 33098 PADERBORN, GERMANY