

Tensor-force effects on nuclear matter in relativistic *ab initio* theory

Sibo Wang,¹ Hui Tong,² Chencan Wang,³ Qiang Zhao,⁴ Peter Ring,⁵ and Jie Meng^{6,7,*}

¹*Department of Physics, Chongqing Key Laboratory for Strongly Coupled Physics, Chongqing University, Chongqing 401331, China*

²*Helmholtz-Institut für Strahlen- und Kernphysik and Bethe Center for Theoretical Physics, Universität Bonn, D-53115 Bonn, Germany*

³*School of Physics and astronomy, Sun Yat-Sen University, Zhuhai 519082, China*

⁴*Center for Exotic Nuclear Studies, Institute for Basic Science, Daejeon 34126, Korea*

⁵*Physik-Department der Technischen Universität München, D-85748 Garching, Germany*

⁶*State Key Laboratory of Nuclear Physics and Technology, School of Physics, Peking University, Beijing 100871, China*

⁷*Yukawa Institute for Theoretical Physics, Kyoto University, Kyoto 606-8502, Japan*

(Dated: March 22, 2024)

Within the relativistic Brueckner-Hartree-Fock theory in the full Dirac space, the tensor-force effects on infinite nuclear matter are elucidated by subtracting the matrix elements of tensor forces from the realistic nucleon-nucleon interaction. The tensor-force effects for the binding energy per particle of symmetric nuclear matter (SNM) as well as the symmetry energy are attractive and are more pronounced around the empirical saturation density, while the tensor forces have little impact on the pure neutron matter. By tuning the tensor-force strength, an infinite (negative) scattering length in the spin-triplet channel is found. This locates the dilute SNM with only the 3S_1 - 3D_1 channel interaction at the unitary limit. Its ground-state energy is found proportional to the energy of a free Fermi gas with a scaling factor 0.38, revealing good universal properties. This work paves the way to study the tensor-force effects in neutron stars as well as finite nuclei from realistic nucleon-nucleon interactions and highlights the role of the tensor force on the deviation of the nuclear physics to the unitary limit.

As a crucial ingredient of the nucleon-nucleon (NN) interaction, the tensor force has an important impact on the structural and dynamical properties of the nuclear many-body systems, particularly the properties of exotic nuclei far away from the stability, which are crucial for understanding the nucleosynthesis in nuclear astrophysics. Many efforts have been devoted to studying the influence of the tensor force in effective NN interaction in nuclear medium within the configuration interaction shell model [1] and the density functional theories (DFTs) [2]. Due to the difference of modeling the effective NN interaction and the difficulty of determining the tensor-force strengths in finite nuclei, the tensor force effects in nuclear physics are not fully settled.

Starting from the realistic NN interactions that are well calibrated in free space, the microscopic *ab initio* studies provide a more promising way to study the tensor-force effects, since no free parameters are introduced throughout the solution of the many-body methods. Progress has been made on the role of tensor force in the binding of light nuclei [3] and the symmetry energy of nuclear matter [4]. Still, insufficient attention has been paid on the studies of tensor-force effects from microscopic *ab initio* calculations.

The relativistic Brueckner-Hartree-Fock (RBHF) theory plays a key role in addressing the challenge of *ab initio* methods rooted in a relativistic framework [6]. The strong correlation in many-body wave function induced

by the realistic NN interaction V is absorbed into the effective G matrix, which is obtained as the solution of the Thompson equation in medium [7]. In addition to the advantages by solving nuclear many-body systems within a relativistic framework [8], the main strength of the RBHF theory lies in its inherent ability to account for a class of three-body forces which are important for satisfactorily reproducing the empirical saturation properties of nuclear matter [7, 9]. The RBHF theory has been successfully applied to describe the properties of nuclear matter and finite nuclei [10, 11].

In the light of the remarkable reliability of the relativistic *ab initio* theory, the many-body predictions from RBHF theory have been utilized as pseudo-data to constrain the parameters in both non-relativistic and relativistic DFTs. By applying the RBHF theory in neutron drops, a systematic and specific pattern in the evolution of spin-orbit (SO) splittings is observed [12, 13], which is found to have strong tensor-force effects. This provides a novel scheme for the determination of the tensor-force strengths and forms an important guide for the microscopic derivations of nuclear DFTs [14, 15].

In this work, we aim to directly study the tensor-force effects in nuclear many-body systems from relativistic *ab initio* theory. As a first step we focus on the infinite nuclear matter, which provides an ideal laboratory accessible to different methods that allows detailed benchmark. The RBHF theory in the full Dirac space




FIG. 1. (Color online) The binding energies per particle for (a) SNM and (b) PNM calculated with (solid lines and full symbols) and without (dotted lines and empty symbols) tensor forces. (c) The symmetry energy difference obtained by performing the RBHF calculations with and without tensor force. The results from Ref. [5] are also shown for comparison. Details can be found in the text.

is adopted, which considers simultaneously the positive- and negative-energy solutions (NESs) of the Dirac equation, thus avoids the uncertainties suffered from the RBHF calculations without NESs. Details can be found in Refs. [10, 16]. To identify the effects of tensor forces, we perform the RBHF calculations, in which the tensor forces in different meson-nucleon couplings are removed from the realistic NN interaction by using the non-relativistic reduced operators [17]

$$\hat{V}_\phi^t = \frac{1}{m_\phi^2 + \mathbf{q}^2} \mathcal{F}_\phi S_{12}. \quad (1)$$

Here ϕ denotes the exchanged meson with mass m_ϕ , $\mathbf{q}$ is the momentum transfer, and S_{12} is the tensor force operator

$$S_{12} \equiv (\boldsymbol{\sigma}_1 \cdot \mathbf{q})(\boldsymbol{\sigma}_2 \cdot \mathbf{q}) - \frac{1}{3}(\boldsymbol{\sigma}_1 \cdot \boldsymbol{\sigma}_2)q^2. \quad (2)$$

The factor $\mathcal{F}_\phi$ in each meson-nucleon coupling can be found in Table 1 of Ref. [17]. This scheme has been initially applied in analyzing the tensor-force effects on the neutron-proton effective mass splitting in neutron-rich matter [18]. To study the tensor-force effects more precisely, a quenching factor λ is introduced in Eq. (1) for each coupling channel $\mathcal{F}_\phi$ to control the strength of the tensor force. The factor $\lambda = 0$ means no tensor force, while $\lambda = 1$ means the tensor forces are considered fully as original.

Figure 1(a) shows the binding energies per particle as a function of the density ρ for symmetric nuclear matter (SNM) calculated within the RBHF theory with and without tensor forces. Clearly the tensor-force effects lead to attraction over a wide range of densities and are more significant at lower densities. The characteristics of attraction can be understood from the Thompson equation, which can be written simply in the operator form

$G = V + V(Q/e)V + \dots$. Here Q and e denote the Pauli operator and the energy denominator, respectively. The tensor force in the realistic interaction V makes no contribution to the binding energy of nuclear matter due to the spin saturation. For the second term $V(Q/e)V$, due to the positive definiteness of quadratic terms of V and the negative energy denominator in the presence of the Pauli operator, the contributions of the tensor force in the second term are always attractive. This implies that the contribution of tensor forces in realistic NN interaction to the nuclear matter binding is reflected through higher-order effects. In addition, among the three parameterizations of the Bonn potential [7], the tensor-force effects of Bonn C are the most pronounced. This is consistent with the well known fact that Bonn C has the strongest tensor force [7], as indicated by the largest D -state probability of the deuteron.

In Fig. 1(b), we illustrate the tensor-force effects on the binding energies per particle in pure neutron matter (PNM) where the total isospin T for two-nucleon state must equal to 1. For the Bonn A potential, the calculations with and without tensor force lead to nearly the same results. This indicates that the total tensor-force effects for Bonn A in isospin triplet with $T = 1$ are marginal. Similar conclusions can be drawn for Bonn B and Bonn C potentials. This can be understood by the fact that the PNM is dominated by 1S_0 partial wave, while the tensor-force contribution appears in P and higher waves.

The results in SNM and PNM allow us to derive the nuclear symmetry energy $E_{\text{sym}}(\rho)$, which is one of the cutting-edge topics in nuclear physics and astrophysics. We depict the differences between the symmetry energies obtained with and without tensor forces in Fig. 1(c). The tensor force plays an essential role around the empirical saturation density $\rho_0 = 0.16 \pm 0.01$ fm $^{-3}$, beyond which




FIG. 2. (Color online) (a) The dimensionless ratio r_t/a_t in spin-triplet (left axis) and the binding energy for the deuteron B_d (right axis) with variation of the tensor force quenching factor λ . The upper (lower) horizontal dashed line denotes the locale of the unitary (bound) limit. (b) The ground-state energy per particle for SNM calculated with Bonn A, B, and C, where the scattering length a_t in spin-triplet channel is tuned to -100 fm. Only contributions from the 3S_1 - 3D_1 channel are considered. The results of a free Fermi gas with a scaling factor 0.38 is also displayed for comparison.

the tensor-force effects become weaker. In comparison to our results, the symmetry energy difference obtained in Ref. [5] is very similar, where the tensor-force effects are analyzed by performing the RBHF calculations without the off-diagonal NN matrix elements in the coupled spin triplet channels. By using the Hellmann-Feynman theorem from non-relativistic *ab initio* method [4], the estimation of the tensor-force contribution to symmetry energy is qualitatively consistent (36 MeV at $\rho = 0.187 \text{ fm}^{-3}$), despite the quantitative discrepancy which may result from relativistic effects in NN interaction and many-body method.

It should be noted that, starting from the effective NN interaction within the DFTs, where only the tree-level interactions are considered, tensor force plays no role in nuclear matter due to the spin saturation. This clearly reveals the advantages from *ab initio* studies that the higher-order effects of the tensor force on nuclear matter can be explored.

The nuclear physics is naturally in proximity to the unitary limit where the two-body scattering length is infinite. This can be recognized by the fact that the scattering lengths in 1S_0 (spin singlet) and 3S_1 (spin triplet) are appreciably large compared to the effective range of the interaction. The system at the unitary limit exhibits universal properties regardless of the nature of the constituent particles and their mutual interactions. In particular, the ground state energy of any fermionic system at the unitary limit is expected to be proportional to the corresponding ones of a non-interacting Fermi gas with the scaling factor called Bertsch parameter [19].

Many efforts have been devoted to studying the relation of the dilute PNM properties with the unitary limit. At low densities, the neutron-neutron (nn) interaction is dominated by the 1S_0 partial wave with the scattering

length $a_s^{nn} = -18.9$ fm. Therefore, the dilute PNM is similar to a unitary Fermi gas, which is an idealized system of spin-1/2 fermions at the unitary limit. It is found that the Bertsch parameter is around 0.3-0.7 [20].

Different to the PNM, the SNM incorporates additionally the spin triplet 3S_1 partial wave, where the scattering length $a_t = 5.4$ fm implying the existence of a bound state for the neutron-proton (np) system, i.e., the deuteron. It is well known that the tensor force plays an important role in the binding energy of the deuteron and leads to the coupling between 3S_1 and 3D_1 partial waves. In Fig. 2(a) we present the ratio of the effective range r_t and the scattering length a_t as a function of the strength of the tensor force. With the continuously decreasing of the quenching factor λ of the tensor force, the ratio r_t/a_t also decreases. Note that the running of r_t/a_t is dominated by the scattering length a_t , while r_t retains in the range of 1.8-5.1 fm. At a certain λ , the scattering length a_t goes to infinity, locating the np system at the unitary limit, which is denoted by the upper horizontal dashed line. This also illustrates that the strong tensor force in the np interaction makes the np system, to some extent, deviate from the unitary limit.

We also show the influence of the tensor force on the binding energy of deuteron in Fig. 2(a). With the decrease of the factor λ , the binding energy of the deuteron B_d also decreases from 2.2246 MeV to zero until meeting the unitary limit. The rapidest case is found for Bonn C, since this potential has the strongest tensor force. The deuteron is not bound any more after the dimensionless ratio r_t/a_t crosses the locale of the unitary limit and the bound limit is denoted by the lower horizontal dashed line.

It is interesting to study how the dilute SNM with only the coupled spin triplet 3S_1 - 3D_1 partial waves be-

haves at the unitary limit. By tuning the quenching factor of the tensor force $\lambda = 0.4029, 0.5624$, and 0.6278 for Bonn A, B, C, respectively, we obtain the scattering length $a_t \approx -100$ fm, which is large enough compared to the effective range r_t and the average distance between nucleons in the dilute SNM. From the modified Bonn potentials near the unitary limit, we calculate the ground-state energies per particle for SNM by only considering the NN matrix elements in the 3S_1 - 3D_1 channel. To avoid the numerical instabilities at extreme low densities, the Thompson equations are solved only once with initial vanishing single-particle potentials. This should be reasonable since the hypothetically converged single-particle potentials are very close to zero at extreme low densities.

In Fig. 2(b), we present the ground-state energy per particle of SNM within the scheme discussed above. The results from Bonn A, B, and C show very good consistency, not sensitive to the details of the interaction. Furthermore, it is found that the results from our calculations can be well simulated with $\xi(E/A)_{\text{free gas}}$, where $(E/A)_{\text{free gas}} = (3k_F^2)/(10M)$ is the energy per particle of the free Fermi gas and the $\xi = 0.38$ is the aforementioned Bertsch parameter. We also calculate the parameter ξ for dilute PNM and SNM in the 1S_0 channel, where the scattering lengths are $a_s^{nn} = -18.9$ fm and $a_s^{np} = -23.7$ fm. The corresponding results are $\xi = 0.60$ and 0.42 , implying the convergence of the Bertsch parameter towards the unitary limit. Besides, the value 0.38 is close to the result 0.372 from quantum Monte Carlo calculations [21] for unitary Fermi gas, and 0.376 measured in ultra cold atoms [22]. With increasing density, the differences among the three potentials and the deviations with respect to the $0.38(E/A)_{\text{free gas}}$ become evident, which means that the universal properties of SNM are getting deteriorated.

In summary, the tensor-force effects on the equation of state of nuclear matter and the symmetry energy have been studied from the realistic Bonn potentials, by invoking the relativistic Brueckner-Hartree-Fock theory with the tensor force matrix elements subtracted. For the binding energies per particle of symmetric nuclear matter (SNM) and the symmetry energy, the tensor-force effects are attractive and are more pronounced around the empirical saturation density. For pure neutron matter, the tensor-force effects are marginal. The impact of the tensor force on the deviation of the np system to the unitary limit has also been studied. By tuning the tensor-force strength, an infinite (negative) scattering length in the spin triplet can be obtained and the dilute SNM is located at the unitary limit. With only the interaction in 3S_1 - 3D_1 channel considered, the ground-state energy of dilute SNM is found proportional to that of a free Fermi gas with a scaling factor 0.38 , which reveals very good universal properties. This work paves the way to study the tensor-force effects in neutron stars as well as

finite nuclei from realistic nucleon-nucleon interactions and highlights the role of the tensor force on the deviation of the nuclear physics to the unitary limit.

S. Wang and Q. Zhao thank S. Aoki for helpful discussions. This work was supported in part by the National Natural Science Foundation of China (NSFC) under Grants No. 12205030, No. 12347101, No. 11935003, No. 11975031, No. 11875075, No. 12070131001 and No. 12047564, the National Key R&D Program of China under Contracts No. 2017YFE0116700 and No. 2018YFA0404400, the Fundamental Research Funds for the Central Universities under Grants No. 2020CDJQY-Z003 and No. 2021CDJZYJH-003, the MOST-RIKEN Joint Project "Ab initio investigation in nuclear physics", the Institute for Basic Science under Grant No. IBS-R031-D1, and the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) under Germany's Excellence Strategy EXC-2094-390783311, ORIGINS. Part of this work was achieved by using the supercomputer OCTOPUS at the Cybermedia Center, Osaka University under the support of Research Center for Nuclear Physics of Osaka University and the High Performance Computing Resources in the Research Solution Center, Institute for Basic Science.

* mengj@pku.edu.cn

- [1] T. Otsuka, A. Gade, O. Sorlin, T. Suzuki, and Y. Utsuno, *Rev. Mod. Phys.* **92**, 015002 (2020).
- [2] H. Sagawa and G. Colò, *Progress in Particle and Nuclear Physics* **76**, 76 (2014).
- [3] B. S. Pudliner, V. R. Pandharipande, J. Carlson, S. C. Pieper, and R. B. Wiringa, *Phys. Rev. C* **56**, 1720 (1997).
- [4] I. Vidaña, A. Polls, and C. m. c. Providência, *Phys. Rev. C* **84**, 062801 (2011).
- [5] C. Wang, J. Hu, Y. Zhang, and H. Shen, *Chinese Physics C* **46**, 064108 (2022).
- [6] S. Shen, H. Liang, W. H. Long, J. Meng, and P. Ring, *Progress in Particle and Nuclear Physics*, 103713 (2019).
- [7] R. Brockmann and R. Machleidt, *Phys. Rev. C* **42**, 1965 (1990).
- [8] P. Ring, *Physica Scripta* **2012**, 014035 (2012).
- [9] G. E. Brown, W. Weise, G. Baym, and J. Speth, *Comm. Nucl. Part. Phys.* **17**, 39 (1987).
- [10] S. Wang, Q. Zhao, P. Ring, and J. Meng, *Phys. Rev. C* **103**, 054319 (2021).
- [11] S. Shen, H. Liang, J. Meng, P. Ring, and S. Zhang, *Phys. Rev. C* **96**, 014316 (2017).
- [12] S. Shen, H. Liang, J. Meng, P. Ring, and S. Zhang, *Phys. Lett. B* **778**, 344 (2018).
- [13] S. Wang, H. Tong, P. Zhao, and J. Meng, *Phys. Rev. C* **100**, 064319 (2019).
- [14] S. Shen, G. Colò, and X. Roca-Maza, *Phys. Rev. C* **99**, 034322 (2019).
- [15] Q. Zhao, P. Zhao, and J. Meng, *Phys. Rev. C* **102**, 034322 (2020).
- [16] S. Wang, H. Tong, Q. Zhao, C. Wang, P. Ring, and

- J. Meng, Phys. Rev. C **106**, L021305 (2022).
- [17] Z. Wang, Q. Zhao, H. Liang, and W. H. Long, Phys. Rev. C **98**, 034313 (2018).
- [18] S. Wang, H. Tong, Q. Zhao, C. Wang, P. Ring, and J. Meng, Phys. Rev. C **108**, L031303 (2023).
- [19] G. A. Baker, Phys. Rev. C **60**, 054311 (1999).
- [20] I. Vidaña, Frontiers in Physics **9** (2021), 10.3389/fphy.2021.660622.
- [21] J. Carlson, S. Gandolfi, K. E. Schmidt, and S. Zhang, Phys. Rev. A **84**, 061602 (2011).
- [22] M. J. H. Ku, A. T. Sommer, L. W. Cheuk, and M. W. Zwierlein, Science **335**, 563 (2012).