

The Flavor Composition of Supernova Neutrinos

Antonio Capanema,^{1,*} Yago Porto,^{2,3,†} and Maria Manuela Saez^{4,5,‡}

¹*Departamento de Física, Pontifícia Universidade Católica do Rio de Janeiro, Rio de Janeiro 22452-970, Brazil*

²*Centro de Ciências Naturais e Humanas, Universidade Federal do ABC, 09210-170, Santo André, SP, Brazil*

³*Instituto de Física Gleb Wataghin, Universidade Estadual de Campinas, 13083-859, Campinas, SP, Brazil*

⁴*RIKEN Interdisciplinary Theoretical and Mathematical Sciences*

Program (iTHEMS), 2-1 Hirosawa, Wako, Saitama 351-0198, Japan

⁵*Department of Physics, University of California, Berkeley, CA 94720, USA*

We show that standard matter effects in the outer layers of core-collapse supernovae significantly constrain the flavor composition of the neutrino flux, even with the enormous uncertainties originating from self-induced flavor conversions in the supernova core. Under certain conditions, the neutrino flux resulting from self-induced conversions can be represented as a combination of flavor eigenstates in an arbitrary flavor ratio configuration. In this scenario, we find that, for the normal mass ordering, the fraction of neutrinos with electron flavor reaching the Earth, denoted as f_{ν_e} , is constrained to be less than 0.5 for all energies throughout the emission phase, whereas, for inverted mass ordering, we anticipate neutrinos arriving in near flavor equipartition ($f_{\nu_e} \approx 1/3$). In case adiabaticity is violated in the region of standard matter effects, the result is flavor equipartition for both mass orderings. Subsequently, we elaborate on the impact of wave-packet decoherence during self-induced conversions and explore alternative scenarios that could affect the aforementioned results.

Introduction.— During a core-collapse supernova (SN) event, the immense gravitational binding energy of the progenitor star is predominantly liberated through a rapid emission of neutrinos, typically lasting around 10 seconds [1, 2]. After being generated in the SN core, these neutrinos traverse the mantle and envelope, where they are thought to be pivotal in driving the explosion [3, 4].

In 1987, the detection of a few tens of SN neutrinos coming from the Large Magellanic Cloud confirmed the basic features of the core-collapse models, such as the mean energy and the duration of the neutrino burst [5–7]. Nevertheless, the intricacies of the SN physics remain a challenge, largely due to the uncertainties surrounding neutrino flavor transformations in the core [8–11]. In such a neutrino-dense environment, the coherent forward scattering of neutrinos onto each other gives rise to collective effects that drive a nonlinear flavor evolution [12–16]. These self-induced conversions are referred to as “slow” or “fast” depending on the characteristic spatial scale in which they manifest. Slow self-induced conversions develop over a length scale governed by the vacuum oscillation frequency, which is on the order of a few kilometers for typical SN neutrino energies (~ 10 MeV) [14, 15]. These conversions take place in an intermediate zone situated ~ 100 km away from the neutrino decoupling region, but before the standard matter-induced resonant neutrino conversion occurs [17–20]. Conversely, fast self-induced conversions have characteristic scales as small as centimeters and could significantly influence the flavor evolution, energy spectrum, and angular distribu-

tion of each neutrino species near the decoupling region [21–25]. The lack of a full solution to the quantum kinetic equations governing this behavior leads to unpredictable consequences for the SN dynamics [10, 11] and, ultimately, for the neutrino signal detected at the Earth [26].

Amidst this conundrum, it is natural to wonder if a meaningful improvement in our understanding of the core-collapse phenomenon and neutrino propagation will be possible, even after the observation of the next galactic SN. In this *Letter*, we show that many features of the neutrino flux coming to the Earth can be understood by adopting a completely agnostic approach to the flavor composition emerging from the SN core, where collective effects dominate. By considering all possible flavor ratio configurations before the onset of standard matter effects, we observe that matter effects alone impose significant constraints on the flavor composition of SN neutrinos. Assuming normal ordering (NO) of neutrino masses, we find that the fraction of electron neutrinos arriving at the Earth, denoted by $f_{\nu_e}^{NO}$, remains consistently below 0.5 for all energies during the emission phase, and gradually converges to near flavor equipartition ($f_{\nu_e}^{NO} \approx 1/3$)¹ during the passage of shock waves. In the inverted ordering (IO) scenario, we predict flavor equipartition ($f_{\nu_e}^{IO} \approx 1/3$) as a generic outcome, which remains robust throughout the entire neutrino burst, even in the presence of shock waves. We specify the circumstances in which the flux resulting from collective effects is accurately described as consisting of flavor eigenstates. In addition, we explore the impact of wave-packet

* E-mail: antoniogalvao@aluno.puc-rio.br

† E-mail: yago.porto@ufabc.edu.br

‡ E-mail: manuela.saez@berkeley.edu

¹ In this *Letter*, “flavor equipartition” simply means $f_{\nu_e} = 1/3$, since discrimination between ν_μ and ν_τ signals in current and next-generation detectors is challenging [27].




FIG. 1. Allowed regions for the flavor composition of supernova neutrinos arriving at the Earth for the cases of normal mass ordering (left) and inverted mass ordering (right). The colors represent three different scenarios. *Red*: The flux emerging from collective effects is composed of flavor eigenstates in an arbitrary flavor ratio configuration, and propagation in the outer layers is adiabatic, $f_{\nu_e}^{NO} \lesssim 0.5$ and $f_{\nu_e}^{IO} \approx 1/3$. *Maroon* (stripes): Flux emerging from collective effects in an arbitrary flavor ratio configuration, but non-adiabatic propagation through the H -resonance (presence of shock wave), $f_{\nu_e}^{NO} \approx f_{\nu_e}^{IO} \approx 1/3$. *Pink*: Flux of matter eigenstates in arbitrary amounts emerging from collective effects (for instance, due to wave packet decoherence), $f_{\nu_e}^{NO} \approx f_{\nu_e}^{IO} \lesssim 0.7$. In this figure, we consider the 3σ range of variability only for the parameters θ_{12} and θ_{13} , which affect the fraction of electron flavor f_{ν_e} . Analogous results for antineutrinos are presented in the Appendix.

decoherence in self-induced conversions, along with other possible scenarios, on our results. If confirmed by the next SN observation, these results will not only validate our fundamental understanding of the SN dynamics and neutrino conversion in the outermost layers, but also elucidate crucial properties of the neutrino fluxes emerging from the region of self-induced conversions.

Standard Matter Effects.— Neutrino-neutrino interactions are expected to play a subdominant role in flavor evolution after the first few hundred kilometers from the neutrino decoupling region [8, 15]. At the outermost layers, neutrino evolution is predominantly affected by the matter potential generated by the coherent forward scattering of neutrinos on electrons, protons and neutrons [17]. In practice, however, only the charged current interactions between electrons and electron (anti)neutrinos influence neutrino oscillations, as neutral current interactions with any of the particles in the medium affect all neutrino flavors equally and can be factored out in the evolution. Consequently, the effective Hamiltonian in the outer SN regions can be written in the flavor basis as follows

$$\mathcal{H}_\nu = \frac{1}{2E} U \begin{pmatrix} 0 & 0 & 0 \\ 0 & \Delta m_{21}^2 & 0 \\ 0 & 0 & \Delta m_{31}^2 \end{pmatrix} U^\dagger + \begin{pmatrix} V_e & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}. \quad (1)$$

Here, U represents the Pontecorvo–Maki–Nakagawa–

Sakata (PMNS) matrix, Δm_{ij}^2 stands for the mass-squared differences and E denotes the neutrino energy. The matter potential affects only the ν_e flavor and has the form $V_e = \sqrt{2}G_F\rho_m Y_e/m_N$, where G_F is the Fermi constant, Y_e is the electron number fraction, m_N is the nucleon mass and ρ_m is the local matter density. It is important to consider the implicit dependence of V_e and \mathcal{H}_ν on time and space, as each neutrino follows a different trajectory inside the SN, and the physical conditions along each trajectory change with time. For antineutrinos, the matter potential has the opposite sign, $\bar{V}_e = -V_e$. Thus, the equation of motion for neutrinos in the SN outer layers is given by

$$i \frac{d\rho_\nu}{dt} = [\mathcal{H}_\nu, \rho_\nu], \quad (2)$$

where ρ_ν is the density matrix describing the neutrino flavor state. Observe that Eq. 2 is a simplified version of the complete expression that takes into account neutrino-neutrino interactions, advection, and collisions [28, 29].

As neutrinos propagate towards vacuum and the matter density (ρ_m) gradually decreases, they encounter two resonance layers where effective flavor conversions can occur [18–20]. These so-called H - and L -resonances occur when the following conditions are satisfied:

$$\sqrt{2}G_F \frac{Y_e}{m_N} \rho_m^{\text{res}} = \frac{\Delta m_{i1}^2 \cos 2\theta_{1i}}{2E}, \quad (3)$$

where $\text{res} = H(L)$ corresponds to $i = 3(2)$. For efficient flavor conversion, adiabaticity must be satisfied at the resonance layers. The degree of adiabaticity can be estimated by considering the parameter

$$\gamma = \frac{\Delta m_{i1}^2 \sin^2 2\theta_{1i}}{2E \cos 2\theta_{1i}} \left(\frac{1}{Y_e \rho_m^{\text{res}}} \left| \frac{d(Y_e \rho_m)}{dr} \right|_{\text{res}} \right)^{-1}, \quad (4)$$

and propagation is adiabatic for $\gamma > 1$ [30].

At densities above ρ_m^H , V_e dominates in Eq. 1 and suppresses vacuum mixing. As a result, the ν_e effectively decouples from the other flavors and behaves as the heaviest eigenstate in matter, which corresponds to ν_3^m in NO and ν_2^m in IO, where the superscript m denotes matter eigenstates [30]. Conversely, the other flavors are equally affected by matter² and oscillate among themselves with frequency given by the vacuum terms and near maximal mixing ($\theta_{23} \approx 49^\circ$). Therefore, the eigenstates of the $\nu_\mu - \nu_\tau$ subspace at $\rho_m \gg \rho_m^H$ are given by approximately equal mixtures of ν_μ and ν_τ . In NO, these matter eigenstates are ν_1^m and ν_2^m , while in IO, they are ν_1^m and ν_3^m . Hence, the emergence of any quantities of ν_μ and ν_τ eigenstates from the region of self-induced flavor conversions suggests that, in the region of standard matter effects, two out of the three matter eigenstates will be present in roughly equal numbers.

Flavor eigenstates emerging from self-induced conversions.— Assuming an emission of 10^{58} neutrinos over the 10-second duration of a galactic SN burst [32], at a distance of 10 kpc, we estimate the flux reaching the Earth to be approximately 10^{15} neutrinos per square meter per second. Before traveling to the Earth, all these neutrinos reach the boundary between collective effects and standard matter effects inside the SN with a quantum state described by a 3×3 density matrix, where the diagonal elements in the flavor basis give the probability for the neutrino to be measured in one of the three flavors, while the off-diagonal elements indicate the ability of the neutrino to interfere with itself (quantum coherence) and generate flavor oscillations. The region of standard matter effects destroys any superposition of electron neutrinos with other flavors initially formed by collective effects, whereas the subspace $\nu_\mu - \nu_\tau$ retains its flavor superposition. As a result, the neutrino state either transitions to ν_e or to a superposition of ν_μ and ν_τ .

The density matrix corresponding to the $\nu_\mu - \nu_\tau$ subspace (sometimes referred to as ν_x) of the k -th neutrino exiting the collective effects region towards the Earth is given by

$$\rho_{\nu_x}^k = \begin{pmatrix} \alpha_k^2 & \alpha_k \beta_k e^{-i\phi_k} \\ \alpha_k \beta_k e^{i\phi_k} & \beta_k^2 \end{pmatrix}. \quad (5)$$

² To be precise, differences between ν_μ and ν_τ arise in matter at the loop level, but are negligible at densities below 10^7 g/cm^3 , where the MSW resonances occur [31].

Here, α_k , β_k , and ϕ_k are real numbers, subject to the condition $\alpha_k^2 + \beta_k^2 = 1$. Furthermore, k ranges from 1 to N , where N represents the number of neutrinos with a specific energy leaving the collective effects region towards the Earth and having collapsed to non-electron flavors. The density matrix for the ensemble of N neutrinos is

$$\rho_{\nu_x} = \sum_{k=1}^N \begin{pmatrix} \alpha_k^2 & \alpha_k \beta_k e^{-i\phi_k} \\ \alpha_k \beta_k e^{i\phi_k} & \beta_k^2 \end{pmatrix}. \quad (6)$$

Since N is very large, as mentioned at the beginning of this section, it is natural to assume that the off-diagonal elements of Eq. 6 average to zero, due to the many oscillatory terms composing the sum³. Note that α_k , β_k , and ϕ_k depend on the evolution of the k -th neutrino, which can be enormously diverse before entering the region of standard matter effects [34]. In such a situation, the superposition between ν_μ and ν_τ is effectively destroyed. Therefore, after collective effects, the neutrino flux density matrix becomes diagonal in the flavor basis and can be described as composed of the various flavor eigenstates in arbitrary proportions.

In what follows, the discussion is centered on neutrinos. For the results related to antineutrinos, see the Appendix.

Results: flavor ratios at the Earth.— In the SN region where neutrino-neutrino potentials are already subdominant and $\rho_m \gg \rho_m^H$, the mixing between the ν_e and the non-electron flavors is suppressed. As previously elucidated, ν_e can be approximated as ν_3^m (ν_2^m) for NO (IO). Meanwhile, inside the subspace $\nu_\mu - \nu_\tau$, the effective mixing is near-maximal and ν_μ and ν_τ are both approximately equal mixtures of the same matter eigenstates: ν_1^m and ν_2^m (ν_1^m and ν_3^m) for NO (IO). Consequently, even if the physical output of the central region of self-induced conversions are flavor eigenstates in random relative proportions, the effective mixing in the $\nu_\mu - \nu_\tau$ subspace ensures that two out of the three matter eigenstates consistently occur in approximately equal amounts.

Assuming adiabaticity of neutrino propagation, the relative proportions between the vacuum eigenstates that reach the Earth are identical to those of matter eigenstates propagating in the SN outer layers ($\nu_1^m \rightarrow \nu_1$, $\nu_2^m \rightarrow \nu_2$, and $\nu_3^m \rightarrow \nu_3$). In the scenario in which at least two of the matter eigenstates are produced in similar numbers, as described in the last paragraph, the possible range of flavor ratios at the Earth can be constrained to be $f_{\nu_e}^{\text{NO}} \lesssim 0.5$ and $f_{\nu_e}^{\text{IO}} \approx 1/3$. This result corresponds to the red-shaded region in Fig. 1. Notably,

³ For simplicity, the density matrix in Eq. 5 corresponds to a pure state, namely $|\nu_x\rangle = \alpha_k |\nu_\mu\rangle + \beta_k e^{i\phi_k} |\nu_\tau\rangle$. However, the same argument applies to states with an intermediate level of coherence, provided that the terms in the off-diagonal sum in Eq. 6 retain some degree of oscillatory behavior [33]. Later, we discuss situations where the assumption of vanishing off-diagonal terms is not valid.

the IO scenario indicates a near equipartition among flavors leaving the SN, irrespective of the flavor output from self-induced conversions.

To better understand the emergence of these constraints, we characterize the evolution within the zone of standard matter effects in terms of flavor ratios $(f_{\nu_e}, f_{\nu_\mu}, f_{\nu_\tau})$. Assuming NO and adiabatic propagation, it follows that an initially produced ν_e arrives at the Earth as ν_3 [35]:

$$(1, 0, 0)_{SN} \rightarrow \left(|U_{e3}|^2, |U_{\mu 3}|^2, |U_{\tau 3}|^2 \right)_\oplus. \quad (7)$$

Conversely, either of the non-electron flavors ν_μ or ν_τ arrive at the Earth approximately as equal mixtures of ν_1 and ν_2 . Therefore,

$$(0, 1, 0)_{SN} \quad \text{or} \quad (0, 0, 1)_{SN} \rightarrow \frac{1}{2} \left(|U_{e1}|^2 + |U_{e2}|^2, |U_{\mu 1}|^2 + |U_{\mu 2}|^2, |U_{\tau 1}|^2 + |U_{\tau 2}|^2 \right)_\oplus. \quad (8)$$

For any initial combination $(a, b, c)_{SN}$, we obtain on Earth the ν_e fraction

$$f_{\nu_e}^{NO} = a |U_{e3}|^2 + b \frac{|U_{e1}|^2 + |U_{e2}|^2}{2} + c \frac{|U_{e1}|^2 + |U_{e2}|^2}{2}. \quad (9)$$

Because of the unitarity of the PMNS matrix and $a + b + c = 1$, Eq. 9 simplifies to

$$f_{\nu_e}^{NO} = \frac{1}{2} \left(1 - |U_{e3}|^2 \right) + \frac{a}{2} \left(3 |U_{e3}|^2 - 1 \right), \quad (10)$$

Adopting $|U_{e3}|^2 \approx 0.02 \ll 1$ [36], we obtain

$$f_{\nu_e}^{NO} \approx \frac{(1-a)}{2} \lesssim 0.5. \quad (11)$$

In the IO scenario, the argument mirrors the one between Eq. 7 and Eq. 10, with the interchange of the PMNS matrix elements $U_{\alpha 3}$ and $U_{\alpha 2}$ for any flavor α . Thus, by replacing U_{e3} with U_{e2} in Eq. 10, we have

$$f_{\nu_e}^{IO} = \frac{1}{2} \left(1 - |U_{e2}|^2 \right) + \frac{a}{2} \left(3 |U_{e2}|^2 - 1 \right). \quad (12)$$

Assuming $|U_{e2}|^2 \approx 1/3$ [36], we obtain⁴

$$f_{\nu_e}^{IO} \approx \frac{1}{3}. \quad (13)$$

The red-shaded regions in Fig. 1 correspond to $f_{\nu_e}^{NO}$ and $f_{\nu_e}^{IO}$ after varying θ_{12} and θ_{13} within their 3σ allowed

range and using the best-fit values for the other oscillation parameters (we have verified that the impact of θ_{23} and δ_{CP} on the resulting f_{ν_e} range is less than 10%) [36]. The uncertainty in $f_{\nu_e}^{IO}$ arises predominantly from the uncertainty in the oscillation parameters, rather than the initial composition $(a, b, c)_{SN}$. As the precision in the determination of the oscillation parameters increases, these constraints should improve.

The results in Eq. 11 and Eq. 13 are applicable to neutrinos of any energy and at any instant in time (for the results related to antineutrinos, see the Appendix). Nonetheless, exceptions may be possible when a shock wave is present, leading to violations of Eq. 11 at certain parts of the energy spectrum.

Results in the presence of shock waves.— When adiabatic propagation cannot be guaranteed, transitions between different matter eigenstates may occur at the H - and L -resonance layers. Furthermore, if the matter profile is affected by shock waves, multiple resonances of the H - and L -types could appear [37, 38]. If an even number of H -resonances are non-adiabatic, results do not change⁵. However, if an odd number of the H -resonances are non-adiabatic, the outcome is modified in the NO scenario once transitions $\nu_3^m \leftrightarrow \nu_2^m$ that happen at one of the resonance layers are not reverted by another transition of the same type. It is straightforward to determine the consequence of an odd number of non-adiabatic H -resonances: transitions of the type $\nu_3^m \leftrightarrow \nu_2^m$ imply that matrix elements $U_{\alpha 3}$ and $U_{\alpha 2}$ should be interchanged in Eqs. 7-10. The argument is similar to the one that leads to Eqs. 12 and Eq. 13, resulting in

$$f_{\nu_e}^{NO}(\text{non-ad } H) \approx \frac{1}{3}. \quad (14)$$

Because the position of the H -resonance is energy dependent, the portion of the energy spectrum influenced by the transient non-adiabaticity provides information about the location of the shock wave front [37, 38]. Thus, we can track the motion of the shock wave through the H -resonance layers using flavor ratios if neutrino masses follow NO: during the shock's transit through that region, f_{ν_e} converges to 1/3 in the corresponding part of the energy spectrum.

In the IO scenario, the H -resonance does not occur in the neutrino channel. Therefore, the result is identical to the adiabatic case:

$$f_{\nu_e}^{IO}(\text{non-ad } H) \approx \frac{1}{3}. \quad (15)$$

The impact of the non-adiabatic H -resonance in neutrino propagation is illustrated by the maroon-striped region in Fig. 1. Once again, we vary only θ_{12} and θ_{13}

⁴ Eqs. 12 and 13 also have some curious implications in the context of solar neutrinos. Namely, they imply that any initial neutrino flavor configuration in the solar core leads to $f_{\nu_e} \approx 1/3$ at the Earth. This fact applies to both orderings, as ν_e is produced as ν_2^m and follows adiabatic propagation in the Sun's interior.

⁵ Here we are neglecting phase effects that develop between a pair of shock waves, as the observation of these effects should be challenging in realistic detectors [39].

within their 3σ uncertainties. We conclude that the presence of shock wave fronts in the H -resonances causes the neutrino system to converge to approximate flavor equipartition at the Earth for both mass orderings (for the results related to antineutrinos, see the Appendix).

Although shock wave fronts can also penetrate the L -resonance layer, adiabaticity is never strongly violated there [15]. Even in cases where such violations may happen, they typically occur at later times (beyond 10 seconds post-bounce) [38], so we do not consider them here.

Wave-packet decoherence during self-induced conversions.— All the derived results so far relied on the assumption that the off-diagonal elements in Eq. 6 are negligible. However, there are physical situations of interest in which Eq. 6 is not diagonal. For instance, the outcome of collective effects could be eigenstates in neutrino-neutrino potentials that adiabatically transform to matter eigenstates in the usual matter potentials. In such a situation, the density matrix for N neutrinos would be diagonal in the matter eigenstates basis, but not in the flavor basis [40]. The matter eigenstates ν_1^m , ν_2^m and ν_3^m would arise in any relative amounts with no correlations between any two of them, and results would differ drastically from previous sections.

Another interesting situation would happen if some unknown physical mechanism restricts the allowed values of all parameters in Eq. 6. In this case, the fractions of ν_1^m , ν_2^m and ν_3^m that enter the region of standard matter effects are strongly dependent on the specific values of α_k , β_k and ϕ_k . If the range of values for these parameters is unknown or varies significantly over time, the initial relative fractions of each matter eigenstate would also appear to be arbitrary.

Within the scenarios described in this section, the outcome of collective effects is better formulated in terms of matter eigenstates rather than flavor eigenstates, and it resembles the usual picture of wave-packet decoherence of neutrino oscillations in vacuum and matter [41]. Nevertheless, we can still constrain the flavor composition on Earth in terms of f_{ν_e} . By considering these eigenstates to leave the collective effects region and, subsequently, arrive at Earth in arbitrary relative proportions, we find that $f_{\nu_e} \lesssim 0.7$ for both mass orderings⁶. These constraints are depicted as pink-shaded regions in the flavor triangles of Fig. 1.

Discussion and Conclusion.— We have shown that the flavor composition of SN neutrinos reaching the Earth can be constrained, even when assuming complete uncertainty about the neutrino state that emerges from the innermost regions dominated by self-induced conversions. Under certain conditions, the neutrino flux can be

viewed as composed of flavor eigenstates in an arbitrary flavor ratio configuration (see Eq. 6 and related discussion). If this scenario occurs, we predict that the proportion of ν_e is always smaller than 0.5 for NO, whereas, for IO, we expect near flavor equipartition ($f_{\nu_e} \approx 1/3$). These results are valid for neutrinos throughout the energy spectrum and duration of the emission (for the results related to antineutrinos, see the Appendix). If the H -resonance is non-adiabatic, for example, due to shock waves, flavor equipartition is the expectation for both mass orderings. However, in the opposite case in which neutrinos come out from self-induced conversions as fluxes of matter eigenstates (for instance, due to wave-packet decoherence), we obtain $f_{\nu_e} \lesssim 0.7$.

Ultimately, this *Letter* advocates for a phenomenological approach to comprehensively address the intricacies of neutrino flavor conversion in SN environments. Experimental challenges along the way include distinguishing between ν_x and $\bar{\nu}_x$ in neutral current scatterings [43]. This approach can also be extended to accommodate non-standard scenarios, potentially unveiling novel phenomena underlying the physics of neutrino masses and mixing [44, 45].

Acknowledgments.— We thank Orlando Peres for useful discussions. AC thanks for the support received by the scholarships CAPES/PROEX No. 88887.511843/2020-00, CNPq No. 140316/2021-3, CAPES-PrInt No. 88887.717489/2022-00 and FAPERJ No. E-26/204.138/2022. The work of YP was supported by the São Paulo Research Foundation (FAPESP) Grant No. 2023/10734-3 and 2023/01467-1, and by the National Council for Scientific and Technological Development (CNPq) Grant No. 151168/2023-7. MMS thanks for the support provided by iTHEMS and ABL from RIKEN, the N3AS Physics Frontier Center, and the NSF under cooperative agreement 2020275. This work would not have been possible without the collaborative environment provided by the Nordic Winter School on Multimessenger Astrophysics. We thank the event organizers, as well as the financial support by Markus Ahlers for AC to attend the event.

APPENDIX

Antineutrinos.— For antineutrinos, the evolution Hamiltonian is similar to Eq. 1, with the modification $V_e \rightarrow -V_e$. Therefore, at $\rho_m \gg \rho_m^H$, $\bar{\nu}_e \approx \bar{\nu}_1^m$ for NO, and the possible fraction of electron antineutrinos on Earth is given by Eq. 10 with $|U_{e1}|^2$ in place of $|U_{e3}|^2$,

$$f_{\bar{\nu}_e}^{NO} = \frac{1}{2} \left(1 - |U_{e1}|^2 \right) + \frac{a}{2} \left(3 |U_{e1}|^2 - 1 \right). \quad (16)$$

Assuming $|U_{e1}|^2 \approx 2/3$ [36], we have

$$\frac{1}{6} \lesssim f_{\bar{\nu}_e}^{NO} \lesssim \frac{2}{3}. \quad (17)$$

⁶ Note that this constraint extends to hydrodynamic turbulence affecting neutrino propagation [42], as well as other alternative scenarios that randomize the quantities of each of the vacuum eigenstates (ν_1 , ν_2 and ν_3) reaching the Earth.

Eq. 17 is applicable even in the presence of shock waves. On the other hand, for IO, $\bar{\nu}_e \approx \bar{\nu}_3^m$ above the H -resonance density, and $f_{\bar{\nu}_e}^{IO}$ is described by Eq. 10,

$$f_{\bar{\nu}_e}^{IO} \lesssim 0.5. \quad (18)$$

If the H -resonance is non-adiabatic, transitions $\bar{\nu}_3^m \rightarrow \bar{\nu}_1^m$ modify the allowed regions for $f_{\bar{\nu}_e}^{IO}$, which will be given

by Eq. 16. Therefore,

$$\frac{1}{6} \lesssim f_{\bar{\nu}_e}^{IO}(\text{non-ad } H) \lesssim \frac{2}{3}. \quad (19)$$

In case antineutrinos emerge from the core as incoherent matter eigenstates, we have $f_{\bar{\nu}_e} \lesssim 0.7$ for both mass orderings. These results are depicted in Fig. 2.

-
- [1] S. A. Colgate and R. H. White, “*The Hydrodynamic Behavior of Supernovae Explosions*,” *Astrophys. J.* **143** (1966) 626.
 - [2] W. D. Arnett, “*Gravitational collapse and weak interactions*,” *Can. J. Phys.* **44** (1966) 2553–2594.
 - [3] J. R. Wilson, “*Supernovae and Post-Collapse Behavior*,” *Numerical Astrophysics* (1985) 422.
 - [4] H. A. Bethe and J. R. Wilson, “*Revival of a stalled supernova shock by neutrino heating*,” *Astrophys. J.* **295** (1985) 14–23.
 - [5] **Kamiokande-II**, K. Hirata *et al.*, “*Observation of a Neutrino Burst from the Supernova SN 1987a*,” *Phys. Rev. Lett.* **58** (1987) 1490–1493.
 - [6] R. M. Bionta *et al.*, “*Observation of a Neutrino Burst in Coincidence with Supernova SN 1987a in the Large Magellanic Cloud*,” *Phys. Rev. Lett.* **58** (1987) 1494.
 - [7] E. N. Alekseev, L. N. Alekseeva, I. V. Krivosheina, and V. I. Volchenko, “*Detection of the Neutrino Signal From SN1987A in the LMC Using the Inr Baksan Underground Scintillation Telescope*,” *Phys. Lett. B* **205** (1988) 209–214.
 - [8] I. Tamborra and S. Shalgar, “*New Developments in Flavor Evolution of a Dense Neutrino Gas*,” *Ann. Rev. Nucl. Part. Sci.* **71** (2021) 165–188, [arXiv:2011.01948](#).
 - [9] J. Ehring, S. Abbar, H.-T. Janka, G. Raffelt, and I. Tamborra, “*Fast neutrino flavor conversion in core-collapse supernovae: A parametric study in 1D models*,” *Phys. Rev. D* **107** (2023) no. 10, 103034, [arXiv:2301.11938](#).
 - [10] J. Ehring, S. Abbar, H.-T. Janka, G. Raffelt, and I. Tamborra, “*Fast Neutrino Flavor Conversions Can Help and Hinder Neutrino-Driven Explosions*,” *Phys. Rev. Lett.* **131** (2023) no. 6, 061401, [arXiv:2305.11207](#).
 - [11] H. Nagakura, “*Roles of Fast Neutrino-Flavor Conversion on the Neutrino-Heating Mechanism of Core-Collapse Supernova*,” *Phys. Rev. Lett.* **130** (2023) no. 21, 211401, [arXiv:2301.10785](#).
 - [12] J. T. Pantaleone, “*Neutrino oscillations at high densities*,” *Phys. Lett. B* **287** (1992) 128–132.
 - [13] J. T. Pantaleone, “*Dirac neutrinos in dense matter*,” *Phys. Rev. D* **46** (1992) 510–523.
 - [14] H. Duan, G. M. Fuller, and Y.-Z. Qian, “*Collective Neutrino Oscillations*,” *Ann. Rev. Nucl. Part. Sci.* **60** (2010) 569–594, [arXiv:1001.2799](#).
 - [15] A. Mirizzi, I. Tamborra, H.-T. Janka, N. Saviano, K. Scholberg, R. Bollig, L. Hudepohl, and S. Chakraborty, “*Supernova Neutrinos: Production, Oscillations and Detection*,” *Riv. Nuovo Cim.* **39** (2016) no. 1-2, 1–112, [arXiv:1508.00785](#).
 - [16] S. Chakraborty, R. Hansen, I. Izaguirre, and G. Raffelt, “*Collective neutrino flavor conversion: Recent developments*,” *Nucl. Phys. B* **908** (2016) 366–381, [arXiv:1602.02766](#).
 - [17] L. Wolfenstein, “*Neutrino Oscillations in Matter*,” *Phys. Rev. D* **17** (1978) 2369–2374.
 - [18] S. P. Mikheyev and A. Y. Smirnov, “*Resonance Amplification of Oscillations in Matter and Spectroscopy of Solar Neutrinos*,” *Sov. J. Nucl. Phys.* **42** (1985) 913–917.
 - [19] S. P. Mikheev and A. Y. Smirnov, “*Resonant amplification of neutrino oscillations in matter and solar neutrino spectroscopy*,” *Nuovo Cim. C* **9** (1986) 17–26.
 - [20] S. P. Mikheev and A. Y. Smirnov, “*Neutrino Oscillations in an Inhomogeneous Medium: Adiabatic Regime*,” *Sov. Phys. JETP* **65** (1987) 230–236.
 - [21] R. F. Sawyer, “*Speed-up of neutrino transformations in a supernova environment*,” *Phys. Rev. D* **72** (2005) 045003, [arXiv:hep-ph/0503013](#).
 - [22] R. F. Sawyer, “*The multi-angle instability in dense neutrino systems*,” *Phys. Rev. D* **79** (2009) 105003, [arXiv:0803.4319](#).
 - [23] R. F. Sawyer, “*Neutrino cloud instabilities just above the neutrino sphere of a supernova*,” *Phys. Rev. Lett.* **116** (2016) no. 8, 081101, [arXiv:1509.03323](#).
 - [24] S. Chakraborty, R. S. Hansen, I. Izaguirre, and G. Raffelt, “*Self-induced neutrino flavor conversion without flavor mixing*,” *JCAP* **03** (2016) 042, [arXiv:1602.00698](#).
 - [25] I. Izaguirre, G. Raffelt, and I. Tamborra, “*Fast Pairwise Conversion of Supernova Neutrinos: A Dispersion-Relation Approach*,” *Phys. Rev. Lett.* **118** (2017) no. 2, 021101, [arXiv:1610.01612](#).
 - [26] S. Abbar and M. C. Volpe, “*Using Bayesian Inference to Distinguish Neutrino Flavor Conversion Scenarios via a Prospective Supernova Neutrino Signal*,” [arXiv:2401.10851](#).
 - [27] E. K. Akhmedov, C. Lunardini, and A. Y. Smirnov, “*Supernova neutrinos: Difference of muon-neutrino - tau-neutrino fluxes and conversion effects*,” *Nucl. Phys. B* **643** (2002) 339–366, [arXiv:hep-ph/0204091](#).
 - [28] G. Sigl and G. Raffelt, “*General kinetic description of relativistic mixed neutrinos*,” *Nucl. Phys. B* **406** (1993) 423–451.
 - [29] C. Y. Cardall, “*Liouville equations for neutrino distribution matrices*,” *Phys. Rev. D* **78** (2008) 085017, [arXiv:0712.1188](#).
 - [30] A. S. Dighe and A. Y. Smirnov, “*Identifying the neutrino mass spectrum from the neutrino burst from a supernova*,” *Phys. Rev. D* **62** (2000) 033007, [arXiv:hep-ph/9907423](#).
 - [31] F. J. Botella, C. S. Lim, and W. J. Marciano,




FIG. 2. Same as Fig. 1, but for antineutrinos.

- “Radiative Corrections to Neutrino Indices of Refraction,” *Phys. Rev. D* **35** (1987) 896.
- [32] M. Sajjad Athar *et al.*, “Status and perspectives of neutrino physics,” *Prog. Part. Nucl. Phys.* **124** (2022) 103947, [arXiv:2111.07586](#).
- [33] J. Kersten and A. Y. Smirnov, “Decoherence and oscillations of supernova neutrinos,” *Eur. Phys. J. C* **76** (2016) no. 6, 339, [arXiv:1512.09068](#).
- [34] R. S. L. Hansen and A. Y. Smirnov, “Neutrino conversion in a neutrino flux: Towards an effective theory of collective oscillations,” *JCAP* **04** (2018) 057, [arXiv:1801.09751](#).
- [35] P. S. B. Dev, S. Jana, and Y. Porto, “Flavor Matters, but Matter Flavors: Matter Effects on Flavor Composition of Astrophysical Neutrinos,” [arXiv:2312.17315](#).
- [36] P. F. de Salas, D. V. Forero, S. Gariazzo, P. Martínez-Miravé, O. Mena, C. A. Ternes, M. Tórtola, and J. W. F. Valle, “2020 global reassessment of the neutrino oscillation picture,” *JHEP* **02** (2021) 071, [arXiv:2006.11237](#).
- [37] R. C. Schirato and G. M. Fuller, “Connection between supernova shocks, flavor transformation, and the neutrino signal,” [arXiv:astro-ph/0205390](#).
- [38] R. Tomas, M. Kachelriess, G. Raffelt, A. Dighe, H. T. Janka, and L. Scheck, “Neutrino signatures of supernova shock and reverse shock propagation,” *JCAP* **09** (2004) 015, [arXiv:astro-ph/0407132](#).
- [39] B. Dasgupta and A. Dighe, “Phase effects in neutrino conversions during a supernova shock wave,” *Phys. Rev. D* **75** (2007) 093002, [arXiv:hep-ph/0510219](#).
- [40] E. Akhmedov, J. Kopp, and M. Lindner, “Collective neutrino oscillations and neutrino wave packets,” *JCAP* **09** (2017) 017, [arXiv:1702.08338](#).
- [41] Y. P. Porto-Silva and A. Y. Smirnov, “Coherence of oscillations in matter and supernova neutrinos,” *JCAP* **06** (2021) 029, [arXiv:2103.10149](#).
- [42] M. Mukhopadhyay and M. Sen, “On probing turbulence in core-collapse supernovae in upcoming neutrino detectors,” [arXiv:2310.08627](#).
- [43] N. Jachowicz, K. Vantournhout, J. Ryckebusch, and K. Heyde, “Identifying neutrinos and antineutrinos in neutral-current scattering reactions,” *Phys. Rev. Lett.* **93** (2004) 082501, [arXiv:nucl-th/0406078](#).
- [44] S. Jana, Y. P. Porto-Silva, and M. Sen, “Exploiting a future galactic supernova to probe neutrino magnetic moments,” *JCAP* **09** (2022) 079, [arXiv:2203.01950](#).
- [45] S. Jana and Y. Porto, “Resonances of Supernova Neutrinos in Twisting Magnetic Fields,” *Phys. Rev. Lett.* **132** (2024) no. 10, 101005, [arXiv:2303.13572](#).