# 3d Modularity Revisited

Miranda C. N. Cheng<sup>\*1,2,3</sup>, Ioana Coman<sup>4</sup>, Piotr Kucharski<sup>5</sup>, Davide Passaro<sup>1</sup>, and Gabriele Sgroi<sup>1</sup>

<sup>1</sup>Institute of Physics, University of Amsterdam, the Netherlands

<sup>2</sup>Institute for Mathematics, Academia Sinica, Taipei, Taiwan

<sup>3</sup>Korteweg-de Vries Institute for Mathematics, University of Amsterdam, the Netherlands

<sup>4</sup>Kavli Institute for the Physics and Mathematics of the Universe, University of Tokyo, Kashiwa, Japan

<sup>5</sup>Institute of Mathematics, University of Warsaw, ul. Banacha 2, 02-097 Warsaw, Poland

### Abstract

The three-manifold topological invariants  $\hat{Z}$  capture the half-index of the three-dimensional theory with  $\mathcal{N}=2$  supersymmetry obtained by compactifying the M5 brane theory on the closed three-manifold. In 2019, surprising general relations between the  $\hat{Z}$ -invariants, quantum modular forms, and vertex algebras, have been proposed. In the meanwhile, an extensive array of examples have been studied, but several general important structural questions remain. First, for many three-manifolds it was observed that the different  $\hat{Z}$ -invariants for the same three-manifolds are quantum modular forms that span a subspace of a Weil representation for the modular group  $SL_2(\mathbb{Z})$ , corresponding to the structure of vector-valued quantum modular forms. We elucidate the meaning of this vector-valued quantum modular form structure by first proposing the analogue  $\hat{Z}$ -invariants with supersymmetric defects, and subsequently showing that the full vector-valued quantum modular form is precisely the object capturing all the Z-invariants, with and without defects, of a given three-manifold. Second, it was expected that matching radial limits is a key feature of  $\hat{Z}$ -invariants when changing the orientation of the plumbed three-manifold, suggesting the relevance of mock modularity. We substantiate the conjecture by providing explicit proposals for such  $\hat{Z}$ -invariants for an infinite family of three-manifolds and verify their mock modularity and limits. Third, we initiate the study of the vertex algebra structure of the mock type invariants by showcasing a systematic way to construct cone vertex operator algebras associated to these invariants, which can be viewed as the partner of logarithmic vertex operator algebras in this context.

<sup>\*</sup>On leave from CNRS, France.

# Contents

1	Introduction, Background, and Summary	3
2	Defects $\hat{Z}$ -invariants and Modularity         2.1       Plumbed Manifolds         2.2       Knot Surgeries         2.3       Modularity	<b>11</b> 11 13 18
3	Mock $\hat{Z}$ -invariants         3.1       Mock, Pure and Mixed         3.2       Going to the Other Side Using Appell-Lerch Sums         3.3       Regularised Indefinite Theta Functions         3.4       Indefinite Theta Functions as Mock $\hat{Z}$ -invariants         3.5       Relation to Surgeries Along Torus Knots	<ul> <li>22</li> <li>22</li> <li>22</li> <li>24</li> <li>26</li> <li>30</li> </ul>
4	Mock Invariants and Vertex Algebras4.1Cone Vertex Algebras4.2Mock VOA Characters	<b>31</b> 32 32
5	Examples5.1Invariants from Plumbing Graphs5.2Invariants from Knot Surgeries5.3More Knot Surgeries Using the Inverted Habiro Series5.4Invariants from a Hyperbolic Knot5.5Mock invariants from Indefinite Theta Functions	<ul> <li>33</li> <li>34</li> <li>37</li> <li>38</li> <li>41</li> <li>42</li> </ul>
6	Conclusion and Discussion	45
Α	Mock Modularity ProofsA.1 The Mock Modular Properties for $\Sigma(2,3,6r+1)$ A.2 Mixed Mock Modularity of $\Sigma(s,t,str+1)$	<b>51</b> 51 54
В	Polynomials $P_n^{p,b}$	55

# 1 Introduction, Background, and Summary

In this beginning section we take a brief journey through the intricate connections between quantum modular forms, vertex operator algebras (VOAs), and three-manifold topological  $\hat{Z}$ - and  $F_K$ - invariants that have been studied in recent years, discussing their context in physics, number theory, and topology, and highlighting some of the aspects that are at present still mysterious. Subsequently, we summarize the main results of this paper in less technical terms and describe the structure of the paper.

### (Mock) Modular Forms

Modular forms famously feature prominently in mathematics and branches of theoretical physics. See for instance [1, 2] for an overview. These functions on the upper-half plane  $\mathbb{H}$  are distinguished by their *modular* symmetry property, which reflects the discrete symmetries of  $\mathbb{H}$ . It is interesting to ask how this symmetry can be broken in natural and meaningful ways. Mock modular forms embody such a natural generalization of modular forms. Here, the modular symmetry only emerges when a non-holomorphic contribution determined by a modular form – the *shadow* – is added to the mock modular form. Since the development of their modern theory two decades ago [3, 4, 5, 6], through a series of rapid developments it has been established that mock modular forms have a similarly prominent role in combinatorics, moonshine, conformal field theory, string theory and more, extending the applications of modular forms in a fascinating way. A partial list of examples can be found in [6, 7, 8, 9, 10, 11, 12, 13, 14]. Specifically, it is known that mock modular forms appear as characters of certain super vertex algebras [15, 16, 17, 18, 13].

Theta functions of lattices with indefinite signatures played an important role in the development of the theory of mock modular forms [3]. Building on previous work by Vignéras [19], Zwegers [3] showed that a regularisation for theta functions of signature (1, n) (one negative direction) leads to a theta function with mock modular properties. Subsequent work has shown that regularised theta functions of general signatures lead to higher-depth mock modular forms [20, 21], and this specific realization of mock modular forms has featured in the study of string theory [22, 23, 24] and umbral moonshine [25, 26]. Generally, indefinite theta functions give rise to mixed-mock modular forms, which are mock-type forms whose completion involves a finite sum of products between usual modular forms and non-holomorphic contributions from the shadows (cf. §3.1).

Apart from its modified modular symmetry property (cf. (1.9)), another earmark of mock theta functions, first pointed out by Ramanujan [27, 28], is their behaviour near the cusps  $\hat{\mathbb{Q}} := \mathbb{Q} \cup \{i\infty\}$  of the upper-half plane. In the modern language, this leads to the related statement that mock modular forms give rise to quantum modular forms (QMFs) [29]. Quantum modular forms, in essence, are functions whose differences with their images under the modular group  $\mathrm{SL}_2(\mathbb{Z})$  enjoy better analytic behaviour when considered near the rationals, compared to the original function. Below we will discuss the relevance of quantum modular forms in the context of three-manifold invariants.

The results of this work further establish the role of (mixed) mock modular forms as topological invariants of three-manifolds, which we will introduce shortly. A preliminary set of results has been reported in [13].

In the discussion of the quantum modularity of three-manifold  $\hat{Z}$ -invariants, a special role will be played by certain Weil representations of the metaplectic group  $\widetilde{SL}_2(\mathbb{Z})$  [30, 31]. In particular, we will encounter the Weil representations  $\Theta^{m+K}$  (2.37, 2.39), which are subrepresentations of the 2*m*-dimensional representation  $\Theta_m$  spanned by the column vector  $\theta_m = (\theta_{m,r})_{r \mod 2m}$  with a positive integer m and theta function components

$$\theta_{m,r}(\tau,z) := \sum_{\ell \equiv r \bmod 2m} q^{\frac{\ell^2}{4m}} e^{2\pi i z \ell} , \qquad q = e^{2\pi i \tau} , \qquad (1.1)$$

labelled by a subgroup K of the group of exact divisors  $Ex_m$  satisfying  $m \notin K$ . The precise definition can be found in §2.3.

#### $\ddot{Z}$ -invariants for Three-Manifolds

Arguably, one of the most prominent open problem in topology is the smooth fourdimensional Poincaré conjecture, which states that there are no exotic spherical smooth structures in four dimensions. Motivated by this, decades ago Crane and Frenkel envisioned that a *categorification* of numerical three-manifold invariants [32], replacing them with more sophisticated structures of (spectral sequences of) vector spaces, could potentially hold the key to defining structures distinguishing exotic smooth structures in four dimensions. A quantum topological invariant for three-manifolds  $M_3$  [33, 34] and the closely related quantum knot invariant  $F_K$  [35] were proposed in physical terms recently, leading to an interesting new approach to such a categorification program. Specifically, the origin of these invariants in physical M-theory lends weight to a possible connection to four-dimensional topology. More precisely, it is proposed to consider the half-index, namely the supersymmetric partition function on the background of cigar times the temporal circle, of the three-dimensional quantum field theory obtained by compactifying the M5 brane theory, or the 6d  $\mathcal{N} = (2,0)$  ADE superconformal field theory to be more precise, on  $M_3$  as the topological invariant of  $M_3$ . Note that the three-dimensional spacetime can also be thought of as a solid torus, with the complex structure  $\tau$  of the boundary torus identified with the argument of the  $\hat{Z}$ -invariants.

However, the physical proposal does not translate into a computational algorithm to compute it for general three-manifolds, due to our insufficient detailed knowledge of M-theory. To move forward, hints can be obtained from their relation to the Witten-Reshetikhin-Turaev invariants they seek to categorify. Following these hints, conjectural expressions for  $\hat{Z}$ -invariants have been proposed in [34] for a particularly simple infinite family of three-manifolds: the weakly negative plumbed three-manifolds. To explain what they are, first recall that plumbed three-manifolds are three-manifolds that can be constructed by taking the boundary of a four-manifold obtained by gluing together disk bundles over  $S^2$ . The data can be encoded in terms of a weighted graph called a plumbing graph, obtained by identifying the set V of vertices with the set of disk bundles, equipped with a weight function  $a: V \to \mathbb{Z}$  whose values are the Euler numbers  $a_v$  of the disk bundles, and connecting the two vertices v and v' with an edge if the corresponding disk bundles are glued. Alternatively, the same data can be captured using the "plumbing matrix" M with the weights  $a_v$  on the diagonal and 1 or 0 for the off-diagonals depending on whether the corresponding pair of vertices are connected. A vertex is said to be a high-degree vertex if it is connected to at least three other vertices,  $deg(v) \geq 3$ . Finally, the plumbed three-manifold is said to be weakly negative if the inverse plumbing matrix  $M^{-1}$  is negative-definite in the subspace generated by the high-degree vertices. In this paper, we mainly focus on Seifert manifolds with three exceptional fibres, which correspond to plumbing graphs with one degree three vertex connected with three rays. We will therefore refer to them as the *negative* and *positive*  $M_3$ , given by the signature of  $M^{-1}$  along the one-dimensional subspace spanned by the central vertex.

Consider such a weakly negative plumbed three-manifold  $M_3$  with plumbing matrix M. Given a choice of  $b \in (\delta + \operatorname{Coker}(\mathsf{M}))$  (cf. (2.3)), corresponding to a choice of  $\operatorname{Spin}^{c}$ -structure on  $M_3$ , it was conjectured that the corresponding 3d half-index is given by the contour integral [34]

$$\hat{Z}_b(M_3;\tau) := q^{\Delta} \oint \prod_{v \in V} \frac{dz_v}{2\pi i z_v} \left( z_v - \frac{1}{z_v} \right)^{2 - \deg(v)} \Theta_b^{\mathsf{M}}(\tau; \mathbf{z})$$
(1.2)

for some  $\Delta \in \mathbb{Q}$ , where the theta function is given by

$$\Theta_b^{\mathsf{M}}(\tau; \mathbf{z}) := \sum_{\ell \in 2\mathsf{M}\mathbb{Z}^{|V|} \pm b} q^{-\ell^T \mathsf{M}^{-1}\ell/4} \, \mathbf{z}^{\ell}.$$
(1.3)

When necessary, the integral is defined as the principal value integration. It was shown that the WRT invariant, which is the Chern-Simons partition function suitably normalized, can indeed be recovered from  $\hat{Z}_b(M_3;\tau)$  defined above, by combining  $\hat{Z}_b(M_3;\tau)$  for different *b* and by taking the limit  $\tau \to 1/k$ , often referred to as the radial limit, from within the upper-half plane [34]. Schematically, we have

$$\hat{Z}_b(M_3;\tau) \xrightarrow[\text{combining } b]{\tau \to \frac{1}{k} + i0^+} \text{WRT}(M_3;k)$$
(1.4)

where  $k \in \mathbb{Z}$  is the (renormalized) Chern-Simons level.

The above consideration can be extended from closed three-manifolds to knot complements. Consider a graph with a distinguished degree one node. It corresponds to a three-manifold Y which can be identified as the complement of a knot K associated with the distinguished node, in a closed three-manifold  $\hat{Y}$  which is the plumbed manifold with the plumbing graph given by the graph that we started with, but with the distinguished node replaced by a regular node. From this construction one can write down the two-variable topological invariant  $F_K(x,\tau)$  associated to the knot complement [35]. Moreover, another closed three-manifold  $Y_{p/r}$  can be obtained via a p/r Dehn surgery along the knot. The  $\hat{Z}$ -invariant of  $Y_{p/r}$  can be obtained via a Laplace-like transformation of the invariant  $F_K(x,\tau)$  (2.8). For some knots K, it was conjectured that there  $F_K$  can be expressed in terms of so-called inverted Habiro series of K [36]. This together with a set of conjectural formulas for Dehn surgeries provides further conjectural methods to compute  $\hat{Z}$ -invariants, sometimes applicable also to closed three-manifolds that are not weakly negative plumbed three-manifolds. All the above (conjectural) ways of obtaining the  $\hat{Z}$ -invariants will be used in our article later.

#### **Defect Operators**

As mentioned above,  $\hat{Z}$ -invariants give information about a three-manifold  $M_3$  by computing the half-index, or the supersymmetric partition function on the background of a cigar times the temporal circle, of the 3d  $\mathcal{N} = 2$  theory arising from compactifying M5branes on  $M_3$ . It is informative to incorporate half-BPS line operators in the 3d  $\mathcal{N} = 2$ theory, arising from M2 branes located at the centre of the cigar and wrap around a link  $K \subset M_3$  ([34], §4). One can then compute the half-index  $\hat{Z}_b(M_3; W; \tau)$  in the presence of these half-BPS line operators W. As before, the parameter  $\tau \in \mathbb{H}$  can be identified with the complex structure of the boundary torus of the 3-dimensional space  $D_2 \times \mathbb{R}_t$  on which the 3d  $\mathcal{N} = 2$  theory lives. In this paper we focus on the gauge group G = SU(2), while a similar defect halfindices can be defined for general ADE groups. See also [37, 38] for prior considerations. Moreover, as in the other parts of the paper, we focus on Seifert manifolds with three exceptional fibres, and links associated with the end nodes of the plumbing graph. We conjecture an explicit expression for  $\hat{Z}_b(M_3; W_{\nu}; \tau)$ , where  $\boldsymbol{\nu} = (\nu_1, \nu_2, \nu_3) \in \mathbb{Z}^3_{\geq 0}$  specifies the highest weight representations, and  $\nu_i$  times the fundamental weight is the highest weight associated to the knot corresponding to the *i*-th end node of the plumbing graph, as well as their formulation in terms of knot surgeries. We will find that these defect halfindices are indispensable in the full understanding of the quantum modular properties, to be discussed below, of the  $\hat{Z}$ -invariants. This is consistent with our expectation that including supersymmetric defects provides important physical insights into the system.



Figure 1: The topics of this paper and their relations.

### $\hat{Z}$ -invariants, Quantum Modular Forms, and Vertex Operator Algebras

The study of the properties of the  $\hat{Z}$ -invariants was initiated in [30]. In particular, the authors of [30] proposed the following

- 1.  $\hat{Z}$ -invariants are closely related to quantum modular forms of some types;
- 2.  $\hat{Z}$ -invariants are related to characters of vertex operator algebras.

The relation to quantum modular objects built on an array of interesting earlier results and observations in a similar context [33, 34, 39, 40, 41, 42, 43].

Since then, it has been an active area of study to extend and verify these broad conjectures [13, 31, 37, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62]. Both of the conjectural properties point to a large system of unexpected symmetries in the underlying physical system and the corresponding topological problem. Consequently, understanding the relation to VOA and quantum modular forms will constitute interesting progress in the understanding of low-dimensional topology and the symmetries of M-theory compactifications.

In the case of VOAs, it is believed that the conjectured property 2. is a consequence of a VOA symmetry acting on the BPS states of the relevant 3d  $\mathcal{N} = 2$  theory. See also [63, 64] for related developments. At the same time, these non-trivial applications of VOAs have the potential of driving further the development of the general theory of (non-rational) vertex operator algebras and quantum modular forms. See Figure 1.

In the case of quantum modular forms, the study can be viewed as a part of the broader effort to understand the mysterious quantum modular behaviour of related invariants in low-dimensional topology [39, 65, 66, 41, 42, 43]. Physically, this type of modified symmetry under the action of the modular group  $SL_2(\mathbb{Z})$  is expected to stem from the  $SL_2(\mathbb{Z})$  action on the torus boundary of the 3d spacetime relevant for the half-index, that is a solid torus. Interestingly, in [30] we have seen for many three-manifolds hints of concrete  $SL_2(\mathbb{Z})$  representations behind the different  $\hat{Z}$ -invariants of the given manifolds, though these invariants appear to only span a subspace of the representation. More concretely, we see that the  $\hat{Z}$ -invariants often correspond to some but not all of the components of a  $SL_2(\mathbb{Z})$  vector-valued quantum modular form. A natural question which we address in this work, is then

# **Question 1:** What is the meaning of the other components of the $SL_2(\mathbb{Z})$ vector-valued quantum modular forms?

We will answer this question by providing an interpretation for the full  $SL_2(\mathbb{Z})$  representations in this work, hence elucidating the action of the modular group in the system. The fact that  $\hat{Z}$ -invariants with different defects form a vector-valued quantum modular form suggests an interesting modular group action on the set of possible supersymmetrypreserving boundary conditions, or equivalently supersymmetric M2 brane configurations.

Note also that the two conjectures are connected in an interesting and non-trivial way. The relevant question bridging the two is: what is the modular-like property of the characters of a general vertex operator algebra? Such properties, if exist, could be viewed as generalizations of the celebrated Zhu's modularity theorem beyond the realm of rational,  $C_2$ -cofinite VOAs [67, 68]. See [69, 70, 71, 72, 73, 74] for some of the related works.

Many questions still remain open for general weakly negative plumbed manifolds and for general ADE gauge groups, including the construction of the vertex operator algebras and the proof of the (higher-depth) quantum modularity properties. But an even more fascinating direction is to explore plumbed manifolds that are not weakly negative, as the "mathematical definition" (1.2) (and the analogue expression for general ADE gauge groups [54]) is not even available in these cases. As pointed out in [13] and will be refined in §3.4, the origin of the difficulty is obvious: the contour integral leads to an infinite sum along the directions of high-degree vertices, and we hence need the bilinear form given by  $-M^{-1}$  to be positive definite when restricted to those directions, in order to lead to a q-series with powers of q bounded from below. In [30], the authors refer to the question of how to execute the  $\tau \mapsto -\tau$  transformation as the question of "going to the other side". Here, the crucial question is

Question 2: How can we take  $\tau \mapsto -\tau$  to turn  $\hat{Z}(M_3)$  into  $\hat{Z}(-M_3)$ ?

Despite the difficulty, in this work we will take a step towards a general definition, exploiting insights from quantum modular forms<sup>1</sup>.

After identifying the invariants  $\hat{Z}(-M_3)$ , the second part of the 3d modularity proposal leads to the following question.

**Question 3:** Which VOAs correspond to the invariants  $\hat{Z}(-M_3)$  on the other side?

<sup>&</sup>lt;sup>1</sup>While this work is being completed, the paper [75] appeared which investigates the problem from an interesting resurgence point of view, complementary to the modular point of view taken in this work.

In this paper we will address the above three questions.

#### The False and the Mock

To understand the phenomenon of "going to the other side", first recall the chirality property of the Chern-Simons theory, which leads to the relation

$$WRT(M_3;k) = WRT(-M_3;-k)$$
(1.5)

between WRT invariants of orientation-reversed three-manifolds. Since the orientation reversal operation does not preserve the weakly negative property of the plumbing graph, the above chirality property, the relation (1.4) between  $\hat{Z}$ -invariants and WRT invariants, and the expression (1.2) for the weakly negative  $\hat{Z}$ -invariants, provide important hints about  $\hat{Z}$ -invariants for plumbed manifolds that are not weakly negative. Naively, one expects

$$\ddot{Z}(-M_3;\tau) = \ddot{Z}(M_3;-\tau).$$

This hint was exploited in [30] and resulted in the so-called False-Mock Conjecture.

Focusing for now on the negative Seifert manifolds with three exceptional fibres. It has been established [30, 44] that their  $\hat{Z}$ -invariants are, up to an overall q-power and the addition of a finite polynomial, false theta functions. By false theta function, we mean any function of the form  $\sum_{k \in \mathbb{Z}} a_k q^{k^2/4m}$ , where  $a_k = a_{k+2m} = -a_{-k}$  for all  $k \in \mathbb{Z}$ , for some positive integer m. In other words, they are linear combination of

$$\tilde{\theta}_{m,r}(\tau) := \sum_{k=r \pmod{2m}} \operatorname{sgn}\left(k\right) q^{\frac{k^2}{4m}} , \ r \in \mathbb{Z}/2m,$$
(1.6)

for a given  $m \in \mathbb{N}$ , where the nomenclature stems from the additional sign factor in the one-dimensional lattice sum.

False theta functions can be viewed as Eichler integrals, which makes obvious their quantum modular properties [29]. More precisely,  $\tilde{\theta}_{m,r}$  is, up to a numerical multiplicative factor, the holomorphic Eichler integral of the unary theta function  $\theta_{m,r}^1(\tau) := \sum_{k=r \pmod{2m}} k q^{\frac{k^2}{4m}}$ . The holomorphic Eichler integral of a weight  $w \in \frac{1}{2}\mathbb{Z}$  cusp form  $g(\tau) = \sum_{n>0} a_g(n)q^n$  is given by

$$\tilde{g}(\tau) := \sum_{n>0} a_g(n) \, n^{1-w} q^n \,, \tag{1.7}$$

or equivalently

$$\tilde{g}(\tau) = C(w) \int_{\tau}^{i\infty} g(\tau')(\tau' - \tau)^{w-2} d\tau'$$
(1.8)

with  $C(w) = (2\pi i)^{w-1}/\Gamma(w-1)$  and a carefully chosen integration path. One similarly defines the non-holomorphic Eichler integral as

$$g^*(\tau) = C(w) \int_{-\bar{\tau}}^{i\infty} \overline{g(-\bar{\tau'})} (\tau' + \tau)^{w-2} d\tau'.$$
(1.9)

In terms of Fourier expansion, we have<sup>2</sup>

$$e(\frac{1-w}{2})\Gamma(w-1)g^*(\tau) = \sum_{n>0} n^{1-w}\bar{a}_g(n)\Gamma(w-1,4\pi n\tau_2)q^{-n}$$
(1.10)

<sup>2</sup>There is in general an additional term  $a_g(0) \frac{(4\pi\tau_2)^{w-1}}{1-w}$  if g is not a cusp form.

where we have written  $\tau_2 = \text{Im } \tau$ ,  $e(x) := e^{2\pi i x}$ , and the incomplete  $\Gamma$  function is given by  $\Gamma(1-k,x) = \int_x^\infty t^{-k} e^{-t}$ . Note that the summand of the second term vanishes as  $e^{-2\pi n \tau_2}$  in the limit  $\tau_2 \to \infty$ .

A weight w (weakly holomorphic) mock modular form  $f : \mathbb{H} \to \mathbb{C}$  with shadow given by a weight 2 - w cusp form g is a (weakly) holomorphic function such that the nonholomorphic function  $\hat{f} := f - g^*$  transforms as a modular form of weight w. From now on we focus on the case of weight w = 1/2 mock modular forms for  $\Gamma \subset SL_2(\mathbb{Z})$  relevant for the context in this paper. Near a cusp  $u \in Q$ , f might have a exponential singularity. In particular, there exists a finite set  $\{f_u\}$  of weakly holomorphic modular forms such that  $f - f_u$  is finite as  $\tau$  approaches u from the upper-half plane [28]. Crucially, the cardinality of this set is necessarily larger than one for any mock modular form f of weight 1/2 with a non-vanishing shadow. Equipped with  $\{f_u\}$ , we can subtract the possible singularities of the mock form f by defining  $f_{u,fin} := f - f_u$ . Then their asymptotic expansions near a cusp and those of the Eichler integral of its shadow agree to infinite order [30, 76]:

$$f_{u,\text{fin}}(u+iv) \sim \sum_{n\geq 0} \alpha_u(n)v^n$$
,  $g^*(-u+iv) \sim \sum_{n\geq 0} \alpha_u(n)(-v)^n$ , (1.11)

Note that the role of the  $f_u$  in the subtraction is just to remove the singularity. So the above relation also holds when  $f_u$  is replaced by f with just the exponential singularities terms subtracted, instead of the whole modular form  $f_u$  (the difference vanishes exponentially near the cusp u). Similar results also generalize to mixed mock modular forms, as we will discuss in §3.1. Note that in the case u = 1/k, the above relation is consistent with the reversal symmetry (1.5) of the WRT invariants. Combined with (1.4), and other evidence, this consideration on the radial limits leads to the following *false-mock conjecture*,<sup>3</sup> which constitutes an important inspiration of the present paper.

**Conjecture 1.** Let  $M_3$  be a three-manifold for which the  $\hat{Z}$ -invariants take the form

$$\hat{Z}_b(M_3;\tau) = q^c \left( \tilde{\vartheta}(\tau) + p(\tau) \right), \qquad (1.12)$$

where  $\tau \in \mathbb{H}$ ,  $c \in \mathbb{Q}$ ,  $\hat{\vartheta}(\tau)$  is the Eichler integral of a unary theta function  $\vartheta(\tau)$  of weight 3/2 and  $p(\tau)$  is a polynomial in  $q = e^{2\pi i \tau}$ . Then the  $\hat{Z}$ -invariant of the manifold with reversed orientation is

$$\hat{Z}_b(-M_3;\tau) = q^{-c} \left( f(\tau) + p(-\tau) \right), \tag{1.13}$$

where  $f(\tau)$  is a weight 1/2 weakly holomorphic (mixed) mock modular form. Moreover, its completion  $\hat{f}$ , transforming as a modular form of weight 1/2 under a certain congruence subgroup of  $SL_2(\mathbb{Z})$ , has the following properties. It is given by

$$\hat{f} = f - \vartheta^* - \sum_{i \in I} g_i \vartheta_i^* \tag{1.14}$$

where I is a finite set,  $\vartheta_i$  is a theta function, and  $g_i$  with any  $i \in I$  is a modular function for some discrete subgroup of  $SL_2(\mathbb{Z})$  that either vanishes or has an exponential singularity at all cusps.

Crucially,  $\hat{\vartheta}$  and the finite part of the shadow  $\vartheta^*$  are the holomorphic resp. non-holomorphic Eichler integrals of the *same* cusp form  $\vartheta$ .

<sup>&</sup>lt;sup>3</sup>Note that this version of the false-mock conjecture is slightly more general than the version in [13, 30]: we have allowed for potential which further requires the mixed mock modular form to be a mock modular form, with I the empty set.

### Summary of Results

In this paper we aim to answer the three questions mentioned above. We will do so in the following steps.

- 1. We propose two expressions, Definition 1 in terms of plumbing graphs and Definition 2 in terms of knot surgeries, as additional  $\hat{Z}$  topological invariants. We conjecture that they give the defect half-indices, capturing the effects of inserting supersymmetric defect operators in the 3d theory  $\mathcal{T}(M_3)$ .
- 2. With the above definition, we propose that, when an appropriately defined set of defects operators are included, the  $\hat{Z}$ -invariants of a given three-manifold are related to a (higher-depth) vector-valued quantum modular form of some type for the modular group  $SL_2(\mathbb{Z})$ .

Focusing on Seifert manifolds with three singular fibres, we can be more concrete about the above expectation which we phrase in terms of the following Modularity Conjecture:

For a given Seifert manifold  $M_3$  with three singular fibres, the vector space spanned by  $\hat{Z}(M_3; W_{\nu})$  is isomorphic to a Weil representation  $\Theta^{m+K}$ , labelled by a positive integer m and a subgroup K of the group of exact divisors of m.

More explicitly, we conjectured that, up to finite polynomials and pre-factors, the vector space spanned by  $\hat{Z}(M_3; W_{\nu})$  is also spanned by the components of a vector-valued quantum modular form for  $\mathrm{SL}_2(\mathbb{Z})$ , transforming under the (dual of the) Weil representation  $\Theta^{m+K}$ . See Conjecture 4 for the precise version of this Modularity Conjecture. Moreover, we show that this conjecture holds when  $M_3$  is any weakly negative Brieskorn sphere  $\Sigma(p,q,r)$  (see Theorem 1).

This answer the  $Question \ 1$  listed above.

3. From analysing the regularization of the contour integral and the knot surgery formulations for  $\hat{Z}$ -invariants, in Conjecture 5 we propose concrete mock modular forms as the (defects)  $\hat{Z}$ -invariants for the inverted Brieskorn spheres  $-\Sigma(2,3,6r+1)$ ,  $-\Sigma(2,3,5)$ . Subsequently, we prove that the Modular Conjecture in the previous item holds for these conjectured invariants of these inverted Brieskorn spheres. See Theorem 2 and Corollary 1. We also analyse the mixed mock modularity of the functions obtained by applying a surgery formula, relevant for the more general manifolds  $-\Sigma(s, t, str + 1)$ .

This provides an answer to *Question 2* listed above in these cases.

- 4. We propose a way to systematically attach vertex operator algebras corresponding to the positive Seifert manifolds  $M_3$  with three singular fibres, when  $\hat{Z}(M_3)$  can be expressed in terms of regularized theta functions for a lattice with indefinite signature. This provides an answer to the Question 3 listed above.
- 5. Using our proposals (Definition 1 and 2) for the defect invariants, in §5 we provide an extended list of concrete examples, furnishing evidence for the Modularity Conjecture 4 as well as the validity of the proposals for the defect and mock invariants themselves.

# 2 Defects $\hat{Z}$ -invariants and Modularity

In this section we will first introduce our proposal for defects  $\hat{Z}$ -invariants in terms of the contour integrals and knot surgeries. Equipped with this larger family of invariants, we will take a fresh look at the quantum modularity of the  $\hat{Z}$ -invariants. In particular, we will shed light on their structure as vector-valued quantum modular forms for the modular group  $\widetilde{SL}_2(\mathbb{Z})^4$ , and the important role of the defects in  $\hat{Z}$ -modularity. The consideration of quantum modularity should hold equally for three-manifolds of both orientations, though as we see the actual methods for computing them will differ on the different sides.

### 2.1 Plumbed Manifolds

First we give more information on the contour integration expression (1.2) for Z-invariants for weakly negative plumbed three-manifolds introduced in §1. Let  $M_3$  be a plumbed three-manifold; for instance, any Seifert manifold with n singular fibres  $M_3(a; \{q_i/p_i\}_{i=1}^n)$ is an example of plumbed three-manifolds. Its plumbing graph contains a central vertex  $v_0$  connected to  $n \geq 3$  legs and the weights  $\alpha_j^{(i)}$  for the nodes on the  $i^{th}$  leg are determined by

$$\frac{q_i}{p_i} = \frac{-1}{\alpha_1^{(i)} - \frac{1}{\alpha_2^{(i)} - \frac{1}{\alpha_3^{(i)} - \dots}}} .$$
(2.1)

For  $M_3$  a weakly negative plumbed three-manifold, using the above notation, the topological invariants  $\hat{Z}_b(M_3; \tau)$  are defined via the principal value |V|-dimensional integral (1.2) [34],

$$\hat{Z}_b(M_3;\tau) := (-1)^{\pi} q^{\frac{3\sigma - \sum_{v \in V} \alpha(v)}{4}} \operatorname{vp} \oint \prod_{v \in V} \frac{dz_v}{2\pi i z_v} \left( z_v - \frac{1}{z_v} \right)^{2 - \operatorname{deg}(v)} \Theta_b^{\mathsf{M}}(\tau, \mathsf{z}) , \qquad (2.2)$$

where  $\sigma$  is the signature of the plumbing matrix M,  $\pi$  denotes the number of positive eigenvalues of M, and the contours of integration are to be the set  $|z_v| = 1$ . The label b can be identified with the elements of the set

$$\operatorname{Spin}^{c}(M_{3}) \cong (2\mathbb{Z}^{|V|} + \delta) / (2M\mathbb{Z}^{|V|}) , \qquad (2.3)$$

where  $\delta \in \mathbb{Z}^{|V|}/2\mathbb{Z}^{|V|}$  is defined by  $\delta_v = \deg(v) \mod 2$ . Labelled by  $b \in \operatorname{Spin}^c(M_3)$  (up to the Weyl group action), the theta function in equation (2.2) is given by (1.3).

Next we propose a generalisation of the Z-invariants, intended to account for the inclusion of supersymmetric defect lines in the 3d  $\mathcal{N} = 2$  quantum field theory. For the purposes of this paper, we only give the explicit expression for SU(2) gauge groups, though the generalization to ADE gauge groups should be straightforward (cf. [37]). Denote by  $\vec{\omega}$  the fundamental weight of the Lie algebra  $\mathfrak{sl}_2$ . We consider defects in the 6-dimensional theory with support on links associated to a collection of nodes,  $v \in V_W$ , in the plumbing graph, with the corresponding highest weight representation with highest weights  $\nu_v \vec{\omega}$ .

 $<sup>^4 \</sup>mathrm{or}$  equivalently something that we will call quantum Jacobi theta functions. See the comment at the end of the section.

**Definition 1.** Consider a weakly negative plumbed manifold  $M_3$ , and defects associated to a collection of nodes  $V_W$  in the plumbing graph, with the highest weight representation with highest weight  $\nu_v \vec{\omega}$ . We define the defect  $\hat{Z}$ -invariant as given by the contour integral

$$Z_{b}(M_{3}; W_{\{\nu_{v}\}_{v \in V_{W}}}; \tau) := (-1)^{\pi} q^{\frac{3\sigma - \sum_{v \in V} \alpha(v)}{4}} \operatorname{vp} \oint \prod_{v \in V} \frac{dz_{v}}{2\pi i z_{v}} \left(z_{v} - \frac{1}{z_{v}}\right)^{2 - deg(v)} \left(\prod_{v \in V_{W}} \chi_{\nu_{v}}(z_{v})\right) \Theta_{b + \{\nu_{v}\}_{v \in V_{W}}}^{\mathsf{M}}(\tau, \mathsf{z}) ,$$

$$(2.4)$$

where  $\chi_{\nu_v}(z_v)$  denotes the  $\mathfrak{sl}_2$  character

$$\chi_{\nu}(z) = \frac{z^{1+\nu} - z^{-1-\nu}}{z - z^{-1}} = \sum_{k=0}^{\nu} z^{\nu-2k} , \qquad (2.5)$$

and the modified theta function  $\Theta_{b+\{\nu_v\}_{v\in V_W}}^{\mathsf{M}}$  is given by modifying  $b \in (2\mathbb{Z}^{|V|}+\delta)/(2M\mathbb{Z}^{|V|})$ in (1.3) by replacing  $b_v \mapsto b_v + \nu_v$  for  $v \in V_W$ .

**Proposition 1.** The defect  $\hat{Z}$ -invariant proposed in Definition 1 is a topological invariant of  $M_3$ , as it is invariant under all the 3d Kirby moves on the plumbing graph of  $M_3$  preserving the nodes with  $\nu_v \neq 0$ .

*Proof.* The proof is analogous to that of Proposition 4.6 of [35].

Moreover, we conjecture that the proposed modification of  $\hat{Z}$  also has the following connection to the physical theory.

**Conjecture 2.** The defect  $\hat{Z}$ -invariant is equal to the half-index of the same  $3d \mathcal{N} = 2$  supersymmetric quantum field theory with the corresponding defect operators included.

- **Remark 1.** The above Conjecture generalizes the expressions in previous work [34, 37] by modifying the theta function  $\Theta_{b+\{\nu_v\}_{v\in V_W}}^{\mathsf{M}}$  in the integrand of the contour integral.
  - In §3 we will discuss how to generalize the above contour integrations for some plumbed three-manifolds which are not weakly negative.
  - In what follows, we will often focus on the case where  $M_3$  is a Seifert manifold with three exceptional fibres, and where Wilson lines are added on links associated to the end nodes of the three legs in the plumbing graph. We will label the corresponding highest weight representations by  $\boldsymbol{\nu} = (\nu_1, \nu_2, \nu_3) \in \mathbb{N}^3$  and denote the corresponding defect  $\hat{Z}$ -invariant by  $\hat{Z}_b(M_3; W_{\boldsymbol{\nu}})$ .
  - Note that the expression (2.4) has the artefact of breaking original the Weyl symmetry, in this case the Weyl symmetry b → −b, of the choice of the (generalized) Spin<sup>c</sup> structure. Instead, the Weyl group acts like b + ν → −(b + ν) now. As a result, for a detailed matching between the physical defect half-indices and the above prescription, the representative of b should be carefully chosen to lie in the right Weyl chamber. See §5 for some examples.

## 2.2 Knot Surgeries

For certain closed manifolds  $M_3$ , the invariants  $\hat{Z}_b(M_3)$  can be computed in a number of ways that are expected to be equivalent. After discussing the contour integral definition using the combinatorial data of the plumbed manifolds, we will now review the proposal for  $\hat{Z}$ -invariants of closed three-manifolds constructed through surgeries of knots.

Starting with a three-manifold Y which is the complement of a tubular neighbourhood of a knot K in  $S^3$ , which has a parametrised torus boundary, we can construct a closed manifold  $S^3_{p/r}(K)$  through surgery by gluing to the boundary  $\partial Y$  a solid torus  $S^1 \times D^2$ which is seen as a tubular neighbourhood of the knot K

$$S^3_{p/r}(K) = Y \cup_{\partial Y} (S^1 \times D^2) , \qquad (2.6)$$

with  $p/r \in \mathbb{Q}$  prescribing the identification of cycles. When the manifold Y admits a plumbing description, we refer to the knot K as a *plumbed knot*. In this case, one can associate to it a two-variable series  $F_K(x;\tau)$  through a contour integral similar to that in (2.2) for closed plumbed manifold, with the only difference being that the plumbing graph has now a distinguished vertex, which we denote by  $v_K$ , corresponding to the boundary  $\partial Y$  [35].

The  $\hat{Z}$ -invariant of the closed plumbed manifolds  $S^3_{p/r}(K)$  is related to  $F_K$  through the surgery formula

$$\hat{Z}_b(S^3_{p/r}(K);\tau) = \epsilon q^d \cdot \mathcal{L}^{(b)}_{p/r} \left[ \left( x^{\frac{1}{2r}} - x^{-\frac{1}{2r}} \right) F_K(x,\tau) \right] , \qquad (2.7)$$

where  $\epsilon$  can be  $\pm 1$ ,  $d \in \mathbb{Q}$  is a number fixed by the manifold  $S^3_{p/r}(K)$ , as defined in Theorem 1.2 of [35]. The map  $\mathcal{L}^{(b)}_{p/r}$  acts as<sup>5</sup>

$$\mathcal{L}_{p/r}^{(b)}(x^{u}) = q^{-u^{2}r/p} \delta_{ru-b/2 \pmod{p}}$$
(2.8)

where for  $x \in \mathbb{Z}, p \in \mathbb{N}$  we define  $\delta_{x \pmod{p}} = 1$  when  $x \equiv 0 \pmod{p}$  and 0 otherwise.

Next we will generalize the above to include Wilson line operators. As before, we can consider defects supported on links associated to the nodes of the plumbing graphs. In what follows we focus on Wilson lines along the knot K corresponding to the distinguished node. The effect of adding Wilson lines associated to other nodes can be computed in a way completely analogous to Definition 1 of the last subsection, leading to a defect  $F_K$ series, which can be denoted by  $F_{K,\nu}$  and whose definition is a simple modification of Definition 2.4.

**Definition 2.** Using the same notation as above, consider the  $F_K(x,\tau)$  series associated to the plumbed knot K with a defect operator along K in the highest weight representation of  $\mathfrak{sl}_2$  with highest weight  $\nu \vec{\omega}$ . We define the corresponding defect invariant for the closed manifold  $S^3_{p/r}(K)$  as

$$\hat{Z}_b(S^3_{p/r}(K); W_{\nu}; \tau) = \epsilon q^d \cdot \mathcal{L}^{(b+\nu)}_{p/r} \left[ \left( x^{\frac{1}{2r}} - x^{-\frac{1}{2r}} \right) F_K(x, \tau) \chi_{\nu}(x^{\frac{1}{2r}}) \right] , \qquad (2.9)$$

where the  $\mathfrak{sl}_2$  character  $\chi_{\nu}$  is as in (2.5), and  $\epsilon$ , d are as in (2.7) where there are no Wilson lines.

<sup>&</sup>lt;sup>5</sup>A relative factor of 1/2 is introduced here when compared to [35], which can be understood as coming from the fact that we use the weight instead of the root basis to write down the highest weight of the Wilson lines as well as in the plumbing prescription (2.4). Relately, one can think about x as  $z^2$  when comparing (2.7) with (2.4).

Note that performing the Laplace transform is equivalent to integrating over the distinguished node (see the proof of Theorem 1.2 of [35]). As a result, the interpretation as the defect half-index of the invariant  $\hat{Z}_b(S^3_{p/r}(K); W_{\nu})$  defined above follows if the interpretation for  $\hat{Z}(M_3; W_{\nu})$  formulated in Conjecture 2 holds true.

**Remark 2.** Note that in (2.4) we have shifted the argument b in the theta function, and in (2.9) the argument in the Laplace transformation. In general, the shift is non-trivial and not just a relabeling since  $b + \nu$  might not belong to the set (2.3) of permissible Spin<sup>c</sup>-structures, when  $\nu$  is odd. The non-trivial spin of the defect operator is likely the reason for this modification of Spin<sup>c</sup>-structures<sup>6</sup>.

Various other methods have been developed for the computation of  $F_K$ -series [35, 36, 77, 78], with each having its advantages. To appreciate the necessity of having other methods, note that the Laplace transforms (2.8) generate q-series with positive powers of the expansion parameter q only when p/r < 0. In what follows we will focus on a method based on the so-called "inverted" Habiro series of the knot K, which unlike (2.8) can in some situations be applied to the p/r > 0 surgeries. This will be especially important when we compute  $\hat{Z}$ -invariants of the mock type in the next section. In the remainder of the subsection, we will focus on integral surgeries where r = 1.

#### **Inverted Habiro Series and Defects**

First recall that for any knot K, there exists a sequence of Laurent polynomials  $a_m(K) \in \mathbb{Z}[q, q^{-1}]$ , which we will refer to as the Habiro coefficients of K, such that the coloured Jones polynomial  $J_K(V_n)$  for the *n*-dimensional irreducible representation  $V_n$  of  $\mathfrak{sl}_2$  can be decomposed as [79]

$$J_K(V_n) = \sum_{m=0}^{\infty} a_m(K;q) \left( \prod_{j=1}^m \left( x + x^{-1} - q^j - q^{-j} \right) \right) \Big|_{x=q^n} .$$
(2.10)

Following this, in [36] it was conjectured that for any knot K with Alexander polynomial  $\Delta_K \neq 1$ , there exist inverted Habiro series with coefficients  $a_{-m}(K;q)$ , such that

$$F_K(x,\tau) = -\left(x^{\frac{1}{2}} - x^{-\frac{1}{2}}\right) \sum_{m=1}^{\infty} \frac{a_{-m}(K;q)}{\prod_{j=0}^{m-1} (x + x^{-1} - q^j - q^{-j})} .$$
 (2.11)

Furthermore, by writing  $F_K(x;\tau) = x^{1/2} \sum_{j=0}^{\infty} f_j(K;q) x^j$ , the inverted Habiro coefficients can be extracted from the two-variable series  $F_K(x,q)$  using the identities [36]

$$f_{j}(K;q) = \sum_{i=0}^{j} \begin{bmatrix} j+i\\ 2i \end{bmatrix} a_{-i-1}(K;q)$$

$$\Leftrightarrow a_{-i-1}(K;q) = \sum_{j=0}^{i} (-1)^{i+j} \begin{bmatrix} 2i\\ i-j \end{bmatrix} \frac{[2j+1]}{[i+j+1]} f_{j}(K;q) ,$$
(2.12)

where we employ the notation

$$[n] = \frac{q^{\frac{n}{2}} - q^{-\frac{n}{2}}}{q^{\frac{1}{2}} - q^{-\frac{1}{2}}}, \qquad [n]! = \prod_{k=1}^{n} [k], \qquad \begin{bmatrix} n\\k \end{bmatrix} = \frac{[n]!}{[k]![n-k]!}. \tag{2.13}$$

<sup>&</sup>lt;sup>6</sup>We thank Mrunmay Jagadale for suggesting the interpretation.

The form (2.11) is particularly useful in the context of  $\pm p$  surgeries. First, when combined with the surgery formula (2.7) for  $p \in \mathbb{Z}_{>0}$  it allows to write the corresponding  $\hat{Z}$ -invariant in the form

$$\hat{Z}_{b}(S^{3}_{-p}(K);\tau) = \epsilon q^{d} \mathcal{L}^{(b)}_{-p} \left[ \left( x^{\frac{1}{2}} - x^{-\frac{1}{2}} \right)^{2} \sum_{m=1}^{\infty} \frac{a_{-m}(K;q)}{\prod_{j=0}^{m-1} (x + x^{-1} - q^{j} - q^{-j})} \right]$$

$$= \epsilon q^{d} \mathcal{L}^{(b)}_{-p} \left[ a_{-1}(K;q) + \sum_{n=1}^{\infty} \frac{a_{-n-1}(K;q)}{\prod_{j=1}^{n} (x + x^{-1} - q^{j} - q^{-j})} \right].$$

$$(2.14)$$

The Habiro coefficients  $a_{-n-1}(K;q)$  are independent of x, so the Laplace transform (2.8) only acts on the product

$$\frac{1}{\prod_{j=1}^{n} (x + x^{-1} - q^{j} - q^{-j})}$$
 (2.15)

Take the (-1)-surgery for instance: the relevant expression is

$$\mathcal{L}_{-1}^{(0)} \left( \frac{1}{\prod_{j=1}^{n} (x + x^{-1} - q^{j} - q^{-j})} \right) = \left( \frac{1}{\prod_{j=1}^{n} (x + x^{-1} - q^{j} - q^{-j})} \right) \Big|_{x^{u} \mapsto q^{u^{2}}} = \frac{q^{n^{2}}}{(q^{n+1};q)_{n}}$$
(2.16)

where  $(x;q)_n = \prod_{i=0}^{n-1} (1-xq^i)$  denotes the q-Pochhammer symbol. Now, the corresponding expression for (-p)-surgery can be written as

$$\mathcal{L}_{-p}^{(b)}\left[\frac{1}{\prod_{j=1}^{n}(x+x^{-1}-q^{j}-q^{-j})}\right] = \frac{q^{n^{2}}}{(q^{n+1};q)_{n}}q^{-\frac{b(2p-b)}{4p}}P_{n}^{p,b}(q^{-1}) , \qquad p \in \mathbb{Z}_{+} .$$
(2.17)

for b even. It turns out that  $P_n^{p,b}$  defined above is a finite polynomial for any  $n \in \mathbb{Z}_+$ . When b is odd, the left-hand side of (2.17) vanishes identically. We record the first few polynomials  $P_n^{p,b}$  in Appendix B.

For (-p)-surgery with  $p \in \mathbb{Z}_+$ , (2.14) and (2.17) gives

$$\hat{Z}_b(S^3_{-p}(K);\tau) = \epsilon q^{d - \frac{b(2p-b)}{4p}} \sum_{n=0}^{\infty} a_{-n-1}(K;q) \frac{q^{n^2}}{(q^{n+1};q)_n} P_n^{p,b}(q^{-1}) .$$
(2.18)

For positive surgeries, taking  $F_K(x,\tau)$  as a series expansion in x and q and directly applying the Laplace transform (2.8) does not generate  $\hat{Z}$ -invariants with convergent qseries in general, as there may be unbounded negative powers of q. The proposal of [36] is to obtain the result of a (+p)-surgery formula by taking q to  $q^{-1}$  in a very specific way that we will now describe. First, assuming that the only factor in the summand of (2.18) that is expanded to be an infinite q-series, namely the factor  $\frac{q^{n^2}}{(q^{n+1};q)_n}$ , is extended outside the unit circle via the q-hypergeometric expression

$$\frac{1}{(q^{-a};q^{-1})_n} = (-1)^n \, \frac{q^{an + \frac{n(n-1)}{2}}}{(q^a;q)_n},\tag{2.19}$$

where we have a = n + 1 for the case of (2.18). Then, the rest of the factors in (2.18) are given by a finite polynomial in q or  $q^{-1}$  and can be transformed in a straightforward way, leading to

$$\mathcal{L}_{+p}^{(b)}\left[\frac{1}{\prod_{j=1}^{n}(x+x^{-1}-q^{j}-q^{-j})}\right] = q^{\frac{b(2p-b)}{4p}}\frac{(-1)^{n}q^{\frac{n(n+1)}{2}}}{(q^{n+1};q)_{n}}P_{n}^{p,b}(q), \ p \in \mathbb{Z}_{+} \ , \ b \ \text{even},$$
(2.20)

which gives<sup>7</sup>

$$\hat{Z}_b(S^3_{+p}(K);\tau) = \epsilon q^{-d + \frac{b(2p-b)}{4p}} \sum_{n=0}^{\infty} a_{-n-1}(K;q) \frac{(-1)^n q^{\frac{n(n+1)}{2}}}{(q^{n+1};q)_n} P_n^{p,b}(q)$$
(2.21)

for even b.

After reviewing conjecture (2.21) for (+p)-surgeries, we discuss how to combine the inverted Habiro description for the  $F_K$  series with the inclusion of defect operators, in accordance with Definition 2. Note that the proposed Wilson line expression (2.9) combined with the expression (2.11) for the  $F_K$ -series leads, for the case of plumbed knot K, to

$$\hat{Z}_{b}(S^{3}_{\pm p}(K); W_{\nu}; \tau) = \epsilon q^{d} \left( a_{-1}(K; q) \mathcal{L}^{(b+\nu)}_{\pm p} \left[ x^{\frac{\nu}{2}} + x^{\frac{\nu}{2}-1} + \dots + x^{-\frac{\nu}{2}} \right] + \sum_{m=1}^{\infty} \mathcal{L}^{(b+\nu)}_{\pm p} \left[ \left( x^{\frac{\nu}{2}} + x^{\frac{\nu}{2}-1} + \dots + x^{-\frac{\nu}{2}} \right) \frac{a_{-m-1}(K; q)}{\prod_{j=1}^{m} (x + x^{-1} - q^{j} - q^{-j})} \right] \right) . \quad (2.22)$$

Below we will work out their explicit expression. For that, first we note the following lemma, relevant for defects with highest weights that are not roots, which can be proven in a way analogous to Proposition 4.1.5. of [80], using the qZeil algorithm [81].

Lemma 1. The following identity holds

$$\frac{x^{1/2} + x^{-1/2}}{\prod_{j=1}^{n} (x + x^{-1} - q^j - q^{-j})} \bigg|_{x^u \mapsto q^{u^2}} = \frac{q^{n^2 - n + \frac{1}{4}}}{(q^n; q)_n} , \qquad (2.23)$$

where  $\Big|_{x^u\mapsto q^{u^2}}$  denotes first series expand in x and then substitute  $x^u$  with  $q^{u^2}$ .

From the above lemma, we get

$$\mathcal{L}_{-1}^{(1)}\left(\frac{x^{1/2} + x^{-1/2}}{\prod_{j=1}^{n}(x + x^{-1} - q^{j} - q^{-j})}\right) = \frac{x^{1/2} + x^{-1/2}}{\prod_{j=1}^{n}(x + x^{-1} - q^{j} - q^{-j})}\Big|_{x^{u} \mapsto q^{u^{2}}} = \frac{q^{n^{2} - n + \frac{1}{4}}}{(q^{n}; q)_{n}}.$$
(2.24)

Again define the (+1)-surgery using the q-hypergeometric expression (2.19), and we obtain

$$\mathcal{L}_{1}^{(1)}\left(\frac{x^{1/2} + x^{-1/2}}{\prod_{j=1}^{n}(x + x^{-1} - q^{j} - q^{-j})}\right) := (-1)^{n} \frac{q^{\frac{n^{2} + n}{2} - \frac{1}{4}}}{(q^{n};q)_{n}} .$$
(2.25)

Similarly to (2.17), we define the polynomials P for odd b via

$$\mathcal{L}_{-p}^{(b)}\left[\frac{x^{1/2} + x^{-1/2}}{\prod_{j=1}^{n}(x + x^{-1} - q^{j} - q^{-j})}\right] = q^{-\frac{b(2p-b)}{4p} - \frac{1}{4}} \mathcal{L}_{-1}^{(b)}\left[\frac{x^{1/2} + x^{-1/2}}{\prod_{j=1}^{n}(x + x^{-1} - q^{j} - q^{-j})}\right] P_{n}^{p,b}(q^{-1})$$
(2.26)

for  $p \in \mathbb{Z}_+$  when b is odd. When b is even, the left-hand side of (2.26) vanishes identically. Using the same logic as before, we define the +p-surgery counterpart through

$$\mathcal{L}_{-p}^{(b)}\left[\frac{x^{1/2} + x^{-1/2}}{\prod_{j=1}^{n}(x + x^{-1} - q^j - q^{-j})}\right] = (-1)^n \frac{q^{\frac{n^2 + n}{2}}}{(q^n; q)_n} q^{\frac{b(2p-b)}{4p}} P_n^{p,b}(q), \ p \in \mathbb{Z}_+, \ b \text{ odd.} \ (2.27)$$

<sup>&</sup>lt;sup>7</sup>Note that here we are changing the surgery from -p to +p without changing the orientation of the knot. Following [35], when flipping the orientation of the knot, it is natural to define the series of the mirror knot m(K) as  $F_{m(K)}(x,\tau) = F_K(x,-\tau)$ .

With the above elements, one can write down the defect generalization of the conjectural expression of the +p-surgery formulas of [36].

**Conjecture 3.** When the inverted Habiro series expression (2.18) and (2.21) leads to the correct  $(\pm p)$ -surgery along a plumbed knot K, inserting a defect operator supported on K in the highest representation of  $\mathfrak{sl}_2$  with highest weight  $\nu \vec{\omega}$  leads to

$$\hat{Z}_{b}(S^{3}_{-p}(K);W_{\nu};\tau) = \epsilon q^{d-\frac{b(2p-b)}{4p}} \sum_{n=0}^{\infty} a_{-n-1}(K;q) \sum_{\ell=1}^{n} \lambda_{\ell}^{n,\nu}(q) \frac{q^{\ell^{2}}}{(q^{\ell+1};q)_{\ell}} P_{\ell}^{p,b+\nu}(q^{-1}) + p(\tau)$$
(2.28)

when  $b \equiv \nu \pmod{2}$  and 0 otherwise. Similarly, when the expression (2.21) leads to the correct (+p)-surgery, we have

$$\hat{Z}_{b}(S^{3}_{+p}(K);W_{\nu};\tau) = \epsilon q^{-d+\frac{b(2p-b)}{4p}} \sum_{n=0}^{\infty} a_{-n-1}(K;q) \sum_{\ell=1}^{n} \lambda_{\ell}^{n,\nu}(q) \frac{(-1)^{\ell} q^{\frac{\ell(\ell+1)}{2}}}{(q^{\ell+1};q)_{\ell}} P_{\ell}^{p,b+\nu}(q) + p(-\tau)$$
(2.29)

when  $b \equiv \nu \pmod{2}$  and 0 otherwise, and  $p(\tau)$  is a finite polynomial. In the above,  $\lambda_{\ell}^{n,\nu}(q) = \lambda_{\ell}^{n,\nu}(q^{-1})$  are finite polynomials, defined via

$$\frac{x^n + x^{n-1} + \dots + x^{-n}}{\prod_{j=1}^m (x + x^{-1} - q^j - q^{-j})} = \lambda'^{m,2n} (x + x^{-1}; q) + \sum_{\ell=1}^m \frac{\lambda_\ell^{m,2n}(q)}{\prod_{j=1}^\ell (x + x^{-1} - q^j - q^{-j})} \quad (2.30)$$

for even  $\nu$  and and similarly

$$\frac{x^{n+\frac{1}{2}} + x^{n-\frac{1}{2}} + \dots + x^{-n-\frac{1}{2}}}{\prod_{j=1}^{m} (x+x^{-1}-q^{j}-q^{-j})} = \lambda^{\prime m,2n+1}(x+x^{-1};q) + \sum_{\ell=1}^{m} \frac{\lambda_{\ell}^{m,2n+1}(q)\left(x^{1/2} + x^{-1/2}\right)}{\prod_{j=1}^{\ell} (x+x^{-1}-q^{j}-q^{-j})}$$
(2.31)

for odd  $\nu$ , where  $\lambda'^{m,k}(X;q)$  is the polynomial in X such that

$$\lambda'^{m,k}(x+x^{-1};q)\prod_{j=1}^{m}(x+x^{-1}-q^{j}-q^{-j})-(x^{\frac{k}{2}}+x^{\frac{k}{2}-1}+\cdots+x^{-\frac{k}{2}})=O(x^{m-\frac{1}{2}}).$$
 (2.32)

The  $\lambda$ -polynomials can be computed explicitly using the recursive relation

$$\frac{\left(x+x^{-1}\right)^m}{\prod_{j=1}^n \left(x+x^{-1}-q^j-q^{-j}\right)} = \frac{\left(q^n+q^{-n}\right)\left(x+x^{-1}\right)^{m-1}}{\prod_{j=1}^n \left(x+x^{-1}-q^j-q^{-j}\right)} + \frac{\left(x+x^{-1}\right)^{m-1}}{\prod_{j=1}^{n-1} \left(x+x^{-1}-q^j-q^{-j}\right)} \,. \tag{2.33}$$

- **Remark 3.** The polynomials  $p(\tau)$  in (2.28) and  $p(-\tau)$  in (2.29) can be identified with the result of Laplace transform of the contribution from the  $\lambda'$  polynomials in (2.30) and (2.31). It is easy to see that it is finite since  $\lambda'^{m,n} = 0$  for n < m.
  - As the reader might have noticed, the proposal on the inverted Habiro series and how to extend it to the cases of Laplace transforms for positive integer surgeries is, from the point of view of the q-series, highly experimental. In this paper we focus on how, when working with the cases where the original proposal of [36] does lead to a correct answer, one can extend the proposal to include supersymmetric defect operators. We will concretely test this proposal in specific examples in §5.

- Note that this method can be used to compute  $\hat{Z}$ -invariants both of the "false" as well as of the "mock" kind, the latter also discussed in §3.
- We find that a similar expression holds for a set of modified  $\lambda$ -polynomials, which leads to interesting results in the case of the figure 8 knot, despite the fact that it is not a plumbed knot and we do not expect Conjecture 3 to apply in that case. See §5.4 for details.

### 2.3 Modularity

After proposing the defect invariants in Definitions 1 and 2, we are ready to discuss the content of the Modularity Conjecture 4, which is really a concrete special case of the general expectation about vector-valued quantum modularity discussed in §1. We will first start with a detailed description of the Weil representation  $\Theta^{m+K}$  featured in the conjecture.

### Weil representations

The  $\hat{Z}$ -invariants have been observed [30] to be related to certain Weil representations  $\Theta^{m+K}$  of the metaplectic group  $\widetilde{\operatorname{SL}}_2(\mathbb{Z})$ , which are subrepresentations of the 2m-dimensional representation  $\Theta_m$  spanned by the column vector  $\theta_m = (\theta_{m,r})_{r \mod 2m}$  with  $m \in \mathbb{Z}_{>1}$  and theta function components (1.1). Derivatives of (1.1) define unary theta functions

$$\theta_{m,r}^{\ell}(\tau) := \left( \left( \frac{1}{2\pi i} \frac{\partial}{\partial z} \right)^{\ell} \theta_{m,r}(\tau, z) \right) \bigg|_{z=0} , \qquad \ell = 0, 1 , \qquad (2.34)$$

with  $\tilde{\theta}_{m,r}(\tau)$  in equation (1.6) proportional to the Eichler integral (1.7) of  $\theta_{m,r}^1(q)$ .

To define the Weil representation  $\Theta^{m+K}$ , one starts with  $\operatorname{Ex}_m$ , the group of exact divisors of m, where a divisor n of m is exact if (n, m/n) = 1 and the group multiplication for  $\operatorname{Ex}_m$  is  $n * n' := nn'/(n, n')^2$ . For a subgroup  $K \subset \operatorname{Ex}_m$  one can construct in the following way a subrepresentation of  $\Theta_m$  denoted  $\Theta^{m+K}$ .

Consider the space of matrices commuting with the S- and T-matrices on  $\Theta_m$ . It is generated by the  $\Omega$ -matrices,  $\Omega_m(n)$  with n|m [82], defined by

$$\Omega_m(n)_{r,r'} = \delta_{r+r' \pmod{2n}} \delta_{r-r' \pmod{2m/n}}$$

$$(2.35)$$

and it represents the group  $\operatorname{Ex}_m$ : for n, n'|m and  $n \in \operatorname{Ex}_m$ , one has  $\Omega_m(n)\Omega_m(n') = \Omega_m(n * n')$ . As a result, one can define the projection operators

$$P_m^{\pm}(n) := \left(\mathbf{1}_m \pm \Omega_m(n)\right)/2 , \qquad n \in \operatorname{Ex}_m , \qquad (2.36)$$

satisfying  $(P_m^{\pm}(n))^2 = P_m^{\pm}(n)$ . For subgroups K with  $m \notin K$ , we define the projector

$$P^{m+K} := \left(\prod_{n \in K} P_m^+(n)\right) P_m^-(m) .$$
 (2.37)

With these,  $\Theta^{m+K}$  can explicitly be constructed from  $\Theta_m$  as spanned by  $(\theta_r^{m+K})_{r\in\sigma^{m+K}}$ 

$$\theta_r^{m+K}(\tau, z) := 2^{\frac{|K|}{2}} \sum_{r' \in \mathbb{Z}/2m} P_{r,r'}^{m+K} \theta_{m,r'}(\tau, z) , \qquad (2.38)$$

where  $\sigma^{m+K} \subset \mathbb{Z}/2m$  is the set which labels the linearly independent functions  $\theta_r^{m+K}(\tau, z)$ . When K is maximal, in the sense that  $\operatorname{Ex}_m = K \cup (m * K)$ , and m is square-free, the Weil representation  $\Theta^{m+K}$  is irreducible. If K is maximal but m is not square-free, then the irreducible representation  $\Theta^{m+K, \operatorname{irred}}$  [83, 84] can be constructed by using the projection operators

$$P^{m+K,\text{irred}} := \left(\prod_{n \in K} P_m^+(n)\right) \left(\prod_{f^2 \mid m} (\mathbf{1}_m - \frac{1}{f}\Omega_m(f))\right) P_m^-(m) .$$
(2.39)

Similarly to equation (2.38), we introduce the notation

$$\widetilde{\theta}_{r}^{m+K}(\tau) := 2^{\frac{|K|}{2}} \sum_{r' \in \mathbb{Z}/2m} P_{r,r'}^{m+K} \widetilde{\theta}_{m,r'}(\tau)$$
(2.40)

and similarly for  $\theta_r^{1,m+K}$  (cf. (2.34)). We also define  $\tilde{\theta}_r^{m+K,\text{irr}}$ ,  $\theta_r^{1,m+K,\text{irr}}$ , and  $\sigma^{m+K,\text{irr}}$  analogously using (2.39). It will be in terms of these Weil representations that we phrase a concrete conjecture (Conjecture 4) as a special realization of the general modularity expectation that we outlined in §1.

#### The Modularity Conjecture

**Conjecture 4.** In this conjecture we say two infinite q-series are equivalent  $f_1 \sim f_2$ , if  $f_1 = q^{\Delta}f_2 + q^{\Delta'}p(q)$ , where  $\Delta, \Delta' \in \mathbb{Q}$  and  $p(q) \in \mathbb{C}[q, q^{-1}]$  is a finite polynomial. Consider a Seifert manifold  $M_3$  with three singular fibres. Define

$$\operatorname{span}(\hat{Z}(M_3)) := \operatorname{span}_{\mathbb{C}}\{\hat{Z}_b(M_3, W_{\boldsymbol{\nu}}; \tau), b \in \operatorname{Spin}^c(M_3) | \boldsymbol{\nu} \in \mathbb{N}^3\}$$

and extend the equivalence between infinite q-series to their spans. Then there exists a Weil representation

$$\Theta^{(M_3)} = \Theta^{m+K} \quad \text{or} \quad \Theta^{(M_3)} = \Theta^{m+K,\text{irr}} \tag{2.41}$$

for some positive integer m and a subgroup  $K \subset Ex_m$ , such that the following is true.

- 1. When  $M_3$  is negative,  $\operatorname{span}(\hat{Z}(M_3))$  is equivalent to  $\operatorname{span}_{\mathbb{C}}\{\widetilde{\theta}_r^{(M_3)}|r\in\mathbb{Z}/2m\}$ .
- 2. When  $M_3$  is positive, there is a  $SL_2(\mathbb{Z})$  vector-valued (mixed) mock modular form  $f^{(M_3)} = (f_r^{(M_3)})$  transforming in the dual representation of  $\Theta^{(M_3)}$ , such that  $\operatorname{span}(\hat{Z}(M_3))$  is equivalent to  $\operatorname{span}_{\mathbb{C}}\{f_r^{(M_3)} | r \in \mathbb{Z}/2m\}$ .

One can prove the above conjecture for all Brieskorn spheres.

**Theorem 1.** The Conjecture 4-1. is true when  $M_3$  is  $\Sigma(p_1, p_2, p_3)$ , with  $m = p_1 p_2 p_3$ ,  $K = \{1, p_1 p_2, p_2 p_3, p_1 p_3\}$ . More precisely, we have

$$\hat{Z}(\Sigma(p_1, p_2, p_3); W_{\nu}; \tau) = cq^{\Delta} \tilde{\theta}_{r_{\nu}}^{m+K} + p(q) \quad , \quad r_{\nu} = m - \sum_i (1 + \nu_i) \bar{p}_i \tag{2.42}$$

where p(q) is a (possibly vanishing) finite polynomial in q and  $q^{-1}$  and  $c \in \mathbb{C}$ ,  $\Delta \in \mathbb{Q}$ .

*Proof.* When  $M = \Sigma(p_1, p_2, p_3)$  is a Brieskorn sphere, we necessarily have  $(p_i, p_j) = 1$  when  $i \neq j$ . Define  $m = p_1 p_2 p_3$  and  $\bar{p}_i := m/p_i$ . From the Chinese remainder theorem, it is easy to see that, for any  $r \in \mathbb{Z}/2m$ , the equations

$$r + r' \equiv 0 \pmod{2p_i}, \ r - r' \equiv 0 \pmod{2\bar{p}_i}$$

$$(2.43)$$

have a unique answer  $r' \in \mathbb{Z}/2m$ . Also, note that there exists  $a_1, a \in \mathbb{Z}$  such that  $1 = a_1\bar{p}_1 + ap_1$ . Another use of the Chinese remainder theorem then implies that there exists  $a_2, a_3 \in \mathbb{Z}$  such that  $a = a_2p_3 + a_3p_2$ , and hence there exists  $(a_1, a_2, a_3) \in \mathbb{Z}^3$  such that  $1 = \sum_{i=1}^3 a_i\bar{p}_i$ . As a result, every  $r \in \mathbb{Z}/2m$  admits an expression in terms of  $A_i \in \mathbb{Z}/p_i$  such that  $r = \sum_{i=1}^3 A_i\bar{p}_i$ .

Putting  $m = p_1 p_2 p_3$ ,  $K = \{1, p_1 p_2, p_2 p_3, p_1 p_3\}$  in the definition (2.37) and (2.40) and using the properties of the  $\Omega$ -matrices, one obtains

$$\widetilde{\theta}_{r}^{m+K} = \sum_{r' \in \mathbb{Z}/2m} \left( \mathbf{1} + \sum_{i=1}^{3} \Omega_{m}(\bar{p}_{i}) \right)_{r,r'} \widetilde{\theta}_{r'}^{m+K}.$$
(2.44)

It is easy to check that, when writing  $r = m - \sum_{i=1}^{3} A_i \bar{p}_i$  (which is always possible from the above argument), we have

$$(\Omega_m(\bar{p}_j))_{r,r'} = \delta_{r' - (m - \sum_i (-1)^{1 + \delta_{i,j}} A_i \bar{p}_i) \pmod{2m}},$$
(2.45)

which, when combined with (2.44), gives

$$\widetilde{\theta}_{r}^{m+K} = \sum_{\substack{(\epsilon_{1},\epsilon_{2},\epsilon_{3}) \in (\mathbb{Z}/2)^{\otimes 3} \\ \sum_{j} \epsilon_{j} \equiv 0 \pmod{2}}} \widetilde{\theta}_{m,m-\sum_{i=1}^{3}(-1)^{\epsilon_{i}}A_{i}\bar{p}_{i}}, \quad r = m - \sum_{i=1}^{3} A_{i}\bar{p}_{i} .$$
(2.46)

Finally, applying Definition 1 to the case where  $M_3 = \Sigma(p_1, p_2, p_3)$  and defect operators corresponding to the end nodes are inserted, one obtains (2.42) after a routine calculation. We therefore conclude that by changing the highest weight  $\nu_i \vec{\omega}$  of the defect operator, we can cover all the components of the Weil representation. Conversely, one can associate a Wilson line configuration to any component labelled by  $r \in \sigma^{m+K}$ .

Later in §3.4, we will also show that the Conjecture 4-2. is compatible with our proposal for the mock  $\hat{Z}$ -invariants of the positive plumbed three-manifolds  $-\Sigma(2,3,6r+1)$ , for all  $r \in \mathbb{N}$ .

Given the prominent role of the Weil representation in Conjecture 4, it might be natural to consider combining the  $\hat{Z}_b(W_{\nu})$ -invariants for different choices of b and  $\nu$  into the Jacobi-like functions  $\sum_r \overline{\hat{\theta}_r^{(M_3)}(\tau)} \theta_{m,r}(\tau,z)$  and  $\sum_r f_r^{(M_3)}(\tau) \theta_{m,r}(\tau,z)$ , for the case of Conjecture 4.1 and 4.2 respectively. We will call them quantum Jacobi theta functions, which can be defined in an obvious way by combining the definition of mock Jacobi theta functions and quantum modular forms. To be specific, let us define the following.

**Definition 3.** We say that  $\phi : \mathbb{H} \times \mathbb{C} \to \mathbb{C}$  with  $\phi(\tau, z) := \sum_{r \in \mathbb{Z}/2m} f_r(\tau) \theta_{m,r}(\tau, z)$  (resp.  $\sum_{r \in \mathbb{Z}/2m} \overline{f_r(\tau)} \theta_{m,r}(\tau, z)$ ) is a weakly holomorphic (resp. skew-holomorphic) quantum Jacobi form of index m and weight w + 1/2, if  $f_r$ ,  $r \in \mathbb{Z}/2m$ , are weakly holomorphic functions on  $\mathbb{H}$ , and they form a vector-valued quantum modular form  $(f_r)_{r \in \mathbb{Z}/2m}$  of weight w for  $\Gamma \subset SL_2(\mathbb{Z})$ , whose character is the dual of that of  $(\theta_{m,r})_{r \in \mathbb{Z}/2m}$  (resp. that of  $(\theta_{m,r})_{r \in \mathbb{Z}/2m}$ ). With the above definition, we can write down a natural corollary for Conjecture 4.

**Proposition 2.** Use the same definition of  $\operatorname{span}(\hat{Z}(M_3))$  and the same equivalence relation. Assuming Conjecture 4 is correct. Then the following is true.

1. When  $M_3$  is negative, there exists a skew-holomorphic quantum Jacobi form

$$\phi(\tau, z) := \sum_{r \in \mathbb{Z}/2m} \overline{f_r(\tau)} \theta_{m,r}(\tau, z)$$

of index m such that  $\operatorname{span}(\hat{Z}(M_3)) = \operatorname{span}_{\mathbb{C}}\{f_r(\tau), r \in \mathbb{Z}/2m\}.$ 

2. When  $M_3$  is positive, there exists a weakly holomorphic quantum Jacobi form

$$\phi(\tau, z) := \sum_{r \in \mathbb{Z}/2m} f_r(\tau) \theta_{m,r}(\tau, z)$$

of index m such that  $\operatorname{span}(\hat{Z}(M_3)) = \operatorname{span}_{\mathbb{C}}\{f_r(\tau), r \in \mathbb{Z}/2m\}.$ 

- **Remark 4.** Here we see the importance of the defect  $\hat{Z}$ -invariants for the understanding of the modularity of  $\hat{Z}$ -invariants. Without including the defect operators, in [30] it was noted that the space spanned by  $\hat{Z}_b(M_3)$  correspond to subspaces of the Weil representation that do not close under the action of the modular group  $\widetilde{SL}_2(\mathbb{Z})$ . The space gets completed once the appropriate defect operators are included.
  - It should also be possible to give a similar proof as that of Theorem 1 using the proposal (Definition 2) for the defect knot surgery for the Brieskorn spheres arising from performing surgeries on torus knots. In this work we only provide examples in §5.2 and leave the general cases as an exercise for the readers.
  - It is easy to see why there are, up to the addition of finite polynomials, finitely many  $\hat{Z}_b(M_3; W_{\nu})$  even though  $\nu \in \mathbb{N}^3$  in principle. From the expression of the contour integration (2.4) and its result (2.42), we see that the equivalence classes (up to the addition of finite polynomials to  $\hat{Z}(W_{\nu})$ ) of Wilson lines are given by  $\nu_i \in \mathbb{Z}/p_i$ , with the finite polynomials in the form of

$$\sum_{\ell \equiv r \pmod{2m}} \operatorname{sgn}(\ell) q^{\ell^2/4m} - \sum_{\ell \equiv r \pmod{2m}} \operatorname{sgn}(\ell + 2mN) q^{\ell^2/4m}$$

for some finite integer N.

- All quantum modular forms we encounter here are the so-called strong quantum modular forms in the language of [66]. We expect this to be true for all quantum modular forms related to  $\hat{Z}$ -invariants.
- Our definition of Jacobi-type quantum modular objects (Definition 3) is different from that in [85] and references therein.
- In Proposition 2, note that  $f_r$  for some  $r \in \mathbb{Z}/2m$  vanish when the corresponding Weil representation is  $\Theta^{m+K} \neq \Theta_m$ . Equivalently, the span is by definition unchanged if instead of taking the set to be  $r \in \mathbb{Z}/2m$  one uses  $r \in \sigma^{m+K}$  resp.  $r \in \sigma^{m+K,irr}$ . The same statement also holds for Conjecture 4.

# 3 Mock $\hat{Z}$ -invariants

In this section we discuss the various considerations on constructing the mock  $\hat{Z}$ -invariants, proposing explicit expressions for specific families of three-manifolds, comparing different approaches, and analysing the modular properties in details.

### 3.1 Mock, Pure and Mixed

In the introduction (around (1.9)), we have reviewed the modern definition of mock modular forms. This definition allows for a few natural generalizations including for instance higher-depth mock modular forms. The generalization we would like to highlight here is that of mixed mock modular forms [86]. To emphasize the difference between mock and mixed mock properties, mock modular forms are also sometimes referred to as "pure" mock modular forms. The novel characteristic of mixed mock modular forms is that they allow a more complex structure in their relation to the completion function (the associated non-holomorphic function that transforms like a modular form). For instance, a simple multiplication of a mock modular form by a modular form leads to a function that is no longer a pure mock modular form but is still a mixed mock modular form.

Formally, a mixed mock modular form is the first function of a triplet  $(f, \{h_j\}_{j \in I}, \{g_j\}_{j \in I})$ of finite sets of functions defined on the upper half plane  $\mathbb{H}$ , such that f is weakly holomorphic and the non-holomorphic function  $\hat{f} := f - \sum_j h_j g_j^*$ , called the completion of f, transforms as a modular form of weight k. In our work, we let the functions  $h_j$  be weakly holomorphic and  $g_j$  are restricted to be cusp forms. Moreover,  $h_i$  and  $g_i$  transform modularly with weights  $\ell$  and  $2 - k + \ell$  respectively, for some  $\ell \in \frac{1}{2} + \mathbb{Z}$ . The mixed mock modular forms we will encounter later all have  $\ell = 0$  and  $h_0 = 1$ , while the other  $h_i$ s, given by the ratio of two cusp forms, either have an exponential singularity or vanish at any cusp. Under these assumptions, using (1.10) and the fact that  $g^* - g^* |\gamma|$  is finite, we conclude that the matching (1.11) of the radial limits holds analogously:

$$f_{u,\text{fin}}(u+iv) \sim \sum_{n\geq 0} \alpha_u(n)v^n$$
,  $g_0^*(-u+iv) \sim \sum_{n\geq 0} \alpha_u(n)(-v)^n$ , (3.1)

with  $f_{u,\text{fin}}$  now given by f with the exponentially divergent part of  $\hat{f}$  – including the contributions from f and from  $h_j g_j^*$ <sup>8</sup> – removed.

### 3.2 Going to the Other Side Using Appell-Lerch Sums

In this subsection, we take a brief detour to discuss a uniform and canonical way to associate a mock modular form to a false theta function  $\tilde{\theta}_{m,r}$ , for all pairs (m,r). This should highlight how tricky the problem of going to the other side is, and in particular that mock modularity alone is not sufficient to fix a unique answer.

Given a false theta function  $\hat{\theta}_{m,r}(\tau)$  (1.6), one can canonically associate to it a partner  $\hat{\theta}_{m,r}^-(\tau)$  as a function on  $\mathbb{H}$ , via the Appell-Lerch sums. Moreover, one can show that  $\hat{\theta}_{m,r}^-$  is a mock modular form and is related to the characters of a vertex algebra [25], constructed as a cone algebra reviewed in §4. Recall that for Seifert manifolds  $M_3$  with three singular fibres, those which are weakly negative have  $\hat{Z}(M_3; W_{\nu})$  invariants that are, up to an overall rational power of q and possibly the addition of a finite q-polynomial,

<sup>&</sup>lt;sup>8</sup>Note that the exponentially divergent terms might have non-trivial  $\bar{\tau}$  dependence in this case.

linear combinations of  $\tilde{\theta}_{m,r}$  for some fixed m [87]. From the false-mock Conjecture 1 and from the above, it might seem that one can now easily compute the corresponding  $\hat{Z}(W_{\nu})$ invariants for the orientation-reversed manifold  $-M_3$  by simply replacing each  $\tilde{\theta}_{m,r}$  with  $\tilde{\theta}_{m,r}^-$  (and flipping  $q \leftrightarrow q^{-1}$  in the overall q-power and the finite polynomial). However, it is rather easy to see that this canonical and uniform method is incompatible with many of the concrete examples computed using various methods in this paper and in [35, 36].

To explain it, recall that by going through the relation to Fine's q-hypergeometric series and the universal mock theta function one arrives at the following partner of the false theta function  $\tilde{\theta}_{m,r}(\tau)$  for any positive integer m and 0 < r < 2m [85]:

$$\tilde{\theta}_{m,r}^{-}(\tau) = -\frac{\eta (2m\tau)}{\eta^2 (4m\tau)} q^{\frac{m}{4}} \times \left[ q^{-\frac{r^2}{4m}} A_2 \left( (r-m)\tau, -2m\tau, 4m\tau \right) - q^{\frac{(r-2m)^2}{4m}} A_2 \left( (m-r)\tau, -2m\tau, 4m\tau \right) \right],$$
(3.2)

where the Appell-Lerch sum is given by

$$A_2(a\tau, -2m\tau; 4m\tau) = q^a \sum_{n \in \mathbb{Z}} \frac{q^{4mn^2 + 2mn}}{1 - q^{a + 4mn}}, \ a \in \mathbb{Q}$$
(3.3)

and is shown to have mock modular properties [88]. A quick computation shows that

$$\tilde{\theta}_{m,r}^{-}(-\tau) = q^{-\frac{r^2}{4m}} \left(1 + O(q)\right) \text{ for } 0 < r < m,$$
(3.4)

which can be extended to all  $r \in \mathbb{Z}$  by the relation  $\tilde{\theta}_{m,r}^- = \tilde{\theta}_{m,r+2m}^- = -\tilde{\theta}_{m,-r}^-$ .

From (3.4) we conclude that the  $\tilde{\theta}_{m,r} \mapsto \tilde{\theta}_{m,r}^-$  swap does not always reproduce the  $\hat{Z}(-M_3)$  that one believes to be correct. For instance, for  $M_3 = \Sigma(2,3,7)$ , the  $\hat{Z}(M_3)$ -invariant is given by

$$\hat{Z}(M_3;\tau) = q^{\frac{83}{168}} \sum_{0 < r < 42} \left(\frac{r}{21}\right) \tilde{\theta}_{42,r}(\tau) = q^{\frac{83}{168}} \left(\tilde{\theta}_{42,1} + \tilde{\theta}_{42,41} - \tilde{\theta}_{42,29} - \tilde{\theta}_{42,13}\right) (\tau).$$
(3.5)

The flipped invariant  $\hat{Z}(-M_3)$ , on the other hand, is expected to be given by the celebrated order 7 mock theta function of Ramanujan [30, 35, 13]:

$$q^{-\frac{1}{168}}F_0(\tau) = q^{-\frac{1}{168}}\sum_{n=0}^{\infty} \frac{q^{n^2}}{(q^{n+1};q)_n}$$
  
=  $-q^{-\frac{1}{168}} \left(1 + q + q^3 + q^4 + q^5 + 2q^7 + q^8 + 2q^9 + q^{10} + 2q^{11} + \mathcal{O}(q^{12})\right)$ .  
(3.6)

It is easy to see that this is not the same as  $\left(\tilde{\theta}_{42,1}^- + \tilde{\theta}_{42,41}^- - \tilde{\theta}_{42,29}^- - \tilde{\theta}_{42,13}^-\right)$ , despite having the same mock modularity and hence essentially the same radial limit<sup>9</sup>, just from the leading behaviour near  $\tau \to i\infty$  using (3.4).

<sup>&</sup>lt;sup>9</sup>Meaning the same radial limit after discarding the exponential singularities (1.11).

### 3.3 Regularised Indefinite Theta Functions

In this subsection we recall the basic construction of indefinite theta functions and their relation to mock modular forms, based on [3] and mostly following the notation of [89, Chapter 8]. We limit our discussion to two-dimensional indefinite lattices that are relevant for the present paper, though the construction is completely analogous for general lattices of signature (1, n). See also [90, 91] for the analysis of general indefinite signature lattices.

For the two-dimensional lattice  $L \cong \mathbb{Z}^2$ , we denote by A the symmetric  $2 \times 2$  matrix with integral entries giving rise to the bilinear form of the lattice, which we will take to have signature (1, 1). One can extend the norm to  $L \otimes_{\mathbb{Z}} \mathbb{R}$  by defining

$$|\mathbf{x}|^2 := \mathbf{x}^T A \mathbf{x} \tag{3.7}$$

for  $x \in \mathbb{R}^2$ . We will also denote the inner product as

$$B(\mathbf{x}, \mathbf{y}) := \mathbf{x}^T A \mathbf{y} \ . \tag{3.8}$$

With these, the set of vectors  $\mathbf{c} \in \mathbb{R}^2$  with negative norm  $|\mathbf{c}|^2 < 0$  splits into two connected components. To specify one component, for a fixed vector  $\mathbf{c}_0$  with  $|\mathbf{c}_0|^2 < 0$ , we denote with

$$C_B(\mathbf{c}_0) := \{ \mathbf{c} \in \mathbb{R}^2 | |\mathbf{c}|^2 < 0, B(\mathbf{c}, \mathbf{c}_0) < 0 \}$$
(3.9)

the component containing  $c_0$ . We also denote with

$$S_B(\mathsf{c}_0) := \{\mathsf{c} = (c_1, c_2) \in \mathbb{Z}^2 | \gcd(c_1, c_2) = 1, |\mathsf{c}|^2 = 0, B(\mathsf{c}, \mathsf{c}_0) < 0 \}$$
(3.10)

the set of integral primitive vectors on the boundary of the component, a set of representatives of the cosets for the cusps of  $C_B(\mathbf{c}_0)/\mathbb{R}_+$  with respect to the orientation-preserving lattice orthogonal group, and  $\bar{C}_B(\mathbf{c}_0) := C_B(\mathbf{c}_0) \cup S_B(\mathbf{c}_0)$  the corresponding compactification. Then, for any  $\mathbf{c}_1, \mathbf{c}_2 \in \bar{C}_B(\mathbf{c}_0) \cap \mathbb{Z}^2$ , one can define the regularization factor as

$$\rho^{\mathbf{c}_1,\mathbf{c}_2}(\mathbf{n}) := \operatorname{sgn}(B(\mathbf{c}_1,\mathbf{n})) - \operatorname{sgn}(B(\mathbf{c}_2,\mathbf{n})) \ . \tag{3.11}$$

It was shown in §2 of [3] that the norm is positive definite when restricted to the support of  $\rho^{c_1,c_2}$ . As a result, for  $a \in \mathcal{R}(c_1) \cap \mathcal{R}(c_2)$ ,  $b \in \mathbb{R}^2$  with

$$\mathcal{R}(\mathsf{c}) := \begin{cases} \mathbb{R}^2 & \text{if } |\mathsf{c}|^2 < 0\\ \{\mathsf{a} \in \mathbb{R}^2 | B(\mathsf{c}, \mathsf{a}) \notin \mathbb{Z}\} & \text{if } |\mathsf{c}|^2 = 0 \end{cases}$$
(3.12)

one can define the regularised indefinite theta function [3]

$$\Theta_{A,\mathbf{a},\mathbf{b},\mathbf{c}_1,\mathbf{c}_2}(\tau) := \sum_{\mathbf{n}\in\mathbf{a}+\mathbb{Z}^2} \rho^{\mathbf{c}_1,\mathbf{c}_2}(\mathbf{n}) \operatorname{e}(\operatorname{B}(\mathbf{n},\mathbf{b})) \operatorname{q}^{\frac{|\mathbf{n}|^2}{2}}, \tag{3.13}$$

which is absolutely convergent on the upper-half plane. It was shown that  $\Theta_{A,a,b,c_1,c_2}$  enjoys a certain mock modular property. In the language of §3.1, it is a mixed mock modular form. Concretely,  $\Theta_{A,a,b,c_1,c_2}$  can be viewed as the holomorphic part of

$$\hat{\Theta}_{A,\mathbf{a},\mathbf{b},\mathbf{c}_1,\mathbf{c}_2}(\tau,\bar{\tau}) := \sum_{\mathbf{n}\in\mathbf{a}+\mathbb{Z}^2} \hat{\rho}^{\mathbf{c}_1,\mathbf{c}_2}(\mathbf{n};\tau) e(B(\mathbf{n},\mathbf{b})) q^{\frac{|\mathbf{n}|^2}{2}},\tag{3.14}$$

where

$$\hat{\rho}^{c_1, c_2}(\mathbf{n}; \tau) := \hat{\rho}^{c_1}(\mathbf{n}; \tau) - \hat{\rho}^{c_2}(\mathbf{n}; \tau)$$

$$\hat{\rho}^{c}(\mathbf{n}; \tau) = \begin{cases} E\left(\frac{B(\mathbf{c}, \mathbf{n})(\operatorname{Im}(\tau))^{\frac{1}{2}}}{\sqrt{-2|c_1|^2}}\right) & \text{if } |\mathbf{c}|^2 < 0 \\ \operatorname{sgn}\left(B(\mathbf{c}, \mathbf{n})\right) & \text{if } |\mathbf{c}|^2 = 0 \end{cases},$$
(3.15)

and the non-holomorphic completion  $\hat{\Theta}_{A,a,b,c_1,c_2}$  transforms like a weight one modular form. In the language of [89, Chapter 8],  $\hat{\Theta}_{A,a,b,c_1,c_2}$  is a weight one mixed harmonic form. In particular, if we consider  $a, b \in L^*$  in the dual lattice, then

$$\hat{\Theta}_{A,\mathsf{a},\mathsf{b},\mathsf{c}_1,\mathsf{c}_2}(-\frac{1}{\tau}) = \frac{\tau e(B(\mathsf{a},\mathsf{b}))}{\sqrt{|\det(A)|}} \sum_{\boldsymbol{\mu} \in L^*} \hat{\Theta}_{A,\boldsymbol{\mu}+\mathsf{b},-\mathsf{a},\mathsf{c}_1,\mathsf{c}_2}(\tau).$$
(3.16)

Its  $\bar{\tau}$ -derivative, specifying the deviation from modularity of the holomorphic function  $\Theta_{A,a,b,c_1,c_2}$ , is given by (see [89, Proposition 8.33])

$$\frac{\overline{\partial}}{\partial \overline{\tau}} (\hat{\Theta}_{A,\mathbf{a},\mathbf{b},\mathbf{c}_{1},\mathbf{c}_{2}}(\tau)) = \frac{1}{i\sqrt{2\operatorname{Im}(\tau)}} \sum_{j=1,2} (-1)^{j} \sum_{\mathbf{l}\in P_{j}} g_{\frac{B(\mathbf{c}_{j},\mathbf{l}+\mathbf{a})}{|\mathbf{c}_{j}|^{2}}, -B(\mathbf{c}_{j},\mathbf{b})} (-|\mathbf{c}_{j}|^{2}\tau) \overline{\sum_{\nu\in(\mathbf{l}+\mathbf{a})_{j}^{\perp}+\langle\mathbf{c}_{j}\rangle_{\mathbb{Z}}^{\perp}}} e(B(\nu,\mathbf{b}))q^{\frac{|\nu|^{2}}{2}}.$$
(3.17)

In the above, we use  $I_j^{\perp} := I - \frac{B(\mathbf{c}_j, \mathbf{l})}{B(\mathbf{c}_j, \mathbf{c}_j)} \mathbf{c}_j$  for the projection of the vector I onto the line that is orthogonal to  $\mathbf{c}_j$  and write

$$\langle \mathbf{c} \rangle_{\mathbb{Z}}^{\perp} := \{ \lambda \in \mathbb{Z}^2 : B(\mathbf{c}, \lambda) = 0 \} .$$
 (3.18)

Denote by  $\mathbf{c}_{j,\perp}$  a generator of  $\langle \mathbf{c}_j \rangle_{\mathbb{Z}}^{\perp}$ , then  $P_j$  can be chosen to be any representative of the quotient  $\mathbb{Z}^2/\mathbb{Z}\mathbf{c}_j \oplus \mathbb{Z}\mathbf{c}_{j,\perp}$  and is in particular independent of  $\mathbf{a}$ .

The function  $g_{\rho,\beta}(\tau)$ 

$$g_{\rho,\beta}(\tau) := \sum_{n \in \rho + \mathbb{Z}} n e^{2\pi i n \beta} q^{\frac{n^2}{2}}, \qquad (3.19)$$

in the case of  $\beta \in \mathbb{Z}$ , is related to a unary theta function as defined in equation (2.34) through

$$g_{\frac{r}{2m},\beta}(\tau) = \sum_{x \in \frac{r}{2m} + \mathbb{Z}} x e(x\beta) q^{\frac{nx^2}{2}} = \frac{e(\frac{r\beta}{2m})}{2m} \theta_{m,r}^1\left(\frac{\tau}{2m}\right) .$$
(3.20)

In this paper we are interested in the special cases where the following additional properties hold:

$$c_j \in \mathbb{Z}^2$$
,  $B(c_j, b) \in \mathbb{Z}$ ,  $-\frac{1}{2} |c_j|^2 = \frac{n_j}{d_j} m$  with  $m \in \mathbb{Z}_+$  for  $j = 1, 2$ , (3.21)

where  $n_j, d_j$  are coprime natural numbers. In later applications, we will choose  $m = -\frac{1}{2}|\mathbf{c}_1|^2$  most of the time. In this case, the shadow simplifies greatly into the following

expression

$$2im\sqrt{2\operatorname{Im}(\tau)} \frac{\partial}{\partial \bar{\tau}} (\hat{\Theta}_{A,\mathsf{a},\mathsf{b},\mathsf{c}_{1},\mathsf{c}_{2}}(\tau)) = \sum_{j=1,2} (-1)^{j} \frac{d_{j}}{n_{j}} \sum_{\mathsf{l}\in P_{j}} \operatorname{e}\left(\frac{d_{j}B(\mathsf{c}_{j},\mathsf{b})B(\mathsf{c}_{j},\mathsf{l}+\mathsf{a})}{2n_{j}m}\right) \\ \times \theta^{1}_{n_{j}m,-d_{j}B(\mathsf{c}_{j},\mathsf{l}+\mathsf{a})} \left(\frac{\tau}{d_{j}}\right) \overline{\sum_{\nu\in(\mathsf{l}+\mathsf{a})_{j}^{\perp}+\mathbb{Z}\mathsf{c}_{j,\perp}} e(B(\nu,\mathsf{b}))q^{\frac{|\nu|^{2}}{2}}} .$$

$$(3.22)$$

# 3.4 Indefinite Theta Functions as Mock $\hat{Z}$ -invariants

We now turn to the second method to construct false-mock pairs, which unlike the inverted Habiro series discussed in  $\S2.2$  is designed to produce topological three-manifold invariants with manifest mock modular properties, making use of the indefinite theta functions discussed in  $\S3.3$ . We will also comment on the relation between the two methods in \$3.5.

For this we start by recalling the proposal for the "false side" Z-invariants for weakly negative three-manifolds in discussed in §2.1. We focus on weakly negative Seifert threemanifolds with three exceptional fibres, corresponding to plumbing graphs with one central node  $v_0$  and three legs. They can be expressed in terms of the Seifert data as  $M(b; \{q_i/p_i\}_{i=1,2,3})$ , which moreover have

$$\mathbf{c} = b + \sum_{i} \frac{q_i}{p_i} < 0. \tag{3.23}$$

Let D be the smallest positive integer such that  $\frac{D}{\mathbf{c}p_i} \in \mathbb{Z}$  for all i, and  $m = -DM_{v_0,v_0}^{-1}$ given by the inverse plumbing matrix. Consider Wilson lines associated to the end nodes of the three legs in the plumbing graph. As before, we label the corresponding highest weight representations with  $\boldsymbol{\nu} = (\nu_1, \nu_2, \nu_3) \in \mathbb{N}^3$  as in §2.3 and denote the corresponding defect  $\hat{Z}$ -invariant with  $\hat{Z}_b(M_3; W_{\boldsymbol{\nu}})$ . The equation (2.4) gives

$$\hat{Z}_b(M_3; W_{\boldsymbol{\nu}}; \tau) = C(q) \sum_{\substack{\hat{\epsilon} = (\epsilon_1, \epsilon_2, \epsilon_3) \\ \in (\mathbb{Z}/2)^{\otimes 3}}} \tilde{\chi}_{\hat{\epsilon}}(\tau) + p_{\hat{\epsilon}}(\tau)$$
(3.24)

after performing the contour integration along all  $z_v$  with  $v \neq v_0$  [37], where  $\tilde{\chi}_{\hat{\epsilon}}$  either vanishes or is given by the false theta function

$$\tilde{\chi}_{\hat{\epsilon}}(\tau) = (-1)^{\hat{\epsilon}} \sum_{\substack{\ell \equiv mv_{\hat{\epsilon}} + a_{\hat{\epsilon}} \\ (\text{mod } 2mD)}} q^{\frac{1}{4Dm}\ell^2} \text{sgn}(\ell)$$
(3.25)

for some  $v_{\hat{\epsilon}}, a_{\hat{\epsilon}} \in \mathbb{Z}$ ,  $p_{\hat{\epsilon}}(\tau)$  a finite polynomial which vanishes when  $0 < a_{\hat{\epsilon}} < m$ , and  $C(q) = cq^{\Delta}$  for some  $c \in \mathbb{C}$  and  $\Delta \in \mathbb{Q}$  is the pre-factor that can be explicitly computed from (2.4) which we will mostly ignore from now on. In what follows, we will often use ~ to denote an equality up to such pre-factors  $cq^{\Delta}$ . We will also use the shorthand notation

$$(-1)^{\hat{\epsilon}} := (-1)^{\sum_i \epsilon_i}.$$

From the arguments reviewed in §1, when mapping  $M_3$  to  $-M_3$  we expect to map  $\tau \to -\tau$ , an action that naively causes (3.25) to cease being a convergent function in  $\mathbb{H}$ . Inspired by the construction of the regularised theta function of indefinite signature

reviewed in §3.3, we propose to regularize the function obtained by  $\tau \to -\tau$  by first turning the sum over a one-dimensional negative-definite lattice sum into a sum over a signature (1, n) lattice, by adding an *n*-dimensional lattice  $\Lambda$  with positive-definite norm. Schematically, we insert into (3.25) the identity

$$1 = \frac{\sum_{v \in \Lambda + \gamma} q^{|v|_{\Lambda}^2/2} \mathbf{e}(\mathbf{B}_{\Lambda}(\mathbf{v}, \rho))}{\theta_{\Lambda, \gamma, \rho}}.$$
(3.26)

Subsequently, we regularize the resulting (1, n) lattice theta function using the method discussed in the previous subsection, restricting the sum over the (1, n) lattice

$$\sum_{\substack{v \in \Lambda + \gamma \\ (\text{mod } 2mD)}} \operatorname{sgn}(\ell) e(B_{\Lambda}(v, \rho)) q^{-\frac{1}{4Dm}\ell^2 + |v|_{\Lambda}^2/2}$$
(3.27)

to a positive-definite cone. To be more concrete, we let n = 1 and consider the lattice  $\sqrt{k\mathbb{Z}}$  with the theta function

$$f_{x,\chi}(\tau) := \sum_{k \in \mathbb{Z}} (-1)^k q^{\frac{x}{2} \left(k - \frac{\chi}{2x}\right)^2} .$$
(3.28)

The restriction to the cone, specified by two negative-norm vectors  $c_1$ ,  $c_2$  as in §3.3, is implemented by including a factor

$$\frac{1}{2}|\operatorname{sgn}(B(\mathsf{c}_1,\mathsf{n})) - \operatorname{sgn}(B(\mathsf{c}_2,\mathsf{n}))|$$
(3.29)

which is equal to one inside either of the upper or the lower components of the cone and zero elsewhere, leading to the restricted lattice sum

$$\tilde{\chi}_{\hat{\epsilon}}(-\tau) := \sum_{\substack{\ell \equiv mv_{\hat{\epsilon}} + a_{\hat{\epsilon}} \\ (\text{mod } 2mD)}} q^{-\frac{1}{4Dm}\ell^2} \operatorname{sgn}(\ell) \\
\times \left( \frac{1}{f_{x,\chi}(\tau)} \sum_{k \in \mathbb{Z}} (-1)^k q^{\frac{x}{2} \left(k - \frac{\chi}{2x}\right)^2} \frac{1}{2} |\operatorname{sgn}(B(\mathsf{c}_1,\mathsf{n})) - \operatorname{sgn}(B(\mathsf{c}_2,\mathsf{n}))| \right)$$
(3.30)

where  $\mathbf{n}^T = (\frac{1}{2mD}\ell, k - \frac{\chi}{2x})$  denotes the vector in the real vector space underlying the two-dimensional lattice with bilinear form (3.7) given by A = diag(-2mD, x), and the factor inside the bracket is the regularizing term. Choosing moreover

$$\mathsf{c}_1 = \begin{pmatrix} 1\\ 0 \end{pmatrix} , \qquad (3.31)$$

we obtain the following expression in terms of the regularized indefinite theta function  $(3.13)^{10}$ 

$$\tilde{\chi}_{\hat{\epsilon}}(-\tau) = \frac{(-1)^{\hat{\epsilon}} \mathbf{e}(\frac{\chi}{4x})}{2f_{x,\chi}(\tau)} \sum_{\mathbf{n}\in\mathbf{a}_{\hat{\epsilon}}+\mathbb{Z}^2} (\operatorname{sgn}(B(\mathbf{c}_1,\mathbf{n})) - \operatorname{sgn}(B(\mathbf{c}_2,\mathbf{n}))) q^{\frac{\mathbf{n}^T A \mathbf{n}}{2}} \mathbf{e}(B(\mathbf{b},\mathbf{n}))$$

$$= \frac{(-1)^{\hat{\epsilon}} \mathbf{e}(\frac{\chi}{4x})}{2f_{x,\chi}(\tau)} \Theta_{A,\mathbf{a}_{\hat{\epsilon}},\mathbf{b},\mathbf{c}_1,\mathbf{c}_2}$$
(3.32)

<sup>10</sup>In all our cases we have  $\chi \neq 0$ ,  $mv_{\hat{\epsilon}} + a_{\hat{\epsilon}} \neq 0$ .

with

$$A = \begin{pmatrix} -2m & 0\\ 0 & x \end{pmatrix}, \ \mathbf{b} = \begin{pmatrix} 0\\ \frac{1}{2x} \end{pmatrix} , \tag{3.33}$$

which follows from our choice of the theta function (3.28), and  $\mathbf{a}_{\hat{\epsilon}}^T = (\frac{mv_{\hat{\epsilon}} + a_{\hat{\epsilon}}}{2mD}, -\frac{\chi}{2x})$ , data that can be read off on the false side (3.25). As a result, for those manifolds  $M_3$  to which the above regularization applies, we can propose

$$\hat{Z}_b(-M_3; W_{\boldsymbol{\nu}}; \tau) = C(q^{-1}) \frac{\mathrm{e}(\frac{\chi}{4x})}{2f_{x,\chi}(\tau)} \sum_{\substack{\hat{\epsilon} = (\epsilon_1, \epsilon_2, \epsilon_3)\\ \in (\mathbb{Z}/2)^{\otimes 3}}} \chi_{\hat{\epsilon}}(-\tau) + p_{\hat{\epsilon}}(-\tau)$$
(3.34)

with  $\chi_{\hat{\epsilon}}(-\tau) = 0$  for those  $\hat{\epsilon}$  with  $\tilde{\chi}_{\hat{\epsilon}} = 0$  and

$$\chi_{\hat{\epsilon}}(-\tau) := (-1)^{\hat{\epsilon}} \frac{\mathrm{e}(\frac{\chi}{4x})}{2f_{x,\chi}(\tau)} \Theta_{A,\mathbf{a}_{\hat{\epsilon}},\mathbf{b},\mathbf{c}_{1},\mathbf{c}_{2}}(\tau)$$
(3.35)

otherwise. For instance, when  $M_3 = \Sigma(p_1, p_2, p_3)$  is a Brieskorn sphere, we have D = 1,  $m = p_1 p_2 p_3$ , and (3.35) holds for all  $\hat{\epsilon} \in (\mathbb{Z}/2)^{\otimes 3}$ . See [37] for details.

From the discussion in the previous subsection, the function (3.34) enjoys a manifest relation to mixed mock modular forms as it is given by indefinite theta functions. To completely specify the function, it remains to specify the second negative-norm vector  $\mathbf{c}_2$  and  $\chi \in \mathbb{Z}$ .

For instance, it is known that the  $\hat{Z}$ -invariant for the homological sphere  $-\Sigma(2,3,7)$ , predicted to be given by Ramanujan's order seven mock theta function in [30] and computed to the leading orders using knot surgery formula in [35], admits expressions in terms of indefinite theta functions [13, 36]. Concretely, for  $M_3 = \Sigma(2,3,7)$  we have for the defect  $\hat{Z}$ -invariants (3.24)

$$\tilde{\chi}_{\hat{\epsilon}}(\tau) = (-1)^{\sum_{i} \epsilon_{i}} \tilde{\theta}_{m, r_{\nu, \hat{\epsilon}}}, \quad r_{\nu, \hat{\epsilon}} = m - \sum_{i=1}^{3} (-1)^{\epsilon_{i}} (1 + \nu_{i}) \bar{p}_{i}.$$
(3.36)

See the proof of Theorem 1 for details. For  $M_3 = -\Sigma(2,3,7)$ , we take

$$\chi = 1, \ x = 3, \ \mathsf{c}_2 = \begin{pmatrix} 3\\ 14 \end{pmatrix}$$
 (3.37)

and obtain

$$\hat{Z}_{b_0}(-M_3; W_{\boldsymbol{\nu}}; \tau) = C(q^{-1}) \frac{\mathrm{e}(\frac{\chi}{4x})}{2f_{x,\chi}(\tau)} \sum_{\substack{\hat{\epsilon} = (\epsilon_1, \epsilon_2, \epsilon_3) \\ \in (\mathbb{Z}/2)^{\otimes 3}}} (-1)^{\hat{\epsilon}} \Theta_{A, \mathbf{a}_{\boldsymbol{\nu}, \hat{\epsilon}}, \mathbf{b}, \mathbf{c}_1, \mathbf{c}_2}(\tau)$$
(3.38)

with  $\mathbf{a}_{\boldsymbol{\nu},\hat{\epsilon}}^T = (\frac{r_{\boldsymbol{\nu},\hat{\epsilon}}}{2m}, -\frac{\chi}{2x})$ . Putting the above proposal for A,  $\mathbf{a}_{\boldsymbol{\nu},\hat{\epsilon}}$ ,  $\mathbf{c}_1$ ,  $\mathbf{c}_2$ ,  $\mathbf{b}$ , and x,  $\chi$  together, one can show that the above expression with no defects ( $\boldsymbol{\nu} = (0, 0, 0)$ ) is given, up to a prefactor, by the order seven mock theta function  $F_0$ :

$$\hat{Z}_{b_0}(-M_3;\tau) \sim \frac{1}{2f_{x,\chi}(\tau)} \sum_{\substack{\hat{\epsilon} = (\epsilon_1, \epsilon_2, \epsilon_3) \\ \in (\mathbb{Z}/2)^{\otimes}3}} (-1)^{\hat{\epsilon}} \Theta_{A,\mathbf{a}_{\hat{\epsilon}},\mathbf{b},\mathbf{c}_1,\mathbf{c}_2}(\tau) 
= H_1^{42+6,14,21}(\tau) = q^{-1/168} F_0(q) = q^{-1/168} \sum_{n \ge 0} \frac{q^{n^2}}{(q^{n+1};q)_n},$$
(3.39)

where  $H_1^{42+6,14,21}(\tau)$  is the mock modular form that is a component of the corresponding optimal mock Jacobi form [92], and is one of Ramanujan's order 7 mock theta function,  $F_0$ , up to a pre-factor  $q^{-1/168}$ . A complete analysis for  $M_3 = \pm \Sigma(2,3,7)$  can be found in §5.2.

This is the first case of our proposal for the mock  $\hat{Z}$ -invariants for the following oneparameter family of three-manifolds.

**Conjecture 5.** Consider the orientation-reversed Brieskorn sphere  $-\Sigma(2, 3, 6r + 1)$ . The defect  $\hat{Z}$ -invariants are given by

$$\hat{Z}_{b_0}(-\Sigma(2,3,6r+1);W_{\boldsymbol{\nu}};\tau) = C(q^{-1}) \frac{\mathrm{e}(\frac{\chi}{4x})}{2f_{x,\chi}(\tau)} \sum_{\substack{\hat{\epsilon} = (\epsilon_1,\epsilon_2,\epsilon_3)\\ \in (\mathbb{Z}/2)^{\otimes 3}}} (-1)^{\hat{\epsilon}} \Theta_{A,\mathbf{a}_{\boldsymbol{\nu},\hat{\epsilon}},\mathbf{b},\mathbf{c}_1,\mathbf{c}_2}(\tau) \quad (3.40)$$

with

$$A = \begin{pmatrix} -2m & 0\\ 0 & x \end{pmatrix}, \ \mathbf{c}_1 = \begin{pmatrix} 1\\ 0 \end{pmatrix}, \ \mathbf{c}_2 = \begin{pmatrix} x\\ 2(6r+1) \end{pmatrix}, \ \mathbf{b} = \begin{pmatrix} 0\\ \frac{1}{2x} \end{pmatrix}, \ \mathbf{a}_{\boldsymbol{\nu},\hat{\epsilon}} = \begin{pmatrix} \frac{r_{\boldsymbol{\nu},\hat{\epsilon}}}{2m}, -\frac{\chi}{2x} \end{pmatrix}$$
(3.41)

where we take x = 2r + 1,  $\chi = 1$ , and we have used the same notation as in the proof of Theorem 1 with the triplet given by  $(p_1, p_2, p_3) = (2, 3, 6r + 1)$ .

Moreover, one can show that they are indeed components of vector-valued mock modular forms for  $SL_2(\mathbb{Z})$  with shadows that are given by the unary theta function, in accordance with the false-mock conjecture.

**Theorem 2.** The conjectural expression (3.40) for the  $\hat{Z}$ -invariants has the property that  $\hat{Z}_{b_0}(-\Sigma(2,3,6r+1);W_{\boldsymbol{\nu}};\tau)$  equals, up to a pre-factor and the addition of a finite polynomial, a mock modular form  $f_{m,r_{\boldsymbol{\nu}}}^{m+K}$  with shadow given by  $\theta_{r_{\boldsymbol{\nu}}}^{1,m+K}$ , where  $r_{\boldsymbol{\nu}} = m - \sum_{i}(1 + \nu_i)\bar{p}_i$ .

**Corollary 1.** The mock modular property of the function in (3.40) is compatible with Conjecture 4.2.

*Proof.* The statement follows from Theorem 2 in combination with the proof of Theorem 1.  $\hfill \square$ 

We refer to Appendix A for the proof of this theorem.

One might wonder about the origin and interpretation of the conjectural expression in Conjecture 5, in particular the choice of the parameters (3.41). Though we do not offer a complete answer to this interesting and important question, we first note in which sense these parameters seem natural to us.

First, the choice of the one-dimensional theta function  $f_{x,\chi}$  is a natural generalization of the Dedekind eta function  $\eta(\tau) = f_{3,1}(\tau) = q^{1/24} \prod_i (1-q^i)$  used in [13], and this choice also determines **b** and the bilinear form A. The choice  $\chi = \pm 1$  guarantees the simple modular transformation (3.16) under the full modular group  $SL_2(\mathbb{Z})$ , as expected on physical grounds. The choice for  $c_1$  is also natural given the sign factor in the false theta function (3.27) for the case of the Seifert manifolds with three singular fibres: it guarantees that the sign factor and the cone support factor (3.29) combines nicely into the regularization factor in indefinite theta functions. The choice of  $c_2$ , which together with  $c_1$  specifies that the positive component of the cone which we sum over in the indefinite lattice is positively generated by the vectors (cf. (4.1))

$$d_1 = c_{1,\perp} = \begin{pmatrix} 0\\1 \end{pmatrix}, \ d_2 = c_{2,\perp} = \begin{pmatrix} 1\\\bar{p}_3 \end{pmatrix},$$
 (3.42)

is the more mysterious choice. In practice, it is found by requiring the resulting mock forms to have the desired mock modular property. See the modularity Theorem 2.

### 3.5 Relation to Surgeries Along Torus Knots

In this subsection we will explain the following fact: from the point of view of knot surgeries, one in fact arrives at the expression in Conjecture 5, for the case without defect line insertions, starting from the conjectural regularized +1/r-surgery formula proposed earlier (Conjecture 5, [36]).

First, recall that, a -1/r-surgery (2.7) along the torus knot T(s,t) leads to the Brieskorn sphere  $S^3_{-1/r}(T(s,t)) = \Sigma(s,t,str+1)$ . Similarly, considering the mirror of the torus knot m(T(s,t)) = T(s,-t), its +1/r-surgery gives the orientation-reversed manifold

$$S_{\pm 1/r}^3(T(s,-t)) = -\Sigma(s,t,str+1).$$
(3.43)

Consider the right-handed trefoil  $\mathbf{3}_1^r = T(2,3)$  and its mirror  $\mathbf{3}_1^l = T(2,-3)$ . From

$$F_{T(s,t)}(x,q) = -q^{\frac{(s-1)(t-1)}{2}} \sum_{k>0} \varepsilon_k q^{\frac{k^2 - (st-s-t)^2}{4st}} (x^{k/2} - x^{-k/2})$$
(3.44)

where  $\varepsilon_k = \sum_{\epsilon \in \mathbb{Z}/2} (-1)^{\epsilon} (\delta_{k+(st+(-1)^{\epsilon}s+t) \pmod{2st}} + \delta_{k-(st+(-1)^{\epsilon}s+t) \pmod{2st}})$ , for the torus knot K = T(s,t), we obtain via  $F_{m(K)}(x,q) = F_K(x,q^{-1})$  the two-variable series for the negative torus knot m(K) = T(s,-t). Now, applying Conjecture 5 of [36] to this case leads to the prediction

$$\tilde{Z}_{0}(S_{1/r}^{3}(T(s,-t));\tau) = \tilde{Z}_{0}(-\Sigma(s,t,str+1);\tau) \\
= C(q) \sum_{k\geq 0} \varepsilon_{k} q^{-\frac{k^{2}-(st-s-t)^{2}}{4st}} \sum_{\epsilon_{2}\in\mathbb{Z}/2} (-1)^{\epsilon_{2}} q^{-\frac{r}{4}(k-(-1)^{\epsilon_{2}}\frac{1}{r})^{2}} \left( \frac{\sum_{\substack{\ell\in\mathbb{Z}\\|\ell|<\frac{k-1}{2}}} (-1)^{\ell} q^{\frac{2r+1}{2}(\ell+\frac{1}{2(2r+1)})^{2}}}{\sum_{\ell\in\mathbb{Z}} (-1)^{\ell} q^{\frac{2r+1}{2}(\ell+\frac{1}{2(2r+1)})^{2}}} \right)$$

$$(3.45)$$

By carefully analysing the region  $|\ell| < \frac{k-1}{2}$  one can show that the above expression is equivalent to (3.40) for the trefoil knot T(s, -t) = T(2, -3), in the case where no Wilson line is inserted.

Now, it is straightforward to generalize the above analysis to the general torus knot T(s, t), and further generalize it to include defect operators in the way discussed in §2. The same manipulation leads to an expression that is totally analogous to that of Conjecture 5 (for the case of no defect); it suggests

$$\hat{Z}_{b_0}(-\Sigma(p_1, p_2, p_3); W_{\boldsymbol{\nu}}; \tau) \sim \frac{\mathrm{e}(\frac{\chi}{4x})}{2f_{x,\chi}(\tau)} \sum_{\substack{\hat{\epsilon} = (\epsilon_1, \epsilon_2, \epsilon_3) \\ \in (\mathbb{Z}/2)^{\otimes 3}}} (-1)^{\hat{\epsilon}} \Theta_{A, \mathbf{a}_{\hat{\epsilon}}, \mathbf{b}, \mathbf{c}_1, \mathbf{c}_2}(\tau)$$
(3.46)

with  $(p_1, p_2, p_3) = (s, t, str + 1)$  and

$$A = \begin{pmatrix} -2m & 0\\ 0 & x \end{pmatrix}, \ \mathsf{c}_1 = \begin{pmatrix} 1\\ 0 \end{pmatrix}, \ \mathsf{c}_2 = \begin{pmatrix} x\\ 2p_3 \end{pmatrix}, \ \mathsf{b} = \begin{pmatrix} 0\\ \frac{1}{2x} \end{pmatrix}, \ \mathsf{a}_{\boldsymbol{\nu},\hat{\epsilon}} = \begin{pmatrix} \frac{r_{\boldsymbol{\nu},\hat{\epsilon}}}{2m}\\ -\frac{\chi}{2x} \end{pmatrix}$$
(3.47)

where  $r_{\nu,\hat{\epsilon}} = m - \sum_i (-1)^{\epsilon_i} (1 + \nu_i) \bar{p}_i$  as before, and we take x = 2r + 1,  $\chi = 1$ . Also, we have used the same notation as in the proof of Theorem 1.

Despite the complete analogy between the T(2,3) case and the general torus knot cases, it turns out that the mock modular properties of the corresponding indefinite theta functions are different: apart from sporadic examples (see Appendix A), only in the case of T(2,3) the resulting indefinite theta functions are pure mock modular forms. In the other cases, generically they are mixed mock modular forms. Nevertheless, it is possible to show that the leaking asymptotic behaviour (3.1) still holds in this case, with  $g_0$  being given by the unary theta function  $\theta_{r_{\nu}}^{1,m+K}$ , and the Weil representation  $\Theta^{m+K}$  as in Theorem 1. This is because, in the notation of §3.1, the modular functions  $h_i$ ,  $i \neq 0$  are given by one-dimensional theta functions divided by  $f_{x,\chi}(\tau)$  and can be shown to either vanish or have exponential singularities near any cusp.

**Proposition 3.** The function

$$\frac{\mathrm{e}(\frac{\Lambda}{4x})}{2f_{x,\chi}(\tau)} \sum_{\substack{\hat{\epsilon} = (\epsilon_1, \epsilon_2, \epsilon_3) \\ \in (\mathbb{Z}/2)^{\otimes 3}}} (-1)^{\hat{\epsilon}} \Theta_{A, \mathbf{a}_{\hat{\epsilon}}, \mathbf{b}, \mathbf{c}_1, \mathbf{c}_2}(\tau)$$

with parameters given in (3.47) above with  $(p_1, p_2, p_3) = (s, t, str+1)$ , is, up to a pre-factor  $c \in \mathbb{C}$ , a mixed mock modular form f such that

$$\hat{f} := f - (\theta_{r_{\nu}}^{m+K})^* - \sum_{\substack{\hat{\epsilon} \in (\mathbb{Z}/2)^{\otimes 3}\\ \ell \in \mathbb{Z}/L\mathbb{Z}}} h_{\hat{\epsilon},\ell} \theta_{\hat{\epsilon},\ell}^*, \quad L = xp_1p_2 - 2p_3$$
(3.48)

transforms as a weight 1/2 modular form, where  $r_{\nu}$  is as in (2.42). Moreover,  $\theta_{\hat{\epsilon}}$  is a onedimensional unary theta function  $\theta_{\hat{\epsilon}} = \theta_{M,R}^1(c\tau)$  for some  $M \in \mathbb{N}$ ,  $R \in \mathbb{Z}/2M$  and  $c \in \mathbb{Q}$ , and  $h_{\hat{\epsilon}}(\tau)$  either has an exponential singularity or vanishes at cusp  $\tau \to \alpha \in \mathbb{Q} \cup \{i\infty\}$ .

We record the proof in Appendix A. Note that there sporadic cases where the additional shadows vanish, leading to  $\sum_{\hat{\epsilon} \in (\mathbb{Z}/2)^{\otimes 3}} h_{\hat{\epsilon},\ell} \theta^*_{\hat{\epsilon},\ell} = 0$  above. This is the case when  $\ell \in \mathbb{Z}/L\mathbb{Z}$ 

 $(p_1, p_2, p_3) = (2, 5, 21)$ . Also, as we will see in §5.5.3, although the manifolds  $-\Sigma(2, 3, 5)$  is not in the family of manifold  $-\Sigma(s, t, str + 1)$  that can be constructed via the (+1/r)-surgery of torus knots, there is an analogous construction of indefinite theta functions that give rise to its pure mock  $-\hat{Z}$ -invariants.

# 4 Mock Invariants and Vertex Algebras

Vertex operator algebras have been shown to be related to 3d supersymmetric conformal field theories, in particular to the  $\hat{Z}$ -invariants, in rich and deep ways. In the context of negative Seifert manifolds with three or four exceptional fibres the  $\hat{Z}$ -invariant has been shown [30, 62] to be given by a linear combination of characters of a class of logarithmic vertex operator algebras. The insertion of defects or a change of the choice of the Spin<sup>c</sup>-structure, while modifying the modules of the vertex algebra computed, does not change

the underlying algebra. In these cases, this has led to a construction of a vertex algebra from a given three-dimensional manifold.

In this section we extend the relation between  $\hat{Z}$ -invariants and vertex operator algebras to the "other side" of mock invariants, by systematically associating vertex operator algebras to the mock  $\hat{Z}$  invariants proposed as indefinite theta function in the previous section. This direct connection to vertex algebras, together with the manifest mock modular property, is one of the most important advantages of having an indefinite theta function expression for  $\hat{Z}$ -invariants for non-weakly-negative three-manifolds.

### 4.1 Cone Vertex Algebras

It is well-known that from an integral lattice L one can construct a Lie algebra and the Lie algebra module  $V_L$ , which has the structure of vertex algebra and is moreover a vertex operator algebra when L is positive definite. We will now briefly describe how one can associate a vertex operator algebra to a positive-definite cone in a lattice of indefinite signature. We will skip many details and simply refer to [25] (§2, §3) for a full account of these cone algebras. For a cone  $C \subset L$  with  $0 \in C$  that is closed under addition, the submodule  $V_C$  of  $V_L$  generated by elements corresponding to  $\lambda \in C$  has the structure of a sub-vertex algebra of  $V_L$ , with the same conformal element. Furthermore,  $V_C$  has the structure of vertex operator algebra when the lattice linear form is positive definite when restricted to C. Given  $\gamma \in L \otimes_{\mathbb{Z}} \mathbb{Q}$ , for any  $C' \subset L + \gamma$  such that  $C' + C \subset C'$ , the corresponding  $V_{C'}$  can be endowed with the structure of a twisted module over  $V_C$ .

Suppose that the cone

$$P := \left\{ \sum_{i=1}^{\operatorname{rk}(L)} \alpha_i \mathsf{d}_i \in L | \alpha_i \ge 0, \ \forall \ i = 1, \dots, \operatorname{rk}(L) \right\}$$
(4.1)

for some  $\mathbf{d}_i \in L$  is such a positive-definite subset of L. Then  $V_P$  has a vertex operator algebra structure. Given a generating set  $\{\boldsymbol{\varepsilon}_1, \ldots, \boldsymbol{\varepsilon}_{\mathrm{rk}(L)}\}$  of L and  $\mathbf{a}^+ = \sum_i a_i \boldsymbol{\varepsilon}_i$ ,  $\mathbf{a} = (a_1, \ldots, a_{\mathrm{rk}(L)}) \in \mathbb{Q}^{\mathrm{rk}(L)}$ , define  $\mathbf{a}^- = -\mathbf{a}^+ + \sum_i \boldsymbol{\varepsilon}_i$  and note that  $C' + P \subset C'$  when  $C' = P + \mathbf{a}^{\pm}$ . The twisted module of  $V_P$  that will be the main object of interest here is

$$V_{\mathsf{a}} := V_{\mathsf{a}^+ + P} \oplus V_{\mathsf{a}^- + P}. \tag{4.2}$$

As usual, this twisted module is simply a module when  $a^+, a^- \in L^*$  are in the dual lattice.

### 4.2 Mock VOA Characters

Next we will establish the relation between certain trace functions of the modules (4.2) and the indefinite theta series of the kind encountered in the last section as proposals for  $\hat{Z}$ -invariants. For the sake of concreteness we will now focus on rank two lattices with signature (1, 1). Generalization to lattices with signature (1, d) is straightforward. Given  $b = \sum_{i=1}^{2} b_i d_i$  with  $b_i \in \mathbb{Q}$ , consider the following automorphism  $g_b$  of the module  $V_a$ : it acts as a multiplication by the phase  $e(B(b, \lambda - a^+))$  on elements corresponding to  $\lambda \in P + a^+$ , and multiplication by the phase  $-e(B(b, -\lambda - a^+))$  on elements corresponding to  $\lambda \in P + a^-$ .

The main object we will be interested in is the trace function

$$T_{\mathsf{a},\mathsf{b}}(\tau) := \operatorname{Tr}_{V_{\mathsf{a}}}\left(g_{\mathsf{b}}q^{L(0)-c/24}\right)$$
(4.3)

which is a well-defined function on  $\tau \in \mathbb{H}$  since the lattice bilinear form is positive definite when restricted to the cone P.

Next we will demonstrate the relation between  $T_{a,b}$  and the indefinite theta functions seen in the previous section, generalizing Lemma 3.1 and Theorem 3.2 of [25].

**Theorem 3.** Consider the lattice bilinear  $A = \text{diag}(-2p\bar{p}, x)$  with  $x, p, \bar{p} \in \mathbb{N}$  satisfying  $\bar{p}x > 2p$ , and the positive-definite cone (4.1) with

$$\mathsf{d}_1 = \mathsf{c}_{1,\perp} = \begin{pmatrix} 0\\1 \end{pmatrix}, \ \mathsf{d}_2 = \mathsf{c}_{2,\perp} = \begin{pmatrix} 1\\\bar{p} \end{pmatrix} . \tag{4.4}$$

Consider  $\tilde{a} = \sum_{i} \tilde{a}_{i} d_{i}$  with  $0 < \tilde{a}_{i} < 1$ . The trace function of the corresponding twisted module  $V_{a}$  is given in terms of the indefinite theta function by

$$T_{\tilde{\mathsf{a}},\mathsf{b}}(\tau) = -\frac{2}{\eta^2(\tau)} e(-B(\mathsf{b},\tilde{\mathsf{a}}))\Theta_{A,\mathsf{a},\mathsf{b},\mathsf{c}_1,\mathsf{c}_2}(\tau)$$
(4.5)

where

$$\mathbf{c}_1 = \begin{pmatrix} 1\\0 \end{pmatrix}, \ \mathbf{c}_2 = \begin{pmatrix} x\\2p \end{pmatrix}, \tag{4.6}$$

and a is any vector satisfying  $a = \tilde{a} \mod \mathbb{Z}d_1 + \mathbb{Z}d_2$ .

*Proof.* Using the definition (4.3), one obtains

$$\begin{split} T_{\tilde{\mathbf{a}},\mathbf{b}}(\tau) &= \frac{1}{\eta^{2}(\tau)} \\ \times \sum_{n_{1},n_{2} \geq 0} \left( e(B(\mathbf{b},\sum_{i} n_{i}\mathbf{d}_{i}))q^{|\sum_{i}(n_{i}+\tilde{a}_{i})\mathbf{d}_{i}|^{2}/2} - e(-B(\mathbf{b},\sum_{i}(n_{i}+1)\mathbf{d}_{i}))q^{|\sum_{i}(n_{i}+1-\tilde{a}_{i})\mathbf{d}_{i}|^{2}/2} \right), \end{split}$$

while plugging in the vector  $c_i$  into (3.13) leads to

$$-\frac{1}{2}\Theta_{A,\mathbf{a},\mathbf{b},\mathbf{c}_{1},\mathbf{c}_{2}}(\tau) = \mathbf{e}(B(\mathbf{b},\tilde{\mathbf{a}}))$$

$$\times \left(\sum_{n_{1},n_{2}\geq0}\mathbf{e}(B(\mathbf{b},\sum_{i}n_{i}\mathbf{d}_{i}))q^{|\sum_{i}(n_{i}+\tilde{a}_{i})\mathbf{d}_{i}|^{2}/2} - \sum_{n_{1},n_{2}>0}\mathbf{e}(-B(\mathbf{b},\sum_{i}n_{i}\mathbf{d}_{i}))q^{|\sum_{i}(-n_{i}+\tilde{a}_{i})\mathbf{d}_{i}|^{2}/2}\right)$$

# 5 Examples

In this section, we will demonstrate the various phenomena discussed in the previous sections, either in the form of conjectures or proven theorems, with various examples of different types.

To simplify the equations, in this section we will not always keep track of pre-factors  $cq^{\Delta}$  for some  $c \in \mathbb{C}$  and  $\Delta \in \mathbb{Q}$  the possible finite polynomials  $p(\tau)$ . In other words, we will often use the same equivalent relation as in the Modularity Conjecture 4, and use  $\sim$  to denote the equivalence defined this way. The data provided in the paper are sufficient for the readers to compute explicitly c,  $\Delta$ , and  $p(\tau)$  in all cases.



Figure 2: Plumbing diagrams

#### 5.1Invariants from Plumbing Graphs

In this subsection, we use the contour integral expressions (1.2) and Definition 1 to compute the examples of Z-invariants, with and without defects. Using these concrete Zinvariants, we provide evidence for the Modularity Conjecture 4, for manifolds for which a general proof is not available (cf. Theorem 1).

So far we have not found a way to compute the relevant  $\widetilde{\mathrm{SL}}_2(\mathbb{Z})$  representation  $\Theta^{m+K}$ for generic Seifert manifolds with three exceptional fibres, despite systematic results for the Brieskorn spheres. This said, we would like to extensively test the Modularity Conjecture 4 beyond the cases in Theorem 1. For a given negative Seifert manifold with three exceptional fibres, we will do this in three steps. First we compute the  $\hat{Z}$ -invariants without defects, for all admissible Spin<sup>c</sup> structures, using the original plumbing proposal (1.2). Second, from those we identify the relevant  $\widetilde{\mathrm{SL}}_2(\mathbb{Z})$  representation  $\Theta^{m+K}$ , using the notation as in (2.38). We will also write  $\sigma^{m+K}$  to denote the set of independent components of  $\Theta^{m+K}$ , in the sense detailed in (2.38). Finally we identify the examples of Wilson line insertions that give rise to the specific components of the corresponding vector-valued quantum modular forms. In all examples in this subsection, invariants will be computed using Definition 2.4.

For concreteness, in all examples in this section we represent one of the Spin<sup>c</sup> structures as

$$b_0 = \{b_{0,v}\}_{v \in V}, \ b_{0,v} = \deg(v) - 2.$$
(5.1)

**5.1.1** 
$$M_3 = \Sigma(2,3,7), m = 42, K = \{1, 6, 14, 21\}$$

Although this Brieskorn sphere case is covered in Theorem 1, we will still give it as an example to illustrate the Theorem, and also for the sake of later comparison with other approaches to this particular manifold.

The plumbing graph for  $\Sigma(2,3,7)$  is depicted in Figure 2a with the weights  $\alpha_i^{(i)}$  on the node labelled by  $v_j^{(i)}$  given by  $(\alpha^{(0)}, \alpha_1^{(1)}, \alpha_1^{(2)}, \alpha_1^{(3)}) = (-1, -2, -3, -7)$  (cf. (2.1)). Since the three-manifold is a homological sphere, we have only one Spin<sup>c</sup>-structure  $b_0$ ,

and we have

$$\hat{Z}_{b_0}(\Sigma(2,3,7);\tau) \sim \tilde{\theta}_1^{m+K}.$$
 (5.2)

The Weil representation  $\Theta^{m+K}$  is a three-dimensional irreducible representation with  $\sigma^{m+K} = \{1, 5, 11\}.$ 

The other two components of the vector-valued quantum modular form

$$\begin{pmatrix} \tilde{\theta}_1^{m+K} \\ \tilde{\theta}_5^{m+K} \\ \tilde{\theta}_{11}^{m+K} \end{pmatrix} (\tau)$$

show up as the following defect invariants

$$\hat{Z}_{b_0}\left(\Sigma\left(2,3,7\right); W_{(0,0,1)}; \tau\right) \sim \tilde{\theta}_5^{m+K}(\tau) 
\hat{Z}_{b_0}\left(\Sigma\left(2,3,7\right); W_{(0,0,2)}; \tau\right) \sim \tilde{\theta}_{11}^{m+K}(\tau).$$
(5.3)

**5.1.2**  $M_3 = M\left(-2; \frac{1}{2}, \frac{2}{3}, \frac{2}{3}\right), \ m = 6, \ K = \{1, 3\}$ 

This example was featured in [30] (Table 13). The corresponding Weil representation  $\Theta^{6+\{1,3\}}$  is a two-dimensional irreducible representation with  $\sigma^{m+K} = \{1,3\}$ . Its plumbing graph is depicted in Figure 2b with the weight assignment  $(\alpha^{(0)}, \alpha_1^{(1)}, \alpha_1^{(2)}, \alpha_2^{(2)}, \alpha_1^{(3)}, \alpha_2^{(3)}) = (-2, -2, -2, -2, -2, -2).$ 

Apart from the trivial Spin<sup>c</sup>-structure  $b_0$ , this manifold permits a second Spin<sup>c</sup>-structure that can be represented by  $b_1 = (-1, 1, 0, 1, 0, 3)$  in the basis defined by the node ordering  $v^{(0)}, v_1^{(1)}, v_1^{(2)}, v_2^{(2)}, v_1^{(3)}, v_2^{(3)}$ . This example has the special property that the two  $\hat{Z}_b$  with no defect modifications already span the vector-valued  $\widetilde{SL}_2(\mathbb{Z})$  quantum modular forms:

$$\hat{Z}_{b_0}\left(M\left(-2;\frac{1}{2},\frac{2}{3},\frac{2}{3}\right);\tau\right) \sim \tilde{\theta}_1^{m+K}(\tau) \\ \hat{Z}_{b_1}\left(M\left(-2;\frac{1}{2},\frac{2}{3},\frac{2}{3}\right);\tau\right) \sim \tilde{\theta}_3^{m+K}(\tau)$$

The inclusion of defects, in this case, gives rise to the same functions up to finite polynomials. For instance, we have

$$\hat{Z}_{b_0}\left(M\left(-2;\frac{1}{2},\frac{2}{3},\frac{2}{3}\right);W_{(0,0,1)};\tau\right)\sim\tilde{\theta}_3^{m+K}(\tau)$$
$$\hat{Z}_{b_1}\left(M\left(-2;\frac{1}{2},\frac{2}{3},\frac{2}{3}\right);W_{(0,0,1)};\tau\right)\sim\tilde{\theta}_1^{m+K}(\tau)$$

consistent with the Modularity Conjecture 4.

**5.1.3** 
$$M_3 = M\left(-1; \frac{1}{2}, \frac{1}{7}, \frac{2}{7}\right), m = 14, K = \{1, 7\}$$

In [30] (Table 13) it was found that this plumbed manifold, with plumbing graph depicted in Figure 2c and the weight assignment  $(\alpha^{(0)}, \alpha_1^{(1)}, \alpha_1^{(2)}, \alpha_1^{(3)}, \alpha_2^{(3)}) = (-1, -2, -7, -4, -2)$ , corresponds to the four-dimensional irreducible Weil representation  $\Theta^{14+\{1,7\}}$  which has  $\sigma^{m+K} = \{1, 3, 5, 7\}$ .

It has two inequivalent Spin<sup>c</sup>-structures  $b_0$  and  $b_1 = (-1, 1, 1, 0, 5)$ . The corresponding  $\hat{Z}$ -invariants correspond to  $\{1, 7\} \subset \sigma^{m+K}$ :

$$\hat{Z}_{b_0}\left(M\left(-1;\frac{1}{2},\frac{1}{7},\frac{2}{7}\right);\tau\right) \sim \tilde{\theta}_3^{m+K}(\tau) \\ \hat{Z}_{b_1}\left(M\left(-1;\frac{1}{2},\frac{1}{7},\frac{2}{7}\right);\tau\right) \sim \tilde{\theta}_7^{m+K}(\tau) .$$

The rest of the components in  $\sigma^{m+K} = \{1, 3, 5, 7\}$  become visible when one considers the defect invariants, consistent with the Modularity Conjecture 4. For instance, we see

$$\begin{aligned} \hat{Z}_{b_0} \left( M\left(-1; \frac{1}{2}, \frac{1}{7}, \frac{2}{7}\right); W_{(0,0,1)}; \tau \right) &\sim \tilde{\theta}_1^{m+K}(\tau) \\ \hat{Z}_{b_0} \left( M\left(-1; \frac{1}{2}, \frac{1}{7}, \frac{2}{7}\right); W_{(0,0,4)}; \tau \right) &\sim \tilde{\theta}_5^{m+K}(\tau) \\ \hat{Z}_{b_0} \left( M\left(-1; \frac{1}{2}, \frac{1}{7}, \frac{2}{7}\right); W_{(0,0,5)}; \tau \right) &\sim \tilde{\theta}_7^{m+K}(\tau) \end{aligned}$$

where we obtain matching after removing certain q-monomials in the latter two cases.

### 5.1.4 Other Examples

In Tables 1 and 2 we compile an extensive list of Seifert manifolds, for which we record the relevant  $\widetilde{\operatorname{SL}}_2(\mathbb{Z})$  representation  $\Theta^{m+K}$  and its corresponding natural components labelled by the elements of the set  $\sigma^{m+K}$ . For all these cases, we have verified the Modularity Conjecture 4. We use the notation  $\sigma_{\hat{Z}}$  to denote the set of components in  $\Theta^{m+K}$  that are seen in the  $\hat{Z}$ -invariants, with and without defects, of the three-manifold in question. The Modularity Conjecture 4 in particular implies  $\sigma^{m+K} = \sigma_{\hat{Z}}$ . In Table 2 we list the cases for which the Weil representation is of the form  $\Theta^{m+K}$  (2.37). As explained in §2.3, sometimes these Weil representations defined through (2.37) are reducible and one needs an additional projection (2.39) to obtain the corresponding irreducible subrepresentation  $\Theta^{m+K,\text{irr}}$ . Interestingly, these also appear as modular representations for  $\hat{Z}$ -invariants for Seifert manifolds with three singular fibres. In Table 2 we record some examples where  $\Theta^{m+K,\text{irr}}$  appear. We use the notation of writing m + K and skip writing the identity element in K (cf. [93]).

$M_3$	m + K	$\sigma^{m+K} = \sigma_{\hat{Z}}$
M(-2;1/2,1/2,1/2)	2	{1}
$M\left(-2;1/2,1/2,2/3 ight)$	3	$\{1,2\}$
$M\left(-2;1/2,1/2,3/4 ight)$	4	$\{1, 2, 3\}$
M(-2; 1/2, 1/2, 4/5)	5	$\{1 \dots 4\}$
M(-2; 1/2, 1/2, 5/6)	6	$\{1 \dots 5\}$
M(-2; 1/2, 2/3, 2/3)	6 + 3	$\{1,3\}$
M(-2; 1/2, 1/2, 6/7)	7	$\{1 \dots 6\}$
M(-2; 1/2, 1/2, 7/8)	8	$\{1,7\}$
M(-2; 1/2, 1/2, 8/9)	9	$\{18\}$
M(-2; 1/2, 1/2, 9/10)	10	$\{1\dots 9\}$
M(-1; 1/2, 1/5, 1/5)	10 + 5	$\{1, 3, 5\}$
M(-4; 1/2, 1/2, 1/2)	10 + 5	$\{1, 3, 5\}$
M(-2; 1/2, 1/2, 11/12)	12	$\{1,11\}$
M(-2; 1/2, 1/2, 12/13)	13	$\{112\}$
M(-5; 1/2, 1/2, 1/2)	14 + 7	$\{1, 3, 5, 7\}$
M(-1; 1/2, 1/7, 2/7)	14 + 7	$\{1, 3, 5, 7\}$
M(-2; 1/2, 1/2, 15/16)	16	$\{115\}$
M(-2; 1/2, 1/2, 17/18)	18	$\{117\}$
M(-6; 1/2, 1/2, 1/2)	18 + 9	$\{1, 3, 5, 7, 9\}$
M(-2; 1/2, 1/2, 24/25)	25	$\{124\}$
Table	e 1: contin	ued on next page

001	unaca no	m provious page
$M_3$	m+K	$\sigma^{m+K} = \sigma_{\hat{Z}}$
M(-7; 1/2, 1/2, 1/2)	22 + 11	$\{1, 3, 5, 7, 9, 11\}$
M(-1; 1/2, 1/11, 4/11)	22 + 11	$\{1, 3, 5, 7, 9, 11\}$
M(-9; 1/2, 1/2, 1/2)	30 + 15	$\{1, 3, 5, 7, 9, 11, 13, 15\}$
M(-1; 1/2, 2/5, 1/15)	30 + 15	$\{1, 3, 5, 7, 9, 11, 13, 15\}$
M(-13; 1/2, 1/2, 1/2)	46 + 23	$\{1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23\}$
M(-2; 1/2, 1/2, 1/3)	6 + 2	$\{1, 2, 4\}$
M(-2; 1/2, 1/2, 3/5)	10 + 2	$\{1 \dots 4, 6, 8\}$
M(-1; 1/3, 1/3, 1/4)	12 + 3	$\{1 \dots 3, 5, 6, 9\}$
M(-2; 1/2, 1/2, 1/4)	12 + 3	$\{1 \dots 3, 5, 6, 9\}$
M(-1; 1/3, 1/5, 2/5)	15 + 5	$\{1, 2, 4, 5, 7, 10\}$
M(-3; 1/2, 1/2, 1/3)	15 + 5	$\{1, 2, 4, 5, 7, 10\}$
M(-2; 1/2, 1/2, 7/9)	18 + 2	$\{1\dots 8, 10, 12, 14, 16\}$
$M\left(-2;1/2,1/2,4/7 ight)$	21 + 3	$\{1 \dots 6, 8, 9, 11, 12, 15, 18\}$
$M\left(-1; 1/4, 1/7, 4/7\right)$	28 + 7	$\{1 \dots 3, 5 \dots 7, 9, 10, 13, 14, 17, 21\}$
$M\left(-1; 1/4, 1/7, 4/7\right)$	28 + 7	$\{1 \dots 3, 5 \dots 7, 9, 10, 13, 14, 17, 21\}$
M(-5; 1/2, 1/2, 1/3)	33 + 11	$\{1, 2, 4, 5, 7, 8, 10, 11, 13, 16, 19, 22\}$
M(-1; 1/3, 1/11, 6/11)	33 + 11	$\{1, 2, 4, 5, 7, 8, 10, 11, 13, 16, 19, 22\}$

Continued from previous page

Table 1: The Seifert manifolds and the associated  $\widetilde{SL}_2(\mathbb{Z})$  representations.

$M_3$	m+K	$\sigma^{m+K,\mathrm{irred}} = \sigma_{\hat{Z}}$
M(-1; 1/2, 1/3, 1/9)	18 + 9	$\{1, 3, 5, 7\}$
M(-2; 1/2, 1/3, 2/3)	18 + 9	$\{1, 3, 5, 7\}$
M(-1; 1/2, 1/4, 1/5)	20 + 4	$\{1, 3, 4, 8, 11\}$
M(-1; 1/2, 1/3, 1/8)	24 + 8	$\{1, 2, 5, 7, 8, 13\}$

Table 2: The Seifert manifolds and the associated  $\widetilde{SL}_2(\mathbb{Z})$  representations.

## 5.2 Invariants from Knot Surgeries

In this subsection, we use Definition 2 to compute the examples of  $\hat{Z}$ -invariants, with and without defects. This provides evidence for the consistency of Definition 2, and the validity of Modularity Conjecture 4.

**5.2.1** 
$$M_3 = \Sigma(2,3,7), m = 42, K = \{1,6,14,21\}$$

Here we will consider two ways in which  $\Sigma(2,3,7)$  can be represented as a knot surgery: either as a (-1)-surgery on a right-handed trefoil  $\mathbf{3}_1^r$  or as a (+1)-surgery on a figure eight knot  $\mathbf{4}_1$ . The two constructions give rise to two distinct types of expressions that evaluate to the same functions. In this subsection we focus on the  $\mathbf{3}_1^r$  construction. The  $F_K$  polynomial for the right-handed trefoil  $\mathbf{3}_1^r = T(2,3)$  reads (3.44)

$$F_{\mathbf{3}_{1}^{r}}(x;q) = \frac{1}{2} \left[ -\left(x^{\frac{1}{2}} - x^{-\frac{1}{2}}\right)q + \left(x^{\frac{5}{2}} - x^{-\frac{5}{2}}\right)q^{2} + \left(x^{\frac{7}{2}} - x^{-\frac{7}{2}}\right)q^{3} \right] + \mathcal{O}\left(q^{6}\right)$$
(5.4)

which leads to the following expression after the Laplace transformation

$$\mathcal{L}_{-1}^{(0)} \left[ \left( x^{\frac{1}{2}} - x^{-\frac{1}{2}} \right) F_{\mathbf{3}_{1}^{r}}(x;\tau) \right] \sim \hat{Z}_{b_{0}} \left( \Sigma \left( 2,3,7 \right);\tau \right) \\ \mathcal{L}_{-1}^{(1)} \left[ \left( x - x^{-1} \right) F_{\mathbf{3}_{1}^{r}}(x;\tau) \right] \sim \hat{Z}_{b_{0}} \left( \Sigma \left( 2,3,7 \right);W_{(0,0,1)};\tau \right) \\ \mathcal{L}_{-1}^{(2)} \left[ \left( x^{\frac{3}{2}} - x^{-\frac{3}{2}} \right) F_{\mathbf{3}_{1}^{r}}(x;\tau) \right] \sim \hat{Z}_{b_{0}} \left( \Sigma \left( 2,3,7 \right);W_{(0,0,2)};\tau \right)$$
(5.5)

which we checked up to  $\mathcal{O}(q^{20})$ . For the sake of comparison, on the right hand side we have used the plumbing notation as in §5.1.1. As shown there, the above  $\hat{Z}$ -invariants are equal, up to an overall q-power and the addition of finite polynomials, to the false theta functions  $\tilde{\theta}_1^{m+K}(\tau)$ ,  $\tilde{\theta}_{11}^{m+K}(\tau)$  and  $\tilde{\theta}_5^{m+K}(\tau)$ .

**5.2.2** 
$$M_3 = M\left(-1; \frac{1}{2}, \frac{1}{3}, \frac{1}{9}\right), \ m = 18, \ K = \{1, 9\}$$

This manifold, also featured in Table 2, can be constructed via a (-3)-surgery on the right-handed trefoil, whose  $F_K(x;\tau)$  invariant is listed in (5.4). The surgery formula gives

$$\mathcal{L}_{-3}^{(0)} \left[ \left( x^{\frac{1}{2}} - x^{-\frac{1}{2}} \right) F_{\mathbf{3}_{1}^{r}}(x;\tau) \right] \sim \hat{Z}_{b_{0}} \left( M \left( -1; \frac{1}{2}, \frac{1}{3}, \frac{1}{9} \right);\tau \right) \\
\mathcal{L}_{-3}^{(2)} \left[ \left( x^{\frac{3}{2}} - x^{-\frac{3}{2}} \right) F_{\mathbf{3}_{1}^{r}}(x;\tau) \right] \sim \hat{Z}_{b_{0}} \left( M \left( -1; \frac{1}{2}, \frac{1}{3}, \frac{1}{9} \right); W_{(0,0,2)};\tau \right) \\
\mathcal{L}_{-3}^{(3)} \left[ \left( x^{2} - x^{-2} \right) F_{\mathbf{3}_{1}^{r}}(x;\tau) \right] \sim \hat{Z}_{b_{0}} \left( M \left( -1; \frac{1}{2}, \frac{1}{3}, \frac{1}{9} \right); W_{(0,0,3)};\tau \right) \\
\mathcal{L}_{-3}^{(4)} \left[ \left( x^{\frac{5}{2}} - x^{-\frac{5}{2}} \right) F_{\mathbf{3}_{1}^{r}}(x;\tau) \right] \sim \hat{Z}_{b_{0}} \left( M \left( -1; \frac{1}{2}, \frac{1}{3}, \frac{1}{9} \right); W_{(0,0,4)};\tau \right) ,$$
(5.6)

which we checked up to  $\mathcal{O}(q^{20})$ . For the sake of comparison, on the right hand side we have used the plumbing notation for the defect operators. The plumbing description is in terms of the plumbing graph as in Figure 2a, with the weight vector given by (-1, -2, -3, -9). They are proportional up to a finite polynomial, to the components of the vector-valued quantum modular form

$$\begin{pmatrix} \tilde{\theta}_{1}^{18+\{1,9\},\text{irr}} \\ \tilde{\theta}_{1}^{18+\{1,9\},\text{irr}} \\ \tilde{\theta}_{3}^{18+\{1,9\},\text{irr}} \\ \tilde{\theta}_{5}^{18+\{1,9\},\text{irr}} \end{pmatrix} (\tau),$$

corresponding to the four-dimensional irreducible Weil representation  $\Theta^{18+\{1,9\},irr}$ .

### 5.3 More Knot Surgeries Using the Inverted Habiro Series

In this subsection we will compute defects  $\hat{Z}$ -invariants using Conjecture 3, generalizing the conjecture in [36] to include defect operators. We present the examples of four Seifert manifolds, displaying different properties while all confirming the Modularity Conjecture 4.

**5.3.1** 
$$M_3 = \pm \Sigma (2, 3, 5), m = 30, K = \{1, 6, 10, 15\}$$

The manifold  $M_3 = \Sigma(2,3,5)$  can be represented as a plumbed manifold with plumbing diagram given in Figure 2a and weights  $(\alpha^{(0)}, \alpha_1^{(1)}, \alpha_1^{(2)}, \alpha_1^{(3)}) = (-1, -2, -3, -5)$ , and also

as the (-1)-surgery of the left-handed trefoil knot. From the latter point of view, we can make use of Conjecture 3 with  $p = \pm 1$  in order to compute the defect  $\hat{Z}$ -invariants for  $\Sigma(2,3,5)$  as well as  $-\Sigma(2,3,5)$ .

For the latter, the results obtained here can be compared to the results obtained as

indefinite theta functions using its description as a plumbed manifold, detailed in 5.5.3. From

$$\Sigma(2,3,5) = S_{-1}^3(\mathbf{3}_1^l) , \qquad (5.7)$$

and the Habiro coefficients

$$a_{-m}(\mathbf{3}_1^l) = (-1)^m q^{\frac{m(m-3)}{2}}, \qquad m \in \mathbb{Z},$$
(5.8)

we arrive at

$$\hat{Z}_{b_0}\left(S_{-1}^3(\mathbf{3}_1^l);\tau\right) \sim \tilde{\theta}_1^{m+K}(\tau)$$

$$\hat{Z}_{b_0}\left(S_{-1}^3(\mathbf{3}_1^l);W_1;\tau\right) \sim \tilde{\theta}_7^{m+K}(\tau).$$
(5.9)

This is exactly what Theorem 1 dictates in this case. Similarly, from

$$-\Sigma(2,3,5) = S_{+1}^3(\mathbf{3}_1^r) , \quad a_{-m}(\mathbf{3}_1^r) = a_{-m}(\mathbf{3}_1^l)|_{q \to q^{-1}}, \tag{5.10}$$

we get

$$\begin{pmatrix} \hat{Z}_{b_0}(-\Sigma(2,3,5))\\ \hat{Z}_{b_0}(-\Sigma(2,3,5);W_1) \end{pmatrix} \sim \begin{pmatrix} H_1^{30+\{1,6,10,15\}}\\ H_5^{30+\{1,6,10,15\}} \end{pmatrix} \sim \begin{pmatrix} \chi_0\\ \chi_1 \end{pmatrix},$$
(5.11)

given by the two order 5 mock theta functions of Ramanujan  $\chi_0$  and  $\chi_1$ , which are also the components of the optimal mock Jacobi theta function with the corresponding m + K [92].

Assuming Conjecture 3 is correct, this example serves also to showcase Conjecture 4.2.

**5.3.2** 
$$M_3 = -M\left(-1; \frac{1}{2}, \frac{1}{3}, \frac{1}{8}\right), \ m = 24, \ K = \{1, 8\}$$

Now we consider

$$M_3 = -M\left(-1; \frac{1}{2}, \frac{1}{3}, \frac{1}{8}\right) = S^3_{+2}(\mathbf{3}^l_1), \tag{5.12}$$

corresponding to the 6-dimensional irreducible Weil representation  $\Theta^{24+\{1,8\},irr}$ , with

$$\sigma^{24+\{1,8\},\mathrm{irr}} = \{1, 2, 5, 7, 8, 13\}.$$

The plumbing description has plumbing graph as in Figure 2a, with weight assignment

$$\left(\alpha^{(0)}, \alpha_1^{(1)}, \alpha_1^{(2)}, \alpha_1^{(3)}\right) = (-1, -2, -3, -8).$$

The two inequivalent Spin<sup>c</sup> structures are represented by  $b_0$  and  $b_1 = (-1, 1, 1, 3)$ .

The prescription in Conjecture 3 leads to q-series which can be observed to coincide, up to a pre-factor and a finite polynomial, the components of the optimal mock Jacobi theta

function with the corresponding m + K, at least to the order  $\mathcal{O}(q^{15})$  we have computed. Explicitly, we have

$$\begin{pmatrix} \hat{Z}_{b_0}(W_0) \\ \hat{Z}_{b_0}(W_1) \\ \hat{Z}_{b_0}(W_2) \\ \hat{Z}_{b_0}(W_3) \\ \hat{Z}_{b_1}(W_0) \\ \hat{Z}_{b_1}(W_2) \end{pmatrix} \sim \begin{pmatrix} H_1^{24+\{1,8\}} \\ H_2^{24+\{1,8\}} \\ H_2^{24+\{1,8\}} \\ H_8^{24+\{1,8\}} \\ H_7^{24+\{1,8\}} \\ H_{13}^{24+\{1,8\}} \end{pmatrix}.$$
(5.13)

We collect in Appendix B the  $P_n^{p,b+\nu}$  and  $P_n^{p,b+\nu}$  polynomials that were used to compute the *q*-series for this example.

**5.3.3** 
$$M_3 = \pm \Sigma (2, 3, 7), m = 42, K = \{1, 6, 14, 21\}$$

From

$$\Sigma(2,3,7) = S_{-1}^3(\mathbf{3}_1^r) , \quad a_{-m}(\mathbf{3}_1^r) = (-1)^m q^{-\frac{m(m-3)}{2}}$$
(5.14)

we get using  $\lambda_n^{n,1}(q) = (q^n + 1 + q^{-n}), \ \lambda_{n-1}^{n,1}(q) = -1 \text{ and } \lambda_{\ell}^{n,1}(q) = 0 \text{ for } \ell \neq n-1, n \text{ the following expressions}$ 

$$\begin{aligned} \hat{Z}_{b_0}(S_{-1}^3(\mathbf{3}_1^r);\tau) &= q^{\frac{1}{2}} \left( 1 + \sum_{n=1}^{\infty} (-1)^n \frac{q^{\frac{n(n+1)}{2}}}{(q^{n+1};q)_n} \right) = q^{\frac{83}{168}} \tilde{\theta}_1^{m+K}(\tau) , \\ \hat{Z}_{b_0}(S_{-1}^3(\mathbf{3}_1^r);W_1;\tau) &= q^{\frac{3}{4}} \left( 2 + \sum_{n=1}^{\infty} (-1)^n \frac{q^{\frac{n(n-1)}{2}}}{(q^n;q)_n} \right) = -q^{\frac{101}{168}} \tilde{\theta}_5^{m+K}(\tau) - p_1(\tau) , \\ \hat{Z}_{b_0}(S_{-1}^3(\mathbf{3}_1^r);W_2;\tau) &= q^{\frac{1}{2}} \left( 1 + 2q + \sum_{n=1}^{\infty} (-1)^n \left( \frac{q^{1+\frac{n(n-3)}{2}}}{(q^n;q)_{n-1}} + (q^n + 1 + q^{-n}) \frac{q^{\frac{n(n+1)}{2}}}{(q^{n+1};q)_n} \right) \right) \\ &= -q^{\frac{131}{168}} \tilde{\theta}_{11}^{m+K}(\tau) - p_2(\tau) , \end{aligned}$$

$$(5.15)$$

where  $p_1(\tau) = -2q^{3/4}$  and  $p_2(\tau) = -2q^{3/2}$ , consistent with the plumbing result in §5.1.1.

Inverting the orientation by inverting the sign of the surgery and taking the mirror knot, namely

$$-\Sigma(2,3,7) = S_{+1}^3(\mathbf{3}_1^l),$$

we get

$$\hat{Z}_{b_0}(S_1^3(\mathbf{3}_1^l);\tau) = q^{-\frac{1}{2}} \left( 1 + \sum_{n=1}^{\infty} \frac{q^{n^2}}{(q^{n+1};q)_n} \right) = q^{-\frac{1}{2}} \left( 1 + q + q^3 + q^4 + q^5 + 2q^7 + \mathcal{O}\left(q^8\right) \right) 
\hat{Z}_{b_0}(S_1^3(\mathbf{3}_1^l);W_1;\tau) = q^{-\frac{3}{4}} \left( 2 + \sum_{n=1}^{\infty} \frac{q^{n^2}}{(q^n;q)_n} \right) = q^{-\frac{3}{4}} \left( 2 + q + q^2 + q^3 + q^4 + q^5 + 2q^6 + \mathcal{O}\left(q^7\right) \right) 
\hat{Z}_{b_0}(S_1^3(\mathbf{3}_1^l);W_2;\tau) = q^{-\frac{1}{2}} \left( 1 + \frac{2}{q} + \sum_{n=1}^{\infty} \left( -\frac{q^{n^2-n}}{(q^n;q)_{n-1}} + \left(q^n + 1 + q^{-n}\right) \frac{q^{n^2}}{(q^{n+1};q)_n} \right) \right) 
= q^{-\frac{3}{2}} \left( 2 + q + q^2 + 2q^3 + q^4 + 2q^5 + \mathcal{O}\left(q^6\right) \right) ,$$
(5.16)

which are the components of the optimal mock Jacobi theta function with the corresponding m + K. In this case, they are proportional to Ramanujan's order 7 mock theta functions  $F_0$ ,  $F_1$ , and  $F_2$ . In other words, we have

$$\begin{pmatrix} \hat{Z}_{b_0} \\ \hat{Z}_{b_0}(W_1) \\ \hat{Z}_{b_0}(W_2) \end{pmatrix} \sim \begin{pmatrix} H_1^{m+K} \\ H_5^{m+K} \\ H_{11}^{m+K} \end{pmatrix} \sim \begin{pmatrix} F_0 \\ F_1 \\ F_2 \end{pmatrix}.$$
 (5.17)

## 5.4 Invariants from a Hyperbolic Knot

Contrary to the trefoil, the figure 8 knot  $\mathbf{4}_1$  is not a plumbed knot. As a result, Conjecture 3 does not apply and we do not expect the  $\lambda_{\ell}^{n,\nu}(q)$  and  $\tilde{\lambda}_{\ell}^{n,\nu}(q)$  polynomials to give the correct invariants.

That said, we know that

$$S_{-1}^{3}(\mathbf{3}_{1}^{r}) = S_{+1}^{3}(\mathbf{4}_{1}) = \Sigma(2,3,7), \ S_{+1}^{3}(\mathbf{3}_{1}^{l}) = S_{-1}^{3}(\mathbf{4}_{1}) = -\Sigma(2,3,7),$$

and the construction using the trefoil knots does give surgeries expressions for the (defect)  $\hat{Z}$ -invariants using Conjecture 3. We hence wonder whether a similar expression could be obtained for surgeries along the  $4_1$  knot.

Using the fact that

$$\frac{\mathcal{L}_{-1}^{(0)}\left(\frac{1}{\prod_{j=1}^{\ell}(x+x^{-1}-q^{j}-q^{-j})}\right)}{\mathcal{L}_{+1}^{(0)}\left(\frac{1}{\prod_{j=1}^{n}(x+x^{-1}-q^{j}-q^{-j})}\right)} = (-1)^{n} \frac{q^{\ell^{2}}}{q^{\frac{n(n+1)}{2}}} \frac{\left(q^{n+1};q\right)_{n}}{\left(q^{\ell+1};q\right)_{\ell}}$$
(5.18)

is a finite polynomial in q and  $q^{-1}$  for all  $\ell \leq n$  and similarly for  $\mathcal{L}_{-1}^{(1)}\left(\frac{x^{1/2}+x^{-1/2}}{\prod_{j=1}^{\ell}(x+x^{-1}-q^{j}-q^{-j})}\right)$ , one can express the invariants  $\hat{Z}(\pm\Sigma(2,3,7);W_{\nu})$  obtained from applying Conjecture 3 to the  $\pm 1$  surgery of the trefoil in the following way

$$\hat{Z}(\Sigma(2,3,7); W_{\nu}, \tau) \sim \sum_{n \ge 0} (-1)^n \frac{q^{\frac{n(n+1)}{2}}}{(q^{n+1};q)_n} \sum_{\ell=0}^n \eta_{\ell}^{n,\nu}(q), \quad \nu \text{ even}$$

$$\hat{Z}(\Sigma(2,3,7); W_{\nu}, \tau) \sim \sum_{n \ge 0} (-1)^n \frac{q^{\frac{n(n+1)}{2} - \frac{1}{4}}}{(q^n;q)_n} \sum_{\ell=0}^n \tilde{\eta}_{\ell}^{n,\nu}(q), \quad \nu \text{ odd}$$
(5.19)

with

$$\eta_{\ell}^{n,\nu}(q) = q^{\ell^2 - n^2 + 1} \frac{(q^{n+1};q)_n}{(q^{\ell+1};q)_{\ell}} \lambda_{\ell}^{n,\nu}(q), \quad \nu \text{ even}$$
  
$$\tilde{\eta}_{\ell}^{n,\nu}(q) = -q^{\ell(\ell-1) - n^2 + \frac{3}{2}} \frac{(q^n;q)_n}{(q^\ell;q)_{\ell}} \lambda_{\ell}^{n,\nu}(q), \quad \nu \text{ odd.}$$
(5.20)

Similarly, considering the (+1)-surgery of the trefoil, or the (-1)-surgery of the figure 8 knot, we obtain

$$\hat{Z}(-\Sigma(2,3,7); W_{\nu}, \tau) \sim \sum_{n \ge 0} \frac{q^{n^2}}{(q^{n+1};q)_n} \sum_{\ell=0}^n \eta_{\ell}^{n,\nu}(q^{-1}), \quad \nu \text{ even}$$

$$\hat{Z}(-\Sigma(2,3,7); W_{\nu}, \tau) \sim \sum_{n \ge 0} \frac{q^{n^2-n+\frac{1}{4}}}{(q^n;q)_n} \sum_{\ell=0}^n \tilde{\eta}_{\ell}^{n,\nu}(q^{-1}), \quad \nu \text{ odd.}$$
(5.21)

One of the possible ways to generalize the above to the  $\pm p$  surgeries of the figure 8 knot, analogous to the expressions in Conjecture 3, leads to

$$\hat{Z}(S_{+p}^{3}(\mathbf{4}_{1}); W_{\nu}, \tau) \stackrel{?}{\sim} \sum_{n \geq 0} P_{n}^{p, b+\nu}(q)(-1)^{n} \frac{q^{\frac{n(n+1)}{2}}}{(q^{n+1}; q)_{n}} \sum_{\ell=0}^{n} \eta_{\ell}^{n, \nu}(q), \quad \nu \text{ even} \\
\hat{Z}(S_{+p}^{3}(\mathbf{4}_{1}); W_{\nu}, \tau) \stackrel{?}{\sim} \sum_{n \geq 0} P_{n}^{p, b+\nu}(q)(-1)^{n} \frac{q^{\frac{n(n+1)}{2}-\frac{1}{4}}}{(q^{n}; q)_{n}} \sum_{\ell=0}^{n} \tilde{\eta}_{\ell}^{n, \nu}(q), \quad \nu \text{ odd} \\
\hat{Z}(S_{-p}^{3}(\mathbf{4}_{1}); W_{\nu}, \tau) \stackrel{?}{\sim} \sum_{n \geq 0} P_{n}^{p, b+\nu}(q^{-1}) \frac{q^{n^{2}}}{(q^{n+1}; q)_{n}} \sum_{\ell=0}^{n} \eta_{\ell}^{n, \nu}(q^{-1}), \quad \nu \text{ even} \\
\hat{Z}(S_{-p}^{3}(\mathbf{4}_{1}); W_{\nu}, \tau) \stackrel{?}{\sim} \sum_{n \geq 0} P_{n}^{p, b+\nu}(q^{-1}) \frac{q^{n^{2}-n+\frac{1}{4}}}{(q^{n}; q)_{n}} \sum_{\ell=0}^{n} \tilde{\eta}_{\ell}^{n, \nu}(q^{-1}), \quad \nu \text{ odd}.$$
(5.22)

In this case we do not have the relation to plumbed manifolds to justify the above procedure, and as a result we put a "?" on top of the equal sign. However, we find it remarkable that the above formula gives rise to very interesting functions, which are consistent with the False-Mock Conjecture 1 and the Modularity Conjecture 4. This justifies recording the highly conjectural equations here for the benefit of the interested readers.

Using the expression (5.22), we obtain the results in Table 3, which states that the right-hand side of (5.22) using the indicated values of b and  $\nu$ , is, up to a pre-factor and possibly a finite polynomial, the same as the false or mock theta function listed in the column " $X_r^{m+K}$ ", where  $H_r^{m+K}$  denotes the component of the optimal mock Jacobi theta function in [92]<sup>11</sup>. In the table we also list the Seifert representation of the resulting three-manifold  $S_{\pm p}(\mathbf{4}_1)$ . In all cases, the plumbing graph is given in Figure 2a, for which we write the weight assignment in the fourth column, labeled by  $(\alpha^{(0)}, \alpha_1^{(1)}, \alpha_1^{(2)}, \alpha_1^{(3)})$ . On the right side of the table, we list the comparison to the results obtained from the plumbing prescription using Definition 1, where we indeed see a non-trivial matching.

**Remark 5.** Note that some of the components, corresponding to the elements of the relevant  $\sigma^{m+K}$  or  $\sigma^{m+K,irr}$ , are missing in the Table. This is because some of the components only appear when more general defect lines, not just those corresponding to the figure 8 knot, are included. For instance,  $\tilde{\theta}_8^{24+\{1,8\},irr}$  appears as  $\hat{Z}_{b_0}(W_{(0,1,2)})$  in the notation of the Table 3, but one doesn't see this false theta function when only Wilson lines corresponding to the distinguished knot is considered. All components should appear when one also considers the defect  $F_K$  series  $F_{K,\nu}$  that we commented on before Definition 2. Similar comments also hold for the case m = 12,  $K = \{1,3\}$ .

### 5.5 Mock invariants from Indefinite Theta Functions

In this subsection we provide specific examples of the discussion in §3.4. In particular, we will showcase two cases,  $M_3 = -\Sigma(2, 3, 6r + 1)$  for r = 1, 2, of the infinite family of the expressions proposed in Conjecture 5, which have been shown to have mock modular property in Theorem 2.

Additionally, we discuss the case of the Poincaré sphere  $M_3 = -\Sigma(2, 3, 5)$ , which goes beyond the scope of Conjecture 5 but possesses the mock modularity property similar to the one stated in Theorem 2.

<sup>&</sup>lt;sup>11</sup>We've checked it up to  $\mathcal{O}(q^{100})$  in the false cases and  $\mathcal{O}(q^{15})$  in the mock cases.

p	$M_3$	m, K	$(\alpha^{(0)}, \alpha_1^{(1)}, \alpha_1^{(2)}, \alpha_1^{(3)})$	b	ν	$X_r^{m+K}$	$( u_1, u_2, u_3)$	$\operatorname{Spin}^{c}$
1	$\Sigma(2,3,7)$	$42, \{1, 6, 14, 21\}$	(-1, -2, -3, -7)	0 0 0	0 1 2	$ \begin{array}{c} \theta_1^{m+K} \\ \theta_5^{m+K} \\ \theta_{11}^{m+K} \end{array} $	(0, 0, 0)(0, 0, 1)(0, 0, 2)	$(-1,1,1,1) \\ (-1,1,1,1) \\ (-1,1,1,1)$
-1	$-\Sigma(2, 3, 7)$	$42, \{1, 6, 14, 21\}$	(1, 2, 3, 7)	0 0 0	0 1 2	$ \begin{array}{c} H_1^{m+K} \\ H_5^{m+K} \\ H_{11}^{m+K} \end{array} $		
2	$M\left(-1;\frac{1}{2},\frac{1}{4},\frac{1}{5}\right)$	$20, \{1, 4\}$	(-1, -2, -4, -5)	0 2 0 0 2	0 0 1 2 2	$ \begin{array}{c} \tilde{\theta}_1^{m+K,\mathrm{irr}} \\ \tilde{\theta}_1^{m+K,\mathrm{irr}} \\ \tilde{\theta}_1^{m+K,\mathrm{irr}} \\ \tilde{\theta}_4^{m+K,\mathrm{irr}} \\ \tilde{\theta}_3^{m+K,\mathrm{irr}} \\ \tilde{\theta}_7^{m+K,\mathrm{irr}} \end{array} $	(0, 0, 0)(0, 0, 0)(0, 1, 3)(0, 2, 6)(0, 2, 6)	$\begin{array}{c} (-1,1,1,1)\\ (-1,1,-1,1)\\ (-1,1,1,1)\\ (-1,1,1,1)\\ (-1,1,-1,1)\end{array}$
-2	$-M\left(-1;\frac{1}{2},\frac{1}{4},\frac{1}{5}\right)$	$20, \{1, 4\}$	(1, 2, 4, 5)	0 2 0 0 2	0 0 1 2 2	$\begin{array}{c} H_1^{m+K} \\ H_1^{m+K,\mathrm{irr}} \\ H_1^{m+K,\mathrm{irr}} \\ H_4^{m+K,\mathrm{irr}} \\ H_2^{m+K,\mathrm{irr}} \\ H_7^{m+K,\mathrm{irr}} \end{array}$		
3	$M\left(-1;\frac{1}{3},\frac{1}{3},\frac{1}{4}\right)$	$12, \{1, 3\}$	(-1, -3, -3, -4)	$     \begin{array}{c}       0 \\       2 \\       0 \\       2     \end{array} $	0 0 1 1	$ \begin{array}{c} \tilde{\theta}_1^{m+K} \\ \tilde{\theta}_9^{m+K} \\ \tilde{\theta}_3^{m+K} \\ \tilde{\theta}_5^{m+K} \end{array} $	$(0,0,0) \\ (0,0,0) \\ (1,0,0) \\ (1,0,0)$	(-1,1,1,1) (-1,1,-1,1) (-1,1,1,1) (-1,1,-1,1)
-3	$-M\left(-1;\frac{1}{3},\frac{1}{3},\frac{1}{4}\right)$	$12, \{1, 3\}$	(1, 3, 3, 4)	$     \begin{array}{c}       0 \\       2 \\       0 \\       2     \end{array} $	0 0 1 1	$\begin{array}{c} \hline H_1^{m+K} \\ H_9^{m+K} \\ H_3^{m+K} \\ H_5^{m+K} \end{array}$		

Table 3: Invariants from Surgeries Along the Figure 8 Knot.

**5.5.1**  $M_3 = -\Sigma(2,3,7), m = 42, K = \{1, 6, 14, 21\}$ 

Using the prescription of Conjecture 5 with r = 1, one obtains the same expression as in (5.16) for the  $\hat{Z}$ -invariants with and without defects.

These are pure mock modular forms, whose shadows coincide with the expectation from the false invariants (5.2)-(5.3), consistent with the False-Mock Conjecture 1 and the Modularity Conjecture 4.2.

**5.5.2** 
$$M_3 = -\Sigma(2, 3, 13), m = 78, K = \{1, 6, 26, 39\}$$

In this case, the corresponding Weil representation has six dimensions and we have  $\sigma^{m+K} = \{1, 5, 7, 11, 17, 23\}$ . All components of the vector-valued false theta function

$$\begin{pmatrix} \tilde{\theta}_{1}^{m+K} \\ \tilde{\theta}_{5}^{m+K} \\ \tilde{\theta}_{5}^{m+K} \\ \tilde{\theta}_{7}^{m+K} \\ \tilde{\theta}_{11}^{m+K} \\ \tilde{\theta}_{23}^{m+K} \end{pmatrix} (\tau)$$

are reproduced in this case by insertion of defects with representations with even highest weights  $\nu_i \in 2\mathbb{Z}$  using the plumbing description for  $\Sigma(2,3,13)$  and Definition 1:

$$q^{-1/2}\hat{Z}_{b_0}(\Sigma(2,3,13);\tau) \sim \tilde{\theta}_7^{m+K} -q^{-3/2}\hat{Z}_{b_0}(\Sigma(2,3,13);W_{(0,0,2)};\tau) \sim \tilde{\theta}_5^{m+K} -q^{-9/2}\hat{Z}_{b_0}(\Sigma(2,3,13);W_{(0,0,4)};\tau) \sim \tilde{\theta}_{17}^{m+K} -q^{-17/2}\hat{Z}_{b_0}(\Sigma(2,3,13);W_{(0,0,6)};\tau) \sim \tilde{\theta}_{23}^{m+K} -q^{-3/2}\hat{Z}_{b_0}(\Sigma(2,3,13);W_{(0,0,8)};\tau) \sim \tilde{\theta}_{11}^{m+K} -q^{-33/2}\hat{Z}_{b_0}(\Sigma(2,3,13);W_{(0,0,10)};\tau) \sim \tilde{\theta}_1^{m+K}.$$
(5.23)

Using the prescription of Conjecture 5 with r = 2, one obtains the following q series

$$\begin{split} \hat{Z}(-\Sigma(2,3,13);\tau) &\sim 1 + q^3 - q^4 + q^5 - q^7 + 2q^8 - 2q^9 + 2q^{10} + \mathcal{O}(q^{11}) \\ \hat{Z}(-\Sigma(2,3,13);W_{(0,0,2)};\tau) &\sim 1 + q^2 + 2q^4 + q^6 + q^7 + q^9 + q^{10} + \mathcal{O}(q^{11}) \\ \hat{Z}(-\Sigma(2,3,13);W_{(0,0,4)};\tau) &\sim 1 + q + 2q^3 - q^4 + 2q^5 + 4q^8 - 2q^9 + 3q^{10} + \mathcal{O}(q^{11}) \\ \hat{Z}(-\Sigma(2,3,13);W_{(0,0,6)};\tau) &\sim 1 + q^2 + q^4 + 2q^5 + 2q^7 + q^9 + 2q^{10} + \mathcal{O}(q^{11}) \\ \hat{Z}(-\Sigma(2,3,13);W_{(0,0,8)};\tau) &\sim 1 + q + q^3 + q^5 + q^6 + 3q^8 - q^9 + 2q^{10} + \mathcal{O}(q^{11}) \\ \hat{Z}(-\Sigma(2,3,13);W_{(0,0,10)};\tau) &\sim 1 + q - q^2 + q^3 + q^4 - q^5 + 2q^6 - q^7 + 2q^8 + q^{10} + \mathcal{O}(q^{11}) \\ \end{split}$$

According to Theorem 2, they are (up to pre-factors) mock modular forms with shadows compatible with the false invariants (5.23), consistent with the False-Mock Conjecture 1 and the Modularity Conjecture 4.2.

It is interesting to note that they do not coincide with the optimal mock theta functions with the same mock modularity as computed in [92] (see Table 18). Consistently, one observes that the q-series above have coefficients growing faster than the "optimal" functions in [92], confirming that the  $\hat{Z}$ -invariants for  $-\Sigma(2,3,13)$  are not optimal in the sense of [92]. **5.5.3**  $M_3 = -\Sigma(2,3,5), m = 30, K = \{1, 6, 10, 15\}$ 

In this case, we propose that

$$\hat{Z}_{b_0}(-\Sigma(2,3,5); W_{\nu}; \tau) = C(q^{-1}) \frac{e(\frac{\chi}{4x})}{2f_{x,\chi}(\tau)} \sum_{\substack{\hat{\epsilon} = (\epsilon_1, \epsilon_2, \epsilon_3) \\ \in (\mathbb{Z}/2)^{\otimes 3}}} (-1)^{\hat{\epsilon}} \Theta_{A, \mathbf{a}_{\hat{\epsilon}}, \mathbf{b}, \mathbf{c}_1, \mathbf{c}_2}(\tau)$$
(5.24)

holds with the regularization given by

$$A = \begin{pmatrix} -60 & 0\\ 0 & 3 \end{pmatrix}, \ \mathsf{c}_1 = \begin{pmatrix} 1\\ 0 \end{pmatrix}, \ \mathsf{c}_2 = \begin{pmatrix} 3\\ 10 \end{pmatrix}, \ \mathsf{b} = \begin{pmatrix} 0\\ \frac{1}{6} \end{pmatrix}, \ \chi = 1 , \tag{5.25}$$

Indeed, one observes that it gives the same leading term expansion as in (5.11). Moreover, one can use the techniques reviewed in §3.3 to compute the shadow of the indefinite theta functions and show that they indeed coincide with the answer in (5.11), obtained using the inverted Habiro series conjecture and coinciding with the order five mock theta functions of Ramanujan<sup>12</sup>. This is of course, consistent with the observation in (5.11). The difference is, the mock modular property is proven and manifest in the present indefinite theta formalism.

# 6 Conclusion and Discussion

In this paper, we shed light on the three important questions regarding the modular and algebraic aspects of the  $\hat{Z}$ -invariants: the  $\widetilde{SL}_2(\mathbb{Z})$  action on the invariants, the construction of mock invariants, and the systematic construction of vertex operator algebras associated with the mock invariants. This work also leads to many more interesting research questions, which we will briefly list in this final section.

- All the proposals, definitions, conjectures, theorems and discussions in this paper should have counterparts for other ADE gauge groups different from SU(2), and three manifolds other than Seifert manifolds with three exceptional fibres. It will be illuminating to develop those explicitly.
- Equipped with the defect invariants introduced in this paper, the resurgence analysis of  $\hat{Z}$ -invariants should reveal more complete structures of complex Chern-Simons theory. In particular, the analysis of the relation to non-Abelian flat connections, analogous to those discussed in [30], should now become more complete with these defect invariants.
- In §3 we compared the indefinite theta function formulation to various other possible approaches to the mock invariants, including the conjectural positive surgery formula of [36] and the Appell-Lerch sum continuation. A very important exercise is to further compare with the interesting approach proposed in [75] using resurgence techniques.

<sup>&</sup>lt;sup>12</sup>An interesting remark concerning the shadow computation is that, unlike in the proof of Theorem 2, where the shadow contribution from the  $c_2$  vector vanishes, here the shadow contribution from the  $c_2$  vector is non-vanishing, being instead proportional to that of  $c_1$ . The same situation is encountered in the case  $-\Sigma(2,5,21)$ .

- In [30], the authors have proposed to understand the associated vertex operator algebras from the point of view of Kazhdan-Lusztig correspondence to quantum groups. In this picture, the VOAs of the false and mock sides should correspond to the negative and positive zones, respectively. It would be interesting to study the cone VOAs we proposed from this perspective. More generally, it would be very illuminating to have a map going directly from the VOA relevant for  $M_3$  to the VOA relevant for  $-M_3$ .
- At least two main mysteries remain surrounding the mock Z-invariants: the exponential singularities near certain cusps that need to be subtracted (cf. (3.1)), the role of the positive-definite lattice (3.26) and the chosen cone. Understanding their physical interpretation will hold the key to unlocking many key aspects of three-dimensional topology and the corresponding 3d SCFTs.

### Acknowledgements

The authors would like to thank Sergei Gukov, Mrunmay Jagadale, Ciprian Manolescu, and Sunghyuk Park for very helpful discussions. We also thank Peter Paule for sharing his implementation of the qZeil algorithm. MC and DP would also like to thank the American Institute of Mathematics for hosting a collaboration meeting on related topics. This work has been supported by the Vidi grant (number 016.Vidi.189.182) from the Dutch Research Council (NWO). The work of P.K. has been supported by the Sonata grant no. 2022/47/D/ST2/02058 funded by the Polish National Science Centre.

# References

- [1] D. Zagier, *Elliptic Modular Forms and Their Applications*. Springer Berlin Heidelberg, Berlin, Heidelberg, 2008.
- [2] E. D'Hoker and J. Kaidi, *Lectures on modular forms and strings*, arXiv:2208.07242.
- [3] S. Zwegers, *Mock Theta Functions*. Ph.d. thesis, Utrecht University, 2002.
- [4] K. Bringmann and K. Ono, The f(q) mock theta function conjecture and partition ranks, Inventiones mathematicae 165 (2006), no. 165 243–266.
- [5] J. Hendrik Bruinier and J. Funke, On Two Geometric Theta Lifts, arXiv Mathematics e-prints (Dec., 2002) math/0212286, [math/0212286].
- [6] D. Zagier, Ramanujan's mock theta functions and their applications (after zwegers and ono-bringmann), Astérisque (2010), no. 326 143–164.
- [7] K. Ono, Unearthing the visions of a master: Harmonic maass forms and number theory, .
- [8] A. Folsom, What is... a mock modular form?, Notices Amer. Math. Soc. 57 (2010), no. 11 1441–1443.
- [9] W. Duke, Almost a century of answering the question: What is a mock theta function?, Notices Amer. Math. Soc. 61 (2014), no. 11 1314–1320.
- [10] M. C. N. Cheng, J. F. R. Duncan, and J. A. Harvey, Umbral moonshine, Commun. Number Theory Phys. 8 (2014), no. 2 101–242.

- [11] M. C. N. Cheng, J. F. R. Duncan, and J. A. Harvey, Umbral moonshine and the niemeier lattices, Research in the Mathematical Sciences 1 (2014), no. 3 1–81.
- [12] K. Bringmann, A. Folsom, K. Ono, and L. Rolen, *Harmonic Maass Forms and Mock Modular Forms: Theory and Applications*, vol. 64. AMS, Colloquium Publications, 2017.
- [13] M. C. N. Cheng, F. Ferrari, and G. Sgroi, Three-manifold quantum invariants and mock theta functions, Phil. Trans. Roy. Soc. Lond. 378 (2019), no. 2163 20180439, [arXiv:1912.07997].
- [14] A. Dabholkar and P. Putrov, Three Avatars of Mock Modularity, Int. J. Mod. Phys. A 36 (2021), no. 36 2130020, [arXiv:2110.09790].
- [15] V. G. Kac and M. Wakimoto, Representations of affine superalgebras and mock theta functions, 2013.
- [16] V. G. Kac and M. Wakimoto, Representations of affine superalgebras and mock theta functions ii, 2014.
- [17] V. G. Kac and M. Wakimoto, Representations of affine superalgebras and mock theta functions. III, Izvestiya: Mathematics 80 (aug, 2016) 693–750.
- [18] C. Alfes and T. Creutzig, The mock modular data of a family of superalgebras, 2012.
- [19] M.-F. Vignéras, Séries theta des formes quadratiques indéfinies, .
- [20] J. Lovejoy and R. Osburn, q-hypergeometric double sums as mock theta functions, Pacific Journal of Mathematics 264 (2013), no. 1 151–162.
- [21] J. Males, A. Mono, and L. Rolen, Higher depth mock theta functions and q-hypergeometric series, in Forum Mathematicum, vol. 33, pp. 857–866, De Gruyter, 2021.
- [22] S. Alexandrov and B. Pioline, Black holes and higher depth mock modular forms, 1808.08479.
- [23] G. Korpas, J. Manschot, G. W. Moore, and I. Nidaiev, Mocking the u-plane integral, arXiv:1910.13410.
- [24] G. Korpas and J. Manschot, Donaldson-Witten theory and indefinite theta functions, JHEP 11 (2017) 083, [arXiv:1707.06235].
- [25] M. C. N. Cheng and G. Sgroi, Cone vertex algebras, mock theta functions, and umbral moonshine modules, arXiv: 2203.03052.
- [26] J. Duncan and J. Harvey, The umbral moonshine module for the unique unimodular niemeier root system, Algebra & Samp; Number Theory 11 (may, 2017) 505–535.
- [27] S. Ramanujan, Collected Papers of Srinivasa Ramanujan. AMS Chelsea Publishing, Providence, RI, 2000.
- [28] M. Griffin, K. Ono, and L. Rolen, Ramanujan's mock theta functions, Proc. Natl. Acad. Sci. USA 110 (2013), no. 15 5765–5768.
- [29] D. Zagier, Quantum modular forms, in Quanta of maths, vol. 11 of Clay Math. Proc., pp. 659–675. Amer. Math. Soc., Providence, RI, 2010.
- [30] M. C. N. Cheng, S. Chun, F. Ferrari, S. Gukov, and S. M. Harrison, 3d modularity, 1809.10148.

- [31] M. C. N. Cheng, I. Coman, D. Passaro, and G. Sgroi, *Quantum Modular*  $\hat{Z}^{G}$ -Invariants, arXiv:2304.03934.
- [32] L. Crane and I. B. Frenkel, Four-dimensional topological quantum field theory, Hopf categories, and the canonical bases, Journal of Mathematical Physics 35 (Oct., 1994) 5136–5154. \_eprint: https://pubs.aip.org/aip/jmp/article-pdf/35/10/5136/8163625/5136\_1\_online.pdf.
- [33] S. Gukov, P. Putrov, and C. Vafa, Fivebranes and 3-manifold homology, JHEP 07 (2017) 071, [arXiv:1602.05302].
- [34] S. Gukov, D. Pei, P. Putrov, and C. Vafa, BPS spectra and 3-manifold invariants, J. Knot Theory Ramif. 29 (2020), no. 02 2040003, [arXiv:1701.06567].
- [35] S. Gukov and C. Manolescu, A two-variable series for knot complements, 1904.06057.
- [36] S. Park, Inverted state sums, inverted habiro series, and indefinite theta functions, arXiv:2106.03942.
- [37] M. C. N. Cheng, S. Chun, B. Feigin, F. Ferrari, S. Gukov, S. M. Harrison, and D. Passaro, 3-Manifolds and VOA Characters, arXiv:2201.04640.
- [38] H.-J. Chung, BPS invariants for a Knot in Seifert manifolds, JHEP 12 (2022) 122, [arXiv:2201.08351].
- [39] R. Lawrence and D. Zagier, Modular forms and quantum invariants of 3-manifolds, Asian Journal of Mathematics 3 (1999) 93–108.
- [40] S. Gukov, M. Marino, and P. Putrov, *Resurgence in complex Chern-Simons theory*, arXiv:1605.07615.
- [41] K. Hikami Regular and Chaotic Dynamics 10 (2005), no. 4 509.
- [42] K. Hikami, On the quantum invariant for the Brieskorn homology spheres, Int. J. Math. 16 (2005) 661–684, [math-ph/0405028].
- [43] K. Hikami, Quantum invariant for torus link and modular forms, Commun. Math. Phys. 246 (2004), no. 2 403–426, [math-ph/0305039].
- [44] K. Bringmann, K. Mahlburg, and A. Milas, Quantum modular forms and plumbing graphs of 3-manifolds, J. Combin. Theory Ser. A 170 (2020) 105145, 32, [arXiv:1810.05612].
- [45] K. Bringmann, J. Kaszian, and A. Milas, Vector-valued higher depth quantum modular forms and higher Mordell integrals, arXiv e-prints (Mar., 2018) arXiv:1803.06261, [arXiv:1803.06261].
- [46] K. Bringmann, K. Mahlburg, and A. Milas, Higher depth quantum modular forms and plumbed 3-manifolds, arXiv:1906.10722.
- [47] K. Bringmann, J. Kaszian, A. Milas, and C. Nazaroglu, Integral Representations of Rank Two False Theta Functions and Their Modularity Properties, arXiv:2101.02902.
- [48] K. Bringmann, J. Kaszian, A. Milas, and C. Nazaroglu, Higher depth false modular forms, Commun. Contemp. Math. 25 (2023), no. 07 2250043, [arXiv:2109.00394].
- [49] J. Chae, Witt invariants from q-series Z, Lett. Math. Phys. 113 (2023), no. 1 3, [arXiv:2204.02794].

- [50] H.-J. Chung, BPS invariants for Seifert manifolds, arXiv:1811.08863.
- [51] P. Kucharski,  $\hat{Z}$  invariants at rational  $\tau$ , JHEP **09** (2019) 092, [arXiv:1906.09768].
- [52] H.-J. Chung, BPS invariants for 3-manifolds at rational level K, arXiv:1906.12344.
- [53] H.-J. Chung, Index for a model of 3d-3d correspondence for plumbed 3-manifolds, arXiv:1912.13486.
- [54] S. Park, Higher rank Z and  $F_K$ , SIGMA 16 (2020) 044, [arXiv:1909.13002].
- [55] F. Costantino, S. Gukov, and P. Putrov, Non-Semisimple TQFT's and BPS q-Series, SIGMA 19 (2023) 010, [arXiv:2107.14238].
- [56] S. Gukov, S. Park, and P. Putrov, Cobordism invariants from BPS q-series, Annales Henri Poincaré 22 (jul, 2021) 4173–4203.
- [57] A. Mori and Y. Murakami, Witten-Reshetikhin-Turaev Invariants, Homological Blocks, and Quantum Modular Forms for Unimodular Plumbing H-Graphs, SIGMA 18 (2022) 034, [arXiv:2110.10958].
- [58] T. Matsusaka and Y. Terashima, Modular transformations of homological blocks for Seifert fibered homology 3-spheres, arXiv:2112.06210.
- [59] Y. Murakami, Witten-Reshetikhin-Turaev invariants and homological blocks for plumbed homology spheres, arXiv:2205.01282.
- [60] Y. Murakami, Witten-Reshetikhin-Turaev invariants and indefinite false theta functions for plumbing indefinite H-graphs, arXiv:2212.09972.
- [61] Y. Murakami, A proof of a conjecture of Gukov-Pei-Putrov-Vafa, arXiv:2302.13526.
- [62] M. C. Cheng, S. Chun, B. Feigin, F. Ferrari, S. Gukov, S. M. Harrison, and D. Passaro, 3-manifolds and voa characters, arXiv preprint arXiv:2201.04640 (2022).
- [63] D. Gaiotto and M. Rapčák, Vertex algebras at the corner, 2018.
- [64] D. Gaiotto and M. Rapcak, Miura operators, degenerate fields and the m2-m5 intersection, 2020.
- [65] D. Zagier, Vassiliev invariants and a strange identity related to the dedekind eta-function, Topology 40 (2001), no. 5 945–960.
- [66] D. Zagier, Quantum modular forms, .
- [67] Y. Zhu, Modular invariance of characters of vertex operator algebras, Journal of the American Mathematical Society 9 (1996), no. 1 237–302.
- [68] C. Dong, H. Li, and G. Mason, Modular-invariance of trace functions in orbifold theory and generalized moonshine, Communications in Mathematical Physics 214 (2000) 1–56.
- [69] T. Creutzig, A. Milas, and S. Wood, On regularised quantum dimensions of the singlet vertex operator algebra and false theta functions, 2016.
- [70] K. Bringmann and A. Milas, W-algebras, higher rank false theta functions, and quantum dimensions, Selecta Math. (N.S.) 23 (2017), no. 2 1249–1278.
- [71] K. Bringmann and A. Milas, W-algebras, false theta functions and quantum modular forms, i, International Mathematics Research Notices 2015 (02, 2015).

- [72] K. Bringmann, J. Kaszian, and A. Milas, Higher depth quantum modular forms, multiple Eichler integrals, and sl<sub>3</sub> false theta functions, Res. Math. Sci. 6 (2019), no. 2 Paper No. 20, 41.
- [73] K. Bringmann, J. Kaszian, A. Milas, and C. Nazaroglu, Higher depth false modular forms, arXiv:2109.00394.
- [74] T. Creutzig and D. Ridout, Logarithmic Conformal Field Theory: Beyond an Introduction, J. Phys. A 46 (2013) 4006, [arXiv:1303.0847].
- [75] O. Costin, G. V. Dunne, A. Gruen, and S. Gukov, Going to the other side via the resurgent bridge, arXiv preprint arXiv:2310.12317 (2023).
- [76] D. Choi, S. Lim, and R. C. Rhoades, Mock modular forms and quantum modular forms, Proc. Amer. Math. Soc. 144 (2016), no. 6 2337–2349.
- [77] T. Ekholm, A. Gruen, S. Gukov, P. Kucharski, S. Park, and P. Sulkowski, Z at large N: from curve counts to quantum modularity, Commun. Math. Phys. 396 (2022), no. 1 143–186, [arXiv:2005.13349].
- [78] T. Ekholm, A. Gruen, S. Gukov, P. Kucharski, S. Park, M. Stosic, and P. Sulkowski, Branches, quivers, and ideals for knot complements, J. Geom. Phys. 177 (2021) 104520, [arXiv:2110.13768].
- [79] K. Habiro, On the quantum sl<sub>2</sub> invariants of knots and integral homology spheres, Geometry & Topology Monographs 4 (2002) 55–68.
- [80] S. Park, 3-manifolds, q-series, and topological strings. PhD thesis, California Institute of Technology, 2022.
- [81] P. Paule and A. Riese, A mathematica q-analogue of zeilberger's algorithm based on an algebraically motivated approach to q-hypergeometric telescoping, Special functions, q-series and related topics 14 (1997) 179–210.
- [82] Cappelli, Andrea and Itzykson, Claude and Zuber, Jean-Bernard, The A-D-E classification of minimal and A<sub>1</sub><sup>(1)</sup> conformal invariant theories, Communications in Mathematical Physics **113** (1987) 1–26.
- [83] N. Skoruppa, Über den Zusammenhang zwischen Jacobiformen und Modulformen halbganzen Gewichts. Bonner mathematische Schriften. Universität Bonn, 1985.
- [84] N.-P. Skoruppa, Jacobi Forms of Critical Weight and Weil Representations, arXiv e-prints (July, 2007) arXiv:0707.0718, [arXiv:0707.0718].
- [85] A. Folsom, Quantum jacobi forms in number theory, topology, and mathematical physics, Research in the Mathematical Sciences 25 (2019).
- [86] A. Dabholkar, S. Murthy, and D. Zagier, Quantum black holes, wall crossing, and mock modular forms, 1208.4074.
- [87] K. Bringmann, K. Mahlburg, and A. Milas, Quantum modular forms and plumbing graphs of three-manifolds, 1810.05612.
- [88] S. Zwegers, Mock Theta Functions, arXiv e-prints (July, 2008) arXiv:0807.4834, [arXiv:0807.4834].
- [89] K. Bringmann, A. Folsom, K. Ono, and L. Rolen, Harmonic Maass Forms and Mock Modular Forms: Theory and Applications, Amer. Math. Soc. Colloq. Publ. 73 (2021), no. 1 108–111.

- [90] S. Alexandrov, S. Banerjee, J. Manschot, and B. Pioline, Indefinite theta series and generalized error functions, Selecta Math. 24 (2018) 3927, [arXiv:1606.05495].
- [91] C. Nazaroglu, r-tuple error functions and indefinite theta series of higher-depth, Communications in Number Theory and Physics 12 (2018), no. 3 581–608.
- [92] M. C. N. Cheng and J. F. R. Duncan, Optimal Mock Jacobi Theta Functions, Adv. Math. 372 (2020) 107284, [arXiv:1605.04480].
- [93] J. H. Conway and S. P. Norton, Monstrous Moonshine, Bull. London Math. Soc. 11 (1979), no. 3 308–339.

# A Mock Modularity Proofs

In this section we will prove Theorem 2 and Proposition 3. As a result, we will be studying the properties of the modular completion  $\hat{\Theta}_{A,\mathbf{a}_{\hat{e}},\mathbf{b},\mathbf{c}_1,\mathbf{c}_2}(\tau)$  of indefinite theta functions  $\Theta_{A,\mathbf{a}_{\hat{e}},\mathbf{b},\mathbf{c}_1,\mathbf{c}_2}(\tau)$ , specified by (3.17), for the choices of quadratic forms, cones, and vectors given in (3.47).

We consider, for  $(p_1, p_2, p_3)$  three co-prime integers,

$$A = \begin{pmatrix} -2m & 0\\ 0 & x \end{pmatrix}, \ \mathsf{c}_1 = \begin{pmatrix} 1\\ 0 \end{pmatrix}, \ \mathsf{c}_2 = \begin{pmatrix} x\\ 2p_3 \end{pmatrix}, \ \mathsf{b} = \begin{pmatrix} 0\\ \frac{1}{2x} \end{pmatrix}, \ \mathsf{a}_{\boldsymbol{\nu},\hat{\epsilon}} = \begin{pmatrix} \frac{r_{\boldsymbol{\nu},\hat{\epsilon}}}{2m}\\ -\frac{\chi}{2x} \end{pmatrix}$$
(A.1)

where  $r_{\boldsymbol{\nu},\hat{\epsilon}} = m - \sum_i (-1)^{\epsilon_i} (1+\nu_i) \bar{p}_i$  and we let  $m = p_1 p_2 p_3$ ,  $\bar{p}_i := m/p_i$  as in the proof of Theorem 1.

## A.1 The Mock Modular Properties for $\Sigma(2,3,6r+1)$

From (3.22), we define

$$S_{j,\mathbf{l},\mathbf{a}} := e\left(\frac{d_j B(\mathbf{c}_j, \mathbf{b}) B(\mathbf{c}_j, \mathbf{l} + \mathbf{a})}{2n_j m}\right) \theta_{n_j m, -d_j B(\mathbf{c}_j, \mathbf{l} + \mathbf{a})}^1 \left(\frac{\tau}{d_j}\right) \frac{1}{\nu \in (\mathbf{l} + \mathbf{a})_j^\perp + \mathbb{Z} \mathbf{c}_{j,\perp}} e(B(\nu, \mathbf{b})) q^{\frac{|\nu|^2}{2}}$$
(A.2)

for  $j \in \{1, 2\}$ ,  $l \in P_j$  (see the text after (3.17) for the definition of  $P_j$ ), so that the function

$$F_{(p_1,p_2,p_3),\boldsymbol{\nu}} := \frac{1}{2f_{x,\chi}(\tau)} \sum_{\substack{\hat{\epsilon} = (\epsilon_1,\epsilon_2,\epsilon_3)\\ \in (\mathbb{Z}/2)^{\otimes 3}}} (-1)^{\hat{\epsilon}} \Theta_{A,\mathbf{a}_{\boldsymbol{\nu},\hat{\epsilon}},\mathbf{b},\mathbf{c}_1,\mathbf{c}_2}(\tau)$$
(A.3)

has completion  $\hat{F}_{(p_1,p_2,p_3),\boldsymbol{\nu}}$  that satisfies

$$2im\sqrt{2\operatorname{Im}(\tau)} \,\overline{\frac{\partial}{\partial\bar{\tau}}} \hat{F}_{(p_1,p_2,p_3),\boldsymbol{\nu}}(\tau) = \frac{1}{2\overline{f_{x,\chi}(\tau)}} \sum_{j=1,2} (-1)^j \frac{d_j}{n_j} \sum_{\substack{\mathsf{l}\in P_j\\\hat{\epsilon}\in(\mathbb{Z}/2)^{\otimes 3}}} (-1)^{\sum_{i=1}^3 \epsilon_i} S_{j,\mathsf{l},\mathsf{a}_{\boldsymbol{\nu},\hat{\epsilon}}}.$$
(A.4)

To analyse the above, we will compute the summand with j = 1 and j = 2 separately. Lemma 2.

$$-\frac{\mathrm{e}\left(-\frac{\chi}{4x}\right)}{2\overline{f_{x,\chi}(\tau)}}\frac{d_j}{n_j}\sum_{\substack{\mathbf{l}\in P_1\\\hat{\epsilon}\in(\mathbb{Z}/2)^{\otimes 3}}}(-1)^{\sum_{i=1}^3\epsilon_i}S_{1,\mathbf{l},\mathbf{a}_{\boldsymbol{\nu},\hat{\epsilon}}}=\theta_{r_{\boldsymbol{\nu}}}^{1,m+K}$$
(A.5)

where  $m, K, r_{\nu}$  are as in Theorem 1.

*Proof.* For  $c_1 = (1,0)^T$ , we have  $n_1 = d_1 = 1$  and  $P_1 \cong \mathbb{Z}^2/(c_j\mathbb{Z} \oplus c_j^{\perp}\mathbb{Z}) \cong \{0\}$ . As a result, the left-hand side of (A.5) reduces to

$$-\frac{\mathrm{e}\left(-\frac{\chi}{4x}\right)}{2\overline{f}_{x,\chi}(\tau)}\sum_{\hat{\epsilon}\in(\mathbb{Z}/2)^{\otimes 3}}(-1)^{\sum_{i=1}^{3}\epsilon_{i}}S_{1,0,\mathbf{a}_{\nu,\hat{\epsilon}}}$$

$$=-\frac{\mathrm{e}\left(-\frac{\chi}{4x}\right)}{2\overline{f}_{x,\chi}(\tau)}\sum_{\hat{\epsilon}\in(\mathbb{Z}/2)^{\otimes 3}}(-1)^{\sum_{i=1}^{3}\epsilon_{i}}\theta_{m,-r_{\nu,\hat{\epsilon}}}^{1}}\overline{\sum_{k\in\mathbb{Z}}(-1)^{k}\mathrm{e}\left(-\frac{\chi}{4x}\right)q^{\frac{x}{2}(k-\frac{\chi}{2x})^{2}}}$$

$$=\sum_{\substack{(\epsilon_{1},\epsilon_{2},\epsilon_{3})\in(\mathbb{Z}/2)^{\otimes 3}\\\sum_{i}\epsilon_{j}\equiv 0\pmod{2}}}\theta_{m,r_{\nu,\hat{\epsilon}}}^{1}=\theta_{m,r_{\nu}}^{1}}$$

where we have used the expression for the theta function  $f_{x,\chi}$  (3.28) in the first equality, and used (2.46) in the last equality.

Next we study the contribution, proportional to

$$\sum_{\substack{\mathbf{l}\in P_2\\\hat{\epsilon}\in (\mathbb{Z}/2)^{\otimes 3}}} (-1)^{\sum_{i=1}^3 \epsilon_i} S_{2,\mathbf{l},\mathbf{a}_{\boldsymbol{\nu},\hat{\epsilon}}}$$
(A.6)

from the j = 2 summand in (A.4). For this purpose, note that we can take

$$\mathsf{c}_{2,\perp} = \begin{pmatrix} 1\\ \bar{p}_3 \end{pmatrix}$$

from which we see

$$P_2 \cong \left\{ \begin{pmatrix} 0\\ \ell \end{pmatrix}, \ell \in \mathbb{Z}/L\mathbb{Z} \right\}, \ L = \bar{p}_3 x - 2p_3 = -\frac{|\mathbf{c}_2|^2}{2p_3 x} = \frac{|\mathbf{c}_{2,\perp}|^2}{\bar{p}_3}.$$

Write

$$S_{2,\mathbf{l},\mathbf{a}} := e\left(\frac{d_2 B(\mathbf{c}_j, \mathbf{b})}{2n_2 m} X_{\ell,\hat{\epsilon}}\right) \theta_{n_2 m, -d_2 X_{\ell,\hat{\epsilon}}}^1 \left(\frac{\tau}{d_2}\right) \overline{e\left(\frac{A_{\ell,\hat{\epsilon}}}{2L}\right) \sum_{k \in \mathbb{Z}} q^{\frac{|\mathbf{c}_{2,\perp}|^2}{2}(k + \frac{A_{\ell,\hat{\epsilon}}}{L\bar{p}_3})^2}}$$
(A.7)

where we have assumed  $\bar{p}_3/2 \in \mathbb{Z}$  (which is the case for Lemma 3), and defined for a given  $\nu$ 

$$X_{\ell,\hat{\epsilon}} := B(\mathsf{c}_2, \mathsf{I} + \mathsf{a}_{\boldsymbol{\nu},\hat{\epsilon}}), \quad A_{\ell,\hat{\epsilon}} := B(\mathsf{c}_{2,\perp}, \mathsf{I} + \mathsf{a}_{\boldsymbol{\nu},\hat{\epsilon}}), \quad \text{with } \mathsf{I} = \begin{pmatrix} 0\\\ell \end{pmatrix}.$$
(A.8)

As a result, the j = 2 piece vanishes if there exists a bijection  $\Gamma$  on  $(\mathbb{Z}/2)^{\otimes 3} \times P_2$  such that

$$\Gamma(\hat{\epsilon}, \ell) = (\Gamma_1(\hat{\epsilon}), \Gamma_2(\ell))$$

has the property that it preserves  $X_{\ell,\hat{\epsilon}}$ , and maps  $\frac{1}{L\bar{p}_3}A_{\ell,\hat{\epsilon}} \mapsto \pm \frac{1}{L\bar{p}_3}A_{\ell,\hat{\epsilon}} \pmod{\mathbb{Z}}$ , and is moreover odd on  $(-1)^{\hat{\epsilon}} e\left(\frac{A_{\ell,\hat{\epsilon}}}{2L}\right)$ .

Lemma 3.

$$\sum_{\substack{\mathbf{l}\in P_2\\\hat{\epsilon}\in (\mathbb{Z}/2)^{\otimes 3}}} (-1)^{\hat{\epsilon}} S_{2,\mathbf{l},\mathbf{a}_{\boldsymbol{\nu},\hat{\epsilon}}} = 0 \tag{A.9}$$

where  $m, K, r_{\nu}$  are as in Theorem 1, when  $p_2 = 2, p_3 = xp_1 - 2$ , and x is an odd positive integer number.

*Proof.* In this case, we have L = 4 and the above expression in A.9 is equal to

$$\sum_{\substack{\ell \in \mathbb{Z}/4\mathbb{Z}\\\hat{\epsilon} \in (\mathbb{Z}/2)^{\otimes 3}}} (-1)^{\hat{\epsilon}} e\left(\frac{d_2 B(\mathsf{c}_j,\mathsf{b})}{2n_2 m} X_{\ell,\hat{\epsilon}}\right) \, \theta_{n_2 m, -d_j X_{\ell,\hat{\epsilon}}}^1\left(\frac{\tau}{d_2}\right) \, e\left(\frac{A_{\ell,\hat{\epsilon}}}{2L}\right) \sum_{k \in \mathbb{Z}} q^{\frac{|\mathsf{c}_{2,\perp}|^2}{2}(k + \frac{A_{\ell,\hat{\epsilon}}}{L\bar{p}_3})^2} = 0.$$

From the above discussion, let us first study the symmetries of  $X_{\ell,\hat{\epsilon}}$ . From

$$X_{\ell,\hat{\epsilon}} = -p_3\chi - mx + x\left(2p_3\ell + \sum_i (-1)^{\epsilon_i} A_i \bar{p}_i\right), \quad A_i = 1 + \nu_i$$

We see that  $X_{\ell,\hat{\epsilon}}$  is invariant under

$$\epsilon_j \mapsto \epsilon_j + \delta_{i,j}, \ \ell \mapsto \ell + \Delta \ell_i$$

for i = 1, 2, with

$$\Delta \ell_i = (-1)^{\epsilon_i} A_i \frac{m}{p_i p_3}.$$

Case 1: odd  $A_2$ From

$$A_{\ell,\hat{\epsilon}} = -m - \sum_{i} (-1)^{\epsilon_i} A_i \bar{p}_i + 2p_1 x\ell - p_1 \chi,$$

it is easy to check that

$$\begin{aligned} A_{\ell,\hat{\epsilon}} + A_{\ell+\Delta\ell_1,(\epsilon_1+1,\epsilon_2,\epsilon_3)} &= 4p_1 \left( -\frac{\chi}{2} + (-1)^{\epsilon_2} \frac{A_2 p_3}{2} - p_3 + \ell x + (-1)^{\epsilon_1} x A_1 + (-1)^{\epsilon_3} A_3 \right) \\ &= 4p_1 \times \\ \left( -\frac{\chi}{2} + (-1)^{\epsilon_2} \frac{x A_2 p_1}{2} - (-1)^{\epsilon_2} A_2 - p_3 + \ell x + (-1)^{\epsilon_1} x A_1 + (-1)^{\epsilon_3} A_3 \right) \\ A_{\ell,\hat{\epsilon}} + A_{\ell+\Delta\ell_1+\Delta\ell_2,(\epsilon_1+1,\epsilon_2+1,\epsilon_3)} &= 4p_1 \left( -\frac{\chi}{2} + (-1)^{\epsilon_2} \frac{x A_2 p_1}{2} - p_3 + \ell x + (-1)^{\epsilon_1} x A_1 + (-1)^{\epsilon_3} A_3 \right) \end{aligned}$$

which shows that, when  $A_2 \equiv 1 \pmod{2}$ , we have

$$A_{\ell,\hat{\epsilon}} + A_{\ell+\Delta\ell_1,(\epsilon_1+1,\epsilon_2,\epsilon_3)} \equiv A_{\ell,\hat{\epsilon}} + A_{\ell+\Delta\ell_1+\Delta\ell_2,(\epsilon_1+1,\epsilon_2+1,\epsilon_3)} \equiv 0 \pmod{4p_1}$$

and

$$\frac{A_{\ell,\hat{\epsilon}} + A_{\ell+\Delta\ell_1,(\epsilon_1+1,\epsilon_2,\epsilon_3)}}{4p_1} + 1 \equiv \frac{A_{\ell,\hat{\epsilon}} + A_{\ell+\Delta\ell_1+\Delta\ell_2,(\epsilon_1+1,\epsilon_2+1,\epsilon_3)}}{4p_1} \pmod{2}.$$

Moreover, from

$$A_{\ell,\hat{\epsilon}} = \frac{1}{2} \left( A_{\ell,\hat{\epsilon}} + A_{\ell+\Delta\ell_1,(\epsilon_1+1,\epsilon_2,\epsilon_3)} \right) - 4(-1)^{\epsilon_1} A_1$$

we see that half of  $\ell \in \mathbb{Z}/4$  satisfies  $A_{\ell,\hat{\epsilon}} + A_{\ell+\Delta\ell_1,(\epsilon_1+1,\epsilon_2,\epsilon_3)} \equiv 0 \pmod{L\bar{p}_3} = 8p_1$ , and simultaneously  $A_{\ell,\hat{\epsilon}} \equiv 0 \pmod{4}$ , while the other satisfies  $A_{\ell,\hat{\epsilon}} + A_{\ell+\Delta\ell_1+\Delta\ell_2,(\epsilon_1+1,\epsilon_2+1,\epsilon_3)} \equiv 0 \pmod{L\bar{p}_3} = 8p_1$  and simultaneously  $A_{\ell,\hat{\epsilon}} \equiv 2 \pmod{4}$ . Hence, we see that in the former case, the transformation  $\Gamma(\hat{\epsilon},\ell) = ((\epsilon_1+1,\epsilon_2,\epsilon_3),\ell+\Delta\ell_1)$  satisfies the condition below (A.8) and in the latter case  $\Gamma(\hat{\epsilon},\ell) = ((\epsilon_1+1,\epsilon_2+1,\epsilon_3),\ell+\Delta\ell_1+\Delta\ell_2)$  does the job. Case 2: even  $A_2$ Note

$$A_{\ell+\Delta\ell_2,(\epsilon_1,\epsilon_2+1,\epsilon_3)} - A_{\ell,\hat{\epsilon}} = 4p_1(-1)^{\epsilon_2}A_2(xp_1-1)$$
(A.10)

so  $A_{\ell+\Delta\ell_2,(\epsilon_1,\epsilon_2+1,\epsilon_3)} \equiv A_{\ell,\hat{\epsilon}} \pmod{\bar{p}_3 L} = 8p_1$  when  $A_2$  is even. We see that the above transformation satisfies the condition below (A.8).

Combining the above, we have proven the statement of the Lemma.

With the two lemmas 2-3, we see that the function (A.3) is a pure mock modular form, and in particular Theorem 2 follows.

## A.2 Mixed Mock Modularity of $\Sigma(s, t, str + 1)$

In this subsection we will prove Proposition 3. Generically, equation (3.17) implies that the indefinite theta series (3.34) will produce mixed mock modular forms. Lemma 2 establishes the contribution from  $c_1$  to the shadow function. The  $c_2$  contribution, on the other hand, is given by

$$\sum_{\substack{\mathbf{l}\in P_2\cong\mathbb{Z}/L\mathbb{Z}\\\hat{\epsilon}\in(\mathbb{Z}/2)^{\otimes 3}}} (-1)^{\sum_{i=1}^3 \epsilon_i} S_{2,\mathbf{l},\mathbf{a}_{\boldsymbol{\nu},\hat{\epsilon}}}$$
(A.11)

with  $S_{2,l,a_{\nu,\ell}}$  given by (A.7). Compared with the notation of Proposition 3, we see that

$$h_{\hat{\epsilon},\ell} = \frac{\sum_{k \in \mathbb{Z}} q^{\frac{|c_{2,\perp}|^2}{2}(k + \frac{A_{\ell,\hat{\epsilon}}}{L\bar{p}_3})^2}}{f_{x,\chi}(\tau)} = \frac{\sum_{k \in \mathbb{Z}} q^{\frac{L\bar{p}_3}{2}(k + \frac{A_{\ell,\hat{\epsilon}}}{L\bar{p}_3})^2}}{f_{x,\chi}(\tau)}$$

and  $\theta_{\hat{\epsilon},\ell}$  is proportional to  $\theta_{n_2m,-d_jX_{\ell,\hat{\epsilon}}}^1\left(\frac{\tau}{d_2}\right)$ .

As discussed in (3.1), we would like to study the radial limit  $\tau \to \alpha + i0^+$  of the function  $h_{\hat{\epsilon},\ell}$  (while  $\theta_{\hat{\epsilon},\ell}$  is manifestly finite in the radial limit). Note that  $h_{\hat{\epsilon},\ell}$  is a ratio of two theta functions of two distinct one-dimensional lattices. From the modular transformation of lattice theta functions, we see that the modular images of the numerator is a linear combination of theta functions of the form  $\sum_{k \in \mathbb{Z}} q^{\frac{L\bar{p}_3}{2}(k+\frac{a}{L\bar{p}_3})^2}$  for  $a \in \mathbb{Z}/(L\bar{p}_3)$ , and the modular images of the denominator is a linear combination of theta functions of the form  $f_{x,b}(\tau) := \sum_{k \in \mathbb{Z}} (-1)^k q^{\frac{x}{2}(k-\frac{b}{2x})^2}$ . Using the fact that  $SL_2(\mathbb{Z})$  acts on the cusps  $\mathbb{Q} \cup \{i\infty\}$  transitively, the asymptotic behaviour of a function near any cusp can be analysed in terms of the asymptotic behaviour near the cusp  $q \to 0$  ( $\tau \to i\infty$ ) of the modular transformation of the same function. From

$$\frac{L\bar{p}_3}{4x} = \frac{st(st-2)}{4(2r+1)}$$

it is easy to see that there is no  $a \in \mathbb{Z}$  such that

$$\frac{a^2}{L\bar{p}_3} = \frac{1}{4x}$$

and therefore  $h_{\hat{e},\ell}$  either vanishes or has exponential singularities near any cusp. This finishes the proof for Proposition 3.

# **B** Polynomials $P_n^{p,b}$

The  $P_n^{p,b}$  polynomials that were used to compute  $\hat{Z}$  invariants through conjecture 3 can be easily computed via

$$P_n^{p,b}(q^{-1}) = q^{\frac{b(2p-b)}{4p} - n^2} \sum_{k=0}^{\infty} \frac{q^{-nk} \left(q^{n+1};q\right)_n \left(q^{k+1};q\right)_{2n}}{(q;q)_{2n}} \times \left(q^{\frac{(n+k)^2}{p}} \delta_{n+k-b/2 \pmod{p}} - q^{\frac{(n+k+1)^2}{p}} \delta_{n+k+1-b/2 \pmod{p}}\right)$$
(B.1)

for even b and

$$P_{n}^{p,b}(q^{-1}) = q^{\frac{b(2p-b)}{4p} - n^{2} - n} \sum_{k=0}^{\infty} \frac{q^{-nk} (q^{n};q)_{n} (q^{k+1};q)_{2n}}{(q;q)_{2n}} \times \left( q^{\frac{(n+k-1/2)^{2}}{p}} \delta_{n+k-1/2-b/2 \pmod{p}} - q^{\frac{(n+k+3/2)^{2}}{p}} \delta_{n+k+3/2-b/2 \pmod{p}} \right)$$
(B.2)

for odd b.

We record the coefficient lists of the first few  $P_n^{p,b}$  in variable q and starting with  $q^0$ . Note that  $P_1^{p,b} = 1$  for all p and b and it is hence not listed in the tables.

Table 4: The coefficient lists of	$P_n^{z,v}$	•
-----------------------------------	-------------	---

b	n	coefficients
	2	1, 0, 1
	3	1, 0, 1, 1, 1
0	4	1, 0, 1, 1, 2, 1, 1, 0, 1
	5	1, 0, 1, 1, 2, 2, 2, 1, 2, 1, 1, 1, 1
	6	1, 0, 1, 1, 2, 2, 3, 2, 3, 2, 3, 2, 3, 2, 3, 2, 2, 1, 1, 0, 1
	2	1,1
	3	1, 1, 1, 1
1	4	1, 1, 1, 2, 1, 1, 1
	5	1, 1, 1, 2, 2, 2, 2, 2, 1, 1, 1
	6	1, 1, 1, 2, 2, 3, 3, 3, 3, 3, 3, 3, 2, 2, 1, 1, 1
	2	1,1
	3	1, 1, 1, 0, 1
2	4	1, 1, 1, 1, 1, 1, 1, 1
	5	1, 1, 1, 1, 2, 1, 2, 2, 2, 1, 1, 0, 1
	6	1, 1, 1, 1, 2, 2, 2, 3, 3, 3, 3, 3, 2, 2, 2, 1, 1, 1, 1
	2	1,1
	3	1, 1, 1, 1
3	4	1, 1, 1, 2, 1, 1, 1
	5	1, 1, 1, 2, 2, 2, 2, 2, 1, 1, 1
	6	1, 1, 1, 2, 2, 3, 3, 3, 3, 3, 3, 3, 2, 2, 1, 1, 1

b	n	coefficients
	2	1, 0, 1, 1
	3	1, 0, 1, 2, 2, 1, 2
0	4	1, 0, 1, 2, 3, 3, 4, 3, 3, 3, 2, 1, 1
	5	1, 0, 1, 2, 3, 4, 6, 5, 7, 7, 8, 7, 8, 5, 6, 4, 3, 2, 2
	6	1, 0, 1, 2, 3, 4, 7, 7, 9, 11, 13, 14, 17, 16, 18, 18, 17, 15, 16, 13, 11, 9, 7, 5, 5, 2, 1, 1
	2	1, 1, 1
	3	1, 1, 2, 2, 2, 1
1	4	1, 1, 2, 3, 4, 4, 4, 3, 3, 1, 1
	5	1, 1, 2, 3, 5, 6, 7, 8, 9, 8, 8, 7, 6, 4, 3, 2, 1
	6	1, 1, 2, 3, 5, 7, 9, 11, 14, 15, 18, 18, 20, 19, 19, 17, 16, 13, 11, 8, 7, 4, 3, 1, 1
	2	1, 1, 1
2	3	1, 1, 2, 1, 2, 1, 1
	4	1, 1, 2, 2, 3, 3, 4, 3, 3, 2, 2, 1
	5	1, 1, 2, 2, 4, 4, 6, 6, 8, 7, 8, 7, 7, 5, 5, 3, 3, 1, 1
	6	1, 1, 2, 2, 4, 5, 7, 8, 11, 12, 15, 15, 17, 17, 18, 17, 17, 15, 14, 11, 10, 8, 6, 4, 3, 2, 1
	2	1, 2
3	3	1, 2, 2, 2, 1, 1
	4	1, 2, 2, 4, 3, 4, 4, 3, 2, 2
	5	1, 2, 2, 4, 5, 6, 7, 9, 8, 9, 8, 6, 5, 5, 2, 1, 1
	6	1, 2, 2, 4, 5, 8, 9, 12, 14, 17, 18, 19, 19, 20, 18, 17, 14, 13, 10, 8, 5, 4, 2, 2

Table 5: The coefficient lists of  $P_n^{3,b}$ .