

# Four-body Semileptonic Decays $B \rightarrow D^* P \ell^+ \nu_\ell$ with the SU(3) Flavor Symmetry

Meng-Yuan Wan, Yuan-Guo Xu<sup>†</sup>, Qi-Lin Jia, Yue-Xin Liu, Yi-Jie Zhang

College of Physics and Communication Electronics, JiangXi Normal University, NanChang, JiangXi 330022, China

<sup>†</sup>Corresponding author. E-mail addresses: yuanguoxu@jxnu.edu.cn

We present a complete study of the  $B \rightarrow D^* P \ell^+ \nu_\ell$  ( $\ell = e, \mu, \tau$ ) decays with the non-resonant, the charmed axial vector resonant and the charmed tensor resonant contributions by using the SU(3) flavor symmetry. Relevant amplitude relations between different decay modes are obtained by the SU(3) flavor symmetry. We then predict non-measured branching ratios of the  $B \rightarrow D^* P \ell^+ \nu_\ell$  decays with the non-resonant and the charmed resonant contributions by using present experimental data of the  $B \rightarrow D^* P \ell'^+ \nu_{\ell'}$  ( $\ell' = e, \mu$ ) decays within  $2\sigma$  errors. We have found that  $B^{0,+} \rightarrow D^* \eta \ell^+ \nu_\ell$ ,  $B^{0,+} \rightarrow D^* \eta' \ell^+ \nu_\ell$ ,  $B_s^0 \rightarrow D_s^* \eta \ell^+ \nu_\ell$ ,  $B_s^0 \rightarrow D_s^* \eta' \ell^+ \nu_\ell$  and  $B^{0,+} \rightarrow D_s^* K \ell^+ \nu_\ell$  decays only receive non-resonant contributions. Decays  $B_s^0 \rightarrow D_s^{*-} \pi^0 \ell^+ \nu_\ell$  only receive the  $D_{s1}'$  resonant contributions. Other decays receive all three kinds of contributions, and three kinds of contributions are important in most of decays.

## I. Introduction

Decays  $B \rightarrow D^* \pi \ell'^+ \nu_{\ell'}$ ,  $B^+ \rightarrow D_s^{*-} K^+ \ell'^+ \nu_{\ell'}$  and their resonant decays  $B \rightarrow D_1^{(*)}(\rightarrow D^* \pi) \ell'^+ \nu_{\ell'}$ ,  $B \rightarrow D_2^{(*)}(\rightarrow D^* \pi) \ell'^+ \nu_{\ell'}$  have been measured by BABAR, Belle and CLEO, for examples, Refs. [1–3]. Present measurements of the  $B \rightarrow D^* P \ell'^+ \nu_{\ell'}$  decays give us an opportunity to additionally test theoretical approaches and to predict many non-measured  $B \rightarrow D^* P \ell^+ \nu_\ell$  decays. And the predictions of the four-body semileptonic decays  $B \rightarrow D^* P \ell^+ \nu_\ell$  can be tested by LHC and Belle-II in the near future.

For the semileptonic decays, since leptons do not participate in the strong interaction, the weak and strong dynamics can be separated, the theoretical description of the semileptonic decays are relatively simple. Moreover, the experimental backgrounds of the semileptonic decays are cleaner than ones of the hadronic decays. The hadronic transition form factors, which are crucial for testing the theoretical calculations of the involved strong interaction, contain all the strong dynamics in the initial and final hadrons. The analytic structure of the  $B \rightarrow D^* P$  form factors is more complicated than for the  $B \rightarrow D^*$  or  $B \rightarrow P$  form factors, as similarly analysed in Ref. [4], and the resonance states also involve in the  $B \rightarrow D^* P \ell^+ \nu_\ell$  decays. There are several methods to calculate the form factors, such as the chiral perturbation theory [5], the unitarized chiral perturbation theory [6, 7], the light-cone sum rules [8–10], and the QCD factorization [11]. Nevertheless, due to our poor understanding of hadronic interactions, the evaluations of the form factors are difficult and often plugged with large uncertainties.

In the absence of reliable calculations, the symmetry analysis can provide very useful information about the decays. SU(3) flavor symmetry, which is independent of the detailed dynamics, provides us an opportunity to relate different

decay modes, nevertheless, it cannot determine the sizes of the amplitudes or the form factors by itself. However, if experimental data are enough, one may use the data to extract the form factors. Even though it is only an approximate symmetry due to up, down, and strange quarks having different masses, it still gives some valuable information about the decays. SU(3) flavor symmetry has been widely used to study hadron decays, including  $b$ -hadron decays [12–25],  $c$ -hadron decays [24–39], and light hadron decays [24, 40–45].

SU(3) flavor breaking effects play a key role in the precise theoretical predictions of the observables and a precise test of the unitarity of the CKM matrix. SU(3) flavor breaking effects due to the mass difference of  $u, d, s$  quarks have also been studied, for instance, in Refs. [46–58]. The SU(3) flavor breaking effects have been found to be small, for examples, in light hyperon semileptonic decays [53, 54] and  $B \rightarrow D\ell\nu_\ell$  decays [46]. Nevertheless, they might be given larger effects in some other processes, for examples,  $D^0 \rightarrow \pi^0 e^- \nu_e$  decay [47] and  $B \rightarrow D^* \ell \nu_\ell$  decays [46]. So the SU(3) flavor breaking effects should be considered for precise theoretical predictions. Just, if considering the SU(3) flavor breaking effects, extra non-perturbative parameters appear, and one needs more experimental data to determine relevant non-perturbative parameters. At present, there are not many relevant experimental data of the  $B \rightarrow D^* P \ell^+ \nu_\ell$  decays. As more and more accurate data are collected in the future, one may study the SU(3) flavor breaking effects in the  $B \rightarrow D_1/D'_1/D_2^* \ell^+ \nu_\ell$  and  $B \rightarrow D^* P \ell^+ \nu_\ell$  decays. So only the SU(3) flavor symmetry contributions will be analyzed in this work.

Some four-body semileptonic decays  $B/D \rightarrow P_1 P_2 \ell^+ \nu_\ell$  and  $B \rightarrow D^{(*)} P \ell^+ \nu_\ell$  have been studied, for instance, in Refs. [4, 10, 11, 59–70]. In this work, we will study the  $B \rightarrow D^* P \ell^+ \nu_\ell$  decays with the non-resonant, the charmed axial vector resonant and the charmed tensor resonant contributions by the SU(3) flavor symmetry. Firstly, we will construct the hadronic amplitude or the form factor relations between different decay modes and use the available data to extract them. Then, we will predict the not-yet-measured modes for further tests in experiments and analyze the contributions with the non-resonance.

This paper is organized as follows. In Sec. II, we calculate the nonresonant contributions of the  $B \rightarrow D^* P \ell \nu_\ell$  decays. In Sec. III, we present the charmed axial vector resonant and the charmed tensor resonant contributions of  $B \rightarrow D^* P \ell^+ \nu_\ell$  decays. Finally, we summarize our work in Sec. IV.

## II. Non-resonant $B \rightarrow D^* P \ell \nu_\ell$ decays

### A. Decay amplitudes

The four-body semileptonic decays  $B \rightarrow D^* P \ell^+ \nu_\ell$  are generated by  $\bar{b} \rightarrow \bar{c} \ell^+ \nu_\ell$  transition, and the effective Hamiltonian is

$$\mathcal{H}_{eff}(\bar{b} \rightarrow \bar{c} \ell^+ \nu_\ell) = \frac{G_F}{\sqrt{2}} V_{cb} \bar{b} \gamma^\mu (1 - \gamma_5) \bar{c} \bar{\nu}_\ell \gamma_\mu (1 - \gamma_5) \ell, \quad (1)$$

where  $G_F$  is the Fermi constant,  $V_{cb}$  is the CKM matrix element. Decay amplitudes of the  $B \rightarrow D^* P \ell^+ \nu_\ell$  decays can be written as

$$\begin{aligned} \mathcal{A}(B \rightarrow D^* P \ell^+ \nu_\ell) &= \langle D^*(k_1) P(k_2) \ell^+(q_1) \nu_\ell(q_2) | \mathcal{H}_{eff}(\bar{b} \rightarrow \bar{c} \ell^+ \nu_\ell) | B(p_B) \rangle \\ &= \frac{G_F}{\sqrt{2}} V_{cb} L_\mu H^\mu, \end{aligned} \quad (2)$$

where  $L_\mu = \bar{\nu}_\ell \gamma_\mu (1 - \gamma_5) \ell$  is the leptonic charged current, and  $H^\mu = \langle D^*(k_1) P(k_2) | \bar{c} \gamma^\mu (1 - \gamma_5) \bar{b} | B(p_B) \rangle$  is the hadronic matrix element.

Usually, the hadronic matrix element  $H^\mu$  can be obtained in terms of the form factors of the  $B \rightarrow D^* P$ . Similar to  $B \rightarrow PP$  form factors given in Ref. [11], the  $B \rightarrow D^* P$  form factors are defined by

$$\langle D^*(k_1) P(k_2) | \bar{c} \gamma^\mu \bar{b} | B(p_B) \rangle = i F_\perp \frac{1}{\sqrt{k^2}} q_\perp^\mu, \quad (3)$$

$$-\langle D^*(k_1) P(k_2) | \bar{c} \gamma^\mu \bar{b} | B(p_B) \rangle = F_t \frac{q^\mu}{\sqrt{q^2}} + F_0 \frac{2\sqrt{q^2}}{\sqrt{\lambda}} k_0^\mu + F_\parallel \frac{1}{\sqrt{k^2}} \bar{k}_\parallel^\mu, \quad (4)$$

where  $k \equiv k_1 + k_2$ ,  $q \equiv q_1 + q_2$ ,  $\lambda = \lambda(m_{D^*}^2, q^2, k^2)$  with  $\lambda(a, b, c) = a^2 + b^2 + c^2 - 2ab - 2bc - 2ac$ , and  $F_0, F_t, F_\perp, F_\parallel$  are the form factors of  $B \rightarrow D^* P$ . In addition,  $q_\perp^\mu$ ,  $k_0^\mu$  and  $\bar{k}_\parallel^\mu$  are defined in Ref. [11].

The differential branching ratios of the nonresonant  $B \rightarrow D^* P \ell^+ \nu_\ell$  decays can be written as

$$\frac{d\mathcal{B}(B \rightarrow D^* P \ell^+ \nu_\ell)_N}{dq^2 dk^2} = \frac{1}{2} \tau_B |\mathcal{N}|^2 \beta_\ell (3 - \beta_\ell) |H_N|^2, \quad (5)$$

with

$$\begin{aligned} |\mathcal{N}|^2 &= G_F^2 |V_{cb}|^2 \frac{\beta_\ell q^2 \sqrt{\lambda}}{3 \times 2^{10} \pi^5 m_B^3}, \quad \text{with} \quad \beta_\ell = 1 - \frac{m_\ell^2}{q^2}. \\ |H_N|^2 &= |F_0|^2 + \frac{2}{3} (|F_\parallel|^2 + |F_\perp|^2) + \frac{3m_\ell^2}{q^2(3 - \beta_\ell)} |F_t|^2, \end{aligned} \quad (6)$$

where  $\tau_M(m_M)$  is lifetime(mass) of  $M$  particle. The ranges of integration are given by  $(m_{D^*} + m_P)^2 \leq k^2 \leq (m_B - m_\ell)^2$  and  $m_\ell^2 \leq q^2 \leq (m_B - \sqrt{k^2})^2$ .

The calculations of the  $F_0, F_t, F_\perp, F_\parallel$  form factors are difficult. If we ignore  $|F_t|^2$  term since it is proportional to  $m_\ell^2$  and it is small when  $\ell = e, \mu$ ,  $|H_N|^2$  is only include the hadronic part. Noted that although  $|F_t|^2$  term might be large when  $\ell = \tau$ , it is difficult to estimate its contribution in this work, so we still ignore it. Then  $|H_N|^2$ , which only includes hadronic part, can be related by the SU(3) flavor symmetry/breaking of  $u, d, s$  quarks.

## B. Hadronic amplitudes based on the SU(3) flavor analysis

Since the SU(3) flavor analysis is based on the SU(3) flavor group, we will give relevant meson multiplets first. Bottom pseudoscalar triplets  $B_i$ , charm vector triplets  $D_i^*$ , and light pseudoscalar octets and singlets  $P_j^i$  under the SU(3) flavor symmetry of  $u, d, s$  quarks are

$$B_i = (B^+(\bar{b}u), B^0(\bar{b}d), B^0(\bar{b}s)), \quad (7)$$

$$D_i^* = (\bar{D}^{*0}(\bar{c}u), D^{*-}(\bar{c}d), D^{*-}(\bar{c}s)), \quad (8)$$

$$P_j^i = \begin{pmatrix} \frac{\pi^0}{\sqrt{2}} + \frac{\eta_8}{\sqrt{6}} + \frac{\eta_1}{\sqrt{3}} & \pi^+ & K^+ \\ \pi^- & -\frac{\pi^0}{\sqrt{2}} + \frac{\eta_8}{\sqrt{6}} + \frac{\eta_1}{\sqrt{3}} & K^0 \\ K^- & \bar{K}^0 & -\frac{2\eta_8}{\sqrt{6}} + \frac{\eta_1}{\sqrt{3}} \end{pmatrix}, \quad (9)$$

where  $i, j = 1, 2, 3$  for  $u, d, s$  quarks, the  $\eta$  and  $\eta'$  are mixtures of  $\eta_1 = \frac{u\bar{u} + d\bar{d} + s\bar{s}}{\sqrt{3}}$  and  $\eta_8 = \frac{u\bar{u} + d\bar{d} - 2s\bar{s}}{\sqrt{6}}$  with the mixing angle  $\theta_P$

$$\begin{pmatrix} \eta \\ \eta' \end{pmatrix} = \begin{pmatrix} \cos\theta_P & -\sin\theta_P \\ \sin\theta_P & \cos\theta_P \end{pmatrix} \begin{pmatrix} \eta_8 \\ \eta_1 \end{pmatrix}. \quad (10)$$




FIG. 1: Diagrams of the non-resonant  $B \rightarrow D^* P \ell^+ \nu_\ell$  decays.

And  $\theta_P = [-20^\circ, -10^\circ]$  from the Particle Data Group (PDG) [1] will be used in our numerical analysis.

Charm axial vector  $A$  and tensor  $T$  contribute to these processes as resonances, which will be discussed next section. There are two types of P-wave charm axial-vector mesons with different quantum numbers  $J^{PC} = 1^{++}$  and  $J^{PC} = 1^{+-}$ . The charm axial vector triplets are

$$D'_{1i} = (\bar{D}'_1(2430)^0, D'_1(2430)^-, D'_{s1}(2460)^-) \quad \text{for } J^{PC} = 1^{++}, \quad (11)$$

$$D_{1i} = (\bar{D}_1(2420)^0, D_1(2420)^-, D_{s1}(2536)^-) \quad \text{for } J^{PC} = 1^{+-}. \quad (12)$$

The charm tensor triplet is

$$D_{2i} = (\bar{D}_2^*(2460)^0, D_2^*(2460)^-, D_{s2}^*(2573)^-). \quad (13)$$

Feynman diagrams for the non-resonant  $B \rightarrow D^* P \ell^+ \nu_\ell$  decays are displayed in Fig. 1. The leptonic charged current is invariant under the SU(3) flavor symmetry, and the hadronic matrix element can be parameterized by the SU(3) flavor symmetry.

The decay amplitudes of the non-resonant  $B \rightarrow D^* P \ell^+ \nu_\ell$  decays in Eq. (2) can be transformed as

$$\mathcal{A}(B \rightarrow D^* P \ell^+ \nu_\ell)_N = \frac{G_F}{\sqrt{2}} V_{cb}^* H(B \rightarrow D^* P)_N \bar{\nu}_\ell \gamma_\mu (1 - \gamma_5) \ell, \quad (14)$$

$$H(B \rightarrow D^* P)_N = a_1 B^i P_i^j D^{*j} + a_2 B^i D^{*i} P_k^k, \quad (15)$$

where  $a_{1,2}$  are the nonperturbative coefficients under the SU(3) flavor symmetry.

The hadronic amplitudes of the non-resonant  $B \rightarrow D^* P \ell^+ \nu_\ell$  decays are given in Tab. I, in which we can see the amplitude relations of them. As given in Fig. 1 (b), the  $a_2$  term in Eq. (15) is suppressed by the Okubo-Zweig-Iizuka (OZI) rule [71–73], and it only appears in the decays with  $\eta, \eta'$  final states. If ignoring the OZI suppressed  $a_2$  term, all hadronic amplitudes can be related by only one nonperturbative coefficient  $a_1$ .

### C. Numerical results for the non-resonant $B \rightarrow D^* P \ell^+ \nu_\ell$ decays

The theoretical input parameters and the experimental data within the  $2\sigma$  errors from PDG [1] will be used in our numerical analysis. The four-body semileptonic  $B^+ \rightarrow D^{*-} \pi^+ \ell^+ \nu_\ell$  and  $B^0 \rightarrow D^{*0} \pi^+ \ell^+ \nu_\ell$  decays have been well

TABLE I: The hadronic amplitudes  $H(B \rightarrow D^* P)_N$  for  $B \rightarrow D^* P \ell^+ \nu_\ell$  decays based on the SU(3) flavor symmetry.

| Decay modes                                          | Hadronic amplitudes                                                              | Decay modes                                          | Hadronic amplitudes                                                                |
|------------------------------------------------------|----------------------------------------------------------------------------------|------------------------------------------------------|------------------------------------------------------------------------------------|
| $B^+ \rightarrow \bar{D}^{*0} \pi^0 \ell^+ \nu_\ell$ | $\frac{a_1}{\sqrt{2}}$                                                           | $B^0 \rightarrow \bar{D}^{*0} \pi^- \ell^+ \nu_\ell$ | $a_1$                                                                              |
| $B^+ \rightarrow \bar{D}^{*0} \eta \ell^+ \nu_\ell$  | $\frac{a_1}{\sqrt{6}} \cos \theta_P - \frac{a_1 + 3a_2}{\sqrt{3}} \sin \theta_P$ | $B^0 \rightarrow D^{*-} \pi^0 \ell^+ \nu_\ell$       | $-\frac{a_1}{\sqrt{2}}$                                                            |
| $B^+ \rightarrow \bar{D}^{*0} \eta' \ell^+ \nu_\ell$ | $\frac{a_1}{\sqrt{6}} \sin \theta_P + \frac{a_1 + 3a_2}{\sqrt{3}} \cos \theta_P$ | $B^0 \rightarrow D^{*-} \eta \ell^+ \nu_\ell$        | $\frac{a_1}{\sqrt{6}} \cos \theta_P - \frac{a_1 + 3a_2}{\sqrt{3}} \sin \theta_P$   |
| $B^+ \rightarrow D^{*-} \pi^+ \ell^+ \nu_\ell$       | $a_1$                                                                            | $B^0 \rightarrow D^{*-} \eta' \ell^+ \nu_\ell$       | $\frac{a_1}{\sqrt{6}} \sin \theta_P + \frac{a_1 + 3a_2}{\sqrt{3}} \cos \theta_P$   |
| $B^+ \rightarrow D_s^{*-} K^+ \ell^+ \nu_\ell$       | $a_1$                                                                            | $B^0 \rightarrow D_s^{*-} K^0 \ell^+ \nu_\ell$       | $a_1$                                                                              |
| $B_s^0 \rightarrow \bar{D}^{*0} K^- \ell^+ \nu_\ell$ | $a_1$                                                                            | $B_s^0 \rightarrow D_s^{*-} \eta \ell^+ \nu_\ell$    | $-\frac{2a_1}{\sqrt{6}} \cos \theta_P - \frac{a_1 + 3a_2}{\sqrt{3}} \sin \theta_P$ |
| $B_s^0 \rightarrow D^{*-} \bar{K}^0 \ell^+ \nu_\ell$ | $a_1$                                                                            | $B_s^0 \rightarrow D_s^{*-} \eta' \ell^+ \nu_\ell$   | $-\frac{2a_1}{\sqrt{6}} \sin \theta_P + \frac{a_1 + 3a_2}{\sqrt{3}} \cos \theta_P$ |

measured [1], and they with the  $2\sigma$  errors are

$$\mathcal{B}(B^+ \rightarrow D^{*-} \pi^+ \ell'^+ \nu_{\ell'})_{all} = (6.0 \pm 0.8) \times 10^{-3}, \quad (16)$$

$$\mathcal{B}(B^+ \rightarrow D^{*-} \pi^+ \ell'^+ \nu_{\ell'})_{\bar{D}_1^0} = (3.03 \pm 0.40) \times 10^{-3}, \quad (17)$$

$$\mathcal{B}(B^+ \rightarrow D^{*-} \pi^+ \ell'^+ \nu_{\ell'})_{\bar{D}_1'^0} = (2.7 \pm 1.2) \times 10^{-3}, \quad (18)$$

$$\mathcal{B}(B^+ \rightarrow D^{*-} \pi^+ \ell'^+ \nu_{\ell'})_{\bar{D}_2^{*0}} = (1.01 \pm 0.48) \times 10^{-3}, \quad (19)$$

$$\mathcal{B}(B^0 \rightarrow D^{*0} \pi^- \ell'^+ \nu_{\ell'})_{all} = (5.8 \pm 1.6) \times 10^{-3}, \quad (20)$$

$$\mathcal{B}(B^0 \rightarrow D^{*0} \pi^- \ell'^+ \nu_{\ell'})_{D_1^-} = (2.80 \pm 0.56) \times 10^{-3}, \quad (21)$$

$$\mathcal{B}(B^0 \rightarrow D^{*0} \pi^- \ell'^+ \nu_{\ell'})_{D_1'^-} = (3.1 \pm 1.8) \times 10^{-3}, \quad (22)$$

$$\mathcal{B}(B^0 \rightarrow D^{*0} \pi^- \ell'^+ \nu_{\ell'})_{D_2^{*-}} = (0.68 \pm 0.24) \times 10^{-3}, \quad (23)$$

where  $\mathcal{B}_{all,R}$  denotes the total and  $R$  resonant branching ratios. From Eqs. (16-23), we obtain the upper limits of non-resonant branching ratios within the  $2\sigma$  errors

$$\mathcal{B}(B^+ \rightarrow D^{*-} \pi^+ \ell'^+ \nu_{\ell'})_N \leq 2.14 \times 10^{-3}, \quad (24)$$

$$\mathcal{B}(B^0 \rightarrow \bar{D}^{*0} \pi^- \ell'^+ \nu_{\ell'})_N \leq 3.42 \times 10^{-3}. \quad (25)$$

In addition,  $B^+ \rightarrow D_s^{*-} K^+ \ell'^+ \nu_{\ell'}$  has also been measured [1]

$$\mathcal{B}(B^+ \rightarrow D_s^{*-} K^+ \ell'^+ \nu_{\ell'})_{all} = (2.9 \pm 3.8) \times 10^{-4}. \quad (26)$$

Since there are no phase spaces for  $\bar{D}_1(2420)^0/\bar{D}_1'(2430)^0/\bar{D}_2(2460)^0 \rightarrow D_s^{*-} K^+$  decays,  $\mathcal{B}(B^+ \rightarrow D_s^{*-} K^+ \ell'^+ \nu_{\ell'})_{all}$  will be considered as only non-resonant branching ratios  $\mathcal{B}(B^+ \rightarrow D_s^{*-} K^+ \ell'^+ \nu_{\ell'})_N$ . The error of  $\mathcal{B}(B^+ \rightarrow D_s^{*-} K^+ \ell'^+ \nu_{\ell'})_{all}$  is very large, all range, that is less than  $6.7 \times 10^{-4}$ , are allowed within  $2\sigma$  error bars.  $\mathcal{B}(B^+ \rightarrow D_s^{*-} K^+ \ell'^+ \nu_{\ell'})_N$  as well as the upper limits of  $\mathcal{B}(B^+ \rightarrow D^{*-} \pi^+ \ell'^+ \nu_{\ell'})_N$  and  $\mathcal{B}(B^0 \rightarrow \bar{D}^{*0} \pi^- \ell'^+ \nu_{\ell'})_N$  in Eqs. (24-26) will be used to constrain the nonperturbative coefficient  $a_1$ . Present non-resonant data only can give the upper limit  $|a_1| \leq 20.70$ . And then one can obtain upper limit predictions of other non-resonant branching ratios, which have not been measured or well measured.

Our upper limit predictions of the non-resonant branching ratios are listed in the second column of Tab. II. For convenient comparison, the available experimental values are also given in Tab. II. And we find that only the

TABLE II: Branching ratios for  $B \rightarrow D^* P \ell^+ \nu_\ell$  decays due to  $\bar{b} \rightarrow \bar{c} \ell^+ \nu_\ell$  within  $2\sigma$  errors. The unit is  $10^{-4}$  for all branching ratios.  $\mathcal{B}_{[R]}$  denotes the R resonant branching ratios. <sup>e</sup>denotes experimental data within  $2\sigma$  errors.

|                                                                         | $\mathcal{B}_N$                    | $\mathcal{B}_{D_1^{(\prime)}}$                                                                                                          | $\mathcal{B}_{D_2^*}$                                         | $\mathcal{B}_{all}$                     |
|-------------------------------------------------------------------------|------------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------|---------------------------------------------------------------|-----------------------------------------|
| $\mathcal{B}(B_u^+ \rightarrow D^{*-} \pi^+ \ell'^+ \nu_{\ell'})$       | $\leq 14.17$                       | $34.38 \pm 4.62_{[D_1^{\prime 0}]}$<br>$30.30 \pm 4.00_{[D_1^0]}$<br>$27.00 \pm 12.00_{[D_1^{\prime 0}]}$<br>$30.30 \pm 4.00_{[D_1^0]}$ | $8.17 \pm 1.52_{[D_2^{*0}]}$<br>$10.10 \pm 4.80_{[D_2^{*0}]}$ | $65.54 \pm 2.46$<br>$60.00 \pm 8.00^e$  |
| $\mathcal{B}(B_u^+ \rightarrow D_s^{*-} K^+ \ell'^+ \nu_{\ell'})$       | $\leq 6.70$<br>$(2.90 \pm 3.80)^e$ | ...                                                                                                                                     | ...                                                           | ...                                     |
| $\mathcal{B}(B_u^+ \rightarrow \bar{D}^{*0} \pi^0 \ell'^+ \nu_{\ell'})$ | $\leq 7.17$                        | $17.44 \pm 2.34_{[D_1^{\prime 0}]}$<br>$15.35 \pm 2.03_{[D_1^0]}$                                                                       | $4.31 \pm 0.80_{[D_2^{*0}]}$                                  | $40.59 \pm 8.18$                        |
| $\mathcal{B}(B_u^+ \rightarrow \bar{D}^{*0} \eta \ell'^+ \nu_{\ell'})$  | $\leq 2.47$                        | ...                                                                                                                                     | ...                                                           | ...                                     |
| $\mathcal{B}(B_u^+ \rightarrow \bar{D}^{*0} \eta' \ell'^+ \nu_{\ell'})$ | $\leq 0.80$                        | ...                                                                                                                                     | ...                                                           | ...                                     |
| $\mathcal{B}(B_d^0 \rightarrow \bar{D}^{*0} \pi^- \ell'^+ \nu_{\ell'})$ | $\leq 13.32$<br>$\leq 34.20^e$     | $32.37 \pm 4.33_{[D_1^{\prime -}]}$<br>$28.40 \pm 3.97_{[D_1^-]}$<br>$31.00 \pm 18.00_{[D_1^{\prime -}]}$<br>$28.00 \pm 5.60_{[D_1^-]}$ | $7.82 \pm 1.38_{[D_2^{*-}]}$<br>$6.80 \pm 2.40_{[D_2^{*-}]}$  | $67.07 \pm 6.93$<br>$58.00 \pm 16.00^e$ |
| $\mathcal{B}(B_d^0 \rightarrow D_s^{*-} K^0 \ell'^+ \nu_{\ell'})$       | $\leq 6.21$                        | ...                                                                                                                                     | ...                                                           | ...                                     |
| $\mathcal{B}(B_d^0 \rightarrow D^{*-} \pi^0 \ell'^+ \nu_{\ell'})$       | $\leq 6.67$                        | $16.11 \pm 2.15_{[D_1^{\prime -}]}$<br>$14.14 \pm 1.98_{[D_1^-]}$                                                                       | $3.83 \pm 0.68_{[D_2^{*-}]}$                                  | $37.17 \pm 7.50$                        |
| $\mathcal{B}(B_d^0 \rightarrow D^{*-} \eta \ell'^+ \nu_{\ell'})$        | $\leq 2.27$                        | ...                                                                                                                                     | ...                                                           | ...                                     |
| $\mathcal{B}(B_d^0 \rightarrow D^{*-} \eta' \ell'^+ \nu_{\ell'})$       | $\leq 0.74$                        | ...                                                                                                                                     | ...                                                           | ...                                     |
| $\mathcal{B}(B_s^0 \rightarrow \bar{D}^{*0} K^- \ell'^+ \nu_{\ell'})$   | $\leq 8.82$                        | $35.60 \pm 15.40_{[D_{s1}^-]}$                                                                                                          | $1.71 \pm 0.51_{[D_{s2}^{*-}]}$                               | $41.70 \pm 20.03$                       |
| $\mathcal{B}(B_s^0 \rightarrow D^{*-} \bar{K}^0 \ell'^+ \nu_{\ell'})$   | $\leq 8.71$                        | $29.67 \pm 13.61_{[D_{s1}^-]}$                                                                                                          | $1.30 \pm 0.40_{[D_{s2}^{*-}]}$                               | $35.40 \pm 17.90$                       |
| $\mathcal{B}(B_s^0 \rightarrow D_s^{*-} \pi^0 \ell'^+ \nu_{\ell'})$     | ...                                | $25.45 \pm 14.11_{[D_{s1}^{\prime -}]}$                                                                                                 | ...                                                           | ...                                     |
| $\mathcal{B}(B_s^0 \rightarrow D_s^{*-} \eta \ell'^+ \nu_{\ell'})$      | $\leq 3.25$                        | ...                                                                                                                                     | ...                                                           | ...                                     |
| $\mathcal{B}(B_s^0 \rightarrow D_s^{*-} \eta' \ell'^+ \nu_{\ell'})$     | $\leq 1.94$                        | ...                                                                                                                                     | ...                                                           | ...                                     |
| $\mathcal{B}(B_u^+ \rightarrow D^{*-} \pi^+ \tau^+ \nu_\tau)$           | $\leq 1.68$                        | $5.85 \pm 0.89_{[D_1^{\prime 0}]}$<br>$5.12 \pm 0.76_{[D_1^0]}$                                                                         | $5.35 \pm 1.53_{[D_2^{*0}]}$                                  | $17.07 \pm 3.64$                        |
| $\mathcal{B}(B_u^+ \rightarrow D_s^{*-} K^+ \tau^+ \nu_\tau)$           | $\leq 0.31$                        | ...                                                                                                                                     | ...                                                           | ...                                     |
| $\mathcal{B}(B_u^+ \rightarrow \bar{D}^{*0} \pi^0 \tau^+ \nu_\tau)$     | $\leq 0.86$                        | $2.97 \pm 0.46_{[D_1^{\prime 0}]}$<br>$2.59 \pm 0.38_{[D_1^0]}$                                                                         | $2.83 \pm 0.81_{[D_2^{*0}]}$                                  | $8.79 \pm 1.90$                         |
| $\mathcal{B}(B_u^+ \rightarrow \bar{D}^{*0} \eta \tau^+ \nu_\tau)$      | $\leq 0.13$                        | ...                                                                                                                                     | ...                                                           | ...                                     |
| $\mathcal{B}(B_u^+ \rightarrow \bar{D}^{*0} \eta' \tau^+ \nu_\tau)$     | $\leq 0.0088$                      | ...                                                                                                                                     | ...                                                           | ...                                     |
| $\mathcal{B}(B_d^0 \rightarrow \bar{D}^{*0} \pi^- \tau^+ \nu_\tau)$     | $\leq 1.58$                        | $5.49 \pm 0.83_{[D_1^{\prime -}]}$<br>$4.79 \pm 0.74_{[D_1^-]}$                                                                         | $5.22 \pm 1.51_{[D_2^{*-}]}$                                  | $16.27 \pm 3.49$                        |
| $\mathcal{B}(B_d^0 \rightarrow D_s^{*-} K^0 \tau^+ \nu_\tau)$           | $\leq 0.29$                        | ...                                                                                                                                     | ...                                                           | ...                                     |
| $\mathcal{B}(B_d^0 \rightarrow D^{*-} \pi^0 \tau^+ \nu_\tau)$           | $\leq 0.79$                        | $2.73 \pm 0.41_{[D_1^{\prime -}]}$<br>$2.38 \pm 0.37_{[D_1^-]}$                                                                         | $2.56 \pm 0.74_{[D_2^{*-}]}$                                  | $8.10 \pm 1.81$                         |
| $\mathcal{B}(B_d^0 \rightarrow D^{*-} \eta \tau^+ \nu_\tau)$            | $\leq 0.12$                        | ...                                                                                                                                     | ...                                                           | ...                                     |
| $\mathcal{B}(B_d^0 \rightarrow D^{*-} \eta' \tau^+ \nu_\tau)$           | $\leq 0.0080$                      | ...                                                                                                                                     | ...                                                           | ...                                     |
| $\mathcal{B}(B_s^0 \rightarrow \bar{D}^{*0} K^- \tau^+ \nu_\tau)$       | $\leq 0.66$                        | $6.10 \pm 2.64_{[D_{s1}^-]}$                                                                                                            | $1.06 \pm 0.37_{[D_{s2}^{*-}]}$                               | $7.44 \pm 3.26$                         |
| $\mathcal{B}(B_s^0 \rightarrow D^{*-} \bar{K}^0 \tau^+ \nu_\tau)$       | $\leq 0.64$                        | $5.08 \pm 2.33_{[D_{s1}^-]}$                                                                                                            | $0.80 \pm 0.28_{[D_{s2}^{*-}]}$                               | $6.18 \pm 2.86$                         |
| $\mathcal{B}(B_s^0 \rightarrow D_s^{*-} \pi^0 \tau^+ \nu_\tau)$         | ...                                | $4.55 \pm 2.52_{[D_{s1}^{\prime -}]}$                                                                                                   | ...                                                           | ...                                     |
| $\mathcal{B}(B_s^0 \rightarrow D_s^{*-} \eta \tau^+ \nu_\tau)$          | $\leq 0.17$                        | ...                                                                                                                                     | ...                                                           | ...                                     |
| $\mathcal{B}(B_s^0 \rightarrow D_s^{*-} \eta' \tau^+ \nu_\tau)$         | $\leq 0.019$                       | ...                                                                                                                                     | ...                                                           | ...                                     |

upper limit of  $\mathcal{B}(B^+ \rightarrow D_s^{*-} K^+ \ell'^+ \nu_{\ell'})_N$  gives the final effective bound. The upper limit predictions of  $\mathcal{B}(B_u^+ \rightarrow D^{*-} \pi^+ \ell'^+ \nu_{\ell'})_N$  and  $\mathcal{B}(B_d^0 \rightarrow \bar{D}^{*0} \pi^- \ell'^+ \nu_{\ell'})_N$  are obviously smaller than their experimental upper limits. Decays  $B_{(s)} \rightarrow D_{(s)}^* \eta \ell^+ \nu_\ell$  and  $B_{(s)} \rightarrow D_{(s)}^* \eta' \ell^+ \nu_\ell$  are suppressed by the narrow phase spaces and the mixing angle  $\theta_P$ , and their upper limits of the branching ratios, especially of  $B_{(s)} \rightarrow D^* \eta' \tau^+ \nu_\tau$ , are obviously smaller than other predictions.

### III. Decays $B \rightarrow D^* P \ell^+ \nu_\ell$ with the $D^{**}$ resonant states

Except for the non-resonant  $B \rightarrow D^* P \ell^+ \nu_\ell$  decays, the  $B \rightarrow D^* P \ell^+ \nu_\ell$  decay with the  $D^{**}$  resonant state is also studied, where  $D^{**}$  is an lightest excited charmed meson. There are the four lightest excited charmed meson states,  $D^{**} = (D_0, D'_1, D_1, D_2^*)$  or  $(D_{s0}, D'_{s1}, D_{s1}, D_{s2}^*)$ . The spin-0 states  $D_0$  and  $D_{s0}$  can only decay to  $DP$  [2], so we will study  $D'_1, D_1$  and  $D_2^*$  decay to  $D^* P$  in this work.

In the case of the decay widths of the resonances are very narrow, the resonant decay branching ratios respect a simple factorization relation

$$\mathcal{B}(B \rightarrow D^{**} \ell^+ \nu_\ell, D^{**} \rightarrow D^* P) = \mathcal{B}(B \rightarrow D^{**} \ell^+ \nu_\ell) \times \mathcal{B}(D^{**} \rightarrow D^* P), \quad (27)$$

and this result is also a good approximation for wider resonances. Above Eq. (27) will be used in our analysis for resonant  $B \rightarrow D^{**}(\rightarrow D^* P) \ell^+ \nu_\ell$  decays. Relevant  $\mathcal{B}(B \rightarrow D^{**} \ell^+ \nu_\ell)$  and  $\mathcal{B}(D^{**} \rightarrow D^* P)$  also can be obtained by the SU(3) flavor symmetry.

#### A. Resonant $B \rightarrow D^* P \ell^+ \nu_\ell$ decays with the charmed axial vectors

##### 1. $B \rightarrow D_1/D'_1 \ell^+ \nu_\ell$ decays

Similar to  $B \rightarrow V \ell^+ \nu_\ell$  decays, the decay amplitudes of the  $B \rightarrow A \ell^+ \nu_\ell$  decays with  $A = D_1/D'_1$  can be written as

$$\mathcal{M}(B \rightarrow A \ell^+ \nu_\ell) = \frac{G_F}{\sqrt{2}} V_{cb} \sum_{mm'} g_{mm'} L_m^{\lambda_\ell \lambda_\nu} H_{m'}^{\lambda_A}, \quad (28)$$

with

$$L_m^{\lambda_\ell \lambda_\nu} = \epsilon_\alpha(m) \bar{\nu}_\ell \gamma^\alpha (1 - \gamma_5) \ell, \quad (29)$$

$$H_{m'}^{\lambda_A} = \epsilon_\beta^*(m') \langle A(p, \varepsilon^*) | \bar{c} \gamma^\beta (1 - \gamma_5) b | B(p_B) \rangle, \quad (30)$$

where the particle helicities  $\lambda_A = 0, \pm 1$ ,  $\lambda_\ell = \pm \frac{1}{2}$  and  $\lambda_\nu = +\frac{1}{2}$ , as well as  $\epsilon_\mu(m)$  is the polarization vectors of the virtual  $W$  with  $m = 0, t, \pm 1$ , and  $\varepsilon^*$  is the polarization vectors of  $A$  meson. Hadronic matrix element can be parameterized by the  $B \rightarrow A$  form factors [74, 75]

$$\begin{aligned} \langle A(p, \varepsilon^*) | \bar{c} \gamma_\mu (1 - \gamma_5) b | B(p_B) \rangle &= \frac{2iA(q'^2)}{m_B - m_A} \epsilon_{\mu\nu\alpha\beta} \varepsilon^{*\nu} p_B^\alpha p^\beta \\ &\quad - \left[ \varepsilon_\mu^*(m_B - m_A) V_1^A(q'^2) - (p_B + p)_\mu (\varepsilon^* \cdot p_B) \frac{V_2^A(q'^2)}{m_B - m_A} \right] \\ &\quad + q'_\mu (\varepsilon^* \cdot p_B) \frac{2m_A}{q'^2} [V_3^A(q'^2) - V_0^A(q'^2)], \end{aligned} \quad (31)$$

with  $q' = p_B - p$ . Then the hadronic helicity amplitudes can be written as [76–79]

$$H_{\pm}^A = (m_B - m_A)V_1^A(q'^2) \mp \frac{2m_B|\vec{p}_c|}{(m_B - m_A)}A(q'^2), \quad (32)$$

$$H_0^A = \frac{1}{2m_A\sqrt{q'^2}} \left[ (m_B^2 - m_A^2 - q'^2)(m_B - m_A)V_1^A(q'^2) - \frac{4m_B^2|\vec{p}_c|^2}{m_B - m_A}V_2^A(q'^2) \right], \quad (33)$$

$$H_t^A = \frac{2m_B|\vec{p}_c|}{\sqrt{q'^2}}V_0^A(q'^2), \quad (34)$$

where  $|\vec{p}_c| \equiv \frac{\sqrt{\lambda}}{2m_B}$  with  $\lambda = m_B^4 + m_A^4 + q'^4 - 2m_B^2m_A^2 - 2q'^2m_B^2 - 2q'^2m_A^2$ .

The differential branching ratios are [80]

$$\frac{d\mathcal{B}(B \rightarrow A\ell^+\nu_\ell)}{dq'^2} = \frac{\tau_B G_F^2 |V_{cb}|^2 \lambda^{1/2} q'^2}{24(2\pi)^3 m_B^3} \left(1 - \frac{m_\ell^2}{q'^2}\right)^2 \left(1 + \frac{m_\ell^2}{2q'^2}\right) \mathcal{H}_{\text{total}}(B \rightarrow A\ell^+\nu_\ell), \quad (35)$$

with

$$\mathcal{H}_{\text{total}}(B \rightarrow A\ell^+\nu_\ell) = \left(|H_+^A|^2 + |H_-^A|^2 + |H_0^A|^2\right) + \frac{3m_\ell^2}{2q'^2} \left(1 + \frac{m_\ell^2}{2q'^2}\right) |H_t^A|^2, \quad (36)$$

where  $m_\ell^2 \leq q'^2 \leq (m_B - m_A)^2$ . Then one can obtain the branching ratios by the Eq. (35) and the form factors. Usually, it is difficult to calculate the form factors, and their results depend on the different approaches.

The branching ratios also can be obtained by using the SU(3) flavor symmetry. In the  $\ell = e, \mu$  cases,  $\frac{3m_\ell^2}{2q'^2}$  is far less than  $1 + \frac{m_\ell^2}{2q'^2}$  in Eq. (36), so the  $|H_t^A|^2$  term in Eq.(36) can ignore safely. In the  $\ell = \tau$  case,  $(\frac{3m_\ell^2}{2q'^2})/(1 + \frac{m_\ell^2}{2q'^2})$  lie in  $[0.48, 1]$ , the  $|H_t^A|^2$  term give non-negligible contribution, but it is difficult to estimate its contribution if we do not depend on any calculation of the form factors. Two cases will be considered in our analysis of the  $B \rightarrow D_1/D_1'\ell^+\nu_\ell$  decays.

$S_1$ : Ignoring  $|H_t^A|^2$  term in the  $\ell = e, \mu, \tau$  decays, then  $\mathcal{H}_{\text{total}}$  only includes the hadronic part that can be related by the SU(3) flavor symmetry as follow.

$$\mathcal{H}_{\text{total}}(B_u^+ \rightarrow \bar{D}_1'^0 \ell^+ \nu_\ell) = \mathcal{H}_{\text{total}}(B_d^0 \rightarrow D_1'^- \ell'^+ \nu_{\ell'}) = \mathcal{H}_{\text{total}}(B_s^0 \rightarrow D_{s1}'^- \ell'^+ \nu_{\ell'}) = |a_0|^2, \quad (37)$$

$$\mathcal{H}_{\text{total}}(B_u^+ \rightarrow \bar{D}_1^0 \ell^+ \nu_\ell) = \mathcal{H}_{\text{total}}(B_d^0 \rightarrow D_1^- \ell^+ \nu_\ell) = \mathcal{H}_{\text{total}}(B_s^0 \rightarrow D_{s1}^- \ell^+ \nu_\ell) = |b_0|^2. \quad (38)$$

where  $a_0$  and  $b_0$  are the nonperturbative coefficients. The actual  $\mathcal{B}(B \rightarrow A\tau^+\nu_\tau)$  might be larger than our later predictions.

$S_2$ :  $\mathcal{H}_{\text{total}}$  are obtained by using the hadronic helicity amplitude expressions in Eqs. (32–34), which are  $q^2$  dependent and can be expressed by the form factors. The form factors of  $B \rightarrow D_1^{1/2}, D_1^{3/2}$  in Ref. [79] are taken (we do not use ones of  $B_s \rightarrow D_{s1}^{1/2}, D_{s1}^{3/2}$ ), we keep  $V_1^A(0)$  as an undetermined constant, other  $F_i(0)$  can be expressed as  $r_F \times V_1^A(0)$ , and  $r_F = \frac{F_i(0)}{V_1^A(0)}$  are taken from Ref. [79]. Since these form factors also preserve the SU(3) flavor symmetry, the same relations in Eqs. (37–38) will be used for  $V_1^A(0)$ .

For the semileptonic  $B \rightarrow D_1^{(\prime)}\ell^+\nu_\ell$  decays, the branching ratios of  $B_u^+ \rightarrow \bar{D}_1'^0 \ell'^+ \nu_{\ell'}$ ,  $B_d^0 \rightarrow D_1'^- \ell'^+ \nu_{\ell'}$ ,  $B_u^+ \rightarrow \bar{D}_1^0 \ell'^+ \nu_{\ell'}$ , and  $B_d^0 \rightarrow D_1^- \ell'^+ \nu_{\ell'}$  decays have been measured by the Belle Collaboration [81], and the experimental data with  $2\sigma$  errors are listed in the second column of Tab. III. Using the experimental data of  $\mathcal{B}(B_u^+ \rightarrow \bar{D}_1'^0 \ell'^+ \nu_{\ell'})$  and  $\mathcal{B}(B_d^0 \rightarrow D_1'^- \ell'^+ \nu_{\ell'})$ , one can constrain the allowed range of  $a_0$  and  $V_1^{D_1}(0)$ , and then one can obtain the predictions of  $\mathcal{B}(B_u^+ \rightarrow \bar{D}_1^0 \ell'^+ \nu_{\ell'})$ ,  $\mathcal{B}(B_d^0 \rightarrow D_1^- \ell'^+ \nu_{\ell'})$  and  $\mathcal{B}(B_s^0 \rightarrow D_{s1}'^- \ell'^+ \nu_{\ell'})$  by the constrained  $a_0$  or  $V_1^{D_1}(0)$ . In a similar way, one

TABLE III: Experimental data and predictions of the branching ratios of the semileptonic  $B \rightarrow D_1^{(\prime)} \ell^+ \nu_\ell$  decays with the  $2\sigma$  errors by the SU(3) flavor symmetry (in units of  $10^{-3}$ ).

|                                                                       | Data [81]     | Predictions in $S_1$ | Predictions in $S_2$ |                                                                   | Predictions in $S_1$ | Predictions in $S_2$ |
|-----------------------------------------------------------------------|---------------|----------------------|----------------------|-------------------------------------------------------------------|----------------------|----------------------|
| $\mathcal{B}(B_u^+ \rightarrow \bar{D}_1^0 \ell'^+ \nu_{\ell'})$      | $4.2 \pm 1.8$ | $5.18 \pm 0.70$      | $5.18 \pm 0.70$      | $\mathcal{B}(B_u^+ \rightarrow \bar{D}_1^0 \tau^+ \nu_\tau)$      | $0.88 \pm 0.13$      | $0.87 \pm 0.65$      |
| $\mathcal{B}(B_d^0 \rightarrow D_1^{\prime-} \ell'^+ \nu_{\ell'})$    | $3.9 \pm 1.6$ | $4.85 \pm 0.65$      | $4.85 \pm 0.65$      | $\mathcal{B}(B_d^0 \rightarrow D_1^{\prime-} \tau^+ \nu_\tau)$    | $0.82 \pm 0.12$      | $0.81 \pm 0.61$      |
| $\mathcal{B}(B_s^0 \rightarrow D_{s1}^{\prime-} \ell'^+ \nu_{\ell'})$ | $\dots$       | $4.99 \pm 0.79$      | $5.15 \pm 0.82$      | $\mathcal{B}(B_s^0 \rightarrow D_{s1}^{\prime-} \tau^+ \nu_\tau)$ | $0.89 \pm 0.14$      | $0.90 \pm 0.67$      |
| $\mathcal{B}(B_u^+ \rightarrow \bar{D}_1^0 \ell'^+ \nu_{\ell'})$      | $6.6 \pm 2.2$ | $6.65 \pm 2.15$      | $6.65 \pm 2.15$      | $\mathcal{B}(B_u^+ \rightarrow \bar{D}_1^0 \tau^+ \nu_\tau)$      | $1.13 \pm 0.38$      | $1.04 \pm 0.55$      |
| $\mathcal{B}(B_d^0 \rightarrow D_1^{\prime-} \ell'^+ \nu_{\ell'})$    | $6.2 \pm 2.0$ | $6.20 \pm 2.00$      | $6.20 \pm 2.00$      | $\mathcal{B}(B_d^0 \rightarrow D_1^{\prime-} \tau^+ \nu_\tau)$    | $1.05 \pm 0.35$      | $0.97 \pm 0.51$      |
| $\mathcal{B}(B_s^0 \rightarrow D_{s1}^{\prime-} \ell'^+ \nu_{\ell'})$ | $\dots$       | $6.29 \pm 2.11$      | $6.60 \pm 2.23$      | $\mathcal{B}(B_s^0 \rightarrow D_{s1}^{\prime-} \tau^+ \nu_\tau)$ | $1.08 \pm 0.36$      | $1.06 \pm 0.56$      |

can constrain  $b_0$  and  $V_1^{D_1}(0)$ , and predict  $\mathcal{B}(B_u^+ \rightarrow \bar{D}_1^0 \tau^+ \nu_\tau)$ ,  $\mathcal{B}(B_d^0 \rightarrow D_1^{\prime-} \tau^+ \nu_\tau)$  and  $\mathcal{B}(B_s^0 \rightarrow D_{s1}^{\prime-} \tau^+ \nu_\tau)$ . We obtain that  $|a_0| = 4.50 \pm 0.52$  and  $|b_0| = 5.08 \pm 1.05$  in the  $S_1$  case, and  $|V_1^{D_1}(0)| = 0.57 \pm 0.32$  and  $|V_1^{D_1}(0)| = 0.51 \pm 0.15$  in the  $S_2$  case. In Ref. [79],  $|V_1^{D_1}(0)| = 0.19 \pm 0.02 \pm 0.01$  and  $|V_1^{D_1}(0)| = 0.58 \pm 0.01_{-0.03}^{+0.02}$ . Our constrained  $|V_1^{D_1}(0)|$  is larger than one in Ref. [79], but it close to the latter within  $2\sigma$  errors. Our constrained  $|V_1^{D_1}(0)|$  is consistent with one in Ref. [79].

Note that, if considering the SU(3) flavor breaking effects from different masses of  $u$ ,  $d$ , and  $s$  quarks, the nonperturbative coefficients of the  $B_s^0 \rightarrow D_{s1}^{\prime-} \ell'^+ \nu_\ell$  decays are different from ones of the  $B_u^+ \rightarrow \bar{D}_1^{(\prime)0} \ell'^+ \nu_\ell$  and  $B_d^0 \rightarrow D_1^{(\prime)-} \ell'^+ \nu_\ell$ . After the  $B_s^0 \rightarrow D_{s1}^{\prime-} \ell'^+ \nu_\ell$  decays are measured, one can estimate the SU(3) flavor breaking effects in the  $B \rightarrow D_1^{(\prime)} \ell'^+ \nu_\ell$  decays. The same situation also appears in the later  $B \rightarrow D_2^* \ell'^+ \nu_\ell$  decays.

Our predictions in the  $S_1$  and  $S_2$  cases are listed in Tab. III. Comparing the  $\mathcal{B}(B \rightarrow D_1^{(\prime)} \ell'^+ \nu_{\ell'})$  predictions in  $S_1$  to ones in  $S_2$ ,  $\mathcal{B}(B_u^+ \rightarrow \bar{D}_1^0 \ell'^+ \nu_{\ell'})$ ,  $\mathcal{B}(B_d^0 \rightarrow D_1^{\prime-} \ell'^+ \nu_{\ell'})$ ,  $\mathcal{B}(B_u^+ \rightarrow \bar{D}_1^0 \ell'^+ \nu_{\ell'})$  and  $\mathcal{B}(B_d^0 \rightarrow D_1^{\prime-} \ell'^+ \nu_{\ell'})$  are exactly same in both  $S_1$  and  $S_2$ .  $\mathcal{B}(B_s^0 \rightarrow D_{s1}^{\prime-} \ell'^+ \nu_{\ell'})$  and  $\mathcal{B}(B_s^0 \rightarrow D_{s1}^{\prime-} \ell'^+ \nu_{\ell'})$  are slightly difference in two cases due to different  $q'^2$  dependence. Comparing our SU(3) flavor symmetry predictions of  $\mathcal{B}(B \rightarrow D_1^{(\prime)} \ell'^+ \nu_{\ell'})$  with their experimental data, one can see that our predictions are coincident with the present data. As given in the sixth and seventh columns of Tab. III, the central values of  $\mathcal{B}(B \rightarrow D_1^{(\prime)} \tau^+ \nu_\tau)$  predictions in  $S_1$  to ones in  $S_2$  are similar to each other, nevertheless, the errors of the predictions, specially for  $\mathcal{B}(B \rightarrow D_1^{\prime-} \tau^+ \nu_\tau)$ , in  $S_2$  case are obviously larger than ones in  $S_1$  case. In later analysis, we will use the predictions of  $\mathcal{B}(B \rightarrow D_1^{(\prime)} \ell'^+ \nu_\ell)$  in  $S_1$  case to give the numerical results of  $\mathcal{B}(B \rightarrow D^* P \ell'^+ \nu_\ell)$  with the  $D_1'/D_1$  resonant states.

## 2. $D_1/D_1' \rightarrow D^* P$ decays

In the SU(3) flavor symmetry limit, the decay amplitudes of the  $D_1^{(\prime)} \rightarrow D^* P$  decays, which decay through the strong or electromagnetic interactions, can be simply parametrized as

$$A(D_1' \rightarrow D^* P) = a^A D_1^i D_j^* P_i^j, \quad A(D_1 \rightarrow D^* P) = a^B D_1^i D_j^* P_i^j, \quad (39)$$

where  $a^{A,B}$  are the nonperturbative coefficients. The amplitude relations of the  $D_1^{(\prime)} \rightarrow D^* P$  decays are given in Tab. IV.

TABLE IV: Decay amplitudes for the  $D_1^{(\prime)} \rightarrow D^*P$  decays by the SU(3) flavor symmetry.

| Decay amplitudes                       | SU(3) amplitudes | Decay amplitudes                       | SU(3) amplitudes |
|----------------------------------------|------------------|----------------------------------------|------------------|
| $A(D_1^0 \rightarrow D^{*0}\pi^0)$     | $a^A/\sqrt{2}$   | $A(D_1^0 \rightarrow D^{*0}\pi^0)$     | $a^B/\sqrt{2}$   |
| $A(D_1^0 \rightarrow D^{*+}\pi^-)$     | $a^A$            | $A(D_1^0 \rightarrow D^{*+}\pi^-)$     | $a^B$            |
| $A(D_1^{'+} \rightarrow D^{*+}\pi^0)$  | $-a^A/\sqrt{2}$  | $A(D_1^{'+} \rightarrow D^{*+}\pi^0)$  | $-a^B/\sqrt{2}$  |
| $A(D_1^{'+} \rightarrow D^{*0}\pi^+)$  | $a^A$            | $A(D_1^{'+} \rightarrow D^{*0}\pi^+)$  | $a^B$            |
| $A(D_{s1}^{'+} \rightarrow D^{*0}K^+)$ | $a^A$            | $A(D_{s1}^{'+} \rightarrow D^{*0}K^+)$ | $a^B$            |
| $A(D_{s1}^{'+} \rightarrow D^{*+}K^0)$ | $a^A$            | $A(D_{s1}^{'+} \rightarrow D^{*+}K^0)$ | $a^B$            |

TABLE V: Branching ratios of the  $D_1^{(\prime)} \rightarrow D^*P$  decays with the  $2\sigma$  errors (in units of  $10^{-2}$ ).

| Decay modes                        | Predictions      | Decay modes                         | Predictions       |
|------------------------------------|------------------|-------------------------------------|-------------------|
| $D_1^0 \rightarrow D^{*0}\pi^0$    | $33.65 \pm 0.03$ | $D_1^0 \rightarrow D^{*0}\pi^0$     | $26.71 \pm 11.44$ |
| $D_1^0 \rightarrow D^{*+}\pi^-$    | $66.35 \pm 0.03$ | $D_1^0 \rightarrow D^{*+}\pi^-$     | $52.73 \pm 22.61$ |
| $D_1^{'+} \rightarrow D^{*+}\pi^0$ | $33.23 \pm 0.01$ | $D_1^{'+} \rightarrow D^{*+}\pi^0$  | $26.48 \pm 11.35$ |
| $D_1^{'+} \rightarrow D^{*0}\pi^+$ | $66.77 \pm 0.01$ | $D_1^{'+} \rightarrow D^{*0}\pi^+$  | $53.19 \pm 22.79$ |
|                                    |                  | $D_{s1}^{'+} \rightarrow D^{*0}K^+$ | $54.98 \pm 7.13$  |
|                                    |                  | $D_{s1}^{'+} \rightarrow D^{*+}K^0$ | $45.02 \pm 7.13$  |

For the  $D_1' \rightarrow D^*P$  decays, they have not been observed yet. Decays  $D_1' \rightarrow D^*\eta$  and  $D_1' \rightarrow D^*\eta'$  are not allowed by the phase spaces. So we assume  $\mathcal{B}(D_1' \rightarrow D^*\pi) = 1$  and use the amplitude relations given in Tab. IV to obtain each branching ratios of the  $D_1' \rightarrow D^*\pi$  decays, which are listed in the second column of Tab. V. Decays  $D_{s1}^{'+} \rightarrow D^{*0}K^+$  and  $D_{s1}^{'+} \rightarrow D^{*+}K^0$  are also not allowed by the phase spaces. Decay  $D_{s1}^{'+} \rightarrow D_s^{*+}\pi^0$  is decayed by different way, its amplitude can not be related by nonperturbative coefficient  $a^A$ . The experimental data  $\mathcal{B}(D_{s1}^{'+} \rightarrow D_s^{*+}\pi^0) = (48 \pm 22)\%$  from PDG [1] will be used in later prediction of  $\mathcal{B}(B_s^0 \rightarrow D_{s1}^{\prime-}\ell^+\nu_\ell, D_{s1}^{\prime-} \rightarrow D_s^{*+}\pi^0)$ .

$D_1$  decays are dominant by  $D_1 \rightarrow D^*\pi$  and  $D_1 \rightarrow D\pi\pi$ . We use  $\mathcal{B}(D_1^- \rightarrow \bar{D}^{*0}\pi^-) = \frac{\mathcal{B}(B^0 \rightarrow D_1^-\ell\nu_\ell, D_1^- \rightarrow \bar{D}^{*0}\pi^-)}{\mathcal{B}(B^0 \rightarrow D_1^-\ell\nu_\ell)}$  to determine the nonperturbative coefficient  $a^B$ , and then give the branching ratios of every relevant decay mode. The results of  $D_1^0$  and  $D_1^{'+}$  decays are listed in the forth column of Tab. V. The decay width of  $D_{s1}^+, \Gamma_{D_{s1}^+} = (0.92 \pm 0.10) \times 10^{-3}$ , is much smaller than ones of  $D_1^{0,+}, \Gamma_{D_1^{0,+}} = (31.3 \pm 3.8) \times 10^{-3}$ . The predictions of  $\mathcal{B}(D_{s1}^+ \rightarrow D^{*0}K^+)$  and  $\mathcal{B}(D_{s1}^+ \rightarrow D^{*+}K^0)$  are much larger than 100% by the constrained  $a^B$  from  $\mathcal{B}(D_1^- \rightarrow \bar{D}^{*0}\pi^-)$ , they are not right, and we will not use them. Using  $\frac{\mathcal{B}(D_{s1}^+ \rightarrow \bar{D}^{*+}K^0)}{\mathcal{B}(D_{s1}^+ \rightarrow \bar{D}^{*+}K^0)} = 0.85 \pm 0.24$  from PDG [1] and assuming  $\mathcal{B}(D_{s1}^+ \rightarrow \bar{D}_s^{*+}K^0) + \mathcal{B}(D_{s1}^+ \rightarrow \bar{D}^{*+}K^0) = 1$ , these branching ratios are obtained, which are given in the forth column of Tab. V.

### 3. $\mathcal{B}(B \rightarrow D^*P\ell^+\nu_\ell)$ with the $D_1'/D_1$ resonant states

In terms of  $\mathcal{B}(B \rightarrow D_1^{(\prime)}\ell^+\nu_\ell)$  in the  $S_1$  case given in Tab. III and  $\mathcal{B}(D_1^{(\prime)} \rightarrow D^*P)$  given in Tab. V, after considering relevant experimental bounds in Eqs. (17-18) and Eqs. (21-22), one can obtain  $\mathcal{B}(B \rightarrow D^*P\ell^+\nu_\ell)$  with the resonant

charmed axial vector. The results are summarized in the third column of Tab. II. There are four experimental data for  $\mathcal{B}(B_u^+ \rightarrow D^{*-}\pi^+\ell'^+\nu_{\ell'})_{[D_1'^0, D_1^0]}$  and  $\mathcal{B}(B_d^0 \rightarrow \bar{D}^{*0}\pi^-\ell'^+\nu_{\ell'})_{[D_1'^-, D_1^-]}$ , which are also listed in the third column of Tab. II. Comparing with the data, one can see that  $\mathcal{B}(B_u^+ \rightarrow D^{*-}\pi^+\ell'^+\nu_{\ell'})_{[D_1'^0, D_1^0]}$  give the further effective constraints. The predictions of  $\mathcal{B}(B_d^0 \rightarrow \bar{D}^{*0}\pi^-\ell'^+\nu_{\ell'})_{[D_1'^-, D_1^-]}$  are consistent with their experimental data but with smaller errors.

## B. Resonant $B \rightarrow D^*P\ell^+\nu_\ell$ decays with the charmed tensors

For the semileptonic  $B \rightarrow D_2^*\ell^+\nu_\ell$  decays, the branching ratios of  $B_u^+ \rightarrow \bar{D}_2^{*0}\ell'^+\nu_{\ell'}$  and  $B_d^0 \rightarrow D_2^{*-}\ell'^+\nu_{\ell'}$  decays have been measured by the Belle Collaboration [81], and the experimental data with  $2\sigma$  errors are listed in the second column of Tab. VI. In similar to  $S_2$  case in Sec. III A, the SU(3) flavor symmetry predictions of the non-measured  $B \rightarrow D_2^*\ell^+\nu_\ell$  decays from the two experimental data are obtained in Ref. [82], and they are listed in the third column of Tab. VI.

For the decay amplitudes of the  $D_2^* \rightarrow D^*P$  decays, they are similar to ones of the  $D_1 \rightarrow D^*P$  decays in Eq. (39) in terms of replacing  $a^B$  by  $a^T$  and replacing  $D_1$  by  $D_2^*$ . The decay amplitudes of the specific  $D_2^* \rightarrow D^*P$  decays are similar to ones in Tab. IV, so we will not show them here. Decays  $D_2^* \rightarrow D^*P$  and  $D_2^* \rightarrow DP$  have not been measured up to now, therefore, we can not constrain  $a^T$  directly. In Ref. [81], they make a assumption that  $\mathcal{B}(D_2^* \rightarrow D^*\pi) + \mathcal{B}(D_2^* \rightarrow D\pi) = 1$ . Using  $\frac{\mathcal{B}(D_2^* \rightarrow D\pi)}{\mathcal{B}(D_2^* \rightarrow D^*\pi)} = 1.52 \pm 0.14$  within  $2\sigma$  errors from PDG [1], and assuming one of  $\mathcal{B}(D_2^{*0} \rightarrow D\pi, D^*\pi)$ ,  $\mathcal{B}(D_2^{*+} \rightarrow D\pi, D^*\pi)$  and  $\mathcal{B}(D_{s2}^{*+} \rightarrow DK, D^*K)$  is equal to one and other two values are less than or equal to one, one can constrain on the nonperturbative coefficients  $a^T$  and  $b^T$  (where  $b^T$  is similar to  $a^T$  but for the  $D_2^* \rightarrow DP$  decays), and they are  $|a^T| = 9.06 \pm 0.85$  and  $|b^T| = 25.14 \pm 1.47$ . And then, the branching ratios of the  $D_2^* \rightarrow D^*P$  decays can be predicted, which are summarized in the second column of Tab. VII. In addition, their decay width predictions and previous width predictions are also given in the forth and fifth columns of Tab. VII, respectively. Our width predictions are larger than ones in Ref. [83].

Using  $\mathcal{B}(B \rightarrow D_2^*\ell^+\nu_\ell)$  given in Tab. VI and  $\mathcal{B}(D_2^* \rightarrow D^*P)$  given in Tab. VII, after considering relevant experimental bounds in Eq. (19) and Eq. (18), one can obtain  $\mathcal{B}(B \rightarrow D^*P\ell^+\nu_\ell)$  with the charmed tensor, and they are summarized in the forth column of Tab. II. Comparing with the data of  $\mathcal{B}(B_u^+ \rightarrow D^{*-}\pi^+\ell'^+\nu_{\ell'})$  and  $\mathcal{B}(B_d^0 \rightarrow \bar{D}^{*0}\pi^-\ell'^+\nu_{\ell'})$ , one can see the upper limit of  $\mathcal{B}(B_d^0 \rightarrow \bar{D}^{*0}\pi^-\ell'^+\nu_{\ell'})$  gives further constraint on the relevant nonperturbative parameters and the branching ratio predictions.

## IV. Global analysis

In this section, we will analyse the numerical results of  $\mathcal{B}(B \rightarrow D^*P\ell^+\nu_\ell)$  given in Tab. II. As given in Tab. II, we can see that  $\mathcal{B}(B \rightarrow D^*P\tau^+\nu_\tau)$  with the non-resonant and the charmed axial vector resonant contributions are much smaller than corresponding  $\mathcal{B}(B \rightarrow D^*P\ell^+\nu_\ell)$  since the former small phase spaces.

We can see that  $B^{0,+} \rightarrow D^*\eta\ell^+\nu_\ell$ ,  $B^{0,+} \rightarrow D^*\eta'\ell^+\nu_\ell$ ,  $B_s^0 \rightarrow D_s^*\eta\ell^+\nu_\ell$ ,  $B_s^0 \rightarrow D_s^*\eta'\ell^+\nu_\ell$  and  $B^{0,+} \rightarrow D_s^*K\ell^+\nu_\ell$  decays only receive the non-resonant contributions. Under the SU(3) flavor symmetry,  $\mathcal{B}(B_s^0 \rightarrow D_s^{*-}\pi^0\ell^+\nu_\ell)_N = 0$ , and  $\mathcal{B}(B_s^0 \rightarrow D_s^{*-}\pi^0\ell^+\nu_\ell)_{all}$  only include the  $D_{s1}^{*-}$  resonant contributions.

Other  $B \rightarrow D^*P\ell^+\nu_\ell$  decays receive the contributions of the non-resonant states, the charmed axial vector resonant

TABLE VI: Branching ratios of the  $B \rightarrow D_2^* \ell^+ \nu_\ell$  decays with the  $2\sigma$  errors.

|                                                                                           | Exp. data [81] | Predictions [82] |
|-------------------------------------------------------------------------------------------|----------------|------------------|
| $\mathcal{B}(B_u^+ \rightarrow \overline{D}_2^{*0} \ell'^+ \nu_{\ell'}) (\times 10^{-3})$ | $2.9 \pm 0.6$  | $3.20 \pm 0.30$  |
| $\mathcal{B}(B_d^0 \rightarrow D_2^{*-} \ell'^+ \nu_{\ell'}) (\times 10^{-3})$            | $2.7 \pm 0.6$  | $2.98 \pm 0.29$  |
| $\mathcal{B}(B_s^0 \rightarrow D_{s2}^{*-} \ell'^+ \nu_{\ell'}) (\times 10^{-3})$         | $\dots$        | $2.72 \pm 0.27$  |
| $\mathcal{B}(B_u^+ \rightarrow \overline{D}_2^{*0} \tau^+ \nu_\tau) (\times 10^{-4})$     | $\dots$        | $2.15 \pm 0.47$  |
| $\mathcal{B}(B_d^0 \rightarrow D_2^{*-} \tau^+ \nu_\tau) (\times 10^{-4})$                | $\dots$        | $1.97 \pm 0.43$  |
| $\mathcal{B}(B_s^0 \rightarrow D_{s2}^{*-} \tau^+ \nu_\tau) (\times 10^{-4})$             | $\dots$        | $1.73 \pm 0.38$  |

TABLE VII: Branching ratios of the  $T \rightarrow D^* P$  decays with the  $2\sigma$  errors.

| Decay modes                                | Branching ratios ( $\times 10^{-2}$ ) | Decay widths (MeV) | Decay widths from Ref. [83] (MeV) |
|--------------------------------------------|---------------------------------------|--------------------|-----------------------------------|
| $D_2^{*0}(2460) \rightarrow D^{*0} \pi^0$  | $12.96 \pm 2.29$                      | $6.19 \pm 1.12$    | $2.40_{-0.94}^{+1.74}$            |
| $D_2^{*0}(2460) \rightarrow D^{*+} \pi^-$  | $24.56 \pm 4.36$                      | $11.71 \pm 2.13$   | $3.99_{-1.56}^{+1.22}$            |
| $D_2^{*+}(2460) \rightarrow D^{*+} \pi^0$  | $12.49 \pm 2.09$                      | $5.98 \pm 1.04$    | $\dots$                           |
| $D_2^{*+}(2460) \rightarrow D^{*0} \pi^+$  | $25.49 \pm 4.25$                      | $12.21 \pm 2.13$   | $\dots$                           |
| $D_{s2}^{*+}(2573) \rightarrow D^{*0} K^+$ | $6.07 \pm 1.54$                       | $1.02 \pm 0.22$    | $0.20_{-0.07}^{+0.09}$            |
| $D_{s2}^{*+}(2573) \rightarrow D^{*+} K^0$ | $4.60 \pm 1.19$                       | $0.78 \pm 0.17$    | $0.15_{-0.05}^{+0.06}$            |

states and the charmed tensor resonant states. In the  $\ell = e, \mu$  decays, the charmed axial vector resonant contributions are largest. But in the  $\ell = \tau$  decays, both the charmed axial vector and tensor resonant contributions are important. We sum all three kinds of contributions as  $\mathcal{B}_{all}$  (for the non-resonant branching ratios, they are taken in the ranges of 0 to their maximal values listed in the second column of Tab. II.), after considering the experimental bounds of  $\mathcal{B}(B^+ \rightarrow D^{*-} \pi^+ \ell'^+ \nu_{\ell'})_{all}$  in Eq. (16) and  $\mathcal{B}(B^0 \rightarrow D^{*0} \pi^+ \ell'^+ \nu_{\ell'})_{all}$  in Eq. (20), the results of  $\mathcal{B}_{all}$  are listed in the last column of Tab. II.

Noted that the interference terms between non-resonant, the axial vector resonant and the tensor resonant contributions exist, but they have not been considered in this work, and they might not be ignored if all three kinds of contributions or two kinds of contributions are important in the decays. And they will be studied in our succeeding work.

## V. Summary

In this paper, we have studied the  $B \rightarrow D^* P \ell^+ \nu_\ell$  decay processes with the non-resonant, the charmed axial vector resonant and the charmed tensor resonant contributions by using the SU(3) flavor symmetry. The amplitude relations of the non-resonant  $B \rightarrow D^* P \ell^+ \nu_\ell$  decays, the semileptonic  $B \rightarrow D^{**} \ell^+ \nu_\ell$  decays and the non-leptonic  $D^{**} \rightarrow D^* P$  decays have been obtained by the SU(3) flavor symmetry. And then using relevant experimental data of the  $B \rightarrow D^* P \ell^+ \nu_\ell$  decays, we have presented a theoretical analysis of these decays. Our main results can be

summarized as follows.

We have found that  $B^{0,+} \rightarrow D^* \eta \ell^+ \nu_\ell$ ,  $B^{0,+} \rightarrow D^* \eta' \ell^+ \nu_\ell$ ,  $B_s^0 \rightarrow D_s^* \eta \ell^+ \nu_\ell$ ,  $B_s^0 \rightarrow D_s^* \eta' \ell^+ \nu_\ell$  and  $B^{0,+} \rightarrow D_s^* K \ell^+ \nu_\ell$  decays only receive non-resonant contributions. Decays  $B_s^0 \rightarrow D_s^{*-} \pi^0 \ell^+ \nu_\ell$  only receive the  $D_{s1}'$  resonant contributions, and the latter contributions might be larger than the former. Other decays receive the non-resonant, the charmed axial vector resonant and the charmed tensor resonant contributions, all three kinds of contributions are important in most of decays. Many branching ratios have been predicted on the order of  $\mathcal{O}(10^{-3})$ , and they might be measured in future experiments.

Although only approximate predictions can be obtained by the SU(3) flavor symmetry, they are still useful for understanding these decays. So far, our predictions under the SU(3) flavor symmetry are quite agree with present experimental data. Our predictions could be tested in future experiments, such as LHCb and Belle II.

## ACKNOWLEDGEMENTS

The work was supported by the National Natural Science Foundation of China (No. 12365014 and No. 12175088).

## VI. References

- 
- [1] R. L. Workman *et al.* [Particle Data Group], PTEP **2022**, 083C01 (2022).
  - [2] F. Meier *et al.* [Belle], Phys. Rev. D **107**, no.9, 092003 (2023) [arXiv:2211.09833 [hep-ex]].
  - [3] P. del Amo Sanchez *et al.* [BaBar], Phys. Rev. Lett. **107**, 041804 (2011) [arXiv:1012.4158 [hep-ex]].
  - [4] T. Feldmann, D. Van Dyk and K. K. Vos, JHEP **10**, 030 (2018) [arXiv:1807.01924 [hep-ph]].
  - [5] X. W. Kang, B. Kubis, C. Hanhart and U. G. Meißner, Phys. Rev. D **89**, 053015 (2014) [arXiv:1312.1193 [hep-ph]].
  - [6] Y. J. Shi, W. Wang and S. Zhao, Eur. Phys. J. C **77**, no.7, 452 (2017) [arXiv:1701.07571 [hep-ph]].
  - [7] Y. J. Shi, C. Y. Seng, F. K. Guo, B. Kubis, U. G. Meißner and W. Wang, JHEP **04**, 086 (2021) [arXiv:2011.00921 [hep-ph]].
  - [8] T. Sekihara and E. Oset, Phys. Rev. D **92**, no.5, 054038 (2015) [arXiv:1507.02026 [hep-ph]].
  - [9] S. Cheng, A. Khodjamirian and J. Virto, JHEP **05**, 157 (2017) [arXiv:1701.01633 [hep-ph]].
  - [10] C. Hambrock and A. Khodjamirian, Nucl. Phys. B **905**, 373-390 (2016) [arXiv:1511.02509 [hep-ph]].
  - [11] P. Böer, T. Feldmann and D. van Dyk, JHEP **02**, 133 (2017) [arXiv:1608.07127 [hep-ph]].
  - [12] X. G. He, Eur. Phys. J. C **9**, 443-448 (1999) [arXiv:hep-ph/9810397 [hep-ph]].
  - [13] X. G. He, Y. K. Hsiao, J. Q. Shi, Y. L. Wu and Y. F. Zhou, Phys. Rev. D **64**, 034002 (2001) [arXiv:hep-ph/0011337 [hep-ph]].
  - [14] H. K. Fu, X. G. He and Y. K. Hsiao, Phys. Rev. D **69**, 074002 (2004) [arXiv:hep-ph/0304242 [hep-ph]].
  - [15] Y. K. Hsiao, C. F. Chang and X. G. He, Phys. Rev. D **93**, no.11, 114002 (2016) [arXiv:1512.09223 [hep-ph]].
  - [16] X. G. He and G. N. Li, Phys. Lett. B **750**, 82-88 (2015) [arXiv:1501.00646 [hep-ph]].
  - [17] M. He, X. G. He and G. N. Li, Phys. Rev. D **92**, no.3, 036010 (2015) [arXiv:1507.07990 [hep-ph]].
  - [18] N. G. Deshpande and X. G. He, Phys. Rev. Lett. **75**, 1703-1706 (1995) [arXiv:hep-ph/9412393 [hep-ph]].
  - [19] M. Gronau, O. F. Hernandez, D. London and J. L. Rosner, Phys. Rev. D **50**, 4529-4543 (1994) [arXiv:hep-ph/9404283 [hep-ph]].

[hep-ph]].

- [20] M. Gronau, O. F. Hernandez, D. London and J. L. Rosner, Phys. Rev. D **52**, 6356-6373 (1995) [arXiv:hep-ph/9504326 [hep-ph]].
- [21] S. Shivashankara, W. Wu and A. Datta, Phys. Rev. D **91**, no.11, 115003 (2015) [arXiv:1502.07230 [hep-ph]].
- [22] S. H. Zhou, Q. A. Zhang, W. R. Lyu and C. D. Lü, Eur. Phys. J. C **77**, no.2, 125 (2017) [arXiv:1608.02819 [hep-ph]].
- [23] H. Y. Cheng, C. W. Chiang and A. L. Kuo, Phys. Rev. D **91**, no.1, 014011 (2015) [arXiv:1409.5026 [hep-ph]].
- [24] R. M. Wang, Y. G. Xu, C. Hua and X. D. Cheng, Phys. Rev. D **103**, no.1, 013007 (2021) [arXiv:2101.02421 [hep-ph]].
- [25] R. M. Wang, X. D. Cheng, Y. Y. Fan, J. L. Zhang and Y. G. Xu, J. Phys. G **48**, no.8, 085001 (2021) [arXiv:2008.06624 [hep-ph]].
- [26] Y. Grossman and D. J. Robinson, JHEP **04**, 067 (2013) [arXiv:1211.3361 [hep-ph]].
- [27] D. Pirtskhalava and P. Uttayarat, Phys. Lett. B **712**, 81-86 (2012) [arXiv:1112.5451 [hep-ph]].
- [28] M. J. Savage and R. P. Springer, Phys. Rev. D **42**, 1527-1543 (1990).
- [29] M. J. Savage, Phys. Lett. B **257**, 414-418 (1991).
- [30] G. Altarelli, N. Cabibbo and L. Maiani, Phys. Lett. B **57**, 277-280 (1975).
- [31] C. D. Lü, W. Wang and F. S. Yu, Phys. Rev. D **93**, no.5, 056008 (2016) [arXiv:1601.04241 [hep-ph]].
- [32] C. Q. Geng, Y. K. Hsiao, Y. H. Lin and L. L. Liu, Phys. Lett. B **776**, 265-269 (2018) [arXiv:1708.02460 [hep-ph]].
- [33] C. Q. Geng, Y. K. Hsiao, C. W. Liu and T. H. Tsai, Phys. Rev. D **97**, no.7, 073006 (2018) [arXiv:1801.03276 [hep-ph]].
- [34] C. Q. Geng, Y. K. Hsiao, C. W. Liu and T. H. Tsai, JHEP **11**, 147 (2017) [arXiv:1709.00808 [hep-ph]].
- [35] C. Q. Geng, C. W. Liu, T. H. Tsai and S. W. Yeh, Phys. Lett. B **792**, 214-218 (2019) [arXiv:1901.05610 [hep-ph]].
- [36] W. Wang, Z. P. Xing and J. Xu, Eur. Phys. J. C **77**, no.11, 800 (2017) [arXiv:1707.06570 [hep-ph]].
- [37] D. Wang, Eur. Phys. J. C **79**, no.5, 429 (2019) [arXiv:1901.01776 [hep-ph]].
- [38] D. Wang, P. F. Guo, W. H. Long and F. S. Yu, JHEP **03**, 066 (2018) [arXiv:1709.09873 [hep-ph]].
- [39] S. Müller, U. Nierste and S. Schacht, Phys. Rev. D **92**, no.1, 014004 (2015) [arXiv:1503.06759 [hep-ph]].
- [40] R. M. Wang, M. Z. Yang, H. B. Li and X. D. Cheng, Phys. Rev. D **100**, no.7, 076008 (2019) [arXiv:1906.08413 [hep-ph]].
- [41] Y. G. Xu, X. D. Cheng, J. L. Zhang and R. M. Wang, J. Phys. G **47**, no.8, 085005 (2020) [arXiv:2001.06907 [hep-ph]].
- [42] H. M. Chang, M. González-Alonso and J. Martin Camalich, Phys. Rev. Lett. **114**, no.16, 161802 (2015) [arXiv:1412.8484 [hep-ph]].
- [43] P. Zenczykowski, Phys. Rev. D **73**, 076005 (2006) [arXiv:hep-ph/0512122 [hep-ph]].
- [44] P. Zenczykowski, Nucl. Phys. B Proc. Suppl. **167**, 54-57 (2007) [arXiv:hep-ph/0610191 [hep-ph]].
- [45] N. Cabibbo, Phys. Rev. Lett. **10**, 531-533 (1963).
- [46] G. Martinelli, M. Naviglio, S. Simula and L. Vittorio, Phys. Rev. D **106**, no.9, 093002 (2022) [arXiv:2204.05925 [hep-ph]].
- [47] R. M. Wang, Y. Qiao, Y. J. Zhang, X. D. Cheng and Y. G. Xu, [arXiv:2301.00090 [hep-ph]].
- [48] M. Imbeault and D. London, Phys. Rev. D **84**, 056002 (2011) [arXiv:1106.2511 [hep-ph]].
- [49] Y. L. Wu and Y. F. Zhou, Phys. Rev. D **72**, 034037 (2005) [arXiv:hep-ph/0503077 [hep-ph]].
- [50] H. Y. Cheng and C. W. Chiang, Phys. Rev. D **86**, 014014 (2012) [arXiv:1205.0580 [hep-ph]].
- [51] A. Dery, M. Ghosh, Y. Grossman and S. Schacht, JHEP **03** (2020), 165 [arXiv:2001.05397 [hep-ph]].
- [52] S. Sasaki and T. Yamazaki, Phys. Rev. D **79** (2009), 074508 [arXiv:0811.1406 [hep-ph]].
- [53] G. S. Yang and H. C. Kim, Phys. Rev. C **92**, 035206 (2015) [arXiv:1504.04453 [hep-ph]].
- [54] T. N. Pham, Phys. Rev. D **87** (2013) no.1, 016002 [arXiv:1210.3981 [hep-ph]].
- [55] C. Q. Geng, Y. K. Hsiao, C. W. Liu and T. H. Tsai, Eur. Phys. J. C **78** (2018) no.7, 593 [arXiv:1804.01666 [hep-ph]].
- [56] R. Flores-Mendieta, E. E. Jenkins and A. V. Manohar, Phys. Rev. D **58** (1998), 094028 [arXiv:hep-ph/9805416 [hep-ph]].
- [57] D. Xu, G. N. Li and X. G. He, Int. J. Mod. Phys. A **29** (2014), 1450011 [arXiv:1307.7186 [hep-ph]].
- [58] X. G. He, G. N. Li and D. Xu, Phys. Rev. D **91** (2015) no.1, 014029 [arXiv:1410.0476 [hep-ph]].

- [59] H. Y. Cheng, C. Y. Cheung, W. Dimm, G. L. Lin, Y. C. Lin, T. M. Yan and H. L. Yu, Phys. Rev. D **48**, 3204-3220 (1993) [arXiv:hep-ph/9305340 [hep-ph]].
- [60] S. Y. Tsai and Y. K. Hsiao, [arXiv:2107.03634 [hep-ph]].
- [61] A. Le Yaouanc, J. P. Leroy and P. Roudeau, Phys. Rev. D **99**, no.7, 073010 (2019) [arXiv:1806.09853 [hep-ph]].
- [62] Y. J. Shi and U. G. Meißner, Eur. Phys. J. C **81**, no.5, 412 (2021) [arXiv:2103.12977 [hep-ph]].
- [63] C. S. Kim, J. Lee and W. Namgung, Phys. Rev. D **60**, 094019 (1999) [arXiv:hep-ph/9904470 [hep-ph]].
- [64] C. S. Kim, J. Lee and W. Namgung, Phys. Rev. D **59**, 114005 (1999) [arXiv:hep-ph/9811395 [hep-ph]].
- [65] N. N. Achasov, A. V. Kiselev and G. N. Shestakov, Phys. Rev. D **102**, no.1, 016022 (2020) [arXiv:2005.06455 [hep-ph]].
- [66] J. Wiss, eConf **C070805**, 34 (2007) [arXiv:0709.3247 [hep-ex]].
- [67] S. Faller, T. Feldmann, A. Khodjamirian, T. Mannel and D. van Dyk, Phys. Rev. D **89**, no.1, 014015 (2014) [arXiv:1310.6660 [hep-ph]].
- [68] W. Wang, Phys. Lett. B **759**, 501-506 (2016) [arXiv:1602.05288 [hep-ph]].
- [69] N. N. Achasov, A. V. Kiselev and G. N. Shestakov, Phys. Rev. D **104**, no.1, 016034 (2021) [arXiv:2106.10670 [hep-ph]].
- [70] E. J. Gustafson, F. Herren, R. S. Van de Water, R. van Tonder and M. L. Wagman, [arXiv:2311.00864 [hep-ph]].
- [71] S. Okubo, Phys. Lett. **5**, 165-168 (1963).
- [72] H. J. Lipkin, Nucl. Phys. B **291**, 720-730 (1987).
- [73] H. J. Lipkin and B. s. Zou, Phys. Rev. D **53**, 6693-6696 (1996).
- [74] H. Y. Cheng and X. W. Kang, Eur. Phys. J. C **77** (2017) no.9, 587 [erratum: Eur. Phys. J. C **77** (2017) no.12, 863] [arXiv:1707.02851 [hep-ph]].
- [75] H. Y. Cheng, C. K. Chua and C. W. Hwang, Phys. Rev. D **69** (2004), 074025 [arXiv:hep-ph/0310359 [hep-ph]].
- [76] P. Colangelo, F. De Fazio and F. Loparco, Phys. Rev. D **100**, no.7, 075037 (2019) [arXiv:1906.07068 [hep-ph]].
- [77] R. H. Li, C. D. Lu and W. Wang, Phys. Rev. D **79**, 034014 (2009) [arXiv:0901.0307 [hep-ph]].
- [78] Y. J. Sun, Z. G. Wang and T. Huang, Chin. Phys. C **36**, 1046-1054 (2012) [arXiv:1106.4915 [hep-ph]].
- [79] R. C. Verma, J. Phys. G **39**, 025005 (2012) [arXiv:1103.2973 [hep-ph]].
- [80] M. A. Ivanov, J. G. Körner, J. N. Pandya, P. Santorelli, N. R. Soni and C. T. Tran, Front. Phys. (Beijing) **14** (2019) no.6, 64401 [arXiv:1904.07740 [hep-ph]].
- [81] F. Abudinén *et al.* [Belle-II], Phys. Rev. D **107**, no.7, 072002 (2023) [arXiv:2205.06372 [hep-ex]].
- [82] Ru-Min Wang, Yi-Jie Zhang, Meng-Yuan Wan, Xiao-Dong Cheng, Yuan-Guo Xu, in preparation.
- [83] Z. Y. Li, Z. G. Wang and G. L. Yu, Mod. Phys. Lett. A **31**, no.06, 1650036 (2016) [arXiv:1506.07761 [hep-ph]].