

GENERALIZED GYROVECTOR SPACES REVISITED

TOSHIKAZU ABE AND OSAMU HATORI

ABSTRACT. We revise a proof of a Mazur-Ulam theorem for generalized gyrovector spaces.

1. INTRODUCTION

Ungar introduced the concept of a gyrovector space, which is discussed in detail in [5, 6]. In contrast, we later proposed the idea of a generalized gyrovector space (GGV) in [2] and proved a Mazur-Ulam theorem for it. However, we have identified some errors in that paper and have revised them in this paper.

We want to clarify that statements of Theorem 13 and Corollary 14 in [2] are correct, but their proofs need some revisions. Specifically, we need to correct an equation on the sixth line of page 398, which should read as $\|\phi(e)\| = 0_{\|\phi(G)\|}$ instead of $\|\phi(e)\| = 0$. It is important to note that $0_{\|\phi(G)\|}$ may not always represent the real number 0, and we have provided an example in Example 3.3. Additionally, we would like to mention that all examples of GGVs presented in [2] and [3] satisfy $\|\phi(e)\| = 0$ (i.e., the real number 0).

Note that not all GGVs (Generalized Gyrovector Spaces) are normed gyrolinear spaces, which the first author defines [1]. However, a GGV that satisfies $\|\phi(e)\| = 0$ can be considered a normed gyrolinear space. We needed to revise inequalities (10) in [2], and now inequality (5.2) is the corrected version. After these revisions, we present revised proofs of the Mazur-Ulam theorem for a GGV in Theorem 4.4 and Corollary 4.5. Our revised proof corrects the errors found in the original proof.

The terminologies are essentially due to that in [2]. See also [5, 6].

2. PRELIMINARY

Before recalling the definition of a generalized gyrovector space, it would be convenient to confirm the following.

Proposition 2.1. *Let (G, \oplus) be a gyrocommutative gyrogroup with the map $\otimes : \mathbb{R} \times G \rightarrow G$. We denote e the unit of (G, \oplus) . Let ϕ be an injection from G into a real normed space $(\mathbb{V}, \|\cdot\|)$. We assume*

(GGV1) $1 \otimes a = a$ for any $a \in G$;

(GGV2) $(r_1 + r_2) \otimes a = (r_1 \otimes a) \oplus (r_2 \otimes a)$ for any $a \in G$, $r_1, r_2 \in \mathbb{R}$;

(GGV3) $(r_1 r_2) \otimes a = r_1 \otimes (r_2 \otimes a)$ for any $a \in G$, $r_1, r_2 \in \mathbb{R}$;

(GGVV) $\|\phi(G)\| = \{\pm\|\phi(a)\| \in \mathbb{R} : a \in G\}$ is a real linear space of dimension one with vector addition \oplus' and scalar multiplication \otimes' .

(GGV7) $\|\phi(r \otimes a)\| = |r| \otimes' \|\phi(a)\|$ for any $a \in G$, $r \in \mathbb{R}$;

2020 Mathematics Subject Classification. 51M10, 20N05.

Key words and phrases. gyrocommutative gyrogroups, generalized gyrovector spaces, normed gyrolinear spaces.

Then we have the following.

- (i) $\|\phi(\mathbf{e})\| = 0_{\|\phi(G)\|}$;
- (ii) $r \otimes \mathbf{e} = \mathbf{e}$ for all $r \in \mathbb{R}$;
- (iii) $0 \otimes \mathbf{a} = \mathbf{e}$ for all $\mathbf{a} \in G$;
- (iv) $r \otimes \mathbf{a} = \mathbf{e}$ for $r \in \mathbb{R}$ and $\mathbf{a} \in G$ implies that $r = 0$ or $\mathbf{a} = \mathbf{e}$;
- (v) $\ominus(\alpha \otimes \mathbf{a}) = (-\alpha) \otimes \mathbf{a}$ for all $\alpha \in \mathbb{R}$, $\mathbf{a} \in G$, in particular, $\ominus \mathbf{a} = (-1) \otimes \mathbf{a}$ for all $\mathbf{a} \in G$
- (vi) $\|\phi(\mathbf{a})\| \neq 0$ for all $\mathbf{a} \in G \setminus \{\mathbf{e}\}$. $\|\phi(r \otimes \mathbf{a})\| \neq 0$ for all $r \in \mathbb{R} \setminus \{0\}$ and $\mathbf{a} \in G \setminus \{\mathbf{e}\}$.

Proof. By the condition (GGV2) we have

$$0 \otimes \mathbf{a} = (0+0) \otimes \mathbf{a} = 0 \otimes \mathbf{a} \oplus 0 \otimes \mathbf{a}$$

for every $\mathbf{a} \in G$. By the left cancelation law for gyrogroups, we infer that $\mathbf{e} = 0 \otimes \mathbf{a}$, proving (iii). In particular, since $\mathbf{e} = 0 \otimes \mathbf{e}$ we obtain

$$\|\phi(\mathbf{e})\| = \|\phi(0 \otimes \mathbf{e})\| = 0 \otimes' \|\phi(\mathbf{e})\| = 0_{\|\phi(G)\|}$$

by (GGV)7, proving (i).

For every $r \in \mathbb{R}$ we have by (GGV3) that

$$r \otimes \mathbf{e} = r \otimes (0 \otimes \mathbf{e}) = (r \times 0) \otimes \mathbf{e} = 0 \otimes \mathbf{e} = \mathbf{e},$$

we are ensuring (ii).

Suppose that $r \otimes \mathbf{a} = \mathbf{e}$ for some $r \in \mathbb{R}$ and $\mathbf{a} \in G$. If $r \neq 0$, then we have by (GGV1) and (GGV3) that

$$\mathbf{a} = 1 \otimes \mathbf{a} = \frac{1}{r} \otimes (r \otimes \mathbf{a}) = \frac{1}{r} \otimes \mathbf{e} = \mathbf{e}.$$

Thus, (iv) holds.

By (GGV2) we have

$$(-\alpha) \otimes \mathbf{a} \oplus \alpha \otimes \mathbf{a} = (-\alpha + \alpha) \otimes \mathbf{a} = 0 \otimes \mathbf{a} = \mathbf{e}.$$

The uniqueness of $\ominus(\alpha \otimes \mathbf{a})$ ensures that $(-\alpha) \otimes \mathbf{a} = \ominus(\alpha \otimes \mathbf{a})$, proving (v).

Suppose that $\|\phi(\mathbf{a})\| = 0$ for some $\mathbf{a} \in G$. Then we obtain

$$\|\phi(\ominus \mathbf{a})\| = \|\phi((-1) \otimes \mathbf{a})\| = |-1| \otimes' \|\phi(\mathbf{a})\| = \|\phi(\mathbf{a})\| = 0.$$

Thus we have $\phi(\ominus \mathbf{a}) = \phi(\mathbf{a}) = 0_{\mathbb{V}}$ as \mathbb{V} is a normed linear space. As ϕ is an injection we infer that $\ominus \mathbf{a} = \mathbf{a}$, so

$$\mathbf{a} = 1 \otimes \mathbf{a} = \frac{1}{2} \otimes (2 \otimes \mathbf{a}) = \frac{1}{2} \otimes ((1+1) \otimes \mathbf{a}) = \frac{1}{2} (\mathbf{a} \oplus \mathbf{a}) = \frac{1}{2} \otimes (\ominus \mathbf{a} \oplus \mathbf{a}) = \frac{1}{2} \otimes \mathbf{e} = \mathbf{e}.$$

We conclude that $\|\phi(\mathbf{a})\| \neq 0$ if $\mathbf{a} \neq \mathbf{e}$. By (iv), we infer that $\|\phi(r \otimes \mathbf{a})\| \neq 0$ if $\mathbf{a} \in G \setminus \{\mathbf{e}\}$ and $r \in \mathbb{R} \setminus \{0\}$. We have proved (vi). □

Note that the proof of Proposition 2.1 shows that $\|\phi(\mathbf{a})\| = 0$ only if $\mathbf{a} = \mathbf{e}$. One might expect $\|\phi(\mathbf{e})\| = 0$ for a GGV, but Example 3.3 illustrates that this is not always the case.

In this paper we simply write $\alpha \otimes \mathbf{a} \oplus \beta \otimes \mathbf{b}$ instead of $(\alpha \otimes \mathbf{a}) \oplus (\beta \otimes \mathbf{b})$, and $\ominus \alpha \otimes \mathbf{a}$ instead of $\ominus(\alpha \otimes \mathbf{a})$ for any $\alpha, \beta \in \mathbb{R}$, $\mathbf{a}, \mathbf{b} \in G$.

3. DEFINITION OF A GENERALIZED GYROVECTOR SPACE REVISITED

We recall the definition of a generalized gyrovector space (cf. [2]).

Definition 3.1 (A generalized gyrovector space). *Let (G, \oplus) be a gyrocommutative gyrogroup with the map $\otimes : \mathbb{R} \times G \rightarrow G$. We denote \mathbf{e} the unit of (G, \oplus) . Let ϕ be an injection from G into a real normed space $(\mathbb{V}, \|\cdot\|)$. We say that $(G, \oplus, \otimes, \phi)$ (or (G, \oplus, \otimes) just for a simple notation) is a generalized gyrovector space or a GGV in short if the following conditions (GGV0) to (GGV8) are fulfilled:*

- (GGV0) $\|\phi(\text{gyr}[\mathbf{u}, \mathbf{v}]\mathbf{a})\| = \|\phi(\mathbf{a})\|$ for any $\mathbf{u}, \mathbf{v}, \mathbf{a} \in G$;
- (GGV1) $1 \otimes \mathbf{a} = \mathbf{a}$ for any $\mathbf{a} \in G$;
- (GGV2) $(r_1 + r_2) \otimes \mathbf{a} = (r_1 \otimes \mathbf{a}) \oplus (r_2 \otimes \mathbf{a})$ for any $\mathbf{a} \in G, r_1, r_2 \in \mathbb{R}$;
- (GGV3) $(r_1 r_2) \otimes \mathbf{a} = r_1 \otimes (r_2 \otimes \mathbf{a})$ for any $\mathbf{a} \in G, r_1, r_2 \in \mathbb{R}$;
- (GGV4) $\phi(|r| \otimes \mathbf{a}) / \|\phi(r \otimes \mathbf{a})\| = \phi(\mathbf{a}) / \|\phi(\mathbf{a})\|$ for any $\mathbf{a} \in G \setminus \{\mathbf{e}\}, r \in \mathbb{R} \setminus \{0\}$, where \mathbf{e} denotes the identity element of the gyrogroup (G, \oplus) ;
- (GGV5) $\text{gyr}[\mathbf{u}, \mathbf{v}](r \otimes \mathbf{a}) = r \otimes \text{gyr}[\mathbf{u}, \mathbf{v}]\mathbf{a}$ for any $\mathbf{u}, \mathbf{v}, \mathbf{a} \in G, r \in \mathbb{R}$;
- (GGV6) $\text{gyr}[r_1 \otimes \mathbf{v}, r_2 \otimes \mathbf{v}] = \text{id}_G$ for any $\mathbf{v} \in G, r_1, r_2 \in \mathbb{R}$, where id_G is the identity map on G ;
- (GGVV) $\|\phi(G)\| = \{\pm \|\phi(\mathbf{a})\| \in \mathbb{R} : \mathbf{a} \in G\}$ is a real linear space of dimension one with vector addition \oplus' and scalar multiplication \otimes' .
- (GGV7) $\|\phi(r \otimes \mathbf{a})\| = |r| \otimes' \|\phi(\mathbf{a})\|$ for any $\mathbf{a} \in G, r \in \mathbb{R}$;
- (GGV8) $\|\phi(\mathbf{a} \oplus \mathbf{b})\| \leq \|\phi(\mathbf{a})\| \oplus' \|\phi(\mathbf{b})\|$ for any $\mathbf{a}, \mathbf{b} \in G$.

Note that the terms in the equations in (GGV4) is well defined in the sense that $\|\phi(\mathbf{a})\| \neq 0$ and $\|\phi(r \otimes \mathbf{a})\| \neq 0$ for every $\mathbf{a} \in G \setminus \{\mathbf{e}\}$ and $r \in \mathbb{R} \setminus \{0\}$, by Proposition 2.1 (vi).

Proposition 3.2. *Suppose that $(G, \oplus, \otimes, \phi)$ is a GGV. Then $\|\phi(\mathbf{a})\| = 0_{\|\phi(G)\|}$ if and only if $\mathbf{a} = \mathbf{e}$, where \mathbf{e} is the unit of G . We have $r \otimes' \|\phi(\mathbf{a})\| = s \otimes' \|\phi(\mathbf{a})\|$ if and only if $r = s$ or $\mathbf{a} = \mathbf{e}$.*

Proof. Suppose that $\mathbf{a} = \mathbf{e}$. Then by (i) of Proposition 2.1 we have $\|\phi(\mathbf{a})\| = 0_{\|\phi(G)\|}$.

Suppose conversely that $\|\phi(\mathbf{a})\| = 0_{\|\phi(G)\|}$. We prove $\mathbf{a} = \mathbf{e}$. Suppose not; $\mathbf{a} \neq \mathbf{e}$. Then we have by (GGV4) that

$$\frac{\phi(2 \otimes \mathbf{a})}{\|\phi(2 \otimes \mathbf{a})\|} = \frac{\phi(\mathbf{a})}{\|\phi(\mathbf{a})\|}.$$

On the other hand

$$\|\phi(2 \otimes \mathbf{a})\| = 2 \otimes' \|\phi(\mathbf{a})\| = 2 \otimes' 0_{\|\phi(G)\|} = 0_{\|\phi(G)\|} = \|\phi(\mathbf{a})\|$$

ensure that

$$\phi(2 \otimes \mathbf{a}) = \phi(\mathbf{a}).$$

As ϕ is injective, we infer that $\mathbf{a} \oplus \mathbf{a} = 2 \otimes \mathbf{a} = \mathbf{a}$, so that $\mathbf{a} = \mathbf{e}$ by the left cancelation law, which is a contradiction since we assume that $\mathbf{a} \neq \mathbf{e}$. Thus, we have that $\mathbf{a} = \mathbf{e}$.

We prove the second statement. Suppose that $r = s$. Then $r \otimes' \|\phi(\mathbf{a})\| = s \otimes' \|\phi(\mathbf{a})\|$ follows. Suppose that $\mathbf{a} = \mathbf{e}$. Then $\|\phi(\mathbf{a})\| = 0_{\|\phi(G)\|}$, so that $r \otimes' \|\phi(\mathbf{a})\| = s \otimes' \|\phi(\mathbf{a})\|$ follows.

Suppose that $r \otimes' \|\phi(\mathbf{a})\| = s \otimes' \|\phi(\mathbf{a})\|$. Then $r = s$ or $\|\phi(\mathbf{a})\| = 0_{\|\phi(G)\|}$. It follows that $r = s$ or $\mathbf{a} = \mathbf{e}$ by the first part. \square

Example 3.3. Let $G = (-\infty, -1) \cup [1, \infty)$. Put a function $\Phi: \mathbb{R} \rightarrow G$ defined by

$$\Phi(x) = \begin{cases} \exp x, & x \geq 0 \\ -\exp(-x), & x < 0. \end{cases}$$

Then Φ is a bijection. Define

$$\mathbf{a} \oplus \mathbf{b} = \Phi(\Phi^{-1}(\mathbf{a}) + \Phi^{-1}(\mathbf{b})), \quad \mathbf{a}, \mathbf{b} \in G.$$

Then the additive group structure of \mathbb{R} is transplanted to G through Φ . Hence (G, \oplus) is a commutative group. Thus (G, \oplus) is a gyrocommutative gyrogroup such that the gyration $\text{gyr}[\mathbf{a}, \mathbf{b}]$ is the identity for all $\mathbf{a}, \mathbf{b} \in G$. The map $\otimes: \mathbb{R} \times G \rightarrow G$ is defined by

$$r \otimes \mathbf{a} = \Phi(r\Phi^{-1}(\mathbf{a})), \quad r \in \mathbb{R}, \mathbf{a} \in G.$$

It is easy to see that (GGV1), (GGV2), (GGV3), (GGV5) and (GGV6) hold. Let $(\mathbb{V}, \|\cdot\|) = (\mathbb{R}, |\cdot|)$. We define the map $\phi: G \rightarrow \mathbb{V}$ by

$$\phi(\mathbf{a}) = \mathbf{a}, \quad \mathbf{a} \in G.$$

Then $\|\phi(\mathbf{a})\| = |\mathbf{a}|$ for every $\mathbf{a} \in G$. It is clear that (GGV0) holds since $\text{gyr}[\cdot, \cdot]$ is the identity. By a simple calculation, we have $\|\phi(r \otimes \mathbf{a})\| = |\mathbf{a}|^{|r|}$ for all $\mathbf{a} \in G$ and $r \in \mathbb{R}$. We also obtain $\phi(|r| \otimes \mathbf{a}) = \mathbf{a}^{|r|}$ if $\mathbf{a} \geq 1$, $\phi(|r| \otimes \mathbf{a}) = -|\mathbf{a}|^{|r|}$ if $\mathbf{a} < -1$. Hence (GGV4) holds. Let

$$\|\phi(G)\|_+ = \{\|\phi(\mathbf{a})\| : \mathbf{a} \in G\},$$

$$\|\phi(G)\|_- = \{-\|\phi(\mathbf{a})\| : \mathbf{a} \in G\}.$$

We infer $\|\phi(G)\|_+ = [1, \infty)$, $\|\phi(G)\|_- = (-\infty, -1]$ and $\|\phi(G)\|_+ \cup \|\phi(G)\|_- = \|\phi(G)\|$. Suppose that $S: (-\infty, 0) \rightarrow (-\infty, -1]$ is a bijection. Let $T: \mathbb{R} \rightarrow \|\phi(G)\|$ be a bijection defined as

$$T(x) = \begin{cases} \exp x, & x \geq 0 \\ S(x), & x < 0. \end{cases}$$

Then, $T([0, \infty)) = \|\phi(G)\|_+$, $T((-\infty, 0)) = \|\phi(G)\|_-$. We define operations \oplus' and \otimes' on $\|\phi(G)\|$ as follows.

$$A \oplus' B = T(T^{-1}(A) + T^{-1}(B)), \quad A, B \in \|\phi(G)\|,$$

$$r \otimes' A = T(rT^{-1}(A)), \quad r \in \mathbb{R}, A \in \|\phi(G)\|.$$

Then $(\|\phi(G)\|, \oplus', \otimes')$ is a real-linear space of dimension one since the algebraic structure is simply transplanted from the usual real linear space \mathbb{R} of dimension one through the map T . By a simple calculation, we have

$$\|\phi(r \otimes \mathbf{a})\| = |\mathbf{a}|^{|r|} = |r| \otimes' \|\phi(\mathbf{a})\|, \quad r \in \mathbb{R}, \mathbf{a} \in G.$$

Thus (GGV7) holds. We have

$$\|\phi(\mathbf{a})\| \oplus' \|\phi(\mathbf{b})\| = T(T^{-1}(|\mathbf{a}|) + T^{-1}(|\mathbf{b}|)) = |\mathbf{a}\mathbf{b}|, \quad \mathbf{a}, \mathbf{b} \in G.$$

We also have

$$\|\phi(\mathbf{a} \oplus \mathbf{b})\| = |\mathbf{a} \oplus \mathbf{b}| = |\Phi(\Phi^{-1}(\mathbf{a}) + \Phi^{-1}(\mathbf{b}))|.$$

We also have $\Phi^{-1}(\mathbf{a}) = \log |\mathbf{a}|$ or $-\log |\mathbf{a}|$ and $\Phi^{-1}(\mathbf{b}) = \log |\mathbf{b}|$ or $-\log |\mathbf{b}|$. As $|\mathbf{a}| \geq 1$ and $|\mathbf{b}| \geq 1$, we infer that

$$-\log |\mathbf{a}| - \log |\mathbf{b}| \leq \Phi^{-1}(\mathbf{a}) + \Phi^{-1}(\mathbf{b}) \leq \log |\mathbf{a}| + \log |\mathbf{b}|.$$

As Φ is increasing, we see that

$$-|\mathbf{ab}| = \Phi(-\log|\mathbf{a}| - \log|\mathbf{b}|) \leq \Phi(\Phi^{-1}(\mathbf{a}) + \Phi^{-1}(\mathbf{b})) \leq \Phi(\log|\mathbf{a}| + \log|\mathbf{b}|) = |\mathbf{ab}|.$$

It follows that (GGV8) holds. We conclude that $(G, \oplus, \otimes, \phi)$ is a GGV, where $\mathbf{e} = 1$ and $0_{\|\phi(G)\|} = \|\phi(\mathbf{e})\| = 1$.

We recall the gyrometric for a GGV which is defined in [2, Definition 6].

Definition 3.4 (gyrometric). *Let $(G, \oplus, \otimes, \phi)$ be a GGV. Put*

$$\varrho(\mathbf{a}, \mathbf{b}) = \|\phi(\mathbf{a} \ominus \mathbf{b})\|, \quad \mathbf{a}, \mathbf{b} \in G.$$

We call $\varrho(\cdot, \cdot)$ a gyrometric on $(G, \oplus, \otimes, \phi)$.

For the convenience of the readers, we state and prove some elementary properties of the gyrometric (as (4) and (8) in [2]).

Proposition 3.5.

$$\varrho(\mathbf{x} \oplus \mathbf{a}, \mathbf{x} \oplus \mathbf{b}) = \varrho(\mathbf{a}, \mathbf{b}) = \varrho(\ominus \mathbf{a}, \ominus \mathbf{b}) = \varrho(\mathbf{b}, \mathbf{a}), \quad \mathbf{x}, \mathbf{a}, \mathbf{b} \in G.$$

Proof. By [6, Theorem 3.13] we have $(\mathbf{x} \oplus \mathbf{a}) \ominus (\mathbf{x} \oplus \mathbf{b}) = \text{gyr}[\mathbf{x}, \mathbf{a}](\mathbf{a} \ominus \mathbf{b})$, hence

$$\varrho(\mathbf{x} \oplus \mathbf{a}, \mathbf{x} \oplus \mathbf{b}) = \|\phi((\mathbf{x} \oplus \mathbf{a}) \ominus (\mathbf{x} \oplus \mathbf{b}))\| = \|\phi(\text{gyr}[\mathbf{x}, \mathbf{a}](\mathbf{a} \ominus \mathbf{b}))\|,$$

by (GGV0)

$$= \|\phi(\mathbf{a} \ominus \mathbf{b})\|.$$

As G is gyrocommutative, we have

$$\varrho(\mathbf{a}, \mathbf{b}) = \|\phi(\mathbf{a} \ominus \mathbf{b})\| = \|\phi(\text{gyr}[\mathbf{a}, \ominus \mathbf{b}](\ominus \mathbf{b} \oplus \mathbf{a}))\|,$$

then by (GGV0),

$$= \|\phi(\ominus \mathbf{b} \oplus \mathbf{a})\| = \varrho(\ominus \mathbf{b}, \ominus \mathbf{a}).$$

We also have by (GGV7) and Proposition 2.1 (v) that

$$\varrho(\mathbf{a}, \mathbf{b}) = \|\phi(\mathbf{a} \ominus \mathbf{b})\| = \|\phi((-1) \otimes (\mathbf{a} \ominus \mathbf{b}))\| = \|\phi(\ominus(\mathbf{a} \ominus \mathbf{b}))\|,$$

then by the gyroautomatic inverse property [6, Theorem 3.2], and the gyrocommutativity, and (GGV0)

$$= \|\phi(\ominus \mathbf{a} \oplus \mathbf{b})\| = \|\phi(\text{gyr}[\ominus \mathbf{a}, \mathbf{b}](\mathbf{b} \ominus \mathbf{a}))\| = \|\phi(\mathbf{b} \ominus \mathbf{a})\| = \varrho(\mathbf{b}, \mathbf{a}).$$

□

Proposition 3.6.

$$(3.1) \quad \varrho(\mathbf{a}, \mathbf{b}) \leq \varrho(\mathbf{a}, \mathbf{c}) \oplus' \varrho(\mathbf{c}, \mathbf{b}), \quad \mathbf{a}, \mathbf{b}, \mathbf{c} \in G.$$

Proof. By Proposition 2.1 (v), Proposition 3.5, (GGV7), and (GGV8), we have

$$\begin{aligned} \varrho(\mathbf{a}, \mathbf{b}) &= \varrho(\ominus \mathbf{a}, \ominus \mathbf{b}) = \varrho(\mathbf{c} \ominus \mathbf{a}, \mathbf{c} \ominus \mathbf{b}) = \|\phi((\mathbf{c} \ominus \mathbf{a}) \ominus (\mathbf{c} \ominus \mathbf{b}))\| \\ &= \|\phi((\mathbf{c} \ominus \mathbf{a}) \oplus (\ominus(\mathbf{c} \ominus \mathbf{b})))\| \leq \|\phi(\mathbf{c} \ominus \mathbf{a})\| \oplus' \|\phi(\ominus(\mathbf{c} \ominus \mathbf{b}))\| \\ &= \|\phi(\mathbf{c} \ominus \mathbf{a})\| \oplus' (|-1| \otimes' \|\phi(\mathbf{c} \ominus \mathbf{b})\|) = \|\phi(\mathbf{c} \ominus \mathbf{a})\| \oplus' \|\phi(\mathbf{c} \ominus \mathbf{b})\| \\ &= \varrho(\mathbf{c}, \mathbf{a}) \oplus' \varrho(\mathbf{c}, \mathbf{b}) = \varrho(\mathbf{a}, \mathbf{c}) \oplus' \varrho(\mathbf{c}, \mathbf{b}). \end{aligned}$$

□

We call the equation (3.1) the gyrotriangle inequality for a GGV.

Definition 3.7 (gyrometric preserving map). *Suppose that $(G_1, \oplus_1, \otimes_1)$ and $(G_2, \oplus_2, \otimes_2)$ be GGVs. Let ϱ_1 and ϱ_2 be gyrometrics of G_1 and G_2 , respectively. We say that a map $T : G_1 \rightarrow G_2$ is gyrometric preserving if the equality*

$$\varrho_2(T\mathbf{a}, T\mathbf{b}) = \varrho_1(\mathbf{a}, \mathbf{b})$$

holds for every pair $\mathbf{a}, \mathbf{b} \in G_1$.

4. A MAZUR-ULAM THEOREM FOR GGVs

The section presents a theorem stating that a surjective gyrometric preserving map preserves gyromidpoints. Gyromidpoints are defined in Definition 4.2. This theorem and its corollary generalize the Mazur-Ulam theorem for real normed spaces.

Definition 4.1 (gyrogroup coaddition). *Let (G, \oplus) be a gyrogroup. The gyrogroup coaddition \boxplus is defined by*

$$\mathbf{a} \boxplus \mathbf{b} = \mathbf{a} \oplus \text{gyr}[\mathbf{a}, \ominus \mathbf{b}]\mathbf{b}, \quad \mathbf{a}, \mathbf{b} \in G.$$

It is known that the gyrogroup coaddition \boxplus is commutative if and only if the gyrogroup is gyrocommutative [6, Theorem 3.4].

Definition 4.2 (gyromidpoint). *Let (G, \oplus, \otimes) is a GGV. The gyromidpoint $\mathbf{P}(\mathbf{a}, \mathbf{b})$ of $\mathbf{a}, \mathbf{b} \in (G, \oplus, \otimes)$ is defined as $\mathbf{P}(\mathbf{a}, \mathbf{b}) = \frac{1}{2} \otimes (\mathbf{a} \boxplus \mathbf{b})$, where \boxplus is the gyrogroup coaddition of the gyrogroup (G, \oplus) .*

Note that $\mathbf{P}(\mathbf{a}, \mathbf{b}) = \mathbf{P}(\mathbf{b}, \mathbf{a})$ as \boxplus is commutative. In addition, as in the same way as the proof of Theorem 6.34 in [6], we have

$$(4.1) \quad \mathbf{P}(\mathbf{a}, \mathbf{b}) = \mathbf{a} \oplus \frac{1}{2} \otimes (\ominus \mathbf{a} \oplus \mathbf{b})$$

(cf. [6, Definition 6.32 and Theorem 6.34]). In particular, $\mathbf{P}(\mathbf{a}, \mathbf{b}) = \frac{1}{2}(\mathbf{a} + \mathbf{b})$ if the GGV (G, \oplus, \otimes) is indeed a real normed space $(G, +, \cdot)$. The equation (9) of Proposition 15 in [2] is

$$(4.2) \quad \varrho(\mathbf{a}, \mathbf{P}(\mathbf{a}, \mathbf{b})) = \varrho(\mathbf{b}, \mathbf{P}(\mathbf{a}, \mathbf{b})) = \frac{1}{2} \otimes' \varrho(\mathbf{a}, \mathbf{b})$$

The following appears as Proposition 17 in [2]. We can similarly prove this as the proof of [6, Theorem 2.28].

Proposition 4.3. *Let (G_1, \oplus_1) and (G_2, \oplus_2) be gyrogroups. Suppose that $T : G_1 \rightarrow G_2$ is a bijection. Then, (I1) and (I2) are equivalent to each other.*

$$(I1): T(\mathbf{a} \oplus_1 \mathbf{b}) = T(\mathbf{a}) \oplus_2 T(\mathbf{b}) \text{ for any } \mathbf{a}, \mathbf{b} \in G_1,$$

$$(I2): T(\mathbf{a} \boxplus_1 \mathbf{b}) = T(\mathbf{a}) \boxplus_2 T(\mathbf{b}) \text{ for any } \mathbf{a}, \mathbf{b} \in G_1.$$

The following theorem and its corollary is a Mazur-Ulam theorem for a GGV.

Theorem 4.4 (Theorem 13 in [2]). *Let $(G_1, \oplus_1, \otimes_1)$ and $(G_2, \oplus_2, \otimes_2)$ be GGVs. Let ϱ_1 and ϱ_2 be gyrometrics of G_1 and G_2 , respectively. Suppose that a surjection $T : G_1 \rightarrow G_2$ satisfies*

$$\varrho_2(T\mathbf{a}, T\mathbf{b}) = \varrho_1(\mathbf{a}, \mathbf{b})$$

for any pair $\mathbf{a}, \mathbf{b} \in G_1$. Then T preserves the gyromidpoints:

$$P(T\mathbf{a}, T\mathbf{b}) = T(P(\mathbf{a}, \mathbf{b}))$$

for any pair $\mathbf{a}, \mathbf{b} \in G_1$.

The next corollary states that a surjective map that preserves the gyrometric structure also preserves the algebraic structure modulo the left gyrotranslations. This implies that any two GGVs with the same gyrometric structure also have the same GGV structure.

Corollary 4.5 (Corollary 14 in [2]). *Let $(G_1, \oplus_1, \otimes_1)$ and $(G_2, \oplus_2, \otimes_2)$ be GGVs. Let ϱ_1 and ϱ_2 be gyrometrics of G_1 and G_2 , respectively. Suppose that a surjection $T : G_1 \rightarrow G_2$ satisfies*

$$\varrho_2(T\mathbf{a}, T\mathbf{b}) = \varrho_1(\mathbf{a}, \mathbf{b})$$

for any pair $\mathbf{a}, \mathbf{b} \in G_1$. Then T is of the form $T = T(\mathbf{e}) \oplus_2 T_0$, where T_0 is an isometrical isomorphism in the sense that it is a bijection such that the equalities

$$(4.3) \quad T_0(\mathbf{a} \oplus_1 \mathbf{b}) = T_0(\mathbf{a}) \oplus_2 T_0(\mathbf{b});$$

$$(4.4) \quad T_0(\alpha \otimes_1 \mathbf{a}) = \alpha \otimes_2 T_0(\mathbf{a});$$

$$(4.5) \quad \varrho_2(T_0(\mathbf{a}), T_0(\mathbf{b})) = \varrho_1(\mathbf{a}, \mathbf{b})$$

for all $\mathbf{a}, \mathbf{b} \in G_1$ and $\alpha \in \mathbb{R}$ hold.

5. PREPARATIONS OF THE PROOF

Let $(G, \oplus, \otimes, \phi)$ be a GGV. Suppose that $\mathbf{a} \in G$ and $0 \leq \alpha \leq \beta$. Then

$$(5.1) \quad \begin{aligned} \alpha \otimes' \|\phi(\mathbf{a})\| &= \|\phi(\alpha \otimes \mathbf{a})\| \\ &= \left\| \phi \left(\left(\frac{\beta + \alpha}{2} - \frac{\beta - \alpha}{2} \right) \otimes \mathbf{a} \right) \right\| \\ &= \left\| \phi \left(\left(\frac{\beta + \alpha}{2} \right) \otimes \mathbf{a} \oplus \left(-\frac{\beta - \alpha}{2} \right) \otimes \mathbf{a} \right) \right\| \\ &\leq \left(\frac{\beta + \alpha}{2} \right) \otimes' \|\phi(\mathbf{a})\| \oplus' \left(\frac{\beta - \alpha}{2} \right) \otimes' \|\phi(\mathbf{a})\| \\ &= \left(\frac{\beta + \alpha}{2} + \frac{\beta - \alpha}{2} \right) \otimes' \|\phi(\mathbf{a})\| \\ &= \beta \otimes' \|\phi(\mathbf{a})\| \end{aligned}$$

If $\alpha \geq 0$, then $\alpha \otimes' \|\phi(\mathbf{a})\| = \|\phi(\alpha \otimes \mathbf{a})\| \geq 0$. By (5.1) and Propositions 2.1 (vi) and 3.2 we have that $0 \leq \alpha < \beta$ implies that

$$0 \leq \|\phi(\alpha \otimes \mathbf{a})\| = \alpha \otimes' \|\phi(\mathbf{a})\| < \beta \otimes' \|\phi(\mathbf{a})\|$$

for any $\mathbf{a} \in G \setminus \{\mathbf{e}\}$. Suppose conversely that

$$0 \leq \alpha \otimes' \|\phi(\mathbf{a})\| < \beta \otimes' \|\phi(\mathbf{a})\|.$$

for some $\alpha \geq 0, \beta \geq 0$ and $\mathbf{a} \in G \setminus \{\mathbf{e}\}$. We prove $\alpha < \beta$. Suppose not: $\alpha \geq \beta$. By (5.1) we have $\beta \otimes' \|\phi(\mathbf{a})\| \leq \alpha \otimes' \|\phi(\mathbf{a})\|$, which is a contradiction proving $\alpha < \beta$. Therefore, we have

$$(5.2) \quad 0 \leq \alpha < \beta \iff 0 \leq \alpha \otimes' \|\phi(\mathbf{a})\| < \beta \otimes' \|\phi(\mathbf{a})\|$$

for any $\alpha, \beta \geq 0$ and $\mathbf{a} \in G \setminus \{\mathbf{e}\}$. (Note that equation (5.2) is a corrected version of (10) in [2], which was partly incorrect. In fact, if $(G, \oplus, \otimes, \phi)$ is a GGV such that $\|\phi(\mathbf{e})\| \neq 0$ (cf. Example 3.3), then $0 < \|\phi(\mathbf{e})\| = 0_{\|\phi(G)\|} = 0 \otimes' \|\phi(\mathbf{a})\|$ for $\mathbf{a} \in G \setminus \{\mathbf{e}\}$ by Proposition 2.1. Proposition 18 in [2] was proved by an application of the statement (10). We revise it as Proposition 5.1. Note also that $\|\phi(\mathbf{e})\| = 0$ which appears on line 6 on page 398 of [2] is a typo, it should be read as $\|\phi(\mathbf{e})\| = 0_{\|\phi(G)\|}$.) The Proposition 18 mentioned in [2] has been revised and corrected as follows.

Proposition 5.1. *Let $(G, \oplus, \otimes, \phi)$ be a GGV. Then there exists a bijection $f : \|\phi(G)\| \rightarrow \mathbb{R}$ that satisfies the following conditions;*

- (F1) $f(A \oplus' B) = f(A) + f(B)$ and $f(r \otimes' A) = rf(A)$ for any $A, B \in \|\phi(G)\|$, $r \in \mathbb{R}$
- (F2) $0 \leq A < B$ if and only if $0 \leq f(A) < f(B)$ for $A, B \in \|\phi(G)\|$ with $A \geq 0$, $B \geq 0$.

Proof. By the condition (GGVV), there exists a bijection $f : \|\phi(G)\| \rightarrow \mathbb{R}$ that satisfies the condition (F1). Needless to say, $-f$ also satisfies the condition (F1). Hence, we may assume that $f(\|\phi(\mathbf{x}_0)\|) > 0$ for some $\mathbf{x}_0 \in G$.

For $0 \leq A, B \in \|\phi(G)\|$, put $X = \|\phi(\mathbf{x}_0)\|$ and $r_A = f(A)/f(X)$, $r_B = f(B)/f(X)$. Then $f(r_A \otimes' X) = r_A f(X) = f(A)$. As f is injective, we infer that $r_A \otimes' X = A$. Then we have $r_A \geq 0$. (Suppose not: $r_A < 0$. Since $(\|\phi(G)\|, \oplus', \otimes')$ is a linear space, $r_A \otimes' X = r_A \otimes' \|\phi(\mathbf{x}_0)\| = -\|\phi((-r_A) \otimes \mathbf{x}_0)\| \leq 0$ by (GGV7). If $\|\phi((-r_A) \otimes \mathbf{x}_0)\| = 0$, then $(-r_A) \otimes \mathbf{x}_0 = \mathbf{e}$ by Proposition 2.1 (vi), so $\mathbf{x}_0 = \mathbf{e}$ as $-r_A \neq 0$. Thus $\|\phi(\mathbf{x}_0)\| = 0_{\|\phi(G)\|}$ by Proposition 2.1 (i). Hence $f(\|\phi(\mathbf{x}_0)\|) = 0$ as $f : \|\phi(G)\| \rightarrow \mathbb{R}$ is linear. This contradicts to our choice of \mathbf{x}_0 ; $f(\|\phi(\mathbf{x}_0)\|) > 0$. We conclude that $-\|\phi((-r_A) \otimes \mathbf{x}_0)\| < 0$, so $A = r_A \otimes' X < 0$, which is a contradiction.) In the same way, we have that $r_B \geq 0$. Then by (5.2) we have

$$0 \leq r_A < r_B \iff 0 \leq A < B$$

since $A = r_A \otimes' X = r_A \otimes' \|\phi(\mathbf{x}_0)\|$ and $B = r_B \otimes' X = r_B \otimes' \|\phi(\mathbf{x}_0)\|$. On the other hand the equations $r_A f(X) = f(A)$ and $r_B f(X) = f(B)$ where $f(X) > 0$ ensures that

$$0 \leq r_A < r_B \iff 0 \leq f(A) < f(B).$$

Hence we conclude that

$$0 \leq A < B \iff 0 \leq f(A) < f(B)$$

for every $A, B \in \|\phi(G)\|$ with $A \geq 0$ and $B \geq 0$. □

Note that $f(A) = 0$ if and only if $A = 0_{\|\phi(\mathbf{e})\|}$ since f is a linear bijection from $\|\phi(G)\|$ onto \mathbb{R} . In this case, A need not be 0 (cf. Example 3.3). It means that $0 < A$ does not always imply $0 < f(A)$.

Proposition 5.2. *Let $(G, \oplus, \otimes, \phi)$ be a GGV. Suppose that $A, B \in \|\phi(G)\|$. Then $0 \leq A < B$ and $0 \leq A' < B'$ imply that $0 \leq A \oplus' A' < B \oplus' B'$.*

Proof. Suppose that $0 \leq A < B$ and $0 \leq A' < B'$. By Proposition 2.1 (F2) we have $0 \leq f(A) < f(B)$ and $0 \leq f(A') < f(B')$. By Proposition 2.1 (F1) we have

$$0 \leq f(A \oplus' A') = f(A) + f(A') < f(B) + f(B') = f(B \oplus' B').$$

Then Proposition 2.1 (F2) ensures $0 \leq A \oplus' A' < B \oplus' B'$. □

Note that the gyrometric on a GGV $(G, \oplus, \otimes, \phi)$ may not be a metric on G . This means that $\varrho(\mathbf{a}, \mathbf{a}) = \|\phi(\mathbf{e})\|$ may not be equal to 0, as shown in Example 3.3. However, it is significant to observe that $f \circ \varrho$ is always a metric on G . Here, f is the linear map described in Proposition 5.1.

6. PROOF OF THEOREM 4.4

In this section, we present a simple proof of Theorem 4.4 by utilizing the defect caused by Nica [4]. We recall Proposition 16 in [2].

Proposition 6.1 (Proposition 16 in [2]). *Let (G, \oplus, \otimes) be a GGV. For $\mathbf{a} \in G$. Let $\varphi: G \rightarrow G$ be defined by $\varphi(\mathbf{x}) = 2 \otimes \mathbf{a} \ominus \mathbf{x}$, $\mathbf{x} \in G$. Then φ is a bijection that satisfies the following.*

- (p1) $\varphi \circ \varphi$ is the identity map on G ;
- (p2) $\varrho(\varphi(\mathbf{x}), \varphi(\mathbf{y})) = \varrho(\mathbf{x}, \mathbf{y})$, $\mathbf{x}, \mathbf{y} \in G$;
- (p3) $\varphi(\mathbf{x}) = \mathbf{x}$ if and only if $\mathbf{x} = \mathbf{a}$;
- (p4) $\varphi(\mathbf{x}) = \mathbf{y}$ and $\varphi(\mathbf{y}) = \mathbf{x}$ if $\mathbf{a} = \mathbf{P}(\mathbf{x}, \mathbf{y})$;
- (p5) $\varrho(\varphi(\mathbf{x}), \mathbf{x}) = 2 \otimes' \varrho(\mathbf{a}, \mathbf{x})$

for all $\mathbf{x}, \mathbf{y} \in G$.

Proof of Theorem 4.4. First, consider the case where G_1 and G_2 are the same GGV. For the first part of the proof, we denote (G, \oplus, \otimes) instead of $(G_i, \oplus_i, \otimes_i)$ to simplify the notations. Let f be the corresponding map for G , which appears in Proposition 5.1. Let $\mathbf{x}_1, \mathbf{x}_2 \in G$ be arbitrary. Put $p = \mathbf{P}(\mathbf{x}_1, \mathbf{x}_2)$, $p' = \mathbf{P}(T(\mathbf{x}_1), T(\mathbf{x}_2))$, and the defect: $d = f(\varrho(T(p), p'))$. We prove $d = 0$. Let $\varphi_1: G \rightarrow G$ be given by $\varphi_1(\mathbf{x}) = 2 \otimes p \ominus \mathbf{x}$ for $\mathbf{x} \in G$ and $\varphi_2: G \rightarrow G$ by $\varphi_2(\mathbf{z}) = 2 \otimes p' \ominus \mathbf{z}$ for $\mathbf{z} \in G$. Put $S = \varphi_1 \circ T^{-1} \circ \varphi_2 \circ T$. Then $S: G \rightarrow G$ is a surjective gyrometric preserving map such that $S(\mathbf{x}_1) = \mathbf{x}_1$ and $S(\mathbf{x}_2) = \mathbf{x}_2$ by Proposition 6.1 (p2) and (p4). We prove

$$(6.1) \quad f(\varrho(S^{2^n}(p), p)) = 2^{n+1}d$$

for every nonnegative integer n . First, we have by Proposition 6.1 (p1) and (p2) that

$$(6.2) \quad \varrho(\varphi_1 \circ S(\mathbf{x}), \mathbf{y}) = \varrho(T^{-1} \circ \varphi_2 \circ T(\mathbf{x}), \mathbf{y}) = \varrho(\mathbf{x}, T^{-1} \circ \varphi_2 \circ T(\mathbf{y})) = \varrho(\varphi_1(\mathbf{x}), S(\mathbf{y})), \quad \mathbf{x}, \mathbf{y} \in G.$$

As $\varphi_1(p) = p$ we have

$$\varrho(S(p), p) = \varrho(\varphi_1 \circ S(p), p) = \varrho(\varphi_2 \circ T(p), T(p)) = 2 \otimes' \varrho(T(p), p').$$

It follows that $f(\varrho(S(p), p)) = 2d$, which is (6.1) for $n = 0$. Suppose that (6.1) holds for n . Since $2 \otimes' \varrho(\mathbf{x}, p) = \varrho(\varphi_1(\mathbf{x}), \mathbf{x})$ for any $\mathbf{x} \in G_1$ by Proposition 6.1 (p5), we have

$$2^{n+2}d = f(2 \otimes' \varrho(S^{2^n}(p), p)) = f(\varrho(\varphi_1 \circ S^{2^n}(p), S^{2^n}(p))),$$

then applying (6.2) 2^n times

$$= f(\varrho(\varphi_1(p), S^{2^{n+1}}(p))) = f(\varrho(p, S^{2^{n+1}}(p))).$$

as $\varphi_1(p) = p$. Then, by induction, we obtain (6.1) for every nonnegative integer n . Since $S(\mathbf{x}_1) = \mathbf{x}_1$ we have by the gyrometric triangle inequality that

$$\varrho(S^{2^n}(p), p) \leq \varrho(S^{2^n}(p), S^{2^n}(\mathbf{x}_1)) \oplus' \varrho(S^{2^n}(\mathbf{x}_1), p) = \varrho(p, \mathbf{x}_1) \oplus' \varrho(\mathbf{x}_1, p)$$

for every nonnegative integer n . Thus

$$2^{n+1}d = f(\varrho(S^{2^n}(p), p)) \leq 2f(\varrho(\mathbf{x}_1, p))$$

for every nonnegative integer n . It follows that $f(\varrho(T(p), p')) = d = 0 = f(0_{\|\phi(G)\|})$. As f is a bijection, we have

$$\|\phi(T(p) \ominus p')\| = \varrho(T(p), p') = \|\phi(\mathbf{e})\| = 0_{\|\phi(G)\|}.$$

Then by Proposition 3.2 we see that $T(p) \ominus p' = \mathbf{e}$. Hence $T(p) = p'$.

We consider the general case. Suppose that $T: G_1 \rightarrow G_2$ satisfies

$$\varrho_2(T\mathbf{a}, T\mathbf{b}) = \varrho_1(\mathbf{a}, \mathbf{b})$$

for any pair $\mathbf{a}, \mathbf{b} \in G_1$. Let $\mathbf{x}_1, \mathbf{x}_2 \in G$ be arbitrary. Put $S = \varphi_1 \circ T^{-1} \circ \varphi_2 \circ T$, where $\varphi_1: G_1 \rightarrow G_1$ is given by $\varphi_1(\mathbf{x}) = 2 \otimes p \ominus \mathbf{x}$ for $\mathbf{x} \in G_1$ and $\varphi_2: G_2 \rightarrow G_2$ is given by $\varphi_2(\mathbf{z}) = 2 \otimes_2 p' \ominus \mathbf{z}$ for $\mathbf{z} \in G_2$, where $p = \mathbf{P}(\mathbf{x}_1, \mathbf{x}_2)$, $p' = \mathbf{P}(T(\mathbf{x}_1), T(\mathbf{x}_2))$. Then $S: G_1 \rightarrow G_1$ is a surjective gyrometric preserving map from G_1 onto itself. By the first part, we obtain that $S(\mathbf{P}(\mathbf{x}_1, \mathbf{x}_2)) = \mathbf{P}(S(\mathbf{x}_1), S(\mathbf{x}_2))$ for every $\mathbf{x}_1, \mathbf{x}_2 \in G_1$. As $S(\mathbf{x}_1) = \mathbf{x}_1$ and $S(\mathbf{x}_2) = \mathbf{x}_2$, we have $S(p) = p$. By this equation, we have $\varphi_2 \circ T(p) = T(p)$ since $\varphi_1(p) = p$. As $\varphi_2(\mathbf{y}) = \mathbf{y}$ if and only if $\mathbf{y} = p'$, we obtain $T(p) = p'$. □

7. PROOF OF COROLLARY 4.5

We prove that a bijective gyrometric preserving map essentially preserves the algebraic structure.

Proof of Corollary 4.2. If G_1 is a singleton, then G_2 is also a singleton. In this case, every statement is clear. In the following, we assume G_1 is not a singleton. Then G_2 is not a singleton as well since T is a bijection. Let $T_0 = \ominus_2 T(\mathbf{e}_1) \oplus_2 T$. Indeed, $T_0: G_1 \rightarrow G_2$ is surjective and $T_0(\mathbf{e}_1) = \mathbf{e}_2$. By the left cancellation law, we have $T = T(\mathbf{e}_1) \oplus_2 T_0$. By Proposition 3.5, T_0 is a gyrometric preserving map because so is T . Applying Theorem 4.4 to T_0 , we have

$$(7.1) \quad T_0\left(\frac{1}{2} \otimes_1 (\mathbf{a} \boxplus_1 \mathbf{b})\right) = \frac{1}{2} \otimes_2 (T_0(\mathbf{a}) \boxplus_2 T_0(\mathbf{b}))$$

for any $\mathbf{a}, \mathbf{b} \in G$. Since $T_0(\mathbf{e}_1) = \mathbf{e}_2$, we have

$$(7.2) \quad \begin{aligned} T_0\left(\frac{1}{2} \otimes_1 \mathbf{x}\right) &= T_0\left(\frac{1}{2} \otimes_1 (\mathbf{x} \boxplus_1 \mathbf{e}_1)\right) \\ &= \frac{1}{2} \otimes_2 (T_0(\mathbf{x}) \boxplus_2 T_0(\mathbf{e}_1)) \\ &= \frac{1}{2} \otimes_2 T_0(\mathbf{x}) \end{aligned}$$

for any $\mathbf{x} \in G_1$. It follows by equations (7.1) and (7.2) that

$$(7.3) \quad T_0(\mathbf{a} \boxplus_1 \mathbf{b}) = T_0(\mathbf{a}) \boxplus_2 T_0(\mathbf{b})$$

for any $\mathbf{a}, \mathbf{b} \in G_1$. Since T_0 is bijective, we have the equation (4.3) by Proposition 4.3.

Next we show that T_0 is homogeneous: $T_0(\alpha \otimes_1 \mathbf{a}) = \alpha \otimes_2 T_0(\mathbf{a})$ for any $\mathbf{a} \in G_1$ and $\alpha \in \mathbb{R}$. First we consider $\alpha = \frac{m}{2^n}$ for some integers m, n . For any $\mathbf{a} \in G_1$ and for any positive integer m , $T_0(m \otimes_1 \mathbf{a}) = m \otimes_2 T_0(\mathbf{a})$ is satisfied by the equation (4.3). We have $\mathbf{e}_2 = T_0(\mathbf{e}_1) = T_0((\mathbf{a} \ominus_1 \mathbf{a}) =$

$T_0(\mathbf{a}) \oplus_2 T_0(\Theta_1 \mathbf{a})$ for any $\mathbf{a} \in G_1$. Hence $T_0(\Theta_1 \mathbf{a}) = \ominus_2 T_0(\mathbf{a})$. It follows by a routine argument that $T_0(m \otimes_1 \mathbf{a}) = m \otimes_2 T_0(\mathbf{a})$ for every integer m and $\mathbf{a} \in G_1$. By the equation (7.2), we have

$$(7.4) \quad T_0\left(\frac{m}{2^n} \otimes_1 \mathbf{a}\right) = \frac{m}{2^n} \otimes_2 T_0(\mathbf{a})$$

for any integers m, n . Next, we consider a general $\alpha \in \mathbb{R}$. To prove the homogeneity of T_0 for a general α , we will show that

$$(7.5) \quad \varrho_2(T_0(\alpha \otimes_1 \mathbf{a}), \alpha \otimes_2 T_0(\mathbf{a})) < \|\phi_2(T_0(\mathbf{b}))\|_2$$

for every $\mathbf{b} \in T_0^{-1}(G_2 \setminus \{\mathbf{e}_2\})$. As

$$\varrho_2(T_0(\alpha \otimes_1 \mathbf{a}), \alpha \otimes_2 T_0(\mathbf{a})) = \|\phi_2(T_0(\alpha \otimes_1 \mathbf{a}) \ominus_2 \alpha \otimes_2 T_0(\mathbf{a}))\|_2$$

(7.5) will be followed by

$$T_0(\alpha \otimes_1 \mathbf{a}) \ominus_2 \alpha \otimes_2 T_0(\mathbf{a}) \neq T_0(\mathbf{b})$$

for every $\mathbf{b} \in T_0^{-1}(G_2 \setminus \{\mathbf{e}_2\})$. As T_0 is surjective we will arrive at

$$T_0(\alpha \otimes_1 \mathbf{a}) \ominus_2 \alpha \otimes_2 T_0(\mathbf{a}) = \mathbf{e}_2,$$

so that

$$T_0(\alpha \otimes_1 \mathbf{a}) = \alpha \otimes_2 T_0(\mathbf{a}),$$

which is the destined equality. We now prove (7.5). Let $\{r_k\}$ be a sequence of rational numbers with the forms $\frac{m}{2^n}$ of integers m and n which converges to α . As $T_0(r_k \otimes_1 \mathbf{a}) = r_k \otimes_2 T_0(\mathbf{a})$ by (7.4) we have by the gyrotriangle inequality Proposition 3.6 that

$$(7.6) \quad \varrho_2(T_0(\alpha \otimes_1 \mathbf{a}), \alpha \otimes_2 T_0(\mathbf{a})) \leq \varrho_2(T_0(r_k \otimes_1 \mathbf{a}), T_0(\alpha \otimes_1 \mathbf{a})) \oplus'_2 \varrho_2(r_k \otimes_2 T_0(\mathbf{a}), \alpha \otimes_2 T_0(\mathbf{a})).$$

We estimate the terms on the right-hand side of the inequality. By Proposition 5.1, there is a bijection $f_i : \|\phi_i(G_i)\|_i \rightarrow \mathbb{R}$ which satisfies the conditions (F1) and (F2) ($i = 1, 2$). Let $\mathbf{b} \in T_0^{-1}(G_2 \setminus \{\mathbf{e}_2\})$ be arbitrarily. Note that $T_0^{-1}(G_2 \setminus \{\mathbf{e}_2\}) \neq \emptyset$ since T_0 is a surjection and G_2 is not a singleton. Then $\mathbf{b} \neq \mathbf{e}_1$ since $T_0(\mathbf{e}_1) = \mathbf{e}_2$. Hence $\|\phi_1(\mathbf{b})\|_1 > 0$ by Proposition 2.1 (vi). Then $f_1(\|\phi_1(\mathbf{b})\|_1) \geq 0$ by Proposition 5.1. In fact, $f_1(\|\phi_1(\mathbf{b})\|_1) > 0$. (Suppose that $f_1(\|\phi_1(\mathbf{b})\|_1) = 0$. Then $\|\phi_1(\mathbf{b})\|_1 = 0_{\|\phi_1(G_1)\|_1}$ since f_1 is a linear bijection from $\|\phi_1(G_1)\|_1$ onto \mathbb{R} . By Proposition 3.2 we have $\mathbf{b} = \mathbf{e}_1$, which is a contradiction.) As $T_0(\mathbf{b}) \neq \mathbf{e}_2$, we also see that $f_2(\|\phi_2(T_0(\mathbf{b}))\|_2) > 0$. There is a positive integer k_0 such that $k > k_0$ implies that

$$2|r_k - \alpha|f_1(\|\phi_1(\mathbf{a})\|_1) < f_1(\|\phi_1(\mathbf{b})\|_1)$$

and

$$2|r_k - \alpha|f_2(\|\phi_2(T_0(\mathbf{a}))\|_2) < f_2(\|\phi_2(T_0(\mathbf{b}))\|_2).$$

As f_1 and f_2 are linear we infer by (GGV7) that

$$(7.7) \quad 0 \leq f_1(\|\phi_1((r_k - \alpha) \otimes_1 \mathbf{a})\|_1) < f_1\left(\frac{1}{2} \otimes'_1 \|\phi_1(\mathbf{b})\|_1\right)$$

and

$$(7.8) \quad 0 \leq f_2(\|\phi_2((r_k - \alpha) \otimes_2 T_0(\mathbf{a}))\|_2) < f_2\left(\frac{1}{2} \otimes'_2 \|\phi_2(T_0(\mathbf{b}))\|_2\right).$$

On the other hand, we have

$$\begin{aligned}
(7.9) \quad \frac{1}{2} \otimes'_1 \|\phi_1(\mathbf{b})\|_1 &= \|\phi_1(\frac{1}{2} \otimes_1 \mathbf{b} \ominus \mathbf{e}_1)\|_1 = \rho_1(\frac{1}{2} \otimes_1 \mathbf{b}, \mathbf{e}_1) \\
&= \varrho_2(T_0(\frac{1}{2} \otimes_1 \mathbf{b}), T_0(\mathbf{e}_1)) = \varrho_2(\frac{1}{2} \otimes_2 T_0(\mathbf{b}), \mathbf{e}_2) \\
&= \|\phi_2(\frac{1}{2} \otimes_2 T_0(\mathbf{b}))\|_2 = \frac{1}{2} \otimes'_2 \|\phi_2(T_0(\mathbf{b}))\|_2.
\end{aligned}$$

Applying Proposition 5.1 (F2) for the inequalities (7.7) and (7.8) we have by (7.9) that

$$(7.10) \quad \|\phi_1((r_k - \alpha) \otimes_1 \mathbf{a})\|_1 < \frac{1}{2} \otimes'_1 \|\phi_1(\mathbf{b})\|_1 = \frac{1}{2} \otimes'_2 \|\phi_2(T_0(\mathbf{b}))\|_2,$$

$$(7.11) \quad \|\phi_2((r_k - \alpha) \otimes_2 T_0(\mathbf{a}))\|_2 < \frac{1}{2} \otimes'_2 \|\phi_2(T_0(\mathbf{b}))\|_2.$$

Applying (7.10) and (7.11) we compute

$$\begin{aligned}
(7.12) \quad \varrho_2(T_0(r_k \otimes_1 \mathbf{a}), T_0(\alpha \otimes_1 \mathbf{a})) &= \varrho_1(r_k \otimes_1 \mathbf{a}, \alpha \otimes_1 \mathbf{a}) \\
&= \|\phi_1(r_k \otimes_1 \mathbf{a} \ominus \alpha \otimes_1 \mathbf{a})\|_1 = \|\phi_1((r_k - \alpha) \otimes_1 \mathbf{a})\|_1 < \frac{1}{2} \otimes'_2 \|\phi_2(T_0(\mathbf{b}))\|_2.
\end{aligned}$$

$$\begin{aligned}
(7.13) \quad \varrho_2(r_k \otimes_2 T_0(\mathbf{a}), \alpha \otimes_2 T_0(\mathbf{a})) &= \|\phi_2(r_k \otimes_2 T_0(\mathbf{a}) \ominus \alpha \otimes_2 T_0(\mathbf{a}))\|_2 \\
&= \|\phi_2((r_k - \alpha) \otimes_2 T_0(\mathbf{a}))\|_2 < \frac{1}{2} \otimes'_2 \|\phi_2(T_0(\mathbf{b}))\|_2
\end{aligned}$$

Applying Proposition 5.2 for (7.12) and (7.13) we have

$$\begin{aligned}
(7.14) \quad \varrho_2(T_0(r_k \otimes_1 \mathbf{a}), T_0(\alpha \otimes_1 \mathbf{a})) \oplus'_2 \varrho_2(r_k \otimes_2 T_0(\mathbf{a}), \alpha \otimes_2 T_0(\mathbf{a})) \\
< \frac{1}{2} \otimes'_2 \|\phi_2(T_0(\mathbf{b}))\|_2 \oplus'_2 \frac{1}{2} \otimes'_2 \|\phi_2(T_0(\mathbf{b}))\|_2 = \|\phi_2(T_0(\mathbf{b}))\|_2.
\end{aligned}$$

Combining (7.6) and (7.14), we obtain (7.5). Therefore we have $T_0(\alpha \otimes_1 \mathbf{a}) = \alpha \otimes_2 T_0(\mathbf{a})$ for every $\mathbf{a} \in G_1$ and $\alpha \in \mathbb{R}$. Proposition 3.5 observes that

$$\begin{aligned}
\varrho_2(T_0(\mathbf{a}), T_0(\mathbf{b})) &= \varrho_2(T(\mathbf{e}) \oplus_2 T_0(\mathbf{a}), T(\mathbf{e}) \oplus_2 T_0(\mathbf{b})) \\
&= \varrho_2(T(\mathbf{a}), T(\mathbf{b})) = \varrho_1(\mathbf{a}, \mathbf{b}), \quad \mathbf{a}, \mathbf{b} \in G_1.
\end{aligned}$$

We prove that T_0 is an injection. Suppose that $T_0(\mathbf{a}) = T_0(\mathbf{b})$ for $\mathbf{a}, \mathbf{b} \in G_1$. Then

$$(7.15) \quad \|\phi_1(\mathbf{a} \ominus \mathbf{b})\|_1 = \|\phi_2(T_0(\mathbf{a}) \ominus T_0(\mathbf{b}))\|_2 = \|\phi_2(\mathbf{e}_2)\|_2 = \|\phi_2(T_0(\mathbf{e}_1))\|_2.$$

In a way similar to (7.9) we see that $\|\phi_2(T_0(\mathbf{e}_1))\|_2 = \|\phi_1(\mathbf{e}_1)\|_1$. Since $\|\phi_1(\mathbf{e}_1)\|_1 = 0_{\|\phi_1(G_1)\|}$, we have by (7.15) that $\|\phi_1(\mathbf{a} \ominus \mathbf{b})\|_1 = 0_{\|\phi_1(G_1)\|}$. Then by Proposition 3.2 we infer that $\mathbf{a} \ominus \mathbf{b} = \mathbf{e}_1$, so $\mathbf{a} = \mathbf{b}$, concluding that T_0 is an injection. Thus T_0 is a bijection. It is easy to see by the left cancelation law that $T = T(\mathbf{e}_1) \oplus_2 T_0$ is an injection. \square

Acknowledgments. The first author was supported by JSPS KAKENHI Grant Numbers JP22K03332. The second author was supported by JSPS KAKENHI Grant Numbers JP19K03536.

REFERENCES

- [1] T. Abe, *Normed gyrolinear spaces: a generalization of normed spaces based on gyrocommutative gyrogroups*, Mathematics Interdisciplinary Research **1** (2016), 143–172
- [2] T. Abe and O. Hatori, *Generalized Gyrovector Spaces and a Mazur-Ulam theorem*, Publ. Math. Debrecen **87** (2015), 393–413; *A Note on the Proof of Theorem 13 in the Paper "Generalized Gyrovector Spaces and a Mazur-Ulam theorem"*, arXiv:1511.05187v1
- [3] O. Hatori, *Examples and applications of generalized gyrovector spaces*, Result. Math. **71**(2017), 295–317
- [4] B. Nica, *The Masur-Ulam theorem*, Expo. Math. **30** (2012), 397–398
- [5] A. A. Ungar, *Analytic Hyperbolic Geometry: Mathematical Foundations and Applications*, World Scientific, (2005)
- [6] A. A. Ungar, *Analytic Hyperbolic Geometry and Albert Einstein's Special Theory of Relativity*, World Scientific, (2008)

FACULTY OF ENGINEERING, IBARAKI UNIVERSITY, HITACHI 316-8511, JAPAN
Email address: toshikazu.abe.bin@vc.ibaraki.ac.jp

INSTITUTE OF SCIENCE AND TECHNOLOGY, NIIGATA UNIVERSITY, NIIGATA 950-2181, JAPAN.
Email address: hatori@math.sc.niigata-u.ac.jp