

QUANTUM GROUP DEFORMATIONS AND QUANTUM R -(CO)MATRICES VS. QUANTUM DUALITY PRINCIPLE

GASTÓN ANDRÉS GARCÍA[♭], FABIO GAVARINI[♯]

ABSTRACT. In this paper we describe the effect on quantum groups — namely, both QUEA’s and QFSHA’s — of deformations by twist and by 2-cocycles, showing how such deformations affect the semiclassical limit.

As a second, more important task, we discuss how these deformation procedures can be “stretched” to a new extent, via a formal variation of the original recipes, using *quasi-twists* and *quasi-2-cocycles*. These recipes seemingly should make no sense at all, yet we prove that they actually work, thus providing well-defined, more general deformation procedures. Later on, we explain the underlying reason that motivates such a result in light of the Quantum Duality Principle, through which every “quasi-twist/2-cocycle” for a given quantum group can be seen as a standard twist/2-cocycle for another quantum group, associated to the original one via the appropriate Drinfeld functor.

As a third task, we consider standard constructions involving R -(co)matrices in the general theory of Hopf algebras. First we adapt them to quantum groups, then we show that they extend to the case of *quasi- R -(co)matrices*, and finally we discuss how these constructions interact with the Quantum Duality Principle. As a byproduct, this yields new special symmetries (isomorphisms) for the underlying pair of dual Poisson (formal) groups that one gets by specialization.

CONTENTS

1. Introduction	2
2. Quantum groups, Quantum Duality Principle, and deformations	4
2.1. Lie bialgebras and Lie deformations	4
2.2. Hopf algebra deformations and R -(co)matrices	6
2.3. Quantum groups	10
2.4. The Quantum Duality Principle	13
3. Deformations of quantum groups	14
3.1. Deformations by twist of QUEA’s	14
3.2. Deformations by 2-cocycle of QFSHA’s	20
3.3. Deformations by quasi-2-cocycle of QUEA’s	28
3.4. Deformations by quasi-twist of QFSHA’s	42
3.5. Duality issues	54
4. Deformations vs. QDP	54
4.1. Some auxiliary results	55
4.2. Drinfeld’s functors and “quasi-deformations”	56
4.3. Drinfeld’s functors and (standard) deformations	57

2020 MSC: 17B37, 17B62 — *Keywords:* Quantum Groups, Quantum Enveloping Algebras.

5.	Morphisms in the “(co)quasitriangular” case	58
5.1.	R -matrices and ϱ -comatrices w.r.t. QDP: quasi-(co)matrices	59
5.2.	Morphisms from R -matrices and ϱ -comatrices	60
5.3.	Morphisms from quasi- R -matrices and quasi- ϱ -comatrices	66
5.4.	Semiclassical morphisms induced by specialization	70
	References	71

1. INTRODUCTION

In Hopf algebra theory, there exists a well-established theory of “deformations” that are produced via specific tools, namely *twists* in one case and *2-cocycles* in the other case (the terminology is not entirely agreed upon, yet our choice of terms seems to be the standard one among Hopf algebraists, at least). Given a Hopf algebra H , a twist for it is a suitable element $\mathcal{F} \in H \otimes H$, while (dually) a 2-cocycle is a suitable 2-form $\sigma \in (H \otimes H)^*$. Deformation by the twist \mathcal{F} provides H with a new Hopf algebra structure, by modifying the coproduct (and the antipode) but not the product, whereas deformation by the 2-cocycle σ endows H with yet another Hopf structure by changing the product alone (and the antipode) but not the coproduct.

Quantum groups are Hopf algebras of special type, which come in two versions: QUEAs (= quantized universal enveloping algebras) and QFSHAs (= quantized formal series Hopf algebras). Roughly speaking, a QUEA is a (topological) Hopf algebra U_{\hbar} over the \mathbb{k} -algebra of formal power series $\mathbb{k}[[\hbar]]$ such that $U_0 := U_{\hbar}/\hbar U_{\hbar}$ is isomorphic to $U(\mathfrak{g})$ for some Lie algebra \mathfrak{g} . It follows then that $U(\mathfrak{g})$ inherits from U_{\hbar} a Poisson cobracket, which makes it into a co-Poisson Hopf algebra, hence \mathfrak{g} bears a Lie cobracket making it into a Lie bialgebra. One then says that U_{\hbar} is a *quantization* of the co-Poisson Hopf algebra $U(\mathfrak{g})$, or just of the Lie bialgebra \mathfrak{g} . Dually, a QFSHA is a (topological) Hopf algebra F_{\hbar} over $\mathbb{k}[[\hbar]]$ such that $F_0 := F_{\hbar}/\hbar F_{\hbar}$ is isomorphic to $F[[G]]$ for some formal algebraic group G . Then $F[[G]]$ inherits from F_{\hbar} a Poisson bracket, which makes it into a Poisson Hopf algebra, thus G bears a Poisson structure which makes it into a formal Poisson (algebraic) group. One says then that F_{\hbar} is a *quantization* of the Poisson Hopf algebra $F[[G]]$, or just of the (formal) Poisson group G .

As a general philosophy, from any Hopf-theoretical notion — at the quantum level — one typically infers a Lie-theoretical counterpart — at the semiclassical level. When dealing with deformations, this leads to devising suitable notions of “twists” and “2-cocycles” for Lie bialgebras as well as “deformations” (of Lie bialgebras) by them. In particular, a deformation by twist yields a new Lie bialgebra structure where only the Lie cobracket is modified, whereas deformation by 2-cocycle defines yet another, similar structure where only the Lie bracket is changed.

Following this recipe, the following should hold: when one deforms (as a Hopf algebra) a quantization U_{\hbar} of \mathfrak{g} by a twist which is trivial modulo \hbar , the outcome is a quantization of \mathfrak{g}' , with the latter being a deformation by twist (as a Lie bialgebra) of \mathfrak{g} : moreover, the (Lie) twist working upon \mathfrak{g} is directly “induced” by the (Hopf)

twist that works upon U_{\hbar} , namely the former (Lie) twist is the “semiclassical limit” of the latter (Hopf) twist.

Dually, the following also should hold: when one deforms (as a Hopf algebra) a quantization F_{\hbar} of G by a 2-cocycle which is trivial modulo \hbar , the outcome is a quantization of G' , the latter being a (formal) Poisson group whose cotangent Lie bialgebra is a deformation by 2-cocycle of the cotangent Lie bialgebra $\mathfrak{g}^* := \text{Lie}(G)^*$ of G : moreover, the (Lie) 2-cocycle acting on \mathfrak{g}^* is “induced” by the (Hopf) 2-cocycle that acts on F_{\hbar} , namely the former (Lie) 2-cocycle is the “semiclassical limit” of the latter (Hopf) 2-cocycle.

Nevertheless, neither of the two results mentioned above seems to be published anywhere in literature (to the best of the authors’ knowledge, say). Therefore, as a first contribution in this paper we provide a full, complete statement and proof for the above sketched results, turning them into well-established theorems.

As a second step — our main contribution in this paper — we extend the notions of (Hopf) twist and 2-cocycle, as well as the construction of (Hopf) deformations by them, to a wider setup. Namely, we introduce the notions of *quasi-twist* for a QFSHA and of *quasi-2-cocycle* for a QUEA: roughly speaking, a quasi-twist for F_{\hbar} has the formal Hopf properties of a twist but has the form $\exp(\hbar^{-1}\varphi)$, while any twist (trivial modulo \hbar) looks like $\exp(\hbar^+1\phi)$ — and similarly for the link between quasi-2-cocycles and 2-cocycles. Thus even the very definition of these “quasi-objects”, at least in this form, seems to be problematic, to say the least — as multiplying by \hbar^{-1} in a $\mathbb{k}[[\hbar]]$ -module is meaningless. In spite of this, we show that the recipe defining deformations by twist, resp. by 2-cocycle, for a QFSHA, resp. for a QUEA, still makes sense if we replace “twists” with “quasi-twists”, resp. “2-cocycles” with “quasi-2-cocycles”. Moreover, we can describe the semiclassical limit of these deformations (by “quasi-objects”), again in terms of deformations of Lie bialgebras by some (Lie) twist, resp. 2-cocycle, that can be read out as the semiclassical limit of the quantum (Hopf) quasi-twist, resp. quasi-2-cocycle, that we started with. In a nutshell, we find the perfect “quasi-versions” of the results above for standard quantum group deformations, i.e. those by twist or by 2-cocycle.

The fact that “deformations by quasi-objects” do make sense, albeit problematic at first sight, can be explained in light of the *Quantum Duality Principle (=QDP)*. In fact, the latter provides functorial recipes — via *Drinfeld’s functors* — which turn any QUEA into a QFSHA and any QFSHA into a QUEA. Then, through the QDP lens, every “quasi-twist” for a QFSHA, resp. every “quasi-2-cocycle” for a QUEA, is just a sheer standard twist, resp. 2-cocycle, for the QUEA, resp. the QFSHA, obtained when applying the appropriate Drinfeld functor. In this way, our deformation procedures “by quasi-objects” turn out to be tightly related with standard ones, but applied to different quantum groups. Nevertheless, one still has to prove that the (standard) deformation applied to the new quantum group can actually be “adapted” (by restriction or by extension, depending on the type of quantum group and Drinfeld functor involved) to the original quantum group; in fact, this still requires a detailed, careful analysis.

As a third contribution, we finally consider some constructions of morphisms that, in the setup of general Hopf algebra theory, are provided by R -matrices or ϱ -comatrices. We apply these constructions to the case of quantum groups, showing

that their outcome is much finer than expected from the general theory, and bringing to light their geometrical meaning at the semiclassical level. In addition, we improve those results as follows: we introduce the notions of *quasi- R -matrices* and *quasi- ϱ -comatrices* (much in the same spirit as with quasi-twists and quasi-2-cocycles), and then we extend the construction of morphisms induced by R -matrices and ϱ -comatrices to the case of quasi- R -matrices and quasi- ϱ -comatrices, with a clear interpretation that once more is provided by the QDP.

The paper is organized as follows.

In §2 we quickly recall the material we work with: Lie bialgebras and their deformations, Hopf algebras and their deformations, quantum groups (in the formal setting, both as QUEAs and as QFSHAs) and the *Quantum Duality Principle*.

In §3 we present the bulk of the paper — its main core, in a sense. Namely, we first study deformations of QUEAs by twist and of QFSHAs by 2-cocycles, then we present the new notions of quasi-2-cocycle (for a QUEA) and quasi-twist (for a QFSHA) and the procedures of deformations by these. Later on, all this material is discussed again in §4, in light of the *Quantum Duality Principle*.

Finally, in §5 we study the morphisms defined through R -matrices or ϱ -comatrices, showing the general results (for any Hopf algebra) actually improve in the case of quantum groups and explaining their meaning at the semiclassical limit. Moreover, we also extend those constructions and results to the newly minted notions of quasi- R -matrices and quasi- ϱ -comatrices.

ACKNOWLEDGEMENTS

This article was partially supported by:

- (1) CONICET, ANPCyT, Secyt (Argentina),
- (2) INdAM research branch GNSAGA (Italy),
- (3) the MIUR *Excellence Department Project MatMod@TOV (CUP E83C23000330006)* awarded to the Department of Mathematics, University of Rome “Tor Vergata” (Italy),
- (4) the European research network *COST Action CaLISTA CA21109*.

The authors thank Benjamin Enriquez for sparking their interest in the topics of §5.

2. QUANTUM GROUPS, QUANTUM DUALITY PRINCIPLE, AND DEFORMATIONS

In this section we recap the basic notions we deal with in this paper: Lie bialgebras, quantum groups, deformations of both, and the Quantum Duality Principle.

2.1. Lie bialgebras and Lie deformations.

In this subsection we recall some definitions and basic facts about Lie bialgebras and their deformations. For a more detailed treatment we refer to [CP], [Mj].

Throughout the paper, \mathbb{k} will be a field of characteristic zero.

2.1.1. Generalities. A *Lie bialgebra* is a triple $(\mathfrak{g}; [\ , \], \delta)$ such that $(\mathfrak{g}, [\ , \])$ is a Lie algebra over \mathbb{k} , (\mathfrak{g}, δ) is a *Lie coalgebra* with *Lie cobracket* $\delta : \mathfrak{g} \longrightarrow \mathfrak{g} \wedge \mathfrak{g}$, i.e. $\delta^* : \mathfrak{g}^* \wedge \mathfrak{g}^* \longrightarrow \mathfrak{g}^*$ is a Lie algebra bracket on \mathfrak{g}^*), and the two structures are linked

by the constraint that δ is a 1-cocycle for the Chevalley-Eilenberg cohomology of the Lie algebra $(\mathfrak{g}; [\ , \], \delta)$ with coefficients in $\mathfrak{g} \wedge \mathfrak{g}$:

$$\begin{aligned} \delta([x, y]) &= \text{ad}_x(\delta(y)) - \text{ad}_y(\delta(x)) = \\ &= [x, y_{[1]}] \otimes y_{[2]} + y_{[1]} \otimes [x, y_{[2]}] - [y, x_{[1]}] \otimes x_{[2]} - x_{[1]} \otimes [y, x_{[2]}] \end{aligned} \quad (2.1)$$

using Sweedler's-like notation $\delta(x) = x_{[1]} \otimes x_{[2]}$ for any $x \in \mathfrak{g}$. We write also $x \wedge y := 2^{-1}(x \otimes y - y \otimes x)$ and thus we identify $\mathfrak{g} \wedge \mathfrak{g}$ with the subspace $(\mathfrak{g} \otimes \mathfrak{g})^{\mathbb{Z}_2}$.

Finite-dimensional Lie bialgebras are *self-dual*, in the sense that $(\mathfrak{g}; [\ , \], \delta)$ is a Lie bialgebra if and only if $(\mathfrak{g}^*; \delta^*, [\ , \]^*)$ is so; the latter is called the *dual* Lie bialgebra to $(\mathfrak{g}; [\ , \], \delta)$. This also holds in the infinite-dimensional case, up to technicalities. We denote a Lie bialgebra simply by \mathfrak{g} , and by \mathfrak{g}^* its dual.

Given $r = r_1 \otimes r_2$ in $\mathfrak{g} \otimes \mathfrak{g}$, we write $r_{2,1} := r_2 \otimes r_1$ and $r_{1,2} := r_1 \otimes r_2 \otimes 1$, $r_{2,3} := 1 \otimes r_1 \otimes r_2$, $r_{1,3} := r_1 \otimes 1 \otimes r_2 \in \mathfrak{g} \otimes \mathfrak{g} \otimes \mathfrak{g}$. For example, given $s = s_1 \otimes s_2 \in \mathfrak{g} \otimes \mathfrak{g}$ we define the element

$$\begin{aligned} [[r, s]] &:= [r_{1,2}, s_{1,3}] + [r_{1,2}, s_{2,3}] + [r_{1,3}, s_{2,3}] \\ &= [r_1, s_1] \otimes r_2 \otimes s_2 + r_1 \otimes [r_2, s_1] \otimes s_2 + r_1 \otimes s_1 \otimes [r_2, s_2]. \end{aligned}$$

2.1.2. Deformations of Lie bialgebras. In this work, we are mainly interested in two kinds of deformations, where either the Lie cobracket or the Lie bracket alone is deformed. A general theory of deformations for Lie bialgebras using cohomology theory exists, see e.g. [CG], [MW], and references therein for more details.

Let $(\mathfrak{g}; [\ , \], \delta)$ be a Lie bialgebra and $c \in \mathfrak{g} \otimes \mathfrak{g}$ be such that

$$\text{ad}_x((\delta \otimes \text{id})(c) + \text{c.p.} + [[c, c]]) = 0, \quad \text{ad}_x(c + c_{2,1}) = 0 \quad \forall x \in \mathfrak{g} \quad (2.2)$$

where ad_x denotes the standard adjoint action of x and c.p. means cyclic permutations on the tensor factors of the previous summand. Then, the map $\delta^c : \mathfrak{g} \longrightarrow \mathfrak{g} \wedge \mathfrak{g}$ defined by

$$\delta^c := \delta - \partial(c), \quad \text{i.e.} \quad \delta^c(x) := \delta(x) - \text{ad}_x(c) \quad \forall x \in \mathfrak{g} \quad (2.3)$$

defines a new Lie cobracket on $(\mathfrak{g}; [\ , \], \delta^c)$ making $(\mathfrak{g}; [\ , \], \delta^c)$ into a new Lie bialgebra (cf. [Mj, Theorem 8.1.7]).

Definition 2.1.3. An element $c \in \mathfrak{g} \otimes \mathfrak{g}$ satisfying (2.2) is called a *twist* of the Lie bialgebra \mathfrak{g} , and the corresponding Lie bialgebra $\mathfrak{g}^c := (\mathfrak{g}; [\ , \], \delta^c)$ is called a *deformation by twist* or *twist deformation* of \mathfrak{g} . \diamond

Remark 2.1.4. To be precise, we are adopting here conventions that are slightly different from those in [Mj], yet equivalent. Indeed, we choose to *define* the deformed Lie cobracket in (2.3) via the formula $\delta^c := \delta - \partial(c)$, whereas Majid's definition is $\delta^c := \delta + \partial(c)$. However, our choice of sign yields better-reading statements for our results that connect, through specialization, deformations by twist at the quantum level with those at the semi-classical level (cf. Theorem 3.1.2 and Theorem 3.4.6 later on); that is why we chose such a different option, lest we should insert an odd-looking sign in those results. The equivalence stands in the fact that any $c \in \mathfrak{g} \otimes \mathfrak{g}$ is a twist element in our sense if and only if its opposite $-c$ is a twist in the sense of Majid, and viceversa.

Now we introduce a deformation of the Lie bracket. Let $(\mathfrak{g}; [\ , \], \delta)$ be a Lie bialgebra and $\gamma \in \text{Hom}_{\mathbb{k}}(\mathfrak{g} \otimes \mathfrak{g}, \mathbb{k})$. For \mathfrak{g} finite-dimensional, we identify $\text{Hom}_{\mathbb{k}}(\mathfrak{g} \otimes \mathfrak{g}, \mathbb{k}) = (\mathfrak{g} \otimes \mathfrak{g})^* = \mathfrak{g}^* \otimes \mathfrak{g}^*$. There exists some technicalities in the infinite-dimensional case, yet the outcome is always the same. Dualizing the notion of twist for \mathfrak{g}^* we obtain the dual notion of 2-cocycle as follows. Condition (2.2) with \mathfrak{g}^* replacing \mathfrak{g} and γ in the role of c yields

$$\text{ad}_{\psi}(\partial_*(\gamma) + [[\gamma, \gamma]]_*) = 0, \quad \text{ad}_{\psi}(\gamma + \gamma_{2,1}) = 0 \quad \forall \psi \in \mathfrak{g}^* \quad (2.4)$$

where $\gamma_{2,1} := \gamma^T$ and $(\partial_*(\gamma))(a, b, c) = \gamma([a, b], c) + \text{c.p.}$. Similarly, $[[\ , \]]$ has the same meaning as above but with respect to \mathfrak{g}^* .

For any γ satisfying (2.4), the map $[\ , \]_{\gamma} : \mathfrak{g} \wedge \mathfrak{g} \longrightarrow \mathfrak{g}$ given by

$$[x, y]_{\gamma} := [x, y] + \gamma(x_{[1]}, y) x_{[2]} - \gamma(y_{[1]}, x) y_{[2]} \quad \forall x, y \in \mathfrak{g} \quad (2.5)$$

defines a new Lie bracket on the Lie coalgebra $(\mathfrak{g}; \delta)$ making $(\mathfrak{g}; [\ , \]_{\gamma}, \delta)$ into a new Lie bialgebra (cf. [Mj, Exercise 8.1.8]).

Definition 2.1.5. Every $\gamma \in \text{Hom}_{\mathbb{k}}(\mathfrak{g} \wedge \mathfrak{g}, \mathbb{k})$ that obeys (2.4) is called a *2-cocycle* of the Lie bialgebra \mathfrak{g} , and the Lie bialgebra $\mathfrak{g}_{\gamma} := (\mathfrak{g}; [\ , \]_{\gamma}, \delta)$ is called a *deformation by 2-cocycle* or *2-cocycle deformation* of \mathfrak{g} . \diamond

Remark 2.1.6. Another observation which is dual to Remark 2.1.4 applies to our given definition of 2-cocycle and of 2-cocycle deformation, in comparison to the original one in [Mj]. Again, our notion of 2-cocycle is equivalent to Majid's because any $\gamma \in (\mathfrak{g} \otimes \mathfrak{g})^*$ is a 2-cocycle in our sense if and only if its opposite $-\gamma$ is a 2-cocycle in Majid's, and viceversa.

The following result, which is standard, formalizes the fact that the notions of “twist” and of “2-cocycle” for Lie bialgebras are devised to be dual to each other.

Proposition 2.1.7. *Let \mathfrak{g} be a Lie bialgebra, and \mathfrak{g}^* the dual Lie bialgebra.*

(a) *Let c be a twist for \mathfrak{g} , and γ_c the image of c in $(\mathfrak{g} \otimes \mathfrak{g})^*$ for the natural composed embedding $\mathfrak{g} \otimes \mathfrak{g} \hookrightarrow \mathfrak{g}^{**} \otimes \mathfrak{g}^{**} \hookrightarrow (\mathfrak{g}^* \otimes \mathfrak{g}^*)^*$. Then γ_c is a 2-cocycle for \mathfrak{g}^* , and there exists a canonical isomorphism $(\mathfrak{g}^*)_{\gamma_c} \cong (\mathfrak{g}^c)^*$.*

(b) *Let γ be a 2-cocycle for \mathfrak{g} ; assume that \mathfrak{g} is finite-dimensional, and let c_{γ} be the image of χ in the natural identification $(\mathfrak{g} \otimes \mathfrak{g})^* = \mathfrak{g}^* \otimes \mathfrak{g}^*$. Then c_{γ} is a twist for \mathfrak{g}^* , and there exists a canonical isomorphism $(\mathfrak{g}^*)^{c_{\gamma}} \cong (\mathfrak{g}_{\gamma})^*$. \square*

2.2. Hopf algebra deformations and R -(co)matrices.

Below we recall some notions on deformations for Hopf algebras. Our main reference for general theory of Hopf algebras is [Ra]. For topological (and complete) Hopf algebras, we refer to [Ks], [CP] and [KS].

Let us fix our notation for Hopf algebra theory (mainly standard, indeed). The comultiplication is denoted Δ , the counit ϵ and the antipode \mathcal{S} ; for the first, we use the Heyneman-Sweedler notation, namely $\Delta(x) = x_{(1)} \otimes x_{(2)}$. The augmentation ideal of any Hopf algebra H is denoted by $H^+ := \text{Ker}(\epsilon)$. In general, we denote

by \mathbb{k} the ground ring of our algebras, coalgebras, etc.; we identify it with its image $\mathbb{k}1_H$ through the unit map of H , with $1_H \in H$ being the unit element.

For a Hopf algebra H (or just bialgebra), we write H^{op} , resp. H^{cop} , for the Hopf algebra (or bialgebra) given by taking in H the opposite product, resp. coproduct.

There exist two standard methods to deform Hopf algebras, leading to so-called “2-cocycle deformations” and to “twist deformations”: hereafter we recall both procedures, and their link via duality. They also adapt to the setup of *topological* Hopf algebras; later on, we apply them to quantum groups.

Definition 2.2.1. Let H be a bialgebra (possibly topological, over some commutative ground ring), and let $\mathcal{F} \in H \otimes H$. Then:

(a) \mathcal{F} is said to be *unitary* if

$$(\epsilon \otimes \text{id})(\mathcal{F}) = 1 = (\text{id} \otimes \epsilon)(\mathcal{F}) \quad (2.6)$$

(b) \mathcal{F} is called a *twist* if it is *invertible* in $H \otimes H$, it is *unitary*, and

$$\mathcal{F}_{12}(\Delta \otimes \text{id})(\mathcal{F}) = \mathcal{F}_{23}(\text{id} \otimes \Delta)(\mathcal{F}) \quad (2.7)$$

(c) \mathcal{F} is called a (*quantum*) *R-matrix* if it is *invertible* in $H \otimes H$ and

$$(\Delta \otimes \text{id})(\mathcal{F}) = \mathcal{F}_{13}\mathcal{F}_{23} \quad , \quad (\text{id} \otimes \Delta)(\mathcal{F}) = \mathcal{F}_{13}\mathcal{F}_{12} \quad (2.8)$$

(d) \mathcal{F} is called a (*quantum*) *R-matrix twist* if it complies both (b) and (c) above

(e) \mathcal{F} is said to be a *solution of the quantum Yang-Baxter equation (=QYBE)* if

$$\mathcal{F}_{12}\mathcal{F}_{13}\mathcal{F}_{23} = \mathcal{F}_{23}\mathcal{F}_{13}\mathcal{F}_{12} \quad (2.9)$$

Remarks 2.2.2. (a) If H is a Hopf algebra (including a topological one) and there exists $\mathcal{F} \in H \otimes H$ which is *invertible* and such that

$$\mathcal{F}\Delta(x)\mathcal{F}^{-1} = \Delta^{\text{op}}(x) \quad \forall x \in H \quad (2.10)$$

then H is said to be *quasicocommutative*. If in addition \mathcal{F} obeys also (2.8), then H itself is said to be *quasitriangular*. Indeed, the standard notion of “*R-matrix*” in literature usually demands the constraint (2.10) besides condition (2.8).

Our notion of “*R-matrix*” as in Definition 2.2.1(c) above is given the name “weak *R-matrix*” in [Ch], Definition 1.1.

(b) Every *R-matrix* in the sense of Definition 2.2.1(c) above is automatically *unitary*, cf. [Ch], Lemma 1.2. Conversely, if \mathcal{F} is *unitary* and enjoys (2.8), then it is *invertible* too, hence it is an *R-matrix*. In short, the conditions “*invertible*” and “*unitary*” for an *R-matrix* are equivalent and interchangeable.

(c) If \mathcal{R} is an *R-matrix* for H , then so is $(\mathcal{R}^{-1})_{21} = (\mathcal{R}_{21})^{-1}$; moreover, \mathcal{R}_{21} and \mathcal{R}^{-1} are *R-matrices* for H^{op} and H^{cop} alike — see [Mj], [Ra].

(d) It follows by definitions that (2.7) and (2.8) jointly imply (2.9), while (2.8) and (2.9) jointly imply (2.7).

2.2.3. Deformations by twist. The importance of twists comes from the following construction. Let H be a bialgebra (over some commutative ring \mathbb{k}), and let $\mathcal{F} \in H \otimes H$ be a twist for it — as in Definition 2.2.1(b) above. Then H bears a second

bialgebra structure, denoted $H^{\mathcal{F}}$ and called *twist deformation* of the old one, with the old product, unit and counit, but with a new “twisted” coproduct $\Delta^{\mathcal{F}}$ given by

$$\Delta^{\mathcal{F}}(x) := \mathcal{F} \Delta(x) \mathcal{F}^{-1} \quad \forall x \in H$$

If in addition H is a Hopf algebra with antipode \mathcal{S} , then this “twisted” bialgebra $H^{\mathcal{F}}$ is again a Hopf algebra with antipode $\mathcal{S}^{\mathcal{F}}$ given by

$$\mathcal{S}^{\mathcal{F}}(x) := v \mathcal{S}(x) v^{-1} \quad \forall x \in H$$

where $v := \sum_{\mathcal{F}} \mathcal{S}(f'_1) f'_2$ — with $\sum_{\mathcal{F}} f'_1 \otimes f'_2 = \mathcal{F}^{-1}$ — is invertible in H (see, [CP, §4.2.E], for further details). When H is in fact a topological bialgebra or Hopf algebra, then the same notions still make sense, and the related results apply again, up to properly reading them.

We present now the dual picture:

Definition 2.2.4. Let H be a bialgebra (possibly topological, over some commutative ground ring), and let $\sigma \in (H^{\otimes 2})^*$. Then:

(a) σ is said to be *unitary* if

$$\sigma(a, 1) = \epsilon(a) = \sigma(1, a) \quad \forall a \in H \quad (2.11)$$

(b) σ is called a *2-cocycle* if it is (*convolution*) *invertible* in $(H^{\otimes 2})^*$, it is unitary, and such that

$$\sigma(a_{(1)}, b_{(1)}) \sigma(a_{(2)} b_{(2)}, c) = \sigma(b_{(1)}, c_{(1)}) \sigma(a, b_{(2)} c_{(2)}) \quad (2.12)$$

for all $a, b, c \in H$ — where we abuse of notation identifying $\sigma \in (H \otimes H)^*$ with the corresponding \mathbb{k} -bilinear map $\rho : H \times H \rightarrow \mathbb{k}$, and we adapt notation accordingly;

(c) σ is called a (*quantum*) ϱ -*comatrix* if it is (*convolution*) *invertible* in $(H^{\otimes 2})^*$ and — for all $a, b, c \in H$ — we have

$$\sigma(ab, c) = \sigma(a, c_{(1)}) \sigma(b, c_{(2)}) \quad , \quad \sigma(a, bc) = \sigma(a_{(1)}, c) \sigma(a_{(2)}, b) \quad (2.13)$$

(d) σ is called a (*quantum*) ϱ -*comatrix 2-cocycle* if it complies with (b) and (c);

(e) σ is said to be a *solution of the quantum Yang-Baxter equation (=QYBE)* if

$$\sigma_{12} * \sigma_{13} * \sigma_{23} = \sigma_{23} * \sigma_{13} * \sigma_{12} \quad (2.14)$$

where hereafter “ $*$ ” denotes the convolution product.

Remarks 2.2.5. (a) If H is a Hopf algebra and there exists $\sigma \in (H \otimes H)^*$ which is (*convolution*) *invertible* and such that

$$\sigma * m * \sigma^{-1} = m_{\text{op}} \quad (2.15)$$

then H is said to be *quasicommutative*. If in addition σ obeys also (2.13), then H itself is said to be *coquasitriangular*. Indeed, the standard notion of “ ϱ -comatrix”, or “dual R -matrix”, in literature usually demands (2.15) besides (2.13).

Following the spirit of [Ch], one might also use such terminology as “weak ϱ -comatrix”, or “weak dual R -matrix”.

(b) Every ϱ -matrix in the sense of Definition 2.2.4(c) above is *unitary*. Conversely, if ρ is *unitary* and enjoys (2.13), then it is (*convolution*) *invertible* too,

hence it is a ϱ -comatrix (cf. [Mj], Lemma 2.2.2). In short, the conditions “invertible” and “unitary” for a ϱ -comatrix are equivalent and interchangeable.

(c) Much like for R -matrices, if ρ is a ϱ -comatrix for H , then so is $(\rho^{-1})_{21} = (\rho_{21})^{-1}$; moreover, ρ_{21} and ρ^{-1} are ϱ -comatrices for H^{op} and H^{cop} alike.

(d) It follows by definitions that (2.12) and (2.13) jointly imply (2.14), while (2.13) and (2.14) jointly imply (2.12).

2.2.6. Deformations by 2-cocycles. We can use 2-cocycles to perform a different type of deformation. Let H be a bialgebra (over some commutative ring \mathbb{k}), and let $\sigma \in (H \otimes H)^*$ be a 2-cocycle for it. Then H bears a second bialgebra structure, denoted H_σ and called *2-cocycle deformation* of the old one, with the old coproduct, counit and unit, but with new product $m_\sigma = \sigma * m * \sigma^{-1} : H \otimes H \rightarrow H$ given by

$$m_\sigma(a, b) = a \cdot_\sigma b = \sigma(a_{(1)}, b_{(1)}) a_{(2)} b_{(2)} \sigma^{-1}(a_{(3)}, b_{(3)}) \quad \forall a, b \in H$$

If in addition H is a Hopf algebra with antipode \mathcal{S} , then this “deformed” bialgebra H_σ is again a Hopf algebra with antipode \mathcal{S}_σ , which in detail reads

$$\mathcal{S}_\sigma(a) = \sigma(a_{(1)}, \mathcal{S}(a_{(2)})) \mathcal{S}(a_{(3)}) \sigma^{-1}(\mathcal{S}(a_{(4)}), a_{(5)}) \quad \forall a \in H$$

(see [Doi] for more details). If H is a *topological bialgebra* or *Hopf algebra*, all this construction applies again, as well as the related results, up to technicalities.

The two notions of “2-cocycle” and of “twist”, as well as the corresponding deformations, are so devised as to be dual to each other with respect to Hopf duality (cf. [Mj]), also in the setup of *topological* Hopf algebras as with QUEA’s and QFSHA’s. The same holds for the notions of “ ϱ -comatrix” and of “ R -matrix”. All this is recorded in the following result, whose proof is trivial (an exercise in Hopf theory):

Proposition 2.2.7. *Let H be a Hopf algebra (possibly topological), and H^* its dual Hopf algebra (possibly in topological sense).*

(a) *Let \mathcal{F} be a twist, resp. an R -matrix, for H , and $\sigma_\mathcal{F}$ the image of \mathcal{F} in $(H \otimes H)^*$ for the natural embedding $H \otimes H \hookrightarrow H^{**} \otimes H^{**} \hookrightarrow (H^* \otimes H^*)^*$. Then $\sigma_\mathcal{F}$ is a 2-cocycle, resp. a ϱ -comatrix, for H^* . Moreover, in the first case there exists a canonical Hopf algebra isomorphism $(H^*)_{\sigma_\mathcal{F}} \cong (H^\mathcal{F})^*$.*

(b) *Let σ be a 2-cocycle, resp. a ϱ -comatrix, for H ; assume that we have a natural identification $(H \otimes H)^* = H^* \otimes H^*$ (e.g., if H is finite-dimensional), and let \mathcal{F}_σ be the image of σ in $H^* \otimes H^*$ via this identification. Then \mathcal{F}_σ is a twist, resp. an R -matrix, for H^* . Moreover, in the first case there exists a canonical Hopf algebra isomorphism $(H^*)^{\mathcal{F}_\sigma} \cong (H_\sigma)^*$. \square*

2.2.8. Hopf morphisms from R -matrices and ϱ -comatrices. Let H be a Hopf algebra, possibly in topological sense. We assume that its (possibly topological) finite dual H^* is a Hopf algebra as well (possibly in a topological sense): the easiest example is when H is defined over a field and it is finite dimensional, but the extension to less trivial cases is straightforward in a wide variety of situations.

Hereafter we recall some well-known constructions, somewhat shortly: further details can be found, e.g., in [CP], [KS] and [Mj].

Proposition 2.2.9.

(a) Every R -matrix $\mathcal{R} = \mathcal{R}_1 \otimes \mathcal{R}_2$ (using Sweedler's-like notation) for H provides two Hopf algebra morphisms

$$\overleftarrow{\Phi}_{\mathcal{R}} : H^* \longrightarrow H^{\text{cop}} \quad (\eta \mapsto \eta(\mathcal{R}_1) \mathcal{R}_2) \quad , \quad \overrightarrow{\Phi}_{\mathcal{R}} : H^* \longrightarrow H^{\text{op}} \quad (\eta \mapsto \mathcal{R}_1 \eta(\mathcal{R}_2))$$

(b) If \mathcal{R} is an R -matrix for H , and \mathcal{R}^{-1} is its inverse, then $\overleftarrow{\Phi}_{\mathcal{R}}$, resp. $\overrightarrow{\Phi}_{\mathcal{R}}$, is convolution invertible, with convolution inverse $\overleftarrow{\Phi}_{\mathcal{R}^{-1}}$, resp. $\overrightarrow{\Phi}_{\mathcal{R}^{-1}}$.

Proof. (a) Consider for instance the case of $\overleftarrow{\Phi}_{\mathcal{R}}$. The left-hand side, resp. the right-hand side, of (2.8) implies that it preserves multiplication, resp. comultiplication. On the other hand, from the right-hand side of (2.7), the identity $(\epsilon \otimes \text{id})(\mathcal{R}) = 1$ implies that $\overleftarrow{\Phi}_{\mathcal{R}}$ preserves the unit, and the identity $(\text{id} \otimes \epsilon)(\mathcal{R}) = 1$ implies that it preserves the counit. The proof for $\overrightarrow{\Phi}_{\mathcal{R}}$ is similar, namely it is left-right symmetric.

(b) This follows directly from definitions, through sheer bookkeeping. \square

The previous result has its dual counterpart, whose proof is again straightforward:

Proposition 2.2.10.

(a) Every ϱ -comatrix ρ for H provides two Hopf algebra morphisms

$$\overleftarrow{\Psi}_{\rho} : H \longrightarrow (H^*)^{\text{cop}} \quad , \quad \ell \mapsto \rho(\ell, -) \quad , \quad \overrightarrow{\Psi}_{\rho} : H \longrightarrow (H^*)^{\text{op}} \quad , \quad \ell \mapsto \rho(-, \ell)$$

(b) If ρ is a ϱ -comatrix for H , and ρ^{-1} is its (convolution) inverse, then $\overleftarrow{\Psi}_{\rho}$, resp. $\overrightarrow{\Psi}_{\rho}$, is convolution invertible, with convolution inverse $\overleftarrow{\Psi}_{\rho^{-1}}$, resp. $\overrightarrow{\Psi}_{\rho^{-1}}$. \square

Remark 2.2.11. Inasmuch as any R -matrix, resp. any ϱ -comatrix, for H is a ϱ -comatrix, resp. an R -matrix, for the dual Hopf algebra H^* — cf. Proposition 4.1.2 — applying Proposition 2.2.9 to H^* we get Proposition 2.2.10, and, conversely, applying Proposition 2.2.10 to H^* we get Proposition 2.2.9. In the same spirit, the following result about Hopf algebras in duality follows from the very definitions:

Proposition 2.2.12. Let K and Γ be two Hopf algebras (over the same ground ring, and possibly topological) that are dual to each other, say $\Gamma = K^*$ and $K = \Gamma^*$ for suitably defined dual functors $(\)^*$ and $(\)^*$. Let also $\mathcal{R} = \rho$ be an R -matrix for K and a ϱ -comatrix for Γ — applying Proposition 2.2.7. Then for the morphisms in Proposition 2.2.9 and Proposition 2.2.10 we have canonical identifications

$$\overleftarrow{\Phi}_{\mathcal{R}} = \overleftarrow{\Psi}_{\rho} : K^* = \Gamma \longrightarrow (\Gamma^* = K)^{\text{cop}} \quad , \quad \overrightarrow{\Phi}_{\mathcal{R}} = \overrightarrow{\Psi}_{\rho} : K^* = \Gamma \longrightarrow (\Gamma^* = K)^{\text{op}} \quad \square$$

2.3. Quantum groups.

We recall hereafter the basic notions on quantum groups, in the shape of either quantized universal enveloping algebras (=QUEA's) or quantized formal series Hopf algebras (=QFSHA's). Both types of “quantum groups” are Hopf algebras in a topological sense, that we shall presently fix.

2.3.1. Classical and quantum preliminaries. Hereafter we fix a base field \mathbb{k} of characteristic zero. We recall the following from [CP].

For any Lie algebra \mathfrak{g} over \mathbb{k} , its universal enveloping algebra $U(\mathfrak{g})$ has a canonical structure of Hopf algebra, which is cocommutative and connected. If \mathfrak{g} is also a Lie bialgebra, with Lie cobracket δ , then δ uniquely extends to define a *Poisson cobracket* $\delta : U(\mathfrak{g}) \rightarrow U(\mathfrak{g}) \otimes U(\mathfrak{g})$, just by imposing that it fulfill the co-Leibnitz identity $\delta(xy) = \delta(x)\Delta(y) + \Delta(x)\delta(y)$. Conversely, if the Hopf algebra $U(\mathfrak{g})$ is actually even a Hopf *co-Poisson* algebra, then its Poisson co-bracket δ maps \mathfrak{g} into $\mathfrak{g} \otimes \mathfrak{g}$, thus yielding a Lie cobracket for \mathfrak{g} that makes the latter into a Lie bialgebra.

Dually, let G be any formal algebraic group G over \mathbb{k} : by this we loosely mean that G is the spectrum of its formal function algebra $F[[G]]$, the latter being a topological Hopf algebra which is commutative and I -adically complete, where $I := \text{Ker}(\epsilon)$ is the augmentation ideal of $F[[G]]$. Then G is a (formal) Poisson group if and only if its formal function algebra $F[[G]]$ is a *Poisson* (formal) Hopf algebra, with respect to some Poisson bracket $\{ , \}$. In this case, the cotangent space I/I^2 of G has a Lie bracket induced by $\{ , \}$ via $[x, y] := \{x', y'\} \pmod{I^2}$ for all $x, y \in I/I^2$ with $x = x' \pmod{I^2}$, $y = y' \pmod{I^2}$: this makes I/I^2 into a Lie algebra, but its dual $\mathfrak{g} = \text{Lie}(G) := (I/I^2)^*$ is also a Lie algebra (the tangent Lie algebra to G) and the two structures are compatible, so that $\mathfrak{g}^* := I/I^2$ is a *Lie bialgebra* indeed.

We come now to *quantizations* of the previous co-Poisson/Poisson structures.

Let $\mathcal{T}_{\widehat{\otimes}}$ be the category whose objects are all topological $\mathbb{k}[[\hbar]]$ -modules which are topologically free (i.e. isomorphic to $V[[\hbar]]$ for some \mathbb{k} -vector space V , with the \hbar -adic topology) and whose morphisms are the $\mathbb{k}[[\hbar]]$ -linear maps (which are automatically continuous). This is a tensor category w.r.t. the tensor product $T_1 \widehat{\otimes} T_2$ defined to be the separated \hbar -adic completion of the algebraic tensor product $T_1 \otimes_{\mathbb{k}[[\hbar]]} T_2$ (for all $T_1, T_2 \in \mathcal{T}_{\widehat{\otimes}}$).

Let $\mathcal{P}_{\widehat{\otimes}}$ be the category whose objects are all topological $\mathbb{k}[[\hbar]]$ -modules isomorphic to modules of the type $\mathbb{k}[[\hbar]]^E$ for some set E : these are complete w.r.t. to the weak topology and whose morphisms in $\mathcal{P}_{\widehat{\otimes}}$ are the $\mathbb{k}[[\hbar]]$ -linear continuous maps. This is a tensor category w.r.t. the tensor product $P_1 \widetilde{\otimes} P_2$ defined to be the completion of the algebraic tensor product $P_1 \otimes_{\mathbb{k}[[\hbar]]} P_2$ w.r.t. the weak topology: therefore $P_i \cong \mathbb{k}[[\hbar]]^{E_i}$ ($i = 1, 2$) yields $P_1 \widetilde{\otimes} P_2 \cong \mathbb{k}[[\hbar]]^{E_1 \times E_2}$ (for all $P_1, P_2 \in \mathcal{P}_{\widehat{\otimes}}$).

Note that the objects of $\mathcal{T}_{\widehat{\otimes}}$ and of $\mathcal{P}_{\widehat{\otimes}}$ are complete and separated w.r.t. the \hbar -adic topology, so one has $X \cong X_0[[\hbar]]$ for every such object X , with $X_0 := X/\hbar X$.

We denote by $\mathcal{HA}_{\widehat{\otimes}}$ the subcategory of $\mathcal{T}_{\widehat{\otimes}}$ whose objects are all the Hopf algebras in $\mathcal{T}_{\widehat{\otimes}}$ and whose morphisms are all the Hopf algebra morphisms in $\mathcal{T}_{\widehat{\otimes}}$. Similarly, we call $\mathcal{HA}_{\widetilde{\otimes}}$ the subcategory of $\mathcal{P}_{\widetilde{\otimes}}$ whose objects are all the Hopf algebras in $\mathcal{P}_{\widetilde{\otimes}}$ and whose morphisms are all the Hopf algebra morphisms in $\mathcal{P}_{\widetilde{\otimes}}$. To simplify notation, we shall usually drop the subscripts “ $\widehat{\otimes}$ ” and “ $\widetilde{\otimes}$ ” from the symbol “ \otimes ”.

Finally, when dealing with any $\mathbb{k}[[\hbar]]$ -module M by such notation as $\mathcal{O}(\hbar^s)$ we shall mean any (unspecified) element belonging to $\hbar^s M$, for all $s, n \in \mathbb{N}$; in other words, for any $x \in M$ by writing $x = \mathcal{O}(\hbar^s)$ we mean that $x \equiv 0 \pmod{\hbar^s M}$.

We are ready now to define quantum groups, in two different incarnations:

2.3.2. Quantized Universal Enveloping Algebras (=QUEA's). Retain notation as in §2.3.1 above.

A *quantized universal enveloping algebra* — or QUEA in short — is a (topological) Hopf algebra U_\hbar in $\mathcal{HA}_{\widehat{\otimes}}$ such that $U_0 := U_\hbar/\hbar U_\hbar$ is a connected, cocommutative Hopf algebra over \mathbb{k} — or, equivalently, U_0 is isomorphic to an enveloping algebra $U(\mathfrak{g})$ for some Lie algebra \mathfrak{g} . Then the formula

$$\delta(x) := \frac{\Delta(x') - \Delta^{\text{op}}(x')}{\hbar} \quad \text{mod } \hbar U_\hbar^{\widehat{\otimes} 2} \quad (2.16)$$

— where $x' \in U_\hbar$ is any lift of $x \in \mathfrak{g}$ — defines a co-Poisson structure on $U_0 = U(\mathfrak{g})$, hence a Lie bialgebra structure on \mathfrak{g} . In this case, we say that U_\hbar is a *quantization* of the co-Poisson Hopf algebra $U(\mathfrak{g})$, or (with a slight abuse of language) of the Lie bialgebra \mathfrak{g} ; conversely, $U(\mathfrak{g})$ — or just \mathfrak{g} alone — is the *semiclassical limit* of U_\hbar . We summarize it writing $U_\hbar(\mathfrak{g}) := U_\hbar$. In the following, we denote by *QUEA* the full subcategory of $\mathcal{HA}_{\widehat{\otimes}}$ whose objects are all of the QUEAs.

2.3.3. Quantized Formal Series Hopf Algebras (=QFSHA's). Retain again notation as in §2.3.1 above.

A *quantized formal series Hopf algebra* — or QFSHA in short — is a (topological) Hopf algebra F_\hbar in $\mathcal{HA}_{\widehat{\otimes}}$ such that $F_0 := F_\hbar/\hbar F_\hbar$ is a commutative, I -adically complete topological Hopf algebra over \mathbb{k} , where I is the augmentation ideal — or, equivalently, F_0 is isomorphic to the algebra of functions for some formal algebraic group $F[[G]]$. Then the formula

$$\{x, y\} := \frac{x' y' - y' x'}{\hbar} \quad \text{mod } \hbar F_\hbar \quad (2.17)$$

— where $x', y' \in F_\hbar$ are lifts of $x, y \in F[[G]]$ — defines a Poisson bracket in $F[[G]]$, thus making G into a (formal) Poisson group. In this case, we say that F_\hbar is a *quantization* of the Poisson Hopf algebra $F[[G]]$, or (stretching a point) of the formal Poisson group G ; conversely, $F[[G]]$ — or just G alone — is the *semiclassical limit* of F_\hbar . We summarize it writing $F_\hbar[[G]] := F_\hbar$. In the following, we denote by *QFSHA* the full subcategory of $\mathcal{HA}_{\widehat{\otimes}}$ whose objects are all of the QFSHAs.

2.3.4. Equivalence and duality between quantizations. If H_1, H_2 , are two QUEA's, respectively two QFSHA's, we say that H_1 is *equivalent* to H_2 , and we write $H_1 \equiv H_2$, if there is an isomorphism $\varphi : H_1 \cong H_2$ (in *QUEA*, resp. in *QFSHA*) such that $\varphi = id \text{ mod } \hbar$. In particular, in both cases the semiclassical limit of either H_1 or H_2 is the same.

By their very construction, the categories *QUEA* and *QFSHA* are dual to each other (w. r. to the natural, topological linear duality functors in both directions). In detail, by *dual* of any $U_\hbar \in \text{QUEA}$, denoted U_\hbar^* , we take the set of all $\mathbb{k}[[\hbar]]$ -linear functions from U_\hbar to $\mathbb{k}[[\hbar]]$ (which are automatically continuous w. r. to the \hbar -adic topology): this is naturally an object in *QFSHA*. On the other hand, by *dual* of any $F_\hbar \in \text{QFSHA}$, denoted F_\hbar^* , we take the set of all maps from F_\hbar to $\mathbb{k}[[\hbar]]$ that are continuous with respect to the \hbar -adic topology on $\mathbb{k}[[\hbar]]$ and to the I_\hbar -adic topology on F_\hbar , with $I_\hbar := \hbar F_\hbar + \text{Ker}(\epsilon_{F_\hbar})$; this F_\hbar^* is an object in *QUEA*. Finally, $()^*$ and $()^*$ are contravariant functors inverse to each other — cf. [Ga1].

We finish this part with a trivial, technical result, that we will use several times:

Lemma 2.3.5. *Let H be a Hopf algebra (possibly topological). We denote by $[,]$ the commutator operation in H , and write $H^+ := \text{Ker}(\epsilon)$. Then:*

(a) *There exists a splitting into direct sum $H = \mathbb{k} \oplus H^+$. With respect to that splitting, every $z \in H$ uniquely splits into $z = \epsilon(z) + z^+$ with $z^+ := z - \epsilon(z) \in H^+$.*

(b) *For any $x, y \in H$ we have $[x, y] = [x^+, y^+]$ — see (a) — so $[H, H] \subseteq H^+$.*

(c) *Assume that $H = F_{\hbar}[[G]]$ is a QFSHA, with $J_{\hbar} := H^+$. Then we have $[H, H] = [J_{\hbar}, J_{\hbar}] \subseteq \hbar J_{\hbar}$, and more in general (for all $k, r_1, r_2, r_3, \dots, r_k, s \in \mathbb{N}_+$)*

$$[J_{\hbar}^{r_1}, [J_{\hbar}^{r_2}, [J_{\hbar}^{r_3}, \dots [J_{\hbar}^{r_k}, J_{\hbar}^s] \dots]]] \subseteq (1 - \delta_{s,0}) \prod_{i=1}^k (1 - \delta_{r_i,0}) \hbar^k J_{\hbar}^{r_1+r_2+r_3+\dots+r_k+s-k}$$

(d) *For any $z \in H$, we have*

$$\Delta(z) = \epsilon(z) \cdot 1 \otimes 1 + z^+ \otimes 1 + 1 \otimes z^+ + (z_{(1)})^+ \otimes (z_{(2)})^+$$

or also

$$\Delta(z) = -\epsilon(z) \cdot 1 \otimes 1 + z \otimes 1 + 1 \otimes z + (z_{(1)})^+ \otimes (z_{(2)})^+$$

in particular, for $\nabla := \Delta - \Delta^{\text{op}}$ this yields $\nabla(z) \in \text{Ker}(H)^{\otimes 2}$, with

$$\nabla(z) = (z_{(1)})^+ \otimes (z_{(2)})^+ - (z_{(2)})^+ \otimes (z_{(1)})^+ \quad \square$$

2.4. The Quantum Duality Principle.

We recall hereafter the main facets of the so-called ‘‘Quantum Duality Principle’’, which establishes an explicit equivalence between the category of QUEA’s and that of QFSHA’s (whereas linear duality provides an *antiequivalence* instead). Further details can be found in [Gal].

Definition 2.4.1. (*Drinfeld’s functors*) We define Drinfeld’s functors from QUEA to QFSHA and viceversa as follows:

(a) Let $U_{\hbar}(\mathfrak{g})$ be any QUEA, and assume for simplicity that \mathfrak{g} is finite-dimensional. Let $\iota : \mathbb{k}[[\hbar]] \longrightarrow U_{\hbar}(\mathfrak{g})$ and $\epsilon : U_{\hbar}(\mathfrak{g}) \longrightarrow \mathbb{k}[[\hbar]]$ be its unit and counit maps; moreover, for every $n \in \mathbb{N}$ set $\delta_n := (\text{id} - \iota \circ \epsilon)^{\otimes n} \circ \Delta^{(n-1)}$ — mapping $U_{\hbar}(\mathfrak{g})$ to $U_{\hbar}(\mathfrak{g})^{\widehat{\otimes} n}$. Then we define

$$U_{\hbar}(\mathfrak{g})' := \left\{ \eta \in U_{\hbar}(\mathfrak{g}) \mid \delta_n(\eta) \in \hbar^n U_{\hbar}(\mathfrak{g})^{\otimes n} \quad \forall n \in \mathbb{N} \right\}$$

This defines the functor $()'$, from QUEA to QFSHA, onto objects: then onto morphisms it is clearly defined by taking restriction.

(b) Let $F_{\hbar}[[G]]$ be any QFSHA, and assume for simplicity that G be finite-dimensional. Let $\epsilon_F : F_{\hbar}[[G]] \longrightarrow \mathbb{k}[[\hbar]]$ be its counit map, and consider also $I_{F_{\hbar}[[G]]} := \hbar F_{\hbar}[[G]] + \text{Ker}(\epsilon_F)$. Then we define

$$F_{\hbar}[[G]]^{\vee} := \hbar\text{-adic completion of } \sum_{n \geq 0} \hbar^{-n} I_{F_{\hbar}[[G]]}^n$$

This defines the functor $()^{\vee}$, from QFSHA to QUEA, onto objects: onto morphisms, we define it via scalar extension — from $\mathbb{k}[[\hbar]]$ to $\mathbb{k}((\hbar))$ — followed by restriction and completion. \diamond

The original recipes for these functors were given in [Dr, §7]; the corresponding proofs (that everything is well-defined, etc.) can be found in [Ga1]. Indeed, the overall result is very strong, involving linear duality for Lie bialgebras, as follows:

Theorem 2.4.2. (“The quantum duality principle”; cf. [Dr], [Ga1])

(a) The assignments $H \mapsto H'$ and $H \mapsto H^\vee$ respectively define functors of tensor categories $QUEA \longrightarrow QFSHA$ and $QFSHA \longrightarrow QUEA$, that are inverse to each other, thus yielding an equivalence of categories.

(b) For all $U_\hbar(\mathfrak{g}) \in QUEA$ and all $F_\hbar[[G]] \in QFSHA$ one has

$$U_\hbar(\mathfrak{g})' / \hbar U_\hbar(\mathfrak{g})' = F[[G^*]] \quad , \quad F_\hbar[[G]]^\vee / \hbar F_\hbar[[G]]^\vee = U(\mathfrak{g}^*)$$

that is, if $U_\hbar(\mathfrak{g})$ is a quantization of $U(\mathfrak{g})$ then $U_\hbar(\mathfrak{g})'$ is a quantization of $F[[G^*]]$, and if $F_\hbar[[G]]$ is a quantization of $F[[G]]$ then $F_\hbar[[G]]^\vee$ is a quantization of $U(\mathfrak{g}^*)$.

(c) Both Drinfeld’s functors preserve equivalence, that is $H_1 \equiv H_2$ implies that $H_1' \equiv H_2'$ and $H_1^\vee \equiv H_2^\vee$ in either case. \square

In a very precise sense, Drinfeld’s functors are dual to each other: namely — cf. [Ga1] — one has (with notation as in §2.3.4)

$$(U_\hbar(\mathfrak{g})^*)^\vee = (U_\hbar(\mathfrak{g})')^* \quad \text{and} \quad (F_\hbar[[G]]^\vee)^* = (F_\hbar[[G]]')^* \quad (2.18)$$

On the other hand, it is worth stressing a strong asymmetry between these functors. Indeed, the definition of $F_\hbar[[G]]^\vee$ is pretty *concrete* (through an explicit generating procedure) whereas that of $U_\hbar(\mathfrak{g})'$ is somewhat *implicit* (it is described as the set of solution of a system of countably many equations), hence way more tough to work out. Nevertheless, an alternative description for $U_\hbar(\mathfrak{g})'$ is available, that we shall make use of in the later on, namely the following (cf. [Ga2, Proposition 3.1.2]):

Proposition 2.4.3. For any \mathbb{k} -basis $\{\bar{y}_i\}_{i \in I}$ of \mathfrak{g} , there are $y_i \in U_\hbar(\mathfrak{g})$ such that:

(a) $\epsilon(y_i) = 0$, $(y_i \bmod \hbar U_\hbar(\mathfrak{g})) = \bar{y}_i$ and $y_i' := \hbar y_i \in U_\hbar(\mathfrak{g})'$ for all $i \in I$;

(b) $U_\hbar(\mathfrak{g})'$ is the completion of the unital $\mathbb{k}[[\hbar]]$ -subalgebra of $U_\hbar(\mathfrak{g})$ generated by all the x_i' ’s with respect to its I'_\hbar -adic topology, where I'_\hbar is the ideal (in that subalgebra) generated by \hbar and all the x_i' ’s, so that $U_\hbar(\mathfrak{g})' = \mathbb{k}[[\{x_i'\}_{i \in I} \cup \{\hbar\}]]$. \square

3. DEFORMATIONS OF QUANTUM GROUPS

This section is dedicated to explore the effect of deformations of quantum groups, either by twist or by 2-cocycle, setting the cases of QUEA’s and QFSHA’s apart.

3.1. Deformations by twist of QUEA’s.

In this subsection we consider deformations by twist of QUEA’s; in some sense, this is the easiest case. We begin with a technical result:

Lemma 3.1.1. Let H be an \hbar -adically complete Hopf algebra over $\mathbb{k}[[\hbar]]$, and let consider an element of the form $\mathcal{F} = \exp(\hbar\varphi) \in H \otimes H$, with $\varphi = \varphi_1 \otimes \varphi_2 \in H^{\otimes 2}$, such that $(\epsilon \otimes \text{id})(\mathcal{F}) = 1 = (\text{id} \otimes \epsilon)(\mathcal{F})$. Then

$$\epsilon(\varphi_1) \otimes \varphi_2 = 0 \quad , \quad \varphi_1 \otimes \epsilon(\varphi_2) = 0 \quad , \quad \epsilon(\varphi_1) \otimes \epsilon(\varphi_2) = 0$$

As a consequence, one can assume $\varphi_1 = \varphi_1^+$, $\varphi_2 = \varphi_2^+ \in \text{Ker}(\epsilon)$, so $\varphi \in \text{Ker}(\epsilon)^{\otimes 2}$.

Proof. By assumption one has $(\epsilon \otimes \text{id})(\mathcal{F}) = 1$. Since $\mathcal{F} = \exp(\hbar\varphi)$, this yields

$$\begin{aligned} \hbar\epsilon(\varphi_1) \otimes \varphi_2 &= (\epsilon \otimes \text{id})(\hbar\varphi) = (\epsilon \otimes \text{id})(\log(\mathcal{F})) = \\ &= (\epsilon \otimes \text{id})\left(\sum_{n>0} \frac{(-1)^{n-1}}{n} (\mathcal{F} - 1)^n\right) = \sum_{n>0} \frac{(-1)^{n-1}}{n} ((\epsilon \otimes \text{id})(\mathcal{F}) - 1)^n = 0 \end{aligned}$$

thus $\hbar\epsilon(\varphi_1) \otimes \varphi_2 = 0$, whence $\epsilon(\varphi_1) \otimes \varphi_2 = 0$ as claimed. Similarly, the condition $(\text{id} \otimes \epsilon)(\mathcal{F}) = 1$ implies $\varphi_1 \otimes \epsilon(\varphi_2) = 0$. Finally, expanding φ_i as $\varphi_i = \epsilon(\varphi_i) + \varphi_i^+$ ($i \in \{1, 2\}$) — cf. Lemma 2.3.5 — and using $\epsilon(\varphi_1) \otimes \varphi_2 = 0 = \varphi_1 \otimes \epsilon(\varphi_2)$ we get

$$\begin{aligned} \varphi &= \varphi_1 \otimes \varphi_2 = \epsilon(\varphi_1) \otimes \epsilon(\varphi_2) + \epsilon(\varphi_1) \otimes \varphi_2^+ + \varphi_1^+ \otimes \epsilon(\varphi_2) + \varphi_1^+ \otimes \varphi_2^+ = \\ &= \epsilon(\varphi_1) \otimes \epsilon(\varphi_2) + \epsilon(\varphi_1) \otimes \varphi_2 - \epsilon(\varphi_1) \otimes \epsilon(\varphi_2) + \varphi_1 \otimes \epsilon(\varphi_2) - \epsilon(\varphi_1) \otimes \epsilon(\varphi_2) + \varphi_1^+ \otimes \varphi_2^+ = \\ &= -\epsilon(\varphi_1) \otimes \epsilon(\varphi_2) + \epsilon(\varphi_1) \otimes \varphi_2 + \varphi_1 \otimes \epsilon(\varphi_2) + \varphi_1^+ \otimes \varphi_2^+ = \\ &= -\epsilon(\varphi_1) \otimes \epsilon(\varphi_2) + \varphi_1^+ \otimes \varphi_2^+ \end{aligned}$$

in short $\varphi = -\epsilon(\varphi_1) \otimes \epsilon(\varphi_2) + \varphi_1^+ \otimes \varphi_2^+$. But $\epsilon(\varphi_1) \otimes \epsilon(\varphi_2) = (\text{id} \otimes \epsilon)(\epsilon(\varphi_1) \otimes \varphi_2) = 0$, hence $\varphi = \varphi_1^+ \otimes \varphi_2^+ \in \text{Ker}(\epsilon)^{\otimes 2}$, q.e.d. \square

We are now ready for our first meaningful result:

Theorem 3.1.2.

Let $U_{\hbar}(\mathfrak{g})$ be a QUEA over the Lie bialgebra $\mathfrak{g} = (\mathfrak{g}; [\ , \], \delta)$. Let $\mathcal{F} \in U_{\hbar}(\mathfrak{g})^{\widehat{\otimes} 2}$ be a twist for $U_{\hbar}(\mathfrak{g})$ such that $\mathcal{F} \equiv 1 \pmod{\hbar U_{\hbar}(\mathfrak{g})^{\widehat{\otimes} 2}}$; then $\kappa := \hbar^{-1} \log(\mathcal{F}) \in U_{\hbar}(\mathfrak{g})^{\widehat{\otimes} 2}$, and $\mathcal{F} = \exp(\hbar\kappa)$. Last, we set $\kappa_a := \kappa - \kappa_{2,1}$. Then we have:

- (a) κ is antisymmetric, i.e. $-\kappa = \kappa_{2,1}$, iff \mathcal{F} is orthogonal, i.e. $\mathcal{F}^{-1} = \mathcal{F}_{2,1}$;
- (b) the element $c := \overline{\kappa_a} = \kappa_a \pmod{\hbar U_{\hbar}(\mathfrak{g})^{\widehat{\otimes} 2}}$ belongs to $\mathfrak{g} \otimes \mathfrak{g}$, and it is an antisymmetric twist element for the Lie bialgebra \mathfrak{g} ;
- (c) the deformation $(U_{\hbar}(\mathfrak{g}))^{\mathcal{F}}$ of $U_{\hbar}(\mathfrak{g})$ by the twist \mathcal{F} is a QUEA for the Lie bialgebra $\mathfrak{g}^c = (\mathfrak{g}; [\ , \], \delta^c)$ which is the deformation of \mathfrak{g} by the twist c ; in a nutshell, we have $(U_{\hbar}(\mathfrak{g}))^{\mathcal{F}} \cong U_{\hbar}(\mathfrak{g}^c)$.

Proof. (a) By construction, this follows from standard identities for exponentials and for logarithms.

(b) We fix hereafter the notation $U_{\hbar} := U_{\hbar}(\mathfrak{g})$ and $J_{\hbar} := \text{Ker}(\epsilon_{U_{\hbar}})$, and we write $\kappa \in U_{\hbar}^{\widehat{\otimes} 2}$ with Sweedler's like σ -notation $\kappa = \kappa_1 \otimes \kappa_2$. By Lemma 3.1.1 we can assume (as we shall do henceforth) that $\kappa_1, \kappa_2 \in J_{\hbar}$, hence $\kappa \in J_{\hbar}^{\widehat{\otimes} 2}$.

Now we consider the identity $\mathcal{F}_{12}(\Delta \otimes \text{id})(\mathcal{F}) = \mathcal{F}_{23}(\text{id} \otimes \Delta)(\mathcal{F})$. Writing $\mathcal{F} = \exp(\hbar\kappa_1 \otimes \kappa_2)$ and $\Delta(\kappa_s) = \kappa_s^{(1)} \otimes \kappa_s^{(2)}$ ($s = 1, 2$) this reads

$$\exp(\hbar\kappa_1 \otimes \kappa_2 \otimes 1) \exp(\hbar\kappa_1^{(1)} \otimes \kappa_1^{(2)} \otimes \kappa_2) = \exp(\hbar 1 \otimes \kappa_1 \otimes \kappa_2) \exp(\hbar\kappa_1 \otimes \kappa_2^{(1)} \otimes \kappa_2^{(2)})$$

Now taking \hbar -adic expansion in both sides of this last identity, at order 0 — in \hbar — we get $1 \otimes 1 \otimes 1 = 1 \otimes 1 \otimes 1$, hence from order 1 we get the non-trivial identity

$$\kappa_1 \otimes \kappa_2 \otimes 1 + \kappa_1^{(1)} \otimes \kappa_1^{(2)} \otimes \kappa_2 \equiv_{\hbar} 1 \otimes \kappa_1 \otimes \kappa_2 + \kappa_1 \otimes \kappa_2^{(1)} \otimes \kappa_2^{(2)} \quad (3.1)$$

where hereafter any symbol \equiv_{\hbar^n} means “congruent modulo $\hbar^n U_{\hbar}^{\otimes 3}$ ” (for any $n \in \mathbb{N}$).

Then taking (3.1) modulo \hbar we get

$$\overline{\kappa_1} \otimes \overline{\kappa_2} \otimes 1 + \overline{\kappa_1}^{(1)} \otimes \overline{\kappa_1}^{(2)} \otimes \overline{\kappa_2} = 1 \otimes \overline{\kappa_1} \otimes \overline{\kappa_2} + \overline{\kappa_1} \otimes \overline{\kappa_2}^{(1)} \otimes \overline{\kappa_2}^{(2)} \quad (3.2)$$

where hereafter $\overline{x} := x \pmod{\hbar}$ and we took into account that $\overline{\kappa_s^{(i)}} = \overline{\kappa_s}^{(i)}$ for all $s, i \in \{1, 2\}$. Now, $\overline{\kappa_s^{(1)}} \otimes \overline{\kappa_s^{(2)}} = \Delta(\overline{\kappa_s})$ with $\overline{\kappa_s} \in U_{\hbar}/\hbar U_{\hbar} = U(\mathfrak{g})$ has the form

$$\overline{\kappa_s}^{(1)} \otimes \overline{\kappa_s}^{(2)} = \overline{\kappa_s} \otimes 1 + 1 \otimes \overline{\kappa_s} + \dot{\overline{\kappa_s}}^{(1)} \otimes \dot{\overline{\kappa_s}}^{(2)} \quad (3.3)$$

for some $\dot{\overline{\kappa_s}}^{(i)} \in \text{Ker}(\epsilon_{U(\mathfrak{g})})$ — $i \in \{1, 2\}$ — having the following property: if we denote by $U(\mathfrak{g})_n$ the n -th piece in the canonical filtration of $U(\mathfrak{g})$ — its coradical filtration, in Hopf theoretical language — and for any $x \in U(\mathfrak{g})_n \setminus U(\mathfrak{g})_{n-1}$ we set $\partial(x) := n$, then in (3.3) we have $\partial(\dot{\overline{\kappa_s}}^{(i)}) \not\leq \partial(\overline{\kappa_s})$. Now, using (3.3) to re-write (3.2) we find, after cancelling out three summands on both sides, the following

$$\dot{\overline{\kappa_1}}^{(1)} \otimes \dot{\overline{\kappa_1}}^{(2)} \otimes \overline{\kappa_2} = \overline{\kappa_1} \otimes \dot{\overline{\kappa_2}}^{(1)} \otimes \dot{\overline{\kappa_2}}^{(2)}$$

and then the condition $\partial(\dot{\overline{\kappa_s}}^{(i)}) \not\leq \partial(\overline{\kappa_s})$ forces $\dot{\overline{\kappa_1}}^{(1)} \otimes \dot{\overline{\kappa_1}}^{(2)} = 0 = \dot{\overline{\kappa_2}}^{(1)} \otimes \dot{\overline{\kappa_2}}^{(2)}$.

Thus (3.3) reads $\Delta(\overline{\kappa_s}) = \overline{\kappa_s}^{(1)} \otimes \overline{\kappa_s}^{(2)} = \overline{\kappa_s} \otimes 1 + 1 \otimes \overline{\kappa_s}$; this means $\overline{\kappa_s} \in \mathfrak{g} \in (\subseteq U(\mathfrak{g}))$ — $s \in \{1, 2\}$ — so $\overline{\kappa} = \overline{\kappa_1} \otimes \overline{\kappa_2} \in \mathfrak{g} \otimes \mathfrak{g}$, hence $c := \overline{\kappa_a} \in \mathfrak{g} \otimes \mathfrak{g}$.

Now we have to prove that c is an *antisymmetric twist* for the Lie bialgebra \mathfrak{g} .

Keeping notation from above, since $\overline{\kappa_s} \in \mathfrak{g}$ we have

$$\Delta(\kappa_s) \equiv_{\hbar^2} \kappa_s \otimes 1 + 1 \otimes \kappa_s + \hbar \kappa_s^{[1]} \otimes \kappa_s^{[2]} \quad (3.4)$$

with $\overline{\kappa_s}^{[1]} \otimes \overline{\kappa_s}^{[2]} - \overline{\kappa_s}^{[2]} \otimes \overline{\kappa_s}^{[1]} = \delta(\overline{\kappa_s})$ being the Lie cobracket of $\overline{\kappa_s}$, by assumption. When we plug (3.4) in the \hbar -adic expansion of the identity

$$\exp(\hbar \kappa_1 \otimes \kappa_2 \otimes 1) \exp\left(\hbar \kappa_1^{(1)} \otimes \kappa_1^{(2)} \otimes \kappa_2\right) = \exp(\hbar 1 \otimes \kappa_1 \otimes \kappa_2) \exp\left(\hbar \kappa_1 \otimes \kappa_2^{(1)} \otimes \kappa_2^{(2)}\right)$$

we find that at order 2 — in \hbar — it implies an identity

$$\overline{\kappa_1}^{[1]} \otimes \overline{\kappa_1}^{[2]} \otimes \overline{\kappa_2} + \overline{\kappa_{1,2}} \cdot \overline{\kappa_{1,3}} + \overline{\kappa_{1,2}} \cdot \overline{\kappa_{2,3}} = \overline{\kappa_1} \otimes \overline{\kappa_2}^{[1]} \otimes \overline{\kappa_2}^{[2]} + \overline{\kappa_{2,3}} \cdot \overline{\kappa_{1,2}} + \overline{\kappa_{2,3}} \cdot \overline{\kappa_{1,3}} \quad (3.5)$$

where each $\overline{\kappa_{i,j}}$, as usual, is the tensor in $\mathfrak{g}^{\otimes 3}$ which sports the κ_1 's in position i , the κ_2 's in position j , and a (repeated) tensor factor 1 in the last remaining position.

Now let us consider $\mathbb{k}[\mathbb{S}_3]$, the group algebra over \mathbb{k} of the symmetric group \mathbb{S}_3 , the “antisymmetrizer” $Alt_3 := (\text{id} - (12) - (23) - (31) + (123) + (321))$ in $\mathbb{k}[\mathbb{S}_3]$, and the natural action of $\mathbb{k}[\mathbb{S}_3]$ onto $U(\mathfrak{g})^{\otimes 3}$. Let Alt_3 act on the identity (3.5): a sheerly straightforward calculation shows that the outcome, using notation $c := \overline{\kappa_a} = \overline{\kappa} - \overline{\kappa_{2,1}}$, eventually is

$$(\delta \otimes \text{id})(c) + \text{c.p.} + [[c, c]] = 0$$

This means exactly that c is a twist for the Lie bialgebra \mathfrak{g} , as in Definition 2.1.3, which is obviously antisymmetric (by construction), q.e.d.

(c) Due to the peculiar form of the twist — namely, its being trivial modulo \hbar — it is easy to see that the Hopf algebra $U_\hbar(\mathfrak{g})^\mathcal{F}$ is again a QUEA, over some bialgebra $\tilde{\mathfrak{g}}$, i.e. $U_\hbar(\mathfrak{g})^\mathcal{F}/\hbar U_\hbar(\mathfrak{g})^\mathcal{F} = U(\tilde{\mathfrak{g}})$, and even that one has $\tilde{\mathfrak{g}} = \mathfrak{g}$ as Lie algebras. In fact, since the twist \mathcal{F} is trivial modulo \hbar , we have that $U_\hbar(\mathfrak{g})/\hbar U_\hbar(\mathfrak{g})$ and $U_\hbar(\mathfrak{g})^\mathcal{F}/\hbar U_\hbar(\mathfrak{g})^\mathcal{F}$ are *isomorphic as Hopf algebras*; in particular, then, $U_\hbar(\mathfrak{g})^\mathcal{F}$ itself is again a QUEA, on the same Lie algebra \mathfrak{g} from $U_\hbar(\mathfrak{g})$ but possibly inducing on \mathfrak{g} a different Lie cobracket. Indeed, what is actually affected, a priori, is the co-Poisson structure on the semiclassical limit — hence the Lie cobracket on \mathfrak{g} — which in general on $U_\hbar(\mathfrak{g})^\mathcal{F}/\hbar U_\hbar(\mathfrak{g})^\mathcal{F}$ will be different from that on $U_\hbar(\mathfrak{g})/\hbar U_\hbar(\mathfrak{g})$.

Let us compute the Lie coalgebra structure of $\tilde{\mathfrak{g}}$ given by (2.16). Given $x \in \tilde{\mathfrak{g}}$, let $x \in U_\hbar(\mathfrak{g})^\mathcal{F}$ be any lift of x : using obvious notation, its twisted coproduct is

$$\begin{aligned} \Delta^\mathcal{F}(x) &= \mathcal{F} \Delta(x) \mathcal{F}^{-1} = e^{\hbar \kappa} \left(x \otimes 1 + 1 \otimes x + \hbar \sum_i x_1^{[i]} \otimes x_2^{[i]} + O(\hbar^2) \right) e^{-\hbar \kappa} = \\ &= (1 \otimes 1 + \hbar \kappa) \left(x \otimes 1 + 1 \otimes x + \hbar \sum_i x_1^{[i]} \otimes x_2^{[i]} \right) (1 \otimes 1 - \hbar \kappa) + O(\hbar^2) = \\ &= x \otimes 1 + 1 \otimes x + \hbar [\kappa, x \otimes 1 + 1 \otimes x] + \hbar \sum_i x_1^{[i]} \otimes x_2^{[i]} + O(\hbar^2) = \\ &= x \otimes 1 + 1 \otimes x + \hbar \left(\sum_i x_1^{[i]} \otimes x_2^{[i]} - \text{ad}_x(\kappa) \right) + O(\hbar^2) \end{aligned}$$

On the other hand, the opposite twisted coproduct is

$$\begin{aligned} (\Delta^\mathcal{F})^{\text{op}}(x) &= (\mathcal{F})_{21} \Delta^{\text{op}}(x) (\mathcal{F})_{21}^{-1} = e^{\hbar \kappa_{2,1}} \Delta^{\text{op}}(x) e^{-\hbar \kappa_{2,1}} = \\ &= x \otimes 1 + 1 \otimes x + \hbar \left(\sum_i x_2^{[i]} \otimes x_1^{[i]} - \text{ad}_x(\kappa_{2,1}) \right) + O(\hbar^2) \end{aligned}$$

Thus, by the very definition of the cobracket — as in (2.16) — we have

$$\delta^\mathcal{F}(x) := \delta(x) + (\text{ad}_x(\kappa_{2,1} - \kappa)) \pmod{\hbar} = \delta(x) - \text{ad}_x(c) =: \delta^c(x)$$

hence $\tilde{\mathfrak{g}}$ is the twist deformation by c of the Lie bialgebra \mathfrak{g} , as claimed. \square

Observation 3.1.3. Let us point out that the twists \mathcal{F} considered in Theorem 3.1.2 above are those of “trivial type”, as they are the identity modulo \hbar . This ensures that twisting $U_\hbar(\mathfrak{g})$ by such an \mathcal{F} does not affect the Hopf structure of the semiclassical limit; in particular, it still is of the form $U(\tilde{\mathfrak{g}})$, with $\tilde{\mathfrak{g}}$ equal to \mathfrak{g} as a Lie algebra but with a different Lie coalgebra structure. A more general twist might be “unfit”, i.e. the deformed Hopf algebra $U_\hbar(\mathfrak{g})^\mathcal{F}$ might no longer be a QUEA.

We present now a concrete example, taken from [GaGa2], where formal “multi-parameter” QUEAs are studied in detail.

Example 3.1.4. Let $n \in \mathbb{N}_+$ and $I := \{1, \dots, n\}$. We fix a free $\mathbb{k}[[\hbar]]$ -module \mathfrak{h} of finite rank t , and we pick subsets $\Pi^\vee := \{T_i^+, T_i^-\}_{i \in I} \subseteq \mathfrak{h}$, $\Pi := \{\alpha_i\}_{i \in I} \subseteq \mathfrak{h}^* := \text{Hom}_{\mathbb{k}[[\hbar]]}(\mathfrak{h}, \mathbb{k}[[\hbar]])$. Let $P \in M_n(\mathbb{k}[[\hbar]])$ be any $(n \times n)$ -matrix with entries in $\mathbb{k}[[\hbar]]$. A *realization* of P over $\mathbb{k}[[\hbar]]$ of rank t is a triple $\mathcal{R} := (\mathfrak{h}, \Pi, \Pi^\vee)$ where $\alpha_j(T_i^+) = p_{ij}$, $\alpha_j(T_i^-) = p_{ji}$ ($\forall i, j \in I$), and $\bar{\Sigma} := \{\bar{S}_i := 2^{-1}(T_i^+ + T_i^-) \pmod{\hbar \mathfrak{h}}\}_{i \in I}$ is \mathbb{k} -linearly independent as a subset in $\bar{\mathfrak{h}} := \mathfrak{h}/\hbar \mathfrak{h}$.

Let $A := (a_{ij})_{i,j \in I} \in M_n(\mathbb{k})$ be a symmetrisable generalized Cartan matrix, with associated diagonal matrix $D := (d_i \delta_{ij})_{i,j \in I}$. We say that a matrix $P \in M_n(\mathbb{k}[[\hbar]])$ is of *Cartan type* with corresponding Cartan matrix A if $P_s := 2^{-1}(P + P^T) = DA$.

A *formal multiparameter quantum universal enveloping algebra* (=FoMpQUEA) with multiparameter P and realization \mathcal{R} is the unital, associative, topological, \hbar -adically complete $\mathbb{k}[[\hbar]]$ -algebra $U_{P,\hbar}^{\mathcal{R}}(\mathfrak{g})$ generated by the $\mathbb{k}[[\hbar]]$ -submodule \mathfrak{h} and all E_i, F_i (for all $i \in I$), with relations (for all $T, T', T'' \in \mathfrak{h}, i, j \in I$)

$$\begin{aligned} T E_j - E_j T &= +\alpha_j(T) E_j, & T F_j - F_j T &= -\alpha_j(T) F_j \\ T' T'' &= T'' T', & E_i F_j - F_j E_i &= \delta_{i,j} \frac{e^{+\hbar T_i^+} - e^{-\hbar T_i^-}}{q_i^{+1} - q_i^{-1}} \\ \sum_{k=0}^{1-a_{ij}} (-1)^k \begin{bmatrix} 1-a_{ij} \\ k \end{bmatrix}_{q_i} q_{ij}^{+k/2} q_{ji}^{-k/2} E_i^{1-a_{ij}-k} E_j E_i^k &= 0 & (i \neq j) \\ \sum_{k=0}^{1-a_{ij}} (-1)^k \begin{bmatrix} 1-a_{ij} \\ k \end{bmatrix}_{q_i} q_{ij}^{+k/2} q_{ji}^{-k/2} F_i^{1-a_{ij}-k} F_j F_i^k &= 0 & (i \neq j) \end{aligned} \quad (3.6)$$

By [GaGa2, Theorem 4.3.2], every FoMpQUEA $U_{P,\hbar}^{\mathcal{R}}(\mathfrak{g})$ bears a structure of topological Hopf algebra over $\mathbb{k}[[\hbar]]$ — with coproduct taking values into the \hbar -adically completed tensor product $U_{P,\hbar}^{\mathcal{R}}(\mathfrak{g}) \widehat{\otimes}_{\mathbb{k}[[\hbar]]} U_{P,\hbar}^{\mathcal{R}}(\mathfrak{g})$ — given by $(\forall T \in \mathfrak{h}, \ell \in I)$

$$\begin{aligned} \Delta(E_\ell) &= E_\ell \otimes 1 + e^{+\hbar T_\ell^+} \otimes E_\ell, & \Delta(T) &= T \otimes 1 + 1 \otimes T, & \Delta(F_\ell) &= F_\ell \otimes e^{-\hbar T_\ell^-} + 1 \otimes F_\ell \\ \epsilon(E_\ell) &= 0, & \epsilon(T) &= 0, & \epsilon(F_\ell) &= 0 \\ \mathcal{S}(E_\ell) &= -e^{-\hbar T_\ell^+} E_\ell, & \mathcal{S}(T) &= -T, & \mathcal{S}(F_\ell) &= -F_\ell e^{+\hbar T_\ell^-} \end{aligned}$$

Furthermore, by [GaGa2, Theorem 6.1.4], $U_{P,\hbar}^{\mathcal{R}}(\mathfrak{g})$ is a *quantized universal enveloping algebra* whose semiclassical limit is $U(\mathfrak{g}_{\bar{P}}^{\mathcal{R}})$, where $\mathfrak{g}_{\bar{P}}^{\mathcal{R}}$ is a Lie *multiparameter* Lie bialgebra. In short, for each pair (P, \mathcal{R}) as above, $U(\mathfrak{g}_{\bar{P}}^{\mathcal{R}})$ is the *specialization* of $U_{P,\hbar}^{\mathcal{R}}(\mathfrak{g})$, or — equivalently — $U_{P,\hbar}^{\mathcal{R}}(\mathfrak{g})$ is a *quantization* of $U(\mathfrak{g}_{\bar{P}}^{\mathcal{R}})$ — or also, by a standard abuse of language, $U_{P,\hbar}^{\mathcal{R}}(\mathfrak{g})$ is a *quantization* of $\mathfrak{g}_{\bar{P}}^{\mathcal{R}}$. In particular, writing again T, E_i and F_i for the “specialized” images of the generators $T \in \mathfrak{h}$ and E_i, F_i ($i \in I$), the Lie algebra structure of $\mathfrak{g}_{\bar{P}}^{\mathcal{R}}$ is given by (3.6) with the commutator replaced by the (Lie) bracket and the quantum Serre relations by the adjoint actions $\text{ad}(E_i)^{1-a_{ij}}(E_j) = 0$ and $\text{ad}(F_i)^{1-a_{ij}}(F_j) = 0$, whereas the coalgebra structure is determined by

$$\delta(T) = 0, \quad \delta(E_i) = 2 T_i^+ \wedge E_i, \quad \delta(F_i) = 2 T_i^- \wedge F_i$$

For example, if we take $P := DA$, $r := \text{rk}(DA)$ and $\mathcal{R} := (\mathfrak{h}, \Pi, \Pi^\vee)$ a realization of DA , where $\text{rk}(\mathfrak{h}) = 2n - r$ and $T_i^+ = T_i^-$ in Π^\vee , for all $i \in I$, one has that $U_{DA,\hbar}^{\mathcal{R}}(\mathfrak{g})$ is the “quantum double version” of the usual Drinfeld’s QUEA $U_\hbar(\mathfrak{g}_A)$ for the Kac-Moody algebra \mathfrak{g}_A associated with the Cartan matrix A ; in particular, its semiclassical limit is $U(\mathfrak{g}_A^{\text{MD}})$, where $\mathfrak{g}_A^{\text{MD}}$ is the “Manin double version” of \mathfrak{g}_A .

Now take any $\mathbb{k}[[\hbar]]$ -basis $\{H_g\}_{g \in \mathcal{G}}$ of \mathfrak{h} where $|\mathcal{G}| = \text{rk}(\mathfrak{h}) = t$. For any anti-symmetric matrix $\Phi = (\phi_{i,j})_{1 \leq i,j \leq n}$ with entries in $\mathbb{k}[[\hbar]]$ we define

$$\mathfrak{J}_\Phi := \sum_{i,j=1}^n \phi_{jk} H_g \otimes H_k \in \mathfrak{h} \otimes \mathfrak{h} \subseteq U_{P,\hbar}^{\mathcal{R}}(\mathfrak{h}) \otimes U_{P,\hbar}^{\mathcal{R}}(\mathfrak{h})$$

By direct check, one sees that the element

$$\mathcal{F}_\Phi := e^{\hbar 2^{-1} \mathfrak{J}_\Phi} = \exp\left(\hbar 2^{-1} \sum_{g,k=1}^t \phi_{gk} H_g \otimes H_k\right)$$

in $U_{P,\hbar}^{\mathcal{R}}(\mathfrak{h}) \widehat{\otimes} U_{P,\hbar}^{\mathcal{R}}(\mathfrak{h})$ is actually a twist for $U_{P,\hbar}^{\mathcal{R}}(\mathfrak{g})$. For $i \in I$, define the elements $\mathcal{L}_{\Phi,i} := e^{+\hbar 2^{-1} \sum_{g,k=1}^t \alpha_i(H_g) \phi_{gk} H_k}$ and $\mathcal{K}_{\Phi,i} := e^{+\hbar 2^{-1} \sum_{g,k=1}^t \alpha_i(H_g) \phi_{kg} H_k}$. Then, the new coproduct in $(U_{P,\hbar}^{\mathcal{R}}(\mathfrak{g}))^{\mathcal{F}_\Phi}$ is given by

$$\begin{aligned} \Delta^\Phi(E_i) &= E_i \otimes \mathcal{L}_{\Phi,i}^+ + e^{+\hbar T_i^+} \mathcal{K}_{\Phi,i}^+ \otimes E_i & (\forall i \in I) \\ \Delta^\Phi(T) &= T \otimes 1 + 1 \otimes T & (\forall T \in \mathfrak{h}) \\ \Delta^\Phi(F_i) &= F_i \otimes \mathcal{L}_{\Phi,i}^- e^{-\hbar T_i^-} + \mathcal{K}_{\Phi,i}^- \otimes F_i & (\forall i \in I) \end{aligned}$$

Similarly, the “twisted” antipode $\mathcal{S}^{\mathcal{F}_\Phi}$ and the counit $\epsilon^\Phi := \epsilon$ are given by

$$\begin{aligned} \mathcal{S}^\Phi(E_i) &= -e^{-\hbar T_i^+} \mathcal{K}_{\Phi,i}^- E_i \mathcal{L}_{\Phi,i}^- , & \epsilon^\Phi(E_i) &= 0 & (\forall i \in I) \\ \mathcal{S}^\Phi(T) &= -T , & \epsilon^\Phi(T) &= 0 & (\forall T \in \mathfrak{h}) \\ \mathcal{S}^\Phi(F_i) &= -\mathcal{K}_{\Phi,i}^+ F_i \mathcal{L}_{\Phi,i}^+ e^{+\hbar T_i^-} , & \epsilon^\Phi(F_i) &= 0 & (\forall i \in I) \end{aligned}$$

A key feature of this family of quantum groups is that the multiparameter encodes different types of deformations: there exists a multiparameter matrix P_Φ and a realization \mathcal{R}_Φ such that $U_{P_\Phi,\hbar}^{\mathcal{R}_\Phi}(\mathfrak{g}) \cong (U_{P,\hbar}^{\mathcal{R}}(\mathfrak{g}))^{\mathcal{F}_\Phi}$ as topological Hopf algebras. In particular, the class of all FoMpQUEAs of any fixed Cartan type and of fixed rank is stable by toral twist deformations, see [GaGa2, Theorem 5.1.4]. Furthermore, it turns out that, under certain restrictions on the realization, every FoMpQUEA can be realized as a toral twist deformation of the “standard” FoMpQUEA by Drinfeld — see [GaGa2, Theorem 5.1.5] for further details.

With respect to the semiclassical limit, $\overline{\mathfrak{J}}_\Phi := \mathfrak{J}_\Phi \pmod{\hbar}$ is actually a (toral) twist for the Lie bialgebra $\mathfrak{g}_{\overline{P}}^{\overline{\mathfrak{R}}}$. The deformed Lie cobracket is given by the formula

$$\delta^{\overline{\mathfrak{J}}_\Phi}(x) := \delta(x) - \text{ad}_x(\overline{\mathfrak{J}}_\Phi) = \delta(x) - \sum_{g,k=1}^t \overline{\phi_{gk}} ([x, H_g] \otimes H_k + H_g \otimes [x, H_k])$$

— for all $x \in \mathfrak{g}_{\overline{P}}^{\overline{\mathfrak{R}}}$, with $\overline{\phi_{gk}} := \phi_{gk} \pmod{\hbar}$ — that on generators reads

$$\delta^{\overline{\mathfrak{J}}_\Phi}(E_i) = 2T_{\Phi,i}^+ \wedge E_i , \quad \delta^{\overline{\mathfrak{J}}_\Phi}(T) = 0 , \quad \delta^{\overline{\mathfrak{J}}_\Phi}(F_i) = 2T_{\Phi,i}^- \wedge F_i , \quad \forall i \in I, T \in \mathfrak{h}$$

where $T_{\Phi,i}^\pm = T_i^\pm \pm \sum_{g,k=1}^t \overline{\phi_{kg}} \alpha_i(H_g) H_k$ for all $i \in I$.

In conclusion, one may consider the deformation $(\mathfrak{g}_{\overline{P}}^{\overline{\mathfrak{R}}})^{\overline{\mathfrak{J}}_\Phi}$ of $\mathfrak{g}_{\overline{P}}^{\overline{\mathfrak{R}}}$ by the (Lie) twist $\overline{\mathfrak{J}}_\Phi$, as well as the deformation $(U_{P,\hbar}^{\mathcal{R}}(\mathfrak{g}))^{\mathcal{F}_\Phi}$ of $U_{P,\hbar}^{\mathcal{R}}(\mathfrak{g})$ by the (Hopf) twist \mathcal{F}_Φ . By [GaGa2, Theorem 6.2.2], we know that $(U_{P,\hbar}^{\mathcal{R}}(\mathfrak{g}))^{\mathcal{F}_\Phi}$ is a QUEA, whose semiclassical limit is $U((\mathfrak{g}_{\overline{P}}^{\overline{\mathfrak{R}}})^{\overline{\mathfrak{J}}_\Phi})$: indeed, we have $(U_{P,\hbar}^{\mathcal{R}}(\mathfrak{g}))^{\mathcal{F}_\Phi} \cong U_{P_\Phi,\hbar}^{\mathcal{R}_\Phi}(\mathfrak{g})$ and $(\mathfrak{g}_{\overline{P}}^{\overline{\mathfrak{R}}})^{\overline{\mathfrak{J}}_\Phi} \cong \mathfrak{g}_{\overline{P}_\Phi}^{\overline{\mathfrak{R}}_\Phi}$.

3.2. Deformations by 2-cocycle of QFSHA's.

We consider now deformations by 2-cocycle of QFSHA's. Due to the (linear) duality between the notions of QUEA and QFSHA, and similarly for those of twist and 2-cocycle, the outcome we find is nothing but the dual counterpart of Theorem 3.1.2 above (and, consistently, it might be deduced from the latter by duality).

Theorem 3.2.1.

Let $F_{\hbar}[[G]]$ be a QFSHA over the Poisson group G , with tangent Lie bialgebra $\mathfrak{g} = (\mathfrak{g}; [\cdot, \cdot], \delta)$. Let σ be a 2-cocycle for $F_{\hbar}[[G]]$ s.t. $\sigma \equiv \epsilon^{\otimes 2} \left(\text{mod } \hbar \left(F_{\hbar}[[G]]^{\tilde{\otimes} 2} \right)^* \right)$; then $\varsigma := \hbar^{-1} \log_*(\sigma) \in \left(F_{\hbar}[[G]]^{\tilde{\otimes} 2} \right)^*$, where “ \log_* ” is the logarithm with respect to the convolution product, and $\sigma = \exp_*(\hbar\varsigma)$. Last, we set $\varsigma_a := \varsigma - \varsigma_{2,1}$. Then:

(a) ς is antisymmetric, i.e. $\varsigma_{2,1} = -\varsigma$, iff σ is orthogonal, i.e. $\sigma_{2,1} = \sigma^{-1}$;

(b) the element $\overline{\varsigma}_a := \varsigma_a \left(\text{mod } \hbar \left(F_{\hbar}[[G]]^{\tilde{\otimes} 2} \right)^* \right)$ provides a well-defined element $\zeta \in (\mathfrak{g}^* \otimes \mathfrak{g}^*)^* = \mathfrak{g} \otimes \mathfrak{g}$ that is an antisymmetric 2-cocycle for the Lie bialgebra \mathfrak{g}^* ;

(c) letting ζ be as in claim (b), the deformation $(F_{\hbar}[[G]])_{\sigma}$ of $F_{\hbar}[[G]]$ by the 2-cocycle σ is a QFSHA for the formal Poisson group G_{σ} with cotangent Lie bialgebra

$$\text{Lie}(G_{\zeta})^* = (\mathfrak{g}^*)_{\zeta} = (\mathfrak{g}^*; ([\cdot, \cdot]_{*})_{\zeta}, \delta_*)$$

which is the deformation of \mathfrak{g}^* by the 2-cocycle ζ ; in short, $(F_{\hbar}[[G]])_{\sigma} \cong F_{\hbar}[[G_{\zeta}]]$.

Proof. (a) This is obvious, just by construction.

(b) As a first step, we have to prove that $\overline{\varsigma}_a := \varsigma_a \left(\text{mod } \hbar \left(F_{\hbar}[[G]]^{\tilde{\otimes} 2} \right)^* \right)$ determines a uniquely defined element $\zeta \in (\mathfrak{g}^* \otimes \mathfrak{g}^*)^* = \mathfrak{g} \otimes \mathfrak{g}$. Hereafter we realize \mathfrak{g}^* as $\mathfrak{g}^* = \mathfrak{m}/\mathfrak{m}^2$ where $\mathfrak{m} := \text{Ker}(\epsilon_{F[[G]])}$; then we have also

$$\mathfrak{g}^* \otimes \mathfrak{g}^* = (\mathfrak{m}/\mathfrak{m}^2) \otimes (\mathfrak{m}/\mathfrak{m}^2) \cong (\mathfrak{m} \otimes \mathfrak{m}) / (\mathfrak{m} \otimes \mathfrak{m}^2 + \mathfrak{m}^2 \otimes \mathfrak{m})$$

thus in the end we have to prove that the function $\overline{\varsigma}_a := \varsigma_a \left(\text{mod } \hbar \right)$ — defined from $F[[G]]^{\tilde{\otimes}} = F_{\hbar}[[G]]^{\tilde{\otimes}} \left(\text{mod } \hbar \right)$ to \mathbb{k} — does vanish onto $\mathfrak{m} \otimes \mathfrak{m}^2 + \mathfrak{m}^2 \otimes \mathfrak{m}$, hence induces ζ defined onto $(\mathfrak{g}^* \otimes \mathfrak{g}^*)^* = (\mathfrak{m} \otimes \mathfrak{m}) / (\mathfrak{m} \otimes \mathfrak{m}^2 + \mathfrak{m}^2 \otimes \mathfrak{m})$ by the canonical recipe $\zeta(\overline{u \otimes v}) := \overline{\varsigma}_a(u \otimes v)$ for each $u, v \in \mathfrak{m}$. In fact, since ς_a is antisymmetric it is enough to prove that $\overline{\varsigma}_a(\mathfrak{m} \otimes \mathfrak{m}^2) = 0$; in turn, this amounts to showing that — writing ς_a as a bilinear function rather than a morphism on a tensor product module — one has

$$\varsigma_a(a, bc) \equiv 0 \quad \forall a, b, c \in J_{\hbar} := \text{Ker}(\epsilon_{F_{\hbar}[[G]])} \quad (3.7)$$

For the given $a, b, c \in J_{\hbar} := \text{Ker}(\epsilon_{F_{\hbar}[[G]])}$, the 2-cocycle nature of σ gives

$$\sigma(b_{(1)}, c_{(1)}) \sigma(a, b_{(2)} c_{(2)}) = \sigma(a_{(1)}, b_{(1)}) \sigma(a_{(2)} b_{(2)}, c) \quad (3.8)$$

Now we expand σ as (cf. §2.3.1 for notation “ $\mathcal{O}(\hbar^2)$ ”)

$$\sigma = \exp_*(\hbar\zeta) = \epsilon^{\otimes 2} + \hbar\zeta + \mathcal{O}(\hbar^2) = \epsilon^{\otimes 2} + \hbar \sum_{\zeta} \zeta' \otimes \zeta'' + \mathcal{O}(\hbar^2)$$

where we used sort of Sweedler’s-like notation $\zeta = \sum_{\zeta} \zeta' \otimes \zeta''$ to denote ζ ; plugging this into (3.8) and expanding everything out, we end up with

$$\begin{aligned} \epsilon(a)\epsilon(b)\epsilon(c) + \hbar \left(\sum_{\zeta} \zeta'(a)\zeta''(bc) + \epsilon(a) \sum_{\zeta} \zeta'(b)\zeta''(c) \right) + \mathcal{O}(\hbar^2) &= \\ = \epsilon(a)\epsilon(b)\epsilon(c) + \hbar \left(\sum_{\zeta} \zeta'(ab)\zeta''(c) + \sum_{\zeta} \zeta'(a)\zeta''(b)\epsilon(c) \right) + \mathcal{O}(\hbar^2) \end{aligned}$$

which implies — since $\epsilon(a) = 0 = \epsilon(c)$ by assumption — also

$$\hbar \sum_{\zeta} \zeta'(a)\zeta''(bc) + \mathcal{O}(\hbar^2) = \hbar \sum_{\zeta} \zeta'(ab)\zeta''(c) + \mathcal{O}(\hbar^2)$$

whence we argue

$$\zeta(a, bc) = \sum_{\zeta} \zeta'(a)\zeta''(bc) \equiv_{\hbar} \sum_{\zeta} \zeta'(ab)\zeta''(c) = \zeta(ab, c) \quad (3.9)$$

Recall also that $F_{\hbar}[[G]]$ is commutative modulo \hbar , so that $xy \equiv_{\hbar} yx$ for all $x, y \in F_{\hbar}[[G]]$. Using this along with several instances of (3.9) one gets

$$\zeta(a, bc) \equiv_{\hbar} \zeta(ab, c) \equiv_{\hbar} \zeta(ba, c) \equiv_{\hbar} \zeta(b, ac) \equiv_{\hbar} \zeta(b, ca) \equiv_{\hbar} \zeta(bc, a)$$

from which we eventually conclude that

$$\zeta_a(a, bc) := \zeta(a, bc) - \zeta(bc, a) \equiv_{\hbar} \zeta(bc, a) - \zeta(bc, a) = 0 \quad , \quad \text{q.e.d.}$$

As a second step, we note that ζ is antisymmetric, by construction, since ζ_a is.

Third, we need to prove that $\zeta : \mathfrak{g}^* \otimes \mathfrak{g}^* \longrightarrow \mathbb{k}$ satisfies the remaining condition of (2.4), so that it is indeed a 2-cocycle. Now, expanding σ up to order 3, as

$$\sigma = \exp_*(\hbar\zeta) = \epsilon^{\otimes 2} + \hbar\zeta + \hbar^2 \zeta^{*2}/2 + \mathcal{O}(\hbar^3)$$

and plugging this into (3.8), we find, for all $a, b, c \in J_{\hbar} := \text{Ker}(\epsilon_{F_{\hbar}[[G]]})$ again,

$$\begin{aligned} & \left(\epsilon(b_{(1)})\epsilon(c_{(1)}) + \hbar\zeta(b_{(1)}, c_{(1)}) + \hbar^2 \zeta(b_{(1)(1)}, c_{(1)(1)}) \zeta(b_{(1)(2)}, c_{(1)(2)}) / 2 + \mathcal{O}(\hbar^3) \right) \cdot \\ & \cdot \left(\epsilon(a)\epsilon(b_{(2)})\epsilon(c_{(2)}) + \hbar\zeta(a, b_{(2)}c_{(2)}) + \right. \\ & \quad \left. + \hbar^2 \zeta(a_{(1)}, b_{(2)(1)}c_{(2)(1)}) \zeta(a_{(2)}, b_{(2)(2)}c_{(2)(2)}) / 2 + \mathcal{O}(\hbar^3) \right) = \\ & = \left(\epsilon(a_{(1)})\epsilon(b_{(1)}) + \hbar\zeta(a_{(1)}, b_{(1)}) + \hbar^2 \zeta(a_{(1)(1)}, b_{(1)(1)}) \zeta(a_{(1)(2)}, b_{(1)(2)}) / 2 + \mathcal{O}(\hbar^3) \right) \cdot \\ & \cdot \left(\epsilon(a_{(2)})\epsilon(b_{(2)})\epsilon(c) + \hbar\zeta(a_{(2)}b_{(2)}, c) + \right. \\ & \quad \left. + \hbar^2 \zeta(a_{(2)(1)}b_{(2)(1)}, c) \zeta(a_{(2)(2)}b_{(2)(2)}, c) / 2 + \mathcal{O}(\hbar^3) \right) \end{aligned}$$

and then performing multiplication and truncating at order 3 we get

$$\begin{aligned} & \epsilon(abc) + \hbar(\varsigma(a, bc) + \epsilon(a)\varsigma(b, c)) + \\ & + \hbar^2\left(\varsigma^{*2}(a, bc)/2 + \varsigma(b_{(1)}, c_{(1)})\varsigma(a, b_{(2)}c_{(2)}) + \epsilon(a)\varsigma^{*2}(b, c)/2\right) + \mathcal{O}(\hbar^3) = \\ = & \epsilon(abc) + \hbar(\varsigma(ab, c) + \varsigma(a, b)\epsilon(c)) + \\ & + \hbar^2\left(\varsigma^{*2}(a, b)\epsilon(c)/2 + \varsigma(a_{(1)}, b_{(1)})\varsigma(a_{(2)}b_{(2)}, c) + \varsigma^{*2}(ab, c)/2\right) + \mathcal{O}(\hbar^3) \end{aligned}$$

which in turn — since $\epsilon(a) = 0 = \epsilon(c)$ by assumption — simplifies into

$$\begin{aligned} & \varsigma(a, bc) + \hbar\left(\varsigma^{*2}(a, bc)/2 + \varsigma(b_{(1)}, c_{(1)})\varsigma(a, b_{(2)}c_{(2)})\right) + \mathcal{O}(\hbar^2) = \\ & = \varsigma(ab, c) + \hbar\left(\varsigma(a_{(1)}, b_{(1)})\varsigma(a_{(2)}b_{(2)}, c) + \varsigma^{*2}(ab, c)/2\right) + \mathcal{O}(\hbar^2) \end{aligned}$$

that we eventually we re-write as

$$\begin{aligned} & \varsigma(a, bc) - \varsigma(ab, c) + \\ & + \hbar\left(\varsigma^{*2}(a, bc)/2 - \varsigma^{*2}(ab, c)/2 + \right. \\ & \left. + \varsigma(b_{(1)}, c_{(1)})\varsigma(a, b_{(2)}c_{(2)}) - \varsigma(a_{(1)}, b_{(1)})\varsigma(a_{(2)}b_{(2)}, c)\right) \equiv_{\hbar^2} 0 \end{aligned} \quad (3.10)$$

Now let $\mathbb{k}[\mathbb{S}_3]$ act onto $F_{\hbar}[[G]]^{\otimes 3}$ and consider in particular the action of the anti-symmetrizer $Alt_3 := (\text{id} - (12) - (23) - (31) + (123) + (321))$ onto the equation in (3.10), which yields a new equation: denoting equation (3.10) by $\otimes = 0$, we will write $Alt_3(\otimes) = 0$ for the newly found equation. To see the latter explicitly, we compute the left-hand member $Alt_3(\otimes)$, starting by computing the action of Alt_3 onto the first line in (3.10): concretely, we find

$$\begin{aligned} & Alt_3(1^{st} \text{ line in (3.10)}) = \\ & = \varsigma(a, bc) - \varsigma(b, ac) - \varsigma(a, cb) - \varsigma(c, ba) + \varsigma(c, ab) + \varsigma(b, ca) - \\ & - \varsigma(ab, c) + \varsigma(ba, c) + \varsigma(ac, b) + \varsigma(cb, a) - \varsigma(ca, b) - \varsigma(bc, a) = \\ & = \varsigma_a(a, [b, c]) + \varsigma_a(b, [c, a]) + \varsigma_a(c, [a, b]) = \varsigma_a(a, [b, c]) + \text{c.p.} \end{aligned} \quad (3.11)$$

where standard notation $[u, v] := uv - vu$ is used to denote the usual commutator. Modulo \hbar , such a commutator in $F_{\hbar}[[G]]$ yields the Poisson bracket in $F[[G]]$, hence we can write $[u, v] = \hbar\{\bar{u}, \bar{v}\}'$ where we use notation $\bar{z} := (z \bmod \hbar F_{\hbar}[[G]])$ for each $z \in F_{\hbar}[[G]]$ and $f' :=$ some lift in $F_{\hbar}[[G]]$ of any $f \in F[[G]]$, i.e. $\bar{f}' = f$; note that f' is only defined up to $\hbar^2 F_{\hbar}[[G]]$, yet that is enough for our purposes. Thanks to this, (3.11) turns into

$$Alt_3(1^{st} \text{ line in (3.10)}) = \hbar\left(\varsigma_a\left(a, \{\bar{b}, \bar{c}\}'\right) + \text{c.p.}\right) \quad (3.12)$$

Looking at (3.10), this entails that the \hbar -adic expansion of $Alt_3(\otimes)$ has zero term at order 0, while at order 1 it also has a contribution coming from (3.12).

Now we go and compute the contribution to $Alt_3(\otimes)$ issuing from the third line in (3.10). Again, direct calculations give

$$Alt_3(3^{rd} \text{ line in (3.10)}) = \varsigma_a(a_{(1)}, b_{(1)})\varsigma_a(c, a_{(2)}b_{(2)}) + \text{c.p.} \quad (3.13)$$

Since one always has $x = \epsilon(x) + x_+$ with $x_+ := (x - \epsilon(x)) \in \text{Ker}(\epsilon)$, applying this to each element $x \in \{a, b, c\}$ occurring in (3.13), then expanding everything and taking into account that $\varsigma_a(J_{\hbar}, J_{\hbar}^2) \stackrel{\equiv}{\hbar} 0 \stackrel{\equiv}{\hbar} \varsigma_a(J_{\hbar}^2, J_{\hbar})$ — cf. (3.7) — we obtain

$$\begin{aligned} \text{Alt}_3(3^{\text{rd}} \text{ line in (3.10)}) &\stackrel{\equiv}{\hbar} \varsigma_a(a, b_{(1)}) \varsigma_a(c, b_{(2)}) + \varsigma_a(a_{(1)}, b) \varsigma_a(c, a_{(2)}) + \text{c.p.} \\ &= \varsigma_a(a, b_{(1)}^{\wedge}) \varsigma_a(c, b_{(2)}^{\wedge}) + \text{c.p.} \end{aligned}$$

where we make use of short-hand notation $x_{(1)}^{\wedge} \otimes x_{(2)}^{\wedge} := x_{(1)} \otimes x_{(2)} - x_{(2)} \otimes x_{(1)}$.

Finally, we go and compute the contribution to $\text{Alt}_3(\otimes)$ issuing from the second line in (3.10). Dropping the coefficients \hbar and $1/2$ we find the following:

$$\begin{aligned} \text{Alt}_3(2^{\text{nd}} \text{ line in (3.10)}) &= \text{Alt}_3(\zeta^{*2}(a, bc) - \zeta^{*2}(b, ac)) = \\ &= \zeta^{*2}(a, bc) - \zeta^{*2}(b, ac) - \zeta^{*2}(a, cb) - \zeta^{*2}(c, ba) + \zeta^{*2}(c, ab) + \zeta^{*2}(b, ca) - \\ &\quad - \zeta^{*2}(ab, c) + \zeta^{*2}(ba, c) + \zeta^{*2}(ac, b) + \zeta^{*2}(cb, a) - \zeta^{*2}(ca, b) - \zeta^{*2}(bc, a) = \\ &= \zeta^{*2}(a, [b, c]) + \zeta^{*2}(b, [c, a]) + \zeta^{*2}(c, [a, b]) - \\ &\quad - \zeta^{*2}([b, c], a) - \zeta^{*2}([c, a], b) - \zeta^{*2}([a, b], c) \end{aligned}$$

which in turn implies $\text{Alt}_3(\zeta^{*2}(a, bc) - \zeta^{*2}(b, ac)) = \mathcal{O}(\hbar)$ — since $[u, v] = \mathcal{O}(\hbar)$ for all $u, v \in F_{\hbar}[[G]]$. The outcome then is that the contribution to $\text{Alt}_3(\otimes)$ given by the second line in (3.10) is trivial modulo \hbar^2 .

Summing up, the outcome of the previous analysis is that

$$\varsigma_a(a, \{\bar{b}, \bar{c}\}') + \text{c.p.} + \varsigma_a(a, b_{(1)}^{\wedge}) \varsigma_a(c, b_{(2)}^{\wedge}) + \text{c.p.} \stackrel{\equiv}{\hbar} 0$$

Taking the latter modulo $\hbar F_{\hbar}[[G]]$ we find

$$\varsigma_a(\bar{a}, \{\bar{b}, \bar{c}\}) + \text{c.p.} + \varsigma_a(\bar{a}, \bar{b}_{(1)}^{\wedge}) \varsigma_a(\bar{c}, \bar{b}_{(2)}^{\wedge}) + \text{c.p.} = 0$$

for the elements $\bar{a}, \bar{b}, \bar{c} \in F[[G]]$.

Now recall that for $x \in J_{\hbar}$ with $\bar{x} := (x \bmod \hbar F_{\hbar}[[G]])$ and $\mathbf{x} := (\bar{x} \bmod \mathfrak{m}^2)$ we have $\delta(\mathbf{x}) := \mathbf{x}_{(1)}^{\wedge} \otimes \mathbf{x}_{(2)}^{\wedge}$ for the induced Lie cobracket of $\mathfrak{g}^* = \mathfrak{m}/\mathfrak{m}^2$ computed on \mathbf{x} , by definition; this means that, using our previously established notation $\delta(\mathbf{x}) := \mathbf{x}_{[1]} \otimes \mathbf{x}_{[2]}$, the last formula above yields

$$\zeta(a, [b, c]) + \text{c.p.} + \zeta(a, b_{[1]}) \zeta(c, b_{[2]}) + \text{c.p.} = 0 \quad (3.14)$$

Finally, the antisymmetry of ζ gives $\zeta(a, [b, c]) + \text{c.p.} = -\zeta([a, b], c) + \text{c.p.}$, while a straightforward check shows that $\zeta(a, b_{[1]}) \zeta(c, b_{[2]}) + \text{c.p.} = -[[\zeta, \zeta]]_*$. Therefore, (3.14) is equivalent to

$$\zeta([a, b], c) + \text{c.p.} + [[\zeta, \zeta]]_* = 0$$

which means that ζ is indeed a (strong type of) 2-cocycle for \mathfrak{g}^* , q.e.d.

(c) Let us consider the deformed algebra $(F_{\hbar}[[G]])_{\sigma}$, which coincides with $F_{\hbar}[[G]]$ as a $\mathbb{k}[[\hbar]]$ -module but is endowed with the deformed multiplication “ \cdot_{σ} ” defined by

$$a \cdot_{\sigma} b := \sigma(a_{(1)}, b_{(1)}) a_{(2)} b_{(2)} \sigma^{-1}(a_{(3)}, b_{(3)}) \quad \forall a, b \in F_{\hbar}[[G]] \quad (3.15)$$

As σ is of the form $\sigma = \exp_*(\hbar\varsigma)$, in particular it is trivial modulo \hbar , it follows from (3.15) that the deformed multiplication “ \cdot_σ ” coincides with the old one modulo \hbar , that is $a \cdot_\sigma b \equiv ab \pmod{\hbar F_\hbar[[G]]}$. Therefore, $(F_\hbar[[G]])_\sigma$ is again commutative modulo \hbar , hence it is (again) a QFSHA, as claimed, say $(F_\hbar[[G]])_\sigma = F_\hbar[[G(\sigma)]]$. Then, in order to prove that the newly found Poisson (formal) group $G(\sigma)$ is indeed G_ζ as claimed it is enough to show that the Lie bracket induced in $\mathfrak{m}_\sigma/\mathfrak{m}_\sigma^2$ — where $\mathfrak{m}_\sigma := \text{Ker}(\epsilon_{F[[G(\sigma)]]})$ — is indeed $[\cdot, \cdot]_\zeta$.

Let us take $a, b \in \mathfrak{m}_\sigma/\mathfrak{m}_\sigma^2$; then we can pick $a, b \in J_\hbar := \text{Ker}(\epsilon_{F_\hbar[[G]]})$ such that $a = a \pmod{(\hbar J_\hbar + J_\hbar^2)}$ and $b = b \pmod{(\hbar J_\hbar + J_\hbar^2)}$. Now, using the expansion $\sigma = \exp_*(\hbar\varsigma) = \epsilon^{\otimes 2} + \hbar\varsigma + \mathcal{O}(\hbar^2)$, formula (3.15) turns into

$$\begin{aligned} a \cdot_\sigma b &= \\ &= (\epsilon(a_{(1)})\epsilon(b_{(1)}) + \hbar\varsigma(a_{(1)}, b_{(1)})) a_{(2)} b_{(2)} (\epsilon(a_{(3)})\epsilon(b_{(3)}) + \hbar\varsigma(a_{(3)}, b_{(3)})) + \mathcal{O}(\hbar^2) = \\ &= \epsilon(a_{(1)}) \epsilon(b_{(1)}) a_{(2)} b_{(2)} \epsilon(a_{(3)}) \epsilon(b_{(3)}) + \\ &+ \hbar \left(\varsigma(a_{(1)}, b_{(1)}) a_{(2)} b_{(2)} \epsilon(a_{(3)}) \epsilon(b_{(3)}) - \epsilon(a_{(1)}) \epsilon(b_{(1)}) a_{(2)} b_{(2)} \varsigma(a_{(3)}, b_{(3)}) \right) + \mathcal{O}(\hbar^2) = \\ &= ab + \hbar \left(\varsigma(a_{(1)}, b_{(1)}) a_{(2)} b_{(2)} - a_{(1)} b_{(1)} \varsigma(a_{(2)}, b_{(2)}) \right) + \mathcal{O}(\hbar^2) \end{aligned}$$

where we took into account coassociativity and counitality properties. Therefore, using “ $[\cdot, \cdot]_\sigma$ ” and “ $[\cdot, \cdot]$ ” to denote the commutator with respect to the new (deformed) and the old (undeformed) multiplication, we also have

$$\begin{aligned} [a, b]_\sigma &:= a \cdot_\sigma b - b \cdot_\sigma a = \\ &= ab + \hbar \left(\varsigma(a_{(1)}, b_{(1)}) a_{(2)} b_{(2)} - a_{(1)} b_{(1)} \varsigma(a_{(2)}, b_{(2)}) \right) + \mathcal{O}(\hbar^2) - \\ &\quad - ba - \hbar \left(\varsigma(b_{(1)}, a_{(1)}) b_{(2)} a_{(2)} - b_{(1)} a_{(1)} \varsigma(b_{(2)}, a_{(2)}) \right) + \mathcal{O}(\hbar^2) = \\ &= ab - ba + \hbar \left(-\varsigma(a_{(2)}, b_{(2)}) a_{(1)} b_{(1)} + \varsigma(b_{(2)}, a_{(2)}) b_{(1)} a_{(1)} - \right. \\ &\quad \left. - \varsigma(b_{(1)}, a_{(1)}) b_{(2)} a_{(2)} + \varsigma(a_{(1)}, b_{(1)}) a_{(2)} b_{(2)} \right) + \mathcal{O}(\hbar^2) = \\ &= [a, b] + \hbar \left(-\varsigma_a(a_{(2)}, b_{(2)}) a_{(1)} b_{(1)} - \varsigma_a(b_{(1)}, a_{(1)}) b_{(2)} a_{(2)} \right) + \mathcal{O}(\hbar^2) \end{aligned}$$

where for the last step we used the fact that $a_{(s)} b_{(s)} \equiv_{\hbar} b_{(s)} a_{(s)}$.

Recall that $[a, b] = \hbar \{\bar{a}, \bar{b}\}'$, where hereafter we write $\bar{x} := x \pmod{\hbar J_\hbar}$ and f' to denote any lift in J_\hbar of some given f in J_\hbar , as we did before; similarly, we have $[a, b]_\sigma = \hbar \{\bar{a}, \bar{b}\}'_\sigma$. Then modulo \hbar our previous computations give

$$\{\bar{a}, \bar{b}\}'_\sigma = \{\bar{a}, \bar{b}\}' - \bar{\varsigma}_a(\bar{a}_{(2)}, \bar{b}_{(2)}) \bar{a}_{(1)} \bar{b}_{(1)} - \bar{\varsigma}_a(\bar{b}_{(1)}, \bar{a}_{(1)}) \bar{b}_{(2)} \bar{a}_{(2)} \quad (3.16)$$

For each $x \in \{\bar{a}_{(s)}, \bar{b}_{(s)} \mid s = 1, 2\}$ we have $x = \epsilon(x) + x^+$ with $x^+ := (x - \epsilon(x)) \in J_\hbar$. Using such expansions in either factor of the products $\bar{a}_{(s)} \bar{b}_{(s)}$ and $\bar{b}_{(s)} \bar{a}_{(s)}$ occurring within (3.16), and noting that $\bar{a}_{(s)}^+ \bar{b}_{(s)}^+ \equiv_{\mathfrak{m}^2} 0 \equiv_{\mathfrak{m}^2} \bar{a}_{(s)}^+ \bar{b}_{(s)}^+$, we get an

equivalence modulo $\mathfrak{m}^2 = \mathfrak{m}_\sigma^2$ (noting that $\mathfrak{m} = \mathfrak{m}_\sigma$ as \mathbb{k} -modules), namely

$$\begin{aligned}
 \{\bar{a}, \bar{b}\}'_\sigma &= \{\bar{a}, \bar{b}\} - \bar{\zeta}_a(\bar{a}_{(2)}, \bar{b}_{(2)}) \bar{a}_{(1)} \bar{b}_{(1)} - \bar{\zeta}_a(\bar{b}_{(1)}, \bar{a}_{(1)}) \bar{b}_{(2)} \bar{a}_{(2)} \stackrel{\equiv}{=} \\
 &\stackrel{\equiv}{=} \{\bar{a}, \bar{b}\} - \bar{\zeta}_a(\bar{a}_{(2)}, \bar{b}_{(2)}) \bar{a}_{(1)} \epsilon(\bar{b}_{(1)}) - \bar{\zeta}_a(\bar{a}_{(2)}, \bar{b}_{(2)}) \epsilon(\bar{a}_{(1)}) \bar{b}_{(1)} - \\
 &\quad - \bar{\zeta}_a(\bar{b}_{(1)}, \bar{a}_{(1)}) \bar{b}_{(2)} \epsilon(\bar{a}_{(2)}) - \bar{\zeta}_a(\bar{b}_{(1)}, \bar{a}_{(1)}) \epsilon(\bar{b}_{(2)}) \bar{a}_{(2)} = \\
 &= \{\bar{a}, \bar{b}\} - \bar{\zeta}_a(\bar{a}_{(2)}, \bar{b}) \bar{a}_{(1)} - \bar{\zeta}_a(\bar{a}, \bar{b}_{(2)}) \bar{b}_{(1)} - \bar{\zeta}_a(\bar{b}_{(1)}, \bar{a}) \bar{b}_{(2)} - \bar{\zeta}_a(\bar{b}, \bar{a}_{(1)}) \bar{a}_{(2)} = \\
 &= \{\bar{a}, \bar{b}\} - \bar{\zeta}_a(\bar{a}_{(2)}, \bar{b}) \bar{a}_{(1)} + \bar{\zeta}_a(\bar{b}_{(2)}, \bar{a}) \bar{b}_{(1)} - \bar{\zeta}_a(\bar{b}_{(1)}, \bar{a}) \bar{b}_{(2)} + \bar{\zeta}_a(\bar{a}_{(1)}, \bar{b}) \bar{a}_{(2)} = \\
 &= \{\bar{a}, \bar{b}\} - (\bar{\zeta}_a(\bar{a}_{(2)}, \bar{b}) \bar{a}_{(1)} - \bar{\zeta}_a(\bar{a}_{(1)}, \bar{b}) \bar{a}_{(2)}) - (\bar{\zeta}_a(\bar{b}_{(1)}, \bar{a}) \bar{b}_{(2)} - \bar{\zeta}_a(\bar{b}_{(2)}, \bar{a}) \bar{b}_{(1)})
 \end{aligned}$$

where the element in last line actually belongs to $\mathfrak{m} = \mathfrak{m}_\sigma$. When we reduce all this modulo $\mathfrak{m}^2 = \mathfrak{m}_\sigma^2$, and recalling the definition of the Lie bracket (for either Lie algebra structure) as being induced by the Poisson bracket, and that of the Lie cobracket in $\mathfrak{m}/\mathfrak{m}^2 = \mathfrak{m}_\sigma/\mathfrak{m}_\sigma^2$ — which coincide as Lie coalgebras — as being induced by $\Delta - \Delta^{\text{op}}$, we eventually end up with

$$[a, b]_{(\sigma)} = [a, b]_* - \zeta(a_{[2]}, b) a_{[1]} - \zeta(b_{[1]}, a) b_{[2]} =: ([a, b]_*)_\zeta$$

thus(cf. Definition 2.5) the Lie bracket we were looking for is just $([,]_*)_\zeta$. \square

Observation 3.2.2. We would better point out that the 2-cocycles σ considered in Theorem 3.2.1 above are those of “trivial-modulo- \hbar -type”, in that they are the identity modulo \hbar . With this assumption, deforming $F_\hbar[[G]]$ by such a σ does not affect the Hopf structure of the semiclassical limit; in particular, it still reads as $F[[\tilde{G}]]$, with \tilde{G} being the same formal group as G but with a different Poisson structure. A more general 2-cocycle might be “unfit”, in that the deformed Hopf algebra $(F_\hbar[[G]])_\sigma$ may no longer be a QFSHA, in general.

Example 3.2.3. Let $G := GL_n(\mathbb{k})$ be the general linear group over \mathbb{k} , and $\mathfrak{g} := \mathfrak{gl}_n(\mathbb{k})$ its tangent Lie algebra. It is well-known — cf. [Dr], [CP] — that a quantization of \mathfrak{g} is provided by the QUEA $U_\hbar(\mathfrak{g}) = U_\hbar(\mathfrak{gl}_n(\mathbb{k}))$ defined as follows: it is the unital, associative, \hbar -adically complete $\mathbb{k}[[\hbar]]$ -algebra with generators

$$F_1, F_2, \dots, F_{n-1}, \Gamma_1, \Gamma_2, \dots, \Gamma_{n-1}, \Gamma_n, E_1, E_2, \dots, E_{n-1}$$

and relations (for all $i, j \in \{1, \dots, n-1\}$, $k, \ell \in \{1, \dots, n\}$)

$$\begin{aligned}
 [\Gamma_k, \Gamma_\ell] &= 0, & [\Gamma_k, F_j] &= -\delta_{k,j} F_j, & [\Gamma_k, E_j] &= +\delta_{k,j} E_j \\
 [E_i, F_j] &= \delta_{i,j} \frac{e^{\hbar(\Gamma_i - \Gamma_{i+1})} - e^{\hbar(\Gamma_{i+1} - \Gamma_i)}}{e^{+\hbar} - e^{-\hbar}} \\
 [E_i, E_j] &= 0, & [F_i, F_j] &= 0 & \forall i, j : |i - j| > 1 \\
 E_i^2 E_j - (q + q^{-1}) E_i E_j E_i + E_j E_i^2 &= 0 & \forall i, j : |i - j| = 1 \\
 F_i^2 F_j - (q + q^{-1}) F_i F_j F_i + F_j F_i^2 &= 0 & \forall i, j : |i - j| = 1.
 \end{aligned}$$

where $[X, Y] := XY - YX$. The (topological) Hopf algebra structure is given by

$$\begin{aligned} \Delta(F_i) &= F_i \otimes e^{\hbar(\Gamma_{i+1} - \Gamma_i)} + 1 \otimes F_i, & S(F_i) &= -F_i e^{\hbar(\Gamma_i - \Gamma_{i+1})}, & \epsilon(F_i) &= 0 \\ \Delta(\Gamma_k) &= \Gamma_k \otimes 1 + 1 \otimes \Gamma_k, & S(\Gamma_k) &= -\Gamma_k, & \epsilon(\Gamma_k) &= 0 \\ \Delta(E_i) &= E_i \otimes 1 + e^{\hbar(\Gamma_i - \Gamma_{i+1})} \otimes E_i, & S(E_i) &= -e^{\hbar(\Gamma_{i+1} - \Gamma_i)} E_i, & \epsilon(E_i) &= 0 \end{aligned}$$

It is also well-known — cf. [Dr], [CP] — that a quantization of $G := GL_n(\mathbb{k})$ is provided by the QFSHA $F_\hbar[[G]] = F_\hbar[[GL_n(\mathbb{k})]]$ defined as follows: it is the unital, associative, I_\hbar -adically complete $\mathbb{k}[[\hbar]]$ -algebra generated by the elements of the set $\{x_{ij} \mid i, j = 1, \dots, n+1\}$ arranged in a q -matrix, with $q := \exp(\hbar)$, with I_\hbar being the ideal generated by $\{\hbar, x_{1,1}, \dots, x_{n,n}\}$; this is a quick way to say that the given generators obey the relations

$$\begin{aligned} x_{ij} x_{ik} &= q x_{ik} x_{ij}, & x_{ik} x_{hk} &= q x_{hk} x_{ik} & \forall j < k, i < h \\ x_{il} x_{jk} &= x_{jk} x_{il}, & x_{ik} x_{jl} - x_{jl} x_{ik} &= (q - q^{-1}) x_{il} x_{jk} & \forall i < j, k < l \end{aligned}$$

whereas the comultiplication Δ , the counit ϵ , and the antipode S are given by matrix formulation

$$\begin{aligned} \Delta\left(\left(x_{ij}\right)_{i=1,\dots,n}^{j=1,\dots,n}\right) &:= \left(x_{ij}\right)_{i=1,\dots,n}^{j=1,\dots,n} \otimes \left(x_{ij}\right)_{i=1,\dots,n}^{j=1,\dots,n} \\ \epsilon\left(\left(x_{ij}\right)_{i=1,\dots,n}^{j=1,\dots,n}\right) &:= \left(\delta_{ij}\right)_{i=1,\dots,n}^{j=1,\dots,n}, & S\left(\left(x_{ij}\right)_{i=1,\dots,n}^{j=1,\dots,n}\right) &:= \left(\left(x_{ij}\right)_{i=1,\dots,n}^{j=1,\dots,n}\right)^{-1} \end{aligned}$$

which in down-to-earth terms read, for all $i, j = 1, \dots, n$,

$$\Delta(x_{ij}) = \sum_{k=1}^n x_{ik} \otimes x_{kj}, \quad \epsilon(x_{ij}) = \delta_{ij}, \quad S(x_{ij}) = (-q)^{j-i} D_q\left(\left(x_{hk}\right)_{h \neq j}^{k \neq i}\right)$$

where D_q denotes the so-called *quantum determinant*, defined on any q -matrix of (square) size ℓ by

$$D_q\left(\left(x_{ij}\right)_{i=1,\dots,\ell}^{j=1,\dots,\ell}\right) := \sum_{\sigma \in S_\ell} (-q)^{l(\sigma)} x_{1,\sigma(1)} x_{2,\sigma(2)} \cdots x_{\ell,\sigma(\ell)}$$

We have also explicit identifications $F_\hbar[[G]] = U_\hbar(\mathfrak{g})^*$ as well as $U_\hbar(\mathfrak{g}) = F_\hbar[[G]]^*$, which can be described via the Hopf pairing $\langle \cdot, \cdot \rangle : F_\hbar[[G]] \times U_\hbar(\mathfrak{g}) \longrightarrow \mathbb{k}[[\hbar]]$ uniquely given by the following values on generators:

$$\langle x_{i,j}, \Gamma_k \rangle = \delta_{i,j} \delta_{i,k}, \quad \langle x_{i,j}, E_t \rangle = \delta_{i+1,j} \delta_{i,t}, \quad \langle x_{i,j}, F_t \rangle = \delta_{i,j+1} \delta_{t,j} \quad (3.17)$$

Now consider in $U_\hbar(\mathfrak{g}) \widehat{\otimes} U_\hbar(\mathfrak{g})$ the element $\mathcal{F} := \exp\left(\hbar 2^{-1} \sum_{k,\ell=1}^n \phi_{t,k} \Gamma_t \otimes \Gamma_k\right)$ that is a twist for $U_\hbar(\mathfrak{g})$ — just as in Example 3.1.4, it trivially follows from the fact that the Γ_t 's are primitive. By Proposition 2.2.7(a), we can see this \mathcal{F} as a 2-cocycle $\sigma_{\mathcal{F}}$ for $U_\hbar(\mathfrak{g})^* = F_\hbar[[G]]$, simply given by *evaluation at \mathcal{F}* , namely

$$\sigma_{\mathcal{F}} : F_\hbar[[G]] \times F_\hbar[[G]] \longrightarrow \mathbb{k}[[\hbar]] \quad , \quad (\varphi, \psi) \mapsto \langle \varphi \otimes \psi, \mathcal{F} \rangle \quad (3.18)$$

Now, from (3.18) and (3.17), direct calculation gives

$$\begin{aligned} \sigma_{\mathcal{F}}(x_{i,r}, x_{\ell,h}) &= \langle x_{i,r} \otimes x_{\ell,h}, \mathcal{F} \rangle = \\ &= \sum_{m=0}^{+\infty} \frac{\hbar^m 2^{-m}}{m!} \left\langle x_{i,r} \otimes x_{\ell,h}, \left(\sum_{t,k=1}^n \phi_{t,k} \Gamma_t \otimes \Gamma_k\right)^m \right\rangle = \\ &= \sum_{m=0}^{+\infty} \frac{\hbar^m 2^{-m}}{m!} \left\langle \Delta^{(m-1)}(x_{i,r} \otimes x_{\ell,h}), \left(\sum_{t,k=1}^n \phi_{t,k} \Gamma_t \otimes \Gamma_k\right)^{\otimes m} \right\rangle \end{aligned}$$

Let us stop and consider $\left\langle \Delta^{(m-1)}(x_{i,r} \otimes x_{\ell,h}), \left(\sum_{t,k=1}^n \phi_{t,k} \Gamma_t \otimes \Gamma_k \right)^{\otimes m} \right\rangle$. Definitions give

$$\begin{aligned} & \left\langle \Delta^{(m-1)}(x_{i,r} \otimes x_{\ell,h}), \left(\sum_{t,k=1}^n \phi_{t,k} \Gamma_t \otimes \Gamma_k \right)^{\otimes m} \right\rangle = \\ &= \sum_{\substack{s_1, \dots, s_{m-1}=1 \\ e_1, \dots, e_{m-1}=1}}^n \left\langle x_{i, s_1} \otimes x_{\ell, e_1} \otimes \dots \otimes x_{s_{m-1}, r} \otimes x_{e_{m-1}, h}, \left(\sum_{t,k=1}^n \phi_{t,k} \Gamma_t \otimes \Gamma_k \right)^{\otimes m} \right\rangle = \\ &= \sum_{\substack{s_1, \dots, s_{m-1}=1 \\ e_1, \dots, e_{m-1}=1}}^n \prod_{c=1}^m \left\langle x_{s_{c-1}, s_c} \otimes x_{e_{c-1}, e_c}, \sum_{t,k=1}^n \phi_{t,k} \Gamma_t \otimes \Gamma_k \right\rangle = \\ &= \sum_{\substack{s_1, \dots, s_{m-1}=1 \\ e_1, \dots, e_{m-1}=1}}^n \prod_{c=1}^m \sum_{t,k=1}^n \phi_{t,k} \langle x_{s_{c-1}, s_c}, \Gamma_t \rangle \langle x_{e_{c-1}, e_c}, \Gamma_k \rangle \end{aligned}$$

where we set $s_0 := i$, $s_m := r$, $e_0 := \ell$, $e_m := h$. Now, the formulas in (3.17) guarantee that $\langle x_{s_{c-1}, s_c}, \Gamma_t \rangle \langle x_{e_{c-1}, e_c}, \Gamma_k \rangle = 0$ whenever $s_{c-1} \neq s_c$ or $e_{c-1} \neq e_c$; therefore, from the previous computation one eventually gets

$$\begin{aligned} \sigma_{\mathcal{F}}(x_{i,r}, x_{\ell,h}) &= \\ &= \delta_{i,r} \delta_{\ell,h} \sum_{m=0}^{+\infty} \frac{\hbar^m 2^{-m}}{m!} \left\langle \Delta^{(m-1)}(x_{i,r} \otimes x_{\ell,h}), \left(\sum_{t,k=1}^n \phi_{t,k} \Gamma_t \otimes \Gamma_k \right)^{\otimes m} \right\rangle = \\ &= \delta_{i,r} \delta_{\ell,h} \sum_{m=0}^{+\infty} \frac{\hbar^m 2^{-m}}{m!} \left(\sum_{t,k=1}^n \phi_{t,k} \langle x_{i,i}, \Gamma_t \rangle \langle x_{\ell,\ell}, \Gamma_k \rangle \right)^m = \\ &= \delta_{i,r} \delta_{\ell,h} \sum_{m=0}^{+\infty} \frac{\hbar^m 2^{-m}}{m!} \left(\sum_{t,k=1}^n \phi_{t,k} \delta_{i,t} \delta_{\ell,k} \right)^m = \\ &= \delta_{i,r} \delta_{\ell,h} \sum_{m=0}^{+\infty} \frac{\hbar^m 2^{-m}}{m!} (\phi_{i,\ell})^m = \delta_{i,r} \delta_{\ell,h} \exp(\hbar 2^{-1} \phi_{i,\ell}) = \delta_{i,r} \delta_{\ell,h} e^{\hbar \phi_{i,\ell}/2} \end{aligned}$$

in short

$$\sigma_{\mathcal{F}}(x_{i,r}, x_{\ell,h}) = \delta_{i,r} \delta_{\ell,h} e^{\hbar \phi_{i,\ell}/2} \quad \forall i, r, \ell, h \in \{1, \dots, n\} \quad (3.19)$$

Using this formula, the deformed product in $F_{\hbar}[[G]]_{\sigma_{\mathcal{F}}}$ can be described as follows:

$$\begin{aligned} x_{i,j} \dot{\sigma}_{\mathcal{F}} x_{\ell,t} &:= \sigma_{\mathcal{F}}((x_{i,j})_{(1)}, (x_{\ell,t})_{(1)}) (x_{i,j})_{(2)} (x_{\ell,t})_{(2)} \sigma_{\mathcal{F}}^{-1}((x_{i,j})_{(3)}, (x_{\ell,t})_{(3)}) = \\ &= \sum_{r,s,h,k=1}^n \sigma_{\mathcal{F}}(x_{i,r}, x_{\ell,h}) x_{r,s} x_{h,k} \sigma_{\mathcal{F}}^{-1}(x_{s,j}, x_{k,t}) = \\ &= \sigma_{\mathcal{F}}(x_{i,i}, x_{\ell,\ell}) x_{i,j} x_{\ell,t} \sigma_{\mathcal{F}}^{-1}(x_{j,j}, x_{t,t}) = e^{\hbar(\phi_{i,\ell} - \phi_{j,t})/2} x_{i,j} x_{\ell,t} \end{aligned}$$

— where we used (3.19); in short, we get

$$x_{i,j} \dot{\sigma}_{\mathcal{F}} x_{\ell,t} = e^{\hbar(\phi_{i,\ell} - \phi_{j,t})/2} x_{i,j} x_{\ell,t} \quad \forall i, j, \ell, t \in \{1, \dots, n\} \quad (3.20)$$

Note that this formula shows how the new, deformed product is equivalent modulo \hbar to the old one: this is a general fact, due to the very construction, namely because we are working with 2-cocycles of the form $\exp(\hbar \varsigma)$ where ς is some bilinear form on the QFSHA to be deformed. By this same reason, *any set of elements which generate, as an algebra, the QFSHA under exam, will also generate it (as an algebra*

again) w.r.t. the new, deformed product. For this reason, the formula (3.20) is enough to describe the deformed algebra $F_{\hbar}[[G]]_{\sigma_{\mathcal{F}}}$ as the latter is (again) generated — w.r.t. the new product — by the $x_{i,j}$'s, just like $F_{\hbar}[[G]]$ was (with the old product).

Let us now see how (3.20) yields a modified Poisson bracket in the semiclassical limit of $F_{\hbar}[[G]]_{\sigma_{\mathcal{F}}}$. Using such notation as $\bar{x}_{r,s} := x_{r,s} \pmod{\hbar F_{\hbar}[[G]]_{\sigma_{\mathcal{F}}}}$, the Poisson bracket inherited from $F_{\hbar}[[G]]_{\sigma_{\mathcal{F}}}$ is given, by definition, by

$$\{\bar{x}_{i,j}, \bar{x}_{\ell,t}\}_{\sigma_{\mathcal{F}}} := \frac{[x_{i,j}, x_{\ell,t}]_{\sigma_{\mathcal{F}}}}{\hbar} \pmod{\hbar F_{\hbar}[[G]]_{\sigma_{\mathcal{F}}}}$$

Now

$$\begin{aligned} [x_{i,j}, x_{\ell,t}]_{\sigma_{\mathcal{F}}} &= x_{i,j} \dot{\sigma}_{\mathcal{F}} x_{\ell,t} - x_{\ell,t} \dot{\sigma}_{\mathcal{F}} x_{i,j} = \\ &= e^{\hbar(\phi_{i,\ell} - \phi_{j,t})/2} x_{i,j} x_{\ell,t} - e^{\hbar(\phi_{j,t} - \phi_{i,\ell})/2} x_{\ell,t} x_{i,j} = \\ &= e^{\hbar(\phi_{i,\ell} - \phi_{j,t})/2} [x_{i,j}, x_{\ell,t}] + (e^{\hbar(\phi_{i,\ell} - \phi_{j,t})/2} - e^{\hbar(\phi_{j,t} - \phi_{i,\ell})/2}) x_{\ell,t} x_{i,j} \end{aligned}$$

hence expanding the exponentials we get

$$[x_{i,j}, x_{\ell,t}]_{\sigma_{\mathcal{F}}} = (1 + \hbar(\phi_{i,\ell} - \phi_{j,t})/2) [x_{i,j}, x_{\ell,t}] + \hbar(\phi_{i,\ell} - \phi_{j,t}) x_{\ell,t} x_{i,j} + \mathcal{O}(\hbar^2)$$

from which we eventually get

$$\{\bar{x}_{i,j}, \bar{x}_{\ell,t}\}_{\sigma_{\mathcal{F}}} := \{\bar{x}_{i,j}, \bar{x}_{\ell,t}\} + (\bar{\phi}_{i,\ell} - \bar{\phi}_{j,t}) \bar{x}_{\ell,t} \bar{x}_{i,j} \quad (3.21)$$

where $\{\bar{x}_{i,j}, \bar{x}_{\ell,t}\}$ denotes the old (undeformed) Poisson bracket and we took into account that $[\bar{x}_{i,j}, \bar{x}_{\ell,t}] = 0$ and that the deformed and undeformed product do coincide modulo \hbar .

In addition, the formula (3.21) for the Poisson bracket also induces, following the general recipe, a concrete description of the modified Lie bracket in the cotangent Lie bialgebra $\mathfrak{g}^* := \mathfrak{m}/\mathfrak{m}^2$, where \mathfrak{m} is the augmentation ideal of $F_{\hbar}[[G]]_{\sigma_{\mathcal{F}}}$. Indeed, the latter has as \mathbb{k} -basis the set of cosets (modulo \mathfrak{m}^2)

$$\left\{ x_{i,j} := (\bar{x}_{i,j} - \delta_{i,j}) \pmod{\mathfrak{m}^2} \mid i, j = 1, \dots, n \right\}$$

and for these elements from (3.21) we deduce the deformed Lie bracket as given by

$$\begin{aligned} [x_{i,j}, x_{\ell,t}]_{\sigma_{\mathcal{F}}} &= [x_{i,j}, x_{\ell,t}] \quad , \quad [x_{i,i}, x_{\ell,\ell}]_{\sigma_{\mathcal{F}}} = [x_{i,i}, x_{\ell,\ell}] \quad \forall i \neq j, \ell \neq t \\ [x_{i,i}, x_{\ell,t}]_{\sigma_{\mathcal{F}}} &= [x_{i,i}, x_{\ell,t}] + (\bar{\phi}_{i,\ell} - \bar{\phi}_{i,t}) x_{\ell,t} \quad \forall \ell \neq t \\ [x_{i,j}, x_{\ell,\ell}]_{\sigma_{\mathcal{F}}} &:= [x_{i,i}, x_{\ell,\ell}] + (\bar{\phi}_{i,\ell} - \bar{\phi}_{j,\ell}) x_{i,j} \quad \forall i \neq j \end{aligned}$$

3.3. Deformations by quasi-2-cocycle of QUEA's.

This subsection is dedicated to deformations by quasi-2-cocycle of QUEA's. In this case the result that we achieve is somewhat surprising, in that we are indeed “stretching the standard recipe”, as the 2-cocycles that we use to deform our Hopf $\mathbb{k}[[\hbar]]$ -algebras are valued in the field $\mathbb{k}((\hbar))$ rather than in our ground ring $\mathbb{k}[[\hbar]]$. Therefore, *a priori* nothing even guarantees that the recipe would just work and produce a new Hopf algebra over $\mathbb{k}[[\hbar]]$; nonetheless, we eventually find quite a

meaningful result, which also says that *the standard procedure of deformation by twist for QUEA's can be extended somewhat beyond its natural borders.*

We begin with two ancillary results.

Lemma 3.3.1.

Let $U_{\hbar} := U_{\hbar}(\mathfrak{g})$ be any QUEA, and $J_{\hbar} := \text{Ker}(\epsilon_{U_{\hbar}})$. For every $z \in U_{\hbar}$, there exists $N \in \mathbb{N}$ such that $\delta_n(z) \in \hbar^{\max(n,N)-N} J_{\hbar}^{\otimes n}$ for every $n \in \mathbb{N}$.

Proof. As Drinfeld's functors are inverse to each other — cf. Theorem 2.4.2(a) — applying $(\)^{\vee}$ after $(\)'$ to the QUEA U_{\hbar} we get $U_{\hbar} = (U_{\hbar}')^{\vee}$: letting $I_{\hbar}' := \hbar U_{\hbar}' + \text{Ker}(\epsilon_{U_{\hbar}'})$, this last identity reads

$$U_{\hbar} = \hbar\text{-adic completion of } \sum_{n \geq 0} \hbar^{-n} (I_{\hbar}')^n = \hbar\text{-adic completion of } \bigcup_{n \geq 0} \hbar^{-n} (I_{\hbar}')^n$$

In particular, this implies that for our $z \in U_{\hbar}(\mathfrak{g})$ there exist some $N \in \mathbb{N}$ and $z' \in (I_{\hbar}')^N$ such that $z \equiv \hbar^{-N} z' \pmod{\hbar U_{\hbar}(\mathfrak{g})}$. Now, given $n \in \mathbb{N}$ we have $\delta_n(z') \in \hbar^n U_{\hbar}^{\otimes n}$ because $z' \in (I_{\hbar}')^N \subseteq U_{\hbar}'$, and also $\delta_n(z') \in \sum_{s_1 + \dots + s_n = N} \otimes_{i=1}^n (I_{\hbar}')^{s_i}$ because I_{\hbar}' is a Hopf ideal; moreover, $\text{Ker}(\epsilon_{U_{\hbar}'}) \subseteq \hbar U_{\hbar}'$ again by construction, hence $I_{\hbar}' \subseteq \hbar U_{\hbar}'$. In the end, all this yields $\delta_n(z') \in \hbar^{\max(n,N)} J_{\hbar}^{\otimes n}$, therefore $\delta_n(z) \in \hbar^{\max(n,N)-N} J_{\hbar}^{\otimes n}$ as claimed. \square

For the second, auxiliary result, we fix some more notation: namely, hereafter by “ \log_* ” and “ \exp_* ” we denote the logarithm and the exponential with respect to the convolution product, whenever defined.

Lemma 3.3.2. Let $U_{\hbar}(\mathfrak{g})$ be any QUEA, and let χ be a $\mathbb{k}[[\hbar]]$ -bilinear form on $U_{\hbar}(\mathfrak{g})$ such that $\chi(z, 1) = 0 = \chi(1, z)$ for any $z \in U_{\hbar}(\mathfrak{g})$; denote also by the same symbol χ the scalar extension of χ to a $\mathbb{k}((\hbar))$ -bilinear form for the $\mathbb{k}((\hbar))$ -vector space $\mathbb{U}_{\hbar}(\mathfrak{g}) := \mathbb{k}((\hbar)) \otimes_{\mathbb{k}[[\hbar]]} U_{\hbar}(\mathfrak{g})$. Then:

(a) the formal expression $\sigma := \exp_*(\hbar^{-1}\chi)$ uniquely provides a well-defined, $\mathbb{k}((\hbar))$ -valued bilinear form for $\mathbb{U}_{\hbar}(\mathfrak{g})$;

(b) $\sigma(z, 1) = \epsilon(z) = \sigma(1, z)$ for any $z \in U_{\hbar}(\mathfrak{g})$;

(c) σ is orthogonal, i.e. $\sigma_{2,1} = \sigma^{-1}$, iff χ is antisymmetric, i.e. $\chi_{2,1} = -\chi$.

Proof. (a) Fix notation $U_{\hbar} := U_{\hbar}(\mathfrak{g})$ and $J_{\hbar} := \text{Ker}(\epsilon_{U_{\hbar}})$. For any $z \in U_{\hbar}$, set

$$\hat{z} := \epsilon(z), \quad z^+ := z - \epsilon(z) = z - \hat{z} \in J_{\hbar}, \quad \text{hence } z = z^+ + \hat{z} \quad (3.22)$$

The assumption $\chi(z, 1) = 0 = \chi(1, z)$ for $z \in U_{\hbar}(\mathfrak{g})$ implies (for all $u, v \in U_{\hbar}$)

$$\chi(u, v) = \chi(u^+ + \hat{u}, v^+ + \hat{v}) = \chi(u^+, v^+) \quad (3.23)$$

Now, for any $a, b \in U_{\hbar}$, the formula $\sigma = \exp_*(\hbar^{-1}\chi) = \sum_{n \geq 0} \hbar^{-n} \chi^{*n} / n!$ gives

$$\begin{aligned} \sigma(a, b) &= \sum_{n \geq 0} \hbar^{-n} \chi^{*n}(a, b) / n! = \\ &= \sum_{n \geq 0} \hbar^{-n} \prod_{i=1}^n \chi(a_{(i)}, b_{(i)}) / n! = \sum_{n \geq 0} \hbar^{-n} \prod_{i=1}^n \chi(a_{(i)}^+, b_{(i)}^+) / n! \end{aligned} \quad (3.24)$$

where we took into account that $\chi^{*k}(u, v) = \prod_{s=1}^k \chi(u_{(s)}, v_{(s)}) = \prod_{s=1}^k \chi(u_{(s)}^+, v_{(s)}^+)$ for each $u, v \in U_{\hbar}$, $k \in \mathbb{N}$, by definitions along with (3.23). Now we notice that $\otimes_{i=1}^n a_{(i)}^+ = \delta_n(a)$ and $\otimes_{i=1}^n b_{(i)}^+ = \delta_n(b)$, hence Lemma 3.3.1 guarantees that

$$h^{-n} \prod_{i=1}^n \chi(a_{(i)}^+, b_{(i)}^+) \in \hbar^{-n + \max(n, A) - A + \max(n, B) - B} \quad \forall n \in \mathbb{N}_+$$

whence in particular

$$\begin{aligned} h^{-n} \prod_{i=1}^n \chi(a_{(i)}^+, b_{(i)}^+) &\in \hbar^{-\min(A, B)} \mathbb{k}[[\hbar]] \quad \forall n \in \mathbb{N}_+ \\ h^{-n} \prod_{i=1}^n \chi(a_{(i)}^+, b_{(i)}^+) &\in \hbar^{n-(A+B)} \mathbb{k}[[\hbar]] \quad \forall n \geq A+B \end{aligned} \quad (3.25)$$

where $A \in \mathbb{N}$, resp. $B \in \mathbb{N}$, plays for a , resp. for b , the role of N for z in Lemma 3.3.1 above; by this, the formal expansion for $\sigma(a, b)$ in (3.24) yields a well defined element in $\mathbb{k}[[\hbar]]$, hence σ is a well-defined $\mathbb{k}((\hbar))$ -bilinear form of $\mathbb{U}_{\hbar}(\mathfrak{g})$ as claimed.

(b-c) Both claims are obvious, by construction, as they follow from standard identities for formal exponentials. \square

The previous result leads us to introduce the following notion:

Definition 3.3.3. Let $U_{\hbar}(\mathfrak{g})$ be a QUEA, and $\mathbb{U}_{\hbar}(\mathfrak{g}) := \mathbb{k}((\hbar)) \otimes_{\mathbb{k}[[\hbar]]} U_{\hbar}(\mathfrak{g})$. Note that $\mathbb{U}_{\hbar}(\mathfrak{g})$ has a natural ‘‘Hopf algebra structure’’ of $\mathbb{U}_{\hbar}(\mathfrak{g})$ induced by scalar extension from $U_{\hbar}(\mathfrak{g})$ — so that, in particular, the ‘‘coproduct’’ takes values in $\mathbb{k}((\hbar)) \otimes_{\mathbb{k}[[\hbar]]} (U_{\hbar}(\mathfrak{g}) \widehat{\otimes}_{\mathbb{k}[[\hbar]]} U_{\hbar}(\mathfrak{g}))$ rather than in $\mathbb{U}_{\hbar}(\mathfrak{g}) \otimes_{\mathbb{k}((\hbar))} \mathbb{U}_{\hbar}(\mathfrak{g})$.

We call *quasi-2-cocycle of $U_{\hbar}(\mathfrak{g})$* any $\mathbb{k}((\hbar))$ -bilinear form σ of $\mathbb{U}_{\hbar}(\mathfrak{g})$ which has the form $\sigma := \exp_*(\hbar^{-1}\chi)$ for some $\mathbb{k}[[\hbar]]$ -bilinear form $\chi \in (U_{\hbar}(\mathfrak{g}) \widehat{\otimes}^2)^*$ of $U_{\hbar}(\mathfrak{g})$ such that $\chi(z, 1) = 0 = \chi(1, z)$ for all $z \in U_{\hbar}(\mathfrak{g})$, and in addition enjoys the 2-cocycle properties with respect to the above ‘‘Hopf algebra structure’’ of $\mathbb{U}_{\hbar}(\mathfrak{g})$.

Remark 3.3.4. The notion of ‘‘quasi-2-cocycle’’ for a QUEA $U_{\hbar}(\mathfrak{g})$ can also be cast in the following, equivalent shape. Recall that $F_{\hbar}[[G]] := U_{\hbar}(\mathfrak{g})^*$ is a QFSHA (cf. §2.3.4), and then $(U_{\hbar}(\mathfrak{g}) \widehat{\otimes}^2)^* = U_{\hbar}(\mathfrak{g})^* \widetilde{\otimes} U_{\hbar}(\mathfrak{g})^* = F_{\hbar}[[G]] \widetilde{\otimes} F_{\hbar}[[G]]$. Given $\chi \in (U_{\hbar}(\mathfrak{g}) \widehat{\otimes}^2)^*$ as in Definition 3.3.3 above, the condition $\chi(z, 1) = 0 = \chi(1, z)$ for all $z \in U_{\hbar}(\mathfrak{g})$ means that $\chi \in J_{F_{\hbar}[[G]]} \widetilde{\otimes} J_{F_{\hbar}[[G]]}$, with $J_{F_{\hbar}[[G]]} := \text{Ker}(\epsilon_{F_{\hbar}[[G]]})$, hence we have $\chi \in \hbar^2 (J_{F_{\hbar}[[G]]}^{\vee}) \widehat{\otimes}^2 \subseteq \hbar^2 (F_{\hbar}[[G]]^{\vee}) \widehat{\otimes}^2$ where $J_{F_{\hbar}[[G]]}^{\vee} := \hbar^{-1} J_{F_{\hbar}[[G]]}$ and $F_{\hbar}[[G]]^{\vee}$ is the QUEA defined in §2.4 out of $F_{\hbar}[[G]]$. Thus, it follows that $\hbar^{-1}\chi \in \hbar (F_{\hbar}[[G]]^{\vee}) \widehat{\otimes}^2$, so $\sigma := \exp_*(\hbar^{-1}\chi)$ is a well-defined element in $(F_{\hbar}[[G]]^{\vee}) \widehat{\otimes}^2$.

Now, the requirement that $\sigma := \exp_*(\hbar^{-1}\chi)$ be a quasi-2-cocycle for $U_{\hbar}(\mathfrak{g})$ in the sense of Definition 3.3.3 above is equivalent to the property of σ being a twist element for $F_{\hbar}[[G]]^{\vee}$ — which makes perfectly sense in sight of Proposition 2.2.7.

Clearly, every 2-cocycle for $U_{\hbar}(\mathfrak{g})$ is a quasi-2-cocycle as well; the converse, instead, is not true, in general (counterexamples do exist). However, the key point is that *every quasi-2-cocycle still provides a well-defined deformation by 2-cocycle of $U_{\hbar}(\mathfrak{g})$* — in short, the construction of *deformations by 2-cocycle* does properly extend to ‘‘deformations by quasi-2-cocycle’’ as well: this is indeed our next result.

Theorem 3.3.5. *Let $U_{\hbar}(\mathfrak{g})$ be a QUEA, and $\sigma = \exp_*(\hbar^{-1}\chi)$ a quasi-2-cocycle for it, as in Definition 3.3.3. Then the procedure of 2-cocycle deformation by σ applied to $U_{\hbar}(\mathfrak{g})$ actually restricts to $U_{\hbar}(\mathfrak{g})$, making the latter into a new QUEA.*

Proof. First of all, we have to explain the statement itself. To begin with, note that, by definitions and by Lemma 3.3.2, we can perform the deformation by the 2-cocycle σ onto the ‘‘Hopf $\mathbb{k}((\hbar))$ -algebra’’ $\mathbb{U}_{\hbar}(\mathfrak{g}) := \mathbb{k}((\hbar)) \widehat{\otimes} U_{\hbar}(\mathfrak{g})$. Our statement then claims the resulting deformed Hopf structure onto $\mathbb{U}_{\hbar}(\mathfrak{g})$ actually ‘‘restricts’’ to a deformation of $U_{\hbar}(\mathfrak{g})$ itself: in turn, this amounts to claiming that $U_{\hbar}(\mathfrak{g})$ is closed for the σ -deformed product in $(\mathbb{U}_{\hbar}(\mathfrak{g}))_{\sigma}$ — so we go and tackle this last problem.

Fix notation $U_{\hbar} := U_{\hbar}(\mathfrak{g})$, $J_{\hbar} := \text{Ker}(U_{\hbar})$, $J'_{\hbar} := \text{Ker}(U'_{\hbar})$ and $\widetilde{J}_{\hbar} := \hbar^{-1}J'_{\hbar}$, where $U'_{\hbar} := U_{\hbar}(\mathfrak{g})'$ is given in Definition 2.4.1(a). As it was mentioned in the proof of Lemma 3.3.1, Theorem 2.4.2(a) implies that $U_{\hbar} = (U'_{\hbar})^{\vee}$, that is

$$U_{\hbar} = \hbar\text{-adic completion of } \sum_{n \geq 0} \hbar^{-n} (I'_{\hbar})^n$$

where $I'_{\hbar} := \hbar U'_{\hbar} + \text{Ker}(\epsilon_{U'_{\hbar}}) = \hbar U'_{\hbar} + J'_{\hbar}$; then a moment's thought shows that the previous expression of U_{\hbar} reads also

$$U_{\hbar} = \hbar\text{-adic completion of } \sum_{n \geq 0} \hbar^{-n} (J'_{\hbar})^n = \hbar\text{-adic completion of } \sum_{n \geq 0} \widetilde{J}_{\hbar}^n \quad (3.26)$$

Note also that clearly J'_{\hbar} is a Hopf ideal in U'_{\hbar} , and moreover $J'_{\hbar} \subseteq \hbar J_{\hbar}$ (by construction); therefore for $z' \in J'^N_{\hbar}$ — with $N \in \mathbb{N}$ — acting like in the proof of Lemma 3.3.1, one sees that

$$\delta_n(z') \in \hbar^n J_{\hbar}^{\otimes n} \cap \left(\sum_{\sum_i N_i = N} \bigotimes_{i=1}^n J'^{N_i}_{\hbar} \right) \subseteq \hbar^{\max(n, N)} J_{\hbar}^{\otimes n} \quad (3.27)$$

Again, for any $z \in U_{\hbar}$ we retain notation as in (3.22) above, that is

$$\widehat{z} := \epsilon(z), \quad z^+ := z - \epsilon(z) = z - \widehat{z} \in J_{\hbar}, \quad \text{hence } z = z^+ + \widehat{z} \quad (3.28)$$

and we recall also that for all $u, v \in U_{\hbar}$ we have

$$\chi(u, v) = \chi(u^+ + \widehat{u}, v^+ + \widehat{v}) = \chi(u^+, v^+) \quad (3.29)$$

Thanks to (3.26), in order to prove that $U_{\hbar}(\mathfrak{g}) =: U_{\hbar}$ is closed for the σ -deformed product \cdot_{σ} it is enough to show that $\widetilde{J}_{\hbar}^A \cdot_{\sigma} \widetilde{J}_{\hbar}^B \subseteq \sum_{n \geq 0} \widetilde{J}_{\hbar}^n$ for any $A, B \in \mathbb{N}_+$.

To begin with, we pick $a \in \widetilde{J}_{\hbar}^A = \hbar^{-A} J'^A_{\hbar}$ and $b \in \widetilde{J}_{\hbar}^B = \hbar^{-B} J'^B_{\hbar}$; by definition,

$$a \cdot_{\sigma} b := \sigma(a_{(1)}, b_{(1)}) a_{(2)} b_{(2)} \sigma^{-1}(a_{(3)}, b_{(3)})$$

whence expanding the formal formula $\sigma = \exp_*(\hbar^{-1}\chi) = \sum_{n \geq 0} \hbar^{-n} \chi^{*n} / n!$ — much like in the proof of Lemma 3.3.2 — we get

$$\begin{aligned} a \cdot_{\sigma} b &= \sum_{t, \ell \geq 0} \hbar^{-(t+\ell)} (-1)^{\ell} (t! \ell!)^{-1} \chi^{*t}(a_{(1)}, b_{(1)}) a_{(2)} b_{(2)} \chi^{*\ell}(a_{(3)}, b_{(3)}) = \\ &= \epsilon(a'_{(1)}) \epsilon(b'_{(1)}) a'_{(2)} b'_{(2)} \epsilon(a'_{(3)}) \epsilon(b'_{(3)}) + \\ &+ \sum_{t+\ell > 0} \hbar^{-(t+\ell)} (-1)^{\ell} (t!)^{-1} (\ell!)^{-1} \chi^{*t}(a_{(1)}, b_{(1)}) a_{(2)} b_{(2)} \chi^{*\ell}(a_{(3)}, b_{(3)}) = \quad (3.30) \\ &= a \cdot b + \\ &+ \sum_{t+\ell > 0} \hbar^{-(t+\ell)} (-1)^{\ell} (t!)^{-1} (\ell!)^{-1} \chi^{*t}(a_{(1)}, b_{(1)}) a_{(2)} b_{(2)} \chi^{*\ell}(a_{(3)}, b_{(3)}) \end{aligned}$$

where we took into account coassociativity and counitality properties.

Let us analyze each summand in the very last line in (3.30). From the identities $\chi^{*k}(u, v) = \prod_{s=1}^k \chi(u_{(s)}, v_{(s)}) = \prod_{s=1}^k \chi(u_{(s)}^+, v_{(s)}^+)$ — cf. (3.29) — we get

$$\begin{aligned} \chi^{*t}(a_{(1)}, b_{(1)}) a_{(2)} b_{(2)} \chi^{*\ell}(a_{(3)}, b_{(3)}) &= \\ &= \prod_{i=1}^t \chi(a_{(i)}, b_{(i)}) a_{(t+1)} b_{(t+1)} \prod_{j=1}^{\ell} \chi(a_{(t+1+j)}, b_{(t+1+j)}) = \\ &= \prod_{i=1}^t \chi(a_{(i)}^+, b_{(i)}^+) a_{(t+1)} b_{(t+1)} \prod_{j=1}^{\ell} \chi(a_{(t+1+j)}^+, b_{(t+1+j)}^+) \end{aligned}$$

Consider the expansion of a as in (3.28). Then, letting $j_{t+1} : U_{\hbar}^{\otimes(t+\ell)} \longrightarrow U_{\hbar}^{\otimes(t+\ell+1)}$ be the map given by $\bigotimes_{s=1}^{t+\ell} x_s \mapsto \left(\bigotimes_{s=1}^t x_s \right) \otimes 1 \otimes \left(\bigotimes_{s=t+1}^{t+\ell} x_s \right)$, we have

$$\begin{aligned} \left(\bigotimes_{i=1}^t a_{(i)}^+ \right) \otimes a_{(t+1)}^+ \otimes \left(\bigotimes_{j=1}^{\ell} a_{(t+1+j)}^+ \right) &= \delta_{t+\ell+1}(a) \in \hbar^{\max(t+\ell+1, A) - A} U_{\hbar}^{\otimes(t+\ell+1)} \\ \left(\bigotimes_{i=1}^t a_{(i)}^+ \right) \otimes \widehat{a}_{(t+1)} \otimes \left(\bigotimes_{j=1}^{\ell} a_{(t+1+j)}^+ \right) &= j_{t+1}(\delta_{t+\ell}(a)) \in \hbar^{\max(t+\ell, A) - A} U_{\hbar}^{\otimes(t+\ell+1)} \end{aligned}$$

so that, summing up,

$$\left(\bigotimes_{i=1}^t a_{(i)}^+ \right) \otimes a_{t+1} \otimes \left(\bigotimes_{j=1}^{\ell} a_{(t+1+j)}^+ \right) = \delta_{t+\ell+1}(a) + j_{t+1}(\delta_{t+\ell}(a)) \in \hbar^{\max(t+\ell, A) - A} U_{\hbar}^{\otimes(t+\ell+1)}$$

— like in the proof of Lemma 3.3.1 — and similarly with b , resp. B , replacing a , resp. A . Eventually, for all $t + \ell > 0$ this gives

$$\begin{aligned} \chi^{*t}(a_{(1)}, b_{(1)}) a_{(2)} b_{(2)} \chi^{*\ell}(a_{(3)}, b_{(3)}) &= \\ &= \chi^{*t}(a_{(1)}, b_{(1)}) \widehat{a}_{(2)} b_{(2)} \chi^{*\ell}(a_{(3)}, b_{(3)}) + \chi^{*t}(a_{(1)}, b_{(1)}) a_{(2)}^+ b_{(2)} \chi^{*\ell}(a_{(3)}, b_{(3)}) \end{aligned} \quad (3.31)$$

where for the two summands in second line, writing $n := t + \ell$, we have

$$\begin{aligned} \chi^{*t}(a_{(1)}, b_{(1)}) \widehat{a}_{(2)} b_{(2)} \chi^{*\ell}(a_{(3)}, b_{(3)}) &= \\ &= \prod_{i=1}^t \chi(a_{(i)}^+, b_{(i)}^+) \widehat{a}_{(t+1)} b_{(t+1)} \prod_{k=t+2}^{n+1} \chi(a_{(k)}^+, b_{(k)}^+) \in \hbar^{\max(n, A) - A + \max(n, B) - B} U_{\hbar}^{\otimes(n+1)} \\ \chi^{*t}(a_{(1)}, b_{(1)}) a_{(2)}^+ b_{(2)} \chi^{*\ell}(a_{(3)}, b_{(3)}) &= \\ &= \prod_{i=1}^t \chi(a_{(i)}^+, b_{(i)}^+) a_{(t+1)}^+ b_{(t+1)} \prod_{k=t+2}^{n+1} \chi(a_{(k)}^+, b_{(k)}^+) \in \hbar^{\max(n+1, A) - A + \max(n, B) - B} U_{\hbar}^{\otimes(n+1)} \end{aligned}$$

Let us now assume that $A := 1$, so that $n := t + \ell > 0$ implies $n := t + \ell \geq 1 = A$. Then the last estimates read

$$\begin{aligned} \chi^{*t}(a_{(1)}, b_{(1)}) \widehat{a}_{(2)} b_{(2)} \chi^{*\ell}(a_{(3)}, b_{(3)}) &\in \hbar^{n-1 + \max(n, B) - B} U_{\hbar}^{\otimes(n+1)} \\ \chi^{*t}(a_{(1)}, b_{(1)}) a_{(2)}^+ b_{(2)} \chi^{*\ell}(a_{(3)}, b_{(3)}) &\in \hbar^{n + \max(n, B) - B} U_{\hbar}^{\otimes(n+1)} \end{aligned} \quad (3.32)$$

The term in the second line, when plugged in (3.31) and then in (3.30), yields a contribution of the form

$$\frac{(-1)^{\ell}}{t! \ell!} \hbar^{-n} \chi^{*t}(a_{(1)}, b_{(1)}) a_{(2)}^+ b_{(2)} \chi^{*\ell}(a_{(3)}, b_{(3)}) \in \hbar^{\max(n, B) - B} U_{\hbar}^{\otimes(n+1)}$$

that belongs to $\hbar^{\max(n,B)-B} U_{\hbar}^{\otimes(n+1)}$, thus for growing n these elements sum up to a convergent series in U_{\hbar} , and we are done.

As to the term in the first line, we split it into

$$\begin{aligned} \chi^{*t}(a_{(1)}, b_{(1)}) \widehat{a}_{(2)} b_{(2)} \chi^{*\ell}(a_{(3)}, b_{(3)}) &= \\ &= \chi^{*t}(a_{(1)}, b_{(1)}) \widehat{a}_{(2)} \widehat{b}_{(2)} \chi^{*\ell}(a_{(3)}, b_{(3)}) + \chi^{*t}(a_{(1)}, b_{(1)}) \widehat{a}_{(2)} b_{(2)}^+ \chi^{*\ell}(a_{(3)}, b_{(3)}) \end{aligned} \quad (3.33)$$

Then for the first summand we have (almost by definition, or acting as before)

$$\chi^{*t}(a_{(1)}, b_{(1)}) \widehat{a}_{(2)} \widehat{b}_{(2)} \chi^{*\ell}(a_{(3)}, b_{(3)}) = \chi^{*(t+\ell)}(a, b)$$

so when we plug every such term in (3.31) and then in (3.30), overall they sum up to give the contribution

$$\begin{aligned} \sum_{t+\ell>0} \hbar^{-(t+\ell)} (-1)^{\ell} (t!)^{-1} (\ell!)^{-1} \chi^{*t}(a_{(1)}, b_{(1)}) \widehat{a}_{(2)} \widehat{b}_{(2)} \chi^{*\ell}(a_{(3)}, b_{(3)}) &= \\ = \sum_{n>0} \sum_{t+\ell=n} \hbar^{-(t+\ell)} (-1)^{\ell} (t!)^{-1} (\ell!)^{-1} \chi^{*(t+\ell)}(a, b) &= \\ = \sum_{n>0} \frac{1}{n!} \hbar^{-n} \left(\sum_{t+\ell=n} (-1)^{\ell} \binom{n}{\ell} \right) \chi^{*n}(a, b) &= 0 \end{aligned}$$

just because of the combinatorial identity $\sum_{t+\ell=n} (-1)^{\ell} \binom{n}{\ell} = 0$.

Finally, we have to dispose of the summands of type

$$\chi^{*t}(a_{(1)}, b_{(1)}) \widehat{a}_{(2)} b_{(2)}^+ \chi^{*\ell}(a_{(3)}, b_{(3)}) \quad (3.34)$$

for which the analogue of the first identity in (3.32) holds true, namely

$$\chi^{*t}(a_{(1)}, b_{(1)}) \widehat{a}_{(2)} b_{(2)}^+ \chi^{*\ell}(a_{(3)}, b_{(3)}) \in \hbar^{n-1+\max(n+1,B)-B} U_{\hbar}^{\otimes(n+1)} \quad (3.35)$$

where $n := t + \ell$, taking into account that $\delta_{n+1}(b) \in \hbar^{\max(n+1,B)-B} U_{\hbar}^{\otimes(n+1)}$.

Then we have to distinguish two cases, depending on $n := t + \ell$.

First we assume $n := t + \ell \geq B$. Then $n - 1 + \max(n + 1, B) - B \geq n$, hence the first identity in (3.32) yields $\chi^{*t}(a_{(1)}, b_{(1)}) \widehat{a}_{(2)} b_{(2)}^+ \chi^{*\ell}(a_{(3)}, b_{(3)}) \in \hbar^n U_{\hbar}^{\otimes(n+1)}$, and then, when plugged in (3.33), and subsequently in (3.31) and in (3.30), this provides to the expansion of $a \cdot b$ a contribution of the form

$$\hbar^{-n} \frac{(-1)^{\ell}}{t! \ell!} \chi^{*t}(a_{(1)}, b_{(1)}) \widehat{a}_{(2)} b_{(2)}^+ \chi^{*\ell}(a_{(3)}, b_{(3)}) \in \hbar^{-n} \hbar^n U_{\hbar}^{\otimes(n+1)} = U_{\hbar}^{\otimes(n+1)}$$

— which is fair! — hence we are done with it.

Then we are left with the case $n := t + \ell \leq B - 1$. Tracking backwards our construction, all these case provide to (3.30) a contribution of the form

$$\begin{aligned} \sum_{t+\ell=1}^{B-1} \hbar^{-(t+\ell)} \frac{(-1)^{\ell}}{t! \ell!} \chi^{*t}(a_{(1)}, b_{(1)}) \widehat{a}_{(2)} b_{(2)}^+ \chi^{*\ell}(a_{(3)}, b_{(3)}) &= \\ = \sum_{n=1}^{B-1} \frac{1}{n!} \hbar^{-n} \sum_{t+\ell=n} (-1)^{\ell} \binom{n}{\ell} \chi^{*t}(a_{(1)}, b_{(1)}) \widehat{a}_{(2)} b_{(2)}^+ \chi^{*\ell}(a_{(3)}, b_{(3)}) \end{aligned} \quad (3.36)$$

With no loss of generality, we can assume that $a \neq 0$, $b \neq 0 \pmod{\hbar U_{\hbar}}$. Then for their corresponding cosets $\bar{a}, \bar{b} \in U_{\hbar} / \hbar U_{\hbar} \cong U(\mathfrak{g})$ we have $\bar{a} \in U(\mathfrak{g})_1$

and $\bar{b} \in U(\mathfrak{g})_B$, where $\{U(\mathfrak{g})_n\}_{n \in \mathbb{N}}$ is the standard, coradical filtration of $U(\mathfrak{g})$, and also $\delta_1(\bar{a}) \neq 0$ as well as $\delta_n(\bar{b}) \neq 0$ for $1 \leq n \leq B$ — cf. [Ga1], Lemma 3.3. Moreover, we recall that $U_\hbar := U_\hbar(\mathfrak{g})$ is *cocommutative modulo $\hbar U_\hbar$* , as it is a QUEA: in particular, this implies that $\delta_n(\bar{b})$ is a *symmetric tensor* — for $1 \leq n \leq B$ — hence we can write $\delta_n(b)$ in the form

$$\delta_n(b) = b_{(1)}^+ \otimes \cdots \otimes b_{(n)}^+ = \beta_{(1)} \otimes \cdots \otimes \beta_{(n)} + \mathcal{O}_n(\hbar^1) \quad (3.37)$$

(for $1 \leq n \leq B$) where $\beta_{(1)} \otimes \cdots \otimes \beta_{(n)}$ — using some σ -notation of sort, as usual — is some *symmetric tensor* in $U_\hbar^{\otimes n}$ and hereafter $\mathcal{O}_n(\hbar^s)$ stands for some element in $\hbar^s U_\hbar^{\otimes n}$, for every $s, n \in \mathbb{N}$. Then plugging (3.37) in (3.34) we find

$$\begin{aligned} \chi^{*t}(a_{(1)}, b_{(1)}) \widehat{a}_{(2)} b_{(2)}^+ \chi^{*\ell}(a_{(3)}, b_{(3)}) &= \prod_{i=1}^t \chi(a_{(i)}^+, b_{(i)}^+) \widehat{a}_{(t+1)} b_{(t+1)}^+ \prod_{k=t+2}^{t+\ell+1} \chi(a_{(k)}^+, b_{(k)}^+) = \\ &= \prod_{i=1}^t \chi(a_{(i)}^+, \beta_{(i)}) \widehat{a}_{(t+1)} \beta_{(t+1)} \prod_{k=t+2}^{t+\ell+1} \chi(a_{(k)}^+, \beta_{(k)}) + \mathcal{O}_1(\hbar^{t+\ell}) \end{aligned}$$

for all $t + \ell \leq B - 1$, with

$$\prod_{i=1}^t \chi(a_{(i)}^+, \beta_{(i)}) \widehat{a}_{(t+1)} \beta_{(t+1)} \prod_{k=t+2}^{t+\ell+1} \chi(a_{(k)}^+, \beta_{(k)}) \in \hbar^{t+\ell-1} U_\hbar$$

Therefore, the contribution to (3.30) given in (3.36) now reads

$$\begin{aligned} \sum_{n=1}^{B-1} \frac{1}{n!} \hbar^{-n} \sum_{t+\ell=n} (-1)^\ell \binom{n}{\ell} \chi^{*t}(a_{(1)}, b_{(1)}) \widehat{a}_{(2)} b_{(2)}^+ \chi^{*\ell}(a_{(3)}, b_{(3)}) &= \\ = \sum_{n=1}^{B-1} \frac{1}{n!} \hbar^{-n} \sum_{t+\ell=n} (-1)^\ell \binom{n}{\ell} \prod_{i=1}^t \chi(a_{(i)}^+, \beta_{(i)}) \widehat{a}_{(t+1)} \beta_{(t+1)} \prod_{k=t+2}^{n+1} \chi(a_{(k)}^+, \beta_{(k)}) &+ \\ &+ \mathcal{O}_1(\hbar^0) \end{aligned}$$

where in the last formula we have

$$\sum_{n=1}^{B-1} \frac{1}{n!} \hbar^{-n} \sum_{t+\ell=n} (-1)^\ell \binom{n}{\ell} \prod_{i=1}^t \chi(a_{(i)}^+, \beta_{(i)}) \widehat{a}_{(t+1)} \beta_{(t+1)} \prod_{k=t+2}^{n+1} \chi(a_{(k)}^+, \beta_{(k)}) \in \hbar^{-1} U_\hbar$$

Now, observe that, setting $n := t + \ell$, we can re-write

$$\prod_{i=1}^t \chi(a_{(i)}^+, \beta_{(i)}) \widehat{a}_{(t+1)} \beta_{(t+1)} \prod_{k=t+2}^{n+1} \chi(a_{(k)}^+, \beta_{(k)}) = \Phi(\delta_n(a) \otimes \beta_{(1)} \otimes \cdots \otimes \beta_{(n+1)}) \quad (3.38)$$

with $\Phi : U_\hbar^{\otimes 2(n+1)} \longrightarrow U_\hbar$ being the map given by the composition

$$\Phi := \mu \circ (\chi^{\otimes t} \otimes \text{id}_{U_\hbar}^{\otimes 2} \otimes \chi^{\otimes (n-t)}) \circ \varsigma_{n+1}$$

where

(1) $\varsigma_{n+1} : U_\hbar^{\otimes (2n+1)} \longrightarrow U_\hbar^{\otimes (2n+1)}$ is the “shuffle” map

$$x_1 \otimes \cdots \otimes x_n \otimes y_1 \otimes \cdots \otimes y_n \otimes y_{n+1} \mapsto x_1 \otimes y_1 \otimes x_2 \otimes y_2 \otimes \cdots \otimes x_n \otimes y_n \otimes y_{n+1}$$

and, considering $\mathbb{k}[[\hbar]]$ as embedded into U_\hbar via the unit map,

(2) $\mu : U_\hbar^{\otimes n} \longrightarrow U_\hbar$ is the obvious (n -fold iterated) multiplication by scalars.

Now recall that $\beta_{(1)} \otimes \cdots \otimes \beta_{(n+1)}$ represents a tensor in σ -notation, so more explicitly we might write $\beta_{(1)} \otimes \cdots \otimes \beta_{(n+1)} = \sum_{s=1}^N \beta_{s,1} \otimes \beta_{s,n+1}$; so in the formula we are

dealing with what is written as a product $\prod_{i=1}^t \chi(a_{(i)}^+, \beta_{(i)}) \widehat{a}_{(t+1)} \beta_{(t+1)} \prod_{k=t+2}^{n+1} \chi(a_{(k)}^+, \beta_{(k)})$ is actually a sum of several products as $\prod_{i=1}^t \chi(a_{(i)}^+, \beta_{s,i}) \widehat{a}_{(t+1)} \beta_{s,t+1} \prod_{k=t+2}^{n+1} \chi(a_{(k)}^+, \beta_{s,k})$. But then recall that this tensor $\beta_{(1)} \otimes \cdots \otimes \beta_{(n+1)} = \sum_{s=1}^N \beta_{s,1} \otimes \beta_{s,n+1}$ is *symmetric*, therefore, the various products $\prod_{i=1}^t \chi(a_{(i)}^+, \beta_{s,i}) \widehat{a}_{(t+1)} \beta_{s,t+1} \prod_{k=t+2}^{n+1} \chi(a_{(k)}^+, \beta_{s,k})$ actually all *coincide*: letting C_n be their “common value”, we deduce that

$$\begin{aligned} \sum_{n=1}^{B-1} \frac{1}{n!} \hbar^{-n} \sum_{t+\ell=n} (-1)^\ell \binom{n}{\ell} \prod_{i=1}^t \chi(a_{(i)}^+, \beta_{(i)}) \widehat{a}_{(t+1)} \beta_{(t+1)} \prod_{k=t+2}^{n+1} \chi(a_{(k)}^+, \beta_{(k)}) &= \\ &= \sum_{n=1}^{B-1} \frac{1}{n!} \hbar^{-n} \left(\sum_{t+\ell=n} (-1)^\ell \binom{n}{\ell} \right) C_n = 0 \end{aligned}$$

again because of the identity $\sum_{t+\ell=n} (-1)^\ell \binom{n}{\ell} = 0$.

Thus, also the last contributions to (3.30) given in (3.36) actually belong to U_{\hbar} .

To sum up, we have proved that

$$\begin{aligned} a \cdot_{\sigma} b &= a \cdot b + z && \text{with } z \in J_{\hbar} \\ \forall a \in \widetilde{J}_{\hbar}^1 &= \hbar^{-1} J'_{\hbar}, \quad b \in \widetilde{J}_{\hbar}^B &= \hbar^{-B} J'^B_{\hbar} \end{aligned} \quad (3.39)$$

and similarly — just switching roles of a and b — also

$$\begin{aligned} b \cdot_{\sigma} a &= b \cdot a + x && \text{with } x \in J_{\hbar} \\ \forall a \in \widetilde{J}_{\hbar}^1 &= \hbar^{-1} J'_{\hbar}, \quad b \in \widetilde{J}_{\hbar}^B &= \hbar^{-B} J'^B_{\hbar} \end{aligned} \quad (3.40)$$

Let $\langle \widetilde{J}_{\hbar} \rangle_{\mathbb{k}[[\hbar]]}^{\sigma}$ the unital $\mathbb{k}[[\hbar]]$ -subalgebra of $(\mathbb{U}_{\hbar}(\mathfrak{g}))_{\sigma} := (\mathbb{k}(\hbar) \otimes_{\mathbb{k}[[\hbar]]} U_{\hbar})_{\sigma}$ generated by \widetilde{J}_{\hbar} . Now recall that $\widetilde{J}_{\hbar} := \hbar^{-1} J'_{\hbar}$ with $J'_{\hbar} := \text{Ker}(U'_{\hbar})$; then $U'_{\hbar} = J'_{\hbar} \oplus \mathbb{k}[[\hbar]] \cdot 1$, which implies $\Delta(J'_{\hbar}) \subseteq J'_{\hbar} \otimes 1 + J'_{\hbar} \otimes J'_{\hbar} + 1 \otimes J'_{\hbar}$. Then we get also $\Delta(\widetilde{J}_{\hbar}) \subseteq \widetilde{J}_{\hbar} \otimes 1 + \widetilde{J}_{\hbar} \otimes \widetilde{J}_{\hbar} + 1 \otimes \widetilde{J}_{\hbar}$. Since the coalgebra structure is the same in $\mathbb{U}_{\hbar}(\mathfrak{g})$ and $(\mathbb{U}_{\hbar}(\mathfrak{g}))_{\sigma}$, it follows eventually from this that $\langle \widetilde{J}_{\hbar} \rangle_{\mathbb{k}[[\hbar]]}^{\sigma}$ is a Hopf $\mathbb{k}[[\hbar]]$ -subalgebra inside $(\mathbb{U}_{\hbar}(\mathfrak{g}))_{\sigma}$.

By repeated use of (3.39) or (3.40) alike, we find that $\langle \widetilde{J}_{\hbar} \rangle_{\mathbb{k}[[\hbar]]}^{\sigma} \subseteq U_{\hbar} = \langle \widetilde{J}_{\hbar} \rangle_{\mathbb{k}[[\hbar]]}$.

Now observe that the original product “ \cdot ” in $U_{\hbar} := U_{\hbar}(\mathfrak{g})$ and $\mathbb{U}_{\hbar}(\mathfrak{g})$ can be obtained from “ \cdot_{σ} ” through deformation via the inverse 2-cocycle σ^{-1} . Thanks to this, we can reverse the roles of $U_{\hbar} = \langle \widetilde{J}_{\hbar} \rangle_{\mathbb{k}[[\hbar]]}$ and $\langle \widetilde{J}_{\hbar} \rangle_{\mathbb{k}[[\hbar]]}^{\sigma}$ in the previous construction (with some care), thus eventually achieving the converse inclusion $\langle \widetilde{J}_{\hbar} \rangle_{\mathbb{k}[[\hbar]]} \subseteq \langle \widetilde{J}_{\hbar} \rangle_{\mathbb{k}[[\hbar]]}^{\sigma}$. Therefore $\langle \widetilde{J}_{\hbar} \rangle_{\mathbb{k}[[\hbar]]} \subseteq \langle \widetilde{J}_{\hbar} \rangle_{\mathbb{k}[[\hbar]]}^{\sigma}$, which in particular implies that $U_{\hbar} = \langle \widetilde{J}_{\hbar} \rangle_{\mathbb{k}[[\hbar]]}$ is closed for the σ -product, q.e.d. \square

Definition 3.3.6. With assumptions as in Theorem 3.3.5, the new QUEA obtained from $U_{\hbar}(\mathfrak{g})$ through the process of 2-cocycle deformation by σ of $\mathbb{U}_{\hbar}(\mathfrak{g})$ followed by restriction will be called *the quasi-2-cocycle deformation of $U_{\hbar}(\mathfrak{g})$ by σ* , and it will be denoted by $U_{\hbar}(\mathfrak{g})_{\sigma}$.

To complete our analysis, next result sheds light onto the new, quasi-2-cocycle deformed QUEA $U_{\hbar}(\mathfrak{g})_{\sigma}$, describing in detail its semiclassical limit:

Theorem 3.3.7. *Let $U_{\hbar}(\mathfrak{g})$ be a QUEA over the Lie bialgebra $\mathfrak{g} = (\mathfrak{g}; [\ , \], \delta)$. Let σ be a quasi-2-cocycle for $U_{\hbar}(\mathfrak{g})$, so $\sigma = \exp_*(\hbar^{-1}\chi)$ for some $\chi \in (U_{\hbar}(\mathfrak{g})^{\widehat{\otimes} 2})^*$ with $\chi(z, 1) = 0 = \chi(1, z)$ for $z \in U_{\hbar}(\mathfrak{g})$. Set also $\chi_a := \chi - \chi_{2,1}$. Then:*

(a) χ is antisymmetric, i.e. $\chi_{2,1} = -\chi$, iff σ is orthogonal, i.e. $\sigma_{2,1} = \sigma^{-1}$;

(b) the \mathbb{k} -linear map $\gamma := \chi_a \left(\text{mod } \hbar (U_{\hbar}(\mathfrak{g})^{\widehat{\otimes} 2})^* \right) \Big|_{\mathfrak{g} \otimes \mathfrak{g}}$ from $\mathfrak{g} \otimes \mathfrak{g}$ to \mathbb{k} is antisymmetric 2-cocycle for the Lie bialgebra \mathfrak{g} ;

(c) the quasi-2-cocycle deformation $(U_{\hbar}(\mathfrak{g}))_{\sigma}$ of $U_{\hbar}(\mathfrak{g})$ is a QUEA for the Lie bialgebra $\mathfrak{g}_{\gamma} = (\mathfrak{g}; [\ , \]_{\gamma}, \delta)$ which is the deformation of \mathfrak{g} by the 2-cocycle γ ; in a nutshell, we have $(U_{\hbar}(\mathfrak{g}))_{\sigma} \cong U_{\hbar}(\mathfrak{g}_{\gamma})$.

In particular, if σ is $\mathbb{k}[[\hbar]]$ -valued — i.e., it is an ordinary 2-cocycle for the Hopf $\mathbb{k}[[\hbar]]$ -algebra $U_{\hbar}(\mathfrak{g})$ — or equivalently $\chi \in \hbar (U_{\hbar}(\mathfrak{g})^{\widehat{\otimes} 2})^*$, then we have just $\gamma = 0$ and $(U_{\hbar}(\mathfrak{g}))_{\sigma} \cong U_{\hbar}(\mathfrak{g}_{\gamma}) = U_{\hbar}(\mathfrak{g})$.

Proof. (a) This follows from claim (c) in Lemma 3.3.2.

(b) We are interested in the restriction to $\mathfrak{g} \otimes \mathfrak{g}$ of the specialization of σ modulo \hbar . So we start with $a, b, c \in \mathfrak{g}$, that we realize as $a = a \pmod{\hbar U_{\hbar}}$, $b = b \pmod{\hbar U_{\hbar}}$ and $c = c \pmod{\hbar U_{\hbar}}$ for some “lifts” $a, b, c \in U_{\hbar}$. By the identity $U_{\hbar} = (U'_{\hbar})^{\vee}$ and by Lemma 3.3 in [Ga1], we can choose the lifts a, b and c belong to $\widetilde{J}_{\hbar} := \hbar^{-1}J'_{\hbar}$, so that $a' := \hbar a$, $b' := \hbar b$ and $c' := \hbar c$ belong to J'_{\hbar} .

As σ is a normalized Hopf 2-cocycle for U_{\hbar} , it must obey the equality

$$\sigma(b'_{(1)}, c'_{(1)}) \sigma(a', b'_{(2)} c'_{(2)}) = \sigma(a'_{(1)}, b'_{(1)}) \sigma(a'_{(2)} b'_{(2)}, c') \quad (3.41)$$

Let us focus on the left hand side of (3.41). Expanding the exponential we get

$$\begin{aligned} \sigma(b'_{(1)}, c'_{(1)}) \sigma(a', b'_{(2)} c'_{(2)}) &= \sum_{n, m \geq 0} \frac{\hbar^{-(n+m)}}{n! m!} \chi^{*n}(b'_{(1)}, c'_{(1)}) \chi^{*m}(a', b'_{(2)} c'_{(2)}) = \\ &= \epsilon(a') \epsilon(b') \epsilon(c') + \hbar^{-1} \chi(b', c') \epsilon(a') + \hbar^{-1} \chi(a', b' c') + \\ &+ \hbar^{-2} \chi(b'_{(1)}, c'_{(1)}) \chi(a', b'_{(2)} c'_{(2)}) + \hbar^{-2} 2^{-1} \chi^{*2}(b', c') \epsilon(a') + \\ &+ \hbar^{-2} 2^{-1} \chi^{*2}(a', b' c') + \sum_{n+m \geq 3} \frac{\hbar^{-(n+m)}}{n! m!} \chi^{*n}(b'_{(1)}, c'_{(1)}) \chi^{*m}(a', b'_{(2)} c'_{(2)}) \end{aligned}$$

Then, noting that $\epsilon(a') = \epsilon(b') = \epsilon(c') = 0$ by construction, and analyzing all other summands as in the proof of claim Theorem 3.3.5, we obtain

$$\begin{aligned} \sigma(b'_{(1)}, c'_{(1)}) \sigma(a', b'_{(2)} c'_{(2)}) &= \hbar^{-1} \chi(a', b' c') + \\ &+ \hbar^{-2} \chi(b'_{(1)}, c'_{(1)}) \chi(a', b'_{(2)} c'_{(2)}) + \hbar^{-2} 2^{-1} \chi^{*2}(a', b' c') + \quad (3.42) \\ &+ (\text{sum of all terms with } n + m \geq 3) \end{aligned}$$

Writing $z' = z'^+ + \epsilon(z')$ and using that $\chi(z, 1) = 0 = \chi(1, z)$ and

$$\chi(x'_{(1)}, y'_{(1)} z'_{(1)}) \chi(x'_{(2)}, y'_{(2)} \epsilon(z'_{(2)})) = \chi(x'_{(1)}, y'_{(1)} z') \chi(x'_{(2)}, y'_{(2)})$$

we have that

$$\begin{aligned} \chi^{*2}(a', b'c') &= \chi(a'_{(1)}, b'_{(1)}c'_{(1)}) \chi(a'_{(2)}, b'_{(2)}c'_{(2)}) = \\ &= \chi(a'^+_{(1)}, b'^+_{(1)}c'^+_{(1)}) \chi(a'^+_{(2)}, b'^+_{(2)}c'^+_{(2)}) + \chi(a'^+_{(1)}, b'^+_{(1)}) \chi(a'^+_{(2)}, b'^+_{(2)}c') + \\ &\quad + \chi(a'^+_{(1)}, c'^+_{(1)}) \chi(a'^+_{(2)}, b'c'^+_{(2)}) + \chi(a'^+_{(1)}, b'c'^+_{(1)}) \chi(a'^+_{(2)}, c'^+_{(2)}) + \\ &\quad + \chi(a'^+_{(1)}, b') \chi(a'^+_{(2)}, c') + \chi(a'^+_{(1)}, b'^+_{(1)}c') \chi(a'^+_{(2)}, b'^+_{(2)}) + \chi(a'^+_{(1)}, c') \chi(a'^+_{(2)}, b') \end{aligned}$$

Now, taking into account that $z' = \hbar z$ and $z'^+_{(i)} \in \hbar U_{\hbar}$, we may re-write the expression above as

$$\chi^{*2}(a', b'c') = \hbar^4 \chi(a_{(1)}, b) \chi(a_{(2)}, c) + \hbar^4 \chi(a_{(1)}, c) \chi(a_{(2)}, b) + \mathcal{O}(\hbar^5)$$

Performing a similar analysis on the term $\chi(b'_{(1)}, c'_{(1)}) \chi(a', b'_{(2)}c'_{(2)})$ we get

$$\chi(b'_{(1)}, c'_{(1)}) \chi(a', b'_{(2)}c'_{(2)}) = \hbar^4 \chi(b, c_{(1)}) \chi(a, c_{(2)}) + \hbar^4 \chi(b_{(1)}, c) \chi(a, b_{(2)}) + \mathcal{O}(\hbar^5)$$

Moreover, with a similar (yet easier) analysis one finds also that

$$\chi^{*n}(b'_{(1)}, c'_{(1)}) \chi^{*m}(a', b'_{(2)}c'_{(2)}) \in \hbar^{+2(n+m)} \mathbb{k}[[\hbar]]$$

for all $n+m \geq 3$, so that the (last) summand “(sum of all terms with $n+m \geq 3$)” in (3.42) is of type $\mathcal{O}(\hbar^{n+m}) = \mathcal{O}(\hbar^3)$. Putting all together in (3.42) we find

$$\begin{aligned} \hbar^3 \sigma(b_{(1)}, c_{(1)}) \sigma(a, b_{(2)}c_{(2)}) &= \\ &= \hbar^2 \chi(a, bc) + \hbar^2 \chi(b, c_{(1)}) \chi(a, c_{(2)}) + \hbar^2 \chi(b_{(1)}, c) \chi(a, b_{(2)}) + \\ &\quad + \hbar^2 2^{-1} \chi(a_{(1)}, b) \chi(a_{(2)}, c) + \hbar^2 2^{-1} \chi(a_{(1)}, c) \chi(a_{(2)}, b) + \mathcal{O}(\hbar^3) \end{aligned}$$

An analogous treatment of the right hand side of (3.41) yields

$$\begin{aligned} \hbar^3 \sigma(a_{(1)}, b_{(1)}) \sigma(a_{(2)}b_{(2)}, c) &= \\ &= \hbar^2 \chi(ab, c) + \hbar^2 \chi(a, b_{(1)}) \chi(b_{(2)}, c) + \hbar^2 \chi(a_{(1)}, b) \chi(a_{(2)}, c) + \\ &\quad + \hbar^2 2^{-1} \chi(a, c_{(1)}) \chi(b, c_{(2)}) + \hbar^2 2^{-1} \chi(b, c_{(1)}) \chi(a, c_{(2)}) + \mathcal{O}(\hbar^3) \end{aligned}$$

Altogether, this implies that

$$\begin{aligned} \chi(a, bc) + \chi(b, c_{(1)}) \chi(a, c_{(2)}) + \chi(b_{(1)}, c) \chi(a, b_{(2)}) + \\ + 2^{-1} \chi(a_{(1)}, b) \chi(a_{(2)}, c) + 2^{-1} \chi(a_{(1)}, c) \chi(a_{(2)}, b) &\equiv_{\hbar} \\ &\equiv_{\hbar} \chi(ab, c) + \chi(a, b_{(1)}) \chi(b_{(2)}, c) + \chi(a_{(1)}, b) \chi(a_{(2)}, c) + \\ &\quad + 2^{-1} \chi(a, c_{(1)}) \chi(b, c_{(2)}) + 2^{-1} \chi(b, c_{(1)}) \chi(a, c_{(2)}) \end{aligned}$$

where (again) \equiv_{\hbar} stands for “congruent modulo $\hbar \mathbb{k}[[\hbar]]$ ”, that we re-write as

$$\begin{aligned} \chi(a, bc) + \chi(b, c_{(1)}) \chi(a, c_{(2)}) + \chi(b_{(1)}, c) \chi(a, b_{(2)}) + \\ + 2^{-1} \chi(a_{(1)}, b) \chi(a_{(2)}, c) + 2^{-1} \chi(a_{(1)}, c) \chi(a_{(2)}, b) - \\ - \chi(ab, c) - \chi(a, b_{(1)}) \chi(b_{(2)}, c) - \chi(a_{(1)}, b) \chi(a_{(2)}, c) - \\ - 2^{-1} \chi(a, c_{(1)}) \chi(b, c_{(2)}) - 2^{-1} \chi(b, c_{(1)}) \chi(a, c_{(2)}) &\equiv_{\hbar} 0 \end{aligned} \tag{3.43}$$

Consider now the action of the group algebra $\mathbb{k}[\mathbb{S}_3]$ of the symmetric group \mathbb{S}_3 on $(U_{\hbar}^{\otimes 3})^*$ given by $\sigma \cdot \varphi(a, b, c) := \varphi(\sigma^{-1} \cdot (a, b, c))$, where the action of $\mathbb{k}[\mathbb{S}_3]$ on $U_{\hbar}^{\otimes 3}$ is the natural one that permutes the tensor factors. Then we let the antisymmetrizer Alt_3 act on both sides of (3.43): using that $\gamma := \chi_a \left(\text{mod } \hbar (U_{\hbar}(\mathfrak{g})^{\otimes 2})^* \right) \Big|_{\mathfrak{g} \otimes \mathfrak{g}}$ and that $a \pmod{\hbar U_{\hbar}} = a$, $b \pmod{\hbar U_{\hbar}} = b$ and $c \pmod{\hbar U_{\hbar}} = c$, a straightforward calculation eventually yields

$$\partial_*(\gamma) + \text{c.p.} + [[\gamma, \gamma]]_* = 0$$

This means exactly that γ is a 2-cocycle for the Lie bialgebra \mathfrak{g} — according to Definition 2.1.5 — that is obviously antisymmetric (by construction), q.e.d.

(c) First of all, we start by noting that $U_{\hbar}(\mathfrak{g})_{\sigma} := (U_{\hbar}(\mathfrak{g}))_{\sigma}$ is equal to $U_{\hbar}(\mathfrak{g})$ as a counital $\mathbb{k}[[\hbar]]$ -coalgebra (by construction), but with the new product defined by

$$m_{\sigma}(a, b) = a \cdot_{\sigma} b = \sigma(a_{(1)}, b_{(1)}) a_{(2)} b_{(2)} \sigma^{-1}(a_{(3)}, b_{(3)}) \quad \forall a, b \in U_{\hbar}(\mathfrak{g})$$

In particular, the $\mathbb{k}[[\hbar]]$ -module $U_{\hbar}(\mathfrak{g})_{\sigma} = U_{\hbar}(\mathfrak{g})$ is still topologically free, so that $U_{\hbar}(\mathfrak{g})_{\sigma}$ is again a Hopf algebra in \mathcal{T}_{\otimes} , cf. §2.3.1. Moreover, its semiclassical limit $\overline{U_{\hbar}(\mathfrak{g})_{\sigma}} := U_{\hbar}(\mathfrak{g})_{\sigma} / \hbar U_{\hbar}(\mathfrak{g})_{\sigma}$ as a coalgebra is the same as that of $U_{\hbar}(\mathfrak{g})$; hence it is again cocommutative connected (as these properties are not affected by 2-cocycle deformations). Thus by Milnor-Moore Theorem we have $\overline{U_{\hbar}(\mathfrak{g})_{\sigma}} = U(\widehat{\mathfrak{g}})$, where $\widehat{\mathfrak{g}} = \text{Prim}(\overline{U_{\hbar}(\mathfrak{g})_{\sigma}})$ is the space of primitive elements in $\overline{U_{\hbar}(\mathfrak{g})_{\sigma}}$, and as such it coincides with $\text{Prim}(\overline{U_{\hbar}(\mathfrak{g})}) = \text{Prim}(U(\mathfrak{g})) = \mathfrak{g}$ as a Lie coalgebra; its Lie algebra structure, on the other hand, does depend on σ . Altogether, this shows that $U_{\hbar}(\mathfrak{g})_{\sigma}$ is indeed a QUEA, whose semiclassical limit is $U(\widehat{\mathfrak{g}})$; then we are only left to prove that the Lie bracket on $\widehat{\mathfrak{g}}$ coincides with that of \mathfrak{g}_{γ} , while also proving that γ is an antisymmetric 2-cocycle for the Lie bialgebra \mathfrak{g} .

The Lie bracket in $\widehat{\mathfrak{g}}$ is given by the commutator inside $U(\widehat{\mathfrak{g}}) = \overline{U_{\hbar}(\mathfrak{g})_{\sigma}}$, so we denote it by $[a, b]_{\sigma} = a \cdot_{\sigma} b - b \cdot_{\sigma} a$ (for all $a, b \in \mathfrak{g}$), where \cdot_{σ} is the product in $U(\widehat{\mathfrak{g}}) = \overline{U_{\hbar}(\mathfrak{g})_{\sigma}}$ induced by the $(\sigma$ -deformed) product in $U_{\hbar}(\mathfrak{g})_{\sigma}$. Therefore, we will compute such a commutator as the coset modulo $\hbar U_{\hbar}$ of a commutator in U_{\hbar} , namely $[a, b]_{\sigma} = a \cdot_{\sigma} b - b \cdot_{\sigma} a = a \cdot_{\sigma} b - b \cdot_{\sigma} a \pmod{\hbar U_{\hbar}}$, where a and b , like in the proof of claim (c), are lifts of a and b — i.e., $a \pmod{\hbar U_{\hbar}} = a$ and $b \pmod{\hbar U_{\hbar}} = b$ — such that $a' := \hbar a \in J'_{\hbar}$ and $b' := \hbar b \in J'_{\hbar}$.

We re-start back from (3.30), which now gives (taking into account all the analysis carried out there, with $A = 1 = B$)

$$\begin{aligned} a \cdot_{\sigma} b - b \cdot_{\sigma} a &\equiv_{\hbar} a \cdot b - b \cdot a + \\ &+ \hbar^{-3} \left(\chi(a'_{(1)}, b'_{(1)}) (\widehat{a}'_{(2)} b'_{(2)} + a'_{(2)} \widehat{b}'_{(2)}) - \chi(b'_{(1)}, a'_{(1)}) (\widehat{b}'_{(2)} a'_{(2)} + b'_{(2)} \widehat{a}'_{(2)}) \right) - \\ &- \hbar^{-3} \left((\widehat{a}'_{(1)} b'_{(1)} + a'_{(1)} \widehat{b}'_{(1)}) \chi(a'_{(2)}, b'_{(2)}) - (\widehat{b}'_{(1)} a'_{(1)} + b'_{(1)} \widehat{a}'_{(1)}) \chi(b'_{(2)}, a'_{(2)}) \right) \end{aligned}$$

Second, letting $\chi_a := \chi - \chi_{2,1}$, the previous formula greatly simplifies into

$$\begin{aligned} a \cdot_{\sigma} b - b \cdot_{\sigma} a &= a \cdot b - b \cdot a + \hbar^{-3} \chi_a(a'_{(1)}^+, b'_{(1)}^+) (\widehat{a}'_{(2)} b'_{(2)} + a'_{(2)} \widehat{b}'_{(2)}) + \\ &\quad + \hbar^{-3} (\widehat{b}'_{(1)} a'_{(1)} + b'_{(1)} \widehat{a}'_{(1)}) \chi_a(b'_{(2)}^+, a'_{(2)}^+) + \mathcal{O}(\hbar) = \\ &= a \cdot b - b \cdot a + \hbar^{-3} \left(\chi_a(a'_{(1)}^+, b') a'_{(2)}^+ - \chi_a(a'_{(2)}^+, b') a'_{(1)}^+ \right) - \\ &\quad - \hbar^{-3} \left(\chi_a(b'_{(1)}^+, a') b'_{(2)}^+ - \chi_a(b'_{(2)}^+, a') b'_{(1)}^+ \right) + \mathcal{O}_1(\hbar) \end{aligned}$$

Now, let us write $z' = \hbar z$ for all $z \in \{a, b\}$: then the last formula turns into

$$\begin{aligned} a \cdot_{\sigma} b - b \cdot_{\sigma} a &= a \cdot b - b \cdot a + \hbar^{-1} \left(\chi_a(a'_{(1)}^+, b) a'_{(2)}^+ - \chi_a(a'_{(2)}^+, b) a'_{(1)}^+ \right) - \\ &\quad - \hbar^{-1} \left(\chi_a(b'_{(1)}^+, a) b'_{(2)}^+ - \chi_a(b'_{(2)}^+, a) b'_{(1)}^+ \right) + \mathcal{O}_1(\hbar) \end{aligned} \quad (3.44)$$

Here we recall that, working with a QUEA, for $c \in \{a, b\}$ we have

$$\Delta(c) = c \otimes 1 + 1 \otimes c + c_{(1)}^+ \otimes c_{(2)}^+ + \mathcal{O}_2(\hbar^2), \quad c_{(1)}^+ \otimes c_{(2)}^+ \in \hbar U_{\hbar}^{\widehat{\otimes} 2}$$

and moreover — for every $c \in \{a, b\}$ and $\mathfrak{c} \in \{\mathfrak{a}, \mathfrak{b}\}$, so that c is a lift of \mathfrak{c} —

$$\overline{\hbar^{-1}(c_{(1)}^+ \otimes c_{(2)}^+ - c_{(2)}^+ \otimes c_{(1)}^+)} = \delta(\mathfrak{c}) =: \mathfrak{c}_{[1]} \otimes \mathfrak{c}_{[2]} \quad (3.45)$$

where hereafter any “overlined” object stands for “its coset modulo \hbar ”; in addition, we recall also that χ_a is antisymmetric. Then (3.44) and (3.45) altogether yield

$$\begin{aligned} [\mathfrak{a}, \mathfrak{b}]_{\sigma} &= \overline{\mathfrak{a} \cdot_{\sigma} \mathfrak{b} - \mathfrak{b} \cdot_{\sigma} \mathfrak{a}} = \overline{\mathfrak{a} \cdot \mathfrak{b} - \mathfrak{b} \cdot \mathfrak{a}} = \\ &= \overline{\mathfrak{a} \cdot \mathfrak{b} - \mathfrak{b} \cdot \mathfrak{a} + \hbar^{-1} \left(\chi_a(a'_{(1)}^+, b) a'_{(2)}^+ - \chi_a(a'_{(2)}^+, b) a'_{(1)}^+ \right) -} \\ &\quad - \overline{\hbar^{-1} \left(\chi_a(b'_{(1)}^+, a) b'_{(2)}^+ - \chi_a(b'_{(2)}^+, a) b'_{(1)}^+ \right)} = \\ &= [\mathfrak{a}, \mathfrak{b}] + \gamma(\mathfrak{a}_{[1]}, \mathfrak{b}) \mathfrak{a}_{[2]} - \gamma(\mathfrak{b}_{[1]}, \mathfrak{a}) \mathfrak{b}_{[2]} = \\ &= [\mathfrak{a}, \mathfrak{b}] - \gamma(\mathfrak{a}_{[2]}, \mathfrak{b}) \mathfrak{a}_{[1]} - \gamma(\mathfrak{b}_{[1]}, \mathfrak{a}) \mathfrak{b}_{[2]} =: [\mathfrak{a}, \mathfrak{b}]_{\gamma} \end{aligned}$$

hence $[\mathfrak{a}, \mathfrak{b}]_{\sigma} = [\mathfrak{a}, \mathfrak{b}]_{\gamma}$ for all $\mathfrak{a}, \mathfrak{b} \in \mathfrak{g}$, in the sense of (2.5), and we are done.

Finally, if in particular σ is $\mathbb{k}[[\hbar]]$ -valued — i.e., it is an ordinary 2-cocycle for the Hopf $\mathbb{k}[[\hbar]]$ -algebra $U_{\hbar}(\mathfrak{g})$ — then we have $\chi = \hbar \log_*(\sigma) \in \hbar (U_{\hbar}(\mathfrak{g})^{\widehat{\otimes} 2})^*$, hence we have just $\gamma = 0$ and $(U_{\hbar}(\mathfrak{g}))_{\sigma} \cong U_{\hbar}(\mathfrak{g}_{\gamma}) = U_{\hbar}(\mathfrak{g})$. \square

Example 3.3.8. Some concrete examples of quasi-2-cocycles and deformation by them are treated in full depth in [GaGa2, Section 5.2]: they concern the wide family of *formal multiparameter QUEAs* that we already treated in Example 3.1.4. We then resume notations and formulas from there.

Fix $n \in \mathbb{N}_+$ and $I := \{1, \dots, n\}$. We choose a multiparameter matrix $P := (p_{i,j})_{i,j \in I} \in M_n(\mathbb{k}[[\hbar]])$ of Cartan type, with associated Cartan matrix A , a realization $\mathcal{R} := (\mathfrak{h}, \Pi, \Pi^{\vee})$ of it and the (topological) Hopf algebra $U_{P, \hbar}^{\mathcal{R}}(\mathfrak{g})$. Let $\{H_g\}_{g \in \mathcal{G}}$ be a $\mathbb{k}[[\hbar]]$ -basis in \mathfrak{h} , where \mathcal{G} is an index set with $|\mathcal{G}| = \text{rk}(\mathfrak{h}) = t$.

We consider special quasi-2-cocycles of $U_{P, \hbar}^{\mathcal{R}}(\mathfrak{g})$, called “toral” as they are induced from the quantum torus. Fix an antisymmetric, $\mathbb{k}[[\hbar]]$ -bilinear map $\chi : \mathfrak{h} \times \mathfrak{h} \longrightarrow$

$\mathbb{k}[[\hbar]]$, that corresponds to $X = (\chi_{g\gamma} = \chi(H_g, H_\gamma))_{g,\gamma \in \mathcal{G}} \in \mathfrak{so}_t(\mathbb{k}[[\hbar]])$. Any such map χ also induces uniquely an antisymmetric, $\mathbb{k}[[\hbar]]$ -bilinear map

$$\tilde{\chi}_U : U_{P,\hbar}^{\mathcal{R}}(\mathfrak{h}) \times U_{P,\hbar}^{\mathcal{R}}(\mathfrak{h}) \longrightarrow \mathbb{k}[[\hbar]]$$

as follows. By definition, $U_{P,\hbar}^{\mathcal{R}}(\mathfrak{h})$ is an \hbar -adically complete topologically free Hopf algebra isomorphic to $\widehat{S}_{\mathbb{k}[[\hbar]]}(\mathfrak{h}) := \widehat{\bigoplus_{n \in \mathbb{N}} S_{\mathbb{k}[[\hbar]]}^n(\mathfrak{h})}$, the \hbar -adic completion of the symmetric algebra $S_{\mathbb{k}[[\hbar]]}(\mathfrak{h}) = \bigoplus_{n \in \mathbb{N}} S_{\mathbb{k}[[\hbar]]}^n(\mathfrak{h})$. Then, $\tilde{\chi}_U$ is defined as the unique $\mathbb{k}[[\hbar]]$ -linear

(hence \hbar -adically continuous) map $U_{P,\hbar}^{\mathcal{R}}(\mathfrak{h}) \otimes U_{P,\hbar}^{\mathcal{R}}(\mathfrak{h}) \xrightarrow{\tilde{\chi}_U} \mathbb{k}[[\hbar]]$ such that

$$\begin{aligned} \tilde{\chi}_U(z, 1) &:= \epsilon(z) =: \tilde{\chi}_U(1, z) & \forall z \in \widehat{S}_{\mathbb{k}[[\hbar]]}(\mathfrak{h}) \\ \tilde{\chi}_U(x, y) &:= \chi(x, y) & \forall x, y \in S_{\mathbb{k}[[\hbar]]}^1(\mathfrak{h}) \\ \tilde{\chi}_U(x, y) &:= 0 & \forall x \in S_{\mathbb{k}}^r(\mathfrak{h}), y \in S_{\mathbb{k}}^s(\mathfrak{h}) : r, s \geq 1, r + s > 2 \end{aligned} \quad (3.46)$$

By construction, $\tilde{\chi}_U$ is a normalized Hochschild 2-cocycle on $U_{P,\hbar}^{\mathcal{R}}(\mathfrak{h})$, that is

$$\epsilon(x) \tilde{\chi}_U(y, z) - \tilde{\chi}_U(xy, z) + \tilde{\chi}_U(x, yz) - \tilde{\chi}_U(x, y) \epsilon(z) = 0 \quad \forall x, y, z \in U_{P,\hbar}^{\mathcal{R}}(\mathfrak{h})$$

By [GaGa2, Lemma 5.2.3], the convolution powers of $\tilde{\chi}_U$ satisfy the following property: for all $H_+, H_- \in \mathfrak{h}$ and $k, \ell, m \in \mathbb{N}_+$, we have

$$\tilde{\chi}_U^{*m}(H_+^k, H_-^\ell) = \begin{cases} \delta_{k,m} \delta_{\ell,m} (m!)^2 \chi(H_+, H_-)^m & \text{for } m \geq 1, \\ \delta_{k,0} \delta_{\ell,0} & \text{for } m = 0. \end{cases}$$

This allows one to define a quasi-2-cocycle χ_U as the unique $\mathbb{k}[[\hbar]]$ -linear map from $U_{P,\hbar}^{\mathcal{R}}(\mathfrak{h}) \otimes_{\mathbb{k}[[\hbar]]} U_{P,\hbar}^{\mathcal{R}}(\mathfrak{h})$ to $\mathbb{k}((\hbar))$ given by the exponentiation of $\hbar^{-1} 2^{-1} \tilde{\chi}_U$, i.e.

$$\chi_U := e^{\hbar^{-1} 2^{-1} \tilde{\chi}_U} = \sum_{m \geq 0} \hbar^{-m} \tilde{\chi}_U^{*m} / 2^m m!$$

By [GaGa2, Lemma 5.2.2], this χ_U is in fact a well-defined quasi-2-cocycle for $U_{P,\hbar}^{\mathcal{R}}(\mathfrak{h})$, in the sense of Definition 3.3.3. Moreover, one has, for all $H_+, H_- \in \mathfrak{h}$, and setting $K_\pm := e^{\hbar H_\pm}$,

$$\chi_U^{\pm 1}(H_+, H_-) = \pm \hbar^{-1} 2^{-1} \chi(H_+, H_-) \quad , \quad \chi_U(K_+, K_-) = e^{\hbar 2^{-1} \chi(H_+, H_-)}$$

Assume now that χ satisfies the additional requirement $\chi(S_i, -) = 0 = \chi(-, S_i)$ for all $i \in I$, where $S_i := 2^{-1}(T_i^+ + T_i^-)$ for all $i \in I$. In particular, one has that $\chi(T_i^+, T) = \chi(-T_i^-, T)$ and $\chi(T, T_i^+) = \chi(T, -T_i^-)$ for all $i \in I$ and $T \in \mathfrak{h}$. Then χ canonically induces a $\mathbb{k}[[\hbar]]$ -bilinear map $\bar{\chi} : \bar{\mathfrak{h}} \times \bar{\mathfrak{h}} \longrightarrow \mathbb{k}[[\hbar]]$, where $\bar{\mathfrak{h}} := \mathfrak{h}/\mathfrak{s}$ with $\mathfrak{s} := \text{Span}_{\mathbb{k}[[\hbar]]}(\{S_i\}_{i \in I})$, given by

$$\bar{\chi}(T' + \mathfrak{s}, T'' + \mathfrak{s}) := \chi(T', T'') \quad \forall T', T'' \in \mathfrak{h}$$

Now, replaying the construction above with $\bar{\mathfrak{h}}$ and $\bar{\chi}$ replacing \mathfrak{h} and χ , we can construct a normalized Hopf quasi-2-cocycle $\bar{\chi}_U : U_{P,\hbar}^{\mathcal{R}}(\bar{\mathfrak{h}}) \times U_{P,\hbar}^{\mathcal{R}}(\bar{\mathfrak{h}}) \longrightarrow \mathbb{k}((\hbar))$. Moreover, since $U_{P,\hbar}^{\mathcal{R}}(\bar{\mathfrak{h}}) \cong \widehat{S}_{\mathbb{k}[[\hbar]]}(\bar{\mathfrak{h}})$, there exists a unique Hopf algebra epimorphism $\pi : U_{P,\hbar}^{\mathcal{R}}(\mathfrak{g}) \longrightarrow U_{P,\hbar}^{\mathcal{R}}(\bar{\mathfrak{h}})$ given by $\pi(E_i) := 0$, $\pi(F_i) := 0$ — for $i \in I$ — and $\pi(T) := (T + \mathfrak{s}) \in \bar{\mathfrak{h}} \subseteq U_{P,\hbar}^{\mathcal{R}}(\bar{\mathfrak{h}})$ — for $T \in \mathfrak{h}$. Then we consider

$$\sigma_\chi := \bar{\chi}_U \circ (\pi \times \pi) : U_{P,\hbar}^{\mathcal{R}}(\mathfrak{g}) \times U_{P,\hbar}^{\mathcal{R}}(\mathfrak{g}) \longrightarrow \mathbb{k}((\hbar))$$

which is *automatically* a normalized, $\mathbb{k}[[\hbar]]$ -valued Hopf quasi-2-cocycle on $U_{P,\hbar}^{\mathcal{R}}(\mathfrak{g})$.

By Theorem 3.3.7, one may define a “deformed product” on $U_{P,\hbar}^{\mathcal{R}}(\mathfrak{g})$ using σ_χ hereafter denoted by $\dot{\sigma}_\chi$. Write $X^{(n)\sigma_\chi} = X \dot{\sigma}_\chi \cdots \dot{\sigma}_\chi X$ for the n -th power of any $X \in U_{P,\hbar}^{\mathcal{R}}(\mathfrak{g})$ with respect to this deformed product.

Directly from definitions, sheer computation yields the following formulas, relating the deformed product with the old one (for all $T', T'', T \in \mathfrak{h}$, $i, j \in I$):

$$\begin{aligned} T' \dot{\sigma}_\chi T'' &= T' T'' \quad , \quad E_i \dot{\sigma}_\chi F_j = E_i F_j \quad , \quad F_j \dot{\sigma}_\chi E_i = F_j E_i \\ T \dot{\sigma}_\chi E_j &= T E_j + 2^{-1} \chi(T, T_j^+) E_j \quad , \quad E_j \dot{\sigma}_\chi T = E_j T + 2^{-1} \chi(T_j^+, T) E_j \\ T \dot{\sigma}_\chi F_j &= T F_j + 2^{-1} \chi(T, T_j^-) F_j \quad , \quad F_j \dot{\sigma}_\chi T = F_j T + 2^{-1} \chi(T_j^-, T) F_j \\ E_i^{(m)\sigma_\chi} &= \prod_{\ell=1}^{m-1} \sigma_\chi \left(e^{+\hbar \ell T_i^+}, e^{+\hbar T_i^+} \right) E_i^m = E_i^m \\ E_i^m \dot{\sigma}_\chi E_j^n &= \sigma_\chi \left(e^{+\hbar m T_i^+}, e^{+\hbar n T_j^+} \right) E_i^m E_j^n = e^{+\hbar m n 2^{-1} \dot{\chi}_{ij}} E_i^m E_j^n \\ E_i^{(m)\sigma_\chi} \dot{\sigma}_\chi E_j \dot{\sigma}_\chi E_k^{(n)\sigma_\chi} &= \left(\prod_{\ell=1}^{m-1} \sigma_\chi \left(e^{+\hbar \ell T_i^+}, e^{+\hbar T_i^+} \right) \right) \left(\prod_{t=1}^{n-1} \sigma_\chi \left(e^{+\hbar t T_k^+}, e^{+\hbar T_k^+} \right) \right) \\ &\quad \cdot \sigma_\chi \left(e^{+\hbar m T_i^+}, e^{+\hbar T_j^+} \right) \sigma_\chi \left(e^{+\hbar(m T_i^+ + T_j^+)}, e^{+\hbar n T_k^+} \right) E_i^m E_j E_k^n \\ F_i^{(m)\sigma_\chi} &= \prod_{\ell=1}^{m-1} \sigma_\chi^{-1} \left(e^{-\hbar \ell T_i^-}, e^{-\hbar T_i^-} \right) F_i^m = F_i^m \\ F_i^m \dot{\sigma}_\chi F_j^n &= \sigma_\chi^{-1} \left(e^{-\hbar m T_i^-}, e^{-\hbar n T_j^-} \right) F_i^m F_j^n = e^{-\hbar m n 2^{-1} \dot{\chi}_{ij}} F_i^m F_j^n \\ F_i^{(m)\sigma_\chi} \dot{\sigma}_\chi F_j \dot{\sigma}_\chi F_k^{(n)\sigma_\chi} &= \left(\prod_{\ell=1}^{m-1} \sigma_\chi^{-1} \left(e^{-\hbar \ell T_i^-}, e^{-\hbar T_i^-} \right) \right) \left(\prod_{t=1}^{n-1} \sigma_\chi^{-1} \left(e^{-\hbar t T_k^-}, e^{-\hbar T_k^-} \right) \right) \\ &\quad \cdot \sigma_\chi^{-1} \left(e^{-\hbar m T_i^-}, e^{-\hbar T_j^-} \right) \sigma_\chi^{-1} \left(e^{-\hbar(m T_i^- + T_j^-)}, e^{-\hbar n T_k^-} \right) F_i^m F_j F_k^n \end{aligned}$$

Fix now $\dot{X} := \left(\dot{\chi}_{ij} = \chi(T_i^+, T_j^+) \right)_{i,j \in I}$ and define the multiparameter matrix

$$P_{(\chi)} := P + \dot{X} = \left(p_{ij}^{(\chi)} := p_{ij} + \dot{\chi}_{ij} \right)_{i,j \in I} \quad , \quad \Pi_{(\chi)} := \left\{ \alpha_i^{(\chi)} := \alpha_i \pm \chi(-, T_i^\pm) \right\}_{i \in I}$$

It turns out that $P_{(\chi)}$ is a matrix of Cartan type — the same of P indeed — and $\mathcal{R}_{(\chi)} = (\mathfrak{h}, \Pi_{(\chi)}, \Pi^\vee)$ is a realization of it. Moreover, by [GaGa2, Theorem 5.2.12], there exists an isomorphism of topological Hopf algebras

$$(U_{P,\hbar}^{\mathcal{R}}(\mathfrak{g}))_{\sigma_\chi} \cong U_{P_{(\chi)},\hbar}^{\mathcal{R}_{(\chi)}}(\mathfrak{g})$$

which is the identity on generators. In short, every toral quasi-2-cocycle deformation of a FoMpQUEA is another FoMpQUEA, whose multiparameter $P_{(\chi)}$ and realization $\mathcal{R}_{(\chi)}$ depend on the original P and \mathcal{R} , as well as on χ . Moreover, under some mild restrictions on the realizations, one proves that the FoMpQUEA $U_{P,\hbar}^{\mathcal{R}}(\mathfrak{g})$ is isomorphic to a toral quasi-2-cocycle deformation of the Drinfeld’s standard double QUEA, see [GaGa2, Theorem 5.2.14].

About the semiclassical limit, we have the following. Taking everything modulo \hbar , the map $\chi : \mathfrak{h} \times \mathfrak{h} \longrightarrow \mathbb{k}[[\hbar]]$ defines a similar antisymmetric, \mathbb{k} -bilinear map $\gamma := (\chi \bmod \hbar) : \mathfrak{h}_0 \times \mathfrak{h}_0 \longrightarrow \mathbb{k}$ — where $\mathfrak{h}_0 := \mathfrak{h} / \hbar \mathfrak{h} = \bar{\mathfrak{h}}$. Out of γ one constructs a toral 2-cocycle $\gamma_{\mathfrak{g}}$ for the Lie bialgebra $\mathfrak{g}_{\bar{P}}^{\bar{\mathcal{R}}}$, and out of it the 2-cocycle deformed Lie bialgebra $(\mathfrak{g}_{\bar{P}}^{\bar{\mathcal{R}}})_{\gamma_{\mathfrak{g}}}$. Similarly as above, out of γ we get the multiparameter

matrix $P_{(\gamma)}$ and its realization $\mathcal{R}_{(\gamma)}$: then by construction $P_{(\gamma)} = \bar{P}_{(x)}$ and $\mathcal{R}_{(\gamma)} = \bar{\mathcal{R}}_{(x)}$. Attached to these we have $U_{P_{(x)}, \hbar}^{\mathcal{R}_{(\gamma)}}(\mathfrak{g})$ and $\mathfrak{g}_{P_{(\gamma)}}^{\mathcal{R}_{(\gamma)}} = \mathfrak{g}_{\bar{P}_{(x)}}^{\bar{\mathcal{R}}_{(x)}}$, again connected via quantization/specialization, and $\mathfrak{g}_{P_{(\gamma)}}^{\mathcal{R}_{(\gamma)}} \cong (\mathfrak{g}_{\bar{P}}^{\bar{\mathcal{R}}})_{\gamma_{\mathfrak{g}}}$ as Lie bialgebras. Actually, one has that “deformation by (quasi-)2-cocycle commutes with specialization”, see [GaGa2, Theorem 6.2.4]: with assumptions as above, we have that $(U_{P, \hbar}^{\mathcal{R}}(\mathfrak{g}))_{\sigma_x}$ is a quantized universal enveloping algebra, with semiclassical limit $U\left((\mathfrak{g}_{\bar{P}}^{\bar{\mathcal{R}}})_{\gamma_{\mathfrak{g}}}\right) \cong U\left(\mathfrak{g}_{P_{(\gamma)}}^{\mathcal{R}_{(\gamma)}}\right)$.

Remark 3.3.9. It is important to stress that *our notion of quasi-2-cocycle did not come out of the blue*, but rather was suggested by the previous example. Indeed, the authors first “met” these objects when studying polynomial-type QUEAs $U_q(\mathfrak{g})$ — i.e., QUEAs “à la Jimbo-Lusztig”, defined over $\mathbb{k}[q, q^{-1}]$: these are standard Hopf algebras (no topology is involved), to which one can apply deformation by 2-cocycles and then obtain some “multiparameter QUEAs” — cf. [GaGa1], §4.2. Every such *polynomial* $U_q(\mathfrak{g})$ can be realized as a Hopf subalgebra of a *formal* $U_{\hbar}(\mathfrak{g})$, hence it makes sense to try and extend the 2-cocycle and the associated deformation procedure used for $U_q(\mathfrak{g})$ to the larger Hopf algebra $U_{\hbar}(\mathfrak{g})$. When we fulfilled this task — in [GaGa2] — what we actually found was that the unique extension of the 2-cocycle of $U_q(\mathfrak{g})$ to $U_{\hbar}(\mathfrak{g})$ actually is a *quasi-2-cocycle* (and not a 2-cocycle any more), yet despite this the deformation procedure does extend from $U_q(\mathfrak{g})$ to the whole $U_{\hbar}(\mathfrak{g})$. Thus the very notion of “quasi-2-cocycle” and the associated deformation procedure showed up as something real from this concrete example.

3.4. Deformations by quasi-twist of QFSHA’s.

In this subsection we consider deformations by twist of QFSHA’s, but again “stretching the standard recipe”, much like in §3.3: in fact, rather than twists in the usual sense we consider some special twist elements belonging to the scalar extension from $\mathbb{k}[[\hbar]]$ to $\mathbb{k}((\hbar))$ of our QFSHA. For these elements — that we call “quasi-twists” — nothing ensures *a priori* that the deformation recipe would properly work on the given QFSHA — as it is defined on $\mathbb{k}[[\hbar]]$; nevertheless, we eventually find that this is indeed the case. In other words, we prove — with a parallel result to Theorem 3.3.5 — that *the standard procedure of deformation by twist for QFSHA’s can be extended (beyond its natural borders) to the case of quasi-twist elements*.

We begin with a couple of technical lemmas:

Lemma 3.4.1. *Let $F_{\hbar}[[G]]$ be a QFSHA, and $F_{\hbar}[[G]]^{\vee}$ the associated QUEA defined in §2.4. Let $\varphi \in J_{\hbar}^2$, with $J_{\hbar} := \text{Ker}(\epsilon_{F_{\hbar}[[G]])}$. Then:*

- (a) $F := \exp(\hbar^{-1}\varphi)$ is a well-defined element in $F_{\hbar}[[G]]^{\vee}$;
- (b) $\text{Ad}(F)(f) := F \cdot f \cdot F^{-1} \in F_{\hbar}[[G]]$ for all $f \in F_{\hbar}[[G]]$, so that the adjoint action of F onto $F_{\hbar}[[G]]^{\vee}$ actually restricts to $F_{\hbar}[[G]]$.

Proof. (a) The assumption $\varphi \in J_{\hbar}^2$ implies $\varphi \in \hbar^2 (J_{\hbar}^{\vee})^2 \subseteq \hbar^2 F_{\hbar}[[G]]^{\vee}$, where $J_{\hbar}^{\vee} := \hbar^{-1}J_{\hbar} \subseteq F_{\hbar}[[G]]^{\vee}$. Therefore $\hbar^{-1}\varphi \in \hbar F_{\hbar}[[G]]^{\vee}$, hence $F := \exp(\hbar^{-1}\varphi)$ is indeed a well-defined element in $F_{\hbar}[[G]]^{\vee}$, q.e.d.

(b) We compute $\text{Ad}(F)(f)$, $f \in F_\hbar[[G]]$: using the identity $\text{Ad}(\exp(X))(Y) = \exp(\text{ad}(X))(Y)$ and expanding the exponential into a power series we get

$$\text{Ad}(F)(f) = \text{Ad}(\exp(\hbar^{-1}\varphi))(f) = \exp(\text{ad}(\hbar^{-1}\varphi))(f) = \sum_{n=0}^{+\infty} \frac{1}{n!} \text{ad}(\hbar^{-1}\varphi)^n(f)$$

Now Lemma 2.3.5(c) and the assumption $\varphi \in J_{F_\hbar}^2$ together guarantee that

$$\text{ad}(\hbar^{-1}\varphi)^n(f) = \text{ad}(\hbar^{-1}\varphi)^n(f_+) \in (1 - \delta_{s,0}) J_{F_\hbar}^{n+s} \quad \forall n \in \mathbb{N}_+$$

with $s \in \mathbb{N}$ such that $f \in J_{F_\hbar}^s$, hence $\text{Ad}(F)(f) = \sum_{n=0}^{+\infty} \frac{1}{n!} \text{ad}(\hbar^{-1}\varphi)^n(f)$ is indeed a well-defined element — a *convergent series!* — of $F_\hbar[[G]]$. \square

Lemma 3.4.2. *Let $F_\hbar[[G]]$ be a QFSHA, and $F_\hbar[[G]]^\vee$ the associated QUEA defined in §2.4. Let $\phi \in F_\hbar[[G]]^{\tilde{\otimes}^2}$ be such that $(\text{id} \otimes \epsilon)(\phi) = 0 = (\epsilon \otimes \text{id})(\phi)$. Then:*

- (a) $\mathcal{F} = \exp(\hbar^{-1}\phi)$ is a well-defined element in $(F_\hbar[[G]]^\vee)^{\hat{\otimes}^2}$;
- (b) $\mathcal{F} \cdot (x \otimes y) \cdot \mathcal{F}^{-1} \in F_\hbar[[G]]^{\tilde{\otimes}^2}$ for all $x, y \in F_\hbar[[G]]$, so that the adjoint action of \mathcal{F} onto $(F_\hbar[[G]]^\vee)^{\hat{\otimes}^2}$ actually restricts to $F_\hbar[[G]]^{\tilde{\otimes}^2}$;
- (c) $(\text{id} \otimes \epsilon)(\mathcal{F}) = 1 = (\epsilon \otimes \text{id})(\mathcal{F})$;
- (d) \mathcal{F} is orthogonal, i.e. $\mathcal{F}_{2,1} = \mathcal{F}^{-1}$, iff ϕ is antisymmetric, i.e. $\phi_{2,1} = -\phi$;

Proof. (a)–(b) The assumption $(\text{id} \otimes \epsilon)(\phi) = 0 = (\epsilon \otimes \text{id})(\phi)$ means that $\phi \in J_\hbar^{\tilde{\otimes}^2}$ where $J_\hbar := \text{Ker}(\epsilon_{F_\hbar[[G]])}$. Now note that $F_\hbar[[G \times G]] := F_\hbar[[G]]^{\tilde{\otimes}^2}$ is in turn a QFSHA, for the group $G \times G$; furthermore, its augmentation ideal is

$$\text{Ker}(\epsilon_{F_\hbar[[G \times G]])} = \text{Ker}(\epsilon_{F_\hbar[[G]])} \otimes F_\hbar[[G]] + F_\hbar[[G]] \otimes \text{Ker}(\epsilon_{F_\hbar[[G]])}$$

so that $J_\hbar^{\tilde{\otimes}^2} \subseteq \text{Ker}(\epsilon_{F_\hbar[[G \times G]])}^2$, and finally, we have also

$$F_\hbar[[G \times G]]^\vee = (F_\hbar[[G]] \tilde{\otimes} F_\hbar[[G]])^\vee = (F_\hbar[[G]]^\vee)^{\hat{\otimes}^2}$$

Therefore, applying Lemma 3.4.1 above to the QFSHA $F_\hbar[[G \times G]]$ and to $\varphi := \phi$ we get both claims (a) and (b).

(c)–(d) Both these claims follow at once from definitions, along with the assumption that $(\text{id} \otimes \epsilon)(\phi) = 0 = (\epsilon \otimes \text{id})(\phi)$. \square

The previous result leads us to introduce the notion of “quasi-twist”, as follows:

Definition 3.4.3. Let $F_\hbar[[G]]$ be a QFSHA, and $F_\hbar[[G]]^\vee$ as in 2.4.1(b). We call *quasi-twist (element) of $F_\hbar[[G]]$* any element in $(F_\hbar[[G]]^{\tilde{\otimes}^2})^\vee = (F_\hbar[[G]]^\vee)^{\hat{\otimes}^2}$ of the form $\mathcal{F} := \exp(\hbar^{-1}\phi)$ — for some $\phi \in F_\hbar[[G]]^{\tilde{\otimes}^2}$ such that $(\text{id} \otimes \epsilon)(\phi) = 0 = (\epsilon \otimes \text{id})(\phi)$ — which have the property of a twist element for the QUEA $F_\hbar[[G]]^\vee$.

Of course, every twist for $F_\hbar[[G]]$ is a quasi-twist too; the converse, instead, in general, is false (counterexamples do exist). However, *every quasi-twist still provides a well-defined deformation by twist of $F_\hbar[[G]]$* — in other words, the construction of *deformations by twist* properly extends to “*deformations by quasi-twist*” as well:

Theorem 3.4.4. *Let $F_{\hbar}[[G]]$ be a QFSHA, and $\mathcal{F} = \exp(\hbar^{-1}\phi)$ a quasi-twist for it, as in Definition 3.4.3. Then the procedure of twist deformation by \mathcal{F} applied to the QUEA $F_{\hbar}[[G]]^{\vee}$ restricts to $F_{\hbar}[[G]]$, making the latter into a new QFSHA.*

Proof. When deforming $F_{\hbar}[[G]]^{\vee}$ by the twist \mathcal{F} one introduces on $F_{\hbar}[[G]]^{\vee}$ the new coproduct $\Delta^{\mathcal{F}}$ given by $\Delta^{\mathcal{F}} := \text{Ad}(\mathcal{F}) \circ \Delta$. Then Lemma 3.4.2(b) ensures that $\Delta^{\mathcal{F}}$ restricts to $F_{\hbar}[[G]]$, in that it maps the latter into $F_{\hbar}[[G]]^{\tilde{\otimes}^2}$. The antipode can be dealt with similarly, whence we conclude that the (deformed) Hopf structure of $(F_{\hbar}[[G]]^{\vee})^{\mathcal{F}}$ does restrict to $F_{\hbar}[[G]]$, q.e.d. \square

Definition 3.4.5. With assumptions as in Theorem 3.4.4, the new QFSHA obtained from $F_{\hbar}[[G]]$ through the process of twist deformation (by \mathcal{F}) of $F_{\hbar}[[G]]^{\vee}$ followed by restriction will be called *the quasi-twist deformation of $F_{\hbar}[[G]]$ by \mathcal{F}* , and denoted by $F_{\hbar}[[G]]^{\mathcal{F}}$.

Finally, the next result describes in detail what exactly is the nature of the new, quasi-twist deformed QFSHA $F_{\hbar}[[G]]^{\mathcal{F}}$, shedding light onto its semiclassical limit:

Theorem 3.4.6. *Let $F_{\hbar}[[G]]$ be a QFSHA over the Lie bialgebra $\mathfrak{g} = (\mathfrak{g}; [\ , \], \delta)$. Set $\mathfrak{m} := \text{Ker}(\epsilon_{F[[G]])}$, so $\mathfrak{m}/\mathfrak{m}^2 \cong \mathfrak{g}^*$ and $(\mathfrak{m} \otimes \mathfrak{m}) / (\mathfrak{m}^2 \otimes \mathfrak{m} + \mathfrak{m} \otimes \mathfrak{m}^2) \cong \mathfrak{g}^* \otimes \mathfrak{g}^*$ as Lie bialgebras. Let \mathcal{F} be a quasi-twist for $F_{\hbar}[[G]]$, of the form $\mathcal{F} = \exp(\hbar^{-1}\phi)$ for some $\phi \in F_{\hbar}[[G]]^{\tilde{\otimes}^2}$, and set also $\phi_a := \phi - \phi_{2,1}$. Then:*

- (a) ϕ is antisymmetric, i.e. $\phi_{2,1} = -\phi$, iff \mathcal{F} is orthogonal, i.e. $\mathcal{F}_{2,1} = \mathcal{F}^{-1}$;
- (b) the element $c := \left(\phi_a \left(\text{mod } \hbar F_{\hbar}[[G]]^{\tilde{\otimes}^2} \right) \text{mod } (\mathfrak{m}^2 \otimes \mathfrak{m} + \mathfrak{m} \otimes \mathfrak{m}^2) \right)$ in $(\mathfrak{m} \otimes \mathfrak{m}) / (\mathfrak{m}^2 \otimes \mathfrak{m} + \mathfrak{m} \otimes \mathfrak{m}^2) \cong \mathfrak{g}^* \otimes \mathfrak{g}^*$ is an antisymmetric twist for \mathfrak{g}^* ;

(c) the quasi-twist deformation $(F_{\hbar}[[G]])^{\mathcal{F}}$ of $F_{\hbar}[[G]]$ is a QFSHA for the Poisson group G^c whose cotangent Lie bialgebra is $\text{Lie}(G^c)^* = (\mathfrak{g}^*)^c = (\mathfrak{g}^*; [\ , \]_*, \delta_*^c)$ that is the deformation of \mathfrak{g}^* by the twist c ; in short, $(F_{\hbar}[[G]])^{\mathcal{F}} \cong F_{\hbar}[[G^c]]$.

In particular, if \mathcal{F} is $\mathbb{k}[[\hbar]]$ -valued — i.e., it is an ordinary twist for the Hopf $\mathbb{k}[[\hbar]]$ -algebra $F_{\hbar}[[G]]$ — or equivalently $\phi \in \hbar F_{\hbar}[[G]]^{\tilde{\otimes}^2}$, then we have just $c = 0$ and $(F_{\hbar}[[G]])^c \cong F_{\hbar}[[G^c]] = F_{\hbar}[[G]]$.

Proof. (a) This is a special case of Lemma 3.4.2(d).

(b) We start from the twist identity $\mathcal{F}_{12}(\Delta \otimes \text{id})(\mathcal{F}) = \mathcal{F}_{23}(\text{id} \otimes \Delta)(\mathcal{F})$ that we re-write in the equivalent form

$$(\Delta \otimes \text{id})(\mathcal{F}) (\text{id} \otimes \Delta)(\mathcal{F})^{-1} = \mathcal{F}_{12}^{-1} \mathcal{F}_{23} \quad (3.47)$$

Replacing $\mathcal{F} = \exp(\hbar^{-1}\phi)$, we find

$$\begin{aligned} (\Delta \otimes \text{id})(\mathcal{F}) \cdot (\text{id} \otimes \Delta)(\mathcal{F})^{-1} &= \\ &= (\Delta \otimes \text{id})(\exp(\hbar^{-1}\phi)) \cdot (\text{id} \otimes \Delta)(\exp(\hbar^{-1}\phi))^{-1} = \\ &= \exp(\hbar^{-1}(\Delta \otimes \text{id})(\phi)) \cdot \exp(-\hbar^{-1}(\text{id} \otimes \Delta)(\phi)) \end{aligned}$$

Now we recall the *Baker-Campbell-Hausdorff's formula*, that is the formal identity

$$\exp(X) \cdot \exp(Y) = \exp(\mathcal{BCH}(X, Y)) \quad (3.48)$$

which allows to express the product of two exponential as a single exponential: in it, $\mathcal{BCH}(X, Y)$ is an explicit formal series given by

$$\begin{aligned} \mathcal{BCH}(X, Y) &:= \log(\exp(X) \exp(Y)) = \\ &= \sum_{n=1}^{+\infty} \frac{(-1)^{n-1}}{n} \sum_{\substack{r_i+s_i>0 \\ 1 \leq i \leq n}} \frac{[X^{\bullet r_1} Y^{\bullet s_1} X^{\bullet r_2} Y^{\bullet s_2} \dots X^{\bullet r_n} Y^{\bullet s_n}]}{(\sum_{i=1}^n (r_i + s_i)) \cdot \prod_{j=1}^n r_j! s_j!} \end{aligned} \quad (3.49)$$

where we use notation

$$\begin{aligned} [X^{\bullet r_1} Y^{\bullet s_1} \dots X^{\bullet r_n} Y^{\bullet s_n}] &:= \\ &= [\underbrace{X, [X, \dots [X, [Y, [Y, \dots [Y, \dots, [X, [X, \dots [X, [Y, [Y, \dots, Y]]]] \dots]] \dots]}_{r_1} \dots]_{s_1} \dots]_{r_n} \dots]_{s_n} \end{aligned}$$

with the silent assumption that the Lie monomial $[X^{\bullet r_1} Y^{\bullet s_1} \dots X^{\bullet r_n} Y^{\bullet s_n}]$ is just X , respectively Y , when $n = 1$ and $s_1 = 0$, respectively $r_1 = 0$, while it is zero whenever $s_n > 1$ or $s_n = 0$ and $r_n > 1$. In words, when $S := \sum_{i=1}^n (r_i + s_i) > 1$ the Lie monomial $[X^{\bullet r_1} Y^{\bullet s_1} \dots X^{\bullet r_n} Y^{\bullet s_n}]$ is the composition of several operators $\text{ad}(X)^{r_i}$ or $\text{ad}(Y)^{s_i}$ to Y — when $s_n = 1$ — or to X — when $s_n = 0$ and $r_n = 1$. Looking up to second order, (3.49) reads

$$\mathcal{BCH}(X, Y) := X + Y + \frac{1}{2} [X, Y] + \mathcal{O}_{\mathcal{L}}(3) \quad (3.50)$$

where $\mathcal{O}_{\mathcal{L}}(3)$ denotes a (formal) infinite linear combination of Lie monomials in X and Y of degree at least 3.

Setting now $X := \hbar^{-1}(\Delta \otimes \text{id})(\phi)$ and $Y := -\hbar^{-1}(\text{id} \otimes \Delta)(\phi)$, the above analysis yields, rewriting (3.48),

$$\begin{aligned} (\Delta \otimes \text{id})(\mathcal{F}) \cdot (\text{id} \otimes \Delta)(\mathcal{F})^{-1} &= \\ &= \exp\left(\mathcal{BCH}\left(\hbar^{-1}(\Delta \otimes \text{id})(\phi), -\hbar^{-1}(\text{id} \otimes \Delta)(\phi)\right)\right) \end{aligned} \quad (3.51)$$

where the BCH series has to be expanded as in (3.49). To this end, writing $\phi = \phi_1 \otimes \phi_2$ (a sum being tacitly intended) with $\phi_1, \phi_2 \in J_{\hbar}$ (by Lemma 3.1.1) we have

$$\begin{aligned} (\Delta \otimes \text{id})(\phi) &= \Delta(\phi_1) \otimes \phi_2 = \phi_1 \otimes 1 \otimes \phi_2 + 1 \otimes \phi_1 \otimes \phi_2 + ((\phi_1)_{(1)})^+ \otimes ((\phi_1)_{(2)})^+ \otimes \phi_2 = \\ &= \phi_{1,3} + \phi_{2,3} + ((\phi_1)_{(1)})^+ \otimes ((\phi_1)_{(2)})^+ \otimes \phi_2 \end{aligned}$$

where we expanded $\Delta(\phi_1)$ as in Lemma 2.3.5(d) and we used that $\epsilon(\phi_1) = 0$. Note that in the expansion of $(\Delta \otimes \text{id})(\phi)$ we have

$$(\phi_{1,3} + \phi_{2,3}) \in J_{\hbar}^{(\otimes 3|2)}, \quad ((\phi_1)_{(1)})^+ \otimes ((\phi_1)_{(2)})^+ \otimes \phi_2 \in J_{\hbar}^{\otimes 3} \quad (3.52)$$

where we introduced the notation $J_{\hbar}^{(\otimes 3|N)} := \sum_{\substack{a,b,c \geq 0 \\ a+b+c \geq N}} J_{\hbar}^a \tilde{\otimes} J_{\hbar}^b \tilde{\otimes} J_{\hbar}^c$ (for $N \in \mathbb{N}$).

A similar analysis for $(\text{id} \otimes \Delta)(\phi)$, just switching the roles of ϕ_1 and ϕ_2 , yields

$$\begin{aligned} (\text{id} \otimes \Delta)(\phi) &= \phi_1 \otimes \Delta(\phi_2) = \phi_1 \otimes \phi_2 \otimes 1 + \phi_1 \otimes 1 \otimes \phi_2 + \phi_1 \otimes ((\phi_2)_{(1)})^+ \otimes ((\phi_2)_{(2)})^+ = \\ &= \phi_{1,2} + \phi_{1,3} + \phi_1 \otimes ((\phi_2)_{(1)})^+ \otimes ((\phi_2)_{(2)})^+ \end{aligned}$$

with

$$(\phi_{1,2} + \phi_{1,3}) \in J_h^{(\otimes 3|2)}, \quad \phi_1 \otimes ((\phi_2)_{(1)})^+ \otimes ((\phi_2)_{(2)})^+ \in J_h^{\otimes 3} \quad (3.53)$$

Now, thanks to Lemma 2.3.5(c), from (3.52) and (3.53) we get

$$\begin{aligned} [\hbar X, \hbar Y] &= [(\Delta \otimes \text{id})(\phi), (\text{id} \otimes \Delta)(\phi)] = \\ &= [\phi_{1,3} + \phi_{2,3}, \phi_{1,2} + \phi_{1,3}] + \hbar \cdot \mathcal{O}\left(J_h^{(\otimes 3|4)}\right) = \\ &= [\phi_{1,3}, \phi_{1,2}] + [\phi_{2,3}, \phi_{1,2}] + [\phi_{2,3}, \phi_{1,3}] + \hbar \cdot \mathcal{O}\left(J_h^{[\otimes 3|4]}\right) \end{aligned}$$

hence

$$[X, Y] = \hbar^{-1} \left(\hbar^{-1} [\phi_{1,3}, \phi_{1,2}] + \hbar^{-1} [\phi_{2,3}, \phi_{1,2}] + \hbar^{-1} [\phi_{2,3}, \phi_{1,3}] + \mathcal{O}\left(J_h^{[\otimes 3|4]}\right) \right) \quad (3.54)$$

for some element $\mathcal{O}\left(J_h^{[\otimes 3|4]}\right) \in J_h^{[\otimes 3|4]}$, where hereafter we use notation

$$J_h^{[\otimes 3|N]} := \sum_{\substack{a,b,c \geq 1 \\ a+b+c \geq N}} J_h^{\otimes a} \otimes J_h^{\otimes b} \otimes J_h^{\otimes c} \subseteq \text{Ker}\left(\epsilon_{F_h[[G]]^{\otimes 3}}\right)^N \quad \forall N \in \mathbb{N}_+ \quad (3.55)$$

Pushing the analysis further on, we find easily that

$$[X^{\bullet r_1} Y^{\bullet s_1} \dots X^{\bullet r_n} Y^{\bullet s_n}] \in \hbar^{-1} J_h^{[\otimes 3|S+1]} \quad \forall S := \sum_{i=1}^n (r_i + s_i) > 1 \quad (3.56)$$

looking at (3.49), this tells us that the expansion of the BCH series occurring in (3.51), when expanded as in (3.49), is actually given by \hbar^{-1} multiplied by a *truly convergent series inside* $F_h[[G]]^{\otimes 3}$. In other words, tidying everything up — from (3.51), (3.49), (3.50), (3.54) and (3.56) altogether — we find that there exists some $\mathcal{Z} \in J_h^{\otimes 3} \subseteq F_h[[G]]^{\otimes 3}$ such that $(\Delta \otimes \text{id})(\mathcal{F}) \cdot (\text{id} \otimes \Delta)(\mathcal{F})^{-1} = \exp(\hbar^{-1} \mathcal{Z})$. Even more, by (3.49) and (3.50) and the previous analysis we do know the expansion of this \mathcal{Z} up to second order, whence we find

$$\begin{aligned} (\Delta \otimes \text{id})(\mathcal{F}) \cdot (\text{id} \otimes \Delta)(\mathcal{F})^{-1} &= \\ &= \exp\left(\hbar^{-1} \left((\Delta \otimes \text{id})(\phi) - (\text{id} \otimes \Delta)(\phi) - \right. \right. \\ &\quad \left. \left. - \hbar^{-1} 2^{-1} [\phi_{1,3}, \phi_{1,2}] - \hbar^{-1} 2^{-1} [\phi_{2,3}, \phi_{1,2}] - \hbar^{-1} 2^{-1} [\phi_{2,3}, \phi_{1,3}] + \mathcal{O}\left(J_h^{[\otimes 3|4]}\right) \right)\right) \end{aligned} \quad (3.57)$$

Now we go and work instead on the right-hand side of (3.47). Again, replacing $\mathcal{F} = \exp(\hbar^{-1} \phi)$, we find

$$\begin{aligned} \mathcal{F}_{12}^{-1} \cdot \mathcal{F}_{23} &= \exp(-\hbar^{-1} \phi \otimes 1) \cdot \exp(\hbar^{-1} 1 \otimes \phi) = \\ &= \exp(-\hbar^{-1} \phi_{1,2}) \cdot \exp(\hbar^{-1} \phi_{2,3}) = \exp\left(\mathcal{BC}\mathcal{H}(-\hbar^{-1} \phi_{1,2}, \hbar^{-1} \phi_{2,3})\right) \end{aligned}$$

Now for the computation of $\mathcal{BC}\mathcal{H}(-\hbar^{-1} \phi_{1,2}, \hbar^{-1} \phi_{2,3})$; to avoid possible confusion, we denote the second, right-hand instance of ϕ by ϕ' . We begin noting that

$$[\phi_{1,2}, \phi'_{2,3}] = [\phi_1 \otimes \phi_2 \otimes 1, 1 \otimes \phi'_1 \otimes \phi'_2] = \phi_1 \otimes [\phi_2, \phi'_1] \otimes \phi'_2 \in \hbar J_h^{[\otimes 3|3]}$$

so that for $X := -\hbar^{-1} \phi_{1,2}$ and $Y := \hbar^{-1} \phi_{2,3}$ we get, using Lemma 2.3.5(c),

$$[X, Y] = [-\hbar^{-1} \phi_{1,2}, \hbar^{-1} \phi'_{2,3}] = -\hbar^{-2} [\phi_{1,2}, \phi'_{2,3}] \in \hbar^{-2} \hbar J_h^{\otimes 3} = \hbar^{-1} J_h^{[\otimes 3|3]}$$

A second, similar step gives (with obvious notation ϕ , ϕ' and ϕ'')

$$\begin{aligned} [\phi_{1,2}, [\phi'_{1,2}, \phi''_{2,3}]] &= [\phi_1 \otimes \phi_2 \otimes 1, \phi'_1 \otimes [\phi'_2, \phi''_1] \otimes \phi''_2] = \\ &= [\phi_1, \phi'_1] \otimes \phi_2 \cdot [\phi'_2, \phi''_1] \otimes \phi''_2 + \phi_1 \cdot \phi'_1 \otimes [\phi_2, [\phi'_2, \phi''_1]] \otimes \phi''_2 \in \hbar^2 J_h^{[\otimes 3|4]} \end{aligned}$$

so that $[X, [X, Y]] \in \hbar^{-3} \hbar^2 J_h^{[\otimes 3|4]} = \hbar^{-1} J_h^{[\otimes 3|4]}$. More in general, iteration yields

$$[X^{\bullet r_1} Y^{\bullet s_1} \dots X^{\bullet r_n} Y^{\bullet s_n}] \in \hbar^{-S} \hbar^{S-1} J_h^{[\otimes 3|S+1]} = \hbar^{-1} J_h^{[\otimes 3|S+1]} \quad (3.58)$$

with notation as before, still using Lemma 2.3.5(c). Tidying everything up we find that there exists $\mathcal{W} \in J_h^{\otimes 3} \subseteq F_h[[G]]^{\otimes 3}$ such that $\mathcal{F}_{12}^{-1} \cdot \mathcal{F}_{23} = \exp(\hbar^{-1} \mathcal{W})$; moreover, by (3.49) and (3.50) along with the previous analysis we can write

$$\mathcal{F}_{1,2}^{-1} \cdot \mathcal{F}_{2,3} = \exp\left(\hbar^{-1} \left(-\phi_{1,2} + \phi_{2,3} - \hbar^{-1} 2^{-1} [\phi_{1,2}, \phi_{2,3}] + \mathcal{O}\left(J_h^{[\otimes 3|4]}\right)\right)\right) \quad (3.59)$$

Finally, comparing (3.57), (3.59) and (3.47) we get the identity in $F_h[[G]]^{\otimes 3}$

$$(\Delta \otimes \text{id})(\phi) - (\text{id} \otimes \Delta)(\phi) -$$

$$\begin{aligned} -\hbar^{-1} 2^{-1} [\phi_{1,3}, \phi_{1,2}] - \hbar^{-1} 2^{-1} [\phi_{2,3}, \phi_{1,2}] - \hbar^{-1} 2^{-1} [\phi_{2,3}, \phi_{1,3}] + \mathcal{O}\left(J_h^{[\otimes 3|4]}\right) = \\ = -\phi_{1,2} + \phi_{2,3} - \hbar^{-1} 2^{-1} [\phi_{1,2}, \phi_{2,3}] + \mathcal{O}\left(J_h^{[\otimes 3|4]}\right) \end{aligned}$$

that in turn, through simplification and reduction modulo $\hbar F_h[[G]]^{\otimes 3}$, yields the following identity inside $F[[G]]^{\otimes 3}$

$$\begin{aligned} (\Delta \otimes \text{id})(\bar{\phi}) - (\text{id} \otimes \Delta)(\bar{\phi}) + \bar{\phi}_{1,2} - \bar{\phi}_{2,3} + \\ + 2^{-1} \{ \bar{\phi}_{1,2}, \bar{\phi}_{1,3} \} + \{ \bar{\phi}_{1,2}, \bar{\phi}_{2,3} \} + 2^{-1} \{ \bar{\phi}_{1,3}, \bar{\phi}_{2,3} \} \equiv_{\mathfrak{m}^{[\otimes 3|4]}} 0 \end{aligned} \quad (3.60)$$

where hereafter we adopt the notation for which $\bar{\varphi}$ denotes the coset modulo \hbar of any element $\varphi \in F_h[[G]]^{\otimes 3}$ with $n \in \mathbb{N}_+$.

Now let $\mathbb{k}[\mathbb{S}_3]$ act onto $F_h[[G]]^{\otimes 3}$ and consider in particular the action of the antisymmetrizer $\text{Alt}_3 := (\text{id} - (12) - (23) - (31) + (123) + (321))$ onto the above identity: this in turn yields a new identity. Within the latter, we have a first contribution of the form

$$\begin{aligned} \text{Alt}_3 \cdot \left((\Delta \otimes \text{id})(\bar{\phi}) - (\text{id} \otimes \Delta)(\bar{\phi}) \right) = \\ = (\nabla \otimes \text{id})(\bar{\phi}) - (\nabla \otimes \text{id})(\bar{\phi}_{2,1}) + \text{c.p.} = (\nabla \otimes \text{id})(\bar{\phi}_a) + \text{c.p.} \end{aligned}$$

and a second contribution of the form

$$\text{Alt}_3 \cdot (\bar{\phi}_{1,2} - \bar{\phi}_{2,3}) = 0$$

The third and last contribution is

$$\text{Alt}_3 \cdot \left(2^{-1} \{ \bar{\phi}_{1,2}, \bar{\phi}_{1,3} \} \right) + \text{Alt}_3 \cdot \left(\{ \bar{\phi}_{1,2}, \bar{\phi}_{2,3} \} \right) + \text{Alt}_3 \cdot \left(2^{-1} \{ \bar{\phi}_{1,3}, \bar{\phi}_{2,3} \} \right)$$

We go and compute the first summand, as follows:

$$\begin{aligned} \text{Alt}_3 \cdot \left(2^{-1} \{ \bar{\phi}_{1,2}, \bar{\phi}_{1,3} \} \right) &= (\text{id} - (23)) \cdot \left(2^{-1} \{ \bar{\phi}_{1,2}, \bar{\phi}_{1,3} \} \right) + \text{c.p.} = \\ &= \left(2^{-1} \{ \bar{\phi}_{1,2}, \bar{\phi}_{1,3} \} - 2^{-1} \{ \bar{\phi}_{1,3}, \bar{\phi}_{1,2} \} \right) + \text{c.p.} = \{ \bar{\phi}_{1,2}, \bar{\phi}_{1,3} \} + \text{c.p.} \end{aligned}$$

A similar analysis applies to the third summand, which yields

$$\begin{aligned} Alt_3 \cdot \left(2^{-1} \{ \bar{\phi}_{1,3}, \bar{\phi}_{2,3} \} \right) &= (\text{id} - (12)) \cdot \left(2^{-1} \{ \bar{\phi}_{1,3}, \bar{\phi}_{2,3} \} \right) + \text{c.p.} = \\ &= \left(2^{-1} \{ \bar{\phi}_{1,3}, \bar{\phi}_{2,3} \} - 2^{-1} \{ \bar{\phi}_{2,3}, \bar{\phi}_{1,3} \} \right) + \text{c.p.} = \{ \bar{\phi}_{1,3}, \bar{\phi}_{2,3} \} + \text{c.p.} \end{aligned}$$

whereas for the second summand instead we get

$$\begin{aligned} Alt_3 \cdot \left(\{ \bar{\phi}_{1,2}, \bar{\phi}_{2,3} \} \right) &= (\text{id} - (13)) \cdot \left(\{ \bar{\phi}_{1,2}, \bar{\phi}_{2,3} \} \right) + \text{c.p.} = \\ &= \{ \bar{\phi}_{1,2}, \bar{\phi}_{2,3} \} - \{ \bar{\phi}_{3,2}, \bar{\phi}_{2,1} \} + \text{c.p.} \end{aligned}$$

Putting all these together we find

$$\begin{aligned} &Alt_3 \cdot \left(2^{-1} \{ \bar{\phi}_{1,2}, \bar{\phi}_{1,3} \} \right) + Alt_3 \cdot \left(\{ \bar{\phi}_{1,2}, \bar{\phi}_{2,3} \} \right) + Alt_3 \cdot \left(2^{-1} \{ \bar{\phi}_{1,3}, \bar{\phi}_{2,3} \} \right) = \\ &= \{ \bar{\phi}_{1,2}, \bar{\phi}_{1,3} \} + \{ \bar{\phi}_{1,2}, \bar{\phi}_{2,3} \} - \{ \bar{\phi}_{3,2}, \bar{\phi}_{2,1} \} + \{ \bar{\phi}_{1,3}, \bar{\phi}_{2,3} \} + \text{c.p.} = \{ \bar{\phi}_a, \bar{\phi}_a \} \end{aligned}$$

where the very last identity follows from a routine calculation. Joint with the previously found identities, the latter gives yet the following, last one, which is the result of letting Alt_3 act onto the congruence (3.60):

$$\left((\nabla \otimes \text{id})(\bar{\phi}_a) + \text{c.p.} \right) + \{ \bar{\phi}_a, \bar{\phi}_a \} \equiv_{\mathfrak{m}^{[4]}} 0$$

At last, recalling that $c := \bar{\phi}_a \pmod{\mathfrak{m}^2}$ in $\mathfrak{m}/\mathfrak{m}^2 = \mathfrak{g}^*$, and that in the latter Lie bialgebra the Lie cobracket, resp. the Lie bracket, is given by ∇ , resp. by $[\ , \]$, reduced modulo \mathfrak{m}^2 , the last formula above — in $\mathfrak{m}^{\otimes 3}$ — implies

$$\left((\delta \otimes \text{id})(c) + \text{c.p.} \right) + [[c, c]] = 0$$

within $(\mathfrak{g}^*)^{\otimes 3}$, which implies exactly that c — which is antisymmetric by construction — is an antisymmetric twist for the Lie bialgebra $\mathfrak{g}^* = \mathfrak{m}/\mathfrak{m}^2$, q.e.d.

(c) We adopt the following notational convention: any element in $F_{\hbar}[[G]]$ will be denoted by an italic letter, say $f \in F_{\hbar}[[G]]$; then its coset modulo $\hbar F_{\hbar}[[G]]$ will be denoted with a line over that letter, say $\bar{f} := (f \pmod{\hbar F_{\hbar}[[G]])$, and finally the coset of the latter modulo \mathfrak{m}^2 will be denoted by the corresponding letter in roman font, say $f := (\bar{f} \pmod{\mathfrak{m}^2})$. Note also that every element in $\mathfrak{g}^* = \mathfrak{m}/\mathfrak{m}^2$ can be written as such an $f = (\bar{f} \pmod{\mathfrak{m}^2})$ for some $f \in J_{\hbar} \in \text{Ker}(\epsilon_{F_{\hbar}[[G]])}$.

Similar notation will be used for elements in $F_{\hbar}[[G]]^{\otimes 2}$ and their coset modulo \hbar and (further on) modulo $\mathfrak{m}^{[\otimes 2|3]} := \mathfrak{m} \otimes \mathfrak{m}^2 + \mathfrak{m}^2 \otimes \mathfrak{m}$.

Recall that the Lie cobracket induced on $\mathfrak{m}/\mathfrak{m}^2 = \mathfrak{g}^*$ by the deformed quantization is defined by

$$\delta^{\mathcal{F}}(f) := \left(\Delta^{\mathcal{F}} - (\Delta^{\mathcal{F}})^{21} \right) (\bar{f}) \pmod{\mathfrak{m}^{[\otimes 2|3]}} = \overline{\Delta^{\mathcal{F}}(f)} - \overline{(\Delta^{\mathcal{F}})^{\text{op}}(f)} \pmod{\mathfrak{m}^{[\otimes 2|3]}}$$

so we start computing $\Delta^{\mathcal{F}}(f)$. Definitions give

$$\begin{aligned}\Delta^{\mathcal{F}}(f) &= \text{Ad}(\mathcal{F})(\Delta(f)) = \text{Ad}(\mathcal{F})(f_{(1)} \otimes f_{(2)}) = \\ &= \text{Ad}(\mathcal{F})\left(\left(f_{(1)} \otimes 1\right) \cdot \left(1 \otimes f_{(2)}\right)\right) = \text{Ad}(\mathcal{F})(f_{(1)} \otimes 1) \cdot \text{Ad}(\mathcal{F})(1 \otimes f_{(2)})\end{aligned}$$

In the last product, we focus on the first factor: thus we get

$$\begin{aligned}\text{Ad}(\mathcal{F})(f_{(1)} \otimes 1) &= \text{Ad}\left(\exp(\hbar^{-1}\phi)\right)(f_{(1)} \otimes 1) = \exp\left(\text{ad}(\hbar^{-1}\phi)\right)(f_{(1)} \otimes 1) = \\ &= \sum_{n=0}^{+\infty} \frac{1}{n!} \text{ad}(\hbar^{-1}\phi)^n(f_{(1)} \otimes 1) = \sum_{n=0}^{+\infty} \frac{1}{n!} \text{ad}(\hbar^{-1}\phi_1 \otimes \phi_2)^n(f_{(1)} \otimes 1) = \\ &= \sum_{n=0}^{+\infty} \frac{1}{n!} \text{ad}(\hbar^{-1}\phi_1)^n(f_{(1)}) \otimes \phi_2^n = f_{(1)} \otimes 1 + [\hbar^{-1}\phi_1, f_{(1)}] \otimes \phi_2 + \mathcal{O}(2)\end{aligned}$$

that is in short

$$\text{Ad}(\mathcal{F})(f_{(1)} \otimes 1) = f_{(1)} \otimes 1 + [\hbar^{-1}\phi_1, f_{(1)}] \otimes \phi_2 + \mathcal{O}(2)$$

where hereafter $\mathcal{O}(2)$ denotes any element in $J_{\hbar}^{[\otimes 2|3]}$. A similar calculation yields

$$\text{Ad}(\mathcal{F})(1 \otimes f_{(2)}) = 1 \otimes f_{(2)} + \phi_1 \otimes [\hbar^{-1}\phi_2, f_{(2)}] + \mathcal{O}(2)$$

Pasting together the last two identities we find

$$\begin{aligned}\Delta^{\mathcal{F}}(f) &= \text{Ad}(\mathcal{F})(f_{(1)} \otimes f_{(2)}) = \text{Ad}(\mathcal{F})(f_{(1)} \otimes 1) \cdot \text{Ad}(\mathcal{F})(1 \otimes f_{(2)}) = \\ &= \left(f_{(1)} \otimes 1 + [\hbar^{-1}\phi_1, f_{(1)}] \otimes \phi_2 + \mathcal{O}(2)\right) \cdot \left(1 \otimes f_{(2)} + \phi_1 \otimes [\hbar^{-1}\phi_2, f_{(2)}] + \mathcal{O}(2)\right) = \\ &= f_{(1)} \otimes f_{(2)} + f_{(1)} \phi_1 \otimes [\hbar^{-1}\phi_2, f_{(2)}] + [\hbar^{-1}\phi_1, f_{(1)}] \otimes \phi_2 f_{(2)} + \mathcal{O}(2) = \\ &= f_{(1)} \otimes f_{(2)} + \epsilon(f_{(1)}) \phi_1 \otimes [\hbar^{-1}\phi_2, f_{(2)}] + [\hbar^{-1}\phi_1, f_{(1)}] \otimes \phi_2 \epsilon(f_{(2)}) + \mathcal{O}(2) = \\ &= f_{(1)} \otimes f_{(2)} + \phi_1 \otimes [\hbar^{-1}\phi_2, f] + [\hbar^{-1}\phi_1, f] \otimes \phi_2 + \mathcal{O}(2) = \\ &= \Delta(f) + \phi_1 \otimes [\hbar^{-1}\phi_2, f] + [\hbar^{-1}\phi_1, f] \otimes \phi_2 + \mathcal{O}(2)\end{aligned}$$

so that, eventually, we get in short

$$\Delta^{\mathcal{F}}(f) \underset{J_{\hbar}^{[\otimes 2|3]}}{\equiv} \Delta(f) + \phi_1 \otimes [\hbar^{-1}\phi_2, f] + [\hbar^{-1}\phi_1, f] \otimes \phi_2$$

Therefore, for $\nabla_{\mathcal{F}} := \Delta^{\mathcal{F}} - (\Delta^{\mathcal{F}})^{21}$ we get

$$\begin{aligned}\nabla_{\mathcal{F}}(f) &\underset{J_{\hbar}^{[\otimes 2|3]}}{\equiv} \Delta(f) + \phi_1 \otimes [\hbar^{-1}\phi_2, f] + [\hbar^{-1}\phi_1, f] \otimes \phi_2 - \\ &\quad - \Delta^{\text{op}}(f) - [\hbar^{-1}\phi_2, f] \otimes \phi_1 - \phi_2 \otimes [\hbar^{-1}\phi_1, f] = \\ &= \nabla(f) + \phi_1^{(a)} \otimes \hbar^{-1}[\phi_2^{(a)}, f] + \hbar^{-1}[\phi_1^{(a)}, f] \otimes \phi_2^{(a)} = \\ &= \nabla(f) - \phi_1^{(a)} \otimes \hbar^{-1}[f, \phi_2^{(a)}] - \hbar^{-1}[f, \phi_1^{(a)}] \otimes \phi_2^{(a)}\end{aligned} \tag{3.61}$$

where we used notation $\phi_a := \phi - \phi_{21} = \phi_1^{(a)} \otimes \phi_2^{(a)}$. When we reduce the last identity in (3.61) modulo $\hbar J_{\hbar}^{\otimes 2}$ we end up with

$$\nabla_{\mathcal{F}}(\bar{f}) \underset{\mathfrak{m}^{[\otimes 2|3]}}{\equiv} \nabla(\bar{f}) - \overline{\phi_1^{(a)}} \otimes \left\{ \bar{f}, \overline{\phi_2^{(a)}} \right\} - \left\{ \bar{f}, \overline{\phi_1^{(a)}} \right\} \otimes \overline{\phi_2^{(a)}}$$

hence reducing the latter modulo $\mathfrak{m}^{[\otimes 2|3]}$ we find in $\mathfrak{m}^{\otimes 2}/\mathfrak{m}^{[\otimes 3]} = \mathfrak{g}^* \otimes \mathfrak{g}^*$ the identity

$$\begin{aligned} \left(\nabla_{\mathcal{F}} \bmod \mathfrak{m}^{[\otimes 2|3]} \right) (f) &= \delta(f) - c_1 \otimes [f, c_2] - [f, c_1] \otimes c_2 = \\ &= \delta(f) - (\text{ad}(f))(c) = (\delta - \partial_c)(f) = \delta^c(f) \end{aligned}$$

which means that the induced Lie cobracket on $\mathfrak{m}/\mathfrak{m}^2 = \mathfrak{g}^*$ is just δ^c , q.e.d. \square

Example 3.4.7. Let $G := GL_n(\mathbb{k})$ be the general linear group over \mathbb{k} , and $\mathfrak{g} := \mathfrak{gl}_n(\mathbb{k})$. We consider the QUEA $U_{\hbar}(\mathfrak{g}) = U_{\hbar}(\mathfrak{gl}_n(\mathbb{k}))$ and the QFSHA $F_{\hbar}[[G]] = F_{\hbar}[[GL_n(\mathbb{k})]]$ introduced in Example 3.2.3. Letting \mathfrak{b}^- and \mathfrak{b}^+ be the Borel Lie subalgebras in \mathfrak{g} of lower triangular and upper triangular matrices, respectively, the subalgebra $U_{\hbar}(\mathfrak{b}^-)$ of $U_{\hbar}(\mathfrak{g})$ generated by the F_i 's and the Γ_k 's is a QUEA for \mathfrak{b}^- , while the subalgebra $U_{\hbar}(\mathfrak{b}^+)$ generated by the E_i 's and the Γ_k 's is a QUEA for \mathfrak{b}^+ — both being also Hopf subalgebras of $U_{\hbar}(\mathfrak{g})$, indeed. Dually, the QFSHA $F_{\hbar}[[B^-]] = U_{\hbar}(\mathfrak{b}^-)^*$ identifies with the Hopf algebra quotient of $F_{\hbar}[[G]]$ obtained by modding out the ideal generated by the $x_{i,j}$'s with $i < j$; similarly, the QFSHA $F_{\hbar}[[B^+]] = U_{\hbar}(\mathfrak{b}^+)^*$ identifies with the Hopf algebra quotient of $F_{\hbar}[[G]]$ obtained by modding out the ideal generated by the $x_{i,j}$'s with $i > j$. Therefore, from the presentation of $F_{\hbar}[[G]]$ in Example 3.2.3 one deduces the following presentations for these quotient Hopf algebras: $F_{\hbar}[[B^-]]$ is generated by the entries of the “lower triangular q -matrix” $(x_{i,j}^-)_{i=1,\dots,n}^{j=1,\dots,n}$; with $x_{i,j}^- := x_{i,j}$ for all $i \geq j$ and $x_{i,j}^- := 0$ for all $i < j$, and similarly $F_{\hbar}[[B^+]]$ is generated by the entries of the “upper triangular q -matrix” $(x_{i,j}^+)_{i=1,\dots,n}^{j=1,\dots,n}$; where $x_{i,j}^+ := x_{i,j}$ when $i \leq j$ and $x_{i,j}^+ := 0$ for $i > j$.

Now we consider a new group G which is “double version” of $GL_n(\mathbb{k})$, in that it is a Manin double of B^- and B^+ ; its tangent Lie algebra \mathfrak{g} then is the Manin double of \mathfrak{b}^- and \mathfrak{b}^+ ; in particular, $G = B^- \times B^+$ as algebraic varieties (not as groups), with B^- and B^+ being embedded as subgroups, whereas $\mathfrak{g} = \mathfrak{b}^- \oplus \mathfrak{b}^+$ as vector spaces, with \mathfrak{b}^- and \mathfrak{b}^+ being embedded as Lie subalgebras (this case is explained in detail in [GaGa2] when G is a “double version” of a semisimple (connected) group: $G = SL_n(\mathbb{k})$ is such an example, and $GL_n(\mathbb{k})$ is just a very slight variation of that).

For these new G and \mathfrak{g} , a QUEA $U_{\hbar}(\mathfrak{g})$ is defined as follows: it is the unital, associative, \hbar -adically complete $\mathbb{k}[[\hbar]]$ -algebra with generators

$$F_1, F_2, \dots, F_{n-1}, \Gamma_1^-, \Gamma_2^-, \dots, \Gamma_{n-1}^-, \Gamma_n^-, \Gamma_1^+, \Gamma_2^+, \dots, \Gamma_{n-1}^+, \Gamma_n^+, E_1, E_2, \dots, E_{n-1}$$

and relations

$$\begin{aligned} [\Gamma_k^{\pm}, \Gamma_{\ell}^{\pm}] &= 0, & [\Gamma_k^{\pm}, F_j] &= -\delta_{k,j} F_j, & [\Gamma_k^{\pm}, E_j] &= +\delta_{k,j} E_j, & [\Gamma_k^{\pm}, \Gamma_{\ell}^{\mp}] &= 0 \\ [E_i, F_j] &= \delta_{i,j} \frac{e^{\hbar(\Gamma_i^+ - \Gamma_{i+1}^+)} - e^{\hbar(\Gamma_{i+1}^- - \Gamma_i^-)}}{e^{+\hbar} - e^{-\hbar}} \\ [E_i, E_j] &= 0, & [F_i, F_j] &= 0 & \forall i, j : |i - j| > 1 \\ E_i^2 E_j - (q + q^{-1}) E_i E_j E_i + E_j E_i^2 &= 0 & \forall i, j : |i - j| = 1 \\ F_i^2 F_j - (q + q^{-1}) F_i F_j F_i + F_j F_i^2 &= 0 & \forall i, j : |i - j| = 1. \end{aligned}$$

where $[X, Y] := XY - YX$ again. The Hopf structure then is given by

$$\begin{aligned} \Delta(F_i) &= F_i \otimes e^{\hbar(\Gamma_{i+1}^- - \Gamma_i^-)} + 1 \otimes F_i, & S(F_i) &= -F_i e^{\hbar(\Gamma_i^- - \Gamma_{i+1}^-)}, & \epsilon(F_i) &= 0 \\ \Delta(\Gamma_k^\pm) &= \Gamma_k^\pm \otimes 1 + 1 \otimes \Gamma_k^\pm, & S(\Gamma_k^\pm) &= -\Gamma_k^\pm, & \epsilon(\Gamma_k^\pm) &= 0 \\ \Delta(E_i) &= E_i \otimes 1 + e^{\hbar(\Gamma_i^+ - \Gamma_{i+1}^+)} \otimes E_i, & S(E_i) &= -e^{\hbar(\Gamma_{i+1}^+ - \Gamma_i^+)} E_i, & \epsilon(E_i) &= 0 \end{aligned}$$

In fact, this $U_\hbar(\mathfrak{g})$ can be realized as a *quantum double* of $U_\hbar(\mathfrak{b}^-)$ and $U_\hbar(\mathfrak{b}^+)$: in particular, then, we have a decomposition $U_\hbar(\mathfrak{g}) = U_\hbar(\mathfrak{b}^-) \widehat{\otimes} U_\hbar(\mathfrak{b}^+)$ as coalgebras. Dually, the latter implies that for the QFSHA $F_\hbar[[G]] := U_\hbar(\mathfrak{g})^*$ we have an identification as algebras

$$F_\hbar[[G]] = (U_\hbar(\mathfrak{b}^-) \widehat{\otimes} U_\hbar(\mathfrak{b}^+))^* = U_\hbar(\mathfrak{b}^-)^* \widetilde{\otimes} U_\hbar(\mathfrak{b}^+)^* = F_\hbar[[B^-]] \widetilde{\otimes} F_\hbar[[B^+]]$$

As a consequence, exploiting the presentations of $F_\hbar[[B^-]]$ and $F_\hbar[[B^+]]$ given above, we find in a nutshell a presentation for $F_\hbar[[G]]$ as being the algebra generated by the entries of the “ q -matrix in blocks”

$$\begin{pmatrix} X^+ & \mathbf{0}_{n \times n} \\ \mathbf{0}_{n \times n} & X^- \end{pmatrix} \quad \text{with} \quad X^\pm := (x_{i,j}^\pm)_{i=1,\dots,n}^{j=1,\dots,n}; \quad \text{as defined above (triangular).}$$

Moreover, explicit identifications $F_\hbar[[G]] = U_\hbar(\mathfrak{g})^*$ and $U_\hbar(\mathfrak{g}) = F_\hbar[[G]]^*$ can be encoded in the Hopf pairing $\langle \cdot, \cdot \rangle : F_\hbar[[G]] \times U_\hbar(\mathfrak{g}) \longrightarrow \mathbb{k}[[\hbar]]$ given on generators by the following values:

$$\begin{aligned} \langle x_{i,j}^-, \prod_{k=1}^n (\Gamma_k^+)^{g_k} \rangle &= 0 = \langle x_{i,j}^-, E_t \rangle, & \langle x_{i,j}^+, F_t \rangle &= 0 = \langle x_{i,j}^+, \prod_{k=1}^n (\Gamma_k^-)^{g_k} \rangle \\ \langle x_{i,j}^-, F_t \rangle &= \delta_{i,j+1} \delta_{t,j}, & \langle x_{i,j}^+, E_t \rangle &= \delta_{i+1,j} \delta_{i,t} \\ \langle x_{i,j}^+, \prod_{k=1}^n (\Gamma_k^+)^{g_k} \rangle &= \delta_{i,j} (1 - \delta_{g_i,0}) \prod_{k \neq i} \delta_{g_k,0} = \langle x_{i,j}^-, \prod_{k=1}^n (\Gamma_k^-)^{g_k} \rangle \end{aligned} \quad (3.62)$$

In particular, from the first line in (3.62) note that if $\underline{\Gamma}_1$ and $\underline{\Gamma}_2$ are two monomials in the Γ_k^\pm 's, then for all $i = 1, \dots, n$ we have

$$\langle x_{i,i}^\pm, \underline{\Gamma}_1 \cdot \underline{\Gamma}_2 \rangle = \langle x_{i,i}^\pm, \underline{\Gamma}_1 \rangle \cdot \langle x_{i,i}^\pm, \underline{\Gamma}_2 \rangle \quad (3.63)$$

Thanks to Proposition 4.1.2 later on, any quasi-twist for $F_\hbar[[G]]$ can be seen as a quasi-2-cocycle for $U_\hbar(\mathfrak{g}) = F_\hbar[[G]]^*$. Now, some examples of the latter were introduced in Example 3.3.8 above for a large class of QUEA, including that for $\mathfrak{g} = \mathfrak{sl}_n(\mathbb{k})$. The same procedure can be applied to the present case, which is a slight variation of that case applied to $\mathfrak{gl}_n(\mathbb{k})$ instead of $\mathfrak{sl}_n(\mathbb{k})$, as follows.

Let \mathfrak{h} be the $\mathbb{k}[[\hbar]]$ -span of $B_\Gamma := \{ \Gamma_k^+, \Gamma_k^- \mid k = 1, \dots, n \}$. Then fix an antisymmetric, $\mathbb{k}[[\hbar]]$ -bilinear map $\chi : \mathfrak{h} \times \mathfrak{h} \longrightarrow \mathbb{k}[[\hbar]]$ whose matrix of values on pairs of elements in the $\mathbb{k}[[\hbar]]$ -basis B_Γ is $X = \left(\chi_{k,t}^{\varepsilon,\eta} = \chi(\Gamma_k^\varepsilon, \Gamma_t^\eta) \right)_{k,t=1,\dots,n}^{\varepsilon,\eta \in \{+,-\}} \in \mathfrak{so}_{2n}(\mathbb{k}[[\hbar]])$. Any such map χ also induces uniquely an antisymmetric, $\mathbb{k}[[\hbar]]$ -bilinear map $\tilde{\chi}_U$ on $U_\hbar(\mathfrak{h}) = \widehat{S}_{\mathbb{k}[[\hbar]]}(\mathfrak{h}) := \bigoplus_{n \in \mathbb{N}} \widehat{S}_{\mathbb{k}[[\hbar]]}^n(\mathfrak{h})$ with values in $\mathbb{k}[[\hbar]]$, by setting

$$\begin{aligned} \tilde{\chi}_U(z, 1) &:= \epsilon(z) =: \tilde{\chi}_U(1, z) & \forall z &\in \widehat{S}_{\mathbb{k}[[\hbar]]}(\mathfrak{h}) \\ \tilde{\chi}_U(x, y) &:= \chi(x, y) & \forall x, y &\in S_{\mathbb{k}[[\hbar]]}^1(\mathfrak{h}) \\ \tilde{\chi}_U(x, y) &:= 0 & \forall x \in S_{\mathbb{k}}^r(\mathfrak{h}), y \in S_{\mathbb{k}}^s(\mathfrak{h}) : r, s \geq 1, r + s > 2 \end{aligned} \quad (3.64)$$

Then we can define a $\mathbb{k}[[\hbar]]$ -linear map $\chi_U := e^{\hbar^{-1}2^{-1}\tilde{\chi}_U} = \sum_{m \geq 0} \hbar^{-m} \tilde{\chi}_U^{*m} / 2^m m!$ from $U_{\hbar}(\mathfrak{h}) \hat{\otimes}_{\mathbb{k}[[\hbar]]} U_{\hbar}(\mathfrak{h})$ to $\mathbb{k}((\hbar))$, which — like in [GaGa2, Lemma 5.2.2] — happens to be a well-defined *quasi-2-cocycle* for $U_{\hbar}(\mathfrak{h})$.

Assume now that χ satisfies the additional constraint $\chi(S_i, -) = 0 = \chi(-, S_i)$ for all $i \in I := \{1, \dots, n-1\}$, where $S_i := 2^{-1}(\Gamma_i^+ - \Gamma_{i+1}^+ + \Gamma_i^- - \Gamma_{i+1}^-)$ for all $i \in I$; note that requiring $\chi(S_i, -) = 0$ for all $i \in I$ is enough (by antisymmetry), and the latter in turn is equivalent to requiring $\chi_{i,t}^{+,\eta} - \chi_{i+1,t}^{+,\eta} + \chi_{i,t}^{-,\eta} - \chi_{i+1,t}^{-,\eta} = 0$ for all $i \in I$, all $t = 1, \dots, n-1$ and all $\eta \in \{+, -\}$. Then χ induces a unique $\mathbb{k}[[\hbar]]$ -bilinear map $\bar{\chi} : \bar{\mathfrak{h}} \times \bar{\mathfrak{h}} \longrightarrow \mathbb{k}[[\hbar]]$, where $\bar{\mathfrak{h}} := \mathfrak{h}/\mathfrak{s}$ with $\mathfrak{s} := \text{Span}_{\mathbb{k}[[\hbar]]}(\{S_i\}_{i=1, \dots, n-1})$, given by $\bar{\chi}(H' + \mathfrak{s}, H'' + \mathfrak{s}) := \chi(H', H'')$ for all $H', H'' \in \mathfrak{h}$.

Now repeat the above construction with $\bar{\mathfrak{h}}$ and $\bar{\chi}$ replacing \mathfrak{h} and χ : this yields a quasi-2-cocycle $\bar{\chi}_U$ for $U_{\hbar}(\bar{\mathfrak{h}})$. But now the additional assumption on χ implies that there exists a *Hopf algebra epimorphism* $\pi : U_{\hbar}(\mathfrak{g}) \longrightarrow \widehat{S}_{\mathbb{k}[[\hbar]]}(\bar{\mathfrak{h}}) \cong U_{\hbar}(\bar{\mathfrak{h}})$ given by $\pi(E_i) := 0$, $\pi(F_i) := 0$ — for $i = 1, \dots, n-1$ — and $\pi(T) := (T + \mathfrak{s}) \in \bar{\mathfrak{h}} \subseteq U_{\hbar}(\bar{\mathfrak{h}})$ — for $T \in \mathfrak{h}$. Finally, we set $\sigma_{\chi} := \bar{\chi}_U \circ (\pi \times \pi)$, which is a well-defined quasi-2-cocycle for $U_{\hbar}(\mathfrak{g})$, again in the sense of Definition 3.3.3. Note that

$$\sigma_{\chi} := \bar{\chi}_U \circ (\pi \times \pi) = \exp(\hbar^{-1}2^{-1}\tilde{\bar{\chi}}_U) \circ (\pi \times \pi) = \exp(\hbar^{-1}2^{-1}\tilde{\bar{\chi}}_U \circ (\pi \times \pi))$$

Now let us re-think the quasi-2-cocycle σ_{χ} for $U_{\hbar}(\mathfrak{g})$ as a quasi-twist for $F_{\hbar}[[G]]$. First of all, comparing (3.63) and (3.64) we deduce that the form $\tilde{\bar{\chi}}_U \circ (\pi \times \pi)$ in $(U_{\hbar}(\mathfrak{g})^{\hat{\otimes} 2})^*$ identifies with $\Phi_{\chi} := \sum_{k,t=1}^n \sum_{\varepsilon, \eta \in \{+, -\}} \chi_{k,t}^{\varepsilon, \eta} y_{k,k}^{\varepsilon} \otimes y_{t,t}^{\eta}$ in $F_{\hbar}[[G]]^{\hat{\otimes} 2}$, where $y_{\ell, \ell}^{\zeta} := \log(x_{\ell, \ell}^{\zeta})$ is a well-defined element in $F_{\hbar}[[G]]$.

Then, exponentiating yields

$$\mathcal{F}_{\chi} := \sigma_{\chi} = \exp(\hbar^{-1}2^{-1}\tilde{\bar{\chi}}_U \circ (\pi \times \pi)) = \exp(\hbar^{-1}2^{-1}\Phi_{\chi})$$

which is exactly the quasi-twist of $F_{\hbar}[[G]]$ we were looking for.

We can also check directly that this \mathcal{F}_{χ} is indeed a quasi-twist by an explicit computation. We see this in the simplest case, when $n = 2$; the other cases are quite similar, but require quite another shot of calculations.

We need to compute the coproduct of the $x_{t,t}^{\varepsilon}$'s in $F_{\hbar}[[G]]$, which is defined (by construction) by the condition $\langle \Delta(x_{t,t}^{\varepsilon}), A \otimes Z \rangle = \langle x_{t,t}^{\varepsilon}, A \cdot Z \rangle$ for all $A, Z \in U_{\hbar}(\mathfrak{g})$; since $U_{\hbar}(\mathfrak{g})$ admits the PBW-type basis

$$\mathcal{B} := \left\{ F^f (\Gamma_1^-)^{g_1^-} (\Gamma_2^-)^{g_2^-} (\Gamma_1^+)^{g_1^+} (\Gamma_2^+)^{g_2^+} E^e \mid f, g_1^-, g_2^-, g_1^+, g_2^+, e \in \mathbb{N} \right\}$$

we can replace A and Z with any two PBW monomials from \mathcal{B} . Now, let us say that a PBW monomial of the form $\mathcal{M} = F^f (\Gamma_1^-)^{g_1^-} (\Gamma_2^-)^{g_2^-} (\Gamma_1^+)^{g_1^+} (\Gamma_2^+)^{g_2^+} E^e$ belongs to the root space $(e - f)\alpha$. Then root/weight considerations easily show that $\langle x_{t,t}^{\varepsilon}, \mathcal{M} \rangle \neq 0$ only for PBW monomials \mathcal{M} in the root space 0, i.e. such that $e = f$. A straightforward computation gives

$$E^e \cdot F^f = \sum_{s=0}^{e \wedge f} ([s]_q!)^2 \begin{bmatrix} e \\ s \end{bmatrix}_q \begin{bmatrix} f \\ s \end{bmatrix}_q F^{f-s} K_{e,f}(s) E^{e-s}$$

where $q := \exp(\hbar)$, $[r]_q := \frac{q^r - q^{-r}}{q - q^{-1}}$, $[m]_q! := \prod_{r=1}^m [r]_q$, $\begin{bmatrix} \ell \\ s \end{bmatrix}_q := \frac{[\ell]_q!}{[s]_q! [\ell - s]_q!}$ and

$$K_{e,f}(s) := \prod_{r=1}^s \frac{q^{2s-e-f+1-r} K_+^{+1} - q^{r-1-2s+e+f} K_-^{-1}}{q^r - q^{-r}}$$

with $K_+^{+1} := 1 \otimes \exp(+\hbar(\Gamma_i^+ - \Gamma_{i+1}^+))^{\pm 1}$ and $K_-^{-1} := \exp(-\hbar(\Gamma_i^- - \Gamma_{i+1}^-))^{\pm 1} \otimes 1$. Then the product of two PBW monomials expands into

$$\begin{aligned} \mathcal{M}' \cdot \mathcal{M}'' &= \\ &= F^{f'}(\Gamma_1^-)^{\dot{g}_1^-}(\Gamma_2^-)^{\dot{g}_2^-}(\Gamma_1^+)^{\dot{g}_1^+}(\Gamma_2^+)^{\dot{g}_2^+} E^{e'} \cdot F^{f''}(\Gamma_1^-)^{\ddot{g}_1^-}(\Gamma_2^-)^{\ddot{g}_2^-}(\Gamma_1^+)^{\ddot{g}_1^+}(\Gamma_2^+)^{\ddot{g}_2^+} E^{e''} = \\ &= \sum_{s=0}^{e' \wedge f''} ([s]_q!)^2 \begin{bmatrix} e \\ s \end{bmatrix}_q \begin{bmatrix} f \\ s \end{bmatrix}_q F^{f'} \underline{\Gamma}^{\dot{g}} F^{f''-s} K_{e',f''}(s) E^{e'-s} \underline{\Gamma}^{\ddot{g}} E^{e''} = \\ &= \sum_{s=0}^{e' \wedge f''} q^{\mathcal{E}(\dot{g}, f'', s, e', \ddot{g})} ([s]_q!)^2 \begin{bmatrix} e' \\ s \end{bmatrix}_q \begin{bmatrix} f'' \\ s \end{bmatrix}_q F^{f'+f''-s} \underline{\Gamma}^{\dot{g}} K_{e',f''}(s) \underline{\Gamma}^{\ddot{g}} E^{e'-s+e''} \end{aligned}$$

where $\underline{\Gamma}^{\dot{g}} := (\Gamma_1^-)^{\dot{g}_1^-}(\Gamma_2^-)^{\dot{g}_2^-}(\Gamma_1^+)^{\dot{g}_1^+}(\Gamma_2^+)^{\dot{g}_2^+}$, $\underline{\Gamma}^{\ddot{g}} := (\Gamma_1^-)^{\ddot{g}_1^-}(\Gamma_2^-)^{\ddot{g}_2^-}(\Gamma_1^+)^{\ddot{g}_1^+}(\Gamma_2^+)^{\ddot{g}_2^+}$ while $\mathcal{E}(\dot{g}, f'', s, e', \ddot{g}) \in \mathbb{Z}$ is a suitable exponent. When we expand the last expression w.r.t. the PBW basis \mathcal{B} , the part given by a linear combination of PBW monomials in the root space $\mathfrak{0}$ is

$$(\mathcal{M}' \cdot \mathcal{M}'')_{\mathfrak{0}} = \delta_{f',0} \delta_{f'',e'} \delta_{e'',0} q^{\mathcal{E}(\dot{g}, e', e', e', \ddot{g})} ([e']_q!)^2 \underline{\Gamma}^{\dot{g}} K_{e',e'}(e') \underline{\Gamma}^{\ddot{g}}$$

Eventually, tidying everything up we find

$$\begin{aligned} \langle \Delta(x_{t,t}^\varepsilon), \mathcal{M}' \otimes \mathcal{M}'' \rangle &= \langle x_{t,t}^\varepsilon, \mathcal{M}' \cdot \mathcal{M}'' \rangle = \langle x_{t,t}^\varepsilon, (\mathcal{M}' \cdot \mathcal{M}'')_{\mathfrak{0}} \rangle = \\ &= \delta_{f',0} \delta_{f'',e'} \delta_{e'',0} q^{\mathcal{E}(\dot{g}, e', e', e', \ddot{g})} ([e']_q!)^2 \langle x_{t,t}^\varepsilon, \underline{\Gamma}^{\dot{g}} K_{e',e'}(e') \underline{\Gamma}^{\ddot{g}} \rangle \end{aligned}$$

Now, a similar analysis yields (notation being obvious, hopefully)

$$\langle x_{t,t}^\varepsilon, \underline{\Gamma}^{\dot{g}} K_{e',e'}(e') \underline{\Gamma}^{\ddot{g}} \rangle = \langle x_{t,t}^\varepsilon, \underline{\Gamma} \rangle^{\dot{g}} \cdot \langle x_{t,t}^\varepsilon, K_{e',e'}(e') \rangle \cdot \langle x_{t,t}^\varepsilon, \underline{\Gamma} \rangle^{\ddot{g}}$$

and finally direct computation gives $\langle x_{t,t}^\varepsilon, K_{e',e'}(e') \rangle = \delta_{e',0}$, which is a direct consequence of the assumption $\chi(S_i, -) = 0 = \chi(-, S_i)$ for $i = 1, \dots, n-1$. Therefore, we end up with

$$\langle \Delta(x_{t,t}^\varepsilon), \mathcal{M}' \otimes \mathcal{M}'' \rangle \neq 0$$

only for monomials with $(f', e') = (0, 0) = (f'', e'')$, and for them we have

$$\begin{aligned} \langle \Delta(x_{t,t}^\varepsilon), \mathcal{M}' \otimes \mathcal{M}'' \rangle &= \langle \Delta(x_{t,t}^\varepsilon), \underline{\Gamma}^{\dot{g}} \otimes \underline{\Gamma}^{\ddot{g}} \rangle = \\ &= \langle x_{t,t}^\varepsilon, \underline{\Gamma}^{\dot{g}} \cdot \underline{\Gamma}^{\ddot{g}} \rangle = \langle x_{t,t}^\varepsilon, \underline{\Gamma}^{\dot{g}} \rangle \cdot \langle x_{t,t}^\varepsilon, \underline{\Gamma}^{\ddot{g}} \rangle = \langle x_{t,t}^\varepsilon, \mathcal{M}' \rangle \cdot \langle x_{t,t}^\varepsilon, \mathcal{M}'' \rangle \end{aligned}$$

which in short means $\Delta(x_{t,t}^\varepsilon) = x_{t,t}^\varepsilon \otimes x_{t,t}^\varepsilon$, i.e. $x_{t,t}^\varepsilon$ is group-like. Therefore, for $y_{t,t}^\varepsilon := \log(x_{t,t}^\varepsilon)$ instead we have $\Delta(y_{t,t}^\varepsilon) = y_{t,t}^\varepsilon \otimes 1 + 1 \otimes y_{t,t}^\varepsilon$, i.e. $y_{t,t}^\varepsilon$ is primitive.

Eventually, as all the $y_{t,t}^\varepsilon$'s are primitive, a trivial computation shows that \mathcal{F}_χ does obey condition (2.7), hence it is indeed a quasi-twist, as claimed.

3.5. Duality issues.

The procedures of deformation by twist or by 2-cocycle, both for Lie bialgebras and for Hopf algebras, are dual to each other, in the sense of Proposition 2.1.7 and Proposition 2.2.7. Because of this, we are lead to compare these two procedures *before* and *after* specialization, as follows.

Proposition 3.5.1. *Let $U_{\hbar}(\mathfrak{g})$ and $F_{\hbar}[[G]]$ be a QUEA and a QFSHA which are dual to each other, that is $F_{\hbar}[[G]] = U_{\hbar}(\mathfrak{g})^*$ and $U_{\hbar}(\mathfrak{g}) = F_{\hbar}[[G]]^*$. Then let \mathcal{F} be a twist for $U_{\hbar}(\mathfrak{g})$, and σ be a 2-cocycle for $F_{\hbar}[[G]]$. Assume that both \mathcal{F} and σ are trivial modulo \hbar , so that there exists a corresponding twist $c_{\mathcal{F}}$ for \mathfrak{g} (induced by \mathcal{F} via Theorem 3.1.2) and a corresponding 2-cocycle ζ_{σ} for \mathfrak{g}^* (induced by σ via Theorem 3.2.1). Finally, we identify twists for $U_{\hbar}(\mathfrak{g})$ and 2-cocycles for $F_{\hbar}[[G]]$ via Proposition 2.2.7, and similarly twists for \mathfrak{g} and 2-cocycles for \mathfrak{g}^* via Proposition 2.1.7.*

Then the following holds: if $\mathcal{F} = \sigma$, then $c_{\mathcal{F}} = \zeta_{\sigma}$. \square

The *proof* of the above claim is trivial — just track the whole construction of both $c_{\mathcal{F}}$ and ζ_{σ} , and compare the outcomes.

A similar duality result holds for deformations by quasi-2-cocycles and deformations by quasi-twists. Indeed, let us first notice that, by the very definitions, if $U_{\hbar}(\mathfrak{g})$ and $F_{\hbar}[[G]]$ are a QUEA and a QFSHA in duality — i.e., $F_{\hbar}[[G]] = U_{\hbar}(\mathfrak{g})^*$ and $U_{\hbar}(\mathfrak{g}) = F_{\hbar}[[G]]^*$ — then *any quasi-2-cocycle for $U_{\hbar}(\mathfrak{g})$ is automatically a quasi-twist for $F_{\hbar}[[G]]$, and viceversa* — see Proposition 4.1.2 later on.

Once this is settled, next result (which mirrors Proposition 3.5.1 above) holds too, whose proof again follows by direct comparison of the two deformation procedures (just tracking the whole construction of γ_{σ} and $c_{\mathcal{F}}$, and comparing the outcomes):

Proposition 3.5.2. *Let $U_{\hbar}(\mathfrak{g})$ and $F_{\hbar}[[G]]$ be a QUEA and a QFSHA which are dual to each other, that is $F_{\hbar}[[G]] = U_{\hbar}(\mathfrak{g})^*$ and $U_{\hbar}(\mathfrak{g}) = F_{\hbar}[[G]]^*$. Then let σ be a quasi-2-cocycle for $U_{\hbar}(\mathfrak{g})$, and \mathcal{F} be a quasi-twist for $F_{\hbar}[[G]]$. Let γ_{σ} be the 2-cocycle for \mathfrak{g} induced by σ via Theorem 3.3.7, and let $c_{\mathcal{F}}$ be the twist for \mathfrak{g}^* induced by \mathcal{F} via Theorem 3.4.6. Finally, we identify quasi-2-cocycles for $U_{\hbar}(\mathfrak{g})$ and quasi-twists for $F_{\hbar}[[G]]$ as mentioned above, and similarly we identify twists for \mathfrak{g} and 2-cocycles for \mathfrak{g}^* via Proposition 2.1.7. Then the following holds: if $\sigma = \mathcal{F}$, then $\gamma_{\sigma} = c_{\mathcal{F}}$. \square*

4. DEFORMATIONS VS. QDP

When we interchange (contravariantly) QUEA's and QFSHA's via linear duality, the interaction of such a process with deformation processes (either by twist or by 2-cocycle) is clear. By Proposition 2.2.7, any twist, resp. 2-cocycle, for a given quantum algebra (a QUEA or QFSHA, say) is automatically a 2-cocycle, resp. a twist, for its dual (a QFSHA or a QUEA, respectively), and the corresponding deformation processes “commute” with dualization.

On the other hand, in this section we investigate how the deformation procedures interact when we interchange (covariantly) QUEA's and QFSHA's via Drinfeld's functors, as in Theorem 2.4.2. In other words, we analyze how deformation procedures behave with respect to the Quantum Duality Principle.

4.1. Some auxiliary results.

We begin with a key observation, which shows a crucial fact: our “quasi-2-cocycles” for any QUEA and “quasi-twists” for any QFSHA are actually *standard* 2-cocycles and twists, respectively, for the QFSHA and for the QUEA that are associated with the original quantum group via Drinfeld’s functors from the QDP.

Here is the precise result:

Lemma 4.1.1.

(a) Let $U_{\hbar}(\mathfrak{g})$ be a QUEA, and let $U_{\hbar}(\mathfrak{g})'$ be its associated QFSHA following Theorem 2.4.2. Let $\sigma := \exp_*(\hbar^{-1}\chi)$ be a quasi-2-cocycle for $U_{\hbar}(\mathfrak{g})$ as in Definition 3.3.3. Then the restriction $\sigma|_{U_{\hbar}(\mathfrak{g})' \times U_{\hbar}(\mathfrak{g})'}$ of σ to $U_{\hbar}(\mathfrak{g})' \times U_{\hbar}(\mathfrak{g})'$ is a well-defined, $\mathbb{k}[[\hbar]]$ -valued bilinear form on $U_{\hbar}(\mathfrak{g})'$, of the form $\sigma' = \exp_*(\hbar^{-1}\chi')$ with $\chi' := (\hbar^{-2}\chi)|_{U_{\hbar}(\mathfrak{g})' \times U_{\hbar}(\mathfrak{g})'}$, and this $\sigma' := \exp_*(\hbar^{-1}\chi')$ is a 2-cocycle for $U_{\hbar}(\mathfrak{g})'$.

(b) Let $F_{\hbar}[[G]]$ be a QFSHA, and let $F_{\hbar}[[G]]^{\vee}$ be its associated QUEA following Theorem 2.4.2. Let $\mathcal{F} := \exp(\hbar^{-1}\phi)$ be a quasi-twist for $F_{\hbar}[[G]]$ as in Definition 3.4.3. Then $\mathcal{F} := \exp(\hbar^{-1}\phi) = \exp(\hbar^{-1}\phi^{\vee})$ with $\phi^{\vee} := \hbar^{-2}\phi \in (F_{\hbar}[[G]]^{\vee})^{\widehat{\otimes}^2}$, and, in these terms, $\mathcal{F}^{\vee} := \exp(\hbar^{-1}\phi^{\vee})$ is a twist for $F_{\hbar}[[G]]^{\vee}$.

Proof. (a) We retain notation from the proof of Lemma 3.3.2, and we proceed along the same lines. Thus we set $U_{\hbar} := U_{\hbar}(\mathfrak{g})$ and $J_{\hbar} := \text{Ker}(U_{\hbar})$, and we write

$$\hat{z} := \epsilon(z), \quad z^+ := z - \epsilon(z) = z - \hat{z} \in J_{\hbar}, \quad \text{hence} \quad z = z^+ + \hat{z} \quad \forall z \in U_{\hbar}$$

We already saw that $\chi(u, v) = \chi(u^+, v^+)$ for all $u, v \in U_{\hbar}$, and then for any $a, b \in U_{\hbar}$ we have

$$\sigma(a, b) = \sum_{n \geq 0} \hbar^{-n} \prod_{i=1}^n \chi(a_{(i)}^+, b_{(i)}^+) / n! \quad (4.1)$$

where $\otimes_{i=1}^n a_{(i)}^+ = \delta_n(a)$ and $\otimes_{i=1}^n b_{(i)}^+ = \delta_n(b)$.

Now, restricting to U_{\hbar}' we get that $a', b' \in U_{\hbar}'$ yields $\delta_n(a'), \delta_n(b') \in \hbar^n U_{\hbar}'^{\widehat{\otimes}^n}$; also, in the sequel we can clearly assume $\epsilon(a') = 0 = \epsilon(b')$. Then we get

$$\prod_{i=1}^n \chi'(a_{(i)}^+, b_{(i)}^+) = \prod_{i=1}^n \hbar^{-2} \chi(a_{(i)}^+, b_{(i)}^+) \in \hbar^{-2n} \hbar^{2n} \mathbb{k}[[\hbar]] = \mathbb{k}[[\hbar]] \quad \forall n \in \mathbb{N}_+$$

whence, just like in (4.1), we eventually get

$$\sigma'(a', b') = \sum_{n \geq 0} \hbar^{+n} \prod_{i=1}^n \chi'(a_{(i)}^+, b_{(i)}^+) / n! \in \mathbb{k}[[\hbar]] \quad \forall a', b' \in U_{\hbar}'$$

which proves the claim.

(b) This follows directly from Definition 3.4.3. \square

As a direct consequence, we have the following significant result:

Proposition 4.1.2. *Let U_{\hbar} be a QUEA and F_{\hbar} be a QFSHA that are dual to each other, i.e. such that $F_{\hbar} = (U_{\hbar})^*$ and $U_{\hbar} = (F_{\hbar})^*$. Then:*

- (a) σ is a quasi-2-cocycle for $U_{\hbar} \iff \sigma$ is a quasi-twist for F_{\hbar} ;
- (b) \mathcal{F} is a quasi-twist for $F_{\hbar} \iff \mathcal{F}$ is a quasi-2-cocycle for U_{\hbar} .

Proof. The proof follows directly from the very definitions of quasi-2-cocycle and quasi-twist, along with the observation that $F_{\hbar} = (U_{\hbar})^*$ and $U_{\hbar} = (F_{\hbar})^*$ imply $F_{\hbar}^{\vee} = (U_{\hbar}')^*$ and $U_{\hbar}' = (F_{\hbar}^{\vee})^*$, by (2.18). \square

4.2. Drinfeld’s functors and “quasi-deformations”.

In this subsection we analyze the interaction between the process of “quasi-deformation” and the action of a Drinfeld’s functor on some quantum group.

We begin with deformations by quasi-twist of a QFSHA and the application to the latter of Drinfeld’s functor $(\)^{\vee}$; then we shall look at deformation by quasi-2-cocycle of a QUEA and the application to the latter of Drinfeld’s functor $(\)'$.

4.2.1. Deformations by quasi-twist under $F_{\hbar}[[G]] \mapsto F_{\hbar}[[G]]^{\vee}$. We look now what happens with deformations by quasi-twist for a QFSHA when the latter is acted upon by the functor $(\)^{\vee}$ which associates with it a QUEA. Here is our result:

Theorem 4.2.2.

Let $F_{\hbar}[[G]]$ be a QFSHA. Let $\mathcal{F} = \exp(\hbar^{-1}\phi)$ be a quasi-twist for $F_{\hbar}[[G]]$, with $\phi \in F_{\hbar}[[G]]^{\otimes 2}$ (cf. Definition 3.4.3). Set $\phi^{\vee} := \hbar^{-1} \log(\mathcal{F}) = \hbar^{-2}\phi$, $\phi_a := \phi - \phi_{2,1}$ and $\phi_a^{\vee} := \phi^{\vee} - \phi_{2,1}^{\vee}$. Then we have:

(a) ϕ is antisymmetric, i.e. $\phi_{2,1} = -\phi$, iff \mathcal{F} is orthogonal, i.e. $\mathcal{F}_{2,1} = \mathcal{F}^{-1}$, iff ϕ^{\vee} is antisymmetric, i.e. $\phi_{2,1}^{\vee} = -\phi^{\vee}$;

(b) $\mathcal{F} = \exp(\hbar\phi^{\vee})$ is a twist element for the QUEA $U_{\hbar}(\mathfrak{g}^*) := F_{\hbar}[[G]]^{\vee}$.

(c) Let c be the antisymmetric twist of \mathfrak{g}^* corresponding to \mathcal{F} as provided by Theorem 3.4.6, and let c^{\vee} be the similar twist provided by Theorem 3.1.2 along with claim (b) above. Then $c = c^{\vee}$.

Proof. (a) This follows directly by construction.

(b) This is granted by Lemma 4.1.1(b).

(c) This follows by a careful — yet entirely straightforward — check, just tracking both constructions involved (of c and of c^{\vee} alike). \square

4.2.3. Deformations by quasi-2-cocycle under $U_{\hbar}(\mathfrak{g}) \mapsto U_{\hbar}(\mathfrak{g})'$. Given a QUEA, we can apply on it Drinfeld’s functor $(\)'$; we now see what happens when a deformation by quasi-2-cocycle is performed. Our result reads as follows:

Theorem 4.2.4.

Let $U_{\hbar}(\mathfrak{g})$ be a QUEA. Let $\sigma = \exp_*(\hbar^{-1}\chi)$ be a quasi-2-cocycle for $U_{\hbar}(\mathfrak{g})$, with $\chi \in (U_{\hbar}(\mathfrak{g})^{\widehat{\otimes} 2})^*$ (cf. Definition 3.3.3). Set $\chi' := \hbar^{-1} \log_*(\sigma) = \hbar^{-2}\chi$, $\chi_a := \chi - \chi_{2,1}$ and $\chi'_a := \chi' - \chi'_{2,1}$. Then we have:

(a) χ is antisymmetric, i.e. $\chi_{2,1} = -\chi$, iff σ is orthogonal, i.e. $\sigma_{2,1} = \sigma^{-1}$, iff χ' is antisymmetric, i.e. $\chi'_{2,1} = -\chi'$;

(b) $\sigma = \exp_*(\hbar\chi')$ is a 2-cocycle for the QFSHA $F_{\hbar}[[G^*]] := U_{\hbar}(\mathfrak{g})'$.

(c) Let γ be the antisymmetric 2-cocycle of \mathfrak{g} corresponding to σ as provided by Theorem 3.3.5, and let γ' be the similar 2-cocycle provided by Theorem 3.2.1 along with claim (b) above — using the identification $\mathfrak{g}^{**} = \mathfrak{g}$. Then $\gamma = \gamma'$.

Proof. (a) This follows directly by construction.

(b) This is true because of Lemma 4.1.1(a).

(c) Much like for Theorem 4.2.2(c), this follows again by a straightforward check, just carefully tracking both constructions involved (of γ and of γ' alike). \square

Remark 4.2.5. Recall that the notions of twist and that of 2-cocycle are dual to each other (cf. Proposition 2.2.7), and the same holds for those of quasi-twist and quasi-2-cocycle (cf. §3.5). Moreover, Drinfeld’s functors are also dual to each other, in the sense of (2.18). Taking all this into account, it turns out easily that Theorem 4.2.2 and Theorem 4.2.4 above are also “dual to each other”, in that either one of these two statements can be deduced from the other by a duality argument.

4.3. Drinfeld’s functors and (standard) deformations.

We analyze now the interaction between the process of deformation — in the standard sense — and the action of a Drinfeld’s functor on some quantum group.

We begin with deformations by twist of a QUEA and the application to the latter of Drinfeld’s functor $(\)'$; then we shall look at deformation by 2-cocycle of any QUEA, to which we apply Drinfeld’s functor $(\)'$. In both cases (obviously related by duality) we have to adopt a slightly stronger assumption, namely that the given twist, resp. 2-cocycle, is additionally an R -matrix, resp. a ϱ -comatrix.

4.3.1. Deformations by twist under $U_{\hbar}(\mathfrak{g}) \mapsto U_{\hbar}(\mathfrak{g})'$. As a first step, we look what happens for deformations by twist of a QUEA when the latter is acted upon by the functor $(\)'$ which associates with it a QFSHA. It turns out that we find a relevant result when we make the stronger assumption that the given twist is in fact a (quantum) R -matrix twist, as in Definition 2.2.1(d). Here is our result:

Theorem 4.3.2.

Let $U_{\hbar}(\mathfrak{g})$ be a QUEA, and let \mathcal{F} be a twist for $U_{\hbar}(\mathfrak{g})$ s.t. $\phi \equiv 1 \pmod{\hbar U_{\hbar}(\mathfrak{g})^{\widehat{\otimes} 2}}$; then $\phi := \hbar^{-1} \log(\mathcal{F}) \in U_{\hbar}(\mathfrak{g})^{\widehat{\otimes} 2}$, and $\mathcal{F} = \exp(\hbar \phi)$. Set also $\phi' := \hbar^{-2} \log(\mathcal{F}) = \hbar^{-2} \phi$, $\phi_a := \phi - \phi_{2,1}$ and $\phi'_a := \phi' - \phi'_{2,1}$. Assume in addition that \mathcal{F} is indeed a (quantum) R -matrix twist, as in Definition 2.2.1(d). Then we have:

(a) ϕ is antisymmetric, i.e. $\phi_{2,1} = -\phi$, iff \mathcal{F} is orthogonal, i.e. $\mathcal{F}_{2,1} = \mathcal{F}^{-1}$, iff ϕ' is antisymmetric, i.e. $\phi'_{2,1} = -\phi'$;

(b) $\mathcal{F} = \exp(\hbar^{-1} \phi')$ is a quasi-twist for the QFSHA $F_{\hbar}[[G^*]] := U_{\hbar}(\mathfrak{g})'$.

(c) Let c be the antisymmetric twist of \mathfrak{g} corresponding to \mathcal{F} as provided by Theorem 3.1.2, and let c' be the similar twist provided by Theorem 3.4.6 along with claim (b) above — using the identification $\mathfrak{g}^{**} = \mathfrak{g}$. Then $c = c'$.

Proof. (a) This follows directly by construction.

(b) This is proved in [EH, Theorem 0.1]. Note that the overall assumption there is that \mathcal{R} be an R -matrix, in the standard sense — so that $U_{\hbar}(\mathfrak{g})$ is quasitriangular.

Nevertheless, all the arguments used there to prove the main result only apply the defining properties of an “ R -matrix” in the sense of Definition 2.2.1, namely (2.8) and the right-hand side of (2.7); the assumption (2.10), instead, is never used. Therefore, the same arguments, and the whole proof, used in [EH] to prove Theorem 0.1 actually do prove also the present statement, that is actually stronger.

(c) Here again, the proof follows from a straightforward, careful checking procedure, keeping track of both constructions involved (of c and of c' alike), much like for Theorem 4.2.2(c) and for Theorem 4.2.4(c). \square

4.3.3. Deformations by 2-cocycle under $F_{\hbar}[[G]] \mapsto F_{\hbar}[[G]]^{\vee}$. As a second step, we look what happens to deformations of a QFSHA by 2-cocycle when we apply Drinfeld’s functor $(\)^{\vee}$. Here again, we get a relevant result under the stronger assumption that the given 2-cocycle is in fact a (quantum) ϱ -comatrix 2-cocycle, as in Definition 2.2.4(d). Our result reads as follows:

Theorem 4.3.4.

Let $F_{\hbar}[[G]]$ be any QFSHA, and let σ be a 2-cocycle for $F_{\hbar}[[G]]$ such that $\sigma \equiv 1 \pmod{\hbar (F_{\hbar}[[G]]^{\otimes 2})^{\star}}$; then $\varsigma := \hbar^{-1} \log(\sigma) \in (F_{\hbar}[[G]]^{\otimes 2})^{\star}$, and $\sigma = \exp(\hbar \varsigma)$. Set also $\varsigma^{\vee} := \hbar \log(\sigma) = \hbar^2 \varsigma$, $\varsigma_a := \varsigma - \varsigma_{2,1}$ and $\varsigma_a^{\vee} := \varsigma^{\vee} - \varsigma_{2,1}^{\vee}$. Assume in addition that σ is a (quantum) ϱ -comatrix 2-cocycle, as in Definition 2.2.4(d).

Then the following holds true:

(a) ς is antisymmetric, i.e. $\varsigma_{2,1} = -\varsigma$, iff σ is orthogonal, i.e. $\sigma_{2,1} = \sigma^{-1}$, iff ς^{\vee} is antisymmetric, i.e. $\varsigma_{2,1}^{\vee} = -\varsigma^{\vee}$;

(b) $\sigma = \exp(\hbar^{-1} \varsigma^{\vee})$ is a quasi-2-cocycle for the QUEA $U_{\hbar}(\mathfrak{g}^*) := F_{\hbar}[[G]]^{\vee}$;

(c) Let γ be the antisymmetric 2-cocycle of \mathfrak{g}^* corresponding to σ as provided by Theorem 3.2.1, and let γ^{\vee} be the similar 2-cocycle provided by 3.3.5 along with claim (b) above. Then $\gamma = \gamma^{\vee}$.

Proof. (a) This is obvious, by standard identities for formal exponentials.

(b) This claim is the dual to Theorem 4.3.2(b), so it follows from that one via a duality argument — involving the results in §4.1, in particular Proposition 4.1.2.

(c) Once more, as in previous cases, the claim follows from direct checking, keeping track of the two involved 2-cocycles — γ and of γ^{\vee} — were constructed. \square

Remark 4.3.5. Much like as we did in Remark 4.2.5, we notice here as well that — by the same reasons as before — Theorem 4.3.2 and Theorem 4.3.4 above are once more “dual to each other”, in that either one of these two statements can be deduced from the other by a duality argument.

5. MORPHISMS IN THE “(CO)QUASITRIANGULAR” CASE

In this section we focus onto R -matrices and ϱ -comatrices.

First, we investigate what happens with R -matrices and ϱ -comatrices with respect to the Quantum Duality Principle, namely when Drinfeld’s functors are applied. This leads us to introduce the weaker notions of “quasi- R -matrix” and “quasi- ϱ -comatrix” which eventually will prove quite significant.

Second, we consider the standard constructions providing morphisms between a Hopf algebra H and its dual when an R -matrix or a ϱ -comatrix is available, looking at how this applies to QUEAs and to QFSHAs. It turns out that the construction then improves, in a way that involves Drinfeld’s functors, again; moreover, we prove that it can also be extended to the case when R -matrices or ρ -comatrices are replaced with their “quasi-” counterparts.

5.1. R -matrices and ϱ -comatrices w.r.t. QDP: quasi-(co)matrices.

In next two results, we explain how R -matrices and ρ -comatrices “behave well” with respect to Drinfeld’s functors and the Quantum Duality Principle. In fact, this leads us to introduce the notions of “quasi- R -matrix” and of “quasi- ρ -comatrix”, which are straight analogue of the notions of “quasi-twist” and of “quasi-2-cocycle”.

We begin introducing some more bare definitions:

Definition 5.1.1.

(a) Let $F_{\hbar}[[G]]$ be a QFSHA. We call “quasi- R -matrix” of $F_{\hbar}[[G]]$ any R -matrix \mathcal{R} for $U_{\hbar}(\mathfrak{g}^*) := F_{\hbar}[[G]]^{\vee}$ such that $\mathcal{R} \equiv 1^{\otimes 2} \pmod{\left(\hbar F_{\hbar}[[G]]^{\vee} \widehat{\otimes} F_{\hbar}[[G]]^{\vee}\right)}$.

(b) Let $U_{\hbar}(\mathfrak{g})$ be a QUEA. We call “quasi- ρ -comatrix” of $U_{\hbar}(\mathfrak{g})$ any ϱ -comatrix ρ for $F_{\hbar}[[G^*]] := U_{\hbar}(\mathfrak{g})'$ such that $\rho \equiv \epsilon^{\otimes 2} \pmod{\left(\hbar (U_{\hbar}(\mathfrak{g})' \widetilde{\otimes} U_{\hbar}(\mathfrak{g})')^{\star}\right)}$.

Remark 5.1.2. In the same spirit of Proposition 2.2.7 and (2.18), it is clear that the two notions of “quasi- R -matrix” and of “quasi- ϱ -comatrix” are dual to each other. Namely, let $U_{\hbar}(\mathfrak{g})$ and $F_{\hbar}[[G]]$ be a QUEA and a QFSHA which are dual to each other, i.e. $F_{\hbar}[[G]] = U_{\hbar}(\mathfrak{g})^{\star}$ and $U_{\hbar}(\mathfrak{g}) = F_{\hbar}[[G]]^{\star}$. Then, through the identification $F_{\hbar}[[G]] \widetilde{\otimes} F_{\hbar}[[G]] = (U_{\hbar}(\mathfrak{g}) \widehat{\otimes} U_{\hbar}(\mathfrak{g}))^{\star}$, we have that $\mathcal{R} = \rho$ is a quasi- R -matrix of $F_{\hbar}[[G]]$ if and only if it is a quasi- ϱ -comatrix of $U_{\hbar}(\mathfrak{g})$.

Observations 5.1.3.

(a) With assumptions as in Definition 5.1.1(a) above, let \mathcal{R} be any quasi- R -matrix for a QFSHA $F_{\hbar}[[G]]$: since $\mathcal{R} \equiv 1^{\otimes 2} \pmod{\hbar}$, we can write \mathcal{R} in the form $\mathcal{R} = \exp(\hbar^{\pm 1} \theta)$ for some $\theta \in F_{\hbar}[[G]]^{\vee} \widehat{\otimes} F_{\hbar}[[G]]^{\vee}$.

Similarly, if ρ is any quasi- ϱ -comatrix for a QUEA $U_{\hbar}(\mathfrak{g})$, then we can write it in the form $\rho = \exp_{\star}(\hbar^{\pm 1} \varsigma)$ for some $\varsigma \in (U_{\hbar}(\mathfrak{g})' \widetilde{\otimes} U_{\hbar}(\mathfrak{g})')^{\star}$.

(b) Note that in the very definition of “quasi- R -matrix”, resp. of “quasi- ϱ -comatrix”, we assume a condition which is quite close, yet weaker, than the one demanded for the definition of “quasi-twist”, resp. of “quasi-2-cocycle”, in Definition 3.3.3, resp. in Definition 3.4.3. In fact, our choice for these definitions about R -matrices and ϱ -comatrices is motivated by Proposition 5.1.4 below, which eventually implies that *when the two setups overlap, the stronger condition for twists/2-cocycles actually holds true* — cf. Theorem 4.3.2 and Theorem 4.3.4.

The key result about quasi- R -matrices and quasi- ϱ -comatrices is the following:

Proposition 5.1.4.

(a) Let $U_{\hbar}(\mathfrak{g})$ be a QUEA, and let $U_{\hbar}(\mathfrak{g})'$ be the QFSHA associated to it by the Quantum Duality Principle, as in Theorem 2.4.2. Then for any R -matrix of $U_{\hbar}(\mathfrak{g})$

of the form $\mathcal{R} = \exp(\hbar\theta)$, with $\theta = \hbar^{-1} \log(\mathcal{R}) \in U_{\hbar}(\mathfrak{g})^{\widehat{\otimes}^2}$, we have

$$\vartheta := \hbar^2 \theta = \hbar^{-1} \log(\mathcal{R}) \in (U_{\hbar}(\mathfrak{g})')^{\widetilde{\otimes}^2}$$

(b) Let $F_{\hbar}[[G]]$ be a QFSHA, and let $F_{\hbar}[[G]]^{\vee}$ be the QUEA associated to it by the Quantum Duality Principle, as in Theorem 2.4.2. Then for any ϱ -comatrix of $F_{\hbar}[[G]]$ of the form $\rho = \exp_*(\hbar\varsigma)$, with $\varsigma = \hbar^{-1} \log_*(\rho) \in (F_{\hbar}[[G]]^{\widehat{\otimes}^2})^*$, we have

$$\zeta := \hbar^2 \varsigma = \hbar^{-1} \log_*(\rho) \in \left((F_{\hbar}[[G]]^{\vee})^{\widehat{\otimes}^2} \right)^*$$

Proof. (a) This is proved in [EH, Theorem 0.1]. Indeed, the overall assumption there is that \mathcal{R} be an R -matrix, in the standard sense — so that $U_{\hbar}(\mathfrak{g})$ is quasitriangular. Nevertheless, all the arguments used there to prove the main result only apply the defining properties of an “ R -matrix” in the sense of Definition 2.2.1, namely (2.8) and the right-hand side of (2.7); the assumption (2.10), instead, is never used. Therefore, the same arguments, and the whole proof, used in [EH] to prove Theorem 0.1 actually do prove also the present, stronger statement.

(b) This follows from claim (a), by duality, using the duality relation (2.18), the fact that $F_{\hbar}[[G]]^{\vee}$ is a QUEA when $F_{\hbar}[[G]]$ is a QFSHA, and Proposition 2.2.7. \square

The previous result has the following, important consequence:

Theorem 5.1.5. *Let $U_{\hbar}(\mathfrak{g})$ be a QUEA and $F_{\hbar}[[G]]$ be a QFSHA; let $F_{\hbar}[[G^*]] := U_{\hbar}(\mathfrak{g})'$ be the QFSHA and $U_{\hbar}(\mathfrak{g}^*) := F_{\hbar}[[G]]^{\vee}$ be the QUEA provided by the Quantum Duality Principle, as in Theorem 2.4.2. Then the following holds:*

(a) every R -matrix for $U_{\hbar}(\mathfrak{g})$ which is congruent to $1^{\otimes 2}$ modulo \hbar is a quasi- R -matrix for $U_{\hbar}(\mathfrak{g})'$;

(b) every ϱ -comatrix for $F_{\hbar}[[G]]$ which is congruent to $\epsilon^{\otimes 2}$ modulo \hbar is a quasi- ϱ -comatrix for $F_{\hbar}[[G]]^{\vee}$;

(c) every ϱ -comatrix for $U_{\hbar}(\mathfrak{g})'$ which is congruent to $\epsilon^{\otimes 2}$ modulo \hbar is a quasi- ϱ -comatrix for $U_{\hbar}(\mathfrak{g})$ itself;

(d) every R -matrix for $F_{\hbar}[[G]]^{\vee}$ which is congruent to $1^{\otimes 2}$ modulo \hbar is a quasi- R -matrix for $F_{\hbar}[[G]]$ itself.

Proof. Claims (a) and (b) follow directly from Proposition 5.1.4, claims (a) and (b), respectively. Now claim (c) follows from claim (b) applied to the QFSHA $F_{\hbar}[[G^*]] := U_{\hbar}(\mathfrak{g})'$, since $F_{\hbar}[[G^*]]^{\vee} = (U_{\hbar}(\mathfrak{g})')^{\vee} = U_{\hbar}(\mathfrak{g})$ by Theorem 2.4.2(a). Similarly, claim (d) follows from (a) applied to the QUEA $U_{\hbar}(\mathfrak{g}^*) := F_{\hbar}[[G]]^{\vee}$, since $U_{\hbar}(\mathfrak{g}^*)' = (F_{\hbar}[[G]]^{\vee})' = F_{\hbar}[[G]]$ by Theorem 2.4.2(a) again. \square

5.2. Morphisms from R -matrices and ϱ -comatrices.

We shall now explore what happens if we apply the general constructions leading to Proposition 2.2.9, resp. to Proposition 2.2.10, is (tentatively) applied to a QUEA, resp. a QFSHA, as the Hopf algebra H to start with, and a suitable R -matrix, resp. ϱ -comatrix, for it. To begin with, we check that Proposition 2.2.9 still makes sense when $H := U_{\hbar}(\mathfrak{g})$ is a QUEA:

Proposition 5.2.1. *Let $U_{\hbar}(\mathfrak{g})$ be a QUEA, and $F_{\hbar}[[G]] := U_{\hbar}(\mathfrak{g})^*$ be its dual QFSHA, as in §2.3.4. Let $\mathcal{R} = \mathcal{R}_1 \otimes \mathcal{R}_2$ (in Sweedler's notation) be an R -matrix for $U_{\hbar}(\mathfrak{g})$. Then there exist two morphisms of topological Hopf algebras*

$$\begin{aligned} \overleftarrow{\Phi}_{\mathcal{R}} : F_{\hbar}[[G]] := U_{\hbar}(\mathfrak{g})^* &\longrightarrow U_{\hbar}(\mathfrak{g})^{\text{cop}}, & \eta &\mapsto \overleftarrow{\Phi}_{\mathcal{R}}(\eta) := \eta(\mathcal{R}_1) \mathcal{R}_2 \\ \overrightarrow{\Phi}_{\mathcal{R}} : F_{\hbar}[[G]] := U_{\hbar}(\mathfrak{g})^* &\longrightarrow U_{\hbar}(\mathfrak{g})^{\text{op}}, & \eta &\mapsto \overrightarrow{\Phi}_{\mathcal{R}}(\eta) := \mathcal{R}_1 \eta(\mathcal{R}_2) \end{aligned}$$

Proof. This is straightforward. Let us write the given R -matrix as an \hbar -adically convergent series $\mathcal{R} = \sum_{n \geq 0} \hbar^n \mathcal{R}'_n \otimes \mathcal{R}''_n$ (where each $\mathcal{R}'_n \otimes \mathcal{R}''_n$ is written in Sweedler's notation). Then $\overleftarrow{\Phi}_{\mathcal{R}}(\eta) := \eta(\mathcal{R}_1) \mathcal{R}_2 = \sum_{n \geq 0} \hbar^n \eta(\mathcal{R}'_n) \otimes \mathcal{R}''_n$ for every $\eta \in U_{\hbar}(\mathfrak{g})^*$, where the last term is a well-defined, \hbar -adically convergent series in $U_{\hbar}(\mathfrak{g})$, so that $\overleftarrow{\Phi}_{\mathcal{R}}(\eta) \in U_{\hbar}(\mathfrak{g})$ and the map $\overleftarrow{\Phi}_{\mathcal{R}}$ is well-defined. The properties making this map $\overleftarrow{\Phi}_{\mathcal{R}}$ into a (topological) Hopf morphism then are proved via a formal check, just like for Proposition 2.2.9. The proof for $\overrightarrow{\Phi}_{\mathcal{R}}$ is pretty similar. \square

Dually, Proposition 2.2.10 still makes sense when $H := F_{\hbar}[[G]]$ is a QFSHA:

Proposition 5.2.2. *Let $F_{\hbar}[[G]]$ be a QFSHA, and $U_{\hbar}(\mathfrak{g}) := F_{\hbar}[[G]]^*$ be its dual QUEA, as in §2.3.4. Let ρ be a ϱ -comatrix for $F_{\hbar}[[G]]$. Then there exist two morphisms of topological Hopf algebras*

$$\begin{aligned} \overleftarrow{\Psi}_{\rho} : F_{\hbar}[[G]] &\longrightarrow (F_{\hbar}[[G]]^*)^{\text{cop}} = U_{\hbar}(\mathfrak{g})^{\text{cop}}, & \ell &\mapsto \overleftarrow{\Psi}_{\rho}(\ell) := \rho(\ell, -) \\ \overrightarrow{\Psi}_{\rho} : F_{\hbar}[[G]] &\longrightarrow (F_{\hbar}[[G]]^*)^{\text{op}} = U_{\hbar}(\mathfrak{g})^{\text{op}}, & \ell &\mapsto \overrightarrow{\Psi}_{\rho}(\ell) := \rho(-, \ell) \end{aligned}$$

Proof. Again, this is straightforward, the main point being to show that the given maps are indeed well-defined. We consider $\overleftarrow{\Psi}_{\rho}$, the case of $\overrightarrow{\Psi}_{\rho}$ is similar.

By definition, $\rho \in (F_{\hbar}[[G]] \otimes F_{\hbar}[[G]])^*$, the latter being the space of all continuous $\mathbb{k}[[\hbar]]$ -linear maps from $F_{\hbar}[[G]] \otimes F_{\hbar}[[G]]$ to $\mathbb{k}[[\hbar]]$. Then for any $\ell \in F_{\hbar}[[G]]$ the “left-hand evaluation” of ρ in ℓ yields clearly a continuous $\mathbb{k}[[\hbar]]$ -linear map $\overleftarrow{\Psi}_{\rho} := \rho(\ell, -) : F_{\hbar}[[G]] \rightarrow \mathbb{k}[[\hbar]]$, $f \mapsto \rho(\ell, f)$; thus $\overleftarrow{\Psi}_{\rho}$ is well defined, q.e.d.

The Hopf properties of this map $\overleftarrow{\Psi}_{\rho}$ are then proved by direct check. \square

Note that $U_{\hbar}(\mathfrak{g})^{\text{cop}}$ is obviously a QUEA for the Lie bialgebra $\mathfrak{g}^{\text{cop}}$ (the “co-opposite” to \mathfrak{g} , i.e. one reverses the Lie cobracket), and similarly $U_{\hbar}(\mathfrak{g})^{\text{op}}$ is a QUEA for the Lie bialgebra \mathfrak{g}^{op} (the “opposite” to \mathfrak{g} , using reversed Lie bracket). Then both previous results provide morphisms between quantum groups of *different* nature, namely a QFSHA as domain and a QUEA as codomain.

Nevertheless, we shall now show that both previous results can be refined, eventually yielding morphisms that connect quantum groups of the *same* nature, namely both QFSHA's in one case and both QUEA's in the other case.

Theorem 5.2.3. *Let $U_{\hbar}(\mathfrak{g})$ be a QUEA, let $F_{\hbar}[[G]] := U_{\hbar}(\mathfrak{g})^*$ be its dual QFSHA, as in §2.3.4, and let $F_{\hbar}[[G^*]] := U_{\hbar}(\mathfrak{g})'$, resp. $U_{\hbar}(\mathfrak{g}^*) := F_{\hbar}[[G]]^{\vee}$, be the QFSHA, resp. the QUEA, introduced in §2.4. Let $\mathcal{R} = \mathcal{R}^i \otimes \mathcal{R}_i$ (sum, possibly infinite, over repeated indices) be an R -matrix for $U_{\hbar}(\mathfrak{g})$, which is congruent to $1^{\otimes 2}$ modulo \hbar .*

Then, for the two morphisms $F_{\hbar}[[G]] \xrightarrow{\overleftarrow{\Phi}_{\mathcal{R}}} U_{\hbar}(\mathfrak{g})^{\text{cop}}$ and $F_{\hbar}[[G]] \xrightarrow{\overrightarrow{\Phi}_{\mathcal{R}}} U_{\hbar}(\mathfrak{g})^{\text{op}}$ in Proposition 5.2.1, the following holds:

(a) they take values inside $U_{\hbar}(\mathfrak{g})'$, and so they corestrict to morphisms

$$\overleftarrow{\Phi}'_{\mathcal{R}} : F_{\hbar}[[G]] \longrightarrow (U_{\hbar}(\mathfrak{g})')^{\text{cop}} = F_{\hbar}[[G^*]]^{\text{cop}}$$

and

$$\overrightarrow{\Phi}'_{\mathcal{R}} : F_{\hbar}[[G]] \longrightarrow (U_{\hbar}(\mathfrak{g})')^{\text{op}} = F_{\hbar}[[G^*]]^{\text{op}}$$

between QFSHA's for mutually dual (formal) Poisson groups;

(b) they uniquely extend to $F_{\hbar}[[G]]^{\vee} = (U_{\hbar}(\mathfrak{g})^*)^{\vee}$, i.e. they extend to morphisms

$$\overleftarrow{\Phi}^{\vee}_{\mathcal{R}} : U_{\hbar}(\mathfrak{g}^*) := F_{\hbar}[[G]]^{\vee} \longrightarrow U_{\hbar}(\mathfrak{g})^{\text{cop}}$$

and

$$\overrightarrow{\Phi}^{\vee}_{\mathcal{R}} : U_{\hbar}(\mathfrak{g}^*) := F_{\hbar}[[G]]^{\vee} \longrightarrow U_{\hbar}(\mathfrak{g})^{\text{op}}$$

between QUEA's for mutually dual Lie bialgebras.

Proof. (a) Recall that $\text{Ker}(\epsilon_{U_{\hbar}(\mathfrak{g})'}) =: J_{U_{\hbar}(\mathfrak{g})'} \subseteq (J_{U_{\hbar}(\mathfrak{g})'} + \mathbb{k}[[\hbar]]1_{U_{\hbar}(\mathfrak{g})'}) =: I_{U_{\hbar}(\mathfrak{g})'}$, and $U_{\hbar}(\mathfrak{g})$ is a topological Hopf algebra with respect to the $I_{U_{\hbar}(\mathfrak{g})}'$ -adic topology.

Recall also that, by Proposition 5.1.4(a), we can write $\mathcal{R} := \exp(\hbar^{-1}\vartheta)$ with $\vartheta \in U_{\hbar}(\mathfrak{g})' \otimes U_{\hbar}(\mathfrak{g})'$; we write the latter as $\vartheta = \vartheta^i \otimes \vartheta_i$ (sum over repeated indices), where $\vartheta^i, \vartheta_i \in J_{U_{\hbar}(\mathfrak{g})}'$. Then we have

$$\hbar^{-1}\vartheta = \hbar^{-1}\vartheta^i \otimes \vartheta_i = (\hbar^{-1}\vartheta^i) \otimes \vartheta_i = \theta^i \otimes \vartheta_i$$

with $\theta^i := \hbar^{-1}\vartheta^i \in \hbar^{-1}J_{U_{\hbar}(\mathfrak{g})}' \subseteq J_{U_{\hbar}(\mathfrak{g})} := \text{Ker}(\epsilon_{U_{\hbar}(\mathfrak{g})})$, where the latter inclusion follows by the basic properties of $U_{\hbar}(\mathfrak{g})'$, cf. [Ga1]. Now writing $(\theta^i \otimes \vartheta_i)^n = \theta_{[n]}^j \otimes \vartheta_{[n],j}$ for each $n \in \mathbb{N}$, we have in particular $\theta_{[n]}^j \in J_{U_{\hbar}(\mathfrak{g})}^n$ and $\vartheta_{[n],j} \in J_{U_{\hbar}(\mathfrak{g})'}^n$, for every $n \in \mathbb{N}$. When we expand \mathcal{R} , by all this we find

$$\mathcal{R} = \exp(\hbar^{-1}\vartheta) = \exp(\theta^i \otimes \vartheta_i) = \sum_{n \geq 0} \frac{1}{n!} (\theta^i \otimes \vartheta_i)^n = \sum_{n \geq 0} \frac{1}{n!} (\theta_{[n]}^j \otimes \vartheta_{[n],j})$$

Therefore, for every $\eta \in F_{\hbar}[[G]] := U_{\hbar}(\mathfrak{g})^*$ we have

$$\overleftarrow{\Phi}_{\mathcal{R}}(\eta) = \eta(\mathcal{R}^s) \mathcal{R}_s = \sum_{n \geq 0} \frac{1}{n!} \eta(\theta_{[n]}^j) \vartheta_{[n],j}$$

which describes a well-defined element (a convergent series, in the relevant topology!) of $(U_{\hbar}(\mathfrak{g})')^{\text{cop}}$ — equal to $U_{\hbar}(\mathfrak{g})'$ as a $\mathbb{k}[[\hbar]]$ -module — exactly because $\vartheta_{[n],j} \in J_{U_{\hbar}(\mathfrak{g})'}^n$ for each $n \in \mathbb{N}$. Thus $\overleftarrow{\Phi}_{\mathcal{R}}$ corestricts to $(U_{\hbar}(\mathfrak{g})')^{\text{cop}} = F_{\hbar}[[G^*]]^{\text{cop}}$ as claimed, q.e.d.

The proof for $\overrightarrow{\Phi}_{\mathcal{R}}$ goes exactly the same, just switching left and right.

(b) We begin acting as in the proof of (a) above, but switching the roles of left and right hand sides. Namely, we write $\hbar\vartheta = \hbar(\vartheta^i \otimes \vartheta_i) = \theta^i \otimes \vartheta_i$ where $\theta_i := \hbar^{-1}\vartheta_i \in \hbar^{-1}J_{U_{\hbar}(\mathfrak{g})}' \subseteq J_{U_{\hbar}(\mathfrak{g})} := \text{Ker}(\epsilon_{U_{\hbar}(\mathfrak{g})})$, and also $(\vartheta^i \otimes \theta_i)^n = \vartheta_{[n]}^j \otimes \theta_{[n],j}$, with $\vartheta_{[n]}^j \in J_{U_{\hbar}(\mathfrak{g})'}^n$ and $\theta_{[n],j} \in J_{U_{\hbar}(\mathfrak{g})}^n$, for all $n \in \mathbb{N}$. Then expanding \mathcal{R} yields

$$\mathcal{R} = \exp(\hbar^{-1}\vartheta) = \exp(\vartheta^i \otimes \theta_i) = \sum_{n \geq 0} \frac{1}{n!} (\vartheta^i \otimes \theta_i)^n = \sum_{n \geq 0} \frac{1}{n!} (\vartheta_{[n]}^j \otimes \theta_{[n],j})$$

hence for every $\mu \in U_{\hbar}(\mathfrak{g})^*$ we have

$$\overleftarrow{\Phi}_{\mathcal{R}}(\mu) := \mu(\mathcal{R}^s) \mathcal{R}_s = \sum_{n \geq 0} \frac{1}{n!} \mu(\vartheta_{[n]}^j) \theta_{[n],j} \quad (5.1)$$

Now, recall that $(U_{\hbar}(\mathfrak{g})^*)^\vee = (U_{\hbar}(\mathfrak{g})')^*$, by (2.18). Then we consider the formula (5.1) for any $\mu \in (U_{\hbar}(\mathfrak{g})^*)^\vee = (U_{\hbar}(\mathfrak{g})')^*$ — which contains $U_{\hbar}(\mathfrak{g})^*$. As all coefficients $\mu(\vartheta'_n)$ belong to $\mathbb{k}[[\hbar]]$, every partial sum in the right-hand side formal series is a well-defined element in $U_{\hbar}(\mathfrak{g})^{\text{cop}}$ — equal to $U_{\hbar}(\mathfrak{g})$ as a $\mathbb{k}[[\hbar]]$ -module. In addition, since $\vartheta_{[n]}^j \in J_{U_{\hbar}(\mathfrak{g})'}^n \subseteq I_{U_{\hbar}(\mathfrak{g})'}^n$ — for each $n \in \mathbb{N}$ — and $\mu : U_{\hbar}(\mathfrak{g})' \rightarrow \mathbb{k}[[\hbar]]$ is *continuous* (with respect to the $I_{U_{\hbar}(\mathfrak{g})'}^n$ -adic topology on the left and the \hbar -adic topology on the right), for every $s \in \mathbb{N}$ there exist n_s such that $\mu(\vartheta_{[n]}^j) \in \hbar^{n_s} \mathbb{k}[[\hbar]]$ for all $n \geq n_s$. This ensures that the formal series in (5.1) is actually convergent in the \hbar -adic topology of $U_{\hbar}(\mathfrak{g})$, thus describing a well-defined element in $U_{\hbar}(\mathfrak{g})$. Letting μ range freely inside $(U_{\hbar}(\mathfrak{g})^*)^\vee$, this proves that $\overleftarrow{\Phi}_{\mathcal{R}}$ does indeed extend from $U_{\hbar}(\mathfrak{g})^*$ to $(U_{\hbar}(\mathfrak{g})^*)^\vee = F_{\hbar}[[G^*]]^\vee =: U_{\hbar}(\mathfrak{g}^*)$, q.e.d.

Switching left and right in the arguments above we get the proof for $\overrightarrow{\Phi}_{\mathcal{R}}$ too. \square

Remark 5.2.4. Claim (a) of Theorem 5.2.3 above appears also in [EK], §4.5.

In the dual framework, the parallel result holds true as well:

Theorem 5.2.5. *Let $F_{\hbar}[[G]]$ be a QFSHA, let $U_{\hbar}(\mathfrak{g}) := F_{\hbar}[[G]]^*$ be its dual QUEA, as in §2.3.4, and let also $U_{\hbar}(\mathfrak{g}^*) := F_{\hbar}[[G]]^\vee$, resp. $F_{\hbar}[[G^*]] := U_{\hbar}(\mathfrak{g})'$, be the QUEA, resp. the QFSHA, introduced in §2.4. Let ρ be a ρ -comatrix for $F_{\hbar}[[G]]$, which is congruent to $\epsilon^{\otimes 2}$ modulo \hbar .*

Then, for the two morphisms $F_{\hbar}[[G]] \xrightarrow{\overleftarrow{\Psi}_{\rho}} U_{\hbar}(\mathfrak{g})^{\text{cop}}$ and $F_{\hbar}[[G]] \xrightarrow{\overrightarrow{\Psi}_{\rho}} U_{\hbar}(\mathfrak{g})^{\text{op}}$ in Proposition 5.2.2, the following holds:

(a) *they take values inside $U_{\hbar}(\mathfrak{g})'$, so they corestrict to morphisms*

$$\overleftarrow{\Psi}'_{\rho} : F_{\hbar}[[G]] \longrightarrow (U_{\hbar}(\mathfrak{g})')^{\text{cop}} = F_{\hbar}[[G^*]]^{\text{cop}}$$

and

$$\overrightarrow{\Psi}'_{\rho} : F_{\hbar}[[G]] \longrightarrow (U_{\hbar}(\mathfrak{g})')^{\text{op}} = F_{\hbar}[[G^*]]^{\text{op}}$$

between QFSHA's for mutually dual (formal) Poisson groups.

(b) *they uniquely extend to $U_{\hbar}(\mathfrak{g}^*) = F_{\hbar}[[G]]^\vee$, i.e. they extend to morphisms*

$$\overleftarrow{\Psi}^{\vee}_{\rho} : U_{\hbar}(\mathfrak{g}^*) = F_{\hbar}[[G]]^\vee \longrightarrow U_{\hbar}(\mathfrak{g})^{\text{cop}}$$

and

$$\overrightarrow{\Psi}^{\vee}_{\rho} : U_{\hbar}(\mathfrak{g}^*) = F_{\hbar}[[G]]^\vee \longrightarrow U_{\hbar}(\mathfrak{g})^{\text{op}}$$

between QUEA's for mutually dual Lie bialgebras.

Proof. (a) By the assumption $\rho \equiv \epsilon^{\otimes 2} \pmod{\hbar}$ and by Proposition 5.1.4(b), we can write ρ in the form $\rho = \exp_*(\hbar^{-1} \zeta)$ for some $\zeta \in \left((F_{\hbar}[[G]]^\vee)^{\widehat{\otimes} 2} \right)^*$. Then

$$\begin{aligned} \zeta \in \left((F_{\hbar}[[G]]^\vee)^{\widehat{\otimes} 2} \right)^* &= (F_{\hbar}[[G]]^\vee \widehat{\otimes} F_{\hbar}[[G]]^\vee)^* = (F_{\hbar}[[G]]^\vee)^* \widehat{\otimes} (F_{\hbar}[[G]]^\vee)^* = \\ &= (F_{\hbar}[[G]]^*)' \widehat{\otimes} (F_{\hbar}[[G]]^*)' = U_{\hbar}(\mathfrak{g}') \widehat{\otimes} U_{\hbar}(\mathfrak{g}') \end{aligned}$$

— thanks to (2.18) — hence $\hbar^{-1} \zeta \in \hbar^{-1} U_{\hbar}(\mathfrak{g}^*)' \widehat{\otimes} U_{\hbar}(\mathfrak{g}^*)'$. Now, the right-hand side of (2.12) for $\sigma := \rho$ implies $\zeta(1, a) = 0$ for all $a \in F_{\hbar}[[G]]$, hence

$\zeta(-, a) \in \text{Ker}(\epsilon_{U_h(\mathfrak{g})'})$ and so $\hbar^{-1}\zeta(-, a) \in (U_h(\mathfrak{g})')^\vee = U_h(\mathfrak{g}) = F_h[[G]]^*$ for all $a \in F_h[[G]]$. This implies that

$$\hbar^{-1}\zeta(-, -) \in (U_h(\mathfrak{g})')^\vee \otimes U_h(\mathfrak{g})' = F_h[[G]]^* \otimes U_h(\mathfrak{g})' \quad (5.2)$$

where hereafter we are being temporarily sloppy with the tensor product — we fix this later on. Clearly, (5.2) implies $\rho = \exp_*(\hbar^{-1}\zeta) \in F_h[[G]]^* \otimes U_h(\mathfrak{g})'$ as well. Therefore we get at once $\Psi_\rho^{\leftarrow}(\ell) := \rho(\ell, -) \in U_h(\mathfrak{g})'$ for all $\ell \in F_h[[G]]$, q.e.d. This proves the claim about $\overleftarrow{\Psi}_\rho$, and that concerning $\overrightarrow{\Psi}_\rho$ is entirely similar.

It remains to “dot your i 's” about the tensor product in (5.2). In fact, *a priori* we have $\rho \in F_h[[G]]^* \widehat{\otimes} F_h[[G]]^* = U_h(\mathfrak{g}) \widehat{\otimes} U_h(\mathfrak{g})$, hence also

$$\hbar^{-1}\zeta \in F_h[[G]]^* \widehat{\otimes} F_h[[G]]^* = U_h(\mathfrak{g}) \widehat{\otimes} U_h(\mathfrak{g})$$

— where the (completed, topological) tensor product “ $\widehat{\otimes}$ ” is considered. On the other hand, we have found that

$$\zeta \in (F_h[[G]]^*)' \widetilde{\otimes} (F_h[[G]]^*)' = U_h(\mathfrak{g})' \widetilde{\otimes} U_h(\mathfrak{g})'$$

— where the (completed, topological) tensor product “ $\widetilde{\otimes}$ ” is used. Then the critical point is: what kind of tensor product “ \otimes ” is taken in (5.2)?

Instead of giving a direct, formal answer to this question, we point out the following. First observe that $U_h(\mathfrak{g})' \widetilde{\otimes} U_h(\mathfrak{g})' = (U_h(\mathfrak{g}) \widehat{\otimes} U_h(\mathfrak{g}))'$ naturally embeds into $U_h(\mathfrak{g}) \widehat{\otimes} U_h(\mathfrak{g})$. Then, when $\hbar^{-1}\zeta \in U_h(\mathfrak{g}) \widehat{\otimes} U_h(\mathfrak{g})$ is expanded into some (suitably convergent) series $\hbar^{-1}\zeta = \beta^i \otimes \beta_i$ (summing over repeated indices) with $\beta^i, \beta_i \in U_h(\mathfrak{g})$ for all i , what we proved above is that we actually have $\beta_i \in U_h(\mathfrak{g})'$ ($\subseteq U_h(\mathfrak{g})$) for all indices i . This is what we loosely wrote as $\hbar^{-1}\zeta \in U_h(\mathfrak{g}) \otimes U_h(\mathfrak{g})' = F_h[[G]]^* \otimes U_h(\mathfrak{g})'$ in (5.2) above.

(b) Acting as in part (a), we find $\rho = \exp_*(\hbar^{-1}\zeta)$ with $\zeta \in (U_h(\mathfrak{g})')^{\widehat{\otimes}^2}$ and $\zeta \in (\text{Ker}(\epsilon_{U_h(\mathfrak{g})}))^{\widehat{\otimes}^2}$ too, so $\zeta \in (\text{Ker}(\epsilon_{U_h(\mathfrak{g})'}))^{\widehat{\otimes}^2}$. Since $\text{Ker}(\epsilon_{U_h(\mathfrak{g})'}) \subseteq \hbar \text{Ker}(\epsilon_{U_h(\mathfrak{g})})$, this implies that, expanding $\hbar^{-1}\zeta$ as a (convergent) series $\hbar^{-1}\zeta = \beta^i \otimes \beta_i$, we can assume $\beta^i \in U_h(\mathfrak{g})'$. As $U_h(\mathfrak{g})' = (F_h[[G]]^*)' = (F_h[[G]]^\vee)^*$, we end up with

$$\hbar^{-1}\zeta = \beta^i \otimes \beta_i \in (F_h[[G]]^\vee)^* \otimes U_h(\mathfrak{g}) \quad (5.3)$$

where again the meaning of the tensor product “ \otimes ” considered in this formula (along with the corresponding convergence issues) is handled just as in part (a). Finally, from (5.3) it follows at once that $\overleftarrow{\Psi}_\rho$ extends from $F_h[[G]]$ to $F_h[[G]]^\vee$ as claimed. This proves our statement for $\overleftarrow{\Psi}_\rho$, and the case of $\overrightarrow{\Psi}_\rho$ is entirely similar. \square

5.2.6. Duality properties. When we deal with a QUEA and a QFSHA which are dual to each other, it makes sense to compare the previous results. The outcome is that Proposition 2.2.12 turns to an enhanced version (with trivial proof), as follows:

Theorem 5.2.7. *Let $U_h(\mathfrak{g})$ be a QUEA, $F_h[[G]]$ a QFSHA, which are dual to each other, i.e. $F_h[[G]] = U_h(\mathfrak{g})^*$ and $U_h(\mathfrak{g}) = F_h[[G]]^*$. Let $\mathcal{R} = \rho$ be an R -matrix for $U_h(\mathfrak{g})$ and a ϱ -comatrix for $F_h[[G]]$, which is trivial modulo \hbar , i.e. congruent to $1^{\otimes 2}$,*

resp. to $\epsilon^{\otimes 2}$, modulo \hbar . Then, for the morphisms in Proposition 5.2.1, Proposition 5.2.2, Theorem 5.2.3 and Theorem 5.2.5 we have the following identifications

$$\overleftarrow{\Phi}_{\mathcal{R}} = \overleftarrow{\Psi}_{\rho}, \quad \overleftarrow{\Phi}'_{\mathcal{R}} = \overleftarrow{\Psi}'_{\rho}, \quad \overleftarrow{\Phi}_{\mathcal{R}}^{\vee} = \overleftarrow{\Psi}_{\rho}^{\vee} \quad \text{and} \quad \overrightarrow{\Phi}_{\mathcal{R}} = \overrightarrow{\Psi}_{\rho}, \quad \overrightarrow{\Phi}'_{\mathcal{R}} = \overrightarrow{\Psi}'_{\rho}, \quad \overrightarrow{\Phi}_{\mathcal{R}}^{\vee} = \overrightarrow{\Psi}_{\rho}^{\vee} \quad \square$$

5.2.8. Comparing morphisms (1). Let us fix assumptions as in Theorem 5.2.3: $U_{\hbar}(\mathfrak{g})$ is a given QUEA, $F_{\hbar}[[G]]$ its dual QFSHA, and $\mathcal{R} = \mathcal{R}^s \otimes \mathcal{R}_s$ is a (quantum) R -matrix of $U_{\hbar}(\mathfrak{g})$. Then from Theorem 5.2.3 we have Hopf algebra morphisms

$$F_{\hbar}[[G]] \xrightarrow{\overleftarrow{\Phi}'_{\mathcal{R}}} F_{\hbar}[[G^*]]^{\text{cop}}, \quad F_{\hbar}[[G]] \xrightarrow{\overrightarrow{\Phi}'_{\mathcal{R}}} F_{\hbar}[[G^*]]^{\text{op}} \quad (5.4)$$

between QFSHA's for mutually dual (formal) Poisson groups, and

$$U_{\hbar}(\mathfrak{g}^*) \xrightarrow{\overleftarrow{\Phi}_{\mathcal{R}}^{\vee}} U_{\hbar}(\mathfrak{g})^{\text{cop}}, \quad U_{\hbar}(\mathfrak{g}^*) \xrightarrow{\overrightarrow{\Phi}_{\mathcal{R}}^{\vee}} U_{\hbar}(\mathfrak{g})^{\text{op}} \quad (5.5)$$

between QUEA's for mutually dual Lie bialgebras, which we re-write in the form

$$U_{\hbar}(\mathfrak{g}^*)^{\text{cop}} \xrightarrow{\overleftarrow{\Phi}_{\mathcal{R}}^{\vee}} U_{\hbar}(\mathfrak{g}) \quad , \quad U_{\hbar}(\mathfrak{g}^*)^{\text{op}} \xrightarrow{\overrightarrow{\Phi}_{\mathcal{R}}^{\vee}} U_{\hbar}(\mathfrak{g}) \quad (5.6)$$

that is entirely equivalent. We now go and compare (5.4) and (5.6).

Recall that $F_{\hbar}[[G^*]] := U_{\hbar}(\mathfrak{g})'$ and $U_{\hbar}(\mathfrak{g}^*) := F_{\hbar}[[G]]^{\vee}$, which are in duality because $U_{\hbar}(\mathfrak{g})$ and $F_{\hbar}[[G]]$ are in duality (by construction) and we can apply (2.18). Then also $F_{\hbar}[[G^*]]^{\text{cop}}$ and $U_{\hbar}(\mathfrak{g}^*)^{\text{op}}$ are in duality, as well as $F_{\hbar}[[G^*]]^{\text{op}}$ and $U_{\hbar}(\mathfrak{g}^*)^{\text{cop}}$.

We are now ready to compare the morphisms in (5.4) with those in (5.6). Namely, we have a couple of diagrams

$$\begin{array}{ccc} F_{\hbar}[[G]] & \xrightarrow{\overleftarrow{\Phi}'_{\mathcal{R}}} & F_{\hbar}[[G^*]]^{\text{cop}} & & F_{\hbar}[[G]] & \xrightarrow{\overrightarrow{\Phi}'_{\mathcal{R}}} & F_{\hbar}[[G^*]]^{\text{op}} & (5.7) \\ \left. \vphantom{F_{\hbar}[[G]]} \right\} & & \left. \vphantom{F_{\hbar}[[G^*]]^{\text{cop}}} \right\} & & \left. \vphantom{F_{\hbar}[[G]]} \right\} & & \left. \vphantom{F_{\hbar}[[G^*]]^{\text{op}}} \right\} & \\ U_{\hbar}(\mathfrak{g}) & \xleftarrow{\overrightarrow{\Phi}_{\mathcal{R}}^{\vee}} & U_{\hbar}(\mathfrak{g}^*)^{\text{op}} & & U_{\hbar}(\mathfrak{g}) & \xleftarrow{\overleftarrow{\Phi}_{\mathcal{R}}^{\vee}} & U_{\hbar}(\mathfrak{g}^*)^{\text{cop}} \end{array}$$

where the vertical, twisting lines denote a relationship of mutual (Hopf) duality. Next result tells us that the link between the morphisms on top row and those underneath is indeed “the best possible one”:

Theorem 5.2.9. *The two morphisms in left-hand side, resp. in right-hand side, of (5.7) are adjoint to each other, that is for all $\eta \in U_{\hbar}(\mathfrak{g}^*)$ and $f \in F_{\hbar}[[G]]$ we have*

$$\left\langle \overleftarrow{\Phi}_{\mathcal{R}}^{\vee}(\eta), f \right\rangle = \left\langle \eta, \overleftarrow{\Phi}'_{\mathcal{R}}(f) \right\rangle \quad \text{and} \quad \left\langle \overrightarrow{\Phi}_{\mathcal{R}}^{\vee}(\eta), f \right\rangle = \left\langle \eta, \overrightarrow{\Phi}'_{\mathcal{R}}(f) \right\rangle$$

where by “ $\langle \ , \ \rangle$ ” we denote the pairing between any two Hopf algebras in duality.

Proof. It is enough to prove half of the claim — the other one being entirely similar — say the right-hand side. Direct computation yields

$$\begin{aligned} \left\langle \overleftarrow{\Phi}_{\mathcal{R}}^{\vee}(\eta), f \right\rangle &= \left\langle \langle \eta, \mathcal{R}^s \rangle \mathcal{R}_s, f \right\rangle = \langle \eta, \mathcal{R}^s \rangle \langle \mathcal{R}_s, f \rangle = \\ &= \langle \mathcal{R}_s, f \rangle \langle \eta, \mathcal{R}^s \rangle = \left\langle \eta, \mathcal{R}^s \langle \mathcal{R}_s, f \rangle \right\rangle = \left\langle \eta, \overrightarrow{\Phi}'_{\mathcal{R}}(f) \right\rangle \end{aligned}$$

for all $\eta \in U_{\hbar}(\mathfrak{g}^*)$ and $f \in F_{\hbar}[[G]]$, hence we are done. \square

As a second step, let now start with assumptions as in Theorem 5.2.5: $F_{\hbar}[[G]]$ is a given QFSHA, $U_{\hbar}(\mathfrak{g})$ its dual QUEA, and ρ is a (quantum) ϱ -comatrix of $F_{\hbar}[[G]]$. Then Theorem 5.2.5 provides Hopf algebra morphisms

$$F_{\hbar}[[G]] \xrightarrow{\overleftarrow{\Psi}'_{\rho}} F_{\hbar}[[G^*]]^{\text{cop}}, \quad F_{\hbar}[[G]] \xrightarrow{\overrightarrow{\Psi}'_{\rho}} F_{\hbar}[[G^*]]^{\text{op}} \quad (5.8)$$

between QFSHA's for mutually dual (formal) Poisson groups, and

$$U_{\hbar}(\mathfrak{g}^*) \xrightarrow{\overleftarrow{\Psi}^{\vee}_{\rho}} U_{\hbar}(\mathfrak{g})^{\text{cop}}, \quad U_{\hbar}(\mathfrak{g}^*) \xrightarrow{\overrightarrow{\Psi}^{\vee}_{\rho}} U_{\hbar}(\mathfrak{g})^{\text{op}} \quad (5.9)$$

between QUEA's for mutually dual Lie bialgebras; we re-write the latter as

$$U_{\hbar}(\mathfrak{g}^*)^{\text{cop}} \xrightarrow{\overleftarrow{\Psi}^{\vee}_{\rho}} U_{\hbar}(\mathfrak{g}), \quad U_{\hbar}(\mathfrak{g}^*)^{\text{op}} \xrightarrow{\overrightarrow{\Psi}^{\vee}_{\rho}} U_{\hbar}(\mathfrak{g}) \quad (5.10)$$

that is entirely equivalent. We now go and compare (5.8) and (5.10).

Acting as before (for the morphisms induced by an R -matrix), we find diagrams

$$\begin{array}{ccc} F_{\hbar}[[G]] \xrightarrow{\overleftarrow{\Psi}'_{\rho}} F_{\hbar}[[G^*]]^{\text{cop}} & & F_{\hbar}[[G]] \xrightarrow{\overrightarrow{\Psi}'_{\rho}} F_{\hbar}[[G^*]]^{\text{op}} \\ \left. \vphantom{F_{\hbar}[[G]]} \right\} & & \left. \vphantom{F_{\hbar}[[G]]} \right\} \\ U_{\hbar}(\mathfrak{g}) \xleftarrow{\overleftarrow{\Psi}^{\vee}_{\rho}} U_{\hbar}(\mathfrak{g}^*)^{\text{op}} & & U_{\hbar}(\mathfrak{g}) \xleftarrow{\overleftarrow{\Psi}^{\vee}_{\rho}} U_{\hbar}(\mathfrak{g}^*)^{\text{cop}} \end{array} \quad (5.11)$$

where the vertical, twisting lines denote a relationship of mutual (Hopf) duality. Again, the link between the morphisms on top row and those underneath turns out to be “the best possible one”, as the following result claims:

Theorem 5.2.10. *The two morphisms in left-hand side, resp. in right-hand side, of (5.7) are adjoint to each other, that is for all $\eta \in U_{\hbar}(\mathfrak{g}^*)$ and $f \in F_{\hbar}[[G]]$ we have*

$$\left\langle \overrightarrow{\Psi}^{\vee}_{\rho}(\eta), f \right\rangle = \left\langle \eta, \overleftarrow{\Psi}'_{\rho}(f) \right\rangle \quad \text{and} \quad \left\langle \overleftarrow{\Psi}'_{\rho}(\eta), f \right\rangle = \left\langle \eta, \overrightarrow{\Psi}'_{\rho}(f) \right\rangle$$

where by “ $\langle \ , \ \rangle$ ” we denote the pairing between any two Hopf algebras in duality.

Proof. We prove the left-hand side of the claim, the other side being similar. By direct computation we find

$$\left\langle \overrightarrow{\Psi}^{\vee}_{\rho}(\eta), f \right\rangle = \left\langle \rho(-, \eta), f \right\rangle = \rho(f, \eta) = \left\langle \eta, \rho(f, -) \right\rangle = \left\langle \eta, \overleftarrow{\Psi}'_{\rho}(f) \right\rangle$$

for all $\eta \in U_{\hbar}(\mathfrak{g}^*)$ and $f \in F_{\hbar}[[G]]$, as requested. \square

5.3. Morphisms from quasi- R -matrices and quasi- ϱ -comatrices.

We shall now explore what happens when the constructions leading to Proposition 2.2.9 or Proposition 2.2.10, respectively, is (tentatively) applied to a QFSHA and a quasi- R -comatrix for it, or to a QUEA and a quasi- ϱ -comatrix for it, respectively. Much like for deformations by quasi-twists or quasi-2-cocycles, we eventually achieve a nice result by “watching through the lens” of the QDP.

As a first result, we find that the construction of Hopf morphisms as in Proposition 2.2.9 can be applied again (though it might not be done through direct application of the abstract, general recipe) when the Hopf algebra under scrutiny is a QFSHA and its R -matrix is indeed only (!) a quasi- R -matrix.

Proposition 5.3.1. *Let $F_\hbar[[G]]$ be a QFSHA, and \mathcal{R} a quasi- R -matrix for it. Then the recipes in Proposition 2.2.9 provide two well-defined morphisms*

$$\begin{aligned} \overleftarrow{\Phi}_{\mathcal{R}} : F_\hbar[[G^*]] &:= (U_\hbar(\mathfrak{g}^*))^* = (F_\hbar[[G]]^\vee)^* \longrightarrow (F_\hbar[[G]]^\vee)^{\text{cop}} = U_\hbar(\mathfrak{g}^*)^{\text{cop}} \\ \overrightarrow{\Phi}_{\mathcal{R}} : F_\hbar[[G^*]] &:= (U_\hbar(\mathfrak{g}^*))^* = (F_\hbar[[G]]^\vee)^* \longrightarrow (F_\hbar[[G]]^\vee)^{\text{op}} = U_\hbar(\mathfrak{g}^*)^{\text{op}} \end{aligned}$$

Proof. This follows from a direct application of Proposition 5.2.1 to the QUEA $U_\hbar(\mathfrak{g}^*) := F_\hbar[[G]]^\vee$ and its R -matrix \mathcal{R} . \square

The previous result provide morphisms from a QFSHA to a QUEA, just like Proposition 5.2.1 did. Following a similar path, we shall now improve such a result — much like we did in §5.2 — finding a couple of morphisms between QFSHA's and another couple between QUEA's.

Theorem 5.3.2. *Assume that \mathcal{R} is a quasi- R -matrix for the QFSHA $F_\hbar[[G]]$, i.e. an R -matrix for the QUEA $F_\hbar[[G]]^\vee := U_\hbar(\mathfrak{g}^*)$, of the form $\mathcal{R} = \exp(\hbar^{-1}r)$ for some $r \in F_\hbar[[G]]^{\otimes 2}$. Then, for the two morphisms $\overleftarrow{\Phi}_{\mathcal{R}}$ and $\overrightarrow{\Phi}_{\mathcal{R}}$ in Proposition 5.3.1 above, the following holds:*

(a) *they corestrict to morphisms*

$$\begin{aligned} \overleftarrow{\Phi}'_{\mathcal{R}} : F_\hbar[[G^*]] &= (F_\hbar[[G]]^\vee)^* \longrightarrow \left((F_\hbar[[G]]^\vee)^{\text{cop}} \right)' = \left((F_\hbar[[G]]^\vee)' \right)^{\text{cop}} = F_\hbar[[G]]^{\text{cop}} \\ \text{and} \quad \overrightarrow{\Phi}'_{\mathcal{R}} : F_\hbar[[G^*]] &= (F_\hbar[[G]]^\vee)^* \longrightarrow \left((F_\hbar[[G]]^\vee)^{\text{op}} \right)' = \left((F_\hbar[[G]]^\vee)' \right)^{\text{op}} = F_\hbar[[G]]^{\text{op}} \end{aligned}$$

between QFSHA's for mutually dual (formal) Poisson groups;

(b) *they extend to morphisms*

$$\begin{aligned} \overleftarrow{\Phi}^\vee_{\mathcal{R}} : U_\hbar(\mathfrak{g}) &= F_\hbar[[G]]^* = \left((F_\hbar[[G]]^\vee)^* \right)^\vee \longrightarrow (F_\hbar[[G]]^\vee)^{\text{cop}} = U_\hbar(\mathfrak{g}^*)^{\text{cop}} \\ \text{and} \quad \overrightarrow{\Phi}^\vee_{\mathcal{R}} : U_\hbar(\mathfrak{g}) &= F_\hbar[[G]]^* = \left((F_\hbar[[G]]^\vee)^* \right)^\vee \longrightarrow (F_\hbar[[G]]^\vee)^{\text{op}} = U_\hbar(\mathfrak{g}^*)^{\text{op}} \end{aligned}$$

between QUEA's for mutually dual Lie bialgebras.

Proof. First of all, note that the chain of identities

$$\left((F_\hbar[[G]]^\vee)^{\text{cop}} \right)' = \left((F_\hbar[[G]]^\vee)' \right)^{\text{cop}} = F_\hbar[[G]]^{\text{cop}}$$

— and similarly with superscript “op” instead of “cop” throughout — is obvious from definitions along with the fact that Drinfeld's functors $(\)'$ and $(\)^\vee$ are inverse to each other. Similarly, it is also obviously true the chain of identities

$$\left((F_\hbar[[G]]^\vee)^* \right)^\vee = \left((F_\hbar[[G]]^\vee)' \right)^* = F_\hbar[[G]]^* = U_\hbar(\mathfrak{g})$$

As to the rest of the claim, everything follows from Theorem 5.2.3 applied to the QUEA $U_\hbar(\mathfrak{g}^*) := F_\hbar[[G]]^\vee$ along with its R -matrix \mathcal{R} . \square

Now we go for the dual constructions, concerning quasi- ρ -comatrices for a QUEA:

Proposition 5.3.3. *Assume that ρ is a quasi- ρ -comatrix for the QUEA $U_\hbar(\mathfrak{g})$, i.e. an element of the form $\rho = \exp_*(\hbar^{-1}\varrho)$ for some $\varrho \in \left(U_\hbar(\mathfrak{g})^{\otimes 2} \right)^*$ — taking into account Lemma 3.3.2 — which obeys (2.13).*

Then the recipes in Proposition 2.2.10 provide two well-defined morphisms

$$\begin{aligned} \overleftarrow{\Psi}_\rho : F_h[[G^*]] := U_h(\mathfrak{g})' &\longrightarrow \left((U_h(\mathfrak{g})')^* \right)^{\text{cop}} = (F_h[[G^*]]^*)^{\text{cop}} = U_h(\mathfrak{g}^*)^{\text{cop}} \\ \text{and} \quad \overrightarrow{\Psi}_\rho : F_h[[G^*]] := U_h(\mathfrak{g})' &\longrightarrow \left((U_h(\mathfrak{g})')^* \right)^{\text{op}} = (F_h[[G^*]]^*)^{\text{op}} = U_h(\mathfrak{g}^*)^{\text{op}} \end{aligned}$$

Proof. Everything follows from definitions once we apply Proposition 5.2.2 to the QFSHA $F_h[[G^*]] := U_h(\mathfrak{g})'$, also taking into the account the chain of (obvious) identities $(U_h(\mathfrak{g})')^* = F[[G^*]]^* = U_h(\mathfrak{g}^*)$. \square

Once again, the previous result provides morphisms from a QFSHA to a QUEA, and now we “enhance” it — like we did with Theorem 5.3.2 — finding morphisms between QFSHA’s and morphisms between QUEA’s:

Theorem 5.3.4. *Assume that ρ is a quasi- ϱ -comatrix for the QUEA $U_h(\mathfrak{g})$, i.e. an element of the form $\rho = \exp_*(\hbar^{-1}\varrho)$ for some $\varrho \in \left(U_h(\mathfrak{g})^{\widehat{\otimes} 2} \right)^*$ — taking into account Lemma 3.3.2 — which obeys (2.13). Then, for the two morphisms $\overleftarrow{\Psi}_\rho$ and $\overrightarrow{\Psi}_\rho$ in Proposition 5.3.3 above, the following holds:*

(a) *they corestrict to morphisms*

$$\begin{aligned} \overleftarrow{\Psi}'_\rho : F_h[[G^*]] = U_h(\mathfrak{g})' &\longrightarrow \left(\left((U_h(\mathfrak{g})')^* \right)^{\text{cop}} \right)' = (U_h(\mathfrak{g}^*)^{\text{cop}})' =: F_h[[G]]^{\text{cop}} \\ \text{and} \quad \overrightarrow{\Psi}'_\rho : F_h[[G^*]] = U_h(\mathfrak{g})' &\longrightarrow \left(\left((U_h(\mathfrak{g})')^* \right)^{\text{op}} \right)' = (U_h(\mathfrak{g}^*)^{\text{op}})' =: F_h[[G]]^{\text{op}} \end{aligned}$$

between QFSHA’s for mutually dual (formal) Poisson groups;

(b) *they extend to morphisms*

$$\begin{aligned} \overleftarrow{\Psi}^\vee_\rho : U_h(\mathfrak{g}) = (U_h(\mathfrak{g})')^\vee &\longrightarrow \left((U_h(\mathfrak{g})')^* \right)^{\text{cop}} = (F_h[[G^*]]^*)^{\text{cop}} = U_h(\mathfrak{g}^*)^{\text{cop}} \\ \text{and} \quad \overrightarrow{\Psi}^\vee_\rho : U_h(\mathfrak{g}) = (U_h(\mathfrak{g})')^\vee &\longrightarrow \left((U_h(\mathfrak{g})')^* \right)^{\text{op}} = (F_h[[G^*]]^*)^{\text{op}} = U_h(\mathfrak{g}^*)^{\text{op}} \end{aligned}$$

between QUEA’s for mutually dual Lie bialgebras.

Proof. As the functors $(\)'$ and $(\)^\vee$ are inverse to each other, definitions yield

$$\left(\left((U_h(\mathfrak{g})')^* \right)^{\text{cop}} \right)' = \left(\left((U_h(\mathfrak{g})')^* \right)' \right)^{\text{cop}} = \left(\left((U_h(\mathfrak{g})')^\vee \right)^* \right)^{\text{cop}} = (U_h(\mathfrak{g}^*)^{\text{cop}})' =: F_h[[G]]^{\text{cop}}$$

— and similarly with superscript “op” instead of “cop” throughout — also thanks to $(\)' \circ (\)^* = (\)^* \circ (\)^\vee$. Basing on this, the entire claim follows at once from Theorem 5.2.5 applied to the QFSHA $F_h[[G^*]] := U_h(\mathfrak{g})'$ and to its ϱ -comatrix ρ . \square

5.3.5. Duality properties. If we consider a QUEA and a QFSHA which are dual to each other, we can compare the previous results: thus we find the following “quasi-analogue” — whose proof is trivial again — of Theorem 5.2.7:

Theorem 5.3.6. *Let $U_h(\mathfrak{g})$ be a QUEA, $F_h[[G]]$ a QFSHA, which are dual to each other, i.e. $F_h[[G]] = U_h(\mathfrak{g})^*$ and $U_h(\mathfrak{g}) = F_h[[G]]^*$. Let $\rho = \mathcal{R}$ be a quasi- ρ -comatrix for $U_h(\mathfrak{g})$ and a quasi- R -matrix for $F_h[[G]]$.*

Then, for the morphisms in Proposition 5.3.1, Proposition 5.3.3, Theorem 5.3.2 and Theorem 5.3.4 we have the following identifications

$$\overleftarrow{\Phi}_{\mathcal{R}} = \overleftarrow{\Psi}_{\rho}, \quad \overleftarrow{\Phi}'_{\mathcal{R}} = \overleftarrow{\Psi}'_{\rho}, \quad \overleftarrow{\Phi}_{\mathcal{R}}^{\vee} = \overleftarrow{\Psi}_{\rho}^{\vee} \quad \text{and} \quad \overrightarrow{\Phi}_{\mathcal{R}} = \overrightarrow{\Psi}_{\rho}, \quad \overrightarrow{\Phi}'_{\mathcal{R}} = \overrightarrow{\Psi}'_{\rho}, \quad \overrightarrow{\Phi}_{\mathcal{R}}^{\vee} = \overrightarrow{\Psi}_{\rho}^{\vee} \quad \square$$

5.3.7. Comparing morphisms (2). We shall now compare morphisms among quantum groups provided by a quasi- R -matrix as above; the analysis is pretty similar to what we did with R -matrices, so we can be quicker.

Let us start with a QFSHA $F_h[[G]]$, with dual QUEA denoted by $U_h(\mathfrak{g})$, and a quasi- R -matrix \mathcal{R} for $F_h[[G]]$. Then Theorem 5.3.2 gives a couple of diagrams

$$\begin{array}{ccc} F_h[[G^*]] & \xrightarrow{\overleftarrow{\Phi}'_{\mathcal{R}}} & F_h[[G]]^{\text{cop}} & & F_h[[G^*]] & \xrightarrow{\overrightarrow{\Phi}'_{\mathcal{R}}} & F_h[[G]]^{\text{op}} & (5.12) \\ \left. \vphantom{F_h[[G^*]]} \right\} & & \left. \vphantom{F_h[[G]]^{\text{cop}}} \right\} & & \left. \vphantom{F_h[[G^*]]} \right\} & & \left. \vphantom{F_h[[G]]^{\text{op}}} \right\} & \\ U_h(\mathfrak{g}^*) & \xleftarrow{\overleftarrow{\Phi}_{\mathcal{R}}^{\vee}} & U_h(\mathfrak{g})^{\text{op}} & & U_h(\mathfrak{g}^*) & \xleftarrow{\overleftarrow{\Phi}_{\mathcal{R}}^{\vee}} & U_h(\mathfrak{g})^{\text{cop}} \end{array}$$

where the vertical, twisting lines denote a relationship of mutual (Hopf) duality while the horizontal arrows are Hopf algebra morphisms. Next result is the “quasi-analogue” of Theorem 5.2.9, telling us that the morphisms on top row and those underneath are “as close as possible”:

Theorem 5.3.8. *The two morphisms in left-hand side, resp. in right-hand side, of (5.12) are adjoint to each other, that is for all $\eta \in U_h(\mathfrak{g})$ and $f \in F_h[[G^*]]$ we have*

$$\left\langle \overrightarrow{\Phi}_{\mathcal{R}}^{\vee}(\eta), f \right\rangle = \left\langle \eta, \overleftarrow{\Phi}'_{\mathcal{R}}(f) \right\rangle \quad \text{and} \quad \left\langle \overleftarrow{\Phi}_{\mathcal{R}}^{\vee}(\eta), f \right\rangle = \left\langle \eta, \overrightarrow{\Phi}'_{\mathcal{R}}(f) \right\rangle$$

where by “ $\langle \ , \ \rangle$ ” we denote the pairing between any two Hopf algebras in duality.

Proof. The proof follows from Theorem 5.2.9 along with Theorem 5.1.5. \square

Similarly, let $U_h(\mathfrak{g})$ be a QUEA, with dual QFSHA denoted by $F_h[[G]]$, and let ρ be a quasi- ρ -comatrix $U_h(\mathfrak{g})$. Then Theorem 5.3.4 yields a couple of diagrams

$$\begin{array}{ccc} F_h[[G^*]] & \xrightarrow{\overleftarrow{\Psi}'_{\rho}} & F_h[[G]]^{\text{cop}} & & F_h[[G^*]] & \xrightarrow{\overrightarrow{\Psi}'_{\rho}} & F_h[[G]]^{\text{op}} & (5.13) \\ \left. \vphantom{F_h[[G^*]]} \right\} & & \left. \vphantom{F_h[[G]]^{\text{cop}}} \right\} & & \left. \vphantom{F_h[[G^*]]} \right\} & & \left. \vphantom{F_h[[G]]^{\text{op}}} \right\} & \\ U_h(\mathfrak{g}^*) & \xleftarrow{\overleftarrow{\Psi}_{\rho}^{\vee}} & U_h(\mathfrak{g})^{\text{op}} & & U_h(\mathfrak{g}^*) & \xleftarrow{\overleftarrow{\Psi}_{\rho}^{\vee}} & U_h(\mathfrak{g})^{\text{cop}} \end{array}$$

where the vertical, twisting lines denote a relationship of mutual (Hopf) duality while the horizontal arrows are Hopf algebra morphisms. We get now the “quasi-analogue” of Theorem 5.2.10, which claims that the morphisms on top row of (5.13) and those underneath are “as close as possible”:

Theorem 5.3.9. *The two morphisms in left-hand side, resp. in right-hand side, of (5.13) are adjoint to each other, that is for all $\eta \in U_h(\mathfrak{g})$ and $f \in F_h[[G^*]]$ we have*

$$\left\langle \overrightarrow{\Psi}_{\rho}^{\vee}(\eta), f \right\rangle = \left\langle \eta, \overleftarrow{\Psi}'_{\rho}(f) \right\rangle \quad \text{and} \quad \left\langle \overleftarrow{\Psi}_{\rho}^{\vee}(\eta), f \right\rangle = \left\langle \eta, \overrightarrow{\Psi}'_{\rho}(f) \right\rangle$$

where “ $\langle \ , \ \rangle$ ” denotes the pairing between any two Hopf algebras in duality.

Proof. Here again, the proof follows from Theorem 5.2.9 and Theorem 5.1.5. \square

5.4. Semiclassical morphisms induced by specialization.

We will now go and study the semiclassical limit of the various morphisms among quantum groups, considered in §§5.2 and 5.3 above, induced by R -matrices, ϱ -comatrices and their “quasi-counterparts”.

First we consider the case of an R -matrix \mathcal{R} for a given QUEA $U_{\hbar}(\mathfrak{g})$, whose dual QFSHA is $F_{\hbar}[[G]]$. With this assumptions, we recall the existence of the Hopf algebra morphisms in (5.7), which by Theorem 5.2.9 are pairwise mutually adjoint.

Specialising \hbar to 0, the left-hand side of (5.7) provides two mutually adjoint morphisms $F[[G]] \xrightarrow{\overleftarrow{\Phi}'_{\mathcal{R}}|_{\hbar=0}} F[[G^*]]^{\text{cop}}$ and $U(\mathfrak{g}^*)^{\text{op}} \xrightarrow{\overrightarrow{\Phi}_{\mathcal{R}}^{\vee}|_{\hbar=0}} U(\mathfrak{g})$, the first being a morphism of *Poisson* Hopf algebras, the second one of *co-Poisson* Hopf algebras. As they are mutually adjoint, each one of them defines one and the same morphism of formal Poisson groups $\phi_{\mathcal{R}}^+ : G_{\text{op}}^* \longrightarrow G$ where G_{op}^* is the *opposite* (i.e., with opposite product) formal Poisson group to G^* . Note that $\phi_{\mathcal{R}}^+ : G_{\text{op}}^* \longrightarrow G$ is directly defined by $\overleftarrow{\Phi}'_{\mathcal{R}}|_{\hbar=0}$, while the morphism of Lie bialgebras $d\phi_{\mathcal{R}}^+ : \mathfrak{g}_{\text{op}}^* \longrightarrow \mathfrak{g}$ can be deduced directly from $\overrightarrow{\Phi}_{\mathcal{R}}^{\vee}|_{\hbar=0}$, by restriction to $\mathfrak{g}_{\text{op}}^*$ and corestriction to \mathfrak{g} .

Similarly, specialising \hbar to 0 the right-hand side of (5.7) yields two mutually adjoint morphisms $F[[G]] \xrightarrow{\overrightarrow{\Phi}'_{\mathcal{R}}|_{\hbar=0}} F[[G^*]]^{\text{op}}$ and $U(\mathfrak{g}^*)^{\text{cop}} \xrightarrow{\overleftarrow{\Phi}_{\mathcal{R}}^{\vee}|_{\hbar=0}} U(\mathfrak{g})$ which in turn defines one single morphism of formal Poisson groups $\phi_{\mathcal{R}}^- : G_{\text{cop}}^* \longrightarrow G$ where now G_{cop}^* denotes the *co-opposite* formal Poisson group to G^* — i.e., with same product but opposite Poisson structure. This goes along with its associated morphism of Lie bialgebras $d\phi_{\mathcal{R}}^- : \mathfrak{g}_{\text{cop}}^* \longrightarrow \mathfrak{g}$. In short, we have pairs of morphisms

$$G_{\text{op}}^* \xrightarrow{\phi_{\mathcal{R}}^+} G, \quad G_{\text{cop}}^* \xrightarrow{\phi_{\mathcal{R}}^-} G \quad \text{and} \quad \mathfrak{g}_{\text{op}}^* \xrightarrow{d\phi_{\mathcal{R}}^+} \mathfrak{g}, \quad \mathfrak{g}_{\text{cop}}^* \xrightarrow{d\phi_{\mathcal{R}}^-} \mathfrak{g} \quad (5.14)$$

of formal Poisson groups and of Lie bialgebras, respectively.

Second, we consider the case of a ϱ -comatrix ρ for a given QFSHA $F_{\hbar}[[G]]$, with dual QUEA $U_{\hbar}(\mathfrak{g})$. In this case, there exist Hopf algebra morphisms as in (5.11), which are pairwise mutually adjoint due to Theorem 5.2.10.

Acting as before, specialising \hbar to 0 we find that the semiclassical limits of these (quantum) morphisms eventually define two pairs of morphisms

$$G_{\text{op}}^* \xrightarrow{\psi_{\rho}^+} G, \quad G_{\text{cop}}^* \xrightarrow{\psi_{\rho}^-} G \quad \text{and} \quad \mathfrak{g}_{\text{op}}^* \xrightarrow{d\psi_{\rho}^+} \mathfrak{g}, \quad \mathfrak{g}_{\text{cop}}^* \xrightarrow{d\psi_{\rho}^-} \mathfrak{g} \quad (5.15)$$

of formal Poisson groups and of Lie bialgebras, respectively.

Third, to compare the two constructions, assume that, given mutually dual quantum groups $U_{\hbar}(\mathfrak{g})$ and $F_{\hbar}[[G]]$, we pick a single element $\mathcal{R} = \rho$, thought of simultaneously as an R -matrix for $U_{\hbar}(\mathfrak{g})$ and as a ϱ -comatrix for $F_{\hbar}[[G]]$, much in the spirit of Proposition 2.2.7 and Theorem 2.2.12. Then morphisms as in (5.14) and (5.15) are defined: but in addition, directly by Theorem 5.2.7 we get at once that

$$\phi_{\mathcal{R}}^+ = \psi_{\rho}^+, \quad \phi_{\mathcal{R}}^- = \psi_{\rho}^- \quad \text{and} \quad d\phi_{\mathcal{R}}^+ = d\psi_{\rho}^+, \quad d\phi_{\mathcal{R}}^- = d\psi_{\rho}^-$$

If one works instead with quasi- R -matrices and quasi- ϱ -comatrices, the roles of G and G^* are reversed, but for the rest the analysis is entirely similar (so we may be more sketchy). Therefore, assume we have dual quantum groups $U_{\hbar}(\mathfrak{g})$ and $F_{\hbar}[[G]]$.

Given a quasi- R -matrix \mathcal{R} for $F_{\hbar}[[G]]$, the Hopf algebra morphisms in Theorem 5.3.2 give rise (through their semiclassical limit) to two pairs of morphisms

$$G_{\text{op}} \xrightarrow{\underline{\phi}_{\mathcal{R}}^+} G^* , \quad G_{\text{cop}} \xrightarrow{\underline{\phi}_{\mathcal{R}}^-} G^* \quad \text{and} \quad \mathfrak{g}_{\text{op}} \xrightarrow{d\underline{\phi}_{\mathcal{R}}^+} \mathfrak{g}^* , \quad \mathfrak{g}_{\text{cop}} \xrightarrow{d\underline{\phi}_{\mathcal{R}}^-} \mathfrak{g}^* \quad (5.16)$$

of formal Poisson groups and of Lie bialgebras, respectively.

Similarly, if ρ is a quasi- ρ -comatrix for $U_{\hbar}(\mathfrak{g})$, the Hopf algebra morphisms in Theorem 5.3.4 define (via their semiclassical limit) two pairs of morphisms

$$G_{\text{op}} \xrightarrow{\underline{\psi}_{\rho}^+} G^* , \quad G_{\text{cop}} \xrightarrow{\underline{\psi}_{\rho}^-} G^* \quad \text{and} \quad \mathfrak{g}_{\text{op}} \xrightarrow{d\underline{\psi}_{\rho}^+} \mathfrak{g}^* , \quad \mathfrak{g}_{\text{cop}} \xrightarrow{d\underline{\psi}_{\rho}^-} \mathfrak{g}^* \quad (5.17)$$

of formal Poisson groups and of Lie bialgebras, respectively.

Finally, if $\mathcal{R} = \rho$ — in the spirit of Proposition 2.2.7, more precisely like in Theorem 5.3.6 — then Theorem 5.3.6 gives at once

$$\underline{\phi}_{\mathcal{R}}^+ = \underline{\psi}_{\rho}^+ , \quad \underline{\phi}_{\mathcal{R}}^- = \underline{\psi}_{\rho}^- \quad \text{and} \quad d\underline{\phi}_{\mathcal{R}}^+ = d\underline{\psi}_{\rho}^+ , \quad d\underline{\phi}_{\mathcal{R}}^- = d\underline{\psi}_{\rho}^-$$

Studying in depth all the morphisms introduced above seems to be quite an interesting problem; we cannot, however, cope with in the present paper — we just finish with a comparison with previous results.

Assume we have an R -matrix \mathcal{R} for a given QUEA $U_{\hbar}(\mathfrak{g})$, whose dual QFSHA is $F_{\hbar}[[G]] := U_{\hbar}(\mathfrak{g})^*$. It is well-known that the “semiclassical limit” of \mathcal{R} , that is $r := \frac{\mathcal{R} - 1^{\otimes 2}}{\hbar} \pmod{\hbar}$, is in turn a “classical r -matrix” for the Lie bialgebra \mathfrak{g} .

Then Lie bialgebra morphisms $\mathfrak{g}_{\text{op}}^* \xrightarrow{\varphi_r^+} \mathfrak{g}$ and $\mathfrak{g}_{\text{cop}}^* \xrightarrow{\varphi_r^-} \mathfrak{g}$ are defined directly through r itself — with no need of \mathcal{R} , nor of $U_{\hbar}(\mathfrak{g})$, nor $F_{\hbar}[[G]]$, cf. [CP], §2.1, or [Mj], §8.1. Tracking the various constructions involved — in particular, the functor $F_{\hbar}[[G]] \mapsto F_{\hbar}[[G]]^{\vee} =: U_{\hbar}(\mathfrak{g}^*)$ — by direct comparison one immediately sees that

$$\varphi_r^+ = d\phi_{\mathcal{R}}^+ \quad \text{and} \quad \varphi_r^- = d\phi_{\mathcal{R}}^-$$

In particular, we get that *the morphisms $d\phi_{\mathcal{R}}^{\pm}$ depend on r alone, rather than on \mathcal{R} , hence the same is true for the morphisms $\phi_{\mathcal{R}}^{\pm}$* ; indeed, both facts can also be easily proved by direct inspection. Similarly, one can prove, via direct analysis again, or by a duality argument from the previous result, that *the morphisms ψ_{ρ}^{\pm} and $d\psi_{\rho}^{\pm}$ depend only on the “classical ρ -comatrix” $\rho_0 := \frac{\rho - \epsilon^{\otimes 2}}{\hbar} \pmod{\hbar}$ alone, rather than on ρ .*

REFERENCES

- [Ch] H.-X. CHEN, *Quasitriangular Structures of Bicrossed Coproducts*, J. Algebra **204** (1998), 504–531.
- [CG] N. CICCOLI, L. GUERRA, *The Variety of Lie Bialgebras*, J. Lie Theory **13** (2003), no. 2, 577–588.
- [CP] V. CHARI, A. PRESSLEY, *A guide to quantum group*, Cambridge University Press, Cambridge, 1995.

- [Doi] Y. DOI, *Braided bialgebras and quadratic bialgebras*, Comm. Algebra **21** (1993), no. 5, 1731–1749.
- [Dr] V. G. DRINFELD, *Quantum groups*, Proc. Int. Congr. Math., Berkeley 1986, vol. **1** (1987), 798–820.
- [EH] B. ENRIQUEZ, G. HALBOUT, *An \hbar -adic valuation property of universal R -matrices*, J. Algebra **261** (2003), no. 2, 434–447.
- [EK] P. ETINGOF, D. KAZHDAN, *Quantization of Lie bialgebras I*, Selecta Math. (N.S.) **2** (1996), no. 1, 1–41.
- [ES] P. ETINGOF, O. SCHIFFMANN, *Lectures on quantum groups*, Second edition. Lectures in Mathematical Physics, International Press, Somerville, MA, 2002, xii+242 pp.
- [Ga1] F. GAVARINI, *The quantum duality principle*, Ann. Inst. Fourier **52** (2002), no. 3, 809–834.
- [Ga2] ———, *Quantum duality principle for quantum continuous Kac-Moody algebras*, J. Lie theory **32** (2022), no. 3, 839–862.
- [GaGa1] G. A. GARCÍA, F. GAVARINI, *Twisted deformations vs. cocycle deformations for quantum groups*, Commun. Contemp. Math. **23** (2021), no. 8, Paper No. 2050084, 56 pp.
- [GaGa2] G. A. GARCÍA, F. GAVARINI, *Formal multiparameter quantum groups, deformations and specializations*, Annales de l’Institut Fourier, 117 pages (to appear) — preprint arXiv:2203.11023 [math.QA] (2022).
- [Ks] C. KASSEL, *Quantum groups*, Graduate Texts in Mathematics **155**, Springer-Verlag, New York, 1995, xii+531 pp.
- [KS] A. KLIMYK, K. SCHMÜDGEN, *Quantum groups and their representations*, Texts and Monographs in Physics, Springer-Verlag, Berlin, 1997, xx+552 pp.
- [Mj] S. MAJID, *Foundations of quantum groups*, Cambridge University Press, Cambridge, 1995.
- [MW] S. MERKULOV, T. WILLWACHER, *Deformation theory of Lie bialgebra properads*, Geometry and physics, Vol. I, 219–247, Oxford Univ. Press, Oxford, 2018.
- [Ra] D. E. RADFORD, *Hopf algebras*, Series on Knots and Everything **49**, World Scientific Publishing Co. Pte. Ltd., Hackensack, NJ, 2012.

DEPARTAMENTO DE MATEMÁTICA, FACULTAD DE CIENCIAS EXACTAS
 UNIVERSIDAD NACIONAL DE LA PLATA — CMALP-CIC-CONICET
 1900 LA PLATA, PCIA. DE BUENOS AIRES, ARGENTINA
 E-MAIL: ggarcia@mate.unlp.edu.ar

DIPARTIMENTO DI MATEMATICA,
 UNIVERSITÀ DEGLI STUDI DI ROMA “TOR VERGATA”
 VIA DELLA RICERCA SCIENTIFICA 1 — I-00133 ROMA, ITALY
 E-MAIL: gavarini@mat.uniroma2.it
 ISTITUTO NAZIONALE DI ALTA MATEMATICA “FRANCESCO SEVERI” — GNSAGA