# Ramsey multiplicity of apices of trees\* Ander Lamaison<sup>§</sup>

Daniel Kráľ<sup>†</sup> Matjaž Krnc<sup>‡</sup>

#### Abstract

A graph H is common if its Ramsey multiplicity, i.e., the minimum number of monochromatic copies of H contained in any 2-edge-coloring of  $K_n$ , is asymptotically the same as the number of monochromatic copies in the random 2-edge-coloring of  $K_n$ . Erdős conjectured that every complete graph is common, which was disproved by Thomason in the 1980s. Till today, a classification of common graphs remains a widely open challenging problem. Grzesik, Lee, Lidický and Volec [Combin. Prob. Comput. 31 (2022), 907–923 conjectured that every k-apex of any connected Sidorenko graph is common. We prove for  $k \leq 5$  that the k-apex of any tree is common.

#### Introduction 1

Ramsey's Theorem [18] asserts that for every graph H, every 2-edge-coloring of a sufficiently large complete graph contains a monochromatic copy of H. In this short paper, we are interested in the quantitative version of the statement: What is the minimum number of monochromatic copies of H in a 2-edge-coloring of  $K_n$ ? The Ramsey multiplicity of a graph H, which is denoted by M(H, n), is the minimum number of monochromatic (labeled) copies of H in a 2-edge-coloring of  $K_n$ , and the Ramsey multiplicity constant of H, denoted by C(H), is the limit

$$C(H) = \lim_{n \to \infty} \frac{M(H, n)}{n^{\nu(H)}}.$$

<sup>\*</sup>The first and third authors were supported by the MUNI Award in Science and Humanities (MUNI/I/1677/2018) of the Grant Agency of Masaryk University. The second author was supported by Slovenian Research and Innovation Agency (research projects J1-4008 and J1-3002).

<sup>&</sup>lt;sup>†</sup>Faculty of Informatics, Masaryk University, Botanická 68A, 602 00 Brno, Czech Republic. E-mail: dkral@fi.muni.cz.

<sup>&</sup>lt;sup>‡</sup>Faculty of Mathematics, Natural Sciences and Information Technologies, University of Primorska, Glagoljaška 8, SI-6000 Koper, Slovenia. E-mail: matjaz.krnc@upr.si.

<sup>&</sup>lt;sup>§</sup>Institute for Basic Science, 55 Expo-ro, Yuseong-gu, 34126 Daejeon, South Korea. E-mail: ander@ibs.re.kr. Previous affiliation: Faculty of Informatics, Masaryk University, Botanická 68A, 602 00 Brno, Czech Republic.

The random 2-edge-coloring yields that  $C(H) \leq 2^{1-e(H)}$  for every graph H, and those graphs H such that  $C(H) = 2^{1-e(H)}$  are called *common*. For example, Goodman's Theorem [10] implies that  $K_3$  is common. Since every graph that has the Sidorenko property is common, the famous conjecture of Sidorenko [20] and of Erdős and Simonovits [8], which asserts that every bipartite graph has the Sidorenko property (the definition is given in Section 2), would imply, if true, that every bipartite graph is common.

The notion of common graphs can be traced to the 1960s when Erdős [7] conjectured that every complete graph is common. Later, Burr and Rosta [2] extended the conjecture to all graphs. Both conjectures turned out to be false: Sidorenko [22, 23] showed that a triangle with a pendant edge is not common and Thomason [25] showed that  $K_4$  is not common. More generally, any graph containing  $K_4$  is not common [13], and any nonbipartite graph can be made to be not common by adding a sufficiently large tree [9].

On the positive side, every graph with the Sidorenko property is common. In particular, families of bipartite graphs known to have the Sidorenko property [1, 3-6, 23, 24] provide examples of bipartite graphs that are common. Examples of non-bipartite common graphs include odd cycles [23] and even wheels [13, 21], and more generally graphs that can be recursively obtained by gluing copies of  $K_3$  by a vertex or an edge [11]. The existence of a common graph with chromatic number at least four was open until 2012 when the flag algebra method of Razborov [19] resulted in a proof that the 5-wheel is common [12]. The existence of common graphs with arbitrarily large chromatic number was established only recently [15]; also see [14] for the construction of such highly connected graphs derived from those presented in [15].

A characterization of common graphs remains an intriguing open problem, which is very far from our reach. Our work is motivated by the following conjecture of Grzesik, Lee, Lidický and Volec [11]. Recall that the k-apex of a graph G is the graph obtained from G by adding k vertices and making each of the new k vertices adjacent to every vertex of G (however, the k new vertices remain to form an independent set).

## **Conjecture 1** (Grzesik, Lee, Lidický and Volec [11, Conjecture 1.3]). If G is a connected graph that has the Sidorenko property, then the k-apex of G is common for every k.

Recall that the above mentioned conjecture of Sidorenko [20] and of Erdős and Simonovits [8] asserts that every bipartite graph has the Sidorenko property, i.e., if both conjectures were true, then every k-apex of every bipartite graph would be common. The assumption in Conjecture 1 that the graph G is connected is necessary: the 1-apex of the union of  $K_1$  and  $K_2$  is a triangle with a pendant edge, which is not common [22, 23].

Conjecture 1 is known for a small number of special classes of graphs: 1-apices of even cycles, which are even wheels [13,21], 1-apices of trees, which are graphs obtained by gluing triangles on edges [13,21], 2-apices of even cycles [11], and k-apices of connected bipartite graphs with at most five vertices [11]. Our main result is a proof of the conjecture for k-apices of trees when  $k \leq 5$ .

**Theorem 1.** Let T be a tree. The k-apex of T for  $k \in \{1, 2, 3, 4, 5\}$  is common.



Figure 1: The graphs  $S_k$  and  $T_k$  for  $k \in \{1, 2, 3, 4\}$ .

Note that Theorem 1 is the first example of an infinite class of bipartite graphs such that k-apices of the graphs from the class are common for  $k \ge 3$ . We also remark that, unlike some results in the area, the proof of Theorem 1 is not computer assisted.

The paper is structured as follows. After reviewing notation in Section 2, we prove Theorem 1 in Section 3. Interestingly, we prove a stronger statement that concerns all *n*vertex graphs with n-1 edges that have the Sidorenko property. In Section 4, we discuss the possibility of extending our arguments to k-apices of trees for k > 5 and present an example that shows that the inequality claimed in Lemma 2, which is essential in our argument, fails for  $k \ge 12$ .

#### 2 Notation

We now review notation used throughout the paper. All graphs considered in this paper are finite and simple. If G is a graph, then V(G) denotes the vertex set of G and E(G)denotes the edge set of G. The number of vertices of a graph G is denoted by v(G) and the number of edges by e(G). Our arguments often involve the following two classes of graphs and so we introduce particular notation for them (see Figure 1 for illustration):  $S_k$  denotes a *star* with k leaves, and  $T_k$  denotes a *triangular book* with k triangles (the complete tripartite  $K_{1,1,k}$ ). Finally, the k-apex of a graph G, which is denoted by  $G^{+k}$ , is the graph obtained from G by adding k new vertices and making each of the new k vertices adjacent to every vertex of G (the k newly added vertices remain to form an independent set). In particular,  $S_k$  is the k-apex of  $K_1$  and  $T_k$  is the k-apex of  $K_2$ .

We treat the notion of common graphs using tools from the theory of graph limits; we refer to the monograph by Lovász [17] for a comprehensive treatment of graph limits. In the theory of graph limits, large graphs are represented by an analytic object called graphon. A graphon is a symmetric measurable function  $W : [0, 1]^2 \rightarrow [0, 1]$ , i.e., W(x, y) = W(y, x)for all  $x, y \in [0, 1]$ . Since a graphon can be thought of as a continuous version of the adjacency matrix of a graph (although a more precise view is as the limit of regularity partitions that approximate large graphs), we will refer to the elements of the domain [0, 1] as vertices of W. The homomorphism density of a graph G in a graphon W is defined as

$$t(G, W) = \int_{[0,1]^{V(G)}} \prod_{uv \in E(G)} W(x_u, x_v) \, \mathrm{d}x_{V(G)}.$$

We say that a graph G has the Sidorenko property if  $t(G, W) \ge t(K_2, W)^{e(G)}$  for every graphon W. Recall that the aforementioned conjecture of Sidorenko [20] and of Erdős and

Simonovits [8] asserts that every bipartite graph has the Sidorenko property. It is interesting to note that a graph G has the Sidorenko property if and only if G is k-common for every k [16], i.e., the random k-edge-coloring of a complete graph asymptotically minimizes the number of monochromatic copies of G among all k-edge-colorings of the host complete graph.

We will often think of a graphon W as representing one color class of edges while the complementary color class is represented by the graphon 1 - W, which is defined as (1-W)(x,y) = 1 - W(x,y). With this view in mind, we define the *monochromatic density* of a graph G in a graphon W as

$$m(G, W) = t(G, W) + t(G, 1 - W).$$

The definition of being common can now be cast as follows: a graph G is common if and only if  $m(G, W) \ge 2^{1-e(G)}$  for every graphon W.

In our arguments, we often consider monochromatic common neighborhoods of k vertices of a graphon. To make our presentation easier, we set  $B_k$  as  $\{0,1\} \times [0,1]^k$  and we think of the first coordinate as representing the color class and the remaining k coordinates as the k-tuple of vertices of a graphon. Using this interpretation, we define the function  $d_k^W : B_k \to [0,1]$  that represents the monochromatic common degree (the size of monochromatic common neighborhood) of a k-tuple of vertices as follows:

$$d_k^W(x_0, x_1, \dots, x_k) = \begin{cases} \int \prod_{i=1}^k W(x_i, y) \, \mathrm{d}y & \text{if } x_0 = 0\\ \int \prod_{i=1}^k (1 - W(x_i, y)) \, \mathrm{d}y & \text{if } x_0 = 1. \end{cases}$$

Observe that

$$m(S_k, W) = \int_{B_k} d_k^W(z) \, \mathrm{d}z$$

for every k. Similarly, we define the function  $\delta_k^W : B_k \to [0, 1]$  that represents the edge density of the monochromatic common neighborhood of a k-tuple of vertices as follows:

$$\delta_k^W(x_0, x_1, \dots, x_k) = \begin{cases} \int W(y, y') \prod_{i=1}^k W(x_i, y) W(x_i, y') \, \mathrm{d}y \, \mathrm{d}y' \\ \frac{[0, 1]^2}{d_k^W(x_0, x_1, \dots, x_k)^2} & \text{if } x_0 = 0 \end{cases}$$

$$\int (1 - W(y, y')) \prod_{i=1}^k (1 - W(x_i, y))(1 - W(x_i, y')) \, \mathrm{d}y \, \mathrm{d}y' \\ \frac{[0, 1]^2}{d_k^W(x_0, x_1, \dots, x_k)^2} & \text{if } x_0 = 1 \end{cases}$$

Observe that

$$m(T_k, W) = \int\limits_{B_k} d_k^W(z)^2 \delta_k^W(z) \,\mathrm{d}z,$$

and more generally, if a graph G has the Sidorenko property, it holds that

$$m(G^{+k}, W) \ge \int_{B_k} d_k^W(z)^{v(G)} \delta_k^W(z)^{e(G)} \,\mathrm{d}z$$
(1)

for every k.

### 3 Main result

In this section, we present the proof of our main result. The following lemma is the core of our argument. In Section 4, we present an example that the statement of the lemma fails for  $k \ge 12$ .

**Lemma 2.** For every  $k \in \{1, \ldots, 5\}$  and every graphon W, it holds that

$$\frac{m(T_k, W)}{m(S_k, W)} \ge 2^{-(k+1)}.$$
(2)

*Proof.* Fix a graphon W. We define a function  $h : [0,1] \to [-1/2, +1/2]$ , which measures how much the degree of a particular vertex x of W differs from 1/2, as follows:

$$h(x) = \int_{[0,1]} W(x,y) \, \mathrm{d}y - \frac{1}{2}$$

Observe that

$$m(S_k, W) = \int_{[0,1]} \left(\frac{1}{2} + h(x)\right)^k + \left(\frac{1}{2} - h(x)\right)^k dx$$
(3)

for every positive integer k. Goodman's formula [10] asserts that

$$m(T_1, W) = \frac{3}{2}m(S_2, W) - \frac{1}{2},$$

which yields that

$$m(T_1, W) = \frac{3}{2} \int_{[0,1]} \left(\frac{1}{2} + h(x)\right)^k + \left(\frac{1}{2} - h(x)\right)^k dx - \frac{1}{2} = \frac{1}{4} + 3 \int_{[0,1]} h(x)^2 dx.$$

The well-known inequality  $m(T_k, W) \ge m(T_1, W)^k$ , which follows from a standard Hölder type argument, implies that

$$m(T_k, W) \ge \left(\frac{1}{4} + 3\int_{[0,1]} h(x)^2 \,\mathrm{d}x\right)^k \ge \frac{1}{2^{2k}} + \frac{3k}{2^{2k-2}}\int_{[0,1]} h(x)^2 \,\mathrm{d}x \tag{4}$$

for every positive integer k.

If k = 1, we derive the desired estimate from (3) and (4) as follows:

$$\frac{m(T_1, W)}{m(S_1, W)} \ge \frac{\frac{1}{4} + 3 \int\limits_{[0,1]} h(x)^2 \, \mathrm{d}x}{\int\limits_{[0,1]} \left(\frac{1}{2} + h(x)\right) + \left(\frac{1}{2} - h(x)\right) \, \mathrm{d}x} = \frac{\frac{1}{4} + 3 \int\limits_{[0,1]} h(x)^2 \, \mathrm{d}x}{1} \ge \frac{1}{4}.$$

If k = 2, we again derive the desired estimate from (3) and (4):

$$\frac{m(T_2, W)}{m(S_2, W)} \ge \frac{\frac{1}{16} + \frac{6}{4} \int\limits_{[0,1]} h(x)^2 \,\mathrm{d}x}{\int\limits_{[0,1]} \left(\frac{1}{2} + h(x)\right)^2 + \left(\frac{1}{2} - h(x)\right)^2 \,\mathrm{d}x} = \frac{\frac{1}{16} + \frac{24}{16} \int\limits_{[0,1]} h(x)^2 \,\mathrm{d}x}{\frac{1}{2} + \frac{4}{2} \int\limits_{[0,1]} h(x)^2 \,\mathrm{d}x} \ge \frac{1}{8}.$$

And if k = 3, we derive the desired estimate from (3) and (4) as follows:

$$\frac{m(T_3, W)}{m(S_3, W)} \ge \frac{\frac{1}{64} + \frac{9}{16} \int\limits_{[0,1]} h(x)^2 \,\mathrm{d}x}{\int\limits_{[0,1]} \left(\frac{1}{2} + h(x)\right)^3 + \left(\frac{1}{2} - h(x)\right)^3 \,\mathrm{d}x} = \frac{\frac{1}{64} + \frac{36}{64} \int\limits_{[0,1]} h(x)^2 \,\mathrm{d}x}{\frac{1}{4} + \frac{12}{4} \int\limits_{[0,1]} h(x)^2 \,\mathrm{d}x} \ge \frac{1}{16}.$$

For  $k \in \{4, 5\}$ , we employ the following estimate, which holds as  $|h(x)| \leq 1/2$ :

$$\int_{[0,1]} h(x)^4 \, \mathrm{d}x \le \frac{1}{4} \int_{[0,1]} h(x)^2 \, \mathrm{d}x \tag{5}$$

If k = 4, we combine (3), (4) and (5) as follows:

$$\frac{m(T_4, W)}{m(S_4, W)} \ge \frac{\frac{1}{256} + \frac{12}{64} \int\limits_{[0,1]} h(x)^2 \,\mathrm{d}x}{\int\limits_{[0,1]} \left(\frac{1}{2} + h(x)\right)^4 + \left(\frac{1}{2} - h(x)\right)^4 \,\mathrm{d}x} \ge \frac{\frac{1}{256} + \frac{48}{256} \int\limits_{[0,1]} h(x)^2 \,\mathrm{d}x}{\frac{1}{8} + \frac{24+4}{8} \int\limits_{[0,1]} h(x)^2 \,\mathrm{d}x} \ge \frac{1}{32}$$

Finally, if k = 5, we obtain using (3), (4) and (5) the following:

$$\frac{m(T_5, W)}{m(S_5, W)} \ge \frac{\frac{1}{1024} + \frac{15}{256} \int\limits_{[0,1]} h(x)^2 \,\mathrm{d}x}{\int\limits_{[0,1]} \left(\frac{1}{2} + h(x)\right)^5 + \left(\frac{1}{2} - h(x)\right)^5 \,\mathrm{d}x} \ge \frac{\frac{1}{1024} + \frac{60}{1024} \int\limits_{[0,1]} h(x)^2 \,\mathrm{d}x}{\frac{1}{16} + \frac{40+20}{16} \int\limits_{[0,1]} h(x)^2 \,\mathrm{d}x} = \frac{1}{64}.$$

The proof of the lemma is now completed.

We are now ready to prove our main result, which is implied by the following more general theorem.

**Theorem 3.** Let G be an n-vertex graph with n-1 edges that has the Sidorenko property. The k-apex  $G^{+k}$  of G for  $k \in \{1, 2, 3, 4, 5\}$  is common.

*Proof.* Fix an *n*-vertex graph G with n-1 edges that has the Sidorenko property, an integer  $k \in \{1, \ldots, 5\}$ , and a graphon W. We need to show that

$$m(G^{+k}, W) \ge 2^{1-e(G^{+k})} = 2^{1-k \cdot v(G) - e(G)} = 2^{2-(k+1)n}.$$
 (6)

Since the graph G has the Sidorenko property, we derive from (1) that

$$m(G^{+k}, W) \ge \int_{B_k} d_k^W(z)^n \delta_k^W(z)^{n-1} \,\mathrm{d}z.$$
 (7)

Hölder's Inequality imply that

$$\int_{B_k} d_k^W(z)^2 \delta_k^W(z) \, \mathrm{d}z \le \left( \int_{B_k} d_k^W(z) \, \mathrm{d}z \right)^{\frac{n-2}{n-1}} \left( \int_{B_k} d_k^W(z)^n \delta_k^W(z)^{n-1} \right)^{\frac{1}{n-1}}.$$
(8)

We next combine (7) and (8) to obtain that

$$m(G^{+k}, W) \ge \frac{\left(\int\limits_{B_k} d_k^W(z)^2 \delta_k^W(z) \, \mathrm{d}z\right)^{n-1}}{\left(\int\limits_{B_k} d_k^W(z) \, \mathrm{d}z\right)^{n-2}} = \frac{m(T_k, W)^{n-1}}{m(S_k, W)^{n-2}}$$
(9)

Since the graph  $T_k$  is common, it holds that  $m(T_k, W) \ge 2^{-2k}$ . We now use Lemma 2 and derive from (9) that

$$m(G^{+k}, W) \ge m(T_k, W) \left(\frac{m(T_k, W)}{m(S_k, W)}\right)^{n-2} \ge 2^{-2k} \cdot 2^{-(n-2)(k+1)} = 2^{2-(k+1)n},$$

which is the estimate (6) required to be established.

#### 4 Concluding remarks

We would like to conclude with two brief remarks on our proof. We find it interesting that Theorem 1 follows from a more general Theorem 3, which does not require the graph G to be connected. We speculate that the assumption on connectivity of a graph G in Conjecture 1 can be replaced with a lower bound on the number of edges of G. Theorem 3 also implies for  $k \in \{1, 2, 3, 4, 5\}$  that the k-apex of any Sidorenko graph is an induced subgraph of a common graph. In particular, the 1-apex of any Sidorenko graph with a suitable number pendant vertices adjacent to the apex vertex is common, however, when a sufficiently large number of such pendant vertices is added, the graph is not common by the result of Fox [9] mentioned in Introduction.



Figure 2: The graphon  $W_{\varepsilon}$  for  $\varepsilon = 1/7$ . The origin of the coordinate system is in the top left corner, the black color represents the value 1, and the white color represents the value 0.

One can wonder whether the statement of Lemma 2 can hold for any k (note that if the statement of the lemma holds for k, then the statement of Theorem 3 holds for the same value of k). Unfortunately, this is not true as shown by the following example. For  $\varepsilon \in [0, 1)$  consider the graphon  $W_{\varepsilon}$  constructed as follows. Split the interval [0, 1] into an interval A of length  $\varepsilon$  and three intervals B, C and D, each of length  $\frac{1-\varepsilon}{3}$ . The graphon  $W_{\varepsilon}$  is equal to one on the sets  $(A \cup B) \setminus A^2$ ,  $(A \cup C) \setminus A^2$  and  $(A \cup D) \setminus A^2$ , and equal to zero elsewhere. The construction is illustrated in Figure 2.

A direct computation yields the following:

$$t(T_k, W_{\varepsilon}) = 2\varepsilon(1-\varepsilon)\left(\frac{1-\varepsilon}{3}\right)^k + 3\left(\frac{1-\varepsilon}{3}\right)^2\left(\frac{1+2\varepsilon}{3}\right)^k$$
$$t(T_k, 1-W_{\varepsilon}) = \varepsilon^{k+2} + 6\left(\frac{1-\varepsilon}{3}\right)^{k+2}$$
$$t(S_k, W_{\varepsilon}) = \varepsilon(1-\varepsilon)^k + (1-\varepsilon)\left(\frac{1+2\varepsilon}{3}\right)^k$$
$$t(S_k, 1-W_{\varepsilon}) = \varepsilon^{k+1} + (1-\varepsilon)\left(\frac{2-2\varepsilon}{3}\right)^k$$

Observe that if  $\varepsilon < 2/5$ , then

$$\lim_{k \to \infty} \frac{1}{k} \log \frac{m(T_k, W_{\varepsilon})}{m(S_k, W_{\varepsilon})} = \log \frac{\max\left\{\frac{1-\varepsilon}{3}, \frac{1+2\varepsilon}{3}, \varepsilon, \frac{1-\varepsilon}{3}\right\}}{\max\left\{1-\varepsilon, \frac{1+2\varepsilon}{3}, \varepsilon, \frac{2-2\varepsilon}{3}\right\}} = \log \frac{\frac{1+2\varepsilon}{3}}{1-\varepsilon} = \log \frac{1+2\varepsilon}{3(1-\varepsilon)}.$$

In particular, the limit is equal to  $\log \frac{10}{21}$  for  $\varepsilon = 1/8$ , however, if Lemma 2 were true for any value of k, this limit would be at least  $\log \frac{1}{2}$ . In fact, this example shows that the statement Lemma 2 fails already for k = 12 as  $m(T_{12}, W_{1/20}) \approx 2.4849 \cdot 10^{-6}$ ,  $m(S_{12}, W_{1/20}) \approx 0.030980$  and so

$$\frac{m(T_{12}, W_{1/20})}{m(S_{12}, W_{1/20})} \approx 0.00008021$$

while  $2^{-13} \approx 0.0001221$ .

### Acknowledgement

The authors would like to thank Andrzej Grzesik and Jan Volec for fruitful discussions concerning Ramsey multiplicity of graphs including the results presented in this paper.

### References

- [1] G. R. Blakley and P. Roy: A Hölder type inequality for symmetric matrices with nonnegative entries, Proc. Amer. Math. Soc. 16 (1965), 1244–1245.
- S. A. Burr and V. Rosta: On the Ramsey multiplicities of graphs—problems and recent results, J. Graph Theory 4 (1980), 347–361.
- [3] D. Conlon, J. Fox and B. Sudakov: An approximate version of Sidorenko's conjecture, Geom. Funct. Anal. 20 (2010), 1354–1366.
- [4] D. Conlon, J. H. Kim, C. Lee and J. Lee: Some advances on Sidorenko's conjecture, J. Lond. Math. Soc. 98 (2018), 593–608.
- [5] D. Conlon and J. Lee: Finite reflection groups and graph norms, Adv. Math. 315 (2017), 130–165.
- [6] D. Conlon and J. Lee: *Sidorenko's conjecture for blow-ups*, Discrete Anal. (2021), paper no. 2, 13pp.
- [7] P. Erdős: On the number of complete subgraphs contained in certain graphs, Magyar Tud. Akad. Mat. Kutató Int. Közl. 7 (1962), 459–464.
- [8] P. Erdős and M. Simonovits: Supersaturated graphs and hypergraphs, Combinatorica 3 (1983), 181–192.
- [9] J. Fox: There exist graphs with super-exponential Ramsey multiplicity constant, J. Graph Theory 57 (2008), 89–98.
- [10] A. W. Goodman: On sets of acquaintances and strangers at any party, Amer. Math. Monthly 66 (1959), 778–783.
- [11] A. Grzesik, J. Lee, B. Lidický and J. Volec: On tripartite common graphs, Combin. Probab. Comput. (2022), 907–923.
- [12] H. Hatami, J. Hladký, D. Kráľ, S. Norine and A. Razborov: Non-three-colourable common graphs exist, Combin. Probab. Comput. 21 (2012), 734–742.
- [13] C. Jagger, P. Stovíček and A. Thomason: *Multiplicities of subgraphs*, Combinatorica 16 (1996), 123–141.

- [14] S. Ko and J. Lee: Common graphs with arbitrary connectivity and chromatic number, J. Combin. Theory Ser. B 162 (2023), 223–230.
- [15] D. Král', J. Volec and F. Wei: Common graphs with arbitrary chromatic number, prepint arXiv:2206.05800.
- [16] D. Král', J. A. Noel, S. Norin, J. Volec and F. Wei: Non-bipartite k-common graphs, Combinatorica 42 (2022), 87–114.
- [17] L. Lovász: Large networks and graph limits, AMS Colloquium Publications, volume 60, 2012.
- [18] F. P. Ramsey: On a problem of formal logic, Proceedings of the London Mathematical Society 2 (1930), 264–286.
- [19] A. A. Razborov: Flag algebras, J. Symbolic Logic **72** (2007), 1239–1282.
- [20] A. Sidorenko: A correlation inequality for bipartite graphs, Graphs Combin. 9 (1993), 201–204.
- [21] A. Sidorenko: Randomness friendly graphs, Random Structures Algorithms 8 (1996), 229–241.
- [22] A. F. Sidorenko: Extremal problems in graph theory and functional-analytic inequalities, Proceedings of the All-Union seminar on discrete mathematics and its applications (Russian) (1986), 99–105.
- [23] A. F. Sidorenko: Cycles in graphs and functional inequalities, Mat. Zametki 46 (1989), 72–79, 104.
- [24] A. F. Sidorenko: Inequalities for functionals generated by bipartite graphs, Diskret. Mat. 3 (1991), 50–65.
- [25] A. Thomason: A disproof of a conjecture of Erdős in Ramsey theory, J. London Math. Soc. (2) 39 (1989), 246–255.