

ADDENDUM TO THE PAPER “REFINED CRITERIA TOWARD BOUNDEDNESS IN AN ATTRACTION-REPULSION CHEMOTAXIS SYSTEM WITH NONLINEAR PRODUCTIONS”

SILVIA FRASSU* AND GIUSEPPE VIGLIALORO

Dipartimento di Matematica e Informatica
Università degli Studi di Cagliari
Via Ospedale 72, 09124. Cagliari (Italy)

ABSTRACT. These notes aim to provide a deeper insight on the specifics of the paper “Refined criteria toward boundedness in an attraction-repulsion chemotaxis system with nonlinear productions” by A. Columbu, S. Frassu and G. Viglialoro [*Appl. Anal.* 2024, 103:2, 415–431].

1. AIM OF THE PAPER

In this report we focus on [1, Theorem 2.2] where an attraction-repulsion chemotaxis model is formulated as follow:

$$(1) \quad \begin{cases} u_t = \Delta u - \chi \nabla \cdot (u \nabla v) + \xi \nabla \cdot (u \nabla w) & \text{in } \Omega \times (0, T_{max}), \\ v_t = \Delta v - \beta v + f(u) & \text{in } \Omega \times (0, T_{max}), \\ w_t = \Delta w - \delta w + g(u) & \text{in } \Omega \times (0, T_{max}), \\ u_\nu = v_\nu = w_\nu = 0 & \text{on } \partial\Omega \times (0, T_{max}), \\ u(x, 0) = u_0(x), v(x, 0) = v_0(x), w(x, 0) = w_0(x) & x \in \bar{\Omega}. \end{cases}$$

Herein, Ω of \mathbb{R}^n , with $n \geq 2$, is a bounded and smooth domain, $\chi, \xi, \beta, \delta > 0$, and $f = f(s)$ and $g = g(s)$ sufficiently regular functions in their argument $s \geq 0$, essentially behaving as s^k and s^l for some $k, l > 0$. Moreover, further regular initial data $u_0(x), v_0(x), w_0(x) \geq 0$ are fixed, u_ν (and similarly v_ν and w_ν) indicates the outward normal derivative of u on $\partial\Omega$, whereas T_{max} identifies the maximum time up to which solutions to the system can be extended.

Once these hypotheses are fixed

$$(2) \quad \begin{cases} f, g \in C^1([0, \infty)) \quad \text{with} \quad 0 \leq f(s) \leq \alpha s^k \text{ and } \gamma_0(1+s)^l \leq g(s) \leq \gamma_1(1+s)^l, & \text{for some } \alpha, k, l > 0, \gamma_1 \geq \gamma_0 > 0, \\ (u_0, v_0, w_0) \in (W^{1,\infty}(\Omega))^3, \text{ with } u_0, v_0, w_0 \geq 0 \text{ on } \bar{\Omega}, \end{cases}$$

[1, Theorem 2.2] establishes that problem (1) admits a unique global and uniformly bounded classical solution (i.e., $T_{max} = \infty$) and there exists $C > 0$ such that $\|u(\cdot, t)\|_{L^\infty(\Omega)} \leq C$ for all $t \in (0, \infty)$ whenever

- (i) $l, k \in (0, \frac{1}{n}]$;
- (ii) $l \in (\frac{1}{n}, \frac{1}{n} + \frac{2}{n^2+4})$ and $k \in (0, \frac{1}{n}]$, or $k \in (\frac{1}{n}, \frac{1}{n} + \frac{2}{n^2+4})$ and $l \in (0, \frac{1}{n}]$;
- (iii) $l, k \in (\frac{1}{n}, \frac{1}{n} + \frac{2}{n^2+4})$.

From the one hand, we mention that the above conditions have been improved in the recent paper [2]; in the specific, [2, Theorem 2.2] ensures boundedness under the more relaxed assumption $k, l \in (0, \frac{2}{n})$.

As to our contribution, we aim at providing a further scenario toward boundedness involving also coefficients connected to g in (2); essentially we will show that

solutions to model (1) are uniformly bounded in time whenever $k < l$ and under a largeness assumption on γ_0 .

Remark 1 (On the origins and the meaning of model (1)). *The interested reader can find motivations connected to biological phenomena described by system (1) exactly in [1], and references therein mentioned. Also known results in close contexts are collected.*

2020 *Mathematics Subject Classification*. Primary: 35A01, 35K55, 35Q92. Secondary: 92C17.

Key words and phrases. Chemotaxis, Global existence, Boundedness, Nonlinear production.

*Corresponding author: silvia.frassu@unica.it.

2. PRESENTATION OF THE MAIN THEOREM

As an essential tool in order to mathematically formulate our main theorem, we have first to recall the following consequence of Maximal Sobolev regularity results ([4] or [3, Theorem 2.3]):

Proposition 1. *For $n \in \mathbb{N}$, let $\Omega \subset \mathbb{R}^n$ be a bounded domain with smooth boundary, $\rho > 0$ and $q > \max\{1, \frac{1}{\rho}\}$. Then there is $C_\rho = C_\rho(\Omega, n, q) > 0$ such that the following holds: Whenever $T \in (0, \infty]$, $I = [0, T)$, $h \in L^q(I; L^q(\Omega))$ and $\psi_0 \in W_{\frac{\rho}{2}}^{2,q}(\Omega) = \{\psi_0 \in W^{2,q}(\Omega) : \partial_\nu \psi_0 = 0 \text{ on } \partial\Omega\}$, every solution $\psi \in W_{loc}^{1,q}(I; L^q(\Omega)) \cap L_{loc}^q(I; W^{2,q}(\Omega))$ of*

$$\psi_t = \Delta \psi - \rho \psi + h \quad \text{in } \Omega \times (0, T); \quad \partial_\nu \psi = 0 \quad \text{on } \partial\Omega \times (0, T); \quad \psi(\cdot, 0) = \psi_0 \quad \text{on } \Omega$$

satisfies

$$\int_0^t e^s \int_\Omega \left(|\psi(\cdot, s)|^q + |\psi_t(\cdot, s)|^q + \left| \frac{\psi(\cdot, s)}{q} \right|^q + |\Delta \psi(\cdot, s)|^q \right) ds \leq 2^{q-1} C_\rho^q \left[\|\psi_0\|_{q, 1-\frac{1}{q}}^q + \int_0^t e^s \int_\Omega |h(\cdot, s)|^q ds \right] \quad \text{for all } t \in (0, T).$$

Proof. The proof is based on the classical result in [8]; for an appropriate adaptation to our case see details, for instance, in [5]. \square

Remark 2 (On the constant C_ρ and the norm $\|\psi_0\|_{q, 1-\frac{1}{q}}$ in Proposition 1). *The key role of Proposition 1 is the existence of the constant C_ρ , which remains defined once n, Ω and q are set. In particular (see [8, Theorem 2.5]), C_ρ does not depend on the initial configuration ψ_0 and the source h .*

As to $\|\psi_0\|_{q, 1-\frac{1}{q}}$, it represents the norm of ψ_0 in the interpolation space $(L^q(\Omega), W_{\frac{\rho}{2}}^{2,q}(\Omega))_{1-\frac{1}{q}, q}$. (See, for instance, [7, §1].)

Exactly in view of what said, we can now give the claim of our

Theorem 2.1. *For $n \in \mathbb{N}$, let Ω be a bounded domain of \mathbb{R}^n with smooth boundary, $0 < k < l$, $\delta, \alpha, \beta > 0$ and*

$$(3) \quad \bar{p} = \max \left\{ \frac{n}{2}, k \left(\frac{1}{\beta} - 1 \right), l \left(\frac{1}{\delta} - 1 \right) \right\} + 1.$$

Additionally, let us set

$$(4) \quad \mathcal{A} = 2^{-\frac{l(\bar{p}+l-1)+\bar{p}}{\bar{p}+l}} \left(\frac{\bar{p}+l}{\bar{p}+2l+\delta(\bar{p}+l)} \right).$$

Then there exists $\mathcal{C} = \mathcal{C}(n, \Omega, l, k, \delta, \beta) > 0$ such that if $\mathcal{C} < \mathcal{A}$, it is possible to find $\gamma_1, \gamma_0 > 0$ fulfilling

$$(5) \quad \gamma_1 \geq \gamma_0 > \mathcal{A}^{-1} \mathcal{C} \gamma_1$$

and with this property: Whenever f, g, u_0, v_0, w_0 are taken as in (2), problem (1) admits a global and uniformly bounded solution $(u, v, w) \in (C^0(\bar{\Omega} \times [0, \infty)) \cap C^{2,1}(\bar{\Omega} \times (0, \infty)))^3$.

3. EXISTENCE OF LOCAL-IN-TIME SOLUTIONS AND A PRINCIPLE FOR BOUNDEDNESS

The arguments concerning the forthcoming local existence issue and the boundedness criterion are standard; details are achievable in [9] and [10, Appendix A.].

3.1. Local existence statement. Once $\chi, \xi, \beta, \delta > 0$ and f, g, u_0, v_0 are fixed as in (2), from here henceforth, with (u, v, w) we will refer to the classical and nonnegative solution to problem (1); u, v, w are defined for all $(x, t) \in \bar{\Omega} \times [0, T_{max})$, for some finite T_{max} .

3.2. Boundedness criterion. As explained in the next lines, if we establish that $u \in L^\infty((0, T_{max}); L^p(\Omega))$, for some $p > \frac{n}{2}$, we can exploit the boundedness criterion below and directly obtain that, indeed, $u \in L^\infty((0, \infty); L^\infty(\Omega))$; as an immediate consequence of that, well-known parabolic regularity results applied to the equations of v and w entail that also v, w belong to $L^\infty((0, \infty); L^\infty(\Omega))$.

Definitely, globality and boundedness of (u, v, w) , in the sense that

$$u, v, w \in C^0(\bar{\Omega} \times [0, \infty)) \cap C^{2,1}(\bar{\Omega} \times (0, \infty)) \cap L^\infty((0, \infty); L^\infty(\Omega))$$

are achieved whenever this *boundedness criterion* applies:

$$(6) \quad \text{If } \exists L > 0, p > \frac{n}{2} \mid \int_\Omega u^p \leq L \text{ on } (0, T_{max}) \Rightarrow (u, v, w) \in (L^\infty((0, \infty); L^\infty(\Omega)))^3.$$

Subsequently, Theorem 2.1 is established once (6) is derived.

4. A PRIORI BOUNDS; PROOF OF THE MAIN RESULT

From now on we will tacitly assume that all the appearing constants below c_i , $i = 1, 2, \dots$, are positive.

4.1. Some preparatory tools. Let us start with this necessary result:

Lemma 4.1. *Let $A, B \geq 0$ and $p \geq 1$. Then we have*

$$(7) \quad (A + B)^p \leq 2^{p-1}(A^p + B^p).$$

Proof. The proof is available [6, Theorem 1]. \square

4.2. Achieving the boundedness criterion. We have this sequence of results, valid for any constant $\Xi > 0$, which will be properly chosen later on, in the proof of our theorem.

The following lemma is valid for a general class of proper functions. Despite that, we contextualize it to the local solution (u, v, w) to problem (1).

Lemma 4.2. *For any $p > 1$ and all $t \in (0, T_{max})$, we have*

$$(p-1)\xi \int_{\Omega} u^p |w_t| \leq (p-1)\xi \Xi \int_{\Omega} \left| w_t + \frac{l}{p+l} w \right|^{\frac{p+l}{p}} + \left[\xi p \frac{p-1}{p+l} \left(\Xi \frac{p+l}{l} \right)^{-\frac{l}{p}} \left(1 + \frac{l}{p+l} \right) \right] \int_{\Omega} u^{p+l} + l \xi \Xi \frac{p-1}{p+l} \int_{\Omega} w^{\frac{p+l}{p}},$$

and

$$(p-1)\xi \delta \int_{\Omega} u^p w \leq (p-1)\xi \delta \Xi \int_{\Omega} w^{\frac{p+l}{p}} + \xi p \delta \frac{p-1}{p+l} \left(\Xi \frac{p+l}{l} \right)^{-\frac{l}{p}} \int_{\Omega} u^{p+l}.$$

Proof. Let, for commodity but also for reasons which will be clearer later, $q = \frac{p+l}{l}$. From the evident relation $|w_t| \leq |w_t + \frac{1}{q} w| + |\frac{1}{q} w|$, we obtain

$$(p-1)\xi \int_{\Omega} u^p |w_t| \leq (p-1)\xi \int_{\Omega} u^p |w_t + \frac{w}{q}| + \xi \frac{p-1}{q} \int_{\Omega} u^p w \quad \text{on } (0, T_{max}),$$

so that thanks to the Young inequality for all $t \in (0, T_{max})$ it is seen

$$(p-1)\xi \int_{\Omega} u^p |w_t| \leq (p-1)\xi \Xi \int_{\Omega} \left| w_t + \frac{w}{q} \right|^q + \frac{(p-1)p\xi}{p+l} (\Xi q)^{-\frac{l}{p}} \int_{\Omega} u^{p+l} + \Xi \frac{p-1}{q} \xi \int_{\Omega} w^q + \xi p \frac{p-1}{q(p+l)} (\Xi q)^{-\frac{l}{p}} \int_{\Omega} u^{p+l},$$

and the first claim is established.

As to the other relation, it can be derived in the same flavor. \square

Lemma 4.3. *For any $p > \max \{1, l(\frac{1}{\delta} - 1)\}$ and $t \in (0, T_{max})$ it holds that*

$$(8) \quad (p-1)\xi \Xi \int_0^t e^s \left(\int_{\Omega} \left| w_t + \frac{l}{p+l} w \right|^{\frac{p+l}{p}} \right) ds \leq (p-1)\xi \Xi 2^{\frac{p}{l}} \mathcal{C}_{\delta}^{\frac{p+l}{l}} \times \left[\|w_0\|_{\frac{p+l}{l}, \frac{p}{p+l}}^{\frac{p+l}{l}} + \gamma_1 \frac{p+l}{l} 2^{p+l-1} \int_0^t e^s \left(\int_{\Omega} u^{p+l} ds \right) + \gamma_1 \frac{p+l}{l} 2^{p+l-1} |\Omega| \int_0^t e^s ds \right],$$

and

$$(9) \quad (p-1)\xi \Xi \left(\frac{l}{p+l} + \delta \right) \int_0^t e^s \left(\int_{\Omega} w^{\frac{p+l}{p}} \right) \leq (p-1)\xi \Xi 2^{\frac{p}{l}} \left(\frac{l}{p+l} + \delta \right) \mathcal{C}_{\delta}^{\frac{p+l}{l}} \times \left[\|w_0\|_{\frac{p+l}{l}, \frac{p}{p+l}}^{\frac{p+l}{l}} + \gamma_1 \frac{p+l}{l} 2^{p+l-1} \int_0^t e^s \left(\int_{\Omega} u^{p+l} ds \right) + \gamma_1 \frac{p+l}{l} 2^{p+l-1} |\Omega| \int_0^t e^s ds \right].$$

Proof. We can derive (8) (and similarly (9)) by invoking Proposition 1 with $\psi = w$, $h = g$ and $\rho = \delta$; indeed, for $q = \frac{p+l}{l}$ as before, it is $q > \max\{1, \frac{1}{\delta}\}$ so that

$$(p-1)\xi \Xi \int_0^t e^s \left(\int_{\Omega} \left| w_t + \frac{w}{q} \right|^q \right) ds \leq (p-1)\xi \Xi \mathcal{C}_{\delta}^q 2^{q-1} \left[\|w_0\|_{q, 1-\frac{1}{q}}^q + \int_0^t e^s \left(\int_{\Omega} g(u)^q \right) ds \right],$$

and the conclusion is attained by virtue of the upper bound (2) for g and (7), in the form $(u+1)^{p+l} \leq 2^{p+l-1}(u^{p+l} + 1)$ (naturally $p+l > 1$). \square

The next two results, indeed, provide properties of local solutions (u, v, w) to model (1) and are based on applications of Proposition 1.

Lemma 4.4. *For any $p > \max \{1, k(\frac{1}{\beta} - 1)\}$ and $t \in (0, T_{max})$ there is c_1 such that*

$$c_1 \int_0^t e^s \left(\int_{\Omega} |\Delta v|^{\frac{p+k}{k}} \right) ds \leq c_1 \mathcal{C}_{\beta}^{\frac{p+k}{k}} \left[\|v_0\|_{\frac{p+k}{k}, \frac{p}{p+k}}^{\frac{p+k}{k}} + \alpha \frac{p+k}{k} \int_0^t e^s \left(\int_{\Omega} u^{p+k} \right) ds \right].$$

Proof. The proof follows from analogous arguments used in Lemma 4.3; in this case, in particular, Proposition 1 is exploited with $q = \frac{p+k}{k} > \max\{1, \frac{1}{\beta}\}$, $\psi = v, h = f$ and $\rho = \beta$. \square

With the aim of ensuring $u \in L^\infty((0, T_{max}); L^p(\Omega))$ for some $p > \frac{n}{2}$, let us study the evolution in time of $t \mapsto \int_\Omega u^p$; this will be done by means of testing procedures.

Lemma 4.5. *For any $p > 1$ and all $t \in (0, T_{max})$ the following relation is satisfied:*

$$(10) \quad \begin{aligned} \frac{d}{dt} \int_\Omega u^p &\leq \int_\Omega u^{p+k} + c_1 \int_\Omega |\Delta v|^{\frac{p+k}{k}} + (p-1)\xi \Xi \int_\Omega |w_t + \frac{l}{p+l}w|^{\frac{p+l}{l}} \\ &+ (p-1)\xi \Xi \left(\delta + \frac{l}{p+l} \right) \int_\Omega w^{\frac{p+l}{l}} + (p-1)\xi \left[\frac{p}{p+l} \left(\frac{p+l}{l} \right)^{-\frac{l}{p}} \Xi^{-\frac{l}{p}} \left(1 + \delta + \frac{l}{p+l} \right) - \gamma_0 \right] \int_\Omega u^{p+l}. \end{aligned}$$

Proof. By testing the first equation of problem (1) with pu^{p-1} , using its boundary conditions and taking into account the second and the third equation, we have thanks to the Young inequality

$$\begin{aligned} \frac{d}{dt} \int_\Omega u^p &= p \int_\Omega u^{p-1} u_t = -p(p-1) \int_\Omega u^{p-2} |\nabla u|^2 - (p-1)\chi \int_\Omega u^p \Delta v + (p-1)\xi \int_\Omega u^p \Delta w \\ &\leq \int_\Omega u^{p+k} + c_1 \int_\Omega |\Delta v|^{\frac{p+k}{k}} + (p-1)\xi \int_\Omega u^p (w_t + \delta w - g(u)) \quad \text{on } (0, T_{max}). \end{aligned}$$

Now, we recall the properties of g given in (2) so to deduce, by using Young's inequality, again the relation $|w_t| \leq |w_t + \frac{l}{p+l}w| + |\frac{l}{p+l}w|$, and Lemma 4.2

$$\begin{aligned} \frac{d}{dt} \int_\Omega u^p &\leq \int_\Omega u^{p+k} + c_1 \int_\Omega |\Delta v|^{\frac{p+k}{k}} + (p-1)\xi \int_\Omega u^p |w_t| + \delta(p-1)\xi \int_\Omega u^p w - \xi \gamma_0 (p-1) \int_\Omega u^{p+l} \\ &\leq \int_\Omega u^{p+k} + c_1 \int_\Omega |\Delta v|^{\frac{p+k}{k}} + (p-1)\xi \Xi \int_\Omega |w_t + \frac{l}{p+l}w|^{\frac{p+l}{l}} \\ &+ \frac{l(p-1)\xi \Xi}{p+l} \int_\Omega w^{\frac{p+l}{l}} + \frac{p(p-1)\xi}{p+l} \left(\Xi \frac{p+l}{l} \right)^{-\frac{l}{p}} \left(1 + \frac{l}{p+l} \right) \int_\Omega u^{p+l} \\ &+ (p-1)\xi \delta \Xi \int_\Omega w^{\frac{p+l}{l}} + \frac{p(p-1)\xi \delta}{p+l} \left(\Xi \frac{p+l}{l} \right)^{-\frac{l}{p}} \int_\Omega u^{p+l} - \xi(p-1)\gamma_0 \int_\Omega u^{p+l} \quad \text{on } (0, T_{max}). \end{aligned}$$

The claim is achieved by collecting terms. \square

Lemma 4.6. *Let $k < l$. Then for every $p > \max\left\{\frac{n}{2}, k\left(\frac{1}{\beta} - 1\right), l\left(\frac{1}{\delta} - 1\right)\right\}$ we have that for all $t < T_{max}$*

$$\begin{aligned} e^t \int_\Omega u^p &\leq c_2 + c_3 \int_0^t e^s ds + \\ &(p-1)\xi \left[\left(1 + \delta + \frac{l}{p+l} \right) \left(\Xi C_\delta^{\frac{p+l}{l}} \gamma_1^{\frac{p+l}{l}} 2^{\frac{p}{l} + p + l - 1} + \frac{p}{p+l} \left(\frac{p+l}{l} \right)^{-\frac{l}{p}} \Xi^{-\frac{l}{p}} \right) + \varepsilon - \gamma_0 \right] \int_0^t e^s \left(\int_\Omega u^{p+l} \right) ds. \end{aligned}$$

Proof. Let us start with these estimates, fruit of the application of Young's inequality: for all $\varepsilon > 0$, $\hat{c} > 0$, $p > 1$ and $0 < k < l$ it holds that

$$(11) \quad \hat{c} \int_\Omega u^p \leq \frac{\varepsilon}{2} \int_\Omega u^{p+l} + c_4 \quad \text{for all } t \in (0, T_{max}),$$

and

$$(12) \quad \hat{c} \int_\Omega u^{p+k} \leq \frac{\varepsilon}{2} \int_\Omega u^{p+l} + c_5 \quad \text{for all } t \in (0, T_{max}).$$

By adding to both sides of relation (10) the term $\int_\Omega u^p$, estimate (11) leads to this inequality, valid on $(0, T_{max})$.

$$(13) \quad \begin{aligned} \frac{d}{dt} \int_\Omega u^p + \int_\Omega u^p &\leq \int_\Omega u^{p+k} + c_1 \int_\Omega |\Delta v|^{\frac{p+k}{k}} + (p-1)\xi \Xi \int_\Omega |w_t + \frac{l}{p+l}w|^{\frac{p+l}{l}} \\ &+ (p-1)\xi \Xi \left(\delta + \frac{l}{p+l} \right) \int_\Omega w^{\frac{p+l}{l}} + (p-1)\xi \left[\frac{p}{p+l} \left(\frac{p+l}{l} \right)^{-\frac{l}{p}} \Xi^{-\frac{l}{p}} \left(1 + \delta + \frac{l}{p+l} \right) + \frac{\varepsilon}{2} - \gamma_0 \right] \int_\Omega u^{p+l} + c_6. \end{aligned}$$

Successively, we multiply (13) by e^t and integrate on $(0, t)$. From the identity $\frac{d}{dt}(e^t \int_\Omega u^p) = e^t \frac{d}{dt} \int_\Omega u^p + e^t \int_\Omega u^p$, we get

$$\begin{aligned} e^t \int_\Omega u^p &\leq \int_\Omega u_0^p + \int_0^t e^s \left\{ \int_\Omega u^{p+k} + c_1 \int_\Omega |\Delta v|^{\frac{p+k}{k}} + (p-1)\xi \Xi \int_\Omega |w_t + \frac{l}{p+l}w|^{\frac{p+l}{l}} \right. \\ &\quad \left. + (p-1)\xi \Xi \left(\delta + \frac{l}{p+l} \right) \int_\Omega w^{\frac{p+l}{l}} + (p-1)\xi \left[\frac{p}{p+l} \left(\frac{p+l}{l} \right)^{-\frac{l}{p}} \Xi^{-\frac{l}{p}} \left(1 + \delta + \frac{l}{p+l} \right) + \frac{\varepsilon}{2} - \gamma_0 \right] \int_\Omega u^{p+l} + c_6 \right\} ds. \end{aligned}$$

The term involving $\int_0^t e^s (\int_\Omega |\Delta v|^{\frac{p+k}{k}}) ds$ can be essentially controlled by $\int_0^t e^s (\int_\Omega u^{p+k}) ds$, thanks to Lemma 4.4; additionally, $\int_\Omega u^{p+k}$ is treated through (12). These two operations provide

$$\begin{aligned} e^t \int_\Omega u^p &\leq \int_\Omega u_0^p + \int_0^t e^s \left\{ (p-1)\xi \Xi \int_\Omega |w_t + \frac{l}{p+l} w|^{\frac{p+l}{l}} + (p-1)\xi \Xi \left(\delta + \frac{l}{p+l} \right) \int_\Omega w^{\frac{p+l}{l}} \right. \\ &\quad \left. + (p-1)\xi \left[\frac{p}{p+l} \left(\frac{p+l}{l} \right)^{-\frac{1}{p}} \Xi^{-\frac{1}{p}} \left(1 + \delta + \frac{l}{p+l} \right) + \varepsilon - \gamma_0 \right] \int_\Omega u^{p+l} + c_7 \right\} ds. \end{aligned}$$

By invoking (8) and (9) we conclude by virtue of a reorganization of the involved terms. \square

4.3. Proof of Theorem 2.1. With the above preparations we are now in a position to establish what anticipated.

Proof. For $0 < k < l$, $\delta, \alpha, \beta > 0$, let \bar{p} be as in (3) and, additionally, for $C_\delta \left(\Omega, n, \frac{\bar{p}+l}{l} \right)$ being the constant provided by Proposition 1, when it is applied to the equation for w in model (1), let also set

$$\mathcal{C} = C_\delta \left(\Omega, n, \frac{\bar{p}+l}{l} \right).$$

Since by assumptions $\mathcal{C} < \mathcal{A}$, where \mathcal{A} is defined in (4), we can find $\gamma_1 \geq \gamma_0$ complying with (5); in these positions, let

$$\Xi = \frac{l}{\bar{p}+l} C_\delta^{-\frac{\bar{p}}{l}} \gamma_1^{-\frac{\bar{p}}{l}} 2^{-\frac{\bar{p}(\bar{p}+(\bar{p}+l-1)l)}{l(\bar{p}+l)}},$$

and let f, g, u_0, v_0 and w_0 obey (2). Some computations show that for proper small $\varepsilon > 0$

$$\gamma_0 > \mathcal{A}^{-1} \mathcal{C} \gamma_1 \Rightarrow \left(1 + \delta + \frac{l}{\bar{p}+l} \right) \left(\Xi C_\delta^{\frac{\bar{p}+l}{l}} \gamma_1^{\frac{\bar{p}+l}{l}} 2^{\frac{\bar{p}}{l} + \bar{p} + l - 1} + \frac{\bar{p}}{\bar{p}+l} \left(\frac{\bar{p}+l}{l} \right)^{-\frac{1}{\bar{p}}} \Xi^{-\frac{1}{\bar{p}}} \right) + \varepsilon - \gamma_0 \leq 0,$$

and henceforth hypothesis (5) allows to exploit Lemma 4.6 and obtain

$$e^t \int_\Omega u^{\bar{p}} \leq c_2 + c_3 \int_0^t e^s ds \quad \text{for all } t \in (0, T_{max}),$$

or also $\int_\Omega u^{\bar{p}} \leq L$ on $(0, T_{max})$. The claim follows from the extensibility criterion (6). \square

Acknowledgments. The authors are members of the Gruppo Nazionale per l'Analisi Matematica, la Probabilità e le loro Applicazioni (GNAMPA) of the Istituto Nazionale di Alta Matematica (INdAM). The authors are partially supported by the research projects *Analysis of PDEs in connection with real phenomena*, funded by Fondazione di Sardegna (2021, CUP F73C22001130007) and MIUR (Italian Ministry of Education, University and Research) Prin 2022 *Nonlinear differential problems with applications to real phenomena* (Grant Number: 2022ZXZTN2). SF also acknowledges financial support by INdAM-GNAMPA project *Problemi non lineari di tipo stazionario ed evolutivo* (CUP E53C23001670001).

REFERENCES

- [1] A. Columbu, S. Frassu, and G. Viglialoro. Refined criteria toward boundedness in an attraction-repulsion chemotaxis system with nonlinear productions. *Appl. Anal.*, 103(2):415–431, 2024.
- [2] A. Columbu, R. D. Fuentes, and S. Frassu. Uniform-in-time boundedness in a class of local and nonlocal nonlinear attraction-repulsion chemotaxis models with logistics. <https://arxiv.org/abs/2311.06526>, 2024.
- [3] Y. Giga and H. Sohr. Abstract L^p estimates for the Cauchy problem with applications to the Navier-Stokes equations in exterior domains. *J. Funct. Anal.*, 102(1):72–94, 1991.
- [4] M. Hieber and J. Prüss. Heat kernels and maximal L^p - L^q estimates for parabolic evolution equations. *Comm. Partial Differential Equations*, 22(9-10):1647–1669, 1997.
- [5] S. Ishida, J. Lankeit, and G. Viglialoro. A Keller–Segel type taxis model with ecological interpretation and boundedness due to gradient nonlinearities. *Discrete Continuous Dyn. Syst. Ser. B.*, doi:10.3934/dcdsb.2024029, 2024.
- [6] G. J. O. Jameson. Some inequalities for $(a+b)^p$ and $(a+b)^p + (a-b)^p$. *Math. Gaz.*, 98(541):96–103, 2014.
- [7] A. Lunardi. *Analytic Semigroups and Optimal Regularity in Parabolic Problems*. Modern Birkhäuser Classics. Springer Basel, 2012.
- [8] J. Prüss and R. Schnaubelt. Solvability and maximal regularity of parabolic evolution equations with coefficients continuous in time. *J. Math. Anal. Appl.*, 256(2):405–430, 2001.
- [9] Y. Tao and Z.-A. Wang. Competing effects of attraction vs. repulsion in chemotaxis. *Math. Models Methods Appl. Sci.*, 23(1):1–36, 2013.
- [10] Y. Tao and M. Winkler. Boundedness in a quasilinear parabolic-parabolic Keller–Segel system with subcritical sensitivity. *J. Differential Equations*, 252(1):692–715, 2012.