ASYMPTOTIC COMPUTATIONS OF TROPICAL REFINED INVARIANTS IN GENUS 0 AND 1

THOMAS BLOMME, GURVAN MÉVEL

ABSTRACT. Block and Göttsche introduced a Laurent polynomial multiplicity to count tropical curves. Itenberg and Mikhalkin then showed that this multiplicity leads to invariant counts called tropical refined invariants. Recently, Brugallé and Jaramillo-Puentes studied the polynomiality properties of the coefficients of these invariants and showed that for fixed genus g, the coefficients ultimately coincide with polynomials in the homology class of the curves we look at. We call the generating series of these polynomials asymptotic refined invariant. In genus 0, the asymptotic refined invariant has been computed by the second author in the h-transverse case.

In this paper, we give a new proof of the formula for the asymptotic refined invariant for g = 0 using variations on the floor diagram algorithm. This technique allows also to compute the asymptotic refined invariant for g = 1. The result exhibits surprising regularity properties related to the generating series of partition numbers and quasi-modular forms.

CONTENTS

ດ

T.	Introduction	2
	1.1. Context	2
	1.2. Overview of results	3
	1.3. Organization of the paper	5
	Acknowledgements	5
2.	Floor diagrams and asymptotic refined invariants	5
	2.1. Toric surfaces, homology classes and polygons.	5
	2.2. Floor diagrams and tropical refined invariants	7
	2.3. Asymptotic refined invariants	10
3.	Generating series in fixed degree	13
	3.1. Degree 0	14
	3.2. Degree 1	15
4.	Asymptotic refined invariant for genus 0	18
	4.1. The case of Hirzebruch surfaces	18
	4.2. The case of <i>h</i> -transverse toric surfaces	23
5.	Asymptotic refined invariant in genus 1	27
	5.1. The case of Hirzebruch surfaces	27
	5.2. The case of h -transverse toric surfaces.	36
References		37

²⁰²⁰ Mathematics Subject Classification. Primary 14T15, 14T90, 05A15; Secondary 14N10.

Key words and phrases. Tropical refined invariants, floor diagrams, generating series, asymptotic behavior.

THOMAS BLOMME, GURVAN MÉVEL

1. INTRODUCTION

1.1. Context.

1.1.1. Enumerative invariants and polynomiality. Given some points in the plane, the problem of determining how many curves of fixed degree and genus pass through these points is a well-known question, which generalizes to other surfaces. Given a non-singular surface X equipped with a sufficiently ample line bundle \mathcal{L} , curves on X may be obtained as zero-sets of sections of \mathcal{L} . For a non-negative integer g, how many such curves of genus g pass through $c_1(X) \cdot \mathcal{L} - 1 + g$ points on X? We denote by $N_g^X(\mathcal{L})$ this number, which is also known as the degree of the corresponding Severi variety. As a consequence of the adjunction formula, we could consider a dual question: given δ a number of nodes, what is the number $N_X^\delta(\mathcal{L})$ of δ -nodal curves passing through the appropriate number of points on X?

Although determining these numbers is a difficult problem, some recursive formulas have been proved in the 90's by Kontsevich [KM94] in the specific case of rational curves, and by Caporaso-Harris [CH98]. In the same decade, Göttsche conjectured in [Göt98] the number $N_X^{\delta}(\mathcal{L})$ to behave polynomially when X, δ are fixed and \mathcal{L} varies. This conjecture has first been proved by Tzeng [Tze12], then also by Kool, Shende and Thomas [KST11], and is as follows: for any δ , there exists a universal polynomial $P_{\delta} \in \mathbb{C}[x, y, z, t]$ such that for any non-singular complex algebraic surface with a sufficiently ample line bundle \mathcal{L} , one has

$$N_X^{\delta}(\mathcal{L}) = P_{\delta}(\mathcal{L}^2, c_1(X) \cdot \mathcal{L}, c_1(X)^2, c_2(X)).$$

Göttsche conjecture also states that the generating series of the $(P_{\delta})_{\delta}$ is multiplicative, in that there exist formal series A_1, \ldots, A_4 such that the generating series is $A_1^x A_2^y A_3^z A_4^t$, with explicit descriptions of some of the A_i .

This polynomial behavior is not satisfied when we fix the genus g. For instance, Di Francesco and Itzykson [DFI95] proved that $\log(N_0^{\mathbb{CP}^2}(d)) \sim 3d \log(d)$. However, Brugallé and Jaramillo-Puentes showed in [BJP22] that we recover polynomiality when looking instead at the coefficients of the *tropical refined invariant*.

1.1.2. Tropical refined invariants and their asymptotic. Tropical refined invariants emanate from Mikhalkin's correspondence theorem [Mik05] which enables the computation of $N_g^X(\mathcal{L})$ for toric surfaces, transforming the previous algebraic problem into a combinatorial enumerative problem dealing with objects called *tropical curves*. The correspondence theorem assigns some integer multiplicities to tropical curves. Block and Göttsche [BG16b] proposed to refine this multiplicity, yielding instead a Laurent polynomial (in a formal variable q) count of tropical curves, which interpolates between Gromov-Witten invariants for q = 1 and tropical Welshinger invariants for q = -1. Itenberg and Mikhalkin [IM13] proved that the enumeration using this new refined multiplicity indeed provides an invariant, known as tropical refined invariant and denoted by $BG_g^X(\mathcal{L})(q)$. In [BJP22], Brugallé and Jaramillo-Puentes showed, in the case of Hirzebruch surfaces

In [BJP22], Brugallé and Jaramillo-Puentes showed, in the case of Hirzebruch surfaces and (weighted) projective spaces, that for any fixed i the coefficient of codegree i of the tropical refined invariant is polynomial, providing that the line bundle is sufficiently ample with respect to i. Results of [BJP22] have been extended in genus 0 to any *h*-transverse toric surface, including singular ones, by the second author [Mév23]. In sight of the multiplicativity part of Göttsche conjecture, [Mév23] also provides explicit formula for the generating series of the polynomials that give the coefficients of fixed codegree. In this paper, we provide a new proof for the generating series of the polynomials in the genus 0 case, as well as formulas for the genus 1 case, that may cast a new mystery toward the nature of tropical refined invariants.

3

1.1.3. Interpretations and applications of tropical refined invariants. Due to their original combinatorial definition, the meaning of tropical refined invariants in classical geometry remained mysterious for quite some time. Up to now, two main interpretations have been proved.

- ▷ Mikhalkin showed [Mik17] that in some situations, genus 0 tropical refined invariants correspond to refined counts of real oriented curves according to the value of a so-called *quantum index*.
- ▷ Bousseau proved [Bou19] that through the change of variable $q = e^{i\hbar}$, tropical refined invariants actually compute generating series of (log-)Gromov Witten invariants with a λ -class.

Since then, results from [Mik17] have been generalized to genus 1 and 2 by Itenberg and Shustin [IS23], leading to real refined invariants. Unfortunately, the correspondence theorem does not relate them to tropical refined invariants defined using the Block-Göttsche multiplicity. Moreover, the tropical refined invariants involved in [Mik17] are of a different kind from the ones considered in the present paper, since its enumerative problem involves *boundary constraints*. Although results from [BJP22] and the present paper do not apply to the invariants from [Mik17], computed in [Blo19], it would be interesting to know if the asymptotic results are true in this boundary setting, which would yield asymptotic information on the real invariants.

Using [Bou19], the results from [BJP22, Mév23] may be interpreted as a subtle asymptotic statement about the (log-)GW invariants with a λ -class. The subtlety is due to the change of variable going from tropical refined invariants to these GW-invariants: $q = e^{i\hbar}$, i.e. $q^{m/2} - q^{-m/2} = 2i \sin\left(\frac{m\hbar}{2}\right)$, so that our results and those from [BJP22, Mév23] give an asymptotic on the Fourier coefficients of the generating series of the GW-invariants with a λ -class. Furthermore, given that $q = e^{i\hbar}$ is also the change of variable occurring when relating GW-invariants and some Donaldson-Thomas invariants, there is also a possibility that the asymptotic of tropical refined invariants is actually a shadow from a property of some DT-invariants.

Finally, in [GS14], the refinement of tropical invariants is conjectured to correspond to the refinement of the Euler characteristic by the Hirzebruch genus χ_{-y} for some relative Hilbert scheme. Some work in this direction has been accomplished in [NPS18]. If such a correspondence was true, the asymptotic results from [BJP22, Mév23] and the present paper may also be interpreted as asymptotic statements on the χ_{-y} -genus of the relative Hilbert schemes.

1.2. Overview of results. In this paper, we pursue the study of tropical refined invariants that was started in [BJP22] and [Mév23], expanding the range of tools and possible computations. We also state some conjectures.

For the surface X, let $AR_{g,i}^X(\beta)$ be the asymptotic polynomial from [BJP22] giving the codegree *i* coefficient of the genus *g* refined invariant obtained by counting curves in the class β . Our results consist in an explicit computation of the $AR_{g,i}^X(\beta)$ in few particular cases, by determining their generating series in *i* or *g*.

It is known from [IM13] that the leading coefficient of the genus g tropical refined invariant is $AR_{g,0}^X = \binom{g_{\max}}{g}$, where g_{\max} is the genus of a non-singular curve in the class β ; by the

adjunction formula one has $g_{\text{max}} = 1 + \frac{1}{2}(\beta^2 + \beta \cdot K_X)$. In other words, one has

$$\sum_{g \geqslant 0} A R_{g,0}^X u^g = (1+u)^{g_{\max}}$$

In this paper we provide a formula for the second term of the refined invariants.

Theorem (3.2). For X a non-singular toric surface associated to a h-transverse and horizontal polygon, one has:

$$\sum_{g \ge 0} AR_{g,1}^X u^g = (1+u)^{g_{\max}} \left[-\beta^2 \frac{u^3}{(1+u)^3} + 2(K_X \cdot \beta) \frac{u^2}{(1+u)^3} + \chi \frac{1}{1+u} - K_X^2 \frac{u}{(1+u)^3} \right].$$

We then give a result concerning generating series when summing over *i*: let $AR_g^X(\beta) = \sum_{i=0}^{\infty} AR_{g,i}^X(\beta)x^i$ be this series, which we call the genus *g* asymptotic refined invariant. In [Mév23], the second author proved that $AR_0^X(\beta) = p(x)^{\chi}$. However, the method used there does not seem to be manageable to deal with higher genus. With a slightly different point of view, we give in Theorem 4.17 a different proof of this fact. The interest is that this point of view also allows to perform the computation in genus 1. A priori, the polynomiality behavior for general *h*-transverse toric surface has not been proven in [BJP22], but in the genus 0 and 1 case it actually follows from our computations.

Theorem (5.18). For X a non-singular toric surface associated to a h-transverse and horizontal polygon and with Euler characteristic χ , the genus 1 asymptotic refined invariant satisfies

$$AR_{1}^{X}(\beta) = p(x)^{\chi} \left(g_{\max} - 12E_{2}(x) \right),$$

where $p(x) = \prod_{j=1}^{\infty} \frac{1}{1-x^j}$ is the generating series of the partition numbers and $E_2(x) = \sum_{a=1}^{\infty} \sigma_1(a) x^a$ is the first Eisenstein series.

To prove the result we use floor diagrams, defined in [BM07, BM08] and adapted in the refined setting in [BG16a]. We start with the case of Hirzebruch surfaces. It turns out most technicalities occur in the latter. It is then quite easy to obtain the result for other h-transverse toric surfaces by using the computation we did in the Hirzebruch case.

The form taken by generating series for fixed genus suggests a nice but subtle regularity of the asymptotic of the refined invariants. However, we are for now limited by the computational techniques at our disposal, as the complexity of the computations increases quite fast with the genus or the codegree. With some time, the techniques presented should be able to provide a result for genus 2 as well as codegree 2, but at the cost of many lengthy computations. Given the above results, we conjecture the following.

Conjecture 1.1. The asymptotic refined invariant AR_q^X has the following form:

$$AR_g^X(eta) = p(x)^\chi \left(egin{pmatrix} g_{\max} \ g \end{pmatrix} + Q_g^X(eta^2,eta\cdot K_X)
ight),$$

where Q_g^X is a polynomial of degree at most g in β^2 and $\beta \cdot K_X$, whose coefficients are quasimodular forms in the x variable vanishing at 0.

As g_{\max} is a polynomial function in β^2 and $\beta \cdot K_X$, so is $\binom{g_{\max}}{g}$. The constant term of the refined invariant has already been computed in [IM13] and is indeed $\binom{g_{\max}}{g}$, so that the conjecture is true modulo x.

The shape given in the conjecture emanates from computations in genus 0 and 1. To give more support to the quasi-modularity claim, we also have the following: in a future paper, we prove that the asymptotic refined invariants for Abelian surfaces is a polynomial in β^2 with coefficients in $\mathbb{Z}[\![x]\!]$ which are quasi-modular forms. This supports the mysterious appearance of quasi-modular forms in this setting. In the examples, the polynomial functions $AR_{g,i}^X(\beta)$ on the lattice $H_2(X,\mathbb{Z})$ seem to be more precisely polynomials in β^2 and $\beta \cdot K_X$, justifying the form of the polynomial Q_g^X . We notice a more general conjecture would deal with the double generating series $\sum_{i,g} AR_{g,i}^X u^g x^i$, and computations suggest to factor out $p(x)^{\chi}$ and $(1+u)^{g_{\text{max}}}$. Theorems 3.2 and 5.18 ensure that we have modulo u^2 :

$$\sum_{i,g} AR_{g,i}^{\chi} u^{g} x^{i} = p(x)^{\chi} (1+u)^{g_{\max}} \left[1 - (\chi + K_{\chi}^{2}) u E_{2}(x) \right] \mod u^{2}.$$

Conjecture 1.1 treats the regularity of the asymptotic invariant for a fixed surface X. Similarly to the Göttsche conjecture, it would be interesting to study the dependence of AR_g^X in the surface X. Hopefully, the polynomials $AR_{g,i}^X(\beta)$ are actually given by a universal polynomial $Q_{g,i}(\beta^2, \beta \cdot K_X, K_X^2, \chi(X))$. Theorems 3.2 and 5.18 prove that it is the case for g = 0, 1 and i = 0, 1.

1.3. Organization of the paper. The precise setting of tropical refined invariants is recalled in Section 2. We also recall how to compute them with floor diagrams. Furthermore, we give a change of variables that transforms the symmetric Laurent polynomials into true polynomials in a new variable x, so that the codegree i coefficient becomes the x^i term. This allows for an easier formulation of the asymptotics.

Section 3 is dedicated to the proof of Theorem 3.2, which amounts to compute the generating series in the genus parameter.

Section 4 introduces words that we will use in section 5. We also prove Theorem 4.17 using this tool to illustrate how it works. Section 5 is devoted to the proof of Theorem 5.18. We start by explaining how to construct floor diagrams of genus 1 from genus 0 floor diagrams. This allows us to express the genus 1 asymptotic refined invariant as a weighted sum over the genus 0 floor diagrams. In both Sections 4 and 5 we start with the Hirzebruch case before going to the case of h-transverse non-singular toric surfaces.

Acknowledgements. Part of the work was accomplished during the stays of T.B. in Nantes in March 2023 and of G.M. in Neuchâtel in October 2023. We both would like to thank the other for the excellent working conditions.

T.B. is partially supported by the SNSF grant 204125. G.M. is supported by the CNRS and conducted his work within the France 2030 framework programme, Centre Henri Lebesgue ANR-11-LABX-0020-01.

2. FLOOR DIAGRAMS AND ASYMPTOTIC REFINED INVARIANTS

In this section, we recall the definition of tropical refined invariants and their computation using floor diagrams. We also reformulate the main result of [BJP22] to introduce *asymptotic* refined invariants.

2.1. Toric surfaces, homology classes and polygons. Let N be a lattice and $M = \text{Hom}(N,\mathbb{Z})$ its dual. We denote by $M_{\mathbb{R}} = M \otimes \mathbb{R}$, $N_{\mathbb{R}} = N \otimes \mathbb{R}$ the associated real vector spaces. Following [Ful93], a compact toric surface X is obtained from a complete fan $\Sigma \subset N_{\mathbb{R}}$, or from a polygon $\Delta \in M_{\mathbb{R}}$ dual to Σ . The toric divisors of X are in bijection with the rays of Σ . We assume that X is smooth, i.e. every cone of Δ is simplicial. The anticanonical class

 $-K_X$ is represented by the sum of toric divisors. The Euler characteristic χ of X is the equal to the number of rays of Σ .

Each complex curve in X realizes some homology class $\beta \in H_2(X, \mathbb{Z})$. Recall that $H_2(X, \mathbb{Z})$ is endowed with the intersection product, which is non-degenerate by Poincaré duality. It is classical to show (see [Ful93]) that the homology group $H_2(X, \mathbb{Z})$ is generated by the classes of the toric divisors in X. In particular, a class $\beta \in H_2(X, \mathbb{Z})$ is fully determined by its intersection numbers with the toric divisors.

Consider the cone $\operatorname{Eff}(X) \subset H_2(X, \mathbb{Z})$ of *effective* classes, i.e. classes β such that $\beta \cdot D > 0$ for any toric divisor D. Let $\beta \in \operatorname{Eff}(X)$ be an effective class. For any ray ρ of the fan Σ , let $n_{\rho} \in N$ be a primitive vector such that $\rho = \mathbb{R}_{\geq 0} \cdot n_{\rho}$, and let D_{ρ} be the toric divisor corresponding to ρ . The multiset

$$\operatorname{trop}(\beta) = \{ n_{\rho}^{\beta \cdot D_{\rho}}, \ \rho \text{ ray of } \Sigma \},\$$

where the notation $n_{\rho}^{\beta \cdot D_{\rho}}$ means that n_{ρ} is taken $\beta \cdot D_{\rho}$ times, is called the *tropical degree* of β . The sum of the vectors of the tropical degree is 0 due to relations in $H_2(X,\beta)$ (see [Ful93]). Therefore, the tropical degree of β determines a convex lattice polygon $\Delta_{\beta} \subset M_{\mathbb{R}}$, with normal fan Σ , and such that the side dual to the ray ρ has integer length $\beta \cdot D_{\rho}$.

Remark 2.1. If $\beta \in \text{Eff}(X)$, the lattice polygon Δ_{β} gives an ample line bundle \mathcal{L}_{β} on X with Chern class $c_1(\mathcal{L}_{\beta}) \in H^2(X, \mathbb{Z})$ Poincaré dual to β . A basis of sections of \mathcal{L}_{β} is indexed by the lattice points of Δ_{β} . For our purpose, we work with the class β instead of the line bundle, as it would be the case in the setting of the Göttsche conjecture. These points of view are equivalent in the case of rational surfaces.

Example 2.2. Take $\Sigma_{\mathbb{P}^2}$ to be the complete fan in \mathbb{R}^2 with three rays spanned by respectively (0, -1), (-1, 0) and (1, 1), giving as toric surface the projective plane \mathbb{P}^2 . Its second homology group $H_2(\mathbb{P}^2, \mathbb{Z})$ is isomorphic to \mathbb{Z} , spanned by the common class L of any of the three toric divisors. The choice of $dL \in H_2(\mathbb{P}^2, \mathbb{Z}) \simeq \mathbb{Z}$ yields the tropical degree

$$\operatorname{trop}(dL) = \{(0, -1)^d, (-1, 0)^d, (1, 1)^d\}.$$

The associated polygon Δ_{dL} is d times the unit triangle, which thus has vertices (0,0), (d,0)and (0,d) (see Figure 1a), with associated line bundle $\mathcal{O}(d)$ on \mathbb{P}^2 .



FIGURE 1

Example 2.3. Take Σ_{δ} , with $\delta \ge 0$, to be the complete fan in \mathbb{R}^2 with rays four spanned by respectively (-1,0), (0,-1), (0,1) and $(1,-\delta)$. Let F, E_{∞} and $E = E_0$ be the classes of the divisors associated to the first three rays. They satisfy $E_0^2 = \delta$, $F^2 = 0$ and $E_0 \cdot F = 1$. The toric surface associated to this fan is the Hirzebruch surface \mathbb{F}_{δ} , which has $H_2(\mathbb{F}_{\delta},\mathbb{Z}) \simeq \mathbb{Z}^2$. We have the relation

$$E_{\infty} = E_0 - \delta F_{\gamma}$$

so that $H_2(\mathbb{F}_{\delta}, \mathbb{Z})$ is spanned by F and E. The toric divisor associated to the last ray lies also in the class F. Let $\beta = aE + bF \in \text{Eff}(\mathbb{F}_{\delta})$ be an effective class, which means that a, b > 0. The associated tropical data is

$$\operatorname{trop}(aE + bF) = \{(0, 1)^{b + \delta a}, (0, -1)^{b}, (-1, 0)^{a}, (1, -\delta)^{a}\}.$$

The corresponding polygon Δ_{aE+bF} is the trapezoid with vertices (0,0), (0,a), (b,0) and $(b + \delta a, a)$, see Figure 1b.

2.2. Floor diagrams and tropical refined invariants. In [IM13] it is shown that the count of genus g tropical curves of degree $\operatorname{trop}(\beta)$ passing through a generic configuration of $-K_X \cdot \beta + g - 1$ points with Block-Göttsche multiplicity does not depend on the choice of the points, yielding the tropical refined invariant $BG_g^X(\beta)(q) \in \mathbb{Z}[q^{\pm 1/2}]$. When X comes from an h-transverse polygon, see the definition below, it is possible to compute them using the floor diagram algorithm from [BG16a], which is the Block-Göttsche version of the floor diagram algorithm from [BM08]. This is the content of [BG16b, Theorem 4.3], stated below as Theorem 2.13 that may be taken as a definition of $BG_{g,\beta}^X(q)$. We now recall how floor diagrams work.

Definition 2.4. A convex lattice polygon Δ is said to be

- ▷ *h*-transverse if any edge of Δ has a direction vector of the form $(\pm 1, 0)$ or $(n, \pm 1)$ for some $n \in \mathbb{Z}$,
- ▷ *horizontal* if it has a top and bottom horizontal edge,
- \triangleright non-singular if the associated toric surface is non-singular.

Given Δ a lattice polygon we set the following notations.

- ▷ The number of interior lattice points of Δ is $g_{\max}(\Delta) = |\mathring{\Delta} \cap \mathbb{Z}^2|$.
- \triangleright The height of Δ is $a(\Delta)$.
- ▷ The length of its top (resp. bottom) edge is $b^{top}(\Delta)$ (resp. $b^{bot}(\Delta)$); these may be 0 if Δ is not horizontal.
- $\triangleright b_{\text{left}}(\Delta)$ (resp. $b_{\text{right}}(\Delta)$) is the multiset of integers k appearing a number of times equal to the integral length of the side of Δ having (-1, k) (resp. (1, k)) as outgoing normal vector.

When no ambiguity is possible we will simply use g_{max} , a, b^{top} , etc.

For a toric surface associated to a h-transverse, horizontal and non-singular polygon, we have the following result.

Lemma 2.5. Let X be a smooth toric surface coming from an h-transverse, with a top and bottom horizontal sides corresponding to divisors D_{top} and D_{bot} . We have

$$D_{\rm top}^2 + D_{\rm bot}^2 + \chi(X) = 4$$

where $\chi(X)$ is the Euler characteristic of X.

Proof. Using the Hodge numbers, the Euler characteristic of the sheaf \mathcal{O} of holomorphic functions satisfies $\chi(\mathcal{O}) = 1$. By Noether's formula, we know that $K_X^2 + \chi(X) = 12$. Moreover, as we have a top and bottom side, the toric surface is endowed with a map to \mathbb{P}^1 provided by the first coordinate in the lattice of monomial. Let F be the class of a fiber of this projection. The fiber over 0 (resp. ∞) is the union of toric divisors coming from the left (resp. right) sides of the polygon. As divisor, it is a linear combination of the corresponding toric divisors. In the general case, the coefficients are the horizontal coordinates of the lattice vectors directing

the corresponding rays in the fan. Since we are in the h-transverse case, the coefficients are 1 so that

$$F = \sum_{D \text{ left side of } \Delta} D = \sum_{D \text{ right side of } \Delta} D.$$

In particular, as the sum of all toric divisors is an anticanonical divisor, $D_{\text{bot}} + D_{\text{top}} + 2F$ is an anticanonical divisor of X. As $D_{\text{top}} \cdot D_{\text{bot}} = 0$, $F^2 = 0$, and $F \cdot D_{\text{top/bot}} = 1$ we get

$$12 - \chi(X) = K_X^2 = (D_{\text{bot}} + D_{\text{top}} + 2F)^2 = D_{\text{bot}}^2 + D_{\text{top}}^2 + 4 + 4.$$

An oriented graph Γ is a collection of vertices $V(\Gamma)$, bounded edges $E^0(\Gamma)$, sinks $E^{\text{top}}(\Gamma)$ and sources $E^{\text{bot}}(\Gamma)$. A bounded edge is a bivalent edge, i.e. adjacent to two vertices. A sink (resp. source) is a univalent edge oriented outward (resp. inward) a vertex. The set of all edges is denoted by $E(\Gamma)$. A weight on Γ is an application $w : E(\Gamma) \to \mathbb{Z}_{>0}$. Every vertex vhas a divergence which is the difference of the total weights entering and leaving the vertex, i.e.

$$\operatorname{div}(v) = \sum_{\stackrel{e}{\to} v} w(e) - \sum_{\stackrel{e}{v \to}} w(e).$$

Definition 2.6. Let Δ be a *h*-transverse polygon. A *floor diagram* \mathcal{D} with Newton polygon Δ is the data of (Γ, w, ℓ, r) , with (Γ, w) a weighted, connected, oriented and acyclic graph satisfying the following conditions :

- \triangleright the graph Γ has $a(\Delta)$ vertices called floors, $b^{\text{top}}(\Delta)$ sinks and $b^{\text{bot}}(\Delta)$ sources,
- \triangleright all sinks and sources have weight 1,
- ▷ the functions $L: V(\Gamma) \to b_{\text{left}}(\Delta)$ and $R: V(\Gamma) \to b_{\text{right}}(\Delta)$ are bijections such that for any vertex v one has $\operatorname{div}(v) = R(v) + L(v)$.

The genus of the floor diagram \mathcal{D} is the first Betti number of the underlying graph Γ . We will often confuse \mathcal{D} and Γ .

Remark 2.7. If $\Delta = \Delta_{\beta}$ for some $\beta \in H_2(X, \mathbb{Z})$, a floor diagram of Newton polygon Δ_{β} is also said to have class β .

Given a non-negative integer n, the quantum integer [n] is the Laurent polynomial in $q^{1/2}$ defined by

$$[n](q) = rac{q^{n/2} - q^{-n/2}}{q^{1/2} - q^{-1/2}} \in \mathbb{Z}_{\geqslant 0}[q^{\pm 1/2}].$$

Definition 2.8. Let \mathcal{D} be a floor diagram. Its refined Block-Göttsche multiplicity is

$$\mu_{BG}(\mathcal{D}) = \prod_{e \in E(\mathcal{D})} [w(e)]^2$$

It is a symmetric Laurent polynomial in q.

Definition 2.9. Let \mathcal{D} be a floor diagram of Newton polygon Δ and genus g. Its *degree* is the degree of the Laurent polynomial $(q^{1/2} - q^{-1/2})^{b^{\text{top}} + b^{\text{bot}} + 2|E^0(\mathcal{D})|} \mu_{BG}(\mathcal{D})$:

$$\deg(\mathcal{D}) = \frac{b^{\text{top}}}{2} + \frac{b^{\text{bot}}}{2} + \sum_{e \in E^0(\mathcal{D})} w(e),$$

and its *codegree* is the complement to the maximal degree of a genus g floor diagram with Newton polygon Δ , i.e. to the euclidean area of Δ_{β} :

$$\operatorname{codeg}(\mathcal{D}) = \operatorname{Area}(\Delta_{\beta}) - \operatorname{deg}(\mathcal{D}).$$

9

 \Diamond

Remark 2.10. From [BM08], we know that floor diagrams actually encode simple tropical curves, which are dual to convex subdivisions of the Newton polygon Δ consisting of triangles and parallelograms. The codegree is actually the area of the parallelograms appearing in the subdivision.

Example 2.11. Consider the polygon Δ of figure 2. It has

$$g_{\text{max}} = 5, \ a = 3, \ b^{\text{top}} = 1, \ b^{\text{bot}} = 3, \ b_{\text{left}} = \{-1, 1, 1\} \text{ and } b_{\text{right}} = \{0, 0, 1\}.$$

The floor diagrams \mathcal{D}_1 , \mathcal{D}_2 and \mathcal{D}_3 have Δ as Newton polygon. We always represent floor diagrams oriented from bottom to top, hence we do not precise orientation on the figures. Besides, we indicate the weight of an edge only if it is at least 2. The genera of \mathcal{D}_1 , \mathcal{D}_2 and \mathcal{D}_3 are respectively 0, 1 and 1, and their codegrees are 0, 2 and 4. Their refined multiplicities are

$$\begin{split} \mu_{BG}(\mathcal{D}_1) &= (q+1+q^{-1})^2 (q^{3/2}+q^{1/2}+q^{-1/2}+q^{-3/2})^2 \\ &= q^5 + 4q^4 + 10q^3 + 18q^2 + 25q + 28 + \dots \\ \mu_{BG}(\mathcal{D}_2) &= (q^{1/2}+q^{-1/2})^2 (q^{1/2}+q^{-1/2})^2 \\ &= q^2 + 4q + 6 + \dots \\ \mu_{BG}(\mathcal{D}_3) &= 1. \end{split}$$



FIGURE 2. Examples of floor diagrams.

Notice that the orientation on a floor diagram \mathcal{D} induces a partial order \prec on $E(\mathcal{D}) \cup V(\mathcal{D})$. We can thus define increasing functions on $E(\mathcal{D}) \cup V(\mathcal{D})$ and the following definition makes sense.

Definition 2.12. A marking \mathfrak{m} of a floor diagram \mathcal{D} is an increasing bijection $E(\mathcal{D}) \cup V(\mathcal{D}) \rightarrow \{1, \ldots, |E(\mathcal{D}) \cup V(\mathcal{D})|\}$. The pair $(\mathcal{D}, \mathfrak{m})$ is a marked floor diagram. Two marked floor diagrams $(\mathcal{D}, \mathfrak{m})$ and $(\mathcal{D}', \mathfrak{m}')$ are isomorphic if there exists an isomorphism $\varphi : \mathcal{D} \to \mathcal{D}'$ of weighted graphs such that $L = L' \circ \varphi$, $R = R' \circ \varphi$ and $\mathfrak{m} = \mathfrak{m}' \circ \varphi$.

The following theorem can be taken as a definition of the tropical refined invariants.

Theorem 2.13 ([BG16a, Theorem 4.3]). Let X be a h-transverse toric surface, $\beta \in H_2(X, \mathbb{Z})$, and $g \in \mathbb{Z}_{\geq 0}$. The tropical refined invariant is given by

$$BG_g^X(eta)(q) = \sum_{(\mathfrak{D},\mathfrak{m})} \mu_{BG}(\mathfrak{D})(q) \in \mathbb{Z}[q^{\pm 1}]$$

where the sums runs over the isomorphism classes of marked floor diagrams with Newton polygon Δ_{β} and genus g.

The tropical refined invariant $BG_g^X(\beta)$ is a symmetric Laurent polynomial in q with integer coefficients. Its degree is $|\mathring{\Delta}_{\beta} \cap \mathbb{Z}^2| - g = g_{\max}(\Delta_{\beta}) - g$.

Lemma 2.14. Let $g \ge 0$ and M, i > 0. Let Δ be a h-transverse polygon. Assume

$$b^{\mathrm{top}}(\Delta), b^{\mathrm{bot}}(\Delta) > M(g+1) + i.$$

Let \mathcal{D} be a floor diagram of genus g and $\operatorname{codeg}(\mathcal{D}) \leq i$. Then for any consecutive floors v_m and v_{m+1} , there is a bounded edge e between them with weight w(e) > M.

Proof. Let ω_m be the weight of the unique edge between the floors m and m+1 in the unique floor diagram of genus 0 and codegree 0 with Newton polygon Δ . It is equal to the integral length of the slice $\Delta \cap (\mathbb{Z} \times \{m\})$ of Δ at height m. As Δ is convex, it is always bigger than $\min(b^{\text{top}}, b^{\text{bot}})$.

Let \mathcal{D} be a genus g marked diagram, and let v_1, \ldots, v_a be the floors of \mathcal{D} , ordered by their marking but not necessarily inside \mathcal{D} . Let $\tilde{\omega}_m$ be the sum of the weights of the bounded edges of \mathcal{D} that link two floors v_m and v_{m+1} . It may be 0 if there are no such edges. In comparison to a diagram of genus g and codegree 0, the codegree of \mathcal{D} comes from two different phenomena:

- ▷ an edge with weight w that skips k floors including v_m or v_{m+1} contributes kw to the codegree, and decreases by w the maximal value of weight $\tilde{\omega}_m$ between v_m and v_{m+1} ,
- ▷ two floors $v_k \prec v_{k+1}$ having $R(v_k) > R(v_{k+1})$ or $L(v_k) > L(v_{k+1})$ contribute at least one to the codegree, and decrease the total weight of the bounded edges between v_k and v_{k+1} by at least 1.

Hence there are at most codeg(D) such phenomena, and therefore

$$\widetilde{\omega}_m \ge \omega_m - \operatorname{codeg}(\mathcal{D}) \ge \omega_m - i > (g+1)M.$$

In particular, there is at least one edge between the floors v_m and v_{m+1} , so that the floors are totally ordered in the diagram. Because \mathcal{D} has genus g, the total weight $\tilde{\omega}_m$ is split into at most g+1 edges. Hence we have at least one of them with w(e) > M.

2.3. Asymptotic refined invariants. In several situations, it is interesting to remove the denominators from the refined Block-Göttsche multiplicities, as in [Mik17] and [Bou19] for instance. In our situation, results also adopt a simpler form if we do so. We will thus forget about the denominators. Then $BG_g^X(\beta)(q)$ becomes a Laurent polynomial obtained by counting marked diagrams with a different multiplicity:

$$\overline{BG}_{g}^{X}(\beta)(q) = \sum_{(\mathcal{D},\mathfrak{m})} \prod_{e \in E^{0}(\mathcal{D})} (q^{w(e)/2} - q^{-w(e)/2})^{2} \prod_{e \in E^{\mathrm{top}}(\mathcal{D}) \cup E^{\mathrm{bot}}(\mathcal{D})} (q^{1/2} - q^{-1/2})$$

The first product is obtained clearing the denominators of each $[w]^2$. The second product comes from the ends: actually, each end contributes $[1] = \frac{q^{1/2}-q^{-1/2}}{q^{1/2}-q^{-1/2}} = 1$ in $BG_g^X(\beta)$, and this becomes $q^{1/2} - q^{-1/2}$ when clearing denominators. The degree of the invariant $\overline{BG}_g^X(\beta)$ with cleared denominators is $\operatorname{Area}(\Delta_\beta)$. This Laurent polynomial is symmetric (resp. antisymmetric) when $\beta \cdot K_X$ is even (resp. odd).

Example 2.15. For the toric surface \mathbb{P}^2 , if L is the class of a toric divisor one has

$$\begin{split} \overline{BG}_{0}^{\mathbb{P}^{2}}(L)(q) &= q^{1/2} - q^{-1/2}, \\ \overline{BG}_{0}^{\mathbb{P}^{2}}(2L)(q) &= (q^{1/2} - q^{-1/2})^{4} \cdot 1 \\ &= q^{2} - 4q + 6 - 4q^{-1} + q^{-2}, \\ \overline{BG}_{0}^{\mathbb{P}^{2}}(3L)(q) &= (q^{1/2} - q^{-1/2})^{7} \cdot (q + 10 + q^{-1}) \\ &= q^{9/2} + 3q^{7/2} - 48q^{5/2} + 168q^{3/2} - 294q^{1/2} + 294q^{-1/2} + \dots - q^{-9/2}. \end{split}$$

The Laurent polynomial $\overline{BG}_{q}^{X}(\beta)(q)$ can be turned into a true polynomial by setting

$$\widetilde{BG}_{g}^{X}(\beta)(x) = x^{\operatorname{Area}(\Delta_{\beta})}\overline{BG}_{g}^{X}(\beta)\left(\frac{1}{x}\right).$$

This way, the codegree *i* coefficient of $\overline{BG}_g^X(\beta)$ (term $q^{\operatorname{Area}(\Delta_\beta)-i}$) becomes the degree *i* coefficient of $\widetilde{BG}_g^X(\beta)$ (term x^i). Thanks to this change of variable, we can now view the refined invariant as a function

$$\widetilde{BG}_{g}^{A} : \operatorname{Eff}(X) \longrightarrow \mathbb{Z}\llbracket x \rrbracket, \ eta & \longmapsto & \widetilde{BG}_{g}^{X}(eta)(x) \end{cases}$$

with values in the ring of formal series with integer coefficients $\mathbb{Z}[x]$, even though for any β the value is polynomial in x of degree $2\text{Area}(\Delta_{\beta})$. The codomain $\mathbb{Z}[x]$ is a valuation ring, and can thus be endowed with the topology coming from the associated ultrametric distance. A basis of neighborhoods of 0 for this topology is given by the ideals $x^n \mathbb{Z}[x]$, so that $f \in \mathbb{Z}[x]$ is close to 0 if $f \equiv 0 \mod x^n$ for $n \in \mathbb{N}$ sufficiently big. We prefer to use $\mathbb{Z}[x]$ as codomain, because it is a complete space, more suited to express our asymptotic result. Meanwhile, we have a notion of neighborhood of infinity in the cone Eff(X): for C > 0, we say that $\beta \succ C$ if $\beta \cdot D \ge C$ for every toric divisor D.

Proposition 2.16. For any h-transverse toric surface X, any $\beta \in H_2(X, \mathbb{Z})$ and any $g \ge 0$, one has

$$\widetilde{BG}_g^X(eta)(x) = \sum_{({\mathbb D},{\mathfrak m})} \mu({\mathbb D})(x)$$

where the sum runs over the isomorphism classes of marked floor diagrams of Newton polygon Δ_{β} and genus g, and with

$$\mu(\mathbb{D})(x) = x^{\operatorname{codeg}(\mathbb{D})} (1-x)^{b^{\operatorname{top}}(\mathbb{D}) + b^{\operatorname{bot}}(\mathbb{D})} \prod_{e \in E^0(\mathbb{D})} \left(1 - x^{w(e)}\right)^2.$$

We call $\mu(\mathcal{D})$ the multiplicity of the floor diagram \mathcal{D} .

Proof. By the definition of the codegree one has

$$\operatorname{Area}(\Delta_{\beta}) = \operatorname{codeg}(\mathcal{D}) + \sum_{e \in E^{0}(\mathcal{D})} w(e) + rac{b^{\operatorname{top}} + b^{\operatorname{bot}}}{2}.$$

Hence

$$\begin{split} \widetilde{BG}_{g}^{X}(\beta) &= x^{\operatorname{Area}(\Delta_{\beta})} \sum_{(\mathcal{D},\mathfrak{m})} \prod_{e \in E^{0}(\mathcal{D})} (x^{-w(e)/2} - x^{w(e)/2})^{2} \prod_{e \in E^{\operatorname{top}}(\mathcal{D}) \cup E^{\operatorname{bot}}(\mathcal{D})} (x^{-1/2} - x^{1/2}) \\ &= \sum_{(D,\mathfrak{m})} x^{\operatorname{codeg}(\mathcal{D})} \prod_{e \in E^{0}(\mathcal{D})} (1 - x^{w(e)})^{2} \prod_{e \in E^{\operatorname{top}}(\mathcal{D}) \cup E^{\operatorname{bot}}(\mathcal{D})} (1 - x) \\ &= \sum_{(D,\mathfrak{m})} \mu(\mathcal{D}). \end{split}$$

The main results of [BJP22] and [Mév23] deal with the polynomiality in terms of β of the coefficient of fixed codegree i of $BG_g^X(\beta)(q)$, which become the x^i coefficient of $\widetilde{BG}_g^X(\beta)(x)$ in our setting. The polynomial behaviour is not affected by the clearing of denominators. Indeed, invariants with or without clearing of denominators differ by multiplication (or division) by $(1-x)^{-K_X \cdot \beta + 2g-2}$, whose coefficients up to degree i are also polynomials in β . We denote the degree i coefficient by $\left(\widetilde{BG}_g^X(\beta)\right)_i$. In [BJP22] the authors show that when X is a Hirzebruch surface or a (weighted) projective space, then for any $i \ge 0$ there exists a polynomial function $AR_{g,i}^X(\beta)$ on the lattice $H_2(X,\mathbb{Z})$ such that for every β with $\beta \cdot D$ large enough with respect to i and g for any D toric divisor, one has

$$\left(\widetilde{BG}_{g}^{X}(\beta)(x)\right)_{i}=AR_{g,i}^{X}(\beta).$$

Taking the generating series in i, the result can be rephrased as follows.

Theorem 2.17. ([BJP22]) Let X be a Hirzebruch surface. For every $g \ge 0$, there exists a function $AR_a^X : H_2(X, \mathbb{Z}) \to \mathbb{Z}[\![x]\!]$ which is a polynomial with coefficients in $\mathbb{Z}[\![x]\!]$, such that

$$\widetilde{BG}_{g}^{X}(\beta) = AR_{g}^{X}(\beta) + o(1) \in \mathbb{Z}\llbracket x \rrbracket.$$

The asymptotic development takes place when $\beta \to \infty$. We call AR_g^X the asymptotic refined invariant.

Remark 2.18. The result is likely to be true for any toric surface, but in [BJP22] the authors restrict the proof to a family of surfaces that includes Hirzebruch surfaces for technical reasons. In genus 0, this result is shown to hold in [Mév23] for any *h*-transverse toric surface, with explicit formula for the polynomials. We give in this paper another proof in genus 0 (Theorem 4.17) and a proof in genus 1 (Theorem 5.18) that holds for any *h*-transverse and non-singular toric surface.

Proof of the formulation using [BJP22]. The result amounts to prove that there exists polynomials $P_{a,i}^X(\beta)$ with degree bounded by a constant in g such that we have

$$\widetilde{BG}_{g}^{X}(\beta) = \sum_{i=0}^{\infty} P_{g,i}^{X}(\beta)x^{i} + o(1).$$

The o(1) means that for every $n \in \mathbb{N}$, there exists C > 0 such that

$$\beta \succ C \Rightarrow \widetilde{BG}_g^X(\beta) - \sum_{i=0}^{\infty} P_{g,i}^X(\beta) x^i \equiv 0 \mod x^n.$$

In other words, $\langle \widetilde{BG}_g^X(\beta) \rangle_i$ is given by $P_{g,i}^X$ if β is sufficiently big. This is the statement from [BJP22] up to the bound on the degrees. Actually, in [BJP22] the degree of $P_{g,i}^X$ is g+i. The dependence in i is due to the fact that the denominators are not removed from the Block-Göttsche multiplicity. If we remove them, [BJP22, Section 3.2] is modified as follows. The function $\Phi_i(k)$ in [BJP22, Corollary 3.6] does not depend on k anymore: it was previously a polynomial of degree i but is now constant, equal to 1 if i = 0, and to 0 if i > 0. Thus, in [BJP22, Corollary 3.7], the degree is also 0. When used in the proof of [BM07, Lemma 5.7], the bound i disappears and yields the fact that the degrees are bounded by g.

One way to interpret the asymptotic is that for any i and any class β large enough (understand $\beta \succ C$ for $C \in \mathbb{N}$ large enough), $AR_g^X(\beta)$ correctly gives the first i coefficients of $\widetilde{BG}_g^X(\beta)$. The strategy to compute the asymptotic refined invariant AR_g^X is thus to fix some $i \in \mathbb{N}$ and to compute $\widetilde{BG}_g^X(\beta)$ modulo x^{i+1} for β big enough. Provided we get an expression that does not depend on i, we can make i go to ∞ to obtain the value of $AR_g^X(\beta) \in \mathbb{Z}[\![x]\!]$.

This formulation as an asymptotic development is in fact inspired by a reformulation of the polynomiality conjecture [Göt98] on the number of curves with a fixed number of nodes δ passing through a suitable number of points. For a surface X with a line bundle \mathcal{L} , Göttsche's conjecture states that the number $N_{\delta}^{X}(\mathcal{L})$ of curves in the linear system $|\mathcal{L}|$ with δ nodes passing through a generic configuration of $h^{0}(X,\mathcal{L})-1-\delta$ points in X is given by a (universal) polynomial $P_{\delta}^{X}(\mathcal{L}^{2},\mathcal{L}\cdot K_{X},K_{X}^{2},c_{2}(X))$ provided \mathcal{L} is sufficiently ample. In other words, if we view P_{δ}^{X} and N_{δ}^{X} as functions

$$P_{\delta}^X, N_{\delta}^X : \operatorname{Amp}(X) \subset H^2(X, \mathbb{Z}) \longrightarrow \mathbb{Z},$$

where $\operatorname{Amp}(X) \subset H^2(X, \mathbb{Z})$ is the ample cone of X, we have

$$N^{\delta}(\mathcal{L}) = P^{\delta}(\mathcal{L}) + o(1)$$

As the topology of \mathbb{Z} is discrete, it means that we have equality if \mathcal{L} is sufficiently ample. The formulation in the refined case is more subtle since the topology on $\mathbb{Z}[x]$ is more sophisticated.

3. Generating series in fixed degree

In this section, we wish to determine the generating series in the genus parameter, i.e. $\sum_{g=0}^{\infty} AR_{g,i}u^g$ for fixed *i*. We provide an explicit expression for i = 0, 1. The i = 0 case amounts to compute the leading coefficient of the tropical refined invariant, which was already known from [IM13]. The main contribution is Theorem 3.2, which gives a closed formula for i = 1.

Recall from [BJP22] or [Mév23] that a diagram \mathcal{D} has codegree 0 if and only if the order is total on its floors, it has no side edge (i.e. an edge bypassing a floor), and the functions Rand L are increasing. Recall also that that when looking at floor diagrams of small codegree, we can assume using Lemma 2.14 that the diagrams have a total order on their vertices, and we can control the number of side edges as well as the monotonicity of the functions L and R.

In all this section, for $\beta \in H_2(X, \mathbb{Z})$ with Δ_β being *h*-transverse, we will refer as \mathcal{D}_0 to be the floor diagram of Figure 3. It is the unique diagram of Newton polygon Δ_β , genus 0 and codegree 0. We denote by ω_m the weight of the edge between the floors v_m and v_{m+1} for $1 \leq m \leq a-1$. Note that



FIGURE 3. The diagram \mathcal{D}_0 .

3.1. **Degree 0.** We start by computing $(AR_{g,0}^X)$, the leading term of the asymptotic refined invariant. This amounts to compute the leading coefficient of the tropical refined invariant, which was already handled in [IM13, Proposition 2.11] using the lattice path algorithm from [Mik05]. We recall a proof here, because it uses a construction starting from \mathcal{D}_0 that will appear several times in subsection 3.2.

Proposition 3.1 ([IM13]). The generating series in the genus parameter of the leading term of the asymptotic refined invariant is given by

$$\sum_{g \ge 0} A R_{g,0}^X u^g = (1+u)^{g_{\max}}.$$

Proof. To construct a marked floor diagram of positive genus and codegree 0, we add g_m edges between the floors v_m and v_{m+1} of \mathcal{D}_0 , marking the new edges increasingly from left to right, and splitting the weight ω_m onto the $g_m + 1$ edges. The genus of the new diagram is $g_1 + \cdots + g_{a-1}$. For each m there are $\binom{\omega_m - 1}{g_m}$ tuples of $g_m + 1$ positive integers with sum ω_m , i.e. ways to distribute ω_m onto the marked edges. Since we only care about the number of marked diagrams of genus g to compute $AR_{q,0}^X$, using the binomial formula one has

$$\sum_{g \ge 0} AR_{g,0}^X u^g = \sum_{\substack{g_1, \dots, g_{a-1} \ge 0}} u^{g_1 + \dots + g_{a-1}} \prod_{m=1}^{a-1} \binom{\omega_m - 1}{g_m}$$
$$= \prod_{m=1}^{a-1} \sum_{\substack{g_m \ge 0}} \binom{\omega_m - 1}{g_m} u^{g_m}$$
$$= \prod_{m=1}^{a-1} (1+u)^{\omega_m - 1} = (1+u)^{g_{\max}}.$$

15

3.2. **Degree 1.** We now compute the generating series of the second terms of the asymptotic refined invariants.

Theorem 3.2. For a h-transverse toric surface X, the asymptotic polynomials yielding the degree 1 coefficient are polynomials in β^2 , $K_X \cdot \beta$, χ , K_X^2 . Moreover, their generating series has the following expression:

$$\sum_{g \ge 0} AR_{g,1}^X u^g = (1+u)^{g_{\max}} \left[-\beta^2 \frac{u^3}{(1+u)^3} + 2(K_X \cdot \beta) \frac{u^2}{(1+u)^3} + \chi \frac{1}{1+u} - K_X^2 \frac{u}{(1+u)^3} \right].$$

Given that the multiplicity takes the form

$$\mu(\mathbb{D}) = x^{\operatorname{codeg}(\mathbb{D})} (1-x)^{b^{\operatorname{top}} + b^{\operatorname{bot}}} \prod_{e \in E_0(\mathbb{D})} (1-x^{w_e})^2 \,,$$

only diagrams with $\operatorname{codeg}(\mathcal{D}) = 0, 1$ contribute to $\sum_g AR_{g,1}^X u^g$. For the unique marked diagram of codegree 0 we need to consider the term in x, while for the marked floor diagrams of codegree 1 we need to consider their number. We subdivide the proof of Theorem 3.2 in four lemmas, each one computing the contribution of a specific family of diagrams to the global sum. To get Theorem 3.2, one only needs to sum the expressions from Lemmas 3.3,3.4,3.5 and 3.6.

Lemma 3.3. The codegree 0 diagrams with Newton polygon Δ_{β} contribute to $\sum_{q} AR_{q,1}^{X} u^{q}$:

$$(1+u)^{g_{\max}}\left[-(b^{\mathrm{top}}+b^{\mathrm{bot}})-2g_{\max}\frac{u^2}{(1+u)^2}-2(a-1)\frac{u(2+u)}{(1+u)^2}\right]$$

Proof. We construct a diagram of genus g and codegree 0 as in the proof of Theorem 3.1. A diagram \mathcal{D} of codegree 0 is counted with the degree 1 term of its multiplicity, that is

$$-(b^{\text{top}} + b^{\text{bot}}) - 2|\{e \in E^0(\mathcal{D}) \mid w_e = 1\}|$$

Hence, the contribution to $\sum_{g} AR_{g,1}^{X} u^{g}$ coming from the first term is $-(b^{\text{top}} + b^{\text{bot}})(1+u)^{g_{\text{max}}}$. We need to compute the contribution coming from the second term, i.e. enumerate the choice of a diagram together with an edge of weight 1. To determine this contribution, we proceed as previously but for any fixed m, we assume one of the $g_m + 1$ edges between v_m and v_{m+1} has weight 1, and it remains a weight $\omega_m - 1$ to split into g_m parts. Forgetting the -2, this gives

$$\begin{split} &\sum_{m=1}^{a-1} \left(\sum_{\substack{g_j \ge 0 \ j \neq m}} \prod_{\substack{g_j \neq m}} {\omega_j - 1 \choose g_j} u^{g_j} \right) \left(\sum_{\substack{g_m \ge 0 \ v_m \to v_{m+1}}} \sum_{\substack{w_m - 2 \\ v_m \to v_{m+1}}} {\omega_m - 2 \choose g_m - 1} u^{g_m} \right) \\ &= \sum_{m=1}^{a-1} (1+u)^{\sum_{j \neq m} [\omega_j - 1]} \sum_{\substack{g_m \ge 0}} (g_m + 1) {\omega_m - 2 \choose g_m - 1} u^{g_m} \\ &= \sum_{m=1}^{a-1} (1+u)^{g_{\max} - (\omega_m - 1)} \left[(\omega_m - 2) u^2 (1+u)^{\omega_m - 3} + 2u (1+u)^{\omega_m - 2} \right] \\ &= (1+u)^{g_{\max}} \left[g_{\max} \frac{u^2}{(1+u)^2} + (a-1) \frac{u(2+u)}{(1+u)^2} \right]. \end{split}$$

We now look at the diagrams of codegree 1. The degree 1 term of the multiplicity of a diagram of codegree 1 is 1, so it suffices to determine the number of marked floor diagrams of codegree 1. There are two possibilities for the codegree being 1: the presence of a side edge, i.e. an edge bypassing a floor, or a slope inversion, i.e. a lack of growth of the divergence function. We investigate all the cases.

Lemma 3.4. The codegree 1 diagrams with an infinite side edge, with Newton polygon Δ_{β} contribute to $\sum_{q} AR_{a,1}^{X} u^{q}$:

$$(1+u)^{g_{\max}}\left[(\omega_1+\omega_{a-1}-2)\frac{u}{(1+u)^2}+(b^{\text{bot}}+b^{\text{top}})\frac{1}{1+u}+2\frac{2+u}{(1+u)^2}\right].$$

Proof. We deal with the case when the side edge is a source ; the case when it is a sink is handled similarly by symmetry.

Let \mathcal{D}_{bot} be the diagram of Figure 4a; it is obtained from \mathcal{D}_0 by putting a source adjacent to v_2 . It has genus 0 and codegree 1. Let $\tilde{\omega}_k$ be the weight of the edge between v_k and v_{k+1} for $1 \leq k \leq a-1$. One has

$$\widetilde{\omega}_1 = \omega_1 - 1 ext{ and } \widetilde{\omega}_k = \omega_k, \ 2 \leqslant k \leqslant a - 1.$$

To create a diagram of genus g, as in Theorem 3.1 we add g_m edges between the floor v_m and v_{m+1} of \mathcal{D}_{bot} , marking the new edges increasingly from left to right, and split the weight w_m onto the $g_m + 1$ edges. The genus of the new diagram is $g_1 + \cdots + g_{a-1}$ and for each m there are $\binom{\widetilde{\omega}_m - 1}{g_m}$ ways to distribute $\widetilde{\omega}_m$ onto the marked edges. To entirely determine the marked floor diagram, it remains to mark the side edge. It is parallel to $(g_1 + 1) + (b^{\text{bot}} - 1)$ edges and 1 floor, hence there are $g_1 + b^{\text{bot}} + 2$ possibilities for its marking. In the end, this case contributes

$$\sum_{g_1,\dots,g_{a-1}\geqslant 0} (g_1+b^{\text{bot}}+2) \prod_{m=1}^{a-1} {\widetilde{\omega}_m - 1 \choose g_m} u^{g_m}$$

= $(1+u)^{g_{\max}-(\omega_1-1)} \sum_{g_1\geqslant 0} (g_1+b^{\text{bot}}+2) {\widetilde{\omega}_1 - 1 \choose g_1} u^{g_1}$
= $(1+u)^{g_{\max}-(\omega_1-1)} \left[(\widetilde{\omega}_1 - 1)u(1+u)^{\widetilde{\omega}_1-2} + (b^{\text{bot}}+2)(1+u)^{\widetilde{\omega}_1-1} \right]$
= $(1+u)^{g_{\max}} \left[(\omega_1 - 1) \frac{u}{(1+u)^2} + b^{\text{bot}} \frac{1}{1+u} + \frac{2+u}{(1+u)^2} \right]$

Similarly, if the side edge is a sink we get

$$(1+u)^{g_{\max}}\left[(\omega_{a-1}-1)\frac{u}{(1+u)^2}+b^{\operatorname{top}}\frac{1}{1+u}+\frac{2+u}{(1+u)^2}\right].$$

We get the result summing the two cases.

Lemma 3.5. The codegree 1 diagrams with a bounded side edge contribute to $\sum_{q} AR_{q,1}^{X} u^{g}$:

$$(1+u)^{g_{\max}}\left[\sum_{j=1}^{a-2} \left(\omega_j + \omega_{j+1} - 2\right) \frac{u^2}{(1+u)^3} + 2(a-2)\frac{u(2+u)}{(1+u)^3}\right].$$

Proof. Start with the diagram \mathcal{D}_j of figure 4b; it has genus 1 and a side edge around the floor v_{j+1} . Let $\tilde{\omega}_m$ be the weight of the edge between v_m and v_{m+1} for $1 \leq m \leq a-1$. One has

$$\widetilde{\omega}_j = \omega_j - 1, \ \widetilde{\omega}_{j+1} = \omega_{j+1} - 1 \text{ and } \widetilde{\omega}_m = \omega_m, \ m \neq j.$$



FIGURE 4

As previously, we add g_m edges between the floor v_m and v_{m+1} of \mathcal{D}_{bot} , mark the new edges increasingly from left to right, and split the weight w_m onto the $g_m + 1$ edges. The created diagram as genus $1 + g_1 + \cdots + g_{a-1}$. The side edge is parallel to 1 floor and $g_j + g_{j+1} + 2$ edges, so there are $g_j + g_{j+1} + 4$ possibilities for its marking. Hence, the contribution in that case is

$$u\sum_{j=1}^{a-2} \sum_{g_1,\dots,g_{a-1} \ge 0} (g_j + g_{j+1} + 4) \prod_{m=1}^{a-1} {\tilde{\omega}_m - 1 \choose g_m} u^{g_m}$$

=
$$\sum_{j=1}^{a-2} (\tilde{\omega}_j + \tilde{\omega}_{j+1} - 2) u^2 (1+u)^{g_{\max}-3} + 4(a-2) u(1+u)^{g_{\max}-2}$$

=
$$(1+u)^{g_{\max}} \left[\sum_{j=1}^{a-2} (\omega_j + \omega_{j+1} - 2) \frac{u^2}{(1+u)^3} + 2(a-2) \frac{u(2+u)}{(1+u)^3} \right].$$

Lemma 3.6. The codegree 1 diagrams with an slope inversion contribute to $\sum_{g} AR_{q,1}^{X} u^{g}$:

 $(\chi - 4)(1+u)^{g_{\max}-1}$.

Proof. To get a floor diagram of codegree 1 with an inversion, the only possibility is the existence of a unique couple (v, v') of adjacent floors such that $v \prec v'$ and R(v) = R(v') + 1 or L(v) = L(v') + 1, and anywhere else in the floor diagram, R and L are increasing. If χ is the number of corners of Δ_{β} , there are $\chi - 4$ such pairs, one for each corner of Δ non-adjacent to a horizontal side. The only difference with the codegree 0 diagram from Figure 3 is that the weight between v_m and v_{m+1} is $\omega_m - 1$ so that the sum of weights yields $g_{\max} - 1$ instead of g_{\max} . In the end, this case contributes

$$(\chi - 4)(1 + u)^{g_{\max} - 1}.$$

We can finally prove Theorem 3.2.

Proof of Theorem 3.2. Summing the contributions of Lemmas 3.3, 3.4, 3.5 and 3.6, and using the relations

$$\begin{cases} g_{\max} = \sum_{\substack{m=1 \ m=1}}^{a-1} (\omega_m - 1), \\ -K_X \cdot \beta = b^{\text{top}} + b^{\text{bot}} + 2a, \\ \beta^2 = 2g_{\max} - 2 + K_X \cdot \beta \\ K_X^2 + \chi = 12 \end{cases}$$

we get

$$\sum_{g \ge 0} AR_{g,1}^X u^g = (1+u)^{g_{\max}} \left[-\beta^2 \frac{u^3}{(1+u)^3} + 2(K_X \cdot \beta) \frac{u^2}{(1+u)^3} + \chi \frac{1}{1+u} - K_X^2 \frac{u}{(1+u)^3} \right].$$

4. Asymptotic refined invariant for genus 0

In this section we compute the asymptotic refined invariant for genus 0 for any *h*-transverse and non-singular polygon having two horizontal sides. This was already done in [Mév23], but we give in this section a different proof to present methods that can be applied when dealing with genus 1 in Section 5. We start with Hirzebruch surfaces before going into the general case. To do so, we use *words* to enumerate marked floor diagrams contributing to the asymptotic count. We will compute $\widetilde{BG}_0^X(\beta)$ modulo x^{i+1} for some *i*, before letting *i* goes to ∞ .

4.1. The case of Hirzebruch surfaces. The tropical refined invariants can be computed by using enumeration of marked floor diagrams. However, as shown in [BJP22, Lemma 4.1], if one cares about the asymptotic of coefficients of fixed codegree only a handful of diagrams contribute. Consider the Hirzebruch surface \mathbb{F}_{δ} , so that all floors have the same divergence. In the genus 0 case, provided that a, b > i, any marked diagram contributing to a coefficient of degree at most *i* satisfies the following:

- \triangleright the floors are totally ordered in the diagram,
- ▷ some of the top (resp. bottom) ends might not be attached to the first (resp. last) floor but to another floor,

Let u_j^{top} (resp. u_j^{bot}) be the number of top (resp. bottom) ends that skip j floors. The codegree of a diagram \mathcal{D} comes from these ends not attached to the extremal floors. It is equal to

$$\operatorname{codeg}(\mathcal{D}) = \sum_{j=1}^{\infty} j(u_j^{\operatorname{top}} + u_j^{\operatorname{bot}}),$$

note that this sum is actually finite. Each diagram is characterized by the numbers $(u_j^{\text{top}}, u_j^{\text{bot}})$. We then have to account for the markings. We restrict to the collection of diagrams for which the floors are totally ordered. Rather than enumerating the latter and count their markings, as done in [BJP22, Section 4] and [Mév23], we directly count these marked marked diagrams, encoding them with words.

4.1.1. From marked diagrams to words. We consider words over the alphabet $\{f, e, b_j, t_j\}_{j \in \mathbb{N}}$. The letters used stand for "floor", "edge/elevator", "bottom end" and "top end". The indices of the letters refer to the number of floors they skip. We first explain how to get a word $W(\mathcal{D})$ from a genus 0 marked diagram \mathcal{D} whose floors are totally ordered. Let $aE + bF \in H_2(\mathbb{F}_{\delta}, \mathbb{Z})$ be the class of the diagram. The floors of \mathcal{D} are labelled from 1 to a. The letters of the word $W(\mathcal{D})$ are in ordered correspondence with the marked points of \mathcal{D} with the following rule:

- \triangleright for a marked point on a floor, the letter is f,
- \triangleright for a marked point on a bounded edge, the letter is e,
- ▷ for a marked point on a top end that skips $j \ge 0$ floors, the letter is t_j ,
- ▷ for a marked point on a bottom end that skips $j \ge 0$ floors, the letter is b_j .



FIGURE 5. The two marked diagrams corresponding to the words from Example 4.1.

Example 4.1. On Figure 5, we have two genus 0 marked diagrams with totally ordered floors corresponding to the words

$$W(\mathcal{D}_1) = b_0 b_0 b_1 b_0 feb_1 feft_0 t_0,$$

$$W(\mathcal{D}_2) = b_0 b_0 fb_1 efefeft_1 t_0 t_0.$$

Note that it is possible to recover the marked diagrams from the words.

This correspondence between diagrams and words is in fact bijective provided we have some assumptions on the words.

 \Diamond

Proposition 4.2. Let \mathcal{D} be a marked floor diagram in the class $\beta = aE + bF \in H_2(\mathbb{F}_{\delta}, \mathbb{Z})$. Then the word $W(\mathcal{D})$ satisfies the following.

(i) Forgetting about the letters b_* and t_* , the word is just

$$(fe)^{a-1}f = fefe \cdots fef.$$

Moreover, there are b letters b_* and $b + \delta a$ letters t_* .

- (ii) Given a letter b_k , assume the word forgetting the e, t_j and remaining b_j is $f^p b_k f^{a-p}$, then we have $k \ge p$.
- (iii) Given a letter t_k , assume the word forgetting the e, b_j and remaining t_j is $f^{a-p}t_k f^p$, then we have $k \ge p$.

Conversely, a word satisfying the above conditions yields a marked floor diagram for which the floors are totally ordered. The set of words satisfying the above conditions is denoted by \mathcal{W}_{aE+bF} .

- *Proof.* (i) The diagram has a floors and they are totally ordered, so that each floor is linked to the next one by a unique elevator. Thus forgetting about t_* and b_* , we get fef \cdots fef. The numbers of floors as well as the number of ends in each direction are fixed by the class aE + bF.
 - (ii) In the word $f^p b_k f^{a-p}$, the marking of the end encoded by b_k lies between the floors p and p+1. Thus, the end being a bottom end, it skips at least the p first floors and is attached to a floor after the (p+1)-th floor, so that $k \ge p$.
 - (iii) The reasoning is the same but with top ends instead of bottom ends.

For the converse construction, let W be a word satisfying (i)-(iii). We start with the ordered graph having a vertices, each linked to the next one by a unique edge, and with a marking. For each b_j (resp. t_j) we insert a bottom (resp. top) end attached to the floor j + 1 (resp. a - j) with a marking lying at the corresponding place in the word. There is a unique way to add weights to the bounded edges so that the diagram is balanced. Condition (i) ensures that the diagram has the right number of floors and ends, and the conditions (ii) and (iii) ensures that it is possible to place the marking of an end on the latter.

4.1.2. Words and codegrees. We define the codegree function on \mathcal{W}_{aE+bF} so that the codegree of a word matches the codegree of the associated diagram. Let \mathcal{W} be the set of all words on the considered alphabet, which is a monoid. The codegree function is actually the restriction of the following morphism of monoids:

$$\begin{array}{ccc} \operatorname{codeg} : & \mathscr{W} & \longrightarrow & \mathbb{N} \\ & \mathsf{t}_j, \ \mathsf{b}_j & \longmapsto & j \\ & \mathsf{e}, \ \mathsf{f} & \longmapsto & 0 \end{array}$$

and by construction we have $\operatorname{codeg}(\mathcal{D}) = \operatorname{codeg}(W(\mathcal{D}))$.

Remark 4.3. The definition of \mathscr{W}_{aE+bF} allows the letters b_* and t_* to interlace, meaning there might be a b_* after a t_* . However, if a b_* lies after a t_* then all floors are skipped by at least one of these two ends so that $\operatorname{codeg}(W(\mathcal{D})) \ge a$. If we restrict to words of codegree at most i and if a > i, then this situation does not appear.

The following lemma describes the shape of the words that have a bounded codegree provided the class is large enough.

Lemma 4.4. Assume $i \ge 1$ and a > 2i. The words in \mathcal{W}_{aE+bF} of codegree at most i are of the following form:

$$\begin{split} & \mathsf{B}_0 \left[\prod_{j=1}^i \mathsf{f} \mathsf{B}_j^{(1)} \mathsf{e} \mathsf{B}_j^{(2)} \right] (\mathsf{f} \mathsf{e})^{a-2i} \left[\prod_{j=1}^i \mathsf{f} \mathsf{T}_{i+1-j}^{(1)} \mathsf{e} \mathsf{T}_{i+1-j}^{(2)} \right] \mathsf{f} \mathsf{T}_0 \\ & = \mathsf{B}_0 \mathsf{f} \mathsf{B}_1^{(1)} \mathsf{e} \mathsf{B}_1^{(2)} \mathsf{f} \mathsf{B}_2^{(1)} \mathsf{e} \mathsf{B}_2^{(2)} \cdots \mathsf{f} \mathsf{e} \mathsf{f} \mathsf{e} \mathsf{f} \mathsf{e} \cdots \mathsf{T}_2^{(1)} \mathsf{e} \mathsf{T}_2^{(2)} \mathsf{f} \mathsf{T}_1^{(1)} \mathsf{e} \mathsf{T}_1^{(2)} \mathsf{f} \mathsf{T}_0, \end{split}$$

where $B_0, B_j^{(k)}$ (resp. $T_0, T_j^{(k)}$) are words in the letters $\{b_*\}_{* \ge 0}$ and $\{b_*\}_{* \ge j}$ (resp. $\{t_*\}_{* \ge 0}$ and $\{t_*\}_{* \ge j}$).

Proof. As a letter b_* put after the first i+1 letters f contributes at least i+1 to the codegree, it cannot appear if the latter is assumed to be smaller than i, and similarly for t_* letters. \Box

Basically, the word has a core $(fe)^{a-1}f$ and we insert a word in the letters b_* (called *B*-word) between each of the 2i consecutive letters on the left, a word in the letters t_* (called *T*-word) similarly on the right. As the roles of *B*-words and *T*-words are symmetric, we call them "end-words". We denote by S the set of sentences, i.e. of families of end-words in s_* where S, s are meant to be replaced by T, t or B, b:

$$\mathbb{S} = \{(\mathsf{S}_0, \mathsf{S}_1^{(1)}, \mathsf{S}_1^{(2)}, \dots, \mathsf{S}_i^{(1)}, \mathsf{S}_i^{(2)}) \mid i \ge 0, \ \mathsf{S}_j^{(k)} \text{ word in } \{\mathsf{s}_*\}_{* \ge j}\}.$$

It is endowed with functions

$$codeg: \mathbb{S} \longrightarrow \mathbb{N},$$
$$\ell_0, \ \ell_j^{(k)}: \mathbb{S} \longrightarrow \mathbb{N},$$
$$\ell: \mathbb{S} \longrightarrow \mathbb{N},$$

that associate to a sentence in S the sum of the codegrees of its words, the length of the words S_0 and $S_j^{(k)}$ (maybe 0), and the sum of their lengths. For $n \ge 0$ we denote by S(n) the set of sentences with total length n. Lemma 4.4 asserts that choosing a word in \mathcal{W}_{aE+bF} having codegree at most i and with a, b large enough amounts to choose :

 \triangleright an element $\mathfrak{b} \in S(b)$ that encodes the *B*-words,

 \triangleright an element $\mathfrak{t} \in \mathfrak{S}(b + \delta a)$ that encodes the *T*-words,

such that $\operatorname{codeg}(\mathfrak{t}) + \operatorname{codeg}(\mathfrak{b}) \leq i$. Essentially, elements of $\mathcal{S}(b)$ and $\mathcal{S}(b + \delta a)$ tell us how to construct half-diagrams which are glued back together. Hence, the computation of a generating series over \mathcal{W}_{aE+bF} will split into the computations of some generating series over $\mathcal{S}(b)$ and $\mathcal{S}(b + \delta a)$.

Definition 4.5. We define the *multiplicity* of a sentence $\mathfrak{s} \in S(n)$ to be

$$\mu_{\mathbb{S}(n)}(\mathfrak{s}) = (1-x)^n x^{\operatorname{codeg}(\mathfrak{s})}.$$

4.1.3. *Enumeration of words.* We now compute the generating series of sentences with their multiplicity.

Lemma 4.6. Assume $i \ge 1$ and a, b > 2i. The multiplicity modulo x^{i+1} of the diagram \mathcal{D} encoded by a word $\mathsf{W} \in \mathscr{W}_{aE+bF}$ is $(1-x)^{2b+\delta a} x^{\operatorname{codeg}(\mathsf{W})}$.

Proof. By definition, the multiplicity is

$$(1-x)^{2b+\delta a}x^{\operatorname{codeg}(\mathsf{W})}\prod_e(1-x^{w(e)})^2,$$

where the product is over the bounded edges e of \mathcal{D} . Assume $\operatorname{codeg}(\mathsf{W}) \leq i$, otherwise there is nothing to prove since we get 0 modulo x^{i+1} . By Lemma 2.14, the unique edge between two consecutive floors has weight bigger than i. Thus, $(1 - x^{w(e)})^2 \equiv 1 \mod x^{i+1}$. \Box

Lemma 4.7. Let $n > i \ge 1$. The generating series of length n sentences counted with their multiplicity is

$$(1-x)^n \sum_{\mathfrak{s} \in \mathbb{S}(n)} x^{\operatorname{codeg}(\mathfrak{s})} \equiv p(x)^2 \mod x^{i+1}.$$

Proof. As we are looking at an equality modulo x^{i+1} , we only care about the elements of S(n)with codegree at most i since the others elements contribute 0. In particular, each sentence contains at most 2i+1 words and the letters involved in each word can only be in $\{s_*\}_{0 \le * \le i+1}$,

so that the sum on the left is well-defined modulo x^{i+1} . Let's fix $(l_0, l_j^{(k)})_{\substack{1 \leq j \leq n \\ k=1,2}}$ a family of integers such that $n = l_0 + \sum_{j,k} l_j^{(k)}$, and look at sentences $\mathfrak{s} = (\mathsf{S}_0, \mathsf{S}_1^{(1)}, \dots, \mathsf{S}_n^{(2)})$ with $\ell_0(\mathfrak{s}) = l_0$ and $\ell_j^{(k)}(\mathfrak{s}) = l_j^{(k)}$. The sum of the multiplicities of such contained in the sentences in

sentences is

$$(1-x)^n \left(\sum_{\ell(\mathsf{S}_0)=l_0} x^{\operatorname{codeg}(\mathsf{S}_0)}\right) \times \prod_{j,k} \left(\sum_{\ell(\mathsf{S}_j^{(k)})=l_j^{(k)}} x^{\operatorname{codeg}(\mathsf{S}_j^{(k)})}\right).$$

Letters in S_0 (resp. $S_j^{(k)}$) can take values in $\{s_*\}_{*\geq 0}$ (resp. $\{s_*\}_{*\geq j}$), so one has

$$\sum_{\ell(\mathsf{S}_{0})=l_{0}} x^{\operatorname{codeg}(\mathsf{S}_{0})} = \left(\sum_{k \ge 0} x^{k}\right)^{l_{0}} = \left(\frac{1}{1-x}\right)^{l_{0}}$$

and
$$\sum_{\ell(\mathsf{S}_{j}^{(k)})=l_{j}^{(k)}} x^{\operatorname{codeg}(\mathsf{S}_{j}^{(k)})} = \left(\sum_{k \ge j} x^{k}\right)^{l_{j}^{(k)}} = \left(\frac{x^{j}}{1-x}\right)^{l_{j}^{(k)}}$$

and the sum of the multiplicities of the sentences with fixed lengths equal to $(l_0, l_i^{(k)})_{j,k}$ is

$$(1-x)^n \left(rac{1}{1-x}
ight)^{l_0} \prod_{j,k} \left(rac{x^j}{1-x}
ight)^{l_j^{(k)}} = \prod_{j,k} x^{j l_j^{(k)}}.$$

It remains to sum over all the possible choices of $(l_0, l_i^{(k)})_{j,k}$. Because the total length of the sentences is n, we can forget about l_0 since it is fully determined by the $l_i^{(k)}$. Moreover we can sum over $l_j^{(k)} \ge 0$ instead of $\sum l_j^{(k)} = n$ because the excess terms will contribute 0 modulo x^n . Therefore, we get

$$\sum_{(l_0, l_j^{(k)})} x^{\sum j l_j^{(k)}} = \prod_{j, k} \sum_{l_j^{(k)} \ge 0} x^{j l_j^{(k)}} = \left(\prod_{j=1}^n \frac{1}{1 - x^j}\right)^2 \equiv p(x)^2 \bmod x^n.$$

Theorem 4.8. The genus 0 asymptotic refined invariant of the Hirzebruch surface \mathbb{F}_{δ} is

$$AR_0^{\mathbb{F}_\delta} = p(x)^4,$$

where p(x) is the generating series of partition numbers.

Proof. We can determine $AR_0^{\mathbb{F}_{\delta}} \mod x^{i+1}$ by summing the multiplicities of the words of \mathscr{W}_{aE+bF} of codegree at most i, and with a, b > 2i. According to Lemma 4.4, choosing a word of codegree at most i amounts to choose sentences $\mathfrak{b} \in \mathfrak{S}(b)$ and $\mathfrak{t} \in \mathfrak{S}(b + \delta a)$ with $\operatorname{codeg}(\mathfrak{b}) + \operatorname{codeg}(\mathfrak{t}) \leq i$. Lemma 4.6 ensures that the multiplicity of the word is

$$(1-x)^b x^{\operatorname{codeg}(\mathfrak{b})} (1-x)^{b+\delta a} x^{\operatorname{codeg}(\mathfrak{t})}.$$

Hence, summing over $S(b) \times S(b + \delta a)$ (and potentially counting terms which contribute 0) modulo x^{i+1}) the generating series factors modulo x^{i+1} :

$$\left((1-x)^b\sum_{\mathfrak{b}\in\mathbb{S}(b)}x^{\operatorname{codeg}(\mathfrak{b})}\right)\left((1-x)^{b+\delta a}\sum_{\mathfrak{t}\in\mathbb{S}(b+\delta a)}x^{\operatorname{codeg}(\mathfrak{t})}\right).$$

Using Lemma 4.7 we get the result modulo x^{i+1} for any *i*, and we conclude.

4.2. The case of *h*-transverse toric surfaces. We now consider the case of a toric surface X associated to a h-transverse, horizontal and non-singular polygon Δ . Let $\beta \in H_2(X,\mathbb{Z})$ be the corresponding homology class. Given a primitive vector α that positively generates a ray of the dual fan of Δ , we denote by D_{α} the corresponding toric divisor.

4.2.1. Words and codegree for h-transverse polygons. Compared with the Hirzebruch case, the marked floor diagrams are modified by incorporating the data (L, R), i.e. assigning a pair of integers called *sloping pair* to each floor. According to [BJP22, Section 3], the codegree coming from the sloping pairs is

$$\operatorname{codeg}(L,R) = \sum_{\substack{v \prec v' \\ \text{s.t. } L(v) > L(v')}} (L(v) - L(v')) + \sum_{\substack{v \prec v' \\ \text{s.t. } R(v) > R(v')}} (R(v) - R(v')) + \sum_{\substack{v \prec v' \\ \text{s.t. } R(v) > R(v')}} (R(v) - R(v')) + \sum_{\substack{v \prec v' \\ \text{s.t. } R(v) > R(v')}} (R(v) - R(v')) + \sum_{\substack{v \prec v' \\ \text{s.t. } R(v) > R(v')}} (R(v) - R(v')) + \sum_{\substack{v \prec v' \\ \text{s.t. } R(v) > R(v')}} (R(v) - R(v')) + \sum_{\substack{v \prec v' \\ \text{s.t. } R(v) > R(v')}} (R(v) - R(v')) + \sum_{\substack{v \prec v' \\ \text{s.t. } R(v) > R(v')}} (R(v) - R(v')) + \sum_{\substack{v \prec v' \\ \text{s.t. } R(v) > R(v')}} (R(v) - R(v')) + \sum_{\substack{v \prec v' \\ \text{s.t. } R(v) > R(v')}} (R(v) - R(v')) + \sum_{\substack{v \prec v' \\ \text{s.t. } R(v) > R(v')}} (R(v) - R(v')) + \sum_{\substack{v \prec v' \\ \text{s.t. } R(v) > R(v')}} (R(v) - R(v')) + \sum_{\substack{v \prec v' \\ \text{s.t. } R(v) > R(v')}} (R(v) - R(v')) + \sum_{\substack{v \prec v' \\ \text{s.t. } R(v) > R(v')}} (R(v) - R(v')) + \sum_{\substack{v \prec v' \\ \text{s.t. } R(v) > R(v')}} (R(v) - R(v')) + \sum_{\substack{v \prec v' \\ \text{s.t. } R(v) > R(v')}} (R(v) - R(v')) + \sum_{\substack{v \prec v' \\ \text{s.t. } R(v) > R(v')}} (R(v) - R(v')) + \sum_{\substack{v \prec v' \\ \text{s.t. } R(v) > R(v')}} (R(v) - R(v')) + \sum_{\substack{v \prec v' \\ \text{s.t. } R(v) > R(v')}} (R(v) - R(v')) + \sum_{\substack{v \prec v' \\ \text{s.t. } R(v) > R(v')}} (R(v) - R(v')) + \sum_{\substack{v \prec v' \\ \text{s.t. } R(v) > R(v')}} (R(v) - R(v')) + \sum_{\substack{v \prec v' \\ \text{s.t. } R(v) > R(v')}} (R(v) - R(v')) + \sum_{\substack{v \prec v' \\ \text{s.t. } R(v) > R(v')}} (R(v) - R(v')) + \sum_{\substack{v \dashv v' \\ \text{s.t. } R(v) > R(v')}} (R(v) - R(v')) + \sum_{\substack{v \dashv v' \\ \text{s.t. } R(v) > R(v')}} (R(v) - R(v')) + \sum_{\substack{v \dashv v' \\ \text{s.t. } R(v) > R(v')}} (R(v) - R(v')) + \sum_{\substack{v \dashv v' \\ \text{s.t. } R(v) > R(v')}} (R(v) - R(v')) + \sum_{\substack{v \dashv v' \\ \text{s.t. } R(v) > R(v')}} (R(v) - R(v')) + \sum_{\substack{v \dashv v' \\ \text{s.t. } R(v') > R(v')}} (R(v) - R(v')) + \sum_{\substack{v \dashv v' \\ \text{s.t. } R(v') > R(v')}} (R(v) - R(v')) + \sum_{\substack{v \dashv v' \\ \text{s.t. } R(v') > R(v')}} (R(v) - R(v')) + \sum_{\substack{v \dashv v' \\ \text{s.t. } R(v') > R(v')}} (R(v) - R(v')) + \sum_{\substack{v \dashv v' \\ \text{s.t. } R(v') > R(v')}} (R(v) - R(v')) + \sum_{\substack{v \dashv v' \\ \text{s.t. } R(v') > R(v')}} (R(v) - R(v')) + \sum_{\substack{v \dashv v' \\ \text{s.t. } R(v') > R(v')}} (R(v) - R(v')) + \sum_{\substack{v \dashv v' \\ \text{s.t. } R(v') > R(v')}} (R(v) - R(v')) + \sum_{\substack{v \dashv v' \\ \text{s.t. } R(v') > R(v')}} ($$

Elements in each of the sums are called *inversions*. In particular, the contribution to the codegree is 0 if L and R are increasing.

To enable the word approach to treat the case of h-transverse polygons, we need to add a sloping pair to each floor. We now consider the alphabet $\{e, f_{*,*}, t_*, b_*\}$ where the indices of $f_{*,*}$ are the members of the sloping pair. Similarly to Proposition 4.2, we have the following lemma that relates words to diagrams.

Proposition 4.9. Let \mathcal{D} be a marked floor diagram in the class $\beta \in H_2(X,\mathbb{Z})$. Then the word $W(\mathcal{D})$ satisfies the following.

(i) Forgetting about b_* , t_* and indices of $f_{*,*}$, the word is $(fe)^{a-1}f$. Moreover, there are b^{top} letters t_* and b^{bot} letters b_* .

(ii)-(iii) from Proposition 4.2 are still satisfied.

(iv) If $k \in b_{left}(\Delta_{\beta})$ (resp. $b_{right}(\Delta_{\beta})$), the number of appearances of k as a L-value (resp. *R*-value) in the sloping pairs is $\beta \cdot D_{\alpha}$, where $\alpha = (-1, k)$ (resp. (1, k)).

We denote by \mathscr{W}_{β} the set of words satisfying the above conditions. Given a word $\mathsf{W} \in \mathscr{W}_{\beta}$, there is a unique way to recover a marked floor diagram in the class β potentially with negative weights.

Proof. The proof of the first three points is verbatim to those of Proposition 4.2. The last one results from the definition of sloping pairs. For the converse construction, we also proceed as in Proposition 4.2. The difference is that when adding the weights of the elevators, we may obtain negative or zero weights. \square

Remark 4.10. During the reconstruction, the weights that appear may be negative. However, for the words of \mathcal{W}_{β} that we will consider all the weights are positive, see Lemma 4.11.

In a word W, we say that two letters $f_{\ell,r}$ and $f_{\ell',r'}$ appearing in that order form a *left* inversion (resp. right inversion) if $\ell > \ell'$ (resp. r > r'). The size of this inversion is the quantity $\ell - \ell'$ (resp. r - r').

The codegree function on \mathcal{W}_{β} is defined to match the codegree of the marked floor diagrams. The difference with the Hirzebruch case is that the codegree comes from the *T*-words and *B*-words, but also from the sloping pairs:

 $\begin{array}{rcl} \mathrm{codeg}: & \mathscr{W}_{\beta} & \longrightarrow & \mathbb{N} \\ & & \mathsf{W} & \longmapsto & \mathrm{codeg}(\mathrm{ft}(\mathsf{W})) + \mathrm{codeg}(L,R) \end{array}$

where ft(W) is the word where we forget the indices of the letters $f_{*,*}$.

Up to the indices of letters $f_{*,*}$, Lemma 4.4 still applies for words in \mathscr{W}_{β} under the hypothesis a > 2i. We deal with the indices of letters $f_{*,*}$ in the following lemma.

Lemma 4.11. Let $i \ge 1$ and assume that for each toric divisor D we have $\beta \cdot D > 2i$. If $W \in \mathcal{W}_{\beta}$ has codegree at most i then:

- (i) all the inversions in the sloping pairs are of size one, i.e. correspond to consecutive sides of the polygon,
- (ii) two letters $f_{*,*}$ part of an inversion are separated by at most i-1 letters $f_{*,*}$,
- (iii) the weights of the elevators in the associated diagram are strictly bigger than i. In particular they are positive, so that W corresponds to a true marked diagram.

Proof. We denote by (L, R) the tuple of sloping pairs of W. We first notice that the tuple L (and similarly for R) differs from the unique tuple of increasing slopes by a finite number of transpositions that switches two consecutive elements. Indeed, this is true for the tuple of codegree 0 since in that case this tuple is increasing. If we consider a tuple of positive codegree then there is a consecutive pair that forms an inversion; if not, the tuple would be increasing. Then, switching both members of the inversion decreases the codegree, and we conclude by induction. Each transposition switching consecutive elements increases the codegree then the increasing tuple by at most i transpositions.

As we assume the lengths of the sides of Δ_{β} to be bigger than 2i, it is not possible to create an inversion of size bigger than 2 with only *i* transpositions, proving (*i*).

Take an inversion $(\ldots, k+1, \ldots, k, \ldots)$ with *i* elements in-between. Any of these *i* elements is either *k* or k+1. If it is a *k* it provides an inversion with the left k+1, and if it is a k+1 it provides an inversion with the right *k*. Hence we get at least 1+i inversion, which is impossible, proving (ii).

Finally, for (iii) if the codegree is 0 then the assumption ensures that the weights of all the elevators are bigger than 2i by Lemma 2.14. If not, each transposition decreases the weight of an edge by 1. Thus, they remain strictly bigger than i after i transpositions.

4.2.2. Encoding the sloping pairs. Proposition 4.9 states that the elements of the sloping pairs are assigned to the floors with some constraints. We use the following objects to encode these assignments. Let \mathcal{P} be the set of non-constant sequences $\mathfrak{p} \in \{\bullet, \circ\}^{\mathbb{Z}}$ up to reindexation by translation of the index such that the set of pairs

$$I(\mathfrak{p}) = \{(k,l) \mid k < l, \mathfrak{p}_k = \circ, \mathfrak{p}_l = \bullet\},\$$

is finite. These pairs are also called *inversions*. We then set $codeg(\mathfrak{p}) = |I(\mathfrak{p})|$.

Example 4.12. We consider the following element, for which the first \circ has index 0:

Since $I(\mathfrak{p}) = \{(0,7), (1,7), (3,7), (6,7), (0,5), (1,5), (3,5), (0,4), (1,4), (3,4), (0,2), (1,2)\},$ it has codegree 12.

We notice that for each element $\mathfrak{p} \in \mathfrak{P}$, as it is non-constant it contains at least a \circ and a •. Since there is a finite number of pairs $\circ \prec \bullet$ (i.e. a pair k < l with $\mathfrak{p}_k = \circ$ and $\mathfrak{p}_l = \bullet$), the sequences is asymptotically constant to \circ near $+\infty$, and \bullet near $-\infty$.

Lemma 4.13. Let $i \ge 1$. There is a finite number of elements of \mathcal{P} with codegree smaller than i, and one has

$$\sum_{\mathfrak{p}\in \mathcal{P}} x^{\mathrm{codeg}(\mathfrak{p})} = p(x).$$

Proof. Let $\mathfrak{p} \in \mathfrak{P}$ be a sequence with codegree smaller than *i*. Choose the reindexation of \mathfrak{p} such that i is the last index whose value is \bullet . As $I(\mathfrak{p})$ is finite, there is a finite number of \circ before index i since each of them yields an inversion. Moreover, none can have negative index otherwise we would have the form

$$\mathfrak{p} = \cdots \bullet \bullet \bullet \cdots \circ [\cdots] \bullet \circ \circ \circ \cdots,$$

and each element in the bracketed zone yields an inversion, leading to more than i + 1 inversions. Thus the set $\{\mathfrak{p} \in \mathcal{P} \mid \operatorname{codeg}(\mathfrak{p}) \leq i\}$ is finite and the generating series is well-defined.

An element $\mathfrak{p} \in \mathcal{P}$ is fully determined by the sequence with finite support $u(\mathfrak{p}) = (u_i)_{i \ge 1}$, with u_j being the number of • with $j \circ$ on their left. The inverse bijection associates to an integer sequence with finite support u the element of \mathcal{P} defined as follows:

- * put a \circ at 0 and \bullet for negative indices,
- * inductively, starting at j = 1, put $u_j \bullet$ and then a new \circ ,
- * as u is of finite support, the algorithm finishes by only putting \circ .

The codegree expresses as

$$\operatorname{codeg}(\mathfrak{p}) = \sum_{j=1}^{\infty} j u_j$$

If $\operatorname{codeg}(\mathfrak{p}) \leq i$ then $u_i = 0$ for j > i. Computing the generating series modulo x^{i+1} , we only care about the p having the sequence u(p) with support in [1; i], and u_i may take any value considered that too large values will contribute 0 modulo x^{i+1} . Thus one has:

$$\sum_{\mathfrak{p}\in\mathfrak{P}} x^{\operatorname{codeg}(\mathfrak{p})} \equiv \sum_{u_1,\dots,u_i=0}^{\infty} x^{\sum ju_j} \mod x^{i+1}$$
$$\equiv \prod_{j=1}^i \left(\sum_{u_j=1}^{\infty} x^{ju_j}\right) = \prod_{j=1}^i \frac{1}{1-x^j} \mod x^{i+1}$$
$$\equiv \prod_{j=1}^{\infty} \frac{1}{1-x^j} = p(x) \mod x^{i+1}.$$

As the congruence is true modulo x^{i+1} for every *i*, we get the desired equality.

Example 4.14. Continuing Example 4.12 one has u(p) = (0, 1, 2, 1, 0, 0, ...).

Lemma 4.15. Let $W \in \mathcal{W}_{\beta}$ with $codeg(W) \leq i$ and $\beta \cdot D > 2i$ for any toric divisor D. Then the data of the sloping pairs (L, R) is equivalent to the data of an element $\mathfrak{p}_c \in \mathfrak{P}$ for any corner of Δ_{β} non-adjacent to a horizontal edge, such that $\operatorname{codeg}(L, R) = \sum_{c} \operatorname{codeg}(\mathfrak{p}_{c})$.

Proof. Let (L,R) be the tuple of sloping pairs of W, and let $\theta \leqslant p \leqslant \theta'$ be the integers such that the edges of the left side of Δ_{β} have outgoing normal vectors (-1, p). Point (i) of lemma 4.11 says that L writes as a concatenation $L = (L_{\theta}, \ldots, L_{\theta'-1})$ where L_p is of the form $(p,\ldots,p,\star,\ldots,\star,p+1,\ldots,p+1)$ with $\star \in \{p,p+1\}$. Given p, let c_p^- be the corner of Δ_β

25

 \Diamond

whose adjacent edges have outgoing normal vectors (-1, p) and (-1, p + 1). Replacing p by • and p+1 by \circ , the tuple L_p gives an element $\mathfrak{p}_{c_p^-} \in \mathcal{P}$. Similarly, R gives elements $\mathfrak{p}_{c_p^+} \in \mathcal{P}$. By construction, one has $\operatorname{codeg}(L, R) = \sum_c \operatorname{codeg}(\mathfrak{p}_c)$, where the sum runs over the corners of Δ_β non-adjacent to a horizontal edge.

Conversely, assume we are given a family $(\mathfrak{p}_c)_c \in \mathfrak{P}^{\chi-4}$. We construct L from the elements $\mathfrak{p}_{c_p^-}$ corresponding to corners of the left side of Δ_β in the following way. For any p, truncate $\mathfrak{p}_{c_p^-}$ just before its first \circ and just after its last \bullet . Replacing \bullet by p and \circ by p+1 gives a tuple \widetilde{L}_p . Then L is the concatenation $L = (\widetilde{L}_{\theta}, \ldots, p, p, \widetilde{L}_p, p+1, p+1, \ldots, L_{\theta'-1})$ where we add sufficiently enough p between \widetilde{L}_{p-1} and \widetilde{L}_p so that the total number of p is the number given by proposition 4.9 (iv). We proceed similarly for R, and by construction one has $\operatorname{codeg}(L, R) = \sum_c \operatorname{codeg}(\mathfrak{p}_c)$.

Example 4.16. To the tuple L = (0, 1, 0, 1, 1, 0, 1, 1, 1, 2, 1, 2, 2) we associate the sequences $\mathfrak{p}_1 = \cdots \bullet \bullet \circ \bullet \circ \circ \circ \circ \circ \circ \cdots$ and $\mathfrak{p}_2 = \cdots \bullet \bullet \circ \bullet \circ \circ \circ \cdots$, where \bullet and \circ correspond to 0 and 1 in \mathfrak{p}_1 (resp. 1 and 2 in \mathfrak{p}_2).

4.2.3. Enumeration of words in the h-transverse setting. We can now compute the asymptotic refined invariant in genus 0 for h-transverse polygons.

Theorem 4.17. Let X be a toric surface associated to a h-transverse, horizontal and nonsingular polygon, with Euler characteristic χ . Then the genus 0 asymptotic refined invariant is

$$AR_0^X = p(x)^{\chi}.$$

Proof. We can determine $AR_0^X \mod x^{i+1}$ by summing the multiplicities of the words of \mathscr{W}_β of codegree at most i, with $\beta \in H_2(X, \mathbb{Z})$ such that for every toric divisor D we have $\beta \cdot D > 2i$.

By Lemma 4.11 the weight of every bounded elevator in the diagram associated to a word $W \in \mathcal{W}_{\beta}$ of codegree at most *i* is strictly bigger than *i*. Hence the multiplicity modulo x^{i+1} is

$$(1-x)^{b^{\mathrm{top}}+b^{\mathrm{bot}}}x^{\mathrm{codeg}(\mathsf{W})}$$

The word is fully determined by the following data:

- ▷ an element $\mathfrak{t} \in S(b^{\text{top}})$ encoding the *T*-words,
- \triangleright an element $\mathfrak{b} \in S(b^{\text{bot}})$ encoding the *B*-words,
- ▷ an element $\mathfrak{p}_c \in \mathcal{P}$ for any of the $\chi 4$ corners c of Δ non-adjacent to a horizontal side,

such that

$$\operatorname{codeg}(\mathsf{W}) = \operatorname{codeg}(\mathfrak{t}) + \operatorname{codeg}(\mathfrak{b}) + \sum_{c} \operatorname{codeg}(\mathfrak{p}_{c}) \leqslant i.$$

The data of t and b are enough to recover the word up to the indices of the letters $f_{*,*}$. The data of the \mathfrak{p}_c allows to recover the sloping pairs (L, R) by lemma 4.15. Hence, summing over $\mathfrak{S}(b^{\mathrm{bot}}) \times \mathfrak{S}(b^{\mathrm{top}}) \times \mathfrak{P}^{\chi-4}$ (and potentially counting terms which contribute 0 modulo x^{i+1}) the generating series of words counted with multiplicity factors modulo x^{i+1} :

$$\left((1-x)^{b^{\mathrm{bot}}} \sum_{\mathfrak{b} \in \mathcal{S}(b^{\mathrm{bot}})} x^{\mathrm{codeg}(\mathfrak{b})} \right) \left((1-x)^{b^{\mathrm{top}}} \sum_{\mathfrak{t} \in \mathcal{S}(b^{\mathrm{top}})} x^{\mathrm{codeg}(\mathfrak{t})} \right) \left(\sum_{\mathfrak{p} \in \mathcal{P}} x^{\mathrm{codeg}(\mathfrak{p})} \right)^{\chi-4}$$

Using Lemmas 4.7 and 4.13 we obtain for the generating series

$$p(x)^2 \cdot p(x)^2 \cdot p(x)^{\chi - 4} = p(x)^{\chi} \mod x^{i+1}.$$

As this is true for every $i \ge 1$ we get the result.

5. Asymptotic refined invariant in genus 1

The idea to compute the genus 1 asymptotic invariant is to construct floor diagrams of genus 1 by adding an edge to a genus 0 diagram. This way, we can group together the genus 1 diagrams obtained from the same genus 0 diagram, so that we reduce the enumeration to the genus 0 case, with a multiplicity corresponding to the weighted count of diagrams. We start with Hirzebruch surfaces before going to *h*-transverse, horizontal and non-singular toric surfaces. The strategy is the same: we compute $\widetilde{BG}_1^X(\beta)$ modulo x^{i+1} and find an expression that does not depend on *i*, before making *i* goes to ∞ .

5.1. The case of Hirzebruch surfaces. To get to the genus 1 case, the idea is that a genus 1 diagram is obtained from a genus 0 diagram by adding one edge, and conversely we get a genus 0 diagram by removing an edge from a genus 1 diagram. However, it might not be clear which edge to remove, and what to do to balance the diagram again. We make this construction precise by introducing the notion of *nerved diagram*.

5.1.1. Nerved diagrams. We already fixed an integer i to bound the codegree of diagrams we look at. Let us fix a second integer $M \ge 1$.

Definition 5.1. Let \mathcal{D} be a genus g diagram in a class aE + bF, with a > 2i and b > (g+1)M + i. Assume $\operatorname{codeg}(\mathcal{D}) \leq i$. A *nerve* for \mathcal{D} is the choice of an edge between each pair of consecutive floors with weight $\geq M$. We call the data of \mathcal{D} with the choice of a nerve a *nerved diagram*. We denote with a tilde the nerved diagrams, e.g. $\widetilde{\mathcal{D}}$.

Remark 5.2. For genus g, provided b > (g+1)M + i and $codeg(\mathcal{D}) \leq i$, Lemma 2.14 ensures the existence of a nerve.

Lemma 5.3. Assume b > i+2M and let \mathcal{D} be a diagram in the class aE+bF with $codeg(\mathcal{D}) \leq i$.

- (i) If \mathcal{D} is of genus 0, there exists a unique choice of nerve.
- (ii) If \mathcal{D} is of genus 1 with an edge skipping some floors, there exists a unique choice of nerve.
- (iii) If \mathcal{D} is of genus 1 with two edges linking consecutive floors, there are one or two possible nerves depending on whether only one of the edges or both have weight bigger than M.
- *Proof.* (i) In the genus 0 case, we already know by [BJP22] that the floors are totally ordered in the diagram. The total weight between the floors m and m + 1 is $b + \delta m$ minus the number of sinks that skip the floor m + 1 and the number of sources that skip the floor m. As the number of ends skipping some floors is bounded by i, the weight of the unique edge between two consecutive floors is bigger than $b i \ge M$, so that there is a unique nerve.
 - (ii) Because $\operatorname{codeg}(\mathcal{D}) \leq i$, the weight of the skipping edge is bounded by *i* and we conclude as in the genus 0 case.
 - (iii) The sum of the weights of the two edges is bigger than b i > 2M, so that at least one of them has weight $\ge M$.

Example 5.4. Assume we chose M = 1, so that there are no condition on the weight of the edges on the nerve. On Figure 6 we depict three nerved diagrams, the nerve consists in thickened edges. The first nerved diagram is the unique nerved diagram associated to the



FIGURE 6. Nerved diagrams of genus 0 and 1.

underlying genus 0 diagram. The remaining two nerved diagrams have the same underlying genus 1 diagram. If we had taken M = 2, only one of the two edges between the second and third floor could have been chosen in the nerve.

We assign to each nerved diagram a multiplicity so that the count of nerved diagram matches the count of diagrams.

Definition 5.5. Let \mathcal{D} be diagram of genus g in the class aE+bF and assume b > i+(g+1)M. Let $N(\mathcal{D})$ be the number of nerves of \mathcal{D} . The *multiplicity* of a nerved diagram $\widetilde{\mathcal{D}}$ is $\mu(\widetilde{\mathcal{D}}) = \frac{1}{N(\mathcal{D})}\mu(\mathcal{D})$.

Remark 5.6. Forgetting about the ends of the diagram, a nerve is a spanning tree of the underlying graph so that there are g bounded edges not belonging to the nerve.

5.1.2. Constructing genus 1 nerved diagrams from genus 0 ones. Let $\widetilde{\mathscr{D}}_g$ be the set of nerved marked diagram of genus g in the class aE + bF. Assume b > i + 2M. We have a map

$$\mathrm{ft}:\widetilde{\mathscr{D}}_1\to\widetilde{\mathscr{D}}_0$$

that forgets the unique bounded edge e not on the nerve and add w(e) to the weights of all the edges between the two vertices to which e was attached. Conversely, we can construct a genus 1 nerved marked diagram from a genus 0 one by adding an edge e with weight w, and removing w to the weights of all the edges between the two vertices to which e is attached. This is possible if we are provided with the weight w of the added edge, the place of its marking between two floors m and m + 1, and the floors it is attached to, encoded by a pair (s_+, s_-) that are the numbers of floors it skips above and below its marking. This data is subject to the following constraints:

$$\triangleright s_{-} \leqslant m-1 \text{ and } s_{+} \leqslant a-m-1,$$

▷ $w \leq \min(w(e)) - M$, where the minimum is over the weights of the edges of the nerve between the floors $m - s_-$ and $m + 1 + s_+$, so that the weights of the nerves are still $\geq M$.



FIGURE 7. On the left a genus 0 nerved marked diagram. On the right, various genus 1 nerved marked diagrams that we can obtain by adding an edge of weight 1 with a marking between the second and third floor.

Example 5.7. Assume M = 2. On Figure 7 we depicted various ways to get a genus 1 nerved marked diagram by adding a dashed edge of weight 1 to \tilde{D}_0 . If we have $(s_+, s_-) = (1, 1)$, we get \tilde{D}_1 because the edge skips one floor above its marking, and one below. Taking (1, 0) or (0, 1) instead, we get \tilde{D}_2 and \tilde{D}_3 . If $s_+ = s_- = 0$, we get \tilde{D}_4 . In all these examples we choose the marking of the added edge to be between the one of the second floor and the one of the bounded edge between the second and third floor.

Let us try to increase the weight w of the dashed edge. For $\tilde{\mathcal{D}}_1, \tilde{\mathcal{D}}_2, \tilde{\mathcal{D}}_3, w$ can also be set equal to 2, but not 3 since one of the edges on the nerve would get weight 1 < M. For $\tilde{\mathcal{D}}_4$, we can take w = 2 or 3, and in that case the underlying diagram has two possible nerves. \Diamond

We now relate the multiplicity of a nerved diagram constructed by the above process to the multiplicity of the initial genus 0 diagram.

Lemma 5.8. Assume M > i and b > i + 2M. Let $\widetilde{\mathbb{D}}$ be a genus 0 nerved marked diagram with $\operatorname{codeg}(\widetilde{\mathbb{D}}) \leq i$ in the class aE + bF, and let $\widetilde{\mathcal{E}}$ be the genus 1 marked nerved diagram constructed by the data of the position of the marking, weight w and (s_+, s_-) . Then we have

$$\mu(\widetilde{\mathcal{E}}) = \frac{1}{1 + \mathbb{1}_{w \ge M}} (1 - x^w)^2 x^{w(s_+ + s_-)} \mu(\widetilde{\mathcal{D}}) \mod x^{i+1}.$$

Proof. If $w \ge M$ then $s_+ = s_- = 0$, otherwise the codegree would be greater than M, and also *i*. Hence in that case the added edge links two consecutives floors and, there are two possible nerves. If w < M, there is a unique nerve. Hence one has $N(\mathcal{E}) = 1 + \mathbb{1}_{w \ge M}$.

By Lemma 2.14, the hypothesis ensures that the sum of weights between consecutive floors in \mathcal{D} is bigger than 2*M*. Thus, the only edge potentially contributing to the multiplicity of \mathcal{E} is the one we add, yielding a factor $(1 - x^w)^2$. The codegree this edge provides is $w(s_+ + s_-)$ since it has weight *w* and skips exactly $s_+ + s_-$ floors. As the weights of the edges in the nerve are still bigger than *M* after we added the new edge, they still do not contribute to the multiplicity modulo x^{i+1} .

Conversely, we can add the multiplicities of the genus 1 nerved marked diagrams constructed from a genus 0 nerved marked diagram. Let $\langle m \rangle = \frac{x^m}{1-x^m}$.

Lemma 5.9. Let $\widetilde{\mathbb{D}}$ be a genus 0 nerved marked diagram and let $1 \leq m \leq a - 1$. Assume M > i, a > 2i and b > i + 2M. Let pos_m be the number of positions where to insert a marking between the floors m and m + 1. Let $\widetilde{\omega}_m$ be the weight of the edge between these floors. The sum of multiplicities of genus 1 nerved marked diagrams obtained by inserting an edge with a marking between these floors is

$$ext{pos}_m \cdot \left(rac{\widetilde{\omega}_m - 1}{2} - d_m
ight) \mu(\widetilde{\mathbb{D}}), \, \, \textit{where} \, \, d_m = \left\{ egin{array}{cc} \langle m
angle & \textit{if} \ m \leqslant i, \ \langle a - m
angle & \textit{if} \ m \geqslant a - i, \ 0 & \textit{else.} \end{array}
ight.$$

Proof. We first choose one of the pos_m possible positions for the marking. We then sum over the possible choices of w, s_{\pm} . There are two possibilities.

- ▷ If $s_+ + s_- > 0$, we can assume the weight w is bounded by i since otherwise, we get multiplicity 0 modulo x^{i+1} .
- ▷ If $s_+ = s_- = 0$, the weight w may takes values from 1 to $\tilde{\omega}_m M$, since the nerve has to keep a weight bigger than M. Furthermore, we start having a factor $\frac{1}{2}$ for the choices of nerves when $w \ge M$. For such a w, we have $(1 x^w)^2 \equiv 1 \mod x^{i+1}$ since $M > 2i \ge i$.

Thus by Lemma 5.8 we have to compute the following:

$$\sum_{w=1}^{i} \sum_{s_{+}+s_{-}>0} (1-x^{w})^{2} x^{w(s_{+}+s_{-})} + \sum_{w=1}^{M-1} (1-x^{w})^{2} + \sum_{w=M}^{\widetilde{\omega}_{m}-M} \frac{1}{2} \mod x^{i+1}.$$

To compute the first sum, we may add the values for w going from i+1 to infinity since they contribute 0 modulo x^{i+1} . If $m \leq i$ we have the bound $s_{-} \leq m-1$, but s_{+} can goes to ∞ since the excess terms contribute 0 modulo x^{i+1} . In that case we get for the first sum

$$\sum_{w=1}^{\infty} (1-x^w)^2 \left(\frac{1-x^{mw}}{(1-x^w)^2} - 1\right) = \sum_{w=1}^{\infty} \left[1 - (1-x^w)^2 - x^{mw}\right]$$
$$= \sum_{w=1}^{\infty} \left[1 - (1-x^w)^2\right] - \langle m \rangle.$$

If $m \ge a-i$ we have the bound $s_+ \le a-m-1$, but s_- can goes to ∞ and the first sum gives

$$\sum_{w=1}^{\infty} \left[1 - (1 - x^w)^2 \right] - \langle a - m \rangle.$$

If i < m < a - i then both s_{-} and s_{+} can go to ∞ so the first sum is

$$\sum_{w=1}^{\infty} \left[1 - (1 - x^w)^2 \right].$$

The others two sums are

$$\sum_{w=1}^{M-1} (1-x^w)^2 + \sum_{w=M}^{\widetilde{\omega}_m - M} \frac{1}{2} = \sum_{w=1}^{M-1} \left[(1-x^w)^2 - 1 \right] + M - 1 + \frac{\widetilde{\omega}_m - M - M + 1}{2}$$
$$\equiv \sum_{w=1}^{\infty} \left[(1-x^w)^2 - 1 \right] + \frac{\widetilde{\omega}_m - 1}{2} \mod x^{i+1}.$$

Putting all sums together, the two sums over w cancel and we get the result.

5.1.3. Integration over the space of genus 0 diagrams. In the computation of the genus 0 asymptotic refined invariants, we encoded marked diagrams with words and proved that the set of words is in bijection with a subset of $S(b) \times S(b + \delta a)$. Elements of S(n) were assigned multiplicities

$$\mu_{\mathbb{S}(n)}(\mathfrak{s}) = (1-x)^n x^{\operatorname{codeg}(\mathfrak{s})}.$$

Recall that we have maps $\ell_0, \ell_j^{(k)} : \mathbb{S} \to \mathbb{Z}_{\geq 0}$ that give the lengths of the words of a sentence. Let \mathcal{L} be the lengths space, i.e. the space of non-negative integer sequences $(l_j^{(k)})_{j,k}$ with finite support, and π be the map $\pi = (\ell_j^{(k)})_{j,k} : \mathbb{S}(n) \to \mathcal{L}$ that maps a sentence to the lengths of its words except the first one. To each element $\mathbf{l} = (l_j^{(k)})_{j,k} \in \mathcal{L}$, we assign a weight $\mu_{\mathcal{L}}(\mathbf{l}) = \prod_{j,k} x^{j l_j^{(k)}}$.

Formally, it is possible to see $\mu_{\mathcal{S}(n)}$ and $\mu_{\mathcal{L}}$ as measures on their corresponding domain, which are discrete spaces. These measures have values in the quotient ring $\mathbb{Z}[x]/(x^{i+1})$ for our choice of *i*. From this point of view, weighted sums become integrals. Moreover, this integral is $\mathbb{Z}[x]/(x^{i+1})$ -linear. There are several reasons for such a consideration: it shortens notations, it becomes easier to see some computational steps, and it formalizes the deletion of diagrams with zero weight. The idea to compute the asymptotic refined invariant in genus 1 is now to integrate the function given by Lemma 5.9 on the space of genus 0 diagrams.

Lemma 4.7 states that $\mu_{\mathbb{S}(n)}$ and $\mu_{\mathcal{L}}$ have total weight $p(x)^2$, so that we may consider the normalized measures $\nu_{\mathbb{S}(n)} = \frac{1}{p(x)^2} \mu_{\mathbb{S}(n)}$ and $\nu_{\mathcal{L}} = \frac{1}{p(x)^2} \mu_{\mathcal{L}}$. For product spaces, we consider the product measures. During the proof of Lemma 4.7, we have

$$\sum_{\substack{\mathfrak{s}\in\mathbb{S}(n)\\\pi(\mathfrak{s})=\mathbf{l}}}\mu_{\mathbb{S}(n)}(\mathfrak{s})=\mu_{\mathbb{S}(n)}(\pi^{-1}(\mathbf{l}))=\prod_{j,k}x^{jl_j^{(k)}}=\mu_{\mathcal{L}}(\mathbf{l}).$$

5.1.4. Some integral computations. Before going through the main computation, we introduce some functions on \mathcal{L} and $\mathcal{S}(n)$, and compute their integrals against the normalized measures. Consider first the lengths functions $\ell_j^{(k)}$, which are the coordinate functions on \mathcal{L} .

Lemma 5.10. We have the following integrals:

$$\int_{\mathcal{L}} \ell_m^{(r)} \mathrm{d}
u_{\mathcal{L}} = \langle m
angle, \; \int_{\mathcal{L}} (\ell_m^{(r)})^2 \mathrm{d}
u_{\mathcal{L}} = \langle m
angle + 2 \langle m
angle^2$$

with $\langle m \rangle = rac{x^m}{1-x^m}.$

Proof. Indeed, by definition, we have

$$\int_{\mathcal{L}} \ell_m^{(r)} \mathrm{d}\nu_{\mathcal{L}} = \frac{1}{p(x)^2} \sum_{\mathbf{l} \in \mathcal{L}} l_m^{(r)} \prod_{j,k} x^{j l_j^{(k)}} = \frac{1}{p(x)^2} \left(\sum_{l_m^{(r)} = 0}^{\infty} l_m^{(r)} x^{m l_m^{(r)}} \right) \prod_{(j,k) \neq (m,r)} \left(\sum_{l_j^{(k)} = 0}^{\infty} x^{j l_j^{(k)}} \right).$$

31

We then use the identity $\sum_{\alpha=0}^{\infty} \alpha y^{\alpha} = \frac{y}{(1-y)^2}$. For the second integral, we use $\sum_{\alpha=0}^{\infty} \alpha^2 y^{\alpha} = \frac{y+y^2}{(1-y)^3}$.

In fact, this method, which is an analog of Fubini's theorem, works for computing the integral of any monomial in the $\ell_j^{(k)}$: the integral of a monomial is equal to the product of integrals over each of the variables appering in the monomial. Hence, it reduces down to the computation of the sums $\sum_{\alpha=0}^{\infty} \alpha^r y^{\alpha}$.

We then set $\ell_m = \ell_m^{(1)} + \ell_m^{(2)}$, so that we now have

$$\int_{\mathcal{L}}\ell_m\mathrm{d}
u_{\mathcal{L}}=2\langle m
angle ext{ and } \int_{\mathcal{L}}\ell_m^2\mathrm{d}
u_{\mathcal{L}}=2\langle m
angle+6\langle m
angle^2.$$

In particular, the following affine function

$$e_m = (1 - x^m) \frac{\ell_m + 2}{2}$$

defined on \mathcal{L} has integral equal to 1.

By composing with $\pi : S(n) \to \mathcal{L}$, it is possible to pull-back functions on \mathcal{L} to get functions on S(n). Due to the normalization by the total weight, their integrals are preserved.

Definition 5.11. We define on S(n) the *leak function* $\phi_m[n](\mathfrak{s})$ equal to the number of letters with an index bigger than m. To get a function of $\mathbf{l} \in \mathcal{L}$, we average over the set $\pi^{-1}(\mathbf{l})$ of sentences with lengths \mathbf{l} :

$$arphi_m[n](\mathbf{l}) = rac{1}{1-x^m} rac{1}{\mu_\mathcal{L}(\mathbf{l})} \int_{\pi^{-1}(\mathbf{l})} \phi_m[n] \mathrm{d} \mu_{\mathbb{S}(n)}.$$

Lemma 5.13 expresses the function $\varphi_m[n](\mathbf{l})$ in terms of the monomials ℓ_j on \mathcal{L} .

Remark 5.12. On the diagram side, the leak function ϕ_m corresponds to the number of ends skipping the floor m. It is also equal to $\omega_m - \tilde{\omega}_m$, which is the complement of the weight between the floors m and m + 1 to the maximal possible weight $\omega_m = b + \delta m$.

Lemma 5.13. We have the following expressions on \mathcal{L} :

Proof. Let $\mathbf{l} \in \mathcal{L}$ and $\mathfrak{s} = (S_0, (S_j^{(k)})_{j,k}) \in \pi^{-1}(\mathbf{l})$ be a sentence. In terms of the letters, the leak function $\phi_m[n]$ is

$$\phi_m[n](\mathfrak{s}) = \sum_{\mathsf{s}\in\mathsf{S}_0}\mathbbm{1}(p\geqslant m ext{ with } \mathsf{s}=\mathsf{s}_p) + \sum_{j,k}\sum_{\mathsf{s}\in\mathsf{S}_j^{(k)}}\mathbbm{1}(p\geqslant m ext{ with } \mathsf{s}=\mathsf{s}_p).$$

Indeed, the leak is due to the ends that skip the floor m, i.e. the letters s_p with an index $p \ge m$. We need to compute $\frac{1}{\mu_{\mathcal{L}}(\mathbf{l})} \int_{\pi^{-1}(\mathbf{l})} \mathbb{1}(p \ge m \text{ with } \mathbf{s} = \mathbf{s}_p) d\mu_{\mathbb{S}(n)}$, for each term $\mathbb{1}(p \ge m \text{ with } \mathbf{s} = \mathbf{s}_p)$ corresponding to a position of the letter \mathbf{s} in the word $\mathbf{S}_j^{(k)}$. To do so, we proceed as in Lemma 4.7. At each letter position \mathbf{s}' in $\mathbf{S}_{j'}^{(k')}$ except the one corresponding to \mathbf{s} , the sum over the possible values of the letter is the geometric series

$$\sum_{p=j'}^{\infty} x^p = \frac{x^{j'}}{1-x}.$$

For the position corresponding to s, because of the condition $(p \ge m \text{ with } s = s_p)$ we have instead

$$\sum_{p=j}^{\infty} \mathbb{1}(p \ge m) x^p = \frac{x^{\max(j,m)}}{1-x} = x^{(m-j)_+} \frac{x^j}{1-x}$$

where $(m-j)_+ = \max(m-j, 0)$. As in Lemma 4.7, we conclude by making the product over all letter positions and we get

. . .

$$\int_{\pi^{-1}(\mathbf{l})} \mathbb{1}(p \geqslant m \text{ with } \mathsf{s} = \mathsf{s}_p) \mathrm{d}\mu_{\mathbb{S}(n)} = x^{(m-j)_+} \prod_{j',k'} x^{j' l_{j'}^{(k')}} = x^{(m-j)_+} \mu_{\mathcal{L}}(\mathbf{l}).$$

Adding the above over all the letter positions in the word, we get

$$\begin{aligned} \frac{1}{\mu_{\mathcal{L}}(\mathbf{l})} \int_{\pi^{-1}(\mathbf{l})} \phi_m[n] \mathrm{d}\nu_{\mathcal{S}(n)} &= \ell_0 x^m + \sum_{j=1}^m (\ell_j^{(1)} + \ell_j^{(2)}) x^{m-j} + \sum_{j=m+1}^\infty (\ell_j^{(1)} + \ell_j^{(2)}) \\ &= \left(n - \sum_{j=1}^\infty \ell_j\right) x^m + \sum_{j=1}^m \ell_j x^{m-j} + \sum_{j=m+1}^\infty \ell_j \\ &= n x^m + \sum_{j=1}^m \ell_j (x^{m-j} - x^m) + (1 - x^m) \sum_{j=m+1}^\infty \ell_j \\ &= (1 - x^m) \left[n \langle m \rangle + \langle m \rangle \sum_{j=1}^m \frac{\ell_j}{\langle j \rangle} + \sum_{j=m+1}^\infty \ell_j \right]. \end{aligned}$$

Lemma 5.14. We have the following integral:

$$\int_{\mathcal{L}} e_m \psi_m \mathrm{d}
u_{\mathcal{L}} = (2m+1) \langle m
angle + 2 \sum_{j=m+1}^\infty \langle j
angle.$$

Proof. We use the expression of ψ_m in terms of the ℓ_j , and the following computations:

$$\int_{\mathcal{L}} e_m \ell_j \mathrm{d}\nu_{\mathcal{L}} = \begin{cases} 2\langle j \rangle & \text{if } j \neq m, \\ 3\langle m \rangle & \text{if } j = m. \end{cases}$$

Hence, $\int e_m \ell_j = \int \ell_j$ except for j = m, where we add $\langle m \rangle$. Thus, we have

$$\begin{split} \int_{\mathcal{L}} e_m \psi_m &= \int_{\mathcal{L}} \psi_m + \langle m \rangle \\ &= \langle m \rangle \sum_{j=1}^m \frac{2\langle j \rangle}{\langle j \rangle} + \sum_{m+1}^\infty 2\langle j \rangle + \langle m \rangle \\ &= (2m+1)\langle m \rangle + 2 \sum_{j=m+1}^\infty \langle j \rangle. \end{split}$$

So far, we defined functions on S(n). The genus 0 marked diagrams of codegree smaller than *i* are in bijection with a subset of $S(b) \times S(b + \delta a)$. By definition, the complement of this subset has measure 0 since it consists of elements with codegree strictly bigger than *i*. Let ρ_1 , ρ_2 be the projections of $S(b) \times S(b + \delta a)$ to S(b) and $S(b + \delta a)$. We can thus pull-back functions by ρ_1 and ρ_2 and obtain the following functions.

33

- \triangleright The number pos_m of positions for a marking between the floors m and m+1 is:
 - equal to $\rho_1^* \ell_m + 2 = \rho_1^* \ell_m^{(1)} + \rho_1^* \ell_m^{(2)} + 2$ if $m \leq i$, where ℓ_m is pull-back from S(b), ◦ equal to 2 if i < m < a - i, since the length functions are 0,
 - equal to $\rho_2^* \ell_{a-m} + 2$ if $m \ge a i$, where ℓ_{a-m} is now pull-back from $S(b + \delta a)$ instead of S(b).
- ▷ We have the same phenomenon for the leak function on a diagram: for $m \leq i$, it is the pull-back of the leak function on S(b), then it is 0 for i < m < a i, and gets pulled-back from $S(b + \delta a)$ for $m \geq a i$.

5.1.5. Computation of the asymptotic refined invariant. For n a positive integer, we consider the function $\sigma_1(n) = \sum_{d|n} d$, and its generating series

$$E_2(x) = \sum_{n \ge 1} \sigma_1(n) x^n.$$

Lemma 5.15. One has

$$E_2(x) = \sum_{n=1}^{\infty} n \frac{x^n}{1-x^n} = \sum_{n=1}^{\infty} \frac{x^n}{(1-x^n)^2} = \sum_{n=1}^{\infty} \sum_{j=n}^{\infty} \frac{x^j}{1-x^j}.$$

Proof. Expanding $\frac{1}{1-x^n}$ yields the first two expressions for $E_2(x)$. The last expression yields the first one when switching the sums over n and j.

Theorem 5.16. The genus 1 asymptotic refined invariant of Hirzebruch surfaces is given by

$$AR_1^{\mathbb{F}_\delta} = p(x)^4 \left(g_{\max} - 12E_2(x)
ight)$$

where $g_{\text{max}} = \frac{1}{2}(a-1)(2b+\delta a-2)$ is the genus of a smooth curve in the class aE + bF.

Proof. The computation of the asymptotic refined invariant goes through three steps: expressing then integrating over $S(b) \times S(b + \delta a)$ the function from Lemma 5.9, and summing these integrals over m from 1 to a - 1.

• First step: expression over $S(b) \times S(b + \delta a)$. By Lemma 5.9, the function giving the sum of multiplicities for insertion of a marking between the floors m and m + 1 is

$$\operatorname{pos}_{m}\left(\frac{\widetilde{\omega}_{m}-1}{2}-d_{m}\right) = \begin{cases} \left(\rho_{1}^{*}\ell_{m}+2\right)\left(\frac{\omega_{m}-\rho_{1}^{*}\phi_{m}[b]-1}{2}-\langle m\rangle\right) & \text{if } m \leqslant i, \\ \omega_{m}-1 & \text{if } i < m < a-i \\ \left(\rho_{2}^{*}\ell_{a-m}+2\right)\left(\frac{\omega_{m}-\rho_{2}^{*}\phi_{a-m}[b+\delta a]-1}{2}-\langle a-m\rangle\right) & \text{if } m \geqslant a-i, \end{cases}$$

where in the first (resp. last) row, functions are pull-back from S(b) (resp. $S(b + \delta a)$). For each value of m, we now need to integrate the above function, and then sum over $1 \leq m \leq a$.

• Second step: integration over $S(b) \times S(b + \delta a)$. If i < m < a - i, we have

$$\int_{\mathbb{S}(b)\times\mathbb{S}(b+\delta a)} (\omega_m - 1) \mathrm{d}\nu = \omega_m - 1.$$

Assume now that $m \leq i$. Since $\int_{S(b+\delta a)} 1 = 1$ we have

$$\int_{\mathbb{S}(b)\times\mathbb{S}(b+\delta a)} (\rho_1^*\ell_m+2) \left(\frac{\omega_m - \rho_1^*\phi_m[b] - 1}{2} - \langle m \rangle\right) = \int_{\mathbb{S}(b)} (\ell_m+2) \left(\frac{\omega_m - \phi_m[b] - 1}{2} - \langle m \rangle\right).$$

Recall that we set $e_m = (1 - x^m) \frac{\ell_m + 2}{2}$ so the integrand rewrites

$$e_m(\omega_m-1)+e_m\left((\omega_m-1)\langle m
angle-rac{\phi_m[b]}{1-x^m}-2rac{x^m}{(1-x^m)^2}
ight)$$

To compute the integral over S(b), we first regroup over each $\pi^{-1}(\mathbf{l})$ considering $\frac{1}{\mu_{\mathcal{L}}(\mathbf{l})} \int_{\pi^{-1}(\mathbf{l})}$. This way, we get a function to integrate over \mathcal{L} . This function is

$$e_m(\omega_m-1)+e_m\left((\omega_m-1)\langle m
angle-arphi_m[b]-2rac{x^m}{(1-x^m)^2}
ight).$$

Because $\int_{\mathcal{L}} e_m = 1$ we have $\int_{\mathcal{L}} e_m(\omega_m - 1) = \omega_m - 1$. It remains to compute the integral of the correction term $e_m\left((\omega_m - 1)\langle m \rangle - \varphi_m[b] - 2\frac{x^m}{(1-x^m)^2}\right)$. As m is close to 1 one has:

$$\varphi_m[b] = b\langle m
angle + \psi_m \text{ and } \omega_m = b + \delta m.$$

We finally get

$$\begin{split} &\int_{\mathcal{L}} e_m \left((\omega_m - 1) \langle m \rangle - \varphi_m[b] - 2 \frac{x^m}{(1 - x^m)^2} \right) d\nu_{\mathcal{L}} \\ &= \int_{\mathcal{L}} e_m \left((b + \delta m - 1) \langle m \rangle - b \langle m \rangle - \psi_m - 2 \frac{x^m}{(1 - x^m)^2} \right) d\nu_{\mathcal{L}} \\ &= \int_{\mathcal{L}} e_m \left((\delta m - 1) \langle m \rangle - \psi_m - 2 \frac{x^m}{(1 - x^m)^2} \right) d\nu_{\mathcal{L}} \\ &= (\delta m - 1) \langle m \rangle - (2m + 1) \langle m \rangle - 2 \sum_{j=m+1}^{\infty} \langle j \rangle - 2 \frac{x^m}{(1 - x^m)^2} \\ &= (\delta - 2) m \frac{x^m}{1 - x^m} - 2 \sum_{j=m}^{\infty} \langle j \rangle - 2 \frac{x^m}{(1 - x^m)^2} \end{split}$$

i.e.

$$\int_{\mathbb{S}(b)\times\mathbb{S}(b+\delta a)} \text{pos}_m\left(\frac{\widetilde{\omega}_m - 1}{2} - d_m\right) = \omega_m - 1 + (\delta - 2)m\frac{x^m}{1 - x^m} - 2\sum_{j=m}^{\infty} \langle j \rangle - 2\frac{x^m}{(1 - x^m)^2}.$$

If $m \ge a - i$, with m' = a - m similar computations lead to

$$\int_{\mathbb{S}(b)\times\mathbb{S}(b+\delta a)} \text{pos}_m\left(\frac{\widetilde{\omega}_m - 1}{2} - d_m\right) = \omega_m - 1 - (\delta + 2)m'\frac{x^{m'}}{1 - x^{m'}} - 2\sum_{j=m'}^{\infty} \langle j \rangle - 2\frac{x^{m'}}{(1 - x^{m'})^2}$$

- \circ Third step: summation over the values of m. We have several sums to compute.
 - \triangleright W hatever the value of m is, the term ω_m-1 appears. We need to sum these terms, and one has

$$\sum_{m=1}^{a-1} (\omega_m - 1) = g_{\max},$$

since it is the number of interior lattice points of the associated Newton polygon.

▷ We have to sum the correction terms for $1 \leq m \leq i$. Since the formula for the correction term gives 0 modulo x^{i+1} when m > i, we let m goes to ∞. By Lemma 5.15, the sum of the correction terms is

$$(\delta - 2)E_2(x) - 2E_2(x) - 2E_2(x) = (\delta - 6)E_2(x)$$

▷ For the correction terms for $a - i \leq m \leq a - 1$, with m' = a - m we sum over m' going from 1 to ∞ and get

$$-(\delta + 6)E_2(x).$$

Adding the three contributions, we obtained $g_{\text{max}} - 12E_2(x)$. Multiplying by the total weight of the space $p(x)^4$ finishes the computation.

5.2. The case of *h*-transverse toric surfaces. The computations made in the Hirzebruch case remain valid with two differences. First, we now need to take into account the sloping pairs to the floors. The marked diagrams of genus 0 and codegree at most *i* are in bijection with a subset of $S(b^{\text{bot}}) \times S(b^{\text{top}}) \times \mathcal{P}^{\chi-4}$, where χ is the number of corners of the polygon and \mathcal{P} , defined in Section 4.2.2, encodes the default of increasingness of the slopes. Second, the self-intersection of the divisors corresponding to the top and bottom horizontal sides, equal to δ and $-\delta$ in the Hirzebruch case, are not opposite anymore, see Lemma 2.5.

Lemma 5.17. We have the following generating series:

$$\sum_{\mathfrak{p}\in\mathfrak{P}}\mathrm{codeg}(\mathfrak{p})x^{\mathrm{codeg}(\mathfrak{p})}=E_2(x)p(x).$$

Proof. We computed in Lemma 4.13 the generating series of $x^{\text{codeg}(p)}$, so we just need to differentiate the relation, multiply by x and use Lemma 5.15:

$$x\frac{d}{dx}\prod_{j=1}^{\infty}\frac{1}{1-x^{j}} = \sum_{m=1}^{\infty}m\frac{x^{m}}{(1-x^{m})^{2}}\prod_{j\neq m}\frac{1}{1-x^{j}}$$
$$= \left(\sum_{m=1}^{\infty}m\frac{x^{m}}{1-x^{m}}\right)\prod_{j=1}^{\infty}\frac{1}{1-x^{j}}$$
$$= E_{2}(x)p(x).$$

Theorem 5.18. Let X be a toric surface with Euler characteristic χ associated to an htransverse, horizontal and non-singular polygon. Let $g_{\max} = 1 + \frac{\beta^2 + K_X \cdot \beta}{2}$ be the polynomial function on $H_2(X,\mathbb{Z})$ that gives the genus of a smooth curve in the class β . The genus 1 asymptotic refined invariant is given by

$$AR_1^X = p(x)^{\chi} \left(g_{\max} - 12E_2(x) \right)$$

Proof. We proceed as in the Hirzebruch case. We assume each $\beta \cdot D$ is large enough for any toric divisor D. Consider $\mathfrak{P} = (\mathfrak{p}_c)_c \in \mathfrak{P}^{\chi-4}$ where we choose an element of \mathfrak{P} for each corner non-adjacent to a horizontal side. For a given \mathfrak{P} , let $\omega_m^{\mathfrak{P}}$ be the maximum weight between the floors m and m + 1 in a diagram obtained with the choice of sloping pairs determined by \mathfrak{P} . It differs from the total weight in the codegree 0 case ω_m in the following way: any element $\mathfrak{p} \in \mathfrak{P}$ is the product of exactly codeg(\mathfrak{p}) transpositions, and each of them reduces the weight at the position of the transposition by 1. So one has

$$\begin{split} \sum_{m=1}^{a-1} \left(\omega_m^{\mathfrak{P}} - 1 \right) &= \sum_{m=1}^{a-1} \left(\omega_m - 1 \right) - \sum_c \operatorname{codeg}(\mathfrak{p}_c) \\ &= g_{\max} - \sum_c \operatorname{codeg}(\mathfrak{p}_c), \end{split}$$

where the sum is indexed by the corners non-adjacent to a horizontal side.

Let δ_{top} (resp. δ_{bot}) be minus the self-intersection of the top (resp. bottom) toric divisor. For a fixed choice of $\mathfrak{P} = (\mathfrak{p}_c)_c \in \mathfrak{P}^{\chi-4}$, the contribution of \mathfrak{P} to the asymptotic refined invariant is computed as in the Hirzebruch case and is

$$x^{\sum_{c} \operatorname{codeg}(\mathfrak{p}_{c})} p(x)^{4} \left[\sum_{m=1}^{a-1} \left(\omega_{m}^{\mathfrak{P}} - 1 \right) - (12 - \delta_{\operatorname{top}} - \delta_{\operatorname{bot}}) E_{2}(x) \right].$$

We now replace the sum of weights by its expression in the $codeg(\mathfrak{p}_c)$ and sum over all the possible $\mathfrak{P} = (\mathfrak{p}_c)_c$. We get modulo x^{i+1} :

$$\sum_{\substack{\mathfrak{P} = (\mathfrak{p}_c)_c \\ \operatorname{codeg}(\mathfrak{p}_c) \leqslant i}} x^{\sum \operatorname{codeg}(\mathfrak{p}_c)} p(x)^4 \left[\left(g_{\max} - \sum_c \operatorname{codeg}(\mathfrak{p}_c) \right) - (12 - \delta_{\operatorname{top}} - \delta_{\operatorname{bot}}) E_2(x) \right].$$

As we only care about the sum modulo x^{i+1} , we may add all the elements in \mathcal{P} since the ones with higher codegree will contribute 0. There are $\chi - 4$ corners where we choose an element $\mathfrak{p} \in \mathcal{P}$. Using Lemma 4.13 and 5.17 to compute the generating series, we get

$$p(x)^{\chi-4}p(x)^4 \left[g_{\max} - (\chi - 4)E_2(x) - (12 - \delta_{top} - \delta_{bot})E_2(x)\right] \mod x^{i+1}.$$

Finally, Lemma 2.5 allows us to conclude.

Remark 5.19. The method can be adapted by adding two additional edges and compute the genus 2 asymptotic refined invariant, and probably more, at the cost of lengthy computations.

•

References

- [BG16a] Florian Block and Lothar Göttsche. Fock spaces and refined severi degrees. International Mathematics Research Notices, 2016(21):6553–6580, 2016.
- [BG16b] Florian Block and Lothar Göttsche. Refined curve counting with tropical geometry. Compositio Mathematica, 152(1):115–151, 2016.
- [BJP22] Erwan Brugallé and Andrés Jaramillo Puentes. Polynomiality properties of tropical refined invariants. Combinatorial Theory, 2, 2022.
- [Blo19] Thomas Blomme. A Caporaso-Harris type formula for relative refined invariants. arXiv preprint arXiv:1912.06453, 2019.
- [BM07] Erwan Brugallé and Grigory Mikhalkin. Enumeration of curves via floor diagrams. Comptes Rendus Mathematique, 345(6):329–334, 2007.
- [BM08] Erwan Brugallé and Grigory Mikhalkin. Floor decompositions of tropical curves: the planar case. arXiv preprint arXiv:0812.3354, 2008.
- [Bou19] Pierrick Bousseau. Tropical refined curve counting from higher genera and lambda classes. *Inventiones mathematicae*, 215(1):1–79, 2019.
- [CH98] Lucia Caporaso and Joe Harris. Counting plane curves of any genus. *Inventiones mathematicae*, 131(2):345–392, 1998.
- [DFI95] Philippe Di Francesco and Claude Itzykson. Quantum intersection rings. In The Moduli Space of Curves, pages 81–148, Boston, MA, 1995. Birkhäuser Boston.
- [Ful93] William Fulton. Introduction to toric varieties. Number 131. Princeton university press, 1993.
- [Göt98] Lothar Göttsche. A conjectural generating function for numbers of curves on surfaces. Communications in mathematical physics, 196(3):523–533, 1998.
- [GS14] Lothar Göttsche and Vivek Shende. Refined curve counting on complex surfaces. Geometry & Topology, 18(4):2245–2307, 2014.
- [IM13] Ilia Itenberg and Grigory Mikhalkin. On Block–Göttsche multiplicities for planar tropical curves. International Mathematics Research Notices, 2013(23):5289–5320, 2013.
- [IS23] Ilia Itenberg and Eugenii Shustin. Real enumerative invariants relative to the anti-canonical divisor and their refinement. arXiv preprint arXiv:2303.06203, 2023.
- [KM94] Maxim Kontsevich and Yu Manin. Gromov-Witten classes, quantum cohomology, and enumerative geometry. *Communications in Mathematical Physics*, 164(3):525–562, 1994.

- [KST11] Martijn Kool, Vivek Shende, and Richard P. Thomas. A short proof of the Göttsche conjecture. Geometry & Topology, 15(1):397–406, 2011.
- [Mév23] Gurvan Mével. Universal polynomials for tropical refined invariants in genus 0. arXiv:2312.16871, 2023.
- [Mik05] Grigory Mikhalkin. Enumerative tropical algebraic geometry in R². Journal of the American Mathematical Society, 18(2):313–377, 2005.
- [Mik17] Grigory Mikhalkin. Quantum indices and refined enumeration of real plane curves. Acta Mathematica, 219(1):135–180, 2017.
- [NPS18] Johannes Nicaise, Sam Payne, and Franziska Schroeter. Tropical refined curve counting via motivic integration. Geometry & Topology, 22(6):3175–3234, 2018.
- [Tze12] Yu-jong Tzeng. A proof of the Göttsche-Yau-Zaslow formula. Journal of Differential Geometry, 90:439–472, 2012.

UNIVERSITÉ DE NEUCHÂTEL, RUE ÉMILE ARGAN 11, NEUCHÂTEL 2000, SUISSE *Email address*: thomas.blomme@unine.ch

CNRS & NANTES UNIVERSITÉ, UMR 6629 LABORATOIRE DE MATHÉMATIQUES JEAN LERAY, 2 RUE DE LA HOUSSINIÈRE, F-44322 NANTES CEDEX 3, FRANCE

Email address: gurvan.mevel@univ-nantes.fr