

# HIGHER ORDER DIFFERENTIAL SUBORDINATIONS FOR CERTAIN STARLIKE FUNCTIONS

NEHA VERMA AND S. SIVAPRASAD KUMAR

**ABSTRACT.** In this paper, we employ a novel second and third-order differential subordination technique to establish the sufficient conditions for functions to belong to the classes  $\mathcal{S}_s^*$  and  $\mathcal{S}_\rho^*$ , where  $\mathcal{S}_s^*$  is the set of all normalized analytic functions  $f$  satisfying  $zf'(z)/f(z) \prec 1 + \sin z$  and  $\mathcal{S}_\rho^*$  is the set of all normalized analytic functions  $f$  satisfying  $zf'(z)/f(z) \prec 1 + \sinh^{-1} z$ .

## 1. INTRODUCTION

Consider the class  $\mathcal{H}(\mathbb{D})$ , comprising analytic functions defined on the unit disk  $\mathbb{D} := \{z \in \mathbb{C} : |z| < 1\}$ . For any positive integer  $n$  and  $a \in \mathbb{C}$ , we define

$$\mathcal{H}[a, n] := \{f \in \mathcal{H} : f(z) = a + a_n z^n + a_{n+1} z^{n+1} + a_{n+2} z^{n+2} + \dots\}.$$

We assume that  $\mathcal{A} := \mathcal{A}_1$  represents the class containing normalized analytic functions defined on  $\mathbb{D}$  in the following format:

$$\mathcal{A}_n = \{f \in \mathcal{H} : f(z) = z + a_{n+1} z^{n+1} + a_{n+2} z^{n+2} + \dots\}.$$

Additionally, we define  $\mathcal{S}$  as a subset of  $\mathcal{A}$  consisting of univalent functions, and  $\mathcal{S}^*$  as a subset of  $\mathcal{S}$  containing all starlike functions. A starlike function's analytic characterization is given by  $\operatorname{Re}(zf'(z)/f(z)) > 0$  for  $z \in \mathbb{D}$ . Now, considering two functions  $f$  and  $g$ , both belonging to the class  $\mathcal{A}$ , we define the subordination relation as  $f \prec g$  if there exists a Schwarz function  $w$  that fulfills the conditions:  $w(0) = 0$  and  $|w(z)| \leq |z|$  such that  $f(z) = g(w(z))$ . Furthermore, if  $g$  is univalent, the subordination  $f \prec g$  holds if and only if  $f(0) = g(0)$  and  $f(\mathbb{D}) \subseteq g(\mathbb{D})$ .

Ma and Minda [12] introduced the class  $\mathcal{S}^*(\varphi)$ , where  $\varphi$  is an analytic univalent function with properties:  $\operatorname{Re} \varphi(z) > 0$ ,  $\varphi(\mathbb{D})$  is symmetric about the real axis and starlike with respect to  $\varphi(0) = 1$  and  $\varphi'(0) > 0$ . This function  $\varphi$  is referred to as the Ma-Minda function. The class  $\mathcal{S}^*(\varphi)$  is defined as:

$$\mathcal{S}^*(\varphi) = \left\{ f \in \mathcal{A} : \frac{zf'(z)}{f(z)} \prec \varphi(z) \right\}. \quad (1.1)$$

Researchers came across a number of interesting subclasses of  $\mathcal{S}^*$  for the proper selections of the function  $\varphi(z)$  in (1.1). We enlist below some of these subclasses in Table 1. In 2019, Cho et al. [3] introduced the class  $\mathcal{S}_s^*$  by choosing  $\varphi(z) = 1 + \sin z$  and studied its radius problems. Within this context, a function  $f$  belongs to  $\mathcal{S}_s^*$  if the ratio  $zf'(z)/f(z)$  resides within  $\Omega_s := \{w \in \mathbb{C} : |\arcsin(w - 1)| < 1\}$ , which is an eight-shaped domain situated in the right half-plane. Similarly, in 2022, Arora and Kumar [2] introduced the class  $\mathcal{S}_\rho^*$  for  $\varphi(z) = 1 + \sinh^{-1} z$ . In this setting, a function  $f$  is categorized as a member of  $\mathcal{S}_\rho^*$  if the quotient  $zf'(z)/f(z)$  resides within  $\Omega_\rho := \{w \in \mathbb{C} : |\sinh(w - 1)| < 1\}$ , a petal-shaped region.

The study of differential subordination is a more generalized version of differential inequalities on the real line to the complex plane. Miller and Mocanu [15] published their monograph

2010 *Mathematics Subject Classification.* 30C45, 30C80.

*Key words and phrases.* Subordination, Sine Function, Petal-Shaped Domain, Starlike Function.

TABLE 1. List of subclasses of  $\mathcal{S}^*$ 

$\mathcal{S}^*(\varphi)$	$\varphi(z)$	Author(s)	Reference
$\mathcal{S}^*[C, D]$	$(1 + Cz)/(1 + Dz)$	Janowski	[6]
$\mathcal{S}_{SG}^*$	$2/(1 + e^{-z})$	Goel and Kumar	[4]
$\mathcal{S}_\rho^*$	$1 + ze^z$	Kumar and Kamaljeet	[10]
$\mathcal{S}_e^*$	$e^z$	Mendiratta et al.	[14]
$\mathcal{S}_q^*$	$z + \sqrt{1 + z^2}$	Raina and Sokół	[18]
$\mathcal{S}_L^*$	$\sqrt{1 + z}$	Sokół and Stankiewicz	[20]

titled “Differential subordination and univalent functions”. Using the concept of admissibility conditions as mentioned in [15], Saliu et al. [19] dealt with first and second-order differential implication for  $p(z) \prec (1 + z)^\alpha$  where  $\alpha \in (0, 1]$ . Similarly, Mushtaq et al. [16] obtained the conditions on  $\beta$  so that the first-order differential subordinations like  $1 + \beta zp'(z)/p^j(z) \prec \sqrt{1 + cz}$  and  $1 + \beta zp'(z)/p^j(z) \prec 1 + \sqrt{2}z + z^2/2$  implies  $p(z) \prec e^z$  for  $j = 0, 1$  and  $2$ . Furthermore, Naz et al. [17] and Goel and Kumar [4] have successfully deduced numerous first-order subordination implications employing admissible functions for the exponential and sigmoid functions, respectively. Also, Madaan et al. [13] established a series of first and second-order differential subordination implications for the lemniscate of Bernoulli, employing admissibility conditions. For a comprehensive exploration of the most recent advancements in this direction, one may see [8, 9, 11, 21]. Hence, drawing inspiration from the aforementioned considerations, this article distinctly directs its attention towards investigating second and third-order subordination implications for the class of starlike functions pertinent to the eight-shaped domain and second-order subordination implications for the class of starlike functions pertinent to the petal-shaped domain in Sections 2 and 3, respectively. Subsequently, we introduce fundamental definitions and notations relevant to differential subordination.

**Definition 1.1.** [1] Let  $\phi(r, s, t, u; z) : \mathbb{C}^4 \times \mathbb{D} \rightarrow \mathbb{C}$  and  $h(z)$  be a univalent function in  $\mathbb{D}$ , if  $p$  is an analytic function in  $\mathbb{D}$  satisfying the third-order differential subordination

$$\phi(p(z), zp'(z), z^2p''(z), z^3p'''(z); z) \prec h(z) \quad (1.2)$$

then  $p$  is called the solution of the differential subordination. The univalent function  $q$  is said to be a dominant of the solutions of the differential subordination if  $p \prec q$  for all  $p$  satisfying (1.2). A dominant  $\bar{q}$  that satisfies  $\bar{q} \prec q$  for all dominants  $q$  of (1.2) is said to be the best dominant of (1.2), which is unique upto the rotations of  $\mathbb{D}$ .

Moreover, suppose  $Q$  be the set of analytic and univalent functions  $q \in \overline{\mathbb{D}} \setminus \mathbb{E}(q)$ , where

$$\mathbb{E}(q) = \{\zeta \in \partial\mathbb{D} : \lim_{z \rightarrow \zeta} q(z) = \infty\}$$

such that  $q'(\zeta) \neq 0$  for  $\zeta \in \partial\mathbb{D} \setminus \mathbb{E}(q)$ . The subclass of  $Q$  for which  $q(0) = a$  is denoted by  $Q(a)$ .

**Lemma 1.2.** [1] Let  $z_0 \in \mathbb{D}$  and  $r_0 = |z_0|$ . Let  $f(z) = \sum_{k=n}^{\infty} a_k z^k$  be continuous on  $\overline{\mathbb{D}_{r_0}}$  and analytic on  $\mathbb{D} \cup \{z_0\}$  with  $f(z) \neq 0$  and  $n \geq 2$ . If  $|f(z_0)| = \max\{|f(z)| : z \in \overline{\mathbb{D}_{r_0}}\}$  and  $|f'(z_0)| = \max\{|f'(z)| : z \in \overline{\mathbb{D}_{r_0}}\}$ , then there exist real constants  $m, k$  and  $l$  such that

$$\frac{z_0 f'(z_0)}{f(z_0)} = m, \quad 1 + \frac{z_0 f''(z_0)}{f'(z_0)} = k \quad \text{and} \quad 2 + \operatorname{Re} \left( \frac{z_0 f'''(z_0)}{f''(z_0)} \right) = l$$

where  $l \geq k \geq m \geq n \geq 2$ .

**Lemma 1.3.** [15, Theorem 2.3b] Let  $\phi \in \Psi_n[\Omega, q]$  with  $q(0) = a$ . If  $p \in \mathcal{H}[a, n]$  satisfies

$$\phi(p(z), zp'(z), z^2p''(z); z) \in \Omega,$$

then  $p \prec q$ .

Note that if  $\Omega \subset \mathbb{C}$  is a simply connected domain, then there exists a conformal mapping  $h$  from  $\mathbb{D}$  onto  $\Omega = h(\mathbb{D})$ . Moreover, if the function  $\phi(p(z), zp'(z), z^2p''(z); z)$  is analytic in  $\mathbb{D}$ , then  $\phi(p(z), zp'(z), z^2p''(z); z) \in \Omega$  can be expressed in terms of subordination as

$$\phi(p(z), zp'(z), z^2p''(z); z) \prec h(z).$$

For the Ma-Minda functions to satisfy the criteria for third-order differential subordination, Kumar and Goel [9] modified the results of Antonino and Miller [1] in 2020. Moreover, they derived the first, second and third-order differential subordination implications for the class  $\mathcal{S}_{SG}^*$ , using these modified results through admissibility conditions. Recently, Verma and Kumar [21] have obtained results pertaining to second and third-order differential subordination for the class  $\mathcal{S}_e^*$ . Thus, this study is motivated to establish parameter-specific conditions enabling the validity of second and third-order subordination implications for the class  $\mathcal{S}_s^*$ , as well as second-order subordination implications for the class  $\mathcal{S}_\rho^*$ .

The modified definition and lemma of the third-order differential subordination implication by Kumar and Goel [9], accompanied by some lemmas are presented below, which are required to deduce our results in the subsequent sections.

**Definition 1.4.** [9] Let  $\Omega$  be a set in  $\mathbb{C}$ ,  $q \in Q$  and  $k \geq m \geq n \geq 2$ . The class of admissible operators  $\Psi_n[\Omega, q]$  consists of those  $\phi : \mathbb{C}^4 \times \mathbb{D} \rightarrow \mathbb{C}$  that satisfy the admissibility conditions

$$\phi(r, s, t, u; z) \notin \Omega \quad \text{whenever} \quad z \in \mathbb{D},$$

$$r = q(\zeta), \quad s = m\zeta q'(\zeta), \quad \operatorname{Re} \left( 1 + \frac{t}{s} \right) \geq m \left( 1 + \operatorname{Re} \frac{\zeta q''(\zeta)}{q'(\zeta)} \right)$$

and

$$\operatorname{Re} \frac{u}{s} \geq m^2 \operatorname{Re} \frac{\zeta^2 q'''(\zeta)}{q'(\zeta)} + 3m(k-1) \operatorname{Re} \frac{\zeta q''(\zeta)}{q'(\zeta)} \quad \text{for} \quad \zeta \in \partial\mathbb{D} \setminus \mathbb{E}(q).$$

**Lemma 1.5.** [9] Let  $p \in \mathcal{H}[a, n]$  with  $m \geq n \geq 2$ , and let  $q \in Q(a)$  such that it satisfies

$$\left| \frac{zp'(z)}{q'(\zeta)} \right| \leq m \quad \text{for} \quad z \in \mathbb{D} \quad \text{and} \quad \zeta \in \partial\mathbb{D} \setminus \mathbb{E}(q).$$

If  $\Omega$  is a set in  $\mathbb{C}$ ,  $\phi \in \Psi_n[\Omega, a]$  and

$$\phi(p(z), zp'(z), z^2p''(z), z^3p'''(z); z) \subset \Omega,$$

then  $p \prec q$ .

**Lemma 1.6.** [4] Let  $r_0 \approx 0.546302$  be the positive root of the equation  $r^2 + 2 \cot(1)r - 1 = 0$ . Then

$$\left| \log \left( \frac{1+z}{1-z} \right) \right| \geq 1 \quad \text{on} \quad |z| = R \quad \text{if and only if} \quad R \geq r_0.$$

**Lemma 1.7.** [9, Lemma 4, Pg No. 192] For any complex number  $z$ , we have

$$|\log(1+z)| \geq 1 \quad \text{if and only if} \quad |z| \geq e-1.$$

## 2. THE CLASS $\mathcal{S}_s^*$

We begin our study with the function  $q(z) := 1 + \sin z$  and define the admissibility class  $\Psi[\Omega, q]$ , where  $\Omega \subset \mathbb{C}$ . We know that  $q(z)$  is analytic and univalent on  $\overline{\mathbb{D}}$ ,  $q(0) = 1$  and it maps  $\mathbb{D}$  onto the domain  $\Omega_s := \{w \in \mathbb{C} : |\arcsin(w-1)| < 1\}$ . Since  $\mathbb{E}(q) = \phi$ , for  $\zeta \in \partial\mathbb{D} \setminus \mathbb{E}(q)$  if and only if  $\zeta = e^{i\theta}$  for  $\theta \in [0, 2\pi]$ . Now, consider

$$|q'(\zeta)| = \sqrt{\cosh^2(\sin \theta) - \sin^2(\cos \theta)} =: n_1(\theta) \quad (2.1)$$

and  $n_1(\theta)$  achieves its minimum value at  $\theta = 0$ , denoted by

$$\nu_0 := n_1(0) = \sqrt{1 - \sin^2 1} \approx 0.540302. \quad (2.2)$$

It is evident that  $\min |q'(\zeta)| > 0$ , which implies that  $q \in Q(1)$ , and consequently, the admissibility class  $\Psi[\Omega, q]$  is well-defined. While considering  $|\zeta| = 1$ , we observe that  $q(\zeta) \in q(\partial\mathbb{D}) = \partial\Omega_s = \{w \in \mathbb{C} : |\arcsin(w - 1)| = 1\}$ . Consequently,  $|\arcsin(q(\zeta) - 1)| = 1$  and  $\arcsin(q(\zeta) - 1) = e^{i\theta}$  for  $\theta \in [0, 2\pi]$ , which implies that  $q(\zeta) = 1 + \sin \zeta$ . Moreover,  $\zeta q'(\zeta) = e^{i\theta} \cos(e^{i\theta})$  and

$$\frac{\zeta q''(\zeta)}{q'(\zeta)} = e^{i\theta} \tan(e^{i\theta}). \quad (2.3)$$

By comparing the real parts on both sides of (2.3), we obtain

$$\operatorname{Re} \left( \frac{\zeta q''(\zeta)}{q'(\zeta)} \right) = \frac{-\cos \theta \sin(2 \cos \theta) + \sin \theta \sinh(2 \sin \theta)}{\cos(2 \cos \theta) + \cosh(2 \sin \theta)} =: n_2(\theta). \quad (2.4)$$

The function  $n_2(\theta)$  as defined in (2.4), attains its minimum at  $\theta = 0$ , denoted by

$$\nu_1 := n_2(0) = -\frac{\sin 2}{1 + \cos 2} \approx -1.55741. \quad (2.5)$$

Moreover, the class  $\Psi[\Omega, 1 + \sin z]$  is precisely defined as the class of all functions  $\phi : \mathbb{C}^3 \times \mathbb{D} \rightarrow \mathbb{C}$  that satisfy the following conditions:

$$\phi(r, s, t; z) \notin \Omega \quad \text{for } z \in \mathbb{D}, \quad \theta \in [0, 2\pi] \quad \text{and} \quad m \geq 1,$$

whenever

$$r = q(\zeta) = 1 + \sin(e^{i\theta}); \quad s = m\zeta q'(\zeta) = me^{i\theta} \cos(e^{i\theta}); \quad \operatorname{Re} \left( 1 + \frac{t}{s} \right) \geq m(1 + n_2(\theta)). \quad (2.6)$$

Furthermore, for  $q(z) = 1 + \sin z$ , we have

$$\zeta^2 \frac{q'''(\zeta)}{q'(\zeta)} = -e^{2i\theta}.$$

On comparing the real parts of both the sides,

$$\operatorname{Re} \left( \zeta^2 \frac{q'''(\zeta)}{q'(\zeta)} \right) = -\cos 2\theta =: n_3(\theta).$$

The minimum value of  $n_3(\theta)$  is  $-1$ , which is attained at  $\theta = 0$ . Thus, if  $\phi : \mathbb{C}^4 \times \mathbb{D} \rightarrow \mathbb{C}$  then  $\phi \in \Psi[\Omega, 1 + \sin z]$ , provided  $\phi$  satisfies the following conditions:

$$\phi(r, s, t, u; z) \notin \Omega \quad \text{for } z \in \mathbb{D}, \quad \theta \in [0, 2\pi] \quad \text{and} \quad k \geq m \geq 2$$

whenever

$$r = q(\zeta) = 1 + \sin e^{i\theta}; \quad s = m\zeta q'(\zeta) = me^{i\theta} \cos(e^{i\theta});$$

$$\operatorname{Re} \left( 1 + \frac{t}{s} \right) \geq m(1 + n_2(\theta)) \quad \text{and} \quad \operatorname{Re} \frac{u}{s} \geq m^2 n_3(\theta) + 3m(k - 1)n_2(\theta).$$

We now find conditions on the parameters  $\beta_1$  and  $\beta_2$  for different choices of  $h(z)$  namely  $\sqrt{1+z}$ ,  $(1+Cz)/(1+Dz)$ ,  $2/(1+e^{-z})$ ,  $z + \sqrt{1+z^2}$ ,  $1 + \sinh^{-1} z$ ,  $1 + ze^z$ ,  $e^z$  and  $1 + \sin z$ , so that the following second order differential subordination implication holds:

$$1 + \beta_1 z p'(z) + \beta_2 z^2 p''(z) \prec h(z) \implies p(z) \prec 1 + \sin z$$

Note that the values of  $\nu_0$  and  $\nu_1$  are given in (2.2) and (2.5) respectively, which are widely used in the upcoming results. We begin with the following Theorem, for the case when  $h(z) = \sqrt{1+z}$ .

**Theorem 2.1.** Let  $\beta_1, \beta_2 > 0$  and  $\nu_0(\beta_1 + \beta_2\nu_1)(\nu_0(\beta_1 + \beta_2\nu_1) - 2) \geq 1$ . Let  $p$  be analytic function in  $\mathbb{D}$  with  $p(0) = 1$  and

$$1 + \beta_1 zp'(z) + \beta_2 z^2 p''(z) \prec \sqrt{1+z}.$$

Then  $p(z) \prec 1 + \sin z$ .

*Proof.* Suppose  $h(z) = \sqrt{1+z}$  for  $z \in \mathbb{D}$ . Then,  $h(\mathbb{D}) = \{w \in \mathbb{C} : |w^2 - 1| < 1\} =: \Omega$ . Let  $\phi : \mathbb{C}^3 \times \mathbb{D} \rightarrow \mathbb{C}$  be defined as  $\phi(r, s, t; z) = 1 + \beta_1 s + \beta_2 t$ . It is clear that  $\phi \in \Psi[\Omega, 1 + \sin z]$  provided  $\phi(r, s, t; z) \notin \Omega$  for  $z \in \mathbb{D}$ . Note that

$$\begin{aligned} |(\phi(r, s, t; z))^2 - 1| &= |(1 + \beta_1 s + \beta_2 t)^2 - 1| \\ &\geq |\beta_1 s + \beta_2 t|(|\beta_1 s + \beta_2 t| - 2) \\ &\geq |\beta_1 s| \operatorname{Re} \left( 1 + \frac{\beta_2 t}{\beta_1 s} \right) \left( |\beta_1 s| \operatorname{Re} \left( 1 + \frac{\beta_2 t}{\beta_1 s} \right) - 2 \right) \\ &\geq L(L - 2), \end{aligned}$$

where  $L = m\beta_1 n_1(\theta) \operatorname{Re}(1 + \beta_2(mn_2(\theta) + m - 1)/\beta_1)$ . Here  $n_1(\theta)$  and  $n_2(\theta)$  are given in (2.1) and (2.4), respectively. Since  $m \geq 1$ , we have

$$\begin{aligned} |(\phi(r, s, t; z))^2 - 1| &\geq \beta_1 n_1(\theta) \operatorname{Re} \left( 1 + \frac{\beta_2}{\beta_1} n_2(\theta) \right) \left( \beta_1 n_1(\theta) \operatorname{Re} \left( 1 + \frac{\beta_2}{\beta_1} n_2(\theta) \right) - 2 \right) \\ &\geq \nu_0(\beta_1 + \beta_2\nu_1)(\nu_0(\beta_1 + \beta_2\nu_1) - 2) \\ &\geq 1. \end{aligned}$$

Therefore,  $\phi(r, s, t; z) \notin \Omega$  and hence  $\phi \in \Psi[\Omega, 1 + \sin z]$ . Now, the result follows at once by an application of Lemma 1.3.  $\blacksquare$

By considering the function  $p(z) = zf'(z)/f(z)$  in Theorem 2.1, we derive the following corollary:

**Corollary 2.2.** Suppose  $\beta_1, \beta_2 > 0$  and  $f \in \mathcal{A}$ . Let

$$S_f(z) := 1 + \beta_1(S_2 - S_1^2 + S_1) + \beta_2(S_3 + 2S_2 + 2S_1^3 - 2S_1^2 - 3S_1S_2), \quad (2.7)$$

where

$$S_1 = \frac{zf'(z)}{f(z)}, \quad S_2 = \frac{z^2 f''(z)}{f(z)} \quad \text{and} \quad S_3 = \frac{z^3 f'''(z)}{f(z)}. \quad (2.8)$$

Then  $f \in \mathcal{S}_s^*$  provided  $S_f(z) \prec \sqrt{1+z}$  and  $\nu_0(\beta_1 + \beta_2\nu_1)(\nu_0(\beta_1 + \beta_2\nu_1) - 2) \geq 1$ .

**Theorem 2.3.** Let  $\beta_1, \beta_2 > 0$  and  $-1 < D < C \leq 1$  with  $\nu_0(\beta_1 + \beta_2\nu_1)(1 - D^2) \geq (C - D)(1 + |D|)$ . Let  $p$  be analytic function in  $\mathbb{D}$  with  $p(0) = 1$  and

$$1 + \beta_1 zp'(z) + \beta_2 z^2 p''(z) \prec \frac{1 + Cz}{1 + Dz}.$$

Then  $p(z) \prec 1 + \sin z$ .

*Proof.* Let  $h(z) = (1 + Cz)/(1 + Dz)$  for  $z \in \mathbb{D}$ , then we have  $h(\mathbb{D}) = \{w \in \mathbb{C} : |w - (1 - CD)/(1 - D^2)| < (C - D)/(1 - D^2)\} =: \Omega$ . Let  $\phi : \mathbb{C}^3 \times \mathbb{D} \rightarrow \mathbb{C}$  be defined as  $\phi(r, s, t; z) = 1 + \beta_1 s + \beta_2 t$ . It

is clear that  $\phi \in \Psi[\Omega, 1 + \sin z]$  only when  $\phi(r, s, t; z) \notin \Omega$  for  $z \in \mathbb{D}$ . Consider

$$\begin{aligned} \left| \phi(r, s, t; z) - \frac{1 - CD}{1 - D^2} \right| &= \left| 1 + \beta_1 s + \beta_2 t - \frac{1 - CD}{1 - D^2} \right| \\ &\geq |\beta_1 s + \beta_2 t| - \frac{|D|(C - D)}{1 - D^2} \\ &\geq |\beta_1 s| \operatorname{Re} \left( 1 + \frac{\beta_2 t}{\beta_1 s} \right) - \frac{|D|(C - D)}{1 - D^2} \\ &\geq m\beta_1 n_1(\theta) \operatorname{Re} \left( 1 + \frac{\beta_2}{\beta_1} (mn_2(\theta) + m - 1) \right) - \frac{|D|(C - D)}{1 - D^2}. \end{aligned}$$

Using the fact that  $m \geq 1$  and proceeding as in the proof of Theorem 2.1, we have

$$\begin{aligned} \left| \phi(r, s, t; z) - \frac{1 - CD}{1 - D^2} \right| &\geq \nu_0(\beta_1 + \beta_2 \nu_1) - \frac{|D|(C - D)}{1 - D^2} \\ &\geq \frac{C - D}{1 - D^2}. \end{aligned}$$

Thus,  $\phi(r, s, t; z) \notin \Omega$  and hence  $\phi \in \Psi[\Omega, 1 + \sin z]$ . Now, the result follows by an application of Lemma 1.3.  $\blacksquare$

**Theorem 2.4.** Suppose  $\beta_1, \beta_2 > 0$  and  $\nu_0(\beta_1 + \beta_2 \nu_1) \geq r_0$ , where  $r_0 \approx 0.546302$  is the positive root of the equation  $r^2 + 2 \cot(1)r - 1 = 0$ . Let  $p$  be analytic function in  $\mathbb{D}$  with  $p(0) = 1$  and

$$1 + \beta_1 z p'(z) + \beta_2 z^2 p''(z) \prec \frac{2}{1 + e^{-z}}.$$

Then  $p(z) \prec 1 + \sin z$ .

*Proof.* Assuming  $h(z) = 2/(1 + e^{-z})$  for  $z \in \mathbb{D}$ . Then,  $h(\mathbb{D}) = \{w \in \mathbb{C} : |\log(w/(2-w))| < 1\} =: \Omega$ . Let  $\phi : \mathbb{C}^3 \times \mathbb{D} \rightarrow \mathbb{C}$  be defined as  $\phi(r, s, t; z) = 1 + \beta_1 s + \beta_2 t$ . We know that  $\phi \in \Psi[\Omega, 1 + \sin z]$  provided  $\phi(r, s, t; z) \notin \Omega$ . First we consider

$$\begin{aligned} |\beta_1 s + \beta_2 t| &= \beta_1 |s| \left| 1 + \frac{\beta_2 t}{\beta_1 s} \right| \\ &\geq \beta_1 |s| \operatorname{Re} \left( 1 + \frac{\beta_2 t}{\beta_1 s} \right) \\ &\geq m\beta_1 n_1(\theta) \left( 1 + \frac{\beta_2}{\beta_1} (mn_2(\theta) + m - 1) \right). \end{aligned}$$

As  $m \geq 1$ , we obtain

$$\begin{aligned} |\beta_1 s + \beta_2 t| &\geq n_1(\theta)(\beta_1 + \beta_2 n_2(\theta)) \\ &\geq \nu_0(\beta_1 + \beta_2 \nu_1) \\ &\geq r_0. \end{aligned} \tag{2.9}$$

Now, we consider

$$\left| \log \left( \frac{\phi(r, s, t; z)}{2 - \phi(r, s, t; z)} \right) \right| = \left| \log \left( \frac{1 + \beta_1 s + \beta_2 t}{1 - (\beta_1 s + \beta_2 t)} \right) \right|.$$

Through Lemma 1.6 and (2.9), we have

$$\left| \log \left( \frac{1 + \beta_1 s + \beta_2 t}{1 - (\beta_1 s + \beta_2 t)} \right) \right| \geq 1,$$

which implies that  $\phi \in \Psi[\Omega, 1 + \sin z]$ . Now, the result follows by an application of Lemma 1.3.  $\blacksquare$

**Theorem 2.5.** Suppose  $\beta_1, \beta_2 > 0$  and  $\nu_0(\beta_1 + \beta_2\nu_1) \geq \sqrt{2}$ . Let  $p$  be analytic function in  $\mathbb{D}$  with  $p(0) = 1$  and

$$1 + \beta_1 zp'(z) + \beta_2 z^2 p''(z) \prec z + \sqrt{1 + z^2}.$$

Then  $p(z) \prec 1 + \sin z$ .

*Proof.* Let  $h(z) = z + \sqrt{1 + z^2}$  for  $z \in \mathbb{D}$ . Then,  $h(\mathbb{D}) = \{w \in \mathbb{C} : |w^2 - 1| < 2|w|\} =: \Omega$ . Let  $\phi : \mathbb{C}^3 \times \mathbb{D} \rightarrow \mathbb{C}$  be defined as  $\phi(r, s, t; z) = 1 + \beta_1 s + \beta_2 t$ . For  $\phi \in \Psi[\Omega, 1 + \sin z]$ , we must have  $\phi(r, s, t; z) \notin \Omega$ . From the geometry of  $z + \sqrt{1 + z^2}$  (see Fig. 1), we note that  $\Omega$  is constructed by the circles

$$C_1 : |z - 1| = \sqrt{2} \quad \text{and} \quad C_2 : |z + 1| = \sqrt{2}.$$




FIGURE 1. Graph of two circles, namely  $C_1$  (blue boundary) and  $C_2$  (orange boundary). While the shaded region represents  $z + \sqrt{1 + z^2}$ .

It is obvious that  $\Omega$  contains the disk enclosed by  $C_1$  and excludes the portion of the disk enclosed by  $C_1 \cap C_2$ . We have

$$|\phi(r, s, t; z) - 1| = |\beta_1 s + \beta_2 t|.$$

Now, proceeding in the similar way as in the proof of Theorem 2.4, we have

$$\begin{aligned} |\beta_1 s + \beta_2 t| &\geq n_1(\theta)(\beta_1 + \beta_2 n_2(\theta)) \\ &\geq \nu_0(\beta_1 + \beta_2 \nu_1) \\ &\geq \sqrt{2}. \end{aligned}$$

The fact that  $\phi(r, s, t; z)$  lies outside the circle  $C_1$  suffices us to deduce that  $\phi(r, s, t; z) \notin \Omega$  and hence  $\phi \in \Psi[\Omega, 1 + \sin z]$ . Now, the result follows by an application of Lemma 1.3.  $\blacksquare$

**Theorem 2.6.** Let  $\beta_1, \beta_2 > 0$  and  $\nu_0(\beta_1 + \beta_2\nu_1) \geq e$ . Let  $p$  be analytic function in  $\mathbb{D}$  with  $p(0) = 1$  and

$$1 + \beta_1 zp'(z) + \beta_2 z^2 p''(z) \prec 1 + ze^z.$$

Then  $p(z) \prec 1 + \sin z$ .

*Proof.* Assuming  $h(z) = 1 + ze^z$  for  $z \in \mathbb{D}$ . Then,  $\Omega := h(\mathbb{D})$ . Let  $\phi : \mathbb{C}^3 \times \mathbb{D} \rightarrow \mathbb{C}$  be defined as  $\phi(r, s, t; z) = 1 + \beta_1 s + \beta_2 t$ . For  $\phi \in \Psi[\Omega, 1 + \sin z]$ , we need to have  $\phi(r, s, t; z) \notin \Omega$ . Through [10, Lemma 3.3], we observe that the smallest disk containing  $\Omega$  is  $\{\delta \in \mathbb{C} : |\delta - 1| < e\}$ . Thus

$$|\phi(r, s, t; z) - 1| = |\beta_1 s + \beta_2 t|.$$

Analogous to Theorem 2.4, we get

$$\begin{aligned} |\beta_1 s + \beta_2 t| &\geq n_1(\theta)(\beta_1 + \beta_2 n_2(\theta)) \\ &\geq \nu_0(\beta_1 + \beta_2 \nu_1) \\ &\geq e. \end{aligned}$$

Clearly,  $\phi(r, s, t; z)$  lies outside the disk  $\{\delta \in \mathbb{C} : |\delta - 1| < e\}$  which suffices us to conclude that  $\phi(r, s, t; z) \notin \Omega$ . Therefore,  $\phi \in \Psi[\Omega, 1 + \sin z]$  and the result follows as an application of Lemma 1.3.  $\blacksquare$

**Theorem 2.7.** Suppose  $\beta_1, \beta_2 > 0$  and  $2\nu_0(\beta_1 + \beta_2 \nu_1) \geq \pi$ . Let  $p$  be analytic function in  $\mathbb{D}$  with  $p(0) = 1$  and

$$1 + \beta_1 z p'(z) + \beta_2 z^2 p''(z) \prec 1 + \sinh^{-1} z.$$

Then  $p(z) \prec 1 + \sin z$ .

*Proof.* Let  $h(z) = 1 + \sinh^{-1} z$  for  $z \in \mathbb{D}$ . Then,  $h(\mathbb{D}) = \{w \in \mathbb{C} : |\sinh(w - 1)| < 1\} =: \Omega_\rho$ . Let  $\phi : \mathbb{C}^3 \times \mathbb{D} \rightarrow \mathbb{C}$  be defined as  $\phi(r, s, t; z) = 1 + \beta_1 s + \beta_2 t$ . For  $\phi \in \Psi[\Omega_\rho, 1 + \sin z]$ , we must have  $\phi(r, s, t; z) \notin \Omega_\rho$ . Through [2, Remark 2.7], we note that the disk  $\{\delta \in \mathbb{C} : |\delta - 1| < \pi/2\}$  is the smallest disk containing  $\Omega_\rho$ . So,

$$|\phi(r, s, t; z) - 1| = |\beta_1 s + \beta_2 t|.$$

Analogous to Theorem 2.4, we have

$$\begin{aligned} |\beta_1 s + \beta_2 t| &\geq n_1(\theta)(\beta_1 + \beta_2 n_2(\theta)) \\ &\geq \nu_0(\beta_1 + \beta_2 \nu_1) \\ &\geq \frac{\pi}{2}. \end{aligned}$$

Clearly,  $\phi(r, s, t; z)$  does not belong to the disk  $\{\delta \in \mathbb{C} : |\delta - 1| < \pi/2\}$  which suffices us to conclude that  $\phi(r, s, t; z) \notin \Omega_\rho$ . Therefore,  $\phi \in \Psi[\Omega_\rho, 1 + \sin z]$  and thus the result follows as an application of Lemma 1.3.  $\blacksquare$

**Theorem 2.8.** Let  $\beta_1, \beta_2 > 0$  and  $\nu_0(\beta_1 + \beta_2 \nu_1) \geq e - 1$ . Let  $p$  be analytic function in  $\mathbb{D}$  with  $p(0) = 1$  and

$$1 + \beta_1 z p'(z) + \beta_2 z^2 p''(z) \prec e^z.$$

Then  $p(z) \prec 1 + \sin z$ .

*Proof.* Suppose  $h(z) = e^z$  for  $z \in \mathbb{D}$ . Then,  $h(\mathbb{D}) = \{w \in \mathbb{C} : |\log w| < 1\} =: \Omega$ . Let  $\phi : \mathbb{C}^3 \times \mathbb{D} \rightarrow \mathbb{C}$  be defined as  $\phi(r, s, t; z) = 1 + \beta_1 s + \beta_2 t$ . For  $\phi \in \Psi[\Omega, 1 + \sin z]$ , we must have  $\phi(r, s, t; z) \notin \Omega$ . So,

$$|\phi(r, s, t; z) - 1| = |\beta_1 s + \beta_2 t|.$$

Proceeding similar to the proof of Theorem 2.4, we have

$$\begin{aligned} |\beta_1 s + \beta_2 t| &\geq n_1(\theta)(\beta_1 + \beta_2 n_2(\theta)) \\ &\geq \nu_0(\beta_1 + \beta_2 \nu_1) \\ &\geq e - 1. \end{aligned} \tag{2.10}$$

Further, we have

$$|\log(\phi(r, s, t; z))| = |\log(1 + \beta_1 s + \beta_2 t)|.$$



Through Lemma 1.7 and (2.10), we have

$$|\log(1 + \beta_1 s + \beta_2 t)| \geq 1,$$

which implies that  $\phi(r, s, t; z) \notin \Omega$ . Therefore,  $\phi \in \Psi[\Omega, 1 + \sin z]$  and thus the result follows as an application of Lemma 1.3.  $\blacksquare$

**Theorem 2.9.** *Suppose  $\beta_1, \beta_2 > 0$  and  $\nu_0(\beta_1 + \beta_2\nu_1) \geq \sinh 1$ . Let  $p$  be analytic function in  $\mathbb{D}$  with  $p(0) = 1$  and*

$$1 + \beta_1 zp'(z) + \beta_2 z^2 p''(z) \prec 1 + \sin z.$$

*Then  $p(z) \prec 1 + \sin z$ .*

*Proof.* Suppose  $h(z) = 1 + \sin z$  for  $z \in \mathbb{D}$ . Then,  $h(\mathbb{D}) = \{w \in \mathbb{C} : |\arcsin(w - 1)| < 1\} =: \Omega_s$ . Let  $\phi : \mathbb{C}^3 \times \mathbb{D} \rightarrow \mathbb{C}$  be defined as  $\phi(r, s, t; z) = 1 + \beta_1 s + \beta_2 t$ . For  $\phi \in \Psi[\Omega_s, 1 + \sin z]$ , we must have  $\phi(r, s, t; z) \notin \Omega_s$ . Through [3, Lemma 3.3], we note that  $\{\delta \in \mathbb{C} : |\delta - 1| < \sinh 1\}$  is the smallest disk containing  $\Omega_s$ . So,

$$|\phi(r, s, t; z) - 1| = |\beta_1 s + \beta_2 t|.$$

Analogous to Theorem 3.3, we have

$$\begin{aligned} |\beta_1 s + \beta_2 t| &\geq n_1(\theta)(\beta_1 + \beta_2 n_2(\theta)) \\ &\geq \nu_0(\beta_1 + \beta_2 \nu_1) \\ &\geq \sinh 1. \end{aligned}$$

Clearly,  $\phi(r, s, t; z)$  lies outside the disk  $\{\delta \in \mathbb{C} : |\delta - 1| < \sinh 1\}$  which is enough to conclude that  $\phi(r, s, t; z) \notin \Omega_s$ . Therefore,  $\phi \in \Psi[\Omega_s, 1 + \sin z]$  and the result follows as an application of Lemma 1.3.  $\blacksquare$

By taking  $p(z) = zf'(z)/f(z)$  in Theorems 2.7-2.12, we obtain the following corollary with  $S_f$  as given by (2.7):

**Corollary 2.10.** *Suppose  $\beta_1, \beta_2 > 0$  and  $f \in \mathcal{A}$ . Then,  $f \in \mathcal{S}_s^*$  if any of these conditions hold:*

- (i)  $S_f(z) \prec (1 + Cz)/(1 + Dz)$  and  $\nu_0(\beta_1 + \beta_2\nu_1)(1 - D^2) \geq (C - D)(1 + |D|)$ , where  $-1 < D < C \leq 1$ .
- (ii)  $S_f(z) \prec 2/(1 + e^{-z})$  and  $\nu_0(\beta_1 + \beta_2\nu_1) \geq r_0$ , where  $r_0 \approx 0.546302$  is the positive root of the equation  $r^2 + 2 \cot(1)r - 1 = 0$ .
- (iii)  $S_f(z) \prec z + \sqrt{1 + z^2}$  and  $\nu_0(\beta_1 + \beta_2\nu_1) \geq \sqrt{2}$ .
- (iv)  $S_f(z) \prec 1 + ze^z$  and  $\nu_0(\beta_1 + \beta_2\nu_1) \geq e$ .
- (v)  $S_f(z) \prec 1 + \sinh^{-1} z$  and  $2\nu_0(\beta_1 + \beta_2\nu_1) \geq \pi$ .
- (vi)  $S_f(z) \prec e^z$  and  $\nu_0(\beta_1 + \beta_2\nu_1) \geq e - 1$ .
- (vii)  $S_f(z) \prec 1 + \sin z$  and  $\nu_0(\beta_1 + \beta_2\nu_1) \geq \sinh 1$ .

Next, we determine the sufficient conditions on the positive constants  $\beta_1, \beta_2$  and  $\beta_3$  for various selections of  $h(z)$ , notably including the case when  $h(z) = 1 + \sin z$  itself. This study aims to establish third-order differential subordination implication, which is as follows:

$$1 + \beta_1 zp'(z) + \beta_2 z^2 p''(z) + \beta_3 z^3 p'''(z) \prec h(z) \implies p(z) \prec 1 + \sin z.$$

The derived conditions involve the variables  $\nu_0$  and  $\nu_1$ , along with the constants  $m$  and  $k$ , which are explicitly defined in (2.2), (2.5) and Lemma 1.2, respectively. We begin with the Theorem below by choosing  $h(z) = \sqrt{1 + z}$  as follows:

**Theorem 2.11.** *Suppose  $\beta_1, \beta_2, \beta_3 > 0$  and satisfy*

$$\nu_0(\beta_1 + \beta_2\nu_1 + \beta_3(-m^2 + 3m(k - 1)\nu_1))(\nu_0(\beta_1 + \beta_2\nu_1 + \beta_3(-m^2 + 3m(k - 1)\nu_1)) - 2) \geq 1.$$

Let  $p$  be analytic function in  $\mathbb{D}$  with  $p(0) = 1$  and

$$1 + \beta_1 z p'(z) + \beta_2 z^2 p''(z) + \beta_3 z^3 p'''(z) \prec \sqrt{1+z}.$$

Then  $p(z) \prec 1 + \sin z$ .

*Proof.* Suppose  $h(z) = \sqrt{1+z}$  for  $z \in \mathbb{D}$ . Then,  $h(\mathbb{D}) = \{w \in \mathbb{C} : |w^2 - 1| < 1\} =: \Omega$ . Let  $\phi : \mathbb{C}^4 \times \mathbb{D} \rightarrow \mathbb{C}$  be defined as  $\phi(r, s, t, u; z) = 1 + \beta_1 s + \beta_2 t + \beta_3 u$ . We know that  $\phi \in \Psi[\Omega, 1 + \sin z]$  provided  $\phi(r, s, t, u; z) \notin \Omega$  for  $z \in \mathbb{D}$ . Consider

$$\begin{aligned} |(\phi(r, s, t, u; z))^2 - 1| &= |(1 + \beta_1 s + \beta_2 t + \beta_3 u)^2 - 1| \\ &\geq |\beta_1 s + \beta_2 t + \beta_3 u| (|\beta_1 s + \beta_2 t + \beta_3 u| - 2) \\ &\geq |\beta_1 s| \operatorname{Re} \left( 1 + \frac{\beta_2 t}{\beta_1 s} + \frac{\beta_3 u}{\beta_1 s} \right) \left( |\beta_1 s| \operatorname{Re} \left( 1 + \frac{\beta_2 t}{\beta_1 s} + \frac{\beta_3 u}{\beta_1 s} \right) - 2 \right) \\ &\geq M(M - 2), \end{aligned}$$

where  $M = m\beta_1 n_1(\theta)(1 + \beta_2(mn_2(\theta) + m - 1)/\beta_1 + \beta_3(m^2 n_3(\theta) + 3m(k - 1)n_2(\theta))/\beta_1)$ . As  $m \geq 1$  and  $\operatorname{Re}(1 + n_2(\theta)) > 0$ , we get

$$\begin{aligned} |(\phi(r, s, t, u; z))^2 - 1| &\geq \beta_1 n_1(\theta) \left( 1 + \frac{\beta_2}{\beta_1} n_2(\theta) + \frac{\beta_3}{\beta_1} (m^2 n_3(\theta) + 3m(k - 1)n_2(\theta)) \right) \left( \beta_1 n_1(\theta) \left( 1 + \frac{\beta_2}{\beta_1} n_2(\theta) + \frac{\beta_3}{\beta_1} (m^2 n_3(\theta) + 3m(k - 1)n_2(\theta)) \right) - 2 \right) \\ &\geq \nu_0(\beta_1 + \beta_2 \nu_1 + \beta_3(-m^2 + 3m(k - 1)\nu_1)) (\nu_0(\beta_1 + \beta_2 \nu_1 + \beta_3(-m^2 + 3m(k - 1)\nu_1)) - 2) \\ &\geq 1. \end{aligned}$$

This implies that  $\phi(r, s, t; z) \notin \Omega$  and thus  $\phi \in \Psi[\Omega, 1 + \sin z]$ . The result follows as a consequence of Lemma 1.5. ■

By considering  $p(z) = zf'(z)/f(z)$  in Theorem 2.11, we have the following:

**Corollary 2.12.** Suppose  $\beta_1, \beta_2, \beta_3 > 0$  and  $f \in \mathcal{A}$ . Let

$$\begin{aligned} \Theta_f(z) &:= 1 + \beta_1 S_1 + (\beta_1 + 2\beta_2)(S_2 - S_1^2) + (\beta_2 + 3\beta_3)(2S_1^3 - 3S_1 S_2 + 3S_3) \\ &\quad + \beta_3(S_4 - 3S_2^2 - 6S_1^4 - 4S_1 S_3 + 12S_1^2 S_2). \end{aligned} \tag{2.11}$$

where  $S_1, S_2$  and  $S_3$  are mentioned in (2.8) and  $S_4 := z^4 f^{(iv)}(z)/f(z)$ . Then,  $f(z) \in \mathcal{S}_s^*$  provided  $\Theta_f(z) \prec \sqrt{1+z}$  and  $\nu_0(\beta_1 + \beta_2 \nu_1 + \beta_3(-m^2 + 3m(k - 1)\nu_1)) - 2\nu_0(\beta_1 + \beta_2 \nu_1 + \beta_3(-m^2 + 3m(k - 1)\nu_1)) \geq 1$ .

**Theorem 2.13.** Suppose  $\beta_1, \beta_2, \beta_3 > 0$  and  $-1 < D < C \leq 1$  with  $\nu_0(\beta_1 + \beta_2 \nu_1 - m^2 \beta_3 + 3m\beta_3(k - 1)\nu_1)(1 - D^2) \geq (1 + |D|)(C - D)$ . Let  $p$  be analytic function in  $\mathbb{D}$  with  $p(0) = 1$  and

$$1 + \beta_1 z p'(z) + \beta_2 z^2 p''(z) + \beta_3 z^3 p'''(z) \prec \frac{1 + Cz}{1 + Dz}.$$

Then  $p(z) \prec 1 + \sin z$ .

*Proof.* Let  $h(z) = (1 + Cz)/(1 + Dz)$  for  $z \in \mathbb{D}$ . Then,  $h(\mathbb{D}) = \{w \in \mathbb{C} : |w - (1 - CD)/(1 - D^2)| < (C - D)/(1 - D^2)\} =: \Omega$ . Let  $\phi : \mathbb{C}^4 \times \mathbb{D} \rightarrow \mathbb{C}$  be defined as  $\phi(r, s, t, u; z) = 1 + \beta_1 s + \beta_2 t + \beta_3 u$ .

For  $\phi \in \Psi[\Omega, 1 + \sin z]$ , it is needed that  $\phi(r, s, t, u; z) \notin \Omega$  for  $z \in \mathbb{D}$ . Consider

$$\begin{aligned} \left| \phi(r, s, t, u; z) - \frac{1 - CD}{1 - D^2} \right| &= \left| 1 + \beta_1 s + \beta_2 t + \beta_3 u - \frac{1 - CD}{1 - D^2} \right| \\ &\geq |\beta_1 s + \beta_2 t + \beta_3 u| - \frac{|D|(C - D)}{1 - D^2} \\ &\geq |\beta_1 s| \operatorname{Re} \left( 1 + \frac{\beta_2 t}{\beta_1 s} + \frac{\beta_3 u}{\beta_1 s} \right) - \frac{|D|(C - D)}{1 - D^2} \\ &\geq m\beta_1 n_1(\theta) \left( 1 + \frac{\beta_2}{\beta_1} (mn_2(\theta) + m - 1) + \frac{\beta_3}{\beta_1} (m^2 n_3(\theta) + 3m(k - 1)n_2(\theta)) \right) \\ &\quad - \frac{|D|(C - D)}{1 - D^2}. \end{aligned}$$

Proceeding as in the proof of Theorem 2.11 and the fact that  $m \geq 1$ , we obtain

$$\begin{aligned} \left| \phi(r, s, t, u; z) - \frac{1 - CD}{1 - D^2} \right| &\geq \nu_0 \left( \beta_1 + \beta_2 \nu_1 + \beta_3 (-m^2 + 3m(k - 1)\nu_1) \right) - \frac{|D|(C - D)}{1 - D^2} \\ &\geq \frac{C - D}{1 - D^2}. \end{aligned}$$

From the above calculations, we conclude that  $\phi \in \Psi[\Omega, 1 + \sin z]$  and hence the result follows as a consequence of Lemma 1.5.  $\blacksquare$

**Theorem 2.14.** Suppose  $\beta_1, \beta_2, \beta_3 > 0$  and  $\nu_0(\beta_1 + \beta_2 \nu_1 - m^2 \beta_3 + 3m\beta_3(k - 1)\nu_1) \geq r_0$ , where  $r_0 \approx 0.546302$  is the positive root of the equation  $r^2 + 2 \cot(1)r - 1 = 0$ . Let  $p$  be analytic function in  $\mathbb{D}$  with  $p(0) = 1$  and

$$1 + \beta_1 z p'(z) + \beta_2 z^2 p''(z) + \beta_3 z^3 p'''(z) \prec \frac{2}{1 + e^{-z}}.$$

Then  $p(z) \prec 1 + \sin z$ .

*Proof.* Assume  $h(z) = 2/(1 + e^{-z})$  for  $z \in \mathbb{D}$ . Then,  $h(\mathbb{D}) = \{w \in \mathbb{C} : |\log(w/(2 - w))| < 1\} =: \Omega$ . Let  $\phi : \mathbb{C}^4 \times \mathbb{D} \rightarrow \mathbb{C}$  be defined as  $\phi(r, s, t, u; z) = 1 + \beta_1 s + \beta_2 t + \beta_3 u$ . For  $\phi \in \Psi[\Omega, 1 + \sin z]$ , we must have  $\phi(r, s, t, u; z) \notin \Omega$ . Consider

$$\begin{aligned} |\beta_1 s + \beta_2 t + \beta_3 u| &= \beta_1 |s| \left| 1 + \frac{\beta_2 t}{\beta_1 s} + \frac{\beta_3 u}{\beta_1 s} \right| \\ &\geq \beta_1 |s| \operatorname{Re} \left( 1 + \frac{\beta_2 t}{\beta_1 s} + \frac{\beta_3 u}{\beta_1 s} \right) \\ &\geq m\beta_1 n_1(\theta) \left( 1 + \frac{\beta_2}{\beta_1} (mn_2(\theta) + m - 1) + \frac{\beta_3}{\beta_1} (m^2 n_3(\theta) + 3m(k - 1)n_2(\theta)) \right). \end{aligned}$$

As  $m \geq 1$  and  $\operatorname{Re}(1 + n_2(\theta)) > 0$ , we have

$$\begin{aligned} |\beta_1 s + \beta_2 t + \beta_3 u| &\geq n_1(\theta) \left( \beta_1 + \beta_2 n_2(\theta) + \beta_3 (m^2 n_3(\theta) + 3m(k - 1)n_2(\theta)) \right) \\ &\geq \nu_0(\beta_1 + \beta_2 \nu_1 - m^2 \beta_3 + 3m\beta_3(k - 1)\nu_1) \\ &\geq r_0. \end{aligned} \tag{2.12}$$

Now, we consider

$$\left| \log \left( \frac{\phi(r, s, t, u; z)}{2 - \phi(r, s, t, u; z)} \right) \right| = \left| \log \left( \frac{1 + \beta_1 s + \beta_2 t + \beta_3 u}{1 - (\beta_1 s + \beta_2 t + \beta_3 u)} \right) \right|.$$

In view of Lemma 1.6 and (2.12), we have

$$\left| \log \left( \frac{1 + \beta_1 s + \beta_2 t + \beta_3 u}{1 - (\beta_1 s + \beta_2 t + \beta_3 u)} \right) \right| \geq 1,$$

which implies that  $\phi \in \Psi[\Omega, 1 + \sin z]$ . Thus, The result follows as an application of Lemma 1.5.  $\blacksquare$

**Theorem 2.15.** *Suppose  $\beta_1, \beta_2, \beta_3 > 0$  and  $\nu_0(\beta_1 + \beta_2\nu_1 - m^2\beta_3 + 3m\beta_3(k-1)\nu_1) \geq \sqrt{2}$ . Let  $p$  be analytic function in  $\mathbb{D}$  with  $p(0) = 1$  and*

$$1 + \beta_1 z p'(z) + \beta_2 z^2 p''(z) + \beta_3 z^3 p'''(z) \prec z + \sqrt{1 + z^2}.$$

*Then  $p(z) \prec 1 + \sin z$ .*

*Proof.* Take  $h(z) = z + \sqrt{1 + z^2}$  for  $z \in \mathbb{D}$ . Then,  $h(\mathbb{D}) = \{w \in \mathbb{C} : |w^2 - 1| < 2|w|\} =: \Omega$ . Let  $\phi : \mathbb{C}^4 \times \mathbb{D} \rightarrow \mathbb{C}$  be defined as  $\phi(r, s, t, u; z) = 1 + \beta_1 s + \beta_2 t + \beta_3 u$ . For  $\phi \in \Psi[\Omega, 1 + \sin z]$ , we must have  $\phi(r, s, t, u; z) \notin \Omega$ . From the geometry of  $z + \sqrt{1 + z^2}$  (see Fig. 1), we note that  $\Omega$  is constructed by two circles

$$C_1 : |z - 1| = \sqrt{2} \quad \text{and} \quad C_2 : |z + 1| = \sqrt{2}.$$

It is obvious that  $\Omega$  contains the disk enclosed by  $C_1$  and excludes the portion of the disk enclosed by  $C_1 \cap C_2$ . We have

$$|\phi(r, s, t, u; z) - 1| = |\beta_1 s + \beta_2 t + \beta_3 u|.$$

Analogous to Theorem 2.14, we have

$$\begin{aligned} |\beta_1 s + \beta_2 t + \beta_3 u| &\geq n_1(\theta) \left( \beta_1 + \beta_2 n_2(\theta) + \beta_3 (m^2 n_3(\theta) + 3m(k-1)n_2(\theta)) \right) \\ &\geq \nu_0(\beta_1 + \beta_2 \nu_1 - m^2 \beta_3 + 3m\beta_3(k-1)\nu_1) \\ &\geq \sqrt{2}. \end{aligned}$$

Thus, we can say that  $\phi(r, s, t, u; z)$  lies outside the circle  $C_1$  which is sufficient to deduce that  $\phi(r, s, t, u; z) \notin \Omega$ . Therefore,  $\phi \in \Psi[\Omega, 1 + \sin z]$  and thus the result follows as a consequence of Lemma 1.5.  $\blacksquare$

**Theorem 2.16.** *Suppose  $\beta_1, \beta_2, \beta_3 > 0$  and  $\nu_0(\beta_1 + \beta_2\nu_1 - m^2\beta_3 + 3m\beta_3(k-1)\nu_1) \geq e$ . Let  $p$  be analytic function in  $\mathbb{D}$  with  $p(0) = 1$  and*

$$1 + \beta_1 z p'(z) + \beta_2 z^2 p''(z) + \beta_3 z^3 p'''(z) \prec 1 + ze^z.$$

*Then  $p(z) \prec 1 + \sin z$ .*

*Proof.* Let  $h(z) = 1 + ze^z$  for  $z \in \mathbb{D}$ , then  $\Omega := h(\mathbb{D})$ . Let  $\phi : \mathbb{C}^4 \times \mathbb{D} \rightarrow \mathbb{C}$  be defined as  $\phi(r, s, t, u; z) = 1 + \beta_1 s + \beta_2 t + \beta_3 u$ . For  $\phi \in \Psi[\Omega, 1 + \sin z]$ , we must have  $\phi(r, s, t, u; z) \notin \Omega$ . From [10, Lemma 3.3], we note that the smallest disk containing  $\Omega$  is  $\{\delta \in \mathbb{C} : |\delta - 1| < e\}$ . Also,

$$|\phi(r, s, t, u; z) - 1| = |\beta_1 s + \beta_2 t + \beta_3 u|.$$

On the similar lines of the proof of Theorem 2.14, we have

$$\begin{aligned} |\beta_1 s + \beta_2 t + \beta_3 u| &\geq n_1(\theta) \left( \beta_1 + \beta_2 n_2(\theta) + \beta_3 (m^2 n_3(\theta) + 3m(k-1)n_2(\theta)) \right) \\ &\geq \nu_0(\beta_1 + \beta_2 \nu_1 - m^2 \beta_3 + 3m\beta_3(k-1)\nu_1) \\ &\geq e. \end{aligned}$$

Clearly,  $\phi(r, s, t, u; z)$  lies outside the disk  $\{\delta \in \mathbb{C} : |\delta - 1| < e\}$  which is enough to conclude that  $\phi(r, s, t, u; z) \notin \Omega$ . Therefore,  $\phi \in \Psi[\Omega, 1 + \sin z]$  and hence, the result follows as an application of Lemma 1.5.  $\blacksquare$

**Theorem 2.17.** Suppose  $\beta_1, \beta_2, \beta_3 > 0$  and  $2\nu_0(\beta_1 + \beta_2\nu_1 - m^2\beta_3 + 3m\beta_3(k-1)\nu_1) \geq \pi$ . Let  $p$  be analytic function in  $\mathbb{D}$  with  $p(0) = 1$  and

$$1 + \beta_1 zp'(z) + \beta_2 z^2 p''(z) + \beta_3 z^3 p'''(z) \prec 1 + \sinh^{-1} z.$$

Then  $p(z) \prec 1 + \sin z$ .

*Proof.* Assume  $h(z) = 1 + \sinh^{-1} z$  for  $z \in \mathbb{D}$ . Then,  $h(\mathbb{D}) = \{w \in \mathbb{C} : |\sinh(w-1)| < 1\} =: \Omega_\rho$ . Let  $\phi : \mathbb{C}^4 \times \mathbb{D} \rightarrow \mathbb{C}$  be defined as  $\phi(r, s, t, u; z) = 1 + \beta_1 s + \beta_2 t + \beta_3 u$ . For  $\phi \in \Psi[\Omega_\rho, 1 + \sin z]$ , we must have  $\phi(r, s, t, u; z) \notin \Omega_\rho$ . Through [2, Remark 2.7], we note that the smallest disk containing  $\Omega_\rho$  is  $\{\delta \in \mathbb{C} : |\delta - 1| < \pi/2\}$ . Thus

$$|\phi(r, s, t, u; z) - 1| = |\beta_1 s + \beta_2 t + \beta_3 u|.$$

Proceeding on the similar lines of Theorem 2.14, we have

$$\begin{aligned} |\beta_1 s + \beta_2 t + \beta_3 u| &\geq n_1(\theta) \left( \beta_1 + \beta_2 n_2(\theta) + \beta_3 (m^2 n_3(\theta) + 3m(k-1)n_2(\theta)) \right) \\ &\geq \nu_0(\beta_1 + \beta_2 \nu_1 - m^2 \beta_3 + 3m\beta_3(k-1)\nu_1) \\ &\geq \frac{\pi}{2}. \end{aligned}$$

Clearly,  $\phi(r, s, t, u; z)$  lies outside the disk  $\{\delta \in \mathbb{C} : |\delta - 1| < \pi/2\}$  which suffices to prove that  $\phi(r, s, t, u; z) \notin \Omega_\rho$ . Therefore,  $\phi \in \Psi[\Omega_\rho, 1 + \sin z]$  and thus the result follows as an application of Lemma 1.5.  $\blacksquare$

**Theorem 2.18.** Suppose  $\beta_1, \beta_2, \beta_3 > 0$  and  $\nu_0(\beta_1 + \beta_2 \nu_1 - m^2 \beta_3 + 3m\beta_3(k-1)\nu_1) \geq e - 1$ . Let  $p$  be analytic function in  $\mathbb{D}$  with  $p(0) = 1$  and

$$1 + \beta_1 zp'(z) + \beta_2 z^2 p''(z) + \beta_3 z^3 p'''(z) \prec e^z.$$

Then  $p(z) \prec 1 + \sin z$ .

*Proof.* Take  $h(z) = e^z$  for  $z \in \mathbb{D}$ . Then,  $h(\mathbb{D}) = \{w \in \mathbb{C} : |\log w| < 1\} =: \Omega$ . Let  $\phi : \mathbb{C}^4 \times \mathbb{D} \rightarrow \mathbb{C}$  be defined as  $\phi(r, s, t, u; z) = 1 + \beta_1 s + \beta_2 t + \beta_3 u$ . For  $\phi \in \Psi[\Omega, 1 + \sin z]$ , we must have  $\phi(r, s, t, u; z) \notin \Omega$ . So,

$$|\phi(r, s, t, u; z) - 1| = |\beta_1 s + \beta_2 t + \beta_3 u|.$$

Proceeding on the similar lines of proof of Theorem 2.14, we get

$$\begin{aligned} |\beta_1 s + \beta_2 t + \beta_3 u| &\geq n_1(\theta) \left( \beta_1 + \beta_2 n_2(\theta) + \beta_3 (m^2 n_3(\theta) + 3m(k-1)n_2(\theta)) \right) \\ &\geq \nu_0(\beta_1 + \beta_2 \nu_1 - m^2 \beta_3 + 3m\beta_3(k-1)\nu_1) \\ &\geq e - 1. \end{aligned} \tag{2.13}$$

Next, we consider

$$|\log(\phi(r, s, t, u; z))| = |\log(1 + \beta_1 s + \beta_2 t + \beta_3 u)|.$$

Through Lemma 1.7 and (2.13), we have

$$|\log(1 + \beta_1 s + \beta_2 t + \beta_3 u)| \geq 1,$$

which implies that  $\phi(r, s, t, u; z) \notin \Omega$ . Therefore,  $\phi \in \Psi[\Omega, 1 + \sin z]$  and the result follows as a consequence of Lemma 1.5.  $\blacksquare$

**Theorem 2.19.** Suppose  $\beta_1, \beta_2, \beta_3 > 0$  and  $\nu_0(\beta_1 + \beta_2 \nu_1 - m^2 \beta_3 + 3m\beta_3(k-1)\nu_1) \geq \sinh 1$ . Let  $p$  be analytic function in  $\mathbb{D}$  with  $p(0) = 1$  and

$$1 + \beta_1 zp'(z) + \beta_2 z^2 p''(z) + \beta_3 z^3 p'''(z) \prec 1 + \sin z.$$

Then  $p(z) \prec 1 + \sin z$ .

*Proof.* Take  $h(z) = 1 + \sin z$  for  $z \in \mathbb{D}$ . Then,  $h(\mathbb{D}) = \{w \in \mathbb{C} : |\arcsin(w - 1)| < 1\} =: \Omega_s$ . Let  $\phi : \mathbb{C}^4 \times \mathbb{D} \rightarrow \mathbb{C}$  be defined as  $\phi(r, s, t, u; z) = 1 + \beta_1 s + \beta_2 t + \beta_3 u$ . For  $\phi \in \Psi[\Omega_s, 1 + \sin z]$ , we must have  $\phi(r, s, t, u; z) \notin \Omega_s$ . Through [3, Lemma 3.3], we note that the smallest disk containing  $\Omega_s$  is  $\{\delta \in \mathbb{C} : |\delta - 1| < \sinh 1\}$ . So,

$$|\phi(r, s, t, u; z) - 1| = |\beta_1 s + \beta_2 t + \beta_3 u|.$$

Similar to the proof of Theorem 2.14, we have

$$\begin{aligned} |\beta_1 s + \beta_2 t + \beta_3 u| &\geq n_1(\theta) \left( \beta_1 + \beta_2 n_2(\theta) + \beta_3 (m^2 n_3(\theta) + 3m(k-1)n_2(\theta)) \right) \\ &\geq \nu_0(\beta_1 + \beta_2 \nu_1 - m^2 \beta_3 + 3m\beta_3(k-1)\nu_1) \\ &\geq \sinh 1. \end{aligned}$$

Clearly,  $\phi(r, s, t, u; z)$  lies outside the disk  $\{\delta \in \mathbb{C} : |\delta - 1| < \sinh 1\}$  which is enough to conclude that  $\phi(r, s, t, u; z) \notin \Omega_s$ . Therefore,  $\phi \in \Psi[\Omega_s, 1 + \sin z]$  and thus the result holds through Lemma 1.5.  $\blacksquare$

Now, we summarize this section by providing a combined conclusion. If  $p(z) = zf'(z)/f(z)$  is considered in Theorems 2.18-2.23 as in Corollary 2.12. We deduce the following

**Corollary 2.20.** *Suppose  $\beta_1, \beta_2, \beta_3 > 0$  and  $f \in \mathcal{A}$ . Then,  $f \in \mathcal{S}_s^*$  if any of the conditions hold:*

- (i)  $\Theta_f(z) \prec (1 + Cz)/(1 + Dz)$  and  $\nu_0(\beta_1 + \beta_2 \nu_1 - m^2 \beta_3 + 3m\beta_3(k-1)\nu_1)(1 - D^2) \geq (1 + |D|)(C - D)$ , where  $-1 < D < C \leq 1$
- (ii)  $\Theta_f(z) \prec 2/(1 + e^{-z})$  and  $\nu_0(\beta_1 + \beta_2 \nu_1 - m^2 \beta_3 + 3m\beta_3(k-1)\nu_1) \geq r_0$ , where  $r_0 \approx 0.546302$  is the positive root of the equation  $r^2 + 2 \cot(1)r - 1 = 0$ .
- (iii)  $\Theta_f(z) \prec z + \sqrt{1 + z^2}$  and  $\nu_0(\beta_1 + \beta_2 \nu_1 - m^2 \beta_3 + 3m\beta_3(k-1)\nu_1) \geq \sqrt{2}$ .
- (iv)  $\Theta_f(z) \prec 1 + ze^z$  and  $\nu_0(\beta_1 + \beta_2 \nu_1 - m^2 \beta_3 + 3m\beta_3(k-1)\nu_1) \geq e$ .
- (v)  $\Theta_f(z) \prec 1 + \sinh^{-1} z$  and  $2\nu_0(\beta_1 + \beta_2 \nu_1 - m^2 \beta_3 + 3m\beta_3(k-1)\nu_1) \geq \pi$ .
- (vi)  $\Theta_f(z) \prec e^z$  and  $\nu_0(\beta_1 + \beta_2 \nu_1 - m^2 \beta_3 + 3m\beta_3(k-1)\nu_1) \geq e - 1$ .
- (vii)  $\Theta_f(z) \prec 1 + \sin z$  and  $\nu_0(\beta_1 + \beta_2 \nu_1 - m^2 \beta_3 + 3m\beta_3(k-1)\nu_1) \geq \sinh 1$ .

### 3. THE CLASS $\mathcal{S}_\rho^*$

Let us consider the function  $q(z) := 1 + \sinh^{-1} z$  and define the admissibility class  $\Psi[\Omega, q]$ , where  $\Omega \subset \mathbb{C}$ . We know that  $q$  is analytic and univalent on  $\overline{\mathbb{D}}$ ,  $q(0) = 1$  and it maps  $\mathbb{D}$  onto the domain  $\Omega_\rho := \{w \in \mathbb{C} : |\sinh(w - 1)| < 1\}$ . Since  $\mathbb{E}(q) = \phi$ , for  $\zeta \in \partial\mathbb{D} \setminus \mathbb{E}(q)$  if and only if  $\zeta = e^{i\theta}$  for  $\theta \in [0, 2\pi]$ . Now, consider the function

$$|q'(\zeta)| = \frac{1}{\sqrt{2}(\cos \theta)^{1/2}} =: n_4(\theta), \quad (3.1)$$

which has a minimum value of  $1/\sqrt{2}$ . It is evident that  $\min |q'(\zeta)| > 0$ , which implies that  $q \in Q(1)$ , and consequently, the admissibility class  $\Psi[\Omega, q]$  is well-defined. While considering  $|\zeta| = 1$ , we observe that  $q(\zeta) \in q(\partial\mathbb{D}) = \partial\Omega_\rho = \{w \in \mathbb{C} : |\sinh(w - 1)| = 1\}$ . Consequently, we have  $|\log q(\zeta)| = 1$  and  $\sinh(q(\zeta) - 1) = e^{i\theta}$  ( $\theta \in [0, 2\pi]$ ), implies that  $q(\zeta) = 1 + \sinh^{-1} \zeta$ . Moreover,  $\zeta q'(\zeta) = e^{i\theta}/(1 + e^{2i\theta})^{1/2}$  and

$$\frac{\zeta q''(\zeta)}{q'(\zeta)} = -\frac{e^{2i\theta}}{1 + e^{2i\theta}}. \quad (3.2)$$

By comparing the real parts on both sides of (3.2), we obtain

$$\operatorname{Re} \left( \frac{\zeta q''(\zeta)}{q'(\zeta)} \right) = -\frac{1}{2} =: n_5(\theta). \quad (3.3)$$

The function  $n_5(\theta)$  defined in (3.3) has minimum value  $-1/2$  being a constant function. Moreover, the class  $\Psi[\Omega, 1 + \sinh^{-1} z]$  is precisely defined as the class of all functions  $\phi : \mathbb{C}^3 \times \mathbb{D} \rightarrow \mathbb{C}$  that satisfy the following conditions:

$$\phi(r, s, t; z) \notin \Omega \quad \text{for } z \in \mathbb{D}, \quad \theta \in [0, 2\pi] \quad \text{and } m \geq 1,$$

whenever

$$r = q(\zeta) = 1 + \sinh^{-1}(e^{i\theta}); \quad s = m\zeta q'(\zeta) = \frac{me^{i\theta}}{\sqrt{1+e^{2i\theta}}}; \quad \operatorname{Re} \left( 1 + \frac{t}{s} \right) \geq m(1 + n_5(\theta)). \quad (3.4)$$

If  $\phi : \mathbb{C}^2 \times \mathbb{D} \rightarrow \mathbb{C}$ , then the admissibility condition (3.4) becomes

$$\phi \left( 1 + \sinh^{-1}(e^{i\theta}), \frac{me^{i\theta}}{\sqrt{1+e^{2i\theta}}}; z \right) \notin \Omega \quad (z \in \mathbb{D}, \theta \in [0, 2\pi], m \geq 1).$$

Now, we discuss the implications of the following second-order differential subordination

$$1 + \beta_1 z p'(z) + \beta_2 z^2 p''(z) \prec h(z) \implies p(z) \prec 1 + \sinh^{-1} z,$$

by finding sufficient conditions on the parameters  $\beta_1$  and  $\beta_2$  for various choices of  $h(z)$  including  $1 + \sinh^{-1} z$  itself. We begin by considering  $h(z) = \sqrt{1+z}$  in the following manner:

**Theorem 3.1.** *Suppose  $\beta_1, \beta_2 > 0$  and  $(2\beta_1 - \beta_2)(2\beta_1 - \beta_2 - 4\sqrt{2}) \geq 8$ . Let  $p$  be analytic function in  $\mathbb{D}$  with  $p(0) = 1$  and*

$$1 + \beta_1 z p'(z) + \beta_2 z^2 p''(z) \prec \sqrt{1+z}.$$

*Then  $p(z) \prec 1 + \sinh^{-1} z$ .*

*Proof.* Consider  $h(z) = \sqrt{1+z}$  for  $z \in \mathbb{D}$ . Then,  $h(\mathbb{D}) = \{w \in \mathbb{C} : |w^2 - 1| < 1\} =: \Omega$ . Let  $\phi : \mathbb{C}^3 \times \mathbb{D} \rightarrow \mathbb{C}$  be defined as  $\phi(r, s, t; z) = 1 + \beta_1 s + \beta_2 t$ . It is clear that  $\phi \in \Psi[\Omega, 1 + \sinh^{-1} z]$  provided  $\phi(r, s, t; z) \notin \Omega$  for  $z \in \mathbb{D}$ . Note that

$$\begin{aligned} |(\phi(r, s, t; z))^2 - 1| &= |(1 + \beta_1 s + \beta_2 t)^2 - 1| \\ &\geq |\beta_1 s + \beta_2 t| (|\beta_1 s + \beta_2 t| - 2) \\ &\geq |\beta_1 s| \operatorname{Re} \left( 1 + \frac{\beta_2 t}{\beta_1 s} \right) \left( |\beta_1 s| \operatorname{Re} \left( 1 + \frac{\beta_2 t}{\beta_1 s} \right) - 2 \right) \\ &\geq m\beta_1 n_4(\theta) \operatorname{Re} \left( 1 + \frac{\beta_2}{\beta_1} (mn_5(\theta) + m - 1) \right) \left( m\beta_1 n_4(\theta) \operatorname{Re} \left( 1 + \frac{\beta_2}{\beta_1} (mn_5(\theta) + m - 1) \right) - 2 \right). \end{aligned}$$

Here  $n_4(\theta)$  and  $n_5(\theta)$  are given in (3.1) and (3.3). Since  $m \geq 1$ , therefore

$$\begin{aligned} |(\phi(r, s, t; z))^2 - 1| &\geq \beta_1 n_4(\theta) \operatorname{Re} \left( 1 + \frac{\beta_2}{\beta_1} n_5(\theta) \right) \left( \beta_1 n_4(\theta) \operatorname{Re} \left( 1 + \frac{\beta_2}{\beta_1} n_5(\theta) \right) - 2 \right) \\ &\geq \frac{1}{\sqrt{2}} \left( \beta_1 - \frac{\beta_2}{2} \right) \left( \frac{1}{\sqrt{2}} \left( \beta_1 - \frac{\beta_2}{2} \right) - 2 \right) \\ &= \frac{(2\beta_1 - \beta_2)(2\beta_1 - \beta_2 - 4\sqrt{2})}{8} \\ &\geq 1. \end{aligned}$$

Therefore,  $\phi(r, s, t; z) \notin \Omega$  and hence  $\phi \in \Psi[\Omega, 1 + \sinh^{-1} z]$ . Thus, result follows as an application of Lemma 1.3. ■

**Theorem 3.2.** Suppose  $\beta_1, \beta_2 > 0$  and  $-1 < D < C \leq 1$  with  $(2\beta_1 - \beta_2)(1 - D^2) \geq 2\sqrt{2}(C - D)(1 + |D|)$ . Let  $p$  be analytic function in  $\mathbb{D}$  with  $p(0) = 1$  and

$$1 + \beta_1 zp'(z) + \beta_2 z^2 p''(z) \prec \frac{1 + Cz}{1 + Dz}.$$

Then  $p(z) \prec 1 + \sinh^{-1} z$ .

*Proof.* Assume  $h(z) = (1 + Cz)/(1 + Dz)$  for  $z \in \mathbb{D}$ . Then,  $h(\mathbb{D}) = \{w \in \mathbb{C} : |w - (1 - CD)/(1 - D^2)| < (C - D)/(1 - D^2)\} =: \Omega$ . Let  $\phi : \mathbb{C}^3 \times \mathbb{D} \rightarrow \mathbb{C}$  be defined as  $\phi(r, s, t; z) = 1 + \beta_1 s + \beta_2 t$ . It is clear that  $\phi \in \Psi[\Omega, 1 + \sinh^{-1} z]$  if  $\phi(r, s, t; z) \notin \Omega$  for  $z \in \mathbb{D}$ . Consider

$$\begin{aligned} \left| \phi(r, s, t; z) - \frac{1 - CD}{1 - D^2} \right| &= \left| 1 + \beta_1 s + \beta_2 t - \frac{1 - CD}{1 - D^2} \right| \\ &\geq |\beta_1 s + \beta_2 t| - \frac{|D|(C - D)}{1 - D^2} \\ &\geq |\beta_1 s| \operatorname{Re} \left( 1 + \frac{\beta_2 t}{\beta_1 s} \right) - \frac{|D|(C - D)}{1 - D^2} \\ &\geq m\beta_1 n_4(\theta) \operatorname{Re} \left( 1 + \frac{\beta_2}{\beta_1} (mn_5(\theta) + m - 1) \right) - \frac{|D|(C - D)}{1 - D^2}. \end{aligned}$$

Using the fact that  $m \geq 1$  and proceeding on the similar lines of the proof of Theorem 3.1, we have

$$\begin{aligned} \left| \phi(r, s, t; z) - \frac{1 - CD}{1 - D^2} \right| &\geq \frac{1}{\sqrt{2}} \left( \beta_1 - \frac{\beta_2}{2} \right) - \frac{|D|(C - D)}{1 - D^2} \\ &\geq \frac{C - D}{1 - D^2}. \end{aligned}$$

Thus,  $\phi \in \Psi[\Omega, 1 + \sinh^{-1} z]$  and the result follows as an application of Lemma 1.3. ■

**Theorem 3.3.** Suppose  $\beta_1, \beta_2 > 0$  and  $2\beta_1 - \beta_2 \geq 2\sqrt{2}r_0$ , where  $r_0 \approx 0.546302$  is the positive root of the equation  $r^2 + 2\cot(1)r - 1 = 0$ . Let  $p$  be analytic function in  $\mathbb{D}$  with  $p(0) = 1$  and

$$1 + \beta_1 zp'(z) + \beta_2 z^2 p''(z) \prec \frac{2}{1 + e^{-z}}.$$

Then  $p(z) \prec 1 + \sinh^{-1} z$ .

*Proof.* Suppose  $h(z) = 2/(1 + e^{-z})$  for  $z \in \mathbb{D}$ . Then,  $h(\mathbb{D}) = \{w \in \mathbb{C} : |\log(w/(2 - w))| < 1\} =: \Omega$ . Let  $\phi : \mathbb{C}^3 \times \mathbb{D} \rightarrow \mathbb{C}$  be defined as  $\phi(r, s, t; z) = 1 + \beta_1 s + \beta_2 t$ . We know that  $\phi \in \Psi[\Omega, 1 + \sinh^{-1} z]$  only when  $\phi(r, s, t; z) \notin \Omega$ . First we consider,

$$\begin{aligned} |\beta_1 s + \beta_2 t| &= \beta_1 |s| \left| 1 + \frac{\beta_2 t}{\beta_1 s} \right| \\ &\geq \beta_1 |s| \operatorname{Re} \left( 1 + \frac{\beta_2 t}{\beta_1 s} \right) \\ &\geq m\beta_1 n_4(\theta) \left( 1 + \frac{\beta_2}{\beta_1} (mn_5(\theta) + m - 1) \right). \end{aligned}$$

As  $m \geq 1$ , we obtain

$$\begin{aligned} |\beta_1 s + \beta_2 t| &\geq n_4(\theta)(\beta_1 + \beta_2 n_5(\theta)) \\ &\geq \frac{1}{\sqrt{2}} \left( \beta_1 - \frac{1}{2}\beta_2 \right) \\ &\geq r_0. \end{aligned} \tag{3.5}$$



Now, we consider

$$\left| \log \left( \frac{\phi(r, s, t; z)}{2 - \phi(r, s, t; z)} \right) \right| = \left| \log \left( \frac{1 + \beta_1 s + \beta_2 t}{1 - (\beta_1 s + \beta_2 t)} \right) \right|.$$

Through Lemma 1.6 and (3.5), we have

$$\left| \log \left( \frac{1 + \beta_1 s + \beta_2 t}{1 - (\beta_1 s + \beta_2 t)} \right) \right| \geq 1,$$

which implies that  $\phi \in \Psi[\Omega, 1 + \sinh^{-1} z]$ . Therefore, result follows as an application of Lemma 1.3.  $\blacksquare$

**Theorem 3.4.** Suppose  $\beta_1, \beta_2 > 0$  and  $2\beta_1 - \beta_2 \geq 4$ . Let  $p$  be analytic function in  $\mathbb{D}$  with  $p(0) = 1$  and

$$1 + \beta_1 z p'(z) + \beta_2 z^2 p''(z) \prec z + \sqrt{1 + z^2}.$$

Then  $p(z) \prec 1 + \sinh^{-1} z$ .

*Proof.* Take  $h(z) = z + \sqrt{1 + z^2}$  for  $z \in \mathbb{D}$ . Then,  $h(\mathbb{D}) = \{w \in \mathbb{C} : |w^2 - 1| < 2|w|\} =: \Omega$ . Let  $\phi : \mathbb{C}^3 \times \mathbb{D} \rightarrow \mathbb{C}$  be defined as  $\phi(r, s, t; z) = 1 + \beta_1 s + \beta_2 t$ . For  $\phi \in \Psi[\Omega, 1 + \sinh^{-1} z]$ , we must have  $\phi(r, s, t; z) \notin \Omega$ . From the geometry of  $z + \sqrt{1 + z^2}$  (see Fig. 1), we note that  $\Omega$  is constructed by the circles

$$C_1 : |z - 1| = \sqrt{2} \quad \text{and} \quad C_2 : |z + 1| = \sqrt{2}.$$

It is obvious that  $\Omega$  contains the disk enclosed by  $C_1$  and excludes the portion of the disk enclosed by  $C_1 \cap C_2$ . We have

$$|\phi(r, s, t; z) - 1| = |\beta_1 s + \beta_2 t|.$$

Similar to the proof of Theorem 3.3, we have

$$\begin{aligned} |\beta_1 s + \beta_2 t| &\geq n_4(\theta)(\beta_1 + \beta_2 n_5(\theta)) \\ &\geq \frac{1}{\sqrt{2}} \left( \beta_1 - \frac{1}{2} \beta_2 \right) \\ &\geq \sqrt{2}. \end{aligned}$$

The fact that  $\phi(r, s, t; z)$  lies outside the circle  $C_1$  suffices us to deduce that  $\phi(r, s, t; z) \notin \Omega$ . Consequently,  $\phi \in \Psi[\Omega, 1 + \sinh^{-1} z]$  and thus the result follows as an application of Lemma 1.3.  $\blacksquare$

**Theorem 3.5.** Suppose  $\beta_1, \beta_2 > 0$  and  $2\beta_1 - \beta_2 \geq 2\sqrt{2} \sinh 1$ . Let  $p$  be analytic function in  $\mathbb{D}$  with  $p(0) = 1$  and

$$1 + \beta_1 z p'(z) + \beta_2 z^2 p''(z) \prec 1 + \sin z.$$

Then  $p(z) \prec 1 + \sinh^{-1} z$ .

*Proof.* Consider  $h(z) = 1 + \sin z$  for  $z \in \mathbb{D}$ . Then,  $h(\mathbb{D}) = \{w \in \mathbb{C} : |\arcsin(w - 1)| < 1\} =: \Omega_s$ . Let  $\phi : \mathbb{C}^3 \times \mathbb{D} \rightarrow \mathbb{C}$  be defined as  $\phi(r, s, t; z) = 1 + \beta_1 s + \beta_2 t$ . For  $\phi \in \Psi[\Omega_s, 1 + \sinh^{-1} z]$ , we must have  $\phi(r, s, t; z) \notin \Omega_s$ . From [3, Lemma 3.3], we note that the smallest disk containing  $\Omega_s$  is  $\{\delta \in \mathbb{C} : |\delta - 1| < \sinh 1\}$ . So,

$$|\phi(r, s, t; z) - 1| = |\beta_1 s + \beta_2 t|.$$

Proceeding on the same lines of the proof of Theorem 3.3, we have

$$\begin{aligned} |\beta_1 s + \beta_2 t| &\geq n_4(\theta)(\beta_1 + \beta_2 n_5(\theta)) \\ &\geq \frac{1}{\sqrt{2}} \left( \beta_1 - \frac{1}{2} \beta_2 \right) \\ &\geq \sinh 1. \end{aligned}$$

Clearly,  $\phi(r, s, t; z)$  lies outside the disk  $\{\delta \in \mathbb{C} : |\delta - 1| < \sinh 1\}$ , which is enough to conclude that  $\phi(r, s, t; z) \notin \Omega_s$ . Therefore,  $\phi \in \Psi[\Omega_s, 1 + \sinh^{-1} z]$  and the result follows as an application of Lemma 1.3.  $\blacksquare$

**Theorem 3.6.** *Let  $\beta_1, \beta_2 > 0$  and  $2\beta_1 - \beta_2 \geq 2\sqrt{2}e$ . Let  $p$  be analytic function in  $\mathbb{D}$  with  $p(0) = 1$  and*

$$1 + \beta_1 z p'(z) + \beta_2 z^2 p''(z) \prec 1 + z e^z.$$

*Then  $p(z) \prec 1 + \sinh^{-1} z$ .*

*Proof.* Let  $h(z) = 1 + z e^z$  for  $z \in \mathbb{D}$ , then  $\Omega := h(\mathbb{D})$ . Let  $\phi : \mathbb{C}^3 \times \mathbb{D} \rightarrow \mathbb{C}$  be defined as  $\phi(r, s, t; z) = 1 + \beta_1 s + \beta_2 t$ . For  $\phi \in \Psi[\Omega, 1 + \sinh^{-1} z]$ , we need to have  $\phi(r, s, t; z) \notin \Omega$ . Through [10, Lemma 3.3], we observe that the smallest disk containing  $\Omega$  is  $\{\delta \in \mathbb{C} : |\delta - 1| < e\}$ . Thus

$$|\phi(r, s, t; z) - 1| = |\beta_1 s + \beta_2 t|.$$

Analogous to Theorem 3.3, we have

$$\begin{aligned} |\beta_1 s + \beta_2 t| &\geq n_4(\theta)(\beta_1 + \beta_2 n_5(\theta)) \\ &\geq \frac{1}{\sqrt{2}} \left( \beta_1 - \frac{1}{2} \beta_2 \right) \\ &\geq e. \end{aligned}$$

Clearly,  $\phi(r, s, t; z)$  lies outside the disk  $\{\delta \in \mathbb{C} : |\delta - 1| < e\}$  which suffices us to conclude that  $\phi(r, s, t; z) \notin \Omega$ . Therefore,  $\phi \in \Psi[\Omega, 1 + \sinh^{-1} z]$  and the result follows as an application of Lemma 1.3.  $\blacksquare$

**Theorem 3.7.** *Suppose  $\beta_1, \beta_2 > 0$  and  $2\beta_1 - \beta_2 \geq 2\sqrt{2}(e - 1)$ . Let  $p$  be analytic function in  $\mathbb{D}$  with  $p(0) = 1$  and*

$$1 + \beta_1 z p'(z) + \beta_2 z^2 p''(z) \prec e^z.$$

*Then  $p(z) \prec 1 + \sinh^{-1} z$ .*

*Proof.* Take  $h(z) = e^z$  for  $z \in \mathbb{D}$ . Then,  $h(\mathbb{D}) = \{w \in \mathbb{C} : |\log w| < 1\} =: \Omega$ . Let  $\phi : \mathbb{C}^3 \times \mathbb{D} \rightarrow \mathbb{C}$  be defined as  $\phi(r, s, t; z) = 1 + \beta_1 s + \beta_2 t$ . For  $\phi \in \Psi[\Omega, 1 + \sinh^{-1} z]$ , we must have  $\phi(r, s, t; z) \notin \Omega$ . So,

$$|\phi(r, s, t; z) - 1| = |\beta_1 s + \beta_2 t|.$$

Similar to the proof of Theorem 3.3, we have

$$\begin{aligned} |\beta_1 s + \beta_2 t| &\geq n_4(\theta)(\beta_1 + \beta_2 n_5(\theta)) \\ &\geq \frac{1}{\sqrt{2}} \left( \beta_1 - \frac{1}{2} \beta_2 \right) \\ &\geq e - 1. \end{aligned} \tag{3.6}$$

Further, we have

$$|\log(\phi(r, s, t; z))| = |\log(1 + \beta_1 s + \beta_2 t)|.$$

Through Lemma 1.7 and (3.6), we have

$$|\log(1 + \beta_1 s + \beta_2 t)| \geq 1,$$

which implies that  $\phi(r, s, t; z) \notin \Omega$ . Therefore,  $\phi \in \Psi[\Omega, 1 + \sinh^{-1} z]$  and thus the result follows as an application of Lemma 1.3.  $\blacksquare$

**Theorem 3.8.** *Suppose  $\beta_1, \beta_2 > 0$  and  $2\beta_1 - \beta_2 \geq \sqrt{2}\pi$ . Let  $p$  be analytic function in  $\mathbb{D}$  with  $p(0) = 1$  and*

$$1 + \beta_1 z p'(z) + \beta_2 z^2 p''(z) \prec 1 + \sinh^{-1} z.$$

*Then  $p(z) \prec 1 + \sinh^{-1} z$ .*

*Proof.* Suppose  $h(z) = 1 + \sinh^{-1} z$  for  $z \in \mathbb{D}$ . Then,  $h(\mathbb{D}) = \{w \in \mathbb{C} : |\sinh(w - 1)| < 1\} =: \Omega_\rho$ . Let  $\phi : \mathbb{C}^3 \times \mathbb{D} \rightarrow \mathbb{C}$  be defined as  $\phi(r, s, t; z) = 1 + \beta_1 s + \beta_2 t$ . For  $\phi \in \Psi[\Omega_\rho, 1 + \sinh^{-1} z]$ , we must have  $\phi(r, s, t; z) \notin \Omega_\rho$ . Through [2, Remark 2.7], we note that the disk  $\{\delta \in \mathbb{C} : |\delta - 1| < \pi/2\}$  is the smallest disk containing  $\Omega_\rho$ . Now

$$|\phi(r, s, t; z) - 1| = |\beta_1 s + \beta_2 t|.$$

Proceeding on the same lines as in proof of Theorem 3.3, we have

$$\begin{aligned} |\beta_1 s + \beta_2 t| &\geq n_4(\theta)(\beta_1 + \beta_2 n_5(\theta)) \\ &\geq \frac{1}{\sqrt{2}} \left( \beta_1 - \frac{1}{2} \beta_2 \right) \\ &\geq \frac{\pi}{2}. \end{aligned}$$

Clearly,  $\phi(r, s, t; z)$  does not belong to the disk  $\{\delta \in \mathbb{C} : |\delta - 1| < \pi/2\}$ , which suffices us to conclude that  $\phi(r, s, t; z) \notin \Omega_\rho$ . Therefore,  $\phi \in \Psi[\Omega_\rho, 1 + \sinh^{-1} z]$  and thus the result follows as an application of Lemma 1.3.  $\blacksquare$

We conclude this article through the following corollary, obtained from Theorems 3.5-3.10. This is accomplished by considering  $p(z) = zf'(z)/f(z)$ , as demonstrated in Corollary 2.2:

**Corollary 3.9.** *Suppose  $\beta_1, \beta_2 > 0$  and  $f \in \mathcal{A}$ . Then,  $f \in \mathcal{S}_\rho^*$  if any of the conditions hold:*

- (i)  $S_f(z) \prec \sqrt{1+z}$  and  $(2\beta_1 - \beta)(2\beta_1 - \beta - 4\sqrt{2}) \geq 8$ .
- (ii)  $S_f(z) \prec (1 + Cz)/(1 + Dz)$  and  $(2\beta_1 - \beta_2)(1 - D^2) \geq 2\sqrt{2}(C - D)(1 + |D|)$ , where  $-1 < D < C \leq 1$ .
- (iii)  $S_f(z) \prec 2/(1 + e^{-z})$  and  $2\beta_1 - \beta_2 \geq 2\sqrt{2}r_0$ , where  $r_0 \approx 0.546302$  is the positive root of the equation  $r^2 + 2\cot(1)r - 1 = 0$ .
- (iv)  $S_f(z) \prec z + \sqrt{1 + z^2}$  and  $2\beta_1 - \beta_2 \geq 4$ .
- (v)  $S_f(z) \prec 1 + \sin z$  and  $2\beta_1 - \beta_2 \geq 2\sqrt{2} \sinh 1$ .
- (vi)  $S_f(z) \prec 1 + ze^z$  and  $2\beta_1 - \beta_2 \geq 2\sqrt{2}e$ .
- (vii)  $S_f(z) \prec e^z$  and  $2\beta_1 - \beta_2 \geq 2\sqrt{2}(e - 1)$ .
- (viii)  $S_f(z) \prec 1 + \sinh^{-1} z$  and  $2\beta_1 - \beta_2 \geq \sqrt{2}\pi$ .

**Acknowledgment.** Neha Verma is thankful to the Department of Applied Mathematics, Delhi Technological University, New Delhi-110042 for providing Research Fellowship.

## REFERENCES

- [1] J. Antonino and S. S. Miller, Third-order differential inequalities and subordinations in the complex plane, *Complex Var. Elliptic Equ.*, **56** (2011), no. 5, 439–454.
- [2] K. Arora and S. S. Kumar, Starlike functions associated with a petal shaped domain, *Bull. Korean Math. Soc.*, **59** (2022), no. 4, 993–1010.
- [3] N. E. Cho, V. Kumar, S. S. Kumar and V. Ravichandran, Radius problems for starlike functions associated with the sine function, *Bull. Iranian Math. Soc.*, **45** (2019), no. 1, 213–232.
- [4] P. Goel and S. S. Kumar, Certain class of starlike functions associated with modified sigmoid function, *Bull. Malays. Math. Sci. Soc.*, **43** (2020), no. 1, 957–991.
- [5] A. W. Goodman, *Univalent functions. Vol. I*, Mariner Publishing Co., Inc., Tampa, FL, 1983.
- [6] W. Janowski, Extremal problems for a family of functions with positive real part and for some related families, *Ann. Polon. Math.*, **23** (1970/71), 159–177.
- [7] S. Kanas, Subordinations for domains bounded by conic sections, *Bull. Belg. Math. Soc. Simon Stevin*, **15** (2008), no. 4, 589–598.
- [8] S. S. Kumar and P. Goel, Application of Pythagorean means and Differential Subordination, *Bull. Belg. Math. Soc. Simon Stevin* **29** (2022), no. 3, 285 - 305.
- [9] S. S. Kumar and P. Goel, Starlike functions and higher order differential subordinations, *Rev. R. Acad. Cienc. Exactas Fís. Nat. Ser. A Mat. RACSAM*, **114** (2020), no. 4, Paper No. 192, 23 pp.

- [10] S. S. Kumar and G. Kamaljeet, A cardioid domain and starlike functions, *Anal. Math. Phys.*, **11** (2021), no. 2, Paper No. 54, 34 pp.
- [11] P. G. Krishnan, V. Ravichandran and P. Saikrishnan, Functions subordinate to the exponential function, *Commun. Korean Math. Soc.*, **38** (2023), no. 1, 163–178.
- [12] W. C. Ma and D. Minda, A unified treatment of some special classes of univalent functions, *Proc. Confer. Complex Anal. (Tianjin, 1992)*, 157–169.
- [13] V. Madaan, A. Kumar and V. Ravichandran, Starlikeness associated with lemniscate of Bernoulli, *Filomat*, **33** (2019), no. 7, 1937–1955.
- [14] R. Mendiratta, S. Nagpal and V. Ravichandran, On a subclass of strongly starlike functions associated with exponential function, *Bull. Malays. Math. Sci. Soc.*, **38** (2015), no. 1, 365–386.
- [15] S. S. Miller and P. T. Mocanu, *Differential subordinations, Monographs and Textbooks in Pure and Applied Mathematics*, 225, Marcel Dekker, Inc., New York, 2000.
- [16] S. Mushtaq, M. Raza and J. Sokół, Differential subordination related with exponential functions, *Quaest. Math.*, **45** (2022), no. 6, 889–899.
- [17] A. Naz, S. Nagpal and V. Ravichandran, Starlikeness associated with the exponential function, *Turkish J. Math.*, **43** (2019), no. 3, 1353–1371.
- [18] R. K. Raina and J. Sokół, Some properties related to a certain class of starlike functions, *C. R. Math. Acad. Sci. Paris*, **353** (2015), no. 11, 973–978.
- [19] A. Saliu, K. Jabeen and V. Ravichandran, Differential subordination for certain strongly starlike functions, *Rendiconti del Circolo Matematico di Palermo Series*, **2** (2023), 1–18.
- [20] J. Sokół and J. Stankiewicz, Radius of convexity of some subclasses of strongly starlike functions, *Zeszyty Nauk. Politech. Rzeszowskiej Mat. No. 19* (1996), 101–105.
- [21] N. Verma and S. S. Kumar, Differential subordination of certain class of starlike functions, *arXiv e-prints*, pp.arXiv:2306.11215(2023) (accepted in *Complex Variables and Elliptic Equations*).

DEPARTMENT OF APPLIED MATHEMATICS, DELHI TECHNOLOGICAL UNIVERSITY, DELHI–110042, INDIA  
*Email address:* `nehaverma1480@gmail.com`

DEPARTMENT OF APPLIED MATHEMATICS, DELHI TECHNOLOGICAL UNIVERSITY, DELHI–110042, INDIA  
*Email address:* `spkumar@dce.ac.in`