

INDECOMPOSABLE SET-THEORETICAL SOLUTIONS TO THE YANG-BAXTER EQUATION OF SIZE p^2

CARSTEN DIETZEL, SILVIA PROPERZI, SENNE TRAPPENIERS

ABSTRACT. The *quantum Yang-Baxter equation* is a braiding condition on complex vector spaces which is of high relevance in several fields of mathematics, such as knot theory and quantum group theory. A combinatorial approach is the investigation of *set-theoretic solutions* to the Yang-Baxter equation and their associated algebraic structures. In this article, we focus on indecomposable set-theoretic solutions to the Yang-Baxter equation. More specifically, we give a full classification of those which are of size p^2 .

INTRODUCTION

The *quantum Yang-Baxter equation*, named after the physicists Chen-Ning Yang and Rodney James Baxter, first appeared in the context of statistical mechanics and integrable systems [3, 21]. Given a dimensional complex vector space V , call a linear operator $R : V^{\otimes 2} \rightarrow V^{\otimes 2}$ a *solution* to the *Yang-Baxter equation* if R is bijective and satisfies the following equation within $\text{End}(V^{\otimes 3})$:

$$(YBE) \quad (R \otimes \text{id}_V)(\text{id}_V \otimes R)(R \otimes \text{id}_V) = (\text{id}_V \otimes R)(R \otimes \text{id}_V)(\text{id}_V \otimes R).$$

Outside of physics, solutions to this equation have been recognized to be of importance in low-dimensional topology, as results of Turaev [20] show that each Yang-Baxter operator gives rise to an associated knot invariant. While these operators are therefore of high importance in knot theory, they are very difficult to construct; initially, the only source of solutions was the theory of quantum groups. However, Drinfeld [8] observed that a particular class of solutions provides a handier way to obtain classes of solutions by demanding that R permutes the tensors $v_i \otimes v_j$ where $(v_i)_{i \in I}$ is a basis of V . By doing so, one gets a discretized version of the Yang-Baxter equation, the *set-theoretic Yang-Baxter equation*, which consists of a set X and a bijection $r : X^2 \rightarrow X^2$ that satisfies the following equation in the symmetric group Sym_{X^3} :

$$(SYBE) \quad (r \times \text{id}_X)(\text{id}_X \times r)(r \times \text{id}_X) = (\text{id}_X \times r)(r \times \text{id}_X)(\text{id}_X \times r).$$

Besides the simpler, combinatorial flavour of this equation, Drinfeld's intention was to submit these solutions to a deformation process from which novel linear solutions can be obtained.

Indeed, set-theoretic solutions admit an analysis via algebraic methods, the first steps being made by Etingof, Schedler and Soloviev [9] and Gateva-Ivanova and Van den Bergh [11].

Two set-theoretic solutions (X, r) and (Y, s) are called *equivalent* if there is a bijection $f : X \rightarrow Y$ such that $(f \times f)r = s(f \times f)$.

Writing a set-theoretic solution (X, r) as $r(x, y) = (\lambda_x(y), \rho_y(x))$ we call it

- (1) *non-degenerate* if the mappings λ_x, ρ_x are bijective for all $x \in X$,
- (2) *involution* if $r^2 = \text{id}_{X^2}$.

By the size of a set-theoretic solution, we mean the size of X . From now on, when talking about *solutions*, we mean a *set-theoretic solution to the Yang–Baxter equation that is finite, involutive and non-degenerate*.

Rump has shown in [17] that the binary operation $x \cdot y = \lambda_x^{-1}(y)$ turns X into a particularly easy algebraic structure; a *(non-degenerate) cycle set*, which is defined as a set X with a binary operation $(x, y) \mapsto x \cdot y$ such that the following axioms are satisfied:

- (C1) $(x \cdot y) \cdot (x \cdot z) = (y \cdot x) \cdot (y \cdot z) \quad \forall x, y, z \in X,$
- (C2) the map $y \mapsto \sigma_x(y) = x \cdot y$ is bijective for all $x \in X,$
- (C3) the square map $x \mapsto \text{Sq}(x) = x \cdot x$ is bijective.

Homomorphisms of cycle sets are defined in the obvious way. Consequently, this holds for all derived notions, such as *isomorphisms*. Note that isomorphism of cycle sets translates to equivalence of set-theoretic solutions under the correspondence described above. As usual, if there is no danger of confusion, a cycle set (X, \cdot) is simply referred to by its underlying set X .

The cycle sets considered in this work will always be assumed to be of finite size and will be referred to as *cycle sets*, without the predicate *nondegenerate*. Cycle sets are in bijective correspondence with finite nondegenerate involutive solutions to the Yang–Baxter equation. More precisely, given a cycle set (X, \cdot) then (X, r_X) is a nondegenerate involutive solution to the YBE, where $r_X : X^2 \rightarrow X^2$ is given by $r_X(x, y) = (\sigma_x^{-1}(y), \sigma_x^{-1}(y) \cdot x)$.

Of particular importance for understanding general cycle sets is the class of *indecomposable* cycle sets. A cycle set X is called *indecomposable* if there are no subsets $X_1, X_2 \neq \emptyset$ closed under the cycle set operation such that $X = X_1 \cup X_2$. In a sense, they are elementary building blocks for cycle sets and turn out to be significantly more well-behaved. This can be seen from computations of Akgün, Mereb and Vendramin [1] that imply that the number of cycle sets of given cardinality grows in an aggressive manner, while the number of indecomposable cycle sets stays quite small. This is a hint that there are strong obstructions for a cycle set to be indecomposable.

Indeed, up to isomorphism, there is only one indecomposable cycle set of size p , p a prime number, that is given by $X = \mathbb{Z}_p$ with the operation $x \cdot y = y + 1$ [9, Theorem 2.13]. Furthermore, Jedlička and Pilitowska [14] have recently classified all indecomposable cycle sets of multipermutation level 2 by group-theoretic means. In particular, this includes all indecomposable cycle sets of order pq that are of *finite multipermutation level*, where p, q are (not necessarily distinct) primes. Cedó and Okniński [6, Section 5] have described a class of simple cycle sets of size p^2 , p a prime. In [7], the same authors have recently proven that all indecomposable cycle sets of squarefree size are of finite multipermutation level.

In this article, we build on these results by providing a full classification of indecomposable cycle sets of size p^2 , p a prime.

Note that a cycle set (X, \cdot) is indecomposable if and only if the corresponding set-theoretic solution (X, r) is indecomposable in the sense that there is no partition $X = A \sqcup B$ ($A, B \neq \emptyset$) such that $r(X^2) = X^2$ and $r(Y^2) = Y^2$.

Using the correspondence between cycle sets and set-theoretic solutions, our final result, Theorem 5.1, reformulated in terms of set-theoretic solutions, will be as follows:

Theorem. *Each indecomposable solution (X, r) to the Yang–Baxter equation of size p^2 for some prime p is equivalent to one of the following solutions:*

- (1) $X = \mathbb{Z}_{p^2}$, with $r(x, y) = (y + 1, x - 1)$.

(2) $X = \mathbb{Z}_p \times \mathbb{Z}_p$, with

$$r \begin{pmatrix} (a, x) \\ (b, y) \end{pmatrix} = \begin{pmatrix} (b-1, y - S\chi_0(b-1) - \Phi(b-1-a)) \\ (a+1, x + S\chi_0(a) + \Phi(a-b+1)) \end{pmatrix}$$

where $\Phi : \mathbb{Z}_p \rightarrow \mathbb{Z}_p$ is a nonconstant map with $\Phi(0) = 0$, $S \in \mathbb{Z}_p$ and

$$\chi_0 : \mathbb{Z}_p \rightarrow \mathbb{Z}_p$$

$$\chi_0(x) = \begin{cases} 1 & x = 0 \\ 0 & x \neq 0 \end{cases}.$$

The parameters S, Φ and S', Φ' define equivalent solutions if and only if $S = S'$ and $\alpha\Phi = \Phi'$ for some $\alpha \in \mathbb{Z}_p^*$.

(3) $X = \mathbb{Z}_p \times \mathbb{Z}_p$, with

$$r \begin{pmatrix} (a, x) \\ (b, y) \end{pmatrix} = \begin{pmatrix} (\alpha^{-1}b - x, \alpha^{-1}y - \Phi(\alpha^{-1}b - x - a)) \\ (\alpha a + y - \Phi(b - \alpha x - \alpha a), \alpha x + \Phi(\alpha a - \alpha x - b)) \end{pmatrix}$$

where $\Phi : \mathbb{Z}_p \rightarrow \mathbb{Z}_p$ is a nonconstant map with $\Phi(x) = \Phi(-x)$ and $\alpha \in \mathbb{Z}_p^*$ is such that $\Phi(\alpha x) = \alpha\Phi(x)$.

The parameters α, Φ and α', Φ' define equivalent cycle sets if and only if $\alpha = \alpha'$ and there is a $\beta \in \mathbb{Z}_p^*$ such that $\beta^{-1}\Phi(\beta x) = \Phi'(x)$ for all $x \in \mathbb{Z}_p$.

Note that the classification of indecomposable retractable solutions of size p^2 follows from the theoretical classification of indecomposable solutions of multipermutation level 2 in [14]. We merely give them in a more explicit form building further upon their results, which also gives a more uniform presentation compared to the irretractable case. This more explicit form is obtained in Section 2, where we also present a relation between the size of any indecomposable cycle set of finite multipermutation level and the order of its permutation group.

The strategy to tackle the irretractable case is to first restrict to the case when the permutation group of a solution is a p -group (Section 3) and then construct all irretractable solutions by an extension procedure (Section 4).

We conclude this article by comparing the found solutions with a family of solutions that have recently been constructed by Cedó and Okniński [6], thus answering [6, Question 7.4] affirmatively. Furthermore, we give an enumeration of the irretractable solutions.

1. PRELIMINARIES

Recall that a *brace* is a triple $(A, +, \circ)$, where $(A, +)$ is an abelian group and (A, \circ) is a group such that for all $a, b, c \in A$ a skew left distributivity holds

$$a \circ (b + c) = a \circ b - a + a \circ c,$$

where $-a$ denotes the inverse of a in $(A, +)$. Similarly, the inverse of an element $a \in A$ with respect to the operation \circ will be denoted by \bar{a} .

The group $A^+ = (A, +)$ is the *additive group* of A and $A^\circ = (A, \circ)$ is the *multiplicative group* of A . We will also denote the conjugation by a in A° as ${}^a b = a \circ b \circ \bar{a}$ for all $a, b \in A$. For $n \in \mathbb{Z}$, $a \in A$, the n -th power of a in A^+ is denoted na and the n -th power in A° is denoted $a^{\circ n}$.

There is an action by automorphisms of A° on A^+ called the λ -action defined by

$$\lambda_a(b) = -a + a \circ b$$

for all $a, b \in A$. Related to this, there is also the operation

$$a * b = \lambda_a(b) - b = -a + a \circ b - b,$$

which is easily verified to be left distributive. The set of fixed points of the λ -action is denoted

$$\text{Fix}(A) = \{a \in A : b + a = b \circ a \text{ for all } b \in A\} = \{a \in A \mid b * a = 0 \text{ for all } b \in A\}$$

A *subbrace* of A is simply a subset I of A that is a subgroup of both A^+ and A° .

A *left ideal* of A is a subset I of A such that I is a subgroup of A^+ , and $\lambda_a(I) \subseteq I$ for all $a \in A$. Note that in particular, every characteristic subgroup of A^+ is a left ideal. If additionally, I is a normal subgroup of A° , one calls I an *ideal*.

The *socle* of A is the ideal

$$\text{Soc}(A) = \ker \lambda = \{a \in A \mid a \circ b = a + b \text{ for all } b \in A\}.$$

Lemma 1.1. *Let A be a brace, then*

$$\lambda_a(b) = -a + {}^a b + \lambda_{a_b}(a) \text{ for all } a, b \in A.$$

In particular, if $a \in \text{Fix}(A)$, then $\lambda_a(b) = {}^a b$ for all $b \in A$ and thus λ_a is a brace automorphism of A .

Proof. Observe that $\lambda_a(b) = -a + a \circ b = -a + {}^a b \circ a = -a + {}^a b + \lambda_{a_b}(a)$. \square

Lemma 1.2. *Let A be a brace. Then $\text{Fix}(A) \cap Z(A^\circ) \subseteq \text{Soc}(A)$.*

Proof. Let $f \in \text{Fix}(A) \cap Z(A^\circ)$, then Lemma 1.1 implies that $\lambda_f(a) = {}^f a = a$ for all $a \in A$, i.e. $f \in \text{Soc}(A)$. \square

Theorem 1.3. *Let A be a brace and B a subbrace such that B° acts trivially on A^+/B^+ under the λ -action. Furthermore, assume that B° is normal in A° . Then B is an ideal of A .*

Proof. Let $b \in B$. We have to show that $\lambda_a(b) \in B$ for all $a \in A$. Note that ${}^a b \in B$ because B is normal, therefore $\lambda_{a_b}(a) \in a + B$ as B° acts trivially on A^+/B^+ . Using Lemma 1.1 and the normality of B^+ , we deduce that

$$\lambda_a(b) = -a + {}^a b + \lambda_{a_b}(a) \in -a + B + a + B = B. \quad \square$$

Given a prime p , we mean by a p -brace a finite brace whose size is a power of p .

Proposition 1.4. *Let A be a p -brace and $B \subseteq A$ a subbrace with $|A| = p|B|$. Then B is an ideal of A .*

Proof. A^+/B^+ is cyclic of order p and therefore is acted upon trivially by B° . Furthermore, as $[A^\circ : B^\circ] = p$ is the smallest prime divisor of $|A|$, the subgroup B° is normal in A° . It follows from Theorem 1.3 that B is an ideal of A . \square

We will not use this consequence but it is worth noting that this line of argumentation proves, more generally, the following statement for skew braces [12]:

Corollary 1.5. *Let A be a finite skew brace and B a subbrace such that B^+ is a normal subgroup of A^+ and $p = |A|/|B|$ is equal to the least prime dividing $|A|$. Then B is an ideal of A .*

Lemma 1.6. *If A is a p -brace, then $\text{Fix}(A) \neq 0$. Moreover, if A° is abelian, then $\text{Soc}(A) \neq 0$.*

Proof. $\text{Fix}(A)$ is the set of fixed points under the action $\lambda : A^\circ \rightarrow \text{Aut}(A^+)$. It is well-known that an action of a p -group by automorphisms of a p -group always has nontrivial fixed points.

If we assume that A° is abelian, Lemma 1.2 implies that $\text{Fix}(A) \subseteq \text{Soc}(A)$, hence $\text{Soc}(A) \neq 0$ as well. \square

Let G be a group acting on a set X . A *system of imprimitivity* is a partition $X = \bigsqcup_{i \in I} X_i$ that is invariant under the group action in the sense that for any $g \in G, i \in I$, there is a $j \in I$ such that $g \cdot X_i = X_j$. The subsets X_i are called *blocks* of the system. A system of imprimitivity is *trivial* if either $|I| = 1$ or $|X_i| = 1$ for all $i \in I$. Else it is nontrivial.

Recall that if X is finite and G is transitive, then G acts transitively on the blocks of a system of imprimitivity, and $|X_i| = |X_j|$ for all $i, j \in X$. In particular, all $|X_i|$ divide $|X|$.

Lemma 1.7. *Let G be a nonabelian p -group with a transitive action on a set X of size p^2 . Then G has at most one nontrivial system of imprimitivity.*

Proof. If there is more than one nontrivial system of imprimitivity, a theorem of Lucchini [16, Theorem 1] implies that $G \leq \text{Sym}_p \times \text{Sym}_p$. But if G is a p -group, this forces G to be isomorphic to a subgroup of $\mathbb{Z}_p \times \mathbb{Z}_p$ and therefore to be abelian. \square

Let the groups G, H act on sets X resp. Y . Then the *wreath product* is defined as the semidirect product $G \wr H = G \ltimes H^X$ where G acts on H^X by $g \cdot (h_x)_{x \in X} = (h_{g^{-1} \cdot x})_{x \in X}$.

$G \wr H$ acts on $X \times Y$ by permutations of the form

$$(x, y) \mapsto (g \cdot x, h_x \cdot y).$$

where $g \in G$, and $(h_x)_{x \in X}$ is a family of elements in H .

Proposition 1.8. *Let $G \leq \text{Sym}_{\mathbb{Z}_p \times \mathbb{Z}_p}$ be a transitive solvable group such that the sets $\{a\} \times \mathbb{Z}_p$, $a \in \mathbb{Z}_p$, form a system of imprimitivity, then G is conjugate to a subgroup of $\text{AGL}(1, p) \wr \text{AGL}(1, p)$. If moreover G is a p -group, then G is conjugate to a subgroup of $\mathbb{Z}_p \wr \mathbb{Z}_p$.*

Proof. Let G be as above. As G respects the given system of imprimitivity, it is well-known that $G \leq \text{Sym}_p \wr \text{Sym}_p$, where the action of the wreath product on $\mathbb{Z}_p \times \mathbb{Z}_p$ is precisely as described above.

As a special case of [13, II, 3.2 Satz], we know that every solvable transitive subgroup of $\text{Sym}_{\mathbb{Z}_p}$ is conjugate to a subgroup of $\text{AGL}(1, p)$. So we conclude that G is conjugate to a subgroup of $\text{AGL}(1, p) \wr \text{AGL}(1, p)$. As $\mathbb{Z}_p \wr \mathbb{Z}_p$ is a p -Sylow subgroup of $\text{AGL}(1, p) \wr \text{AGL}(1, p)$ also the last part of the statement follows. \square

Note that elements of $\mathbb{Z}_p \wr \mathbb{Z}_p$ are precisely permutations of the form

$$(a, x) \mapsto (a + \beta, x + \gamma_a),$$

with $\beta, \gamma_1, \dots, \gamma_p \in \mathbb{Z}_p$.

On any brace A we have a canonical cycle set structure (A, \cdot) where $a \cdot b = \lambda_a^{-1}(b)$. A subset $X \subseteq A$ is a *cycle base* of A if it is a sub-cycle set of (A, \cdot) and also X generates the groups A^+ and A° . A cycle base is called *transitive* if X forms an orbit under the λ -action of A , or equivalently if X is an indecomposable cycle set.

To a cycle set (X, \cdot) we associate the group

$$(G(X), \circ) = \langle X \mid x \circ y = \sigma_x^{-1}(y) \circ (\sigma_x^{-1}(y) \cdot x) \text{ for all } x, y \in X \rangle.$$

By defining $x + y = x \circ \sigma_x(y)$, we then obtain a brace $(G(X), +, \circ)$. The canonical map $\iota : (X, \cdot) \rightarrow (G(X), \cdot) : x \mapsto x$ is always an injective cycle set homomorphism, so we can identify X with $\iota(X)$. Note that it follows immediately from the definition that $\iota(X)$ is a cycle base of $G(X)$.

To any cycle set X we associate its *permutation group*, defined as

$$\mathcal{G}(X) = \langle \sigma_x \mid x \in X \rangle \subseteq \text{Sym}_X.$$

Recall that a cycle set X is indecomposable if and only if $\mathcal{G}(X)$ acts transitively on X . The above group structure will also be denoted by $(\mathcal{G}(X), \circ)$ and we obtain, as

described in [5], a brace structure $(\mathcal{G}(X), +, \circ)$ by defining $\sigma_x^{-1} + \sigma_y^{-1} = \sigma_x^{-1} \circ \sigma_{\sigma_x(y)}^{-1}$. Note that this means that we obtain a surjective brace homomorphism $G(X) \rightarrow \mathcal{G}(X) : x \mapsto \sigma_x^{-1}$, where the kernel is precisely $\text{Soc}(G(X))$. As a result, the canonical map

$$\pi : (X, \cdot) \mapsto (\mathcal{G}(X), \cdot); \quad x \mapsto \sigma_x^{-1},$$

becomes a cycle set homomorphism. The image $\pi(X)$ is called the retraction of X , denoted $\text{Ret}(X)$. If π is injective, hence (X, \cdot) is isomorphic to a sub-cycle set of $(\mathcal{G}(X), \cdot)$, then we say that X is *irretractable*. When π is not injective, (X, \cdot) is *retractable*. Note that $\pi(x) = \pi(y)$ if and only if $\sigma_x = \sigma_y$. If moreover, we can obtain a one-element cycle set by iteratively repeating the retraction process on a cycle set X , we say that X has *finite multipermutation level*.

By a result of Cedó and Okniński [7, Lemma 3.3] we have:

Proposition 1.9. *Let $f : X \rightarrow Y$ be a surjective cycle set homomorphism. If X is indecomposable, then Y is also indecomposable and the fibres $f^{-1}(y)$ all have the same cardinality. In particular, $|Y|$ divides $|X|$.*

2. INDECOMPOSABLE CYCLE SETS OF FINITE MULTIPERMUTATION LEVEL

Let e_i be the canonical basis of $\bigoplus_{i \in \mathbb{Z}} \mathbb{Z}$. We define

$$c_k = \begin{cases} \sum_{i=1}^{-k} -e_{1-i} & k < 0 \\ 0 & k = 0 \\ \sum_{i=1}^k e_i & k > 0 \end{cases}$$

For the reader's convenience, we first recall a special case of the construction of indecomposable cycle sets of multipermutation level 2 from the following result of Jedlička and Pilitowska [14, Prop. 5.1, Prop. 5.7].

Theorem 2.1. *Let X be an indecomposable cycle set of multipermutation level 2 such that $|\text{Ret}(X)| = m$, then there exist*

- (1) *a subgroup $H \leq \bigoplus_{i \in \mathbb{Z}} \mathbb{Z}$ such that $c_i - c_{i+m} \in H$ for all $i \in \mathbb{Z}$,*
- (2) *$s \in (\bigoplus_{i \in \mathbb{Z}} \mathbb{Z}) / H$,*

such that X is isomorphic to a cycle set of the form $X = (\mathbb{Z} \times \bigoplus_{i \in \mathbb{Z}} \mathbb{Z}) / \sim$ where

$$(a, x) \sim (b, y) \iff a - b \equiv 0 \pmod{m} \text{ and } x - y \equiv \frac{a - b}{m} s \pmod{H}$$

and

$$[(a, x)] \cdot [(b, y)] = [(b - 1, y - c_{a-b} + c_{-b})].$$

Moreover, different choices of H and s yield non-isomorphic cycle sets.

We are now able to give a more explicit version of the above result.

Theorem 2.2. *Let X be an indecomposable cycle set of multipermutation level 2 such that $|\text{Ret}(X)| = m$, then X is isomorphic to a cycle set of the form $X = \mathbb{Z}_m \times A$ with the multiplication*

$$(a, x) \cdot (b, y) = (b + 1, y + S\chi_0(b) + \Phi(b - a)),$$

where $(A, +)$ is an abelian group and $\Phi : \mathbb{Z}_m \rightarrow A$ is a non-constant map such that $\Phi(0) = 0$ and $S \in A$. Two such cycle sets, given by (A, Φ, S) and (B, Φ', S') , are isomorphic if and only if there exists a group isomorphism $f : A \rightarrow B$ such that $\Phi' = f\Phi$ and $f(S) = S'$.

Proof. Note that the elements $\{(a, x) \mid 1 \leq a \leq m, x \in A\}$ form a system of representatives for the relation \sim . We define the map

$$\psi : (\mathbb{Z} \times \bigoplus_{i \in \mathbb{Z}} \mathbb{Z}) / \sim \rightarrow \mathbb{Z}_m \times (\bigoplus_{i \in \mathbb{Z}} \mathbb{Z}) / H,$$

as $\psi([(a, x)]) = (a, x)$ for $1 \leq a \leq m$. In particular,

$$\psi([(0, x)]) = \psi([(m, x + s)]) = (m, x + s).$$

Under this identification, we find that the cycle set as given in Theorem 2.1 is isomorphic to the cycle set on $\mathbb{Z}_m \times (\bigoplus_{i \in \mathbb{Z}} \mathbb{Z}) / H$ given by

$$(a, x) \cdot (b, y) = (b - 1, y - s\chi_0(b) - c_{a-b} + c_{-b}).$$

Now instead of starting from $H \leq \bigoplus_{i \in \mathbb{Z}} \mathbb{Z}$ and $s \in \bigoplus_{i \in \mathbb{Z}} \mathbb{Z} / H$ we can also start with an abelian group A , $s \in A$ and a group homomorphism $\phi : \bigoplus_{i \in \mathbb{Z}} \mathbb{Z} \rightarrow A$, we then let $H = \ker \phi$. As the c_i , $i \neq 0$, form a basis of $\bigoplus_{i \in \mathbb{Z}} \mathbb{Z}$, we can freely choose the images $\phi(c_i) \in A$ as long as $\phi(c_i) = \phi(c_{i+m})$. If we denote $\phi(c_i) = \Phi(i)$ we see that every such ϕ uniquely corresponds to a map $\Phi : \mathbb{Z}_m \rightarrow A$ such that $\Phi(0) = 0$. Using ϕ to identify $\bigoplus_{i \in \mathbb{Z}} \mathbb{Z} / H$ and A we find a cycle set structure on $\mathbb{Z}_m \times A$ given by

$$(a, x) \cdot (b, y) = (b - 1, y - s\chi_0(b) - \Phi(a - b) + \Phi(-b)).$$

Recall that different choices of H and s give non-isomorphic cycle sets. It is clear that for two abelian groups A, B , $s \in A$, $s' \in B$, and maps $\Phi : \mathbb{Z}_m \rightarrow A$ and $\Phi' : \mathbb{Z}_m \rightarrow B$, the associated homomorphism ϕ and ϕ' have the same kernel H if and only if there exists a group isomorphism $f : A \rightarrow B$ such that $\phi' = f\phi$, or equivalently $\Phi' = f\Phi$. Moreover, s and s' correspond to the same element in $\bigoplus_{i \in \mathbb{Z}} \mathbb{Z} / H$ precisely if $f(s) = s'$.

At last, define $g : \mathbb{Z}_m \rightarrow A$ as $g(b) = \sum_{i=0}^{b-1} \Phi(i)$ for $1 \leq b \leq m$. If $b \neq 0$, then $g(b+1) - g(b) = \Phi(b)$ and if $b = 0$ then $g(b+1) - g(b) = -\sum_{i=1}^{m-1} \Phi(i)$. Under the permutation $\theta : (a, x) \mapsto (-a, -x + g(a))$ the cycle set structure now becomes

$$\begin{aligned} \theta^{-1}(\theta(a, x) \cdot \theta(b, y)) &= \theta^{-1}((-a, -x + g(a)) \cdot (-b, -y + g(b))) \\ &= \theta^{-1}(-b - 1, -y + g(b) - s\chi_0(b) - \Phi(b - a) + \Phi(b)) \\ &= (b + 1, y - g(b) + s\chi_0(b) + \Phi(b - a) - \Phi(b) + g(b + 1)) \\ &= (b + 1, y + S\chi_0(b) + \Phi(b - a)), \end{aligned}$$

where $S = s - \sum_{i=0}^{m-1} \Phi(i)$. To conclude the proof, note that if we are given an abelian group B , an isomorphism $f : A \rightarrow B$ and we set $\Phi' = f\Phi$, then $f(s) = s'$ if and only if $f(S) = f(s - \sum_{i=0}^{m-1} \Phi(i)) = s' - \sum_{i=1}^{m-1} \Phi'(i) = S'$. \square

Remark 2.3. Note that the solutions of the YBE on $X = \mathbb{Z}_p \times A$ corresponding to the cycle sets in Theorem 2.2 are given by

$$r \begin{pmatrix} (a, x) \\ (b, y) \end{pmatrix} = \begin{pmatrix} (b - 1, y - S\chi_0(b - 1) - \Phi(b - 1 - a)) \\ (a + 1, x + S\chi_0(a) + \Phi(a - b + 1)) \end{pmatrix}.$$

Corollary 2.4. Let X be a retractable indecomposable cycle set of size p^2 , p a prime, then X has finite multipermutation level and is isomorphic to one of the following.

- (1) $X = \mathbb{Z}_{p^2}$ with $x \cdot y = y + 1$.
- (2) $X = \mathbb{Z}_p \times \mathbb{Z}_p$, with

$$(a, x) \cdot (b, y) = (b + 1, y + S\chi_0(b) + \Phi(b - a)),$$

where $\Phi : \mathbb{Z}_p \rightarrow \mathbb{Z}_p$ is a non-constant map such that $\Phi(0) = 0$ and $S \in \mathbb{Z}_p$. The parameters S, Φ and S', Φ' define isomorphic cycle sets if and only if $S' = \alpha S$ and $\Phi' = \alpha \Phi$ for some $\alpha \in \mathbb{Z}_p^*$.

Proof. It follows from Proposition 1.9 that $|\text{Ret}(X)| \in \{1, p\}$. If $|\text{Ret}(X)| = 1$ then X clearly has finite multipermutation level and is isomorphic to the given cyclic cycle set on \mathbb{Z}_{p^2} . If $|\text{Ret}(X)| = p$, then we know that $\text{Ret}(X)$ is isomorphic to the cycle set on \mathbb{Z}_p with $x \cdot y = y + 1$ [9, Theorem 2.13]. In particular, $|\text{Ret}(\text{Ret}(X))| = 1$ and thus X has multipermutation level 2. The statement now follows directly from Theorem 2.2. \square

One easily verifies the following result:

Lemma 2.5. *Let $\psi : X \rightarrow Y$ be a surjective homomorphism of cycle sets. Then there is a unique surjective brace homomorphism $\mathcal{G}(\psi) : \mathcal{G}(X) \rightarrow \mathcal{G}(Y)$ such that $\mathcal{G}(\psi)(\sigma_x) = \sigma_{\psi(x)}$.*

For any integer n , we denote by $\pi(n)$ the set of prime divisors of n .

Lemma 2.6. *Let $\psi : X \rightarrow Y$ be a surjective homomorphism of indecomposable cycle sets. Let K be the kernel of the induced brace homomorphism $\mathcal{G}(\psi) : \mathcal{G}(X) \rightarrow \mathcal{G}(Y)$. If K° is nilpotent, then on the orbits \tilde{X} of K° we have a well-defined cycle set structure and surjective cycle set homomorphisms $X \xrightarrow{f} \tilde{X} \xrightarrow{g} Y$ such that*

- (1) $\psi = gf$,
- (2) the induced homomorphism $\mathcal{G}(g) : \mathcal{G}(\tilde{X}) \rightarrow \mathcal{G}(Y)$ is an isomorphism,
- (3) $\pi(|X|/|\tilde{X}|) = \pi(|K|)$.

Proof. Recall that under the canonical embedding, X embeds as a transitive cycle base into the structure brace $G(X)$. Let the ideal K' be the inverse image of K under the canonical homomorphism $G(X) \rightarrow \mathcal{G}(X)$. Then it follows from [18, Theorem 1] that the orbits of X under the λ -action of K' (which are the same as under the action of K) form a congruence relation on X . Let us denote the resulting cycle set \tilde{X} with factor map $f : X \rightarrow \tilde{X}$. As elements of the same orbit of K get identified under ψ , we find a well-defined surjective homomorphism $g : \tilde{X} \rightarrow Y$ such that $\psi = gf$. Now let I denote the kernel of $\mathcal{G}(f) : \mathcal{G}(X) \rightarrow \mathcal{G}(\tilde{X})$. As $\mathcal{G}(\psi) = \mathcal{G}(g)\mathcal{G}(f)$, we know that $I \leq K$. On the other hand, for any $\sigma \in K$, its permutation of X induces the identity mapping on \tilde{X} , hence we see that $\sigma \in I$. We conclude that $I = K$ and therefore $\mathcal{G}(g)$ is an isomorphism.

As K is normal in $\mathcal{G}(X)^\circ$, its orbits all have the same size $|X|/|\tilde{X}|$. Clearly $\pi(|X|/|\tilde{X}|) \subseteq \pi(|K|)$. On the other hand, let p be a prime not contained in $\pi(|X|/|\tilde{X}|)$ and denote K_p the unique Sylow p -subgroup of K° . Then for every $x \in X$, the size of its orbit is coprime to p hence K_p is contained in the stabiliser of x under the action of K . As K° acts faithfully on X , we conclude that $K_p = 0$ hence $p \notin \pi(|K|)$. \square

Theorem 2.7. *Let X be a finite indecomposable cycle set of finite multipermutation level, then $\pi(|X|) = \pi(|\mathcal{G}(X)|)$.*

Proof. We prove this by induction on $|X|$. If $|X| = 1$ then $\mathcal{G}(X)$ is trivial, hence the statement holds. Now assume that $|X| > 1$ and consider the canonical retraction homomorphism $\psi : X \rightarrow \text{Ret}(X)$. From [4, Corollary 3.9] we know that the kernel K of $\mathcal{G}(X) \rightarrow \mathcal{G}(\text{Ret}(X))$ is precisely $\text{Soc}(\mathcal{G}(X))$. By Lemma 2.6 we know that the orbits of K° yield a cycle set \tilde{X} and because K° is abelian we get that $\pi(|X|/|\tilde{X}|) = \pi(|K|)$. As $|\tilde{X}| < |X|$ we know that $\pi(|\tilde{X}|) = \pi(|\mathcal{G}(\tilde{X})|) = \pi(|\mathcal{G}(X)/K|)$, from which we conclude that

$$\pi(|X|) = \pi(|X|/|\tilde{X}|) \cup \pi(|\tilde{X}|) = \pi(|K|) \cup \pi(|\mathcal{G}(X)/K|) = \pi(|\mathcal{G}(X)|). \quad \square$$

Corollary 2.8. *Let X be an indecomposable cycle set of finite multipermutation level and size p^n for some prime p , then $\mathcal{G}(X)$ is a p -group.*

Proof. This follows directly from Theorem 2.7. \square

3. INDECOMPOSABLE IRRETRACTABLE CYCLE SETS OF SIZE p^2 - THE p -GROUP CASE

In this section, we will solve the following problem:

Problem 3.1. Let p be a prime. Find all isomorphism classes of indecomposable irretractable cycle sets (X, \cdot) of size p^2 such that $\mathcal{G}(X)$ is a p -group.

We first construct these solutions in Section 3.1. In Section 3.2 we then study when such solutions are isomorphic and moreover we determine their automorphism groups.

3.1. Constructing the solutions. We first observe the following:

Proposition 3.2. *Let X be an irretractable cycle set, then $\text{Soc}(\mathcal{G}(X)) = 0$ and $\mathcal{G}(X)^\circ$ is not abelian.*

Proof. As X is irretractable, it follows from [2, Lemma 2.1.] that $\text{Soc}(\mathcal{G}(X)) = 0$. From Lemma 1.6 we then find that $\mathcal{G}(X)^\circ$ is not abelian. \square

In the remainder of the section, we let X be an indecomposable irretractable cycle set of size p^2 and assume that $\mathcal{G} = \mathcal{G}(X)$ is a p -group. We associate X with its image in \mathcal{G} , which is a transitive cycle base of \mathcal{G} .

By Lemma 1.7 and Proposition 3.2 we have a unique system of imprimitivity, for $x \in X$ we denote by \mathcal{B}_x the block containing x . We denote by $\mathcal{A} = \mathcal{G} \cap \mathbb{Z}_p^p$ the abelian subgroup which fixes the blocks setwise. Note that $[\mathcal{G}^\circ : \mathcal{A}^\circ] = p$.

Proposition 3.3. $\text{Fix}(\mathcal{G}) \cap \mathcal{A} = 0$.

Proof. Suppose that $0 \neq f \in \text{Fix}(\mathcal{G}) \cap \mathcal{A}$, without loss of generality we may assume that $f^{\circ p} = 0$. By Lemma 1.1 and the abelianity of \mathcal{A} , we see that $\lambda_f(g) = f g = g$ for all $g \in \mathcal{A}$. Thus λ_f fixes all elements in \mathcal{A} . Moreover, by Lemma 1.2, $\text{Fix}(\mathcal{G}) \cap Z(\mathcal{G}^\circ) \subseteq \text{Soc}(\mathcal{G}^\circ) = 0$, so $f \notin Z(\mathcal{G}^\circ)$. Hence, knowing also that $|\mathcal{G}|/|\mathcal{A}| = p$, we see that $\mathcal{A} = \{g \in \mathcal{G} \mid f g = g\}$. As λ_f is an automorphism, \mathcal{A} is a subbrace of index p in \mathcal{G} . By Proposition 1.4, \mathcal{A} is an ideal of \mathcal{G} .

As \mathcal{G}/\mathcal{A} is a brace of order p , it must be trivial and the canonical map $\mathcal{G} \rightarrow \mathcal{G}/\mathcal{A}$ maps the transitive cycle base X to a single element. Hence there exists some $g \in \mathcal{G}$ such that for all $x \in X$ there exists some $a_x \in \mathcal{A}$ such that $x = g + a_x$. In particular, if we set $\gamma = f * g \in \mathcal{A}$, we find that also $f * x = f * g + f * a_x = \gamma$. Hence inductively for all $n \in \mathbb{N}$ we find $f^{\circ n} * x = n\gamma$ since

$$\begin{aligned} f^{\circ n} * x &= \lambda_{f^{\circ n}}(x) - x = \lambda_f(\lambda_{f^{\circ n-1}}(x)) - \lambda_f(x) + \lambda_f(x) - x \\ &= \lambda_f(f^{\circ n-1} * x) + f * x = \lambda_f((n-1)\gamma) + \gamma = n\gamma \end{aligned}$$

In particular, this implies that $p\gamma = 0$. If $\gamma = 0$, then $\lambda_f(x) = x$ for all $x \in X$ and therefore $\lambda_f = \text{id}_{\mathcal{G}}$, which would imply that $f \in \text{Soc}(\mathcal{G})$. It therefore follows that $\gamma \neq 0$, which in turn implies that λ_f has no fixed points on X . In particular, for each $a \in \mathcal{A}$ and $x \in X$, there exists some $n \in \mathbb{N}$ such that $\lambda_a(x) = \lambda_f^n(x)$, hence $a * x = f^{\circ n} * x = n\gamma$.

Now consider I be the subgroup of \mathcal{G}_+ generated by $\{a * g \mid a \in \mathcal{A}, g \in \mathcal{G}\}$, which is an ideal by [18, Corollary after Proposition 6]. Because X generates \mathcal{G}^+ and $*$ is left distributive, I is the subgroup of \mathcal{G}_+ generated by $\{a * x \mid a \in \mathcal{A}, x \in X\}$. By the previous discussion, $I = \{0, \gamma, \dots, (p-1)\gamma\}$. As $|I| = p$, we find that $I \subseteq \text{Fix}(\mathcal{G})$. Since I is a minimal normal subgroup of the nilpotent group \mathcal{G}° , we also find that $I \subseteq Z(\mathcal{G}^\circ)$, hence Lemma 1.2 yields a contradiction with the assumption that $\text{Soc}(\mathcal{G}) = 0$. \square

Now that we know that $\text{Fix}(\mathcal{G}) \cap \mathcal{A} = 0$, it follows from Lemma 1.6 that $|\text{Fix}(\mathcal{G})| = p$. Also, we have a semidirect product $\mathcal{G}^\circ = \mathcal{A}^\circ \rtimes \text{Fix}(\mathcal{G})^\circ$. We use this for the following construction: for all $g \in \mathcal{G}$, we have an equality $g \circ \text{Fix}(\mathcal{G}) = g + \lambda_g(\text{Fix}(\mathcal{G})) = g + \text{Fix}(\mathcal{G})$. Therefore, \mathcal{A} forms a system of representatives for $\mathcal{G}^\circ / \text{Fix}(\mathcal{G})^\circ$ and $\mathcal{G}^+ / \text{Fix}(\mathcal{G})^+$. We keep the multiplication on \mathcal{A} as it is, but as \mathcal{A} is not necessarily closed under $+$, we define $g \oplus h$ as the unique element in $\mathcal{A} \cap (g + h + \text{Fix}(\mathcal{G}))$.

Proposition 3.4. \mathcal{A} is a brace under \circ and \oplus .

Proof. For $a, b, c \in \mathcal{A}$, we calculate

$$\begin{aligned} \{a \circ b \ominus a \oplus a \circ c\} &= \mathcal{A} \cap (a \circ b - a + a \circ c + \text{Fix}(\mathcal{G})) = \mathcal{A} \cap (a \circ (b + c) + \text{Fix}(\mathcal{G})) \\ &= \mathcal{A} \cap ((a + \text{Fix}(\mathcal{G})) \circ (b + c + \text{Fix}(\mathcal{G}))) \\ &= (\mathcal{A} \cap (a + \text{Fix}(\mathcal{G}))) \circ (\mathcal{A} \cap (b + c + \text{Fix}(\mathcal{G}))) \\ &= (\mathcal{A} \cap (a + \text{Fix}(\mathcal{G}))) \circ \{b \oplus c\} = \{a \circ (b \oplus c)\}. \end{aligned}$$

Hence $a \circ (b \oplus c) = a \circ b \ominus a \oplus a \circ c$. \square

Denote by $\tilde{\mathcal{A}}$ the thus constructed brace on \mathcal{A} . We define the *transition* map

$$\begin{aligned} \tau : \mathcal{G} &\rightarrow \tilde{\mathcal{A}} \\ g &\mapsto [g], \text{ where } \{[g]\} = \mathcal{A} \cap (g \circ \text{Fix}(\mathcal{G})). \end{aligned}$$

Note that τ is a group homomorphism $\mathcal{G}^+ \rightarrow \tilde{\mathcal{A}}^\oplus$ but this is not necessarily true for $\mathcal{G}^\circ \rightarrow \tilde{\mathcal{A}}^\circ$. With this notation, we also see that $g \oplus h = [g + h]$. Furthermore, on $\tilde{\mathcal{A}}$, the λ -action changes to $\tilde{\lambda}_g(h) = [\lambda_g(h)]$ which implies that the image $[X] \subseteq \tilde{\mathcal{A}}$ is invariant under its $\tilde{\lambda}$ -action (which is not necessarily transitive). Therefore, it is again a cycle set under the operation $[x] \cdot [y] = \tilde{\lambda}_{[x]}^{-1}([y])$. Denote this cycle set by \tilde{X} .

Proposition 3.5. $|\tilde{X}| = p$.

Proof. Note that \mathcal{G}° still acts transitively on \tilde{X} by $\lambda_g([x]) = [\lambda_g(x)]$ and that $\tau^{-1}([x]) = (x + \text{Fix}(\mathcal{G})) \cap X$. In particular, $|\tau^{-1}([x])| \leq p$ and thus $|\tilde{X}| \in \{p, p^2\}$.

Suppose that $\tau : X \rightarrow \tilde{X}$ is injective, then $\mathcal{G}(\tilde{X})^\circ$ is isomorphic to \mathcal{A}° and each element of \tilde{X} acts differently on \tilde{X} by the $\tilde{\lambda}$ -action, so \tilde{X} is irretractable. Hence, this contradicts Proposition 3.2. Therefore, $|\tilde{X}| = p$. \square

Proposition 3.6. For all $x \in X$, we have the equality $\mathcal{B}_x = x + \text{Fix}(\mathcal{G}) = x \circ \text{Fix}(\mathcal{G})$. In particular, each block intersects \mathcal{A} in precisely one element.

Proof. By Proposition 3.5, for all $x \in X$, we have $x + \text{Fix}(\mathcal{G}) \subseteq X$. As \mathcal{G}° leaves $\text{Fix}(\mathcal{G})$ invariant under the λ -action, we deduce that the cosets $x + \text{Fix}(\mathcal{G})$ form a nontrivial system of imprimitivity for the λ -action of \mathcal{G}° on X . The uniqueness of such a system (Lemma 1.7) implies that $\mathcal{B}_x = x + \text{Fix}(\mathcal{G})$.

Moreover, it was observed earlier that $\mathcal{A} \cap (g + \text{Fix}(\mathcal{G}))$ is a singleton for each $g \in \mathcal{G}$. Together with Proposition 3.6 this gives the last part of the statement. \square

Proposition 3.7. Let $x \in X$, then $X = \text{Fix}(\mathcal{G}) \circ x \circ \text{Fix}(\mathcal{G})$.

Proof. Let $0 \neq f \in \text{Fix}(\mathcal{G})$. Then ${}^f \mathcal{B}_x = \lambda_f(\mathcal{B}_x) \neq \mathcal{B}_x$ as $f \notin \mathcal{A}$. Therefore, $\text{Fix}(\mathcal{G})$ acts transitively on the system of blocks by conjugation. Furthermore, by Proposition 3.6, $\text{Fix}(\mathcal{G})$ acts transitively on every single block by right-multiplication. Therefore, each $y \in X$ is of the form $y = {}^f x \circ f'$ for some $f, f' \in \text{Fix}(\mathcal{G})$, which can be rewritten as $y = f \circ x \circ f' \circ \bar{f}$. \square

From Proposition 3.7 it therefore follows that we can coordinatize the elements of X in the following way: by Proposition 3.6 we can choose an element $\alpha_{0,0} \in X \cap \mathcal{A}$ and some $0 \neq f \in \text{Fix}(\mathcal{G})$. For $(a, x) \in \mathbb{Z}_p \times \mathbb{Z}_p$, we then set $\alpha_{a,x} = (\bar{f}^{\circ a} \alpha_{0,0}) \circ f^{\circ x}$. By Proposition 3.7, this gives a unique coordinatization of the elements in X .

$\lambda_{\alpha_{0,0}}$ leaves all blocks invariant, therefore we can write $\lambda_{\alpha_{0,0}}(\alpha_{b,0}) = \alpha_{b,-\Phi(b)}$ for some map $\Phi : \mathbb{Z}_p \rightarrow \mathbb{Z}_p$. We can now calculate:

$$\begin{aligned} \alpha_{0,0} \cdot \alpha_{b,y} &= \lambda_{\alpha_{0,0}}^{-1}(\alpha_{b,-\Phi(b)} + (y + \Phi(b))f) \\ &= \alpha_{b,0} + (y + \Phi(b))f \\ &= \alpha_{b,y+\Phi(b)}. \end{aligned}$$

Also, we need to determine

$$f^{\circ a} \cdot \alpha_{b,y} = \lambda_{\bar{f}^{\circ a}}(\bar{f}^{\circ b} \alpha_{0,0} \circ f^{\circ y}) = \bar{f}^{\circ(a+b)} \alpha_{0,0} \circ f^{\circ y} = \alpha_{a+b,y}.$$

We write $\alpha_{a,x} = \bar{f}^{\circ a} \circ \alpha_{0,0} \circ f^{\circ(a+x)}$ and calculate:

$$\begin{aligned} \alpha_{a,x} \cdot \alpha_{b,y} &= f^{\circ(a+x)} \cdot (\alpha_{0,0} \cdot (\bar{f}^{\circ a} \cdot \alpha_{b,y})) \\ &= f^{\circ(a+x)} \cdot (\alpha_{0,0} \cdot \alpha_{b-a,y}) \\ &= f^{\circ(a+x)} \cdot \alpha_{b-a,y+\Phi(b-a)} \\ &= \alpha_{b+x,y+\Phi(b-a)}. \end{aligned}$$

Theorem 3.8. *Let X be an indecomposable, irretractable cycle set with $\mathcal{G}(X)$ a p -group, then X is isomorphic to a cycle set of the form $X = \mathbb{Z}_p \times \mathbb{Z}_p$ with the multiplication*

$$(a, x) \cdot (b, y) = (b + x, y + \Phi(b - a))$$

where $\Phi : \mathbb{Z}_p \rightarrow \mathbb{Z}_p$ is a nonconstant map with $\Phi(A) = \Phi(-A)$ for all $A \in \mathbb{Z}_p$. Vice versa, this construction always results in an indecomposable, irretractable cycle set with $\mathcal{G}(X)$ a p -group.

Proof. In the preceding calculations, we have already established that the given multiplication rule is necessary. We now determine

$$\begin{aligned} ((a, x) \cdot (b, y)) \cdot ((a, x) \cdot (c, z)) &= (b + x, y + \Phi(b - a)) \cdot (c + x, z + \Phi(c - a)) \\ &= (c + x + y + \Phi(b - a), z + \Phi(c - a) + \Phi(c - b)). \end{aligned}$$

Similarly,

$$((b, y) \cdot (a, x)) \cdot ((b, y) \cdot (c, z)) = (c + y + x + \Phi(a - b), z + \Phi(c - b) + \Phi(c - a)).$$

A comparison shows that in order for X to satisfy Eq. (C1), $\Phi(b - a) = \Phi(a - b)$ must hold for all $a, b \in \mathbb{Z}_p$ which amounts to saying that $\Phi(A) = \Phi(-A)$ for all $A \in \mathbb{Z}_p$. By the same calculation, one sees that this obstruction to Φ is sufficient for X to satisfy Eq. (C1). By construction, all maps $\sigma_{(a,x)}$ are bijective. Furthermore, the square map

$$\text{Sq}(a, x) = (a, x) \cdot (a, x) = (a + x, x + \Phi(0))$$

is also quickly seen to be bijective. Finally, irretractability is the same as saying that for any $a, a' \in \mathbb{Z}_p$, there is at least one $b \in \mathbb{Z}_p$ such that $\Phi(b - a) = \Phi(b - a')$. But this is clearly equivalent to Φ not being constant.

Finally, note that $\mathcal{G}(X)$ clearly acts transitively on the system of blocks $\{a\} \times \mathbb{Z}_p$. Furthermore, for a $b \in \mathbb{Z}_p$ with $\Phi(b) \neq 0$, we clearly have $(0, 0) \cdot (b, y) = (b, y + \Phi(b)) \neq (b, y)$ which shows that the $\mathcal{G}(X)$ -orbit of (b, y) contains at least the block $\{b\} \times \mathbb{Z}_p$. This implies that $\mathcal{G}(X)$ acts transitively on X , hence X is indecomposable. Also note that $\mathcal{G}(X)$ is contained in $\mathbb{Z}_p \wr \mathbb{Z}_p \leq \text{Sym}_{\mathbb{Z}_p \times \mathbb{Z}_p}$, hence $\mathcal{G}(X)$ is a p -group. \square

We note the following corollary, which will be useful later in Section 4:

Corollary 3.9. *The elements in $X \cap \mathcal{A}$ generate the whole cycle set.*

Proof. Note that, using the explicit form in Theorem 3.8, we are considering the set $X \cap \mathcal{A} = \{(a, 0) \mid a \in \mathbb{Z}_p\}$. As Φ is nonconstant, we easily see that these elements generate the whole cycle set X . \square

3.2. Getting rid of redundancy and determining automorphisms. The aim of this subsection is to determine unique representatives for the irretractable cycle sets determined in Section 3 and moreover, describe their automorphism groups.

Let \mathcal{F}_p be the set of all nonconstant functions $\Phi : \mathbb{Z}_p \rightarrow \mathbb{Z}_p$ with the property that $\Phi(A) = \Phi(-A)$ for all $A \in \mathbb{Z}_p$. \mathcal{F}_p is acted upon by \mathbb{Z}_p^* via $({}^\alpha\Phi)(A) = \alpha^{-1}\Phi(\alpha A)$. From now on, let \mathcal{R}_p be a fixed system of representatives for this action.

Recall from Theorem 3.8 that the cycle sets in the considered case are described as $\mathbb{Z}_p \times \mathbb{Z}_p$ with the operation

$$(a, x) \cdot (b, y) = (b + x, y + \Phi(b - a))$$

where $\Phi \in \mathcal{F}_p$.

By Proposition 3.2, $\mathcal{G}(X)$ is nonabelian and, by Lemma 1.7, has a unique non-trivial system of imprimitivity that consists of the blocks $\{a\} \times \mathbb{Z}_p$. Assume that $X = \mathbb{Z}_p \times \mathbb{Z}_p$ comes with two cycle set operations \cdot, \cdot' that are given by the parameters $\Phi, \Phi' \in \mathcal{F}_p$ and that $\phi : (X, \cdot) \rightarrow (X, \cdot')$ is an isomorphism. Then $\phi\mathcal{G}(X, \cdot)\phi^{-1} = \mathcal{G}(X, \cdot')$ so in particular ϕ must normalize the cyclic permutation action on the blocks and thus be of the form $\phi(a, x) = (\alpha a + \beta, \pi_a(x))$ for some $\pi_a \in \text{Sym}_{\mathbb{Z}_p}$, $\alpha \in \mathbb{Z}_p^*$, $\beta \in \mathbb{Z}_p$. We now calculate

$$\begin{aligned} \phi((a, x) \cdot (b, y)) &= (\alpha(b + x) + \beta, \pi_{b+x}(y + \Phi(b - a))), \\ \phi(a, x) \cdot' \phi(b, y) &= (\alpha b + \beta + \pi_a(x), \pi_b(y) + \Phi'(\alpha(b - a))). \end{aligned}$$

Equating these terms results in $\pi_a(x) = \alpha x$, considering the first coordinate. Taking this into account when considering the second coordinate leaves us with the equation

$$\alpha(y + \Phi(b - a)) = \alpha y + \Phi'(\alpha(b - a)) \Leftrightarrow \Phi(b - a) = \alpha^{-1}\Phi'(\alpha(b - a)).$$

This shows that Φ, Φ' define isomorphic cycle sets if and only if there is an $\alpha \in \mathbb{Z}_p^*$ such that $\Phi = {}^\alpha\Phi'$. Putting $\Phi = \Phi'$, the same considerations prove that ϕ provides an automorphism of a solution with parameter Φ if and only if $\phi(a, x) = (\alpha x + \beta, \alpha x)$ for $\alpha \in \mathbb{Z}_p^*$, $\beta \in \mathbb{Z}_p$ with ${}^\alpha\Phi = \Phi$.

We conclude:

Theorem 3.10. *Let X be an indecomposable irretractable cycle set of size p^2 where p is a prime.*

- (1) *If $\mathcal{G}(X)$ is a p -group, then there is a unique $\Phi \in \mathcal{R}_p$ such that X is isomorphic to the cycle set on $X = \mathbb{Z}_p \times \mathbb{Z}_p$ with multiplication*

$$(a, x) \cdot (b, y) = (b + x, y + \Phi(b - a)).$$

- (2) *Let X, Φ be as in the previous item. Then any automorphism of X is of the form $(a, x) \mapsto (\alpha a + \beta, \alpha x)$ for some $\alpha \in \mathbb{Z}_p^*$, $\beta \in \mathbb{Z}_p$ with ${}^\alpha\Phi = \Phi$.*

4. INDECOMPOSABLE IRRETRACTABLE CYCLE SETS OF SIZE p^2 - THE GENERAL CASE

In this section we will now obtain a solution to the general problem:

Problem 4.1. Let p be a prime. Find all indecomposable irretractable cycle sets X of size p^2 .

Before restricting to this specific case, we prove a useful lemma. Given a brace A and a subset $S \subseteq A$ we define

$$\text{Fix}_A(S) = \{a \in A \mid \lambda_s(a) = a \text{ for all } s \in S\}.$$

In fact, it follows from Lemma 1.1 that $\lambda_{\bar{a}}(\bar{a}s) = -a + s + \lambda_s(a)$, so $\lambda_s(a) = a$ if and only if $\lambda_a(\bar{a}s) = s$, which is equivalent to $\lambda_{\bar{a}}(s) = \bar{a}s$. So alternatively

$$\text{Fix}_A(S) = \{a \in A \mid \lambda_{\bar{a}}(s) = \bar{a}s \text{ for all } s \in S\}.$$

Lemma 4.2. *Let A be a finite brace, L a left ideal of A and G a normal subgroup of A° , then $\text{Fix}_A(L \cap G)$ is a subbrace of A contained in the normaliser of $L \cap G$ in A°*

Proof. From the original definition of $\text{Fix}_A(L \cap G)$ we see that $\text{Fix}_A(L \cap G)^+$ is a group. Let $a \in \text{Fix}_A(L \cap G)$ and $b \in L \cap G$. Then $\lambda_{\bar{a}}(b) \in L$, but as $\lambda_{\bar{a}}(b) = \bar{a}b$, also $\lambda_{\bar{a}}(b) \in G$. We conclude that $\lambda_{\bar{a}}(b) \in L \cap G$. Using the alternative description of $\text{Fix}_A(L \cap G)$ we now see that $\text{Fix}_A(L \cap G)^\circ$ is multiplicatively closed and non-empty. As A is finite, we conclude that $\text{Fix}_A(L \cap G)^\circ$ is a subgroup of A° . Therefore $\text{Fix}_A(L \cap G)$ is a subbrace. As $\bar{a}b \in L \cap G$ for all $a \in \text{Fix}_A(L \cap G)$, $b \in L \cap G$ we find indeed that $\text{Fix}_A(L \cap G)$ is contained in the normaliser of $L \cap G$ in A° . \square

In the remainder of the section, we let X be an indecomposable irretractable cycle set of size p^2 and $\mathcal{G} = \mathcal{G}(X)$. As we already covered the case where \mathcal{G} is a p -group in Section 3, we can assume that \mathcal{G} is not a p -group. We associate X with its image in \mathcal{G} , which is a transitive cycle base. As X is irretractable, it follows from [2, Lemma 2.1.] that $\text{Soc}(\mathcal{G}) = 0$.

Let \mathcal{G}_p be the Sylow p -subgroup of \mathcal{G}^+ and let $\mathcal{G}_{p'}$ be the Hall p' -subgroup of \mathcal{G}^+ , both are characteristic in \mathcal{G}^+ hence they are left ideals of \mathcal{G} . We denote $X = \{x_1, \dots, x_{p^2}\}$ and for $1 \leq i \leq p^2$ we define $y_i \in \mathcal{G}_p$ and $z_i \in \mathcal{G}_{p'}$ such that $x_i = y_i + z_i$. As the λ -action of \mathcal{G}_p is transitive on X , and therefore also on $Y = \{y_i \mid 1 \leq i \leq p^2\}$, we find that Y is a transitive cycle base of the brace \mathcal{G}_p . In particular, this implies that $|Y| \in \{1, p, p^2\}$.

Let $q \neq p$ be prime and \mathcal{G}_q the q -Sylow subgroup of \mathcal{G}^+ which is a left ideal by the same argument as above. $\text{Fix}_{\mathcal{G}}(\mathcal{G}_q)$ is a brace by Lemma 4.2 and $Y \cap \text{Fix}_{\mathcal{G}}(\mathcal{G}_q)$ is a sub-cycle set of Y . Now if Y has finite multipermutation level, then $Y \cap \text{Fix}_{\mathcal{G}}(\mathcal{G}_q)$ is either empty or equal to Y by [4, Theorem 5.1]. As $|Y|$ is a p -power, \mathcal{G}_q fixes at least one point in Y under the λ -action. This means that $Y \cap \text{Fix}_{\mathcal{G}}(\mathcal{G}_q) = Y$ and thus $\mathcal{G}_p \subseteq \text{Fix}_{\mathcal{G}}(\mathcal{G}_q)$. It follows that \mathcal{G}_p normalizes \mathcal{G}_q but as \mathcal{G}_p acts faithful and transitive on a set of size p^2 , this implies that $\mathcal{G}_q = 0$ and hence, $\mathcal{G}_{p'} = 0$ but this contradicts the assumption that \mathcal{G} is not a p -group. We therefore deduce that Y is not of finite multipermutation level. Together with the earlier observation that $|Y| \in \{1, p, p^2\}$, we conclude that Y is irretractable of size p^2 and therefore as described in Theorem 3.8. In particular, we find that \mathcal{G}_p° is not abelian.

From now on we consider the unique block system of X under the action of \mathcal{G} , and recall that the uniqueness is guaranteed by Lemma 1.7. As before we denote this by $\{\mathcal{B}_x \mid x \in X\}$. We denote the subgroup of \mathcal{G}° that fixes the blocks setwise by \mathcal{A} . Also, we define $\mathcal{A}_p = \mathcal{A} \cap \mathcal{G}_p$ and $\mathcal{A}_{p'} = \mathcal{A} \cap \mathcal{G}_{p'}$. Note that in particular, \mathcal{A}_p is a p -Sylow subgroup of \mathcal{A}° and $\mathcal{A}_{p'}$ is a Hall p' -subgroup of \mathcal{A}° . In particular, \mathcal{A}_p is normal in \mathcal{G}° by Proposition 1.8.

Let $a \in \mathcal{A}_p$ and $g \in \mathcal{G}_{p'}$, then Lemma 1.1 yields $\lambda_g(\bar{g}a) = -g + a + \lambda_a(g)$ hence $-a + \lambda_g(\bar{g}a) = -g + \lambda_a(g)$. As $-a + \lambda_g(\bar{g}a)$ is contained in \mathcal{G}_p and $-g + \lambda_a(g)$ is contained in $\mathcal{G}_{p'}$, we find that $\lambda_a(g) = g$. By Lemma 1.1 this implies that $^ga = \lambda_g(a)$, so the λ -action of $\mathcal{G}_{p'}$ restricts to \mathcal{A}_p , so also to $Y \cap \mathcal{A}_p$.

By Proposition 3.6 we know that \mathcal{A}_p contains a unique representative of each block in the block system of Y under the action of $\mathcal{G}_{p'}^\circ$, hence also under the action

of \mathcal{G}° . This means that $\mathcal{A}_{p'}$ acts trivially on the set $Y \cap \mathcal{A}_p$ and thus $Y \cap \mathcal{A}_p \subseteq \text{Fix}_{\mathcal{G}}(\mathcal{A}_{p'})$. From Lemma 4.2 we know that $\text{Fix}_{\mathcal{G}}(\mathcal{A}_{p'})$ is a subbrace, since $\mathcal{A}_{p'} = \mathcal{A} \cap \mathcal{G}_{p'}$. In particular, $\text{Fix}_{\mathcal{G}}(\mathcal{A}_{p'}) \cap Y$ is a sub-cycle set of Y which contains $Y \cap \mathcal{A}_p$, but from Corollary 3.9 it then follows that $\text{Fix}_{\mathcal{G}}(\mathcal{A}_{p'}) \cap Y = Y$. However, as the λ -action of \mathcal{G} on Y is faithful, this implies that $\mathcal{A}_{p'} = 0$.

In particular, from Proposition 1.8 we find that $|\mathcal{G}_{p'}| < p$ and thus the λ -action of \mathcal{G}_p on $\mathcal{G}_{p'}$ is trivial. As this same action is transitive on Z we find $|Z| = 1$. Let $z \in Z$, then λ_z is a brace automorphism of \mathcal{G}_p by Lemma 1.1 and therefore also its restriction to Y yields a cycle set automorphism. For any $x_i, x_j \in X$ we find

$$x_i \cdot x_j = (y_i + z) \cdot (y_j + z) = \lambda_{y_i+z}^{-1}(y_j + z) = \lambda_z^{-1}(y_i \cdot y_j) + z.$$

We find that the cycle set structure on X is obtained by deforming the cycle set structure on Y by an automorphism of Y . Lemma 4.5 shows that such a deformation is always possible.

Remark 4.3. We remark that the idea of starting from a cycle set X and then considering its projection Y onto the p -Sylow subgroup \mathcal{G}_p , is strongly related to the notion of cabling, see [10, 15]. Recall that for $k \in \mathbb{Z}$, the k -cabled cycle set is defined as (X, \cdot_k) with $x \cdot_k y = (k\sigma_x)(y)$, where $k\sigma_x \in \mathcal{G}(X)$ is the k -th power of σ_x in $(\mathcal{G}(X), +)$.

In particular, $\mathcal{G}(X, \cdot_k)$ is a subgroup of $\mathcal{G}(X, \cdot)^\circ$. We now claim that it is even a subbrace. If we denote the σ -maps of (X, \cdot_k) by σ'_x , then we find that the addition $+_k$ in $\mathcal{G}(X, \cdot_k)$ is given by

$$\begin{aligned} \sigma'_x +_k \sigma'_y &= \sigma'_x \circ \sigma'_{x \cdot_k y} \\ &= (k\sigma_x) \circ (k\sigma_{(k\sigma_x)(y)}) \\ &= (k\sigma_x) + \lambda_{k\sigma_x}(k\sigma_{(k\sigma_x)(y)}) \\ &= (k\sigma_x) + k\lambda_{k\sigma_x}(\sigma_{(k\sigma_x)(y)}) \\ &= (k\sigma_x) + (k\sigma_y) = \sigma'_x + \sigma'_y, \end{aligned}$$

where $+$ denotes the addition in $\mathcal{G}(X, \cdot)$.

Let $|\mathcal{G}(X, \cdot)| = p^r m$ where $(p, m) = 1$. As the additive p -Sylow subgroup is $\mathcal{G}(X, \cdot)_p = \{kg \mid g \in \mathcal{G}\}$ for any non-zero multiple k of m , we find that $\mathcal{G}(X, \cdot)_p = \mathcal{G}(X, \cdot_k)$. In particular, $\text{Ret}(X, \cdot_k)$ is isomorphic to the cycle set $\{k\sigma_x \mid x \in X\} \subseteq \mathcal{G}(X, \cdot)$ where the equivalence class of x is mapped to $k\sigma_x$. If we let k be such that $m \equiv 1 \pmod{p^r}$ and $m \equiv 0 \pmod{k}$ then we find that the $k\sigma_x$ is precisely the projection of σ_x onto $\mathcal{G}(X, \cdot)_p$, hence in this case we find that $Y \cong \text{Ret}(X, \cdot_k)$.

Proposition 4.4. *Let (X, \cdot) be an indecomposable cycle set of order p^n , with p a prime. Let k be the largest divisor of $|\mathcal{G}(X, \cdot)|$ coprime to p . If (X, \cdot_k) has finite multipermutation level, then $\mathcal{G}(X, \cdot)$ is a p -group and thus $k = 1$.*

Proof. The proof is essentially the same as how we proved earlier in this section that Y is not of finite multipermutation level.

First of all, as $\mathcal{G}(X, \cdot_k)^\circ$ is a p -Sylow subgroup, (X, \cdot_k) is still an indecomposable cycle set. Now let $Y = \{k\sigma_x \mid x \in X\} \cong \text{Ret}(X, \cdot_k)$. As (X, \cdot_k) has finite multipermutation level, so does Y .

Let $q \neq p$ be a prime and let $\mathcal{G}(X, \cdot)_q$ be the q -Sylow subgroup of $\mathcal{G}(X, \cdot)^+$. If we consider the λ -action of $\mathcal{G}(X, \cdot)_q$ on Y we find that it has fixed points. By Lemma 4.2 we find that $\text{Fix}(\mathcal{G}(X, \cdot)_q)$ is a brace thus $\text{Fix}(\mathcal{G}(X, \cdot)_q) \cap Y$ is a sub-cycle set. As this sub-cycle set is non-empty, [4, Theorem 5.1] implies that $Y \subseteq \text{Fix}(\mathcal{G}(X, \cdot)_q)$ and hence $\mathcal{G}(X, \cdot)_p \subseteq \text{Fix}(\mathcal{G}(X, \cdot)_q)$. As a result, $\mathcal{G}(X, \cdot)_q$ is a normal subgroup of $\mathcal{G}(X, \cdot)^\circ$, but this is impossible as $\mathcal{G}(X, \cdot)^\circ$ acts transitively and faithfully on a set of p -power order. We conclude that $\mathcal{G}(X, \cdot)$ is a p -group and therefore $k = 1$. \square

Lemma 4.5. *Let (X, \cdot) be a cycle set and ϕ be an automorphism of (X, \cdot) . Then the following statements hold:*

(1) *X is a cycle set for the operation*

$$x \cdot_\phi y = \phi(x \cdot y).$$

(2) *If (X, \cdot) is irretractable, then so is (X, \cdot_ϕ) .*

Proof. One can verify directly that (X, \cdot_ϕ) satisfies (C1)-(C3). However, as this is useful in the proof of Lemma 4.6 we construct (X, \cdot_ϕ) as the sub-cycle set of a cycle set coming from a brace.

By the functoriality of $G(X, r)$ we get an induced automorphism ϕ' of $G(X, r)$ which restricts to ϕ on the generating set $X \subseteq G(X, \cdot)$. We let \mathbb{Z} denote the trivial brace on \mathbb{Z} and we let \mathbb{Z}° act on $G(X, \cdot)$ by $1 \cdot g = (\phi')^{-1}(g)$ for $g \in G(X, \cdot)$. Hence we can construct the semi-direct product $G(X, \cdot) \rtimes \mathbb{Z}$ in the sense of [19, Corollary 3.37]. The set $X + 1 \subseteq G(X, \cdot) \rtimes \mathbb{Z}$ is closed under the λ -action. Hence it is a sub-cycle set which is precisely the cycle set in the statement under the correspondence $x \mapsto x + 1$.

If (X, \cdot) is irretractable, then also (X, \cdot_ϕ) is irretractable as it follows directly that $x \cdot z = y \cdot z$ if and only if $x \cdot_\phi z = y \cdot_\phi z$ for $x, y, z \in X$. \square

Lemma 4.6. *Let (X, \cdot) be a cycle set and ϕ an automorphism of (X, \cdot) of order m coprime to $|\mathcal{G}(X)|$ such that ϕ has a fixed point. Then $\mathcal{G}(X, \cdot_\phi)^\circ = \mathcal{G}(X, \cdot)^\circ \rtimes \langle \phi \rangle$ as subgroups of Sym_X and $\mathcal{G}(X, \cdot_\phi) \cong \mathcal{G}(X, \cdot) \rtimes \mathbb{Z}_m$ as braces where \mathbb{Z}_m is given the trivial brace structure and acts on $\mathcal{G}(X, \cdot)$ by ϕ .*

In particular, if (X, \cdot) is indecomposable then so is (X, \cdot_ϕ) and $(X, \cdot) = (X, (\cdot_\phi)_k)$ for any $k \in \mathbb{Z}$ such that $k \equiv 0 \pmod{m}$ and $k \equiv 1 \pmod{|\mathcal{G}(X)|}$.

Proof. Let $G(X, \cdot) \rtimes \mathbb{Z}$ be as described in the proof of Lemma 4.5. The universal property of $G(X, \cdot_\phi)$ provides a brace homomorphism $f : G(X, \cdot_\phi) \rightarrow G(X, \cdot) \rtimes \mathbb{Z}$ mapping $x \mapsto x + 1$. The canonical map $g : (G(X, \cdot) \rtimes \mathbb{Z})^\circ \rightarrow \text{Sym}_X$ sending $x \mapsto \sigma_x^{-1}$ and $1 \mapsto \phi^{-1}$ is a group homomorphism. Clearly $\ker gf = \text{Soc}(G(X, \cdot_\phi))$ and the image is precisely $\mathcal{G}(X, \cdot_\phi)$. As $\mathcal{G}(X, \cdot) \cap \langle \phi \rangle = \{\text{id}\}$ we find that $\mathcal{G}(X, \cdot_\phi)^\circ = \mathcal{G}(X, \cdot)^\circ \rtimes \langle \phi \rangle$. Also, as $\ker g = \text{Soc}(G(X, \cdot)) \rtimes m\mathbb{Z}$, we find that $\mathcal{G}(X, \cdot_\phi) \cong \mathcal{G}(X, \cdot) \rtimes \mathbb{Z}_m$ as braces.

It follows directly that if $\mathcal{G}(X, \cdot)$ acts transitively on X then so does $\mathcal{G}(X, \cdot_\phi)$.

As $\mathcal{G}(X, \cdot)$ is a Hall-subgroup of $\mathcal{G}(X, \cdot_\phi)^+$, the last part of the statement follows from Remark 4.3. \square

For an element g of a finite group G , we denote its order by $o(g)$.

Lemma 4.7. *Let $(X, \cdot), (X, \cdot')$ be cycle sets and $\phi \in \text{Aut}(X, \cdot), \psi \in \text{Aut}(X, \cdot')$ such that $\gcd(|\mathcal{G}(X, \cdot)|, |\mathcal{G}(X, \cdot')|, o(\phi)o(\psi)) = 1$. Then $f : (X, \cdot_\phi) \rightarrow (X, \cdot'_\psi)$ is an isomorphism if and only if $f : (X, \cdot) \rightarrow (X, \cdot')$ is an isomorphism and $\psi = f\phi f^{-1}$. In particular, $\text{Aut}(X, \cdot_\phi)$ is precisely the centraliser of ϕ in $\text{Aut}(X, \cdot)$.*

Proof. Assume that $f : (X, \cdot_\phi) \rightarrow (X, \cdot'_\psi)$ is an isomorphism. From Lemma 4.6 and the assumptions we find the existence of some $k \in \mathbb{Z}$ such that $(X, (\cdot_\phi)_k) = (X, \cdot)$ and $(X, (\cdot'_\psi)_k) = (X, \cdot')$. The functoriality of cabling now yields that f induces an isomorphism $f : (X, \cdot) \rightarrow (X, \cdot')$. Now for any $x, y \in X$ we find $f(x \cdot_\phi y) = f\phi(x \cdot y)$ and $f(x \cdot_\psi y) = \psi f(x \cdot y)$, hence $f(x \cdot_\phi y) = f(x) \cdot_\psi f(y)$ if and only if $\psi = f\phi f^{-1}$. This proves one implication of the statement.

Assume that $f : (X, \cdot) \rightarrow (X, \cdot')$ is an isomorphism and $\psi = f\phi f^{-1}$. Then

$$f(x \cdot_\phi y) = f\phi(x \cdot y) = \psi(f(x \cdot y)) = \psi(f(x) \cdot' f(y)) = f(x) \cdot'_\psi f(y),$$

for all $x, y \in X$. \square

Theorem 4.8. *Let X be an irretractable cycle set of size p^2 where p is a prime. Then there exists a unique $\Phi \in \mathcal{R}_p$ and $\alpha \in \mathbb{Z}_p^*$ satisfying ${}^\alpha\Phi = \Phi$ such that X is isomorphic to the cycle set on $\mathbb{Z}_p \times \mathbb{Z}_p$ with multiplication*

$$(a, x) \cdot (b, y) = (\alpha b + \alpha x, \alpha y + \alpha \Phi(b - a)).$$

If $\alpha = 1$, then the cycle sets are the ones that appear in Theorem 3.10. If $\alpha \neq 1$, then any automorphism of (X, \cdot) is of the form $(a, x) \mapsto (\gamma a, \gamma x)$ for some $\gamma \in \mathbb{Z}_p^$ with ${}^\gamma\Phi = \Phi$.*

Proof. From the discussion preceding Lemma 4.5 we know that (X, \cdot) can be obtained by starting from a cycle set structure on X whose permutation group is a p -group and deforming such cycle set by an automorphism of order coprime to p , in the sense of Lemma 4.5. From Theorem 3.10 it follows that, up to a cycle set isomorphism, $X = \mathbb{Z}_p \times \mathbb{Z}_p$ and

$$(a, x) \cdot (b, y) = (\alpha b + \alpha x + \beta, \alpha y + \alpha \Phi(b - a)),$$

for some $\Phi \in \mathcal{R}_p$, $\alpha \in \mathbb{Z}_p^*$ and $\beta \in \mathbb{Z}_p$, satisfying ${}^\alpha\Phi = \Phi$. By Lemma 4.7 we may even assume $\beta = 0$. We therefore get that up to isomorphism the multiplication on X is precisely as in the statement.

Conversely, it follows directly from Lemma 4.5 and Lemma 4.6 that $\mathbb{Z}_p \times \mathbb{Z}_p$ with the given multiplication always yields an indecomposable irretractable cycle set. As a consequence of Lemma 4.7 we find that different choices of α and Φ yield non-isomorphic solutions and also that the automorphisms are the ones described in the statement. \square

5. SUMMARY

We summarize our classification result in the following theorem.

Theorem 5.1. *Let X be an indecomposable cycle set of size p^2 . Then X is isomorphic to a cycle set of one of the following forms:*

- (1) $X = \mathbb{Z}_{p^2}$, $x \cdot y = y + 1$,
- (2) $X = \mathbb{Z}_p \times \mathbb{Z}_p$,

$$(a, x) \cdot (b, y) = (b + 1, y + S\chi_0(b) + \Phi(b - a))$$

where $\Phi : \mathbb{Z}_p \rightarrow \mathbb{Z}_p$ is a nonconstant map with $\Phi(0) = 0$, $S \in \mathbb{Z}_p$ and

$$\chi_0 : \mathbb{Z}_p \rightarrow \mathbb{Z}_p$$

$$\chi_0(x) = \begin{cases} 1 & x = 0 \\ 0 & x \neq 0 \end{cases}.$$

The parameters S, Φ and S', Φ' define isomorphic cycle sets if and only if $S = S'$ and $\alpha\Phi = \Phi'$ for some $\alpha \in \mathbb{Z}_p^$.*

- (3) $X = \mathbb{Z}_p \times \mathbb{Z}_p$,

$$(a, x) \cdot (b, y) = (\alpha b + \alpha x, \alpha y + \alpha \Phi(b - a))$$

where $\Phi : \mathbb{Z}_p \rightarrow \mathbb{Z}_p$ is a nonconstant map with $\Phi(x) = \Phi(-x)$ and $\alpha \in \mathbb{Z}_p^*$ is such that $\Phi(\alpha x) = \alpha\Phi(x)$.

The parameters α, Φ and α', Φ' define isomorphic cycle sets if and only if $\alpha = \alpha'$ and there is a $\beta \in \mathbb{Z}_p^$ such that $\beta^{-1}\Phi(\beta x) = \Phi'(x)$ for all $x \in \mathbb{Z}_p$.*

These three cases are mutually exclusive.

Proof. Corollary 2.4 tells us that the indecomposable cycle sets of size p^2 that have finite multipermutation level 1 and 2, are exactly the ones described in (1) resp. (2). On the other hand, the irretractable cycle sets are classified, up to isomorphism in Theorem 4.8 and make up case (3). \square

5.1. Indecomposable set-theoretic solutions of size p^2 . Recall that given a cycle set on X , the associated solution is given by

$$r_X(x, y) = (\sigma_x^{-1}(y), \sigma_x^{-1}(y) \cdot x).$$

Therefore we can obtain all indecomposable solutions of size p^2 simply translating the cycle sets obtained in Theorem 5.1 to set-theoretical solutions. In case (1) we obtain solutions of the form $r(x, y) = (y - 1, x + 1)$, as $\sigma_x : y \mapsto y + 1$, thus $\lambda_x = \sigma_x^{-1} : y \mapsto y - 1$ and $\rho_y : x \mapsto \sigma_x^{-1}(y) \cdot x = (y - 1) \cdot x = x + 1$.

In case (2) we have that $\sigma_{(a,x)} : (b, y) \mapsto (b + 1, y + S\chi_0(b) + \Phi(b - a))$, hence

$$\lambda_{(a,x)}(b, y) = \sigma_{(a,x)}^{-1}(b, y) = (b - 1, y - S\chi_0(b - 1) - \Phi(b - 1 - a))$$

and

$$\begin{aligned} \rho_{(b,y)}(a, x) &= \sigma_{(a,x)}^{-1}(b, y) \cdot (a, x) = (b - 1, y - S\chi_0(b - 1) - \Phi(b - 1 - a)) \cdot (a, x) \\ &= (a + 1, x + S\chi_0(a) + \Phi(a - b + 1)). \end{aligned}$$

Thus the associated solution is

$$r \begin{pmatrix} (a, x) \\ (b, y) \end{pmatrix} = \begin{pmatrix} (b - 1, y - S\chi_0(b - 1) - \Phi(b - 1 - a)) \\ (a + 1, x + S\chi_0(a) + \Phi(a - b + 1)) \end{pmatrix}$$

Finally in case (3) we have $\sigma_{(a,x)} : (b, y) \mapsto (\alpha b + \alpha x, \alpha y + \alpha \Phi(b - a))$, hence, using that $\Phi(\alpha x) = \alpha \Phi(x)$ for all $x \in \mathbb{Z}_p$,

$$\lambda_{(a,x)}(b, y) = \sigma_{(a,x)}^{-1}(b, y) = (\alpha^{-1}b - x, \alpha^{-1}y - \Phi(\alpha^{-1}b - x - a))$$

and

$$\begin{aligned} \rho_{(b,y)}(a, x) &= \sigma_{(a,x)}^{-1}(b, y) \cdot (a, x) = (\alpha^{-1}b - x, \alpha^{-1}y - \Phi(\alpha^{-1}b - x - a)) \cdot (a, x) \\ &= (\alpha a + \alpha(\alpha^{-1}(y - \Phi(b - \alpha x - \alpha a))), \alpha x + \alpha \Phi(a - \alpha^{-1}(b - \alpha x))) \\ &= (\alpha a + y - \Phi(b - \alpha x - \alpha a), \alpha x + \Phi(\alpha a - b + \alpha x)). \end{aligned}$$

Thus the associated solution is

$$r \begin{pmatrix} (a, x) \\ (b, y) \end{pmatrix} = \begin{pmatrix} (\alpha^{-1}b - x, \alpha^{-1}y - \Phi(\alpha^{-1}b - x - a)) \\ (\alpha a + y - \Phi(b - \alpha x - \alpha a), \alpha x + \Phi(\alpha a - \alpha x - b)) \end{pmatrix}$$

Note that these solutions are the same constructed in [6, Theorem 5.1].

Theorem 5.2 ([6, Theorem 5.1]). *Let p be a prime number. Let $t \in \mathbb{Z}_p$ be a nonzero element. Let $f : \mathbb{Z}_p \rightarrow \mathbb{Z}_p$ be a map such that*

- (S1) $f(i) = f(-i)$ for all $i \in \mathbb{Z}_p$,
- (S2) $f(t^s i) = t^s f(i) - (t^s - 1)f(0)$ for all $i \in \mathbb{Z}_p$ and $s \in \mathbb{Z}$,
- (S3) f is not a constant map.

Let $X = \mathbb{Z}_p \times \mathbb{Z}_p$ and $r : X^2 \rightarrow X^2$ be the map $r \begin{pmatrix} (i, j) \\ (k, l) \end{pmatrix} = \begin{pmatrix} \lambda_{(i,j)}(k, l) \\ \lambda_{\lambda_{(i,j)}(k, l)}^{-1}(i, j) \end{pmatrix}$, where $\lambda_{(i,j)}(k, l) = (tk + j, t(l - f(tk + j - i)))$. Then (X, r) is a simple solution of the YBE in the following sense: if (Y, s) is a solution and $f : X \rightarrow Y$ is a surjection with $f(r \times r) = (s \times s)f$, then $|Y| \in \{1, |X|\}$ ¹.

We will denote the solution associated with cycle sets of the form (3) in Theorem 5.1 with parameters Φ, α as $r^{\alpha, \Phi}$ with first component

$$\lambda_{(a,x)}^{\Phi, \alpha} : (b, y) \mapsto (\alpha^{-1}b - x, \alpha^{-1}y - \Phi(\alpha^{-1}b - x - a)).$$

¹This means that the corresponding cycle sets are simple in the sense that there are no non-trivial quotients.

Similarly, we will denote the solution constructed in Theorem 5.2 with parameters f, t as $r^{f,t}$ with first component

$$\ell_{(i,j)}^{f,t} : (k, l) \mapsto (tk + j, t(l - f(tk + j - i))).$$

With this notation and fixing $X = \mathbb{Z}_p \times \mathbb{Z}_p$, it is easy to prove that the map $\Psi(i, j) = (i, -j)$ is an isomorphism of solutions $\Psi : (X, r^{\Phi, \alpha}) \rightarrow (X, r^{f_{\Phi, \alpha}, \alpha^{-1}})$, where $f_{\Phi, \alpha} : i \mapsto -\Phi(\alpha i)$. In fact,

$$\begin{aligned} \lambda_{\Psi(i,j)}^{\Phi, \alpha}(\Psi(k, l)) &= \lambda_{(i,-j)}^{\Phi, \alpha}((k, -l)) = (\alpha^{-1}(k + \alpha j), \alpha^{-1}(-l - \Phi(k + \alpha j - \alpha i))) \\ &= (\alpha^{-1}k + j, \alpha^{-1}(-l + f_{\Phi, \alpha}(\alpha^{-1}k + j - i))) \\ &= \Psi(\alpha^{-1}k + j, \alpha^{-1}(l - f_{\Psi, \alpha}(k + \alpha j - \alpha i))) \\ &= \Psi\left(\ell_{(i,j)}^{f_{\Phi, \alpha}, \alpha^{-1}}(k, l)\right). \end{aligned}$$

It remains to show that, with the conditions for Φ and α given in Theorem 5.1, the parameters $f = f_{\Phi, \alpha}$ and $t = \alpha^{-1}$ satisfy the properties required by Theorem 5.2. Since Φ satisfies (S1) and (S3), so does $f_{\Phi, \alpha}$. Moreover, since $\Phi(\alpha i) = \alpha\Phi(i)$, we have that $f_{\Phi, \alpha}(\alpha^{-s}i) = -\Phi(\alpha\alpha^{-s}i) = -\alpha^{-s}\Phi(\alpha i) = \alpha^{-s}f_{\Phi, \alpha}(i)$. Hence $f_{\Phi, \alpha}$ satisfies (S2) if and only if $(\alpha^{-s} - 1)f_{\Phi, \alpha}(0) = 0$ for all $s \in \mathbb{Z}$, which is equivalent to $(\alpha - 1)\Phi(0) = 0$. But the latter is a consequence of the properties of Φ and α as $\Phi(0) = \Phi(\alpha 0) = \alpha\Phi(0)$.

In particular, we have answered [6, Question 7.4] affirmatively.

5.2. Enumeration of indecomposable, irretractable cycle sets of size p^2 .

In this subsection, we will use the following convention: for a group G acting on a set X by an action $(g, x) \mapsto {}^g x$, we denote the set of fixed points by

$$\text{Fix}_X(G) = \{x \in X \mid {}^g x = x \text{ for all } g \in G\}.$$

Recall that \mathcal{F}_p has been defined as the set of all nonconstant maps $\Phi : \mathbb{Z}_p \rightarrow \mathbb{Z}_p$ such that $\Phi(-A) = \Phi(A)$.

By Theorem 5.1, every irretractable cycle sets can be described by a pair (Φ, α) where $\alpha \in \mathbb{Z}_p^*$ satisfies ${}^\alpha \Phi = \Phi$ for all $A \in \mathbb{Z}_p$, where ${}^\alpha \Phi : A \mapsto \alpha^{-1}\Phi(\alpha A)$. Note that this is an action of $\mathbb{Z}_p^* \cong \mathbb{Z}_{p-1}$ on \mathcal{F}_p . Furthermore, (Φ, α) and (Φ', α') define isomorphic cycle sets, if and only if $\alpha = \alpha'$ and $\Phi' = {}^\beta \Phi$ for some $\beta \in \mathbb{Z}_p^*$.

Assume $p \neq 2$ from now on.

For a pair (Φ, α) with $\Phi(0) \neq 0$, the condition $\Phi(\alpha A) = \alpha\Phi(A)$ forces $\alpha = 1$ and Φ has to be nonconstant and even. As $\Phi(0) \neq 0$, there is exactly one $\beta \in \mathbb{Z}_p^*$ such that ${}^\beta \Phi(0) = 1$. Therefore, each cycle set with parameters (Φ, α) , $\Phi(0) \neq 0$, is isomorphic to a unique cycle set with parameters $(\tilde{\Phi}, 1)$ where $\tilde{\Phi}(0) = 1$, and these parameters define mutually nonisomorphic cycle sets. We therefore find $n_p = p^{\frac{p-1}{2}} - 1$ such cycle sets.

Now assume $\Phi(0) = 0$. The condition $p \neq 2$ excludes $\alpha = -1$, as $\Phi(A) = \Phi(-A)$. More general, $-1 \notin \langle \alpha \rangle \leq \mathbb{Z}_p^*$. Writing $p - 1 = 2^k l$, with $2 \nmid l$ we see that

$$\alpha \in \{x \in \mathbb{Z}_p^* : x^l = 1\} =: \zeta_l \cong \mathbb{Z}_l.$$

As ζ_l is cyclic, there are $\varphi(d)$ elements $\alpha \in \zeta_l$ with multiplicative order $\text{ord}(\alpha) = d$, where φ denotes the Euler φ -function.

In order to count the number of orbits under the action of \mathbb{Z}_p^* , we use Burnside's lemma. Note that only the elements of ζ_l have fixed points in \mathcal{F}_p , so we can restrict to those. Given $\alpha \in \zeta_l$ with $\text{ord}(\alpha) = d$, a function $\Phi \in \mathcal{F}_p$ that satisfies ${}^\alpha \Phi = \Phi$, is already defined by its values on coset representatives of $\mathbb{Z}_p^* / \langle -1, \alpha \rangle$. Under the restriction that Φ is nonconstant and $\Phi(0) = 0$, there are $p^{\frac{p-1}{2d}} - 1$ choices for Φ .

As $\langle \alpha \rangle$ leaves each of these options invariant, we only need to count orbits with respect to the induced action of $\mathbb{Z}_p^*/\langle \alpha \rangle$ on $\text{Fix}_{\mathcal{F}_p}(\alpha)$. Given an element $[\beta] \in \mathbb{Z}_p^*/\langle \alpha \rangle$, it has fixed points in $\text{Fix}_{\mathcal{F}_p}(\alpha)$ if and only if $\beta \in \zeta_l$. Assuming the latter, and letting $c = \text{ord}_{\mathbb{Z}_p^*}(\beta)$, we get $p^{\frac{p-1}{2c}} - 1$ elements in $\text{Fix}_{\mathcal{F}_p}([\beta])$.

Using Burnside's lemma and the fact that there are $\varphi(\frac{c}{d})$ elements $[\beta] \in \zeta_l/\langle \alpha \rangle$ with $\text{ord}_{\mathbb{Z}_p^*}(\beta) = c$, we get that there are

$$\frac{d}{p-1} \sum_{c; d|c|l} \varphi\left(\frac{c}{d}\right) (p^{\frac{p-1}{2c}} - 1)$$

equivalence classes for parameters of the form (Φ, α) where $\text{ord}(\alpha) = d$. Considering that there are $\varphi(d)$ such $\alpha \in \zeta_l$, we get the following number of nonisomorphic cycle sets with parameters (Φ, α) , $\Phi(0) \neq 0$:

$$n'_p = \frac{1}{p-1} \sum_{c, d; d|c|l} d \varphi(d) \varphi\left(\frac{c}{d}\right) (p^{\frac{p-1}{2c}} - 1) = \frac{1}{p-1} \sum_{c, d; d|c|l} d \varphi(d) \varphi\left(\frac{c}{d}\right) (p^{2^{k-1} \frac{l}{c}} - 1).$$

Note that the function

$$\psi(n) = \sum_{d|n} d \varphi(d) \varphi\left(\frac{n}{d}\right)$$

is a convolution of multiplicative functions² and therefore also a multiplicative function which evaluates on prime powers q^ν , $\nu \geq 1$, as

$$\begin{aligned} \psi(q^\nu) &= \sum_{k=0}^{\nu} q^k \varphi(q^k) \varphi(q^{\nu-k}) \\ &= q^\nu (q-1) q^{\nu-1} + (q-1) q^{\nu-1} + \sum_{k=1}^{\nu-1} (q-1)^2 q^{\nu+k-2} \\ &= (q-1) \left(q^{2\nu-1} + q^{\nu-1} + (q-1) q^{\nu-1} \sum_{k=1}^{\nu-1} q^{k-1} \right) \\ &= (q-1) (q^{2\nu-1} + q^{\nu-1} + q^{\nu-1} (q^{\nu-1} - 1)) \\ &= (q-1) (q^{2\nu-1} + q^{2\nu-2}) \\ &= (q^2 - 1) q^{2\nu-2}. \end{aligned}$$

For a number with prime factorization $n = \prod_i q_i^{\nu_i}$, we therefore get

$$\psi(n) = \prod_i (q_i^2 - 1) q_i^{2\nu_i-2}.$$

The total number of indecomposable, nonisomorphic, irretractable cycle sets of size p^2 can therefore be described as:

$$n_p + n'_p = p^{\frac{p-1}{2}} - 1 + \sum_{d|l} \psi\left(\frac{l}{d}\right) \frac{p^{2^{k-1}d} - 1}{p-1}$$

where $p-1 = 2^k l$ with $2 \nmid l$.

²Here, *multiplicative* means $\psi(mn) = \psi(m)\psi(n)$ for *coprime* positive integers m, n .

ACKNOWLEDGEMENTS

This work is partially supported by the project OZR3762 of Vrije Universiteit Brussel and by Fonds Wetenschappelijk Onderzoek - Vlaanderen, via the Senior Research Project G004124N.

The first author expresses his gratitude to the Alexander Humboldt Foundation which funds, by means of a Feodor Lynen fellowship, the research project that encompasses the research conducted for this article.

The second author is supported by Fonds Wetenschappelijk Onderzoek - Vlaanderen, via a PhD Fellowship fundamental research, grant 11PIO24N.

The third author is supported by Fonds Wetenschappelijk Onderzoek - Vlaanderen, via a PhD Fellowship fundamental research, grant 1160522N.

REFERENCES

- [1] O. Akgun, M. Mereb, and L. Vendramin. Enumeration of set-theoretic solutions to the Yang-Baxter equation. *Math. Comp.*, 91:1469–1481, 2020.
- [2] D. Bachiller, F. Cedó, E. Jespers, and J. Okniński. A family of irretractable square-free solutions of the Yang-Baxter equation. *Forum Math.*, 29(6):1291–1306, 2017.
- [3] R. J. Baxter. Partition function of the eight-vertex lattice model. *Ann. Physics*, 70(1):193–228, 1972.
- [4] M. Castelli and S. Trappeniens. Studying solutions of the yang-baxter equation through skew braces, with an application to indecomposable involutive solutions with abelian permutation group, 2023.
- [5] F. Cedó, E. Jespers, and J. Okniński. Braces and the Yang-Baxter equation. *Comm. Math. Phys.*, 327(1):101–116, 2014.
- [6] F. Cedó and J. Okniński. Constructing finite simple solutions of the Yang-Baxter equation. *Adv. Math.*, 391:Paper No. 107968, 40, 2021.
- [7] F. Cedó and J. Okniński. Indecomposable solutions of the Yang-Baxter equation of square-free cardinality. *arXiv.2212.06753*, 2022.
- [8] V. G. Drinfeld. On some unsolved problems in quantum group theory. In P. P. Kulish, editor, *Quantum Groups*, pages 1–8, Berlin, Heidelberg, 1992. Springer Berlin Heidelberg.
- [9] P. Etingof, T. Schedler, and A. Soloviev. Set-theoretical solutions to the quantum Yang-Baxter equation. *Duke Math. J.*, 100(2):169 – 209, 1999.
- [10] E. Feingessicht. Dehornoy’s class and Sylows for set-theoretical solutions of the Yang-Baxter equation. *arXiv.2302.02652*, 2023.
- [11] T. Gateva-Ivanova and M. Van den Bergh. Semigroups of I-type. *J. Algebra*, 206:97–112, 1998.
- [12] L. Guarnieri and L. Vendramin. Skew braces and the Yang-Baxter equation. *Math. Comp.*, 86(307):pp. 2519–2534, 2017.
- [13] B. Huppert. *Endliche Gruppen*. Finite groups / B. Huppert, N. Blackburn. Springer, 1983.
- [14] P. Jedlička and A. Pilitowska. Indecomposable involutive solutions of the Yang-Baxter equation of multipermutation level 2 with non-abelian permutation group. *Journal of Combinatorial Theory, Series A*, 197:105753, 2023.
- [15] V. Lebed, S. Ramírez, and L. Vendramin. Involutive Yang-Baxter: cabling, decomposability, and Dehornoy class. *Revista Matemática Iberoamericana*, pages 1–13, 2023.
- [16] A. Lucchini. On imprimitive groups with small degree. *Rend. Semin. Mat. Univ. Padova*, 86:131–142, 1991.
- [17] W. Rump. A decomposition theorem for square-free unitary solutions of the quantum yang-baxter equation. *Adv. Math.*, 193(1):40–55, 2005.
- [18] W. Rump. Braces, radical rings, and the quantum Yang-Baxter equation. *J. Algebra*, 307(1):153–170, 2007.
- [19] A. Smoktunowicz and L. Vendramin. On skew braces (with an appendix by N. Byott and L. Vendramin). *J. Comb. Algebra*, 2(1):47–86, 2018.
- [20] V. G. Turaev. The Yang-Baxter equation and invariants of links. *Invent. Math.*, 92:527–553, 1988.
- [21] C. N. Yang. Some exact results for the many-body problem in one dimension with repulsive delta-function interaction. *Phys. Rev. Lett.*, 19:1312–1315, Dec 1967.

(Carsten Dietzel) DEPARTMENT OF MATHEMATICS AND DATA SCIENCE, VRIJE UNIVERSITEIT BRUSSEL, PLEINLAAN 2, 1050 BRUSSEL, BELGIUM

Email address: Carsten.Dietzel@vub.be

(Silvia Properzi) DEPARTMENT OF MATHEMATICS AND DATA SCIENCE, VRIJE UNIVERSITEIT
BRUSSEL, PLEINLAAN 2, 1050 BRUSSEL, BELGIUM

Email address: **Silvia.Properzi@vub.be**

(Senne Trappeniers) DEPARTMENT OF MATHEMATICS AND DATA SCIENCE, VRIJE UNIVERSITEIT
BRUSSEL, PLEINLAAN 2, 1050 BRUSSEL, BELGIUM

Email address: **Senne.Trappeniers@vub.be**