Long gamma-ray burst light curves as the result of a common stochastic pulse-avalanche process

L. Bazzanini^{1,2,*}, L. Ferro^{1,2}, C. Guidorzi^{1,2,3}, G. Angora^{1,4}, L. Amati², M. Brescia^{4,5}, M. Bulla^{1,3,6}, F. Frontera^{1,2}, R. Maccary¹, M. Maistrello¹, P. Rosati^{1,2,3}, and A. Tsvetkova^{7,2,8}

Department of Physics and Earth Science, University of Ferrara, via Saragat 1, I-44122, Ferrara, Italy

2 INAF - Osservatorio di Astrofisica e Scienza dello Spazio di Bologna, Via Piero Gobetti 101, I-40129 Bologna, Italy

INFN - Sezione di Ferrara, via Saragat 1, I-44122, Ferrara, Italy

4 INAF - Osservatorio Astronomico di Capodimonte, Salita Moiariello 16, I-80131 Napoli, Italy

Dipartimento di Fisica "E. Pancini", Università di Napoli "Federico II", Via Cinthia 21, I-80126 Napoli, Italy

INAF, Osservatorio Astronomico d'Abruzzo, via Mentore Maggini snc, 64100 Teramo, Italy

Department of Physics, University of Cagliari, SP Monserrato-Sestu, km 0.7, 09042 Monserrato, Italy

Ioffe Institute, Politekhnicheskaya 26, 194021 St. Petersburg, Russia

Received xxx; accepted xxx

ABSTRACT

Context. The complexity and variety exhibited by the light curves of long gamma-ray bursts (GRBs) enclose a wealth of information that still awaits being fully deciphered. Despite the tremendous advance in the knowledge of the energetics, structure, and composition of the relativistic jet that results from the core collapse of the progenitor star, the nature of the inner engine, how it powers the relativistic outflow, and the dissipation mechanisms remain open issues.

Aims. A promising way to gain insights is describing GRB light curves as the result of a common stochastic process. In the Burst And Transient Source Experiment (BATSE) era, a stochastic pulse avalanche model was proposed and tested through the comparison of ensemble-average properties of simulated and real light curves. Here we aim to revive and further test this model.

Methods. We apply it to two independent data sets, BATSE and Swift/BAT, through a machine learning approach: the model parameters are optimised using a genetic algorithm.

Results. The average properties are successfully reproduced. Notwithstanding the different populations and passbands of both data sets, the corresponding optimal parameters are interestingly similar. In particular, for both sets the dynamics appears to be close to a critical state, which is key to reproduce the observed variety of time profiles.

Conclusions. Our results propel the avalanche character in a critical regime as a key trait of the energy release in GRB engines, which underpins some kind of instability.

Key words. Gamma-ray burst: general – Methods: statistical – machine learning – genetic algorithms

Gamma-ray bursts (GRBs) are the most powerful explosions on stellar scale in the Universe. At least two kinds of progenitors are known: (i) so-called 'collapsar' (Woosley 1993; Paczyński 1998; MacFadyen & Woosley 1999), that is a hydrogen-stripped massive star, whose core collapses to a compact object, which launches a relativistic ($\Gamma \sim 10^2 - 10^3$) jet; (ii) merger of a compact binary (Eichler et al. 1989; Paczynski 1991; Narayan et al. 1992), where at least one of the two objects is supposed to be a neutron star (NS) and which also results in a short-lived relativistic jet (see Kumar & Zhang 2015; Zhang 2018 for recent reviews). There are alternative models to the collapsar, such as the binary-driven hypernova model (BdHN; Rueda & Ruffini 2012; Becerra et al. 2019), in which the final collapse of a CO core of a massive star can trigger the collapse of a companion neutron star. Most GRBs due to (i) manifest themselves as long GRBs (LGRBs), lasting longer than $\sim 2 s^1$, while (ii) usually exhibit a subsecond spike

occasionally followed by weak, long-lasting emission (Norris & Bonnell 2006), and are commonly referred to as short GRBs (SGRBs). Actually, the emerging picture is more complicated, as shown by the increasing number of cases with deceptive time profiles found in both classes (Gehrels et al. 2006; Rastinejad et al. 2022; Gompertz et al. 2023; Yang et al. 2022; Troja et al. 2022; Ahumada et al. 2021; Zhang et al. 2021; Rossi et al. 2022; Levan et al. 2023; Levan et al. 2024).

The nature of the dissipation mechanism that is responsible for the GRB prompt emission is still an open issue. The great variety observed in the light curves (LCs) of LGRBs is thought to be the result of the variability imprinted to the relativistic outflow by the inner engine left over by the collapsar, either a millisecond magnetised NS or a black hole (BH), along with the effects of the propagation of the jet within the stellar envelope (e.g., Morsony et al. 2010; Geng et al. 2016; Gottlieb et al. 2020b,a, 2021b,a), although some models ascribe the possible presence of subsecond variability to magnetic reconnection events taking place at larger radii (Zhang & Yan 2011). Some correlations were found between variability and minimum variability timescale on one side, as defined in a number of ways, and luminosity

^{*} bzzlnz[at]unife[dot]it

¹ This boundary value is from CGRO/BATSE GRB catalogue and slightly depends on the detector's passband.

and initial Lorentz factor of the outflow on the other side (e.g., see Camisasca et al. 2023 and references therein). However, apart from the study of average and of individual Fourier power density spectra of GRBs (Beloborodov et al. 1998; Guidorzi et al. 2012, 2016; Dichiara et al. 2013), the study of the waiting time distribution between pulses (Ramirez-Ruiz et al. 2001; Nakar & Piran 2002; Quilligan et al. 2002; Guidorzi et al. 2015), and the distribution of the number of peaks per GRB (Guidorzi et al. 2024), little progress has been made in deciphering and characterising the variety of LGRB LCs within a unifying scheme that could explain the large diversity (in terms of duration, number of pulses, distribution of energy and waiting times between pulses) and relate it to other key properties. Recent investigations found possible evidence that GRB engines emit as self-organised critical (SOC) systems (Wang & Dai 2013; Yi et al. 2017; Lyu et al. 2020; Wei 2023; Li & Yang 2023; Maccary et al. 2024), in which energy is released through avalanches whenever the system naturally reaches a critical point. Yet, the interpretation is not straightforward, since SOC dynamics is usually invoked for systems that are continuously fed by some energy input and are not characterised by the kind of irreversible evolution expected for a GRB inner engine. A successful description of the inner engine variability would help constrain the mechanism that powers the jetted outflow in GRBs and, ultimately, the nature of the compact object. Furthermore, it would provide the community with a reliable tool to simulate credible GRB LCs as they would be measured by future experiments, avoiding the pitfalls of using real noisy LCs (e.g., Sanna et al. 2020).

In this respect, an interesting attempt was laid out by Stern & Svensson (1996, hereafter SS96) in the Compton Gamma–Ray Observatory era (CGRO; 1991-2000) on the GRB catalogue of one of its experiments, the Burst And Transient Source Experiment (BATSE). These authors proposed a common stochastic process built on a pulse avalanche mechanism and tried to reproduce some of the observed distributions of BATSE GRB LCs. At that time, the cosmological distances of GRBs and the progenitors' nature of the two classes were yet to be firmly established, with the first afterglow discoveries starting from 1997 (Costa et al. 1997). By manually guessing the values of the seven model parameters, SS96 came up with a process operating in a nearly critical regime and capable of reproducing the variety of observed GRB LCs, as long as the chosen metrics are concerned. This approach of simulating GRB LCs was adopted in Greiner et al. (2022) to assess localisation capabilities of a proposed network of GRB detectors on the global navigation satellite system Galileo G2.

In the big data era, advanced statistical and machine learning (ML) techniques applied to astrophysics have become routine (e.g., see Feigelson et al. 2021 for a review). In this paper we aim to verify and improve the results obtained by SS96 on the BATSE data and, for the first time, apply their model to a sample from another detector operating in a softer energy band, such as the Burst Alert Telescope (BAT; Barthelmy et al. 2005) aboard the Neil Gehrels Swift Observatory (Gehrels et al. 2004). Specifically, we aim at optimising the model parameters using a genetic algorithm (GA; Rojas 1996).

A similar technique, in which the parameters of a physical model were optimised through the application of a GA, was recently applied by Vargas et al. (2022) to model the shock propagation in the supernova SN2014C progenitor star and ejecta, in which the GA was used to optimise a hydrodynamic and radiation transfer model.

Unlike SS96, we restrict our analysis to LGRBs, whose progenitor is thought to be a collapsar, to preserve as much as possible the homogeneity of the putative GRB inner engines. For $\sim 30\%$ of the *Swift* sample with measured redshift, in principle it is possible to carry out the same analysis in the GRB rest frame. However, we did not consider this option, since the cosmological dilation correction by (1 + z) is partly counteracted by other energy-dependent effects, which make the final correction milder and less obvious (see Camisasca et al. 2023 and references therein for a detailed explanation).

In this paper, we report the main results and implications. A companion and more ML-oriented paper will report all the technical details. The present work is organised as follows: in Section 2 we describe the data analysis and sample selection, while in Section 3 we illustrate the methods underpinning the avalanche model and the implementation of the genetic algorithm. Section 4 reports the results, whose discussion and conclusions are laid out in Section 5.

2. Data Analysis

2.1. Sample selection

From the BATSE 4B catalogue (Paciesas et al. 1999) we took the 64-ms time profiles that were made available by the BATSE team². Observed with the BATSE eight Large Area Detectors (LADs), these data are the result of a concatenation of three standard BATSE types, DISCLA, PREB, and DISCSC, available in four energy channels: 25–55, 55–110, 110–320, and > 320 keV. We used the total passband LCs. For each GRB the background was interpolated with polynomials of up to fourth degree as prescribed by the BATSE team. In our analysis, we used the backgroundsubtracted LCs.

From an initial sample of 2024 GRBs we selected only those that satisfy the following requirements:

- $T_{90} > 2$ s, that is, only *long* GRBs;
- data available for at least 150 s after the brightest peak;
 signal-to-noise ratio (S/N) of the total net counts within the duration of the event greater than 70.

In order to estimate the S/N, following SS96, rather than the commonly used T_{90} , we used as a proxy of the GRB duration the time interval from the first to the last time bin whose counts exceed the threshold of 20% of the peak counts, henceforth called $T_{20\%}$. Before evaluating the $T_{20\%}$, the LCs were first convolved with a Savitzky-Golay smoothing filter (Savitzky & Golay 1964), using a second order interpolating polynomial, and a moving window of size $T_{90}/15$. Accordingly, we defined the S/N of a GRB as the sum of the net counts in the whole $T_{20\%}$ interval, divided by the corresponding error. The value of the S/N threshold was the result of a trade-off between the number of GRBs and the statistical quality of the LCs in the sample. Furthermore, the $T_{20\%}$ is also used to compute the duration distribution of the LCs (Section 2.2).

We ended up with 585 long GRBs satisfying the aforementioned properties. Hereafter, this will be referred to as the BATSE sample.

As a second dataset, we considered the GRBs detected by Swift/BAT from January 2005 to November 2023 and covered in burst mode. We used the total 15-150 keV passband LCs, with 64-ms bin time; these were extracted as mask-weighted background-subtracted LCs, following the standard procedure recommended by the BAT team.³ From an initial sample of 1389

https://heasarc.gsfc.nasa.gov/FTP/compton/data/ batse/ascii_data/64ms/

³ https://swift.gsfc.nasa.gov/analysis/threads/bat_ threads.html.

GRBs observed in burst mode, 531 passed the selection based on the same criteria adopted for BATSE, except for the value of the S/N threshold, which was lowered to 15 to obtain a sample of comparable size to the BATSE one, but still ensuring the required statistical quality. Hereafter, this will be referred to as the *Swift* sample.

2.2. Statistical metrics

We considered the following four metrics, which were also used by **SS96**:

- the average peak-aligned post-peak time profile (Mitrofanov 1996), in the time range 0–150 s *after* the brightest peak. It is evaluated by averaging the normalised count rate of all the LCs in the sample, i.e. ⟨F/F_p⟩, F_p being the peak count rate. Further details are given in Stern (1996);
- the average peak-aligned third moment of post-peak time profiles ⟨(F/F_p)³⟩, evaluated analogously to the first moment;
 the average auto-correlation function (ACF). For both data
- the average auto-correlation function (ACF). For both data samples the ACF is corrected for the counting statistics noise as in Link et al. (1993) and is computed in the 0–150 s interval;
- 4. the $T_{20\%}$ distribution, with $T_{20\%}$ used as a proxy of the duration.

As in SS96, (1)-(4) are used as metrics to evaluate the degree of similarity between the real and the simulated LCs.

3. Methods

3.1. Light curve simulations

The stochastic process conceived by SS96 belongs to the class of so-called 'branching' processes, which describe the development of a population whose members reproduce according to some random process (Harris 1963). We outline its key features below, and refer the reader to SS96 for more details.

The **SS96** model is based on the assumptions that (i) GRB LCs can be viewed as distinct random realisations of a common stochastic process, within narrow parameter ranges; (ii) the stochastic process should be scale invariant in time, and (iii) it operates close to a critical state. With this model, each LC consists of a series of spontaneous primary (or *parent*) pulses, each of which can give rise to secondary (or *child*) pulses, which can then further generate pulses until the process reaches subcritical conditions and stops. Each pulse, which acts as a building block, is described by a Gaussian rise followed by a simple exponential decay:

$$f(t) = \begin{cases} A \exp\left\{-(t - t_p)^2 / \tau_r^2\right\}, & \text{for } t < t_p \\ A \exp\left\{-(t - t_p) / \tau\right\}, & \text{for } t > t_p \end{cases}$$
(1)

where τ is roughly the pulse width, t_p is the peak time, A is the amplitude, and we assume $\tau_r = \tau/2$ (Norris et al. 1996). Differently from SS96, we do not sample A from a uniform distribution $\mathcal{U}[0, 1]$, rather, for each GRB we sample the value A_{max} from the distribution of the peak count rates of the real observed LCs, and then the amplitude of each pulse composing that LC is sampled from $\mathcal{U}[0, A_{\text{max}}]$.

The model is described by seven parameters:

- μ_0 rules the number μ_s of spontaneous initial pulses per GRB, which is sampled from a Poisson distribution with μ_0 as expected value:

$$p(\mu_s|\mu_0) = \frac{\mu_0^{\mu_s} \exp(-\mu_0)}{\mu_s!} \quad .$$
 (2)

- μ rules the number of child pulses μ_c generated by each parent pulse, which is sampled from a Poisson distribution with μ as expected value:

$$p(\mu_c|\mu) = \frac{\mu^{\mu_c} \exp(-\mu)}{\mu_c!} \quad .$$
 (3)

- α rules the delay Δt between a child and its parent. This delay is exponentially distributed, with e-folding time given by $(\alpha \tau)$, where τ is the time constant of the child pulse:

$$p(\Delta t) = (\alpha \tau)^{-1} \exp(-\Delta t / \alpha \tau) \quad . \tag{4}$$

Moreover, the spontaneous μ_s primary pulses are all assumed to be delayed with respect to a common invisible trigger event; the probability distribution of such delay *t* is exponentially distributed:

$$p(t) = (\alpha \tau_0)^{-1} \exp(-t/\alpha \tau_0) \quad , \tag{5}$$

 τ_0 being the time constant of the primary pulse.

- τ_{\min} and τ_{\max} define the boundaries for the constant τ_0 of the primary spontaneous pulses and whose probability density function is $p(\tau_0) \propto 1/\tau_0$, equivalent to a uniform distribution of log τ_0 :

$$p(\log \tau_0) = \left[\log \tau_{\max} - \log \tau_{\min}\right]^{-1} , \qquad (6)$$

where τ_{\min} has to be shorter than the time resolution of the instrument. Varying τ_{\max} is equivalent to rescaling all average avalanche properties in time.

- δ_1 and δ_2 define the boundaries, $[\delta_1, \delta_2]$, of a uniform distribution assumed for the logarithm of the ratio between τ of the child and τ_p of its parent:

$$p[\log(\tau/\tau_p)] = |\delta_2 - \delta_1|^{-1} \quad , \tag{7}$$

with
$$\delta_1 < 0$$
, $\delta_2 \ge 0$, and $|\delta_1| > |\delta_2|$.

Each of the μ_s spontaneous initial pulses gives rise to a pulse avalanche, acting as a parent, spawning another set of child pulses, in a recurrent way. Finally, it is the superposition of all the parent and child pulses generated during the avalanche that shapes the LC of an individual GRB.

The stochastic pulse avalanche model was used to simulate both BATSE and *Swift*/BAT LCs. The statistical noise depends on the total counts in each time bin, which requires the knowledge of the typical background count rate for a given instrument. For BATSE, which consisted of NaI(Tl) scintillators, we assumed a constant background rate of 2.9 cnt s⁻¹cm⁻², which corresponds to the median of the distribution of the measured error rates. Each final simulated LC was the result of a Poisson realisation, assuming for each time bin the total counts (that is, noise-free simulated profile plus background) as expected value. Lastly, the background was removed.

Swift/BAT is a coded mask coupled with a CZT detection array. Its background-subtracted LCs are the result of the deconvolution of the detection with the pattern of the mask, so the rate in each time bin can be modelled as a Gaussian variable. To simulate BAT LCs, the rate of each time bin was sampled from a Gaussian distribution centred on the LC (noise-free) model obtained with the pulse avalanche model, and with standard deviation randomly sampled from the errors measured in the real *Swift*/BAT LCs.

All the simulations were carried out using an open-source Python⁴ implementation⁵ by one of the authors.

⁴ https://www.python.org/

⁵ https://github.com/anastasia-tsvetkova/lc_pulse_ avalanche

3.2. Genetic algorithm

SS96 proposed a set of values for the seven model parameters as the result of an educated guess. The optimisation of these parameters, however, is an ideal task for nowadays routinely used ML techniques.

GAs are a specific type of algorithms in the larger family of the so-called evolutionary algorithms (Russell & Norvig 2021; Rojas 1996; Aggarwal 2021; Hurbans 2020), where a Darwinian evolution process is simulated to find the parameters that maximise a function.

In GAs, each solution to an optimisation problem can be seen as an *individual*, with the "fitness" of that individual being determined by the objective function value of the corresponding solution. These solutions are points in the domain of the function to be optimised. In our work, each individual is represented by a genome made of seven genes, which are the parameters of the SS96 model described in Section 3.1.

At each generation, a new set of individuals is created. Over time, the points belonging to the new generations gradually converge towards local maxima of the fitness function. In order to improve over successive generations the overall fitness of the population, GAs incorporate three fundamental processes: selection, crossover, and mutation.

The typical life cycle of a GA, made up of a succession of the so-called *generations*, includes the following steps:

- 1. Population initialisation: Generating randomly a population of potential solutions;
- 2. Evaluating fitness: Assessing the quality of each individual by employing a fitness function that assigns scores to evaluate their fitness;
- 3. Parent selection: Choosing pairs of parents for reproduction based on their fitness score;
- 4. Offspring creation: Producing offspring by combining genetic information from parents, and introducing random mutations;
- 5. Generation advancement: Selecting individuals and offspring from the population to progress to the next generation.

GAs are particularly useful in situations where there is no available information about the function's gradient at the evaluated points. Indeed, GA can effectively handle functions that are not continuous or differentiable (Rojas 1996).

3.3. Parameter optimisation

The GA has been implemented using PyGAD⁶, an open-source Python library containing a collection of several ML algorithms (Gad 2023).

We constrain the seven parameters of the model within the intervals shown in Table 1.

The GA evolves through a sequence of generations consisting of a population with $N_{pop} = 2000$ individual sets of seven parameters. Each set of parameters, hereafter referred to as an individual, is then used to generate $N_{grb} = 2000$ LCs. The very same three constraints, mentioned in Section 2.1, and used for the selection of BATSE and *Swift*/BAT dataset, are applied also on the simulated GRB LCs, the generated ones not satisfying such constraints being discarded. For each of the N_{pop} individuals, we evaluate the same four metrics defined in SS96 over the corresponding N_{grb} LCs (cfr. Section 2.2), and compare them with the values obtained from the real datasets, by computing the *L*2 loss between these four observables. The final loss associated with a

Table 1: Region of exploration during the GA optimisation of the seven parameters of the **SS96** stochastic model.

Parameter	Lower bound	Upper bound
μ	0.80	1.7
μ_0	0.80	1.7
α	1	15
δ_1	-1.5	-0.30
δ_2	0	0.30
$ au_{ m min}$	0.01 s	<pre>bin_time s</pre>
$ au_{ m max}$	1 s	60 s

given individual is simply defined as the average of these four quantities, the fitness score being the inverse of this value.

Individuals are then ranked based on their loss. The next generation of individuals is obtained by mixing the genes (i.e. the values of the seven parameters) of the fittest individuals in the current generation. No individuals are instead automatically kept in the next generation, that is, we set to zero the so-called "elitism"⁷. The offspring is obtained by randomly sampling two individuals among the top 15% in the current generation and assigning to each gene the value of the seven parameters from one of the two parents, with equal probability.

Finally, we include the possibility for genetic random mutations to occur. During the mating step, each one of the seven parameters has a 4% probability of undergoing mutation, meaning that the value of the parameters is not inherited from one of the two parents, but instead, it is randomly sampled from the exploration range of the parameter (Table 1).

The optimisation process is stopped when convergence of the loss, and thus of the value of the seven parameters, is reached.

4. Results

In Table 2 we compare the values of the seven model parameters suggested in SS96 with the results of our GA optimisation on the BATSE and *Swift* training datasets. The final optimised values of the seven parameters are obtained as the median value in the whole population of the last GA generation. We also list the achieved values of the loss function evaluated on the training set (both in terms of best parameter configuration and by averaging on the last population) as well as on the test set (i.e. estimated by using 5000 newly simulated GRB LCs). In the bottom, we resolve the individual contribution of each component to the test loss.

Figure 1 displays the comparison of the four observables, described in Section 2.2, between the real BATSE curves and simulated ones (test set). In particular, the panels show the average profiles obtained from the 585 useful BATSE events (blue), the ones estimated from 5000 simulated GRBs with optimised parameters (red), and the ones estimated from 5000 LCs simulated using the parameter values guessed by SS96 (green). Figure 2 shows the analogous comparison between the simulated and the real *Swift*/BAT curves.

We find an excellent agreement for three out of the four metrics computed from real and simulated BATSE LCs, in particular for the average post-peak time profile, its third moment, and the

⁶ https://github.com/ahmedfgad/GeneticAlgorithmPython

 $^{^{7}}$ Due to the stochastic nature of the SS96 algorithm, the same set of parameters will never produce a set of LCs with the same loss; therefore keeping a set of individuals in the next generation is not helping, since in reality, given seven fixed parameters, there are fluctuations in the value of the corresponding loss.



Fig. 1: Average distributions of real (blue), simulated GA-optimised (red), and simulated SS96 (green) BATSE GRB profiles, estimated on the test set (see Table 2). *Top left*: average peak-aligned post-peak normalised time profile, together with the r.m.s. deviation of the individual peak-aligned time profiles, $F_{\rm rms} \equiv [\langle (F/F_p)^2 \rangle - \langle F/F_p \rangle^2]^{1/2}$. *Top right*: average peak-aligned third moment test. *Bottom left*: Average ACF of the GRBs. *Bottom right*: distribution of duration, measured at a level of 20% of the peak amplitude ($T_{20\%}$). In *top left* and *top right* panels, both real and simulated averaged curves were smoothed with a Savitzky-Golay filter to reduce the effect of Poisson noise. In *bottom right* panel, a Gaussian kernel convolution has been applied to both real and simulated distributions.

average auto-correlation, whose L2 loss values are smaller than the corresponding ones estimated with SS96 non-optimised parameters, as can be seen from the bottom part of Table 2. For instance, the average ACF metric shows a relative improvement of ~ 74% after the optimisation. The $T_{20\%}$ distribution holds the largest contribution to the loss; yet, it slightly improves the SS96 performance (~ 8% relative improvement). Overall, compared with SS96, our GA-optimised results on BATSE data better reproduce the observed distributions.

As can be inferred from Table 2, and graphically from Figure 2, according to the loss function the avalanche model appears to work even better in the case of *Swift* data: the results on the average ACF and third moment of peak-aligned profiles are comparably good, whereas the average peak-aligned profile and duration distributions are significantly improved with respect to the BATSE case, with a relative loss decrease of $\sim 43\%$ and $\sim 58\%$, respectively.

Notably, the two sets of best-fitting parameters obtained with BATSE and with *Swift*/BAT are very similar and, surprisingly,

overall not too different from the one guessed by SS96. In Section 5 we discuss the relevance of this result in more detail.

The parameters for which our optimal values for both sets are somewhat different from those of SS96 are (δ_1, δ_2) , τ_{max} , and α . The former pair defines the dynamic range of the child-toparent pulse duration ratio: our values turn into broader dynamical ranges than SS96, and, at variance with those authors, they admit the possibility of children lasting longer than parents, being $\delta_2 >$ 0. While SS96 assumed $\tau_{max} = 26$ s as the maximum value for the duration of parent pulses, our optimised solution favours the possibility of longer parent pulses: 40.2 s for BATSE and 56.8 s for *Swift*/BAT. Finally, our best-fit values for α , which rules the time delay between parent and child, lean towards slightly shorter intervals than SS96.

As in SS96, Figure 3 presents the comparison of four real BATSE time profiles with four simulated LCs of similar morphology and complexity, sampled from the test set, generated using the best-fitting set of BATSE model parameters given above. This qualitative plot shows the ability of the SS96 stochastic model to



Fig. 2: Comparison between the real *Swift*/BAT dataset and the corresponding simulated dataset on the same four metrics defined for the BATSE dataset, analogously to Figure 1.

reproduce the different morphological classes of LCs discussed in the literature from the earliest observations (Fishman & Meegan 1995).

5. Discussion and conclusions

For the first time in the GRB literature, here we developed and implemented an ML technique to optimise the parameters of a stochastic model capable of generating ex-novo realistic GRB LCs. Our work confirmed the soundness of the insight by SS96: a simple toy model like the stochastic pulse avalanche one is able to generate populations of LCs, whose average behaviour closely resembles that of the real populations of BATSE and *Swift*/BAT long GRBs. With the implementation of GA, we found the two best fit sets of the seven parameters of the model that best reproduce the average behaviours of the two datasets, thus (i) progressing from the educated-guessed values of the original paper to a real fit of the model on BATSE data, and (ii) applying it for the first time to an independent catalogue of GRB LCs like *Swift*/BAT, whose data differ from the BATSE one in many aspects, as detailed below.

In light of our GA-optimised results on BATSE data, the educated guess by SS96 turns out to be surprisingly good. In particular, the finding that μ , that is the average number of child-pulses generated by each parent-pulse, must be close to unity (our

 $1.10^{+0.03}_{-0.02}$ vs. 1.20 of SS96) confirms the insightful Ansatz by SS96 that the model must operate very close to a critical regime ($\mu = 1$), to account for the observed variety of GRB profiles. Interestingly, the same clue is also obtained in the GA-optimised parameters of the Swift/BAT sample. This result is far from obvious for three main reasons: (i) the passband of the two experiments is significantly different, with Swift/BAT profiles being softer and, as such, less spiky (e.g., Fenimore et al. 1995); (ii) the average S/N of the two sets is also remarkably different, with a minimum value of 70 for BATSE, to be compared with the poorer lower threshold of 15 for Swift/BAT (see Section 2); (iii) the GRB populations seen by the two experiments are likely different: thanks to its larger effective area at low energies, longer trigger accumulation times and much more complex trigger algorithms, Swift/BAT detects more high-redshift GRBs (Band 2006; Lien et al. 2014; Wanderman & Piran 2010). Therefore, our results provide additional evidence for a nearly critical regime in which GRB engines would work, in agreement with other independent investigations (Maccary et al. 2024; Guidorzi et al. 2024).

In addition to providing new clues on the dynamical behaviour of LGRB inner engines or, more generally, on the way some kind of energy is dissipated into gamma-rays, this model offers the practical possibility of simulating realistic GRB profiles with future experiments, such as HERMES (Fiore et al. 2020) and



Fig. 3: Four examples of as many classes of GRB LCs from the BATSE real sample (*left*) along with their trigger number, and the corresponding simulated one (*right*). Following the same qualitative classification adopted by SS96, from top to bottom the four classes are "single pulse", "blending of some pulses", "moderately structured", and "highly erratic". On the top right of each subplot is shown the average error on the counts of the corresponding LC.

possibly the X/Gamma-ray Imaging Spectrometer (XGIS; Amati et al. 2022) aboard ESA/M7 candidate THESEUS (Amati et al. 2021) currently selected for a phase A study. The task of simulating realistic GRB time profiles, as they will be seen by forthcoming detectors, is far from obvious: the alternative option of renormalising real LCs observed with different instruments is inevitably hampered by the presence of counting statistics (Poisson) noise, which cannot be merely rescaled without altering its nature. A filtering procedure would be then required, which in turn assumes that the uncorrelated Poisson noise can be disentangled from the genuine (unknown) variance of GRBs, which also requires substantial effort.

Summing up, the present work showcases the potential of a simple toy model like the avalanche one conceived by SS96, once it is properly bolstered with ML techniques. Moreover, it paves the way to further optimisation of the model in different directions: (i) by adding further metrics, such as the distributions of the following observables: GRB S/N, duration of observed pulses, or the number of peaks per GRB (Guidorzi et al. 2024); (ii) by studying in more detail the dependence of the model parameters on the energy channels; (iii) by carrying out the same study in the comoving frame of a sample of GRBs with known redshift, assuming the luminosity and released energy distributions of individual pulses (Maccary et al. 2024). Eventually, these efforts should end up with a reliable and accessible machine for simulating credible LGRB profiles with any experiment. In parallel, the final outcome would be a detailed characterisation of the dynamics that rules long GRB prompt emission, possibly disclosing the nature of long GRB engines.

The source code of our algorithm, alongside all the scripts used to perform the data analysis and produce the plots, will be publicly released on GitHub⁸ upon publication.

Acknowledgements. LB is indebted to the communities behind the multiple free, libre, and open-source software packages on which we all depend. AT acknowledges financial support from ASI-INAF Accordo Attuativo HERMES Pathfinder operazioni n. 2022-25-HH.0 and the basic funding program of the Ioffe Institute FFUG-2024-0002.

CG conceived the research based on the original code developed by AT according to the SS96, LB, GA, and CG developed the methodology, LB, LF and CG performed all the numerical work, including software development, investigation, and validation. All authors contributed to the discussion of the results, the editing and revision of the paper.

This work uses the following software packages: Python (Van Rossum & Drake 2009), PyGAD (Gad 2023), NumPy (van der Walt et al. 2011; Harris et al. 2020), SciPy (Virtanen et al. 2020) matplotlib (Hunter 2007), seaborn (Waskom 2021), MEPSA (Guidorzi 2015), gnuplot (Williams et al. 2023), bash (GNU 2007).

References

Aggarwal, C. C. 2021, Artificial Intelligence: A Textbook (Springer)

- Ahumada, T., Singer, L. P., Anand, S., et al. 2021, Nature Astronomy, 5, 917
- Amati, L., Labanti, C., Mereghetti, S., et al. 2022, in Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series, Vol. 12181, Space Telescopes and Instrumentation 2022: Ultraviolet to Gamma Ray, ed. J.-W. A. den Herder, S. Nikzad, & K. Nakazawa, 1218126

Amati, L., O'Brien, P. T., Götz, D., et al. 2021, Experimental Astronomy, 52, 183 Band, D. L. 2006, ApJ, 644, 378

Barthelmy, S. D., Barbier, L. M., Cummings, J. R., et al. 2005, Space Sci. Rev., 120, 143

⁸ https://github.com/LBasz/geneticgrbs

Parameter	SS96	BATSE	Swift/BAT
μ	1.20	$1.10^{+0.03}_{-0.02}$	$1.34^{+0.03}_{-0.02}$
μ_0	1.00	$0.91\substack{+0.06 \\ -0.07}$	$1.16^{+0.18}_{-0.10}$
α	4.00	$2.57^{+0.07}_{-0.52}$	$2.53^{+0.25}_{-0.01}$
δ_1	-0.50	$-1.28^{+0.16}_{-0.05}$	$-0.75^{+0.11}_{-0.29}$
δ_2	0	$0.28^{+0.01}_{-0.03}$	$0.27^{+0.01}_{-0.02}$
$ au_{ m min}$	0.02 s	$0.02^{+0.02}_{-0.01}\mathrm{s}$	$0.03^{+0.02}_{-0.02} \mathrm{s}$
$ au_{ m max}$	26.0 s	$40.2^{+0.9}_{-1.2}\mathrm{s}$	$56.8^{+0.4}_{-1.3}$ s
Loss (Train best)	-	0.72	0.38
Loss (Train avg.)	-	0.98	0.66
Loss (Test)	1.47	0.88	0.56
Loss (<i>Test</i> : $\langle F/F_p \rangle$)	1.01	0.67	0.46
Loss (<i>Test</i> : $\langle (F/F_p)^3 \rangle$)	0.40	0.20	0.20
Loss (Test: (ACF))	2.24	0.64	0.49
Loss (<i>Test</i> : <i>T</i> _{20%})	2.22	2.04	1.08

Table 2: Results of the GA optimisation on the BATSE and Swift datasets.

Notes. Col. 2 presents the parameters given by SS96 (for the BATSE dataset), while Col. 3 and Col. 4 show the optimised ones obtained after 30 generations of the GA for BATSE and Swift/BAT, respectively. From the distribution of the seven parameters in the last generation we estimated their best-fitting values as the median, and their corresponding errors as the 16-th and 84-th percentiles. "Train best" is the loss of the best generation, while "Train avg." is the average loss in the last generation. The test set is a newly produced set of 5000 simulated LCs; the last four rows show all the single contributions to the "Test" loss.

- Becerra, L., Ellinger, C. L., Fryer, C. L., Rueda, J. A., & Ruffini, R. 2019, ApJ, 871, 14
- Beloborodov, A. M., Stern, B. E., & Svensson, R. 1998, ApJ, 508, L25
- Camisasca, A. E., Guidorzi, C., Amati, L., et al. 2023, A&A, 671, A112
- Costa, E., Frontera, F., Heise, J., et al. 1997, Nature, 387, 783
- Dichiara, S., Guidorzi, C., Amati, L., & Frontera, F. 2013, MNRAS, 431, 3608
- Eichler, D., Livio, M., Piran, T., & Schramm, D. N. 1989, Nature, 340, 126
- Feigelson, E. D., de Souza, R. S., Ishida, E. E. O., & Jogesh Babu, G. 2021, Annual Review of Statistics and Its Application, 8, 493
- Fenimore, E. E., in 't Zand, J. J. M., Norris, J. P., Bonnell, J. T., & Nemiroff, R. J. 1995, ApJ, 448, L101
- Fiore, F., Burderi, L., Lavagna, M., et al. 2020, in Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series, Vol. 11444, Space Telescopes and Instrumentation 2020: Ultraviolet to Gamma Ray, ed. J.-W. A. den Herder, S. Nikzad, & K. Nakazawa, 114441R
- Fishman, G. J. & Meegan, C. A. 1995, ARA&A, 33, 415
- Gad, A. F. 2023, Multimedia Tools and Applications, 1
- Gehrels, N., Chincarini, G., Giommi, P., et al. 2004, ApJ, 611, 1005
- Gehrels, N., Norris, J. P., Barthelmy, S. D., et al. 2006, Nature, 444, 1044
- Geng, J.-J., Zhang, B., & Kuiper, R. 2016, ApJ, 833, 116
- GNU, P. 2007, Free Software Foundation. Bash (3.2. 48)[Unix shell program] Gompertz, B. P., Ravasio, M. E., Nicholl, M., et al. 2023, Nature Astronomy, 7,
- 67
- Gottlieb, O., Bromberg, O., Levinson, A., & Nakar, E. 2021a, MNRAS, 504, 3947
- Gottlieb, O., Bromberg, O., Singh, C. B., & Nakar, E. 2020a, MNRAS, 498, 3320
- Gottlieb, O., Levinson, A., & Nakar, E. 2020b, MNRAS, 495, 570
- Gottlieb, O., Nakar, E., & Bromberg, O. 2021b, MNRAS, 500, 3511
- Greiner, J., Hugentobler, U., Burgess, J. M., et al. 2022, A&A, 664, A131
- Guidorzi, C. 2015, Astronomy and computing, 10, 54 Guidorzi, C., Dichiara, S., & Amati, L. 2016, A&A, 589, A98
- Guidorzi, C., Dichiara, S., Frontera, F., et al. 2015, ApJ, 801, 57
- Guidorzi, C., Margutti, R., Amati, L., et al. 2012, MNRAS, 422, 1785
- Guidorzi, C., Sartori, M., Maccary, R., et al. 2024, A&A, in press, arXiv:2402.17282
- Harris, C. R., Millman, K. J., van der Walt, S. J., et al. 2020, Nature, 585, 357-362

Article number, page 8 of 8

- Harris, T. E. 1963, The theory of branching processes, Die Grundlehren der Mathematischen Wissenschaften, Bd. 119 (Berlin: Springer-Verlag), xiv+230 Hunter, J. D. 2007, Computing in Science & Engineering, 9, 90
- Hurbans, R. 2020, Grokking Artificial Intelligence Algorithms (Manning)
- Kumar, P. & Zhang, B. 2015, Phys. Rep., 561, 1
- Levan, A. J., Gompertz, B. P., Salafia, O. S., et al. 2024, Nature, 626, 737
- Levan, A. J., Malesani, D. B., Gompertz, B. P., et al. 2023, Nature Astronomy, 7, 976
- Li, X.-J. & Yang, Y.-P. 2023, ApJ, 955, L34
- Lien, A., Sakamoto, T., Gehrels, N., et al. 2014, ApJ, 783, 24
- Link, B., Epstein, R. I., & Priedhorsky, W. C. 1993, ApJ, 408, L81
- Lyu, F., Li, Y.-P., Hou, S.-J., et al. 2020, Frontiers of Physics, 16, 14501
- Maccary, R., Guidorzi, C., Amati, L., et al. 2024, ApJ, in press, arXiv:2401.14063 MacFadyen, A. I. & Woosley, S. E. 1999, ApJ, 524, 262
- Mitrofanov, I. G. 1996, Mem. Soc. Astron. Italiana, 67, 417
- Morsony, B. J., Lazzati, D., & Begelman, M. C. 2010, ApJ, 723, 267
- Nakar, E. & Piran, T. 2002, MNRAS, 331, 40
- Narayan, R., Paczynski, B., & Piran, T. 1992, ApJ, 395, L83
- Norris, J. P. & Bonnell, J. T. 2006, ApJ, 643, 266
- Norris, J. P., Nemiroff, R. J., Bonnell, J. T., et al. 1996, ApJ, 459, 393
- Paciesas, W. S., Meegan, C. A., Pendleton, G. N., et al. 1999, ApJS, 122, 465
- Paczynski, B. 1991, Acta Astron., 41, 257
- Paczyński, B. 1998, ApJ, 494, L45
- Quilligan, F., McBreen, B., Hanlon, L., et al. 2002, A&A, 385, 377
- Ramirez-Ruiz, E., Merloni, A., & Rees, M. J. 2001, MNRAS, 324, 1147
- Rastinejad, J. C., Gompertz, B. P., Levan, A. J., et al. 2022, Nature, 612, 223
- Rojas, R. 1996, Neural Networks: A Systematic Introduction (Springer Nature)
- Rossi, A., Rothberg, B., Palazzi, E., et al. 2022, ApJ, 932, 1
- Rueda, J. A. & Ruffini, R. 2012, ApJ, 758, L7
- Russell, S. J. & Norvig, P. 2021, Artificial Intelligence: a modern approach, 4th edn. (Pearson)
- Sanna, A., Burderi, L., Di Salvo, T., et al. 2020, in Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series, Vol. 11444, Space Telescopes and Instrumentation 2020: Ultraviolet to Gamma Ray, ed. J.-W. A. den Herder, S. Nikzad, & K. Nakazawa, 114444X
- Savitzky, A. & Golay, M. J. E. 1964, Analytical Chemistry, 36, 1627 Stern, B. E. 1996, ApJ, 464, L111
- Stern, B. E. & Svensson, R. 1996, ApJ, 469, L109
- Troja, E., Fryer, C. L., O'Connor, B., et al. 2022, Nature, 612, 228
- van der Walt, S., Colbert, S. C., & Varoquaux, G. 2011, Computing in Science Engineering, 13, 22
- Van Rossum, G. & Drake, F. L. 2009, Python 3 Reference Manual (Scotts Valley, CA: CreateSpace)
- Vargas, F., De Colle, F., Brethauer, D., Margutti, R., & Bernal, C. G. 2022, The Astrophysical Journal, 930, 150

- Waskom, M. L. 2021, Journal of Open Source Software, 6, 3021
- Wei, J.-J. 2023, Physical Review Research, 5, 013019
- Williams, T., Kelley, C., & many others. 2023, Gnuplot 6.0: an interactive plotting
- Woosley, S. E. 1993, ApJ, 405, 273
- Yang, J., Ai, S., Zhang, B. B., et al. 2022, Nature, 612, 232
- Yi, S.-X., Lei, W.-H., Zhang, B., et al. 2017, Journal of High Energy Astrophysics, 13.1
- Zhang, B. 2018, The Physics of Gamma-Ray Bursts
- Zhang, B. & Yan, H. 2011, ApJ, 726, 90
- Zhang, B. B., Liu, Z. K., Peng, Z. K., et al. 2021, Nature Astronomy, 5, 911

Virtanen, P., Gommers, R., Oliphant, T. E., et al. 2020, Nature Methods, 17, 261 Wanderman, D. & Piran, T. 2010, MNRAS, 406, 1944

- Wang, F. Y. & Dai, Z. G. 2013, Nature Physics, 9, 465

- program