

Introducing scalar leptoquarks into a 3-3-1 model to solve the $(g - 2)_\mu$ puzzle

A. Doff^{b*} and C. A. de S. Pires^{a†}

^a*Departamento de Física, Universidade Federal da Paraíba,*

Caixa Postal 5008, 58051-970, João Pessoa, PB, Brazil and

^b*Universidade Tecnológica Federal do Paraná - UTFPR - DAFIS,*

R. Doutor Washington Subtil Chueire,

330 - Jardim Carvalho, 84017-220, Ponta Grossa, PR, Brazil

(Dated: March 29, 2024)

Abstract

In this work we introduce scalar leptoquarks into the 3-3-1 model with right-handed neutrinos with the aim of solving the $(g - 2)_\mu$ puzzle. We show that besides the model supports leptoquarks in the octet, sextet, triplet and singlet representations, we identified that only one specif leptoquark in the singlet representation leads to flip of chirality as required to generate positive and robust contribution to the $(g - 2)_\mu$. Then we calculate its contributions to $(g - 2)_\mu$ and to the decay process $\mu \rightarrow e\gamma$ and discuss the results.

* agomes@utfpr.edu.br

† cpires@fisica.ufpb.br

I. INTRODUCTION

The anomalous magnetic moment of the muon, $a_\mu = (\frac{g-2}{2})_\mu$, has long stood as a tantalizing puzzle in the particle physics[1, 2]. Experimental measurements of a_μ have consistently shown a deviation from the predictions of the Standard Model (SM) of the order of the electroweak contribution[3],

$$\begin{aligned} a_\mu^{Exp} &= 116592059(22) \times 10^{-11}, \\ a_\mu^{SM} &= 116591802(2) \times 10^{-11}, \\ \Delta a_\mu &= a_\mu^{Exp} - a_\mu^{SM} = 251(59) \times 10^{-11}. \end{aligned} \quad (1)$$

Such result has sparked an intense theoretical and experimental scrutiny to unravel its underlying causes[4–6].

One intriguing avenue of investigation of a_μ lies in the realm of leptoquark models. Recent theoretical studies have demonstrated the viability of chirality flip effects mediated by leptoquarks(LQ) in addressing the muon anomalous magnetic moment problem, potentially reconciling the observed anomaly with experimental measurements[6–8].

Leptoquarks are hypothetical particles in the form of scalars or vectors that have gained significant attention in the last decades[9]. These particles combine features of both leptons and quarks and have fractional electric charge[10, 11]. All these features have important impact in flavor physics[9].

Besides the standard model of particle physics does not predict leptoquarks, they emerge as essential components in various extensions to the Standard Model, including grand unified theories (GUTs)[12], supersymmetry[13], and composite models [14]. However the symmetries of the SM support the introduction of leptoquarks into it in the triplet, doublet and singlet representations. The last two representations are capables of generating flip of chirality among the fermions which generate an enhancement of $(g - 2)_\mu$ by the factor $\frac{m_t}{m_\mu}$ [15].

As in the SM case, the $SU(3)_C \times SU(3)_L \times U(1)_N$ (3-3-1) models face, too, serious difficulties in accommodating the current value of a_μ at one-loop[16–22]¹. In view of this, we introduce leptoquarks to the original particle content of a version of the 3-3-1 models that features right-handed neutrinos as third component of the lepton triplets (331RHN for short)[23, 24]. As it is the first

¹It was showed in Ref. [21] that the extension of the model to generate the inverse seesaw mechanism is not able to explain Δa_μ anomaly

work in which leptoquarks are introduced into this model.², we first identify the representation content of leptoquarks the model supports. Next we determine which leptoquark representation leads to flip of chirality among fermions in the 331RHN. We show that the model allows leptoquarks in the octet, sextet, triplet and singlet representation but only one specific leptoquark in the singlet representation is capable of generating flip of chirality among the fermions. We then focus on this leptoquark and calculate and discuss its contributions at one loop to $(g - 2)_\mu$ with the aim of obtaining which form of enhancement this leptoquark promote. We also discuss the contribution of such leptoquark to the $\mu \rightarrow e\gamma$ decay and if it discriminata family.

This paper is organized in the following way: in Sec. II we present the main aspects of the model. In Sec. III we introduce leptoquarks identifying representation content and Yukawa interactions. In Sec. IV we review the one loop contributions of the original 331RHN to $(g - 2)_\mu$ and calculate the contributions due to the singlet of leptoquark. In Sec. V we calculate the contribution of the singlet of leptoquark to the $\mu \rightarrow e\gamma$. In Sec. VI we discuss all the findings and in Sec. VII we present our conclusions.

II. THE MAIN ASPECTS OF THE MODEL

In the 331RHN right-handed neutrinos compose the third component of the leptonic triplets[23, 24],

$$L_{aL} = \begin{pmatrix} \nu_a \\ e_a \\ \nu_a^c \end{pmatrix}_L \sim (1, 3, -1/3), \quad e_{aR} \sim (1, 1, -1), \quad (2)$$

with $a = 1, 2, 3$ representing the three SM generations of leptons.

In the hadronic sector anomaly cancellation demands that one family transforms differently from the other two. Here we chose the first two families of quarks transforming as anti-triplet

²In fact leptoquarks were first introduced in [25] where it was considered the possibility of including leptoquarks in the minimal 3-3-1 model with the aim of evading the Landau pole

while the third one transforming as triplet by $SU(3)_L$,

$$\begin{aligned}
Q_{iL} &= \begin{pmatrix} d_i \\ -u_i \\ d'_i \end{pmatrix}_L \sim (3, \bar{3}, 0), u_{iR} \sim (3, 1, 2/3), \\
d_{iR} &\sim (3, 1, -1/3), \quad d'_{iR} \sim (3, 1, -1/3), \\
Q_{3L} &= \begin{pmatrix} u_3 \\ d_3 \\ u'_3 \end{pmatrix}_L \sim (3, 3, 1/3), u_{3R} \sim (3, 1, 2/3), \\
d_{3R} &\sim (3, 1, -1/3), \quad u'_{3R} \sim (3, 1, 2/3),
\end{aligned} \tag{3}$$

where the index $i = 1, 2$ is restricted to only two generations. The negative signal in the anti-triplet Q_{iL} is just to standardise the signals of the charged current interactions with the gauge bosons. The primed quarks are new heavy quarks with the usual $(+\frac{2}{3}, -\frac{1}{3})$ electric charges.

The original scalar sector of the model is composed by three triplets of scalars

$$\eta = \begin{pmatrix} \eta^0 \\ \eta^- \\ \eta'^0 \end{pmatrix}, \quad \rho = \begin{pmatrix} \rho^+ \\ \rho^0 \\ \rho'^+ \end{pmatrix}, \quad \chi = \begin{pmatrix} \chi^0 \\ \chi^- \\ \chi'^0 \end{pmatrix}, \tag{4}$$

with η and χ transforming as $(1, 3, -1/3)$ and ρ as $(1, 3, 2/3)$. After spontaneous breaking of the 3-3-1 symmetry ($\langle \eta^0 \rangle = v_\eta$, $\langle \rho^0 \rangle = v_\rho$, $\langle \chi'^0 \rangle = v_{\chi'}$ with $v_{\chi'} \gg v_\eta, v_\rho$) the scalar sector³ get composed by three CP-even scalars, h_1 , h_2 , and H where the first recovers the features of the standard Higgs and the other two are heavy scalars with mass proportional to $v_{\chi'}$ with $m_{h_1}^2 \approx \frac{vv_{\chi'}}{2}$ and $m_H^2 \sim v_{\chi'}^2$. The model also has in its spectrum of scalars one pseudoscalar we call A , whose mass is given by $m_A^2 \approx \frac{vv_{\chi'}}{2}$ and two singly charged scalars h_1^+ and h_2^+ with $m_{h_1^+}^2 \approx vv_{\chi'}$ and $m_{h_2^+}^2 \sim v_{\chi'}^2$. Except by h_1 that must recover the 125 GeV standard Higgs, all the other scalars have their masses determined by $v_{\chi'}$ which means that the natural energy scale of the masses of these scalars is the TeV scale⁴.

Here the most economical Yukawa interactions that generate the correct mass for all standard quarks and leptons are composed by the terms

$$-\mathcal{L}_Y \supset g_{ia} \bar{Q}_{iL} \eta^* d_{aR} + h_{3a} \bar{Q}_{3L} \eta u_{aR} + g_{3a} \bar{Q}_{3L} \rho d_{aR} + h_{ia} \bar{Q}_{iL} \rho^* u_{aR} + Y \bar{L} \rho e_R + \text{H.c.}, \tag{5}$$

³Here we follow Ref.[17], and assume $v_\eta = v_\rho = v$

⁴For a detailed discussion of the scalar sector, see Refs. [26–28]

where $a = 1, 2, 3$ and the parameters g_{ab} , h_{ab} and Y are Yukawa couplings that, for sake of simplification, we consider reals⁵.

The gauge sector of the model involves nine gauge bosons related to the electroweak sector, where four of them are the standard gauge bosons ($W^\pm$, Z^0 , γ) and the other five ($W'^\pm$, U^0 , $U^{0\dagger}$, Z') are the typical 3-3-1 gauge bosons, and eight gluons g related to the strong force. The standard gauge bosons recover the physics of the SM while the new gauge bosons have mass belonging to the TeV scale and interactions that connect the standard particle content with the specific 3-3-1 particle content⁶. The mass of W' and Z' are given by,

$$M_{Z'}^2 \approx \frac{g^2 c_W^2}{(3 - 4s_w^2)} v_{\chi'}^2, \quad M_{W'}^2 = \frac{g^2}{4} \left(\frac{v^2}{2} + v_{\chi'}^2 \right). \quad (6)$$

Observe that $v_{\chi'}$ practically determine the masses of the spectrum of new particle characteristic of the 3-3-1.

III. LEPTOQUARK: REPRESENTATION CONTENT AND YUKAWA INTERACTIONS

With all this in hand we are ready to determine the possible set of leptoquarks that the 331RHN supports. Observe that from the quark and lepton content of the model we can build the following bilinears,

$$\begin{aligned} \bar{L}_{aL}^C Q_{3L} &\sim (1, 3, -1/3) \times (3, 3, 1/3) \sim (3, 3^* \oplus 6, 0), \\ \bar{L}_{aL}^C Q_{iL} &\sim (1, 3, -1/3) \times (3, 3^*, 0) \sim (3, 1 \oplus 8, -1/3), \\ \bar{L}_{aL} d_{bR} &\sim (1, 3^*, 1/3) \times (3, 1, -1/3) \sim (3, 3^*, 0), \\ \bar{L}_{aL} u_{bR} &\sim (1, 3^*, 1/3) \times (3, 1, 2/3) \sim (3, 3^*, 1), \\ \bar{e}_{aR}^C d_{bR} &\sim (1, 1, -1) \times (3, 1, -1/3) \sim (3, 1, -4/3), \\ \bar{e}_{aR}^C u_{bR} &\sim (1, 1, -1) \times (3, 1, 2/3) \sim (3, 1, -1/3), \end{aligned} \quad (7)$$

where in the singlet of quarks above we are considering the exotic ones. According to these bilinears the 331RHN supports the following set of leptoquarks:

$$\begin{aligned} S_8 &\sim (3, 8, 1/3), & S_6 &\sim (3, 6, 0), \\ S_3 &\sim (3, 3, 0), & \tilde{S}_3 &\sim (3, 3, 1), \\ S &\sim (3, 1, 1/3), & \tilde{S} &\sim (3, 1, 4/3). \end{aligned} \quad (8)$$

⁵For a more comprehensive discussion of the Yukawa sector, see Ref. [29]

⁶Concerning gauge bosons of the model see Ref. [30, 31]

As we see, we have six categories of leptoquarks with each one of them leading to a complex phenomenology in regard to flavor physics.

According with the bilinears above, with the leptoquarks S_8 and S_6 we compose the following Yukawa interactions,

$$\mathcal{L} \supset y_{a3}^{LQ} \bar{L}_{aL}^C S_6 Q_{3L} + y_{ai}^{LQ} \bar{L}_{aL}^C S_8 Q_{iL} + H.c., \quad (9)$$

where the S_8 and S_6 components are given by

$$S_6 = \begin{pmatrix} \tilde{R}_1^{-5/3} & R_2^{-2/3} & R_3^{-5/3} \\ R_2^{-2/3} & \tilde{R}_2^{-2/3} & R_4^{-5/3} \\ R_3^{-5/3} & R_4^{-5/3} & \tilde{R}_3^{-1/3} \end{pmatrix}, \quad S_8 = \begin{pmatrix} \tilde{T}_1^{-2/3} & T_2^{-5/3} & T_3^{-2/3} \\ T_4^{-2/3} & \tilde{T}_2^{-5/3} & T_4^{-2/3} \\ T_5^{4/3} & T_6^{1/3} & \tilde{T}_3^{4/3} \end{pmatrix}. \quad (10)$$

The triplet S_3 and $\tilde{S}_3$ are composed by the following set of leptoquarks,

$$S_3 = \begin{pmatrix} S_3^{+1/3} \\ S_3^{+2/3} \\ S_3^{\prime+1/3} \end{pmatrix}_L, \quad \tilde{S}_3 = \begin{pmatrix} \tilde{S}_3^{+2/3} \\ \tilde{S}_3^{+4/3} \\ \tilde{S}_3^{\prime+2/3} \end{pmatrix}_L \quad (11)$$

The third components of these triplets have $F(S_3^{1/3}, \tilde{S}_3^{2/3}) = -2$ while the other components have $F = 0$ where $F = 3B + L$. Their Yukawa interactions are given by

$$\mathcal{L} \supset \bar{Y}_{a3}^{LQ} \epsilon_{ijk} (\bar{L}_{aL}^C)_i (S_3)_j (Q_{3L})_k + \tilde{Y}_{ab}^{LQ} \bar{L}_{aL} \tilde{S}_3 d_{bR} + H.c. \quad (12)$$

Concerning the allowed singlets of leptoquarks, we have that $\tilde{S}$ has $F = -2$ and, like the triplets above, do not leads to the flip of chirality. This means that they do not generate positive and robust contributions to the $g - 2$ of the muon.

The last leptoquark is the singlet S . It has $F = -2$ and compose the following Yukawa interactions with quarks and leptons,

$$\mathcal{L} \supset \tilde{g}_{ia}^{LQ} \bar{Q}_{iL}^C L_{aL} S + h_{ab}^{LQ} \bar{u}_{aR}^C e_{bR} S + H.c., \quad (13)$$

with $i = 1, 2$ and $a, b = 1, 2, 3$. Opening the first term above we get,

$$\bar{Q}_{iL}^C L_{aL} = \bar{d}_{iL}^C \nu_{aL} - \bar{u}_{iL}^C e_{aL} + \bar{\nu}_{aR} d'_{iL}. \quad (14)$$

Observe that according with the Yukawa interactions above we identify that S generates flip of chirality. It is due to this fact that from now on we focus our attention exclusively in developing some physical aspects of this leptoquark.

For reasons of simplification, and without any loss for the model, we make the following set of assumptions: charged leptons, neutrinos, right-handed quarks and the new quarks come in a diagonal basis. With this we obtain the Yukawa interactions among mass eigenstates of leptons and quarks with the leptoquark S ,

$$\mathcal{L} = \bar{u}_a^C (g_{ab}^{LQ} P_L + h_{ab}^{LQ} P_R) e_b S + \bar{d}_{aL}^C g_{ab}^{'LQ} \nu_{aL} S + \bar{\nu}_{aR} \tilde{g}_{ia}^{LQ} d_{iL}' S + H.c., \quad (15)$$

with $g_{ab}^{LQ} = -\tilde{g}_{ib}^{LQ} (V_L^u)_{bi}$ and $g_{ab}^{'LQ} = \tilde{g}_{ib}^{LQ} (V_L^d)_{bi}$. The physical quarks $\hat{u}_a = (u, c, t)$ and $\hat{d}_a = (d, s, b)$ are related to the symmetric ones by means of the transformations $\hat{u}_L = V_L^{\dagger u} u_L$ and $\hat{d}_L = V_L^{\dagger d} d_L$. The first term in Eq. (15) is the one that generates the flip of chirality among the interaction of S with the charged fermions. The interest in this interaction is because it provides a positive and robust contribution to the $g - 2$ of the muon as we see below.

As we said in the introduction, the 331RHN with its original particle content, and even modified to implement the inverse seesaw mechanism, is not able to accommodate the current muon $g - 2$. But now with the introduction of singlet of leptoquark S we may cure this deficiency of the model. For this we made a brief review the contributions of the original particle content of the model to the muon $g - 2$ at one-loop.

IV. REVIEW OF THE ONE LOOP CONTRIBUTION TO Δa_μ IN THE 331RHN AND THE CONTRIBUTION OF THE LEPTOQUARK S

A. 331RHN original particle content contribution to Δa_μ

We review the main contributions at one loop to the muon anomalous magnetic moment $a_\mu = \frac{(g-2)_\mu}{2}$ in the 331RHN. For this we follow Ref. [20] where it is took $v_\eta = v_\rho = v = 174$ GeV. We are interested in the discrepancy Δa_μ produced by the new charged gauge boson $W'^\pm$, Z' , the single charged scalar h_1^+ , the neutral scalars h_2 and the pseudo Goldstone A^7 .

The interactions of interest are given by

$$\mathcal{L}_l^{CC} = -\frac{g}{2\sqrt{2}} [\bar{\nu}_R^c \gamma^\mu (1 - \gamma_5) \bar{l} W_\mu'^-], \quad (16)$$

⁷The charged scalar h_2^+ and the neutral scalar H do not contribute to $(g - 2)_\mu$ [28]




FIG. 1. One-loop contributions to the $g - 2$ of the muon in the 331RHN where ϕ and ϕ^- represent the neutral and charged scalar contributions

$$\mathcal{L}^{NC} = \bar{f} \gamma^\mu [g_V(f) + g_A(f)\gamma_5] f Z'_\mu, \quad (17)$$

$$\mathcal{L}_Y^{h_1^+} = \sqrt{2} \frac{m_l}{v} \bar{\nu}_{lL} l_R h_1^+, \quad (18)$$

$$\mathcal{L}^{h_2+A} = \frac{m_\mu}{\sqrt{2}v} \bar{\mu} \mu h_2 + i \frac{m_\mu}{v} \bar{\mu} \gamma_5 \mu A, \quad (19)$$

where g_V and g_A are the vector and axial coupling constants and are given by,

$$g'_V(l) = \frac{g}{4c_W} \frac{(1 - 4s_W^2)}{\sqrt{3 - 4s_W^2}}, \quad g'_A(l) = -\frac{g}{4c_W \sqrt{3 - 4s_W^2}}, \quad (20)$$

The Feynman diagrams displaying the contributions of these interactions to $(g - 2)_\mu$ are showed in FIG. 1.

The evaluation of these contributions at one-loop to Δa_μ generated by these set of particles follow the procedure of Ref. [20]. To avoid being repetitive we simply quote them in the heavy mediator regime. With that being said, we have the following contributions of the 331RHN to the Δa_μ ,

$$\Delta a_\mu(\nu, W') \simeq \frac{1}{4\pi^2} \frac{m_\mu^2}{M_{W'}^2} \left| \frac{g}{2\sqrt{2}} \right|^2 \left(\frac{5}{3} \right), \quad (21)$$

$$\Delta a_\mu(\mu, Z') \simeq \frac{-1}{4\pi^2} \frac{m_\mu^2}{M_{Z'}^2} \frac{1}{3} \left| -\frac{g}{4c_W \sqrt{3 - 4s_W^2}} \right|^2 \left[-|1 - 4s_W^2|^2 + 5 \right], \quad (22)$$

$$\Delta a_\mu(h_1^+) \simeq \frac{-1}{4\pi^2} \frac{m_\mu^2}{M_{h_1^+}^2} \left| \frac{m_\mu \sqrt{2}}{2v_\eta} \right|^2 \frac{1}{6}, \quad (23)$$

$$\Delta a_\mu(h_2) \simeq \frac{1}{4\pi^2} \frac{m_\mu^2}{M_{h_2}^2} \left(\frac{m_\mu \sqrt{2}}{2v_\eta} \right)^2 \left[\frac{1}{6} + \left(\frac{3}{4} + \log \left(\frac{m_\mu}{M_{h_2}} \right) \right) \right]. \quad (24)$$




FIG. 2. Overall contribution to Δa_μ from Eqs.(21)-(25) to the 331RHN model. The dash-dotted bands in red are delimited by the current 1σ bound.

$$\Delta a_\mu(A) = -\frac{m_\mu^2}{8\pi^2 M_A^2} \left(\frac{g A_\mu m_\mu}{2M_W} \right)^2 H \left(\frac{m_\mu^2}{M_A^2} \right), \quad (25)$$

with $H(y) = \int_0^1 \frac{dx x^3}{1-x+x^2 y}$ where $y = \frac{m_\mu^2}{M_A^2}$. In summary, the 331RHN yields five contributions to Δa_μ with only W' and h_2 contributions being positive. In view of this, according to Ref. [17], the only way the one-loop contribution would accommodate the current prediction of the Δa_μ was if the positive contributions of the W' and h_2 could surpass the other ones. This could happen for W' having mass at the electroweak scale which is completely discarded since current LHC bounds demands the gauge bosons of the 331RHN have mass at the TeV scale. To be precise LHC demands $m_{Z'}$ with mass in the range $(3 - 4)$ TeVs[32], depending if Z' may decay in the new quarks or don't[33]. This implies $v_{\chi'}$ in the range $10 - 12$ TeVs. In FIG.2 we present the prediction of the 331RHN to Δa_μ in function of $v_{\chi'}$ varying in the range $1 - 12$ TeVs. As we can see in FIG. 2 one-loop contributions to Δa_μ in the 331RHN are not able to accommodate the current Δa_μ . Observe that for $v_{\chi'}$ around 10^4 GeV the prediction to Δa_μ lies three order of magnitude below the experimental prediction. Thus the one loop contribution to Δa_μ in the 331RHN is practically negligible. As sad in the introduction, it is hard to find modifications of the 331RHN that accommodate the current Δa_μ value. A simple route to accommodate Δa_μ into the 331RHN is via the introduction of vectorlike leptons and inert scalar triplets to the model[17]. What we are going to show next is that the addition of only one singlet of leptoquark resolves the $(g - 2)_\mu$ anomaly.




FIG. 3. One-loop diagrams contributing to Δa_μ and $\mu \rightarrow e\gamma$ decay induced by S leptoquarks: To the contributions to Δa_μ in this figure $l_i = l_j = l_2 = \mu$, while to muon decay we have $l_i = l_2$ and $l_j = l_1 = e$.

B. The contribution of S to Δa_μ in the 331RHN

The one-loop contributions to Δa_μ due to the leptoquark S , depicted in the FIG. 3 and in according to the interactions in Eq. (15) are given by[6, 9]

$$\Delta a_\mu^{LQ} = \frac{m_\mu^2}{48\pi^2 m_S^2} f(m_\mu) \left(\frac{m_q}{m_\mu} g_{a2}^{LQ} h_{a2}^{LQ} L_1(x_q) + \frac{(g_{a2}^{LQ})^2 + (h_{a2}^{LQ})^2}{4} L_2(x_q) \right), \quad (26)$$

where $x_q = \frac{m_q^2}{m_S^2}$ and

$$\begin{aligned} L_1(x) &= 4F_F(x) - F_C(x), \quad L_2(x) = 2F_E(x) - F_B(x) \\ F_F(x) &= \frac{3(-3 + 4x - x^2 - 2\ln x)}{2(1-x)^3}, \quad F_E(x) = \frac{2(2 + 3x - 6x^2 + x^3 + 6x\ln x)}{(1-x)^4}, \\ F_C(x) &= \frac{3(1 - x^2 + 2x\ln x)}{(1-x)^3}, \quad F_B(x) = \frac{2(1 - 6x + 3x^2 + 2x^3 - 6x^2\ln x)}{(1-x)^4}. \\ f(m_\mu) &= \left(1 + \frac{4\alpha}{\pi} \ln\left(\frac{m_\mu}{m_S}\right)\right). \end{aligned} \quad (27)$$

We consider only the maximum contribution given by Eq. (26) which is due to the top quark ($a = 3$ in Eq. (26)). The free parameters involved in such contributions are the Yukawa couplings g_{32}^{LQ} , h_{32}^{LQ} and the mass of S which we called m_S . The current LHC bounds demands the lowest leptoquark to have mass in the range $O(0.98-1.73)TeV$, depending on the spin of leptoquarks and its couplings to SM fermions[34]. Here we adopt the current assumption $m_S = 1.8TeV$ and extract constraint on the Yukawa couplings g_{32}^{LQ} and h_{32}^{LQ} .

Our numerical result is displayed in FIG. 4. As we can see there the existence of a simple singlet of leptoquark S presenting flip of chirality give positive and robust contributions to Δa_μ explaining




FIG. 4. In this figure we show in parameter space $(g_{32}^{LQ} - h_{32}^{LQ})$ the allowed values to Δa_μ . The dash-dotted line corresponds to central value obtained to Δa_μ . The respective bands $1(2)\sigma$ are represented in yellow(green)

easily the observed value of Δa_μ . That is a very nice result in what concern the development of the 331RHN. In the figure the dash-dotted line corresponds to central value $\Delta a_\mu = 25.1 \times 10^{-10}$. The Yellow(Green) bands corresponds to the current $1\sigma(2\sigma)$ bounds found by requiring $\Delta a_\mu \lesssim 59 \times 10^{-11}$. In the next section, we complement this result with the study of the decay $\mu \rightarrow e\gamma$ once the contributions are the same being necessary just to take of-shell photon in FIG. 3.

V. CONTRIBUTIONS OF S LEPTOQUARKS TO $\mu \rightarrow e\gamma$ DECAY

In the FIG. 3 we show the contributions of S leptoquarks to the muon decay $\mu \rightarrow e\gamma$. These contributions are similar to the corrections of S to Δa_μ , and in this Feynman diagrams we consider only the chirality enhanced terms, given by Eq.(13), that explain the observed Δa_μ . To this decay the branching ratio presents the following form[35]

$$BR(\mu \rightarrow e\gamma) = \frac{m_\mu^5}{16\pi\Gamma_\mu} (|A_2^L|^2 + |A_2^R|^2) \quad (28)$$




FIG. 5. Allowed parameter regions for the $\mu \rightarrow e\gamma$ decay, assuming that Δa_μ is explained by top quark chirally enhanced term. The contextualization of the curves behavior are described in the text.

where $\Gamma_\mu = 2.996 \times 10^{-19} \text{GeV}$ and the structure of the dipole form factors $A_{L,R}^2$ are given by

$$\begin{aligned} A_2^L &= -\frac{1}{16\pi^2} \frac{e}{6m_S^2} \left(\frac{m_q}{m_\mu} g_{a1}^{LQ} h_{a2}^{LQ} L_1(x_q) + \frac{1}{4} h_{a1}^{LQ} h_{a2}^{LQ} L_2(x_q) \right), \\ A_2^R &= -\frac{1}{16\pi^2} \frac{e}{6m_S^2} \left(\frac{m_q}{m_\mu} h_{a1}^{LQ} g_{a2}^{LQ} L_1(x_q) + \frac{1}{4} g_{a1}^{LQ} g_{a2}^{LQ} L_2(x_q) \right). \end{aligned} \quad (29)$$

Following the same line of the previous section, we will consider the maximum contribution to $\mu \rightarrow e\gamma$ decay that is due to the top quark ($a = 3$).

Considering the comparison with the experimental results to Δa_μ , in the FIG. 4 we show the bounds on products of g_{32}^{LQ} and h_{32}^{LQ} couplings in parameter space. In this figure the band in Green restricts the product of these couplings, i. e. the left and handed leptoquarks couplings of top with the muon. As consequence the 2σ band results in the following allowed range of possible couplings

$$3.1 \times 10^{-3} < g_{32}^{LQ} * h_{32}^{LQ} < 9.3 \times 10^{-3}. \quad (30)$$

In the Eq.(29) the left and handed leptoquarks couplings of top with the electron are g_{31}^{LQ} and h_{31}^{LQ} . In order to find the allowed parameter space in the $(g_{31}^{LQ} - h_{31}^{LQ})$ plane, currently permitted by experimental bounds[36][37], we can consider the above result, combined with the expression to the branching ratio given by Eq.(28).

Therefore, considering the Eqs.(28), (29) and (30) we can write the following inequality

$$\left(|h_{32}^{LQ}|^2 |h_{31}^{LQ}|^2 + |g_{32}^{LQ}|^2 |g_{31}^{LQ}|^2 \right) < 3 \times 10^{-2} BR(\mu \rightarrow e\gamma). \quad (31)$$

In order to generate the parameter region allowed, we will consider the following possibilities of

top couplings

$$\begin{aligned}
& \text{(a) Symmetric leptoquark coupling : } h_{32}^{LQ} = g_{32}^{LQ}, \\
& \text{(b) Asymmetric leptoquark coupling : } h_{32}^{LQ} > g_{32}^{LQ}, \\
& \text{(c) Asymmetric leptoquark coupling : } h_{32}^{LQ} < g_{32}^{LQ}.
\end{aligned} \tag{32}$$

Assuming the lower bound ($3.1 \times 10^{-3} < g_{32}^{LQ} * h_{32}^{LQ}$), we obtain to Case (a) ($|h_{32}^{LQ}|^2 = |g_{32}^{LQ}|^2 > 3.1 \times 10^{-3}$), while that to $h_{32}^{LQ} = 2$, we have to Case (b) ($g_{32}^{LQ} > 1.55 \times 10^{-3}$) and similarly to $g_{32}^{LQ} = 2$ the Case (c) corresponds to ($h_{32}^{LQ} > 1.55 \times 10^{-3}$).

Currently the experimental limits to $BR(\mu \rightarrow e\gamma)$ correspond to $BR(\mu \rightarrow e\gamma) = 4.2 \times 10^{-13}$ [36] and $BR(\mu \rightarrow e\gamma) = 6 \times 10^{-14}$ [37]. In FIG. 5 we show in the plane ($g_{31}^{LQ} - h_{31}^{LQ}$) the allowed values to $g_{31}^{LQ} * h_{31}^{LQ}$, assuming the considerations indicated in the paragraph above.

In FIG. (5a), corresponding to the symmetric case, the region filled in blue leads to the upper bound to the allowed values of couplings as $g_{31}^{LQ} * h_{31}^{LQ} < 2.056 \times 10^{-12}$, while for region filled in green we have an more restrictive upper bound $g_{31}^{LQ} * h_{31}^{LQ} < 2.93 \times 10^{-13}$.

For the asymmetric cases corresponding to FIG. (5b) and (5c), the limits obtained are smaller compared to case (a), so that they are within the region covered in this situation. Considering the combination of results presented in these figures, that illustrate the allowed values to S leptoquarks couplings permitted by actual experimental bounds, we found that leptoquarks couplings to electrons must be more than 4 orders of magnitude smaller than the corresponding couplings to muons.

VI. DISCUSSION

Anomalies cancellation inside 3-3-1 models demand that at least one family of quarks transforms differently from the other two[38–40]. This means that for each version of 3-3-1 models there are three variants as discussed in Ref.[41, 42]. Each variant has its proper implications in flavor physics. However, in adding leptoquarks to the 331RHN, besides the Yukawa interactions among leptoquarks and fermions seems to discriminate families, as expressed in Eq. (13), when we go to the mass eigenstates we see that leptoquarks do not distinguish families, as expressed in Eq. (15). This is so because multiplet of leptoquarks do not contribute to the masses of the fermions after spontaneous breaking of the symmetry. Then we can always redefine the Yukawa couplings by means of the mixing matrix that transform fermions eigenstate of symmetry into the

mass eigenstates. Thus the fact we have chosen the third family transforming as triplet and the first and second ones transforming as anti-triplet was a mere question of convenience.

Another point we would like to call the attention is that while the standard model supports two representation of leptoquarks inducing flip of chirality[15], namely a singlet and a doublet of leptoquarks, in the 331RHN, besides it supports a more extensive set of multiplet of leptoquarks (octet, sextet, triplet and singlet), however only a specific leptoquark transforming as singlet by the 3-3-1 symmetry leads to the flip of chirality. Nevertheless, such singlet is sufficient to do the job of explaining Δa_μ .

Observe that the leptoquark S in eq. (15) enhances the contribution to $(g - 2)_\mu$ by a factor $\frac{m_t}{m_\mu}$. However if we switch on the VEVs of χ^0 and η^0 we have, as main consequence, a mixing among the standard quarks with the new ones. As main implication to $(g - 2)_\mu$ we will have an enhancement of Δa_μ by a factor of $\frac{m_{u'}}{m_\mu}$. A $m_{u'}$ at TeV scale would give a very robust contribution to Δa_μ rendering constraints in the mixing among standard and new quarks which translate in constraints on the VEVs v_χ and $v_{\eta'}$.

In FIG. 3 we took $q = t$ which implies an enhancement of the contributions to Δa_μ by a factor of $\frac{m_t}{m_\mu}$. This is the reason why people consider in the standard model only the interaction of leptoquarks with the third generation of quarks. We can not do this here because the symmetries do not allow. The leptoquark S has an intriguing set of Yukawa interactions as we can see in Eq. (13). The interaction with the quark top arises due to the mixing among the quarks given by the mixing matrix $V_L^{u,d}$.

VII. CONCLUSION

In this work we introduced leptoquarks into the 3-3-1 model with right-handed neutrinos with the aim of resolving the problem of the Δa_μ anomaly since the original version of the model gives a negligible contribution to Δa_μ . We saw that the model supports multiplet of leptoquarks in the representation of octet, sextet, triplet and singlet. We determined the particle content of these multiplets and presented they Yukawa interactions with the fermions of the model.

Besides this large spectrum of leptoquarks, only a specific singlet S is capable of generate a robust contribution to Δa_μ since only it leads to flip of chirality among the fermions. We then developed the main aspects of the leptoquark S in what concern its contributions to Δa_μ and to the muon decay channel $\mu \rightarrow e\gamma$. Our main result is that singlet of leptoquark is not only viable

but also the most economical extension of the 3-3-1 model that explain the Δa_μ anomaly.

ACKNOWLEDGMENTS

C.A.S.P was supported by the CNPq research grants No. 311936/2021-0.

-
- [1] H. N. Brown *et al.* (Muon g-2), Phys. Rev. Lett. **86**, 2227 (2001), arXiv:hep-ex/0102017.
 - [2] F. Jegerlehner, *The Anomalous Magnetic Moment of the Muon*, Vol. 274 (Springer, Cham, 2017).
 - [3] G. W. Bennett *et al.* (Muon g-2), Phys. Rev. Lett. **89**, 101804 (2002), [Erratum: Phys.Rev.Lett. 89, 129903 (2002)], arXiv:hep-ex/0208001.
 - [4] M. Lindner, M. Platscher, and F. S. Queiroz, Phys. Rept. **731**, 1 (2018), arXiv:1610.06587 [hep-ph].
 - [5] T. Aoyama *et al.*, Phys. Rept. **887**, 1 (2020), arXiv:2006.04822 [hep-ph].
 - [6] P. Athron, C. Balázs, D. H. J. Jacob, W. Kotlarski, D. Stöckinger, and H. Stöckinger-Kim, JHEP **09**, 080 (2021), arXiv:2104.03691 [hep-ph].
 - [7] A. Crivellin and M. Hoferichter, JHEP **07**, 135 (2021), [Erratum: JHEP 10, 030 (2022)], arXiv:2104.03202 [hep-ph].
 - [8] D. Stöckinger and H. Stöckinger-Kim, Front. in Phys. **10**, 944614 (2022).
 - [9] I. Doršner, S. Fajfer, A. Greljo, J. F. Kamenik, and N. Košnik, Phys. Rept. **641**, 1 (2016), arXiv:1603.04993 [hep-ph].
 - [10] J. C. Pati and A. Salam, Phys. Rev. D **10**, 275 (1974), [Erratum: Phys.Rev.D 11, 703–703 (1975)].
 - [11] H. Georgi and S. L. Glashow, Phys. Rev. Lett. **32**, 438 (1974).
 - [12] J. C. Pati and A. Salam, Phys. Rev. D **8**, 1240 (1973).
 - [13] J. L. Hewett and T. G. Rizzo, Phys. Rept. **183**, 193 (1989).
 - [14] B. Schrempp and F. Schrempp, Phys. Lett. B **153**, 101 (1985).
 - [15] E. Coluccio Leskow, G. D’Ambrosio, A. Crivellin, and D. Müller, Phys. Rev. D **95**, 055018 (2017), arXiv:1612.06858 [hep-ph].
 - [16] C. Kelso, H. N. Long, R. Martinez, and F. S. Queiroz, Phys. Rev. D **90**, 113011 (2014), arXiv:1408.6203 [hep-ph].
 - [17] A. S. De Jesus, S. Kovalenko, F. S. Queiroz, C. Siqueira, and K. Sinha, Phys. Rev. D **102**, 035004 (2020), arXiv:2004.01200 [hep-ph].

- [18] A. E. Cárcamo Hernández, Y. Hidalgo Velásquez, S. Kovalenko, H. N. Long, N. A. Pérez-Julve, and V. V. Vien, *Eur. Phys. J. C* **81**, 191 (2021), arXiv:2002.07347 [hep-ph].
- [19] N. A. Ky, H. N. Long, and D. Van Soa, *Phys. Lett. B* **486**, 140 (2000), arXiv:hep-ph/0007010.
- [20] A. S. de Jesus, S. Kovalenko, F. S. Queiroz, C. A. de S. Pires, and Y. S. Villamizar, *Phys. Lett. B* **809**, 135689 (2020), arXiv:2003.06440 [hep-ph].
- [21] J. a. P. Pinheiro, C. A. de S. Pires, F. S. Queiroz, and Y. S. Villamizar, *Phys. Lett. B* **823**, 136764 (2021), arXiv:2107.01315 [hep-ph].
- [22] T. T. Hong, N. H. T. Nha, T. P. Nguyen, L. T. T. Phuong, and L. T. Hue, *PTEP* **2022**, 093B05 (2022), arXiv:2206.08028 [hep-ph].
- [23] R. Foot, H. N. Long, and T. A. Tran, *Phys. Rev. D* **50**, R34 (1994), arXiv:hep-ph/9402243.
- [24] J. C. Montero, F. Pisano, and V. Pleitez, *Phys. Rev. D* **47**, 2918 (1993), arXiv:hep-ph/9212271.
- [25] A. Doff and C. A. de S. Pires, *Nucl. Phys. B* **992**, 116254 (2023), arXiv:2302.08578 [hep-ph].
- [26] H. N. Long, *Mod. Phys. Lett. A* **13**, 1865 (1998), arXiv:hep-ph/9711204.
- [27] R. A. Diaz, R. Martinez, and F. Ochoa, *Phys. Rev. D* **69**, 095009 (2004), arXiv:hep-ph/0309280.
- [28] J. P. Pinheiro and C. A. de S. Pires, *Phys. Lett. B* **836**, 137584 (2023), arXiv:2210.05426 [hep-ph].
- [29] A. Doff, C. A. de S. Pires, and P. S. R. da Silva, *Phys. Rev. D* **74**, 015014 (2006), arXiv:hep-ph/0604021.
- [30] H. N. Long, *Phys. Rev. D* **54**, 4691 (1996), arXiv:hep-ph/9607439.
- [31] Q.-H. Cao and D.-M. Zhang, (2016), arXiv:1611.09337 [hep-ph].
- [32] Y. A. Coutinho, V. Salustino Guimarães, and A. A. Nepomuceno, *Phys. Rev. D* **87**, 115014 (2013), arXiv:1304.7907 [hep-ph].
- [33] A. Alves, L. Duarte, S. Kovalenko, Y. M. Oviedo-Torres, F. S. Queiroz, and Y. S. Villamizar, *Phys. Rev. D* **106**, 055027 (2022), arXiv:2203.02520 [hep-ph].
- [34] A. Hayrapetyan *et al.* (CMS), (2024), arXiv:2402.08668 [hep-ex].
- [35] K. T. J. Hisano, T. Moroi and M. Yamaguchi, *Phys. Rev. D* **53**, 2442 (1996), arXiv:9510309 [hep-ph].
- [36] A. M. Baldini *et al.* (MEGI collaboration), *Eur. Phys. J. C* **80**, 858 (2020), arXiv:[2005.00339] [hep-ph].
- [37] A. M. Baldini *et al.* (MEGII collaboration), *Eur. Phys. J. C* **78**, 380 (2018), arXiv:[1801.04688] [hep-ph].
- [38] P. H. Frampton, *Phys. Rev. Lett.* **69**, 2889 (1992).
- [39] D. Ng, *Phys. Rev. D* **49**, 4805 (1994), arXiv:hep-ph/9212284.

- [40] J. T. Liu and D. Ng, Phys. Rev. D **50**, 548 (1994), arXiv:hep-ph/9401228.
- [41] V. Oliveira and C. A. de S. Pires, Phys. Lett. B **846**, 138216 (2023), arXiv:2211.03835 [hep-ph].
- [42] V. Oliveira and C. A. d. S. Pires, J. Phys. G **50**, 115002 (2023), arXiv:2208.00420 [hep-ph].