On (θ, Θ) -cyclic codes and their applications in constructing QECCs

Awadhesh Kumar Shukla¹, Sachin Pathak², Om Prakash Pandey¹, Vipul Mishra¹, Ashish Kumar Upadhyay¹

¹Department of Mathematics, Banaras Hindu University, Varanasi, Uttar Pradesh 221005, India.

²Department of Mathematics and Basic Sciences, NIIT University, Neemrana 301705, India. April 2, 2024

Abstract

Let \mathbb{F}_q be a finite field, where q is an odd prime power. Let $R = \mathbb{F}_q + u\mathbb{F}_q + v\mathbb{F}_q + uv\mathbb{F}_q$ with $u^2 = u, v^2 = v, uv = vu$. In this paper, we study the algebraic structure of (θ, Θ) -cyclic codes of block length (r, s) over $\mathbb{F}_q R$. Specifically, we analyze the structure of these codes as left $R[x:\Theta]$ -submodules of $\mathfrak{R}_{r,s} = \frac{\mathbb{F}_q[x:\theta]}{\langle x^r-1 \rangle} \times \frac{R[x:\Theta]}{\langle x^s-1 \rangle}$. Our investigation involves determining generator polynomials and minimal generating sets for this family of codes. Further, we discuss the algebraic structure of separable codes. A relationship between the generator polynomials of (θ, Θ) -cyclic codes over $\mathbb{F}_q R$ and their duals is established. Moreover, we calculate the generator polynomials of dual of (θ, Θ) -cyclic codes. As an application of our study, we provide a construction of quantum error-correcting codes (QECCs) from (θ, Θ) -cyclic codes of block length (r, s) over $\mathbb{F}_q R$. We support our theoretical results with illustrative examples.

Keywords: (θ, Θ) -cyclic code, Generator polynomials, Dual codes, Separable Codes, QECCs.

1 Introduction

Cyclic codes have been extensively researched and analyzed by researchers due to their abundant algebraic structure. This inherent mathematical richness makes cyclic codes one of the most significant classes of error-correcting codes. In 1994, Hammon et al. [11] presented a novel viewpoint by constructing non-linear codes over \mathbb{Z}_2 using the Gray map derived from codes over \mathbb{Z}_4 . Over the last three decades, scholars have dedicated significant attention to the exploration of the characteristics of cyclic codes, particularly their generator polynomials.

In 1997, Rifà and Pojul [30] pioneered introducing codes over mixed alphabets. Subsequently, Browers et al. [10] examined the codes over the mixed alphabets \mathbb{Z}_2 and \mathbb{Z}_3 , focusing on deriving the maximum size of error-correcting codes. After that many scholars have concentrated extensively on mixed alphabets. In 2010, Borges et al. [6] delved into the realm of mixed alphabets and examined the $\mathbb{Z}_2\mathbb{Z}_4$ -additive codes. In their work, they focused not only

Email: shuklaawadhesh@bhu.ac.in (A. K. Shukla), sachiniitk93@gmail.com (Sachin Pathak), opandey1302@bhu.ac.in (O. P. Pandey), mishravipul10@bhu.ac.in (V. Mishra), upadhyay@bhu.ac.in (A. K. Upadhyay).

on finding the generator matrices but also the parity check matrices and other parameters for these codes. Building upon prior research, Aydogdu and Siap [4] delved deeper into the realm of $\mathbb{Z}_2\mathbb{Z}_{2^s}$ -additive codes. Abualrub et al. [1] examined $\mathbb{Z}_2\mathbb{Z}_4$ -additive cyclic codes, with a particular focus on identifying their generator polynomials and minimum spanning sets in 2014. This research led to the construction of several Maximum Distance Separable (MDS) codes.

The double cyclic codes over \mathbb{Z}_2 were analyzed and examined in 2018 by Borges et al. [7] for the first time. Their study involved considering \mathbb{Z}_2 -double cyclic codes as $\mathbb{Z}_2[x]$ -submodules of $\frac{\mathbb{Z}_2[x]}{\langle x^r-1 \rangle} \times \frac{\mathbb{Z}_2[x]}{\langle x^s-1 \rangle}$. Within this research, they successfully determined the generating sets and derived numerous optimal codes. The structural characteristics of double cyclic codes over \mathbb{Z}_4 was thoroughly studied by Gao et al. [20]. In their research, they focused on generator polynomials and minimum spanning sets and derived corresponding dual codes. Furthermore, they find several non-linear optimal codes over \mathbb{Z}_2 . In 2020, Dinh et al. [16] examined the structural characteristics of cyclic codes over $\mathbb{F}_q R$ and their corresponding duals. Additionally, they applied their findings to construct several novel QECCs.

In 2007, D. Bouchers et al. [8] introduced skew cyclic codes. They derived improved linear codes with better parameters in [9]. Shortly thereafter, in 2011, Siap et al. [32] extended the study of [9] and studied the skew cyclic codes of arbitrary lengths. Subsequent to their groundbreaking work, several researchers delved into the exploration of skew rings of the form $R = \mathbb{F}_q + v\mathbb{F}_q$, where $v^2 = v$, employing various automorphisms on R [see [19, 21, 27]]. In 2015, Yao et al. [13] conducted a comprehensive study on skew cyclic codes over the ring $R = \mathbb{F}_q + u\mathbb{F}_q + v\mathbb{F}_q + uv\mathbb{F}_q$, with $u^2 = u, v^2 = v$ and uv = vu.

A pivotal moment in the development of quantum error-correction occurred in 1995 when Shor [31] introduced the first QECC. Following this, Calderbank et al. [12] introduced a technique for constructing QECCs in 1998. They achieved this by leveraging classical errorcorrecting codes and successfully established their existence through proof. Building upon the concept introduced by Calderbank et al. [12], numerous QECCs have been obtained by adapting cyclic codes and their extensions, working within the realms of both finite rings and finite fields.

Recently, Aydogdu et al. studied double skew cyclic code over \mathbb{F}_q in [3]. Within this research paper, they examined the generator polynomials of this family of codes and their corresponding dual codes. Skew polynomial rings allow for numerous factorizations of every polynomial, they do not have unique factorization features. This makes the study of codes over non-commutative rings especially important.

After being motivated from the study of [3], in our paper, we consider $R = \mathbb{F}_q + u\mathbb{F}_q + v\mathbb{F}_q + uv\mathbb{F}_q$, with $q = p^m$, where p is an odd prime and $u^2 = u, v^2 = v$ and uv = vu, m is natural number and study structural properties of (θ, Θ) -cyclic code over $\mathbb{F}qR$ and their application in constructing in QECCs. This paper is outlined as follows: Section 2 provides fundamental concepts, notations, and prior findings relevant to the study of (θ, Θ) -cyclic codes of block length (r,s) over $\mathbb{F}_q R$. Moreover, a naturally extended Gray map from $\mathbb{F}_q^r \times R^s \to \mathbb{F}_q^{r+4s}$ is defined, and some properties of this map are established. In Section 3, we explore the generator polynomials by Theorem 3.12 and minimal generating set by Theorem 3.19 of this family of codes. Moreover, we discuss the generator polynomials of separable code in Theorem 3.18. Additionally, we find some near-optimal and optimal codes over \mathbb{F}_9 in Table 1. In Section 4, we conduct the study on the dual of (θ, Θ) -cyclic codes of block length (r, s) over $\mathbb{F}_q R$ and determine the generator polynomials of dual codes in Theorem 4.11. Section 5 focuses on constructing QECCs from separable (θ, Θ) -cyclic codes over $\mathbb{F}_q R$. In Examples 5.8, 5.9 and 5.10, we provide an explanation of constructing QECCs from separable (θ, Θ) -cyclic codes of block length (r, s)over $\mathbb{F}_q R$. In Table 2, we obtain several maximum distance separable (MDS) QECCs from θ -cyclic code over \mathbb{F}_q and new QECCs having better parameters than existing QECCs from Θ -cyclic code over R and (θ, Θ) -cyclic code over $\mathbb{F}_q R$ are given in tables 3 and 4. In Section 6, this paper is concluded.

2 Preliminaries

Let \mathbb{F}_q be a finite field, where q is a power of an odd prime p. The set \mathbb{F}_q^r consists of tuples of length r, where elements can be added and multiplied by scalars following standard rules and forms a vector space over \mathbb{F}_q . A non-empty subspace \mathfrak{C}_r of \mathbb{F}_q^r is called a linear code of length r over \mathbb{F}_q , with its dimension indicating the size of \mathfrak{C}_r . Elements in \mathfrak{C}_r are termed codewords over \mathbb{F}_q . The Hamming weight of any codeword $\mathbf{c} = (c_0, c_1, \ldots, c_{r-1}) \in \mathfrak{C}_r$, denoted as $w_H(\mathbf{c})$, counts the number of non-zero components. Furthermore, the Hamming distance between any two codewords, \mathbf{c} and \mathbf{c}' , is defined as the number of positions where they differ, given by $d_H(\mathbf{c}, \mathbf{c}') = w_H(\mathbf{c} - \mathbf{c}')$. The Hamming distance of a linear code \mathfrak{C}_r is determined by

 $d_H((\mathfrak{C}_r) = \min \{ d_H(\mathbf{c}, \mathbf{c}'), \forall \mathbf{c}, \mathbf{c}' \in \mathfrak{C}_r, \mathbf{c} \neq \mathbf{c}' \}.$

Definition 2.1. Let $\operatorname{Aut}(\mathbb{F}_q)$ denote the set of all automorphism on \mathbb{F}_q and $\theta \in \operatorname{Aut}(\mathbb{F}_q)$. Then we define the set

$$\mathbb{F}_q[x:\theta] = \{g_0 + g_1 x + g_2 x^2 + \dots + g_n x^n, g_i \in \mathbb{F}_q, \ \theta \in \operatorname{Aut}(\mathbb{F}_q)\}.$$

If θ is not an identity automorphism, then under the conventional addition and multiplication of the polynomials given by $(gx^i)(g'x^j) = g\theta^i(g')x^{i+j}$, the set $\mathbb{F}_q[x:\theta]$ constitutes a noncommutative ring and known as skew polynomial ring over \mathbb{F}_q . we refer [17] for additional details.

As the set of all automorphism on \mathbb{F}_q constitutes a group under the composition of maps, let $\theta \in \operatorname{Aut}(\mathbb{F}_q)$ be an automorphism. The smallest positive integer t satisfying $\theta^t(g) = g$ for all $g \in \mathbb{F}_q$, is called the order of θ and denoted by $\operatorname{ord}(\theta)$.

Proposition 2.2. [24] In $\mathbb{F}_q[x:\theta]$, the following are equivalent :

- 1. $x^r 1 \in Z(\mathbb{F}_q[x:\theta])$, where $Z(\mathbb{F}_q[x:\theta])$ denote the center of the ring $\mathbb{F}_q[x:\theta]$.
- 2. $\langle x^r 1 \rangle$ forms a two-sided ideal.
- 3. $\operatorname{ord}(\theta)$ divides r.

We now extend our above discussion over a finite ring. We begin with following definition to expanding our study.

Definition 2.3. Suppose a finite commutative ring R and the set of all automorphism on R denoted by $\operatorname{Aut}(R)$. Assume $\Theta \in \operatorname{Aut}(R)$, then

$$R[x:\Theta] = \{b_0 + b_1 x + b_2 x^2 + \dots + b_n x^n \mid b_i \in R, \ \Theta \in \text{Aut}(R)\}.$$

If Θ is not an identity automorphism then under the conventional addition and multiplication of the polynomials given by $(bx^i)(b'x^j) = b\Theta^i(b')x^{i+j}$, the set $R[x:\Theta]$ constitutes a noncommutative ring over R.

Definition 2.4. A non-empty R-submodule \mathfrak{C}_s of \mathbb{R}^s is referred to a linear code of length S over \mathbb{R} .

In this context, if $\operatorname{ord}(\Theta)$ divides s, the quotient space $\frac{R[x:\Theta]}{\langle x^s-1 \rangle}$ forms a ring and if $\operatorname{ord}(\Theta)$ does not divide s, then $\frac{R[x:\Theta]}{\langle x^s-1 \rangle}$ does not form a ring. However, it still behaves as a left $R[x:\Theta]$ -module in the later case. The scalar multiplication in this context can be given as follows: suppose $s(x), t(x) \in R[x:\Theta]$,

$$s(x)(t(x) + \langle x^s - 1 \rangle) = s(x)t(x) + \langle x^s - 1 \rangle.$$

Definition 2.5. Suppose $\Theta \in \operatorname{Aut}(R)$ and \mathfrak{C}_s is a linear code of length s. If for any $\mathbf{b} = (b_0, b_1, \dots, b_{s-1}) \in \mathfrak{C}_s$, implies $(\Theta(b_{s-1}), \Theta(b_0), \dots, \Theta(b_{s-2})) \in \mathfrak{C}_s$, then it refers to a Θ -cyclic code over R.

It is possible to identify any codeword $\mathbf{b} = (b_0, b_1, \dots, b_{s-1}) \in \mathfrak{C}_s$ as a polynomial $(b_0 + b_1 x + b_2 x^2 + \dots + b_{s-1} x^{s-1}) \in \frac{R[x:\Theta]}{\langle x^s-1 \rangle}$. Consequently, any Θ -cyclic code of length s It is called a Θ -cyclic code over R considered as a left $R[x:\Theta]$ -submodule of $\frac{R[x:\Theta]}{\langle x^s-1 \rangle}$.

Throughout this work, we examine the ring $R = \mathbb{F}_q + u\mathbb{F}_q + v\mathbb{F}_q + uv\mathbb{F}_q$ with $u^2 = u$, $v^2 = v$, and uv = vu, where $q = p^m$ for an odd prime p and m being a positive integer. Recall that, $\theta \in \operatorname{Aut}(\mathbb{F}_q)$ is given by $\theta(g) = g^{p^i}$, where $1 \leq i \leq m$. Let us define a map Θ on R in the following way:

$$\Theta(a_0 + ua_1 + va_2 + uva_3) = \theta(a_0) + u\theta(a_1) + v\theta(a_2) + uv\theta(a_3)$$
$$= a_0^{p^i} + ua_1^{p^i} + va_2^{p^i} + uva_3^{p^i},$$

where $a_0, a_1, a_2, a_3 \in \mathbb{F}_q$. By the definition of θ , it can be seen that $\Theta \in \operatorname{Aut}(R)$.

Suppose

$$\kappa_1 = 1 - u - v + uv, \ \kappa_2 = u - uv,$$

$$\kappa_3 = v - uv, \ \kappa_4 = uv,$$

then

1. $\kappa_1 + \kappa_2 + \kappa_3 + \kappa_4 = 1$. 2. $\kappa_i^2 = \kappa_i$. 3. $\kappa_i \cdot \kappa_j = 0$ for $i \neq j$.

It is clear that $\{\kappa_1, \kappa_2, \kappa_3, \kappa_4\}$ forms an idempotent set of R. By using this idempotent set, we can express $R = \kappa_1 R \oplus \kappa_2 R \oplus \kappa_3 R \oplus \kappa_4 R$. If $b \in R$, then

$$b = a_0 + ua_1 + va_2 + uva_3 = \kappa_1 n_1 + \kappa_2 n_2 + \kappa_3 n_3 + \kappa_4 n_4,$$

where $n_1 = a_0$, $n_2 = a_0 + a_1$, $n_3 = a_0 + a_2$ and $n_4 = a_0 + a_1 + a_2 + a_3$.

Suppose \mathfrak{C}_s is a linear codes of length s over R and consider the sets

$$egin{aligned} \mathfrak{C}_{s,1} &= \{m{n}_1 \in \mathbb{F}_q^s, \kappa_1 m{n}_1 + \kappa_2 m{n}_2 + \kappa_3 m{n}_3 + \kappa_4 m{n}_4 \in \mathfrak{C}_s: \ m{n}_2, m{n}_3, m{n}_4 \in \mathbb{F}_q^s\}, \ \mathfrak{C}_{s,2} &= \{m{n}_2 \in \mathbb{F}_q^s, \kappa_1 m{n}_1 + \kappa_2 m{n}_2 + \kappa_3 m{n}_3 + \kappa_4 m{n}_4 \in \mathfrak{C}_s: \ m{n}_1, m{n}_3, m{n}_4 \in \mathbb{F}_q^s\}, \ \mathfrak{C}_{s,3} &= \{m{n}_3 \in \mathbb{F}_q^s, \kappa_1 m{n}_1 + \kappa_2 m{n}_2 + \kappa_3 m{n}_3 + \kappa_4 m{k}_4 \in \mathfrak{C}_s: \ m{n}_1, m{n}_2, m{n}_4 \in \mathbb{F}_q^s\}, \ \mathfrak{C}_{s,4} &= \{m{n}_4 \in \mathbb{F}_q^s, \kappa_1 m{n}_1 + \kappa_2 m{n}_2 + \kappa_3 m{n}_3 + \kappa_4 m{n}_4 \in \mathfrak{C}_s: \ m{n}_1, m{n}_2, m{n}_4 \in \mathbb{F}_q^s\}, \end{aligned}$$

By the definition of $\mathfrak{C}_{s,i}$, for i = 1, 2, 3, 4, it is clear that $\mathfrak{C}_{s,i}$ are linear codes of length s over \mathbb{F}_q . Moreover,

$$\mathfrak{C}_s = \kappa_1 \mathfrak{C}_{s,1} \oplus \kappa_2 \mathfrak{C}_{s,2} \oplus \kappa_3 \mathfrak{C}_{s,3} \oplus \kappa_4 \mathfrak{C}_{s,4}.$$

Now, let us introduce a Gray map $\phi_1 : R \to \mathbb{F}_q^4$ given by

$$\phi(b) = \phi_1(\kappa_1 n_1 + \kappa_2 n_2 + \kappa_3 n_3 + \kappa_4 n_4) = (n_1, n_2, n_3, n_4)M_2$$

where $M \in \text{GL}(4, \mathbb{F}_q)$ satisfying $MM^T = \lambda I_4, \lambda \neq 0 \in \mathbb{F}_q$, I_4 is identity matrix of order 4 and M^T denotes the transpose of matrix M. For convenience, we write $(n_1, n_2, n_3, n_4)M = bM$.

It is straightforward to confirm that ϕ is \mathbb{F}_q -linear. A natural extension of this Gray map is Φ_1 from \mathbb{R}^s to \mathbb{F}_q^{4s} , which is given by

$$\Phi_1(b_0, b_1, \dots, b_{s-1}) = (n_{1,0}, n_{2,0}, n_{3,0}, n_{4,0}, n_{1,1}, n_{2,1}, n_{3,1}, n_{4,1}, \dots, n_{1,s-1}, n_{2,s-1}, n_{3,s-1}, n_{4,s-1})M$$

= $(b_0M, b_1M, \dots, b_{s-1}M),$

where $b_i = \kappa_1 n_{1,i} + \kappa_2 n_{2,i} + \kappa_3 n_{3,i} + \kappa_4 n_{4,i} \in \mathbb{R}$, for i = 0, 1, 2..., s - 1.

Let w_L and w_H denote the Lee and Hamming Weight, respectively. Then the Lee weight of any arbitrary element $b = \kappa_1 n_1 + \kappa_2 n_2 + \kappa_3 n_3 + \kappa_4 n_4 \in \mathbb{R}$ is defined as

$$w_L(b) = w_L(\kappa_1 n_1 + \kappa_2 n_2 + \kappa_3 n_3 + \kappa_4 n_4)$$

= $w_H(\phi_1(\kappa_1 n_1 + \kappa_2 n_2 + \kappa_3 n_3 + \kappa_4 n_4))$
= $w_H(n_1 M) + w_H(n_2 M) + w_H(n_3 M) + w_H(n_4 M).$

Furthermore, if **b** and **b'** are any two distinct elements in \mathbb{R}^s , then Lee distance is defined as

$$d_L(\boldsymbol{b}, \boldsymbol{b}') = w_L(\boldsymbol{b} - \boldsymbol{b}') = w_H(\Phi_1(\boldsymbol{b} - \boldsymbol{b}')) = d_H(\Phi_1(\boldsymbol{b}), \Phi_1(\boldsymbol{b}')).$$

The Lee weight of any linear code \mathfrak{C}_s over R is the least Lee weight codeword among all the possible pair of distinct codewords in \mathfrak{C}_s .

We define

$$\mathfrak{R}_{r,s} = \frac{\mathbb{F}_q[x:\theta]}{\langle x^r - 1 \rangle} \times \frac{R[x:\Theta]}{\langle x^s - 1 \rangle} \quad \text{and} \quad \mathbb{F}_q R = \{(g,b), g \in \mathbb{F}_q, \ b \in R\}.$$

The set $\mathbb{F}_q R$ constitutes a ring when addition and multiplication are operated with componentwise. Now, a homomorphism $\eta : R \to \mathbb{F}_q$ is given by

$$\eta(a_0 + ua_1 + va_2 + uva_3) = a_0.$$

where $a + ub + vc + uvw \in R$. Moreover, R acts on $\mathbb{F}_q R$ by multiplication \star as

$$r \star (g, b) = (\eta(r)g, rb).$$

We can extend this scalar multiplication \star on $\mathbb{F}_q^r \times \mathbb{R}^s$ as

 $r \star d = (\eta(r)g_0, \eta(r)g_1, \dots, \eta(r)g_{r-1}, rb_0, rb_1, \dots, rb_{s-1}),$

where $\boldsymbol{d} = (g_0, g_1, \dots, g_{r-1}, b_0, b_1, \dots, b_{s-1}) \in \mathbb{F}_q^r \times R^s$. Indeed, the multiplication \star makes $\mathbb{F}_q^r \times R^s$ an *R*-module.

Definition 2.6. Suppose \mathfrak{D} is a non empty R submodule of $\mathbb{F}_q^r \times R^s$, then it refers to a linear code of block length (r, s) over $\mathbb{F}_q R$.

Let $(g_0, g_1, \ldots, g_{r-1}, b_0, b_1, \ldots, b_{s-1}) \in \mathbb{F}_q^r \times R^s$ and $\rho_{(\theta,\Theta)}$ be a map from $\mathbb{F}_q^r \times R^s$ to $\mathbb{F}_q^r \times R^s$ given by

 $\rho_{(\theta,\Theta)}(g_0, g_1, \dots, g_{r-1}, b_0, b_1, \dots, b_{s-1}) = (\theta(g_{r-1}), \theta(g_0), \dots, \theta(g_{r-2}), \Theta(b_{s-1}), \Theta(b_0), \dots, \Theta(b_{s-2})).$

We refer to this map as the (θ, Θ) -cyclic shift on $\mathbb{F}_q^r \times \mathbb{R}^s$.

Definition 2.7. A linear code \mathfrak{D} of block length (r, s) over $\mathbb{F}_q R$, which is closed under the map $\rho_{(\theta,\Theta)}$ is refers to a (θ,Θ) -cyclic code, i.e., if any $\mathbf{d} = (g_0, g_1, ..., g_{r-1}, b_0, b_1, ..., b_{s-1}) \in \mathfrak{D}$, implies

 $\rho_{\theta,\Theta}(d) = (\theta(g_{r-1}), \theta(g_0), ..., \theta(g_{r-2}), \Theta(b_{s-1}), \Theta(b_0), ..., \Theta(b_{s-2})) \in \mathfrak{D}.$

Now, We shall give a Gray map over $\mathbb{F}_q R$, along with providing a natural extension of this map from $\mathbb{F}_q^r \times R^s \to \mathbb{F}_q^{r+4s}$.

Suppose $(g,b) = (g,\kappa_1n_1 + \kappa_2n_2 + \kappa_3n_3 + \kappa_4n_4) \in \mathbb{F}_q R$. Consider the map ϕ from $\mathbb{F}_q R$ to \mathbb{F}_q^5 given by

$$\phi(g,b) = (g,\eta(b)) = (g,(n_1,n_2,n_3,n_4)M).$$

This map is \mathbb{F}_q -linear. Now, a natural extension map Φ from $\mathbb{F}_q^r \times \mathbb{R}^s$ to \mathbb{F}_q^{r+4s} is given by

$$(g_0, g_1, \dots, g_{r-1}, b_0, b_1, \dots, b_{s-1}) \rightarrow (g_0, g_1, \dots, g_{r-1}, (n_{1,0}, n_{2,0}, n_{3,0}, n_{4,0}, n_{1,1}, n_{2,1}, n_{3,1}, n_{4,1}, \dots, n_{1,s-1}, n_{2,s-1}, n_{3,s-1}, n_{4,s-1})M) = (g_0, g_1, \dots, g_{r-1}, b_0M, b_1M, \dots, b_{s-1}M),$$

where $b_i = \kappa_1 n_{1,i} + \kappa_2 n_{2,i} + \kappa_3 n_{3,i} + \kappa_4 n_{4,i}$, for $i = 0, 1, \dots, s - 1$, $(g_0, g_1, \dots, g_{r-1}) \in \mathbb{F}_q^r$ and $(b_0, b_1, \dots, b_{s-1}) \in \mathbb{R}^s$.

Let $(\boldsymbol{g}, \boldsymbol{b}) \in \mathbb{F}_q^r \times R^s$. Then $w_L(\boldsymbol{g}, \boldsymbol{b}) = w_H(\boldsymbol{g}) + w_L(\boldsymbol{b})$. Let $\boldsymbol{d}_1, \boldsymbol{d}_2 \in \mathbb{F}_q^r \times R^s$ be any two codewords in \mathfrak{D} . Then the Lee distance is given by

$$d_L(d_1, d_2) = w_L(d_1 - d_2) = w_H(\Phi(d_1 - d_2)) = d_H(\Phi(d_1), \Phi(d_2)).$$

$$MM^T = I_{4 \times 4}$$

Proposition 2.8. [16] Let the Gray map Φ .

- 1. Then Φ is a distance-preserving and \mathbb{F}_q -linear.
- 2. $\Phi(\mathfrak{D})$ is a $[r+4s, k, d_H]$ linear code, where $d_L = d_H$ over \mathbb{F}_q , for a linear code \mathfrak{D} of block length (r, s) over $\mathbb{F}_q R$ with $|\mathfrak{D}| = q^k$.

Let $d = (g_0, g_1, \ldots, g_{r-1}, b_0, b_1, \ldots, b_{s-1}) \in \mathbb{F}_q^r \times R^s$. Every element of $\mathbb{F}_q^r \times R^s$ can be identified by a pair of polynomials in $\mathfrak{R}_{r,s}$ as follows:

$$(g_0, g_1, \dots, g_{r-1}, b_0, b_1, \dots, b_{s-1}) \to (g_0 + g_1 x + \dots + g_{r-1} x^{r-1}, b_0 + b_1 x + \dots + b_{s-1} x^{s-1})$$

= $(g(x), b(x)),$

where $g(x) \in \frac{\mathbb{F}_q[x:\Theta]}{\langle x^r - 1 \rangle}$ and $b(x) \in \frac{R[x:\Theta]}{\langle x^s - 1 \rangle}$. This map is called the polynomial identification of elements from $\mathbb{F}_q^r \times \mathbb{R}^s$ to $\mathfrak{R}_{r,s}$. This shows that there is a bijection between $\mathbb{F}_q^r \times \mathbb{R}^s$ and $\mathfrak{R}_{r,s}$.

Let $(g(x), b(x)) \in \mathfrak{R}_{r,s}$ and $r(x) = r_0 + r_1 x + \cdots + r_{n-1} x^{n-1} \in R[x : \Theta]$. The scalar multiplication on $\mathfrak{R}_{r,s}$ is given by

$$r(x) \star (g(x), b(x)) = (\eta(r(x))g(x), r(x)b(x)),$$

where $\eta(r(x)) = \eta(r_0) + \eta(r_1)x + \cdots + \eta(r_{n-1})x^{n-1}$. Moreover, when we consider the scalar multiplication " \star " the ring $\Re_{r,s}$ forms a left $R[x:\Theta]$ -module.

Proposition 2.9. Assume \mathfrak{D} is a linear code of block length (r, s) over $\mathbb{F}_q R$. If \mathfrak{D} identifies as a polynomial such that it is a left $R[x : \Theta]$ -submodule of $\mathfrak{R}_{r,s}$, then \mathfrak{D} is called (θ, Θ) -cyclic code over $\mathbb{F}_q R$.

Proof. Consider any arbitrary element $d = (g_0, g_1, \ldots, g_{r-1}, b_0, b_1, \ldots, b_{s-1}) \in \mathfrak{D} \subseteq \mathbb{F}_q^r \times \mathbb{R}^s$. So d can be identified by polynomial $(g_0 + g_1x + \cdots + g_{r-1}x^{r-1}, b_0 + b_1x + \cdots + b_{s-1}x^{s-1}) \in \mathfrak{R}_{r,s}$. Now, we have

$$x \star d(x) = (xg_0 + xg_1x + \dots + xg_{r-1}x^{r-1}, xb_0 + xb_1x + \dots + xb_{s-1}x^{s-1})$$

= $(\theta(g_{r-1}) + \theta(g_0)x + \dots + \theta(g_{r-2})x^{r-1}, \Theta(b_{s-1}) + \Theta(b_0)x + \dots + \Theta(b_{s-2})x^{s-1}) \in \mathfrak{R}_{r,s}$

Clearly, the corresponding codeword of $x \star d(x)$ is $(\theta(g_{r-1}), \theta(g_0), ..., \theta(g_{r-2}), \Theta(b_{s-1}), \Theta(b_0), ..., \Theta(b_{s-2}))$ and it is a (θ, Θ) -cyclic shift of codeword $d \in \mathfrak{D}$. Hence, by the linearity of \mathfrak{D} , it can be easily seen that the polynomial representation of \mathfrak{D} is a $R[x : \Theta]$ -submodule of $\mathfrak{R}_{r,s}$.

Let $d = (g_0, g_1, \dots, g_{r-1}, b_0, b_1, \dots, b_{s-1})$ and $d' = (g'_0, g'_1, \dots, g'_{r-1}, b'_0, b'_1, \dots, b'_{s-1}) \in \mathbb{F}_q^r \times \mathbb{R}^s$. The inner product is defined as

$$\boldsymbol{d} \cdot \boldsymbol{d}' = \kappa_1 \sum_{i=0}^{r-1} g_i g_i' + \sum_{j=0}^{s-1} b_j b_j' \in R.$$

The dual of a linear code \mathfrak{D} is denoted by \mathfrak{D}^{\perp} , and defined as

$$\mathfrak{D}^{\perp} = \{ \boldsymbol{d}' \in \mathbb{F}_q^r \times R^s \mid \boldsymbol{d} \cdot \boldsymbol{d}' = 0 \ \forall \ \boldsymbol{d} \in \mathfrak{D} \}.$$

Moreover, if \mathfrak{D} is equal to \mathfrak{D}^{\perp} , then \mathfrak{D} is self-dual and if $\mathfrak{D} \subseteq \mathfrak{D}^{\perp}$, then \mathfrak{D} is called self-orthogonal. Now we establish the relation between \mathfrak{D} and its dual \mathfrak{D}^{\perp} .

Proposition 2.10. Suppose $\operatorname{ord}(\theta) \mid r$, $\operatorname{ord}(\Theta) \mid s$ and \mathfrak{D} is a (θ, Θ) -cyclic code of block length (r, s) over $\mathbb{F}_q R$, then \mathfrak{D}^{\perp} is also a (θ, Θ) -cyclic code of the same block length over $\mathbb{F}_q R$.

Proof. Suppose $d = (g_0, g_1, \ldots, g_{r-1}, b_0, b_1, \ldots, b_{s-1}) \in \mathfrak{D}$ and since \mathfrak{D} is a (θ, Θ) -cyclic code, so that $(\theta(g_{r-1}), \theta(g_0), \ldots, \theta(g_{r-2}), \Theta(b_{s-1}), \Theta(b_0), \ldots, \Theta(b_{s-2})) \in \mathfrak{D}$. Now, consider $d' = (g'_0, g'_1, \ldots, g'_{r-1}, b'_0, b'_1, \ldots, b'_{s-1}) \in \mathfrak{D}^{\perp}$. Then $d \cdot d' = 0$.

In order to check that \mathfrak{D}^{\perp} is also a (θ, Θ) -cyclic code of block length (r, s) over $\mathbb{F}_q R$, it is enough to prove that

$$\rho_{(\theta,\Theta)}(\mathbf{d}') = (\theta(g'_{r-1}), \dots, \theta(g'_{r-2}), \Theta(b'_{s-1}), \dots, \Theta(b'_{s-2})) \in \mathfrak{D}^{\perp}.$$

In other words, we need to have

$$\boldsymbol{d} \cdot \rho_{(\boldsymbol{\theta},\Theta)}(\boldsymbol{d}') = 0,$$

i.e.,

 $\kappa_1 g_0 \theta(g_{r-1}) + \kappa_1 g_1 \theta(g_0) + \dots + \kappa_1 g_{r-1} \theta(g_{r-2}) + b_0 \Theta(b'_{s-1}) + b_1 \Theta(b'_0) + \dots + b_{s-1} \Theta(b'_{s-2}) = 0.$ As we have \mathfrak{D} is a (θ, Θ) -cyclic code. So, we have

$$\rho_{(\theta,\Theta)}(\boldsymbol{d}) \in \mathfrak{D} \text{ and } \rho_{(\theta,\Theta)}(\boldsymbol{d}) \cdot \boldsymbol{d}' = 0.$$

Let $l = \operatorname{lcm}(r, s)$. Then

$$\rho_{(\theta,\Theta)}^{l-1}(\mathbf{d}) = (\theta^{l-1}(g_1), \dots, \theta^{l-1}(g_0), \Theta^{l-1}(b_1), \dots, \Theta^{l-1}(b_0)) \in \mathfrak{D}.$$

Now, we get

$$0 = \rho_{(\theta,\Theta)}^{l-1}(\mathbf{d}) \cdot \mathbf{d}'$$

= $\kappa_1(\theta^{l-1}(g_1)g_0' + \theta^{l-1}(g_2)g_1' + \dots + \theta^{l-1}(g_{r-1})g_{r-2} + \theta^{l-1}(g_0)g_{r-1}')$
+ $(\Theta^{l-1}(b_1)b_0 + \Theta^{l-1}(b_2)b_1' + \dots + \Theta^{l-1}(b_{s-1})b_{s-2}' + \Theta^{l-1}(b_0)b_{s-1}).$

Since $\operatorname{ord}(\theta) | r$ and $\operatorname{ord}(\Theta) | s$, we have θ^l and Θ^l are identity automorphisms and $\theta = \Theta$ over \mathbb{F}_q . Now, applying Θ both side in the above equation

$$\kappa_1(g_1\theta(g'_0)+g_2\theta(g'_1)+\dots+g_{r-2}\theta(g'_{r-1})+g_{r-1}\theta(g'_0))+(b_1\Theta(b'_0)+\dots+b_{s-2}\Theta(b'_{s-1})+b_{s-1}\Theta(b'_0))=0.$$

Hence, $\boldsymbol{d} \cdot \rho_{(\theta,\Theta)}(\boldsymbol{d}')=0.$ Thus, we infer that $\rho_{(\theta,\Theta)}(\boldsymbol{d}) \in \mathfrak{D}^{\perp}.$ This completes our proof. \Box

3 Algebraic structure of (θ, Θ) -cyclic codes over $\mathbb{F}_{q}R$

Let us begin by providing the decomposed structure of linear codes \mathfrak{D} of block length (r, s) over $\mathbb{F}_q R$. Let \mathcal{D}_i be sets, for i = 1, 2, 3, 4. We define

$$\mathcal{D}_1 \oplus \mathcal{D}_2 \oplus \mathcal{D}_3 \oplus \mathcal{D}_4 = \{ d_1 + d_2 + d_3 + d_4 \mid d_i \in \mathcal{D}_i \text{ for } i = 1, 2, 3, 4 \}.$$

Since, $\sum_{i=1}^{4} \kappa_i = 1$, so any $\boldsymbol{d} = (\boldsymbol{g}, \boldsymbol{b}) \in \mathbb{F}_q^r \times R^s$ can be expressed as $\boldsymbol{d} = \sum_{i=1}^{4} \kappa_i(\boldsymbol{g}, \boldsymbol{b}) \in \mathbb{F}_q^r \times R^s$, where $\boldsymbol{g} \in \mathbb{F}_q^r$ and $\boldsymbol{b} = \kappa_1 \boldsymbol{n}_1 + \kappa_2 \boldsymbol{n}_2 + \kappa_3 \boldsymbol{n}_3 + \kappa_4 \boldsymbol{n}_4 \in R^s$. We define

 $egin{aligned} \mathfrak{D}_1 &= \{(oldsymbol{g},oldsymbol{n}_1) \in \mathbb{F}_q^r imes \mathbb{F}_q^s \mid oldsymbol{g} \in \mathbb{F}_q^r, \ oldsymbol{n}_1 \in \mathfrak{C}_{s,1}\}, \ \mathfrak{D}_2 &= \{(oldsymbol{g},oldsymbol{n}_2) \in \mathbb{F}_q^r imes \mathbb{F}_q^s \mid oldsymbol{g} \in \mathbb{F}_q^r, \ oldsymbol{n}_2 \in \mathfrak{C}_{s,2}\}, \ \mathfrak{D}_3 &= \{(oldsymbol{g},oldsymbol{n}_3) \in \mathbb{F}_q^r imes \mathbb{F}_q^s \mid oldsymbol{g} \in \mathbb{F}_q^r, \ oldsymbol{n}_3 \in \mathfrak{C}_{s,3}\}, \ \mathfrak{D}_4 &= \{(oldsymbol{g},oldsymbol{n}_4) \in \mathbb{F}_q^r imes \mathbb{F}_q^s \mid oldsymbol{g} \in \mathbb{F}_q^r, \ oldsymbol{n}_4 \in \mathfrak{C}_{s,4}\}. \end{aligned}$

It is evident that \mathfrak{D}_i , for i = 1, 2, 3, 4 is a linear code of block length (r, s) over \mathbb{F}_q . Consequently, any linear code \mathfrak{D} of block length (r, s) over $\mathbb{F}_q R$ has a unique expression as

$$\mathfrak{D} = \kappa_1 \mathfrak{D}_1 \oplus \kappa_2 \mathfrak{D}_2 \oplus \kappa_3 \mathfrak{D}_3 \oplus \kappa_4 \mathfrak{D}_4.$$

Let c(x), $a(x) \neq 0$ be two polynomials in $\mathbb{F}_q[x : \theta]$. Then a(x) is called the right divisor of c(x), if c(x) = d(x)a(x), where $d(x) \in \mathbb{F}_q[x : \theta]$ and denoted by $a(x) \mid_r c(x)$. Similarly, we can define left divisor of c(x).

Definition 3.1. The polynomial c(x) is central polynomial in $\mathbb{F}_q[x:\theta]$, if xc(x) = c(x)x.

Proposition 3.2. [24, Proposition-2.3] Suppose c(x), $d(x) \in \mathbb{F}_q[x : \theta]$ are monic central polynomials. Then d(x)c(x) = c(x)d(x).

Lemma 3.3. [17, Theorem-II.11] Suppose c(x) and $a(x)(monic) \in \mathbb{F}_q[x:\theta]$. Then $c(x) = q_1(x)a(x)+q_2(x)$, where $q_2 = 0$ or $\deg(q_2(x)) < \deg(a(x))$, for some $q_1(x)$ and $q_2(x)$ in $\mathbb{F}_q[x:\theta]$.

In the context of $R[x:\Theta]$, we now introduce right division algorithm.

Proposition 3.4. Let s(x) and $t(x) \in R[x : \Theta]$ be two polynomials such that t(x) is monic. Then $s(x) = q'_1(x)t(x) + q'_2(x)$, where $q'_2(x) = 0$ or $\deg(q'_2(x)) < \deg(t(x))$, for some $q'_1(x)$, $q'_2(x) \in R[x : \Theta]$.

Proof. Let $s(x) = \sum_{i=0}^{m} s_i x^i$ and $t(x) = \sum_{j=0}^{k-1} t_j x^j + x^k$. Then $\deg(s(x) - s_m x^{m-k} t(x))$ less than $\deg(s(x))$. By repeatedly applying this procedure with successive polynomials, we can get the desired $q'_1(x)$ and $q'_2(x)$. Now, we prove the uniqueness of $q'_1(x)$ and $q'_2(x)$. Suppose

$$s(x) = q'_1(x)t(x) + q'_2(x) = q''_1(x)t(x) + q''_2(x),$$

then

$$(q_1'(x) - q_1''(x))t(x) = q_2''(x) - q_2'(x).$$

Here $q'_1(x) - q''_1(x) \neq 0$, implying that the right polynomial must have a degree equal to or greater than the degree of t(x) and at the same time, the degree of right polynomial not be more than deg(t(x)) - 1. This leads us to a contradiction. Therefore, $q'_1(x)$ and $q'_2(x)$ are unique.

Now, we present the definition of least common multiple (lcm) and greatest common divisor (gcd), which are going to be used frequently throughout this paper.

Definition 3.5. Suppose polynomials a(x), c(x), $d(x) \in \mathbb{F}_q[x : \theta]$ with $d(x) \mid_r a(x)$ and $d(x) \mid_r c(x)$. If d'(x) is another polynomial with the same property, then $d'(x) \mid_r d(x)$. Consequently, d(x) is said to be right gcd and denoted by $\gcd_r(a(x), c(x))$. Similarly, we can define the left gcd (\gcd_l) .

Definition 3.6. Suppose polynomials $m(x), a(x), c(x) \in \mathbb{F}_q[x : \Theta]$ with $a(x) \mid_l m(x)$ and $c(x) \mid_l m(x)$ and $c(x) \mid_l m(x)$ and if any other polynomial $m'(x) \in \mathbb{F}_q[x : \theta]$ with same property implies $m(x) \mid_l m'(x)$. Then m(x) is said to be left lcm and denoted by $\operatorname{lcm}_l(a(x), c(x))$.

Lemma 3.7. [26, pp.486] Let d(x) and $c(x) \in \mathbb{F}_q[x:\theta]$. Then $\deg(\operatorname{lcm}_l(d(x), c(x)) = \deg(d(x)) + \deg(c(x)) - \deg(\gcd_r(d(x), c(x)))$.

Lemma 3.8. [3, Lemma-2.7] Suppose $d_1(x), d_2(x), d_3(x)$ and $d(x) \in \mathbb{F}_q[x : \theta]$, where d(x) is a central polynomial with

$$d_1(x)d_2(x) \equiv 0 \pmod{d(x)}$$

and

$$d_1(x)d_3(x) \equiv 0 \pmod{d(x)}.$$

Then

$$d_1(x)\operatorname{gcd}_l(d_2(x), d_3(x)) \equiv 0 \pmod{d(x)}$$

In the upcoming two propositions, we articulate the findings discussed by T. Yao et al. [33], wherein they characterized Θ -cyclic codes over R.

Proposition 3.9. [33, Theorem-4.3] Let $\mathfrak{C}_s = \kappa_1 \mathfrak{C}_{s,1} \oplus \kappa_2 \mathfrak{C}_{s,2} \oplus \kappa_3 \mathfrak{C}_{s,3} \oplus \kappa_4 \mathfrak{C}_{s,4}$ be a linear code of length s over R. Then \mathfrak{C}_s is Θ -cyclic code over R if and only if $\mathfrak{C}_{s,i}$ are θ -cyclic codes over \mathbb{F}_q .

Proposition 3.10. [33, Theorem-4.5] Let $\mathfrak{C}_s = \kappa_1 \mathfrak{C}_{s,1} \oplus \kappa_2 \mathfrak{C}_{s,2} \oplus \kappa_3 \mathfrak{C}_{s,3} \oplus \kappa_4 \mathfrak{C}_{s,4}$ and $\mathfrak{C}_{s,i} = \langle t_i(x) \rangle$ with $t_i(x) \mid_r x^s - 1$ in $R[x:\Theta]$ for i = 1, 2, 3, 4. Then $\mathfrak{C}_s = \langle t(x) \rangle$, where $t(x) = \kappa_1 t_1(x) + \kappa_2 t_2(x) + \kappa_3 t_3(x) + \kappa_4 t_4(x)$ with $t(x) \mid_r x^s - 1$ in $R[x:\Theta]$. Additionally, $|\mathfrak{C}_s| = q^{4s - \sum_{i=1}^4 \deg(t_i(x))}$.

In next result, we scrutinize the decompose phenomena of (θ, Θ) -cyclic code over $\mathbb{F}_q R$.

Proposition 3.11. Let $\mathfrak{D} = \kappa_1 \mathfrak{D}_1 \oplus \kappa_2 \mathfrak{D}_2 \oplus \kappa_3 \mathfrak{D}_3 \oplus \kappa_4 \mathfrak{D}_4$ be a linear code of block length (r, s) over $\mathbb{F}_q R$. Then \mathfrak{D} is (θ, Θ) -cyclic code if and only if each \mathfrak{D}_i is a θ -cyclic code of block length (r, s) over \mathbb{F}_q for i = 1, 2, 3, 4.

Proof. Suppose $d = (g_0, g_1, \ldots, g_{r-1}, b_0, b_1, \ldots, b_{s-1}) \in \mathfrak{D}$, where $b_i = \kappa_1 n_{1,i} + \kappa_2 n_{2,i} + \kappa_3 n_{3,i} + \kappa_4 n_{4,i}$ for $i = 0, 1, \ldots, s - 1$. Then we have

 $\begin{aligned} & (g_0, g_1, \dots, g_{r-1}, n_{1,0}, n_{1,1}, \dots, n_{1,s-1}) \in \mathfrak{D}_1 \\ & (g_0, g_1, \dots, g_{r-1}, n_{2,0}, n_{2,1}, \dots, n_{2,s-1}) \in \mathfrak{D}_2 \\ & (g_0, g_1, \dots, g_{r-1}, n_{3,0}, n_{3,1}, \dots, n_{3,s-1}) \in \mathfrak{D}_3 \\ & (g_0, g_1, \dots, g_{r-1}, n_{4,0}, n_{4,1}, \dots, n_{4,s-1}) \in \mathfrak{D}_4. \end{aligned}$

Now, consider if \mathfrak{D} is a (θ, Θ) -cyclic code, then

$$\rho_{(\theta,\Theta)}(\boldsymbol{d}) = (\theta(g_{r-1}), \theta(g_0), \dots, \theta(g_{r-1}), \Theta(b_{s-1}), \Theta(b_0), \Theta(b_1), \dots, \Theta(b_{s-2})) \in \mathfrak{D}.$$

Therefore,

$$\begin{aligned} & (\theta(g_{r-1}), \theta(g_0), \dots, \theta(g_{r-2}), \theta(n_{1,s-1}), \theta(n_{1,0}), \dots, \theta(n_{1,s-2})) \in \mathfrak{D}_1 \\ & (\theta(g_{r-1}), \theta(g_0), \dots, \theta(g_{r-2}), \theta(n_{2,s-1}), \theta(n_{2,0}), \dots, \theta(n_{2,s-2})) \in \mathfrak{D}_2 \\ & (\theta(g_{r-1}), \theta(g_0), \dots, \theta(g_{r-2}), \theta(n_{3,s-1}), \theta(n_{3,0}), \dots, \theta(n_{3,s-2})) \in \mathfrak{D}_3 \\ & (\theta(g_{r-1}), \theta(g_0), \dots, \theta(g_{r-2}), \theta(n_{4,s-1}), \theta(n_{4,0}), \dots, \theta(n_{4,s-2})) \in \mathfrak{D}_4 \end{aligned}$$

Consequently, it follows that \mathfrak{D}_i for i = 1, 2, 3, 4 is θ -cyclic code of block length (r, s) over \mathbb{F}_q .

The converse part can be proved by using similar arguments.

Now, we determine the generator polynomials for (θ, Θ) -cyclic codes of block length (r, s) over $\mathbb{F}_q R$. Additionally, we shall investigate the size of this family of codes and examine the minimum generating set.

Theorem 3.12. Let \mathfrak{D} be a (θ, Θ) -cyclic code of block length (r, s) over $\mathbb{F}_q R$. Then

$$\mathfrak{D} = \langle (\ell(x), 0), (s(x), t(x)) \rangle,$$

where the polynomials $\ell(x), s(x) \in \mathbb{F}_q[x:\theta]$ with $\ell(x) \mid_r x^r - 1$ and $t(x) = \kappa_1 t_1(x) + \kappa_2 t_2(x) + \kappa_3 t_3(x) + \kappa_4 t_4(x), \quad t_i(x) \in \mathbb{F}_q[x:\theta]$ with $t_i(x) \mid_r (x^s - 1)$ in $\mathbb{F}_q[x:\theta]$, for i = 1, 2, 3, 4 and $t(x) \mid_r x^s - 1$ in $R[x:\Theta]$.

Proof. As both \mathfrak{D} and $\frac{R[x:\Theta]}{\langle x^s-1\rangle}$ are left $R[x:\Theta]$ -submodule of $\mathfrak{R}_{r,s}$. We define a left $R[x:\Theta]$ -module homomorphism π as follows:

$$\pi:\mathfrak{D}\to \frac{R[x:\Theta]}{\langle x^s-1\rangle}$$

and given by

$$\pi(d_1(x), d_2(x)) = d_2(x).$$

Clearly, $\pi(\mathfrak{D})$ is a left-submodule of $\frac{R[x:\Theta]}{\langle x^s-1 \rangle}$. Then by Proposition 3.10, we get that $\pi(\mathfrak{D}) = \langle t(x) \rangle$. Also, $\operatorname{Ker}(\pi) = \{ (d_1(x), 0) \in \mathfrak{R}_{r,s} \mid (d_1(x), d_2(x)) \in \mathfrak{D} \}$. Consider a set

$$K = \{ f(x) \in \mathbb{F}_q[x:\theta] \mid (f(x),0) \in \operatorname{Ker}(\pi) \}.$$

The set K is an left-submodule of $\frac{\mathbb{F}_q[x:\theta]}{\langle x^r-1\rangle}$. Therefore $K = \langle \ell(x) \rangle$ with $\ell(x) \mid_r (x^r - 1)$. Hence, any $(f(x), 0) \in \operatorname{Ker}(\pi)$ and $f(x) = \mu(x)\ell(x)$, where $\mu(x) \in \mathbb{F}_q[x:\theta]$. Consequently, we can express (f(x), 0) as $\lambda(x) \star (\ell(x), 0)$. Thus, $\operatorname{Ker}(\pi) = \langle \ell(x), 0 \rangle$ is a left $R[x:\Theta]$ -submodule of \mathfrak{D} . Moreover,

$$\frac{\mathfrak{D}}{\operatorname{Ker}(\pi)} \cong \pi(\mathfrak{D}) = \langle t(x) \rangle$$

Suppose $(s(x), t(x)) \in \mathfrak{D}$, then $\pi(s(x), t(x)) = t(x)$. Therefore, we conclude that the generator polynomials of (θ, Θ) -cyclic code of block length (r, s) over $\mathbb{F}_q R$ are $(\ell(x), 0)$ and (s(x), t(x)). More precisely, $\mathfrak{D} = \langle (\ell(x), 0), (s(x), t(x) \rangle$.

Lemma 3.13. Let $\mathfrak{D} = \langle (\ell(x), 0), (s(x), t(x)) \rangle$, where $x^r - 1 = f(x)\ell(x)$, $t(x) = \kappa_1 t_1(x) + \kappa_2 t_2(x) + \kappa_3 t_3(x) + \kappa_4 t_4(x)$ and $x^s - 1 = h_i(x)t_i(x)$ for i = 1, 2, 3, 4. Then deg $s(x) < \deg \ell(x)$, $\ell(x) \mid_r h_1(x)s(x)$ and lcm_l $(s(x), \ell(x)) \mid_r h_1(x)s(x)$.

Proof. Assume deg(s(x)) >deg $(\ell(x))$, then by using division algorithm in $\mathbb{F}_q[x:\theta]$, we have

$$s(x) = \mu_1(x)\ell(x) + \mu_2(x),$$

where $\mu_2(x) = 0$ or $\deg(\mu_2(x)) < \deg(\ell(x))$.

$$(\mu_2(x), t(x)) = (s(x) - \mu_1(x)\ell(x), t(x))$$

= $(s(x), t(x)) - \mu_1(x)(\ell(x), 0) \in \mathfrak{D}$
 $(s(x), t(x)) = (\mu_2(x), t(x)) + \mu_1(x)(\ell(x), 0).$

Hence, from this argument, it is clear that $\deg(s(x)) < \deg(\ell(x))$. Now,

$$(\kappa_1 h_1(x) + \kappa_2 h_2(x) + \kappa_3 h_3(x) + \kappa_4 h_4(x)) \star (s(x), t(x)) = (h_1(x) s(x), 0)$$

= $\langle (\ell(x), 0) \rangle,$

where $t(x) = \kappa_1 t_1(x) + \kappa_2 t_2(x) + \kappa_3 t_3(x) + \kappa_4 t_4(x)$. Thus, $\ell(x) \mid_r h_1(x) s(x)$.

Since $\ell(x) \mid_r (h_1(x)s(x))$ and $s(x) \mid_r (h_1(x)s(x))$. Hence, $\operatorname{lcm}_l (\ell(x), s(x)) \mid_r h_1(x)s(x)$.

Now, we deliberate algebraic structure of separable codes.

Definition 3.14. Let \mathfrak{D} be a linear code of block length (r, s) over $\mathbb{F}_q R$. Suppose the set \mathfrak{C}_r is obtained by deleting last s coordinates and \mathfrak{C}_s is obtained by deleting first r coordinates from \mathfrak{D} . It is worth noting that \mathfrak{C}_r and C_s are linear code of over \mathbb{F}_q and R, respectively, then \mathfrak{D} is referred to as separable code if $\mathfrak{D} = \mathfrak{C}_r \times \mathfrak{C}_s$.

Theorem 3.15. Let $\mathfrak{D} = \mathfrak{C}_r \times \mathfrak{C}_s$ be a separable code. Then \mathfrak{D} is called separable (θ, Θ) -cyclic code if and only if \mathfrak{C}_r and \mathfrak{C}_s are θ and Θ -cyclic code over \mathbb{F}_q and R, respectively.

Proof. Assume that \mathfrak{D} is a separable (θ, Θ) -cyclic code over $\mathbb{F}_q R$ and $(g_0, g_1, \ldots, g_{r-1}, b_0, b_1, \ldots, b_{s-1}) \in \mathfrak{D}$, then $(g_0, g_1, \ldots, g_{r-1}) \in \mathfrak{C}_r$, $(b_0, b_1, \ldots, b_{s-1}) \in \mathfrak{C}_s$. So

 $(\theta(g_{r-1}), \theta(g_0), \dots, \theta(g_{r-2}), \Theta(b_{s-1}), \Theta(b_0), \dots, \Theta(b_{s-2})) \in \mathfrak{D},$

which implies $(\theta(g_{r-1}), \theta(g_0), \ldots, \theta(g_{r-2})) \in \mathfrak{C}_r$ and $(\Theta(b_{s-1}), \Theta(b_0), \ldots, \theta(b_{s-2})) \in \mathfrak{C}_s$. Therefore, the result follows for one side.

For converse, assume that \mathfrak{C}_r and \mathfrak{C}_s are θ and Θ -cyclic code over \mathbb{F}_q and R, respectively and $(g_0, g_1, \ldots, g_{r-1}) \in \mathfrak{C}_r, (b_0, b_1, \ldots, b_{s-1}) \in \mathfrak{C}_s$, then $(\theta(g_{r-1}), \theta(g_0), \ldots, \theta(g_{r-2})) \in \mathfrak{C}_r$ and $(\Theta(b_{s-1}), \Theta(b_0), \ldots, \Theta(b_{s-2})) \in \mathfrak{C}_s$. Therefore,

$$(\theta(g_{r-1}), \theta(g_0), \dots, \theta(g_{r-2}), \Theta(b_{s-1}), \Theta(b_0), \dots, \theta(b_{s-2})) \in \mathfrak{D} = \mathfrak{C}_r \times \mathfrak{C}_s.$$

Therefore, it confirms that \mathfrak{D} is a separable (θ, Θ) -cyclic code over $\mathbb{F}_q R$.

Now, we explicitly present the generator polynomials of \mathfrak{C}_r and \mathfrak{C}_s .

Theorem 3.16. Let
$$\mathfrak{D} = \langle (\ell(x), 0), (s(x), t(x)) \rangle$$
. Then $\mathfrak{C}_r = \langle \gcd_r(\ell(x), s(x)) \rangle$ and $\mathfrak{C}_s = \langle t(x) \rangle$.

 $\begin{array}{l} \textit{Proof. Suppose } c(x) \in \mathfrak{C}_r, \ \text{then for some } e(x) \in \frac{R[x:\Theta]}{\langle x^s-1\rangle}, \ \text{we have } (c(x),e(x)) \in \mathfrak{D} \ . \ \text{Since } \\ \mathfrak{D} = \langle (\ell(x),0),(s(x),t(x)\rangle \ , \ \text{then } (c(x),e(x)) = \mu(x) \star (\ell(x),0) + \nu(x) \star (s(x),t(x)) \ \text{for some } \\ \mu(x),\nu(x) \in R[x:\Theta]. \ \text{This shows that } c(x) = \eta(\mu(x))\ell(x) + \eta(\nu(x))s(x). \ \text{It follows that } \\ \gcd_r(\ell(x),s(x)) \mid_r \ c(x). \ \text{Consequently}, \ c(x) \in \langle \gcd_r(\ell(x),s(x)) \rangle. \ \text{Thus, } \mathfrak{C}_r \subseteq \langle \gcd_r(\ell(x),s(x)) \rangle. \end{array}$

Conversely, since $gcd_r(\ell(x), s(x)) = \mu(x)\ell(x) + \nu(x)s(x)$ for some $\mu(x), \ \nu(x) \in \mathbb{F}_q[x : \theta]$. Therefore,

$$\operatorname{gcd}_r(\ell(x), s(x)), \nu(x)s(x)) = \mu(x) \star (\ell(x), 0) + \nu(x) \star (s(x), t(x)) \in \mathfrak{D}.$$

This infer that $\langle \gcd_r(\ell(x), s(x)) \rangle \subseteq \mathfrak{C}_r$. Hence, we deduce that $\mathfrak{C}_r = \langle \gcd_r(\ell(x), s(x)) \rangle$. We can determine $\mathfrak{C}_s = \langle t(x) \rangle$ by using similar steps.

Proposition 3.17. Let $\mathfrak{D} = \langle (\ell(x), 0), (s(x), t(x)) \rangle$. Then $\ell(x) \mid_r s(x)$ if and only if s(x) = 0.

Proof. Assume s(x) = 0, then it clearly holds.

Conversely, suppose that $\ell(x) \mid_r s(x)$. So $s(x) = \nu(x)\ell(x)$ for some polynomial $\nu(x) \in \mathbb{F}_q[x : \theta]$. Now, let us assume

$$\mathfrak{D}' = \langle (\ell(x), 0), (0, t(x)) \rangle.$$

So, $(0, t(x)) = (s(x), t(x)) - \nu(x) \star (\ell(x), 0) \in \mathfrak{D}$, implies that $\mathfrak{D}' \subseteq \mathfrak{D}$. Moreover, $(s(x), t(x)) = \nu(x) \star (\ell(x), 0)) + (0, t(x)) \in \mathfrak{D}'$, indicating that $\mathfrak{D} \subseteq \mathfrak{D}'$. Therefore, we conclude that s(x) = 0.

We sum up our earlier findings in our following theorem.

Theorem 3.18. Let $\mathfrak{D} = \langle (\ell(x), 0), (s(x), t(x)) \rangle$. The following are equivalent:

1.
$$\mathfrak{C}_r = \langle \ell(x) \rangle, \mathfrak{C}_s = \langle t(x) \rangle;$$

- 2. $\ell(x) \mid_r s(x);$
- 3. \mathfrak{D} is a separable code;
- 4. $\mathfrak{D} = \langle (\ell(x), 0), (0, t(x)) \rangle.$

Proof. Theorem 3.16 and Proposition 3.17 together provide the proof.

In the subsequent theorem, we aim to study the minimal generating sets for (θ, Θ) -cyclic codes over $\mathbb{F}_q R$. This investigation is crucial for understanding the fundamental properties and structures of these codes. By characterizing the minimal generating sets, we can gain deeper insights into the algebraic and combinatorial aspects of (θ, Θ) -cyclic codes.

Theorem 3.19. Let $\mathfrak{D} = \langle (\ell(x), 0), (s(x), t(x)) \rangle$, where $x^r - 1 = f(x)\ell(x)$, $t(x) = \kappa_1 t_1(x) + \kappa_2 t_2(x) + \kappa_3 t_3(x) + \kappa_4 t_4(x)$ and $x^s - 1 = h_i(x)t_i(x)$, for i = 1, 2, 3, 4. Suppose

$$\begin{split} G_1 &= \bigcup_{i=0}^{\deg(f(x))-1} \{x^i \star (\ell(x), 0)\}, \\ G_2 &= \bigcup_{i=0}^{\deg(h_1(x))-1} \{x^i \star (s(x), \kappa_1 t_1(x))\}, \\ G_3 &= \bigcup_{i=0}^{\deg(h_2(x))-1} \{x^i \star (0, \kappa_2 t_2(x))\}, \\ G_4 &= \bigcup_{i=0}^{\deg(h_3(x))-1} \{x^i \star (0, \kappa_3 t_3(x))\}, \\ G_5 &= \bigcup_{i=0}^{\deg(h_4(x))-1} \{x^i \star (0, \kappa_4 t_4(x))\}. \end{split}$$

Then $G = G_1 \cup G_2 \cup G_3 \cup G_4 \cup G_5$ constitutes a minimal generating set of \mathfrak{D} . Additionally, $|\mathfrak{D}| = q^{\deg(f(x)) + \sum_{i=1}^4 \deg(h_i(x))}$.

Proof. Suppose $d(x) \in \mathfrak{D}$, then

$$d(x) = m_1(x) \star (\ell(x), 0) + m_2(x) \star (s(x), \kappa_1 t_1(x) + \kappa_2 t_2(x) + \kappa_3 t_3(x) + \kappa_4 t_4(x)),$$

for some polynomials $m_1(x), m_2(x) \in R[x : \Theta]$. Let $m_1(x) = m'_0 + m'_1 x + \dots + m'_\alpha x^\alpha$, where $m'_i = \kappa_1 n'_{1,i} + \kappa_2 n'_{2,i} + \kappa_3 n'_{3,i} + \kappa_4 n'_{4,i}$ for $i = 0, 1, \dots, \alpha$. Then $m_1(x)$ can be written as

$$m_1(x) = \kappa_1 (n'_{1,0} + n'_{1,1}x + \dots + n'_{1,\alpha}x^{\alpha}) + \kappa_2 (n'_{2,0} + n'_{2,1}x + \dots + n'_{2,\alpha}x^{\alpha}) + \kappa_3 (n'_{3,0} + n'_{3,1}x + \dots + n'_{3,\alpha}x^{\alpha}) + \kappa_4 (n'_{4,0} + n'_{4,1}x + \dots + n'_{4,\alpha}x^{\alpha}) = \kappa_1 n'_1(x) + \kappa_2 n'_2(x) + \kappa_3 n'_3(x) + \kappa_4 n'_4(x),$$

where $n'_{i}(x) = n'_{i,0} + n'_{i,1}x + \dots + n'_{i,\alpha}x^{\alpha}$ for i = 1, 2, 3, 4. Now,

$$m_{1}(x) \star (\ell(x), 0) = (\kappa_{1}n'_{1}(x) + \kappa_{2}n'_{2}(x) + \kappa_{3}n'_{3}(x) + \kappa_{4}n'_{4}(x)) \star (\ell(x), 0)$$

= $n'_{1}(x) \star (\eta(\kappa_{1})\ell(x), 0) + n'_{2}(x) \star (\eta(\kappa_{2})\ell(x), 0)$
+ $n'_{3}(x) \star (\eta(\kappa_{3})\ell(x), 0) + n'_{4}(x) \star (\eta(\kappa_{4})\ell(x), 0)$
= $n'_{1}(x) \star (\ell(x), 0).$

In scenario, $\deg(n'_1(x)) < \deg(f(x))$ implies $n'_1(x) \star (\ell(x), 0) \in \operatorname{Span}(G_1)$. In another scenario, by division algorithm, we have $n'_1(x) = \mu(x)f(x) + \nu(x)$ such that $\nu(x) = 0$ or $\deg(\nu(x)) < \deg(f(x))$, for $\mu(x), \nu(x) \in \mathbb{F}_q[x:\theta]$. Therefore, we get

$$\begin{aligned} n_1'(x) \star (\ell(x), 0) &= (\mu(x)f(x) + \nu(x)) \star (\ell(x), 0) \\ &= \mu(x)f(x) \star (\ell(x), 0) + \nu(x) \star (\ell(x), 0) \\ &= 0 + \nu(x) \star (\ell(x), 0). \end{aligned}$$

Thus, it concludes $n'_1(x) \star (\ell(x), 0) \in \text{Span}(G_1)$. Now, it remains to prove that

$$m_2(x) \star (s(x), \kappa_1 t_1(x) + \kappa_2 t_2(x) + \kappa_3 t_3(x) + \kappa_4 t_4(x)) \in \text{Span}(G_1 \cup G_2 \cup G_3 \cup G_4 \cup G_5).$$

Let $m_2(x) = m_0'' + m_1''x + \dots + m_\beta''x^\beta$, where $m_i'' = \kappa_1 n_{1,i}'' + \kappa_2 n_{2,i}'' + \kappa_3 n_{3,i}'' + \kappa_4 n_{4,i}''$ for $i = 0, 1, \dots, \beta$. So,

$$m_{2}(x) = \kappa_{1}(n_{1,0}'' + n_{1,1}'' x + \dots + n_{1,\beta}'' x^{\beta}) + \kappa_{2}(n_{2,0}'' + n_{2,1}'' x + \dots + n_{2,\beta}'' x^{\beta}) + \kappa_{3}(n_{3,0}'' + n_{3,1}'' x + \dots + n_{3,\beta}'' x^{\beta}) + \kappa_{4}(n_{4,0}'' + n_{4,0}'' x + \dots + n_{4,1}'' x^{\beta}) = \kappa_{1}n_{1}''(x) + \kappa_{2}n_{2}''(x) + \kappa_{3}n_{3}''(x) + \kappa_{4}n_{4}''(x),$$

where $n''_{i}(x) = n''_{i,0} + n''_{i,1}x + \dots + n''_{i,\alpha}x^{\alpha}$ for i = 1, 2, 3, 4. Now,

$$\begin{split} m_2(x) \star (s(x), \kappa_1 t_1(x) + \kappa_2 t_2(x) + \kappa_3 t_3(x) + \kappa_4 t_4(x)) \\ &= (\kappa_1 n_1''(x) + \kappa_2 n_2''(x) + \kappa_3 n_3''(x) + \kappa_4 n_4''(x)) \star (s(x), \kappa_1 t_1(x) + \kappa_2 t_2(x) + \kappa_3 t_3(x) + \kappa_4 t_4(x))) \\ &= n_1''(x) \star (s(x), \kappa_1 t_1(x)) + n_2''(x) \star (0, \kappa_2 t_2(x)) + n_3''(x) \star (0, \kappa_3 t_3(x)) + n_4''(x) \star (0, \kappa_4 t_4(x)). \end{split}$$

In scenario, $\deg(n_2''(x)) < \deg(h_2(x))$ implies $n_2''(x) \star (0, \kappa_2 t_2(x)) \in \operatorname{Span}(G_3)$. In another scenario, by division algorithm, we have $n_2''(x) = q_2(x)h_2(x) + r_2(x)$ such that $r_2(x) = 0$ or $\deg(r_2(x)) < \deg(h_2(x))$, for $q_2(x), r_2(x) \in R[x : \Theta]$. Therefore, we get

$$n_2''(x) \star (0, \kappa_2 t_2(x)) = (q_2(x)h_2(x) + r_2(x)) \star (0, \kappa_2 t_2(x))$$

= $q_2(x)h_2(x) \star (0, \kappa_2 t_2(x)) + r_2(x) \star (0, \kappa_2 t_2(x))$
= $0 + r_2(x) \star (0, \kappa_2 t_2(x)).$

Hence, we get $n_2''(x) \star (0, \kappa_2 t_2(x)) \in \text{Span}(G_3)$. In similar manner, we can prove that $n_3''(x) \star (0, \kappa_3 t_3(x)) \in \text{Span}(G_4)$ and $n_4''(x) \star (0, \kappa_4 t_4(x)) \in \text{Span}(G_5)$.

Now, consider $n''_1(x) \star (s(x), \kappa_1 t_1(x))$. In scenario, $\deg(n''_1(x)) < \deg(h_1(x))$ implies $n''_1(x) \star (s(x), \kappa_1 t_1(x)) \in \operatorname{Span}(G_2)$. In another scenario, by division algorithm, we have $n''_1(x) = q_1(x)h_1(x) + r_1(x)$ such that $r_1(x) = 0$ or $\deg(r_1(x)) < \deg(h_1(x))$, for $q_1(x), r_1(x) \in R[x : \theta]$. Therefore, we get

$$n_1''(x) \star (s(x), \kappa_1 t_1(x)) = (q_1(x)h_1(x)) \star (s(x), \kappa_1 t_1(x)) + r_1(x) \star (s(x), \kappa_1 t_1(x)))$$

= $q_1(x) \star ((h_1(x) \star s(x)), 0) + r_1(x) \star (s(x), \kappa_1 t_1(x)),$

and $r_1(x) \star (s(x), \kappa_1 t_1(x)) \in \operatorname{Span}(G_2)$. From Lemma 3.13, we have $\ell(x) \mid_r h_1(x)s(x)$ which implies, $q_1(x) \star ((h_1(x) \star s(x)), 0) \in \operatorname{Span}(G_1)$. Thus, we get $n''_1(x) \star (s(x), \kappa_1 t_1(x)) \in \operatorname{Span}(G_1 \cup G_2)$. Hence, we have $d(x) \in \operatorname{Span}(G_1 \cup G_2 \cup G_3 \cup G_4 \cup G_5)$. It is worth noting that the elements in span of $G_1 \cup G_2 \cup G_3 \cup G_4 \cup G_5$ are linearly independent as $R[x : \Theta]$ -submodule. This implies that $G = \bigcup_{i=1}^5 G_i$ forms a minimal spanning set of \mathfrak{D} . Moreover, $|\mathfrak{D}| = q^{\operatorname{deg}(f(x) + \sum_{i=1}^4 (h_i(x)) - \sum$

Now, we provide a detailed example to elaborate the previous results.

Example 3.20. Let $q = 27, r = 3 = s, R = \mathbb{F}_{27} + u\mathbb{F}_{27} + u\mathbb{F}_{27} + uv\mathbb{F}_{27}$, where $\mathbb{F}_{27} = \mathbb{F}_3[\omega]$ with $\omega^3 + 2\omega + 2 = 0$ and Frobenius automorphism $\theta(\alpha) = \alpha^3$ for all $\alpha \in \mathbb{F}_{27}$. We have order of θ , Θ is 3 such that $\operatorname{ord}(\theta) \mid r, \operatorname{ord}(\Theta) \mid s$. Since $\mathbb{F}_{27}[x : \theta]$ is not a UFD, so the polynomial $x^3 - 1$ has more than one factorizations. Consider $\mathfrak{D} = \langle (\ell(x), 0), (s(x), t(x)) \rangle$ is a (θ, Θ) -cyclic code of block length (3, 3) over $\mathbb{F}_{27}R$, where

$$\ell(x) = x + \omega^{17}, t_1(x) = t_2(x) = x + \omega^5, t_3(x) = t_4(x) = x^2 + \omega^6 x + \omega^8, s(x) = \omega.$$

Next, we have

$$f(x) = x^{2} + \omega^{6}x + \omega^{8},$$

$$h_{1}(x) = h_{2}(x) = x^{2} + \omega^{18}x + \omega^{22},$$

$$h_{3}(x) = h_{4}(x) = x + \omega^{17}.$$

Now, \mathfrak{D} possesses the generator matrix G as shown below,

$$G = \begin{pmatrix} \omega^{17} & 1 & 0 & 0 & 0 & 0 \\ 0 & \omega^{25} & 1 & 0 & 0 & 0 \\ \omega & 0 & 0 & \omega^5 \kappa_1 & \kappa_1 & 0 \\ 0 & \omega^3 & 0 & 0 & \omega^{15} \kappa_1 & \kappa_1 \\ 0 & 0 & 0 & \omega^5 \kappa_2 & \kappa_2 & 0 \\ 0 & 0 & 0 & \omega^5 \kappa_2 & \kappa_2 & 0 \\ 0 & 0 & 0 & \omega^{15} \kappa_2 & \kappa_2 \\ 0 & 0 & 0 & \omega^8 \kappa_3 & \omega^6 \kappa_3 & \kappa_3 \\ 0 & 0 & 0 & \omega^8 \kappa_4 & \omega^6 \kappa_4 & \kappa_4 \end{pmatrix},$$

and according to Theorem 3.19, \mathfrak{D} has $= 27^8$ codewords.

In Table 1, we provide examples to demonstrate our findings, where we obtained some nearoptimal, marked as **, and optimal codes, marked as * over the field \mathbb{F}_9 , where $\mathbb{F}_9 = \mathbb{F}_3[\omega]$ with $\omega^2 + \omega + 1 = 0$. In rest of Table 1, we use Frobenius automorphism over the field \mathbb{F}_9 defined as $\theta(\alpha) = \alpha^3$ for all $\alpha \in \mathbb{F}_9$ and the coefficients of generator polynomials set in the ascending order for instance, $\omega^7 \omega^3 1$ represents the polynomial $\omega^7 + \omega^3 x + x^2$.

Table 1: Near-optimal and optimal codes from Θ -cyclic code \mathfrak{C}_s over R

q	n	$t_1(x)$	$t_2(x)$	$t_3(x)$	$t_4(x)$	$\Phi_1(\mathfrak{C}_s)$
9	4	$\omega^7 \omega^3 1$	11	$\omega^{5}1$	ω	$[16, 12, 4]^*$
9	6	$\omega^2 \omega 1$	11	$\omega^6 1$	1	$[24, 20, 3]^{**}$
9	6	$2\omega^5\omega^3 1$	$\omega^2 1$	21	ω	$[24, 19, 4]^*$
9	8	$\omega^7 \omega^6 \omega 1$	$\omega^6 \omega^3 1 \omega^6 \omega^7 1$	$\omega^7 01$	$\omega^3 21$	$[32, 19, 8]^{**}$
9	8	$\omega^7 \omega^6 \omega 1$	$\omega\omega^3 2\omega^6 1$	$1\omega^3 1$	$\omega 11$	$[32, 21, 7]^{**}$
9	8	$\omega^3 \omega^7 \omega^2 1$	$\omega^5 \omega^7 1$	$\omega^7 \omega 1$	11	$[32, 24, 6]^*$
9	8	$\omega^7 \omega^5 21$	$\omega^7 1$	$\omega 1$	1	$[32, 27, 4]^*$
9	12	$2\omega^6\omega 1$	$\omega^2 1$	$\omega^{3}1$	1	$[48, 43, 3]^{**}$
9	12	$\omega^5 \omega^6 \omega^6 21$	$\omega^2 1$	$\omega^{3}1$	1	$[48, 42, 4]^{**}$

4 Duality of (θ, Θ) -cyclic codes over $\mathbb{F}_q R$

This section focuses on the algebraic structural characteristics of the dual of (θ, Θ) -cyclic code of block length (r, s) over $\mathbb{F}_q R$. Moreover, we determine the relation between (θ, Θ) -cyclic code and their dual.

In preliminaries, it has shown that the dual of (θ, Θ) -cyclic code \mathfrak{D} of block length (r, s) over $\mathbb{F}_q R$ again a (θ, Θ) -cyclic code. Therefore,

$$\mathfrak{D}^{\perp} = \langle (\bar{\ell}(x), 0), (\bar{s}(x), \bar{t}(x)) \rangle,$$

where $\bar{\ell}(x)$ and $\bar{s}(x) \in \frac{\mathbb{F}_q[x:\theta]}{\langle x^r-1 \rangle}$, $\bar{t}(x) \in \frac{R[x:\Theta]}{\langle x^s-1 \rangle}$, and $\deg(\bar{\ell}(x) < \deg(\bar{s}(x))$.

Define $\Gamma_n(x) = \sum_{i=0}^{n-1} x^i$. We give the following lemma without a formal proof.

Lemma 4.1. Suppose that $\mathfrak{m}, \mathfrak{n} \in \mathbb{N}$. Then

$$x^{\mathfrak{mn}-1} = (x^{\mathfrak{m}} - 1)\Gamma_{\mathfrak{n}}(x^{\mathfrak{m}}) = \Gamma_{\mathfrak{n}}(x^{\mathfrak{m}})(x^{\mathfrak{m}} - 1).$$

Let a non-zero polynomial $\gamma(x) = a_0 + a_1 x + a_2 x^2 + \dots + a_t x^t$ in $\mathbb{F}_q[x:\theta]$. Then

$$\gamma^{\dagger}(x) = a_t + \theta(a_{t-1})x + \theta^2(a_{t-2})x^2 + \dots + \theta^t(a_0)x^t$$

is referred to as reciprocal polynomial of $\gamma(x)$. Alternatively, we can express the reciprocal as $\gamma^{\dagger}(x) = \sum_{i=0}^{t} \theta^{i}(a_{t-i})x^{i}$.

Lemma 4.2. [13, Lemma 5] Suppose Ψ_1 is a map defined as

$$\Psi_1: \mathbb{F}_q[x:\theta] \to \mathbb{F}_q[x:\theta],$$

and given by

$$\sum_{k=0}^{r} g_k x^k \to \sum_{k=0}^{r} \theta(g_k) x^k.$$

Then Ψ_1 is a ring homomorphism.

Proposition 4.3. Suppose $\mu(x), \nu(x) \in \mathbb{F}_q[x:\theta]$ are any two polynomials. Then

1. If $\deg(\mu(x)) \ge \deg(\nu(x))$, then $(\mu(x) + \nu(x))^{\dagger} = \mu^{\dagger}(x) + x^{\deg(\mu(x)) - \deg(\nu(x))}\nu^{\dagger}(x)$.

2.
$$(\mu(x)^{\dagger})^{\dagger} = \Psi_1^r(\mu(x)), \text{ where } \deg(\mu(x)) = r.$$

3.
$$(\mu(x)\nu(x))^{\dagger} = \Psi_1^{\deg\mu(x)}(\nu^{\dagger}(x))\mu^{\dagger}(x).$$

Proof. Similar concepts are used to prove this proposition as given in [22, Lemma 2.8]. \Box

Lemma 4.4. [13, Lemma 5] Suppose Ψ_2 is a map defined as

$$\Psi_2: R[x:\Theta] \to R[x:\Theta]$$

and given by

$$\sum_{j=0}^{s} b_j x^j \to \sum_{j=0}^{s} \Theta(b_j) x^j,$$

where $b_j \in R$. Then Ψ_2 is a ring homomorphism.

Proposition 4.5. Suppose $\mu_1(x), \nu_1(x) \in \mathbb{R}[x:\Theta]$ are any two polynomials. Then

- 1. $(\mu_1(x)\nu_1(x))^{\dagger} = \Psi_2^{\deg \mu_1(x)}(\nu_1^{\dagger}(x))\mu_1^{\dagger}(x).$
- 2. $(\mu_1(x)^{\dagger})^{\dagger} = \Psi_2^s(\mu(x)), \text{ where } \deg(\mu_1(x)) = s.$
- 3. If $\deg(\mu_1(x)) \ge \deg(\nu_1(x))$, then $(\mu_1(x) + \nu_1(x))^{\dagger} = \mu_1^{\dagger}(x) + x^{\deg(\mu_1(x)) \deg(\nu_1(x))} \nu_1^{\dagger}(x)$.

Proof. Similar concepts are used to prove this proposition as given in [22, Lemma 5].

We assume $\mathfrak{m} = r \times s$ throughout this paper.

Definition 4.6. Let $d(x) = (g(x), b(x)), d'(x) = (g'(x)), b'(x)) \in \mathfrak{R}_{r,s}$. Define the map

$$o: \mathfrak{R}_{r,s} \times \mathfrak{R}_{r,s} \to \frac{R[x:\Theta]}{\langle x^{\mathfrak{m}} - 1 \rangle},$$

such that

$$o(d(x), d'(x)) = \kappa_1 g(x) \Psi_1^{\mathfrak{m}-\deg(g'(x))} (g'^{\dagger}(x)) x^{\mathfrak{m}-1-\deg(g'(x))} \Gamma_{\frac{\mathfrak{m}}{r}}(x^r) + b(x) \Psi_2^{\mathfrak{m}-\deg(b'(x))} (b'^{\dagger}(x)) x^{\mathfrak{m}-1-\deg(b'(x))} \Gamma_{\frac{\mathfrak{m}}{s}}(x^s) \pmod{(x^{\mathfrak{m}}-1)}.$$

Clearly, between left $R[x : \Theta]$ -modules, the map "o" is bilinear. For more detailed information about this map, we refer [3, Definition 4.2]. We denote o(d(x), d'(x)) by d(x) o d'(x) for convenience.

Proposition 4.7. Let d and d' be two vectors in $\mathbb{F}_q^r \times \mathbb{R}^s$ and their associated polynomials are d(x) = (g(x), b(x)) and $d'(x) = (g'(x), b'(x)) \in \mathfrak{R}_{r,s}$. Then d and all its (θ, Θ) -cyclic shift are orthogonal to d' if and only if d(x) o $d'(x) \equiv 0 \pmod{x^m - 1}$.

Proof. Suppose $\boldsymbol{d} = (g_0, g_1, \dots, g_{r-1}b_0, b_1, \dots, b_{s-1})$ and $\boldsymbol{d}' = (g'_0, g'_1, \dots, g'_{r-1}, b'_0, b'_1, \dots, b'_{s-1}) \in \mathbb{F}_q^r \times R^s$. For $0 \leq i \leq \mathfrak{m} - 1$, the i^{th} (θ, Θ) -cyclic shift of \boldsymbol{d} be represented by the vector $\boldsymbol{d}^{(i)} = (\theta^i(g_{0-i}), \theta^i(g_{1-i}), \dots, \theta^i(g_{r-1-i}), \Theta^i(b_{0-i}), \Theta^i(b_{1-i}), \dots, \Theta^i(b_{s-1-i}))$. Now, $\boldsymbol{d}^{(i)} \cdot \boldsymbol{d}' \equiv 0$ (mod $x^{\mathfrak{m}} - 1$) if and only if

$$\kappa_1 \sum_{j=0}^{r-1} \theta^i(g_{j-i}) g'_j + \sum_{\nu=0}^{s-1} \Theta^i(b'_{\nu-i}) b'_{\nu} = 0.$$

Let $S_i = \kappa_1 \sum_{j=0}^{r-1} \theta^i(g_{j-i})g'_j + \sum_{\nu=0}^{s-1} \Theta^i(b'_{\nu-i})b'_{\nu}$. Then we have

$$d(x) \ o \ d'(x) = \kappa_1 \left[\sum_{\mu=0}^{r-1} \sum_{j=\mu}^{r-1} g_{j-\mu} \theta^{\mathfrak{m}-\mu}(g'_j) x^{\mathfrak{m}-1-\mu} + \sum_{\mu=1}^{r-1} \sum_{j=\mu}^{r-1} g_j \theta^{\mu}(g'_{j-\mu}) x^{\mathfrak{m}-1+\mu} \right] \Gamma_{\frac{\mathfrak{m}}{r}}(x^r) + \left[\sum_{k=0}^{s-1} \sum_{\nu=k}^{s-1} b_{\nu-k} \Theta^{\mathfrak{m}-k}(b'_{\nu}) x^{\mathfrak{m}-1-k} + \sum_{k=1}^{s-1} \sum_{\nu=k}^{s-1} b_k \Theta^k(b'_{\nu-k}) x^{\mathfrak{m}-1+k} \right] \Gamma_{\frac{\mathfrak{m}}{s}}(x^s) = \sum_{i=0}^{\mathfrak{m}-1} \rho_{(\theta,\Theta)}^{\mathfrak{m}-i} \ S_i x^{\mathfrak{m}-1-i} \pmod{x^{\mathfrak{m}}-1}.$$

Hence, $d(x) \circ d'(x) \equiv 0 \pmod{x^{\mathfrak{m}} - 1}$ if and only if $S_i = 0$ for $i = 0, 1, \dots, \mathfrak{m} - 1$.

We give the following proposition to find the generator polynomials for \mathfrak{D}^{\perp} based on the above discussion.

Proposition 4.8. Let $d(x) = (g(x), b(x)), d'(x) = (g'(x), b'(x)) \in \mathfrak{R}_{r,s}$ and d(x) o $d'(x) \equiv 0 \pmod{x^{\mathfrak{m}} - 1}$. Then

1.
$$g(x)\Psi_1^{\mathfrak{m}-\deg(g'(x))}(g'^{\dagger}(x)) \equiv 0 \pmod{x^r - 1}$$
, if either $b(x) = 0$ or $b'(x) = 0$.
2. $b(x)\Psi_2^{\mathfrak{m}-\deg(b'(x))}(b'^{\dagger}(x)) \equiv 0 \pmod{x^s - 1}$, if either $g(x) = 0$ or $g'(x) = 0$.

Proof. Assume that b(x) = 0 or b'(x) = 0, then

$$d(x) \ o \ d'(x) = \kappa_1 g(x) \Psi_1^{\mathfrak{m} - \deg(g'(x))}(g'^{\dagger}(x)) x^{\mathfrak{m} - 1 - \deg(g'(x))} \Gamma_{\frac{\mathfrak{m}}{r}}(x^r) + 0 \equiv 0 \pmod{x^{\mathfrak{m}} - 1}.$$

Implying the existence of $\lambda(x) \in \mathbb{F}_q[x:\theta]$ such that

$$\kappa_1 g(x) \Psi_1^{\mathfrak{m}-\deg(g'(x))}(g'^{\dagger}(x)) x^{\mathfrak{m}-1-\deg(g'(x))} \Gamma_{\frac{\mathfrak{m}}{r}}(x^r) = \kappa_1 \lambda(x) x^{\mathfrak{m}} - 1.$$

Since $\Gamma_{\frac{m}{r}}(x^r) = \frac{x^m - 1}{x^r - 1}$ and $(x^m - 1)(x^r - 1) = (x^r - 1)(x^m - 1)$. Hence,

$$\kappa_1 g(x) \Psi_1^{\mathfrak{m}-\deg(g'(x))}(g'^{\dagger}(x)) x^{\mathfrak{m}} = \kappa_1 \lambda(x) x^{(\deg(g'(x))+1)}(x^r-1).$$

Consequently,

$$g(x)\Psi_1^{\mathfrak{m}-\deg(g'(x))}(g'^{\dagger}(x)) \equiv 0 \pmod{x^r-1}.$$

The second case can be proven using the same argument.

Proposition 4.9. Let $\mathfrak{D} = \langle (\ell(x), 0), (s(x), t(x)) \rangle$, where $t(x) = \kappa_1 t_1(x) + \kappa_2 t_2(x) + \kappa_3 t_3(x) + \kappa_4 t_4(x)$. Then

$$\begin{split} |\mathfrak{C}_{r}| &= q^{r-\deg(\gcd_{r}(\ell(x),s(x)))}, \ |\mathfrak{C}_{s}| = q^{4s-\sum_{i=1}^{4} \deg(t_{i}(x))}, \\ |\kappa_{1}\mathfrak{C}_{s}| &= q^{s-\deg(t_{1}(x))}, \ |\kappa_{2}\mathfrak{C}_{s}| = q^{s-\deg(t_{2}(x))}, \\ |\kappa_{3}\mathfrak{C}_{s}| &= q^{s-\deg(t_{3}(x))}, \ |\kappa_{4}\mathfrak{C}_{s}| = q^{s-\deg(t_{4}(x))}, \\ |(\mathfrak{C}_{r})^{\perp}| &= q^{\deg(\gcd_{r}(\ell(x),s(x)))}, \ |(\mathfrak{C}_{s})^{\perp}| = q^{\sum_{i=1}^{4} \deg(t_{i}(x))}, \\ |(\mathfrak{C}^{\perp})_{r}| &= q^{\deg(\ell(x))}, \ |(\mathfrak{C}^{\perp})_{s}| = q^{\sum_{i=1}^{4} \deg(t_{i}(x)) + \deg(\ell(x)) - \deg(\gcd_{r}(\ell(x),s(x)))}, \\ |\kappa_{1}(\mathfrak{C}_{s})^{\perp}| &= q^{\deg(t_{1}(x))}, \ |\kappa_{2}(\mathfrak{C}_{s})^{\perp}| = q^{\deg(t_{2}(x))}, \\ |\kappa_{3}(\mathfrak{C}_{s})^{\perp}| &= q^{\deg(t_{1}(x)) + \deg(\ell(x) - \deg(\gcd_{r}(\ell(x),s(x))))}, \ |\kappa_{2}(\mathfrak{C}^{\perp})_{s}| = q^{\deg(t_{2}(x))}, \\ |\kappa_{3}(\mathfrak{C}^{\perp})_{s}| &= q^{\deg(t_{3}(x))}, \ |\kappa_{4}(\mathfrak{C}^{\perp})_{s}| = q^{\deg(t_{4}(x))}, \\ |\kappa_{3}(\mathfrak{C}^{\perp})_{s}| &= q^{\deg(t_{3}(x))}, \ |\kappa_{4}(\mathfrak{C}^{\perp})_{s}| = q^{\deg(t_{4}(x))}. \end{split}$$

Proof. Since we have proved in Theorem 3.16 that $\mathfrak{C}_r = \langle \gcd_r(\ell(x), s(x)) \rangle$ and $\mathfrak{C}_s = \langle t(x) \rangle = \langle \kappa_1 t_1(x) + \kappa_2 t_2(x) + \kappa_3 t_3(x) + \kappa_4 t_4(x) \rangle$ Therefore, $|\mathfrak{C}_r| = q^{r-\deg(\gcd_r(\ell(x), s(x)))}$ and $|\mathfrak{C}_s| = q^{4s - \sum_{i=1}^4 \deg(t_i(x))}$. We can demonstrate the other results in a similar way.

In our next proposition, we shall determine the degree of generator polynomials for \mathfrak{D}^{\perp} .

Proposition 4.10. Let $\mathfrak{D} = \langle (\ell(x), 0), (s(x), t(x)) \rangle$, where $t(x) = \kappa_1 t_1(x) + \kappa_2 t_2(x) + \kappa_3 t_3(x) + \kappa_4 t_4(x)$ and $\mathfrak{D}^{\perp} = \langle (\bar{\ell}(x), 0), (\bar{s}(x), \bar{t}(x)) \rangle$. Then

$$deg(\ell(x)) = r - deg(gcd_r(\ell(x), s(x))),$$

$$deg(\bar{t}_1(x)) = s - deg(t_1(x)) - deg(\ell(x)) + deg(gcd_r(s(x), \ell(x))),$$

$$deg(\bar{t}_j(x)) = s - deg(t_j(x)), \text{ for } j = 2, 3, 4.$$

Proof. It can be easily established that $(\mathfrak{C}_r)^{\perp} = \langle \bar{\ell}(x) \rangle$. So $|(\mathfrak{C}_r)^{\perp}| = q^{r-\deg(\bar{\ell}(x))}$. By Proposition 4.9, we get that $|(\mathfrak{C}_r)^{\perp}| = q^{\deg(\gcd_r(\ell(x),s(x)))}$. Hence, $\deg(\bar{\ell}(x)) = r - \deg(\gcd_r(\ell(x),s(x)))$. Again, it can be easily seen that that $\kappa_1(\mathfrak{C}^{\perp})_s = \langle \kappa_1 \bar{t}_1(x) \rangle$ is θ -cyclic code over \mathbb{F}_q . Therefore, $|\kappa_1(\mathfrak{C}^{\perp})_s| = q^{s-\deg(\bar{t}_1(x))}$. Now, by using Proposition 4.9, we have $|\kappa_1(\mathfrak{C}^{\perp})_s| = q^{\deg(t_1(x))+\deg(\ell(x))-\deg(\gcd_r(\ell(x),s(x)))}$. This shows that $\deg(\bar{t}_1(x)) = s - \deg(t_1(x)) - \deg(\ell(x)) + \deg(\gcd_r(\ell(x),s(x)))$. Similarly, the remaining parts can be proved.

In the next theorem, using the above results, we shall calculate the generator polynomial for \mathfrak{D}^{\perp} .

Theorem 4.11. Let $\mathfrak{D} = \langle (\ell(x), 0), (s(x), t(x)) \rangle$. Then

$$\mathfrak{D}^{\perp} = \langle (\bar{\ell}(x), 0), (\bar{s}(x), \bar{t}(x)) \rangle,$$

 θ],

such that

$$1. \ \bar{\ell}(x) = \frac{x^{r-1}}{\gcd_l\left(\Psi_1^{\mathfrak{m}-\deg(\ell(x))}(\ell^{\dagger}(x)),\Psi_2^{\mathfrak{m}-\deg(s(x))}(s^{\dagger}(x))\right)},$$

$$2. \ \bar{s}(x) = \frac{\lambda(x)(x^{r}-1)}{\Psi_1^{\mathfrak{m}-\deg(\ell(x))}(\ell^{\dagger}(x))}, \ where \ \lambda(x) \in \mathbb{F}_q[x:\theta]$$

$$3. \ \bar{t}_1(x) = \frac{x^{s-1}}{\Psi_2^{\mathfrak{m}-\deg\left(\frac{\ell(x)t_1(x)}{\gcd_l(\ell(x),s(x))}\right)\left(\frac{\ell(x)t_1(x)}{\gcd_l(\ell(x),s(x))}\right)^{\dagger}},$$

$$4. \ \bar{t}_2(x) = \frac{x^{s-1}}{\Psi_2^{\mathfrak{m}-\deg(t_2(x)}(t^{\dagger}_2(x))},$$

$$5. \ \bar{t}_3(x) = \frac{x^{s-1}}{\Psi_2^{\mathfrak{m}-\deg(t_3(x)}(t^{\dagger}_3(x))},$$

$$6. \ \bar{t}_4(x) = \frac{x^{s-1}}{\Psi_2^{\mathfrak{m}-\deg(t_4(x)}(t^{\dagger}_4(x))}.$$

Proof. (1) Since $(\bar{\ell}(x), 0) \in \mathfrak{D}^{\perp}$. So $(\bar{\ell}(x), 0)$ o $(\ell(x), 0) \equiv 0 \pmod{x^{\mathfrak{m}} - 1}$ and $(\bar{\ell}(x), 0)$ o $(s(x), t(x)) \equiv 0 \pmod{x^{\mathfrak{m}} - 1}$.

Now, using Proposition 4.8, we obtain

$$\bar{\ell}(x)\Psi_1^{\mathfrak{m}-\deg(\ell(x))}(\ell^{\dagger}(x)) \equiv 0 \pmod{x^r - 1}$$

and

$$\bar{\ell}(x)\Psi_1^{\mathfrak{m}-\deg(s(x))}(s^{\dagger}(x)) \equiv 0 \pmod{x^r - 1}$$

Since $(x^r - 1) \in Z((\mathbb{F}_q[x:\theta]))$, so using Lemma 3.8, we obtain

$$\bar{\ell}(x)\operatorname{gcd}_l\left(\Psi_1^{\mathfrak{m}-\operatorname{deg}(\ell(x)}(\ell^{\dagger}(x)),\Psi_1^{\mathfrak{m}-\operatorname{deg}(s(x))}(s^{\dagger}(x))\right) \equiv 0 \pmod{x^r - 1}$$

Implying

$$\bar{\ell}(x)\operatorname{gcd}_l\left(\Psi_1^{\mathfrak{m}-\operatorname{deg}(\ell(x)}(\ell^{\dagger}(x)),\Psi_1^{\mathfrak{m}-\operatorname{deg}(s(x))}(s^{\dagger}(x))\right) = \mu(x)(x^r-1),$$

for $\mu(x) \in \mathbb{F}_q[x:\theta]$. As $\deg(\bar{\ell}(x)) = r - \deg(\gcd_r(\ell(x), s(x)))$, so we conclude that $\mu(x) = 1$. Hence,

$$\bar{\ell}(x) = \frac{x^{\prime} - 1}{\gcd_l\left(\Psi_1^{\mathfrak{m}-\deg(\ell(x))}(\ell^{\dagger}(x)), \Psi_1^{\mathfrak{m}-\deg(s(x))}(s^{\dagger}(x)\right)\right)}$$

(2) As $d'(x) = (\bar{s}(x), \bar{t}(x)) \in \mathfrak{D}^{\perp}$, then

$$d'(x)o(\ell(x),0) = (\bar{s}(x),\bar{t}(x)) \ o \ (\ell(x),0)$$

by the bilinearity of map "o", we have

$$\bar{s}(x)\Psi_1^{\mathfrak{m}-\deg(\ell(x))}(\ell^{\dagger}(x)) \equiv 0 \pmod{x^r-1}.$$

Consequently, for some $\lambda(x) \in \mathbb{F}_q[x:\theta]$, we have

$$\bar{s}(x)\Psi_1^{\mathfrak{m}-\deg(\ell(x))}(\ell^{\dagger}(x)) = \lambda(x)(x^r-1).$$

As a result,

$$\bar{s}(x) = \frac{\lambda(x)(x^r - 1)}{\Psi_1^{\mathfrak{m}-\deg(\ell(x))}(\ell^{\dagger}(x))}$$

(3) Since

$$\begin{split} \kappa_1 \left(\frac{\ell(x)}{\gcd_r(\ell(x), s(x))} \right) \star (s(x), \kappa_1 t_1(x) + \kappa_2 t_2(x) + \kappa_3 t_3(x) + \kappa_4 t_4(x)) \\ &- \frac{s(x)}{\gcd_r(\ell(x), s(x))} \star (\ell(x), 0) \\ &= \left(0, \frac{\kappa_1 \ell(x) t_1(x)}{\gcd_r(\ell(x), s(x))} \right) \in \mathfrak{D}. \end{split}$$

Further, $\kappa_1 \star (\bar{s}(x), \kappa_1 \bar{t}_1(x) + \kappa_2 \bar{t}_2(x) + \kappa_3 \bar{t}_3(x) + \kappa_4 \bar{t}_4(x)) = (\bar{s}(x), \kappa_1 \bar{t}_1(x)) \in \mathfrak{D}^{\perp}$. Therefore,

$$(\bar{s}(x),\kappa_1\bar{t}_1(x))\ o\left(0,\kappa_1\frac{\ell(x)t_1(x)}{\gcd_r(\ell(x),s(x))}\right) = 0.$$

Using Lemma 4.8, we get

$$\kappa_1 \bar{t}_1(x) \Psi_2^{\mathfrak{m}-\deg\left(\frac{\ell(x)t_1(x)}{\gcd_r(\ell(x),s(x))}\right)} \left(\frac{\ell(x)t_1(x)}{\gcd_r(\ell(x),s(x))}\right)^{\dagger} \equiv 0 \pmod{x^s - 1}.$$

Hence, $\kappa_1 \overline{t}_1(x) \Psi_2^{\mathfrak{m}-\deg\left(\frac{\ell(x)t_1(x)}{\gcd_r(\ell(x),s(x))}\right)} \left(\frac{\ell(x)t_1(x)}{\gcd_r(\ell(x),s(x))}\right)^{\dagger} = \kappa_1 \gamma(x)(x^s-1), \text{ where } \gamma(x) \in R[x:\Theta].$ So,

$$\bar{t}_1(x) = \frac{\gamma(x)(x^s - 1)}{\Psi_2^{\mathfrak{m}-\deg\left(\frac{\ell(x)t_1(x)}{\gcd_r(\ell(x), s(x))}\right)} \left(\frac{\ell(x)t_1(x)}{\gcd_r(\ell(x), s(x))}\right)^{\dagger}}$$

Now, Proposition 4.10 implies that $\deg(\bar{t}_1(x)) = s - \deg(t_1(x)) - \deg(\ell(x)) + \deg(\gcd_r(\ell(x), s(x)))$, so $\gamma(x) = 1$.

Therefore,
$$\bar{t}_1(x) = \frac{x^s - 1}{\Psi_2^{\mathfrak{m} - \deg\left(\frac{\ell(x)t_1(x)}{\gcd_r(\ell(x), s(x))}\right)} \left(\frac{\ell(x)t_1(x)}{\gcd_r(\ell(x), s(x))}\right)^{\dagger}}.$$

(4) Since $\kappa_2 \star (\bar{s}(x), \kappa_1 \bar{t}_1(x) + \kappa_2 \bar{t}_2(x) + \kappa_3 \bar{t}_3(x) + \kappa_4 \bar{t}_4(x)) = (0, \kappa_2 \bar{t}_2(x)) \in \mathfrak{D}^{\perp}$. Then

 $(0, \kappa_2 \bar{t}_2(x)) \ o \ (s(x), \kappa_1 \bar{t}_1(x) + \kappa_2 \bar{t}_2(x) + \kappa_3 \bar{t}_3(x) + \kappa_4 \bar{t}_4(x)) \equiv 0 \pmod{x^{\mathfrak{m}} - 1}.$

Lemma 4.8 implies that

$$\kappa_2 \bar{t}_2(x) \Psi_2^{\mathfrak{m}-\deg(t_2(x))}(t_2^{\dagger}(x)) \equiv 0 \pmod{x^s - 1}.$$

So, for some $\nu(x) \in R[x:\Theta]$, we have

$$\kappa_2 \bar{t}_2(x) \Psi_1^{\mathfrak{m}-\deg t_2(x)}(t_2^{\dagger}(x)) = \nu(x)(x^s-1).$$

Next, by Proposition 4.10, it follows that $\deg(\overline{t}_2(x)) = s - \deg(t_2(x))$. Therefore, $\nu(x) = 1$. Consequently, we have

$$\bar{t}_2(x) = \frac{x^s - 1}{\Psi_2^{\mathfrak{m} - \deg t_2(x)}(t_2^{\dagger}(x))}$$

By using similar argument as above we can calculate,

$$\bar{t}_3(x) = \frac{x^s - 1}{\Psi_2^{\mathfrak{m} - \deg t_3(x)}(t_3^{\dagger}(x))}$$

and

$$\bar{t}_4(x) = \frac{x^s - 1}{\Psi_2^{\mathfrak{m} - \deg t_4(x)}(t_4^{\dagger}(x))}$$

5 QECCs from (θ, Θ) -cyclic codes over $\mathbb{F}_q R$

Quantum computers can potentially solve intricate problems at a significantly accelerated pace compared to classical computers. The field of quantum computing employs QECCs to shield quantum information from potential errors stemming from factors such as decoherence and quantum noise.

This section examines the method to construct QECCs from separable (θ, Θ) -cyclic code over $\mathbb{F}_q R$. Let $(\mathbb{C}^q)^{\otimes n}$ denote a q^n -dimensional Hilbert space over the complex field \mathbb{C} . Then a q-ary $[[n, k, d]]_q$ QECC is K-dimensional subspace of $(\mathbb{C}^q)^{\otimes n}$, where $k = \log_q K$ denote the dimension, n is length and d denote the Hamming distance of QECC, respectively.

Definition 5.1. [25](Quantum singleton bound) Consider \mathcal{Q} as an $[[n, k, d]]_q$ QECC. According to the Singleton bound, it follows that $2d \leq n - k + 2$. Additionally, \mathcal{Q} is termed a Maximum Distance Separable (MDS) QECC if 2d = n - k + 2.

We now present some results which will be useful in constructing QECCs.

Theorem 5.2. Let \mathfrak{D} be a linear code of block length (r, s) over $\mathbb{F}_q R$. Then $\Phi(\mathfrak{D}^{\perp}) = \Phi(\mathfrak{D})^{\perp}$. Moreover, $\Phi(\mathfrak{D})$ is self-dual if and only if \mathfrak{D} is self-dual.

Proof. Let $\mathbf{d}_1 = (g_0, g_1, \dots, g_{r-1}, b_0, b_1, \dots, b_{s-1}) \in \mathfrak{D}, \ \mathbf{d}_2 = (g'_0, g'_1, \dots, g'_{r-1}, b'_0, b'_1, \dots, b'_{s-1}) \in \mathfrak{D}^{\perp}$, where $b_i = \kappa_1 n_{1,i} + \kappa_2 n_{2,i} + \kappa_3 n_{3,i} + \kappa_4 n_{4,i}$ and $b'_i = \kappa_1 n'_{1,i} + \kappa_2 n'_{2,i} + \kappa_3 n'_{3,i} + \kappa_4 n'_{4,i}$ for $i = 0, 1, 2, \dots, s - 1$. Now,

$$d_1 \cdot d_2 = \kappa_1 \sum_{j=0}^{r-1} g_j g'_j + \sum_{i=0}^{s-1} b_i b'_i = 0,$$

which implies

$$\kappa_{1} \sum_{j=0}^{r-1} g_{j} g_{j}' + \sum_{i=0}^{s-1} (\kappa_{1} n_{1,i} + \kappa_{2} n_{2,i} + \kappa_{3} n_{3,i} + \kappa_{4} n_{4,i}) (\kappa_{1} n_{1,i}' + \kappa_{2} n_{2,i}' + \kappa_{3} n_{3,i}' + \kappa_{4} n_{4,i}') = 0$$

$$\kappa_{1} \sum_{j=0}^{r-1} g_{i} g_{i}' + \sum_{i=0}^{s-1} (\kappa_{1} n_{1,i} n_{1,i}' + \kappa_{2} n_{2,i} n_{2,i}' + \kappa_{3} n_{3,i} n_{3,i}' + \kappa_{4} n_{4,i} n_{4,i}') = 0$$

Now, by comparing the coefficients of κ_1 , κ_2 , κ_3 , and κ_4 on both the sides, we get

$$\sum_{j=0}^{r-1} g_j g'_j + \sum_{i=0}^{s-1} n_{1,i} n'_{1,i} = 0,$$
$$\sum_{i=0}^{s-1} n_{2,i} n'_{2,i} = 0,$$
$$\sum_{i=0}^{s-1} n_{3,i} n'_{3,i} = 0,$$
$$\sum_{i=0}^{s-1} n_{4,i} n'_{4,i} = 0.$$

Next, we have

$$\Phi(d_1) \cdot \Phi(d_2) = \sum_{j=0}^{r-1} g_j g'_j + \sum_{i=0}^{s-1} b_i M M^T b'_i,$$

=
$$\sum_{j=0}^{r-1} g_j g'_j + \sum_{i=0}^{s-1} (n_{1,i} n'_{1,i} + n_{2,i} n'_{2,i} + n_{3,i} n'_{3,i} + n_{4,i} n'_{4,i}).$$

Since $MM^T = I_{4\times 4}$. Thus, from the above equation, we have $\sum_{j=0}^{r-1} g_j g'_j + \sum_{i=0}^{s-1} (n_{1,i}n'_{1,i} + n_{2,i}n'_{2,i} + n_{3,i}n'_{3,i} + n_{4,i}n'_{4,i}) = 0$. This implies that $\Phi(\mathbf{d}_2) \in \Phi(\mathfrak{D})^{\perp}$ for $\Phi(\mathbf{d}_1) \in \Phi(\mathfrak{D})$. Therefore, $\Phi(\mathfrak{D}^{\perp}) \subseteq \Phi(\mathfrak{D})^{\perp}$. Since Φ is a bijection, so we have $|\Phi(\mathfrak{D}^{\perp})| = |\Phi(\mathfrak{D})^{\perp}|$. Thus, we can conclude that $\Phi(\mathfrak{D}^{\perp}) = \Phi(\mathfrak{D})^{\perp}$. Furthermore, if \mathfrak{D} is a self-dual code, then $\Phi(\mathfrak{D}) = \Phi(\mathfrak{D}^{\perp}) = \Phi(\mathfrak{D})^{\perp}$. Consequently, we can conclude that $\Phi(\mathfrak{D})$ is also self-dual.

To construct QECCs in our setup, we present a well-known result, called CSS construction, discussed by Calderbank, Shor and Stean in 1998.

Theorem 5.3. [12, CSS construction] Let $\mathfrak{D}_1 = [n, k_1, d_1]$ and $\mathfrak{D}_2 = [n, k_2, d_2]$ are linear codes over \mathbb{F}_q with $\mathfrak{D}_2^{\perp} \subseteq \mathfrak{D}_1$. Then a $[[n, k_1 + k_2 - n, min\{d_1, d_2\}]]_q$ QECC exists. Particularly, if $\mathfrak{D}_1^{\perp} \subseteq \mathfrak{D}_1$, then a $[[n, 2k_1 - n, d_1]]_q$ QECC can be constructed.

Our next theorems present the dual containing property of θ -cyclic codes over \mathbb{F}_q , Θ -cyclic codes over R and separable (θ, Θ) -cyclic code over $\mathbb{F}_q R$. Here, we consider $\operatorname{ord}(\theta) \mid r$ and $\operatorname{ord}(\Theta) \mid s$.

Theorem 5.4. [15] Let $\mathfrak{C}_r = \langle \ell(x) \rangle$ be a θ -cyclic code of length r over \mathbb{F}_q . Then $\mathfrak{C}_r^{\perp} \subseteq \mathfrak{C}_r$ if and only if $x^r - 1 \mid_r f^{\dagger}(x) f(x)$, where $x^r - 1 = f(x)\ell(x)$.

Theorem 5.5. Suppose $\mathfrak{C}_s = \langle \kappa_1 t_1 + \kappa_2 t_2 + \kappa_3 t_3 + \kappa_4 t_4 \rangle$ is a Θ -cyclic code of length s over R. Then $\mathfrak{C}_s^{\perp} \subseteq \mathfrak{C}_s$ if and only if $x^s - 1 \mid_r h_i^{\dagger}(x)h_i(x)$, where $h_i(x)t_i(x) = x^s - 1$ for i = 1, 2, 3, 4.

Proof. We know that $\mathfrak{C}_s = \kappa_1 \mathfrak{C}_{s,1} \oplus \kappa_2 \mathfrak{C}_{s,2} \oplus \kappa_3 \mathfrak{C}_{s,3} \oplus \kappa_4 \mathfrak{C}_{s,4}$, is Θ -cyclic code of length s over R if and only if each $\mathfrak{C}_{s,i}$ is θ -cyclic code of length s over \mathbb{F}_q . Also, $\mathfrak{C}_s^{\perp} \subseteq \mathfrak{C}_s$ if and only if $\mathfrak{C}_{s,i}^{\perp} \subseteq \mathfrak{C}_{s,i}$. Further $\mathfrak{C}_{s,i} = \langle t_i(x) \rangle$ such that $x^s - 1 = h_i(x)t_i(x)$ for i = 1, 2, 3, 4. Now, Theorem 5.4 infer that $\mathfrak{C}_{s,i}^{\perp} \subseteq \mathfrak{C}_{s,i}$ if and only if $x^s - 1 \mid_r h_i^{\dagger}(x)h_i(x)$.

Theorem 5.6. Suppose \mathfrak{D} is a separable (θ, Θ) -cyclic code over $\mathbb{F}_q R$. Then $\mathfrak{D}^{\perp} \subseteq \mathfrak{D}$ if and only if $\mathfrak{C}_r^{\perp} \subseteq \mathfrak{C}_r$ and $\mathfrak{C}_s^{\perp} \subseteq \mathfrak{C}_s$.

Proof. Assume $\mathfrak{D}^{\perp} \subseteq \mathfrak{D} = \mathfrak{C}_r \times \mathfrak{C}_s$, then $\mathfrak{C}_r^{\perp} \times \mathfrak{C}_s^{\perp} \subseteq \mathfrak{C}_r \times \mathfrak{C}_s$. Which implies $\mathfrak{C}_r^{\perp} \subseteq \mathfrak{C}_r$ and $\mathfrak{C}_s^{\perp} \subseteq \mathfrak{C}_s$.

Same argument can be used to prove the converse.

Now, by considering the above discussion, we present the key result.

Theorem 5.7. Let $\mathfrak{D} = \mathfrak{C}_r \times \mathfrak{C}_s$ be a separable (θ, Θ) -cyclic code of block length (r, s) over $\mathbb{F}_q R$. If $\mathfrak{C}_r^{\perp} \subseteq \mathfrak{C}_r$ and $\mathfrak{C}_s^{\perp} \subseteq \mathfrak{C}_s$, then exists a $[[r+4s, 2k-(r+4s), d_H]]_q$ QECC over \mathbb{F}_q , where k is dimension and d_H is Hamming distance of $\Phi(\mathfrak{D})$.

Proof. Suppose $\mathfrak{C}_r^{\perp} \subseteq \mathfrak{C}_r$ and $\mathfrak{C}_s^{\perp} \subseteq \mathfrak{C}_s$, then by Theorem 5.6, we get $\mathfrak{D}^{\perp} \subseteq \mathfrak{D}$. Now, by Theorem 5.2, we have $|\Phi(\mathfrak{D}^{\perp})| = |\Phi(\mathfrak{D})^{\perp}|$, then $\Phi(\mathfrak{D})^{\perp} \subseteq \Phi(\mathfrak{D})$. Again by Theorem 5.3, this infer that there exists a $[[r+4s, 2k-(r+4s), d_H]]_q$ QECC over \mathbb{F}_q .

In order to support our findings, we present some illustrative examples of constructing QECCs.

Example 5.8. Let q = 9, r = 26, s = 6 and $R = \mathbb{F}_9 + u\mathbb{F}_9 + v\mathbb{F}_9 + uv\mathbb{F}_9$, where $\mathbb{F}_9 = \mathbb{F}_3[\omega]$ with $\omega^2 + 2\omega + 2 = 0$. The Frobenius automorphism $\theta : \mathbb{F}_9 \to \mathbb{F}_9$ is defined as $\theta(\alpha) = \alpha^3$ for all $\alpha \in \mathbb{F}_9$. We have $\operatorname{ord}(\theta) = 2 = \operatorname{ord}(\Theta)$ such that $\operatorname{ord}(\theta) \mid r$, $\operatorname{ord}(\Theta) \mid s$. Since $\mathbb{F}_9[x : \theta]$ is not a UFD, so both the polynomials $x^{26} - 1$ and $x^6 - 1$ have more than one factorization. Let us delve into one of these factors.

$$\begin{aligned} x^{26} - 1 &= (x^3 + \omega x^2 + 2x\omega + 2)(x^3 + \omega^7 x^2 + 2x + 1)(x^3 + 2x^2 + \omega^7 x + 1)(x^3 + x^2 + \omega^7 x + 2) \times \\ &\quad (x^3 + \omega x^2 + \omega^6 x + 1)(x^3 + \omega^7 x^2 + \omega^6 x + 2)(x^3 + x^2 + \omega^3 x + \omega^6) \times \\ &\quad (x^3 + 2x^2 + \omega^3 x + \omega^2)(x + \omega^2)^2 \end{aligned}$$

and

$$x^{6} - 1 = (x + \omega^{6})(x + 1)(x + \omega^{2})^{2}(x + 2)(x + \omega^{2}).$$

Let $\ell(x) = (x^3 + 2x^2 + \omega^3 x + \omega^2)(x + \omega^2)$. Then

$$\begin{split} f(x) &= x^{22} + \omega^7 x^{21} + \omega^6 x^{20} + \omega^3 x^{19} + \omega^6 x^{18} + \omega^2 x^{16} + \omega^3 x^{15} + 2x^{14} + \omega^7 x^{13} + \omega^3 x^{12} + \\ & \omega^7 x^{11} + x^{10} + \omega^3 x^9 + \omega^2 x^8 + \omega^7 x^7 + \omega x^6 + \omega^3 x^5 + \omega^3 x^3 + \omega^7 x^2 + \omega^3 x + 1 \\ f^{\dagger}(x) &= x^{22} + \omega x^{21} + \omega^7 x^{20} + \omega x^{19} + \omega x^{17} + \omega x^{16} + \omega^5 x^{15} + \omega^2 x^{14} + \omega x^{13} + x^{12} + \omega^5 x^{11} + \\ & \omega^3 x^{10} + \omega^5 x^9 + 2x^8 + \omega x^7 + \omega^2 x^6 + \omega^6 x^4 + \omega x^3 + \omega^6 x^2 + \omega^5 x + 1 \\ f^{\dagger}(x) f(x) &= (x^{18} + \omega^6 x^{17} + \omega^5 x^{16} + \omega^2 x^{15} + \omega^7 x^{14} + \omega^2 x^{12} + 2x^{10} + \omega^2 x^9 + x^8 + \omega^6 x^6 + \\ & \omega^3 x^4 + \omega^2 x^3 + \omega x^2 + \omega^6 x + 1)(x^{26} - 1). \end{split}$$

Next, let us choose $t_1(x) = (x+1)(x+\omega^6)$, then

$$s_1(x) = x^4 + \omega^3 x^3 + \omega x^2 + \omega^7 x + \omega^6$$

$$s_1^{\dagger}(x) = \omega^6 x^4 + \omega^5 x^3 + \omega x^2 + \omega x + 1$$

$$s_1^{\dagger}(x)s_1(x) = (\omega^6 x^2 + \omega^2)(x^6 - 1).$$

Finally, we choose $t_i(x) = (x + w^6)$ for i = 2, 3, 4, then

$$s_i(x) = x^5 + \omega^6 x^4 + x^3 + \omega^6 x^2 + x + \omega^6$$

$$s_i^{\dagger}(x) = \omega^2 x^5 + x^4 + \omega^2 x^3 + x^2 + \omega^2 x + 1$$

$$s_i^{\dagger}(x)s_i(x) = (\omega^2 x^4 + \omega^2 x^2 + \omega^2)(x^6 - 1).$$

Here, $\mathfrak{C}_r = \langle \ell(x) \rangle$ is a θ -cyclic code having parameters [26, 22, 3] over \mathbb{F}_9 and $\mathfrak{C}_s = \langle \kappa_1 t_1(x) + \kappa_2 t_2(x) + \kappa_3 t_3(x) + \kappa_4 t_4(x) \rangle$ is a Θ -cyclic code over R such that $\Phi_1(\mathfrak{C}_s)$ has parameters [24, 19, 3] over \mathbb{F}_9 . Therefore, by Proposition 2.8, we get $\Phi(\mathfrak{D})$ is a [50, 41, 3] linear code \mathbb{F}_9 . From the above equation, we have $x^{26} - 1 \mid_r (f^{\dagger}(x)f(x))$ and $x^6 - 1 \mid_r (s_i^{\dagger}(x)s_i(x))$ for i = 1, 2, 3, 4. Therefore, by Theorems 5.4 and 5.5, we get that $\mathfrak{C}_r^{\perp} \subseteq \mathfrak{C}_r$ and $\mathfrak{C}_s^{\perp} \subseteq \mathfrak{C}_s$. Hence, from Theorem 5.6, we conclude that $\mathfrak{D}^{\perp} \subseteq \mathfrak{D}$. Thus, Theorems 5.7 enables us to construct a [[50, 32, 3]]_9 QECC. This QECC has better parameters than [[50, 30, 3]]_9 as given in [18].

Example 5.9. Let q = 25, r = 8, s = 10 and $R = \mathbb{F}_{25} + u\mathbb{F}_{25} + uv\mathbb{F}_{25}$, where $\mathbb{F}_{25} = \mathbb{F}_5[\omega]$ with $\omega^2 + 4\omega + 2 = 0$. The Frobenius automorphism $\theta : \mathbb{F}_{25} \to \mathbb{F}_{25}$ is defined as $\theta(\alpha) = \alpha^5$ for all $\alpha \in \mathbb{F}_{25}$. We have $\operatorname{ord}(\theta) = 2 = \operatorname{ord}(\Theta)$ such that $\operatorname{ord}(\theta) \mid r$, $\operatorname{ord}(\Theta) \mid s$. Since $\mathbb{F}_{25}[x : \theta]$ is not a UFD, so both the polynomials $x^8 - 1$ and $x^{10} - 1$ have more than one factorization. Let us delve into one of these factors.

$$x^{8} - 1 = (x+1)(x+4)(x+\omega^{9})^{2}(x+\omega^{3})^{2}(x+3)(x+2)$$

and

$$x^{10} - 1 = (x+4)(x+\omega^{20})(x+\omega^{16})(x+4)(x+1)^2(x+\omega^8)(x+\omega^{16})(x+\omega^{20})(x+\omega^4)(x+\omega^{16})$$

Let $\ell(x) = (x + \omega^9)(x + 2)$. Then

$$f(x) = x^{6} + \omega^{4}x^{5} + \omega^{10}x^{4} + \omega^{4}x^{3} + \omega^{22}x + \omega^{21}$$

$$f^{\dagger}(x) = \omega^{21}x^{6} + \omega^{14}x^{5} + \omega^{5}x^{4} + \omega^{20}x^{3} + \omega^{10}x^{2} + \omega^{20}x + 1$$

$$f^{\dagger}(x)f(x) = (\omega^{21}x^{4} + 2x^{3} + 3x + w^{9})(x^{8} - 1).$$

Next, let us choose $t_1(x) = (x + \omega^8)(x + \omega^{16})(x + \omega^{20})$, then

$$s_1(x) = x^7 + \omega^4 x^6 + 2x^5 + \omega^{10} x^4 + 3x^3 + \omega^{22} x^2 + 4x + \omega^{16}$$

$$s_1^{\dagger}(x) = \omega^8 x^7 + 4x^6 + \omega^{14} x^5 + 3x^4 + \omega^2 x^3 + 2x^2 + \omega^{20} x + 1$$

$$s_1^{\dagger}(x)s_1(x) = (\omega^8 x^4 + \omega^8 x^3 + \omega^{15} x^2 + \omega^{20} x + \omega^4)(x^{10} - 1).$$

Finally, we choose $t_i(x) = (x + w^{20})$ for i = 2, 3, 4, then

$$s_{i}(x) = x^{9} + \omega^{16}x^{8} + x^{7} + \omega^{16}x^{6} + x^{5} + \omega^{16}x^{4} + x^{3} + \omega^{16}x^{2} + x + \omega^{16}$$

$$s_{i}^{\dagger}(x) = \omega^{8}x^{9} + x^{8} + \omega^{8}x^{7} + x^{6} + \omega^{8}x^{5} + x^{4} + \omega^{8}x^{3} + x^{2} + \omega^{8}x + 1$$

$$s_{i}^{\dagger}(x)s_{i}(x) = (\omega^{8}x^{8} + \omega^{20}x^{7} + \omega^{7}x^{6} + \omega^{2}x^{5} + \omega^{21}x^{4} + \omega^{14}x^{3} + \omega^{23}x^{2} + \omega^{8}x + \omega^{4})(x^{10} - 1).$$

Here, $\mathfrak{C}_r = \langle \ell(x) \rangle$ is a θ -cyclic code having parameters [8, 6, 3] over \mathbb{F}_{25} , $\mathfrak{C}_s = \langle \kappa_1 t_1(x) + \kappa_2 t_2(x) + \kappa_3 t_3(x) + \kappa_4 t_4(x) \rangle$ is a Θ -cyclic code over R such that $\Phi_1(\mathfrak{C}_s)$ has parameters [40, 34, 3] over \mathbb{F}_{25} . Therefore by Proposition 2.8, we get $\Phi(\mathfrak{D})$ is a [48, 40, 3] linear code over \mathbb{F}_{25} . From the above, we have $x^8 - 1 \mid_r (f^{\dagger}(x)f(x))$ and $x^{10} - 1 \mid_r (s_i^{\dagger}(x)s_i(x))$ for i = 1, 2, 3, 4. Therefore, by Theorems 5.4 and 5.5, we get that $\mathfrak{C}_r^{\perp} \subseteq \mathfrak{C}_r$ and $\mathfrak{C}_s^{\perp} \subseteq \mathfrak{C}_s$. Hence, from Theorem 5.6, we get that $\mathfrak{D}^{\perp} \subseteq \mathfrak{D}$. Thus, Theorem 5.7 enables us to construct a [[48, 32, 3]]_{25} QECC. This QECC has better parameters than [[48, 16, 3]]_{25} as given in [14].

Example 5.10. Let $q = 27, r = 9, s = 3, R = \mathbb{F}_{27} + u\mathbb{F}_{27} + u\mathbb{F}_{27} + uv\mathbb{F}_{27}$, where $\mathbb{F}_{27} = \mathbb{F}_3[\omega]$ with $\omega^3 + 2\omega + 2 = 0$ and Frobenius automorphism $\theta(\alpha) = \alpha^3$ for all $\alpha \in \mathbb{F}_{27}$. We have order of θ , Θ is 3 such that $\operatorname{ord}(\theta) \mid r$, $\operatorname{ord}(\Theta) \mid s$. Since $\mathbb{F}_{27}[x : \theta]$ is not a UFD, so both the polynomials $x^9 - 1$ and $x^3 - 1$ have more than one factorization. Let us delve into one of these factorizations.

$$x^9 - 1 = (x + \omega^{21})(x + \omega^9)(x + \omega^{15})(x + \omega^5)(x + \omega^{25})(x + \omega)(x + \omega^{27})(x + \omega^{21})(x + 2)$$

and

$$x^{3} - 1 = (x + \omega^{9})(x + \omega^{3})(x + \omega)$$

Let $\ell(x) = (x + \omega^9)(x + 2)$. Then

$$\begin{split} f(x) &= x^7 + \omega^{16} x^6 + \omega^2 x^5 + \omega^7 x^4 + \omega^{12} x^3 + \omega^{15} x^2 + \omega^{17} x + \omega^{17} \\ f^{\dagger}(x) &= \omega^{25} x^7 + \omega^{17} x^6 + \omega^5 x^5 + \omega^{10} x^4 + \omega^7 x^3 + \omega^{18} x^2 + \omega^{22} x + 1 \\ f^{\dagger}(x) f(x) &= (\omega^{25} x^5 + \omega^9 x^4 + \omega^{17} x^3 + \omega^{12} x^2 + \omega^{22} x + \omega^4) (x^9 - 1). \end{split}$$

Next, let us choose $t_i(x) = (x + \omega)$ for i = 1, 2, then

$$s_i(x) = x^2 + \omega^{22}x + \omega^{12}$$

$$s_i^*(x) = \omega^4 x^2 + \omega^{14} x + 1$$

$$s_i^{\dagger}(x)s_i(x) = (\omega^4 x + \omega^{25})(x^{10} - 1)$$

Finally, let us choose $t_j(x) = (x + w^3)$ for j = 3, 4 then

$$s_j(x) = x^2 + \omega^{14}x + \omega^{10}$$

$$s_j^{\dagger}(x) = \omega^{12}x^2 + \omega^{16}x + 1$$

$$s_j^*(x)s_j(x) = (\omega^{12}x + \omega^{23})(x^3 - 1).$$

Here, $\mathfrak{C}_r = \langle \ell(x) \rangle$ is a θ -cyclic code over having parameters [9,7,3] over \mathbb{F}_{27} , $\mathfrak{C}_s = \langle \kappa_1 t_1 + \kappa_2 t_2 + \kappa_3 t_3 + \kappa_4 t_4 \rangle$ is a Θ -cyclic code over R such that $\Phi_1(\mathfrak{C}_s)$ has parameters [12,8,3] over \mathbb{F}_{27} . Then, $\Phi(\mathfrak{D})$ is a [21,15,3] linear code over \mathbb{F}_{27} . From the above, we have $x^9 - 1 \mid_r (f^{\dagger}(x)f(x))$ and $x^3 - 1 \mid_r (s_i^{\dagger}(x)s_i(x))$ for i = 1, 2, 3, 4. Therefor, by Theorems 5.4 and 5.5, we can conclude that $\mathfrak{C}_r^{\perp} \subseteq \mathfrak{C}_r$ and $\mathfrak{C}_s^{\perp} \subseteq \mathfrak{C}_s$. Moreover, Theorem 5.6 leads to the conclusion that $\mathfrak{D}^{\perp} \subseteq \mathfrak{D}$. Hence, Theorem 5.7 enables us to construct a QECC with parameters [[21,9,3]]_{27}.

As an application of our study, Table 2 showcases several MDS QECCs obtained from θ cyclic codes over \mathbb{F}_q . The coefficients of generator polynomials are presented in ascending order. For instance, $\omega^7 \omega^5 21$ corresponds to the polynomial $w^7 + w^5 x + 2x^2 + x^3$. In Table 3, we obtain some new QECCs from Θ -cyclic code over R, which have better parameters than the existing quantum codes. Furthermore, in Table 4, we construct some new QECCs from (θ, Θ) -cyclic code over $\mathbb{F}_q R$.

q	n	$\ell(x)$	$[[n,k,d]]_q$
3^{2}	6	$\omega^6 \omega^7 1$	$[[6, 2, 3]]_{3^2}$
3^4	8	$\omega^{69}\omega^{28}1$	$[[8, 4, 3]]_{3^4}$
5^{2}	8	$\omega^{23}\omega^{10}1$	$[[8, 4, 3]]_{5^2}$
13^{2}	8	$\omega^{117}\omega^{109}1$	$[[8, 4, 3]]_{13^2}$
3^{3}	9	$\omega^{18}\omega^{19}1$	$[[9, 5, 3]]_{3^3}$
5^{3}	12	$\omega^{99}\omega^{38}1$	$[[12, 8, 3]]_{5^3}$
5^{3}	15	$\omega^2 3 \omega^{76} 1$	$[[15, 9, 4]]_{5^3}$
7^{5}	15	$\omega^{295}\omega^{96}$	$[[15, 11, 3]]_{7^5}$
3^{3}	18	$\omega^{18}\omega^{19}1$	$[[18, 14, 3]]_{3^3}$
7^{3}	18	$\omega^{339}\omega^{212}1$	$[[18, 14, 3]]_{7^3}$
5^{4}	20	$\omega^{332}\omega^{292}1$	$[[20, 16, 3]]_{5^4}$
7^{3}	21	$\omega^{261}\omega^{17}\omega^{178}1$	$[[21, 15, 4]]_{7^3}$
5^{5}	25	$\omega^{1374}\omega^{3116}\omega^{3015}1$	$[[25, 19, 4]]_{5^5}$
13^2	26	$1\omega^{21}\omega^{148}1$	$[[26, 20, 4]]_{13^2}$

Table 2: MDS QECCs from θ -cyclic code over \mathbb{F}_q

q	n	$t_1(x)$	$t_2(x)$	$t_3(x)$	$t_4(x)$	$\Phi_1(\mathfrak{C}_s)$	New QECCS	Existing QECCs
9	6	$1 \ \omega^5 \omega^5 1$	$\omega^2 1$	$\omega^6 1$	11	[24, 18, 4]	$[[24, 12, 4]]_9$	$[[24, 10, 4]]_9[23]$
9	6	$\omega^6 \omega^3 1$	21	$\omega^2 1$	$\omega^6 1$	[24, 19, 3]	$[[24, 14, 3]]_9$	$[[24, 8, 2]]_9[2]$
27	9	$\omega^{18}\omega^{19}1$	ω^{91}	$\omega^{3}1$	1	[36, 32, 3]	$[[36, 28, 3]]_{27}$	$[[36, 20, 3]]_{27}[29]$
25	10	$\omega^4 4 \omega^{16} 1$	$\omega^{20}1$	$\omega^{8}1$	41	[40, 34, 3]	$[[40, 28, 3]]_{25}$	$[[40, 24, 3]]_{25}[5]$
9	12	$\omega^6 \omega^3 \omega^5 \omega^3 1$	$\omega^{3}1$	$\omega^{3}1$	$\omega^{3}1$	[48, 41, 3]	$[[48, 34, 3]]_9$	$[[48, 30, 3]]_9[28]$
9	12	$\omega^3 \omega 121$	$\omega^{6}1$	11	1	[48, 42, 4]	$[[48, 36, 4]]_9$	$[[48, 34, 4]]_9[29]$
49	14	$\omega^{18}\omega^{15}1$	$\omega^{6}1$	$\omega^6 1$	$\omega^6 1$	[56, 51, 3]	$[[56, 48, 3]]_{49}$	$[[56, 44, 3]]_{49}[28]$
49	14	$6 \omega \omega^{46} \omega^{47} 1$	$\omega^{18}1$	$\omega^6 1$	11	[56, 49, 4]	$[[56, 42, 4]]_{49}$	$[[56, 40, 4]]_{49}[28]$
9	18	$\omega^6 \omega^3 \omega^2 \ 1 \ 1$	$\omega^6 1$	$\omega^2 1$	1	[72, 66, 3]	$[[72, 60, 3]]_9$	$[[72, 54, 3]]_9[29]$
121	22	$\omega^{100}\omega^{80}\omega^{38}\omega^{20}1$	$\omega^{40}1$	$\omega^{80}1$	11	[88, 81, 4]	$[[88, 74, 4]]_{121}$	$[[80, 64, 4]]_{121}[5]$
25	20	$\omega^{22} 3 \omega^{13} 1$	$\omega^{22}1$	$\omega^{8}1$	1	[80, 75, 3]	$[[80, 70, 3]]_{25}$	$[[80, 56, 3]]_{25}[5]$
121	$\overline{22}$	$\omega^{20}\omega^{107}\omega^{32}1$	11	$\omega^{80}1$	$\omega^{40}1$	[88, 82, 3]	$[[88, 76, 3]]_{121}$	$[[88, 72, 3]]_{121}[5]$
169	26	$1\omega^{148}\omega^{21}1$	$\omega^{60}1$	$\omega^{48}1$	1	[104, 99, 4]	$[[104, 94, 4]]_{169}$	$[[104, 80, 4]]_{169}[5]$

Table 3: New QECCs constructed from Θ -cyclic code \mathfrak{C}_s over R

q	(r,s)	$\ell(x)$	$t_1(x)$	$t_2(x)$	$t_3(x)$	$t_4(x)$	$\Phi(\mathfrak{D})$	New QECCS	Existing QECCs
9	(12, 6)	$\omega^3 \omega 121$	$1 \ \omega^5 \omega^5 1$	$\omega^2 1$	$\omega^6 1$	ω	[36, 27, 4]	$[[36, 18, 4]]_9$	$[[36, 6, 4]]_9[18]$
25	(8, 8)	$\omega^{23}\omega^{10}1$	$\omega^{23}\omega^{10}1$	ω^5	$\omega^{3}1$	1	[40, 34, 3]	$[[40, 28, 3]]_{25}$	$[[40, 24, 3]]_{25}[5]$
27	(18, 9)	$\omega^{18}\omega^{19}1$	$\omega^{18}\omega^{19}1$	$\omega^{9}1$	$\omega^{3}1$	1	[54, 48, 3]	$[[54, 42, 3]]_{27}$	$[[54, 36, 3]]_{27}[29]$
25	(40, 10)	$2\omega^4\omega^81$	$\omega^{20}4\omega^81$	$\omega^{20}1$	$\omega^{20}1$	1	[80, 72, 3]	$[[80, 64, 3]]_{25}$	$[[80, 56, 3]]_{25}[5]$
25	(8, 10)	$\omega^{15}\omega^{16}1$	$\omega^{20}4\omega^81$	$\omega^{20}1$	$\omega^{20}1$	1	[48, 41, 3]	$[[48, 34, 3]]_{25}$	$[[48, 16, 3]]_{25}[14]$
9	(56, 6)	$1022\omega^6 1$	$\omega^6 \omega^3 1$	21	$\omega^2 1$	$\omega^6 1$	[80, 70, 4]	$[[80, 60, 4]]_9$	$[[80, 48, 4]]_9[2]$
169	(8, 24)	$\omega^{117}\omega^{109}1$	$\omega^{15}\omega^{3}1$	(11)1	$\omega^{85}1$	1	[104, 98, 3]	$[[104, 92, 3]]_{169}$	$[[104, 88, 3]]_{169}[5]$

Table 4: New QECCs constructed from (θ, Θ) -cyclic code \mathfrak{D} over $\mathbb{F}_q R$

6 Conclusion

In summary, this paper investigates the algebraic structure of (θ, Θ) -cyclic codes with block length (r, s) over the ring $\mathbb{F}_q R$, where $R = \mathbb{F}_q + u\mathbb{F}_q + v\mathbb{F}_q + uv\mathbb{F}_q$ with $u^2 = u, v^2 = v, uv = vu$, and q is an odd prime power. Our exploration begins with the decomposition of the ring R into idempotent, followed by an in-depth examination of linear codes over the ring $\mathbb{F}_q R$. We define a Gray map from $\mathbb{F}_q^r \times \mathbb{R}^s \to \mathbb{F}_q^{r+4s}$ and investigate its fundamental properties. The algebraic structure of (θ, Θ) -cyclic codes with block length (r, s) over $\mathbb{F}_q R$ is thoroughly determined, and we establish generator polynomials for this family of codes in Theorem 3.12. The algebraic structure of separable codes is discussed in detail by Theorem 3.16, and the minimal generating set of (θ, Θ) -cyclic codes over $\mathbb{F}_q R$ is presented in Theorem 3.19. The relationship between generator polynomials of (θ, Θ) -cyclic codes and their duals is established in Theorem 4.11. As an application of our findings, we demonstrate the construction of QECCs from separable (θ, Θ) cyclic codes over $\mathbb{F}_q R$ in Theorem 5.7. Additionally, detailed examples (Examples 5.8, 5.9, 5.10) are provided to illustrate the process of constructing QECCs. Our study yields optimal and near-optimal codes from Θ -cyclic codes over R (Table 1), MDS QECCs from θ -cyclic codes over \mathbb{F}_q (Table 2), and new QECCs with improved parameters compared to existing codes (Tables 3 and 4).

References

- T. Abualrub, I. Siap and N. Aydin, Z₂Z₄-Additive Cyclic Codes, *IEEE Trans. Inform.* Theory 60(3), pp. 1508–1514 (2014).
- [2] M. Ashraf and G. Mohammad, Quantum codes from cyclic codes over $\mathbb{F}_q + u\mathbb{F}_q + v\mathbb{F}_q + uv\mathbb{F}_q$, Quantum Inf. Process. 15(10), pp. 4089-4098 (2016).
- [3] I. Aydogdu, R. M. Hesari and K. Samei, Double skew cyclic codes over \mathbb{F}_q , Comput. Appl. Math. 41(3), pp. 1-17 (2022).
- [4] I. Aydogdu and I. Siap, The structure of Z₂Z^s₂-additive codes: Bounds on the Minimum Distance, Appl. Math. Inf. Sci. 7(6), pp. 2271-2278 (2013).
- [5] T. Bag, H. Q. Dinh, A. K. Upadhyay, R. K. Bandi and W. Yamaka, Quantum codes from skew constacyclic codes over the ring $\mathbb{F}_q[u, v]/\langle u^2 1, v^2 1, uv vu \rangle$, Discrete Math. 343(3), pp. 1-17 (2020).
- [6] J. Borges, C. Fernández-Córdoba, J. Pujol, J. Rifá and M. Villanueva, Z₂Z₄-linear codes: generator matrices and duality, *Des. Codes Cryptogr.* 54, pp. 167-179 (2010).
- [7] J. Borgers, C. Fernández-Córdob and R. Ten-valls, Z₂-double cyclic codes, *Des. codes cryptogr.* 86, pp. 463-479 (2018).
- [8] D. Boucher, W. Geiselmann and F. Ulmer, Skew-cyclic codes, Appl. Algebra Eng. Commun. Comput. 18, pp. 379-389 (2007).
- [9] D. Boucher, and F. Ulmer, Coding with skew polynomial rings, J. Symbolic Comput. 44(12), pp. 1644-1656 (2009).
- [10] A. E. Brouwer, H.O. Hamalainen, P. R. Ostergard and N. J. A. Sloane, Boundes on mixed binary/ternary codes, *IEEE Trans. Inform. Theory* 44(1), pp. 140-161 (1998).
- [11] A. R. Calderbank, A. R. Hammons, P. V. Kumar, N. J. A. Sloane and P. Solé, The Z₄linearity of Kerdock, Preparata, Goethals and related codes, *IEEE Trans. Inform. Theory* 40(2), pp. 301-309 (1994).

- [12] A. R. Calderbank, E. M. Rains, P. M. Shor and N. J. A. Sloane, Quantum error correction via codes over GF(4), *IEEE Trans. Inform. Theory* 44(4), pp. 1369-1387 (1998).
- [13] L. Chausde, P. Laidreau and F. Ulmer, skew cyclic code of prescribed distance or rank, Des. Codes Cryptogr. 50(3), pp. 267-284 (2009).
- [14] H. Q. Dinh, T. Bag, K. Abdukhalikov, S. Pathak, A. K. Upadhyay, R. Bandi and W. Chinnakum, On a class of skew constacyclic codes over mixed alphabets and applications in constructing optimal and quantum codes, *Cryptogr. Commun.* 15(1), pp. 171-198 (2023).
- [15] H. Q. Dinh, T. Bag, A. K. Upadhyay, R. K. Bandi and R. Tansuchat, A class of skew cyclic code and application in quantum codes construction, *Discrete Math.* 344(2), pp. 1-13 (2021).
- [16] H. Q. Dinh, S. Pathak, T. Bag, A. K. Upadhyay and W. Chinnakum, A Study of $\mathbb{F}_q R$ -cyclic codes and their applications in constructing quantum codes, *IEEE Access* 8, pp. 190049-190063 (2020).
- [17] BR. Mc Donald, Finite ring with identity, Marcel Dekker, New york, 1974.
- [18] Y. Edel, Some good quantum twisted codes (2020). https://www.mathi.uniheidelberg.de/ yves/Matritzen/QTBCH/QTBCHTab9.html
- [19] J. Gao, Skew cyclic codes over $\mathbb{F}_p + v\mathbb{F}_p$, J. Appl. Math. Inform. 31(3-4), pp. 337-342 (2013).
- [20] J. Gao, M. J. Shi, T. Wu and F. W. Fu, On double cyclic codes ove Z4, *Finite Fields Appl.* 39, pp. 233–250 (2016).
- [21] F. Gursoy, I. Siap and B. Yildiz, Construction of skew cyclic codes over $\mathbb{F}_q + v\mathbb{F}_q$, Adv. Math. Commun. 8(3), pp. 313-322 (2014).
- [22] R. M. Hesari, R. Rezaei and K. Samei, On self-dual skew cyclic codes of length p^s over $\mathbb{F}_{p^m} + u\mathbb{F}_{p^m}$, Discrete Math. 344(11), pp. 1-16 (2021).
- [23] H. Islam, O. Prakash and R. K. Verma, New quantum codes from constacyclic codes over the ring $R_{k,m}$, Adv. Math. Commun. 16(1), pp. 17-35 (2022).
- [24] Jitmans, Lingss and P. Udomkavanich, Skew constacyclic codes over finite chain ring, Adv. Math. Commun. 6, pp. 39-62 (2012).
- [25] A. Ketkar, A. Klappenecker, S. Kumar and P. K. Sarvepalli, Nonbinary Stabilizer Codes Over Finite Fields, *IEEE Trans. Inf. Theory* 52(11), pp. 4892-4914 2006.
- [26] O. Ore, Theory of non-commutative polynomials, Ann. of Math. pp. 480-508, (1933).
- [27] M. Özen, N. Tuğba Özzaim and H. İnce, Skew quasi cyclic codes over $\mathbb{F}_q + u\mathbb{F}_q$, J. Algebra Appl. 18(4), (2019).
- [28] O. Prakash, H. Islam, S. Patel and et al. New Quantum Codes from Skew Constacyclic Codes Over a Class Of Non-Chain Rings $R_{e, q}$, Int. J. Theor. Phys. 60 (3), pp. 334–3352 (2021).
- [29] P. Rai, B. Singh and A. Gupta, Quantum and LCD codes from Skew Constacyclic Codes over a General Class of Non-chain Rings, *Preprint*, doi.org/10.21203/rs.3.rs-3777147/v1
- [30] J. RifA and J. Pujol, Translations-invariant propelinear codes, *IEEE Trans. Inform. Theory* 43(2), pp. 590-598 (1997).

- [31] P. W. Shor, Scheme for reducing decoherence in quantum computer memory, *Phys. Rev.* A 52(4), pp. R2493–R2496 (1995).
- [32] I. Siap, T. Abualrub, N. Aydin and P. Seneviratne, Skew cyclic codes of arbitrary length, Int. J. Inf. Coding Theory 2(1), pp. 10-20 (2011).
- [33] T. Yao, M. Shi and P. Solé, Skew cyclic codes over $\mathbb{F}_q + u\mathbb{F}_q + v\mathbb{F}_q + uv\mathbb{F}_q$, J. Algebra Comb. Discrete Struct. Appl. 3(2), pp. 163-168 (2015).