# Consistency of the bootstrap for asymptotically linear estimators based on machine learning

Zhou Tang Ted Westling Department of Mathematics and Statistics University of Massachusetts Amherst

April 19, 2024

#### Abstract

arXiv:2404.03064v2 [math.ST] 18 Apr 2024

The bootstrap is a popular method of constructing confidence intervals due to its ease of use and broad applicability. Theoretical properties of bootstrap procedures have been established in a variety of settings. However, there is limited theoretical research on the use of the bootstrap in the context of estimation of a differentiable functional in a nonparametric or semiparametric model when nuisance functions are estimated using machine learning. In this article, we provide general conditions for consistency of the bootstrap in such scenarios. Our results cover a range of estimator constructions, nuisance estimation methods, bootstrap sampling distributions, and bootstrap confidence interval types. We provide refined results for the empirical bootstrap and smoothed bootstraps, and for one-step estimators, plug-in estimators, empirical mean plug-in estimators, and estimating equationsbased estimators. We illustrate the use of our general results by demonstrating the asymptotic validity of bootstrap confidence intervals for the average density value and G-computed conditional mean parameters, and compare their performance in finite samples using numerical studies. Throughout, we emphasize whether and how the bootstrap can produce asymptotically valid confidence intervals when standard methods fail to do so.

1	Introduction	3
2	Estimation and bootstrap framework         2.1       Statistical setup	<b>5</b> 5 6 7 9
3	General results         3.1       Asymptotic linearity of the estimator	<ul> <li>10</li> <li>10</li> <li>13</li> <li>16</li> <li>19</li> <li>19</li> <li>22</li> <li>23</li> </ul>
4	Remainder calculations for specific estimator constructions         4.1       One-step estimator         4.2       Plug-in estimator         4.3       Empirical mean of a nuisance-dependent function         4.4       Estimating equations-based estimator	24 24 26 28 29
5	Applications of the general theory5.1Average density value5.2G-computed conditional mean	<b>33</b> 33 38
6	Numerical study	41
7	Conclusion	54
Sι	applement A Proof of results in Section 2	61
Sι	applement B Proof of results in Section 3	62
Sι	applement C Proof of results in Section 4	79
Sι	applement D Proof of results in Section 5	86
Sι	upplement E Lemmas supporting the proof of Proposition 5.3	100
Sι	applement F Lemmas supporting the proof of Proposition 5.5	119

# 1 Introduction

In many problems in statistics, interest focuses on a parameter that depends on one or more unknown infinite-dimensional nuisance functions. For example, treatment effects in causal inference often depend on conditional mean functions (Robins, 1986; Holland, 1986; Gill & Robins, 2001), average derivative parameters depend on derivatives of conditional mean functions (Powell et al., 1989), and survival functions with informative censoring depend on conditional distribution functions (Dabrowska, 1989). Estimating such a parameter typically involves estimating one or more infinite-dimensional nuisance parameters. In order to reduce potential bias due to model misspecification, researchers may choose to use a data-adaptive estimator for the nuisance, such as a tree-based estimator, a neural network, splines, or an ensemble of many such estimators. However, directly plugging a data-adaptive nuisance estimator into the parameter mapping often results in asymptotic bias that hinders valid statistical inference for the parameter of interest. If the parameter is a smooth function of the underlying data-generating distribution, it is possible to mitigate this bias enough for the estimator to be asymptotically linear under a sufficient rate of convergence of the nuisance estimator (Pfanzagl, 1982; van der Vaart, 1991). There are several general approaches for constructing asymptotically linear estimators in the presence of dataadaptive nuisance estimators, including the one-step construction (Bickel, 1982; Pfanzagl, 1982), sieve and series approximations (Geman & Hwang, 1982), under-smoothing (Newey et al., 1998), estimating equations (Liang & Zeger, 1986; Hardin & Hilbe, 2002), and targeted minimum loss-based estimators (van der Laan & Rubin, 2006; van der Laan & Rose, 2011).

If an asymptotically linear estimator can be constructed, it can often be used to conduct asymptotically valid inference. The most popular method of constructing confidence intervals based on an asymptotically linear estimator is the Wald interval using a normal approximation to the sampling distribution and the so-called *influence function-based* variance estimator. The bootstrap (Efron, 1982; Efron & Tibshirani, 1994) is an alternative method of constructing confidence intervals that has several potential advantages over Wald intervals. First, the bootstrap has been shown in some settings to have higher-order accuracy, and hence better finite-sample coverage, than Wald intervals (Hall, 1988, 1992; Diciccio & Romano, 1988). Second, bootstrap confidence intervals can sometimes automatically correct bias, and can thus be asymptotically valid under weaker conditions (Cattaneo & Jansson, 2018, 2022). Third, in some cases, an estimator of the asymptotic variance is not readily available because, for instance, the influence function of the estimator does not have a closed form (e.g., Geskus & Groeneboom, 1999; Quale et al., 2006). Finally, Wald intervals are not guaranteed to respect constraints on the parameter space, whereas some types of bootstrap intervals are.

The theoretical properties of the bootstrap have been studied for many problems. For example, Hall (1988, 1992) established higher-order properties of bootstrap confidence intervals; Giné & Zinn (1990) and van der Vaart & Wellner (1996) established uniform bootstrap central limit theorems; Chernozhukov et al. (2017, 2022) established properties of the bootstrap for high-dimensional data; and Han et al. (2018), Austern & Syrgkanis (2020), and Cattaneo et al. (2020) studied properties of the bootstrap for non-Gaussian limits. Several authors have studied bootstrap methods for problems involving nuisance parameters. Kosorok et al. (2004) and Dixon et al. (2005) proved consistency of bootstrap procedures in semiparametric models when the nuisance parameter can be estimated at the  $n^{-1/2}$  rate. Ma & Kosorok (2005) and Cheng & Huang (2010) demonstrated consistency of the weighted empirical bootstrap for M-estimators in semiparametric models, permitting the nuisance to converge at a rate slower than  $n^{-1/2}$ . Coyle & van der Laan (2018) proposed methods of bootstrap to achieve better finite sample coverage using the highly adaptive lasso targeted minimum loss-based estimator. However, to the best of our knowledge, a comprehensive study of bootstrap procedures for asymptotically linear estimators with data-adaptive nuisance estimators does not yet exist.

In this article, we provide results for consistency of bootstrap methods for asymptotically linear estimators involving data-adaptive nuisance estimation. Several notable contributions of our work include: (1) we propose a general framework that allows us to study a variety of asymptotically linear estimator constructions, and we study several specific constructions in depth; (2) we provide conditions under which the bootstrap estimator is conditionally asymptotically linear; (3) we use our framework to provide conditions for consistency of bootstrap confidence intervals, highlighting in particular settings in which automatic bias correction is and is not possible; and (4) we cover a variety of bootstrap sampling distributions, including both the empirical bootstrap and smooth bootstraps, and a variety of methods of bootstrap nuisance estimation.

This last contribution is especially important in the context of data-adaptive nuisance estimators. As noted in Bickel et al. (1997) and Coyle & van der Laan (2018), the empirical bootstrap can fail if the estimator is sensitive to ties in the data. For example, a nuisance estimator that uses cross-validation to select tuning parameters or as part of an ensemble learning strategy may not behave as expected when applied to an empirical bootstrap sample because duplicate observations can appear in both the training and testing folds. Smooth bootstraps—i.e., bootstrap distributions that are dominated by Lebesgue measure—can resolve this issue by producing bootstrap samples without duplicates. Although smooth bootstraps have received considerably less theoretical attention than the empirical bootstrap, asymptotic properties of the smooth bootstrap have been established in, e.g., Hall et al. (1989), Cuevas & Romo (1997), and Gaenssler & Rost (2003). Our results for smooth bootstraps in Section 2.4 build on these works. Another way to avoid issues with duplicate observations is to alter the way that the nuisance is estimated for the bootstrap sample. For example, rather than using the bootstrap sample to choose tuning parameters, some authors have proposed fixing tuning parameters such as bandwidths at the values selected by the original data when constructing the bootstrap nuisance estimator (Hall & Kang, 2001). Alternatively, the entire bootstrap nuisance estimator could be fixed at the value estimated by the original data. Our framework permits these types of approaches.

The remainder of the article is organized as follows. In Section 2, we define the statistical setting we work in and outline our estimation and bootstrap frameworks. In Section 3, we provide general theoretical results, including conditional asymptotic linearity and weak convergence of the bootstrap estimator and consistency of bootstrap confidence intervals. In Section 4, we provide refined conditions implying a key condition of our main results for four estimator constructions. In Section 5, we illustrate the use of our theoretical results by studying various candidate bootstrap procedures for two parameters, deriving new results for both parameters.. In Section 6, we present a simulation study for the methods studied in Section 5. Section 7 presents a brief discussion. Proofs of all theorems and additional technical details are provided in Supplementary Material.

# 2 Estimation and bootstrap framework

# 2.1 Statistical setup

We suppose that  $X_1, \ldots, X_n \in \mathbb{R}^d$  is an IID sample from a probability measure  $P_0$  on a measurable space  $(\mathfrak{X}, \mathfrak{B})$ . We assume that  $P_0$  is known to lie in a statistical model  $\mathfrak{M}$ , which is a nonparametric or semiparametric model in our motivating applications. We will use subscript 0 to indicate that an object depends on  $P_0$ . We let  $\mathbb{P}_n$  be the empirical distribution of  $X_1, \ldots, X_n$ . For any measure Pand P-integrable function f, we define  $Pf := \int f \, dP$ . We define the *empirical process* evaluated at a  $P_0$ -integrable function f as  $\mathbb{G}_n f := n^{1/2} (\mathbb{P}_n - P_0) f$ .

For a set of functions  $\mathcal{F}$ , we define  $\ell^{\infty}(\mathcal{F})$  as the Banach space of real-valued bounded functions

 $z: \mathcal{F} \mapsto \mathbb{R}$  equipped with the supremum norm  $||z||_{\mathcal{F}} := \sup_{f \in \mathcal{F}} |z(f)|$ . To characterize weak convergence in the space  $\ell^{\infty}(\mathcal{F})$ , we utilize the bounded dual Lipschitz distance based on outer expectations. For an arbitrary metric space  $(\mathbb{D}, d)$  (frequently,  $\mathbb{D} = \ell^{\infty}(\mathcal{F})$  and  $d(z_1, z_2) = \sup_{f \in \mathcal{F}} |z_1(f) - z_2(f)|$ ), we denote  $C_b(\mathbb{D})$  as the set of bounded and continuous functions from  $\mathbb{D}$  to  $\mathbb{R}$ , and we denote  $\mathrm{BL}_1(\mathbb{D})$  as  $h: \mathbb{D} \mapsto [-1, 1]$  such that h is 1-Lipschitz; i.e.,  $\sup_{z_1, z_2 \in \mathbb{D}, z_1 \neq z_2} |h(z_1) - h(z_2)| / d(z_1, z_2) \leq 1$ . We then say that a possibly non-measurable sequence of stochastic processes  $G_n$  on  $\mathbb{D}$  converges weakly to a Borel measurable limit G in  $\mathbb{D}$ , denoted as  $G_n \rightsquigarrow G$ , if  $E_0^*h(G_n) \to E_0h(G)$  for every  $h \in C_b(\mathbb{D})$ . Here  $E_0^*$  is the *outer expectation*, which is used to accommodate non-measurable  $G_n$ . If G is a separable process (i.e., there exists  $S \subseteq \mathbb{D}$  such that  $P(G \in S) = 1$  and S has a countable dense subset), then  $G_n \rightsquigarrow G$  if and only if  $\sup_{h \in \mathrm{BL}_1(\mathbb{D})} |E_0^*h(G_n) - E_0h(G)| \to 0$ . We refer the reader to van der Vaart & Wellner (1996) for a review of outer expectation and weak convergence.

A class  $\mathcal{F}$  of measurable functions  $f : \mathfrak{X} \mapsto \mathbb{R}$  is called  $P_0$ -Donsker if the sequence of empirical processes  $\{\mathbb{G}_n f : f \in \mathcal{F}\}$  converges weakly in  $\ell^{\infty}(\mathcal{F})$  to a tight, Borel measurable limit process  $\mathbb{G}_0$ , or  $\sup_{h \in \mathrm{BL}_1(\ell^{\infty}(\mathcal{F}))} |E_0^*h(\mathbb{G}_n) - E_0h(\mathbb{G}_0)| \to 0$  (van der Vaart & Wellner, 1996, Chapter 1.12). The limit process  $\mathbb{G}_0$  is necessarily a Gaussian process with zero mean and covariance function  $\mathrm{Cov}(\mathbb{G}_0 f, \mathbb{G}_0 g) =$  $P_0(fg) - P_0 f P_0 g$  for  $f, g \in \mathcal{F}$ , which is known as  $P_0$ -Brownian bridge process. Implicitly, the Donsker property requires that the sample paths  $f \mapsto \mathbb{G}_n f$  are almost surely uniformly bounded for every n, so that  $\mathbb{G}_n$  may be regarded as a map from the underlying product measurable space  $(\mathfrak{X}^{\infty}, \mathfrak{B}^{\infty})$  to  $\ell^{\infty}(\mathcal{F})$ . This is the case if  $\sup_{f \in \mathcal{F}} |f(x) - P_0 f| < \infty$  for all  $x \in \mathfrak{X}$ .

# 2.2 Asymptotically linear estimator framework

We are interested in inference for a real-valued target parameter  $\psi : \mathcal{M} \to \mathbb{R}$ . All the results in this article extend to Euclidean parameters  $\psi : \mathcal{M} \to \mathbb{R}^p$  for  $p < \infty$  fixed, but we assume p = 1 for simplicity of exposition. We assume that  $\psi$  is a smooth enough mapping on the model  $\mathcal{M}$  to permit construction of an asymptotically linear estimator  $\psi_n$  with influence function  $\phi_0 := \phi_{P_0}$ , meaning that  $\psi_n = \psi_0 + \mathbb{P}_n \phi_0 + o_{P_0^*}(n^{-1/2})$ , where  $\psi_0 := \psi(P_0)$ . For any  $P \in \mathcal{M}$ ,  $\phi_P : \mathcal{X} \to \mathbb{R}$  is a function satisfying  $P\phi_P = 0$  and  $P\phi_P^2 < \infty$ . Conditions under which this is possible and derivation of influence functions is not our main focus here, but we refer the interested reader to Pfanzagl (1982), Klaassen (1987), Pfanzagl (1990), and van der Vaart (1991). Here, we will not require that  $\phi_0$  is the efficient influence function, but rather allow it to be any influence function. In addition, we do not explicitly require that  $\psi_n$  is a regular estimator. The premise of this article is that the parameter  $\psi$  depends on P through an infinite-dimensional nuisance parameter  $\eta : \mathcal{M} \to \mathcal{H}$ . For instance,  $\eta$  may be a (conditional) density function, a (conditional) cumulative distribution function, a regression function, or some combination of these. A nuisance estimator  $\eta_n$  may then be used in the construction of an estimator  $\psi_n$  of  $\psi_0$ , However, if  $\eta_n$  is constructed without consideration of  $\psi$  or its influence function, then the resulting  $\psi_n$  may have excess bias inherited from  $\eta_n$  that precludes asymptotic linearity. There are several existing remedies. One approach is to construct the nuisance estimator to reduce the bias of the plug-in estimator. Undersmoothing (Newey et al., 1998), twicing kernels (Newey et al., 2004), sieves (Geman & Hwang, 1982; Shen, 1997), and TMLE (van der Laan & Rubin, 2006; van der Laan & Rose, 2011) are examples of this approach. Another approach is to abandon plug-in estimation and instead use the influence function to target the parameter of interest. One-step estimators (Bickel, 1982; Pfanzagl, 1982) and estimating equationsbased estimators (Liang & Zeger, 1986) are examples of this approach.

We now introduce a general framework that encompasses many approaches to constructing an asymptotically linear estimator. This will allow us to study these approaches in a unified manner. We assume there exists a function  $T : \mathcal{H} \times \mathcal{M}^+$ , where  $\mathcal{M}^+$  is the union of  $\mathcal{M}$  and the set of finite discrete probability measures on  $(\mathfrak{X}, \mathfrak{B})$ , such that  $\psi_0 = T(\eta_0, P_0)$  and  $\psi_n = T(\eta_n, \mathbb{P}_n)$ for a nuisance estimator  $\eta_n \in \mathcal{H}$ . We note that for a given estimator  $\psi_n$ , there may be multiple representations of  $\psi_n$  in terms of different functions T because  $\eta_n$  also typically depends on  $\mathbb{P}_n$ . To illustrate this general framework, we provide two brief examples of estimator constructions T. These examples and others will be discussed further in Section 4. The *one-step* construction is defined as  $T(\eta, P) = \psi(\eta) + P\phi_{\eta}$ , where it is assumed that the parameter  $\psi(P)$  and its influence function  $\phi_P$  depend on P only through  $\eta_P$ . The mean-zero property of influence functions implies that  $T(\eta_0, P_0) = \psi_0$ . A one-step estimator is then given by  $T(\eta_n, \mathbb{P}_n) = \psi(\eta_n) + \mathbb{P}_n \phi_{\eta_n}$ . Alternatively, the *empirical mean plug-in* construction is defined as  $T(\eta, P) := \int g(x, \eta) \, dP(x)$  for a function  $g: \mathfrak{X} \times \mathfrak{H} \to \mathbb{R}$ , which can be used when  $\psi(P) = \int g(x, \eta_P) \, dP(x)$ . The resulting estimator is then given by  $T(\eta_n, \mathbb{P}_n) = \int g(x, \eta_n) \, d\mathbb{P}_n(x) = \frac{1}{n} \sum_{i=1}^n g(X_i, \eta_n)$ .

#### 2.3 Bootstrap framework

We now introduce the class of bootstrap schemes that we will consider. At a high level, the bootstrap schemes we will consider involve three steps. First, bootstrap samples  $X_1^*, \ldots, X_n^*$  are generated in some manner based on the data  $X_1, \ldots, X_n$ . Second, a version of the estimation procedure is applied to the bootstrap data to produce a bootstrap estimate  $\psi_n^*$ . Third, this process is repeated *B* times to approximate the sampling distribution of  $\psi_n^* - \psi_n$  given the data. Throughout, we will ignore the effect of approximating the distribution of  $\psi_n^* - \psi_n$  using a finite number of repetitions. The distribution of  $\psi_n^* - \psi_n$  given the data is used to approximate the sampling distribution of  $\psi_n - \psi_0$ , and ultimately to construct confidence intervals for  $\psi_0$ . There are many specific approaches to each of these three steps, and as discussed in the introduction, our goal is to provide results that cover a broad set of these approaches. In this section, we precisely define the approaches to the first two steps that we will consider; procedures for constructing confidence intervals are discussed in Section 3.4.

We assume that, given the data  $X_1, \ldots, X_n$ , the bootstrap sample  $X_1^*, \ldots, X_n^*$  is drawn IID from an estimate  $\hat{P}_n$  of  $P_0$  based on  $X_1, \ldots, X_n$ . We refer to  $\hat{P}_n$  as the bootstrap sampling distribution. Taking  $\hat{P}_n = \mathbb{P}_n$  corresponds to the empirical bootstrap, in which case the bootstrap sample consists of nsamples drawn IID (i.e., with replacement) from  $X_1, \ldots, X_n$ . Another common approach to defining  $\hat{P}_n$  is through smoothing methods such as kernel density estimation. This will be discussed at more length in Section 2.4. Notably, since we assume that n samples are drawn independently from  $\hat{P}_n$ , we exclude the exchangeable, weighted, and m-out-of-n bootstraps. Our results could be generalized to the weighted bootstrap if the nuisance estimator utilizes sample weights, and to the exchangeable bootstrap if the nuisance estimator does not depend on the independence of the data. We define the *bootstrap empirical distribution*  $\mathbb{P}_n^*$  as the empirical distribution of the bootstrap data  $X_1^*, \ldots, X_n^*$  and the *bootstrap empirical process*  $\mathbb{G}_n^*$  as  $\mathbb{G}_n^* := n^{1/2}(\mathbb{P}_n^* - \hat{P}_n)$ .

Once the process for generating bootstrap data has been defined, the next step is to define the process for constructing the estimator using the bootstrap data. Since our definition of the original estimator is  $\psi_n = T(\eta_n, \mathbb{P}_n)$ , we will define  $\psi_n^* := T(\eta_n^*, \mathbb{P}_n^*)$  as the estimator using the bootstrap data, where  $\eta_n^*$  is an estimator of the nuisance parameter  $\eta_0$  based on the bootstrap data and original data, and  $\mathbb{P}_n^*$  is the bootstrap empirical distribution as previously defined. The spirit of bootstrap estimation would suggest that  $\eta_n^*$  be constructed using the bootstrap data in the exact same manner as  $\eta_n$  was constructed using the original data. However, we will not require this — instead, we will be agnostic about the way  $\eta_n^*$  is constructed. There are several reasons motivating this increase in generality. First, in many of our intended applications,  $\eta_n$  is estimated using machine learning, which may be very computationally intensive. Repeating a computationally intensive procedure for every bootstrap sample may be infeasible since the number of bootstrap samples B is typically in the hundreds or thousands. Second, as discussed in the introduction, there are certain cases where it is not advisable

to exactly mirror the estimation of  $\eta_n$  when constructing  $\eta_n^*$ . For instance, many machine learning algorithms involve cross-validation. However, if the bootstrap sampling process produces replicates in the bootstrap data, which is the case for the empirical bootstrap, cross-validation may not perform as expected (see, e.g., Silverman, 1986, Page 51 and Coyle & van der Laan, 2018, Chapter 28). In this case, modifications of the estimation procedure of  $\eta_n^*$  have been proposed to avoid these issues. In addition, some authors have proposed fixing tuning parameters such as bandwidths at the values selected by the original data when constructing  $\eta_n^*$  using the bootstrap data (Hall & Kang, 2001).

Our asymptotic results presented in Section 3 will require high-level conditions about  $\eta_n^*$ : consistency, a Donsker condition, and negligibility of a remainder term. This mirrors the high-level conditions required of  $\eta_n$  for asymptotic linearity of  $\psi_n$ . Our conditions can be satisfied if the construction of  $\eta_n^*$ mirrors that of  $\eta_n$  completely or partially. In some cases, our conditions will also permit  $\eta_n^* = \eta_n$ . That is, we will permit that the nuisance is not re-estimated at all using the bootstrap data, but rather the estimator computed on the original data is used when constructing the bootstrap estimator. This is discussed more in Section 4.

#### 2.4 Additional bootstrap notation

Crucial to bootstrap theory is taking conditional expectations of the bootstrap data given the original data. To do so precisely, we make the following definitions, which are common in the bootstrap literature (see, e.g., van der Vaart & Wellner, 1996; van der Vaart, 2000; Kosorok, 2008). We suppose that  $W_n = (W_{n1}, \ldots, W_{nn})$  is an IID sample independent of  $X_1, \ldots, X_n$ , where each  $W_{ni}$  is a random vector with distribution  $Q_n$  on a measurable space  $(W_n, C_n)$ . We then assume that for each  $i \in \{1, \ldots, n\}, X_i^* = \gamma_n(X_1, \ldots, X_n, W_{ni})$ , where  $\gamma_n$  is a fixed measurable function. Hence,  $W_{ni}$  represents the additional source of randomness used in generating the bootstrap observation  $X_i^*$ . Since  $W_{n1}, \ldots, W_{nn}$  are assumed to be IID and independent of  $X_1, \ldots, X_n, X_1^*, \ldots, X_n^*$  are IID conditional on  $X_1, \ldots, X_n$ . With this setup,  $\hat{P}_n$  is defined as the conditional distribution of  $X_i^*$  given  $X_1, \ldots, X_n$ . The bootstrap sample  $X_1^*, \ldots, X_n^*$  lies in the product probability space  $(\mathfrak{X}, \mathfrak{B}, P_0)^n \times (\mathfrak{W}_n, \mathfrak{C}_n, Q_n)^n$ .

We provide two concrete examples to illustrate the above definitions. For the empirical bootstrap, we can let  $W_{ni}$  have a categorical distribution  $Q_n$  on  $\{1, \ldots, n\}$  with event probabilities  $(1/n, \ldots, 1/n)$ , and set  $X_i^* = \gamma_n(X_1, \ldots, X_n, W_{ni}) = X_{W_{ni}}$ . For one-dimensional observations  $X_i$  and any bootstrap, we can let  $W_{ni}$  be IID Uniform(0, 1), and set  $X_i^* := F_n^{-1}(W_{ni})$  for  $F_n^{-1}$  the quantile function corresponding to  $\hat{P}_n$ . Here,  $F_n^{-1}$  is assumed to be a measurable function of  $X_1, \ldots, X_n$ . We now define conditional expectations given the data  $X_1, \ldots, X_n$ . Since we will be dealing with processes that may not be conditionally measurable, we will use outer expectations. For any real-valued function  $h : \mathfrak{X}^n \times \mathfrak{W}_n^n \to \mathbb{R}$ , we define the *conditional outer expectation* of h given  $X_1, \ldots, X_n$  with respect to  $Q_n$  as

$$(E_W^*h)(X_1,\ldots,X_n) := \inf_U \left\{ \int_{\mathcal{W}_n^n} U(X_1,\ldots,X_n,w_1,\ldots,w_n) \,\mathrm{d}Q_n^n(w_1,\ldots,w_n) \right\},\$$

where the infimum is taken over all functions U such that  $w_1, \ldots, w_n \mapsto U(X_1, \ldots, X_n, w_1, \ldots, w_n)$ is measurable,  $U \ge h$  almost surely, and  $\int U \, dQ_n^n$  exists. Since each  $X_i^*$  is a measurable function of  $X_1, \ldots, X_n$  and  $W_{ni}$ , this can also be used to define conditional expectations of functions of  $X_1, \ldots, X_n$ and  $X_1^*, \ldots, X_n^*$ . We then define the *conditional outer probability* given  $X_1, \ldots, X_n$  as  $P_W^*(A) := E_W^* \mathbf{1}_A$ for any  $A \in \mathcal{B}^n \times \mathcal{C}_n^n$ .

We say that a (possibly non-measurable) sequence of random elements  $G_n: \mathfrak{X}^n \times \mathfrak{W}_n^n \to \mathbb{D}$  for a metric space  $\mathbb{D}$  conditionally weakly converges to a tight, Borel measurable limit G in  $\mathbb{D}$  given  $X_1, \ldots, X_n$  in outer probability if  $\sup_{h \in \mathrm{BL}_1(\mathbb{D})} |E_W^*h(G_n) - E_0h(G)| \xrightarrow{P_0^*} 0$ , and we denote this as  $G_n \overset{P_0^*}{\underset{W}{W}} G$ . In addition, we say  $\mathfrak{F}$  is  $P_0$ -Donsker in probability if it holds that  $\sup_{h \in \mathrm{BL}_1(\ell^\infty(\mathfrak{F}))} |E_W^*h(\mathbb{G}_n^*) - E_0h(\mathbb{G})| = o_{P_0^*}(1)$ (Giné & Zinn, 1990, Section 3). We say a sequence of random variables  $Y_n: \mathfrak{X}^n \times \mathfrak{W}_n^n \to \mathbb{R}$  conditionally converge to 0 in probability if for any  $\varepsilon > 0$ ,  $P_W^*(|Y_n| \ge \varepsilon) \xrightarrow{P_0^*} 0$ , and we denote this as  $Y_n = o_{P_W^*}(1)$ . We say  $Y_n$  is conditionally stochastically bounded if for any  $\varepsilon > 0$ , there exist  $M \in (0, \infty)$  such that  $P_0^*(P_W^*(|Y_n| \ge M) \ge \epsilon) \to 0$ , and we denote this as  $Y_n = O_{P_W^*}(1)$ .

# 3 General results

### 3.1 Asymptotic linearity of the estimator

Before moving to our general bootstrap consistency results, we provide general conditions under which  $T(\eta_n, \mathbb{P}_n)$  is an asymptotically linear estimator of  $\psi_0$  with influence function  $\phi_0$ . While the result is simple and based on well-known ideas, it is important background because our theoretical study of the bootstrap will focus on consistency of the sampling distribution of the bootstrap estimator, which requires knowing the distribution it should be consistent for — i.e., the asymptotic distribution of the bootstrap estimator. In addition, the conditions for conditional asymptotic linearity of the bootstrap estimator will mirror the conditions for asymptotic linearity of  $T(\eta_n, \mathbb{P}_n)$ . Hence, these conditions will

shed more light on the bootstrap conditions presented below.

For each n, we define  $\phi_n : \mathfrak{X} \to \mathbb{R}$  as an estimator of the influence function  $\phi_0$ . For the one-step and estimating equations-based estimators,  $\phi_n$  will be the influence function with the estimated nuisance parameter  $\eta_n$ . For estimators such as the plug-in estimator where the influence function is not explicitly estimated as part of the estimation procedure, the role of  $\phi_n$  will be discussed more below. We also define  $R_n := T(\eta_n, \mathbb{P}_n) - T(\eta_0, P_0) - (\mathbb{P}_n - P_0)\phi_n$ . We then introduce the following conditions.

- (A1) There exists a class  $\mathcal{F}$  of measurable functions from  $(\mathfrak{X}, \mathcal{B})$  to  $\mathbb{R}$  such that:
  - (a)  $\phi_0 \in \mathcal{F}$  and  $P_0^*(\phi_n \in \mathcal{F}) \to 1$ , and
  - (b)  $\mathbb{G}_n \rightsquigarrow \mathbb{G}_0$  in  $\ell^{\infty}(\mathcal{F})$ , where  $\mathbb{G}_0$  is the  $P_0$ -Brownian bridge process.
- (A2) It holds that  $\|\phi_n \phi_0\|_{L_2(P_0)} = o_{P_0^*}(1)$ .
- (A3) The remainder satisfies  $R_n = o_{P_0^*}(n^{-1/2})$ .

Under these conditions, we have the following result regarding asymptotic linearity of  $T(\eta_n, \mathbb{P}_n)$ .

**Theorem 3.1.** If conditions (A1)-(A3) hold, then  $T(\eta_n, \mathbb{P}_n)$  is asymptotically linear in the sense that  $T(\eta_n, \mathbb{P}_n) = T(\eta_0, P_0) + \mathbb{P}_n \phi_0 + o_{P_0^*}(n^{-1/2})$ , which implies that  $n^{1/2} [T(\eta_n, \mathbb{P}_n) - T(\eta_0, P_0)] \rightsquigarrow \mathbb{G}_0 \phi_0$ .

Theorem 3.1 provides conditions under which  $\psi_n = T(\eta_n, \mathbb{P}_n)$  is an asymptotically linear estimator of  $\psi_0 = T(\eta_0, P_0)$ . This implies convergence in distribution of  $n^{1/2} [T(\eta_n, \mathbb{P}_n) - T(\eta_0, P_0)]$  to  $\mathbb{G}_0 \phi_0$ , which follows a normal distribution with mean zero and variance  $P_0 \phi_0^2$ . In particular, this gives a method for constructing asymptotically valid confidence intervals. If  $\sigma_n$  is a consistent estimator of  $(P_0 \phi_0^2)^{1/2}$ , then the two-sided Wald confidence interval

$$[T(\eta_n, \mathbb{P}_n) + z_\beta \sigma_n / n^{1/2}, T(\eta_n, \mathbb{P}_n) + z_{1-\alpha} \sigma_n / n^{1/2}]$$

$$\tag{1}$$

has asymptotic level  $1 - \alpha - \beta$  by Slutsky's lemma. The simplest variance estimator is the *influence* function-based variance given by  $\sigma_n^2 := \mathbb{P}_n \phi_n^2$ . The next proposition provides general conditions under which this variance estimator is consistent and demonstrates that conditions (A1)–(A2) in particular imply that it is consistent.

**Proposition 3.2.** If (i)  $P_0^*(\phi_n^2 \in \mathfrak{G}) \to 1$  for a  $P_0$ -Glivenko Cantelli class of measurable functions  $\mathfrak{G}$ , and (ii)  $P_0(\phi_n^2 - \phi_0^2) \xrightarrow{P_0^*} 0$ , then  $\mathbb{P}_n \phi_n^2 \xrightarrow{P_0^*} \sigma_0^2$ . Furthermore, if  $\sup_{f \in \mathfrak{F}} |P_0 f| < \infty$ , condition (A1) implies (i) and condition (A2) implies (ii). Theorem 3.1 and Proposition 3.2 yield the following result for consistency of the Wald-type confidence interval with influence function-based variance estimator.

**Corollary 3.3.** If  $\sup_{f \in \mathcal{F}} |P_0 f| < \infty$ , conditions (A1)-(A3) hold, and  $\sigma_0^2 > 0$ , then the Wald-type confidence interval defined in (1) with  $\sigma_n^2 = \mathbb{P}_n \phi_n^2$  has asymptotic confidence level  $1 - \alpha - \beta$ .

We now discuss conditions (A1)–(A3). Theorem 3.1 is based on the first-order expansion

$$T(\eta_n, \mathbb{P}_n) = T(\eta_0, P_0) + \mathbb{P}_n \phi_0 + R_n + S_n,$$

where  $S_n := n^{-1/2} \mathbb{G}_n(\phi_n - \phi_0)$  is an *empirical process term*. Conditions (A1) and (A2) together imply that  $S_n = o_{P_0^*}(n^{-1/2})$ . Condition (A1) requires that the estimated influence function  $\phi_n$  falls in a  $P_0$ -Donsker class with probability tending to one. Satisfying this condition typically requires restricting the complexity of the function class that the nuisance estimator  $\eta_n$ , and hence  $\phi_n$ , falls in. A main way this is accomplished is by using bracketing entropy or uniform entropy (Chapters 2.6 and 2.7 of van der Vaart & Wellner, 1996). Condition (A2) requires that  $\phi_n$  is a consistent estimator of  $\phi_0$  in the  $L_2(P_0)$ norm. When  $\phi_n$  and  $\phi_0$  depend on n and  $P_0$  through  $\eta_n$  and  $\eta_0$ , respectively, this is typically implied by consistency of  $\eta_n$  for  $\eta_0$  in an appropriate sense coupled with continuity of  $\phi$  as a function of  $\eta$ .

Condition (A3) controls the remainder term  $R_n$  in the above expansion. Other authors have used analogous conditions in related work (see, e.g., the smoothness property discussed in Section 4.1 of Shen, 1997 and the quadratic functional in Section 3.2 of Cattaneo & Jansson, 2018). This remainder term can typically be decomposed into further remainders, including the so-called *second-order remainder* and the bias term  $-\mathbb{P}_n\phi_n$ . The exact way that this remainder decomposes depends on the form of T. Estimators such as one-step and estimating equations-based estimators that explicitly use the influence function as part of the construction control the asymptotic bias as part of the estimation procedure. On the other hand, plug-in estimators typically control the bias term  $-\mathbb{P}_n\phi_n$  through construction of the nuisance estimator  $\eta_n$ . Several approaches to constructing nuisance estimators that yield  $-\mathbb{P}_n\phi_n = o_{\mathbb{P}_0^*}(n^{-1/2})$ are sieve estimators (Shen, 1997), under-smoothing (Newey et al., 1998), twicing kernels (Newey et al., 2004), and TMLE. Simpler sufficient conditions for (A3) for several different estimator constructions will be discussed at more length in Section 4.

# 3.2 Conditional asymptotic linearity of the bootstrap estimator

We now present our general result regarding conditional asymptotic linearity of the bootstrap procedure defined in Section 2.3. As mentioned above,  $\phi_n^*$  is a bootstrap influence function estimator. We begin by introducing conditions we will rely upon. We define  $R_n^* := T(\eta_n^*, \mathbb{P}_n^*) - T(\eta_n, \hat{P}_n) - (\mathbb{P}_n^* - \hat{P}_n)\phi_n^*$ .

(B1) There exists a class  $\mathcal{F}$  of measurable functions from  $(\mathfrak{X}, \mathfrak{B})$  to  $\mathbb{R}$  such that

- (a)  $\phi_0 \in \mathcal{F} \text{ and } P_W^*(\phi_n^* \in \mathcal{F}) \xrightarrow{P_0^*} 1$ , and
- (b)  $\mathbb{G}_n^* \overset{P_0^*}{\underset{W}{\longrightarrow}} \mathbb{G}_0$  in  $\ell^{\infty}(\mathcal{F})$ , where  $\mathbb{G}_0$  is the  $P_0$ -Brownian bridge process.
- **(B2)** It holds that  $\|\phi_n^* \phi_0\|_{L_2(P_0)} = o_{P_W^*}(1)$ .
- **(B3)** The remainder term satisfies  $R_n^* = o_{P_W^*}(n^{-1/2})$ .

Under these conditions, we have the following result.

**Theorem 3.4.** If conditions (B1)-(B3) hold, then  $T(\eta_n^*, \mathbb{P}_n^*)$  is conditionally asymptotically linear in the sense that  $T(\eta_n^*, \mathbb{P}_n^*) = T(\eta_n, \hat{P}_n) + (\mathbb{P}_n^* - \hat{P}_n)\phi_0 + o_{P_W^*}(n^{-1/2})$ , which implies that

$$n^{1/2} \left[ T\left(\eta_n^*, \mathbb{P}_n^*\right) - T(\eta_n, \hat{P}_n) \right] \overset{P_0^*}{\underset{W}{\longrightarrow}} \mathbb{G}_0(\phi_0).$$

Theorem 3.4 establishes general conditions under which the bootstrap estimator  $T(\eta_n^*, \mathbb{P}_n^*)$  is conditionally asymptotically linear. Theorem 3.4 is notable for its generality: conditions (B1)–(B3) cover a variety of estimator constructions T, bootstrap nuisance estimators  $\eta_n^*$ , and bootstrap sampling distributions  $\hat{P}_n$ . We expect this generality to increase the range of potential applications of the result.

The most important implication of conditional asymptotic linearity is that the bootstrap provides a consistent approximation to the sampling distribution of  $n^{1/2}(\psi_n - \psi_0)$ . Here, consistency means that the conditional distribution of the centered and scaled bootstrap estimator  $n^{1/2}[T(\eta_n^*, \mathbb{P}_n^*) - T(\eta_n, \hat{P}_n)]$  converges weakly in outer probability to the same limit as  $n^{1/2}(\psi_n - \psi_0)$ . In Section 3.4, we will discuss how this can be used to demonstrate asymptotic validity of bootstrap confidence intervals. However, as with asymptotic linearity, conditional asymptotic linearity offers additional utility beyond conditional weak convergence. In particular, conditional asymptotic linearity of multiple bootstrap estimators implies *joint* conditional asymptotic normality of the estimators, which is useful for constructing simultaneous

confidence regions and confidence regions for functions of two or more parameters. In contrast, such joint asymptotic behavior cannot be determined by marginal weak convergence results.

Theorem 3.4 centers the bootstrap estimator  $T(\eta_n^*, \mathbb{P}_n^*)$  around  $T(\eta_n, \hat{P}_n)$  rather than  $T(\eta_n, \mathbb{P}_n) = \psi_n$ . There is no difference between these two possibilities when using the empirical bootstrap, so that  $\hat{P}_n = \mathbb{P}_n$ . However, when  $\hat{P}_n \neq \mathbb{P}_n$ , such as when utilizing a smooth bootstrap procedure, the two are not necessarily the same. Intuitively, we center around  $T(\eta_n, \hat{P}_n)$  because  $\hat{P}_n$  is the distribution used to generate the bootstrap data upon which the bootstrap empirical distribution  $\mathbb{P}_n^*$  is based. Hence,  $\hat{P}_n$  plays the role of  $P_0$  in the bootstrap. The potential consequences of this for the construction of confidence intervals will be further explored in Section 3.4.

Conditions (B1)–(B3) mirror conditions (A1)–(A3) used to demonstrate asymptotic linearity of the estimator  $T(\eta_n, \mathbb{P}_n)$  in Theorem 3.1. However, since (B1)–(B3) concern the bootstrap estimator, they require convergence conditional on the original observations. In some cases, conditional asymptotic linearity of the bootstrap estimator is actually implied by the conditions of Theorem 3.1. We will discuss this more in Section 4. This is related to Beran (1997), who showed that for locally asymptotically normal parametric models, conditional weak convergence of the parametric and nonparametric bootstraps is equivalent to regularity of the estimator. For pathwise differentiable parameters, an asymptotically linear estimator is regular if and only if its influence function is a gradient of  $\psi$  relative to  $\mathcal{M}$  at  $P_0$ (Pfanzagl, 1982, 1990; van der Vaart, 1991). We typically construct the estimator so that  $\phi_0$  is indeed a gradient. Furthermore, negligibility of the second-order remainder term for condition (A3) is often established using conditions that also imply that  $\phi_0$  is a gradient, as we will discuss in Section 4. Hence, while we do not explicitly require regularity, the estimator is regular in most of our intended applications. However, it is not entirely clear if regularity of the estimator plays as strong a role in our setting as it does in that of Beran (1997) for parametric models.

Theorem 3.4 is based on the bootstrap first-order expansion

$$T(\eta_n^*, \mathbb{P}_n^*) = T(\eta_n, \hat{P}_n) + (\mathbb{P}_n^* - \hat{P}_n)\phi_0 + R_n^* + S_n^*,$$

where  $S_n^* = n^{-1/2} \mathbb{G}_n^*(\phi_n^* - \phi_0)$  is a bootstrap empirical process remainder term. Conditions (B1)–(B2) are used to control  $S_n^*$ . This is analogous to how conditions (A1)–(A2) were used to control the ordinary empirical process remainder  $S_n$ . In particular, condition (B1)(b) requires conditional weak convergence in outer probability of the bootstrap empirical process  $\mathbb{G}_n^* = n^{1/2}(\mathbb{P}_n^* - \hat{P}_n)$  in the space  $\ell^{\infty}(\mathcal{F})$  to a  $P_0$ -Brownian bridge process. For the empirical bootstrap where  $\hat{P}_n = \mathbb{P}_n$ , this holds as long as  $\mathcal{F}$  is  $P_0$ -Donsker (Giné & Zinn, 1990). However, for other types of bootstraps, this condition is more difficult to verify. This will be discussed in depth and further sufficient conditions for smooth bootstrap sampling distributions will be provided in Section 3.3.

Condition (B2) requires conditional weak consistency of the bootstrap nuisance estimator  $\eta_n^*$  to the true nuisance  $\eta_0$ . If  $\eta_n^*$  is constructed in an exactly analogous manner using the bootstrap data as  $\eta_n$  is constructed using the original data, the bootstrap data has replicated observations, and the method of constructing  $\eta_n$  is sensitive to ties in the data, (B2) may not be satisfied. As discussed in Section 2.3, for this reason and others we do not require that  $\eta_n^*$  be constructed in an exactly analogous manner to  $\eta_n$ , so these issues can be avoided. In particular, the simplest approach for constructing  $\eta_n^*$  is to define  $\eta_n^* = \eta_n$ . This approach is appealing in its computational simplicity because it does not require re-fitting the nuisance estimator using the bootstrap data, which can be computationally intensive when machine learning estimators are used to construct  $\eta_n$ . Furthermore, when  $\eta_n^* = \eta_n$  and  $\phi_P$  only depends on P through  $\eta_P$ , condition (B2) reduces to the requirement that  $\|\phi_n - \phi_0\|_{L_2(P_0)} \xrightarrow{P_0^*} 0$ , which was already required for asymptotic linearity of  $\psi_n$  in Theorem 3.1. Intuitively, the precise behavior of the nuisance estimator  $\eta_n$  does not play a role in the first-order asymptotic behavior of  $\psi_n$  as long as the high-level conditions (A1)–(A3) hold, and similarly the precise behavior of  $\eta_n^*$  does not play a role in the first-order asymptotic behavior of the bootstrap estimator as long as the high-level conditions (B1)-(B3) hold. However, setting  $\eta_n^* = \eta_n$  may yield worse finite-sample coverage, and does not yield valid bootstrap confidence intervals when conditions (A3) and/or (B3) do not hold, as we will discuss in Section 3.4.

Our proof technique for Theorem 3.4 could be adapted to other tight and Borel-measurable limit processes in condition (B1)(b). This is not relevant to demonstrating consistency of the bootstrap when the estimator is asymptotically Gaussian, as is the case here, but it may be of interest in other situations where the limit distribution is not Gaussian. However, if  $\mathbb{G}_0$  were a non-Gaussian process in condition (B1)(b) and the sample paths of  $\mathbb{G}_0$  were not almost surely uniformly continuous in the  $L_2(P_0)$ metric, it would be necessary to replace condition (B2) with the requirement that  $\rho(\phi_n^*, \phi_0) = o_{P_W^*}(1)$ for a semimetric  $\rho$  on  $\ell^{\infty}(\mathcal{F})$  such that the sample paths of  $\mathbb{G}_0$  are almost surely uniformly  $\rho$ -continuous.

Condition (B3) requires that the bootstrap analogue of the remainder term in (A3) be sufficiently negligible. As discussed after Theorem 3.1, this remainder term is again a combination of two remainders: the bootstrap bias term  $-\mathbb{P}_n^*\phi_n^*$  and the bootstrap second-order remainder term. Further sufficient conditions for (B3) for specific estimators T will be provided in Section 4.

Theorem 3.4 also relies on the following bootstrap version of Lemma 19.24 of van der Vaart (2000), which is useful in its own right.

**Lemma 3.5.** Suppose that  $\mathcal{F}$  is a class of measurable functions such that  $\mathbb{G}_n^* \overset{P_0^*}{\underset{W}{\leftrightarrow}} \mathbb{G}_0$  in  $\ell^{\infty}(\mathcal{F})$ . Let  $\phi_n^*$  be a sequence of random functions possibly depending on the bootstrap sample such that  $P_W^* (\phi_n^* \in \mathcal{F}) \xrightarrow{P_0^*} 1$ . If  $\rho(\phi_n^*, \phi_\infty) = o_{P_W^*}(1)$  for some  $\phi_\infty \in \mathcal{F}$  and a semimetric  $\rho$  on  $\mathcal{F}$  for which the sample paths of  $\mathbb{G}_0$  are almost surely uniformly  $\rho$ -continuous, then  $\mathbb{G}_n^*(\phi_n^* - \phi_\infty) = o_{P_W^*}(1)$ .

### 3.3 Conditional weak convergence of the bootstrap empirical process

We now provide further sufficient conditions for conditional weak convergence of the bootstrap empirical process  $\mathbb{G}_n^*$  required by condition (B1)(b). We first discuss the case of the empirical bootstrap, where  $\hat{P}_n = \mathbb{P}_n$ . The properties of the empirical bootstrap have been extensively studied by Efron (1982), Giné & Zinn (1990, 1991), Praestgaard & Wellner (1993) and van der Vaart & Wellner (1996), among others. In particular, Theorem 3.1 of Giné & Zinn (1990) and Theorem 3.6.1 of van der Vaart & Wellner (1996) provided the following necessary and sufficient condition for (B1)(b) in the case of the empirical bootstrap.

**Lemma 3.6** (Theorem 3.6.1 of van der Vaart & Wellner, 1996). If  $\hat{P}_n = \mathbb{P}_n$  and  $\mathcal{F}$  is a class of measurable function with finite envelope function, then  $\mathcal{F}$  is  $P_0$ -Donsker if and only if condition (B1)(b) holds and  $\mathbb{G}_n^*$  is asymptotically measurable.

We now turn to the case where  $\hat{P}_n$  is not the empirical distribution. We first provide a general set of sufficient conditions for (B1)(b) based on the notion of *uniform Donsker classes*. For a probability measure P on  $(\mathfrak{X}, \mathfrak{B})$ , we denote  $\mathbb{G}_{n,P} := n^{1/2}(\mathbb{P}_n - P)$  as the empirical process centered at P and  $\mathbb{G}_P$ as the P-Brownian bridge process. Following Giné & Zinn (1991) and Sheehy & Wellner (1992), for a set  $\mathfrak{P}$  of probability measures on  $(\mathfrak{X}, \mathfrak{B})$ , we then say  $\mathfrak{F}$  is *Donsker uniformly in*  $P \in \mathfrak{P}$  if

$$\sup_{P \in \mathcal{P}} \sup_{h \in \mathrm{BL}_1(\ell^{\infty}(\mathcal{F}))} |E^*h(\mathbb{G}_{n,P}) - Eh(\mathbb{G}_P)| \to 0,$$

and  $\mathbb{G}_P$  satisfies  $\sup_{P \in \mathcal{M}} E \|\mathbb{G}_P\|_{\mathcal{F}} < \infty$  and

 $\lim_{\delta \to 0} \sup_{P \in \mathcal{P}} E \sup_{\rho_P(f,g) < \delta} |\mathbb{G}_P(f) - \mathbb{G}_P(g)| = 0,$ 

where  $\rho_P : (f,g) \in \mathcal{F} \times \mathcal{F} \to [P(f-g)^2]^{1/2}$  is the *P*-standard deviation semi-metric on  $\mathcal{F} \times \mathcal{F}$ . Theorem 4.5 of Sheehy & Wellner (1992) provides general sufficient conditions for an almost sure convergence version of (B1)(b). Below, we restate their result relaxed to convergence in outer probability.

**Lemma 3.7** (Theorem 4.5 of Sheehy & Wellner, 1992). If  $\mathcal{F}$  is a class of measurable functions with envelope function F such that: (i)  $\mathcal{F}$  is square integrable uniformly in  $P \in \mathcal{P}$  in the sense that  $\lim_{M\to\infty} \sup_{P\in\mathcal{P}} PF^2 1\{F > M\} = 0$ ; (ii)  $\mathcal{F}$  is Donkser uniformly in  $P \in \mathcal{P}$  where  $\mathcal{P}$  is such that  $P_0^*(\hat{P}_n \in \mathcal{P}) \to 1$ ; and (iii) the semi-metric  $L_2(\hat{P}_n)$  converges uniformly to  $L_2(P_0)$  in the sense that

$$\sup_{f,g\in\mathcal{F}} \left| \|f - g\|_{L_2(\hat{P}_n)} - \|f - g\|_{L_2(P_0)} \right| \xrightarrow{P_0^*} 0, \tag{2}$$

then  $\mathbb{G}_n^* \overset{P_0^*}{\underset{W}{\leftrightarrow}} \mathbb{G}_0$  in  $\ell^{\infty}(\mathfrak{F})$ .

Lemma 3.7 is a bootstrap version of Lemma 2.8.7 of van der Vaart & Wellner (1996). Theorems 2.8.9 and 2.8.10 of van der Vaart & Wellner (1996) provide sufficient conditions for a class  $\mathcal{F}$  to be uniform Donkser using uniform entropy and bracketing entropy conditions, respectively. We also note that if Fis constant, then uniform square integrability holds, and, by Theorem 2.8.3 of van der Vaart & Wellner (1996), if some measurability conditions are satisfied, then  $\mathcal{F}$  is Donsker uniformly in  $P \in \mathcal{P}$  provided the uniform entropy integral is finite:

$$\int_{0}^{\infty} \sup_{Q} \left\{ \log N(\varepsilon \|F\|_{L_{2}(Q)}, \mathcal{F}, L_{2}(Q)) \right\}^{1/2} \mathrm{d}\varepsilon < \infty,$$
(3)

where the supremum is taken over all finitely discrete probability measures Q on  $\mathfrak{X}$  with  $\int F^2 dQ > 0$ . Finally, we note that

$$\sup_{f,g\in\mathcal{F}} \left| \|f - g\|_{L_2(\hat{P}_n)} - \|f - g\|_{L_2(P_0)} \right| \le \sup_{f,g\in\mathcal{F}} \left\{ \left| (\hat{P}_n - P_0)(f - g)^2 \right| \right\}^{1/2} \le 2 \sup_{f,g\in\mathcal{F}} \left\{ \left| (\hat{P}_n - P_0)(fg) \right| \right\}^{1/2},$$

so  $\sup_{f,g\in\mathcal{F}} \left| (\hat{P}_n - P_0)(fg) \right| \xrightarrow{P_0^*} 0$  implies (2). Hence, as noted in Coyle & van der Laan (2018), in many cases it is not necessary for the bootstrap sampling distribution  $\hat{P}_n$  to be globally consistent for  $P_0$ ; rather, it is sufficient that means of products or squared differences of influence functions under  $\hat{P}_n$  be consistent for means of the same under  $P_0$  uniformly over the class  $\mathcal{F}$  induced by the nuisance estimators  $\eta_n$  and  $\eta_n^*$ .

We now use Lemma 3.7 to show that bootstrap distributions obtained via smoothing through con-

volution satisfy (B1)(b). Specifically, for a sequence of probability measures  $L_n$ , which we will require converges weakly to 0, we say  $\hat{P}_n$  is obtained by convolution of  $\mathbb{P}_n$  with  $L_n$ , and we write  $\hat{P}_n = \mathbb{P}_n * L_n$ , if for any  $B \in \mathcal{B}$ ,

$$\hat{P}_n(B) := \int_{\mathcal{X}} \int_{\mathcal{X}} \mathbf{1}_B(x+y) \, \mathrm{d}\mathbb{P}_n(x) \, \mathrm{d}L_n(y) = \frac{1}{n} \sum_{i=1}^n \int_{\mathcal{X}} \mathbf{1}_B(X_i+y) \, \mathrm{d}L_n(y) = \frac{1}{n} \sum_{i=1}^n L_n(B-X_i).$$
(4)

The most well-known example of smoothing through convolution is the kernel density estimator. Let  $K : \mathfrak{X} \to \mathbb{R}$  be a fixed kernel function and  $h_n > 0$  a possibly random sequence of bandwidths. If  $L_n(B) := \int_B h_n^{-d} K(h_n^{-1}x) dx$  for any  $B \in \mathcal{B}$ , then  $\hat{P}_n := \mathbb{P}_n * L_n$  defines a kernel density estimator with kernel K and bandwidth  $h_n$ .

Properties of smoothing through convolution estimators, including weak convergence of  $n^{1/2}(\hat{P}_n - P_0)$ , were studied in Yukich (1992), van der Vaart (1994), Rost (2000), Radulović & Wegkamp (2000); Gaenssler & Rost (2003), and Beutner & Zähle (2023), among others. In their Theorem 2.1, Gaenssler & Rost (2003) demonstrated that the bootstrap empirical process  $n^{1/2}(\mathbb{P}_n^* - \hat{P}_n)$  converges weakly to  $\mathbb{G}_0$  in  $\ell^{\infty}(\mathcal{F})$  if  $\mathcal{F}$  is equicontinuous and other conditions hold. However, equicontinuity is a strong assumption, and may not hold in some of our applications of interest. For instance, in some cases, the influence function involves indicator functions, which are not continuous. Equicontinuity is used in their result to show that  $\sup_{f \in \mathcal{F}} |(P_0 * L_n)f^2 - P_0f^2| \to 0$ . Weak convergence of  $L_n$  does not generally imply this result, as shown in Example 2.3 of Gaenssler & Rost (2000). However, if  $P_0$  is absolutely continuous, then weak convergence of  $L_n$  does imply that  $\sup_{f \in \mathcal{F}} |(P_0 * L_n)f^2 - P_0f^2| \to 0$ . While the condition that  $P_0$  is dominated by Lebesgue measure  $\lambda$  and the corresponding density function is Lipschitz continuous, then weak convergence of  $L_n$  does imply that  $\sup_{f \in \mathcal{F}} |(P_0 * L_n)f^2 - P_0f^2| \to 0$ . While the condition that  $P_0$  is dominated by Lebesgue measure is strong, it is typically assumed when using kernel density estimation. Proposition 3.8 below formalizes this to provide general conditions under which smoothing through convolution estimators  $\hat{P}_n$  satisfy condition (B1)(b) without assuming equicontinuity of  $\mathcal{F}$ .

**Proposition 3.8.** Suppose  $\mathcal{F}$  is a class of Borel measurable functions with uniformly bounded envelope function F and finite uniform entropy integral as in (3) such that  $\mathcal{F}_{\delta,P} = \{f-g: f, g \in \mathcal{F}, \|f-g\|_{L_2(P)} < \delta\}$  and  $\mathcal{F}_{\infty}^2 = \{(f-g)^2: f, g \in \mathcal{F}\}$  are P-measurable for every  $\delta > 0$  and  $P \in \mathcal{M}$ . If  $P_0$  is absolutely continuous with respect to Lebesgue measure  $\lambda$  with uniformly bounded and Lipschitz continuous density function  $p_0$ , and  $\hat{P}_n = \mathbb{P}_n * L_n$  for a sequence of random measures  $L_n$  converging weakly to Dirac measure at 0, then the conditions of Proposition 3.7 hold, so that  $\mathbb{G}_n^* \overset{P_0}{\underset{W}{\longrightarrow}} \mathbb{G}_0$  in  $\ell^{\infty}(\mathcal{F})$ .

As discussed in Sheehy & Wellner (1992), if  $L_n(B) = \int_B h_n^{-d} K(h_n^{-1}x) dx$  is a kernel density esti-

mator with bandwidth  $h_n$  satisfying  $h_n \to 0$ ,  $nh_n^d/\log(n) \to \infty$  and  $\| d\hat{P}_n/d\lambda - dP_0/d\lambda \|_{\infty} \xrightarrow{\text{a.s.}^*} 0$ , then  $L_n$  converges weakly to Dirac measure at 0. Furthermore,  $\| d\hat{P}_n/d\lambda - dP_0/d\lambda \|_{\infty} \xrightarrow{\text{a.s.}^*} 0$  follows if  $dP_0/d\lambda$  is uniformly continuous,  $nh_n^d/\log(h_n^{-1}) \to \infty$ ,  $|\log h_n|/\log\log(n) \to \infty$ , and  $h_n^d \le ch_{2n}^d$  for some constant c > 0 by Theorem 2.3 of Giné & Guillou (2002).

#### 3.4 Consistency of bootstrap confidence intervals

We now discuss general conditions for asymptotic validity of bootstrap confidence intervals. Conditional asymptotic linearity of the bootstrap is sufficient for asymptotic validity of many bootstrap confidence intervals. Hence, the conditions of Theorem 3.4 in many cases imply that associated bootstrap confidence intervals are asymptotically valid. However, conditional asymptotic linearity or conditional weak convergence of the bootstrap estimator are not *necessary* for asymptotic validity of bootstrap confidence intervals. In some cases, bootstrap confidence intervals are asymptotically confidence intervals are asymptotically valid. We illustrate this phenomenon in detail for several types of bootstrap confidence intervals.

#### **3.4.1** Percentile and percentile *t*-methods

We first consider the percentile and percentile *t*-methods. We note that we are using the terminology of van der Vaart (2000), but that in other literature, what we are calling the percentile method is called the "basic" or "reverse percentile" method. We suppose that  $\sigma_n^{*2}$  is an estimator of  $\sigma_0^2$  based on the bootstrap data. We then define  $\xi_{n,p}^*$  as the *p*th quantile of the conditional distribution of  $[T(\eta_n^*, \mathbb{P}_n^*) - T(\eta_n, \hat{P}_n)]/\sigma_n^*$  given the data, i.e.,

$$\xi_{n,p}^* := \inf \left\{ \xi \in \mathbb{R} : P_W^* \left( \frac{T(\eta_n^*, \mathbb{P}_n^*) - T(\eta_n, \hat{P}_n)}{\sigma_n^*} \le \xi \right) \ge p \right\}.$$

We emphasize that  $[T(\eta_n^*, \mathbb{P}_n^*) - T(\eta_n, \hat{P}_n)]/\sigma_n^*$  is centered around  $T(\eta_n, \hat{P}_n)$  rather than  $T(\eta_n, \mathbb{P}_n) = \psi_n$ for reasons discussed following Theorem 3.4. A two-sided  $(1 - \alpha - \beta)$ -level bootstrap percentile t-method confidence interval is then given by

$$\left\{\psi:\xi_{n,\beta}^* \le \frac{T(\eta_n,\mathbb{P}_n)-\psi}{\sigma_n} \le \xi_{n,1-\alpha}^*\right\} = \left[T(\eta_n,\mathbb{P}_n)-\xi_{n,1-\alpha}^*\sigma_n, T(\eta_n,\mathbb{P}_n)-\xi_{n,\beta}^*\sigma_n\right].$$
(5)

This interval is based on the *t*-statistic  $n^{1/2}[T(\eta_n, \mathbb{P}_n) - T(\eta_0, P_0)]/\sigma_n$ . Typically,  $\alpha$  and  $\beta$  are chosen to be equal, resulting in an equi-tailed confidence interval. Setting  $\sigma_n = \sigma_n^* = 1$  yields the *bootstrap*  percentile method confidence interval

$$\left[T(\eta_n, \mathbb{P}_n) - \xi_{n,1-\alpha}^*, T(\eta_n, \mathbb{P}_n) - \xi_{n,\beta}^*\right].$$
(6)

The percentile *t*-method has been shown to be more accurate than the percentile method in many cases because the studentized statistic is asymptotically pivotal (Hall, 1992). The next result provides conditions under which the bootstrap percentile and percentile *t*-intervals are asymptotically valid.

**Theorem 3.9.** Suppose that  $\hat{P}_n \phi_0^2 \xrightarrow{P_0^*} P_0 \phi_0^2$  and  $(\hat{P}_n - P_0)[\phi_0^2 1\{|\phi_0| > M\}] \xrightarrow{P_0^*} 0$  for every M > 0. If  $S_n^* - S_n = o_{P_W^*}(n^{-1/2})$  and  $R_n^* - R_n = o_{P_W^*}(n^{-1/2})$ , then

$$\sup_{t \in \mathbb{R}} \left| P_W^* \left( T(\eta_n^*, \mathbb{P}_n^*) - T(\eta_n, \hat{P}_n) \le t \right) - P_0^* \left( T(\eta_n, \mathbb{P}_n) - T(\eta_0, P_0) \le t \right) \right| \xrightarrow{P_0^*} 0.$$

and the bootstrap percentile confidence interval defined in (6) has asymptotic confidence level  $1 - \alpha - \beta$ . If in addition  $\sigma_n^2 \xrightarrow{P_0^*} \sigma_0^2$ ,  $\sigma_n^{*2} \xrightarrow{P_0^*} \sigma_0^2$  and  $(S_n + R_n)(\sigma_n^* - \sigma_n) = o_{P_W^*}(n^{-1/2})$ , then

$$\sup_{t \in \mathbb{R}} \left| P_W^* \left( [T(\eta_n^*, \mathbb{P}_n^*) - T(\eta_n, \hat{P}_n)] / \sigma_n^* \le t \right) - P_0^* \left( [T(\eta_n, \mathbb{P}_n) - T(\eta_0, P_0)] / \sigma_n \le t \right) \right| \xrightarrow{P_0^*} 0,$$

and the percentile t-confidence interval defined in (5) has asymptotic confidence level  $1 - \alpha - \beta$ .

Conditions (A1)–(A3) imply that  $S_n$  and  $R_n$  are  $o_{P_0^*}(n^{-1/2})$ , and conditions (B1)–(B3) imply that  $o_{P_W^*}(n^{-1/2})$ . This yields the following Corollary.

**Corollary 3.10.** If conditions (A1)–(A3) and (B1)–(B3) hold, then the bootstrap percentile confidence interval has asymptotic confidence level  $1 - \alpha - \beta$ . If in addition  $\sigma_n^2 \xrightarrow{P_0^*} \sigma_0^2$  and  $\sigma_n^{*2} \xrightarrow{P_0^*} \sigma_0^2$ , then the percentile t-confidence interval defined in (5) has asymptotic confidence level  $1 - \alpha - \beta$ .

Corollary 3.10 demonstrates that if the conditions of Theorems 3.1 and 3.4 hold, so that the estimator is asymptotically linear and the bootstrap estimator is conditionally asymptotically linear, then bootstrap confidence intervals using the percentile and percentile *t*-methods are asymptotically valid. However, Theorem 3.9 demonstrates that the percentile and percentile *t*-methods can yield valid confidence intervals even if (A1)–(A3) and (B1)–(B3) do not hold. Specifically, even if  $S_n$  and  $R_n$  are not  $o_{P_0^*}(n^{-1/2})$ , and  $S_n^*$  and  $R_n^*$  are not  $o_{P_W^*}(n^{-1/2})$ , as long as  $S_n^*$  and  $R_n^*$  are sufficiently good approximations of  $S_n$  and  $R_n$ , respectively, the percentile methods can yield asymptotically valid confidence intervals. This phenomenon was studied in Cattaneo & Jansson (2018, 2022) for kernel-based nuisance estimators. In particular,  $R_n^* - R_n$  can be  $o_{P_W^*}(n^{-1/2})$  under slower rates of convergence of nuisance estimators than those used to demonstrate that (A3) and (B3) hold, or if the estimator is not targeted toward  $\psi$ . Similarly,  $S_n^* - S_n$  can be  $o_{P_W^*}(n^{-1/2})$  under weaker entropy conditions than those used to demonstrate that (A1) and (B1) hold. Hence, in some cases bootstrap confidence intervals are asymptotically valid even when the estimator is not asymptotically linear because bootstrap intervals can automatically correct excess bias in the estimator. We will discuss this in more depth in Sections 4 and 5.

Theorem 3.9 also requires consistency of certain moments of the bootstrap sampling distribution, which is used to establish that  $\mathbb{G}_n^* \phi_0 \stackrel{P_0^*}{\underset{W}{\longrightarrow}} \mathbb{G}_0 \phi_0$ . For the empirical bootstrap, these conditions are implied by the law of large numbers. For smooth bootstraps, these conditions are satisfied if  $\left\| \frac{\mathrm{d}\hat{P}_n}{\mathrm{d}\lambda} - \frac{\mathrm{d}P_0}{\mathrm{d}\lambda} \right\|_{\infty} \stackrel{\mathrm{a.s.}^*}{\longrightarrow} 0.$ 

Theorem 3.9 and Corollary 3.10 both require that  $\sigma_n^*$  is a conditionally consistent estimator of  $\sigma_0$  for validity of the percentile *t*-method. A bootstrap analogue of the influence function-based variance estimator defined in Section 3.1 is  $\sigma_n^{*2} := \mathbb{P}_n^* \phi_n^{*2}$ . In the following lemma, we show that for the empirical bootstrap, conditions (B1) and (B2) imply that the bootstrap influence function-based variance estimator  $\sigma_n^{*2}$  is conditionally consistent. The situation is not quite as straightforward for other types of bootstraps, but it is still the case that conditions (B1)(a), (B2), and the sufficient conditions for (B1)(b) established in Proposition 3.7 together imply conditional consistency of  $\sigma_n^{*2}$ .

**Lemma 3.11.** If there exists a class of measurable functions  $\mathcal{G}$  such that (i)  $P_W^*(\phi_n^{*2} \in \mathcal{G}) \xrightarrow{P_0^*} 1$ , (ii)  $\sup_{g \in \mathcal{G}} |(\hat{P}_n - P_0)g| = o_{P_0^*}(1)$ , (iii)  $\sup_{g \in \mathcal{G}} |(\mathbb{P}_n^* - \hat{P}_n)g| = o_{P_W^*}(1)$ , and (iv)  $P_0(\phi_n^{*2} - \phi_0^2) = o_{P_W^*}(1)$ , then  $\mathbb{P}_n^*\phi_n^{*2} - \sigma_0^2 = o_{P_W^*}(1)$ . Furthermore, condition (B2) implies (iv). For the empirical bootstrap where  $\hat{P}_n = \mathbb{P}_n$ , condition (B1) implies condition (i)-(iii). For any bootstrap  $\hat{P}_n$ , condition (B1)(a) and the conditions of Proposition 3.7 imply conditions (i)-(iii).

Lastly, Theorem 3.9 requires that  $(S_n + R_n)(\sigma_n^* - \sigma_n) = o_{P_W^*}(n^{-1/2})$  for validity of the percentile *t*-method. This is satisfied if conditions (A1)–(A3) hold and  $\sigma_n^* \xrightarrow{P_0^*} \sigma_0$  and  $\sigma_n \xrightarrow{P_0^*} \sigma_0$ . However, if condition (A1) or (A3) do not hold, then a faster rate of convergence of  $\sigma_n^* - \sigma_n$  may be required. For example, Coyle & van der Laan (2018) suggested using targeted estimators for  $\sigma_n^*$  and  $\sigma_n$ , which can yield  $\sigma_n^* - \sigma_0 = O_{P_W^*}(n^{-1/2})$  and  $\sigma_n - \sigma_0 = O_{P_0^*}(n^{-1/2})$  under suitable conditions, so that  $(S_n + R_n)(\sigma_n^* - \sigma_n) = o_{P_W^*}(n^{-1/2})$  as long as  $S_n$  and  $R_n$  are  $o_{P_0^*}(1)$ .

#### 3.4.2 Efron's percentile method

A third method of constructing bootstrap confidence intervals is *Efron's percentile method*, which is sometimes called the percentile method. In this case, the confidence interval is given by  $[\zeta_{n,\beta}^*, \zeta_{n,1-\alpha}^*]$  for  $\zeta_{n,p}^*$  equal to the lower *p*th quantile of the distribution of  $T(\eta_n^*, \mathbb{P}_n^*)$  given the data; i.e.,  $\zeta_{n,p}^* := \inf\{\zeta \in \mathbb{R} :$  $P_W^*(T(\eta_n^*, \mathbb{P}_n^*) \leq \zeta) \geq p\}$ . The next result provides conditions under which Efron's percentile method yields an asymptotically valid confidence interval.

**Theorem 3.12.** Suppose that  $\hat{P}_n \phi_0^2 \xrightarrow{P_0^*} P_0 \phi_0^2$  and  $(\hat{P}_n - P_0)[\phi_0^2 1\{|\phi_0| > M\}] \xrightarrow{P_0^*} 0$  for every M > 0. If  $[R_n + S_n] + [R_n^* + S_n^*] + [T(\eta_n, \hat{P}_n) - T(\eta_n, \mathbb{P}_n)] = o_{P_W^*}(n^{-1/2})$ , then

$$\sup_{t\in\mathbb{R}}|P_W^*\left(T(\eta_n^*,\mathbb{P}_n^*)-T(\eta_n,\mathbb{P}_n)\leq t\right)-P_0^*\left(-\left[T(\eta_n,\mathbb{P}_n)-T(\eta_0,P_0)\right]\leq t\right)|\xrightarrow{P_0^*}0,$$

and Efron's percentile confidence interval  $[\zeta_{n,\beta}^*, \zeta_{n,1-\alpha}^*]$  has asymptotic confidence level  $1 - \alpha - \beta$ .

As with the percentile and percentile *t*-intervals, conditions (A1)–(A3) and (B1)–(B3) imply that  $S_n, R_n, S_n^*$ , and  $R_n^*$  are all  $o_{P_W^*}(n^{-1/2})$ . This yields the following Corollary.

**Corollary 3.13.** If (A1)-(A3) and (B1)-(B3) hold, and  $T(\eta_n, \hat{P}_n) - T(\eta_n, \mathbb{P}_n) = o_{P_0^*}(n^{-1/2})$ , then Efron's percentile confidence interval  $[\zeta_{n,\beta}^*, \zeta_{n,1-\alpha}^*]$  has asymptotic confidence level  $1 - \alpha - \beta$ .

Corollary 3.13 demonstrates that as with the percentile and percentile *t*-methods, if the conditions of Theorems 3.1 and 3.4 hold, then Efron's percentile intervals are asymptotically valid. However, the conditions of Theorem 3.12 differ substantially from those of Theorem 3.9. First, the sums  $S_n^* + S_n$ and  $R_n^* + R_n$  appear in the condition for Efron's percentile method, in contrast with the the differences  $S_n^* - S_n$  and  $R_n^* - R_n$  in the condition for the other percentile methods. Typically,  $S_n^* + S_n + R_n^* + R_n =$  $o_{P_W^*}(n^{-1/2})$  will be established by showing that each summand is  $o_{P_W^*}(n^{-1/2})$ . Hence, Efron's percentile method generally does not have the potential for automatic bias correction. This was also noted in Cattaneo & Jansson (2022). Furthermore, even if Efron's percentile method is asymptotically valid, the appearance of the sums rather than differences means that it can have worse finite sample behavior than the percentile and percentile *t*-methods. This will be investigated further in Sections 4 and 6. Second, the term  $T(\eta_n, \hat{P}_n) - T(\eta_n, \mathbb{P}_n)$  appears in the conditions for Efron's percentile method, but not for the other percentile methods. This is because Efron's percentile method uses the bootstrap distribution of  $T(\eta_n^*, \mathbb{P}_n^*)$  directly without centering. The distribution of  $T(\eta_n^*, \mathbb{P}_n^*)$  is asymptotically symmetric around  $T(\eta_n, \hat{P}_n)$ , which does not equal the original estimator  $\psi_n = T(\eta_n, \mathbb{P}_n)$  for nonempirical bootstraps. If the term  $T(\eta_n, \hat{P}_n) - T(\eta_n, \mathbb{P}_n)$  is not  $o_{\mathbb{P}_0^*}(n^{-1/2})$  then Efron's percentile method may not have asymptotically valid coverage. This may be the case, for instance, if  $\hat{P}_n$  is a distribution based on a kernel density estimator that is not targeted toward the parameter of interest.

#### 3.4.3 Bootstrap Wald method

The final method of constructing bootstrap confidence intervals that we will discuss is the *bootstrap Wald* method, which is based on a normal approximation. We define  $\bar{\sigma}_n^2 := E_W^* \{ n^{1/2} [T(\eta_n^*, \mathbb{P}_n^*) - T(\eta_n, \hat{P}_n)] \}^2$ as the variance of the centered and scaled bootstrap estimator distribution given the data (not to be confused with  $\sigma_n^{*2}$ ). The two-sided  $(1 - \alpha - \beta)$ -level bootstrap Wald confidence interval is then given by

$$\left[T(\eta_n, \mathbb{P}_n) + z_\beta \bar{\sigma}_n n^{-1/2}, T(\eta_n, \mathbb{P}_n) + z_{1-\alpha} \bar{\sigma}_n n^{-1/2}\right],\tag{7}$$

where  $z_p$  is the lower-*p* quantile of the standard normal distribution. In practice, the bootstrap quantiles  $\xi_{n,p}^*$  and  $\zeta_{n,p}^*$  and the bootstrap variance  $\bar{\sigma}_n^2$  are approximated using empirical analogues based on a large number of bootstrap samples. As mentioned in Section 2.3, we ignore the effect of this approximation. The final result of this section provides conditions under which the bootstrap Wald interval is asymptotically valid.

**Theorem 3.14.** Denote  $T_n^* := n^{1/2} [T(\eta_n^*, \mathbb{P}_n^*) - T(\eta_n, \hat{P}_n)]$ . If conditions (A1)-(A3) and (B1)-(B3) hold and  $T_n^*$  is asymptotically uniformly square-integrable in the sense that

$$\lim_{m\to\infty}\limsup_{n\to\infty}E_0^*E_W^*\left[T_n^{*2}\mathbf{1}\{T_n^{*2}\geq m\}\right]=0,$$

then  $\bar{\sigma}_n \xrightarrow{P_0^*} \sigma_0$ , so the bootstrap Wald confidence interval defined in (7) has asymptotic confidence level  $1 - \alpha - \beta$ .

Since the bootstrap Wald method is based on a normal approximation, its asymptotic validity requires conditional weak convergence of the bootstrap to the same normal limit as the original estimator. Hence, this method does not have the same possibility of automatic bias correction as the percentile and percentile t methods. Furthermore, since weak convergence does not imply convergence of moments, Theorem 3.14 also requires asymptotically uniform integrability of the centered and scaled bootstrap estimator.

# 4 Remainder calculations for specific estimator constructions

### 4.1 One-step estimator

In this section, we explain how (A3), (B3), and  $R_n^* - R_n = o_{P_W^*}(n^{-1/2})$  can be verified for several specific estimator constructions T. Throughout this section, if  $\hat{P}_n$  is in the domain of  $\eta$ , then we denote  $\hat{\eta}_n := \eta(\hat{P}_n)$ .

We first consider the one-step construction. Suppose that the parameter of interest  $\psi(P)$  and its influence function  $\phi_P$  depend on P only through  $\eta_P$ , so that with some abuse of notation we can write  $\psi(P) = \psi(\eta_P)$  and  $\phi_P = \phi_{\eta_P}$ . A one-step estimator of  $\psi_0$  based on nuisance estimator  $\eta_n$  is then defined as  $\psi_n := \psi(\eta_n) + \mathbb{P}_n \phi_{\eta_n}$ , which can be represented as  $\psi_n = T(\eta_n, \mathbb{P}_n)$  for  $T(\eta, P) := \psi(\eta) + P\phi_\eta$ . The mean-zero property of influence functions implies that  $T(\eta_0, P_0) = \psi_0$ .

For the one-step estimator,  $R_n = \psi(\eta_n) - \psi(\eta_0) + P_0 \phi_{\eta_n}$ , which is known as the *second-order re*mainder. For so-called strongly differentiable parameters, the second-order remainder term is  $o_{P_0^*}(n^{-1/2})$ under conditions on the rate of convergence of  $\eta_n - \eta_0$  (see, e.g., Chapter 4 of Pfanzagl, 1982). Hence, a benefit of the one-step construction is that  $\eta_n$  does not typically need to be tailored to  $\psi$  for the one-step estimator to be asymptotically linear, though  $\eta_n$  does usually need to satisfy rate and entropy requirements.

We define the bootstrap one-step estimator based on bootstrap nuisance estimator  $\eta_n^*$  as  $\psi_n^* := T(\eta_n^*, \mathbb{P}_n^*) = \psi(\eta_n^*) + \mathbb{P}_n^* \phi_{\eta_n^*}$ . We then have  $R_n^* = \psi(\eta_n^*) - \psi(\eta_n) + \hat{P}_n(\phi_{\eta_n^*} - \phi_{\eta_n})$ . To demonstrate that (B3) holds, we can decompose  $R_n^*$  in two ways:

$$R_{n}^{*} = \left[\psi(\eta_{n}^{*}) - \psi(\eta_{0}) + P_{0}\phi_{\eta_{n}^{*}}\right] - \left[\psi(\eta_{n}) - \psi(\eta_{0}) + P_{0}\phi_{\eta_{n}}\right] + (\hat{P}_{n} - P_{0})(\phi_{\eta_{n}^{*}} - \phi_{\eta_{n}})$$

$$= \left[\psi(\eta_{n}^{*}) - \psi(\hat{\eta}_{n}) + \hat{P}_{n}\phi_{\eta_{n}^{*}}\right] - \left[\psi(\eta_{n}) - \psi(\hat{\eta}_{n}) + \hat{P}_{n}\phi_{\eta_{n}}\right].$$
(8)

For the first decomposition, we note that  $\psi(\eta_n) - \psi(\eta_0) + P_0\phi_{\eta_n}$  is the second-order remainder term defined above, and  $\psi(\eta_n^*) - \psi(\eta_0) + P_0\phi_{\eta_n^*}$  is a second-order remainder term with  $\eta_n^*$  playing the role of  $\eta_n$ . Hence, this term will typically be  $o_{P_W^*}(n^{-1/2})$  under conditions on the conditional rate of convergence of  $\eta_n^* - \eta_0$ . Similarly, the terms  $\psi(\eta_n^*) - \psi(\hat{\eta}_n) + \hat{P}_n\phi_{\eta_n^*}$  and  $\psi(\eta_n) - \psi(\hat{\eta}_n) + \hat{P}_n\phi_{\eta_n}$  in the second decomposition are second-order remainder terms that will typically be  $o_{P_W^*}(n^{-1/2})$  under conditional rates of convergence of  $\eta_n^* - \hat{\eta}_n$  and  $\eta_n - \hat{\eta}_n$ , respectively.

To demonstrate that bootstrap percentile intervals are asymptotically valid, we can decompose

 $R_n^* - R_n$  in two analogous ways:

$$R_{n}^{*} - R_{n} = \left[\psi(\eta_{n}^{*}) - \psi(\eta_{0}) + P_{0}\phi_{\eta_{n}^{*}}\right] - 2\left[\psi(\eta_{n}) - \psi(\eta_{0}) + P_{0}\phi_{\eta_{n}}\right] + (\hat{P}_{n} - P_{0})(\phi_{\eta_{n}^{*}} - \phi_{\eta_{n}})$$

$$= \left[\psi(\eta_{n}^{*}) - \psi(\hat{\eta}_{n}) + \hat{P}_{n}\phi_{\eta_{n}^{*}}\right] - \left[\psi(\eta_{n}) - \psi(\eta_{0}) + P_{0}\phi_{\eta_{n}}\right] - \left[\psi(\eta_{n}) - \psi(\hat{\eta}_{n}) + \hat{P}_{n}\phi_{\eta_{n}}\right].$$
(9)

As discussed in Section 3.4, these two expressions can be  $o_{P_W^*}(n^{-1/2})$  even if  $R_n$  and/or  $R_n^*$  are not. For instance, in the first decomposition,  $\psi(\eta_n^*) - \psi(\eta_0) + P_0 \phi_{\eta_n^*}$  may be within  $o_{P_W^*}(n^{-1/2})$  of  $2[\psi(\eta_n) - \psi(\eta_0) + P_0 \phi_{\eta_n}]$  even if each of these terms individually is not  $o_{P_W^*}(n^{-1/2})$ . Similarly, in the second decomposition,  $\psi(\eta_n^*) - \psi(\hat{\eta}_n) + \hat{P}_n \phi_{\eta_n^*}$  may be within  $o_{P_W^*}(n^{-1/2})$  of  $\psi(\eta_n) - \psi(\eta_0) + P_0 \phi_{\eta_n}$  even if each of these terms individually is not  $o_{P_W^*}(n^{-1/2})$  of  $\psi(\eta_n) - \psi(\eta_0) + P_0 \phi_{\eta_n}$  even if each of these terms individually is not  $o_{P_W^*}(n^{-1/2})$ .

The bootstrap sampling distribution is a primary consideration when determining which of the two decompositions in (8) and (9) should be used. The first decomposition is more suitable for the empirical bootstrap  $\hat{P}_n = \mathbb{P}_n$ . This is because for most of our applications of interest, empirical distributions are not in the domain of the nuisance parameter  $\eta$ , so that  $\hat{\eta}_n := \eta(\mathbb{P}_n)$  does not exist. For example, this is the case if  $\eta(P)$  is the Lebesgue density of P. Hence, for the empirical bootstrap, the first set of conditions, which do not involve  $\hat{\eta}_n$ , should typically be used. In this case,  $n^{1/2}(\hat{P}_n - P_0)(\phi_{\eta_n^*} - \phi_{\eta_n}) = \mathbb{G}_n(\phi_{\eta_n^*} - \phi_{\eta_n}) = o_{P_W^*}(1)$  under (A1), (A2), (B1)(a), and (B2). Hence, the empirical bootstrap one-step estimator is conditionally asymptotically linear if (A1)–(A3), (B1)(a), and (B2) hold and  $\psi(\eta_n^*) - \psi(\eta_0) + P_0\phi_{\eta_n^*} = o_{P_W^*}(n^{-1/2})$ . Alternatively, bootstrap percentile confidence intervals based on the empirical bootstrap one-step estimator are consistent if (A1)–(A2), (B1)(a), and (B2) hold and  $2[\psi(\eta_n) - \psi(\eta_0) + P_0\phi_{\eta_n}] - [\psi(\eta_n^*) - \psi(\eta_0) + P_0\phi_{\eta_n^*}] = o_{P_W^*}(n^{-1/2}).$ 

For non-empirical bootstraps where  $\hat{\eta}_n$  is well-defined, we expect the second decomposition in (8) and (9) to be easier to verify. This is because in many cases when  $\hat{P}_n \neq \mathbb{P}_n$ , it may be difficult to show that  $(\hat{P}_n - P_0)(\phi_{\eta_n^*} - \phi_{\eta_n}) = o_{P_W^*}(n^{-1/2})$ . For the special case of  $\hat{P}_n$  equal to the distribution corresponding to a kernel density estimator, Section 3.2 of Giné & Nickl (2008) and Theorem 10 of Radulović & Wegkamp (2009) establish asymptotic uniform equicontinuity of  $\{n^{1/2}(\hat{P}_n - P_0)f : f \in \mathcal{F}\}$ under conditions on the bandwidth and smoothness of functions  $f \in \mathcal{F}$ . This implies  $(\hat{P}_n - P_0)(\phi_{\eta_n^*} - \phi_{\eta_n}) = o_{P_W^*}(n^{-1/2})$  under conditions (A1), (A2), (B1)(a), and (B2). In cases where asymptotic uniform equicontinuity of  $\{n^{1/2}(\hat{P}_n - P_0)f : f \in \mathcal{F}\}$  is hard to establish, but  $\hat{P}_n$  and  $P_0$  are dominated by a fixed measure  $\lambda$ , a simple but crude approach to showing (B2) is to use the Cauchy-Schwartz inequality:  $|(\hat{P}_n - P_0)(\phi_{\eta_n^*} - \phi_{\eta_n})| \leq ||d\hat{P}_n - dP_0||_{L_2(\lambda)}||\phi_{\eta_n^*} - \phi_{\eta_n}||_{L_2(\lambda)}$ . Alternatively, a more direct calculation may be employed.

An important special case is  $\eta(\hat{P}_n) = \eta_n$ ; i.e., the bootstrap sampling distribution is based on the original nuisance estimator. For instance, if  $\eta_n$  is a Lebesgue density estimator, this would correspond to drawing bootstrap samples from the distribution corresponding to  $\eta_n$ . If  $\eta(\hat{P}_n) = \eta_n$  then  $\hat{\eta}_n = \eta_n$ , and hence the second decomposition in (8) reduces to  $\psi(\eta_n^*) - \psi(\hat{\eta}_n) + \hat{P}_n \phi_{\eta_n^*}$ , and the second decomposition in (9) reduces to  $[\psi(\eta_n) - \psi(\eta_0) + P_0\phi_{\eta_n}] - [\psi(\eta_n^*) - \psi(\hat{\eta}_n) + \hat{P}_n\phi_{\eta_n^*}].$ 

If we use the original nuisance estimator for the bootstrap estimator, i.e.  $\eta_n^* = \eta_n$ , then  $R_n^* = 0$ . This leads to the following corollary to Theorem 3.4.

**Corollary 4.1.** For the bootstrap one-step estimator with  $\eta_n^* = \eta_n$ , if conditions (A1)(a), (B1)(b) and (A2) hold, then  $T(\eta_n, \mathbb{P}_n^*) = T(\eta_n, \hat{P}_n) + (\mathbb{P}_n^* - \hat{P}_n)\phi_0 + o_{P_W^*}(n^{-1/2})$ . In particular, for the empirical bootstrap  $\hat{P}_n = \mathbb{P}_n$ , conditions (A1) and (A2) imply the result.

Corollary 4.1 indicates that a subset of the conditions for asymptotic linearity implies consistency of the empirical bootstrap for the one-step estimator with the original nuisance estimator. This is convenient as it means that in this case, no additional work needs to be done to establish asymptotic validity of bootstrap confidence intervals beyond that for establishing asymptotic linearity of the estimator. However, it also means that consistency of the percentile and percentile *t*-intervals requires that  $R_n = o_{P_0^*}(n^{-1/2})$ , so automatic bias correction of bootstrap confidence intervals does not occur.

#### 4.2 Plug-in estimator

Our next example of an estimator construction is the plug-in estimator. Suppose that the parameter of interest  $\psi(P)$  depends on P only through  $\eta(P)$ , so that, with some abuse of notation, we can write  $\psi(P) = \psi(\eta_P)$ . A plug-in estimator is then given by  $\psi_n = \psi(\eta_n)$ , which can be represented as  $\psi_n = T(\eta_n, \mathbb{P}_n)$  for  $T(\eta, P) = \psi(\eta)$ . In this case, T is a function of  $\eta$  alone, but for consistency of notation, we will continue to write it as a function of P as well.

For the plug-in estimator,  $R_n = [\psi(\eta_n) - \psi(\eta_0) + P_0\phi_n] - \mathbb{P}_n\phi_n$ . If the influence function  $\phi_P = \phi_{\eta_P,\pi_P}$ depends on  $\eta_P$  and an additional nuisance parameter  $\pi_P$  and we set  $\phi_n = \phi_{\eta_n,\pi_n}$  for  $\pi_n$  an estimator of  $\pi_P$ , then  $\psi(\eta_n) - \psi(\eta_0) + P_0\phi_{\eta_n,\pi_n}$  is a second-order remainder term that will typically be  $o_{P_0^*}(n^{-1/2})$ under sufficient rates of convergence of  $\eta_n$  to  $\eta_0$  and  $\pi_n$  to  $\pi_0$  in appropriate semi-metrics, as discussed in Section 4.1. Plug-in estimators also require that  $\mathbb{P}_n\phi_n = o_{P_0^*}(n^{-1/2})$ , which typically requires careful construction of  $\eta_n$ , as discussed in Section 3.1. We let  $\eta_n^*$  be a bootstrap nuisance estimator, and we define  $\psi_n^* = T(\eta_n^*, \mathbb{P}_n^*) = \psi(\eta_n^*)$  as the bootstrap plug-in estimator. We then have  $R_n^* = [\psi(\eta_n^*) - \psi(\eta_n) + \hat{P}_n \phi_n^*] - \mathbb{P}_n^* \phi_n^*$ . To demonstrate that (B3) holds, we can decompose  $R_n^*$  in two ways:

$$R_{n}^{*} = [\psi(\eta_{n}^{*}) - \psi(\eta_{0}) + P_{0}\phi_{n}^{*}] - [\psi(\eta_{n}) - \psi(\eta_{0}) + P_{0}\phi_{n}] + (\hat{P}_{n} - P_{0})(\phi_{n}^{*} - \phi_{n}) + [\hat{P}_{n}\phi_{n} - \mathbb{P}_{n}^{*}\phi_{n}^{*}]$$

$$= [\psi(\eta_{n}^{*}) - \psi(\hat{\eta}_{n}) + \hat{P}_{n}\phi_{n}^{*}] - [\psi(\eta_{n}) - \psi(\hat{\eta}_{n}) + \hat{P}_{n}\phi_{n}] + [\hat{P}_{n}\phi_{n} - \mathbb{P}_{n}^{*}\phi_{n}^{*}]$$
(10)

If  $\phi_n = \phi_{\eta_n,\pi_n}$  and  $\phi_n^* = \phi_{\eta_n^*,\pi_n^*}$ , then the first three terms in square braces of the first decomposition and the first two terms in square braces of the second decomposition are second-order remainder terms. Negligibility of these terms was discussed following (8). Compared with (8), both decompositions in (10) additionally involve the term  $\hat{P}_n \phi_n - \mathbb{P}_n^* \phi_n^*$ . Ensuring that this term is  $o_{P_W^*}(n^{-1/2})$  typically requires careful construction of  $\eta_n^*$  and  $\eta_n$ , as discussed in Section 3.1. If  $\hat{P}_n = \mathbb{P}_n$  is the empirical bootstrap, then  $\hat{P}_n \phi_n = o_{P_0^*}(n^{-1/2})$  is typically required for (A3) to hold, as discussed above. If this holds, then it is sufficient that  $\mathbb{P}_n^* \phi_n^* = o_{P_W^*}(n^{-1/2})$ , which is the bootstrap analogue of  $\mathbb{P}_n \phi_n = o_{P_0^*}(n^{-1/2})$ . Alternatively, if  $\phi_n = \phi_{\eta_n,\pi_n}$  and  $\hat{P}_n$  is based on  $(\eta_n,\pi_n)$ , then  $\hat{P}_n \phi_n = 0$ , so it is again sufficient that  $\mathbb{P}_n^* \phi_n^* = o_{P_W^*}(n^{-1/2})$ .

To demonstrate that bootstrap percentile intervals are asymptotically valid, we can decompose  $R_n^* - R_n$  in two analogous ways:

$$R_{n}^{*} - R_{n} = [\psi(\eta_{n}^{*}) - \psi(\eta_{0}) + P_{0}\phi_{n}^{*}] - 2[\psi(\eta_{n}) - \psi(\eta_{0}) + P_{0}\phi_{n}] + (\hat{P}_{n} - P_{0})(\phi_{n}^{*} - \phi_{n}) + [\hat{P}_{n}\phi_{n} + \mathbb{P}_{n}\phi_{n} - \mathbb{P}_{n}^{*}\phi_{n}^{*}] = [\psi(\eta_{n}^{*}) - \psi(\hat{\eta}_{n}) + \hat{P}_{n}\phi_{n}^{*}] - [\psi(\eta_{n}) - \psi(\eta_{0}) + P_{0}\phi_{n}] - [\psi(\eta_{n}) - \psi(\hat{\eta}_{n}) + \hat{P}_{n}\phi_{n}] + [\hat{P}_{n}\phi_{n} + \mathbb{P}_{n}\phi_{n} - \mathbb{P}_{n}^{*}\phi_{n}^{*}]$$
(11)

As mentioned in Section 3.4 and Section 4.1, these two expressions can be  $o_{P_W^*}(n^{-1/2})$  even if  $R_n$  and/or  $R_n^*$  are not. The first three terms in square braces of both decompositions were discussed in Section 4.1. The bootstrap sampling distribution is a primary consideration when determining which of the two decompositions in (10) and (11) should be used. We refer readers to Section 4.1 for further discussion.

To show that  $R_n^* - R_n = o_{P_W^*}(n^{-1/2})$  using either decomposition in (11), it may be necessary to additionally show that  $\hat{P}_n \phi_n + \mathbb{P}_n \phi_n - \mathbb{P}_n^* \phi_n^* = o_{P_W^*}(n^{-1/2})$ . For the empirical bootstrap  $\hat{P}_n = \mathbb{P}_n$ , this is the case if  $\mathbb{P}_n^* \phi_n^*$  is within  $o_{P_W^*}(n^{-1/2})$  of  $2\mathbb{P}_n \phi_n$ . If instead  $\hat{P}_n$  is based on  $(\eta_n, \pi_n)$ , then it is sufficient that  $\mathbb{P}_n^* \phi_n^*$  is within  $o_{P_W^*}(n^{-1/2})$  of  $\mathbb{P}_n \phi_n$ . In both cases, the conditions can be met even if  $\mathbb{P}_n \phi_n$  and/or  $\mathbb{P}_n^* \phi_n^*$  are not  $o_{P_0^*}(n^{-1/2})$  and  $o_{P_W^*}(n^{-1/2})$ , respectively. Hence, bootstrap percentile confidence intervals based on a plug-in estimator can be asymptotically valid even when the estimator is not targeted toward the functional of interest.

Lastly, we note that neither (B3) nor  $R_n^* - R_n = o_{P_W^*}(n^{-1/2})$  hold for the bootstrap plug-in estimator if we set  $\eta_n^* = \eta_n$  because this would result in  $\psi_n^* = \psi(\eta_n^*) = \psi(\eta_n) = \psi_n$  with  $P_W^*$  probability 1; i.e., a bootstrap distribution equal to a point mass at the original estimator.

#### 4.3 Empirical mean of a nuisance-dependent function

Our next example of an estimator construction is the empirical mean of a nuisance-dependent function, or empirical mean plug-in estimator for brevity. Suppose that the parameter of interest can be written as  $\psi(P) := Ph_{\eta_P}$  for some known function  $h_{\eta} : \mathfrak{X} \to \mathbb{R}$  depending on  $\eta \in \mathcal{H}$ . A simple estimator is then given by  $\psi_n = \mathbb{P}_n h_{\eta_n}$ , which can be represented as  $\psi_n = T(\eta_n, \mathbb{P}_n)$  for  $T(\eta, P) = Ph_{\eta}$ . In some cases, this estimator and the plug-in estimator considered in Section 4.2 are the same. However, the representations of the two estimators in terms of T are different, which leads to different conditions for asymptotic linearity and especially for consistency of the bootstrap.

We can write the influence function as  $\phi_P(x) = h_{\eta_P}(x) + \gamma_P(x) - \psi(P)$  for  $\gamma_P(x) := \phi_P(x) - h_{\eta_P}(x) + \psi(P)$ , and we note that  $P\gamma_P = 0$ . Heuristically,  $\gamma_P$  can be viewed as the contribution to the influence function of fluctuating  $\eta_P$  within the model. We write  $\phi_n$  as  $\phi_n(x) = h_{\eta_n}(x) + \gamma_n(x) - \psi_n$  for some estimator  $\gamma_n$  of  $\gamma_0$ . We then have  $R_n = P_0(h_{\eta_n} - h_{\eta_0} + \gamma_n) - \mathbb{P}_n \gamma_n$ , which leads to the following conditions under which (A3) holds. If  $\eta$  and  $\gamma$  are compatible in the sense that there exists  $P' \in \mathcal{M}$  such that  $\eta_{P'} = \eta$  and  $\gamma_{P'} = \gamma$ , then  $\psi(P') - \psi(P) + P\phi_{P'} = P(h_\eta - h_{\eta_P} + \gamma)$  is the second-order remainder term discussed in Section 4.1. Hence, if there exists  $P_n$  such that  $\eta_{P_n} = \eta_n$  and  $\gamma_{P_n} = \gamma_n$ , then  $P_0(h_{\eta_n} - h_{\eta_0} + \gamma_n)$  is a second-order remainder and can be expected to be  $o_{P_0^*}(n^{-1/2})$  if  $(\eta_n, \gamma_n)$  converges to  $(\eta_0, \gamma_0)$  at a sufficient rate in an appropriate semi-metric. The condition that  $\mathbb{P}_n \gamma_n = o_{P_0^*}(n^{-1/2})$  typically requires careful construction of  $\eta_n$ , as discussed in Section 3.1.

As usual, we let  $\psi_n^* = T(\eta_n^*, \mathbb{P}_n^*)$  be the bootstrap empirical mean plug-in estimator for a bootstrap nuisance estimator  $\eta_n^*$ . We also define the bootstrap influence function estimator as  $\phi_n^*(x) = h_{\eta_n^*}(x) + \gamma_n^*(x) - \psi_n^*$ , where  $\gamma_n^*$  is the bootstrap estimator of  $\gamma_0$ . We then have  $R_n^* = \hat{P}_n(h_{\eta_n^*} - h_{\eta_n} + \gamma_n^*) - \mathbb{P}_n^* \gamma_n^*$ . To demonstrate that (B3) holds, we can decompose  $R_n^*$  in two ways:

$$R_{n}^{*} = P_{0}(h_{\eta_{n}^{*}} - h_{\eta_{0}} + \gamma_{n}^{*}) - P_{0}(h_{\eta_{n}} - h_{\eta_{0}} + \gamma_{n})$$

$$+ (\hat{P}_{n} - P_{0})(h_{\eta_{n}^{*}} - h_{\eta_{n}} + \gamma_{n}^{*} - \gamma_{n}) + (\hat{P}_{n}\gamma_{n} - \mathbb{P}_{n}^{*}\gamma_{n}^{*})$$

$$= \hat{P}_{n}(h_{\eta_{n}^{*}} - h_{\hat{\eta}_{n}} + \gamma_{n}^{*}) - \hat{P}_{n}(h_{\eta_{n}} - h_{\hat{\eta}_{n}} + \gamma_{n}) + (\hat{P}_{n}\gamma_{n} - \mathbb{P}_{n}^{*}\gamma_{n}^{*}t).$$
(12)

If  $\eta_n^*$  and  $\gamma_n^*$  are compatible and  $\eta_n$  and  $\gamma_n$  are compatible, then the first two terms of both decompositions are second-order remainders. We discussed negligibility of these terms and of the third term in the first decomposition in Section 4.1. Both decompositions in (12) also have the term  $\hat{P}_n\gamma_n - \mathbb{P}_n^*\gamma_n^*$ , and negligibility of this term typically requires careful construction of  $\eta_n^*$  and  $\eta_n$ , as discussed in Section 4.2.

We can decompose  $R_n^* - R_n$  in two analogous ways:

$$R_{n}^{*} - R_{n} = P_{0}(h_{\eta_{n}^{*}} - h_{\eta_{0}} + \gamma_{n}^{*}) - 2P_{0}(h_{\eta_{n}} - h_{\eta_{0}} + \gamma_{n}) + (\hat{P}_{n} - P_{0})(h_{\eta_{n}^{*}} - h_{\eta_{n}} + \gamma_{n}^{*} - \gamma_{n}) + (\hat{P}_{n}\gamma_{n} + \mathbb{P}_{n}\gamma_{n} - \mathbb{P}_{n}^{*}\gamma_{n}^{*}) = \hat{P}_{n}(h_{\eta_{n}^{*}} - h_{\hat{\eta}_{n}} + \gamma_{n}^{*}) - P_{0}(h_{\eta_{n}} - h_{\eta_{0}} + \gamma_{n}) - \hat{P}_{n}(h_{\eta_{n}} - h_{\hat{\eta}_{n}} + \gamma_{n}) + (\hat{P}_{n}\gamma_{n} + \mathbb{P}_{n}\gamma_{n} - \mathbb{P}_{n}^{*}\gamma_{n}^{*}).$$
(13)

As with the one-step and plug-in estimators, these two expressions can be  $o_{P_W^*}(n^{-1/2})$  even if  $R_n$  and/or  $R_n^*$  are not; we refer readers to the discussions following (9) and (11).

Lastly, we note that whether condition (B3) holds if we set  $\eta_n^* = \eta_n$  depends on the particular problem. If  $\gamma_P$  is a function of P through  $\eta_P$ , then condition (B3) usually will not hold for either the empirical or non-empirical bootstraps. This is because in this case,  $\eta_n^* = \eta_n$  implies that  $\gamma_n^* = \gamma_n$ , so that even if  $\mathbb{P}_n \gamma_n = o_{P_0^*}(n^{-1/2})$ , it does not typically follow that  $\mathbb{P}_n^* \gamma_n^* = \mathbb{P}_n^* \gamma_n = o_{P_W^*}(n^{-1/2})$ . Similarly,  $\hat{P}_n \gamma_n + \mathbb{P}_n \gamma_n - \mathbb{P}_n^* \gamma_n^* = \mathbb{P}_n \gamma_n - n^{-1/2} \mathbb{G}_n^* \gamma_n$  will not be typically be  $o_{P_W^*}(n^{-1/2})$ , so that  $R_n^* - R_n = o_{P_W^*}(n^{-1/2})$  may not hold. However, if  $\gamma_P$  involves additional summaries of P beyond  $\eta_P$ , then it may be possible to construct  $\eta_n$  in such a way that both  $\mathbb{P}_n \gamma_n = o_{P_0^*}(n^{-1/2})$  and  $\mathbb{P}_n^* \gamma_n^* = o_{P_W^*}(n^{-1/2})$ , or such that  $R_n^* - R_n = o_{P_W^*}(n^{-1/2})$ .

#### 4.4 Estimating equations-based estimator

We next consider estimating equations-based estimators. Suppose that the influence function  $\phi_P$  depends on P through  $\psi(P)$  and  $\eta_P$ , so that we can write  $\phi_P = \phi_{\psi(P),\eta_P}$ . For each  $P \in \mathcal{M}^+$  and

 $\eta \in \mathcal{H}$ , we define the estimating function  $G_{P,\eta} : \psi \mapsto P\phi_{\psi,\eta}$ , and we assume for simplicity that for each  $(P,\eta) \in \mathcal{M}^+ \times \mathcal{H}$ , there is a unique  $\psi \in \mathbb{R}$  such that  $G_{P,\eta}(\psi) = 0$ . For each  $\eta \in \mathcal{H}$ , we define the true population estimating function as  $G_{0,\eta} := G_{P_0,\eta}$ , and we define the sample estimating function as  $G_{n,\eta_n} := G_{\mathbb{P}_n,\eta_n}$ , where  $\eta_n$  is a nuisance estimator. If  $\psi_0$  is the unique solution to  $G_{0,\eta_0}(\psi_0) = 0$ , then an estimating equation-based estimator  $\psi_n$  of  $\psi_0$  is defined as the solution to  $G_{n,\eta_n}(\psi_n) = 0$ . We can then write  $\psi_n = T(\eta_n, \mathbb{P}_n)$  for  $T(\eta, P)$  defined as the solution to  $G_{P,\eta} = 0$ . We then have  $R_n = \psi_n - \psi_0 + P_0 \phi_{\psi_n,\eta_n}$ . We also note that  $\psi_n$  does not need to solve the equation exactly; it suffices that  $G_{n,\eta_n}(\psi_n) = o_{P_0^*}(n^{-1/2})$ . The derivations for estimating equations-based estimators are more complicated than those for other estimator constructions we have considered, so here we provide theorems for clarity.

Estimating equations-based estimators have been studied in a variety of contexts. Estimating equations often arise in semiparametric models indexed by the pair  $(\psi, \eta) \in \mathbb{R} \times \mathcal{H}$ . We do not explicitly require this, but our results can be applied in this setting. It is sometimes assumed that  $\psi_n$  approximately solves a "profile" estimating equation  $\psi \mapsto \mathbb{P}_n \phi_{\psi,\eta_n(\psi)} = 0$ , where  $\eta_n(\psi)$  is a solution to  $\eta \mapsto \mathbb{P}_n \phi_{\psi,\eta}$ (see, e.g., Murphy & van der Vaart, 2000 and Chapter 21 of Kosorok, 2008). This is the case when  $(\psi_n, \eta_n)$  are defined as the joint optimizers of a criterion function such as a likelihood. In contrast, our goal is to permit  $\eta_n$  to be an arbitrary nuisance estimator satisfying certain rate and entropy conditions, and to use the estimating equation provided by the influence function to mitigate the asymptotic bias, as the one-step estimator does. This approach is more in line with van der Laan & Robins (2003) and Chapter 5 of van der Vaart (2000), among others. In particular, Theorem 5.31 of van der Vaart (2000) provides conditions for asymptotic linearity of an estimating equations-based estimator  $\psi_n$  as we have defined it above. We now provide a slightly reformulated version of this result under which (A3) holds for an estimating equations-based estimator. Following van der Vaart (2000) and others, we say that  $\psi_0$ is a *well-separated* solution of the population estimating equation if  $P_0\phi_{\psi_0,\eta_0} = 0 < \inf_{|\psi-\psi_0|>\delta} |P_0\phi_{\psi,\eta_0}|$ for every  $\delta > 0$ .

Lemma 4.2. If condition (A1) holds for  $\phi_n := \phi_{\psi_n,\eta_n}$ ,  $\psi_0$  is a well-separated solution of the population estimating equation,  $\psi_n = O_{P_0^*}(1)$ , there exists a map  $G'_{0,\eta} : \mathbb{R} \to \mathbb{R}$  depending on  $\eta \in \mathcal{H}$  such that  $\Gamma_{0,\eta} :$  $\psi \mapsto G_{0,\eta}(\psi) - G_{0,\eta}(\psi_0) - G'_{0,\eta}(\psi_0)(\psi - \psi_0)$  satisfies  $\sup_{|\psi| \le M} |\Gamma_{0,\eta_n}(\psi) - \Gamma_{0,\eta_0}(\psi)| = o_{P_0^*}(1)$  for every M > 0 and  $\sup_{\eta: ||\eta - \eta_0||_{\mathcal{H}} < \delta} |\Gamma_{0,\eta}(\psi)| = o(|\psi - \psi_0|)$  for some  $\delta > 0$ ,  $G'_{0,\eta}$  satisfies  $\lim_{\eta \to \eta_0} G'_{0,\eta}(\psi_0) =$  $G'_{0,\eta_0}(\psi_0) = -1$ ,  $||\eta_n - \eta_0||_{\mathcal{H}} = o_{P_0^*}(1)$ , and  $P_0\phi_{\psi_0,\eta_n} = o_{P_0^*}(n^{-1/2})$ , then (A3) holds for the estimating As in Theorem 5.31 of van der Vaart (2000), the assumption that  $\psi_0$  is a well-separated solution of  $\psi \mapsto P_0 \phi_{\psi,\eta_0}$  is used to establish consistency of  $\psi_n$ , and the differentiability assumption is used to ensure that  $n^{1/2}(\psi_n - \psi_0)$  can be linearized. The requirement that  $G'_{0,\eta_0}(\psi_0) = -1$  is not present in Theorem 5.31 of van der Vaart (2000); this is because we are assuming the estimating function *is* the influence function, which requires proper scaling. Theorem 5.31 also permits that the "drift term"  $P_0\phi_{\psi_0,\eta_n}$  contributes to the asymptotic distribution of  $n^{1/2}(\psi_n - \psi_0)$ , while we assume it is negligible since our goal is to establish asymptotic linearity of  $\psi_n$  with influence function  $\phi_0$ . Negligibility of this term is implied by a sufficient rate of convergence of  $\eta_n$  to  $\eta_0$  if  $\eta \mapsto P_0\phi_{\psi_0,\eta}$  is differentiable near  $\eta_0$ in an appropriate sense with derivative map equal to zero (Chernozhukov et al., 2018). However, this is not always the case, and when it fails, the requirement that  $P_0\phi_{\psi_0,\eta_n} = o_{P_0^*}(n^{-1/2})$  requires undersmoothing or otherwise targeting  $\eta_n$ . Hence, estimating equations do not always sufficiently control the asymptotic bias, unlike the one-step estimator. We will see an example of this in Section 5.

We define the bootstrap estimating equations-based estimator as  $\psi_n^* = T(\eta_n^*, \mathbb{P}_n^*)$ ; i.e.,  $\psi_n^*$  is the solution to  $G_{n,\eta_n^*}^*(\psi) := \mathbb{P}_n^* \phi_{\psi,\eta_n^*} = 0$ . As above, it is sufficient that  $G_{n,\eta_n^*}^*(\psi_n^*) = o_{P_W^*}(n^{-1/2})$ . We provide separate sufficient conditions for (B3) for the empirical and smooth bootstraps. First, for the empirical bootstrap  $\hat{P}_n = \mathbb{P}_n$ , we have  $R_n^* = \psi_n^* - \psi_n + \mathbb{P}_n \phi_{\psi_n^*,\eta_n^*}$ , and we have the following result.

**Lemma 4.3.** Let  $\hat{P}_n = \mathbb{P}_n$  be the empirical bootstrap. Suppose the conditions of Lemma 4.2 hold, condition (B1) holds for  $\phi_n^* = \phi_{\psi_n^*,\eta_n^*}$ ,  $\psi_n^* = O_{P_W^*}(1)$ ,  $\sup_{|\psi| \leq M} |\Gamma_{0,\eta_n^*}(\psi) - \Gamma_{0,\eta_0}(\psi_0)| = o_{P_W^*}(1)$  for every M > 0,  $\|\eta_n^* - \eta_0\|_{\mathcal{H}} = o_{P_W^*}(1)$ , and  $P_0\phi_{\psi_0,\eta_n^*} = o_{P_W^*}(1)$ . If  $P_0\phi_{\psi_0,\eta_n^*} = o_{P_W^*}(n^{-1/2})$ , then (B3) holds for the estimating equations-based estimator. If  $P_0\phi_{\psi_0,\eta_n^*} - 2P_0\phi_{\psi_0,\eta_n} = o_{P_W^*}(n^{-1/2})$ , then  $R_n^* - R_n = o_{P_W^*}(n^{-1/2})$ holds for the estimating equations-based estimator.

As discussed above,  $P_0\phi_{\psi_0,\eta_n^*} = o_{P_W^*}(n^{-1/2})$  is sometimes implied by a sufficient rate of convergence of  $\eta_n^*$  to  $\eta_0$ . In particular, if  $\eta_n^* = \eta_n$ , then this condition is implied by  $P_0\phi_{\psi_0,\eta_n} = o_{P_0^*}(n^{-1/2})$ . Hence, similar to Corollary 4.1 for the bootstrap one-step estimator with fixed nuisance estimator, the empirical bootstrap estimating equations-based estimator with fixed nuisance parameter is consistent under the same conditions used for asymptotic linearity of the estimating equations-based estimator. Lemma 4.3 also shows that  $R_n^* - R_n = o_{P_W^*}(n^{-1/2})$  can hold even when  $P_0\phi_{\psi_0,\eta_n}$  is not  $o_{P_0^*}(n^{-1/2})$  or  $P_0\phi_{\psi_0,\eta_n^*}$  is not  $o_{P_W^*}(n^{-1/2})$ , as long as  $P_0\phi_{\psi_0,\eta_n^*}$  is within  $o_{P_W^*}(n^{-1/2})$  of  $2P_0\phi_{\psi_0,\eta_n}$ . This is analogous to discussions about validity of bootstrap percentile intervals above. Consistency of the empirical bootstrap for estimating equations-based estimators was also studied in Wellner & Zhan (1996) and Cheng & Huang (2010). Wellner & Zhan (1996) generalized Theorem 3.3.1 of van der Vaart & Wellner (1996) to weighted bootstraps. Cheng & Huang (2010) focused on the case of a semiparametric model where  $\psi_n^*$  solves a profile bootstrap estimating equation. As discussed above, our results can be used in this context, but also permit  $\eta_n^*$  to be constructed in a different manner than  $\eta_n$ , and can be applied in nonparametric models as well. In addition, Lemma 4.3 also addresses consistency of certain bootstrap confidence intervals without conditional weak convergence, which was not studied in these earlier papers.

The next result provides sufficient conditions for (B3) for the bootstrap estimating equations-based estimator that is applicable to non-empirical bootstrap distributions. We denote  $\hat{\psi}_n := T(\hat{\eta}_n, \hat{P}_n)$ . By definition, we have  $R_n^* = \psi_n^* - \psi_n^\circ + \hat{P}_n \phi_{\psi_n^*, \eta_n^*}$ , where  $\psi_n^\circ := T(\eta_n, \hat{P}_n)$ . Below, we define  $\hat{G}_{n,\eta}(\psi) := \hat{P}_n \phi_{\psi,\eta}$ .

Lemma 4.4. Let  $\hat{P}_n$  be the smooth bootstrap. Suppose the conditions of Lemma 4.2 hold, condition (B1) holds for  $\phi_n^* = \phi_{\psi_n^*,\eta_n^*}, \psi_n^* = O_{P_W^*}(1), \|\eta_n^* - \eta_0\|_{\mathcal{H}} = o_{P_W^*}(1), \sup_{|\psi| \leq M} |\Gamma_{0,\hat{\eta}_n}(\psi) - \Gamma_{0,\eta_0}(\psi)| = o_{P_0^*}(1)$  for every M > 0, there exists a map  $\hat{G}'_{n,\eta} : \mathbb{R} \to \mathbb{R}$  such that  $\hat{\Gamma}_{n,\eta} : \psi \mapsto \hat{G}_{n,\eta}(\psi) - \hat{G}_{n,\eta}(\hat{\psi}_n) - \hat{G}'_{n,\eta}(\hat{\psi}_n)(\psi - \hat{\psi}_n)$  satisfies  $\sup_{|\psi| \leq M} |\hat{\Gamma}_{n,\eta_n^*}(\psi) - \hat{\Gamma}_{n,\hat{\eta}_n}(\hat{\psi}_n)| = o_{P_W^*}(1)$  for every M > 0 and  $\sup_{\eta: \|\eta - \hat{\eta}_n\|_{\mathcal{H}} < \delta} |\hat{\Gamma}_{n,\eta}(\psi)| = o_{P_0^*}(\psi - \hat{\psi}_n)$  for some  $\delta > 0$ ,  $\hat{G}'_{n,\eta}$  satisfies  $\hat{G}'_{n,\eta_n^*}(\hat{\psi}_n) + 1 = o_{P_W^*}(1)$ , there exists a P<sub>0</sub>-Glivenko Cantelli class  $\mathcal{F}$  such that  $\phi_{\hat{\psi}_n,\hat{\eta}_n}$  is contained in  $\mathcal{F}$  with probability tending to one,  $\phi_{\psi_n^*,\eta_n^*}$  is contained in  $\mathcal{F}$  with conditional probability tending to one, and  $\|\hat{P}_n - P_0\|_{\mathcal{F}} = o_{P_W^*}(1)$ ,  $\hat{\psi}_n = O_{P_0^*}(1)$ ,  $P_0\phi_{\psi_0,\hat{\eta}_n} = o_{P_0^*}(1)$ ,  $\|\hat{\eta}_n - \eta_0\|_{\mathcal{H}} = o_{P_0^*}(1)$ , and  $\psi_n^\circ - \hat{\psi}_n = o_{P_0^*}(n^{-1/2})$ . If  $\hat{P}_n\phi_{\hat{\psi}_n,\eta_n^*} = o_{P_W^*}(n^{-1/2})$ , then (B3) holds for the estimating equations-based estimator. If  $\hat{P}_n\phi_{\hat{\psi}_n,\eta_n^*} - P_0\phi_{\psi_0,\eta_n} = o_{P_W^*}(n^{-1/2})$ , then  $R_n^* - R_n = o_{P_W^*}(n^{-1/2})$ holds for the estimating equations-based estimator.

Lemma 4.4 requires  $\hat{P}_n \phi_{\hat{\psi}_n, \eta_n^*} = o_{P_W^*}(n^{-1/2})$  for (B3), which as discussed above is sometimes implied by sufficient rates of convergence of  $\eta_n^* - \hat{\eta}_n$ . However, as in previous results,  $R_n^* - R_n = o_{P_W^*}(n^{-1/2})$  can hold even if  $R_n^*$  is not  $o_{P_W^*}(n^{-1/2})$  and/or  $R_n$  is not  $o_{P_0^*}(n^{-1/2})$ . The conditions  $\|\hat{P}_n - P_0\|_{\mathcal{F}} = o_{P_0^*}(1)$ and  $P_0^*(\phi_{\hat{\psi}_n,\hat{\eta}_n} \in \mathcal{F}) \to 1$  are used to guarantee conditional consistency of  $(\psi_n^*, \eta_n^*)$ . As discussed in Section 3.3, if  $\hat{P}_n$  is a smoothing through convolution estimator, then  $\|\hat{P}_n - P_0\|_{\mathcal{F}} = o_{P_0^*}(1)$  is implied by the conditions of Proposition 3.8. Lastly, Lemma 4.4 requires  $\psi_n^\circ - \hat{\psi}_n = o_{P_0^*}(n^{-1/2})$ . If  $\hat{P}_n$  is based on the original nuisance estimator  $\eta_n$  so that  $\hat{\eta}_n = \eta_n$ , then  $\psi_n^\circ = \hat{\psi}_n$ . Otherwise,  $\psi_n^\circ - \hat{\psi}_n = o_{P_0^*}(n^{-1/2})$ follows if  $P_0^*(\phi_{\psi_n^\circ,\eta_n} \in \mathcal{F}) \to 1$ ,  $\|\phi_{\psi_n^\circ,\eta_n} - \phi_{\psi_0,\eta_0}\|_{L_2(P_0)} = o_{P_0^*}(1)$ , and  $n^{1/2}(\hat{P}_n - P_0)$  converges weakly to a tight measurable limit in  $\ell^{\infty}(\mathcal{F})$ . For the special case of  $\hat{P}_n$  equal to the distribution corresponding to a kernel density estimator, Section 3.2 of Giné & Nickl (2008) and Theorem 10 of Radulović & Wegkamp (2009) establish weak convergence of  $n^{1/2}(\hat{P}_n - P_0)$  under conditions on the bandwidth and smoothness of functions  $f \in \mathcal{F}$ .

# 5 Applications of the general theory

#### 5.1 Average density value

We now illustrate the use of our general results for two bootstrap strategies and three estimators of the average density value parameter. This example has been used extensively as a test case for semiparametric theory and methods (see, e.g., Carone et al., 2019; Cai & van der Laan, 2020, and Cattaneo & Jansson, 2022, among others). We suppose that  $\mathfrak{X} \subseteq \mathbb{R}^d$ , and we let  $\mathfrak{M}$  be the set of probability measures on  $\mathfrak{X}$  dominated by Lebesgue measure  $\lambda$ . For  $P \in \mathfrak{M}$  we let  $\eta_P := dP/d\lambda$  be the Lebesgue density function of P. We then define the average density value parameter as  $\psi(P) :=$  $\int_{\mathfrak{X}} \eta_P(x)^2 dx$ . In this example, the nuisance parameter is the density function  $\eta_P \in \mathfrak{H} := \{h \in L_1(\lambda) :$  $h \geq 0, \int_{\mathfrak{X}} h(x) dx = 1\}$ . The nonparametric efficient influence function of  $\psi$  is  $\phi_P(x) := 2\eta_P(x) - 2\psi(P)$ . For any  $\alpha := (\alpha_1, \ldots, \alpha_d) \in \mathbb{N}^d$  and  $x \in \mathbb{R}^d$ , we define  $|\alpha| := \sum_{i=1}^d \alpha_i, \alpha! := \prod_{i=1}^d \alpha_i, \text{ and } x^{\alpha} := \prod_{i=1}^d x_i^{\alpha_i}$ . For any suitable function  $f : \mathbb{R}^d \to \mathbb{R}$ , we denote  $D^{\alpha}f := \frac{\partial^{\alpha}f}{\partial x_1^{\alpha_1} \cdots \partial x_d^{\alpha_d}}$ . For deterministic sequences  $r_n$ and  $s_n$ , we say  $r_n \prec s_n$  if  $r_n/s_n = o(1)$ .

We consider three approaches to constructing an asymptotically linear estimator of  $\psi_0$ . We let  $\eta_n$ be an estimator of the density  $\eta_0$ . First, we consider the one-step estimator discussed in Section 4.1. In this case,  $T_1(\eta, P) = \psi(\eta) + P\phi_\eta = 2P\eta - \int \eta^2$ , so that the one-step estimator is  $\psi_{n,1} := T_1(\eta_n, \mathbb{P}_n) =$  $2\mathbb{P}_n\eta_n - \int \eta_n^2$ . Second, we consider the plug-in estimator discussed in Section 4.2. In this case,  $T_2(\eta, P) =$  $\int \eta(x)^2 dx$ , so that the plug-in estimator is given by  $\psi_{n,2} := T_2(\eta_n, \mathbb{P}_n) = \int \eta_n(x)^2 dx$ . Third, we consider the mean of a nuisance-dependent function discussed in Section 4.3. In this case,  $g_\eta(x) := \eta(x)$  and  $T_3(\eta, P) = P\eta$ , so that the the estimator is given by  $\psi_{n,3} := T_3(\eta_n, \mathbb{P}_n) = \mathbb{P}_n\eta_n$ . We note that  $\psi_{n,3}$  can also be viewed as an estimating equations-based estimator (discussed in Section 4.4) because it solves  $\psi \mapsto \mathbb{P}_n \phi_{\psi,\eta_n} = 2\mathbb{P}_n\eta_n - 2\psi = 0$ .

The next result uses Theorem 3.1 and the derivations in Section 4 to provide conditions on  $\eta_n$  under which these three estimators are asymptotically linear. **Proposition 5.1.** Suppose that  $\eta_n$  falls in a  $P_0$ -Donsker class with probability tending to one, there exists  $M \in (0,\infty)$  such that  $\|\eta_0\|_{\infty} < M$  and  $\|\eta_n\|_{\infty} < M$  with probability tending to one, and  $\|\eta_n - \eta_0\|_{L_2(\lambda)} = o_{P_0^*}(n^{-1/4})$ . Then  $\psi_{n,1}$  is asymptotically linear with influence function  $\phi_0$ . If in addition  $\int \eta_n^2 - \mathbb{P}_n \eta_n = o_{P_0^*}(n^{-1/2})$ , then  $\psi_{n,2}$  and  $\psi_{n,3}$  are asymptotically linear as well.

We now discuss the conditions of Proposition 5.1 for the specific case where  $\eta_n$  is a kernel density estimator (KDE) with bandwidth  $h \in \mathbb{R}$  (depending on n) and kernel function  $K : \mathbb{R}^d \to \mathbb{R}$ . For KDEs, the Donsker condition is satisfied if  $\eta_n$  falls in a class of functions with uniformly bounded partial derivatives up to order  $\ell > d/2$  and  $P_0$  satisfies a tail bound (van der Vaart, 2000, Example 19.9). If  $\eta_0$  is m > d/2 times differentiable,  $\int [D^{\alpha}\eta_0(x)]^2 dx < \infty$  for all  $\alpha \in \mathbb{N}^d$  with  $|\alpha| = m$  and mth order kernels are used, then  $\|\eta_n - \eta_0\|_{L_2(\lambda)} = O_{P_0^*}(\{nh^d\}^{-1/2} + h^m)$ . Hence, if  $n^{-1/(2d)} \prec h \prec n^{-1/(4m)}$ , then  $\|\eta_n - \eta_0\|_{L_2(\lambda)} = o_{P_0^*}(n^{-1/4})$ . These conditions are satisfied if h is selected at the optimal rate  $h \propto n^{-1/(2m+d)}$ , which yields  $\|\eta_n - \eta_0\|_{L_2(\lambda)} = O_{P_0^*}(n^{-m/(2m+d)})$ .

Proposition 5.1 requires the extra condition that  $\int \eta_n^2 - \mathbb{P}_n \eta_n = o_{P_0^*}(n^{-1/2})$  for asymptotic linearity of the plug-in and empirical mean plug-in estimators. This illustrates the excess bias sometimes incurred by these methods. For  $\eta_n$  a KDE, if  $\eta_0$  is m times differentiable and mth order kernels are used, then  $\int \eta_n^2 - \mathbb{P}_n \eta_n = O_{P_0^*}(\{nh^d\}^{-1} + h^m)$ . Hence, the condition is satisfied if m > d and  $n^{-1/(2d)} \prec h \prec$  $n^{-1/(2m)}$ . This requires that h go to zero faster than the optimal rate  $h \propto n^{-1/(2m+d)}$ , which is why this method is known as *under-smoothing*. It also requires more smoothness of  $\eta_0$ . If  $\eta_0$  is only assumed to be  $m \in (0, d]$  times differentiable, then there is no choice of h that makes  $\int \eta_n^2 - \mathbb{P}_n \eta_n = o_{P_0^*}(n^{-1/2})$ , so in this case it may not be possible to make the bias term for  $\psi_{n,2}$  or  $\psi_{n,3}$  asymptotically negligible using a standard KDE.

We now turn to methods of bootstrapping  $\psi_{n,1}$ ,  $\psi_{n,2}$ , and  $\psi_{n,3}$ . We let  $\eta_n^*$  be a bootstrap estimator of  $\eta_0$ , and we consider the bootstrap estimators  $\psi_{n,1}^* = T_1(\eta_n^*, \mathbb{P}_n^*) = 2\mathbb{P}_n^*\eta_n^* - \int \eta_n^{*2}$ ,  $\psi_{n,2}^* = T_2(\eta_n^*, \mathbb{P}_n^*) =$  $\int \eta_n^{*2}(x) \, dx$ , and  $\psi_{n,3}^* = T_3(\eta_n^*, \mathbb{P}_n^*) = \mathbb{P}_n^*\eta_n^*$ . The next result uses Theorem 3.4 and the derivations in Section 4 to provide conditions for conditional asymptotic linearity of these three estimators.

**Proposition 5.2.** Suppose  $\hat{P}_n = \mathbb{P}_n$  is the empirical bootstrap and the assumptions of Proposition 5.1 hold. For  $\mathcal{F}$  and M defined in Proposition 5.1, suppose  $P_W^*(\eta_n^* \in \mathcal{F}) \xrightarrow{P_0^*} 1$ ,  $P_W^*(||\eta_n^*||_{\infty} \ge M) = o_{P_0^*}(1)$ , and  $||\eta_n^* - \eta_n||_{L_2(\lambda)} = o_{P_W^*}(n^{-1/4})$ . Then  $\psi_{n,1}^* = \psi_{n,1} + (\mathbb{P}_n^* - \mathbb{P}_n)\phi_0 + o_{P_W^*}(n^{-1/2})$ . If in addition  $\int \eta_n^{*2} - \mathbb{P}_n^*\eta_n^* = o_{P_W^*}(n^{-1/2})$ , then  $\psi_{n,2}^* = \psi_{n,2} + (\mathbb{P}_n^* - \mathbb{P}_n)\phi_0 + o_{P_W^*}(n^{-1/2})$  and  $\psi_{n,3}^* = \psi_{n,3} + (\mathbb{P}_n^* - \mathbb{P}_n)\phi_0 + o_{P_W^*}(n^{-1/2})$ . If  $\eta_n$  and  $\eta_n^*$  are KDEs with the same kernel and bandwidth h such that  $n^{-1/(2d)} \prec h$ ,

then  $\|\eta_n^* - \eta_n\|_{L_2(\lambda)} = o_{P_W^*}(n^{-1/4}).$ 

Proposition 5.2 requires that  $\eta_n^*$  converge fast enough to  $\eta_n$  conditional on the data. If  $\eta_n^* = \eta_n$ , then this condition holds automatically. Hence, the empirical bootstrap for the one-step estimator with fixed nuisance is consistent if the conditions of Proposition 5.1 hold, as guaranteed by Corollary 4.1. If  $\eta_n^*$  is estimated using the bootstrap sample, some care must be taken to ensure that  $\|\eta_n^* - \eta_n\|_{L_2(\lambda)} =$  $o_{P_W^*}(n^{-1/4})$ . If the bootstrap nuisance estimator is sensitive to ties in the data, this condition might not hold. However, Proposition 5.2 demonstrates that for KDEs, fixing the bandwidth at the value selected by the original data, as suggested by Hall & Kang (2001), Cattaneo & Jansson (2022), and others, yields  $\|\eta_n^* - \eta_n\|_{L_2(\lambda)} = o_{P_W^*}(n^{-1/4})$ .

Proposition 5.2 requires the extra condition  $\int \eta_n^{*2} - \mathbb{P}_n^* \eta_n^* = o_{P_W^*}(n^{-1/2})$  for conditional asymptotic linearity of the empirical bootstrap plug-in and empirical mean estimators, which is analogous to the condition required by Proposition 5.1 for asymptotic linearity of these estimators. If  $\eta_n^* = \eta_n$ , then this condition reduces to  $\int \eta_n^2 - \mathbb{P}_n^* \eta_n = o_{P_W^*}(n^{-1/2})$ , which does not hold because we can write  $n^{1/2} \left( \int \eta_n^2 - \mathbb{P}_n^* \eta_n \right) = n^{1/2} \left( \int \eta_n^2 - \mathbb{P}_n \eta_n \right) - \mathbb{G}_n^* \eta_n$ , and  $\mathbb{G}_n^* \eta_n$  converges weakly to a non-degenerate limit conditional on the data. Hence, fixing the bootstrap nuisance does not yield a conditionally asymptotically linear plug-in or empirical mean plug-in estimator for the empirical bootstrap for this parameter. If  $\eta_0$  is m > d/2 times differentiable with  $\int [D^{\alpha} \eta_0(x)]^2 dx < \infty$  for all  $|\alpha| = m$ ,  $P_0^*(\eta_n \in \mathcal{F}) \to 1$ , and  $\eta_n^*$  is a KDE based on the bootstrap data with deterministic bandwidth h and mth order kernel functions, then  $\int \eta_n^{*2} - \mathbb{P}_n^* \eta_n^* = O_{P_W^*}(\{nh^d\}^{-1} + h^m\}$ . Hence,  $\int \eta_n^{*2} - \mathbb{P}_n^* \eta_n^* = o_{P_W^*}(n^{-1/2})$  if  $n^{-1/(2d)} \prec h \prec n^{-1/(2m)}$ . This again requires m > d and under-smoothing the bootstrap nuisance estimator.

The next result provides conditions under which the empirical bootstrap percentile method is asymptotically valid.

**Proposition 5.3.** Suppose  $\hat{P}_n = \mathbb{P}_n$  is the empirical bootstrap,  $\eta_0$  is uniformly bounded and m-times continuously differentiable with  $\int [D^{\alpha}\eta_0(x)]^2 dx < \infty$  for all  $|\alpha| = m$ , both  $\eta_n$  and  $\eta_n^*$  are KDEs with common symmetric mth order kernel function and common bandwidth h such that  $n^{-1/d} \prec h$ . If  $h \prec n^{-1/(4m)}$ , then bootstrap percentile intervals based on  $\psi_{n,1}^*$  are asymptotically valid. If  $h \prec n^{-1/(2m)}$ , then bootstrap percentile intervals based on  $\psi_{n,2}^*$  and  $\psi_{n,3}^*$  are asymptotically valid.

The results of Proposition 5.3 agree with those of Cattaneo & Jansson (2022), though they consider a different one-step estimator than ours. The conditions for the one-step estimator in Proposition 5.3 can be satisfied if m > d/4, and the conditions for the plug-in and empirical mean plug-in estimators can be satisfied if m > d/2. Both of these conditions are weaker than the conditions for (conditional) asymptotic linearity from Proposition 5.2, and the conditions for the one-step estimator are again weaker than the plug-in estimators. The conditions for the plug-in and empirical mean plug-in estimators still require undersmoothing. For the one-step estimator, if  $m \in (d/4, d/2]$ , then satisfying  $h \prec n^{-1/(4m)}$ also requires under-smoothing.

In the common case that  $\eta_0$  is assumed to be m = 2 times differentiable and a second-order kernel is used, the (empirical bootstrap) one-step estimator is (conditionally) asymptotically linear for  $d \leq 3$ if  $n^{-1/(2d)} \prec h \prec n^{-1/8}$ , so that the optimal bandwidth may be used. However, bootstrap percentile intervals based on the one-step estimator are asymptotically valid for  $d \leq 7$  as long as  $n^{-1/d} \prec h \prec n^{-1/8}$ , which requires under-smoothing for  $4 \leq d \leq 7$ , but not for  $d \leq 3$ . The (empirical bootstrap) plug-in and empirical mean plug-in estimators are (conditionally) asymptotically linear only for d = 1 if  $n^{-1/2} \prec h \prec$  $n^{-1/4}$ , which requires under-smoothing. Empirical bootstrap percentile intervals based on the plug-in and empirical mean plug-in estimators are asymptotically valid for  $d \leq 3$  if  $n^{-1/d} \prec h \prec n^{-1/4}$ , which again requires under-smoothing.

The next result provides conditions for conditional asymptotic linearity of the three bootstrap estimators when  $\hat{P}_n$  is a smooth bootstrap distribution.

**Proposition 5.4.** Suppose  $\hat{P}_n$  possesses Lebesgue density function  $\hat{\eta}_n$  and the assumptions of Proposition 3.7 and Proposition 5.1 hold. For  $\mathcal{F}$  and M defined in Proposition 5.1, assume that  $\mathcal{F}$  is a  $\mathcal{M}$ uniform Donsker class such that  $P_W^*(\eta_n^* \in \mathcal{F}) \xrightarrow{P_0^*} 1$  and  $P_W^*(\|\hat{\eta}_n\|_{\infty} \ge M) = o_{P_0^*}(1)$ . If  $\|\eta_n^* - \hat{\eta}_n\|_{L_2(\lambda)} = o_{P_W^*}(n^{-1/4})$  and  $\|\hat{\eta}_n - \eta_n\|_{L_2(\lambda)} = o_{P_0^*}(n^{-1/4})$ , then  $\psi_{n,1}^* = \psi_{n,1} + (\mathbb{P}_n^* - \hat{P}_n)\phi_0 + o_{P_0^*}(n^{-1/2})$ . If in addition  $\int \eta_n^{*2} - \mathbb{P}_n^* \eta_n^* = o_{P_W^*}(n^{-1/2}) \text{ and } \int \eta_n^2 - \hat{P}_n \eta_n = o_{P_0^*}(n^{-1/2}), \text{ then } \psi_{n,2}^* = \psi_{n,2} + (\mathbb{P}_n^* - \hat{P}_n)\phi_0 + o_{P_0^*}(n^{-1/2})$ and  $\psi_{n,3}^* = \psi_{n,3} + n^{1/2}(\mathbb{P}_n^* - \hat{P}_n)\phi_0 + o_{P_0^*}(n^{-1/2}).$ 

Proposition 5.4 requires that  $\eta_n^*$  converge fast enough given the data to the density  $\hat{\eta}_n$  used for generating the bootstrap data. If  $\eta_n^*$  is an estimator based on the bootstrap sample, then this can again be achieved by many nonparametric estimators under mild smoothness conditions. For example, if  $\eta_0$ is m > d/2 times differentiable with  $\int [D^{\alpha}\eta_0(x)]^2 dx < \infty$  for all  $|\alpha| = m$ ,  $P_0^*(\hat{\eta}_n \in \mathcal{F}) \to 1$ , and  $\eta_n^*$  is a kernel density estimator with bandwidth  $h^*$  and mth order kernel functions, then  $\|\eta_n^* - \hat{\eta}_n\|_{L_2(\lambda)} = O_{P_W^*}(\{nh^{*d}\}^{-1/2} + h^{*m})$ . Hence, if  $n^{1/(2d)} \prec h^* \prec n^{-1/(4m)}$ , then  $\|\eta_n^* - \hat{\eta}_n\|_{L_2(\lambda)} = o_{P_W^*}(n^{-1/4})$ . Proposition 5.4 also requires that  $\eta_n^*$  falls in a  $\mathcal{P}$ -uniform Donsker class. Furthermore, Proposition 5.4 requires that the conditions of Proposition 3.7 hold. If  $\hat{\eta}_n$  is a kernel density estimator, then Proposition 3.8 can
be used to establish the conditions of Proposition 3.7 and the uniform Donsker condition.

Proposition 5.4 requires the extra conditions  $\int \eta_n^{*2} - \mathbb{P}_n^* \eta_n^* = o_{P_W^*}(n^{-1/2})$  and  $\int \eta_n^2 - \hat{P}_n \eta_n = o_{P_0^*}(n^{-1/2})$  for consistency of the smooth bootstrap plug-in and empirical mean plug-in estimators. If  $\eta_0$  is  $m \ge 2$  times differentiable,  $\eta_n^*$  and  $\hat{\eta}_n$  are KDEs with bandwidths  $h^*$  and  $\hat{h}$ , respectively, an *m*th order kernel function is used for  $\eta_n^*$ ,  $n^{-1/(2d)} \prec h^* \prec n^{-1/(4m)}$ ,  $\hat{h} \prec n^{-1/(2m)}$ , and  $P_0^*(\eta_n \in \mathcal{F}) \to 1$ , then  $\int \eta_n^{*2} - \mathbb{P}_n^* \eta_n^* = o_{P_W^*}(n^{-1/2})$  and  $\int \eta_n^2 - \hat{P}_n \eta_n = o_{P_0^*}(n^{-1/2})$ . Hence, consistency of the smooth bootstrap for the plug-in or empirical mean plug-in estimators with an under-smoothed nuisance estimator also requires under-smoothing the bootstrap sampling distribution.

The next result provides conditions under which the bootstrap percentile method is asymptotically valid when  $\hat{P}_n$  is the distribution corresponding to the kernel density estimator  $\eta_n$ .

**Proposition 5.5.** Suppose that  $\eta_0$  is uniformly bounded and m-times continuously differentiable, and for all  $|\alpha| = m$ ,  $D^{\alpha}\eta_0$  is uniformly bounded and  $\int [D^{\alpha}\eta_0(x)]^2 dx < \infty$ . If both  $\eta_n$  and  $\eta_n^*$  are kernel density estimators with common uniformly bounded symmetric mth order kernel function K and common bandwidth h,  $\hat{P}_n$  is the distribution corresponding to  $\eta_n$ , and  $n^{-1/(2d)} \prec h \prec n^{-1/(4m)}$ , then bootstrap percentile intervals based on  $\psi_{n,1}^*$ ,  $\psi_{n,2}^*$ , and  $\psi_{n,3}^*$  are asymptotically valid.

To the best of our knowledge, Proposition 5.5 is the first result establishing automatic bias correction of bootstrap confidence intervals using the smooth bootstrap. The bandwidth condition  $n^{-1/(2d)} \prec h \prec$  $n^{-1/(4m)}$  in Proposition 5.5 can be satisfied if m > d/2. However, the condition is different from the bandwidth condition of Proposition 5.3 for the empirical bootstrap in several interesting ways. For the one-step estimator, the conditions of Proposition 5.5 are the same as those used for (conditional) asymptotic linearity of the one-step estimator. Hence, unlike the empirical bootstrap, it does not appear that the smooth bootstrap produces valid confidence intervals based on the one-step estimator under weaker smoothness or dimension requirements than non-bootstrap Wald intervals. However, for the plug-in and empirical mean plug-in estimators, the requirements of Proposition 5.5 are weaker than those required for (conditional) asymptotic linearity of the estimators because they require that  $h \prec n^{-1/(4m)}$  rather than  $h \prec n^{-1/(2m)}$ . Hence, both empirical and smooth bootstrap percentile confidence intervals based on the plug-in estimators can be valid if m > d/2. However, the conditions for the plug-in estimators based on the smooth bootstrap are satisfied if the optimal bandwidth is used, while the conditions for the empirical bootstrap require under-smoothing. Thus, in this case, the empirical bootstrap is preferable for the one-step estimator, while the smooth bootstrap is preferable for the plug-in and empirical mean plug-in estimators. We emphasize that it is not presently clear whether these conclusions would remain true with other nuisance estimators or smooth bootstrap sampling distributions, or for other parameter mappings.

#### 5.2 G-computed conditional mean

The second parameter we will use to illustrate the use of our general results is the G-computed conditional mean. Suppose that  $\mathcal{X} = \mathbb{R} \times \{0,1\} \times \mathbb{R}^d$  and X = (Y, A, Z), where  $Y \in \mathbb{R}$  is an outcome of interest,  $A \in \{0,1\}$  is a binary treatment or exposure, and  $Z \in \mathbb{R}^d$  is a vector of adjustment covariates. We then define the G-computed conditional mean as  $\psi(P) = E_P[\mu_P(Z) \mid A = 1]$ , where  $\mu_P(z) := E_P(Y \mid A = 0, Z = z)$ . Under the no unobserved confounding causal model,  $\psi_0$  corresponds to the mean outcome among treated units (i.e., those with A = 1) had they been assigned to receive control A = 0 (Robins, 1986; Gill & Robins, 2001). We use this parameter as an example rather than the simpler G-computed mean,  $E_P[\mu_P(Z)]$  because the one-step and estimating equations-based estimators are different for the conditional mean, which gives us the chance to illustrate the use of our results for estimating equations-based estimators.

The efficient influence function of  $\psi$  at P relative to a nonparametric model is given by

$$\phi_P(y, a, z) = \frac{I(a=0)g_P(z)}{\pi_P \left[1 - g_P(z)\right]} \left[y - \mu_P(z)\right] + \frac{I(a=1)}{\pi_P} \left[\mu_P(z) - \psi_P\right],$$

where  $g_P(z) := P(A = 1 | Z = z)$  is the propensity score function and  $\pi_P := P(A = 1)$ . In this example, the nuisance parameter is  $\eta_P = (\mu_P, g_P, Q_P)$ , where  $Q_P$  is the marginal distribution of Z under P, and  $\psi_P$  and  $\pi_P$  are defined through  $\eta_P$  as  $\psi_P = \int \mu_P(z)g_P(z)\pi_P^{-1} dQ_P(z)$  and  $\pi_P = \int g_P dQ_P$ .

We consider two approaches to constructing an asymptotically linear estimator of  $\psi_0$ . We let  $\eta_n = (\mu_n, g_n, Q_n)$  be an estimator of the nuisance  $\eta_0$ , where  $Q_n$  is the marginal empirical distribution of Z. First, we consider the one-step estimator discussed in Section 4.1, which is given by

$$\psi_{n,1} = \frac{1}{n} \sum_{i=1}^{n} \left\{ \frac{I(A_i = 0)g_n(Z_i)}{\pi_n [1 - g_n(Z_i)]} [Y_i - \mu_n(Z_i)] + \left[2 - \frac{\bar{\pi}_n}{\pi_n}\right] \frac{I(A_i = 1)}{\pi_n} \mu_n(Z_i) \right\},$$

where  $\pi_n := \int g_n dQ_n$  and  $\bar{\pi}_n := \frac{1}{n} \sum_{i=1}^n A_i$ . Second, we consider the estimating equations-based estimator discussed in Section 4.4. We define the estimating function  $G_{P,\eta}(\psi) := P\phi_{\psi,\eta}$ , and we note

that with  $\eta = (\mu, g, Q)$  and  $\pi := \int g \, \mathrm{d}Q$ ,

$$G_{0,\eta}(\psi) = \pi^{-1} P_0 \left[ \frac{(g - g_0)(\mu - \mu_0)}{1 - g} \right] + \frac{\pi_0}{\pi} (\psi_0 - \psi).$$

In particular,  $G_0(\psi) = \psi_0 - \psi$ , so  $\psi_0$  is the unique solution to the population estimating equation, and  $G'_{0,\eta}(\psi) = -\pi_0/\pi$ , which approaches -1 as  $\eta \to \eta_0$ . We then define the estimating equations-based estimator  $\psi_{n,2} = T_2(\eta_n, \mathbb{P}_n)$  as the solution to the sample estimating function  $G_{n,\eta_n}(\psi) := \mathbb{P}_n \phi_{\psi,\eta_n}$ , which it is easy to see equals

$$\psi_{n,2} = \frac{1}{n} \sum_{i=1}^{n} \left\{ \frac{I(A_i = 0)g_n(Z_i)}{\bar{\pi}_n [1 - g_n(Z_i)]} [Y_i - \mu_n(Z_i)] + \frac{I(A_i = 1)}{\bar{\pi}_n} \mu_n(Z_i) \right\}.$$

We note that if  $\pi_n = \bar{\pi}_n$ , then  $\psi_{n,1} = \psi_{n,2}$ . The next result provides conditions under which these two estimators are asymptotically linear using Theorem 3.1.

**Proposition 5.6.** If  $\mu_n$ ,  $g_n$ , and  $(y, a, z) \mapsto (1 - a)yg_n(z)/[1 - g_n(z)]$  fall in  $P_0$ -Donsker classes with probability tending to 1,  $E_0(Y^2) < \infty$ , there exists constants 0 < a < b < 1 such that  $P_0(g_0(Z) \in (a, b)) =$  $1, P_0(g_n(Z) \in (a, b)) = 1$ , and  $P_0(|\mu_n(Z)| \le b) = 1$ ,  $||g_n - g_0||_{L_2(P_0)} = o_{P_0^*}(1)$ ,  $||\mu_n - \mu_0||_{L_2(P_0)} = o_{P_0^*}(1)$ , and  $P_0\{(g_n - g_0)(\mu_n - \mu_0)/(1 - g_n)\} = o_{P_0^*}(n^{-1/2})$ , then  $\psi_{n,2}$  is asymptotically linear with influence function  $\phi_0$ . If in addition  $(\pi_n - \pi_0)(\psi_n - \psi_0) = o_{P_0^*}(n^{-1/2})$ , then  $\psi_{n,1}$  is asymptotically linear with influence function  $\phi_0$ .

We now turn to methods of bootstrapping  $\psi_{n,1}$  and  $\psi_{n,2}$ . We define  $\eta_n^* = (\mu_n^*, g_n^*, Q_n^*)$  as a bootstrap estimator of  $\eta_0$  based on n bootstrap observations, where  $Q_n^*$  is the empirical distribution of the bootstrap covariates. We then consider  $\psi_{n,1}^* = T_1(\eta_n^*, \mathbb{P}_n^*)$  and  $\psi_{n,2}^* = T_2(\eta_n^*, \mathbb{P}_n^*)$ . The next result provides conditions under which these estimators are conditionally asymptotically linear for the empirical bootstrap.

**Proposition 5.7.** Suppose  $\hat{P}_n = \mathbb{P}_n$  is the empirical bootstrap and the conditions of Proposition 5.6 hold. If  $\mu_n^*$ ,  $g_n^*$ , and  $(y, a, z) \mapsto (1-a)yg_n^*(z)/[1-g_n^*(z)]$  fall in  $P_0$ -Donsker classes with conditional probability tending to 1, there exist constants 0 < a < b < 1 such that  $P_W^*(g_n^*(Z) \in (a, b)) = 1$ , and  $P_W^*(|\mu_n^*(Z)| \le b) = 1$ ,  $||g_n^* - g_0||_{L_2(P_0)} = o_{P_W^*}(1)$ ,  $||\mu_n^* - \mu_0||_{L_2(P_0)} = o_{P_W^*}(1)$ , and  $P_0\left\{(g_n^* - g_0)(\mu_n^* - \mu_0)/(1-g_n^*)\right\} = o_{P_W^*}(n^{-1/2})$ , then  $\psi_{n,2}^*$  is conditionally asymptotically linear with influence function  $\phi_0$ . If in addition  $(\pi_n^* - \pi_0)(\psi_n^* - \psi_0) = o_{P_W^*}(n^{-1/2})$ , then  $\psi_{n,1}^*$  is conditionally asymptotically linear with influence function  $\phi_0$ . Proposition 5.7 requires that  $\mu_n^*$  and  $g_n^*$  converge fast enough to  $\mu_0$  and  $g_0$ , respectively, conditional on the data. We also note that, as above, percentile confidence intervals based on the empirical bootstrap might be consistent under weaker conditions than Proposition 5.7.

The next result addresses the case where  $\hat{P}_n$  is a non-empirical bootstrap sampling distribution. We let  $\hat{Q}_n$  be the marginal distribution of Z under  $\hat{P}_n$ ,  $\hat{g}_n(z) := \hat{P}_n(A = 1|Z = z)$ ,  $\hat{\mu}_n(z) := E_{\hat{P}_n}(Y|A = 0, Z = z)$ , and  $\hat{\sigma}_n^2(z) := \operatorname{Var}_{\hat{P}_n}(Y \mid A = 0, Z = z)$ , all of which we assume are well-defined. We note that  $\hat{Q}_n$  need not be a smooth distribution. We also define  $\sigma_0^2(z) := \operatorname{Var}_0(Y \mid A = 0, Z = z)$ . We then have the following result regarding conditional asymptotic linearity of the bootstrap one-step and estimating equations-based estimators when sampling from  $\hat{P}_n$ .

**Proposition 5.8.** Suppose  $P_0^*(\hat{P}_n \in \mathfrak{P}) \to 1$ , where  $\mathfrak{P}$  is such that  $\lim_{M\to\infty} \sup_{P\in\mathfrak{P}} E_P[Y^2I(Y^2 > M)] = 0$ ,  $\hat{\mu}_n \in \mathfrak{F}_\mu$  and  $\hat{g}_n \in \mathfrak{F}_g$  with probability tending to one, where  $\mathfrak{F}_\mu$  is uniformly bounded,  $\mathfrak{F}_g$  is uniformly bounded away from zero, and  $\mathfrak{F}_\mu$  and  $\mathfrak{F}_g$  possess finite uniform entropy integrals,  $\|\hat{g}_n - g_0\|_{L_2(P_0)}$ ,  $\|\hat{\mu}_n - \mu_0\|_{L_2(P_0)}$ ,  $\|\hat{\sigma}_n^2 - \sigma_0^2\|_{L_2(P_0)}$  are each  $o_{P_0^*}(1)$ , and each of the following is  $o_{P_0^*}(1)$ :

$$\sup_{g,\bar{g}} \left| (\hat{Q}_n - Q_0) \left[ \frac{g\bar{g}(1 - \hat{g}_n)}{(1 - g)(1 - \bar{g})} (\hat{\sigma}_n^2 + \hat{\mu}_n^2) \right] \right|, \quad \sup_{g,\bar{g},\mu,\bar{\mu}} \left| (\hat{Q}_n - Q_0) \left[ \frac{g\bar{g}(1 - \hat{g}_n)}{(1 - g)(1 - \bar{g})} \mu \bar{\mu} \right] \right|, \quad (14)$$

$$\sup_{\mu} \left| (\hat{Q}_n - Q_0) \left[ \mu \hat{g}_n \right] \right|, \quad \sup_{\mu,\bar{\mu}} \left| (\hat{Q}_n - Q_0) \left[ \mu \bar{\mu} \hat{g}_n \right] \right|, \quad \sup_{\mu,g} \left| (\hat{Q}_n - Q_0) \left[ \frac{g(1 - \hat{g}_n)}{1 - g} \mu \right] \right|, \quad (\hat{Q}_n - Q_0) \hat{g}_n.$$

where the suprema over  $\mu$  and  $\bar{\mu}$  are taken over  $\mathfrak{F}_{\mu}$  and the suprema over g and  $\bar{g}$  are taken over  $\mathfrak{F}_{g}$ , and  $\hat{Q}_{n} \{(\hat{g}_{n} - g_{n})(\hat{\mu}_{n} - \mu_{n})/(1 - g_{n})\} = o_{P_{0}^{*}}(n^{-1/2})$ . Suppose also that  $\mu_{n}^{*} \in \mathfrak{F}_{\mu}$  and  $g_{n}^{*} \in \mathfrak{F}_{g}$  with conditional probability tending to one,  $\|g_{n}^{*} - g_{0}\|_{L_{2}(P_{0})} = o_{P_{W}^{*}}(1)$ ,  $\|\mu_{n}^{*} - \mu_{0}\|_{L_{2}(P_{0})} = o_{P_{W}^{*}}(1)$ , and  $\hat{Q}_{n} \{(g_{n}^{*} - \hat{g}_{n})(\mu_{n}^{*} - \hat{\mu}_{n})/(1 - g_{n}^{*})\} = o_{P_{W}^{*}}(n^{-1/2})$ . If  $(\pi_{n}^{*} - \hat{\pi}_{n})(\psi_{n}^{*} - \hat{\psi}_{n}) = o_{P_{W}^{*}}(n^{-1/2})$  and  $(\pi_{n} - \hat{\pi}_{n})(\psi_{n} - \hat{\psi}_{n}) = o_{P_{0}^{*}}(n^{-1/2})$ , then  $\psi_{n,1}^{*}$  is conditionally asymptotically linear with influence function  $\phi_{0}$ . If the conditions of Proposition 5.6 hold and  $\pi_{n}^{*} - \hat{\pi}_{n} = o_{P_{W}^{*}}(1)$ , then  $\psi_{n,2}^{*}$  is conditionally asymptotically linear with influence function  $\phi_{0}$ .

Proposition 5.8 illustrates that the bootstrap sampling distribution  $\hat{P}_n$  can produce valid bootstrap confidence intervals even if it is not globally consistent. In this case, it is sufficient that the propensity score, conditional mean, and conditional variance functions induced by  $\hat{P}_n$  be consistent, and that certain means of the marginal distribution of the covariates  $\hat{Q}_n$  be consistent. If  $\hat{Q}_n = Q_n$  is the empirical distribution of the observed covariates, then the conditions in (14) hold by the assumption that  $\mathcal{F}_{\mu}$  and  $\mathcal{F}_{g}$  possess finite uniform entropy integrals. However, it may be of interest to use something other than the empirical distribution for  $\hat{Q}_n$  in order to, for instance, produce unique bootstrap covariate values. We also note that it is possible that stronger notions of consistency of  $\hat{P}_n$  have implications for higher-order properties of bootstrap confidence intervals. Finally, we note that some of the conditions of Proposition 5.8 hold automatically if  $\hat{\mu}_n = \mu_n$  or  $\hat{g}_n = g_n$ .

#### 6 Numerical study

We conducted a simulation study to assess the finite-sample performance of the methods of inference for the average density value parameter studied in Section 5.1. We set  $P_0$  as the standard normal distribution. For each sample size  $n \in \{50, 100, 200, 300, 400, 500, 1000, 2000, 3000, 4000, 5000\}$ , we simulated 1000 datasets of n independent and identically distributed observations from  $P_0$ . For each dataset, we considered the three estimator constructions defined in Section 5: the one-step estimator  $\psi_{n,1} = T_1(\eta_n, \mathbb{P}_n)$ , the plug-in estimator  $\psi_{n,2} = T_2(\eta_n, \mathbb{P}_n)$ , and the empirical mean plug-in estimator  $\psi_{n,3} = T_3(\eta_n, \mathbb{P}_n)$ . For each estimator, we used three different nuisance estimators  $\eta_n$ : (1) a KDE with Gaussian kernel and bandwidth h selected at the optimal rate  $h \propto n^{-1/5}$  for twice-differentiable densities using the method of Sheather & Jones (1991); (2) a KDE with under-smoothed bandwidth  $h/n^{1/10}$ ; and (3) TMLE using (1) as the initial estimator. Hence, we constructed a total of nine distinct estimators for each dataset.

We considered four bootstrap sampling distributions: the empirical distribution, and three smooth distributions corresponding to the three density estimators defined above. For each dataset, we generated B = 1000 bootstrap datasets using these four bootstrap distributions. For each bootstrap dataset, we then considered the same three estimator constructions using the bootstrap data:  $\psi_{n,1}^* = T_1(\eta_n^*, \mathbb{P}_n^*)$ ,  $\psi_{n,2}^* = T_2(\eta_n^*, \mathbb{P}_n^*)$ , and  $\psi_{n,3}^* = T_3(\eta_n^*, \mathbb{P}_n^*)$ . We considered two bootstrap nuisance estimators  $\eta_n^*$ : the same nuisance estimation procedure used for the original data applied to the bootstrap sample, with bandwidth fixed at the value selected using the original data, and using the fixed nuisance estimator obtained from the original data, i.e.  $\eta_n^* = \eta_n$ . Finally, we used all four methods of constructing bootstrap confidence intervals defined in Section 3.4 to construct two-sided, equi-tailed 95% confidence intervals for  $\psi_0$  based on each bootstrap sample. For the percentile *t*-method, we used the influence function-based variance estimator. For comparison, we also constructed ordinary Wald-style confidence intervals using the influence function-based variance estimator. We evaluated the performance of these confidence intervals by computing their empirical coverage and average width over the 1000 simulations.



Figure 1: Empirical coverage of 95% confidence intervals based on the bootstrap one-step estimator when re-estimating the nuisance using the bootstrap sample. "KDE" stands for kernel density estimator; "US" stands for under-smoothed; TMLE stands for targeted maximum likelihood estimator.



Figure 2: Scaled average width of 95% confidence intervals based on the bootstrap one-step estimator when re-estimating the nuisance using the bootstrap sample. Abbreviations as in Figure 1.

We now turn to the results of the simulation study. Figures 1, 3, and 4 display empirical coverage and Figures 2, 5 and 6 display the average width of 95% confidence intervals based on the one-step, plug-in, and empirical mean plug-in estimators, respectively, when the bootstrap nuisance was re-estimated using the bootstrap sample. Figures 7 and 8 display empirical coverage and Figures 9 and 10 display average width based on the one-step and empirical mean plug-in constructions, respectively, when the bootstrap nuisance was fixed. The results for the plug-in construction with the nuisance fixed are not displayed because the coverage in this case was always zero, as discussed in Section 4.2. In each figure, the rows represent the bootstrap sampling distribution and the columns represent the method of construction of the nuisance estimator. For example, the top left panel of Figure 1 shows the empirical coverage rate of confidence intervals based on the one-step estimator where the bootstrap sampling distribution was the empirical distribution  $\mathbb{P}_n$  and the nuisance estimator was the KDE with optimal bandwidth.

We first discuss the results displayed in Figure 1 for the one-step estimator with re-estimated nuisance. Efron's percentile method did not yield valid confidence intervals at large sample sizes when the bootstrap sampling distribution was based on a KDE with optimal bandwidth (second row from the top). This was expected based on the results of Section 3.4. The bias  $T_1(\eta_n, \hat{P}_n) - T_1(\eta_n, \mathbb{P}_n) = (\mathbb{P}_n - \hat{P}_n)\phi_{\eta_n}$ in this case is not  $o_{P_0^*}(n^{-1/2})$  because  $\hat{P}_n$  was not under-smoothed. The coverage of all other confidence intervals for the bootstrap one-step estimator approached 95% as the sample size increased, which is in line with Propositions 5.2 and 5.4. Efron's percentile confidence intervals had poor coverage for small and moderate sample sizes in some cases, which we hypothesize is due to excess bias in this method, as discussed in Section 3.4. The coverage of (non-bootstrap) Wald-type confidence intervals approached 95% in all cases, as expected, though its performance for smaller sample sizes was not always good. An exception was when the nuisance estimator was TMLE (third column from the left), in which case the coverage of the Wald-type estimator was excellent at all sample sizes considered. Otherwise, there was no clear and consistent best nuisance estimator or bootstrap sampling distribution. The average widths scaled by  $n^{1/2}$  displayed in Figure 2 all converge to the same value.

We next discuss the results displayed in Figures 3 and 4 for the plug-in and empirical mean plug-in estimators with re-estimated nuisance. Using the KDE with optimal bandwidth as nuisance estimator did not generally yield valid Wald or bootstrap confidence intervals because neither estimator is asymptotically linear in this case (first column from the left). However, the percentile and percentile-t confidence intervals had close to nominal coverage in large samples for both estimators when both the nuisance estimator and bootstrap sampling distribution were the KDE with optimal bandwidth (second



Figure 3: Empirical coverage of 95% confidence intervals based on the bootstrap plug-in estimator when re-estimating the nuisance using the bootstrap sample. Abbreviations as in Figure 1.



Figure 4: Empirical coverage of 95% confidence intervals based on the bootstrap empirical mean plug-in estimator when re-estimating the nuisance using the bootstrap sample. Abbreviations as in Figure 1.



Figure 5: Scaled average width of 95% confidence intervals based on the bootstrap plug-in estimator when re-estimating the nuisance using the bootstrap sample. Abbreviations as in Figure 1.



Figure 6: Scaled average width of 95% confidence intervals based on the bootstrap empirical mean plug-in estimator when re-estimating the nuisance using the bootstrap sample. Abbreviations as in Figure 1.

row from the top and first column from the left). This was expected based on Proposition 5.5. Besides the bootstrap Wald method, using the KDE with optimal bandwidth as the bootstrap sampling distribution did not yield valid bootstrap confidence intervals in large samples for other nuisance estimators (second row from the top and second and third columns from the left). All other confidence intervals based on the plug-in and empirical mean plug-in estimators with re-estimated nuisance had close to 95% coverage for large sample sizes, which aligns with Propositions 5.1, 5.2, and 5.4. Among the methods with good large-sample coverage, using TMLE as the nuisance estimator and bootstrap sampling distribution along with the Wald, bootstrap Wald, percentile, or percentile *t*-confidence interval yielded the best coverage for small and moderate sample sizes. In many cases, Efron's method again had poor coverage in small and moderate samples. The average widths scaled by  $n^{1/2}$  displayed in Figures 5 and 6 appear to again converge to roughly the same value.

We now turn to the results displayed in Figure 7 for the one-step estimator with fixed bootstrap nuisance. Effon's percentile method again did not yield valid coverage in large samples when the bootstrap sampling distribution was based on a KDE with optimal bandwidth (second row from the top). As discussed following Figure 1, this was expected based on the results of Section 3.4. All other confidence intervals for the bootstrap one-step estimator with fixed bootstrap nuisance estimator had good coverage in large samples, as expected based on Corollary 4.1. Compared to the results in Figure 1 for the bootstrap one-step estimator with re-estimated nuisance, the coverage of bootstrap Wald, percentile, and percentile t-confidence intervals were mostly worse when fixing the bootstrap nuisance than when re-estimating it. However, interestingly, the coverage of Efron's percentile intervals was often better. For the empirical bootstrap, percentile t-confidence intervals had slightly better coverage in small samples than other bootstrap intervals, but in many other cases the various bootstrap methods had very similar coverage when the nuisance was fixed. For many of the cases considered, bootstrap intervals had better coverage in small and moderate samples than Wald-type intervals, indicating there may still be a benefit of the bootstrap even when fixing the bootstrap nuisance. There was not a substantial difference between the bootstrap sampling distributions, but the KDE with optimal bandwidth had the best overall performance in small and moderate samples. As with the bootstrap one-step estimator with the nuisance re-estimated, TMLE had the best performance among the nuisance estimators considered, with excellent coverage even for small sample sizes. The average widths scaled by  $n^{1/2}$  displayed in Figures 9 appear to again converge to roughly the same value.

Finally, from Figures 8 and 10, all of the bootstrap confidence intervals for the bootstrap empirical



Figure 7: Empirical coverage of 95% confidence intervals based on the bootstrap one-step estimator when the nuisance was not re-estimated using the bootstrap sample. Abbreviations as in Figure 1.



Figure 8: Empirical coverage of 95% confidence intervals based on the bootstrap empirical mean plugin estimator when the nuisance was not re-estimated using the bootstrap sample. Abbreviations as in Figure 1.



Figure 9: Scaled average width of 95% confidence intervals based on the bootstrap one-step when the nuisance was not re-estimated using the bootstrap sample. Abbreviations as in Figure 1.



Figure 10: Scaled average width of 95% confidence intervals based on the bootstrap empirical mean plug-in estimator when the nuisance was not re-estimated using the bootstrap sample. Notes as in Figure 1.

mean plug-in estimator with fixed bootstrap nuisance had poor coverage at all sample sizes because they were on average too narrow. This is in line with the results and discussion of Section 4.3.

### 7 Conclusion

In this paper, we studied the problem of bootstrapping asymptotically estimators that rely on a dataadaptive nuisance estimator. We proposed a framework that encompasses many approaches to constructing asymptotically linear estimators and bootstrapping them, and we provided high-level conditions for consistency of the bootstrap in this framework. We also provided more detailed conditions for several bootstrap distributions and estimator constructions. We used our general results to demonstrate that a wide variety of bootstrap confidence intervals are asymptotically valid in this setting, and our simulation study confirmed this. It is our hope that the generality of our framework and theory ensures that there are many potential future applications of our results.

An important area of future research is establishing rates of convergence for the bootstrap methods considered here. We focused on consistency of the bootstrap because it is an important first step, and because we expect that rates of convergence will require stronger assumptions than we used here. However, understanding how the different components of the original estimator and the bootstrap sampling scheme contribute to the accuracy of bootstrap confidence intervals is crucial for deciding which method to use in practice. For instance, while we showed that the precise behavior of the bootstrap nuisance estimator does not play a role in the first-order asymptotic behavior of the bootstrap estimator as long as our high-level conditions hold, we expect that it plays an important role in the finite-sample accuracy of the bootstrap. This was sometimes, but not always, the case in our numerical studies. Similarly, while we showed that both the empirical and smooth bootstraps can yield asymptotically valid bootstrap confidence intervals, our results did not reveal which approach will yield better finitesample coverage.

### Acknowledgements

The authors gratefully acknowledge support from the University of Massachusetts Amherst Department of Mathematics and Statistics startup fund and NSF Award 2113171.

# References

- Austern, M. & Syrgkanis, V. (2020). Asymptotics of the empirical bootstrap method beyond asymptotic normality. arXiv preprint arXiv:2011.11248.
- Beran, R. (1997). Diagnosing bootstrap success. Annals of the Institute of Statistical Mathematics, 49, 1–24.
- Beutner, E. & Zähle, H. (2023). Donsker results for the empirical process indexed by functions of locally bounded variation and applications to the smoothed empirical process. *Bernoulli*, 29(1), 205–228.
- Bickel, P. J. (1982). On adaptive estimation. The Annals of Statistics, 10(3), 647–671.
- Bickel, P. J., Götze, F., & van Zwet, W. R. (1997). Resampling fewer than *n* observations: gains, losses, and remedies for losses. *Statistica Sinica*, 7(1), 1–31.
- Cai, W. & van der Laan, M. (2020). Nonparametric bootstrap inference for the targeted highly adaptive least absolute shrinkage and selection operator (LASSO) estimator. The International Journal of Biostatistics, 16(2), 20170070.
- Carone, M., Luedtke, A. R., & van der Laan, M. J. (2019). Toward computerized efficient estimation in infinite-dimensional models. *Journal of the American Statistical Association*, 114(527), 1174–1190. PMID: 32405108.
- Cattaneo, M. D. & Jansson, M. (2018). Kernel-based semiparametric estimators: Small bandwidth asymptotics and bootstrap consistency. *Econometrica*, 86(3), 955–995.
- Cattaneo, M. D. & Jansson, M. (2022). Average density estimators: Efficiency and bootstrap consistency. *Econometric Theory*, 38(6), 1140–1174.
- Cattaneo, M. D., Jansson, M., & Nagasawa, K. (2020). Bootstrap-based inference for cube root asymptotics. *Econometrica*, 88(5), 2203–2219.
- Cheng, G. & Huang, J. Z. (2010). Bootstrap consistency for general semiparametric M-estimation. The Annals of Statistics, 38(5), 2884–2915.
- Chernozhukov, V., Chetverikov, D., Demirer, M., Duflo, E., Hansen, C., Newey, W., & Robins, J. (2018). Double/debiased machine learning for treatment and structural parameters. *The Econometrics Journal*, 21(1), C1–C68.

- Chernozhukov, V., Chetverikov, D., & Kato, K. (2017). Central limit theorems and bootstrap in high dimensions. The Annals of Probability, 45(4), 2309–2352.
- Chernozhukov, V., Chetverikov, D., Kato, K., & Koike, Y. (2022). Improved central limit theorem and bootstrap approximations in high dimensions. *The Annals of Statistics*, 50(5), 2562–2586.
- Coyle, J. & van der Laan, M. J. (2018). Targeted bootstrap. In *Targeted learning in data science* chapter 28, (pp. 523–539). Springer Cham.
- Cuevas, A. & Romo, J. (1997). Differentiable functionals and smoothed bootstrap. Annals of the Institute of Statistical Mathematics, 49(2), 355–370.
- Dabrowska, D. M. (1989). Uniform consistency of the kernel conditional kaplan-meier estimate. The Annals of Statistics, 17(3), 1157–1167.
- Diciccio, T. J. & Romano, J. P. (1988). A review of bootstrap confidence intervals. Journal of the Royal Statistical Society: Series B (Methodological), 50(3), 338–354.
- Dixon, J. R., Kosorok, M. R., & Lee, B. L. (2005). Functional inference in semiparametric models using the piggyback bootstrap. Annals of the Institute of statistical Mathematics, 57(2), 255–277.
- Efron, B. (1982). *The Jackknife, the Bootstrap and Other Resampling Plans*. Society for Industrial and Applied Mathematics.
- Efron, B. & Tibshirani, R. J. (1994). An introduction to the Bootstrap. Chapman and Hall/CRC.
- Gaenssler, P. & Rost, D. (2000). On uniform laws of large numbers for smoothed empirical measures. In
  E. Giné, D. M. Mason, & J. A. Wellner (Eds.), *High Dimensional Probability II* (pp. 79–87). Boston,
  MA: Birkhäuser Boston.
- Gaenssler, P. & Rost, D. (2003). Smoothed empirical processes and the bootstrap. In J. Hoffmann-Jørgensen, J. A. Wellner, & M. B. Marcus (Eds.), *High Dimensional Probability III* (pp. 313–320). Basel: Birkhäuser Basel.
- Geman, S. & Hwang, C.-R. (1982). Nonparametric maximum likelihood estimation by the method of sieves. The Annals of Statistics, 10(2), 401–414.

- Geskus, R. & Groeneboom, P. (1999). Asymptotically optimal estimation of smooth functionals for interval censoring, case 2. The Annals of Statistics, 27(2), 627–674.
- Gill, R. D. & Robins, J. M. (2001). Causal Inference for Complex Longitudinal Data: The Continuous Case. The Annals of Statistics, 29(6), 1785–1811.
- Giné, E. & Nickl, R. (2008). Uniform central limit theorems for kernel density estimators. Probability Theory and Related Fields, 141(3), 333–387.
- Giné, E. & Zinn, J. (1990). Bootstrapping general empirical measures. The Annals of Probability, 18(2), 851–869.
- Giné, E. & Zinn, J. (1991). Gaussian characterization of uniform donsker classes of functions. The Annals of Probability, 19(2), 758–782.
- Giné, E. & Guillou, A. (2002). Rates of strong uniform consistency for multivariate kernel density estimators. Annales de l'Institut Henri Poincare (B) Probability and Statistics, 38(6), 907–921.
- Hall, P. (1988). Theoretical comparison of bootstrap confidence intervals. The Annals of Statistics, 16(3), 927–953.
- Hall, P. (1992). The Bootstrap and Edgeworth Expansion. Springer New York, NY.
- Hall, P., DiCiccio, T. J., & Romano, J. P. (1989). On smoothing and the bootstrap. The Annals of Statistics, 17(2), 692–704.
- Hall, P. & Kang, K.-H. (2001). Bootstrapping nonparametric density estimators with empirically chosen bandwidths. *The Annals of Statistics*, 29(5), 1443–1468.
- Han, F., Xu, S., & Zhou, W.-X. (2018). On Gaussian comparison inequality and its application to spectral analysis of large random matrices. *Bernoulli*, 24(3), 1787–1833.
- Hardin, J. W. & Hilbe, J. M. (2002). Generalized Estimating Equations. Chapman and Hall/CRC.
- Holland, P. W. (1986). Statistics and causal inference. Journal of the American Statistical Association, 81(396), 945–960.
- Klaassen, C. A. J. (1987). Consistent estimation of the influence function of locally asymptotically linear estimators. *The Annals of Statistics*, 15(4), 1548–1562.

- Kosorok, M. R. (2008). Introduction to Empirical Processes and Semiparametric Inference. Springer New York, NY.
- Kosorok, M. R., Lee, B. L., & Fine, J. P. (2004). Robust inference for univariate proportional hazards frailty regression models. *The Annals of Statistics*, 32(4), 1448 1491.
- Liang, K.-Y. & Zeger, S. L. (1986). Longitudinal data analysis using generalized linear models. Biometrika, 73(1), 13–22.
- Ma, S. & Kosorok, M. R. (2005). Robust semiparametric *m*-estimation and the weighted bootstrap. Journal of Multivariate Analysis, 96(1), 190–217.
- Murphy, S. A. & van der Vaart, A. W. (2000). On profile likelihood. Journal of the American Statistical Association, 95(450), 449–465.
- Newey, W. K., Hsieh, F., & Robins, J. M. (1998). Undersmoothing and bias corrected functional estimation. *Working papers*.
- Newey, W. K., Hsieh, F., & Robins, J. M. (2004). Twicing kernels and a small bias property of semiparametric estimators. *Econometrica*, 72(3), 947–962.
- Pfanzagl, J. (1982). Contributions to A General Asymptotic Statistical Theory. Springer New York, NY.
- Pfanzagl, J. (1990). Estimation in Semiparametric Models. Springer New York, NY.
- Powell, J. L., Stock, J. H., & Stoker, T. M. (1989). Semiparametric estimation of index coefficients. Econometrica: Journal of the Econometric Society, 57(6), 1403–1430.
- Praestgaard, J. & Wellner, J. A. (1993). Exchangeably weighted bootstraps of the general empirical process. The Annals of Probability, 21(4), 2053 – 2086.
- Quale, C. M., van der Laan, M. J., & Robins, J. R. (2006). Locally efficient estimation with bivariate right-censored data. *Journal of the American Statistical Association*, 101(475), 1076–1084.
- Radulović, D. & Wegkamp, M. (2000). Weak convergence of smoothed empirical processes: Beyond donsker classes. In E. Giné, D. M. Mason, & J. A. Wellner (Eds.), *High Dimensional Probability II* (pp. 89–105). Boston, MA: Birkhäuser Boston.

- Radulović, D. & Wegkamp, M. (2009). Uniform central limit theorems for pregaussian classes of functions. In C. Houdré, V. Koltchinskii, D. M. Mason, & M. Peligrad (Eds.), *High Dimensional Probability* V: The Luminy Volume (pp. 84–102).
- Robins, J. (1986). A new approach to causal inference in mortality studies with a sustained exposure period—application to control of the healthy worker survivor effect. *Mathematical Modelling*, 7(9), 1393–1512.
- Rost, D. (2000). Limit theorems for smoothed empirical processes. In E. Giné, D. M. Mason, & J. A.
  Wellner (Eds.), *High Dimensional Probability II* (pp. 107–113). Boston, MA: Birkhäuser Boston.
- Sheather, S. J. & Jones, M. C. (1991). A Reliable Data-Based Bandwidth Selection Method for Kernel Density Estimation. Journal of the Royal Statistical Society: Series B (Methodological), 53(3), 683– 690.
- Sheehy, A. & Wellner, J. A. (1992). Uniform donsker classes of functions. The Annals of Probability, 20(4), 1983–2030.
- Shen, X. (1997). On methods of sieves and penalization. The Annals of Statistics, 25(6), 2555–2591.
- Silverman, B. W. (1986). Density Estimation for Statistics and Data Analysis. London: Chapman and Hall.
- van der Laan, M. J. & Robins, J. M. (2003). Unified Methods for Censored Longitudinal Data and Causality. Springer New York, NY.
- van der Laan, M. J. & Rose, S. (2011). Targeted Learning: Causal Inference for Observational and Experimental Data. Springer New York, NY.
- van der Laan, M. J. & Rubin, D. (2006). Targeted maximum likelihood learning. *The International Journal of Biostatistics*, 2(1).
- van der Vaart, A. W. (1991). On differentiable functionals. The Annals of Statistics, 19(1), 178–204.
- van der Vaart, A. W. (1994). Weak convergence of smoothed empirical processes. *Scandinavian Journal* of *Statistics*, 21(4), 501–504.

van der Vaart, A. W. (2000). Asymptotic Statistics, volume 3. Cambridge University Press.

- van der Vaart, A. W. & Wellner, J. A. (1996). Weak Convergence and Empirical Processes: With Applications to Statistics. Springer New York, NY.
- Wellner, J. A. & Zhan, Y. (1996). Bootstrapping z-estimators. University of Washington Department of Statistics Technical Report, 308(5).
- Yukich, J. E. (1992). Weak convergence of smoothed empirical processes. Scandinavian Journal of Statistics, 19(3), 271–279.

# Supplementary Material

## Supplement A Proof of results in Section 2

**Theorem 3.1.** If conditions (A1)-(A3) hold, then  $T(\eta_n, \mathbb{P}_n)$  is asymptotically linear in the sense that  $T(\eta_n, \mathbb{P}_n) = T(\eta_0, P_0) + \mathbb{P}_n \phi_0 + o_{P_0^*}(n^{-1/2})$ , which implies that  $n^{1/2} [T(\eta_n, \mathbb{P}_n) - T(\eta_0, P_0)] \rightsquigarrow \mathbb{G}_0 \phi_0$ .

Proof of Theorem 3.1. Consider the decomposition

$$T(\eta_n, \mathbb{P}_n) - T(\eta_0, P_0) = \mathbb{P}_n \phi_0 + (\mathbb{P}_n - P_0)(\phi_n - \phi_0) + [T(\eta_n, \mathbb{P}_n) - T(\eta_0, P_0) - (\mathbb{P}_n - P_0)\phi_n].$$

By conditions (A1)–(A2) and Lemma 19.24 of van der Vaart (2000),  $(\mathbb{P}_n - P_0)(\phi_n - \phi_0) = o_{P_0^*}(n^{-1/2})$ . By condition (A3), the remainder term is  $o_{P_0^*}(n^{-1/2})$ . The second statement follows by the central limit theorem since  $P_0\phi_0 = 0$  and  $P_0\phi_0^2 < \infty$  by assumption.

**Proposition 3.2.** If (i)  $P_0^*(\phi_n^2 \in \mathfrak{G}) \to 1$  for a  $P_0$ -Glivenko Cantelli class of measurable functions  $\mathfrak{G}$ , and (ii)  $P_0(\phi_n^2 - \phi_0^2) \xrightarrow{P_0^*} 0$ , then  $\mathbb{P}_n \phi_n^2 \xrightarrow{P_0^*} \sigma_0^2$ . Furthermore, if  $\sup_{f \in \mathcal{F}} |P_0 f| < \infty$ , condition (A1) implies (i) and condition (A2) implies (ii).

Proof of Proposition 3.2. By adding and subtracting terms, we have

$$\mathbb{P}_n \phi_n^2 - \sigma_0^2 = (\mathbb{P}_n - P_0)\phi_n^2 + P_0(\phi_n^2 - \phi_0^2).$$

The second term is  $o_{P_0^*}(1)$  by assumption. For the first term, for any  $\varepsilon > 0$ , we have

$$\begin{aligned} P_0^*\left(\left|(\mathbb{P}_n - P_0)\phi_n^2\right| \ge \varepsilon\right) &= P_0^*\left(\left|(\mathbb{P}_n - P_0)\phi_n^2\right| \ge \epsilon, \, \phi_n^2 \in \mathcal{G}\right) + P_0^*\left(\left|(\mathbb{P}_n - P_0)\phi_n^2\right| \ge \varepsilon, \, \phi_n^2 \notin \mathcal{G}\right) \\ &\le P_0^*\left(\sup_{g \in \mathcal{G}} \left|(\mathbb{P}_n - P_0)g\right| \ge \varepsilon\right) + P_0^*\left(\phi_n^2 \notin \mathcal{G}\right). \end{aligned}$$

Since  $\mathcal{G}$  is a  $P_0$ -Glivenko Cantelli class, the first term is o(1), and since  $\phi_n^2 \in \mathcal{G}$  with probability tending to one the second term is o(1). Hence,  $(\mathbb{P}_n - P_0)\phi_n^2 = o_{P_0^*}(1)$ , which completes the proof of the first statement.

By Theorem 2.10.14 of van der Vaart & Wellner (1996), since  $\sup_{f \in \mathcal{F}} |P_0 f| < \infty$ , condition (A1) implies that  $\mathcal{F}^2 = \{f^2 : f \in \mathcal{F}\}$  is  $P_0$ -Glivenko-Cantelli in probability, and since  $P_0^*(\phi_n^2 \in \mathcal{F}^2) \to 1$ , this implies (i). By Cauchy-Schwarz and Minkowski's inequalities, we also have

$$\begin{aligned} \left| P_0(\phi_n^2 - \phi_0^2) \right| &\leq P_0 |\phi_n^2 - \phi_0^2| \\ &= P_0 \left| (\phi_n - \phi_0) (\phi_n + \phi_0) \right| \\ &\leq \left\| \phi_n - \phi_0 \right\|_{L_2(P_0)} \left\| \phi_n + \phi_0 \right\|_{L_2(P_0)} \\ &\leq \left\| \phi_n - \phi_0 \right\|_{L_2(P_0)} \left[ \left\| \phi_n - \phi_0 \right\|_{L_2(P_0)} + 2 \left\| \phi_0 \right\|_{L_2(P_0)} \right] \end{aligned}$$

which is  $o_{P_0^*}(1)$  by (A2).

# Supplement B Proof of results in Section 3

**Lemma 3.5.** Suppose that  $\mathcal{F}$  is a class of measurable functions such that  $\mathbb{G}_n^* \overset{P_0^*}{\underset{W}{\leftrightarrow}} \mathbb{G}_0$  in  $\ell^{\infty}(\mathcal{F})$ . Let  $\phi_n^*$  be a sequence of random functions possibly depending on the bootstrap sample such that  $P_W^*(\phi_n^* \in \mathcal{F}) \xrightarrow{P_0^*} 1$ . If  $\rho(\phi_n^*, \phi_\infty) = o_{P_W^*}(1)$  for some  $\phi_\infty \in \mathcal{F}$  and a semimetric  $\rho$  on  $\mathcal{F}$  for which the sample paths of  $\mathbb{G}_0$  are almost surely uniformly  $\rho$ -continuous, then  $\mathbb{G}_n^*(\phi_n^* - \phi_\infty) = o_{P_W^*}(1)$ .

**Proof of Lemma 3.5.** A standard way to prove results of this type outside the setting of the bootstrap is the continuous mapping theorem (see, e.g., van der Vaart, 2000, Theorem 19.24). However, the continuous mapping theorem might not be applicable in the bootstrap setting as the map  $(w_1, \ldots, w_n) \mapsto$  $h(\mathbb{G}_n^*(x_1, \ldots, x_n, w_1, \ldots, w_n))$  might not be measurable given almost  $x_1, \ldots, x_n$  for all  $h \in C_b(\ell^{\infty}(\mathcal{F}))$ .

For any  $\tau > 0$ , we define  $\mathfrak{F}_{\tau} = \{f - f_{\infty} : f \in \mathfrak{F}, \rho(f, f_{\infty}) < \tau\}$ . We then have

$$\begin{aligned} \{|\mathbb{G}_n^*(f_n^* - f_\infty)| \ge \varepsilon\} &\subseteq \{||\mathbb{G}_n^*(f_n^* - f_\infty)| \ge \varepsilon, \ \rho(f_n^*, f_\infty) < \tau, \ f_n^* \in \mathcal{F}\} \\ &\cup \{\rho(f_n^*, f_\infty) \ge \tau \text{ or } f_n^* \notin \mathcal{F}\} \\ &\subseteq \{||\mathbb{G}_n^*||_{\mathcal{F}_\tau} \ge \varepsilon\} \cup \{\rho(f_n^*, f_\infty) \ge \tau\} \cup \{f_n^* \notin \mathcal{F}\} \end{aligned}$$

for any  $\varepsilon, \tau > 0$ . Thus,

$$P_W^*\left(|\mathbb{G}_n^*(f_n^* - f_\infty)| \ge \varepsilon\right) \le P_W^*\left(||\mathbb{G}_n^*||_{\mathcal{F}_\tau} \ge \varepsilon\right) + P_W^*\left(\rho(f_n^*, f_\infty) \ge \tau\right) + P_W^*\left(f_n^* \notin \mathcal{F}\right).$$

The second and third terms on the right-hand side conditionally converge to zero in outer probability for any  $\tau > 0$  by assumption. Hence, the proof is complete if we can show that for all  $\varepsilon, \delta, \gamma > 0$  there exists  $\tau > 0$  such that  $P_0^* \left( P_W^* \left( \| \mathbb{G}_n^* \|_{\mathcal{F}_{\tau}} \ge \varepsilon \right) > \delta \right) < \gamma$  for all n large enough.

For each  $\varepsilon, \tau > 0$  there exists a sequence of functions  $h_m : \ell^{\infty}(\mathcal{F}) \to \mathbb{R}$  such that  $h_m$  is *m*-Lipschitz for all  $m, 1\{||z||_{\mathcal{F}_{\tau}} \ge \varepsilon\} \le h_m(z) \le 1$  for all  $z \in \ell^{\infty}(\mathcal{F})$  and m, and  $h_m(z)$  monotonically decreases to  $1\{||z||_{\mathcal{F}_{\tau}} \ge \varepsilon\}$  as  $m \to \infty$  for each  $z \in \ell^{\infty}(\mathcal{F})$ . For instance,  $h_m : z \mapsto \min\{\max(m[||z||_{\mathcal{F}_{\tau}} - \varepsilon] + 1, 0), 1\}$ satisfies these criteria. For any  $\varepsilon, \tau > 0$  and  $m \in \{1, 2, ...\}$ , we can now write

$$\begin{aligned} P_W^*\left(\|\mathbb{G}_n^*\|_{\mathcal{F}_{\tau}} \ge \varepsilon\right) &= \left[E_W^* \mathbf{1}\left\{\|\mathbb{G}_n^*\|_{\mathcal{F}_{\tau}} \ge \varepsilon\right\} - E_W^* h_m(\mathbb{G}_n^*)\right] + \left[E_W^* h_m(\mathbb{G}_n^*) - E_0 h_m(\mathbb{G}_0)\right] \\ &+ \left[E_0 h_m(\mathbb{G}_0) - E_0 \mathbf{1}\left\{\|\mathbb{G}_0\|_{\mathcal{F}_{\tau}} \ge \varepsilon\right\}\right] + P_0\left(\|\mathbb{G}_0\|_{\mathcal{F}_{\tau}} \ge \varepsilon\right) \\ &\leq \left[E_W^* h_m(\mathbb{G}_n^*) - E_0 h_m(\mathbb{G}_0)\right] + \left[E_0 h_m(\mathbb{G}_0) - E_0 \mathbf{1}\left\{\|\mathbb{G}_0\|_{\mathcal{F}_{\tau}} \ge \varepsilon\right\}\right] \\ &+ P_0\left(\|\mathbb{G}_0\|_{\mathcal{F}_{\tau}} \ge \varepsilon\right),\end{aligned}$$

where the second inequality follows because  $E_W^* 1 \{ \| \mathbb{G}_n^* \|_{\mathcal{F}_\tau} \ge \varepsilon \} \le E_W^* h_m(\mathbb{G}_n^*)$  for all  $\varepsilon, \tau > 0$  and  $m \in \{1, 2, ...\}$  by assumption. For the final term on the right-hand side, since almost all sample paths of  $\mathbb{G}_0$  are uniformly continuous in  $\ell^{\infty}(\mathcal{F})$  with respect to  $\rho$ , for any  $\varepsilon, \delta > 0$ , we can choose  $\tau > 0$  such that  $P_0(\|\mathbb{G}_0\|_{\mathcal{F}_\tau} \ge \varepsilon) < \delta$ . For the second term on the right-hand side, by the monotone convergence theorem,  $\lim_{m\to\infty} E_0 h_m(\mathbb{G}_0) = E_0 1\{\|\mathbb{G}_0\|_{\mathcal{F}_\tau} \ge \varepsilon\}$ . Hence, for any  $\varepsilon, \delta, \tau > 0$ , we can choose m such that  $E_0 h_m(\mathbb{G}_0) - E_0 1\{\|\mathbb{G}_0\|_{\mathcal{F}_\tau} \ge \varepsilon\} < \delta$ . Finally, for the first term on the right-hand side, since  $h_m$  is bounded and m-Lipschitz,  $h_m/m \in \mathrm{BL}_1(\ell^{\infty}(\mathcal{F}))$  for each m. Therefore, by the assumed conditional weak convergence of  $\mathbb{G}_n^*$  to  $\mathbb{G}_0$  in  $\ell^{\infty}(\mathcal{F}), E_W^* \frac{1}{m} h_m(\mathbb{G}_n^*) \xrightarrow{P_0^*} E_0 \frac{1}{m} h_m(\mathbb{G}_0)$  for each m, so that  $E_W^* h_m(\mathbb{G}_n^*) \xrightarrow{P_0^*} E_0 h_m(\mathbb{G}_0)$  as well. Hence, for any  $\varepsilon, \delta, \gamma, \tau > 0$  and  $m \in \{1, 2, ...\}, P_0^*(E_W^* h_m(\mathbb{G}_n^*) - E_0 h_m(\mathbb{G}_0) > \delta\} < \gamma$  for all n large enough. This completes the proof.

**Theorem 3.4.** If conditions (B1)-(B3) hold, then  $T(\eta_n^*, \mathbb{P}_n^*)$  is conditionally asymptotically linear in the sense that  $T(\eta_n^*, \mathbb{P}_n^*) = T(\eta_n, \hat{P}_n) + (\mathbb{P}_n^* - \hat{P}_n)\phi_0 + o_{P_W^*}(n^{-1/2})$ , which implies that

$$n^{1/2} \left[ T\left(\eta_n^*, \mathbb{P}_n^*\right) - T(\eta_n, \hat{P}_n) \right] \underset{W}{\overset{P_0^*}{\rightsquigarrow}} \mathbb{G}_0(\phi_0).$$

**Proof of Theorem 3.4.** We let h be an arbitrary element of  $BL_1(\mathbb{R})$ . By adding and subtracting

terms and the triangle inequality, we have

$$\begin{aligned} \left| E_W^* h\left( n^{1/2} \left[ T(\eta_n^*, \mathbb{P}_n^*) - T(\eta_n, \hat{P}_n) \right] \right) - E_0 h\left( \mathbb{G}_0 \phi_0 \right) \right| \\ &\leq E_W^* \left| h\left( n^{1/2} \left[ T(\eta_n^*, \mathbb{P}_n^*) - T(\eta_n, \hat{P}_n) \right] \right) - h\left( \mathbb{G}_n^* \phi_0 \right) \right| + \left| E_W^* h\left( \mathbb{G}_n^* \phi_0 \right) - E_0 h\left( \mathbb{G}_0 \phi_0 \right) \right|. \end{aligned} \tag{15}$$

For the first term on the right-hand side of (15), we note that for any  $x_1, x_2 \in \mathbb{R}$ ,  $|h(x_1) - h(x_2)| \le 2 \land |x_1 - x_2|$ , and for any  $\varepsilon > 0$ ,  $2 \land |x_1 - x_2| \le \varepsilon + 2I(|x_1 - x_2| > \varepsilon)$ , which implies that for any  $\varepsilon > 0$ ,

$$E_W^* \left| h \left( n^{1/2} \left[ T(\eta_n^*, \mathbb{P}_n^*) - T(\eta_n, \hat{P}_n) \right] \right) - h \left( \mathbb{G}_n^* \phi_0 \right) \right|$$

$$\leq \varepsilon + 2P_W^* \left( \left| n^{1/2} \left[ T(\eta_n^*, \mathbb{P}_n^*) - T(\eta_n, \hat{P}_n) \right] - \mathbb{G}_n^* \phi_0 \right| \geq \varepsilon \right).$$

$$\tag{16}$$

We now write

$$n^{1/2} \left[ T(\eta_n^*, \mathbb{P}_n^*) - T(\eta_n, \hat{P}_n) \right] - \mathbb{G}_n^* \phi_0 = \mathbb{G}_n^* (\phi_n^* - \phi_0) + n^{1/2} \left[ T(\eta_n^*, \mathbb{P}_n^*) - T(\eta_n, \hat{P}_n) - (\mathbb{P}_n^* - \hat{P}_n) \phi_n^* \right]$$
$$= \mathbb{G}_n^* (\phi_n^* - \phi_0) + n^{1/2} R_n^*.$$

Note that  $\mathbb{G}_n^*(\phi_n^* - \phi_0) = o_{P_W^*}(1)$  by conditions (B1)–(B2) and Lemma 3.5, and  $n^{1/2}R_n^* = o_{P_W^*}(1)$  by condition (B3). This implies the first term on the right-hand side of (15) is  $o_{P_0^*}(1)$ . For the second term on the right hand side of (15), we define the function  $g : \ell^{\infty}(\mathcal{F}) \to \mathbb{R}$  as  $g(z) := h(z(\phi_0))$ . For any  $z_1, z_2 \in \ell^{\infty}(\mathcal{F})$  with  $||z_1 - z_2||_{\mathcal{F}} > 0$ , we have

$$\frac{|g(z_1) - g(z_2)|}{\|z_1 - z_2\|_{\mathcal{F}}} = \frac{|h(z_1(\phi_0)) - h(z_2(\phi_0))|}{\|z_1 - z_2\|_{\mathcal{F}}} \le \frac{|z_1(\phi_0) - z_2(\phi_0)|}{\|z_1 - z_2\|_{\mathcal{F}}} \le 1$$

because  $h \in BL_1(\mathbb{R})$  and  $\phi_0 \in \mathcal{F}$ . Hence,  $g \in BL_1(\ell^{\infty}(\mathcal{F}))$ . Therefore,

$$\left|E_{W}^{*}h\left(\mathbb{G}_{n}^{*}\phi_{0}\right)-E_{0}h\left(\mathbb{G}_{0}\phi_{0}\right)\right|\leq\sup_{g\in\mathrm{BL}_{1}\left(\ell^{\infty}\left(\mathfrak{F}\right)\right)}\left|E_{W}^{*}g\left(\mathbb{G}_{n}^{*}\right)-E_{0}g\left(\mathbb{G}_{0}\right)\right|\xrightarrow{P_{0}^{*}}0$$

by condition (B1). This implies conditional asymptotic linearity of  $T(\eta_n^*, \mathbb{P}_n^*)$ . Conditional weak convergence of  $n^{1/2} \left[ T(\eta_n^*, \mathbb{P}_n^*) - T(\eta_n, \hat{P}_n) \right]$  is then implied by (16).

**Lemma 3.7** (Theorem 4.5 of Sheehy & Wellner, 1992). If  $\mathcal{F}$  is a class of measurable functions with envelope function F such that: (i)  $\mathcal{F}$  is square integrable uniformly in  $P \in \mathcal{P}$  in the sense that  $\lim_{M\to\infty} \sup_{P\in\mathcal{P}} PF^2 1\{F > M\} = 0$ ; (ii)  $\mathcal{F}$  is Donkser uniformly in  $P \in \mathcal{P}$  where  $\mathcal{P}$  is such that  $P_0^*(\hat{P}_n \in \mathfrak{P}) \to 1$ ; and (iii) the semi-metric  $L_2(\hat{P}_n)$  converges uniformly to  $L_2(P_0)$  in the sense that

$$\sup_{f,g\in\mathcal{F}} \left| \|f - g\|_{L_2(\hat{P}_n)} - \|f - g\|_{L_2(P_0)} \right| \xrightarrow{P_0^*} 0, \tag{2}$$

then  $\mathbb{G}_n^* \overset{P_0^*}{\underset{W}{\longrightarrow}} \mathbb{G}_0$  in  $\ell^{\infty}(\mathfrak{F})$ .

**Proof of Lemma 3.7.** We argue along subsequences using a standard argument structure. By Lemma 1.9.2 of van der Vaart & Wellner (1996), every subsequence  $\{n_k : k = 1, 2, ...\}$  has a further subsequence  $\{n_{k_l} : l = 1, 2, ...\}$  such

$$\sup_{f,g\in\mathcal{F}} \left\| \|f - g\|_{L_2(\hat{P}_{n_{k_l}})} - \|f - g\|_{L_2(P_0)} \right\| \xrightarrow{\text{a.s.}^*} 0.$$

By Theorem 4.5 of Sheehy & Wellner (1992), we then have  $n_{k_l}^{1/2}(\mathbb{P}_{n_{k_l}}^* - \hat{P}_{n_{k_l}}) \xrightarrow{\text{a.s.}^*}_{W} \mathbb{G}_0$ . Hence, every  $\{n_k\}$  has a further subsequence  $\{n_{k_l}\}$  such that

$$\sup_{h\in \mathrm{BL}_1(\ell^{\infty}(\mathfrak{F}))} \left| E_W^* h(\mathbb{G}_{n_{k_l}}^*) - E_0 h(\mathbb{G}_0) \right| \xrightarrow{\mathrm{a.s.}^*} 0,$$

which by Lemma 1.9.2 of van der Vaart & Wellner (1996) again implies that

$$\sup_{h \in \mathrm{BL}_1(\ell^{\infty}(\mathcal{F}))} |E_W^* h(\mathbb{G}_n^*) - E_0 h(\mathbb{G}_0)| \xrightarrow{P_0^*} 0,$$

i.e.,  $\mathbb{G}_n^* \overset{P_0^*}{\underset{W}{\longrightarrow}} \mathbb{G}_0.$ 

**Proposition 3.8.** Suppose  $\mathcal{F}$  is a class of Borel measurable functions with uniformly bounded envelope function F and finite uniform entropy integral as in (3) such that  $\mathcal{F}_{\delta,P} = \{f-g: f, g \in \mathcal{F}, \|f-g\|_{L_2(P)} < \delta\}$  and  $\mathcal{F}_{\infty}^2 = \{(f-g)^2: f, g \in \mathcal{F}\}$  are P-measurable for every  $\delta > 0$  and  $P \in \mathcal{M}$ . If  $P_0$  is absolutely continuous with respect to Lebesgue measure  $\lambda$  with uniformly bounded and Lipschitz continuous density function  $p_0$ , and  $\hat{P}_n = \mathbb{P}_n * L_n$  for a sequence of random measures  $L_n$  converging weakly to Dirac measure at 0, then the conditions of Proposition 3.7 hold, so that  $\mathbb{G}_n^* \overset{P_0^*}{\underset{W}{\longrightarrow}} \mathbb{G}_0$  in  $\ell^{\infty}(\mathcal{F})$ .

**Proof of Proposition 3.8.** Since  $\mathcal{F}$  has uniformly bounded envelope function and finite uniform entropy integral, by Theorem 2.8.3 of van der Vaart & Wellner (1996), conditions (i)–(ii) of Lemma 3.7 are satisfied. We next show the uniform convergence of semi-metric. For simplicity, we denote  $\mathcal{F}_{\infty} :=$ 

 $\{f-g: f,g\in \mathcal{F}\} \text{ and } \mathcal{F}^2_\infty := \{(f-g)^2: f,g\in \mathcal{F}\}. \text{ Since } |\sqrt{x}-\sqrt{y}| \leq \sqrt{|x-y|} \text{ for all } x,y\in \mathbb{R}, \text{ we have } x \in \mathbb{R}, x \in \mathbb{R}\}.$ 

$$\sup_{f,g\in\mathcal{F}} \left| \|f - g\|_{L_2(\hat{P}_n)} - \|f - g\|_{L_2(P_0)} \right| \le \sup_{f,g\in\mathcal{F}} \left| \int (f - g)^2 \,\mathrm{d}(\hat{P}_n - P_0) \right|^{1/2}$$

Hence, condition (iii) of Lemma 3.7 is satisfied if  $\|\hat{P}_n - P_0\|_{\mathcal{F}^2_{\infty}} = o_{P_0^*}(1)$ .

We show that  $P_0^*(\|\hat{P}_n - P_0\|_{\mathcal{F}^2_{\infty}} \geq \delta) \to 0$  for all  $\delta > 0$  using Theorem 2.2 of Gaenssler & Rost (2000). To do so, we verify their conditions (2.2)–(2.5) with  $\mu_n = L_n$  and  $\nu_n = \mathbb{P}_n$ . Since  $\mathcal{F}^2_{\infty}$ is uniformly bounded by the uniform boundedness of  $\mathcal{F}$ , conditions (2.2) and (2.3) are satisfied, as discussed following Theorem 2.2 of Gaenssler & Rost (2000). To show condition (2.4), we firstly note that following inequality holds for all  $\varepsilon > 0$  and measures Q on  $(\mathfrak{X}, \mathcal{B})$ ,

$$N(4\varepsilon, \mathcal{F}_{\infty}^2, L_1(Q)) \le N(4\varepsilon, \mathcal{F}_{\infty}^2, L_2(Q)) \le N(2\varepsilon, \mathcal{F}_{\infty}, L_2(Q)) \le N(\varepsilon, \mathcal{F}, L_2(Q))^2.$$

The first inequality is because  $\|\cdot\|_{L_1(Q)} \leq \|\cdot\|_{L_2(Q)}$  by Hölder's inequality. The second inequality is because for any  $f_{\infty}, g_{\infty} \in \mathcal{F}_{\infty}$ , we have  $Q(f_{\infty}^2 - g_{\infty}^2)^2 = Q(f_{\infty} - g_{\infty})^2(f_{\infty} + g_{\infty})^2 \leq 4Q(f_{\infty} - g_{\infty})^2$ by uniform boundedness of  $\mathcal{F}$ . The third inequality is because for any  $f_1, g_1, f_2, g_2 \in \mathcal{F}$ , we have  $Q((f_1-g_1)-(f_2-g_2))^2 \leq Q(f_1-f_2)^2 + Q(g_1-g_2)^2 + 2\|f_1-f_2\|_{L_2(Q)}\|g_1-g_2\|_{L_2(Q)}$ . Since  $\mathcal{F}$  has uniformly integrable  $L_2$  entropy, we have  $\int_0^\infty \sup_Q \{\log N(\varepsilon, \mathcal{F}, L_2(Q))\}^{1/2} d\varepsilon < \infty$ , where the supremum is taken over all finitely discrete probability measures Q on  $(\mathcal{X}, \mathcal{B})$ . By exercise 1 in Chapter 2.5 of van der Vaart & Wellner (1996), the supremum over finitely discrete Q can be replaced by the supremum over all probability measures Q such that  $0 < QF^2 < \infty$  without changing the assumption of uniformly integrable  $L_2$  entropy of  $\mathcal{F}$ . So, we have

$$\int_{0}^{\infty} \sup_{Q} \left\{ \log N(\varepsilon, \mathcal{F}_{\infty}^{2}, L_{1}(Q)) \right\}^{1/2} \, \mathrm{d}\varepsilon < \infty,$$
(17)

where the supremum is taken over all probability measures Q on  $(\mathfrak{X}, \mathfrak{B})$ . Since  $N(\varepsilon, \mathcal{F}_{\infty}^2, L_1(\hat{P}_n)) \leq \sup_Q N(\varepsilon, \mathcal{F}_{\infty}^2, L_1(Q))$  almost surely, and the latter is finite for  $\varepsilon > 0$  by (17),  $N(\varepsilon, \mathcal{F}_{\infty}^2, L_1(\hat{P}_n))$  is stochastically bounded for all  $\varepsilon > 0$ . Furthermore, since  $\|\cdot\|_{d_{\tilde{\nu}_n}^{(1)}} \leq \|\cdot\|_{L_1(\hat{P}_n)}$  for  $d_{\tilde{\nu}_n}^{(1)}$  defined in Gaenssler & Rost (2000), this further implies that condition (2.4) of Theorem 2.2 of Gaenssler & Rost (2000) holds. Lastly, to show condition (2.5) in Theorem 2.2 of Gaenssler & Rost (2000) holds, we show that

$$\lim_{n \to \infty} \sup_{f_{\infty} \in \mathcal{F}_{\infty}} \left| P_0 * L_n(f_{\infty}^2) - P_0(f_{\infty}^2) \right|$$
$$= \lim_{n \to \infty} \sup_{f_{\infty} \in \mathcal{F}_{\infty}} \left| \iint f_{\infty}(x+y)^2 \, \mathrm{d}P_0(x) \, \mathrm{d}L_n(y) - \iint f_{\infty}(x+y)^2 \, \mathrm{d}P_0(x) \, \mathrm{d}\delta_0(y) \right| = 0.$$

The function  $y \mapsto \int f_{\infty}^2(x+y) dP_0(x)$  is uniformly bounded by 1 since  $\mathcal{F}_{\infty}^2$  is uniformly bounded by 1. Furthermore, for any  $y_1, y_2 \in \mathfrak{X}$  and  $f_{\infty} \in \mathcal{F}_{\infty}$ , using the assumption that  $p_0$  is Lipschitz, we have

$$\begin{aligned} \left| \int f_{\infty}(x+y_{1})^{2} dP_{0}(x) - \int f_{\infty}(x+y_{2})^{2} dP_{0}(x) \right| \\ &= \left| \int f_{\infty}(x+y_{1})^{2} p_{0}(x) d\lambda(x) - \int f_{\infty}(x+y_{2})^{2} p_{0}(x) d\lambda(x) \right| \\ &= \left| \int f_{\infty}(z)^{2} [p_{0}(z-y_{1}) - p_{0}(z-y_{2})] d\lambda(z) \right| \\ &\leq \gamma |y_{1} - y_{2}| \int f_{\infty}^{2} d\lambda \\ &\leq 4\gamma |y_{1} - y_{2}| \int F^{2} d\lambda. \end{aligned}$$

for some  $\gamma > 0$ . Since  $\int F^2 d\lambda < \infty$  by assumption, this implies  $y \mapsto \int f_{\infty}^2(x+y) dP_0(x)$  is uniformly bounded by 1 and  $\gamma'$ -Lipschitz for  $\gamma' := 4\gamma \int F^2 d\lambda$  for all  $f_{\infty} \in \mathcal{F}_{\infty}$ . Thus, the function  $\min(\gamma'^{-1}, 1) \times \int f_{\infty}^2(x+\cdot) dP_0(x) \in BL_1(\mathbb{R})$ , which implies that

$$\sup_{f_{\infty}\in\mathfrak{F}_{\infty}}\left|\iint f_{\infty}(x+y)^{2} \,\mathrm{d}P_{0}(x) \,\mathrm{d}L_{n}(y) - \iint f_{\infty}(x+y)^{2} \,\mathrm{d}P_{0}(x) \,\mathrm{d}\delta_{0}(y)\right|$$
$$\leq \frac{1}{\min(\gamma'^{-1},1)} \sup_{h\in\mathrm{BL}_{1}(\mathfrak{X})}\left|\int h(y) \,\mathrm{d}L_{n}(y) - \int h(y) \,\mathrm{d}\delta_{0}(y)\right|,$$

which goes to 0 as  $n \to \infty$  because  $L_n$  converges weakly to  $\delta_0$  by assumption. We have now checked all the conditions of Theorem 2.2 of Gaenssler & Rost, 2000, so we conclude that  $E_0^* \|\hat{P}_n - P_0\|_{\mathcal{F}^2_\infty}^p \to 0$  for all  $p \ge 1$ . This demonstrates that condition (iii) of Lemma 3.7 holds, so the result follows.

**Lemma 3.11.** If there exists a class of measurable functions  $\mathcal{G}$  such that (i)  $P_W^*(\phi_n^{*2} \in \mathcal{G}) \xrightarrow{P_0^*} 1$ , (ii)  $\sup_{g \in \mathcal{G}} |(\hat{P}_n - P_0)g| = o_{P_0^*}(1)$ , (iii)  $\sup_{g \in \mathcal{G}} |(\mathbb{P}_n^* - \hat{P}_n)g| = o_{P_W^*}(1)$ , and (iv)  $P_0(\phi_n^{*2} - \phi_0^2) = o_{P_W^*}(1)$ , then  $\mathbb{P}_n^*\phi_n^{*2} - \sigma_0^2 = o_{P_W^*}(1)$ . Furthermore, condition (B2) implies (iv). For the empirical bootstrap where  $\hat{P}_n = \mathbb{P}_n$ , condition (B1) implies condition (i)–(iii). For any bootstrap  $\hat{P}_n$ , condition (B1)(a) and the conditions of Proposition 3.7 imply conditions (i)-(iii).

Proof of Lemma 3.11. By adding and subtracting terms,

$$\mathbb{P}_{n}^{*}\phi_{n}^{*2} - \sigma_{0}^{2} = (\mathbb{P}_{n}^{*} - \hat{P}_{n})\phi_{n}^{*2} + (\hat{P}_{n} - P_{0})\phi_{n}^{*2} + P_{0}(\phi_{n}^{*2} - \phi_{0}^{2}).$$
(18)

For the first term on the right-hand side of (18), for any  $\varepsilon > 0$ , we have

$$\begin{aligned} P_W^*\left(\left|(\mathbb{P}_n^* - \hat{P}_n)\phi_n^{*2}\right| \ge \varepsilon\right) &= P_W^*\left(\left|(\mathbb{P}_n^* - \hat{P}_n)\phi_n^{*2}\right| \ge \varepsilon, \ \phi_n^{*2} \in \mathfrak{G}\right) \\ &+ P_W^*\left(\left|(\mathbb{P}_n^* - \hat{P}_n)\phi_n^{*2}\right| \ge \varepsilon, \ \phi_n^{*2} \notin \mathfrak{G}\right) \\ &\le P_W^*\left(\sup_{g \in \mathfrak{G}}\left|(\mathbb{P}_n^* - \hat{P}_n)g\right| \ge \varepsilon\right) + P_W^*\left(\phi_n^{*2} \notin \mathfrak{G}\right) \\ &= o_{P_0^*}(1). \end{aligned}$$

For the second term on the right-hand side of (18), for any  $\varepsilon > 0$ , we have

$$\begin{aligned} P_W^*\left(\left|(\hat{P}_n - P_0)\phi_n^{*2}\right| \ge \varepsilon\right) &= P_W^*\left(\left|(\hat{P}_n - P_0)\phi_n^{*2}\right| \ge \varepsilon, \ \phi_n^{*2} \in \mathfrak{G}\right) \\ &+ P_W^*\left(\left|(\hat{P}_n - P_0)\phi_n^{*2}\right| \ge \varepsilon, \ \phi_n^{*2} \notin \mathfrak{G}\right) \\ &\le P_W^*\left(\sup_{g\in\mathfrak{G}}\left|(\hat{P}_n - P_0)g\right| \ge \varepsilon\right) + P_W^*\left(\phi_n^{*2} \notin \mathfrak{G}\right) \\ &= o_{P_0^*}(1). \end{aligned}$$

Therefore,  $\mathbb{P}_n^* \phi_n^* - \sigma_0^2 = o_{P_W^*}(1)$  by assumption. Furthermore,  $P_0(\phi_n^{*2} - \phi_0^2) = o_{P_W^*}(1)$  is implied by condition (B2). This is because by Cauchy-Schwarz and Minkowski's inequalities,

$$\begin{aligned} \left| P_0(\phi_n^{*2} - \phi_0^2) \right| &\leq P_0 |\phi_n^{*2} - \phi_0^2| \\ &= P_0 \left| (\phi_n^* - \phi_0) (\phi_n^* + \phi_0) \right| \\ &\leq \left\| \phi_n^* - \phi_0 \right\|_{L_2(P_0)} \left\| \phi_n^* + \phi_0 \right\|_{L_2(P_0)} \\ &\leq \left\| \phi_n^* - \phi_0 \right\|_{L_2(P_0)} \left[ \left\| \phi_n^* - \phi_0 \right\|_{L_2(P_0)} + 2 \left\| \phi_0 \right\|_{L_2(P_0)} \right] \\ &= o_{P_W^*}(1). \end{aligned}$$

Specifically, for the empirical bootstrap  $\hat{P}_n = \mathbb{P}_n$ , condition (B1)(b) implies  $\mathcal{F}$  is  $P_0$ -Donsker by The-

orem 3.6.1 of van der Vaart & Wellner (1996). Hence,  $\mathcal{G} = \{f^2 : f \in \mathcal{F}\}$  is  $P_0$ -Glivenko-Cantelli in probability by Lemma 2.10.14 of van der Vaart (1991), which implies (ii) and (iii) by Theorem 2.6 of Giné & Zinn (1990). Meanwhile, (i) is implied by condition (B1)(a), where  $\mathcal{G} = \{f^2 : f \in \mathcal{F}\}$ .

**Lemma B.1** (Bootstrap Slutsky's Theorem). Let  $X_n^*$  and  $Y_n^*$  be two sequences of real-valued random variables defined on the product probability space  $(\mathfrak{X}, \mathfrak{B}, P_0)^n \times (\mathfrak{W}_n, \mathfrak{C}_n, Q_n)^n$ . Assume that  $X_n^* \overset{P_0^*}{\underset{W}{\to}} X$  and  $Y_n^* \overset{P_0^*}{\underset{W}{\to}} c$  in  $\mathbb{R}$ , where X is a tight, Borel measurable limit and c is a constant. We then have (i)  $Y_n^* X_n^* \overset{P_0^*}{\underset{W}{\to}} cX$  in  $\mathbb{R}$ , (ii)  $(Y_n^*)^{-1} X_n^* \overset{P_0^*}{\underset{W}{\to}} c^{-1} X$  in  $\mathbb{R}$  provided  $c \neq 0$ , and (iii)  $X_n^* + Y_n^* \overset{P_0^*}{\underset{W}{\to}} X + Y$  in  $\mathbb{R}$ .

**Proof of Lemma B.1.** We only show (i); a similar argument can be applied to yield (ii) and (iii). As in the proof of Lemma 3.5, a standard way to prove results of this type outside the setting of the bootstrap is the continuous mapping theorem (e.g., Lemma 2.8 of van der Vaart, 2000). However, the continuous mapping theorem might not be applicable in the bootstrap setting as the map  $(w_1, \ldots, w_n) \mapsto h(\mathbb{G}_n^*(x_1, \ldots, x_n, w_1, \ldots, w_n))$  might not be measurable given almost every  $x_1, \ldots, x_n$  for all  $h \in C_b(\ell^{\infty}(\mathcal{F}))$ .

We let h be an arbitrary element of  $BL_1(\mathbb{R})$ . By adding and subtracting terms and the triangle inequality, for any  $\varepsilon > 0$ ,

$$|E_W^*h(X_n^*Y_n^*) - E_0h(Xc)| \le |E_W^*h(X_n^*Y_n^*) - E_W^*h(X_n^*c)| + |E_W^*h(X_n^*c) - E_0h(Xc)|$$
  
$$\le \varepsilon + 2P_W^*(|X_n^*Y_n^* - X_n^*c| \ge \varepsilon) + |E_W^*h(X_n^*c) - E_0h(Xc)|, \qquad (19)$$

where the second inequality is because  $|h(x_1) - h(x_2)| \le 2 \land |x_1 - x_2| \le \varepsilon + 2I(|x_1 - x_2| > \varepsilon)$  for any  $x_1, x_2 \in \mathbb{R}$ . By defining  $g : \mathbb{R} \to \mathbb{R}$  as g(x) = cx, we note that  $\min(|c|-1, 1) \times (h \circ g) \in BL_1(\mathbb{R})$  because for any  $x_1, x_2 \in \mathbb{R}$  and  $x_1 \neq x_2$ ,

$$\frac{|\min(|c|^{-1}, 1) \times (h \circ g)(x_1) - \min(|c|^{-1}, 1) \times (h \circ g)(x_2)|}{|x_1 - x_2|}$$
  
=  $\min(|c|^{-1}, 1) \frac{|(h \circ g)(x_1) - (h \circ g)(x_2)|}{|g(x_1) - g(x_2)|} \frac{|g(x_1) - g(x_2)|}{|x_1 - x_2|}$   
 $\leq \min(|c|, 1) \frac{|(h \circ g)(x_1) - (h \circ g)(x_2)|}{|g(x_1) - g(x_2)|}$   
 $\leq 1$ 

because  $h \in BL_1(\mathbb{R})$ . Since in addition  $|\min(|c|^{-1}, 1) \times (h \circ g)| \le 1$ ,  $\min(|c|^{-1}, 1) \times (h \circ g) \in BL_1(\mathbb{R})$ .

Hence, for the third term on the right-hand side of (19), we have

$$\begin{split} |E_W^*h(X_n^*c) - E_0h(Xc)| &= \frac{1}{\min(|c|^{-1}, 1)} \left| E_W^*(\min(|c|^{-1}, 1) \times (h \circ g))(X_n^*) - E_0(\min(|c|^{-1}, 1) \times (h \circ g))(X) \right| \\ &\leq \frac{1}{\min(|c|^{-1}, 1)} \sup_{h \in \mathrm{BL}_1(\mathbb{R})} |E_W^*h(X_n^*) - E_0h(X)|, \end{split}$$

which converges to 0 in outer probability since  $X_n^* \overset{P_0^*}{\underset{W}{\longrightarrow}} X$  by assumption. For the second term on the right-hand side of (19), for any M > 0,

$$\begin{split} P_W^*(|X_n^*Y_n^* - X_n^*c| \ge \varepsilon) &= P_W^*(|X_n^*Y_n^* - X_n^*c| \ge \varepsilon, |X_n^*| < M) \\ &+ P_W^*(|X_n^*Y_n^* - X_n^*c| \ge \varepsilon, |X_n^*| \ge M) \\ &\le P_W^*(|Y_n^* - c| \ge \varepsilon/M) + P_W^*(|X_n^*| \ge M). \end{split}$$

By assumption,  $Y_n^* \xrightarrow{P_0^*} c$  so that  $P_W^*(|Y_n^* - c| \ge \varepsilon/M) = o_{P_0^*}(1)$ . Hence, the proof is complete if we can show that given any  $\varepsilon, \gamma > 0$ , there exists M > 0 such that  $P_0^*(P_W^*(|X_n^*| \ge M) \ge \varepsilon) < \gamma$  for all n large enough.

For each  $\varepsilon, \gamma > 0$  there exists a sequence of functions  $h_m : \mathbb{R} \to \mathbb{R}$  such that  $h_m$  is *m*-Lipschitz for all  $m, 1\{|x| \ge M\} \le h_m(z) \le 1$  for all  $x \in \mathbb{R}$  and m, and  $h_m(z)$  monotonically decreases to  $1\{|x| \ge M\}$  as  $m \to \infty$  for each  $x \in \mathbb{R}$ . For instance,  $h_m : x \mapsto \min\{\max(m[|x| - M] + 1, 0), 1\}$  satisfies these criteria. For any  $\varepsilon > 0$  and  $m \in \{1, 2, ...\}$ , we can now write

$$P_W^* \left( |X_n^*| \ge M \right) = \left[ E_W^* 1\left\{ |X_n^*| \ge M \right\} - E_W^* h_m \left( X_n^* \right) \right] + \left[ E_W^* h_m \left( X_n^* \right) - E_0 h_m(X) \right] \\ + \left[ E_0 h_m(X) - E_0 1\left\{ |X| \ge M \right\} \right] + P_0 \left( |X| \ge M \right) \\ \le \left[ E_W^* h_m \left( X_n^* \right) - E_0 h_m(X) \right] + \left[ E_0 h_m(X) - E_0 1\left\{ |X| \ge M \right\} \right] \\ + P_0 \left( |X| \ge M \right),$$

where the second inequality follows because  $E_W^* 1\{|X_n^*| \ge M\} \le E_W^* h_m(X_n^*)$  for all  $\varepsilon > 0$  and  $m \in \{1, 2, ...\}$  by assumption. We therefore have

$$P_0^*(P_W^*(|X_n^*| \ge M) \ge \varepsilon)$$
  
$$\le P_0^*([E_W^*h_m(X_n^*) - E_0h_m(X)] \ge \varepsilon/3) + P_0^*([E_0h_m(X) - E_01\{|X| \ge M\}] \ge \varepsilon/3)$$

$$+ P_0^*(P_0(|X| \ge M) \ge \varepsilon/3)$$

$$= P_0^*([E_W^*h_m(X_n^*) - E_0h_m(X)] \ge \varepsilon/3) + 1\{[E_0h_m(X) - E_01\{|X| \ge M\}] \ge \varepsilon/3\}$$

$$+ 1\{P_0(|X| \ge M) \ge \varepsilon/3\}.$$
(20)

First, since X is tight by assumption, we can choose M sub that the third term on the right-hand side of (20) is 0. Next, for this choice of M, by the monotone convergence theorem we can choose m large enough so that the second term on the right-hand side of (20) is 0. Finally, for these choices of m and M, for the first term on the right-hand side of (20), we have  $\min(m^{-1}, 1) \times h_m \in BL_1(\mathbb{R})$  because  $h_m$  is bounded and m-Lipschitz. Therefore, by the assumed conditional weak convergence of  $X_n^*$  to X in  $\mathbb{R}$ ,  $E_W^*[\min(m^{-1}, 1) \times h_m(X_n^*)] \xrightarrow{P_0^*} E_0[\min(m^{-1}, 1) \times h_m(X)]$  for each m, so that  $E_W^*h_m(X_n^*) \xrightarrow{P_0^*} E_0h_m(X)$ as well. Hence, the first term on the right-hand side of (20) converges to 0 for fixed m and  $\varepsilon$ , so we can make it as small as we like for n large enough. Hence, for any  $\varepsilon$ , there exist M > 0 and  $m \in \{1, 2, ...\}$ such that  $P_W^*(|X_n^*| \ge M) = o_{P_0^*}(1)$  for n large enough.

We next have a result that generalizes Theorem 23.4 of van der Vaart (2000) to include smooth bootstraps.

**Lemma B.2** (Conditional CLT). Let  $\phi_0 : \mathbb{R}^d \mapsto \mathbb{R}$  satisfy  $P_0\phi_0 = 0$  and  $P_0\phi_0^2 < \infty$ . Let  $X_1, X_2, \ldots$  be IID random vectors with mean  $\mu$  and covariance matrix  $\Sigma$ . If  $\hat{P}_n\phi_0^2 \xrightarrow{P_0^*} P_0\phi_0^2$  and  $(\hat{P}_n - P_0)[\phi_0^2 1\{|\phi_0| > M\}] \xrightarrow{P_0^*} 0$  for any M > 0, then conditionally on every sequence  $X_1, X_2, \ldots$ , in outer probabiliy,

$$n^{1/2} \left( \mathbb{P}_n^* - \hat{P}_n \right) \phi_0 \underset{W}{\overset{P_0^*}{\leadsto}} N(0, P_0 \phi_0^2).$$

**Proof of Lemma B.2.** We use the Lindeberg-Feller CLT for triangular arrays (e.g., Proposition 2.27 of van der Vaart, 2000). Since the bootstrap sample  $X_1^*, \ldots, X_n^*$  is drawn IID from  $\hat{P}_n$  given  $X_1, \ldots, X_n$ , the conditional mean and variance of  $\phi_0(X_i^*)$  are given by

$$E_W \phi_0(X_i^*) = \int \phi_0(x) \, \mathrm{d}\hat{P}_n(x) = \hat{P}_n \phi_0, \text{ and}$$
$$E_W [\phi_0(X_i^*) - E_W \phi_0(X_i^*)]^2 = E_W \phi_0^2(X_i^*) - [E_W \phi_0(X_i^*)]^2 = \hat{P}_n \phi_0^2 - (\hat{P}_n \phi_0)^2 \xrightarrow{P_0^*} P_0 \phi_0^2.$$

We next verify the Lindeberg condition. For any  $\varepsilon, \gamma > 0$ , there exists M > 0 such that  $\int \phi_0(x)^2 1\{|\phi_0(x)| > 0\}$ 

M d $P_0(x) < \varepsilon/2$  and for all n large enough,  $M \le \gamma \sqrt{n}$ . Therefore,

$$\begin{aligned} P_0\left(E_W[\phi_0(X_i^*)^2 1\{|\phi_0(X_i^*)| > \gamma\sqrt{n}\}] \ge \varepsilon\right) \\ &= P_0\left(\hat{P}_n[\phi_0^2 1\{|\phi_0| > \gamma\sqrt{n}\}] \ge \varepsilon\right) \\ &\le P_0\left(\hat{P}_n[\phi_0^2 1\{|\phi_0| > M\}] \ge \varepsilon\right) \\ &\le P_0\left(|(\hat{P}_n - P_0)[\phi_0^2 1\{|\phi_0| > M\}]| \ge \varepsilon/2\right) + P_0\left(P_0[\phi_0^2 1\{|\phi_0| > M\}] \ge \varepsilon/2\right) \\ &= P_0\left(|(\hat{P}_n - P_0)[\phi_0^2 1\{|\phi_0| > M\}]| \ge \varepsilon/2\right) + 1\left\{P_0[\phi_0^2 1\{|\phi_0| > M\}] \ge \varepsilon/2\right\} \\ &= P_0\left(|(\hat{P}_n - P_0)[\phi_0^2 1\{|\phi_0| > M\}]| \ge \varepsilon/2\right) \\ &\longrightarrow 0 \end{aligned}$$

Thus,  $E_W[\phi_0(X_i^*)^2 1\{|\phi_0(X_i^*)| > \varepsilon \sqrt{n}\}] \xrightarrow{P_0^*} 0$ . Therefore, the conditions of the Lindeberg-Feller CLT holds conditionally on every sequence  $X_1, X_2, \ldots$ , in outer probability, which yields the result. **Theorem 3.9.** Suppose that  $\hat{P}_n \phi_0^2 \xrightarrow{P_0^*} P_0 \phi_0^2$  and  $(\hat{P}_n - P_0)[\phi_0^2 1\{|\phi_0| > M\}] \xrightarrow{P_0^*} 0$  for every M > 0. If  $S_n^* - S_n = o_{P_W^*}(n^{-1/2})$  and  $R_n^* - R_n = o_{P_W^*}(n^{-1/2})$ , then

$$\sup_{t\in\mathbb{R}} \left| P_W^*\left( T(\eta_n^*, \mathbb{P}_n^*) - T(\eta_n, \hat{P}_n) \le t \right) - P_0^*\left( T(\eta_n, \mathbb{P}_n) - T(\eta_0, P_0) \le t \right) \right| \xrightarrow{P_0^*} 0,$$

and the bootstrap percentile confidence interval defined in (6) has asymptotic confidence level  $1 - \alpha - \beta$ . If in addition  $\sigma_n^2 \xrightarrow{P_0^*} \sigma_0^2$ ,  $\sigma_n^{*2} \xrightarrow{P_0^*} \sigma_0^2$  and  $(S_n + R_n)(\sigma_n^* - \sigma_n) = o_{P_W^*}(n^{-1/2})$ , then

$$\sup_{t\in\mathbb{R}} \left| P_W^*\left( \left[ T(\eta_n^*, \mathbb{P}_n^*) - T(\eta_n, \hat{P}_n) \right] / \sigma_n^* \le t \right) - P_0^*\left( \left[ T(\eta_n, \mathbb{P}_n) - T(\eta_0, P_0) \right] / \sigma_n \le t \right) \right| \xrightarrow{P_0^*} 0,$$

and the percentile t-confidence interval defined in (5) has asymptotic confidence level  $1 - \alpha - \beta$ .

**Proof of Theorem 3.9.** We prove the result for the percentile *t*-method; the result for the percentile method follows the same argument setting  $\sigma_n = \sigma_n^* = \sigma_0 = 1$  (so that the additional conditions for the percentile *t*-method hold automatically).

We first show that  $\sigma_n^{*-1}(S_n^* + R_n^*) - \sigma_n^{-1}(S_n + R_n) = o_{P_W^*}(n^{-1/2})$ . By adding and subtracting terms, we note that

$$\frac{S_n^* + R_n^*}{\sigma_n^*} - \frac{S_n + R_n}{\sigma_n} = \frac{\sigma_n (S_n^* + R_n^*) - \sigma_n (S_n + R_n) + \sigma_n (S_n + R_n) - \sigma_n^* (S_n + R_n)}{\sigma_n \sigma_n^*}$$
$$=\frac{(S_n^* - S_n) + (R_n^* - R_n)}{\sigma_n^*} + \frac{(\sigma_n - \sigma_n^*)(S_n + R_n)}{\sigma_n \sigma_n^*}.$$
 (21)

Since  $\sigma_n^* \xrightarrow{P_0^*} \sigma_0 > 0$ , we can show that  $\sigma_n^{*-1} = O_{P_W^*}(1)$  using the same logic as the unconditional result. Hence, the above equals  $o_{P_W^*}(n^{-1/2})O_{P_W^*}(1) + o_{P_W^*}(n^{-1/2})O_{P_W^*}(1)$  by assumption.

For any  $t \in \mathbb{R}$ , by definition of  $S_n$  and  $R_n$ , we have

$$P_0^* \left( \frac{T(\eta_n, \mathbb{P}_n) - T(\eta_0, P_0)}{\sigma_n} \le t \right) = P_0^* \left( \sigma_n^{-1} \mathbb{G}_n \phi_0 + n^{1/2} \sigma_n^{-1} \left[ S_n + R_n \right] \le n^{1/2} t \right)$$
$$= P_0^* \left( \sigma_n^{-1} \mathbb{G}_n \phi_0 \le T_n \right),$$

where  $T_n := n^{1/2}t - n^{1/2}\sigma_n^{-1}[S_n + R_n]$ . Thus, by definition of  $S_n^*$  and  $R_n^*$ ,

$$P_W^* \left( \frac{T(\eta_n^*, \mathbb{P}_n^*) - T(\eta_n, \hat{P}_n)}{\sigma_n^*} \le t \right) = P_W^* \left( \sigma_n^{*-1} \mathbb{G}_n^* \phi_0 + n^{1/2} \sigma_n^{*-1} \left[ S_n^* + R_n^* \right] \le n^{1/2} t \right)$$
$$= P_W^* \left( \sigma_n^{*-1} \mathbb{G}_n^* \phi_0 + n^{1/2} \sigma_n^{*-1} \left[ S_n^* + R_n^* \right] \le T_n + n^{1/2} \sigma_n^{-1} \left[ S_n + R_n \right] \right)$$
$$= P_W^* \left( \sigma_n^{*-1} \mathbb{G}_n^* \phi_0 + A_n^* \le T_n \right),$$

where  $A_n^* := n^{1/2} \{ \sigma_n^{*-1} [S_n^* + R_n^*] - \sigma_n^{-1} [S_n + R_n] \}$ . Thus,

$$\begin{split} \sup_{t \in \mathbb{R}} \left| P_W^* \left( \frac{T(\eta_n^*, \mathbb{P}_n^*) - T(\eta_n, \hat{P}_n)}{\sigma_n^*} \le t \right) - P_0^* \left( \frac{T(\eta_n, \mathbb{P}_n) - T(\eta_0, P_0)}{\sigma_n} \le t \right) \right| \\ &= \sup_{t \in \mathbb{R}} \left| P_W^* \left( \sigma_n^{*-1} \mathbb{G}_n^* \phi_0 + A_n^* \le t \right) - P_0^* \left( \sigma_n^{-1} \mathbb{G}_n \phi_0 \le t \right) \right| \\ &\le \sup_{t \in \mathbb{R}} \left| P_W^* \left( \sigma_n^{*-1} \mathbb{G}_n^* \phi_0 + A_n^* \le t \right) - P_0^* \left( \sigma_0^{-1} \mathbb{G}_0 \phi_0 \le t \right) \right| + \sup_{\xi \in \mathbb{R}} \left| P_0^* \left( \sigma_n^{-1} \mathbb{G}_n \phi_0 \le t \right) - P_0^* \left( \sigma_0^{-1} \mathbb{G}_0 \phi_0 \le t \right) \right| \,. \end{split}$$

By Lemma B.2, we have  $\mathbb{G}_n^* \phi_0 \overset{P_0^*}{\underset{W}{\longrightarrow}} \mathbb{G}_0 \phi_0$ . Since  $\sigma_n^{*2} \overset{P_0^*}{\underset{W}{\longrightarrow}} \sigma_0^2$ ,  $\sigma_n^{*-1} \mathbb{G}_n^* \phi_0 \overset{P_0^*}{\underset{W}{\longrightarrow}} \sigma_0^{-1} \mathbb{G}_0 \phi_0$  by Lemma B.1. Since  $A_n^* = o_{P_W^*}(1)$ , it follows that  $\sigma_n^{*-1} \mathbb{G}_n^* \phi_0 + A_n^* \overset{P_0^*}{\underset{W}{\longrightarrow}} \sigma_0^{-1} \mathbb{G}_0 \phi_0$ . Hence, by the Portmanteau theorem (e.g., Lemma 1.3.4 of van der Vaart & Wellner, 1996),

$$\sup_{t\in\mathbb{R}} \left| P_W^* \left( \sigma_n^{*-1} \mathbb{G}_n^* \phi_0 + A_n^* \le t \right) - P_0^* \left( \sigma_0^{-1} \mathbb{G}_0 \phi_0 \le t \right) \right| \xrightarrow{P_0^*} 0.$$

Similarly,  $\sigma_n^{-1} \mathbb{G}_n \phi_0 \rightsquigarrow \sigma_0^{-1} \mathbb{G}_0 \phi_0$ , so

$$\sup_{t \in \mathbb{R}} \left| P_0^* \left( \sigma_n^{-1} \mathbb{G}_n \phi_0 \le \xi \right) - P_0^* \left( \sigma_0^{-1} \mathbb{G}_0 \phi_0 \le t \right) \right| \to 0.$$

The first statement follows.

By definition of  $\xi_{n,p}^*$ ,  $S_n^*$ , and  $R_n^*$ , we have

$$\begin{aligned} \xi_{n,p}^* &:= \inf \left\{ \xi \in \mathbb{R} : P_W^* \left( \frac{T(\eta_n^*, \mathbb{P}_n^*) - T(\eta_n, \hat{P}_n)}{\sigma_n^*} \le \xi \right) \ge p \right\} \\ &= \inf \left\{ \xi \in \mathbb{R} : P_W^* \left( n^{-1/2} \sigma_n^{*-1} \mathbb{G}_n^* \phi_0 + \sigma_n^{*-1} \left[ S_n^* + R_n^* \right] \le \xi \right) \ge p \right\}. \end{aligned}$$

Since  $\sigma_n^{-1} \left[ S_n + R_n \right]$  is a function of the original data, we have

$$n^{1/2} \left( \xi_{n,p}^* - \sigma_n^{-1} \left[ S_n + R_n \right] \right)$$
  
=  $\inf \left\{ \xi \in \mathbb{R} : P_W^* \left( n^{-1/2} \sigma_n^{*-1} \mathbb{G}_n^* \phi_0 + \sigma_n^{*-1} \left[ S_n^* + R_n^* \right] \le n^{-1/2} \xi + \sigma_n^{-1} \left[ S_n + R_n \right] \right) \ge p \right\}$   
=  $\inf \left\{ \xi \in \mathbb{R} : P_W^* \left( \sigma_n^{*-1} \mathbb{G}_n^* \phi_0 + A_n^* \le \xi \right) \ge p \right\}.$ 

By the above derivations and van der Vaart, 2000, Lemma 21.2, for any  $p \in (0, 1)$ , we then have

$$n^{1/2} \left(\xi_{n,p}^* - \sigma_n^{-1} \left[S_n + R_n\right]\right) \xrightarrow{P_0^*} \Phi^{-1}(p).$$

Therefore,

$$P_{0}^{*}\left(T(\eta_{n}, \mathbb{P}_{n}) - \xi_{n,1-\alpha}^{*}\sigma_{n} \leq T(\eta_{0}, P_{0}) \leq T(\eta_{n}, \mathbb{P}_{n}) - \xi_{n,\beta}^{*}\sigma_{n}\right)$$

$$= P_{0}^{*}\left(n^{1/2}\xi_{n,\beta}^{*} \leq n^{1/2}[T(\eta_{n}, \mathbb{P}_{n}) - T(\eta_{0}, P_{0})]/\sigma_{n} \leq n^{1/2}\xi_{n,1-\alpha}^{*}\right)$$

$$= P_{0}^{*}\left(n^{1/2}\left[\xi_{n,\beta}^{*} - \sigma_{n}^{-1}(S_{n} + R_{n})\right] \leq \sigma_{n}^{-1}\mathbb{G}_{n}\phi_{0} \leq n^{1/2}\left[\xi_{n,1-\alpha}^{*} - \sigma_{n}^{-1}(S_{n} + R_{n})\right]\right)$$

$$\longrightarrow P_{0}^{*}\left(\Phi^{-1}(\beta) \leq \sigma_{0}^{-1}\mathbb{G}_{0}\phi_{0} \leq \Phi^{-1}(1-\alpha)\right)$$

$$= 1 - \alpha - \beta.$$

Corollary 3.10. If conditions (A1)-(A3) and (B1)-(B3) hold, then the bootstrap percentile confidence

interval has asymptotic confidence level  $1 - \alpha - \beta$ . If in addition  $\sigma_n^2 \xrightarrow{P_0^*} \sigma_0^2$  and  $\sigma_n^{*2} \xrightarrow{P_0^*} \sigma_0^2$ , then the percentile t-confidence interval defined in (5) has asymptotic confidence level  $1 - \alpha - \beta$ .

**Proof of Corollary 3.10.** We note that in the proof of Theorem 3.9, the conditions  $\hat{P}_n \phi_0^2 \xrightarrow{P_0^*} P_0 \phi_0^2$ and  $(\hat{P}_n - P_0)[\phi_0^2 1\{|\phi_0| > M\}] \xrightarrow{P_0^*} 0$  were only used to establish that  $\mathbb{G}_n^* \phi_0 \overset{P_0^*}{\underset{W}{\longrightarrow}} \mathbb{G}_0 \phi_0$  using Lemma B.2. However, condition (B1) directly implies  $\mathbb{G}_n^* \phi_0 \overset{P_0^*}{\underset{W}{\longrightarrow}} \mathbb{G}_0 \phi_0$ , so it suffices to check the remainder of the conditions of Theorem 3.9. By conditions (A1)–(A3), we have  $S_n = o_{P_0^*}(n^{-1/2})$  and  $R_n = o_{P_0^*}(n^{-1/2})$ . By conditions (B1)–(B3), we have  $S_n^* = o_{P_W^*}(n^{-1/2})$  and  $R_n^* = o_{P_W^*}(n^{-1/2})$ . Hence,  $S_n^* - S_n = o_{P_W^*}(n^{-1/2})$ and  $R_n^* - R_n = o_{P_W^*}(n^{-1/2})$ , which implies the consistency of the bootstrap percentile confidence interval by Theorem 3.9. For the percentile t interval,  $\sigma_n^* \xrightarrow[W]{P_0^*} \sigma_0$  and  $\sigma_n \xrightarrow[W]{P_0^*} \sigma_0$  by assumption, which also implies  $(S_n + R_n)(\sigma_n^* - \sigma_n) = o_{P_W^*}(n^{-1/2})o_{P_W^*}(1) = o_{P_W^*}(n^{-1/2}).$ 

**Theorem 3.12.** Suppose that  $\hat{P}_n \phi_0^2 \xrightarrow{P_0^*} P_0 \phi_0^2$  and  $(\hat{P}_n - P_0)[\phi_0^2 1\{|\phi_0| > M\}] \xrightarrow{P_0^*} 0$  for every M > 0. If  $[R_n + S_n] + [R_n^* + S_n^*] + [T(\eta_n, \hat{P}_n) - T(\eta_n, \mathbb{P}_n)] = o_{P_W^*}(n^{-1/2})$ , then

$$\sup_{t\in\mathbb{R}}|P_W^*\left(T(\eta_n^*,\mathbb{P}_n^*)-T(\eta_n,\mathbb{P}_n)\leq t\right)-P_0^*\left(-\left[T(\eta_n,\mathbb{P}_n)-T(\eta_0,P_0)\right]\leq t\right)|\xrightarrow{P_0^*}0,$$

and Efron's percentile confidence interval  $[\zeta_{n,\beta}^*, \zeta_{n,1-\alpha}^*]$  has asymptotic confidence level  $1 - \alpha - \beta$ . **Proof of Theorem 3.12.** For any  $t \in \mathbb{R}$ , by definition of  $S_n$  and  $R_n$ , we have

$$P_0^* \left( - \left[ T(\eta_n, \mathbb{P}_n) - T(\eta_0, P_0) \right] \le t \right) = P_0^* \left( -\mathbb{G}_n \phi_0 - n^{1/2} \left[ S_n + R_n \right] \le n^{1/2} t \right) = P_0^* \left( -\mathbb{G}_n \phi_0 \le T_n \right),$$

where  $T_n := n^{1/2}t + n^{1/2}[S_n + R_n]$ . Thus, by definition of  $S_n^*$  and  $R_n^*$ ,

$$P_W^* \left( T(\eta_n^*, \mathbb{P}_n^*) - T(\eta_n, \mathbb{P}_n) \le t \right) = P_W^* \left( \mathbb{G}_n^* \phi_0 + n^{1/2} \left[ S_n^* + R_n^* + T(\eta_n, \hat{P}_n) - T(\eta_n, \mathbb{P}_n) \right] \le n^{1/2} t \right)$$
$$= P_W^* \left( \mathbb{G}_n^* \phi_0 + B_n^* \le T_n \right),$$

where  $B_n^* := n^{1/2} \left[ S_n^* + S_n + R_n^* + R_n + T(\eta_n, \hat{P}_n) - T(\eta_n, \mathbb{P}_n) \right]$ . Thus,

t

$$\sup_{t \in \mathbb{R}} |P_W^* \left( T(\eta_n^*, \mathbb{P}_n^*) - T(\eta_n, \mathbb{P}_n) \le t \right) - P_0^* \left( - \left[ T(\eta_n, \mathbb{P}_n) - T(\eta_0, P_0) \right] \le t \right) |$$
  
$$= \sup_{t \in \mathbb{R}} |P_W^* \left( \mathbb{G}_n^* \phi_0 + B_n^* \le t \right) - P_0^* \left( - \mathbb{G}_n \phi_0 \le t \right) |$$
  
$$\le \sup_{t \in \mathbb{R}} |P_W^* \left( \mathbb{G}_n^* \phi_0 + B_n^* \le t \right) - P_0^* \left( \mathbb{G}_0 \phi_0 \le t \right) | + \sup_{t \in \mathbb{R}} |P_0^* \left( - \mathbb{G}_n \phi_0 \le t \right) - P_0^* \left( \mathbb{G}_0 \phi_0 \le t \right) |$$

By Lemma B.2 and since  $B_n^* = o_{P_W^*}(1)$  by assumption, we have  $\mathbb{G}_n^* \phi_0 + B_n^* \overset{P_0^*}{\underset{W}{\longrightarrow}} \mathbb{G}_0 \phi_0$ . Hence, by the Portmanteau theorem (e.g., Lemma 1.3.4 of van der Vaart & Wellner, 1996),

$$\sup_{t \in \mathbb{R}} |P_W^* \left( \mathbb{G}_n^* \phi_0 + B_n^* \le t \right) - P_0^* \left( \mathbb{G}_0 \phi_0 \le t \right)| \xrightarrow{P_0^*} 0.$$

Similarly,  $\mathbb{G}_n \phi_0 \rightsquigarrow \mathbb{G}_0 \phi_0$ , and since  $\mathbb{G}_0 \phi_0$  follows a mean-zero normal distribution,  $\mathbb{G}_n \phi_0 \rightsquigarrow -\mathbb{G}_0 \phi_0$  as well. Therefore,

$$\sup_{t \in \mathbb{R}} |P_0^* \left( \mathbb{G}_n \phi_0 \le t \right) - P_0^* \left( -\mathbb{G}_0 \phi_0 \le t \right)| \to 0.$$

The first statement follows.

By definition of  $\zeta_{n,p}^*, S_n^*$  and  $R_n^*$ , we have

$$\begin{aligned} \zeta_{n,p}^* &:= \inf \left\{ \zeta \in \mathbb{R} : P_W^* \left( T(\eta_n^*, \mathbb{P}_n^*) \le \zeta \right) \ge p \right\} \\ &= \inf \left\{ \zeta \in \mathbb{R} : P_W^* \left( n^{-1/2} \mathbb{G}_n^* \phi_0 + (S_n^* + R_n^*) + T(\eta_n, \hat{P}_n) \le \zeta \right) \ge p \right\}. \end{aligned}$$

Since  $[S_n + R_n - T(\eta_n, \mathbb{P}_n)]$  is a function of the original data, we have

$$n^{1/2} \left\{ \zeta_{n,p}^* + [S_n + R_n - T(\eta_n, \mathbb{P}_n)] \right\}$$
  
=  $\inf \left\{ \zeta \in \mathbb{R} : P_W^* \left( n^{-1/2} \mathbb{G}_n^* \phi_0 + (S_n^* + R_n^*) + T(\eta_n, \hat{P}_n) \le n^{-1/2} \zeta - [S_n + R_n - T(\eta_n, \mathbb{P}_n)] \right) \ge p \right\}$   
=  $\inf \left\{ \zeta \in \mathbb{R} : P_W^* \left( \mathbb{G}_n^* \phi_0 + B_n^* \le \zeta \right) \ge p \right\},$ 

As argued above,  $\mathbb{G}_n^* \phi_0 + B_n^* \overset{P_0^*}{\underset{W}{\longrightarrow}} \mathbb{G}_0 \phi_0$ , so by Lemma 21.2 of van der Vaart (2000), for any  $p \in (0, 1)$ , we then have

$$n^{1/2}\left\{\zeta_{n,p}^* + \left[S_n + R_n - T(\eta_n, \mathbb{P}_n)\right]\right\} \xrightarrow{P_0^*} \sigma_0 \Phi^{-1}(p).$$

Therefore,

$$P_{0}^{*}\left(\zeta_{n,1-\alpha}^{*} \leq T(\eta_{0},P_{0}) \leq \zeta_{n,\beta}^{*}\right)$$

$$= P_{0}^{*}\left(-n^{1/2}[\zeta_{n,\beta}^{*} - T(\eta_{n},\mathbb{P}_{n})] \leq n^{1/2}[T(\eta_{n},\mathbb{P}_{n}) - T(\eta_{0},P_{0})] \leq -n^{1/2}[\zeta_{n,1-\alpha}^{*} - T(\eta_{n},\mathbb{P}_{n})]\right)$$

$$= P_{0}^{*}\left(-n^{1/2}[\zeta_{n,\beta}^{*} - T(\eta_{n},\mathbb{P}_{n})] \leq \mathbb{G}_{n}\phi_{0} + n^{1/2}(S_{n} + R_{n}) \leq -n^{1/2}[\zeta_{n,1-\alpha}^{*} - T(\eta_{n},\mathbb{P}_{n})]\right)$$

$$= P_{0}^{*}\left(-n^{1/2}[\zeta_{n,\beta}^{*} + S_{n} + R_{n} - T(\eta_{n},\mathbb{P}_{n})] \leq \mathbb{G}_{n}\phi_{0} \leq -n^{1/2}[\zeta_{n,1-\alpha}^{*} + S_{n} + R_{n} - T(\eta_{n},\mathbb{P}_{n})]\right)$$

$$\longrightarrow P_0^* \left( -\Phi^{-1}(\beta) \le \sigma_0^{-1} \mathbb{G}_0 \phi_0 \le -\Phi^{-1}(1-\alpha) \right)$$
$$= 1 - \alpha - \beta,$$

where the last equality is due to the symmetry of the standard normal distribution. This shows the consistency of Efron's percentile method.

**Corollary 3.13.** If (A1)-(A3) and (B1)-(B3) hold, and  $T(\eta_n, \hat{P}_n) - T(\eta_n, \mathbb{P}_n) = o_{P_0^*}(n^{-1/2})$ , then Efron's percentile confidence interval  $[\zeta_{n,\beta}^*, \zeta_{n,1-\alpha}^*]$  has asymptotic confidence level  $1 - \alpha - \beta$ .

**Proof of Corollary 3.13.** As in the proof of Corollary 3.10, we note that in the proof of Theorem 3.12, the conditions  $\hat{P}_n \phi_0^2 \xrightarrow{P_0^*} P_0 \phi_0^2$  and  $(\hat{P}_n - P_0)[\phi_0^2 1\{|\phi_0| > M\}] \xrightarrow{P_0^*} 0$  were only used to establish that  $\mathbb{G}_n^* \phi_0 \xrightarrow{P_0^*} \mathbb{G}_0 \phi_0$  using Lemma B.2. However, condition (B1) directly implies  $\mathbb{G}_n^* \phi_0 \xrightarrow{P_0^*} \mathbb{G}_0 \phi_0$ , so it suffices to check the remainder of the conditions of Theorem 3.12.

By conditions (A1)–(A3), we have  $S_n = o_{P_0^*}(n^{-1/2})$  and  $R_n = o_{P_0^*}(n^{-1/2})$ . By conditions (B1)– (B3), we have  $S_n^* = o_{P_W^*}(n^{-1/2})$  and  $R_n^* = o_{P_W^*}(n^{-1/2})$ . Since  $T(\eta_n, \hat{P}_n) - T(\eta_n, \mathbb{P}_n) = o_{P_0^*}(n^{-1/2})$  by assumption, we then have  $S_n^* + S_n + R_n^* + R_n + T(\eta_n, \hat{P}_n) - T(\eta_n, \mathbb{P}_n) = o_{P_W^*}(n^{-1/2})$ . Hence, the result follows by Theorem 3.12.

**Theorem 3.14.** Denote  $T_n^* := n^{1/2} [T(\eta_n^*, \mathbb{P}_n^*) - T(\eta_n, \hat{P}_n)]$ . If conditions (A1)-(A3) and (B1)-(B3) hold and  $T_n^*$  is asymptotically uniformly square-integrable in the sense that

$$\lim_{m\to\infty}\limsup_{n\to\infty}E_0^*E_W^*\left[T_n^{*2}\mathbf{1}\{T_n^{*2}\geq m\}\right]=0,$$

then  $\bar{\sigma}_n \xrightarrow{P_0^*} \sigma_0$ , so the bootstrap Wald confidence interval defined in (7) has asymptotic confidence level  $1 - \alpha - \beta$ .

**Proof of Theorem 3.14.** Let  $h(x) := x^2$ . There exists a sequence of functions  $h_m : \mathbb{R} \to \mathbb{R}$ ,  $m = 1, 2, \ldots$  such that  $h_m \in BL_1(\mathbb{R})$  for all m and  $mh_m$  monotonically increases to h as  $m \to \infty$ . For instance,  $h_m(x) := \min\{x^2/(m \lor 4), 1\}$  satisfies these criteria. We then write

$$E_{W}^{*}h(T_{n}^{*}) - E_{0}h(\mathbb{G}_{0}\phi_{0}) = \left[E_{W}^{*}h(T_{n}^{*}) - mE_{W}^{*}h_{m}(T_{n}^{*})\right] + \left[mE_{W}^{*}h_{m}(T_{n}^{*}) - mE_{0}h_{m}(\mathbb{G}_{0}\phi_{0})\right]$$

 $+ \left[ m E_0 h_m \left( \mathbb{G}_0 \phi_0 \right) - E_0 h(\mathbb{G}_0 \phi_0) \right].$ 

Hence, for any  $\varepsilon > 0$ , we have

$$\begin{aligned} P_0^* \left( |E_W^* h\left(T_n^*\right) - E_0 h(\mathbb{G}_0 \phi_0)| \ge \varepsilon \right) \\ &\le P_0^* \left( |E_W^* h\left(T_n^*\right) - m E_W^* h_m\left(T_n^*\right)| \ge \varepsilon/3 \right) + P_0^* \left( |E_W^* h_m\left(T_n^*\right) - E_0 h_m\left(\mathbb{G}_0 \phi_0\right)| \ge \varepsilon/(3m) \right) \\ &+ P_0^* \left( |m E_0 h_m\left(\mathbb{G}_0 \phi_0\right) - E_0 h(\mathbb{G}_0 \phi_0)| \ge \varepsilon/3 \right) \\ &\le P_0^* \left( |E_W^* h\left(T_n^*\right) - m E_W^* h_m\left(T_n^*\right)| \ge \varepsilon/3 \right) + P_0^* \left( \sup_{g \in \mathrm{BL}_1(\mathbb{R})} |E_W^* g\left(T_n^*\right) - E_0 g\left(\mathbb{G}_0 \phi_0\right)| \ge \varepsilon/(3m) \right) \\ &+ 1 \left\{ E_0 \left| m h_m\left(\mathbb{G}_0 \phi_0\right) - h(\mathbb{G}_0 \phi_0) \right| \ge \varepsilon/3 \right\}. \end{aligned}$$

By conditions (B1)-(B3), we have  $E_W^*T_n^* \overset{P_0^*}{\underset{W}{\longrightarrow}} \mathbb{G}_0\phi_0$ , i.e.,  $\sup_{g\in \mathrm{BL}_1(\mathbb{R})} |E_W^*g(T_n^*) - E_0g(\mathbb{G}_0\phi_0)| \xrightarrow{P_0^*} 0$ . Therefore, for each fixed m, the second term on the right-hand side of previous display converges to 0. By the monotone convergence theorem, the third term on the right-hand side of previous display display converges to 0 as  $m \longrightarrow \infty$ . For the first term on the right hand side of previous display, we note that

$$\begin{aligned} P_0^* \left( |E_W^* h\left(T_n^*\right) - m E_W^* h_m\left(T_n^*\right)| \ge \varepsilon/3 \right) &= P_0^* \left( \left|E_W^* T_n^{*2} - E_W^* \min\{T_n^{*2}, m\}\right| \ge \varepsilon/3 \right) \\ &= P_0^* \left( \left|E_W^* (T_n^{*2} - m) 1\{T_n^{*2} \ge m\}\right| \ge \varepsilon/3 \right) \\ &\le P_0^* \left( \left|E_W^* T_n^{*2} 1\{T_n^{*2} \ge m\}\right| \ge \varepsilon/3 \right) \\ &\le E_0^* E_W^* T_n^{*2} 1\{T_n^{*2} \ge m\} / (\varepsilon/3), \end{aligned}$$

and  $\lim_{m\to\infty}\limsup_{n\to\infty}$  of this latter expression is 0 by assumption. Therefore, we have

$$\limsup_{n \to \infty} P_0^* \left( \left| E_W^* h\left( T_n^* \right) - E_0 h(\mathbb{G}_0 \phi_0) \right| \ge \varepsilon \right) = \lim_{m \to \infty} \limsup_{n \to \infty} P_0^* \left( \left| E_W^* h\left( T_n^* \right) - E_0 h(\mathbb{G}_0 \phi_0) \right| \ge \varepsilon \right) = 0$$

for every  $\varepsilon > 0$ . Hence,  $\bar{\sigma}_n^2 = E_W^* T_n^{*2} \xrightarrow{P_0^*} E_0(\mathbb{G}_0 \phi_0)^2 = \sigma_0^2$ . By conditions (A1)–(A3), we have  $n^{1/2}[T(\eta_n, \mathbb{P}_n) - T(\eta_0, P_0)] \rightsquigarrow \mathbb{G}_0 \phi_0$ . By Slutsky's theorem, we have  $n^{1/2} \bar{\sigma}_n^{-1}[T(\eta_n, \mathbb{P}_n) - T(\eta_0, P_0)] \rightsquigarrow \sigma_0^{-1} \mathbb{G}_0 \phi_0$ . We then have

$$P_0\left(T(\eta_n, \mathbb{P}_n) + z_\beta \bar{\sigma}_n n^{-1/2} \le T(\eta_0, P_0) \le T(\eta_n, \mathbb{P}_n) + z_{1-\alpha} \bar{\sigma}_n n^{-1/2}\right)$$
$$= P_0\left(-z_{1-\alpha} \le n^{1/2} [T(\eta_n, \mathbb{P}_n) - T(\eta_0, P_0)] / \bar{\sigma}_n \le -z_\beta\right)$$

$$\rightarrow P_0 \left( -z_{1-\alpha} \le \mathbb{G}_0 \phi_0 / \sigma_0 \le -z_\beta \right)$$
$$= 1 - \alpha - \beta,$$

and the result follows.

## Supplement C Proof of results in Section 4

We first introduce a Lemma providing conditions under which the (bootstrap) estimating equationsbased estimator is (conditionally) consistent, which may be useful in its own right.

Lemma C.1. If  $\psi_0$  is a well-separated solution of  $\psi \mapsto G_{0,\eta_0}(\psi)$ ,  $\mathbb{P}_n \phi_{\psi_n,\eta_n} = o_{P_0^*}(1)$ ,  $\phi_{\psi_n,\eta_n}$  is contained in a  $P_0$ -Glivenko Cantelli class with probability tending to one,  $\sup_{|\psi| \leq M} |P_0(\phi_{\psi,\eta_n} - \phi_{\psi,\eta_0})| = o_{P_0^*}(1)$ for all M > 0, and  $\psi_n = O_{P_0^*}(1)$ , then  $\psi_n - \psi_0 = o_{P_0^*}(1)$ . If  $\psi_0$  is a well-separated solution of  $\psi \mapsto G_{0,\eta_0}(\psi)$ ,  $\mathbb{P}_n^* \phi_{\psi_n^*,\eta_n^*} = o_{P_W^*}(1)$ ,  $\phi_{\psi_n^*,\eta_n^*}$  is contained in a  $P_0$ -Glivenko Cantelli class  $\mathfrak{F}$  with conditional probability tending to one,  $\sup_{|\psi| \leq M} |P_0(\phi_{\psi,\eta_n^*} - \phi_{\psi,\eta_0})| = o_{P_W^*}(1)$  for all M > 0,  $\psi_n^* = O_{P_W^*}(1)$ , and  $\|\hat{P}_n - P_0\|_{\mathfrak{F}} = o_{P_0^*}(1)$ , then  $\psi_n^* - \psi_0 = o_{P_W^*}(1)$ .

**Proof of Lemma C.1.** By the well-separated assumption, for every  $\varepsilon > 0$  there exists  $\delta > 0$  such that  $|P_0\phi_{\psi,\eta_0}| \ge \delta$  for every  $\psi \in \mathbb{R}$  such that  $|\psi - \psi_0| \ge \varepsilon$ . Therefore, the event  $\{|\psi_n - \psi_0| \ge \varepsilon\}$  is contained in the event  $\{|P_0\phi_{\psi_n,\eta_0}| \ge \delta\}$ . We then write

$$P_0\phi_{\psi_n,\eta_0} = -(\mathbb{P}_n - P_0)\phi_{\psi_n,\eta_n} + \mathbb{P}_n\phi_{\psi_n,\eta_n} - P_0\left(\phi_{\psi_n,\eta_n} - \phi_{\psi_n,\eta_0}\right).$$

By the assumption that  $\phi_{\psi_n,\eta_n}$  is contained in a  $P_0$ -Glivenko Cantelli class with probability tending to one,  $(\mathbb{P}_n - P_0)\phi_{\psi_n,\eta_n} = o_{P_0^*}(1)$ , and  $\mathbb{P}_n\phi_{\psi_n,\eta_n} = o_{P_0^*}(1)$  by assumption. Since  $\psi_n = O_{P_0^*}(1)$ , for all  $\gamma > 0$ there exists M such that  $P_0^*(|\psi_n| > M) < \gamma$  for all n large enough. Thus,

$$P_0\left(|P_0(\phi_{\psi_n,\eta_n} - \phi_{\psi_n,\eta_0})| \ge \delta\right) \le P_0\left(\sup_{|\psi| \le M} |P_0(\phi_{\psi,\eta_n} - \phi_{\psi,\eta_0})| \ge \delta\right) + P_0(|\psi_n| > M),$$

which is less than  $2\gamma$  for all *n* large enough since  $\sup_{|\psi| \le M} |P_0(\phi_{\psi,\eta_n} - \phi_{\psi,\eta_0})| = o_{P_0^*}(1)$  by assumption.

Therefore, for any  $\varepsilon > 0$ ,

$$P_0^*\left(|\psi_n - \psi_0| \ge \varepsilon\right) \le P_0^*\left(|P_0\phi_{\psi_n,\eta_0}| \ge \delta\right) \longrightarrow 0.$$

We show that  $\psi_n^* \xrightarrow{P_0^*} \psi_0$  using a similar method. By the well-separated condition, the event  $\{|\psi_n^* - \psi_0| \ge \varepsilon\}$  is contained in the event  $\{|P_0\phi_{\psi_n^*,\eta_0}| \ge \delta\}$ . We then write

$$P_0\phi_{\psi_n^*,\eta_0} = -(\mathbb{P}_n^* - \hat{P}_n)\phi_{\psi_n^*,\eta_n^*} - (\hat{P}_n - P_0)\phi_{\psi_n^*,\eta_n^*} + \mathbb{P}_n^*\phi_{\psi_n^*,\eta_n^*} - P_0\left(\phi_{\psi_n^*,\eta_n^*} - \phi_{\psi_n^*,\eta_0}\right)$$

By the assumption that  $\phi_{\psi_n^*,\eta_n^*}$  is contained in a  $P_0$ -Glivenko Cantelli class  $\mathcal{F}$  with conditional probability tending to one,  $(\mathbb{P}_n^* - \hat{P}_n)\phi_{\psi_n^*,\eta_n^*} = o_{P_W^*}(1)$  and  $|(\hat{P}_n - P_0)\phi_{\psi_n^*,\eta_n^*}| \leq ||\hat{P}_n - P_0||_{\mathcal{F}} = o_{P_W^*}(1)$ , and  $\mathbb{P}_n^*\phi_{\psi_n^*,\eta_n^*} = o_{P_W^*}(1)$  by assumption. Since  $\psi_n^* = O_{P_W^*}(1)$ , for all  $\gamma > 0$  there exists M such that  $P_0^*(P_W^*(|\psi_n^*| > M) \geq \gamma/2) = o(1)$ . Thus,

$$\begin{split} P_{0}^{*}\left(P_{W}^{*}\left(\left|P_{0}\left(\phi_{\psi_{n}^{*},\eta_{n}^{*}}-\phi_{\psi_{n}^{*},\eta_{0}}\right)\right|\geq\delta\right)\geq\gamma\right)\leq P_{0}^{*}\left(P_{W}^{*}\left(\sup_{|\psi|\leq M}\left|P_{0}\left(\phi_{\psi,\eta_{n}^{*}}-\phi_{\psi,\eta_{0}}\right)\right|\geq\delta\right)>\gamma/2\right)\right.\\ &+P_{0}^{*}\left(P_{W}^{*}(|\psi_{n}^{*}|>M)>\gamma/2\right),\end{split}$$

and both terms are o(1). The result follows.

Lemma 4.2. If condition (A1) holds for  $\phi_n := \phi_{\psi_n,\eta_n}$ ,  $\psi_0$  is a well-separated solution of the population estimating equation,  $\psi_n = O_{P_0^*}(1)$ , there exists a map  $G'_{0,\eta} : \mathbb{R} \to \mathbb{R}$  depending on  $\eta \in \mathcal{H}$  such that  $\Gamma_{0,\eta} :$  $\psi \mapsto G_{0,\eta}(\psi) - G_{0,\eta}(\psi_0) - G'_{0,\eta}(\psi_0)(\psi - \psi_0)$  satisfies  $\sup_{|\psi| \le M} |\Gamma_{0,\eta_n}(\psi) - \Gamma_{0,\eta_0}(\psi)| = o_{P_0^*}(1)$  for every M > 0 and  $\sup_{\eta: ||\eta - \eta_0||_{\mathcal{H}} < \delta} |\Gamma_{0,\eta}(\psi)| = o(|\psi - \psi_0|)$  for some  $\delta > 0$ ,  $G'_{0,\eta}$  satisfies  $\lim_{\eta \to \eta_0} G'_{0,\eta}(\psi_0) =$  $G'_{0,\eta_0}(\psi_0) = -1$ ,  $||\eta_n - \eta_0||_{\mathcal{H}} = o_{P_0^*}(1)$ , and  $P_0\phi_{\psi_0,\eta_n} = o_{P_0^*}(n^{-1/2})$ , then (A3) holds for the estimating equations-based estimator.

**Proof of Lemma 4.2.** We first show that the assumptions imply that  $\psi_n \xrightarrow{P_0^*} \psi_0$ . By assumption,  $\psi_0$  is a well-separated solution of  $\psi \mapsto G_{0,\eta_0}(\psi)$  and  $\mathbb{P}_n \phi_{\psi_n,\eta_n} = o_{P_0^*}(1)$ . By condition (A1),  $\phi_n := \phi_{\psi_n,\eta_n}$  is contained in a  $P_0$ -Donsker class with probability tending to one, which implies that it is contained in a  $P_0$ -Glivenko Cantelli class with probability tending to one. By adding and subtracting terms, we can write

$$P_0(\phi_{\psi,\eta_n} - \phi_{\psi,\eta_0}) = \Gamma_{0,\eta_n}(\psi) - \Gamma_{0,\eta_0}(\psi) + P_0\phi_{\psi_0,\eta_n} + \left[G'_{0,\eta_n}(\psi_0) + 1\right](\psi - \psi_0)$$

By assumption,  $\sup_{|\psi| \leq M} |\Gamma_{0,\eta_n}(\psi) - \Gamma_{0,\eta_0}(\psi)| = o_{P_0^*}(1), P_0\phi_{\psi_0,\eta_n} = o_{P_0^*}(1)$ , and

$$\sup_{|\psi| \le M} \left| \left[ G'_{0,\eta_n}(\psi_0) + 1 \right] (\psi - \psi_0) \right| = o_{P_0^*}(1)$$

for every M > 0 since  $\eta_n \xrightarrow{P_0^*} \eta_0$ . Finally,  $\psi_n = O_{P_0^*}(1)$  by assumption. Thus, the conditions of Lemma C.1 are satisfied, so  $\psi_n \xrightarrow{P_0^*} \psi_0$ .

We next show that  $\psi_n - \psi_0 = O_{P_0^*}(n^{-1/2})$  under the assumptions of the lemma. We have

$$n^{1/2}(\psi_n - \psi_0) = \mathbb{G}_n \phi_{\psi_n, \eta_n} - n^{1/2} \mathbb{P}_n \phi_{\psi_n, \eta_n} + n^{1/2} P_0 \phi_{\psi_0, \eta_n} + n^{1/2} \left[ P_0(\phi_{\psi_n, \eta_n} - \phi_{\psi_0, \eta_n}) + (\psi_n - \psi_0) \right].$$

Now,  $\mathbb{G}_n \phi_{\psi_n,\eta_n} = O_{P_0^*}(1)$  by (A1), and  $n^{1/2} \mathbb{P}_n \phi_{\psi_n,\eta_n}$  and  $n^{1/2} P_0 \phi_{\psi_0,\eta_n}$  are both  $o_{P_0^*}(1)$  by assumption. Since  $\|\eta_n - \eta_0\|_{\mathcal{H}} = o_{P_0^*}(1)$  by assumption, with probability tending to one it holds that

$$\begin{aligned} |P_0(\phi_{\psi_n,\eta_n} - \phi_{\psi_0,\eta_n}) + (\psi_n - \psi_0)| &= |G_{0,\eta_n}(\psi_n) - G_{0,\eta_n}(\psi_0) + (\psi_n - \psi_0)| \\ &\leq |\Gamma_{0,\eta_n}(\psi_n)| + |G'_{0,\eta_n}(\psi_0) + 1| |\psi_n - \psi_0| \\ &\leq \sup_{\eta: \|\eta - \eta_0\|_{\mathcal{H}} < \delta} |\Gamma_{0,\eta}(\psi_n)| + |G'_{0,\eta_n}(\psi_0) + 1| |\psi_n - \psi_0| .\end{aligned}$$

Now since  $\|\eta_n - \eta_0\|_{\mathcal{H}} = o_{P_0^*}(1)$  and  $G_{0,\eta}(\psi_0)$  is continuous in  $\eta$  at  $\eta_0$  with  $G_{0,\eta}(\psi_0) = -1$ ,  $\left|G'_{0,\eta_n}(\psi_0) + 1\right| = o_{P_0^*}(1)$ . Since  $\psi_n \xrightarrow{P_0^*} \psi_0$  as established above, we then have  $\sup_{\eta: \|\eta - \eta_0\|_{\mathcal{H}} < \delta} |\Gamma_{0,\eta}(\psi_n)| = o_{P_0^*}(|\psi_n - \psi_0|)$  by the assumed differentiability. Hence, we have

$$n^{1/2}|\psi_n - \psi_0| = O_{P_0^*}(1) + n^{1/2}o_{P_0^*}(|\psi_n - \psi_0|),$$

which implies that  $\psi_n - \psi_0 = O_{P_0^*}(n^{-1/2}).$ 

Now since  $G'_{0,\eta_n}(\psi_0) + 1 = o_{P_0^*}(1)$ ,  $P_0\phi_{\psi_0,\eta_n} = o_{P_0^*}(n^{-1/2})$ , and  $\psi_n - \psi_0 = O_{P_0^*}(n^{-1/2})$ , we have

$$\psi_n - \psi_0 + P_0 \phi_{\psi_n,\eta_n} = \left[ P_0(\phi_{\psi_n,\eta_n} - \phi_{\psi_0,\eta_n}) - G'_{0,\eta_n}(\psi_0)(\psi_n - \psi_0) \right] \\ + \left[ G'_{0,\eta_n}(\psi_0) + 1 \right] (\psi_n - \psi_0) + P_0 \phi_{\psi_0,\eta_n} \\ = \Gamma_{0,\eta_n}(\psi_n) + o_{P_0^*}(n^{-1/2}).$$

The first term is  $o_{P_0^*}(n^{-1/2})$  by the differentiability assumption since  $\|\eta_n - \eta_0\|_{\mathcal{H}} = o_{P_0^*}(1)$  and  $\psi_n - \psi_0 = O_{P_0^*}(n^{-1/2})$  as discussed above. The result follows.

Lemma 4.3. Let  $\hat{P}_n = \mathbb{P}_n$  be the empirical bootstrap. Suppose the conditions of Lemma 4.2 hold, condition (B1) holds for  $\phi_n^* = \phi_{\psi_n^*,\eta_n^*}$ ,  $\psi_n^* = O_{P_W^*}(1)$ ,  $\sup_{|\psi| \leq M} |\Gamma_{0,\eta_n^*}(\psi) - \Gamma_{0,\eta_0}(\psi_0)| = o_{P_W^*}(1)$  for every M > 0,  $\|\eta_n^* - \eta_0\|_{\mathcal{H}} = o_{P_W^*}(1)$ , and  $P_0\phi_{\psi_0,\eta_n^*} = o_{P_W^*}(1)$ . If  $P_0\phi_{\psi_0,\eta_n^*} = o_{P_W^*}(n^{-1/2})$ , then (B3) holds for the estimating equations-based estimator. If  $P_0\phi_{\psi_0,\eta_n^*} - 2P_0\phi_{\psi_0,\eta_n} = o_{P_W^*}(n^{-1/2})$ , then  $R_n^* - R_n = o_{P_W^*}(n^{-1/2})$ holds for the estimating equations-based estimator.

**Proof of Lemma 4.3.** We first show that the assumptions imply that  $\psi_n^* \xrightarrow{P_0^*}_W \psi_0$ . By assumption,  $\psi_0$  is a well-separated solution of  $\psi \mapsto G_{0,\eta_0}(\psi)$  and  $\mathbb{P}_n^* \phi_{\psi_n^*,\eta_n^*} = o_{P_0^*}(1)$ . By condition (A1),  $\phi_n^* := \phi_{\psi_n^*,\eta_n^*}$  is contained in a  $P_0$ -Donsker class with probability tending to one, which implies that it is contained in a  $P_0$ -Glivenko Cantelli class with probability tending to one. Adding and subtracting terms, we can write

$$P_0\left(\phi_{\psi,\eta_n^*} - \phi_{\psi,\eta_0}\right) = \Gamma_{0,\eta_n^*}(\psi) - \Gamma_{0,\eta_0}(\psi) + P_0\phi_{\psi_0,\eta_n^*} + \left[G'_{0,\eta_n^*}(\psi_0) + 1\right](\psi - \psi_0)$$

By assumption,  $\sup_{|\psi| \le M} \left| \Gamma_{0,\eta_n^*}(\psi) - \Gamma_{0,\eta_0}(\psi_0) \right| = o_{P_W^*}(1), P_0\phi_{\psi_0,\eta_n^*} = o_{P_0^*}(1)$ , and

$$\sup_{|\psi| \le M} \left| \left[ G'_{0,\eta_n^*}(\psi_0) + 1 \right] (\psi - \psi_0) \right| = o_{P_0^*}(1)$$

for every M > 0 since  $\eta_n^* \xrightarrow{P_0^*} \eta_0$ . Finally,  $\psi_n^* = O_{P_W^*}(1)$  and  $\|\mathbb{P}_n - P_0\|_{\mathcal{F}} = o_{P_W^*}(1)$  by condition (A1). Thus, the conditions of Lemma C.1 are satisfied, so  $\psi_n^* \xrightarrow{P_0^*} \psi_0$ .

We next show that  $|P_0(\phi_{\psi_n^*,\eta_n^*} - \phi_{\psi_0,\eta_n^*}) + (\psi_n^* - \psi_0)| = o_{P_W^*}(|\psi_n^* - \psi_0|)$  under the assumptions of the lemma. Since  $\|\eta_n^* - \eta_0\|_{\mathcal{H}} = o_{P_W^*}(1)$  by assumption, with conditional probability tending to one it holds that

$$\begin{split} \left| P_0(\phi_{\psi_n^*,\eta_n^*} - \phi_{\psi_0,\eta_n^*}) + (\psi_n^* - \psi_0) \right| &= \left| G_{0,\eta_n^*}(\psi_n^*) - G_{0,\eta_n^*}(\psi_0) + (\psi_n^* - \psi_0) \right| \\ &\leq \left| \Gamma_{0,\eta_n^*}(\psi_n^*) \right| + \left| G'_{0,\eta_n^*}(\psi_0) + 1 \right| \left| \psi_n^* - \psi_0 \right| \\ &\leq \sup_{\eta: \|\eta - \eta_0\|_{\mathcal{H}} < \delta} \left| \Gamma_{0,\eta}(\psi_n^*) \right| + \left| G'_{0,\eta_n^*}(\psi_0) + 1 \right| \left| \psi_n^* - \psi_0 \right| . \end{split}$$

Now since  $\|\eta_n^* - \eta_0\|_{\mathcal{H}} = o_{P_W^*}(1)$  and  $G_{0,\eta}(\psi_0)$  is continuous in  $\eta$  at  $\eta_0$  with  $G_{0,\eta_0}(\psi_0) = -1$ ,  $\left|G'_{0,\eta_n^*}(\psi_0) + 1\right| = o_{P_W^*}(1)$ . Since  $\psi_n^* \xrightarrow{P_0^*}_W \psi_0$  as shown above, we also have  $\sup_{\eta: \|\eta - \eta_0\|_{\mathcal{H}} < \delta} |\Gamma_{0,\eta}(\psi_n^*)| = o_{P_W^*}(|\psi_n^* - \psi_0|)$ . The result follows.

We now show that  $\psi_n^* - \psi_0 = O_{P_W^*}(n^{-1/2})$  under the assumptions of the lemma. We have

$$n^{1/2}(\psi_n^* - \psi_0) = \mathbb{G}_n^* \phi_{\psi_n^*, \eta_n^*} + \mathbb{G}_n \phi_{\psi_n^*, \eta_n^*} - n^{1/2} \mathbb{P}_n^* \phi_{\psi_n^*, \eta_n^*} + n^{1/2} P_0 \phi_{\psi_0, \eta_n^*} + n^{1/2} \left[ P_0(\phi_{\psi_n^*, \eta_n^*} - \phi_{\psi_0, \eta_n^*}) + (\psi_n^* - \psi_0) \right].$$

Now,  $\mathbb{G}_n^* \phi_{\psi_n^*, \eta_n^*} = O_{P_W^*}(1)$  by (B1),  $\mathbb{G}_n \phi_{\psi_n^*, \eta_n^*} = O_{P_W^*}(1)$  by (A1), and  $n^{1/2} \mathbb{P}_n^* \phi_{\psi_n^*, \eta_n^*}$  and  $n^{1/2} P_0 \phi_{\psi_0, \eta_n^*}$ are both  $o_{P_W^*}(1)$  by assumption. We therefore have

$$\begin{split} n^{1/2}|\psi_n^* - \psi_0| &= O_{P_W^*}(1) + n^{1/2} \left| P_0(\phi_{\psi_n^*,\eta_n^*} - \phi_{\psi_0,\eta_n^*}) + (\psi_n^* - \psi_0) \right| \\ &= O_{P_W^*}(1) + n^{1/2} o_{P_W^*}(|\psi_n^* - \psi_0|), \end{split}$$

which implies that  $\psi_n^* - \psi_0 = O_{P_W^*}(n^{-1/2}).$ 

We now show the first statement of Lemma 4.3. We write

$$\psi_{n}^{*} - \psi_{n} - (\mathbb{P}_{n}^{*} - \mathbb{P}_{n})\phi_{\psi_{n}^{*},\eta_{n}^{*}} = [\psi_{n}^{*} - \psi_{0} + P_{0}\phi_{\psi_{n}^{*},\eta_{n}^{*}}] - [\psi_{n} - \psi_{0} + P_{0}\phi_{\psi_{n},\eta_{n}}] + (\mathbb{P}_{n} - P_{0})(\phi_{\psi_{n}^{*},\eta_{n}^{*}} - \phi_{\psi_{n},\eta_{n}}) + \mathbb{P}_{n}\phi_{\psi_{n},\eta_{n}} - \mathbb{P}_{n}^{*}\phi_{\psi_{n}^{*},\eta_{n}^{*}}.$$
(22)

By Lemma 4.2, we have  $\psi_n - \psi_0 + P_0 \phi_{\psi_n,\eta_n} = o_{P_0^*}(n^{-1/2})$ . By the definitions of  $\psi_n$  and  $\psi_n^*$ , we have  $\mathbb{P}_n \phi_{\psi_n,\eta_n} = o_{P_W^*}(n^{-1/2})$  and  $\mathbb{P}_n^* \phi_{\psi_n,\eta_n^*} = o_{P_W^*}(n^{-1/2})$ . By conditions (A1) and (A2) and Lemma 19.24 of van der Vaart (2000), we have  $(\mathbb{P}_n - P_0)(\phi_{\psi_n,\eta_n} - \phi_{\psi_0,\eta_0}) = o_{P_0^*}(n^{-1/2})$ . Similarly, by (A1)(b), (B1)(a), (B2) and a minor modification of Lemma 19.24 van der Vaart (2000), we have  $(\mathbb{P}_n - P_0)(\phi_{\psi_n^*,\eta_n^*} - \phi_{\psi_0,\eta_0}) = o_{P_W^*}(n^{-1/2})$ . Hence,  $(\mathbb{P}_n - P_0)(\phi_{\psi_n^*,\eta_n^*} - \phi_{\psi_n,\eta_n}) = o_{P_W^*}(n^{-1/2})$ . Finally, since  $G'_{0,\eta_n^*}(\psi_0) + 1 = o_{P_W^*}(1)$ ,  $P_0\phi_{\psi_0,\eta_n^*} = o_{P_W^*}(n^{-1/2})$ , and  $\psi_n^* - \psi_0 = O_{P_W^*}(n^{-1/2})$ , we have

$$\begin{aligned} |\psi_n^* - \psi_0 + P_0 \phi_{\psi_n^*, \eta_n^*}| &\leq \left| P_0(\phi_{\psi_n^*, \eta_n^*} - \phi_{\psi_0, \eta_n^*}) + (\psi_n^* - \psi_0) \right| + \left| P_0 \phi_{\psi_0, \eta_n^*} \right| \\ &= o_{P_W^*}(|\psi_n^* - \psi_0|) + o_{P_W^*}(n^{-1/2}) \\ &= o_{P_W^*}(n^{-1/2}). \end{aligned}$$

The result follows.

We next show the second statement in Lemma 4.3. Since  $R_n = \psi_n - \psi_0 - (\mathbb{P}_n - P_0)\phi_{\psi_n,\eta_n}$  and

 $R_n^* = \psi_n^* - \psi_n - (\mathbb{P}_n^* - \mathbb{P}_n)\phi_{\psi_n^*,\eta_n^*}$ , by adding and subtracting terms, we have

$$R_n^* - R_n = \left(\psi_n^* - \psi_0 + P_0 \phi_{\psi_n,\eta_n^*} - P_0 \phi_{\psi_0,\eta_n^*}\right) - 2\left(\psi_n - \psi_0 + P_0 \phi_{\psi_n,\eta_n} - P_0 \phi_{\psi_0,\eta_n}\right) \\ + \left(P_0 \phi_{\psi_0,\eta_n^*} - 2P_0 \phi_{\psi_0,\eta_n}\right) + \left(\mathbb{P}_n - P_0\right)\left(\phi_{\psi_n^*,\eta_n^*} - \phi_{\psi_n,\eta_n}\right) + \left(2\mathbb{P}_n \phi_{\psi_n,\eta_n} - \mathbb{P}_n^* \phi_{\psi_n^*,\eta_n^*}\right).$$

As we showed above,  $\psi_n^* - \psi_0 + P_0 \phi_{\psi_n,\eta_n^*} - P_0 \phi_{\psi_0,\eta_n^*} = o_{P_W^*}(n^{-1/2})$  and  $(\mathbb{P}_n - P_0)(\phi_{\psi_n,\eta_n^*} - \phi_{\psi_n,\eta_n}) = o_{P_W^*}(n^{-1/2})$ . By assumption,  $P_0 \phi_{\psi_0,\eta_n^*} - 2P_0 \phi_{\psi_0,\eta_n} = o_{P_W^*}(n^{-1/2})$ . By the definitions of  $\psi_n^*$  and  $\psi_n$ ,  $2\mathbb{P}_n \phi_{\psi_n,\eta_n}$  and  $\mathbb{P}_n^* \phi_{\psi_n^*,\eta_n^*}$  are both  $o_{P_W^*}(n^{-1/2})$ . Therefore,

$$R_n^* - R_n = -2\left(\psi_n - \psi_0 + P_0\phi_{\psi_n,\eta_n} - P_0\phi_{\psi_0,\eta_n}\right) + o_{P_W^*}(n^{-1/2})$$
$$= -2\Gamma_{0,\eta_n}(\psi_n) - 2\left[G'_{0,\eta_n}(\psi_0) + 1\right](\psi_n - \psi_0) + o_{P_W^*}(n^{-1/2}).$$

The first two terms are  $o_{P_0^*}(n^{-1/2})$  as discussed above. The result follows.

Lemma 4.4. Let  $\hat{P}_n$  be the smooth bootstrap. Suppose the conditions of Lemma 4.2 hold, condition (B1) holds for  $\phi_n^* = \phi_{\psi_n^*,\eta_n^*}, \psi_n^* = O_{P_W^*}(1), \|\eta_n^* - \eta_0\|_{\mathcal{H}} = o_{P_W^*}(1), \sup_{|\psi| \leq M} |\Gamma_{0,\hat{\eta}_n}(\psi) - \Gamma_{0,\eta_0}(\psi)| = o_{P_0^*}(1)$  for every M > 0, there exists a map  $\hat{G}'_{n,\eta} : \mathbb{R} \to \mathbb{R}$  such that  $\hat{\Gamma}_{n,\eta} : \psi \mapsto \hat{G}_{n,\eta}(\psi) - \hat{G}_{n,\eta}(\hat{\psi}_n) - \hat{G}'_{n,\eta}(\hat{\psi}_n)(\psi - \hat{\psi}_n)$  satisfies  $\sup_{|\psi| \leq M} |\hat{\Gamma}_{n,\eta_n^*}(\psi) - \hat{\Gamma}_{n,\hat{\eta}_n}(\hat{\psi}_n)| = o_{P_W^*}(1)$  for every M > 0 and  $\sup_{\eta: \|\eta - \hat{\eta}_n\|_{\mathcal{H}} < \delta} |\hat{\Gamma}_{n,\eta}(\psi)| = o_{P_0^*}(\psi - \hat{\psi}_n)$  for some  $\delta > 0$ ,  $\hat{G}'_{n,\eta}$  satisfies  $\hat{G}'_{n,\eta_n^*}(\hat{\psi}_n) + 1 = o_{P_W^*}(1)$ , there exists a P<sub>0</sub>-Glivenko Cantelli class  $\mathfrak{F}$  such that  $\phi_{\hat{\psi}_n,\hat{\eta}_n}$  is contained in  $\mathfrak{F}$  with probability tending to one,  $\phi_{\psi_n^*,\eta_n^*}$  is contained in  $\mathfrak{F}$  with conditional probability tending to one, and  $\|\hat{P}_n - P_0\|_{\mathfrak{F}} = o_{P_W^*}(1)$ ,  $\hat{\psi}_n = O_{P_0^*}(1)$ ,  $P_0\phi_{\psi_0,\hat{\eta}_n} = o_{P_0^*}(1)$ ,  $\|\hat{\eta}_n - \eta_0\|_{\mathfrak{H}} = o_{P_0^*}(1)$ , and  $\psi_n^\circ - \hat{\psi}_n = o_{P_0^*}(n^{-1/2})$ . If  $\hat{P}_n\phi_{\hat{\psi}_n,\eta_n^*} = o_{P_W^*}(n^{-1/2})$ , then (B3) holds for the estimating equations-based estimator. If  $\hat{P}_n\phi_{\hat{\psi}_n,\eta_n^*} - P_0\phi_{\psi_0,\eta_n} = o_{P_W^*}(n^{-1/2})$ , then  $R_n^* - R_n = o_{P_W^*}(n^{-1/2})$ holds for the estimating equations-based estimator.

**Proof of Lemma 4.4.** By the same argument as used in Lemma 4.3, and since  $\|\hat{P}_n - P_0\|_{\mathcal{F}} = o_{P_W^*}(1)$  by assumption, we can show that the conditions of Lemma C.1 are satisfied, so that  $\psi_n^* \xrightarrow[W]{W}{W} \psi_0$ .

We recall that  $\hat{\psi}_n := T(\hat{\eta}_n, \hat{P}_n)$ . By the same argument as used in Lemma 4.3,  $P_0 \phi_{\hat{\psi}_n, \eta_0} = o_{P_0^*}(1)$ implies that  $\hat{\psi}_n - \psi_0 = o_{P_0^*}(1)$ . We write

$$P_{0}\phi_{\hat{\psi}_{n},\eta_{0}} = \hat{P}_{n}\phi_{\hat{\psi}_{n},\hat{\eta}_{n}} - \left(\hat{P}_{n} - P_{0}\right)\left(\phi_{\hat{\psi}_{n},\hat{\eta}_{n}}\right) - P_{0}\left(\phi_{\hat{\psi}_{n},\hat{\eta}_{n}} - \phi_{\hat{\psi}_{n},\eta_{0}}\right).$$

By the definition of  $\hat{\psi}_n$ , we have  $\hat{P}_n \phi_{\hat{\psi}_n, \hat{\eta}_n} = 0$ . Under the assumptions  $\|\hat{P}_n - P_0\|_{\mathcal{F}} = o_{P_0^*}(1)$  and  $\phi_{\hat{\psi}_n, \hat{\eta}_n}$  is contained in  $\mathcal{F}$  with probability tending to one, we have  $(\hat{P}_n - P_0)\phi_{\hat{\psi}_n, \hat{\eta}_n} = o_{P_0^*}(1)$ . Finally, since  $\hat{\psi}_n = O_{P_0^*}(1)$  and  $\sup_{|\psi| \leq M} |\Gamma_{0, \hat{\eta}_n}(\psi) - \Gamma_{0, \eta_0}(\psi)| = o_{P_0^*}(1)$  for every M > 0,  $P_0\phi_{\psi_0, \hat{\eta}_n} = o_{P_0^*}(1)$ , and  $\|\hat{\eta}_n - \eta_0\|_{\mathcal{H}} = o_{P_0^*}(1)$ , by the same argument as in Lemma 4.2,  $P_0\left(\phi_{\hat{\psi}_n, \hat{\eta}_n} - \phi_{\hat{\psi}_n, \eta_0}\right) = o_{P_0^*}(1)$ . Thus,  $P_0\phi_{\hat{\psi}_n, \eta_0} = o_{P_0^*}(1)$ , so that  $\hat{\psi}_n \xrightarrow{P_0^*} \psi_0$ . Hence, we also have  $\psi_n^* - \hat{\psi}_n = (\psi_n^* - \psi_0) - (\hat{\psi}_n - \psi_0) \xrightarrow{P_0^*} W_0$ .

We now show that  $\hat{P}_n\left(\phi_{\psi_n^*,\eta_n^*} - \phi_{\hat{\psi}_n,\eta_n^*}\right) + (\psi_n^* - \hat{\psi}_n) = o_{P_W^*}(\psi_n^* - \hat{\psi}_n)$  under the assumptions of the lemma. With conditional probability tending to one, it holds that

$$\begin{aligned} \left| \hat{P}_{n} \left( \phi_{\psi_{n}^{*}, \eta_{n}^{*}} - \phi_{\hat{\psi}_{n}, \eta_{n}^{*}} \right) + \left( \psi_{n}^{*} - \hat{\psi}_{n} \right) \right| \\ &= \left| \hat{G}_{n, \eta_{n}^{*}} (\psi_{n}^{*}) - \hat{G}_{n, \eta_{n}^{*}} (\hat{\psi}_{n}) + \left( \psi_{n}^{*} - \hat{\psi}_{n} \right) \right| \\ &\leq \left| \hat{G}_{n, \eta_{n}^{*}} (\psi_{n}^{*}) - \hat{G}_{n, \eta_{n}^{*}} (\hat{\psi}_{n}) - \hat{G}'_{n, \eta_{n}^{*}} (\hat{\psi}_{n}) (\psi_{n}^{*} - \hat{\psi}_{n}) \right| + \left| \hat{G}'_{n, \eta_{n}^{*}} (\hat{\psi}_{n}) + 1 \right| \left| \psi_{n}^{*} - \hat{\psi}_{n} \right| \\ &\leq \sup_{\eta: \|\eta - \hat{\eta}_{n}\|_{\mathcal{H}} < \delta} \left| \hat{\Gamma}_{n, \eta} (\psi_{n}^{*}) \right| + \left| \hat{G}'_{n, \eta_{n}^{*}} (\hat{\psi}_{n}) + 1 \right| \left| \psi_{n}^{*} - \hat{\psi}_{n} \right| \end{aligned}$$

Since by assumption  $\hat{G}'_{n,\eta^*_n}(\psi_n) + 1 = o_{P^*_W}(1)$ , the second term on the right hand side of previous display is  $o_{P^*_W}(\psi^*_n - \hat{\psi}_n)$ . The result follows.

We next show that  $\psi_n^* - \hat{\psi}_n = O_{P_W^*}(n^{-1/2})$  under the assumptions of the lemma. We have

$$n^{1/2}(\psi_n^* - \hat{\psi}_n) = \mathbb{G}_n^* \phi_{\psi_n^*, \eta_n^*} - n^{1/2} \mathbb{P}_n^* \phi_{\psi_n^*, \eta_n^*} + n^{1/2} \hat{P}_n \phi_{\hat{\psi}_n, \eta_n^*} + n^{1/2} \left[ \hat{P}_n \left( \phi_{\psi_n^*, \eta_n^*} - \phi_{\hat{\psi}_n, \eta_n^*} \right) + \psi_n^* - \hat{\psi}_n \right].$$

Now,  $\mathbb{G}_n^* \phi_{\psi_n,\eta_n^*} = O_{P_W^*}(1)$  because (B1) holds for  $\phi_n^* = \phi_{\psi_n^*,\eta_n^*}$  by assumption, and  $n^{1/2} \mathbb{P}_n^* \phi_{\psi_n,\eta_n^*}$  and  $n^{1/2} \hat{P}_n \phi_{\hat{\psi}_n,\eta_n^*}$  are both  $o_{P_W^*}(1)$  by assumption. As discussed above,  $\hat{P}_n \left( \phi_{\psi_n^*,\eta_n^*} - \phi_{\hat{\psi}_n,\eta_n^*} \right) + \psi_n^* - \hat{\psi}_n = o_{P_W^*}(\psi_n^* - \hat{\psi}_n)$ . Hence, we have

$$n^{1/2}|\psi_n^* - \hat{\psi}_n| = O_{P_W^*}(1) + n^{1/2}o_{P_W^*}(|\psi_n^* - \hat{\psi}_n|),$$

which implies that  $\psi_n^* - \hat{\psi}_n = O_{P_W^*}(n^{-1/2}).$ 

We can now show the first statement of Lemma 4.4. We recall that  $\psi_n^{\circ} := T(\eta_n, \hat{P}_n)$ . We can write

$$\begin{aligned} R_n^* &= \psi_n^* - \psi_n^\circ - (\mathbb{P}_n^* - \hat{P}_n)\phi_{\psi_n^*,\eta_n^*} \\ &= \left(\psi_n^* - \hat{\psi}_n + \hat{P}_n\phi_{\psi_n^*,\eta_n^*} - \hat{P}_n\phi_{\hat{\psi}_n,\eta_n^*}\right) - \left(\psi_n^\circ - \hat{\psi}_n\right) + \hat{P}_n\phi_{\hat{\psi}_n,\eta_n^*} - \mathbb{P}_n^*\phi_{\psi_n^*,\eta_n^*}; \end{aligned}$$

We have  $\psi_n^{\circ} - \hat{\psi}_n = o_{P_0^*}(n^{-1/2})$  and  $\hat{P}_n \phi_{\hat{\psi}_n, \eta_n^*} = o_{P_W^*}(n^{-1/2})$  by assumption,  $\mathbb{P}_n^* \phi_{\psi_n^*, \eta_n^*} = o_{P_W^*}(n^{-1/2})$  by definition of  $\psi_n^*$ , and

$$\psi_n^* - \hat{\psi}_n + \hat{P}_n \phi_{\psi_n^*, \eta_n^*} - \hat{P}_n \phi_{\hat{\psi}_n, \eta_n^*} = o_{P_W^*} \left( \psi_n^* - \hat{\psi}_n \right) = o_{P_W^*} (n^{-1/2})$$

by the derivations above. Hence,  $R_n^* = o_{P_W^*}(n^{-1/2})$ .

Lastly, we show the second statement in Lemma 4.4. Recall that  $R_n = \psi_n - \psi_0 - (\mathbb{P}_n - P_0)\phi_{\psi_n,\eta_n}$ and  $R_n^* = \psi_n^* - \psi_n^\circ - (\mathbb{P}_n^* - \hat{P}_n)\phi_{\psi_n^*,\eta_n^*}$ . By adding and subtracting terms, we can write

$$R_{n}^{*} - R_{n} = \left(\psi_{n}^{*} - \hat{\psi}_{n} + \hat{P}_{n}\phi_{\psi_{n},\eta_{n}^{*}} - \hat{P}_{n}\phi_{\hat{\psi}_{n},\eta_{n}^{*}}\right) - \left(\psi_{n} - \psi_{0} + P_{0}\phi_{\psi_{n},\eta_{n}} - P_{0}\phi_{\psi_{0},\eta_{n}}\right) \\ + \left(\hat{P}_{n}\phi_{\hat{\psi}_{n},\eta_{n}^{*}} - P_{0}\phi_{\psi_{0},\eta_{n}}\right) + \left(\mathbb{P}_{n}\phi_{\psi_{n},\eta_{n}} - \mathbb{P}_{n}^{*}\phi_{\psi_{n}^{*},\eta_{n}^{*}}\right) + \left(\hat{\psi}_{n} - \psi_{n}^{\circ}\right).$$

We have  $\psi_n^* - \hat{\psi}_n + \hat{P}_n \phi_{\psi_n,\eta_n^*} - \hat{P}_n \phi_{\hat{\psi}_n,\eta_n^*} = o_{P_W^*}(n^{-1/2})$  by the derivations above,  $\psi_n - \psi_0 + P_0 \phi_{\psi_n,\eta_n} - P_0 \phi_{\psi_0,\eta_n} = o_{P_0^*}(n^{-1/2})$  by Lemma 4.2,  $\hat{\psi}_n - \psi_n^\circ = o_{P_W^*}(n^{-1/2})$  and  $\hat{P}_n \phi_{\hat{\psi}_n,\eta_n^*} - P_0 \phi_{\psi_0,\eta_n} = o_{P_W^*}(n^{-1/2})$  by assumption, and  $\mathbb{P}_n \phi_{\psi_n,\eta_n} - \mathbb{P}_n^* \phi_{\psi_n,\eta_n^*} = o_{P_W^*}(n^{-1/2})$  by the definitions of  $\psi_n$  and  $\psi_n^*$ .

## Supplement D Proof of results in Section 5

**Proposition 5.1.** Suppose that  $\eta_n$  falls in a  $P_0$ -Donsker class with probability tending to one, there exists  $M \in (0,\infty)$  such that  $\|\eta_0\|_{\infty} < M$  and  $\|\eta_n\|_{\infty} < M$  with probability tending to one, and  $\|\eta_n - \eta_0\|_{L_2(\lambda)} = o_{P_0^*}(n^{-1/4})$ . Then  $\psi_{n,1}$  is asymptotically linear with influence function  $\phi_0$ . If in addition  $\int \eta_n^2 - \mathbb{P}_n \eta_n = o_{P_0^*}(n^{-1/2})$ , then  $\psi_{n,2}$  and  $\psi_{n,3}$  are asymptotically linear as well.

**Proof of Proposition 5.1.** We use Theorem 3.1. Condition (A1) holds because  $\eta_n$  falls in a  $P_0$ -Donsker class by assumption. For any  $\varepsilon > 0$ , by the triangle and Cauchy–Schwarz inequalities, we have

$$\begin{aligned} P_{0}^{*} \left( \left\| \phi_{n} - \phi_{0} \right\|_{L_{2}(P_{0})} \geq \varepsilon \right) \\ &\leq P_{0}^{*} \left( 2 \left\| \eta_{n} - \eta_{0} \right\|_{L_{2}(P_{0})} \geq \varepsilon/2 \right) + P_{0}^{*} \left( 2 \left| \int \eta_{n}^{2} - \int \eta_{0}^{2} \right| \geq \varepsilon/2 \right) \\ &\leq P_{0}^{*} \left( 2M \left\| \eta_{n} - \eta_{0} \right\|_{L_{2}(\lambda)} \geq \varepsilon/2 \right) + P_{0}^{*} \left( 2 \left\| \eta_{n} - \eta_{0} \right\|_{L_{2}(\lambda)} \left\| \eta_{n} + \eta_{0} \right\|_{L_{2}(\lambda)} \geq \varepsilon/2 \right) \\ &\leq P_{0}^{*} \left( 2M \left\| \eta_{n} - \eta_{0} \right\|_{L_{2}(\lambda)} \geq \varepsilon/2 \right) + P_{0}^{*} \left( 4M \left\| \eta_{n} - \eta_{0} \right\|_{L_{2}(\lambda)} \geq \varepsilon/2 \right) + P_{0}^{*} \left( \left\| \eta_{n} \right\|_{L_{2}(\lambda)} \geq \varepsilon/2 \right) + P_{0}^{*} \left( \left\| \eta_{n} \right\|_{L_{2}(\lambda)} \geq M \right). \end{aligned}$$

Each term on the right-hand side converges to 0, which implies condition (A2).

For condition (A3) for  $\psi_{n,1}$ , we have

$$T_{1}(\eta_{n}, \mathbb{P}_{n}) - T_{1}(\eta_{0}, P_{0}) - (\mathbb{P}_{n} - P_{0})\phi_{n} = \left(2\mathbb{P}_{n}\eta_{n} - \int \eta_{n}^{2}\right) - \int \eta_{0}^{2} - 2(\mathbb{P}_{n} - P_{0})\eta_{n}$$
$$= -\int (\eta_{n} - \eta_{0})^{2}$$
$$= -\|\eta_{n} - \eta_{0}\|_{L_{2}(\lambda)}^{2}$$

which is  $o_{P_0^*}(n^{1/2})$  by assumption. For  $\psi_{n,2}$  and  $\psi_{n,3}$ , we have

$$T_{2}(\eta_{n}, \mathbb{P}_{n}) - T_{2}(\eta_{0}, P_{0}) - (\mathbb{P}_{n} - P_{0})\phi_{n} = \int \eta_{n}^{2} - \int \eta_{0}^{2} - 2(\mathbb{P}_{n} - P_{0})\eta_{n}$$
  
$$= -\int (\eta_{n} - \eta_{0})^{2} + 2\left[\int \eta_{n}^{2} - \mathbb{P}_{n}\eta_{n}\right]$$
  
$$T_{3}(\eta_{n}, \mathbb{P}_{n}) - T_{3}(\eta_{0}, P_{0}) - (\mathbb{P}_{n} - P_{0})\phi_{n} = \mathbb{P}_{n}\eta_{n} - P_{0}\eta_{0} - 2(\mathbb{P}_{n} - P_{0})\eta_{n}$$
  
$$= -\int (\eta_{n} - \eta_{0})^{2} + \left[\int \eta_{n}^{2} - \mathbb{P}_{n}\eta_{n}\right],$$

which are both  $o_{P_0^*}(n^{-1/2})$  when  $\int \eta_n^2 - \mathbb{P}_n \eta_n = o_{P_0^*}(n^{-1/2})$  and  $\|\eta_n - \eta_0\|_{L_2(\lambda)}^2 = o_{P_0^*}(n^{-1/4}).$ 

**Proposition 5.2.** Suppose  $\hat{P}_n = \mathbb{P}_n$  is the empirical bootstrap and the assumptions of Proposition 5.1 hold. For  $\mathcal{F}$  and M defined in Proposition 5.1, suppose  $P_W^*(\eta_n^* \in \mathcal{F}) \xrightarrow{P_0^*} 1$ ,  $P_W^*(||\eta_n^*||_{\infty} \ge M) = o_{P_0^*}(1)$ , and  $||\eta_n^* - \eta_n||_{L_2(\lambda)} = o_{P_W^*}(n^{-1/4})$ . Then  $\psi_{n,1}^* = \psi_{n,1} + (\mathbb{P}_n^* - \mathbb{P}_n)\phi_0 + o_{P_W^*}(n^{-1/2})$ . If in addition  $\int \eta_n^{*2} - \mathbb{P}_n^*\eta_n^* = o_{P_W^*}(n^{-1/2})$ , then  $\psi_{n,2}^* = \psi_{n,2} + (\mathbb{P}_n^* - \mathbb{P}_n)\phi_0 + o_{P_W^*}(n^{-1/2})$  and  $\psi_{n,3}^* = \psi_{n,3} + (\mathbb{P}_n^* - \mathbb{P}_n)\phi_0 + o_{P_W^*}(n^{-1/2})$ . If  $\eta_n$  and  $\eta_n^*$  are KDEs with the same kernel and bandwidth h such that  $n^{-1/(2d)} \prec h$ , then  $||\eta_n^* - \eta_n||_{L_2(\lambda)} = o_{P_W^*}(n^{-1/4})$ .

**Proof of Proposition 5.2.** We use Theorem 3.4. Condition (B1) holds by Lemma 3.6 because  $\eta_n^*$  is contained in a  $P_0$ -Donsker class with  $P_W^*$  probability tending to one. For any  $\varepsilon > 0$ , by the triangle and Cauchy–Schwarz inequalities, we have

$$\begin{split} P_W^* \left( \|\phi_n^* - \phi_0\|_{L_2(P_0)} \ge \varepsilon \right) \\ &\leq P_W^* \left( 2 \|\eta_n^* - \eta_0\|_{L_2(P_0)} \ge \varepsilon/2 \right) + P_W^* \left( 2 \left| \int \eta_n^{*2} - \int \eta_0^2 \right| \ge \varepsilon/2 \right) \\ &\leq P_W^* \left( 2M \|\eta_n^* - \eta_0\|_{L_2(\lambda)} \ge \varepsilon/2 \right) + P_W^* \left( 2 \|\eta_n^* - \eta_0\|_{L_2(\lambda)} \|\eta_n^* + \eta_0\|_{L_2(\lambda)} \ge \varepsilon/2 \right) \\ &\leq P_W^* \left( 2M \|\eta_n^* - \eta_0\|_{L_2(\lambda)} \ge \varepsilon/2 \right) + P_W^* \left( 4M \|\eta_n^* - \eta_0\|_{L_2(\lambda)} \ge \varepsilon/2 \right) + P_W^* \left( \|\eta_n^*\|_{L_2(\lambda)} \ge M \right), \end{split}$$

Each term on the right-hand side is  $o_{P_0^*}(1)$ , which implies condition (B2).

For condition (B3) for  $\psi_{n,1}^*$ , we write

$$\begin{aligned} |T_{1}(\eta_{n}^{*},\mathbb{P}_{n}^{*}) - T_{1}(\eta_{n},\mathbb{P}_{n}) - (\mathbb{P}_{n}^{*} - \mathbb{P}_{n})\phi_{n}^{*}| \\ &= \left| \left( 2\mathbb{P}_{n}^{*}\eta_{n}^{*} - \int \eta_{n}^{*2} \right) - \left( 2\mathbb{P}_{n}\eta_{n} - \int \eta_{n}^{2} \right) - (\mathbb{P}_{n}^{*} - \mathbb{P}_{n})2\eta_{n}^{*} \right| \\ &= \left| -\int (\eta_{n}^{*} - \eta_{n})^{2} - 2\int \eta_{n}(\eta_{n}^{*} - \eta_{n}) + 2(\mathbb{P}_{n} - P_{0})(\eta_{n}^{*} - \eta_{n}) + 2P_{0}(\eta_{n}^{*} - \eta_{n}) \right| \\ &= \left| -\int (\eta_{n}^{*} - \eta_{n})^{2} - 2\int (\eta_{n} - \eta_{0})(\eta_{n}^{*} - \eta_{n}) + 2(\mathbb{P}_{n} - P_{0})(\eta_{n}^{*} - \eta_{n}) \right| \\ &\leq \left\| \eta_{n}^{*} - \eta_{n} \right\|_{L_{2}(\lambda)}^{2} + 2\left\| \eta_{n} - \eta_{0} \right\|_{L_{2}(\lambda)} \left\| \eta_{n}^{*} - \eta_{n} \right\|_{L_{2}(\lambda)} + 2n^{-1/2} \left| \mathbb{G}_{n}(\eta_{n}^{*} - \eta_{n}) \right|. \end{aligned}$$

The first two terms are  $o_{P_W^*}(n^{-1/2})$  by assumption. In addition,  $\mathbb{G}_n(\eta_n^* - \eta_0) = o_{P_W^*}(1)$  because  $\|\eta_n^* - \eta_0\|_{L_2(P_0)} = o_{P_W^*}(1)$  and  $\eta_n^*$  is contained in a  $P_0$ -Donsker class with  $P_W^*$ -probability tending to one. Hence, condition (B3) holds for  $\psi_{n,1}^*$ , and the conditional asymptotic linearity result for  $\psi_{n,1}^*$  follows.

For condition (B3) for  $\psi_{n,2}^*$  and  $\psi_{n,3}^*$ , we have

$$\begin{split} T_{2}(\eta_{n}^{*},\mathbb{P}_{n}^{*}) - T_{2}(\eta_{n},\mathbb{P}_{n}) - (\mathbb{P}_{n}^{*} - \mathbb{P}_{n})\phi_{n}^{*} &= \int \eta_{n}^{*2} - \int \eta_{n}^{2} - 2(\mathbb{P}_{n}^{*} - \mathbb{P}_{n})\eta_{n}^{*} \\ &= -\int (\eta_{n}^{*} - \eta_{n})^{2} + 2\left(\int \eta_{n}^{*2} - \mathbb{P}_{n}^{*}\eta_{n}^{*}\right) - 2\left(\int \eta_{n}\eta_{n}^{*} - \mathbb{P}_{n}\eta_{n}^{*}\right) \\ T_{3}(\eta_{n}^{*},\mathbb{P}_{n}^{*}) - T_{3}(\eta_{n},\mathbb{P}_{n}) - (\mathbb{P}_{n}^{*} - \mathbb{P}_{n})\phi_{n}^{*} &= \mathbb{P}_{n}^{*}\eta_{n}^{*} - \mathbb{P}_{n}\eta_{n} - 2(\mathbb{P}_{n}^{*} - \mathbb{P}_{n})\eta_{n}^{*} \\ &= -\int (\eta_{n}^{*} - \eta_{n})^{2} + \left(\int \eta_{n}^{*2} - \mathbb{P}_{n}^{*}\eta_{n}^{*}\right) + \left(\int \eta_{n}^{2} - \mathbb{P}_{n}\eta_{n}\right) \\ &- 2\left(\int \eta_{n}\eta_{n}^{*} - \mathbb{P}_{n}\eta_{n}^{*}\right). \end{split}$$

Now,  $\int (\eta_n^* - \eta_n)^2$  and  $\int \eta_n^{*2} - \mathbb{P}_n^* \eta_n^*$  are both  $o_{P_W^*}(n^{-1/2})$  by assumption, and  $\int \eta_n^2 - \mathbb{P}_n \eta_n = o_{P_0^*}(n^{-1/2})$  by assumption. In addition,

$$\begin{aligned} \left| \int \eta_n \eta_n^* - \mathbb{P}_n \eta_n^* \right| &= \left| \int (\eta_n - \eta_0) (\eta_n^* - \eta_n) - (\mathbb{P}_n - P_0) (\eta_n^* - \eta_n) + \int \eta_n^2 - \mathbb{P}_n \eta_n \right| \\ &\leq \|\eta_n - \eta_0\|_{L_2(\lambda)} \|\eta_n^* - \eta_n\|_{L_2(\lambda)} + n^{-1/2} |\mathbb{G}_n (\eta_n^* - \eta_n)| + \left| \int \eta_n^2 - \mathbb{P}_n \eta_n \right|, \end{aligned}$$

which is  $o_{P_W^*}(n^{-1/2})$ . Therefore, condition (B3) holds for  $\psi_{n,2}^*$  and  $\psi_{n,3}^*$ .

We now turn to the setting where  $\eta_n$  and  $\eta_n^*$  are kernel density estimators using the same bandwidth

h. We note that for any x, since  $\hat{P}_n$  is the empirical bootstrap,

$$E_W^*\left[\eta_n^*(x)\right] = E_W^*\left[\mathbb{P}_n^* h^{-d} K\left(\frac{X_i^* - x}{h}\right)\right] = \mathbb{P}_n h^{-d} K\left(\frac{X_i - x}{h}\right) = \eta_n(x).$$

We therefore have

$$\begin{split} E_W^* \left\| \eta_n^* - \eta_n \right\|_{L_2(\lambda)}^2 &= \int E_W^* [\eta_n^*(x) - \eta_n(x)]^2 \, \mathrm{d}x = \int \operatorname{Var}_W^* (\eta_n^*(x)) \, \mathrm{d}x \\ &= \int \frac{1}{nh^{2d}} \operatorname{Var}_W^* \left[ K\left(\frac{X_1^* - x}{h}\right) \right] \, \mathrm{d}x \\ &\leq \frac{1}{nh^{2d}} \int E_W^* \left[ K\left(\frac{X_1^* - x}{h}\right) \right]^2 \, \mathrm{d}x \\ &= \frac{1}{nh^{2d}} \int \mathbb{P}_n \left[ K\left(\frac{X_i - x}{h}\right) \right]^2 \, \mathrm{d}x \\ &= \frac{1}{nh^d} \int K^2(t) \, \mathrm{d}t. \end{split}$$

Hence, by Chebyshev's inequality, for any  $\varepsilon > 0$ , we have

$$P_{W}^{*}\left(n^{1/4} \|\eta_{n}^{*} - \eta_{n}\|_{L_{2}(\lambda)} \ge \varepsilon\right) \le n^{1/2} E_{W}^{*} \|\eta_{n}^{*} - \eta_{n}\|_{L_{2}(\lambda)}^{2} / \varepsilon^{2} \\ \le \left(nh^{2d}\right)^{-1/2} \varepsilon^{-2} \int K^{2}(t) \, \mathrm{d}t,$$

which goes to zero since  $nh^{2d} \longrightarrow \infty$ . Therefore  $\|\eta_n^* - \eta_n\|_{L_2(\lambda)} = o_{P_W^*}(n^{-1/4})$ .

The proof of Proposition 5.3 relies on technical lemmas that are stated and proved in Appendix E.

**Proposition 5.3.** Suppose  $\hat{P}_n = \mathbb{P}_n$  is the empirical bootstrap,  $\eta_0$  is uniformly bounded and m-times continuously differentiable with  $\int [D^{\alpha}\eta_0(x)]^2 dx < \infty$  for all  $|\alpha| = m$ , both  $\eta_n$  and  $\eta_n^*$  are KDEs with common symmetric mth order kernel function and common bandwidth h such that  $n^{-1/d} \prec h$ . If  $h \prec n^{-1/(4m)}$ , then bootstrap percentile intervals based on  $\psi_{n,1}^*$  are asymptotically valid. If  $h \prec n^{-1/(2m)}$ , then bootstrap percentile intervals based on  $\psi_{n,2}^*$  and  $\psi_{n,3}^*$  are asymptotically valid.

**Proof of Proposition 5.3.** Since  $\phi_P = 2(\eta_P - \psi_P)$ , we then have  $S_n^* = (\mathbb{P}_n^* - \mathbb{P}_n)(\phi_n^* - \phi_0) = 2(\mathbb{P}_n^* - \mathbb{P}_n)(\eta_n^* - \eta_0)$  and  $S_n = 2(\mathbb{P}_n - P_0)(\eta_n - \eta_0)$ . By Lemma E.7, we have  $S_n^* - S_n = o_{P_W^*}(n^{-1/2})$  under the stated assumptions.

We have  $R_n^* = \psi_n^* - \psi_n - (\mathbb{P}_n^* - \mathbb{P}_n)\phi_n^* = \psi_n^* - \psi_n - 2(\mathbb{P}_n^* - \mathbb{P}_n)\eta_n^*$ . For the one-step estimator  $\psi_{n,1}^*$ ,

we have  $R_n^* = -\int (\eta_n^* - \eta_0)^2 + \int (\eta_n - \eta_0)^2 + (\mathbb{P}_n - P_0)(\phi_n^* - \phi_n)$  and  $R_n = -\int (\eta_n - \eta_0)^2$ . Hence

$$R_n^* - R_n = -\int (\eta_n^* - \eta_0)^2 + 2\int (\eta_n - \eta_0)^2 + (\mathbb{P}_n - P_0)(\phi_n^* - \phi_n).$$

If  $nh^{4m} \to 0$  and  $nh^d \to \infty$  hold, then  $\int (\eta_n^* - \eta_0)^2 - 2 \int (\eta_n - \eta_0)^2 = o_{P_W^*}(n^{-1/2})$  by Lemma E.8, and  $(\mathbb{P}_n - P_0)(\phi_n^* - \phi_n) = o_{P_W^*}(n^{-1/2})$  by Lemma E.7. This implies bootstrap percentile confidence intervals based on  $\psi_{n,1}^*$  are asymptotically valid by Theorem 3.9.

For the plug-in estimator  $\psi_{n,2}^*$ , we have  $R_n^* = -\int (\eta_n^* - \eta_0)^2 + \int (\eta_n - \eta_0)^2 - \mathbb{P}_n^* \phi_n^* + \mathbb{P}_n \phi_n + (\mathbb{P}_n - P_0)(\phi_n^* - \phi_n)$  and  $R_n = -\int (\eta_n - \eta_0)^2 - \mathbb{P}_n \phi_n$ . Hence,

$$R_n^* - R_n = -\int (\eta_n^* - \eta_0)^2 + 2\int (\eta_n - \eta_0)^2 - \mathbb{P}_n^* \phi_n^* + 2\mathbb{P}_n \phi_n + (\mathbb{P}_n - P_0)(\phi_n^* - \phi_n).$$

Since  $nh^{2m} \to 0$  and  $nh^d \to \infty$  hold, we have  $\int (\eta_n^* - \eta_0)^2 - 2 \int (\eta_n - \eta_0)^2 = o_{P_W^*}(n^{-1/2})$  by Lemma E.8,  $\mathbb{P}_n^* \phi_n^* - 2\mathbb{P}_n \phi_n = o_{P_W^*}(n^{-1/2})$  by Lemma E.9 and  $(\mathbb{P}_n - P_0)(\phi_n^* - \phi_n) = o_{P_W^*}(n^{-1/2})$  by Lemma E.7. This implies bootstrap percentile confidence intervals based on  $\psi_{n,2}^*$  are asymptotically valid by Theorem 3.9.

For the empirical mean plug-in estimator  $\psi_{n,3}^*$ , we have  $R_n^* = -\int (\eta_n^* - \eta_0)^2 + \int (\eta_n - \eta_0)^2 - \mathbb{P}_n^* \phi_n^*/2 + \mathbb{P}_n \phi_n/2 + (\mathbb{P}_n - P_0)(\phi_n^* - \phi_n)$  and  $R_n = -\int (\eta_n - \eta_0)^2 - \mathbb{P}_n \phi_n/2$ . Hence,

$$R_n^* - R_n = -\int (\eta_n^* - \eta_0)^2 + 2\int (\eta_n - \eta_0)^2 - \mathbb{P}_n^* \phi_n^* / 2 + \mathbb{P}_n \phi_n + (\mathbb{P}_n - P_0)(\phi_n^* - \phi_n).$$

Since  $nh^{2m} \to 0$  and  $nh^d \to \infty$  hold, we have  $\int (\eta_n^* - \eta_0)^2 = 2 \int (\eta_n - \eta_0)^2 + o_{P_W^*}(n^{-1/2})$  by Lemma E.8,  $\mathbb{P}_n^* \phi_n^* = 2\mathbb{P}_n \phi_n + o_{P_W^*}(n^{-1/2})$  by Lemma E.9 and  $(\mathbb{P}_n - P_0)(\phi_n^* - \phi_n) = o_{P_W^*}(n^{-1/2})$  by Lemma E.7. This implies bootstrap percentile confidence intervals based on  $\psi_{n,3}^*$  are asymptotically valid by Theorem 3.9.

**Proposition 5.4.** Suppose  $\hat{P}_n$  possesses Lebesgue density function  $\hat{\eta}_n$  and the assumptions of Proposition 3.7 and Proposition 5.1 hold. For  $\mathcal{F}$  and M defined in Proposition 5.1, assume that  $\mathcal{F}$  is a  $\mathcal{M}$ uniform Donsker class such that  $P_W^*(\eta_n^* \in \mathcal{F}) \xrightarrow{P_0^*} 1$  and  $P_W^*(\|\hat{\eta}_n\|_{\infty} \ge M) = o_{P_0^*}(1)$ . If  $\|\eta_n^* - \hat{\eta}_n\|_{L_2(\lambda)} = o_{P_W^*}(n^{-1/4})$  and  $\|\hat{\eta}_n - \eta_n\|_{L_2(\lambda)} = o_{P_0^*}(n^{-1/4})$ , then  $\psi_{n,1}^* = \psi_{n,1} + (\mathbb{P}_n^* - \hat{P}_n)\phi_0 + o_{P_0^*}(n^{-1/2})$ . If in addition  $\int \eta_n^{*2} - \mathbb{P}_n^* \eta_n^* = o_{P_W^*}(n^{-1/2}) \text{ and } \int \eta_n^2 - \hat{P}_n \eta_n = o_{P_0^*}(n^{-1/2}), \text{ then } \psi_{n,2}^* = \psi_{n,2} + (\mathbb{P}_n^* - \hat{P}_n)\phi_0 + o_{P_0^*}(n^{-1/2})$ and  $\psi_{n,3}^* = \psi_{n,3} + n^{1/2}(\mathbb{P}_n^* - \hat{P}_n)\phi_0 + o_{P_0^*}(n^{-1/2}).$ 

**Proof of Proposition 5.4.** We use Theorem 3.4. Condition (B1) holds because  $P_W(\eta_n^* \in \mathcal{F}) \xrightarrow{P_0^*} 1$ 

and the conditions of Lemma 3.7 hold by assumption. For any  $\varepsilon > 0$ , by the triangle and Cauchy–Schwarz inequalities, we have

$$\begin{split} P_W^* \left( \|\phi_n^* - \phi_0\|_{L_2(P_0)} \ge \varepsilon \right) \\ &\leq P_W^* \left( 2 \|\eta_n^* - \eta_0\|_{L_2(P_0)} \ge \varepsilon/2 \right) + P_W^* \left( 2 \left| \int \eta_n^{*2} - \int \eta_0^2 \right| \ge \varepsilon/2 \right) \\ &\leq P_W^* \left( 2M \|\eta_n^* - \eta_0\|_{L_2(\lambda)} \ge \varepsilon/2 \right) + P_W^* \left( 2 \|\eta_n^* - \eta_0\|_{L_2(\lambda)} \|\eta_n^* + \eta_0\|_{L_2(\lambda)} \ge \varepsilon/2 \right) \\ &\leq P_W^* \left( 2M \|\eta_n^* - \eta_0\|_{L_2(\lambda)} \ge \varepsilon/2 \right) + P_W^* \left( 4M \|\eta_n^* - \eta_0\|_{L_2(\lambda)} \ge \varepsilon/2 \right) + P_W^* \left( \|\eta_n^*\|_{L_2(\lambda)} \ge M \right). \end{split}$$

Since  $\|\eta_n^* - \eta_0\|_{L_2(\lambda)} \leq \|\eta_n^* - \hat{\eta}_n\|_{L_2(\lambda)} + \|\hat{\eta}_n - \eta_0\|_{L_2(\lambda)}$  and  $\|\eta_n^*\|_{L_2(\lambda)} \leq \|\eta_n^*\|_{\infty}$ , each term on the right-hand side of the previous display is  $o_{P_0^*}(1)$ , which implies condition (B2).

For condition (B3), we have

$$T_1(\eta_n^*, \mathbb{P}_n^*) - T_1(\eta_n, \hat{P}_n) - (\mathbb{P}_n^* - \hat{P}_n)\phi_n^* = -\int (\eta_n^* - \hat{\eta}_n)^2 + \int (\hat{\eta}_n - \eta_n)^2,$$

which is  $o_{P_W^*}(n^{-1/2})$  by assumption. Hence, condition (B3) holds for  $\psi_{n,1}^*$ . We next have

$$\begin{split} T_2(\eta_n^*, \mathbb{P}_n^*) - T_2(\eta_n, \hat{P}_n) - (\mathbb{P}_n^* - \hat{P}_n)\phi_n^* &= \int \eta_n^{*2} - \int \eta_n^2 - 2(\mathbb{P}_n^* - \hat{P}_n)\eta_n^* \\ &= -\int (\eta_n^* - \hat{\eta}_n)^2 + \int (\hat{\eta}_n - \eta_n)^2 + 2\left[\int \eta_n^{*2} - \mathbb{P}_n^* \eta_n^*\right] \\ &- 2\left[\int \eta_n^2 - \hat{P}_n \eta_n\right], \\ T_3(\eta_n^*, \mathbb{P}_n^*) - T_3(\eta_n, \hat{P}_n) - (\mathbb{P}_n^* - \hat{P}_n)\phi_n^* &= \mathbb{P}_n^* \eta_n^* - \hat{P}_n \eta_n - 2(\mathbb{P}_n^* - \hat{P}_n)\eta_n^* \\ &= -\int (\eta_n^* - \hat{\eta}_n)^2 + \int (\hat{\eta}_n - \eta_n)^2 + \left[\int \eta_n^{*2} - \mathbb{P}_n^* \eta_n^*\right] \\ &- \left[\int \eta_n^2 - \hat{P}_n \eta_n\right], \end{split}$$

both of which are  $o_{P_W^*}(n^{-1/2})$  by assumption. Hence, condition (B3) holds for  $\psi_{n,2}^*$  and  $\psi_{n,3}^*$ .

The proof of Proposition 5.5 relies on technical lemmas that are stated and proved in Appendix F.

**Proposition 5.5.** Suppose that  $\eta_0$  is uniformly bounded and m-times continuously differentiable, and for all  $|\alpha| = m$ ,  $D^{\alpha}\eta_0$  is uniformly bounded and  $\int [D^{\alpha}\eta_0(x)]^2 dx < \infty$ . If both  $\eta_n$  and  $\eta_n^*$  are kernel density estimators with common uniformly bounded symmetric mth order kernel function K and common bandwidth h,  $\hat{P}_n$  is the distribution corresponding to  $\eta_n$ , and  $n^{-1/(2d)} \prec h \prec n^{-1/(4m)}$ , then bootstrap percentile intervals based on  $\psi_{n,1}^*$ ,  $\psi_{n,2}^*$ , and  $\psi_{n,3}^*$  are asymptotically valid.

**Proof of Proposition 5.5.** Since  $\phi_P = 2(\eta_P - \psi_P)$ , we then have  $S_n^* = (\mathbb{P}_n^* - \hat{P}_n)(\phi_n^* - \phi_0) = 2(\mathbb{P}_n^* - \hat{P}_n)(\eta_n^* - \eta_0)$  and  $S_n = 2(\mathbb{P}_n - P_0)(\eta_n - \eta_0)$ . By Lemma F.8, we have  $S_n^* - S_n = o_{P_W^*}(n^{-1/2})$  under the stated assumptions.

We have  $R_n^* = \psi_n^* - T(\eta_n, \hat{P}_n) - (\mathbb{P}_n^* - \hat{P}_n)\phi_n^* = \psi_n^* - T(\eta_n, \hat{P}_n) - 2(\mathbb{P}_n^* - \hat{P}_n)\eta_n^*$ . For the one-step estimator  $\psi_{n,1}^*$ , we have  $R_n^* = -\int (\eta_n^* - \eta_n)^2$  and  $R_n = -\int (\eta_n - \eta_0)^2$ . Hence

$$R_n^* - R_n = -\int (\eta_n^* - \eta_n)^2 + \int (\eta_n - \eta_0)^2$$

If  $nh^{4m} \longrightarrow 0$  and  $nh^{2d} \longrightarrow \infty$  hold, then  $\int (\eta_n^* - \eta_n)^2 - \int (\eta_n - \eta_0)^2 = o_{P_W^*}(n^{-1/2})$  by Lemma F.9. This implies bootstrap percentile confidence intervals based on  $\psi_{n,1}^*$  are asymptotically valid by Theorem 3.9.

For the plug-in estimator  $\psi_{n,2}^*$ , we have  $R_n^* = -\int (\eta_n^* - \eta_n)^2 - \mathbb{P}_n^* \phi_n^*$  and  $R_n = -\int (\eta_n - \eta_0)^2 - \mathbb{P}_n \phi_n$ . Hence,

$$R_{n}^{*} - R_{n} = -\int (\eta_{n}^{*} - \eta_{n})^{2} + \int (\eta_{n} - \eta_{0})^{2} - \mathbb{P}_{n}^{*} \phi_{n}^{*} + \mathbb{P}_{n} \phi_{n}.$$

Since  $nh^{4m} \longrightarrow 0$  and  $nh^{2d} \longrightarrow \infty$  hold, we have  $\int (\eta_n^* - \eta_n)^2 - \int (\eta_n - \eta_0)^2 = o_{P_W^*}(n^{-1/2})$  by Lemma F.9,  $\mathbb{P}_n^* \phi_n^* - \mathbb{P}_n \phi_n = o_{P_W^*}(n^{-1/2})$  by Lemma F.10. This implies bootstrap percentile confidence intervals based on  $\psi_{n,2}^*$  are asymptotically valid by Theorem 3.9.

For the empirical mean plug-in estimator  $\psi_{n,3}^*$ , we have  $R_n^* = -\int (\eta_n^* - \eta_n)^2 - \mathbb{P}_n^* \phi_n^*/2$  and  $R_n = -\int (\eta_n - \eta_0)^2 - \mathbb{P}_n \phi_n/2$ . Hence,

$$R_n^* - R_n = -\int (\eta_n^* - \eta_n)^2 + \int (\eta_n - \eta_0)^2 - \mathbb{P}_n^* \phi_n^* / 2 + \mathbb{P}_n \phi_n / 2.$$

Since  $nh^{4m} \longrightarrow 0$  and  $nh^{2d} \longrightarrow \infty$  hold, we have  $\int (\eta_n^* - \eta_n)^2 - \int (\eta_n - \eta_0)^2 = o_{P_W^*}(n^{-1/2})$  by Lemma F.9,  $\mathbb{P}_n^* \phi_n^* - \mathbb{P}_n \phi_n = o_{P_W^*}(n^{-1/2})$  by Lemma F.10. This implies bootstrap percentile confidence intervals based on  $\psi_{n,3}^*$  are asymptotically valid by Theorem 3.9.

Proposition 5.6. If  $\mu_n$ ,  $g_n$ , and  $(y, a, z) \mapsto (1 - a)yg_n(z)/[1 - g_n(z)]$  fall in  $P_0$ -Donsker classes with probability tending to 1,  $E_0(Y^2) < \infty$ , there exists constants 0 < a < b < 1 such that  $P_0(g_0(Z) \in (a, b)) = 1$ ,  $P_0(g_n(Z) \in (a, b)) = 1$ ,  $\|g_n - g_0\|_{L_2(P_0)} = o_{P_0^*}(1)$ ,  $\|\mu_n - \mu_0\|_{L_2(P_0)} = o_{P_0^*}(1)$ , and  $P_0\{(g_n - g_0)(\mu_n - \mu_0)/(1 - g_n)\} = o_{P_0^*}(n^{-1/2})$ , then  $\psi_{n,2}$  is asymptotically linear with influence function  $\phi_0$ . If in addition  $(\pi_n - \pi_0)(\psi_n - \psi_0) = o_{P_0^*}(n^{-1/2})$ , then  $\psi_{n,1}$  is asymptotically linear with influence function  $\phi_0$ .

**Proof of Proposition 5.6.** Condition (A1) holds by the Donsker assumptions, the boundedness assumptions on  $\mu_n$  and  $g_n$ , and  $E(Y^2) < \infty$  together with preservation of the Donsker condition under Lipschitz transformations (e.g., Theorem 2.10.6 of van der Vaart & Wellner, 1996).

We next address condition (A2). By adding and subtracting terms, we have

$$\begin{split} \phi_n(y,a,z) - \phi_0(y,a,z) &= (1-a) \left[ y - \mu_0(z) \right] \frac{g_n(z)}{1 - g_n(z)} \frac{1}{\pi_n \pi_0} \left[ \pi_0 - \pi_n \right] + a \left[ \mu_0(z) - \psi_0 \right] \frac{1}{\pi_n \pi_0} \left[ \pi_0 - \pi_n \right] \\ &+ \frac{(1-a) \left[ y - \mu_0(z) \right]}{\pi_0 \left[ 1 - g_n(z) \right] \left[ 1 - g_0(z) \right]} \left[ g_n(z) - g_0(z) \right] \\ &+ \frac{1}{\pi_n} \left[ \frac{a}{g_n(z)} - 1 \right] \frac{g_n(z)}{1 - g_n(z)} \left[ \mu_n(z) - \mu_0(z) \right] - \frac{a}{\pi_n} \left[ \psi_n - \psi_0 \right]. \end{split}$$

Therefore, by the assumed bounds and the triangle inequality,  $\|\phi_n - \phi_0\|_{L_2(P_0)}$  is bounded up to a constant by

$$|\pi_n - \pi_0| + ||g_n - g_0||_{L_2(P_0)} + ||\mu_n - \mu_0||_{L_2(P_0)} + |\psi_n - \psi_0|.$$

We note that consistency of  $\mu_n$  and  $g_n$  implies consistency of  $\pi_n$  and  $\psi_n$ . Therefore, condition (A2) holds.

For condition (A3) for the one-step estimator  $\psi_{n,1} = T_1(\eta_n, \mathbb{P}_n)$ , we have  $T_1(\eta_n, \mathbb{P}_n) - T_1(\eta_0, P_0) - (\mathbb{P}_n - P_0)\phi_n = \psi_n - \psi_0 + P_0\phi_n$ . Using the law of total expectation, a straightforward calculation shows that

$$P_0\phi_{\psi,\eta} = P_0\left\{\frac{(g_0-g)(\mu-\mu_0)}{\pi(1-g)}\right\} + \frac{\pi_0}{\pi}(\psi_0-\psi).$$

Therefore,

$$\psi_n - \psi_0 + P_0 \phi_n = P_0 \left\{ \frac{(g_0 - g_n)(\mu_n - \mu_0)}{\pi_n (1 - g_n)} \right\} + \frac{\pi_n - \pi_0}{\pi_n} (\psi_n - \psi_0).$$

These terms are both  $o_{P_0^*}(n^{-1/2})$  by the assumed rates and boundedness conditions. The result for the one-step estimator follows.

For the estimating equations-based estimator, we use Lemma 4.2. We have already established that

condition (A1) holds for  $\phi_n = \phi_{\psi_n,\eta_n}$ . We have that the population estimating equation is  $G_{0,\eta_0}(\psi) = \psi - \psi_0$ , so  $G_{0,\eta_0}(\psi_0) = 0$  and  $\inf_{|\psi-\psi_0|>\delta} |G_{0,\eta_0}(\psi)| = \delta$ , and hence  $\psi_0$  is a well-separated solution of the population estimating equation. Furthermore, it is straightforward to see that the boundedness conditions and the assumption that  $E_0(Y^2) < \infty$  imply that  $\psi_{n,2} = O_{P_0^*}(1)$ . Next, defining  $G'_{0,\eta}(\psi_0) = -\pi_0\pi^{-1}$ , we then have  $G'_{0,\eta_0}(\psi_0) = -1$  and  $\lim_{\eta\to\eta_0} G'_{0,\eta}(\psi_0) = \lim_{\eta\to\eta_0} -\pi_0\pi^{-1} = -1$ . In addition,

$$\begin{split} \Gamma_{0,\eta}(\psi) &= G_{0,\eta}(\psi) - G_{0,\eta}(\psi_0) - G'_{0,\eta}(\psi_0)(\psi - \psi_0) \\ &= P_0 \phi_{\eta,\psi} - P_0 \phi_{\eta,\psi_0} + \frac{\pi_0}{\pi}(\psi - \psi_0) \\ &= P_0 \left\{ \frac{g_0(Z) - g(Z)}{\pi \left[1 - g(Z)\right]} \left[ \mu(Z) - \mu_0(Z) \right] \right\} + \frac{\pi_0}{\pi}(\psi - \psi_0) \\ &- P_0 \left\{ \frac{g_0(Z) - g(Z)}{\pi \left[1 - g(Z)\right]} \left[ \mu(Z) - \mu_0(Z) \right] \right\} - \frac{\pi_0}{\pi}(\psi_0 - \psi_0) \\ &+ \frac{\pi_0}{\pi}(\psi_0 - \psi) \\ &= 0. \end{split}$$

Therefore, both conditions about  $\Gamma_{0,\eta}$  hold. Finally,

$$P_0 \phi_{\psi_0,\eta_n} = P_0 \left\{ \frac{g_0 - g_n}{\pi_n [1 - g_n]} [\mu_n - \mu_0] \right\} - \frac{\pi_0}{\pi_n} (\psi_0 - \psi_0)$$
$$= P_0 \left\{ \frac{g_0 - g_n}{\pi_n [1 - g_n]} [\mu_n - \mu_0] \right\},$$

which is  $o_{P_0^*}(n^{-1/2})$  by assumption. Hence, condition (A3) holds for the estimating equations-based estimator.

**Proposition 5.7.** Suppose  $\hat{P}_n = \mathbb{P}_n$  is the empirical bootstrap and the conditions of Proposition 5.6 hold. If  $\mu_n^*$ ,  $g_n^*$ , and  $(y, a, z) \mapsto (1-a)yg_n^*(z)/[1-g_n^*(z)]$  fall in  $P_0$ -Donsker classes with conditional probability tending to 1, there exist constants 0 < a < b < 1 such that  $P_W^*(g_n^*(Z) \in (a, b)) = 1$ , and  $P_W^*(|\mu_n^*(Z)| \le b) = 1$ ,  $||g_n^* - g_0||_{L_2(P_0)} = o_{P_W^*}(1)$ ,  $||\mu_n^* - \mu_0||_{L_2(P_0)} = o_{P_W^*}(1)$ , and  $P_0\{(g_n^* - g_0)(\mu_n^* - \mu_0)/(1-g_n^*)\} = o_{P_W^*}(n^{-1/2})$ , then  $\psi_{n,2}^*$  is conditionally asymptotically linear with influence function  $\phi_0$ . If in addition  $(\pi_n^* - \pi_0)(\psi_n^* - \psi_0) = o_{P_W^*}(n^{-1/2})$ , then  $\psi_{n,1}^*$  is conditionally asymptotically linear with influence function  $\phi_0$ .

**Proof of Proposition 5.7.** By Lemma 3.6, condition (B1) holds by the Donsker assumptions, the boundedness assumptions on  $\mu_n^*$  and  $g_n^*$ , and  $E(Y^2) < \infty$  together with preservation of the Donsker

condition under Lipschitz transformations.

As in the proof of Proposition 5.6, we can show that  $\|\phi_n^* - \phi_0\|_{L_2(P_0)}$  is bounded up to a constant by

$$|\pi_n^* - \pi_0| + ||g_n^* - g_0||_{L_2(P_0)} + ||\mu_n^* - \mu_0||_{L_2(P_0)} + |\psi_n^* - \psi_0|.$$

Therefore, condition (B2) holds under the conditional weak  $L_2(P_0)$  conditional consistency of  $g_n^*$  and  $\mu_n^*$ , which also implies weak conditional consistency of  $\psi_n^*$  and  $\pi_n^*$ .

For condition (B3) for the one-step estimator  $\psi_{n,1}^* = T_1(\eta_n^*, \mathbb{P}_n^*)$ , as in equation (8), we have

$$R_n^* = [\psi_n^* - \psi_0 + P_0\phi_n^*] - [\psi_n - \psi_0 + P_0\phi_n] + (\mathbb{P}_n - P_0)(\phi_n^* - \phi_n)$$

By the proof of Proposition 5.6,  $\psi_n - \psi_0 + P_0\phi_n = o_{P_0^*}(n^{-1/2})$ , and

$$n^{1/2}(\mathbb{P}_n - P_0)(\phi_n^* - \phi_n) = \mathbb{G}_n(\phi_n^* - \phi_0) - \mathbb{G}_n(\phi_n - \phi_0),$$

which is  $o_{P_W^*}(n^{-1/2})$  by Proposition 5.6 and the above. By the derivations in Proposition 5.6, we also have

$$\psi_n^* - \psi_0 + P_0 \phi_n^* = P_0 \left\{ \frac{(g_0 - g_n^*) (\mu_n^* - \mu_0)}{\pi_n^* (1 - g_n^*)} \right\} + \frac{\pi_n^* - \pi_0}{\pi_n^*} (\psi_n^* - \psi_0) + \frac{\pi_0}{\pi_n^*} (\psi_n^* - \psi_0$$

This is  $o_{P_W^*}(n^{-1/2})$  by assumption.

For condition (B3) for the estimating equations-based estimator  $\psi_{n,2}^*$ , we use Lemma 4.3. We note that the conditions of Lemma 4.2 hold by Proposition 5.6 and condition (B1) holds by the above. We have  $\psi_{n,2}^* = O_{P_W^*}(1)$  by the boundedness conditions and because  $E_0(Y^2) < \infty$ . As in the proof of Proposition 5.6, we have  $\Gamma_{0,\eta}(\psi) = 0$  for all  $\eta \in \mathcal{H}$  and  $\psi \in \mathbb{R}$ , so that  $\sup_{|\psi| \leq M} |\Gamma_{0,\eta_n^*}(\psi) - \Gamma_{0,\eta_0}(\psi_0)| = 0$  for every M > 0. Finally,

$$P_0\phi_{\psi_0,\eta_n^*} = P_0\left\{\frac{(g_0 - g_n^*)(\mu_n - \mu_0)}{\pi_n^*(1 - g_n^*)}\right\}.$$

which is  $o_{P_W^*}(n^{-1/2})$  by assumption.

**Proposition 5.8.** Suppose  $P_0^*(\hat{P}_n \in \mathfrak{P}) \to 1$ , where  $\mathfrak{P}$  is such that  $\lim_{M\to\infty} \sup_{P\in\mathfrak{P}} E_P[Y^2I(Y^2 > M)] = 0$ ,  $\hat{\mu}_n \in \mathfrak{F}_\mu$  and  $\hat{g}_n \in \mathfrak{F}_g$  with probability tending to one, where  $\mathfrak{F}_\mu$  is uniformly bounded,  $\mathfrak{F}_g$ 

is uniformly bounded away from zero, and  $\mathcal{F}_{\mu}$  and  $\mathcal{F}_{g}$  possess finite uniform entropy integrals,  $\|\hat{g}_{n} - g_{0}\|_{L_{2}(P_{0})}$ ,  $\|\hat{\mu}_{n} - \mu_{0}\|_{L_{2}(P_{0})}$ ,  $\|\hat{\sigma}_{n}^{2} - \sigma_{0}^{2}\|_{L_{2}(P_{0})}$  are each  $o_{P_{0}^{*}}(1)$ , and each of the following is  $o_{P_{0}^{*}}(1)$ :

$$\sup_{g,\bar{g}} \left| (\hat{Q}_n - Q_0) \left[ \frac{g\bar{g}(1 - \hat{g}_n)}{(1 - g)(1 - \bar{g})} (\hat{\sigma}_n^2 + \hat{\mu}_n^2) \right] \right|, \quad \sup_{g,\bar{g},\mu,\bar{\mu}} \left| (\hat{Q}_n - Q_0) \left[ \frac{g\bar{g}(1 - \hat{g}_n)}{(1 - g)(1 - \bar{g})} \mu \bar{\mu} \right] \right|, \quad (14)$$

$$\sup_{\mu} \left| (\hat{Q}_n - Q_0) \left[ \mu \hat{g}_n \right] \right|, \quad \sup_{\mu,\bar{\mu}} \left| (\hat{Q}_n - Q_0) \left[ \mu \bar{\mu} \hat{g}_n \right] \right|, \quad \sup_{\mu,g} \left| (\hat{Q}_n - Q_0) \left[ \frac{g(1 - \hat{g}_n)}{1 - g} \mu \right] \right|, \quad (\hat{Q}_n - Q_0) \hat{g}_n.$$

where the suprema over  $\mu$  and  $\bar{\mu}$  are taken over  $\mathcal{F}_{\mu}$  and the suprema over g and  $\bar{g}$  are taken over  $\mathcal{F}_{g}$ , and  $\hat{Q}_{n} \{(\hat{g}_{n} - g_{n})(\hat{\mu}_{n} - \mu_{n})/(1 - g_{n})\} = o_{P_{0}^{*}}(n^{-1/2})$ . Suppose also that  $\mu_{n}^{*} \in \mathcal{F}_{\mu}$  and  $g_{n}^{*} \in \mathcal{F}_{g}$  with conditional probability tending to one,  $\|g_{n}^{*} - g_{0}\|_{L_{2}(P_{0})} = o_{P_{W}^{*}}(1)$ ,  $\|\mu_{n}^{*} - \mu_{0}\|_{L_{2}(P_{0})} = o_{P_{W}^{*}}(1)$ , and  $\hat{Q}_{n} \{(g_{n}^{*} - \hat{g}_{n})(\mu_{n}^{*} - \hat{\mu}_{n})/(1 - g_{n}^{*})\} = o_{P_{W}^{*}}(n^{-1/2})$ . If  $(\pi_{n}^{*} - \hat{\pi}_{n})(\psi_{n}^{*} - \hat{\psi}_{n}) = o_{P_{W}^{*}}(n^{-1/2})$  and  $(\pi_{n} - \hat{\pi}_{n})(\psi_{n} - \hat{\psi}_{n}) = o_{P_{0}^{*}}(n^{-1/2})$ , then  $\psi_{n,1}^{*}$  is conditionally asymptotically linear with influence function  $\phi_{0}$ . If the conditions of Proposition 5.6 hold and  $\pi_{n}^{*} - \hat{\pi}_{n} = o_{P_{W}^{*}}(1)$ , then  $\psi_{n,2}^{*}$  is conditionmally asymptotically linear with influence function  $\phi_{0}$ .

**Proof of Proposition 5.8.** We use Lemma 3.7 to show condition (B1) holds. By the assumed classes of  $\mu_n^*$  and  $g_n^*$ , with conditional probability tending to 1,  $\phi_n^*$  falls in a class  $\mathcal{F}$  with envelope F(y, a, z) := c|Y| + d for some fixed  $c, d \in (0, \infty)$ . Thus, condition (i) of Lemma 3.7 holds by the assumption that  $\lim_{M\to\infty} \sup_{P\in\mathcal{P}} E_P[Y^2I(Y^2 > M)] = 0$ . Next, by preservation of the finite uniform entropy integral condition under Lipschitz transformations, (see, e.g., Lemma 9.17 of Kosorok, 2008),  $\mathcal{F}$  possesses finite uniform entropy integral. Hence, condition (ii) of Lemma 3.7 is satisfied by Theorem 2.8.3 of van der Vaart & Wellner (1996). Next, as noted following Lemma 3.7,  $\sup_{f,g\in\mathcal{F}} \left| (\hat{P}_n - P_0)(fg) \right| = o_{P_0^*}(1)$  implies (2). Let  $\mu, \bar{\mu} \in \mathcal{F}_{\mu}$  and  $g, \bar{g} \in \mathcal{F}_g$ , and let  $\pi, \bar{\pi}$  be implied by  $g, \bar{g}$  and  $\psi, \bar{\psi}$  be implied by  $\mu, g$  and  $\bar{\mu}, \bar{g}$ . Using the fact that a(1 - a) = 0 and the tower property, we then have

$$\begin{split} (\hat{P}_n - P_0) \left( \phi_{\mu,g} \phi_{\bar{\mu},\bar{g}} \right) \\ &= \int \left[ \frac{(1-a)g\bar{g}}{\pi\bar{\pi}(1-g)(1-\bar{g})} (y-\mu)(y-\bar{\mu}) + \frac{a}{\pi\bar{\pi}}(\mu-\psi)(\bar{\mu}-\bar{\psi}) \right] d(\hat{P}_n - P_0) \\ &= \int \left[ \frac{(1-a)g\bar{g}}{\pi\bar{\pi}(1-g)(1-\bar{g})} \left\{ y^2 - (\mu+\bar{\mu})y + \mu\bar{\mu} \right\} \right] d(\hat{P}_n - P_0) \\ &+ \int \left[ \frac{a}{\pi\bar{\pi}}(\mu-\psi)(\bar{\mu}-\bar{\psi}) \right] d(\hat{P}_n - P_0) \\ &= \frac{1}{\pi\bar{\pi}} \int \frac{g\bar{g}}{(1-g)(1-\bar{g})} \left\{ (1-\hat{g}_n) \left[ (\hat{\sigma}_n^2 + \hat{\mu}_n^2) - (\mu+\bar{\mu})\hat{\mu}_n + \mu\bar{\mu} \right] d\hat{Q}_n \\ &- (1-g_0) \left[ (\sigma_0^2 + \mu_0^2) - (\mu+\bar{\mu})\mu_0 + \mu\bar{\mu} \right] dQ_0 \right\} \end{split}$$

$$\begin{aligned} &+ \frac{1}{\pi\bar{\pi}} \int (\mu - \psi)(\bar{\mu} - \bar{\psi}) \left[ \hat{g}_n \, \mathrm{d}\hat{Q}_n - g_0 \, \mathrm{d}Q_0 \right] \\ &= \frac{1}{\pi\bar{\pi}} \int \frac{g\bar{g}}{(1 - g)(1 - \bar{g})} \left\{ (1 - \hat{g}_n) \left[ (\hat{\sigma}_n^2 + \hat{\mu}_n^2) - (\mu + \bar{\mu})\hat{\mu}_n + \mu\bar{\mu} \right] \right. \\ &- (1 - g_0) \left[ (\sigma_0^2 + \mu_0^2) - (\mu + \bar{\mu})\mu_0 + \mu\bar{\mu} \right] \right\} \, \mathrm{d}Q_0 \\ &+ \frac{1}{\pi\bar{\pi}} \int \frac{g\bar{g}(1 - \hat{g}_n)}{(1 - g)(1 - \bar{g})} \left[ (\hat{\sigma}_n^2 + \hat{\mu}_n^2) - (\mu + \bar{\mu})\hat{\mu}_n + \mu\bar{\mu} \right] \, \mathrm{d}(\hat{Q}_n - Q_0) \\ &+ \frac{1}{\pi\bar{\pi}} \int (\mu - \psi)(\bar{\mu} - \bar{\psi})(\hat{g}_n - g_0) \, \mathrm{d}Q_0 + \frac{1}{\pi\bar{\pi}} \int (\mu - \psi)(\bar{\mu} - \bar{\psi})\hat{g}_n \, \mathrm{d}(\hat{Q}_n - Q_0). \end{aligned}$$

By adding and subtracting terms and using the assumed bounds, we have

$$\begin{split} \sup_{\mu,\bar{\mu},g,\bar{g}} \left| (\hat{P}_n - P_0) \left( \phi_{\mu,g} \phi_{\bar{\mu},\bar{g}} \right) \right| &\lesssim \|\hat{g}_n - g_0\|_{L_2(Q_0)} + \|\hat{\mu}_n - \mu_0\|_{L_2(Q_0)} + \|\hat{\sigma}_n^2 - \sigma_0^2\|_{L_2(Q_0)} \\ &+ \sup_{g,\bar{g}} \left| \int \frac{g\bar{g}(1-\hat{g}_n)}{(1-g)(1-\bar{g})} (\hat{\sigma}_n^2 + \hat{\mu}_n^2) \operatorname{d}(\hat{Q}_n - Q_0) \right| \\ &+ \sup_{g,\bar{g},\mu} \left| \int \frac{g\bar{g}(1-\hat{g}_n)}{(1-g)(1-\bar{g})} \mu \hat{\mu}_n \operatorname{d}(\hat{Q}_n - Q_0) \right| \\ &+ \sup_{g,\bar{g},\mu,\bar{\mu}} \left| \int \frac{g\bar{g}(1-\hat{g}_n)}{(1-g)(1-\bar{g})} \mu \bar{\mu} \operatorname{d}(\hat{Q}_n - Q_0) \right| \\ &+ \sup_{\mu} \left| \int \mu \hat{g}_n \operatorname{d}(\hat{Q}_n - Q_0) \right| + \sup_{\mu,\bar{\mu}} \left| \int \mu \bar{\mu} \hat{g}_n \operatorname{d}(\hat{Q}_n - Q_0) \right|. \end{split}$$

Since  $\hat{\mu}_n \in \mathcal{F}_{\mu}$  with probability tending to one, by assumption, each of these terms is  $o_{P_0^*}(1)$ , which implies (2). Hence, the conditions of Lemma 3.7 hold, which implies condition (B1).

We next show condition (B2) holds. As in the proof of Proposition 5.6, we can show that  $\|\phi_n^* - \phi_0\|_{L_2(P_0)}$  is bounded up to a constant by

$$|\pi_n^* - \pi_0| + ||g_n^* - g_0||_{L_2(P_0)} + ||\mu_n^* - \mu_0||_{L_2(P_0)} + |\psi_n^* - \psi_0|.$$

Each of these terms is  $o_{P_W^*}(1)$ .

Finally, we turn to condition (B3). By the tower property, we can show that

$$P\phi_{\mu,g,\psi} = Q_P \left[ \frac{(g_P - g)(\mu - \mu_P)}{(1 - g)\pi} \right] + \frac{\pi_P}{\pi} \left[ \psi(P) - \psi \right]$$

for  $\pi := \int g \, dQ$ . Thus,

$$\psi(\eta) - \psi(P) + P\phi_{\eta} = Q_P \left[ \frac{(g_P - g)(\mu - \mu_P)}{(1 - g)\pi(\eta)} \right] + \frac{\pi(\eta) - \pi_P}{\pi(\eta)} \left[ \psi(\eta) - \psi(P) \right].$$

Hence, for the one-step estimator  $\psi_{n,1}^* = T_1(\eta_n^*, \mathbb{P}_n^*)$ , as in equation (8), we have

$$\begin{aligned} R_n^* &= \left[ \psi(\eta_n^*) - \psi(\hat{\eta}_n) + \hat{P}_n \phi_n^* \right] - \left[ \psi(\eta_n) - \psi(\hat{\eta}_n) + \hat{P}_n \phi_n \right] \\ &= \left[ \hat{Q}_n \left\{ \frac{(\hat{g}_n - g_n^*)(\mu_n^* - \hat{\mu}_n)}{\pi_n^* (1 - g_n^*)} \right\} + \frac{\pi_n^* - \hat{\pi}_n}{\pi_n^*} (\psi_n^* - \hat{\psi}_n) \right] \\ &- \left[ \hat{Q}_n \left\{ \frac{(\hat{g}_n - g_n)(\mu_n - \hat{\mu}_n)}{\pi_n (1 - g_n)} \right\} + \frac{\pi_n - \hat{\pi}_n}{\pi_n} (\psi_n - \hat{\psi}_n) \right]. \end{aligned}$$

Each term is  $o_{P_W^*}(n^{-1/2})$  by assumption. The result for the one-step estimator follows.

For condition (B3) for the estimating equations-based estimator  $\psi_{n,2}^*$ , we use Lemma 4.4. Since the conditions of Proposition 5.6 hold, the conditions of Lemma 4.2 hold as well. We have  $\psi_{n,2}^* = O_{P_W^*}(1)$  by the boundedness conditions on  $\mu_n^*$ ,  $g_n^*$ , and  $\hat{g}_n$ , and the moment condition for  $\hat{P}_n$ . A similar argument shows that  $\hat{\psi}_n = O_{P_0^*}(1)$ . As in the proof of Proposition 5.6, we have  $\Gamma_{0,\eta}(\psi) = 0$  for any  $\eta \in \mathcal{H}$  and  $\psi \in \mathbb{R}$ , so that the condition  $\sup_{|\psi| \leq M} |\Gamma_{0,\hat{\eta}_n}(\psi) - \Gamma_{0,\eta_0}(\psi)| = o_{P_0^*}(1)$  holds. Next, defining  $\hat{G}'_{n,\eta}(\hat{\psi}_n) = -\hat{\pi}_n \pi^{-1}$ , we then have  $\hat{G}'_{n,\hat{\eta}_n^*}(\hat{\psi}_n) + 1 = 1 - \hat{\pi}_n/\pi_n^* = o_{P_W^*}(1)$  because  $\pi_n^* - \hat{\pi}_n = o_{P_W^*}(1)$  and  $g_n^*$  is bounded away from zero. In addition,

$$\begin{split} \hat{\Gamma}_{n,\eta}(\psi) &= \hat{G}_{n,\eta}(\psi) - \hat{G}_{n,\eta}(\hat{\psi}_n) - \hat{G}'_{n,\eta}(\hat{\psi}_n)(\psi - \hat{\psi}_n) \\ &= \hat{P}_n \phi_{\eta,\psi} - \hat{P}_n \phi_{\eta,\hat{\psi}_n} + \frac{\hat{\pi}_n}{\pi}(\psi - \hat{\psi}_n) \\ &= \hat{P}_n \left\{ \frac{\hat{g}_n - g}{\pi \left[ 1 - g \right]} \left[ \mu - \hat{\mu}_n \right] \right\} + \frac{\hat{\pi}_n}{\pi}(\psi - \hat{\psi}_n) - \hat{P}_n \left\{ \frac{\hat{g}_n - g}{\pi \left[ 1 - g \right]} \left[ \mu - \hat{\mu}_n \right] \right\} - \frac{\hat{\pi}_n}{\pi}(\hat{\psi}_n - \hat{\psi}_n) \\ &+ \frac{\hat{\pi}_n}{\pi}(\hat{\psi}_n - \psi) \\ &= 0. \end{split}$$

Hence, the conditions about  $|\hat{\Gamma}_{n,\eta_n^*}(\psi) - \hat{\Gamma}_{n,\hat{\eta}_n}(\hat{\psi})|$  and  $|\hat{\Gamma}_{n,\eta}(\psi)|$  hold. Next,  $\phi_{\hat{\psi}_n,\hat{\eta}_n}$  and  $\phi_{\psi_n^*,\hat{\eta}_n^*}$  fall in a  $P_0$ -Glivenko Cantelli class with probability to tending one by the finite entropy integral and boundedness assumptions. Next, we verify that  $\|\hat{P}_n - P_0\|_{\mathcal{F}} = o_{P_0^*}(1)$  holds under the stated assumptions.

$$\int \phi_{\mu,g}(y,a,w) \,\mathrm{d}(\hat{P}_n - P_0) = \int \left[ \frac{(1-a)g}{\pi(1-g)} (y-\mu) + \frac{a}{\pi} (\mu-\psi) \right] \,\mathrm{d}(\hat{P}_n - P_0)$$

$$= \frac{1}{\pi} \int \frac{g}{(1-g)} \left\{ (1-\hat{g}_n)(\hat{\mu}_n - \mu) \,\mathrm{d}\hat{Q}_n - (1-g_0)(\mu_0 - \mu) \,\mathrm{d}Q_0 \right\}$$

$$+ \frac{1}{\pi} \int (\mu-\psi) \left[ \hat{g}_n \,\mathrm{d}\hat{Q}_n - g_0 \,\mathrm{d}Q_0 \right]$$

$$= \frac{1}{\pi} \int \frac{g}{(1-g)} \left\{ (1-\hat{g}_n)(\hat{\mu}_n - \mu) - (1-g_0)(\mu_0 - \mu) \right\} \,\mathrm{d}Q_0$$

$$+\frac{1}{\pi}\int \frac{g}{(1-g)}(1-\hat{g}_n)(\hat{\mu}_n-\mu) \,\mathrm{d}(\hat{Q}_n-Q_0) \\ +\frac{1}{\pi}\int (\mu-\psi)\left(\hat{g}_n-g_0\right) \,\mathrm{d}Q_0 +\frac{1}{\pi}\int (\mu-\psi)\hat{g}_n \,\mathrm{d}(\hat{Q}_n-Q_0).$$

Therefore, adding and subtracting terms and using assumed bounds,

$$\begin{split} \sup_{\mu,g} \left| \int \phi_{\mu,g} \,\mathrm{d}(\hat{P}_n - P_0) \right| &\lesssim \|\hat{g}_n - g_0\|_{L_2(Q_0)} + \|\hat{\mu}_n - \mu_0\|_{L_2(Q_0)} \\ &+ \sup_{\mu,g} \left| \int \frac{g}{(1-g)} (1-\hat{g}_n)(\hat{\mu}_n - \mu) \,\mathrm{d}(\hat{Q}_n - Q_0) \right| \\ &+ \sup_{\mu} \left| \int (\mu - \psi) \hat{g}_n \,\mathrm{d}(\hat{Q}_n - Q_0) \right|. \end{split}$$

Each of these terms is  $o_{P_0^*}(1)$  by assumption. Thus,  $\|\hat{P}_n - P_0\|_{\mathcal{F}} = o_{P_0^*}(1)$ .

Next, we have

$$P_0\phi_{\psi_0,\hat{\eta}_n} = P_0\left\{\frac{(g_0 - \hat{g}_n)(\hat{\mu}_n - \mu_0)}{\hat{\pi}_n(1 - \hat{g}_n)}\right\}.$$

which is  $o_{P_0^*}(1)$  by the assumed boundedness of  $\hat{g}_n$  away from zero and the consistency of  $\hat{\mu}_n$  and  $\hat{g}_n$ .

Next, we show that  $\psi_n^{\circ} - \hat{\psi}_n = o_{P_0^*}(n^{-1/2})$ . Use the definition of the estimating equations-based estimator construction  $T_2$ , we can see that

$$\begin{split} \psi_n^{\circ} &:= T_2(\eta_n, \hat{P}_n) = \hat{Q}_n \left\{ \frac{g_n \left(1 - \hat{g}_n\right)}{\hat{\pi}_n (1 - g_n)} \left( \hat{\mu}_n - \mu_n \right) + \frac{\hat{g}_n}{\hat{\pi}_n} \mu_n \right\}, \\ \hat{\psi}_n &:= T_2(\hat{\eta}_n, \hat{P}_n) = \hat{Q}_n \left\{ \frac{\hat{g}_n}{\hat{\pi}_n} \hat{\mu}_n \right\}, \\ \psi_n^{\circ} - \hat{\psi}_n &= \frac{1}{\hat{\pi}_n} \hat{Q}_n \left\{ \frac{g_n - \hat{g}_n}{1 - g_n} \left( \hat{\mu}_n - \mu_n \right) \right\}, \end{split}$$

which is  $o_{P_0^*}(n^{-1/2})$  by assumption. Finally,

$$\begin{split} \hat{P}_n \phi_{\hat{\psi}_n, \eta_n^*} &= \hat{Q}_n \left\{ \frac{\hat{g}_n - g_n^*}{\pi_n^* \left[ 1 - g_n^* \right]} \left[ \mu_n^* - \hat{\mu}_n \right] \right\} - \frac{\hat{\pi}_n}{\pi_n^*} (\hat{\psi}_n - \hat{\psi}_n) \\ &= \hat{Q}_n \left\{ \frac{\hat{g}_n - g_n^*}{\pi_n^* \left[ 1 - g_n^* \right]} \left[ \mu_n^* - \hat{\mu}_n \right] \right\}, \end{split}$$

which is  $o_{P_0^*}(n^{-1/2})$  by assumption as discussed above. Hence, condition (B3) holds for the estimating equations-based estimator.

## Supplement E Lemmas supporting the proof of Proposition 5.3

We first present several simple algebraic identities that we will use repeatedly.

**Lemma E.1.** For any  $\{Z_{ij} \in \mathbb{R} : i, j = 1, 2, ..., n\}$  such that  $Z_{ij} = Z_{ji}$  and  $Z_{ii} = \tau_f$  for all i, j, we have

$$\left[\sum_{i\neq j} Z_{ij}\right]^2 = 4 \sum_{i\neq j, j\neq k, k\neq i} Z_{ij} Z_{ik} + 2 \sum_{i\neq j} Z_{ij}^2 + \sum_{\substack{i\neq j, i\neq k, i\neq l\\ j\neq k, j\neq l, k\neq l}} Z_{ij} Z_{kl}, \text{ and}$$
(23)

$$\sum_{i,j,k} Z_{ij} Z_{ik} = 2\tau_f \sum_{i \neq j} Z_{ij} + \sum_{i \neq j} Z_{ij}^2 + n\tau_f^2 + \sum_{i \neq j, i \neq k, j \neq k} Z_{ij} Z_{ik}.$$
(24)

Proof of Lemma E.1. We have

$$\begin{bmatrix} \sum_{i \neq j} Z_{ij} \end{bmatrix}^{2} = \sum_{i \neq j, k \neq l} Z_{ij} Z_{kl}$$

$$= \sum_{\substack{i \neq j, k \neq l \\ i = k, j \neq l}} Z_{ij} Z_{kl} + \sum_{\substack{i \neq j, k \neq l \\ i = l, j \neq k}} Z_{ij} Z_{kl} + \sum_{\substack{i \neq j, k \neq l \\ i \neq l, j = k}} Z_{ij} Z_{kl} + \sum_{\substack{i \neq j, k \neq l \\ i \neq k, j = l}} Z_{ij} Z_{kl} + \sum_{\substack{i \neq j, k \neq l \\ i = l, j = k}} Z_{ij} Z_{kl} + \sum_{\substack{i \neq j, k \neq l \\ i = l, j = k}} Z_{ij} Z_{kl} + \sum_{\substack{i \neq j, i \neq k, i \neq l \\ j \neq k, j \neq l, k \neq l}} Z_{ij} Z_{kl}$$

$$= 4 \sum_{\substack{i \neq j, j \neq k, k \neq i}} Z_{ij} Z_{ik} + 2 \sum_{\substack{i \neq j}} Z_{ij}^{2} + \sum_{\substack{i \neq j, i \neq k, i \neq l \\ j \neq k, j \neq l, k \neq l}} Z_{ij} Z_{kl},$$

which proves (23). We also have

$$\begin{split} \sum_{i,j,k} Z_{ij} Z_{ik} &= \sum_{i=j \neq k} Z_{ij} Z_{ik} + \sum_{i=k \neq j} Z_{ij} Z_{ik} + \sum_{i \neq j=k} Z_{ij} Z_{ik} + \sum_{i=j=k} Z_{ij} Z_{ik} + \sum_{i \neq j, i \neq k, j \neq k} Z_{ij} Z_{ik} \\ &= 2\tau_f \sum_{i \neq j} Z_{ij} + \sum_{i \neq j} Z_{ij}^2 + n\tau_f^2 + \sum_{i \neq j, i \neq k, j \neq k} Z_{ij} Z_{ik}, \end{split}$$

which proves (24).

We next present several non-asymptotic bounds for empirical and bootstrap empirical V-processes for a generic function  $f : \mathbb{R}^d \times \mathbb{R}^d \mapsto \mathbb{R}$  satisfying f(x, y) = f(y, x) and  $f(x, x) = \tau_f \in \mathbb{R}$  for all  $x, y \in \mathbb{R}^d$ . Here, f may depend on n, but is assumed to be deterministic. We use the symbol  $\lesssim$  to mean "less than or equal up to a constant not depending on n." **Lemma E.2.** If  $f : \mathbb{R}^d \times \mathbb{R}^d \mapsto \mathbb{R}$  satisfies f(x, y) = f(y, x) and  $f(x, x) = \tau_f \in \mathbb{R}$  for all  $x, y \in \mathbb{R}^d$ , then

$$E_0 \left\{ \int f \, \mathrm{d}[(\mathbb{P}_n - P_0) \times (\mathbb{P}_n - P_0)] - n^{-1} \tau_f \right\}^2 \lesssim n^{-3} \int \left[ \int f(x, y) \, \mathrm{d}P_0(x) \right]^2 \, \mathrm{d}P_0(y) + n^{-2} \, \|f\|_{L_2(P_0 \times P_0)}^2 \, \mathrm{d}$$

**Proof of Lemma E.2.** For simplicity, we denote  $V_0(x, y) := \int f d [(\delta_x - P_0) \times (\delta_y - P_0)]$  for any  $x, y \in \mathbb{R}^d$ . By the law of total expectation, for any  $i \neq j$ , we note that

$$E_0[V_0(X_i, X_j) \mid X_j] = E_0 \left[ f(X_i, X_j) - \int f(X_i, y) \, \mathrm{d}P_0(y) - \int f(x, X_j) \, \mathrm{d}P_0(x) + \int f \, \mathrm{d}(P_0 \times P_0) \mid X_j \right]$$
  
=  $\int f(x, X_j) \, \mathrm{d}P_0(x) - \int f \, \mathrm{d}(P_0 \times P_0) - \int f(x, X_j) \, \mathrm{d}P_0(x) + \int f \, \mathrm{d}(P_0 \times P_0)$   
= 0

Hence,  $E_0[V_0(X_i, X_j)] = 0$  for  $i \neq j$  and  $E_0[V_0(X_i, X_j)V_0(X_i, X_k)] = 0$  for  $i \neq j \neq k$ .

By definition, we have

$$E_{0} \left\{ \int f d[(\mathbb{P}_{n} - P_{0}) \times (\mathbb{P}_{n} - P_{0})] - n^{-1} \tau_{f} \right\}^{2}$$

$$= E_{0} \left\{ \frac{1}{n^{2}} \sum_{i=j} V_{0}(X_{i}, X_{j}) + \frac{1}{n^{2}} \sum_{i \neq j} V_{0}(X_{i}, X_{j}) - n^{-1} \tau_{f} \right\}^{2}$$

$$\lesssim E_{0} \left\{ \frac{1}{n^{2}} \sum_{i=j} V_{0}(X_{i}, X_{j}) - n^{-1} \tau_{f} \right\}^{2} + E_{0} \left\{ \frac{1}{n^{2}} \sum_{i \neq j} V_{0}(X_{i}, X_{j}) \right\}^{2}.$$
(25)

For the first term on the right-hand side of (25), by the symmetry of f and the definition of  $\tau_f$ ,

$$E_{0}\left\{\frac{1}{n^{2}}\sum_{i=j}V_{0}(X_{i},X_{j})-\frac{\tau_{f}}{n}\right\}^{2} = E_{0}\left\{-\frac{2}{n}\int f\,\mathrm{d}(P_{0}\times\mathbb{P}_{n})+\frac{1}{n}\int f\,\mathrm{d}(P_{0}\times P_{0})\right\}^{2}$$

$$= E_{0}\left\{-\frac{2}{n}\int \left[\int f(x,y)\,\mathrm{d}P_{0}(x)\right]\,\mathrm{d}(\mathbb{P}_{n}-P_{0})(y)-\frac{1}{n}\int f\,\mathrm{d}(P_{0}\times P_{0})\right\}^{2}$$

$$\lesssim E_{0}\left\{\frac{1}{n}\int \left[\int f(x,y)\,\mathrm{d}P_{0}(x)\right]\,\mathrm{d}(\mathbb{P}_{n}-P_{0})(y)\right\}^{2}+\frac{1}{n^{2}}\|f\|_{L_{1}(P_{0}\times P_{0})}^{2}$$

$$=\frac{1}{n^{3}}\mathrm{Var}\left(\left[\int f(x,X_{1})\,\mathrm{d}P_{0}(x)\right]^{2}\,\mathrm{d}P_{0}(y)+\frac{1}{n^{2}}\|f\|_{L_{1}(P_{0}\times P_{0})}^{2}.$$

For the second term on the right-hand side of (25), by Lemma E.1 and properties of  $V_0$ ,

$$\begin{split} E_{0} \left[ \frac{1}{n^{2}} \sum_{i \neq j} V_{0}(X_{i}, X_{j}) \right]^{2} \\ &= E_{0} \left[ \frac{4}{n^{4}} \sum_{i \neq j, j \neq k, k \neq i} V_{0}(X_{i}, X_{j}) V_{0}(X_{i}, X_{k}) + \frac{2}{n^{4}} \sum_{i \neq j} V_{0}(X_{i}, X_{j})^{2} \right. \\ &\left. + \frac{1}{n^{4}} \sum_{\substack{i \neq j, i \neq k, i \neq l \\ j \neq k, j \neq l, k \neq l}} V_{0}(X_{i}, X_{j}) V_{0}(X_{k}, X_{l}) \right] \\ &= E_{0} \left[ \frac{2}{n^{4}} \sum_{i \neq j} V_{0}(X_{i}, X_{j})^{2} \right] \\ &\leq \frac{2}{n^{2}} E_{0} \left[ V_{0}(X_{1}, X_{2})^{2} \right] \\ &= \frac{2}{n^{2}} E_{0} \left[ f(X_{1}, X_{2}) - \int f(X_{1}, y) \, dP_{0}(y) - \int f(x, X_{2}) \, dP_{0}(x) + \int f \, d(P_{0} \times P_{0}) \right]^{2} \\ &= \frac{2}{n^{2}} E_{0} \left[ f(X_{1}, X_{2}) - \int f(X_{1}, y) \, dP_{0}(y) - \int f(x, X_{2}) \, dP_{0}(x) \right]^{2} - \frac{2}{n^{2}} \left[ \int f \, d(P_{0} \times P_{0}) \right]^{2} \\ &\leq \frac{2}{n^{2}} E_{0} \left[ f(X_{1}, X_{2}) - \int f(X_{1}, y) \, dP_{0}(y) - \int f(x, X_{2}) \, dP_{0}(x) \right]^{2} - \frac{2}{n^{2}} \left[ \int f \, d(P_{0} \times P_{0}) \right]^{2} \\ &\leq \frac{2}{n^{2}} E_{0} \left[ f(X_{1}, X_{2}) - \int f(X_{1}, y) \, dP_{0}(y) - \int f(x, X_{2}) \, dP_{0}(x) \right]^{2}. \end{split}$$

For the right hand side of previous display, we note that

$$E_{0} \left[ f(X_{1}, X_{2}) - \int f(X_{1}, y) \, \mathrm{d}P_{0}(y) - \int f(x, X_{2}) \, \mathrm{d}P_{0}(x) \right]^{2}$$

$$= E_{0}[f(X_{1}, X_{2})^{2}] + E_{0} \left[ \int f(X_{1}, y) \, \mathrm{d}P_{0}(y) \right]^{2} + E_{0} \left[ \int f(x, X_{2}) \, \mathrm{d}P_{0}(x) \right]^{2}$$

$$- 2E_{0} \left[ \int f(X_{1}, y) \, \mathrm{d}P_{0}(y) \right]^{2} - 2E_{0} \left[ \int f(x, X_{2}) \, \mathrm{d}P_{0}(x) \right]^{2}$$

$$+ 2E_{0} \left[ \int f(X_{1}, y) \, \mathrm{d}P_{0}(y) \right] \left[ \int f(x, X_{2}) \, \mathrm{d}P_{0}(x) \right]$$

$$= E_{0}[f(X_{1}, X_{2})^{2}] - E_{0} \left[ \int f(X_{1}, y) \, \mathrm{d}P_{0}(y) \right]^{2} - E_{0} \left[ \int f(x, X_{2}) \, \mathrm{d}P_{0}(x) \right]^{2} + 2[(P_{0} \times P_{0})f]^{2}$$

$$\leq E_{0}[f(X_{1}, X_{2})^{2}],$$

where the last inequality is because  $[(P_0 \times P_0)f]^2 \leq E_0 \left[\int f(X_1, y) dP_0(y)\right]^2$  and  $[(P_0 \times P_0)f]^2 \leq E_0 \left[\int f(x, X_2) dP_0(x)\right]^2$  by the Cauchy–Schwarz inequality. This implies  $E_0 \left[n^{-2} \sum_{i \neq j} V_0(X_i, X_j)\right]^2 \lesssim n^{-2} \|f\|_{L_2(P_0 \times P_0)}^2$ .

**Lemma E.3.** If  $f : \mathbb{R}^d \times \mathbb{R}^d \mapsto \mathbb{R}$  satisfies f(x, y) = f(y, x) and  $f(x, x) = \tau_f$  for all  $x, y \in \mathbb{R}^d$ , then for the empirical bootstrap  $\hat{P}_n = \mathbb{P}_n$ ,  $E_0 E_W \left[ \int f d[(\mathbb{P}_n - P_0) \times (\mathbb{P}_n^* - \mathbb{P}_n)] \right]^2 \lesssim n^{-3} \tau_f^2 + n^{-3} \tau_f ||f||_{L_1(P_0 \times P_0)} + n^{-2} ||f||_{L_2(P_0 \times P_0)}^2$ .

**Proof of Lemma E.3.** For simplicity, we denote  $g_n(y) := \int f(x,y) d(\mathbb{P}_n - P_0)(x)$ . We then have

$$\int f d[(\mathbb{P}_n - P_0) \times (\mathbb{P}_n^* - \mathbb{P}_n)] = (\mathbb{P}_n^* - \mathbb{P}_n)g_n = \frac{1}{n} \sum_{i=1}^n [g_n(X_i^*) - \mathbb{P}_n g_n]$$

Since  $X_1^*, \ldots, X_n^* \stackrel{\text{iid}}{\sim} \mathbb{P}_n, E_W[g_n(X_i^*) - \mathbb{P}_n g_n][g_n(X_j^*) - \mathbb{P}_n g_n] = 0$  for any  $i \neq j$ , so

$$E_{W}\left[\frac{1}{n}\sum_{i=1}^{n}[g_{n}(X_{i}^{*})-\mathbb{P}_{n}g_{n}]\right]^{2} = E_{W}\left[\frac{1}{n^{2}}\sum_{i,j}^{n}[g_{n}(X_{i}^{*})-\mathbb{P}_{n}g_{n}][g_{n}(X_{j}^{*})-\mathbb{P}_{n}g_{n}]\right]$$
$$= E_{W}\left[\frac{1}{n^{2}}\sum_{i=j}^{n}[g_{n}(X_{i}^{*})-\mathbb{P}_{n}g_{n}][g_{n}(X_{j}^{*})-\mathbb{P}_{n}g_{n}]\right]$$
$$= \frac{1}{n}E_{W}[g_{n}(X_{i}^{*})-\mathbb{P}_{n}g_{n}]^{2}$$
$$= \frac{1}{n}\left[\mathbb{P}_{n}g_{n}^{2}-(\mathbb{P}_{n}g_{n})^{2}\right]$$
$$\leq \frac{1}{n}\mathbb{P}_{n}g_{n}^{2}.$$

We then have

$$\mathbb{P}_{n}g_{n}^{2} = \frac{1}{n^{3}}\sum_{i,j,k} f(X_{i}, X_{j})f(X_{i}, X_{k}) - \frac{2}{n^{2}}\sum_{i,j} f(X_{i}, X_{j}) \left[ \int f(x, X_{j}) \,\mathrm{d}P_{0}(x) \right] \\ + \frac{1}{n}\sum_{i=1}^{n} \left[ \int f(x, X_{j}) \,\mathrm{d}P_{0}(x) \right]^{2}.$$
(26)

By Lemma E.1, the expectation of the first term on the right-hand side of (26) is

$$E_{0}\left[\frac{1}{n^{3}}\sum_{i,j,k}f(X_{i},X_{j})f(X_{i},X_{k})\right]$$

$$=E_{0}\left[\frac{2\tau_{f}}{n^{3}}\sum_{i\neq j}f(X_{i},X_{j})+\frac{1}{n^{3}}\sum_{i\neq j}f(X_{i},X_{j})^{2}+\frac{\tau_{f}^{2}}{n^{2}}+\frac{1}{n^{3}}\sum_{i\neq j,i\neq k,j\neq k}f(X_{i},X_{j})f(X_{i},X_{k})\right]$$

$$=\frac{2(n-1)\tau_{f}}{n^{2}}E_{0}\left[f(X_{1},X_{2})\right]+\frac{n-1}{n^{2}}E_{0}\left[f(X_{1},X_{2})^{2}\right]+\frac{\tau_{f}^{2}}{n^{2}}+\frac{(n-1)(n-2)}{n^{2}}E_{0}[f(X_{1},X_{2})f(X_{1},X_{3})].$$

The expectation of the second term on the right-hand side of (26) is

$$E_{0}\left\{\frac{2}{n^{2}}\sum_{i,j}f(X_{i},X_{j})\left[\int f(x,X_{j}) dP_{0}(x)\right]\right\}$$
  
=  $E_{0}\left\{\frac{2}{n^{2}}\sum_{i=j}f(X_{i},X_{j})\left[\int f(x,X_{j}) dP_{0}(x)\right]\right\} + E_{0}\left\{\frac{2}{n^{2}}\sum_{i\neq j}f(X_{i},X_{j})\left[\int f(x,X_{j}) dP_{0}(x)\right]\right\}$   
=  $\frac{2\tau_{f}}{n}E_{0}\left[f(X_{1},X_{2})\right] + \frac{2(n-1)}{n}E_{0}\left[f(X_{1},X_{2})f(X_{1},X_{3})\right].$ 

The expectation of the third term on the right-hand side of (26) is

$$E_0\left\{\frac{1}{n}\sum_{j=1}\left[\int f(x,X_j)\,\mathrm{d}P_0(x)\right]^2\right\} = E_0\left[f(X_1,X_2)f(X_1,X_3)\right].$$

Combining these calculations, we get

$$\begin{aligned} \left| \frac{1}{n} E_0 \left( \mathbb{P}_n g_n^2 \right) \right| &= \left| -\frac{2\tau_f}{n^3} E_0 \left[ f(X_1, X_2) \right] + \frac{n-1}{n^3} E_0 \left[ f(X_1, X_2)^2 \right] + \frac{\tau_f^2}{n^3} - \frac{n-2}{n^3} E_0 \left[ f(X_1, X_2) f(X_1, X_3) \right] \right. \\ &\leq \frac{2\tau_f}{n^3} E_0 |f(X_1, X_2)| + \frac{1}{n^2} E_0 \left[ f(X_1, X_2)^2 \right] + \frac{\tau_f^2}{n^3} + \frac{1}{n^2} E_0 |f(X_1, X_2) f(X_1, X_3)| \\ &\leq \frac{2\tau_f}{n^3} E_0 |f(X_1, X_2)| + \frac{2}{n^2} E_0 \left[ f(X_1, X_2)^2 \right] + \frac{\tau_f^2}{n^3}, \end{aligned}$$

where the last inequality is due to

$$E_{0}|f(X_{1}, X_{2})f(X_{1}, X_{3})| = E_{0} \{E_{0}[|f(X_{1}, X_{2})| | X_{1}]E_{0}[|f(X_{1}, X_{3})| | X_{1}]\}$$
  
$$= E_{0} \{(E_{0}[|f(X_{1}, X_{2})| | X_{1}])^{2}\}$$
  
$$\leq E_{0} \{E_{0} [f(X_{1}, X_{2})^{2} | X_{1}]\}$$
  
$$= E_{0} [f(X_{1}, X_{2})^{2}].$$

The result follows.

**Lemma E.4.** If  $f : \mathbb{R}^d \times \mathbb{R}^d \mapsto \mathbb{R}$  satisfies f(x, y) = f(y, x) and  $f(x, x) = \tau_f \in \mathbb{R}$  for all  $x, y \in \mathbb{R}^d$ , then for the empirical bootstrap  $\hat{P}_n = \mathbb{P}_n$ ,

$$E_0 E_W \left\{ \int f d[(\mathbb{P}_n^* - \mathbb{P}_n) \times (\mathbb{P}_n^* - \mathbb{P}_n)] - n^{-1} \tau_f \right\}^2$$

$$\lesssim n^{-3}\tau_f^2 + n^{-3}\tau_f \|f\|_{L_1(P_0 \times P_0)} + n^{-2} \|f\|_{L_2(P_0 \times P_0)}^2 + n^{-3} \int \left[\int f(x,y) \, \mathrm{d}P_0(x)\right]^2 P_0(y).$$

Proof of Lemma E.4. We define

$$V_n(x,y) := \int f \, d \left[ (\delta_x - \mathbb{P}_n) \times (\delta_y - \mathbb{P}_n) \right]$$
  
=  $f(x,y) - \frac{1}{n} \sum_{k=1}^n \left[ f(x, X_k) + f(y, X_k) \right] + \frac{1}{n^2} \sum_{k,l=1}^n f(X_k, X_l).$ 

By the law of total expectation, for any  $i\neq j,$ 

$$E_W \left[ V_n(X_i^*, X_j^*) \mid X_j^* \right] = E_W \left[ f(X_i^*, X_j^*) - \frac{1}{n} \sum_{k=1}^n \left\{ f(X_i^*, X_k) + f(X_j^*, X_k) \right\} + \frac{1}{n^2} \sum_{k,l=1}^n f(X_k, X_l) \mid X_j^* \right]$$
$$= \frac{1}{n} \sum_{i=1}^n f(X_i, X_j^*) - \frac{1}{n^2} \sum_{i,k=1}^n f(X_i, X_k) - \frac{1}{n} \sum_{k=1}^n f(X_j^*, X_k) + \frac{1}{n^2} \sum_{k,l=1}^n f(X_k, X_l)$$
$$= 0.$$

Hence,  $E_W[V_n(X_i^*, X_j^*)] = 0$  for  $i \neq j$  and  $E_W[V_n(X_i^*, X_j^*)V_n(X_i^*, X_k^*)] = 0$  for  $i \neq j \neq k$ .

By definition, we have

$$E_{0}E_{W}\left\{\int f d[(\mathbb{P}_{n}^{*}-\mathbb{P}_{n})\times(\mathbb{P}_{n}^{*}-\mathbb{P}_{n})] - \frac{\tau_{f}}{n}\right\}^{2}$$

$$= E_{0}E_{W}\left\{\frac{1}{n^{2}}\sum_{i=j}V_{n}(X_{i}^{*},X_{j}^{*}) + \frac{1}{n^{2}}\sum_{i\neq j}V_{n}(X_{i}^{*},X_{j}^{*}) - \frac{\tau_{f}}{n}\right\}^{2}$$

$$\lesssim E_{0}E_{W}\left\{\frac{1}{n^{2}}\sum_{i=j}V_{n}(X_{i}^{*},X_{j}^{*}) - \frac{\tau_{f}}{n}\right\}^{2} + E_{0}E_{W}\left\{\frac{1}{n^{2}}\sum_{i\neq j}V_{n}(X_{i}^{*},X_{j}^{*})\right\}^{2}$$
(27)

For the first term on the right-hand side of (27),

$$\frac{1}{n^2} \sum_{i=j} V_n(X_i^*, X_j^*) - \frac{\tau_f}{n}$$

$$= \frac{1}{n^2} \sum_{i=1}^n \int f \, d[(\delta_{X_i^*} - \mathbb{P}_n) \times (\delta_{X_i^*} - \mathbb{P}_n)] - \frac{\tau_f}{n}$$

$$= -\frac{2}{n} \int f \, d(\mathbb{P}_n \times \mathbb{P}_n^*) + \frac{1}{n} \int f \, d(\mathbb{P}_n \times \mathbb{P}_n)$$

$$= -\frac{2}{n} \int f \, d[\mathbb{P}_n \times (\mathbb{P}_n^* - \mathbb{P}_n)] - \frac{1}{n} \int f \, d(\mathbb{P}_n \times \mathbb{P}_n)$$

$$= -\frac{2}{n} \int f \,\mathrm{d}[(\mathbb{P}_n - P_0) \times (\mathbb{P}_n^* - \mathbb{P}_n)] - \frac{2}{n} \int f \,\mathrm{d}[P_0 \times (\mathbb{P}_n^* - \mathbb{P}_n)] - \frac{1}{n} \int f \,\mathrm{d}(\mathbb{P}_n \times \mathbb{P}_n).$$

Hence,

$$E_{0}E_{W}\left\{\frac{1}{n^{2}}\sum_{i=j}V_{n}(X_{i}^{*},X_{j}^{*})-\frac{\tau_{f}}{n}\right\}^{2}$$

$$\lesssim E_{0}E_{W}\left\{\frac{1}{n}\int f\,\mathrm{d}[(\mathbb{P}_{n}-P_{0})\times(\mathbb{P}_{n}^{*}-\mathbb{P}_{n})]\right\}^{2}+E_{0}E_{W}\left\{\frac{1}{n}\int f\,\mathrm{d}[P_{0}\times(\mathbb{P}_{n}^{*}-\mathbb{P}_{n})]\right\}^{2} \qquad (28)$$

$$+E_{0}\left\{\frac{1}{n}\int f\,\mathrm{d}(\mathbb{P}_{n}\times\mathbb{P}_{n})\right\}^{2}.$$

By Lemma E.3, the first term on the right-hand side of (28) is bounded up to a constant by  $n^{-5}\tau_f^2 + n^{-5}\tau_f \|f\|_{L_1(P_0 \times P_0)} + n^{-4} \|f\|_{L_2(P_0 \times P_0)}^2$ . For the second term on the right-hand side of (28),

$$E_0 E_W \left\{ \frac{1}{n} \int f \, \mathrm{d}[P_0 \times (\mathbb{P}_n^* - \mathbb{P}_n)] \right\}^2 = \frac{1}{n^3} E_0 \left[ \operatorname{Var}_W \left( \int f(x, X_1^*) \, \mathrm{d}P_0(x) \right) \right]$$
$$\leq \frac{1}{n^3} E_0 \left[ \frac{1}{n} \sum_{i=1}^n \left( \int f(x, X_i) \, \mathrm{d}P_0(x) \right)^2 \right]$$
$$\leq \frac{1}{n^3} \int \left( \int f(x, y) \, \mathrm{d}P_0(x) \right)^2 \, \mathrm{d}P_0(y).$$

For the last term on the right-hand side of (28), we have

$$E_{0}\left\{\frac{1}{n}\int f d(\mathbb{P}_{n}\times\mathbb{P}_{n})\right\}^{2} = \frac{1}{n^{2}}E_{0}\left\{\frac{1}{n^{2}}\sum_{i=j}f(X_{i},X_{j}) + \frac{1}{n^{2}}\sum_{i\neq j}f(X_{i},X_{j})\right\}^{2}$$
$$= \frac{1}{n^{2}}E_{0}\left\{\frac{\tau_{f}}{n} + \frac{1}{n^{2}}\sum_{i\neq j}f(X_{i},X_{j})\right\}^{2}$$
$$= \frac{1}{n^{2}}\left\{\frac{\tau_{f}^{2}}{n^{2}} + \frac{2\tau_{f}}{n^{3}}E_{0}\sum_{i\neq j}f(X_{i},X_{j}) + \frac{1}{n^{4}}E_{0}\left[\sum_{i\neq j}f(X_{i},X_{j})\right]^{2}\right\}$$
$$\lesssim \frac{\tau_{f}^{2}}{n^{4}} + \frac{2\tau_{f}}{n^{3}}E_{0}f(X_{1},X_{2}) + \frac{1}{n^{6}}E_{0}\left[\sum_{i\neq j}f(X_{i},X_{j})\right]^{2}$$
$$\leq \frac{\tau_{f}^{2}}{n^{4}} + \frac{2\tau_{f}}{n^{3}}\|f\|_{L_{1}(P_{0}\times P_{0})} + \frac{1}{n^{6}}E_{0}\left[\sum_{i\neq j}f(X_{i},X_{j})\right]^{2}.$$

By Lemma E.1, we note that

$$E_{0}\left[\sum_{i\neq j}f(X_{i},X_{j})\right]^{2} = E_{0}\left[4\sum_{i\neq j,j\neq k,k\neq i}f(X_{i},X_{j})f(X_{i},X_{k}) + 2\sum_{i\neq j}f(X_{i},X_{j})^{2} + \sum_{\substack{i\neq j,i\neq k,i\neq l\\ j\neq k,j\neq l,k\neq l}}f(X_{i},X_{j})f(X_{k},X_{l})\right]$$
$$\lesssim n^{3}E_{0}f(X_{1},X_{2})f(X_{1},X_{3}) + n^{2}E_{0}f(X_{1},X_{2})^{2} + n^{4}[E_{0}f(X_{1},X_{2})]^{2}$$
$$\leq n^{3}\int\left(\int f(x,y)\,\mathrm{d}P_{0}(x)\right)^{2}\,\mathrm{d}P_{0}(y) + n^{2}\,\|f\|_{L_{2}(P_{0}\times P_{0})}^{2} + n^{4}\,\|f\|_{L_{1}(P_{0}\times P_{0})}^{2}.$$

Combining these calculations, we get

$$E_0 \left\{ \frac{1}{n} \int f \, \mathrm{d}(\mathbb{P}_n \times \mathbb{P}_n) \right\}^2 \lesssim \frac{\tau_f^2}{n^4} + \frac{\tau_f}{n^3} \|f\|_{L_1(P_0 \times P_0)} + \frac{1}{n^2} \|f\|_{L_1(P_0 \times P_0)}^2 + \frac{1}{n^4} \|f\|_{L_2(P_0 \times P_0)}^2 + \frac{1}{n^3} \int \left( \int f(x, y) \, \mathrm{d}P_0(x) \right)^2 \, \mathrm{d}P_0(y).$$

Therefore,

$$E_0 E_W \left\{ \frac{1}{n^2} \sum_{i=j} V_n(X_i^*, X_j^*) - \frac{\tau_f}{n} \right\}^2 \lesssim \tau_f^2 n^{-4} + n^{-3} \tau_f \|f\|_{L_1(P_0 \times P_0)} + n^{-2} \|f\|_{L_1(P_0 \times P_0)}^2$$
$$+ n^{-4} \|f\|_{L_2(P_0 \times P_0)}^2 + n^{-3} \int \left( \int f(x, y) \, \mathrm{d}P_0(x) \right)^2 \, \mathrm{d}P_0(y).$$

For the second term on the right-hand side of (27), by Lemma E.1 and the properties of  $V_n$  derived above,

$$E_{W}\left[\frac{1}{n^{2}}\sum_{i\neq j}V_{n}(X_{i}^{*},X_{j}^{*})\right]^{2}$$

$$=E_{W}\left[\frac{4}{n^{4}}\sum_{\substack{i\neq j, j\neq k, k\neq i\\ j\neq k, j\neq l, k\neq l}}V_{n}(X_{i}^{*},X_{j}^{*})V_{n}(X_{i}^{*},X_{k}^{*})+\frac{2}{n^{4}}\sum_{\substack{i\neq j\\ i\neq j}}V_{n}(X_{i}^{*},X_{j}^{*})^{2}\right]$$

$$+\frac{1}{n^{4}}\sum_{\substack{i\neq j, i\neq k, i\neq l\\ j\neq k, j\neq l, k\neq l}}V_{n}(X_{i}^{*},X_{j}^{*})V_{n}(X_{k}^{*},X_{l}^{*})\right]$$

$$= E_{W} \left[ \frac{2}{n^{4}} \sum_{i \neq j} V_{n}(X_{i}^{*}, X_{j}^{*})^{2} \right]$$

$$\leq \frac{2}{n^{2}} E_{W} \left[ V_{n}(X_{1}^{*}, X_{2}^{*})^{2} \right]$$

$$= \frac{2}{n^{2}} E_{W} \left[ f(X_{1}^{*}, X_{2}^{*}) - \int f(X_{1}^{*}, y) \, \mathrm{d}\mathbb{P}_{n}(y) - \int f(x, X_{2}^{*}) \, \mathrm{d}\mathbb{P}_{n}(x) + (\mathbb{P}_{n} \times \mathbb{P}_{n}) f \right]^{2}$$

$$= \frac{2}{n^{2}} E_{W} \left[ f(X_{1}^{*}, X_{2}^{*}) - \int f(X_{1}^{*}, y) \, \mathrm{d}\mathbb{P}_{n}(y) - \int f(x, X_{2}^{*}) \, \mathrm{d}\mathbb{P}_{n}(x) \right]^{2} - \frac{2}{n^{2}} \left[ (\mathbb{P}_{n} \times \mathbb{P}_{n}) f \right]^{2}$$

$$\leq \frac{2}{n^{2}} E_{W} \left[ f(X_{1}^{*}, X_{2}^{*}) - \int f(X_{1}^{*}, y) \, \mathrm{d}\mathbb{P}_{n}(y) - \int f(x, X_{2}^{*}) \, \mathrm{d}\mathbb{P}_{n}(x) \right]^{2}.$$

Note that

$$\begin{split} E_W \left[ f(X_1^*, X_2^*) - \int f(X_1^*, y) \, d\mathbb{P}_n(y) - \int f(x, X_2^*) \, d\mathbb{P}_n(x) \right]^2 \\ &= E_W \left\{ f(X_1^*, X_2^*)^2 + \left[ \int f(X_1^*, y) \, d\mathbb{P}_n(y) \right]^2 + \left[ \int f(x, X_2^*) \, d\mathbb{P}_n(x) \right]^2 \\ &\quad - 2f(X_1^*, X_2^*) \left[ \int f(X_1^*, y) \, d\mathbb{P}_n(y) \right] - 2f(X_1^*, X_2^*) \left[ \int f(x, X_2^*) \, d\mathbb{P}_n(x) \right] \\ &\quad + 2 \left[ \int f(X_1^*, y) \, d\mathbb{P}_n(y) \right] \left[ \int f(x, X_2^*) \, d\mathbb{P}_n(x) \right] \right\} \\ &= E_W \left\{ f(X_1^*, X_2^*)^2 - \left[ \int f(X_1^*, y) \, d\mathbb{P}_n(y) \right]^2 - \left[ \int f(x, X_2^*) \, d\mathbb{P}_n(x) \right]^2 + 2 \left[ \int f \, d(\mathbb{P}_n \times \mathbb{P}_n) \right]^2 \right\} \\ &\leq \int f^2 \, d(\mathbb{P}_n \times \mathbb{P}_n), \end{split}$$

where the last inequality is because  $\left[n^{-2}\sum_{i,j} f(X_i, X_j)\right]^2 \leq n^{-1}\sum_j \left[n^{-1}\sum_i f(X_i, X_j)\right]^2$  by Jensen's inequality. Hence,

$$E_{0}E_{W}\left[\frac{1}{n^{2}}\sum_{i\neq j}V_{n}(X_{i}^{*},X_{j}^{*})\right]^{2} \leq \frac{2}{n^{2}}E_{0}\int f^{2} d(\mathbb{P}_{n}\times\mathbb{P}_{n})$$
$$=\frac{2}{n^{2}}E_{0}\left[\frac{1}{n^{2}}\sum_{i=j}f(X_{i},X_{j})^{2}+\frac{1}{n^{2}}\sum_{i\neq j}f(X_{i},X_{j})^{2}\right]$$
$$\lesssim \frac{\tau_{f}^{2}}{n^{3}}+\frac{1}{n^{2}}\int f^{2} d(P_{0}\times P_{0}),$$

and the result follows.

In the next few results, we assume that  $K : \mathbb{R}^d \to \mathbb{R}$  is a kernel function satisfying K(-x) = K(x)
for all  $x \in \mathbb{R}^d$ ,  $\int K(x) dx = 1$ , and  $\int K(x)^2 dx < \infty$ . Notably, K need not be non-negative or have compact support. We also let  $h = h_n > 0$  be a sequence of bandwidths. For simplicity, we assume the bandwidth matrix is  $hI_d$ , though the results can be extended to more general bandwidth matrices. For some results, we require that K is an mth order kernel function, by which we mean that  $\int z^{\alpha} K(z) dz = 0$ for all  $\alpha$  such that  $1 \leq |\alpha| \leq m - 1$  and  $\int |z^{\alpha} K(z)| dz < \infty$  for all  $\alpha$  such that  $|\alpha| = m$ .

As in the main text, we define  $K_h(x, y) := h^{-d}K(h^{-1}(x-y))$  for  $x, y \in \mathbb{R}^d$ . We define the functions  $f_1$  and  $f_2$  from  $\mathbb{R}^d$  to  $\mathbb{R}$ , each depending on K and h, as  $f_1 : (x, y) \mapsto K_h(x, y)$  and  $f_2 : (x, y) \mapsto \int K_h(x, z)K_h(y, z) \, dz$ .

**Corollary E.5.** If  $P_0$  possesses uniformly bounded Lebesgue density function  $\eta_0$ , then  $\int f d[(\mathbb{P}_n - P_0) \times (\mathbb{P}_n^* - \mathbb{P}_n)] = o_{P_W^*}(n^{-1/2})$  for each  $f \in \{f_1, f_2\}$ . If in addition  $nh^d \longrightarrow \infty$ , then  $\int f d[(\mathbb{P}_n - P_0) \times (\mathbb{P}_n - P_0)] = n^{-1}\tau_f + o_{P_0^*}(n^{-1/2})$  and  $\int f d[(\mathbb{P}_n^* - \mathbb{P}_n) \times (\mathbb{P}_n^* - \mathbb{P}_n)] = n^{-1}\tau_f + o_{P_W^*}(n^{-1/2})$ .

**Proof of Corollary E.5.** We first show that  $||f||_{L_1(P_0 \times P_0)} = O(1)$ ,  $\int \left\{ \int f(x, y) dP_0(x) \right\}^2 dP_0(y) = O(1)$ , and  $||f||_{L_2(P_0 \times P_0)} = O(h^{-d/2})$  for each  $f \in \{f_1, f_2\}$ . Note that

$$\begin{split} \|f_1\|_{L_1(P_0 \times P_0)} &= \iint \frac{1}{h^d} \left| K\left(\frac{x-y}{h}\right) \right| \, \mathrm{d}P_0(x) \, \mathrm{d}P_0(y) \\ &= \iint |K(s)| \, \eta_0(sh+y) \, \mathrm{d}s \, \mathrm{d}P_0(y) \\ &\lesssim \int |K(s)| \, \mathrm{d}s \leq \|K\|_{L_2(\lambda)} < \infty, \text{ and} \\ \|f_2\|_{L_1(P_0 \times P_0)} &= \iint \frac{1}{h^{2d}} \left| \int K\left(\frac{x-z}{h}\right) K\left(\frac{y-z}{h}\right) \, \mathrm{d}z \right| \, \mathrm{d}P_0(x) \, \mathrm{d}P_0(y) \\ &\leq \iint |K(s) \, K(t)| \left[ \int \eta_0(z+hs) \eta_0(z+ht) \, \mathrm{d}z \right] \, \mathrm{d}s \, \mathrm{d}t \\ &\lesssim \left[ \int |K(s)| \, \mathrm{d}s \right]^2 \leq \|K\|_{L_2(\lambda)}^2 < \infty. \end{split}$$

Next,

$$\int \left[ \int f_1(x,y) \, \mathrm{d}P_0(x) \right]^2 \, \mathrm{d}P_0(y) = \int \left[ \frac{1}{h^d} \int K\left(\frac{x-y}{h}\right) \, \mathrm{d}P_0(x) \right]^2 \, \mathrm{d}P_0(y)$$
$$= \int \left[ \int K(s) \, \eta_0(sh+y) \, \mathrm{d}s \right]^2 \, \mathrm{d}P_0(y)$$
$$\lesssim \|K\|_{L_2(\lambda)}^2 < \infty, \text{ and}$$
$$\int \left[ \int f_2(x,y) \, \mathrm{d}P_0(x) \right]^2 \, \mathrm{d}P_0(y) = \int \left[ \frac{1}{h^{2d}} \iint K\left(\frac{x-z}{h}\right) K\left(\frac{y-z}{h}\right) \, \mathrm{d}z \, \mathrm{d}P_0(x) \right]^2 \, \mathrm{d}P_0(y)$$

$$\leq \int \left[ \iint K(s) K(t) \eta_0(y - ht + hs) \, \mathrm{d}s \, \mathrm{d}t \right]^2 \, \mathrm{d}P_0(y)$$
  
$$\lesssim \|K\|_{L_2(\lambda)}^2 < \infty.$$

Similarly,

$$\begin{split} \|f_1\|_{L_2(P_0 \times P_0)}^2 &= \frac{1}{h^{2d}} \iint K^2 \left(\frac{x-y}{h}\right) \, \mathrm{d}P_0(x) \, \mathrm{d}P_0(y) \\ &= \frac{1}{h^d} \iint K^2(s) \, \eta_0(sh+y) \, \mathrm{d}s \, \mathrm{d}P_0(y) \\ &\lesssim \frac{1}{h^d} \|K\|_{L_2(\lambda)}^2, \end{split}$$

and

$$\begin{split} \|f_2\|_{L_2(P_0 \times P_0)}^2 &= \iint \left[\frac{1}{h^{2d}} \int K\left(\frac{x-z}{h}\right) K\left(\frac{y-z}{h}\right) dz\right]^2 dP_0(x) dP_0(y) \\ &= \frac{1}{h^{4d}} \iiint K\left(\frac{x-z}{h}\right) K\left(\frac{y-z}{h}\right) K\left(\frac{x-w}{h}\right) K\left(\frac{y-w}{h}\right) dz \, dw \, dP_0(x) \, dP_0(y) \\ &= \frac{1}{h^{2d}} \iiint K\left(s\right) K\left(s\right) K\left(s + \frac{y-x}{h}\right) K\left(t\right) K\left(t + \frac{y-x}{h}\right) ds \, dt \, dP_0(x) \, dP_0(y) \\ &= \frac{1}{h^d} \iiint K\left(s\right) K\left(r\right) K\left(t\right) K\left(t + r - s\right) \eta_0(y - h(r - s)) \, ds \, dt \, dP_0(y) \\ &\lesssim \frac{1}{h^d} \iiint |K\left(s\right) K\left(r\right) K\left(t\right) K\left(t + r - s\right)| \, ds \, dt \, dr \\ &\leq \frac{1}{h^d} \iint |K\left(s\right) K\left(r\right)| \left[\int K^2\left(t\right) \, dt\right]^{1/2} \left[\int K^2\left(t + r - s\right) \, dt\right]^{1/2} \, ds \, dr \\ &= \frac{1}{h^d} \left[\int |K\left(s\right)| \, ds\right]^2 \left[\int K^2\left(t\right) \, dt\right] \\ &\leq \frac{1}{h^d} \|K\|_{L_2(\lambda)}^4, \end{split}$$

which implies that  $||f||_{L_2(P_0 \times P_0)} = O(h^{-d/2})$  for each  $f \in \{f_1, f_2\}$  as claimed.

We note that f(x,y) = f(y,x) for each  $f \in \{f_1, f_2\}$  because K(-u) = K(u) by assumption, and  $\tau_{f_1} = h^{-d}K(0)$  and  $\tau_{f_2} = h^{-d}\int K^2(u) du$  for all  $x \in \mathbb{R}^d$ , which are both  $O(h^{-d})$ . Hence, using the results above, by Lemma E.3, for each  $f \in \{f_1, f_2\}$ ,

$$E_0 E_W \left\{ \int f d[(\mathbb{P}_n - P_0) \times (\mathbb{P}_n^* - \mathbb{P}_n)] \right\}^2 \lesssim h^{-2d} n^{-3} + h^{-d} n^{-3} \|f\|_{L_1(P_0 \times P_0)} + n^{-2} \|f\|_{L_2(P_0 \times P_0)}^2$$
$$\lesssim h^{-2d} n^{-3} + h^{-d} n^{-3} + n^{-2} h^{-d}$$

$$\lesssim n^{-1} (nh^d)^{-2} + n^{-1} (nh^d)^{-1}$$

which is  $o(n^{-1})$  if  $nh^d \longrightarrow \infty$ . This implies that  $\int f d[(\mathbb{P}_n - P_0) \times (\mathbb{P}_n^* - \mathbb{P}_n)] = o_{P_W^*}(n^{-1/2})$  if  $nh^d \longrightarrow \infty$ . Since  $\int \left[\int f(x, y) dP_0(x)\right]^2 dP_0(y) = O(1)$ , by Lemma E.2, we have

$$E_0 \left\{ \int f d[(\mathbb{P}_n - P_0) \times (\mathbb{P}_n - P_0)] - n^{-1} \tau_f \right\}^2 \lesssim n^{-3} \int \left[ \int f(x, y) dP_0(x) \right]^2 dP_0(y) + n^{-2} \|f\|_{L_2(P_0 \times P_0)}^2$$
  
$$\lesssim n^{-3} + n^{-1} (nh^d)^{-1},$$

which is  $o(n^{-1})$  if  $nh^d \longrightarrow \infty$ . This implies that  $\int f d[(\mathbb{P}_n - P_0) \times (\mathbb{P}_n - P_0)] = n^{-1}\tau_f + o_{P_W^*}(n^{-1/2})$  if  $nh^d \longrightarrow \infty$ . Since  $\int \left[\int f(x, y) dP_0(x)\right]^2 dP_0(y) = O(1)$ , by Lemma E.4, we have

$$E_{0}E_{W}\left\{\int f d[(\mathbb{P}_{n}^{*}-\mathbb{P}_{n})\times(\mathbb{P}_{n}^{*}-\mathbb{P}_{n})]-n^{-1}\tau_{f}\right\}^{2}$$
  
$$\lesssim n^{-3}\tau_{f}^{2}+n^{-3}\tau_{f}\|f\|_{L_{1}(P_{0}\times P_{0})}+n^{-2}\|f\|_{L_{2}(P_{0}\times P_{0})}^{2}+n^{-3}\int \left[\int f(x,y) dP_{0}(x)\right]^{2} dP_{0}(y)$$
  
$$\lesssim n^{-3}h^{-2d}+n^{-3}h^{-d}+n^{-2}h^{-d}+n^{-3},$$

which is  $o(n^{-1})$  if  $nh^d \longrightarrow \infty$ . This implies that  $\int f d[(\mathbb{P}_n^* - \mathbb{P}_n) \times (\mathbb{P}_n^* - \mathbb{P}_n)] = n^{-1}\tau_f + o_{P_W^*}(n^{-1/2})$  if  $nh^d \longrightarrow \infty$ .

**Lemma E.6.** Suppose that the density function  $\eta_0$  of  $P_0$  is m-times continuously differentiable with  $\int [(D^{\alpha}\eta_0)(x)]^2 dx < \infty$  for all  $\alpha$  such that  $|\alpha| = m$ , and K is an mth order kernel. Then for each  $f \in \{f_1, f_2\}$ 

$$\int \left[\int f(x,y) \,\mathrm{d}P_0(y) - \eta_0(x)\right]^2 \,\mathrm{d}x = O(h^{2m}).$$

If in addition  $\sup_{x \in \mathbb{R}^d} |D^{\alpha} \eta_0(x)| < \infty$ , then  $\sup_{x \in \mathbb{R}^d} \left| \int f(x, y) \, \mathrm{d}P_0(y) - \eta_0(x) \right| = O(h^m)$ .

**Proof of Lemma E.6.** Since  $\eta_0$  is *m*-times continuously differentiable, for all *u*, a Taylor expansion with the Laplacian representation of the remainder gives

$$\eta_0(x+u) - \eta_0(x) = \sum_{1 \le |\alpha| \le m-1} \frac{1}{\alpha!} u^{\alpha} (D^{\alpha} \eta_0)(x) + \sum_{|\alpha|=m} \frac{m}{\alpha!} u^{\alpha} \int_0^1 (1-r)^{m-1} (D^{\alpha} \eta_0)(x+ru) \, \mathrm{d}r.$$
<sup>(29)</sup>

Hence, for  $f = f_1$  and any  $x \in \mathbb{R}^d$ , we have

$$\begin{split} \int f_1(x,y) \, \mathrm{d}P_0(y) &- \eta_0(x) = \int \frac{1}{h^d} K\left(\frac{x-y}{h}\right) \eta_0(y) \, \mathrm{d}y - \eta_0(x) \\ &= \int K(s) [\eta_0(x+hs) - \eta_0(x)] \, \mathrm{d}s \\ &= \sum_{1 \le |\alpha| \le m-1} \frac{1}{\alpha!} h^{|\alpha|} \int s^{\alpha} K(s) \, \mathrm{d}s (D^{\alpha} \eta_0)(x) \\ &+ \sum_{|\alpha|=m} \frac{m}{\alpha!} h^{|\alpha|} \iint_0^1 s^{\alpha} K(s) (1-r)^{m-1} (D^{\alpha} \eta_0)(x+rhs) \, \mathrm{d}r \, \mathrm{d}s. \end{split}$$

Since K is an *m*th order kernel function, the first term on the right hand size is zero. Defining  $H(x, s, \alpha) := \int_0^1 (1 - r)^{m-1} (D^{\alpha} \eta_0) (x + rhs) \, \mathrm{d}r$ , we then have

By the Cauchy–Schwarz inequality, we have

$$\int H(x,s,\alpha)^{2} dx = \int \left[ \int_{0}^{1} (1-r)^{m-1} (D^{\alpha}\eta_{0})(x+rhs) dr \right]^{2} dx$$
  

$$\leq \int \left[ \int_{0}^{1} (1-r)^{2(m-1)} dr \right] \left[ \int_{0}^{1} (D^{\alpha}\eta_{0})(x+rhs)^{2} dr \right] dx$$
  

$$\leq \iint_{0}^{1} (D^{\alpha}\eta_{0})(x+rhs)^{2} dr dx$$
  

$$= \int [(D^{\alpha}\eta_{0})(x)]^{2} dx,$$

which is finite by assumption. Hence,

$$\int \left[ \int f_1(x,y) \, \mathrm{d}P_0(y) - \eta_0(x) \right]^2 \, \mathrm{d}x \lesssim h^{2m} \sum_{|\alpha|,|\beta|=m} \iint \left| s^{\alpha} t^{\beta} K(s) K(t) \right| \, \mathrm{d}s \, \mathrm{d}t$$

$$= h^{2m} \left[ \sum_{|\alpha|=m} \int |s^{\alpha} K(s)| \, \mathrm{d}s \right]^2 = O(h^{2m}).$$

If  $D^{\alpha}\eta_0$  is uniformly bounded, then

$$\begin{split} \left| \int f_1(x,y) \, \mathrm{d}P_0(y) - \eta_0(x) \right| \\ &= \left| \sum_{|\alpha|=m} \frac{m}{\alpha!} h^{|\alpha|} \int \!\!\!\int_0^1 s^\alpha K(s) (1-r)^{m-1} (D^\alpha \eta_0) (x+rhs) \, \mathrm{d}r \, \mathrm{d}s \right| \\ &\leq \|D^\alpha \eta_0\|_\infty \sum_{|\alpha|=m} \frac{mh^m}{\alpha!} \int |s^\alpha K(s)| \left[ \int_0^1 (1-r)^{m-1} \, \mathrm{d}r \right] \, \mathrm{d}s \\ &= \|D^\alpha \eta_0\|_\infty \sum_{|\alpha|=m} \frac{h^m}{\alpha!} \int |s^\alpha K(s)| \, \mathrm{d}s, \end{split}$$

which is independent of x and order of  $h^m$ .

For  $f = f_2$ , we have

$$\begin{split} \int f_2(x,y) \, \mathrm{d}P_0(y) &- \eta_0(x) \\ &= \iint \frac{1}{h^{2d}} K\left(\frac{x-z}{h}\right) K\left(\frac{y-z}{h}\right) \, \mathrm{d}z \, \mathrm{d}P_0(y) - \eta_0(x) \\ &= \iint \frac{1}{h^d} K\left(s\right) K\left(s + \frac{y-x}{h}\right) \, \mathrm{d}s \, \mathrm{d}P_0(y) - \eta_0(x) \\ &= \iint K(s) K(t) \eta_0(x+h(t-s)) \, \mathrm{d}s \, \mathrm{d}t - \eta_0(x) \\ &= \iint K(s) K(t) [\eta_0(x+h(t-s)) - \eta_0(x)] \, \mathrm{d}s \, \mathrm{d}t \\ &= \sum_{1 \le |\alpha| \le m-1} \frac{1}{\alpha!} h^{|\alpha|} \iint (t-s)^{\alpha} K(s) K(t) \, \mathrm{d}s \, \mathrm{d}t (D^{\alpha} \eta_0)(x) \\ &\quad + \sum_{|\alpha|=m} \frac{m}{\alpha!} h^{|\alpha|} \iiint^1 (t-s)^{\alpha} K(s) K(t) (1-r)^{m-1} (D^{\alpha} \eta_0)(x+rh(t-s)) \, \mathrm{d}r \, \mathrm{d}s \, \mathrm{d}t, \end{split}$$

where the first term on the right hand size is zero because K is an mth order kernel function. Hence, we have

$$\int \left[ \int f_2(x,y) \, \mathrm{d}P_0(y) - \eta_0(x) \right]^2 \, \mathrm{d}x$$
$$= \int \left[ \sum_{|\alpha|=m} \frac{m}{\alpha!} h^{|\alpha|} \iint (t-s)^{\alpha} K(s) K(t) H(x,s-t,\alpha) \, \mathrm{d}s \, \mathrm{d}t \right]^2 \, \mathrm{d}x$$

$$\begin{split} &= \sum_{|\alpha|,|\beta|=m} \frac{m^2}{\alpha!\beta!} h^{|\alpha|} h^{|\beta|} \iiint (t-s)^{\alpha} (t'-s')^{\beta} K(s) K(s') K(t) K(t') \\ &\quad \times \left[ \int H(x,s-t,\alpha) H(x,s'-t',\beta) \, \mathrm{d}x \right] \, \mathrm{d}s \, \mathrm{d}t \, \mathrm{d}s' \, \mathrm{d}t' \\ &\lesssim \sum_{|\alpha|,|\beta|=m} h^{2m} \iiint \left| (t-s)^{\alpha} (t'-s')^{\beta} K(s) K(s') K(t) K(t') \right| \\ &\quad \times \left[ \int H(x,s-t,\alpha)^2 \, \mathrm{d}x \right]^{1/2} \left[ \int H(x,s'-t',\beta)^2 \, \mathrm{d}x \right]^{1/2} \, \mathrm{d}s \, \mathrm{d}t \, \mathrm{d}s' \, \mathrm{d}t' \\ &\lesssim h^{2m} \left[ \sum_{|\alpha|=m} \iint \left| (t-s)^{\alpha} K(s) K(t) \right| \, \mathrm{d}s \, \mathrm{d}t \right]^2 \\ &\lesssim h^{2m} \left[ \sum_{|\alpha|=m} \iint \left| \prod_{i=1}^d \left( \sum_{k_i=0}^{\alpha_i} t_i^{k_i} s_i^{\alpha_i-k_i} \right) K(s) K(t) \right| \, \mathrm{d}s \, \mathrm{d}t \right]^2 \\ &\leq h^{2m} \left[ \sum_{|\alpha|=m} \iint \left| \sum_{\substack{0 \le \beta \le \alpha \\ 0 \le \gamma \le \alpha}} t^{\beta} s^{\gamma} K(s) K(t) \right| \, \mathrm{d}s \, \mathrm{d}t \right]^2 \\ &\leq h^{2m} \left[ \sum_{\substack{0 \le |\beta| \le m \\ 0 \le |\gamma| \le m}} \iint \left| t^{\beta} s^{\gamma} K(s) K(t) \right| \, \mathrm{d}s \, \mathrm{d}t \right]^2 \\ &= h^{2m} \left[ \sum_{\substack{0 \le |\beta| \le m \\ 0 \le |\gamma| \le m}} \int \left| t^{\beta} K(t) \right| \, \mathrm{d}t \right]^4 , \end{split}$$

which is  $O(h^{2m})$  because  $\sum_{|\beta|=m} \int |t^{\beta}K(t)| dt < \infty$  by assumption, and for any  $0 \le \beta < m$  there exists  $\alpha$  with  $|\alpha| = m$  such that

$$\int \left| t^{\beta} K(t) \right| \, \mathrm{d}t = \int_{[-1,1]^d} \left| t^{\beta} K(t) \right| \, \mathrm{d}t + \int_{\mathbb{R}^d \setminus [-1,1]^d} \left| t^{\beta} K(t) \right| \, \mathrm{d}t$$
$$\leq \int |K(t)| \, \mathrm{d}t + \int |t^{\alpha} K(t)| \, \mathrm{d}t,$$

which is finite by assumption.

If  $D^{\alpha}\eta_0$  is uniformly bounded, then

$$\begin{aligned} \left| \int f_2(x,y) \, \mathrm{d}P_0(y) - \eta_0(x) \right| \\ &\leq \|D^{\alpha} \eta_0\|_{\infty} \sum_{|\alpha|=m} \frac{m}{\alpha!} h^{|\alpha|} \iint |(t-s)^{\alpha} K(s) K(t)| \, \mathrm{d}s \, \mathrm{d}t \int_0^1 (1-r)^{m-1} \, \mathrm{d}r \end{aligned}$$

$$= \|D^{\alpha}\eta_0\|_{\infty} \sum_{|\alpha|=m} \frac{h^m}{\alpha!} \iint |(t-s)^{\alpha} K(s)K(t)| \, \mathrm{d}s \, \mathrm{d}t.$$

We showed the integral in the final expression is finite above, so the expression is independent of x and order of  $h^m$ .

We now consider kernel density estimators  $\eta_n$  and  $\eta_n^*$  with common bandwidth h and kernel K, i.e.  $\eta_n(x) = n^{-1} \sum_{i=1}^n K_h(x, X_i)$  and  $\eta_n^*(x) = n^{-1} \sum_{i=1}^n K_h(x, X_i^*)$ .

**Lemma E.7.** Suppose that the density function  $\eta_0$  of  $P_0$  is uniformly bounded and m-times continuously differentiable with  $\int [(D^{\alpha}\eta_0)(x)]^2 dx < \infty$  for all  $\alpha$  such that  $|\alpha| = m$  and K is an m-th order kernel function. Then  $(\mathbb{P}_n - P_0)(\eta_n^* - \eta_n) = o_{P_W^*}(n^{-1/2})$ . If in addition  $nh^d \longrightarrow \infty$  holds, then  $(\mathbb{P}_n^* - \mathbb{P}_n)(\eta_n^* - \eta_0) - (\mathbb{P}_n - P_0)(\eta_n - \eta_0) = o_{P_W^*}(n^{-1/2})$ .

Proof of Lemma E.7. To show the first statement, we note that

$$(\mathbb{P}_n - P_0)(\eta_n^* - \eta_n) = \int K_h \,\mathrm{d}[(\mathbb{P}_n^* - \mathbb{P}_n) \times (\mathbb{P}_n - P_0)],$$

which is  $o_{P_W^*}(n^{-1/2})$  by Corollary E.5 with  $f = f_1$ .

To show the second statement, by adding and subtracting terms and defining  $\eta_{0,h}(x) = \int K_h(x,y) \, dP_0(y)$ , we have

$$\begin{aligned} (\mathbb{P}_{n}^{*} - \mathbb{P}_{n})(\eta_{n}^{*} - \eta_{0}) - (\mathbb{P}_{n} - P_{0})(\eta_{n} - \eta_{0}) \\ &= (\mathbb{P}_{n}^{*} - \mathbb{P}_{n})(\eta_{n}^{*} - \eta_{n}) + (\mathbb{P}_{n}^{*} - \mathbb{P}_{n})(\eta_{n} - \eta_{0,h}) + (\mathbb{P}_{n}^{*} - \mathbb{P}_{n})(\eta_{0,h} - \eta_{0}) - (\mathbb{P}_{n} - P_{0})(\eta_{n} - \eta_{0,h}) \\ &- (\mathbb{P}_{n} - P_{0})(\eta_{0,h} - \eta_{0}) \\ &= \int K_{h} d[(\mathbb{P}_{n}^{*} - \mathbb{P}_{n}) \times (\mathbb{P}_{n}^{*} - \mathbb{P}_{n})] + \int K_{h} d[(\mathbb{P}_{n} - P_{0}) \times (\mathbb{P}_{n}^{*} - \mathbb{P}_{n})] \\ &+ \int [K_{h}(x, y) - \eta_{0}(x)] d(\mathbb{P}_{n}^{*} - \mathbb{P}_{n})(x) dP_{0}(y)] - \int K_{h} d[(\mathbb{P}_{n} - P_{0}) \times (\mathbb{P}_{n} - P_{0})] \\ &- \int [K_{h}(x, y) - \eta_{0}(x)] d(\mathbb{P}_{n} - P_{0})(x) dP_{0}(y)]. \end{aligned}$$

By Corollary E.5 with  $f = f_1$ ,

$$\int K_h \, d[(\mathbb{P}_n - P_0) \times (\mathbb{P}_n^* - \mathbb{P}_n)] = o_{P_W^*}(n^{-1/2}),$$

and if  $nh^d \longrightarrow 0$ , then

$$\int K_h \,\mathrm{d}[(\mathbb{P}_n^* - \mathbb{P}_n) \times (\mathbb{P}_n^* - \mathbb{P}_n)] - \int K_h \,\mathrm{d}[(\mathbb{P}_n - P_0) \times (\mathbb{P}_n - P_0)] = o_{P_W^*}(n^{-1/2}).$$

Finally, we write

$$\int [K_h(x,y) - \eta_0(x)] d(\mathbb{P}_n - P_0)(x) dP_0(y)] = (\mathbb{P}_n - P_0)g_h, \text{ and} \\ \int [K_h(x,y) - \eta_0(x)] d(\mathbb{P}_n^* - \mathbb{P}_n)(x) dP_0(y)] = (\mathbb{P}_n^* - \mathbb{P}_n)g_h$$

for  $g_h(x) := \int [K_h(x,y) - \eta_0(x)] dP_0(y)$ . As in the proofs of Lemma E.3 and Lemma E.4, we have

$$E_0 \left[ (\mathbb{P}_n - P_0)g_h \right]^2 \le n^{-1} \|g_h\|_{L_2(P_0)}^2, \text{ and}$$
$$E_0 E_W \left[ (\mathbb{P}_n^* - \mathbb{P}_n)g_h \right]^2 \le n^{-1} E_0 \|g_h\|_{L_2(\mathbb{P}_n)}^2 = n^{-1} \|g_h\|_{L_2(P_0)}^2$$

Since  $\int g_h(x)^2 dx = O(h^{2m})$  by Lemma E.6, we have  $\|g_h\|_{L_2(P_0)}^2 \longrightarrow 0$  as long as  $h \longrightarrow 0$  and  $P_0$  possesses uniformly bounded density. This implies that  $\int [K_h(x,y) - \eta_0(x)] d(\mathbb{P}_n - P_0)(x) dP_0(y)] = o_{P_0^*}(n^{-1/2})$ and  $\int [K_h(x,y) - \eta_0(x)] d(\mathbb{P}_n^* - \mathbb{P}_n)(x) dP_0(y)] = o_{P_W^*}(n^{-1/2}).$ 

**Lemma E.8.** Suppose that the density function  $\eta_0$  of  $P_0$  is uniformly bounded and m-times continuously differentiable with  $\int [(D^{\alpha}\eta_0)(x)]^2 dx < \infty$  for all  $\alpha$  such that  $|\alpha| = m$  and K is an m-th order kernel function. If  $nh^d \longrightarrow \infty$  and  $nh^{4m} \longrightarrow 0$ , then  $2 \int [\eta_n(x) - \eta_0(x)]^2 dx - \int [\eta_n^*(x) - \eta_0(x)]^2 dx = o_{P_W^*}(n^{-1/2})$  holds for the empirical bootstrap.

**Proof of Lemma E.8.** Recall that  $f_1(x, y) := K_h(x, y)$  and  $f_2(x, y) := \int K_h(x, z) K_h(y, z) dz$ . We denote  $\eta_{0,h}(x) := \int K_h(x, y) dP_0(y)$  for simplicity. We then note that

$$\int \eta_n^2 = \int f_2 d(\mathbb{P}_n \times \mathbb{P}_n), \ \int \eta_n^{*2} = \int f_2 d(\mathbb{P}_n^* \times \mathbb{P}_n^*), \ \int \eta_{0,h}^2 = \int f_2 d(P_0 \times P_0)$$
$$\int \eta_n^* \eta_n = \int f_2 d(\mathbb{P}_n^* \times \mathbb{P}_n), \ \int \eta_n^* \eta_{0,h} = \int f_2 d(\mathbb{P}_n^* \times P_0), \ \int \eta_n \eta_{0,h} = \int f_2 d(\mathbb{P}_n \times P_0).$$

Hence, by adding and subtracting term, we have

$$\int (\eta_n^* - \eta_0)^2 - 2 \int (\eta_n - \eta_0)^2$$

$$\begin{split} &= \int (\eta_n^* - \eta_n + \eta_n - \eta_{0,h} + \eta_{0,h} - \eta_0)^2 - 2 \int (\eta_n - \eta_{0,h} + \eta_{0,h} - \eta_0)^2 \\ &= \int (\eta_n^* - \eta_n)^2 - \int (\eta_n - \eta_{0,h})^2 - \int (\eta_{0,h} - \eta_0)^2 + 2 \int (\eta_n^* - \eta_n) (\eta_n - \eta_{0,h}) \\ &+ 2 \int (\eta_n^* - \eta_n) (\eta_{0,h} - \eta_0) - 2 \int (\eta_n - \eta_{0,h}) (\eta_{0,h} - \eta_0) \\ &= \int f_2 d[(\mathbb{P}_n^* - \mathbb{P}_n) \times (\mathbb{P}_n^* - \mathbb{P}_n)] - \int f_2 d[(\mathbb{P}_n - P_0) \times (\mathbb{P}_n - P_0)] - \int (\eta_{0,h} - \eta_0)^2 \\ &+ 2 \int f_2 d[(\mathbb{P}_n^* - \mathbb{P}_n) \times (\mathbb{P}_n - P_0)] + 2 \int [f_2(x, y) - \eta_{0,h}(x)] d(\mathbb{P}_n^* - \mathbb{P}_n)(x) dP_0(y) \\ &- 2 \int [f_2(x, y) - \eta_{0,h}(x)] d(\mathbb{P}_n - P_0)(x) dP_0(y). \end{split}$$

By Corollary E.5,  $2 \int f_2 d[(\mathbb{P}_n^* - \mathbb{P}_n) \times (\mathbb{P}_n - P_0)] = o_{P_W^*}(n^{-1/2})$ , and if  $nh^d \longrightarrow 0$ , then

$$\int f_2 d[(\mathbb{P}_n^* - \mathbb{P}_n) \times (\mathbb{P}_n^* - \mathbb{P}_n)] - \int f_2 d[(\mathbb{P}_n - P_0) \times (\mathbb{P}_n - P_0)] = o_{P_W^*}(n^{-1/2})$$

By Lemma E.6, the third term on the right is  $O(h^{2m})$ , which is  $o(n^{-1/2})$  if  $nh^{4m} \longrightarrow 0$ . Finally, we write

$$\int [f_2(x,y) - \eta_{0,h}(x)] d(\mathbb{P}_n - P_0)(x) dP_0(y) = (\mathbb{P}_n - P_0)g_h, \text{ and} \\ \int [f_2(x,y) - \eta_{0,h}(x)] d(\mathbb{P}_n^* - \mathbb{P}_n)(x) dP_0(y) = (\mathbb{P}_n^* - \mathbb{P}_n)g_h$$

for  $g_h(x) := \int [f_2(x,y) - \eta_{0,h}(x)] dP_0(y) = \int [f_2(x,y) - f_1(x,y)] dP_0(y)$ . As in the proofs of Lemma E.3 and Lemma E.4, we have

$$E_0 \left[ (\mathbb{P}_n - P_0) g_h \right]^2 \le n^{-1} \|g_h\|_{L_2(P_0)}^2, \text{ and}$$
$$E_0 E_W \left[ (\mathbb{P}_n^* - \mathbb{P}_n) g_h \right]^2 \le n^{-1} E_0 \|g_h\|_{L_2(\mathbb{P}_n)}^2 = n^{-1} \|g_h\|_{L_2(P_0)}^2$$

By Lemma E.6, we have

$$\begin{split} \int \left\{ \int [f_2(x,y) - f_1(x,y)] \, \mathrm{d}P_0(y) \right\}^2 \, \mathrm{d}x \\ &\leq 2 \int \left\{ \int f_2(x,y) \, \mathrm{d}P_0(y) - \eta_0(x) \right\}^2 \, \mathrm{d}x + 2 \int \left\{ \int f_1(x,y) \, \mathrm{d}P_0(y) - \eta_0(x) \right\}^2 \, \mathrm{d}x = O(h^{2m}). \end{split}$$

This implies  $||g_h||^2_{L_2(P_0)} = O(h^{2m}) = o(n^{-1/2})$  since  $nh^{4m} \longrightarrow 0$  and  $P_0$  possesses uniformly bounded density. Hence,  $\int [f_2(x,y) - \eta_{0,h}(x)] d(\mathbb{P}_n - P_0)(x) dP_0(y) = o_{P_0^*}(n^{-3/4})$  and  $\int [f_2(x,y) - \eta_{0,h}(x)] d(\mathbb{P}_n^* - P_0)(x) dP_0(y) = o_{P_0^*}(n^{-3/4})$ 

$$\mathbb{P}_n(x) \,\mathrm{d}P_0(y) = o_{P_W^*}(n^{-3/4}).$$

**Lemma E.9.** Suppose that the density function  $\eta_0$  of  $P_0$  is uniformly bounded and m-times continuously differentiable with  $\int [(D^{\alpha}\eta_0)(x)]^2 dx < \infty$  for all  $\alpha$  such that  $|\alpha| = m$  and K is an m-th order kernel function. If  $nh^d \longrightarrow \infty$  and  $nh^{2m} \longrightarrow 0$ , then  $2\mathbb{P}_n\phi_n - \mathbb{P}_n^*\phi_n^* = o_{P_W}(n^{-1/2})$  holds for the empirical bootstrap.

**Proof of Lemma E.9.** Recall that  $f_1(x,y) := K_h(x,y)$  and  $f_2(x,y) := \int K_h(x,z)K_h(y,z) dz$ . We define  $f_3 := f_1 - f_2$ . We then have

$$\mathbb{P}_n \phi_n = 2 \int f_3 d(\mathbb{P}_n \times \mathbb{P}_n), \text{ and } \mathbb{P}_n^* \phi_n^* = 2 \int f_3 d(\mathbb{P}_n^* \times \mathbb{P}_n^*).$$

Hence, by adding and subtracting terms,

$$\begin{aligned} \left(\mathbb{P}_{n}^{*}\phi_{n}^{*}-2\mathbb{P}_{n}\phi_{n}\right)/2 \\ &=\int f_{3}\,\mathrm{d}[\left(\mathbb{P}_{n}^{*}-\mathbb{P}_{n}+\mathbb{P}_{n}-P_{0}+P_{0}\right)\times\left(\mathbb{P}_{n}^{*}-\mathbb{P}_{n}+\mathbb{P}_{n}-P_{0}+P_{0}\right)] \\ &\quad -2\int f_{3}\,\mathrm{d}[\left(\mathbb{P}_{n}-P_{0}+P_{0}\right)\times\left(\mathbb{P}_{n}-P_{0}+P_{0}\right)] \\ &=\int f_{3}\,\mathrm{d}[\left(\mathbb{P}_{n}^{*}-\mathbb{P}_{n}\right)\times\left(\mathbb{P}_{n}^{*}-\mathbb{P}_{n}\right)] -\int f_{3}\,\mathrm{d}[\left(\mathbb{P}_{n}-P_{0}\right)\times\left(\mathbb{P}_{n}-P_{0}\right)] -\int f_{3}\,\mathrm{d}[P_{0}\times P_{0}] \\ &\quad +2\int f_{3}\,\mathrm{d}[\left(\mathbb{P}_{n}^{*}-\mathbb{P}_{n}\right)\times\left(\mathbb{P}_{n}-P_{0}\right)] +2\int f_{3}\,\mathrm{d}[\left(\mathbb{P}_{n}^{*}-\mathbb{P}_{n}\right)\times P_{0}] \\ &\quad -2\int f_{3}\,\mathrm{d}[\left(\mathbb{P}_{n}-P_{0}\right)\times P_{0}]. \end{aligned}$$

By Corollary E.5,

$$\int f_3 d[(\mathbb{P}_n^* - \mathbb{P}_n) \times (\mathbb{P}_n - P_0)] = \int f_1 d[(\mathbb{P}_n^* - \mathbb{P}_n) \times (\mathbb{P}_n - P_0)] - \int f_2 d[(\mathbb{P}_n^* - \mathbb{P}_n) \times (\mathbb{P}_n - P_0)] = o_{P_W^*}(n^{-1/2})$$

and if  $nh^d \longrightarrow 0$ , then

$$\int f_3 d[(\mathbb{P}_n^* - \mathbb{P}_n) \times (\mathbb{P}_n^* - \mathbb{P}_n)] - \int f_3 d[(\mathbb{P}_n - P_0) \times (\mathbb{P}_n - P_0)]$$
  
=  $\left\{ \int f_1 d[(\mathbb{P}_n^* - \mathbb{P}_n) \times (\mathbb{P}_n^* - \mathbb{P}_n)] - \int f_1 d[(\mathbb{P}_n - P_0) \times (\mathbb{P}_n - P_0)] \right\}$   
 $- \left\{ \int f_2 d[(\mathbb{P}_n^* - \mathbb{P}_n) \times (\mathbb{P}_n^* - \mathbb{P}_n)] - \int f_2 d[(\mathbb{P}_n - P_0) \times (\mathbb{P}_n - P_0)] \right\}$   
=  $o_{P_W^*}(n^{-1/2}).$ 

Finally, we define  $g_h(x) := \int f_3(x, y) \, dP_0(y)$ . As in the proofs of Lemma E.3 and Lemma E.4, we have

$$E_0 \left[ (\mathbb{P}_n - P_0)g_h \right]^2 \le n^{-1} \|g_h\|_{L_2(P_0)}^2, \text{ and}$$
$$E_0 E_W \left[ (\mathbb{P}_n^* - \mathbb{P}_n)g_h \right]^2 \le n^{-1} E_0 \|g_h\|_{L_2(\mathbb{P}_n)}^2 = n^{-1} \|g_h\|_{L_2(P_0)}^2.$$

By Lemma E.6, we have

$$\int g_h(x)^2 dx = \int \left\{ \int [f_1(x,y) - f_2(x,y)] dP_0(y) \right\}^2 dx$$

$$\leq 2 \int \left\{ \int f_1(x,y) dP_0(y) - \eta_0(x) \right\}^2 dx + 2 \int \left\{ \int f_2(x,y) dP_0(y) - \eta_0(x) \right\}^2 dx,$$
(30)

which is  $O(h^{2m})$ . This implies that  $(\mathbb{P}_n - P_0)g_h$  and  $(\mathbb{P}_n^* - \mathbb{P}_n)g_h$  are  $o_{P_0^*}(n^{-1})$  and  $o_{P_W^*}(n^{-1})$ , respectively, since  $nh^{2m} \longrightarrow 0$  and  $P_0$  possesses uniformly bounded density. Finally, (30) also implies that  $\int f_3 d(P_0 \times P_0) = O(h^m) = o(n^{-1/2})$  since  $nh^{2m} \longrightarrow 0$ .

## Supplement F Lemmas supporting the proof of Proposition 5.5

For any  $f : \mathbb{R}^d \times \mathbb{R}^d \mapsto \mathbb{R}$ , we define

$$G[f](x,y) := \iint f(s,t)K_h(s,x)K_h(t,y)\,\mathrm{d}s\,\mathrm{d}t = \iint f(x+hu,y+hv)K(u)K(v)\,\mathrm{d}u\,\mathrm{d}v.$$

If f is symmetric, which is the case for  $f \in \{f_1, f_2\}$ , then G[f] is symmetric as well. If f(x+c, y+c) = f(x, y) holds for any  $c \in \mathbb{R}^d$ , which is also the case for  $f \in \{f_1, f_2\}$ , then

$$G[f](x,x) = \iint f(x+hu, x+hv)K(u)K(v) \,\mathrm{d} u \,\mathrm{d} v = \iint f(hu,hv)K(u)K(v) \,\mathrm{d} u \,\mathrm{d} v,$$

which does not depend on x. For simplicity, we denote  $\tau_{G[f]} = G[f](x, x)$  for any  $x \in \mathbb{R}^d$ .

**Lemma F.1.** Suppose that  $P_0$  possesses uniformly bounded density. If  $f : \mathbb{R}^d \times \mathbb{R}^d \mapsto \mathbb{R}$  satisfies  $f(x + c, y + c) = f(x, y) \in \mathbb{R}$  for all  $x, y, c \in \mathbb{R}^d$ , and  $\int [K(x)]^2 dx < \infty$ , then  $\|G[f]\|_{L_1(P_0 \times P_0)} \lesssim \|f\|_{L_1(\lambda \times P_0)}$ ,  $\|G[f]\|_{L_2(P_0 \times P_0)} \lesssim \|f\|_{L_2(\lambda \times P_0)}$ , and  $\int \left[\int G[f](x, y) dP_0(x)\right]^2 dP_0(y) \lesssim \int \left[\int |f(x, y)| dx\right]^2 dP_0(y)$ , where the constants depend on K and  $P_0$ . For  $f \in \{f_1, f_2\}$ ,  $\|f\|_{L_1(\lambda \times P_0)} = O(1)$ ,  $\|f\|_{L_2(\lambda \times \nu)}^2 = O(\nu(\mathbb{R}^d)h^{-d})$ , and  $\int \left[\int |f(x, y)| dx\right]^2 d\nu(y) \lesssim \nu(\mathbb{R}^d)$  for a constant only depending on K and any finite measure  $\nu$ .

**Proof of Lemma F.1.** By the property of f, the boundedness of the density of  $P_0$ , and the assumption

that  $\int K^2 < \infty$ ,

$$\begin{split} \|G[f]\|_{L_{1}(P_{0}\times P_{0})} &= \iint |G[f](x,y)| \, \mathrm{d}P_{0}(x) \, \mathrm{d}P_{0}(y) \\ &= \iint \left| \iint f(x+sh,y+th)K(s)K(t) \, \mathrm{d}s \, \mathrm{d}t \right| \, \mathrm{d}P_{0}(x) \, \mathrm{d}P_{0}(y) \\ &\leq \iint \left[ \iint |f(x+sh,y+th)| \, \mathrm{d}P_{0}(x) \, \mathrm{d}P_{0}(y) \right] |K(s)K(t)| \, \mathrm{d}s \, \mathrm{d}t \\ &= \iint \left[ \iint |f(x+sh-th,y)| \, \mathrm{d}P_{0}(x) \, \mathrm{d}P_{0}(y) \right] |K(s)K(t)| \, \mathrm{d}s \, \mathrm{d}t \\ &\lesssim \iint \left[ \iint |f(x',y)| \, \mathrm{d}x' \, \mathrm{d}P_{0}(y) \right] |K(s)K(t)| \, \mathrm{d}s \, \mathrm{d}t \\ &= \|f\|_{L_{1}(\lambda \times P_{0})} \left[ \int |K(s)| \, \mathrm{d}s \right]^{2} \\ &\leq \|f\|_{L_{1}(\lambda \times P_{0})} \int [K(s)]^{2} \, \mathrm{d}s \\ &\lesssim \|f\|_{L_{1}(\lambda \times P_{0})} \, . \end{split}$$

Similarly,

$$\begin{split} \|G[f]\|_{L_{2}(P_{0}\times P_{0})}^{2} &= \iint |G[f](x,y)|^{2} \, \mathrm{d}P_{0}(x) \, \mathrm{d}P_{0}(y) \\ &= \iint \left| \iint f(x+sh,y+th)K(s)K(t) \, \mathrm{d}s \, \mathrm{d}t \right|^{2} \, \mathrm{d}P_{0}(x) \, \mathrm{d}P_{0}(y) \\ &\leq \iint \left[ \iint |f(x+sh,y+th)K(s)K(t)|^{2} \, \mathrm{d}P_{0}(x) \, \mathrm{d}P_{0}(y) \right] \, \mathrm{d}s \, \mathrm{d}t \\ &= \iint \left[ \iint |f(x+sh-th,y)|^{2} \, \mathrm{d}P_{0}(x) \, \mathrm{d}P_{0}(y) \right] \left[ K(s)K(t) \right]^{2} \, \mathrm{d}s \, \mathrm{d}t \\ &\lesssim \iint \left[ \iint |f(x',y)|^{2} \, \mathrm{d}x' \, \mathrm{d}P_{0}(y) \right] \left[ K(s)K(t) \right]^{2} \, \mathrm{d}s \, \mathrm{d}t \\ &= \|f\|_{L_{2}(\lambda \times P_{0})}^{2} \left[ \int \{K(s)\}^{2} \, \mathrm{d}s \right]^{2} \\ &\lesssim \|f\|_{L_{2}(\lambda \times P_{0})}^{2} \, . \end{split}$$

Finally,

$$\int \left[ \int G[f](x,y) \, \mathrm{d}P_0(x) \right]^2 \, \mathrm{d}P_0(y) = \int \left[ \iiint f(x+sh,y+th)K(s)K(t) \, \mathrm{d}s \, \mathrm{d}t \, \mathrm{d}P_0(x) \right]^2 \, \mathrm{d}P_0(y)$$

$$\leq \iint \left\{ \int \left[ \int f(x+sh,y+th) \, \mathrm{d}P_0(x) \right]^2 \, \mathrm{d}P_0(y) \right\} [K(s)K(t)]^2 \, \mathrm{d}s \, \mathrm{d}t \, \mathrm{d}P_0(y)$$

$$= \iint \left\{ \int \left[ \int f(x+sh-th,y) \, \mathrm{d}P_0(x) \right]^2 \, \mathrm{d}P_0(y) \right\} [K(s)K(t)]^2 \, \mathrm{d}s \, \mathrm{d}t$$
  
$$\lesssim \iint \left\{ \int \left[ \int |f(x+sh-th,y)| \, \mathrm{d}x \right]^2 \, \mathrm{d}P_0(y) \right\} [K(s)K(t)]^2 \, \mathrm{d}s \, \mathrm{d}t$$
  
$$= \left\{ \int \left[ \int |f(x',y)| \, \mathrm{d}x' \right]^2 \, \mathrm{d}P_0(y) \right\} \left\{ \iint [K(s)K(t)]^2 \, \mathrm{d}s \, \mathrm{d}t \right\}$$
  
$$\lesssim \int \left[ \int |f(x',y)| \, \mathrm{d}x' \right]^2 \, \mathrm{d}P_0(y).$$

Next, we note that

$$\begin{split} \|f_1\|_{L_1(\lambda \times P_0)} &= \iint \frac{1}{h^d} \left| K\left(\frac{x-y}{h}\right) \right| \, \mathrm{d}x \, \mathrm{d}P_0(y) \\ &= \iint \left| K(x') \right| \, \mathrm{d}x' \, \mathrm{d}P_0(y) \\ &= \int \left| K(x') \right| \, \mathrm{d}x', \text{ and} \\ \|f_2\|_{L_1(\lambda \times P_0)} &= \iint \frac{1}{h^{2d}} \left| \int K\left(\frac{x-z}{h}\right) K\left(\frac{y-z}{h}\right) \, \mathrm{d}z \right| \, \mathrm{d}x \, \mathrm{d}P_0(y) \\ &\leq \iint \left| K\left(x'\right) K\left(y'\right) \right| \left[ \int \eta_0(z+hy') \, \mathrm{d}z \right] \, \mathrm{d}x' \, \mathrm{d}y' \\ &= \left[ \int \left| K\left(x'\right) \right| \, \mathrm{d}x' \right]^2. \end{split}$$

Both of these are finite constants depending only on K. Next,

$$\begin{split} \int \left[ \int |f_1(x,y)| \, \mathrm{d}x \right]^2 \, \mathrm{d}\nu(y) &= \int \left[ \frac{1}{h^d} \int \left| K\left(\frac{x-y}{h}\right) \right| \, \mathrm{d}x \right]^2 \, \mathrm{d}\nu(y) \\ &= \int \left[ \int |K\left(x'\right)| \, \mathrm{d}x' \right]^2 \, \mathrm{d}\nu(y) \\ &= \left[ \int |K\left(x'\right)| \, \mathrm{d}x' \right]^2 \nu(\mathbb{R}^d), \text{ and} \\ \int \left[ \int |f_2(x,y)| \, \mathrm{d}x \right]^2 \, \mathrm{d}\nu(y) &= \int \left[ \int \left| \frac{1}{h^{2d}} \int K\left(\frac{x-z}{h}\right) K\left(\frac{y-z}{h}\right) \, \mathrm{d}z \right| \, \mathrm{d}x \right]^2 \, \mathrm{d}\nu(y) \\ &\leq \int \left[ \frac{1}{h^{2d}} \iint \left| K\left(\frac{x-z}{h}\right) K\left(\frac{y-z}{h}\right) \right| \, \mathrm{d}z \, \mathrm{d}x \right]^2 \, \mathrm{d}\nu(y) \\ &= \int \left[ \iint |K\left(x'\right)| \, \mathrm{d}x' \right]^4 \nu(\mathbb{R}^d). \end{split}$$

Similarly,

$$\begin{split} \|f_1\|_{L_2(\lambda \times \nu)}^2 &= \frac{1}{h^{2d}} \iint \left[ K\left(\frac{x-y}{h}\right) \right]^2 \, \mathrm{d}x \, \mathrm{d}\nu(y) \\ &= \frac{1}{h^d} \iint \left[ K\left(x'\right) \right]^2 \, \mathrm{d}x' \, \mathrm{d}\nu(y) \\ &= \frac{1}{h^d} \|K\|_{L_2(\lambda)}^2 \nu(\mathbb{R}^d), \end{split}$$

and

$$\begin{split} \|f_{2}\|_{L_{2}(\lambda\times\nu)}^{2} &= \iint \left[\frac{1}{h^{2d}} \int K\left(\frac{x-z}{h}\right) K\left(\frac{y-z}{h}\right) dz\right]^{2} dx \, d\nu(y) \\ &= \frac{1}{h^{4d}} \iiint K\left(\frac{x-z}{h}\right) K\left(\frac{y-z}{h}\right) K\left(\frac{x-w}{h}\right) K\left(\frac{y-w}{h}\right) dz \, dw \, dx \, d\nu(y) \\ &= \frac{1}{h^{2d}} \iiint K\left(z'\right) K\left(z'\right) K\left(z'+\frac{y-x}{h}\right) K\left(w'\right) K\left(w'+\frac{y-x}{h}\right) dz' \, dw' \, dx \, d\nu(y) \\ &= \frac{1}{h^{d}} \iiint K\left(z'\right) K\left(x'\right) K\left(w'\right) K\left(w'+x'-z'\right) \, dz' \, dw' \, dx' \, d\nu(y) \\ &= \frac{1}{h^{d}} \nu(\mathbb{R}^{d}) \iiint K\left(z'\right) K\left(z'\right) K\left(x'\right) \left[\int \left\{K\left(w'\right)\right\}^{2} \, dw'\right]^{1/2} \left[\int \left\{K\left(w'+x'-z'\right)\right\}^{2} \, dw'\right]^{1/2} \, dz' \, dx' \, dx'$$

which yields the result.

**Corollary F.2.** If  $\hat{P}_n$  is the distribution corresponding to a kernel density estimator with bandwidth h and kernel K satisfying  $\int K^2 < \infty$ ,  $P_0$  possesses uniformly bounded density, and  $f : \mathbb{R}^d \times \mathbb{R}^d \mapsto \mathbb{R}$ satisfies f(x, y) = f(y, x),  $f(x, x) = \tau_f \in \mathbb{R}$  and  $f(x + c, y + c) = f(x, y) \in \mathbb{R}$  for all  $x, y, c \in \mathbb{R}^d$ , then

$$E_0 \left\{ \int f \, \mathrm{d} \left[ (\hat{P}_n - P_{0,h}) \times (\hat{P}_n - P_{0,h}) \right] - n^{-1} \tau_{G[f]} \right\}^2 \lesssim n^{-3} \int \left[ \int |f(x,y)| \, \mathrm{d}x \right]^2 \, \mathrm{d}P_0(y) + n^{-2} \, \|f\|_{L_2(\lambda \times P_0)}^2$$

for a constant depending on K and  $P_0$ .

**Proof of Corollary F.2.** Recall that  $G[f](x,y) := \int f(s,t)K_h(s,x)K_h(t,y)\,\mathrm{d}s\,\mathrm{d}t$ . Under the as-

sumptions on f, we have G[f](x,y) = G[f](y,x) and  $G[f](x,x) = \tau_{G[f]}$  for all  $x,y \in \mathbb{R}^d$ . Since  $(\hat{p}_n - \eta_{0,h})(x) = \int K_h(s,x) d(\mathbb{P}_n - P_0)(s)$ , we can write

$$\int f \operatorname{d}\left[(\hat{P}_n - P_{0,h}) \times (\hat{P}_n - P_{0,h})\right] = \int G[f] \operatorname{d}\left[(\mathbb{P}_n - P_0) \times (\mathbb{P}_n - P_0)\right]$$

By Lemma E.2, we then have

$$E_0 \left\{ \int f \, \mathrm{d} \left[ (\hat{P}_n - P_{0,h}) \times (\hat{P}_n - P_{0,h}) \right] - n^{-1} \tau_{G[f]} \right\}^2 = E_0 \left\{ \int G[f] \, \mathrm{d} \left[ (\mathbb{P}_n - P_0) \times (\mathbb{P}_n - P_0) \right] - n^{-1} \tau_{G[f]} \right\}^2$$
$$\lesssim n^{-3} \int \left[ \int G[f](x, y) \, \mathrm{d}P_0(x) \right]^2 \, \mathrm{d}P_0(y)$$
$$+ n^{-2} \left\| G[f] \right\|_{L_2(P_0 \times P_0)}^2.$$

The result follows by Lemma F.1.

**Lemma F.3.** If  $P_0$  possesses uniformly bounded and m-times continuously differentiable density  $\eta_0$ with  $\int [(D^{\alpha}\eta_0)(x)]^2 dx < \infty$  for all  $\alpha$  such that  $|\alpha| = m$ ,  $\hat{P}_n$  is the distribution corresponding to a kernel density estimator with bandwidth h and kernel K satisfying  $\int K^2 < \infty$ , and  $f : \mathbb{R}^d \times \mathbb{R}^d \to \mathbb{R}$  satisfies f(x,y) = f(y,x) and  $f(x,x) = \tau_f$  for all  $x, y \in \mathbb{R}^d$ , then

$$E_0 E_W \left\{ \int f \, \mathrm{d} \left[ (P_{0,h} - P_0) \times (\mathbb{P}_n^* - \hat{P}_n) \right] \right\}^2 \lesssim n^{-1} h^{2m} \, \|f\|_{L_2(\lambda \times P_{0,h})}^2$$

**Proof of Lemma F.3.** We denote  $f^{\circ}(y) := \int f(x,y) d(P_{0,h} - P_0)(x)$ . We then have

$$\int f d[(P_{0,h} - P_0) \times (\mathbb{P}_n^* - \hat{P}_n)] = (\mathbb{P}_n^* - \hat{P}_n)f^{\circ} = \frac{1}{n} \sum_{i=1}^n \left[ f^{\circ}(X_i^*) - \hat{P}_n f^{\circ} \right].$$

Since  $X_1^*, \ldots, X_n^* \stackrel{\text{iid}}{\sim} \hat{P}_n, E_W \left[ f^{\circ}(X_i^*) - \hat{P}_n f^{\circ} \right] \left[ f^{\circ}(X_j^*) - \hat{P}_n f^{\circ} \right] = 0$  for any  $i \neq j$ , so

$$E_{W}\left\{\frac{1}{n}\sum_{i=1}^{n}\left[f^{\circ}(X_{i}^{*})-\hat{P}_{n}f^{\circ}\right]\right\}^{2}=E_{W}\left\{\frac{1}{n^{2}}\sum_{i,j=1}^{n}\left[f^{\circ}(X_{i}^{*})-\hat{P}_{n}f^{\circ}\right]\left[f^{\circ}(X_{j}^{*})-\hat{P}_{n}f^{\circ}\right]\right\}$$
$$=E_{W}\left\{\frac{1}{n^{2}}\sum_{i=1}^{n}\left[f^{\circ}(X_{i}^{*})-\hat{P}_{n}f^{\circ}\right]^{2}\right\}$$
$$=\frac{1}{n}E_{W}\left[f^{\circ}(X_{i}^{*})-\hat{P}_{n}f^{\circ}\right]^{2}$$
$$=\frac{1}{n}\left[\hat{P}_{n}(f^{\circ})^{2}-(\hat{P}_{n}f^{\circ})^{2}\right]$$

$$\leq \frac{1}{n}\hat{P}_n(f^\circ)^2.$$

By Lemma E.6, we have

$$\begin{split} E_0 \left[ n^{-1} \hat{P}_n(f^\circ)^2 \right] &= n^{-1} P_{0,h}(f^\circ)^2 \\ &= n^{-1} \int \left[ \int f(x,y) [\eta_{0,h}(x) - \eta_0(x)] \, \mathrm{d}x \right]^2 \, \mathrm{d}P_{0,h}(y) \\ &\leq n^{-1} \int \left[ \int |f(x,y)| \, |\eta_{0,h}(x) - \eta_0(x)| \, \, \mathrm{d}x \right]^2 \, \mathrm{d}P_{0,h}(y) \\ &\leq n^{-1} \iint |f(x,y)|^2 \, \mathrm{d}x \, \mathrm{d}P_{0,h}(y) \int [\eta_{0,h}(x) - \eta_0(x)]^2 \, \mathrm{d}x \\ &\lesssim n^{-1} h^{2m} \, \|f\|_{L_2(\lambda \times P_{0,h})}^2 \, . \end{split}$$

**Lemma F.4.** If  $\hat{P}_n$  is the distribution corresponding to a kernel density estimator with bandwidth h and uniformly bounded kernel K,  $P_0$  possesses uniformly bounded density, then for any fixed  $f : \mathbb{R}^d \times \mathbb{R}^d \mapsto \mathbb{R}$ ,

$$E_0 E_W \left\{ \int f \, \mathrm{d}[(\hat{P}_n - P_{0,h}) \times (\mathbb{P}_n^* - \hat{P}_n)] \right\}^2 \lesssim n^{-1} \left[ (nh^d)^{-1} + (nh^d)^{-2} \right] \int \left[ \int |f(x,y)| \, \mathrm{d}x \right]^2 \, \mathrm{d}P_{0,h}(y),$$

where the constant in the bound depends on  $P_0$  and K.

**Proof of Lemma F.4.** We note that

$$\begin{split} E_W \left\{ \int f \, \mathrm{d}[(\hat{P}_n - P_{0,h}) \times (\mathbb{P}_n^* - \hat{P}_n)] \right\}^2 &= E_W \left\{ \int \left[ \int f(x,z) \, \mathrm{d}(\hat{P}_n - P_{0,h})(x) \right] \, \mathrm{d}(\mathbb{P}_n^* - \hat{P}_n)(z) \right\}^2 \\ &= \frac{1}{n} \mathrm{Var}_W \left( \int f(x,Z) \, \mathrm{d}(\hat{P}_n - P_{0,h})(x) \right) \\ &\leq \frac{1}{n} E_W \left[ \int f(x,Z) \, \mathrm{d}(\hat{P}_n - P_{0,h})(x) \right]^2 \\ &= \frac{1}{n} \int \left[ \int f(x,z) \, \mathrm{d}(\hat{P}_n - P_{0,h})(x) \right]^2 \, \mathrm{d}\hat{P}_n(z) \\ &= \frac{1}{n} \int \left[ \int f(x,z) \, \mathrm{d}(\hat{P}_n - P_{0,h})(x) \right]^2 \, \mathrm{d}P_{0,h}(z) \\ &\quad + \frac{1}{n} \int \left[ \int f(x,z) \, \mathrm{d}(\hat{P}_n - P_{0,h})(x) \right]^2 \, \mathrm{d}(\hat{P}_n - P_{0,h})(z). \end{split}$$

We bound the two terms in the final expression separately. We can write

$$E_{0} \left\{ \int \left[ \int f(x,z) d(\hat{P}_{n} - P_{0,h})(x) \right]^{2} dP_{0,h}(z) \right\}$$
  
=  $E_{0} \left\{ \iiint f(x,z) f(y,z) d(\hat{P}_{n} - P_{0,h})(x) d(\hat{P}_{n} - P_{0,h})(y) dP_{0,h}(z) \right\}$   
=  $E_{0} \left\{ \iiint f(x,z) f(y,z) (\hat{p}_{n} - \eta_{0,h})(x) (\hat{p}_{n} - \eta_{0,h})(y) dx dy dP_{0,h}(z) \right\}$   
 $\leq \iiint |f(x,z) f(y,z)| \left\{ E_{0} [\hat{p}_{n}(x) - \eta_{0,h}(x)]^{2} \right\}^{1/2} \left\{ E_{0} [\hat{p}_{n}(y) - \eta_{0,h}(y)]^{2} \right\}^{1/2} dx dy dP_{0,h}(z)$ 

Since  $\eta_0$  is uniformly bounded and  $\int K^2 < \infty$ ,

$$E_0 \left[ \hat{p}_n(x) - \eta_{0,h}(x) \right]^2 = E_0 \left[ \int K_h(s,x) \, \mathrm{d}(\mathbb{P}_n - P_0)(s) \right]^2$$
$$= \frac{1}{n} \operatorname{Var}_0 \left[ K_h(X,x) \right]$$
$$\leq \frac{1}{nh^{2d}} E_0 \left[ \left\{ K \left( \frac{X-x}{h} \right) \right\}^2 \right]$$
$$= \frac{1}{nh^{2d}} \int \left\{ K \left( \frac{t-x}{h} \right) \right\}^2 \eta_0(t) \, \mathrm{d}t$$
$$= \frac{1}{nh^d} \int \left\{ K(u) \right\}^2 \eta_0(x+hu) \, \mathrm{d}u$$
$$\lesssim \frac{1}{nh^d}$$

for each  $x \in \mathbb{R}^d$ . Thus,

$$E_0\left\{\int \left[\int f(x,z) \operatorname{d}(\hat{P}_n - P_{0,h})(x)\right]^2 \operatorname{d}P_{0,h}(z)\right\} \lesssim \frac{1}{nh^d} \iiint |f(x,z)f(y,z)| \operatorname{d}x \operatorname{d}y \operatorname{d}P_{0,h}(z)$$
$$= \frac{1}{nh^d} \int \left[\int |f(x,z)| \operatorname{d}x\right]^2 \operatorname{d}P_{0,h}(z).$$

Next, we note that for each  $x, y, z \in \mathbb{R}^d$ , since  $X_1, \ldots, X_n$  are IID,

$$E_{0} \{ [\hat{p}_{n}(x) - \eta_{0,h}(x)] [\hat{p}_{n}(y) - \eta_{0,h}(y)] [\hat{p}_{n}(z) - \eta_{0,h}(z)] \}$$

$$= E_{0} \left[ \int K_{h}(s,x) d(\mathbb{P}_{n} - P_{0})(s) \int K_{h}(t,y) d(\mathbb{P}_{n} - P_{0})(t) \int K_{h}(u,z) d(\mathbb{P}_{n} - P_{0})(u) \right]$$

$$= E_{0} \left\{ \frac{1}{n^{3}} \sum_{i,j,k} [K_{h}(X_{i},x) - \eta_{0,h}(x)] [K_{h}(X_{j},y) - \eta_{0,h}(y)] [K_{h}(X_{k},z) - \eta_{0,h}(z)] \right\}$$

$$= E_0 \left\{ \frac{1}{n^3} \sum_{i=1}^n \left[ K_h(X_i, x) - \eta_{0,h}(x) \right] \left[ K_h(X_i, y) - \eta_{0,h}(y) \right] \left[ K_h(X_i, z) - \eta_{0,h}(z) \right] \right\}$$
$$= \frac{1}{n^2} E_0 \left\{ \left[ K_h(X, x) - \eta_{0,h}(x) \right] \left[ K_h(X, y) - \eta_{0,h}(y) \right] \left[ K_h(X, z) - \eta_{0,h}(z) \right] \right\}.$$

Hence,

$$\begin{split} E_0 \left\{ \int \left[ \int f(x,z) \,\mathrm{d}(\hat{P}_n - P_{0,h})(x) \right]^2 \,\mathrm{d}(\hat{P}_n - P_{0,h})(z) \right\} \\ &= E_0 \left\{ \iiint f(x,z) f(y,z) \left[ \hat{p}_n(x) - \eta_{0,h}(x) \right] \left[ \hat{p}_n(y) - \eta_{0,h}(y) \right] \left[ \hat{p}_n(z) - \eta_{0,h}(z) \right] \,\mathrm{d}x \,\mathrm{d}y \,\mathrm{d}z \right\} \\ &= \frac{1}{n^2} E_0 \iiint f(x,z) f(y,z) \left[ K_h(X,x) - \eta_{0,h}(x) \right] \left[ K_h(X,y) - \eta_{0,h}(y) \right] \left[ K_h(X,z) - \eta_{0,h}(z) \right] \,\mathrm{d}x \,\mathrm{d}y \,\mathrm{d}z \\ &= \frac{1}{n^2} E_0 \iint \left\{ \int f(x,z) \left[ K_h(X,x) - \eta_{0,h}(x) \right] \,\mathrm{d}x \right\}^2 \left[ K_h(X,z) - \eta_{0,h}(z) \right] \,\mathrm{d}z. \end{split}$$

Since K is uniformly bounded, we have  $|K_h(x,y) - \eta_{0,h}(x)| \leq h^{-d}$  for any  $x, y \in \mathbb{R}^d$ . Therefore,

$$\begin{aligned} \frac{1}{n^2} E_0 \int \left\{ \int f(x,z) \left[ K_h(X,x) - \eta_{0,h}(x) \right] \, \mathrm{d}x \right\}^2 \left[ K_h(X,z) - \eta_{0,h}(z) \right] \, \mathrm{d}z \right| \\ & \leq \left| \frac{1}{n^2} E_0 \int \left\{ \int f(x,z) \left[ K_h(X,x) - \eta_{0,h}(x) \right] \, \mathrm{d}x \right\}^2 K_h(X,z) \, \mathrm{d}z \right| \\ & + \left| \frac{1}{n^2} E_0 \int \left\{ \int f(x,z) \left[ K_h(X,x) - \eta_{0,h}(x) \right] \, \mathrm{d}x \right\}^2 \eta_{0,h}(z) \, \mathrm{d}z \right| \\ & \lesssim \frac{1}{n^2 h^{2d}} \left| E_0 \int \left\{ \int \left| f(x,z) \right| \, \mathrm{d}x \right\}^2 K_h(X,z) \, \mathrm{d}z \right| + \frac{1}{n^2 h^{2d}} \left| \int \left\{ \int \left| f(x,z) \right| \, \mathrm{d}x \right\}^2 \eta_{0,h}(z) \, \mathrm{d}z \right| \\ & = \frac{2}{n^2 h^{2d}} \int \left\{ \int \left| f(x,z) \right| \, \mathrm{d}x \right\}^2 \, \mathrm{d}P_{0,h}(z). \end{aligned}$$

Putting the pieces together completes the result.

**Lemma F.5.** If  $\hat{P}_n$  is the distribution corresponding to a kernel density estimator with bandwidth h and uniformly bounded kernel K,  $P_0$  possesses a Lebesgue density function, and  $f : \mathbb{R}^d \times \mathbb{R}^d \mapsto \mathbb{R}$  satisfies f(x,y) = f(y,x) for all  $x, y \in \mathbb{R}^d$  then

$$E_0 E_W \left\{ \int f \, d[(\mathbb{P}_n - P_0) \times (\mathbb{P}_n^* - \hat{P}_n)] \right\}^2 \lesssim \frac{1}{n^2} \|f\|_{L_2(P_0 \times P_{0,h})}^2 + \frac{1}{n^3 h^d} \|f\|_{L_2(\lambda \times P_0)}^2,$$

where the constant in the bound depends on K only.

**Proof of Lemma F.5.** As in the proof of Lemma F.4, we have

$$E_{W} \left\{ \int f d[(\mathbb{P}_{n} - P_{0}) \times (\mathbb{P}_{n}^{*} - \hat{P}_{n})] \right\}^{2} \leq \frac{1}{n} \int \left[ \int f(x, z) d(\mathbb{P}_{n} - P_{0})(x) \right]^{2} dP_{0,h}(z) + \frac{1}{n} \int \left[ \int f(x, z) d(\mathbb{P}_{n} - P_{0})(x) \right]^{2} d(\hat{P}_{n} - P_{0,h})(z).$$

We first bound the first term. We have

$$E_0 \left\{ \int \left[ \int f(x,z) \, \mathrm{d}(\mathbb{P}_n - P_0)(x) \right]^2 \, \mathrm{d}P_{0,h}(z) \right\} = \int E_0 \left[ \int f(x,z) \, \mathrm{d}(\mathbb{P}_n - P_0)(x) \right]^2 \, \mathrm{d}P_{0,h}(z)$$
$$= \frac{1}{n} \int \operatorname{Var}_0(f(X,z)) \, \mathrm{d}P_{0,h}(z)$$
$$\leq \frac{1}{n} \int E_0 \left[ f(X,z) \right]^2 \, \mathrm{d}P_{0,h}(z)$$
$$= \frac{1}{n} \|f\|_{L_2(P_0 \times P_{0,h})}^2.$$

Next, we note that

$$E_{0}\left\{ \int \left[ \int f(x,z) d(\mathbb{P}_{n} - P_{0})(x) \right]^{2} d(\hat{P}_{n} - P_{0,h})(z) \right\}$$

$$= E_{0}\left\{ \int \int f(x,z) d(\mathbb{P}_{n} - P_{0})(x) \int f(y,z) d(\mathbb{P}_{n} - P_{0})(y) \int K_{h}(w,z) d(\mathbb{P}_{n} - P_{0})(w) dz \right\}$$

$$= \int E_{0}\left\{ \frac{1}{n^{3}} \sum_{i,j,k} \left[ f(X_{i},z) - P_{0}f(\cdot,z) \right] \left[ f(X_{j},z) - P_{0}f(\cdot,z) \right] \left[ K_{h}(X_{k},z) - \eta_{0,h}(z) \right] \right\} dz$$

$$= \int E_{0}\left\{ \frac{1}{n^{3}} \sum_{i=1}^{n} \left[ f(X_{i},z) - P_{0}f(\cdot,z) \right]^{2} \left[ K_{h}(X_{i},z) - \eta_{0,h}(z) \right] \right\} dz$$

$$= \frac{1}{n^{2}} \int E_{0}\left\{ \left[ f(X,z) - P_{0}f(\cdot,z) \right]^{2} \left[ K_{h}(X,z) - \eta_{0,h}(z) \right] \right\} dz.$$

Since K is uniformly bounded, we have  $|K_h(x,z) - \eta_{0,h}(z)| \leq h^{-d}$  for all  $x, z \in \mathbb{R}^d$ . Thus,

$$\left| E_0 \left\{ \int \left[ \int f(x,z) \, \mathrm{d}(\mathbb{P}_n - P_0)(x) \right]^2 \, \mathrm{d}(\hat{P}_n - P_{0,h})(z) \right\} \right|$$
  
$$\lesssim \frac{1}{n^2 h^d} \int E_0 \left[ f(X,z) - P_0 f(\cdot,z) \right]^2 \, \mathrm{d}z$$
  
$$\leq \frac{1}{n^2 h^d} \int E_0 \left[ f(X,z) \right]^2 \, \mathrm{d}z$$
  
$$= \frac{1}{n^2 h^d} \left\| f \right\|_{L_2(\lambda \times P_0)}^2.$$

**Lemma F.6.** If  $\hat{P}_n$  is the distribution corresponding to a kernel density estimator with bandwidth hand uniformly bounded kernel K satisfying  $\int K^2 < \infty$ ,  $P_0$  possesses uniformly bounded density, and  $f : \mathbb{R}^d \times \mathbb{R}^d \mapsto \mathbb{R}$  satisfies f(x,y) = f(y,x),  $f(x,x) = \tau_f \in \mathbb{R}$  and f(x+c,y+c) = f(x,y) for all  $x, y, c \in \mathbb{R}^d$ , then

$$E_{0}E_{W}\left\{\int f d[(\mathbb{P}_{n}^{*}-\hat{P}_{n})\times(\mathbb{P}_{n}^{*}-\hat{P}_{n})]-n^{-1}\tau_{f}\right\}^{2}$$

$$\lesssim \frac{\tau_{G[f]}^{2}}{n^{4}}+\frac{\tau_{G[f^{2}]}}{n^{3}}+\frac{\tau_{G[f]}}{n^{3}}\|f\|_{L_{1}(\lambda\times P_{0})}+\frac{1}{n^{2}}\|f\|_{L_{2}(\lambda\times P_{0})}^{2}+\frac{1}{n^{3}}\int \left[\int |f(x,y)| dx\right]^{2} dP_{0}(y)$$

$$+\frac{1}{n^{3}}\left[1+(nh^{d})^{-1}+(nh^{d})^{-2}\right]\int \left(\int |f(x,y)| dx\right)^{2} dP_{0,h}(y)$$

**Proof of Lemma F.6.** We define  $V_n(x,y) := \int f d\left[ (\delta_x - \hat{P}_n) \times (\delta_y - \hat{P}_n) \right]$ . We then have

$$E_{0}E_{W}\left\{\int f d[(\mathbb{P}_{n}^{*}-\hat{P}_{n})\times(\mathbb{P}_{n}^{*}-\hat{P}_{n})]-\frac{\tau_{f}}{n}\right\}^{2}$$

$$=E_{0}E_{W}\left\{\frac{1}{n^{2}}\sum_{i=1}^{n}V_{n}(X_{i}^{*},X_{i}^{*})+\frac{1}{n^{2}}\sum_{i\neq j}V_{n}(X_{i}^{*},X_{j}^{*})-\frac{\tau_{f}}{n}\right\}^{2}$$

$$\leq 2E_{0}E_{W}\left\{\frac{1}{n^{2}}\sum_{i=1}^{n}V_{n}(X_{i}^{*},X_{i}^{*})-\frac{\tau_{f}}{n}\right\}^{2}+2E_{0}E_{W}\left\{\frac{1}{n^{2}}\sum_{i\neq j}V_{n}(X_{i}^{*},X_{j}^{*})\right\}^{2}.$$
(31)

For the first term on the right-hand side of (31), by definition of  $\tau_f$ , symmetry of f, and adding and subtracting terms,

$$\begin{split} \frac{1}{n^2} \sum_{i=1}^n V_n(X_i^*, X_i^*) &- \frac{\tau_f}{n} \\ &= \frac{1}{n^2} \sum_{i=1}^n \int f \, \mathrm{d}[(\delta_{X_i^*} - \hat{P}_n) \times (\delta_{X_i^*} - \hat{P}_n)] - \frac{\tau_f}{n} \\ &= -\frac{2}{n} \int f \, \mathrm{d}(\hat{P}_n \times \mathbb{P}_n^*) + \frac{1}{n} \int f \, \mathrm{d}(\hat{P}_n \times \hat{P}_n) \\ &= -\frac{2}{n} \int f \, \mathrm{d}[\hat{P}_n \times (\mathbb{P}_n^* - \hat{P}_n)] - \frac{1}{n} \int f \, \mathrm{d}(\hat{P}_n \times \hat{P}_n) \\ &= -\frac{2}{n} \int f \, \mathrm{d}[\hat{P}_n \times (\mathbb{P}_n^* - \hat{P}_n)] - \frac{1}{n} \int f \, \mathrm{d}(\hat{P}_n \times \hat{P}_n) \\ &= -\frac{2}{n} \int f \, \mathrm{d}[(\hat{P}_n - P_{0,h}) \times (\mathbb{P}_n^* - \hat{P}_n)] - \frac{2}{n} \int f \, \mathrm{d}[P_{0,h} \times (\mathbb{P}_n^* - \hat{P}_n)] - \frac{1}{n} \int f \, \mathrm{d}(\hat{P}_n \times \hat{P}_n). \end{split}$$

Hence,

$$E_{0}E_{W}\left\{\frac{1}{n^{2}}\sum_{i=j}V_{n}(X_{i}^{*},X_{j}^{*})-\frac{\tau_{f}}{n}\right\}^{2}$$

$$\lesssim E_{0}E_{W}\left\{\frac{1}{n}\int f\,\mathrm{d}[(\hat{P}_{n}-P_{0,h})\times(\mathbb{P}_{n}^{*}-\hat{P}_{n})]\right\}^{2}+E_{0}E_{W}\left\{\frac{1}{n}\int f\,\mathrm{d}[P_{0,h}\times(\mathbb{P}_{n}^{*}-\hat{P}_{n})]\right\}^{2} \quad (32)$$

$$+E_{0}\left\{\frac{1}{n}\int f\,\mathrm{d}(\hat{P}_{n}\times\hat{P}_{n})\right\}^{2}.$$

By Lemma F.4, the first term on the right-hand side of (32) is bounded up to a constant by

$$n^{-3} \left[ (nh^d)^{-1} + (nh^d)^{-2} \right] \int \left[ \int |f(x,y)| \, \mathrm{d}x \right]^2 \, \mathrm{d}P_{0,h}(y).$$

For the second term on the right-hand side of (32),

$$\begin{split} E_0 E_W \left\{ \frac{1}{n} \int f \, \mathrm{d}[P_{0,h} \times (\mathbb{P}_n^* - \hat{P}_n)] \right\}^2 &= \frac{1}{n^3} E_0 \left[ \operatorname{Var}_W \left( \int f(x, X_1^*) \, \mathrm{d}P_{0,h}(x) \right) \right] \\ &\leq \frac{1}{n^3} E_0 \left[ \int \left( \int f(x, y) \, \mathrm{d}P_{0,h}(x) \right)^2 \, \mathrm{d}\hat{P}_n(y) \right] \\ &= \frac{1}{n^3} \int \left( \int f(x, y) \, \mathrm{d}P_{0,h}(x) \right)^2 \, \mathrm{d}P_{0,h}(y) \\ &\lesssim \frac{1}{n^3} \int \left( \int |f(x, y)| \, \mathrm{d}x \right)^2 \, \mathrm{d}P_{0,h}(y), \end{split}$$

where the last inequality is because K is uniformly bounded. For the last term on the right-hand side of (32), we have

$$E_{0}\left\{\frac{1}{n}\int f d(\hat{P}_{n} \times \hat{P}_{n})\right\}^{2} = E_{0}\left\{\frac{1}{n^{3}}\sum_{i,j}G[f](X_{i}, X_{j})\right\}^{2}$$
$$= \frac{1}{n^{2}}E_{0}\left\{\frac{1}{n^{2}}\sum_{i=1}^{n}G[f](X_{i}, X_{i}) + \frac{1}{n^{2}}\sum_{i\neq j}G[f](X_{i}, X_{j})\right\}^{2}$$
$$= \frac{1}{n^{2}}E_{0}\left\{\frac{\tau_{G}[f]}{n} + \frac{1}{n^{2}}\sum_{i\neq j}G[f](X_{i}, X_{j})\right\}^{2}$$
$$= \frac{1}{n^{2}}\left\{\frac{\tau_{G}^{2}[f]}{n^{2}} + \frac{2\tau_{G}[f]}{n^{3}}\sum_{i\neq j}E_{0}\left[G[f](X_{i}, X_{j})\right] + \frac{1}{n^{4}}E_{0}\left[\sum_{i\neq j}G[f](X_{i}, X_{j})\right]^{2}\right\}$$

$$\lesssim \frac{\tau_{G[f]}^2}{n^4} + \frac{\tau_{G[f]}}{n^3} E_0 \left[ G[f](X_1, X_2) \right] + \frac{1}{n^6} E_0 \left[ \sum_{i \neq j} G[f](X_i, X_j) \right]^2.$$

By Lemma F.1, we have  $|E_0[G[f](X_1, X_2)]| \leq ||f||_{L_1(\lambda \times P_0)}$ . By Lemma E.1 and Lemma F.1, we note that

$$\begin{split} E_0 \left[ \sum_{i \neq j} G[f](X_i, X_j) \right]^2 &= E_0 \left[ 4 \sum_{i \neq j, j \neq k, k \neq i} G[f](X_i, X_j) G[f](X_i, X_k) + 2 \sum_{i \neq j} [G[f](X_i, X_j)]^2 \\ &+ \sum_{\substack{i \neq j, i \neq k, i \neq l \\ j \neq k, j \neq l, k \neq l}} G[f](X_i, X_j) G[f](X_k, X_l) \right] \\ &\lesssim n^3 E_0 \left[ G[f](X_1, X_2) G[f](X_1, X_3) \right] + n^2 E_0 \left[ G[f](X_1, X_2) \right]^2 + n^4 [E_0 G[f](X_1, X_2)]^2 \\ &\leq n^3 \int \left( \int G[f](x, y) \, \mathrm{d}P_0(x) \right)^2 \, \mathrm{d}P_0(y) + n^2 \, \|G[f]\|_{L_2(P_0 \times P_0)}^2 + n^4 \, \|G[f]\|_{L_1(P_0 \times P_0)}^2 \\ &\lesssim n^3 \int \left( \int |f(x, y)| \, \mathrm{d}x \right)^2 \, \mathrm{d}P_0(y) + n^2 \, \|f\|_{L_2(\lambda \times P_0)}^2 + n^4 \, \|f\|_{L_1(\lambda \times P_0)}^2 \, . \end{split}$$

Combining these calculations, we get

$$E_0 \left\{ \frac{1}{n} \int f \,\mathrm{d}(\hat{P}_n \times \hat{P}_n) \right\}^2 \lesssim \frac{\tau_{G[f]}^2}{n^4} + \frac{\tau_{G[f]}}{n^3} \|f\|_{L_1(\lambda \times P_0)} + \frac{1}{n^2} \|f\|_{L_1(\lambda \times P_0)}^2 + \frac{1}{n^4} \|f\|_{L_2(\lambda \times P_0)}^2 + \frac{1}{n^3} \int \left( \int |f(x,y)| \,\mathrm{d}x \right)^2 \,\mathrm{d}P_0(y).$$

Therefore,

$$E_{0}E_{W}\left\{\frac{1}{n^{2}}\sum_{i=j}V_{n}(X_{i}^{*},X_{j}^{*})-\frac{\tau_{f}}{n}\right\}^{2} \lesssim \frac{\tau_{G[f]}^{2}}{n^{4}}+\frac{\tau_{G[f]}}{n^{3}}\|f\|_{L_{1}(\lambda\times P_{0})}+\frac{1}{n^{2}}\|f\|_{L_{1}(\lambda\times P_{0})}^{2}+\frac{1}{n^{4}}\|f\|_{L_{2}(\lambda\times P_{0})}^{2}$$
$$+\frac{1}{n^{3}}\int\left(\int|f(x,y)|\,\mathrm{d}x\right)^{2}\,\mathrm{d}P_{0}(y)$$
$$+\frac{1}{n^{3}}\left[1+(nh^{d})^{-1}+(nh^{d})^{-2}\right]\int\left(\int|f(x,y)|\,\mathrm{d}x\right)^{2}\,\mathrm{d}P_{0,h}(y).$$

For the second term on the right-hand side of (31), we note that for any  $i \neq j$ ,

$$E_W \left[ V_n(X_i^*, X_j^*) \mid X_j^* \right]$$
  
=  $E_W \left[ f(X_i^*, X_j^*) - \int f(X_i^*, y) \, \mathrm{d}\hat{P}_n(y) - \int f(x, X_j^*) \, \mathrm{d}\hat{P}_n(x) + (\hat{P}_n \times \hat{P}_n) f \mid X_j^* \right]$ 

$$= \int f(x, X_j^*) \,\mathrm{d}\hat{P}_n(x) - (\hat{P}_n \times \hat{P}_n)f - \int f(x, X_j^*) \,\mathrm{d}\hat{P}_n(x) + (\hat{P}_n \times \hat{P}_n)f = 0$$

Hence, by the law of total expectation,  $E_W[V_n(X_i^*, X_j^*)] = 0$  for all  $i \neq j$ , and by symmetry of f, which implies symmetry of  $V_n$ ,  $E_W[V_n(X_i^*, X_j^*)V_n(X_i^*, X_k^*)] = 0$  for all  $i \neq j \neq k$ . Thus, by Lemma E.1,

$$\begin{split} E_W \left[ \frac{1}{n^2} \sum_{i \neq j} V_n(X_i^*, X_j^*) \right]^2 \\ &= E_W \left[ \frac{4}{n^4} \sum_{i \neq j, j \neq k, k \neq i} V_n(X_i^*, X_j^*) V_n(X_i^*, X_k^*) + \frac{2}{n^4} \sum_{i \neq j} \left\{ V_n(X_i^*, X_j^*) \right\}^2 \\ &\quad + \frac{1}{n^4} \sum_{\substack{i \neq j, i \neq k, i \neq i \\ j \neq k, j \neq l, k \neq l}} V_n(X_i^*, X_j^*) V_n(X_k^*, X_l^*) \right] \\ &= E_W \left[ \frac{2}{n^4} \sum_{i \neq j} \left\{ V_n(X_i^*, X_j^*) \right\}^2 \right] \\ &\lesssim \frac{1}{n^2} E_W \left[ V_n(X_i^*, X_2^*) \right]^2 \\ &= \frac{1}{n^2} E_W \left[ f(X_1^*, X_2^*) - \int f(X_1^*, y) \, d\hat{P}_n(y) - \int f(x, X_2^*) \, d\hat{P}_n(x) + (\hat{P}_n \times \hat{P}_n) f \right]^2 \\ &= \frac{1}{n^2} Var_W \left[ f(X_1^*, X_2^*) - \int f(X_1^*, y) \, d\hat{P}_n(y) - \int f(x, X_2^*) \, d\hat{P}_n(x) \right] \\ &\leq \frac{1}{n^2} E_W \left[ f(X_1^*, X_2^*) - \int f(X_1^*, y) \, d\hat{P}_n(y) - \int f(x, X_2^*) \, d\hat{P}_n(x) \right]^2 \\ &= \frac{1}{n^2} E_W \left\{ f(X_1^*, X_2^*) - \int f(X_1^*, y) \, d\hat{P}_n(y) \right]^2 + \left[ \int f(x, X_2^*) \, d\hat{P}_n(x) \right]^2 \\ &= \frac{1}{n^2} E_W \left\{ f(X_1^*, X_2^*)^2 + \left[ \int f(X_1^*, y) \, d\hat{P}_n(y) \right]^2 + \left[ \int f(x, X_2^*) \, d\hat{P}_n(x) \right]^2 \\ &\quad + 2 \left[ \int f(X_1^*, y) \, d\hat{P}_n(y) \right] \left[ \int f(x, X_2^*) \, d\hat{P}_n(x) \right] \right\} \\ &= \frac{1}{n^2} E_W \left\{ f(X_1^*, X_2^*)^2 - 2 \left[ \int f(X_1^*, y) \, d\hat{P}_n(y) \right]^2 + 2 \left[ \int f \, d(\hat{P}_n \times \hat{P}_n) \right]^2 \right\} \\ &\leq \frac{1}{n^2} \int f^2 \, d(\hat{P}_n \times \hat{P}_n), \end{split}$$

where the last inequality is because  $[\int f d(\hat{P}_n \times \hat{P}_n)]^2 \leq \int [\int f d\hat{P}_n]^2 d\hat{P}_n$  by Jensen's inequality. Hence,

by the assumption that f(x + c, y + c) = f(x, y),

$$\begin{split} E_{0}E_{W}\left[\frac{1}{n^{2}}\sum_{i\neq j}V_{n}(X_{i}^{*},X_{j}^{*})\right]^{2} &\lesssim \frac{1}{n^{2}}E_{0}\left[\int f^{2}\,\mathrm{d}(\hat{P}_{n}\times\hat{P}_{n})\right] \\ &= \frac{1}{n^{2}}E_{0}\left[\frac{1}{n^{2}h^{2d}}\sum_{i,j}\int\!\!\!\int\{f(s,t)\}^{2}\,K\left(\frac{s-X_{i}}{h}\right)K\left(\frac{t-X_{j}}{h}\right)\,\mathrm{d}s\,\mathrm{d}t\right] \\ &= \frac{1}{n^{2}}E_{0}\left[\frac{1}{n^{2}}\sum_{i,j}\int\!\!\!\int f(X_{i}+hs',X_{j}+ht')^{2}K(s')K(t')\,\mathrm{d}s'\,\mathrm{d}t'\right] \\ &= \frac{1}{n^{2}}E_{0}\left[\frac{1}{n^{2}}\sum_{i=1}^{n}\int\!\!\!\int\{f(X_{i}+hs',X_{i}+ht')\}^{2}\,K(s')K(t')\,\mathrm{d}s'\,\mathrm{d}t'\right] \\ &\quad + \frac{1}{n^{2}}E_{0}\left[\frac{1}{n^{2}}\sum_{i\neq j}\int\!\!\!\int\{f(X_{i}+hs',X_{j}+ht')\}^{2}\,K(s')K(t')\,\mathrm{d}s'\,\mathrm{d}t'\right] \\ &= \frac{1}{n^{3}}\int\!\!\!\int\{f(hs',ht')\}^{2}K(s')K(t')\,\mathrm{d}s'\,\mathrm{d}t' \\ &\quad + \frac{n-1}{n^{3}}\int\!\!\!\int E_{0}\left[f(X_{i}+hs'-ht',X_{j})^{2}\right]K(s')K(t')\,\mathrm{d}s'\,\mathrm{d}t' \\ &\lesssim \frac{1}{n^{3}}\int\!\!\!\int\{f(hs',ht')\}^{2}K(s')K(t')\,\mathrm{d}s'\,\mathrm{d}t' + \frac{1}{n^{2}}\|f\|_{L_{2}(\lambda\times P_{0})}^{2} \\ &= \frac{1}{n^{3}}\tau_{G[f^{2}]} + \frac{1}{n^{2}}\|f\|_{L_{2}(\lambda\times P_{0})}^{2}, \end{split}$$

where the last equality is because f(x + c, y + c) = f(x, y) for all  $x, y, c \in \mathbb{R}^d$  and the definition of  $\tau_{G[f]}$ . The result follows.

**Corollary F.7.** If  $\hat{P}_n$  is the distribution corresponding to a kernel density estimator with bandwidth h and uniformly bounded kernel K satisfying  $\int K^2 < \infty$ ,  $P_0$  possesses uniformly bounded density, and  $nh^d \longrightarrow \infty$ , then for each  $f \in \{f_1, f_2\}$ , we have  $\int f d[(\hat{P}_n - P_{0,h}) \times (\mathbb{P}_n^* - \hat{P}_n)] = o_{P_W^*}(n^{-1/2}),$  $\int f d[(\mathbb{P}_n - P_0) \times (\mathbb{P}_n - P_0)] = n^{-1} \tau_f + o_{P_0^*}(n^{-1/2}), \text{ and } \int f d[(\mathbb{P}_n^* - \hat{P}_n) \times (\mathbb{P}_n^* - \hat{P}_n)] = n^{-1} \tau_f + o_{P_W^*}(n^{-1/2}).$ 

**Proof of Corollary F.7.** We note that  $\int \left\{ \int |f(x,y)| \, dx \right\}^2 dP_{0,h}(y) = O(1)$  for each  $f \in \{f_1, f_2\}$  by Lemma F.1 since  $P_{0,h}(\mathbb{R}^d) = 1$ . Hence, by Lemma F.4, we then have

$$E_0 E_W \left\{ \int f \, \mathrm{d}[(\hat{P}_n - P_{0,h}) \times (\mathbb{P}_n^* - \hat{P}_n)] \right\}^2 \lesssim n^{-1} \left[ (nh^d)^{-1} + (nh^d)^{-2} \right] \int \left[ \int |f(x,y)| \, \mathrm{d}x \right]^2 \, \mathrm{d}P_{0,h}(y) \\ \lesssim n^{-1} \left[ (nh^d)^{-1} + (nh^d)^{-2} \right],$$

which is  $o(n^{-1})$  if  $nh^d \longrightarrow \infty$ . This shows the first statement.

Next, we note that  $\int \left\{ \int |f(x,y)| \, dx \right\}^2 dP_0(y) = O(1)$  and  $\|f\|_{L_2(\lambda \times P_0)} = O(h^{-d/2})$  for each  $f \in \{f_1, f_2\}$  by Lemma F.1. Hence, since  $P_0$  possesses uniformly bounded Lebesgue density function, by Lemma E.2, we have

$$\begin{split} E_0 \left\{ \int f \, \mathrm{d}[(\mathbb{P}_n - P_0) \times (\mathbb{P}_n - P_0)] - n^{-1} \tau_f \right\}^2 &\lesssim n^{-3} \int \left[ \int f(x, y) \, \mathrm{d}P_0(x) \right]^2 \, \mathrm{d}P_0(y) + n^{-2} \, \|f\|_{L_2(P_0 \times P_0)}^2 \\ &\leq n^{-3} \int \left[ \int |f(x, y)| \, \mathrm{d}x \right]^2 \, \mathrm{d}P_0(y) + n^{-2} \, \|f\|_{L_2(\lambda \times P_0)}^2 \\ &\lesssim n^{-3} + n^{-1} (nh^d)^{-1}, \end{split}$$

which is  $o(n^{-1})$  if  $nh^d \to \infty$ . This implies that  $\int f d[(\mathbb{P}_n - P_0) \times (\mathbb{P}_n - P_0)] = n^{-1}\tau_f + o_{P_W^*}(n^{-1/2})$  if  $nh^d \to \infty$ .

Finally, since for each  $f \in \{f_1, f_2\}$ , we have f(x, y) = f(y, x),  $f(x, x) = \tau_f$  and f(x+c, y+c) = f(x, y) for all  $x, y, c \in \mathbb{R}^d$ , under the stated conditions, by Lemma F.6,

$$\begin{split} E_0 E_W \left\{ \int f \, \mathrm{d}[(\mathbb{P}_n^* - \hat{P}_n) \times (\mathbb{P}_n^* - \hat{P}_n)] - n^{-1} \tau_f \right\}^2 \\ &\lesssim \frac{\tau_G^2[f]}{n^4} + \frac{\tau_G[f^2]}{n^3} + \frac{\tau_G[f]}{n^3} \, \|f\|_{L_1(\lambda \times P_0)} + \frac{1}{n^2} \, \|f\|_{L_2(\lambda \times P_0)}^2 + \frac{1}{n^3} \int \left[ \int |f(x, y)| \, \mathrm{d}x \right]^2 \, \mathrm{d}P_0(y) \\ &\quad + \frac{1}{n^3} \left[ 1 + (nh^d)^{-1} + (nh^d)^{-2} \right] \int \left( \int |f(x, y)| \, \mathrm{d}x \right)^2 \, \mathrm{d}P_{0,h}(y). \end{split}$$

By Lemma F.1, we have  $||f||_{L_1(\lambda \times P_0)} = O(1)$ ,  $||f||_{L_2(\lambda \times P_0)} = O(h^{-d/2})$ , and  $\int \left\{ \int |f(x,y)| \, dx \right\}^2 \, dP_0(y) = O(1)$  for each  $f \in \{f_1, f_2\}$ . We also note that  $\tau_{G[f]} := \iint f(hu, hv)K(u)K(v) \, du \, dv \lesssim h^{-d}$  and  $\tau_{G[f^2]} := \iint \{f(hu, hv)\}^2 K(u)K(v) \, du \, dv \lesssim h^{-2d}$  for each  $f \in \{f_1, f_2\}$ , where the constants only depend on K. Thus,

$$E_0 E_W \left\{ \int f d[(\mathbb{P}_n^* - \hat{P}_n) \times (\mathbb{P}_n^* - \hat{P}_n)] - n^{-1} \tau_f \right\}^2 \lesssim \frac{h^{-2d}}{n^4} + \frac{h^{-2d}}{n^3} + \frac{h^{-d}}{n^3} + \frac{h^{-d}}{n^2} + \frac{1}{n^3} + \frac{h^{-2d}}{n^5} \\ = n^{-2} (nh^d)^{-2} + n^{-1} (nh^d)^{-2} + n^{-2} (nh^d)^{-1} \\ + n^{-1} (nh^d)^{-1} + n^{-3} + n^{-3} (nh^d)^{-2},$$

which is again  $o(n^{-1})$  if  $nh^d \longrightarrow \infty$ .

**Lemma F.8.** If  $P_0$  possesses uniformly bounded and continuously differentiable density function  $\eta_0$ 

with  $\int [(D^{\alpha}\eta_0)(x)]^2 dx < \infty$  for all  $\alpha$  such that  $|\alpha| = 1$ ,  $\hat{P}_n$  is a kernel density estimator with uniformly bounded kernel function K satisfying  $\int K^2 < \infty$  and bandwidth h satisfying  $nh^d \longrightarrow \infty$  and  $h \longrightarrow 0$ ,  $\eta_n$  is the density corresponding to  $\hat{P}_n$ , and  $\eta_n^*$  is a kernel density estimator based on the bootstrap data with the same kernel and bandwidth, then  $(\mathbb{P}_n^* - \hat{P}_n)(\eta_n^* - \eta_n) - (\mathbb{P}_n - P_0)(\eta_n - \eta_0) = o_{P_W^*}(n^{-1/2}).$ 

**Proof of Lemma F.8.** As above, we define  $\eta_{0,h}(x) := \int K_h(x,y) \, dP_0(y)$  and  $\eta_{n,h}(x) := \int K_h(x,y) \, d\hat{P}_n(y)$ . By adding and subtracting terms, we then have

$$\begin{split} (\mathbb{P}_{n}^{*} - \hat{P}_{n})(\eta_{n}^{*} - \eta_{0}) &- (\mathbb{P}_{n} - P_{0})(\eta_{n} - \eta_{0}) \\ &= (\mathbb{P}_{n}^{*} - \hat{P}_{n})(\eta_{n}^{*} - \eta_{n,h}) + (\mathbb{P}_{n}^{*} - \hat{P}_{n})(\eta_{n,h} - \eta_{0,h}) + (\mathbb{P}_{n}^{*} - \hat{P}_{n})(\eta_{0,h} - \eta_{0}) \\ &- (\mathbb{P}_{n} - P_{0})(\eta_{n} - \eta_{0,h}) - (\mathbb{P}_{n} - P_{0})(\eta_{0,h} - \eta_{0}) \\ &= \int K_{h} d[(\mathbb{P}_{n}^{*} - \hat{P}_{n}) \times (\mathbb{P}_{n}^{*} - \hat{P}_{n})] + \int K_{h} d[(\mathbb{P}_{n}^{*} - \hat{P}_{n}) \times (\hat{P}_{n} - P_{0,h})] \\ &+ \iint [K_{h}(x, y) - \eta_{0}(x)] d(\mathbb{P}_{n}^{*} - \hat{P}_{n})(x) dP_{0}(y)] - \int K_{h} d[(\mathbb{P}_{n} - P_{0}) \times (\mathbb{P}_{n} - P_{0})] \\ &- \iint [K_{h}(x, y) - \eta_{0}(x)] d(\mathbb{P}_{n} - P_{0})(x) dP_{0}(y)] \end{split}$$

By Corollary F.7, if  $nh^d \longrightarrow \infty$ , then

$$\int K_h \,\mathrm{d}[(\mathbb{P}_n^* - \hat{P}_n) \times (\hat{P}_n - P_{0,h})] = o_{P_W^*}(n^{-1/2})$$

and

$$\int K_h d[(\mathbb{P}_n^* - \hat{P}_n) \times (\mathbb{P}_n^* - \hat{P}_n)] - \int K_h d[(\mathbb{P}_n - P_0) \times (\mathbb{P}_n - P_0)]$$
$$= \left[ n^{-1} \tau_{f_1} + o_{P_W^*}(n^{-1/2}) \right] - \left[ n^{-1} \tau_{f_1} + o_{P_0^*}(n^{-1/2}) \right]$$
$$= o_{P_W^*}(n^{-1/2}).$$

Finally, we write

$$\iint [K_h(x,y) - \eta_0(x)] d(\mathbb{P}_n^* - \hat{P}_n)(x) dP_0(y)] = (\mathbb{P}_n^* - \hat{P}_n)g_h, \text{ and}$$
$$\iint [K_h(x,y) - \eta_0(x)] d(\mathbb{P}_n - P_0)(x) dP_0(y)] = (\mathbb{P}_n - P_0)g_h$$

for  $g_h(x) := \int [K_h(x,y) - \eta_0(x)] dP_0(y)$ . As in the proofs of Lemma F.3 and Lemma F.6, we have

 $E_0 [(\mathbb{P}_n - P_0)g_h]^2 \leq n^{-1} ||g_h||_{L_2(P_0)}^2$ . By Lemma E.6,  $\int g_h(x)^2 dx = O(h^2)$ , so  $||g_h||_{L_2(P_0)}^2 = o(1)$  as long as  $h \longrightarrow 0$  since  $P_0$  has uniformly bounded density. This implies that  $\iint [K_h(x, y) - \eta_0(x)] d(\mathbb{P}_n - P_0)(x) dP_0(y) = o_{P_0^*}(n^{-1/2})$ . Furthermore,

$$\begin{split} E_0 E_W \left[ (\mathbb{P}_n^* - \hat{P}_n) g_h \right]^2 &\leq n^{-1} E_0 \|g_h\|_{L_2(\hat{P}_n)}^2 \\ &= n^{-1} \int [g_h(x)]^2 \, \mathrm{d}P_{0,h}(x) \\ &= n^{-1} \iint [g_h(x)]^2 \, K_h(x,y) \eta_0(y) \, \mathrm{d}y \, \mathrm{d}x \\ &= n^{-1} \iint [g_h(x)]^2 \, K(u) \eta_0(x+hu) \, \mathrm{d}u \, \mathrm{d}x \\ &\lesssim n^{-1} \int [g_h(x)]^2 \, \mathrm{d}x. \end{split}$$

By Lemma E.6,  $\int g_h(x)^2 dx = O(h^2) = o(1)$ . Thus,  $\int [K_h(x,y) - \eta_0(x)] d(\mathbb{P}_n^* - \hat{P}_n)(x) dP_0(y)] = o_{P_W^*}(n^{-1/2}).$ 

Lemma F.9. If  $P_0$  possesses uniformly bounded and m-times continuously differentiable density  $\eta_0$ with  $\int [(D^{\alpha}\eta_0)(x)]^2 dx < \infty$  for all  $\alpha$  such that  $|\alpha| = m$ ,  $\hat{P}_n$  is a kernel density estimator with uniformly bounded mth order kernel function K satisfying  $\int K^2 < \infty$  and bandwidth h satisfying  $nh^{2d} \longrightarrow \infty$  and  $nh^{4m} \longrightarrow 0$ ,  $\eta_n$  is the density corresponding to  $\hat{P}_n$ , and  $\eta_n^*$  is a kernel density estimator based on the bootstrap data with the same kernel and bandwidth, then  $\int [\eta_n(x) - \eta_0(x)]^2 dx - \int [\eta_n^*(x) - \eta_n(x)]^2 dx =$  $o_{P_W^*}(n^{-1/2}).$ 

**Proof of Lemma F.9.** Recall that  $f_1(x,y) := K_h(x,y)$  and  $f_2(x,y) := \int K_h(x,z)K_h(y,z) dz$ . As above, we denote  $\eta_{0,h}(x) := \int K_h(x,y) dP_0(y)$  and  $\eta_{n,h}(x) = \int K_h(x,y) d\hat{P}_n(y)$ . We then note that

$$\int \eta_n^2 = \int f_2 \,\mathrm{d}(\mathbb{P}_n \times \mathbb{P}_n), \ \int (\eta_n^*)^2 = \int f_2 \,\mathrm{d}(\mathbb{P}_n^* \times \mathbb{P}_n^*), \ \int \eta_{0,h}^2 = \int f_2 \,\mathrm{d}(P_0 \times P_0)$$
$$\int \eta_n^* \eta_n = \int f_2 \,\mathrm{d}(\mathbb{P}_n^* \times \mathbb{P}_n), \ \int \eta_n^* \eta_{0,h} = \int f_2 \,\mathrm{d}(\mathbb{P}_n^* \times P_0), \ \int \eta_n \eta_{0,h} = \int f_2 \,\mathrm{d}(\mathbb{P}_n \times P_0)$$
$$\int \eta_{n,h}^2 = \int f_2 \,\mathrm{d}(\hat{P}_n \times \hat{P}_n), \ \int \eta_n^* \eta_{n,h} = \int f_2 \,\mathrm{d}(\mathbb{P}_n^* \times \hat{P}_n), \ \int \eta_n \eta_{n,h} = \int f_2 \,\mathrm{d}(\mathbb{P}_n \times \hat{P}_n)$$

Hence, by adding and subtracting term, we have

$$\int (\eta_n^* - \eta_n)^2 - \int (\eta_n - \eta_0)^2$$
  
=  $\int (\eta_n^* - \eta_{n,h} + \eta_{n,h} - \eta_n)^2 - \int (\eta_n - \eta_{0,h} + \eta_{0,h} - \eta_0)^2$ 

$$= \int (\eta_n^* - \eta_{n,h})^2 + \int (\eta_{n,h} - \eta_n)^2 + 2 \int (\eta_n^* - \eta_{n,h}) (\eta_{n,h} - \eta_n) - \int (\eta_n - \eta_{0,h})^2 - \int (\eta_{0,h} - \eta_0)^2 - 2 \int (\eta_n - \eta_{0,h}) (\eta_{0,h} - \eta_0) = \left\{ \int f_2 d[(\mathbb{P}_n^* - \hat{P}_n) \times (\mathbb{P}_n^* - \hat{P}_n)] - \int f_2 d[(\mathbb{P}_n - P_0) \times (\mathbb{P}_n - P_0)] \right\} + \int (\eta_{n,h} - \eta_n)^2 - \int (\eta_{0,h} - \eta_0)^2 + 2 \int f_2 d[(\mathbb{P}_n^* - \hat{P}_n) \times (\hat{P}_n - \mathbb{P}_n)] - 2 \int (f_2 - f_1) d[(\mathbb{P}_n - P_0) \times P_0].$$

By Corollary F.7, we have

$$\int f_2 d[(\mathbb{P}_n^* - \hat{P}_n) \times (\mathbb{P}_n^* - \hat{P}_n)] - \int f_2 d[(\mathbb{P}_n - P_0) \times (\mathbb{P}_n - P_0)] = o_{P_W^*}(n^{-1/2}).$$

By Lemma E.6,  $\int (\eta_{0,h} - \eta_0)^2 = O(h^{2m})$ , which is  $o(n^{-1/2})$  since  $nh^{4m} \longrightarrow 0$ . Next, we note that

$$\begin{split} \eta_{n,h}(x) - \eta_n(x) &= \int K_h(x,z) \,\mathrm{d}\hat{P}_n(z) - \int K_h(x,y) \,\mathrm{d}\mathbb{P}_n(y) \\ &= \int K_h(x,z)\eta_n(z) \,\mathrm{d}z - \int K_h(x,y) \,\mathrm{d}\mathbb{P}_n(y) \\ &= \int \left[ \int K_h(x,z)K_h(y,z) \,\mathrm{d}z \right] \,\mathrm{d}\mathbb{P}_n(y) - \int K_h(x,y) \,\mathrm{d}\mathbb{P}_n(y) \\ &= \int \left[ f_2(x,y) - f_1(x,y) \right] \,\mathrm{d}\mathbb{P}_n(y) \\ &= \int f_3(x,y) \,\mathrm{d}\mathbb{P}_n(y), \end{split}$$

where  $f_3 = f_2 - f_1$ . Therefore,

$$\begin{split} E_0 \int (\eta_{n,h} - \eta_n)^2 &= E_0 \int \left[ \int f_3(x,y) \, \mathrm{d}\mathbb{P}_n(y) \right]^2 \, \mathrm{d}x \\ &= E_0 \int \left\{ \frac{1}{n^2} \sum_{i,j} f_3(x,X_i) f_3(x,X_j) \right\} \, \mathrm{d}x \\ &= E_0 \int \left\{ \frac{1}{n^2} \sum_{i=1}^n f_3(x,X_i) f_3(x,X_i) \right\} \, \mathrm{d}x + E_0 \int \left\{ \frac{1}{n^2} \sum_{i\neq j} f_3(x,X_i) f_3(x,X_j) \right\} \, \mathrm{d}x \\ &= \frac{1}{n} \int E_0 \left[ f_3^2(x,X) \right] \, \mathrm{d}x + \frac{n-1}{n} \int \left\{ E_0 \left[ f_3(x,X) \right] \right\}^2 \, \mathrm{d}x \\ &= \frac{1}{n} \left\| f_3 \right\|_{L_2(\lambda \times P_0)}^2 + \frac{n-1}{n} \int \left\{ E_0 \left[ f_3(x,X) \right] \right\}^2 \, \mathrm{d}x. \end{split}$$

By Lemma F.1, we have  $||f_3||_{L_2(\lambda \times P_0)} \le ||f_1||_{L_2(\lambda \times P_0)} + ||f_2||_{L_2(\lambda \times P_0)} = O(h^{-d/2})$ . By Lemma E.6, we have

$$\int \left\{ E_0 \left[ f_3(x,X) \right] \right\}^2 \, \mathrm{d}x \le 2 \int \left\{ E_0 \left[ f_1(x,X) \right] - \eta_0(x) \right\}^2 \, \mathrm{d}x + 2 \int \left\{ E_0 \left[ f_2(x,X) \right] - \eta_0(x) \right\}^2 \, \mathrm{d}x = O(h^{2m}).$$

Hence,  $\int (\eta_{n,h} - \eta_n)^2 = O_{P_0^*}(\{nh^d\}^{-1} + h^{2m})$ , which is  $o_{P_0^*}(n^{-1/2})$  under the conditions  $nh^{4m} \longrightarrow 0$  and  $nh^{2d} \longrightarrow \infty$ .

Next, we have

$$\int f_2 d[(\mathbb{P}_n^* - \hat{P}_n) \times (\hat{P}_n - \mathbb{P}_n)]$$
  
=  $\int f_2 d[(\mathbb{P}_n^* - \hat{P}_n) \times (\hat{P}_n - P_{0,h})] + \int f_2 d[(\mathbb{P}_n^* - \hat{P}_n) \times (P_{0,h} - P_0)]$   
-  $\int f_2 d[(\mathbb{P}_n^* - \hat{P}_n) \times (\mathbb{P}_n - P_0)].$ 

By Lemma F.4, we have

$$E_0 E_W \left\{ \int f_2 d[(\hat{P}_n - P_{0,h}) \times (\mathbb{P}_n^* - \hat{P}_n)] \right\}^2 \lesssim n^{-1} \left[ (nh^d)^{-1} + (nh^d)^{-2} \right] \int \left[ \int |f_2(x,y)| \, \mathrm{d}x \right]^2 \, \mathrm{d}P_{0,h}(y),$$

which is  $o(n^{-1})$  under the condition  $nh^d \longrightarrow \infty$  by Lemma F.1. By Lemma F.3, we have

$$E_0 E_W \left[ \int f_2 d[(P_{0,h} - P_0) \times (\mathbb{P}_n^* - \hat{P}_n)] \right]^2 \lesssim n^{-1} h^{2m} \|f_2\|_{L_2(\lambda \times P_{0,h})}^2,$$

which is  $O(n^{-1}h^{2m-d})$  by Lemma F.1, which is  $o(n^{-1})$  by assumption. By Lemma F.1 and Lemma F.5, we have

$$\begin{split} E_0 E_W \left\{ \int f_2 \, \mathrm{d}[(\mathbb{P}_n - P_0) \times (\mathbb{P}_n^* - \hat{P}_n)] \right\}^2 &\lesssim \frac{1}{n^2} \, \|f_2\|_{L_2(P_0 \times P_{0,h})}^2 + \frac{1}{n^3 h^d} \, \|f_2\|_{L_2(\lambda \times P_0)}^2 \\ &\lesssim \frac{1}{n^2} \, \|f_2\|_{L_2(P_0 \times P_{0,h})}^2 + \frac{1}{n^3 h^{2d}} \\ &\lesssim \frac{1}{n^2 h^d} + \frac{1}{n^3 h^{2d}}, \end{split}$$

where the last inequality is because

$$\iint f_2^2(x,z) \, \mathrm{d}P_0(x) \, \mathrm{d}P_{0,h}(z) = \iint \left[ \int K_h(x,s) K_h(z,s) \, \mathrm{d}s \right]^2 \, \mathrm{d}P_0(x) \, \mathrm{d}P_{0,h}(z)$$

$$= \iint \left[ \iint K_h(x,s) K_h(z,s) K_h(x,t) K_h(z,t) \, \mathrm{d}s \, \mathrm{d}t \right] \, \mathrm{d}P_0(x) \, \mathrm{d}P_{0,h}(z)$$

$$= \frac{1}{h^{4d}} \iiint K\left(\frac{x-s}{h}\right) K\left(\frac{z-s}{h}\right) K\left(\frac{x-t}{h}\right)$$

$$\times K\left(\frac{z-t}{h}\right) \, \mathrm{d}s \, \mathrm{d}t \, \mathrm{d}P_0(x) \, \mathrm{d}P_{0,h}(z)$$

$$= \frac{1}{h^{2d}} \left| \iiint K\left(s'\right) K\left(s'\right) K\left(s'+\frac{z-x}{h}\right) K\left(t'\right)$$

$$\times K\left(t'+\frac{z-x}{h}\right) \, \mathrm{d}s' \, \mathrm{d}t' \, \mathrm{d}P_0(x) \, \mathrm{d}P_{0,h}(z) \right|$$

$$= \frac{1}{h^d} \left| \iiint K\left(s'\right) K\left(x'\right) K\left(t'\right)$$

$$\times K\left(t'+x'-s'\right) \eta_0(z-(x'-s')h) \, \mathrm{d}s' \, \mathrm{d}t' \, \mathrm{d}P_{0,h}(z) \right|$$

$$\lesssim \frac{1}{h^d} \iiint |K\left(s'\right) K\left(x'\right) K\left(t'\right) K\left(t'+x'-s'\right)| \, \mathrm{d}s' \, \mathrm{d}t' \, \mathrm{d}P_{0,h}(z)$$

$$\leq \frac{1}{h^d} \left[ \int K^2(x) \, \mathrm{d}x \right]^2.$$

This implies that  $E_0 E_W \left\{ \int f_2 d[(\mathbb{P}_n - P_0) \times (\mathbb{P}_n^* - \hat{P}_n)] \right\}^2 = o(n^{-1})$  under the assumption  $nh^d \longrightarrow \infty$ . Hence,  $\int f_2 d[(\mathbb{P}_n^* - \hat{P}_n) \times (\hat{P}_n - \mathbb{P}_n)] = o_{P_W^*}(n^{-1/2})$  if  $nh^d \longrightarrow \infty$ .

Finally, by Lemma E.6, we have

$$E_0 \left[ \int (f_2 - f_1) d[(\mathbb{P}_n - P_0) \times P_0] \right]^2$$
  

$$\leq \frac{1}{n} E_0 \left[ \int \{ f_2(X, y) - f_1(X, y) \} dP_0(y) \right]^2$$
  

$$\leq \frac{2}{n} E_0 \left[ \int \{ f_1(X, y) - \eta_0(X) \} dP_0(y) \right]^2 + \frac{2}{n} E_0 \left[ \int \{ f_2(X, y) - \eta_0(X) \} dP_0(y) \right]^2$$
  

$$= O(n^{-1}h^{2m}),$$

which is  $o(n^{-1})$ .

**Lemma F.10.** If  $P_0$  possesses uniformly bounded and m-times continuously differentiable density  $\eta_0$ with  $\int [(D^{\alpha}\eta_0)(x)]^2 dx < \infty$  for all  $\alpha$  such that  $|\alpha| = m$ ,  $\hat{P}_n$  is a kernel density estimator with uniformly bounded mth order kernel function K satisfying  $\int K^2 < \infty$  and bandwidth h satisfying  $nh^{2d} \longrightarrow \infty$  and  $nh^{4m} \longrightarrow 0$ ,  $\eta_n$  is the density corresponding to  $\hat{P}_n$ , and  $\eta_n^*$  is a kernel density estimator based on the bootstrap data with the same kernel and bandwidth, then  $\mathbb{P}_n\phi_n - \mathbb{P}_n^*\phi_n^* = o_{P_W}(n^{-1/2})$ .

**Proof of Lemma F.10.** Recall that  $f_1(x,y) := K_h(x,y)$  and  $f_2(x,y) := \int K_h(x,z)K_h(y,z) dz$ . We

define  $f_3 := f_1 - f_2$ . We then have  $\mathbb{P}_n \phi_n = 2 \int f_3 d(\mathbb{P}_n \times \mathbb{P}_n)$  and  $\mathbb{P}_n^* \phi_n^* = 2 \int f_3 d(\mathbb{P}_n^* \times \mathbb{P}_n^*)$ . Hence, by adding and subtracting terms and the symmetry of  $f_3$ ,

$$\begin{split} \left(\mathbb{P}_{n}^{*}\phi_{n}^{*}-\mathbb{P}_{n}\phi_{n}\right)/2 \\ &=\int f_{3}\,\mathrm{d}[\left(\mathbb{P}_{n}^{*}-\hat{P}_{n}+\hat{P}_{n}-P_{0,h}+P_{0,h}\right)\times\left(\mathbb{P}_{n}^{*}-\hat{P}_{n}+\hat{P}_{n}-P_{0,h}+P_{0,h}\right)] \\ &-\int f_{3}\,\mathrm{d}[\left(\mathbb{P}_{n}-P_{0}+P_{0}\right)\times\left(\mathbb{P}_{n}-P_{0}+P_{0}\right)] \\ &=\left\{\int f_{3}\,\mathrm{d}[\left(\mathbb{P}_{n}^{*}-\hat{P}_{n}\right)\times\left(\mathbb{P}_{n}^{*}-\hat{P}_{n}\right)]-\int f_{3}\,\mathrm{d}[\left(\mathbb{P}_{n}-P_{0}\right)\times\left(\mathbb{P}_{n}-P_{0}\right)]\right\} \\ &+\int f_{3}\,\mathrm{d}[\left(\hat{P}_{n}-P_{0,h}\right)\times\left(\hat{P}_{n}-P_{0,h}\right)]+\left\{\int f_{3}\,\mathrm{d}[P_{0,h}\times P_{0,h}]-\int f_{3}\,\mathrm{d}[P_{0}\times P_{0}]\right\} \\ &+2\int f_{3}\,\mathrm{d}[\left(\mathbb{P}_{n}^{*}-\hat{P}_{n}\right)\times\left(\hat{P}_{n}-P_{0,h}\right)]+2\int f_{3}\,\mathrm{d}[\left(\mathbb{P}_{n}^{*}-\hat{P}_{n}\right)\times\left(P_{0,h}-P_{0}\right)] \\ &+2\int f_{3}\,\mathrm{d}[\left(\mathbb{P}_{n}^{*}-\hat{P}_{n}\right)\times P_{0}]+2\int f_{3}\,\mathrm{d}[(\hat{P}_{n}-P_{0,h})\times P_{0,h}]-2\int f_{3}\,\mathrm{d}[(\mathbb{P}_{n}-P_{0})\times P_{0}]. \end{split}$$

We show each term on the right hand side of previous display is  $o_{P_W^*}(n^{-1/2})$ . First, we can use the same logic as in the proof of Lemma F.9 to show that

$$\int f_3 d[(\mathbb{P}_n^* - \hat{P}_n) \times (\mathbb{P}_n^* - \hat{P}_n)] - \int f_3 d[(\mathbb{P}_n - P_0) \times (\mathbb{P}_n - P_0)] = o_{P_W^*}(n^{-1/2}),$$
  
$$\int f_3 d[(\mathbb{P}_n^* - \hat{P}_n) \times (\hat{P}_n - P_{0,h})] = o_{P_W^*}(n^{-1/2}),$$
  
$$\int f_3 d[(\mathbb{P}_n^* - \hat{P}_n) \times (P_{0,h} - P_0)] = o_{P_W^*}(n^{-1/2}), \text{ and}$$
  
$$\int f_3 d[(\mathbb{P}_n - P_0) \times P_0] = o_{P_0^*}(n^{-1/2}).$$

By Lemma F.1 and Corollary F.2, we have

$$E_0 \left\{ \int f \, d[(\hat{P}_n - P_{0,h}) \times (\hat{P}_n - P_{0,h})] - n^{-1} \tau_{G[f]} \right\}^2$$
  
$$\lesssim n^{-3} \int \left[ \int |f(x,y)| \, dx \right]^2 \, dP_0(y) + n^{-2} \, \|f\|_{L_2(\lambda \times P_0)}^2$$
  
$$= O(n^{-3}) + O(n^{-2}h^{-d})$$

for  $f \in \{f_1, f_2\}$ , which is  $o(n^{-1})$  because  $nh^d \longrightarrow \infty$ . Hence,

$$\int f_3 \,\mathrm{d}[(\hat{P}_n - P_{0,h}) \times (\hat{P}_n - P_{0,h})]$$

$$= \int f_1 d[(\hat{P}_n - P_{0,h}) \times (\hat{P}_n - P_{0,h})] - \int f_2 d[(\hat{P}_n - P_{0,h}) \times (\hat{P}_n - P_{0,h})]$$
$$= n^{-1} \left( \tau_{G[f_1]} + \tau_{G[f_2]} \right) + o_{P_0^*} (n^{-1/2}).$$

Furthermore, we have  $\tau_{G[f]} = O(h^{-d})$  for  $f \in \{f_1, f_2\}$  as shown above. Since  $nh^{2d} \longrightarrow \infty$ , we then have  $n^{-1}(\tau_{G[f_1]} + \tau_{G[f_2]}) = O(\{nh^d\}^{-1}) = O(n^{-1/2}\{nh^{2d}\}^{-1/2}) = o(n^{-1/2}).$ 

Next, we note that

$$\begin{split} \left| \int f_{3} d(P_{0,h} \times P_{0,h}) - \int f_{3} d(P_{0} \times P_{0}) \right| \\ &= \left| \iint \left[ \iint f_{3}(s,t) K_{h}(x,s) K_{h}(y,t) ds dt \right] dP_{0}(x) dP_{0}(y) - \int f_{3} d(P_{0} \times P_{0}) \right| \\ &= \left| \iint f_{3}(x+sh,y+th) K(s) K(t) ds dt \right] dP_{0}(x) dP_{0}(y) - \int f_{3} d(P_{0} \times P_{0}) \right| \\ &= \left| \iiint f_{3}(x,y+t'h-s'h) K(s') K(t') dP_{0}(x) dP_{0}(y) ds' dt' - \int f_{3} d(P_{0} \times P_{0}) \right| \\ &= \left| \iiint f_{3}(x,y') K(s') K(t') dP_{0}(x) \eta_{0}(y'+s'h-t'h) dy' ds' dt' - \int f_{3} d(P_{0} \times P_{0}) \right| \\ &= \left| \int \left[ \int f_{3}(x,y) dP_{0}(x) \right] \left[ \iint \{\eta_{0}(y+sh-th) - \eta_{0}(y)\} K(s) K(t) ds dt \right] dy \right| \\ &\leq \left\{ \int \left[ \int f_{3}(x,y) dP_{0}(x) \right]^{2} dy \right\}^{1/2} \left\{ \int \left[ \iint \{\eta_{0}(y+sh-th) - \eta_{0}(y)\} K(s) K(t) ds dt \right]^{2} dP_{0}(y) \right\}^{1/2} \end{split}$$

We can write

$$\int \left[ \int f_3(x,y) \, \mathrm{d}P_0(y) \right]^2 \, \mathrm{d}x$$
  
$$\leq 2 \int \left[ \int f_1(x,y) \, \mathrm{d}P_0(y) - \eta_0(x) \right]^2 \, \mathrm{d}x + 2 \int \left[ \int f_2(x,y) \, \mathrm{d}P_0(y) - \eta_0(x) \right]^2 \, \mathrm{d}x,$$

which is  $O(h^{2m})$  by Lemma E.6. Since  $\eta_0$  is *m*-times continuously differentiable, for all *u*, a Taylor expansion with the Laplacian representation of the remainder gives

$$\begin{split} \iint \left\{ \eta_0(y + sh - th) - \eta_0(y) \right\} K(s) K(t) \, \mathrm{d}s \, \mathrm{d}t \\ &= \iint \left\{ \sum_{1 \le |\alpha| \le m-1} \frac{1}{\alpha!} (sh - th)^{\alpha} (D^{\alpha} \eta_0)(y) \right\} K(s) K(t) \, \mathrm{d}s \, \mathrm{d}t \\ &+ \iint \left\{ \sum_{|\alpha| = m} \frac{mh^m}{\alpha!} (s - t)^{\alpha} \int_0^1 (1 - r)^{m-1} (D^{\alpha} \eta_0) (y + rh[s - t]) \, \mathrm{d}r \right\} K(s) K(t) \, \mathrm{d}s \, \mathrm{d}t \end{split}$$

$$= \sum_{1 \le |\alpha| \le m-1} \frac{1}{\alpha!} (D^{\alpha} \eta_0)(y) \iint (sh - th)^{\alpha} K(s) K(t) \, \mathrm{d}s \, \mathrm{d}t \\ + \sum_{|\alpha|=m} \frac{mh^m}{\alpha!} \iint (s - t)^{\alpha} \int_0^1 (1 - r)^{m-1} (D^{\alpha} \eta_0)(y + rh[s - t]) \, \mathrm{d}r K(s) K(t) \, \mathrm{d}s \, \mathrm{d}t.$$

Since K is an *m*th order kernel function,  $\iint (sh - th)^{\alpha} K(s) K(t) \, ds \, dt = 0$  for all  $\alpha$  such that  $1 \le |\alpha| \le m - 1$ . Defining  $H(y, u, \alpha) := \int_0^1 (1 - r)^{m-1} (D^{\alpha} \eta_0) (y + rhu) \, dr$ , we then have

$$\begin{split} &\int \left[ \iint \left\{ \eta_0(y+sh-th) - \eta_0(y) \right\} K(s) K(t) \, \mathrm{d}s \, \mathrm{d}t \right]^2 \, \mathrm{d}P_0(y) \\ &= \int \left[ \sum_{|\alpha|=m} \frac{mh^m}{\alpha!} \iint (s-t)^{\alpha} H(y,s-t,\alpha) K(s) K(t) \, \mathrm{d}s \, \mathrm{d}t \right]^2 \, \mathrm{d}P_0(y) \\ &= \int \sum_{|\alpha|,|\beta|=m} \frac{m^2 h^{2m}}{\alpha!\beta!} \iiint (s-t)^{\alpha} (s'-t')^{\beta} H(y,s-t,\alpha) H(y,s'-t',\beta) \\ &\quad \times K(s) K(t) K(s') K(t') \, \mathrm{d}s \, \mathrm{d}t \, \mathrm{d}s' \, \mathrm{d}t' \, \mathrm{d}P_0(y). \end{split}$$

The same method as in the proof of Lemma E.6 can be used to show that this is  $O(h^{2m})$ . Hence,  $\int f_3 d(P_{0,h} \times P_{0,h}) - \int f_3 d(P_0 \times P_0) = O(h^{2m})$ , which is  $o(n^{-1/2})$  if  $nh^{4m} \longrightarrow 0$ .

We next consider the term  $\int f_3 d[(\mathbb{P}_n^* - \hat{P}_n) \times P_0]$ . As in the proofs of Lemma E.3 and Lemma E.4,

$$E_0 E_W \left\{ \int \left[ \int f_3(x, y) \, \mathrm{d}P_0(y) \right] \, \mathrm{d}(\mathbb{P}_n^* - \hat{P}_n)(x) \right\}^2$$
  
$$\leq E_0 \left\{ \frac{1}{n} \int \left[ \int f_3(x, y) \, \mathrm{d}P_0(y) \right]^2 \, \mathrm{d}\hat{P}_n(x) \right\}$$
  
$$= \frac{1}{n} \int \left[ \int f_3(x, y) \, \mathrm{d}P_0(y) \right]^2 \, \mathrm{d}P_{0,h}(x)$$
  
$$\lesssim \frac{1}{n} \int \left[ \int f_3(x, y) \, \mathrm{d}P_0(y) \right]^2 \, \mathrm{d}x,$$

where the last inequality is because  $P_{0,h}$  possesses uniformly bounded density. The last expression is  $O(n^{-1}h^{2m})$  by Lemma E.6 as shown above. Hence,  $\int f_3 d[(\mathbb{P}_n^* - \hat{P}_n) \times P_0] = O_{P_W^*}(n^{-1/2}h^m) = o_{P_W^*}(n^{-1/2}).$ 

Finally, we consider the term  $\int f_3 d[(\hat{P}_n - P_{0,h}) \times P_{0,h}]$ . We note that

$$\int f_3 \,\mathrm{d}[(\hat{P}_n - P_{0,h}) \times P_{0,h}] = \iiint f_3(s,t) K_h(x,s) K_h(y,t) \,\mathrm{d}s \,\mathrm{d}t \,\mathrm{d}P_0(y) \,\mathrm{d}(\mathbb{P}_n - P_0)(x).$$

Hence,

$$\begin{split} E_0 \left\{ \int f_3 \,\mathrm{d}[(\hat{P}_n - P_{0,h}) \times P_{0,h}] \right\}^2 \\ &\leq \frac{1}{n} \int \left[ \iiint f_3(s,t) K_h(x,s) K_h(y,t) \,\mathrm{d}s \,\mathrm{d}t \,\mathrm{d}P_0(y) \right]^2 \,\mathrm{d}P_0(x) \\ &= \frac{1}{n} \int \left[ \iiint f_3(x+s'h,y+t'h) K(s') K(t') \,\mathrm{d}s' \,\mathrm{d}t' \,\mathrm{d}P_0(y) \right]^2 \,\mathrm{d}P_0(x) \\ &\leq \frac{1}{n} \iiint \left[ \int f_3(x+s'h-t'h,y) \,\mathrm{d}P_0(y) \right]^2 K(s') K(t') \,\mathrm{d}s' \,\mathrm{d}t' \,\mathrm{d}P_0(x) \\ &\lesssim \frac{1}{n} \int \left[ \int f_3(x,y) \,\mathrm{d}P_0(y) \right]^2 \,\mathrm{d}x. \end{split}$$

As above, the last expression is  $O(n^{-1}h^{2m})$ , so that  $\int f_3 d[(\hat{P}_n - P_{0,h}) \times P_{0,h}] = O_{P_0^*}(n^{-1/2}h^m) = O_{P_0^*}(n^{-1/2}).$