

On checking L^p -admissibility for parabolic control systems

Philip Preußler^{†‡}

Felix L. Schwenninger[§]

April 9, 2024

Abstract

In this note we discuss the difficulty of verifying L^p -admissibility for $p \neq 2$ —that even manifests in the presence of a self-adjoint semigroup generator on a Hilbert space—and survey tests for L^p -admissibility of given control operators. These tests are obtained by virtue of either mapping properties of boundary trace operators, yielding a characterization of admissibility via abstract interpolation spaces; or through Laplace–Carleson embeddings, slightly extending results from Jacob, Partington and Pott [31] to a class of systems which are not necessarily diagonal with respect to sequence spaces. Special focus is laid on illustrating the theory by means of examples based on the heat equation on various domains.

Keywords: admissible operator, infinite-dimensional system, Laplace–Carleson embedding, Weiss conjecture.

Mathematics Subject Classification (2020): 93B28, 93C05, 93C25, 47N70.

1 Introduction

1.1 State space systems and admissibility

We consider abstract linear systems of the form

$$\dot{x}(t) = Ax(t) + Bu(t), \quad t > 0, \quad (1)$$

on an infinite-dimensional Banach space X which we call the *state space*. Here we require that the operator $A: X \supset \mathcal{D}(A) \rightarrow X$ generates a strongly continuous semigroup of linear operators, or C_0 -semigroup, on X , denoted by $T = (T(t))_{t \geq 0}$. The *state trajectory*, defined for $t \geq 0$, is denoted by x and the *input function* (or *control function*), also defined for $t \geq 0$, is denoted by u . We require u to be U -valued, where U is a Banach space that we call the *input space*. Moreover, inputs u enter the system through the *control operator* B only.

[†]Corresponding author

[‡]Department of Applied Mathematics, University of Twente, P.O. Box 217, 7500 AE Enschede, The Netherlands, p.n.preusler@utwente.nl

[§]Department of Applied Mathematics, University of Twente, P.O. Box 217, 7500 AE Enschede, The Netherlands, f.l.schwenninger@utwente.nl

In the field of infinite-dimensional systems theory there is a wide array of literature dealing with bounded control operators; that is to say, systems with $B \in \mathcal{L}(U, X)$. However, many systems arising from controlled partial differential equations lead to control operators that fail to be bounded as maps from U to X ; for example in the case of control acting on the boundary. This leads us to the study of unbounded control operators B .

Here B is called an *unbounded* control operator if B is a linear operator which is bounded as a map from U to X_{-1} , but not in $\mathcal{L}(U, X)$. The extrapolation space $X_{-1} \supset X$ is defined as the completion of X with the weaker norm

$$\|z\|_{X_{-1}} := \|(sI - A)^{-1}z\|_X$$

for some complex number s in the resolvent set $\rho(A)$.

This allows us to consider (1) in the ambient space X_{-1} . To that end, consider a fixed initial value $x(0) \in X$ and an unbounded control operator B . The mild solution of the controlled system (1) is given by

$$x(t) = T(t)x(0) + \int_0^t T(t-s)Bu(s) \, ds, \quad t \geq 0, \quad (2)$$

and takes values which a priori only lie in X_{-1} . As we are interested in the case with continuous state trajectories taking values in X —such as in the case of bounded B —we want to find conditions on B such that (2) lies in X for all $t \geq 0$, bringing us to the property of admissibility, [28, 50, 51].

Definition 1.1 (L^p -admissible control operators). For $1 \leq p \leq \infty$, the control operator $B \in \mathcal{L}(U, X_{-1})$ is called *finite-time L^p -admissible for T (or for A)* if the convolution type Bochner integral

$$\int_0^t T(t-s)Bu(s) \, ds$$

is an element of X for any $t \geq 0$ and there exists $C > 0$ such that

$$\left\| \int_0^t T(t-s)Bu(s) \, ds \right\|_X \leq C\|u\|_{L^p([0,t],U)}$$

holds for all input functions $u \in L^p([0,t],U)$.

The operator B is called *infinite-time L^p -admissible for T (or for A)* if the constant $C > 0$ from above can be chosen independently of t .

Remark 1.2. Some remarks on the definition of L^p -admissibility of control operators:

1. If the context is clear, one can also drop the reference to the semigroup T or the generator A when referring to admissibility. Moreover, one can also drop the letter L and speak of p -admissibility of control operators. Lastly, when the distinction between finite-time and infinite-time admissibility is not made, one generally speaks about infinite-time admissibility.
2. We also note that—due to translation invariance of the set of input functions—one can replace $T(t-s)$ by $T(s)$ in the integrals above. Moreover, infinite-time p -admissibility is equivalent to the existence of $C > 0$ such that

$$\left\| \int_0^\infty T(s)Bu(s) \, ds \right\|_X \leq C\|u\|_{L^p([0,\infty),U)}$$

holds for all $u \in L^p([0,\infty),U)$.

3. The property that $B \in \mathcal{L}(U, X_{-1})$ is p -admissible can be rephrased by saying that the mapping $\Phi_t: u \mapsto \int_0^t T(t-s)Bu(s) ds$ is bounded from L^p to X —rather than only mapping to X_{-1} —for some and hence all $t > 0$. Dualizing this statement, we arrive at an admissibility notion for *observation operators* $C: \mathcal{D}(A) \rightarrow Y$, where $\mathcal{D}(A)$ is equipped with the graph norm of A and the Banach space Y is called the *output space*.

More precisely, we fix $p \in (1, \infty)$ and denote the Hölder conjugate of p by p' . Then, provided that the dual semigroup $T' = (T(t)')_{t \geq 0}$ is strongly continuous on the Banach space dual X' , $B: U \rightarrow X_{-1}$ is p -admissible if and only if the operator $C = B': \mathcal{D}(A') \rightarrow U'$ satisfies

$$\int_0^t \|B'T(t)'x\|_{Y'}^{p'} dt \leq K \|x\|_{X'}^p, \quad x \in \mathcal{D}(A')$$

for some $K > 0$, see e.g. [51, Theorem 6.9].

1.2 Boundary control systems

The most natural form of many systems of partial differential equations with control on the boundary of the domain is not the state space form as in (1) above. Rather, in these systems the control enters via the boundary condition; for example in the Dirichlet trace sense $x|_{\partial\Omega} = u$. This then leads to the type of systems known as *boundary control systems*, see e.g. Salamon [39] and Tucsnak and Weiss [46]. Boundary control systems take the form

$$\begin{cases} \dot{x}(t) = \mathfrak{A}x(t) & \text{on } (0, \infty), \\ \mathfrak{B}x(t) = u(t) & \text{on } (0, \infty), \\ x(0) = x_0. \end{cases} \quad (3)$$

Typically, $\mathfrak{A}$ is a differential operator and $\mathfrak{B}$ is a boundary trace operator, such that the system corresponds to a (partial) differential equation with the control input acting via the boundary of the spatial domain. To relate this formulation to the state space form, we need some well-posedness assumptions.

Definition 1.3 (boundary control systems). Given Banach spaces X and U and closed operators

$$\mathfrak{A}: X \supset \mathcal{D}(\mathfrak{A}) \rightarrow X, \quad \mathfrak{B}: X \supset \mathcal{D}(\mathfrak{B}) = \mathcal{D}(\mathfrak{A}) \rightarrow U,$$

where we have normed the space $\mathcal{D}(\mathfrak{A})$ with the graph norm. If now the restriction of $\mathfrak{A}$ to $\ker \mathfrak{B}$ is the generator of a C_0 -semigroup on X and $\mathfrak{B}$ has a right inverse $B_0 \in \mathcal{L}(U, \mathcal{D}(\mathfrak{A}))$, then we call the pair of operators $(\mathfrak{A}, \mathfrak{B})$ and the corresponding equations (3) a *(linear) boundary control system*.

Remark 1.4. One can translate a boundary control system on X into a state space system on X by setting

$$A := \mathfrak{A}|_{\ker \mathfrak{B}}, \quad B = (\mathfrak{A} - A_{-1})B_0, \quad (4)$$

where $A_{-1}: X_{-1} \supset \mathcal{D}(A_{-1}) \rightarrow X_{-1}$ is the unique isometric extension of A to the space X_{-1} with domain $\mathcal{D}(A_{-1}) = X$, see e.g. [9, 42, 46]. More on the rewriting procedure for boundary control systems into state space form can e.g. be found in [41] or [46, Chapter 10].

This correspondence allows us to transfer concepts formulated in the state space system framework—including admissibility—to the boundary control context. The intended meaning here is that the operator B that results from $\mathfrak{B}$ through the rewriting procedure is admissible for $A = \mathfrak{A}|_{\ker \mathfrak{B}}$. While the right inverse involved in the construction of B is generally not unique, the operator B is uniquely determined. Yet, computing B from the expression given in (4) above is usually not feasible. However, on Hilbert spaces, the relation

$$\langle \mathfrak{A}x, y \rangle - \langle x, A^*y \rangle = \langle \mathfrak{B}x, B^*y \rangle, \quad (5)$$

see e.g. [41, Proposition 2.9], holds for all $x \in \mathcal{D}(\mathfrak{A})$ and $y \in \mathcal{D}(A^*)$. With this identity, we can identify the control operator B , simplifying the computation in contrast to (4).

1.3 Current state of L^p -admissibility theory

When verifying p -admissibility, there is a vast difference between the cases $p = 2$ and $p \neq 2$; a result of the missing Hilbert space structure in the latter case. This discrepancy is reflected in the amount of results available in the literature on 2-admissibility compared to p -admissibility, see also the survey article [28] by Jacob and Partington that deals mostly with the case $p = 2$. As an example, for $p = 2$ characterizations of admissibility using Lyapunov equations and Lyapunov inequalities are available, compare also [46, Chapter 5]. These methods do not generalize to $p \neq 2$ as they rely on the Hilbert space structure of the state space X , of the input space U and of the L^2 space in the domain of the input functions; illustrating the added difficulty of checking for p -admissibility.

This disparity could lead to the question of practical relevance of the more general concept of p -admissibility compared to 2-admissibility. In other words, one could ask if there are any common systems that are not 2-admissible but only p -admissible for some $p > 2$. Concretely, we are interested in finding examples of systems arising from partial differential equations—on a natural state space such as $L^2(\Omega)$ for some domain $\Omega \subset \mathbb{R}^n$ —such that the corresponding control operator B is not 2-admissible. Examples of this type exist; for instance the heat equation on the spatial domain $\Omega = [0, 1]$ with Dirichlet boundary control and scalar input space $U = \mathbb{C}$. Here the control operator B resulting from rewriting the boundary control system into state space form turns out to be $B = \delta'_0$, which is p -admissible for all $p > 4$ and not p -admissible for $p \leq 4$, see [34] and also Section 2.2 in the sequel. Moreover, p -admissibility (in a weighted variant) appears as a prerequisite to the abstract Kato method in the article [17] by Haak and Kunstmann to derive results for Navier–Stokes equations.

In the context of abstract linear systems, p -admissibility for $p \neq 2$ has been studied in the operator-theoretic framework by and since Weiss’s [50, 51] seminal works on observation and control operators. Weiss also justifies the assumption $B \in \mathcal{L}(U, X_{-1})$ in the definition of an admissible operator by showing that each “admissible” operator B (that would be bounded only from U to some space V in which X lies dense) is equivalent to an operator $\hat{B}$ that is bounded $U \rightarrow X_{-1}$ in the sense that they lead to the same abstract linear system. Furthermore, tests for p -admissibility are given for some special cases, such as 1-admissibility on reflexive X , as Weiss [50] shows that B is admissible in this situation if and only if $B \in \mathcal{L}(U, X)$.

It should be mentioned that the concept of admissibility—with respect to L^2 —had previously appeared in Salamon [39] as hypothesis (S2). Connections between state space systems and boundary control systems are also discussed in the aforementioned work by Salamon, and we also mention the work by Curtain and Pritchard [7] for more on this topic.

Among approaches to characterizing admissibility, important examples are Carleson measure criteria for diagonal systems, which are known as such due to being based on certain discrete measures with weights depending on the system in question being Carleson measures. A method based on these conditions for checking 2-admissibility with scalar input space $U = \mathbb{C}$ appears in Ho and Russell [24], was extended to an equivalent condition by Weiss [49] and was further expanded to control operators defined on $\ell^2(\mathbb{N})$ by Hansen and Weiss [21]. Generalizations of these results to the context of L^p -admissibility for diagonal semigroups on state spaces of ℓ^q -type using Laplace–Carleson embeddings can be found in works by Haak [15], by Unteregge [47] by Jacob, Partington and Pott [31] and by Jacob, Partington, Pott, Rydhe and Schwenninger [32]. In this frame of reference we also mention that extensions to normal semigroups have appeared in Weiss [48], Hansen and Weiss [21], Unteregge [47] and Staffans [42].

Another property of note which is used to derive results for admissibility is the Weiss property and its L^p -version. For $p \in [1, \infty]$ the *p-Weiss property for control operators* for the generator A is the statement that for any Banach space U and any $B \in \mathcal{L}(U, X_{-1})$ we have the equivalence

$$B \text{ is infinite-time } p\text{-admissible} \Leftrightarrow \sup_{\operatorname{Re} \lambda > 0} (\operatorname{Re} \lambda)^{\frac{1}{p}} \|R(\lambda, A)B\| < \infty;$$

for the analogous formulation for observation operators see Section 5 in the sequel.

The property originates from conjectures formulated by Weiss in [52], which postulated the equivalence of a resolvent condition and admissibility. In [52] the author remarked that the conjecture is not true on general state spaces X that are not Hilbert spaces. Later it was shown that the conjecture is false even in the Hilbert space setting. For counterexamples see Jacob, Partington and Pott [30] based on the right shift semigroup; compare with Zwart and Jacob [54] and Zwart, Jacob and Staffans [55] with counterexamples where the semigroups are even analytic. However—in special cases—positive results on the Weiss property were also found, e.g. by Jacob and Partington [29] for contraction semigroups with $p = 2$ and finite-dimensional input spaces, and by Le Merdy [35] in the case of 2-admissibility for bounded analytic semigroups satisfying a square function estimate and arbitrary input spaces.

In conjunction with Le Merdy’s article [35] the connection with functional calculus methods and square function estimates must also be mentioned, which are used as a powerful tool to generate estimates in the admissibility context. With these methods, Le Merdy showed that if A has dense range and generates a bounded analytic semigroup, then admissibility of $(-A)^{1/2}$ in the observation operator sense—which is a square function estimate for A —is equivalent to A having the 2-Weiss property. That is to say that $C: X_1 \rightarrow Y$ is 2-observation-admissible for A if and only if

$$\sup_{\operatorname{Re} \lambda > 0} \sqrt{\operatorname{Re} \lambda} \|CR(\lambda, A)\| < \infty.$$

Note that for Hilbert spaces, the Weiss property is specifically satisfied if A is densely defined and generates an analytic contraction semigroup.

Further results in this direction appear in the work [20] by Haak and Le Merdy, where the H^∞ functional calculus is used to show the equivalence of certain square function estimates and weighted 2-admissibility for analytic semigroups. The functional calculus also is an integral part of the dissertation [14] by Haak, where extensions of Le Merdy’s results to L^p -admissibility are treated with L^p estimates in lieu of the square function estimates that appear in the L^2 setting. In particular, p -admissibility is characterized under the assumption that $(-A)^{1/p}$ is p -admissible for A . The article [19] by Haak and Kunstmann features generalized results in

this direction. Specifically, the authors investigate the more general property of weighted L^p -admissibility of type α . Here so-called L^p_* functional calculus estimates are used, generalizing the square function estimates appearing in the condition of admissibility of $(-A)^{1/2}$. With this assumption, the authors prove that the Weiss resolvent condition in the L^p version is equivalent to p -admissibility of B . They also show that $-A$ has L^p_* estimates if and only if the state space X embeds into the real interpolation space $(\dot{X}_{-1}, \dot{X}_1)_{1/2, p}$. This leads to results for concrete example systems if full expressions for these interpolation spaces are known, such as the Neumann controlled heat equation on L^q spaces or zero smoothness Besov spaces $B^0_{q, p}$. Furthermore, a test for the resolvent condition in the Weiss property via real interpolation spaces is given.

For more applications of functional calculus methods to proofs of admissibility, we refer to Haak [16], where the interplay of the Weiss condition and admissibility with respect to the weak-type space $L^{2, \infty}$ is discussed. Adding to the positive results on the Weiss conjecture, in [4] the authors Bounit, Driouich and El-Mennaoui characterize the p -Weiss condition for bounded analytic semigroups by admissibility of $(-A)^{1/p}$ without resorting to the H^∞ functional calculus.

Finally we remark that the above list is not exhaustive, as there are even more methods of showing L^p -admissibility such as semigroup generation criteria. These are methods based on block operator matrices generating C_0 -semigroups; see Grabowski and Callier [11], Engel [8] and the recent article [25] by Hofeld, Jacob and Schwenninger, which treats the more general case of admissibility with respect to Orlicz spaces. We further stress that ∞ -admissibility, which is formally very closely related to the notion of input-to-state stability (ISS), see e.g. the survey article [37], is not the focus of the present note. However, as p -admissibility for any finite p implies ∞ -admissibility, the discussed results serve as natural sufficient conditions for ISS.

1.4 Aims

The methods of verifying p -admissibility—for $p \neq 2$ —that we set our focus on can be grouped into

- (i) sufficient conditions for analytic semigroups using mapping properties of boundary trace operators, see e.g. [34, 36, 41],
- (ii) tests arising from Laplace–Carleson embeddings for diagonal semigroups with finite-dimensional input spaces, see e.g. [15, 31, 32, 47],
- (iii) applying p -Weiss property type results in cases where the property is valid, see e.g. [4, 14, 18, 19, 35, 52].

Our aim is to provide a comparison—with some extensions—of the aforementioned methods of verifying p -admissibility while reviewing applicability to infinite-dimensional input spaces and potential problems in applications. In this setting, we also comment on the case $p > 2$ posing even more difficulties than the case $p \leq 2$.

Elaborating on this, we discuss that (i) even leads to a sort of characterization of p -admissibility for values of p strictly greater than a certain threshold, which depends on the interpolation space that the control operator B maps into, see also [36].

For (ii) we note that L^p -admissibility of the input element $b \in \mathcal{L}(\mathbb{C}, X_{-1})$ is equivalent to boundedness of the Laplace transform operator $\mathcal{L}$, which is to be understood as a mapping

$$\mathcal{L}: L^p([0, \infty)) \rightarrow L^2(\mathbb{C}_+, \mu).$$

The measure μ is given by the discrete measure $\mu = \sum_k |b_k|^2 \delta_{-\lambda_k}$ in the diagonal case, i.e. in the presence of a Riesz basis of eigenvectors of the generator with eigenvalues $\{\lambda_k\}_{k \in \mathbb{N}}$, cf. results by Jacob, Partington and Pott [31]. However, the conditions given in their work are not applicable if the generator does not have compact resolvents as the spectrum is then not discrete. For instance, if the generator A is a normal operator, the spectral theorem yields a spectral measure E (that is not necessarily discrete) associated to A , admitting the representation

$$A = \int_{\sigma(A)} z \, dE(z).$$

Using this fact, we are able to generalize the diagonal results. The extensions are established using a suitable generalization of the discrete measure μ related to the measure $\langle Eb, b \rangle$; see also similar considerations for the special case $p = 2$ in [21, 42, 47, 48]. We also discuss a related result for systems with semigroups generated by multiplication operators on L^q spaces, generalizing the setting where one has a q -Riesz basis. This is related to diagonal systems on ℓ^q sequence spaces as in [31].

As an outlook, and relating to (iii), we provide an argument based on the p -Weiss property which allows the extension of results for scalar input spaces to infinite-dimensional input spaces. In certain special cases where the p -Weiss property does hold, such as for negative semidefinite self-adjoint generators when $p \leq 2$, this provides a tool to assess p -admissibility for the critical value of p ; that is to say the value of p that defines the threshold between admissibility and non-admissibility.

2 Interpolation spaces, extrapolation spaces and admissibility

2.1 Mapping properties and their relation to admissibility

Here and throughout the rest of the paper, we denote the resolvent of an operator A by $R(z, A) := (zI - A)^{-1}$ and the range of A by the symbol $\mathcal{R}(A)$. Moreover, we denote the *growth bound* of the semigroup T by $\omega(T)$ and we recall that T is called *exponentially stable* if $\omega(T) < 0$. A C_0 -semigroup T is called *analytic* if there exists $\varphi \in (0, \pi/2]$ such that T can be extended to an analytic map on the open sector of angle 2φ around the positive real line defined by $S_\varphi := \{z \in \mathbb{C} \setminus \{0\} : |\operatorname{Arg} z| < \varphi\}$. In the presence of an analytic semigroup T and a generator A such that $0 \in \rho(A)$, we define a scale of interpolation and extrapolation spaces X_β for $\beta \in (-1, 1)$. These spaces lie between the spaces $X_1 = \mathcal{D}(A)$, $X_0 := X$ and X_{-1} . Using this scale of spaces allows for the description of finer mapping properties of control operators. This is meant in the sense that if B maps into a space $X_{-\beta}$ for some $\beta \in (0, 1)$, then we can use this information to make statements about admissibility. In defining these spaces, we make use of fractional powers $(-A)^{-\beta}: X \rightarrow X$ which are defined by the operator-valued integral

$$(-A)^{-\beta} := \int_{\gamma} z^{-\beta} R(z, -A) \, dz,$$

where the curve γ is the boundary of a sector of the form S_φ as above that contains $\sigma(-A)$, see also [9, Chapter II.5] and [41]. The corresponding positive power $(-A)^\beta$ is then defined as the inverse of $(-A)^{-\beta}$, i.e.

$$(-A)^\beta := ((-A)^{-\beta})^{-1} : \mathcal{R}((-A)^{-\beta}) \rightarrow X.$$

Let A generate an analytic semigroup on X with growth bound less than 0. For any $\beta \in (0, 1)$ define the space $X_{-\beta}$ as the completion of X with respect to the norm given on X by $\|x\|_{-\beta} := \|(-A)^{-\beta}x\|_X$. Furthermore, we define the space X_β as the set $\mathcal{D}((-A)^\beta) = \mathcal{R}((-A)^{-\beta})$ with the norm $\|x\|_\beta := \|x\|_X + \|(-A)^\beta x\|_X$. These spaces are also known as *abstract Sobolev spaces*. For general analytic semigroups the spaces X_β are defined by considering the shifted generator $A - \lambda I$ for sufficiently large λ , noting that this definition is independent of the choice of λ and hence also consistent.

In the following we present a characterization of p -admissibility that is derived from the mapping property that B be bounded into $X_{-\beta}$ for some $\beta \in (0, 1)$, compare also [36]. We also record that for $p \in [1, \infty]$ we use the notation p' for the Hölder conjugate (or conjugate exponent) satisfying $1/p + 1/p' = 1$ with the usual convention for the conjugated exponents of 1 and ∞ . The following lemma is essentially well-known in the context of admissibility for analytic semigroups. For the sake of the reader we provide the argument.

Lemma 2.1. *Let A be the generator of an analytic semigroup T on X . Fix $0 < \beta < 1$. If B is bounded as an operator $U \rightarrow X_{-\alpha}$ for all $\alpha > \beta$, then B is finite-time p -admissible for all $p > \frac{1}{1-\beta} = (\beta^{-1})'$.*

Proof. We fix $0 < \beta < 1$ and let $B \in \mathcal{L}(U, X_{-1})$. Without loss of generality, assume that T is exponentially stable. Assume $B \in \mathcal{L}(U, X_{-\alpha})$ for an arbitrary $\alpha > \beta$. Then $(-A)^{-\alpha}B \in \mathcal{L}(U, X)$. Note that for A generating an analytic semigroup we have

$$\|(-A)^\alpha T(t)\|_{\mathcal{L}(X)} \leq C_\alpha t^{-\alpha} e^{t\omega}$$

for $t > 0$ and $\omega > \omega(T)$ and some constant $C_\alpha > 0$, see [38, Theorem 6.13]. Using these properties, we can deduce that

$$\begin{aligned} \|T(t)B\|_{\mathcal{L}(U, X)} &= \|T(t)(-A)^\alpha (-A)^{-\alpha}B\|_{\mathcal{L}(U, X)} \\ &\leq \|T(t)(-A)^\alpha\|_{\mathcal{L}(X)} \|(-A)^{-\alpha}B\|_{\mathcal{L}(U, X)} \\ &\leq C_\alpha t^{-\alpha} e^{t\omega} \|(-A)^{-\alpha}B\|_{\mathcal{L}(U, X)} \\ &\leq C_\alpha t^{-\alpha} e^{t\omega} \|B\|_{\mathcal{L}(U, X_{-\alpha})} \end{aligned}$$

holds for $t > 0$ and $\omega > \omega(T)$. Applying this to the left-hand side of the defining inequality for L^p -admissibility, we get

$$\begin{aligned} \left\| \int_0^t T(t-s)Bu(s) \, ds \right\|_X &\leq \int_0^t \|T(t-s)Bu(s)\|_X \, ds \\ &\leq C_\alpha \int_0^t (t-s)^{-\alpha} e^{(t-s)\omega} \|B\|_{\mathcal{L}(U, X_{-\alpha})} \|u(s)\|_U \, ds. \end{aligned}$$

We can now use Hölder's inequality to further estimate this quantity and find

$$\int_0^t (t-s)^{-\alpha} e^{(t-s)\omega} \|u(s)\|_U \, ds \leq \|u\|_{L^p([0, t], U)} \left(\int_0^t ((t-s)^{-\alpha} e^{(t-s)\omega})^{p'} \, ds \right)^{1/p'}.$$

The right-hand side is finite if and only if $p' < \frac{1}{\alpha}$. Combining the two above inequalities, we get

$$\left\| \int_0^t T(t-s)Bu(s) \, ds \right\|_X \leq K \|u\|_{L^p([0,t],U)}$$

for $p' < \frac{1}{\alpha}$ and some constant $K > 0$. Thus B is p -admissible for all $p > \frac{1}{1-\beta}$. $\square$

We want to show that the converse of this statement also holds. To do this, we need to introduce the scale of Favard spaces, which are another useful class of intermediate spaces as there are embeddings that allow us to derive L^p estimates. We give a short summary of the needed results and properties of these spaces.

Definition 2.2 (Favard spaces [9]). Let T be an exponentially stable C_0 -semigroup. For $0 < \alpha \leq 1$ we define the α -Favard norm by

$$\|x\|_{F_\alpha} := \sup_{t>0} t^{-\alpha} \|T(t)x - x\|_X$$

and the Favard space F_α as the set of all $x \in X$ with finite α -Favard norm.

To extend the definition to arbitrary real indices α , we decompose the number $\alpha \in \mathbb{R}$ into an integer part and a fractional part; that is to say we write $\alpha = k + \beta$ (uniquely) with $k \in \mathbb{Z}$ and $0 < \beta \leq 1$ and define F_α to be the Favard space of order β for the extended (resp. restricted) semigroup T_k . Moreover, if T is not exponentially stable, we define F_α as the α -Favard space corresponding to the semigroup $(e^{-\omega t}T(t))_{t \geq 0}$ for some $\omega > \omega(T)$.

Note that this definition yields the same space for any such choice of ω ; for more on this topic we refer to Engel and Nagel [9, Chapter II.5].

We now quote results about Favard spaces needed in the sequel.

Remark 2.3. For A generating a C_0 -semigroup T the following hold:

1. If $\alpha > \beta$, we have

$$X_\alpha \subset F_\alpha \hookrightarrow X_\beta \subset F_\beta,$$

cf. [9, Proposition II.5.14, Proposition II.5.33]; note that X_α is defined alternatively there.

2. If T is exponentially stable, the α -Favard norm with $0 < \alpha \leq 1$ is equivalent to

$$\|x\|_{F_\alpha} := \sup_{\lambda>0} \lambda^\alpha \|AR(\lambda, A)x\|_X,$$

cf. [9, Proposition II.5.12].

We give a characterization of the property that B maps into extrapolation spaces of the form $X_{-\beta}$ for $\beta \in (0, 1)$ by admissibility of B , providing a counterpart to Lemma 2.1 above. In this context we note that similar results have appeared in [36].

Lemma 2.4. *Let A generate an analytic semigroup T on X . Let $\beta \in (0, 1)$ and assume that for all $p > \frac{1}{1-\beta}$ the resolvent condition in the p -Weiss condition holds for A and a given $B \in \mathcal{L}(U, X_{-1})$, i.e.*

$$\exists \alpha > \omega(T): \sup_{\operatorname{Re} \lambda > \alpha} (\operatorname{Re} \lambda)^{1/p} \|R(\lambda, A)B\|_{\mathcal{L}(U, X)} < \infty. \quad (6)$$

Then B is bounded as a mapping $B: U \rightarrow X_{-\alpha}$ for all $\alpha > \beta$.

Proof. Again we assume without loss of generality that T is exponentially stable. For a fixed $q > \frac{1}{1-\beta}$ we have by the resolvent condition (6) that

$$\|R(\lambda, A)Bu_0\|_X \leq K(\operatorname{Re} \lambda)^{-1/q}\|u_0\|_U, \quad \operatorname{Re} \lambda > 0.$$

Using the extension A_{-1} of the generator A , this implies

$$(\operatorname{Re} \lambda)^{1/q}\|AR(\lambda, A)Bu_0\|_{X_{-1}} \leq K\|u_0\|_U \quad (7)$$

for all $\lambda \in \mathbb{C}$ with $\operatorname{Re} \lambda > 0$. By exponential stability of T and $0 < \gamma < 1$, Item 2 in Remark 2.3 implies that we can consider the norm

$$\|x\|_{F_{-\gamma}} = \sup_{\lambda > 0} \lambda^{1-\gamma}\|AR(\lambda, A)x\|_{X_{-1}}, \quad x \in X_{-1},$$

which is equivalent to the usual norm on $F_{-\gamma} \subset X_{-1}$. Recall that for negative parameters the Favard space was defined in terms of the extended generator on X_{-1} . Using (7) we get

$$\|Bu_0\|_{F_{-(1-1/q)}} = \sup_{\lambda > 0} \lambda^{1-(1-1/q)}\|AR(\lambda, A)Bu_0\|_{X_{-1}} \leq K\|u_0\|_U.$$

Relating this to abstract Sobolev spaces, notice that for any choice of $\alpha \in (\beta, 1]$ there exists a $q > 1$ such that $\alpha > 1 - \frac{1}{q} > \beta$. In fact, a possible choice is $q = (1 - \frac{\alpha+\beta}{2})^{-1}$. Embedding relations between Favard and abstract Sobolev spaces from Remark 2.3 then imply $F_{-(1-1/q)} \hookrightarrow X_{-\alpha}$. As a consequence, we also have

$$\|Bu_0\|_{X_{-\alpha}} \leq K_1\|Bu_0\|_{F_{-(1-1/q)}} \leq K_2\|u_0\|_U$$

for all $\alpha > \beta$. Hence we have shown that $B \in \mathcal{L}(U, X_{-\alpha})$. $\square$

Note that Lemma 2.4 is in particular applicable if B is even infinite-time p -admissible for all $p > \frac{1}{1-\beta}$. Indeed, specifying the input function $u = e^{-\lambda(\cdot)}u_0$ for $u_0 \in U$ and using the representation of the resolvent as the Laplace transform of the semigroup, the definition of infinite-time p -admissibility yields

$$\|R(\lambda, A)Bu_0\|_X \leq K(\operatorname{Re} \lambda)^{-1/p}\|u_0\|_U, \quad \lambda \in \rho(A).$$

The next theorem combines Lemmas 2.1 and 2.4 into a characterization of p -admissibility of B for p greater than a threshold in terms of boundedness of B into spaces of type $X_{-\alpha}$.

Theorem 2.5. *Let A be the generator of an analytic and exponentially stable semigroup T on X and let $B \in \mathcal{L}(U, X_{-1})$ be a control operator. Then the following are equivalent for $\beta \in (0, 1)$:*

- (a) B is finite-time L^p -admissible for all $p > \frac{1}{1-\beta}$.
- (b) B is a bounded map into $X_{-\alpha}$ for all $\alpha > \beta$.

Proof. If $B \in \mathcal{L}(U, X_{-1})$ is L^p -admissible for all $p > \frac{1}{1-\beta}$, then B is also infinite-time L^p -admissible for all $p > \frac{1}{1-\beta}$ due to the assumption of exponential stability. Since in this case the resolvent condition (6) holds for all $p > \frac{1}{1-\beta}$, we have $B \in \mathcal{L}(U, X_{-\alpha})$ for $\alpha > \beta$ by Lemma 2.4. The implication (b) $\Rightarrow$ (a) directly follows from an application of Lemma 2.1. $\square$

The statement in the previous theorem does not hold in the endpoint case $p = \frac{1}{1-\beta}$ and $\alpha = \beta$, as the following result shows.

Proposition 2.6. *For any $p \in (1, \infty)$ there exists an analytic semigroup generator on $X = \ell^2$ and a p -admissible $B : \mathbb{C} \rightarrow X_{-1}$ that is not in $\mathcal{L}(\mathbb{C}, X_{-1/p'})$.*

Proof. We adapt an example given in [46, Example 5.3.11] to $p \neq 2$. In fact, we let $1 < p < \infty$ be arbitrary. As in [46, Example 5.3.11], we consider the semigroup generated by the diagonal operator A , that is, $Ae_k = \lambda_k e_k$, $k \in \mathbb{N}$; with $\{e_k\}_{k \in \mathbb{N}}$ referring to the canonical basis and with eigenvalue sequence defined by $\lambda_k = -2^k$ for $k \in \mathbb{N}$.

Due to the identification of $\mathcal{L}(\mathbb{C}, X_{-1})$ with the space X_{-1} itself, the claim amounts to existence of a sequence $b \in X_{-1}$ such that b is p -admissible but fails to be an element of $X_{-1/p'}$. We consider the cases $1 < p \leq 2$ and $2 < p < \infty$ separately and use the appropriate Carleson measure criteria from [31, Theorem 3.2, Theorem 3.5]; see also Theorem 3.2 below.

Let $1 < p \leq 2$ and set $b = \{2^{k/p'}\}_{k \in \mathbb{N}}$. This choice of b represents an element of X_{-1} as

$$\sum_{k=1}^{\infty} \frac{|b_k|^2}{|\lambda_k|^2} = \sum_{k=1}^{\infty} 2^{2k/p'} 2^{-2k} = \sum_{k=1}^{\infty} 2^{-2k/p} < \infty.$$

To prove p -admissibility using the Carleson measure criterion we need to show that there exists $K > 0$ such that $\mu(Q_I) \leq K|I|^{2/p'}$, where $\mu = \sum_k |b_k|^2 \delta_{-\lambda_k}$ and Q_I is the Carleson square defined for an interval I on the imaginary axis symmetric about 0 (an interval of the form $i[-a, a]$ for some $a > 0$) by

$$Q_I := \{z \in \mathbb{C} : i \operatorname{Im} z \in I, 0 < \operatorname{Re} z < |I|\},$$

see also Figure 1 below.




Figure 1: The Carleson square Q_I for the interval $I = i[-a, a]$

We therefore proceed by showing that there exists $K > 0$ with

$$\sum_{k \in \mathbb{N}} 2^{2k/p'} \delta_{2^k}((0, |I|)) \leq K|I|^{2/p'}$$

for all choices of I as above. Since $|I| \geq 2^{\lfloor \log_2 |I| \rfloor}$ we set $m = \lfloor \log_2 |I| \rfloor$ and get

$$\begin{aligned} \sum_{k \in \mathbb{N}} 2^{2k/p'} \delta_{2^k}((0, |I|)) &= \sum_{k=1}^m (2^{2/p'})^k = \frac{2^{2/p'}}{2^{2/p'} - 1} ((2^m)^{2/p'} - 1) \\ &\leq K_{p'} |I|^{2/p'}, \end{aligned}$$

which implies p -admissibility of b .

Now, if b were in $X_{-1/p'}$, then we would have $(-A)^{-1/p'}b \in X$. But due to $((-A)^{-1/p'}b)_k \equiv 1$, this condition is violated and it follows that $b \notin \mathcal{L}(\mathbb{C}, X_{-1/p'})$.

In the case $p > 2$ we consider the same semigroup, but now we let b be the control operator represented by $b_k = k^{-1/2}2^{k/p'}$, $k \in \mathbb{N}$. We have $b \in X_{-1}$ since

$$\sum_{k=1}^{\infty} \frac{|b_k|^2}{|\lambda_k|^2} = \sum_{k=1}^{\infty} k^{-1} 2^{2k/p'} 2^{-2k} \leq \sum_{k=1}^{\infty} 2^{-2k/p} < \infty.$$

In this case, the Carleson measure criterion reads

$$\{2^{-2n/p'} \mu(S_n)\}_{n \in \mathbb{N}} \in \ell^{p/(p-2)}(\mathbb{N})$$

where $\mu = \sum_{k \in \mathbb{N}} |b_k|^2 \delta_{-\lambda_k}$ and $S_n = \{z \in \mathbb{C} : \operatorname{Re} z \in (2^{n-1}, 2^n]\}$.

For this choice of semigroup and control operator the condition means that

$$\sum_{k=1}^{\infty} \left| k^{-1/2} 2^{k/p'} \right|^{\frac{2p}{p-2}} \cdot 2^{-\frac{2k}{p'} \frac{p}{p-2}} < \infty,$$

which holds true here as

$$\sum_{k=1}^{\infty} \left| k^{-1/2} 2^{k/p'} \right|^{\frac{2p}{p-2}} \cdot 2^{-\frac{2k}{p'} \frac{p}{p-2}} = \sum_{k=1}^{\infty} k^{-\frac{p}{p-2}} < \infty.$$

However,

$$\|(-A)^{-1/p'} b\|_{\ell^2} = \sum_{k=1}^{\infty} 2^{-2k/p'} |k^{-1/2} 2^{k/p'}|^2 = \sum_{k=1}^{\infty} k^{-1}$$

with a divergent sum; thereby showing that b is unbounded as an operator $\mathbb{C} \rightarrow X_{-1/p'}$. $\square$

2.2 Example: admissibility for the heat equation via interpolation

As indicated before, we are interested in L^p -admissibility for the heat equation on the spatial domain $[0, 1]$ with Dirichlet control at the boundary point 1, given through the system's equations

$$\begin{cases} x_t = x_{\xi\xi} & \text{in } (0, 1) \times (0, \infty), \\ x = 0 & \text{on } \{\xi = 0\} \times (0, \infty), \\ x = u & \text{on } \{\xi = 1\} \times (0, \infty), \\ x = x_0 & \text{on } (0, 1) \times \{t = 0\}, \end{cases}$$

with initial heat distribution given by x_0 and heat injection with magnitude $u(t)$ at the point 1.

In this simple example, all properties of the system can basically be derived from the solution formulae given through the spectral decomposition. Yet, it serves as the most natural and well-known example where 2-admissibility is not satisfied if the state space is chosen to be $X = L^2([0, 1])$, see also [34]. We formulate this as a boundary control system by defining

$$\mathfrak{A}: X \supset \mathcal{D}(\mathfrak{A}) \rightarrow X, \quad f \mapsto f''$$

on the domain $\mathcal{D}(\mathfrak{A}) = \{f \in H^2([0, 1]) : f(0) = 0\}$. The abstract boundary operator is then given by

$$\mathfrak{B} = \delta_1 : X \supset \mathcal{D}(\mathfrak{A}) \rightarrow U, \quad f \mapsto f(1).$$

The Sobolev embedding theorem (see e.g. [1, Theorem 4.12]) implies that each H^2 function in spatial dimension $n = 1$ is continuous, ensuring that $\mathfrak{B}$ is well-defined. We want to translate this system into state space form to make statements about admissibility. Define the operator A by

$$A = \mathfrak{A}|_{\ker \mathfrak{B}} : \mathcal{D}(A) \rightarrow X, \quad f \mapsto f''$$

on the domain $\mathcal{D}(A) = \{f \in H^2([0, 1]) : f(0) = f(1) = 0\}$. Using the identity (5)—as both X and U are Hilbert spaces—one integrates by parts twice to find that $\mathfrak{B}x \cdot B^*y = -x(1)y'(1)$ for all $x \in \mathcal{D}(\mathfrak{A})$ and $y \in \mathcal{D}(A)$. Thus we can identify that B is given by

$$B = \delta'_1 : \mathbb{C} \rightarrow X_{-1}, \quad z \mapsto z\delta'_1,$$

where $\delta'_1(f) = -f'(1)$.

To proceed, we consider the orthonormal basis $\{e_n : n \in \mathbb{N}\}$ with $e_n(\xi) = \sqrt{2} \sin(n\pi\xi)$. Then the diagonal representation $Ae_n = \lambda_n e_n$ holds with $\lambda_n = -n^2\pi^2$. The control operator B can be represented by the sequence $\{b_n\}_{n \in \mathbb{N}}$ for $b_n = (-1)^n \sqrt{2}n\pi$ as in [27, Example 5.1].

Using the representation of the system in diagonal form, we show that $b = \{b_n\}_{n \in \mathbb{N}}$ is an element of the interpolation space $X_{-\beta}$ for $\beta > 3/4$. By Lemma 2.1, this then will imply that the control operator is p -admissible for all $p > \frac{1}{1-3/4} = 4$. It remains to show that

$$\sum_{n=1}^{\infty} \frac{|b_n|^2}{|\lambda_n|^{2\beta}} < \infty$$

for all $\beta > \frac{3}{4}$. We have

$$\sum_{n \in \mathbb{N}} \frac{|b_n|^2}{|\lambda_n|^{2\beta}} = \sum_{n \in \mathbb{N}} \frac{2n^2\pi^2}{\pi^{4\beta}n^{4\beta}} = \frac{2}{\pi^{4\beta-2}} \sum_{k \in \mathbb{N}} \frac{1}{n^{4\beta-2}},$$

which is finite for $\beta > \frac{3}{4}$. Invoking Lemma 2.1 then implies p -admissibility for all $p > 4$.

Remark 2.7. One can also use the interpolation space argument to show that the Dirichlet controlled heat equation is admissible for $p > 4$ in multiple spatial dimensions. To show this, we follow [41] and fix a dimension $n \geq 2$ and a domain $\Omega \subset \mathbb{R}^n$ with sufficiently smooth boundary $\partial\Omega$. The heat equation with Dirichlet boundary control is represented by the system

$$\begin{cases} \dot{x}(t) = \Delta x(t) \\ \gamma_0 x = u, \\ x(0) = x_0 \in X \end{cases}$$

on $X = L^2(\Omega)$ with input space $U = L^2(\partial\Omega)$, where γ_0 is the Dirichlet trace operator on $\partial\Omega$. Relating the above to abstract boundary control systems, we set

$$\mathfrak{A} = \Delta, \quad \mathcal{D}(\mathfrak{A}) = H^2(\Omega) + DL^2(\partial\Omega), \quad \mathfrak{B} = \gamma_0,$$

where $D \in \mathcal{L}(L^2(\partial\Omega), L^2(\Omega))$ denotes the Dirichlet map which maps $g \in L^2(\partial\Omega)$ to the unique solution $h \in L^2(\Omega)$ of

$$\Delta h = 0, \quad \gamma_0 h = g;$$

see [46, Proposition 10.6.6]. Since by [46, Proposition 10.6.4] the composition $\gamma_0 D$ yields the identity on $L^2(\partial\Omega)$, we have $\ker \mathfrak{B} = H^2(\Omega) \cap H_0^1(\Omega)$. The general rewriting procedure for boundary control systems leads to the self-adjoint operator

$$A = \mathfrak{A}|_{\ker \mathfrak{B}} = \Delta, \quad \mathcal{D}(A) = H^2(\Omega) \cap H_0^1(\Omega).$$

As X and U are Hilbert spaces, we use the formula (5) and Green's first identity to deduce that

$$\langle \mathfrak{A}x, y \rangle_X - \langle x, Ay \rangle_X = \langle \mathfrak{B}x, B^*y \rangle_U = -\langle \mathfrak{B}x, \frac{\partial y}{\partial \nu} \rangle_U,$$

whereupon we find that B^*y is the normal derivative operator $\frac{\partial y}{\partial \nu}$ on $\partial\Omega$. By [34, Lemma 3.1.1], B^* is bounded $H^\beta(\Omega) \cap H_0^1(\Omega) \rightarrow L^2(\partial\Omega)$ for all $\beta > \frac{3}{2}$. Hence, for all $\beta > 3/4$ this yields

$$B^* \in \mathcal{L}(X_\beta, U^*) = \mathcal{L}(H^{2\beta}(\Omega) \cap H_0^1(\Omega), L^2(\partial\Omega)), \quad B^*: f \mapsto \frac{\partial f}{\partial \nu} \Big|_{\partial\Omega}.$$

By duality—as $X = L^2(\Omega)$ is reflexive—we conclude that $(-A)^{-\beta} B$ is bounded as an operator $U \rightarrow X$ for all $\beta > 3/4$. Lemma 2.1 then implies that B is p -admissible for all $p > 4$.

Remark 2.8 (Notes on Neumann control). Although it does not appear using the notion of *admissibility* explicitly, we remark that the heat equation on the state space $X = L^2(\Omega)$ with Neumann control acting via the boundary—and inputs in $U = L^2(\partial\Omega)$ —is featured in the book by Lasiecka and Triggiani [34, Remark 3.3.1.4] (and the result is folklore). There the control operator B is computed and shown to satisfy $(-A)^{-(1/4+\varepsilon)} B \in \mathcal{L}(U, X)$ for any $\varepsilon > 0$. This allows the application of the interpolation space argument as in Section 2.1.

Furthermore, p -admissibility of the Neumann controlled heat equation is characterized by an embedding argument due to Haak and Kunstmann, see [19, Proposition 2.4]. We also refer to Byrnes, Gilliam, Shubov and Weiss [5] for the case $p = 2$. Denoting the Neumann Laplacian on $\Omega \subset \mathbb{R}^n$ by Δ_N , it follows from Haak and Kunstmann's results that the control operator $B = \gamma_0'$ with input space $L^2(\partial\Omega)$ —resulting from the Neumann controlled heat equation on $X = L^2(\Omega)$ —is p -admissible for $A = -\Delta_N + I$ if and only if $p \geq \frac{4}{3}$. This is a special case of a more general result they obtain in the context of weighted admissibility on state spaces such as $L^q(\Omega)$. In the unweighted case that we consider here, the result yields p -admissibility whenever U embeds into the Besov space $B_{q,\infty}^{2/p-1-1/q}(\partial\Omega)$. We employ the Besov space embedding

$$B_{2,2}^0 \hookrightarrow B_{2,\infty}^{-s}, \quad s \geq 0,$$

see e.g. [45, Section 2.3.2, Remark 2] or [40, Section 2.3], such that in the case of the Neumann controlled heat equation on $X = L^2(\Omega)$ with $U = L^2(\partial\Omega)$ one arrives at the condition

$$U = B_{2,2}^0(\partial\Omega) \hookrightarrow B_{2,\infty}^{2/p-3/2}(\partial\Omega)$$

which is satisfied if and only if $p \geq \frac{4}{3}$, as claimed.

3 Characterizations for multiplier-like systems

We are interested in testing p -admissibility also in cases where the system is not necessarily of diagonal form. In this section, we formulate extensions of Laplace–Carleson type results to normal semigroups and multiplication semigroups. For this, recall the following result due to Jacob, Partington and Pott for diagonal semigroups.

Theorem 3.1 ([31, Theorem 2.1]). *Let A generate a diagonal semigroup on $X = \ell^q(\mathbb{Z})^1$ with eigenvalues given by $\{\lambda_k\}_{k \in \mathbb{Z}}$. Let $p \in [1, \infty)$. Then $B = \{b_k\}_{k \in \mathbb{Z}} \in \mathcal{L}(\mathbb{C}, X_{-1})$ is infinite-time L^p -admissible if and only if the Laplace transform*

$$\mathcal{L}: L^p([0, \infty)) \rightarrow L^q(\mathbb{C}_+, \mu), \quad f \mapsto \left(z \mapsto \int_0^\infty e^{-zt} f(t) dt \right)$$

is a bounded operator for the measure

$$\mu = \sum_{k \in \mathbb{Z}} |b_k|^q \delta_{-\lambda_k}.$$

Note that by using the special form of a diagonal semigroup the proof of the above result is rather direct and the difficulty in assessing admissibility only transfers to checking the boundedness of the Laplace–Carleson embedding. This is where arguments from harmonic analysis, and in particular from the theory of Carleson measures, take effect—at least under further assumptions on the range of p and the location of the eigenvalues (in sectors or strips). The latter situations reflect whether the diagonal semigroup is analytic or even a group. In those cases, admissibility indeed reduces to elementary conditions on certain sequences lying in corresponding ℓ^r -spaces, such as employed in Section 2.1 above and Section 3.1 below.

For convenience, we recall the admissibility results for diagonal semigroups on ℓ^q spaces from [31].

Proposition 3.2 ([31, Theorem 3.2, Theorem 3.5]). *Fix $1 \leq q < \infty$ and assume that $A: \ell^q(\mathbb{Z}) \supset \mathcal{D}(A) \rightarrow \ell^q(\mathbb{Z})$ generates a diagonal semigroup on $X = \ell^q(\mathbb{Z})$ with eigenvalues $\{\lambda_k\}_{k \in \mathbb{Z}}$ lying in the left half plane such that $-\lambda_k \in S_\theta$ for some angle $\theta \in (0, \pi/2)$ and all $k \in \mathbb{Z}$. Consider the operator $B \in \mathcal{L}(\mathbb{C}, X_{-1})$ represented by the sequence $\{b_k\}_{k \in \mathbb{N}}$.*

If $1 < p \leq q$, then the following statements are equivalent:

1. *B is p -admissible.*
2. *There exists $K > 0$ such that*

$$\mu(Q_I) \leq K |I|^{q/p'}$$

holds for all intervals $I \subset i\mathbb{R}$ symmetric about 0; i.e. intervals of the form $i[-a, a]$ for some $a > 0$, where $Q_I := \{z \in \mathbb{C} : i \operatorname{Im} z \in I, 0 < \operatorname{Re} z < |I|\}$.

3. *There exists $K > 0$ such that $\|R(z, A)B\|_X \leq K z^{-1/p}$ for all $z > 0$.*

Here $Q_I = \{z \in \mathbb{C} : i \operatorname{Im} z \in I, 0 < \operatorname{Re} z < |I|\}$.

If instead $q < p < \infty$, the following are equivalent:

1. *B is p -admissible.*
2. *$\{2^{-nq/p'} \mu(S_n)\}_{k \in \mathbb{Z}} \in \ell^{p/(p-q)}(\mathbb{Z})$.*
3. *$\{2^{n/p} \|R(2^n, A)B\|_{\ell^q}\}_{k \in \mathbb{Z}} \in \ell^{p/(p-q)}(\mathbb{Z})$.*

Here $\mu = \sum_{k \in \mathbb{N}} |b_k|^2 \delta_{-\lambda_k}$ and $S_n = \{z \in \mathbb{C} : \operatorname{Re} z \in (2^{n-1}, 2^n]\}$.

¹More generally, one can assume that the semigroup is diagonal with respect to a q -Riesz basis on an arbitrary Banach space X .

Thus, in this section we are looking for (more general) measures μ such that the equivalence

$$\mathcal{L} \in \mathcal{L}(L^p([0, \infty)), L^q(\mathbb{C}_+, \mu)) \Leftrightarrow \left\| \int_0^\infty T(s)Bu(s) \, ds \right\|_X \leq C \|u\|_{L^p([0, \infty))}$$

holds for normal semigroups and multiplication semigroups.

3.1 Non-admissibility for a Dirichlet controlled heat equation

As an application of the Laplace–Carleson method from [31] we want to show that the Dirichlet controlled heat equation in one spatial dimension as above is not L^p -admissible for any $p \leq 4$.

It is enough to show the claim for $p = 4$. To show that Dirichlet boundary control does not lead to a 4-admissible control operator for this system, we apply the necessary and sufficient condition given by the Laplace–Carleson method from [31, Theorem 3.5], as above in Proposition 2.6 and Proposition 3.2. Keeping the same notation, i.e. $\mu = \sum_{k \in \mathbb{N}} |b_k|^2 \delta_{-\lambda_k}$ and $S_n = \{z \in \mathbb{C} : \operatorname{Re} z \in (2^{n-1}, 2^n]\}$, we insert $p = 4$ into the condition, which yields

$$B \text{ 4-admissible} \Leftrightarrow s := \{2^{-3n/2} \mu(S_n)\}_{n \in \mathbb{Z}} \in \ell^2(\mathbb{Z}).$$

In showing that s is not square-summable, we use the following expression for $\mu(S_n)$: Recall that $\lambda_k = -k^2 \pi^2$ ([27, Example 5.1]). For $n \in \mathbb{Z}$ we have

$$\begin{aligned} 2^{-2n/p'} \mu(S_n) &= 2^{-2n/p'} \sum_{k \in \mathbb{N}} |\sqrt{2}k\pi|^2 \delta_{k^2 \pi^2}(\{z \in \mathbb{C} : \operatorname{Re} z \in (2^{n-1}, 2^n]\}) \\ &= 2^{1-2n/p'} \pi^2 \sum_{k \in \mathbb{N} \cap \left(\frac{\sqrt{2^{n-1}}}{\pi}, \frac{\sqrt{2^n}}{\pi}\right]} k^2. \end{aligned} \tag{8}$$

Combining (8) with the fact that $k \mapsto k^2$ is an increasing function on $[0, \infty)$, we estimate

$$\mu(S_n) = 2\pi^2 \sum_{k \in \mathbb{N} \cap \left(\frac{\sqrt{2^{n-1}}}{\pi}, \frac{\sqrt{2^n}}{\pi}\right]} k^2 \geq 2\pi^2 \int_{[2^{(n-1)/2}/\pi]}^{[2^{n/2}/\pi]} k^2 \, dk = \frac{2\pi^2}{3} \left(\left\lfloor \frac{2^{n/2}}{\pi} \right\rfloor^3 - \left\lfloor \frac{2^{(n-1)/2}}{\pi} \right\rfloor^3 \right).$$

Then we get

$$\mu(S_n) \geq \frac{2\pi^2}{3} \left(\left\lfloor \frac{2^{n/2}}{\pi} \right\rfloor^3 - \left\lfloor \frac{2^{(n-1)/2}}{\pi} \right\rfloor^3 \right)$$

which, in combination with the estimate $\lfloor b \rfloor^3 - \lfloor a \rfloor^3 \geq (b-1)^3 - a^3$, gives

$$\mu(S_n) \geq \frac{2\pi^2}{3} \left(\frac{(1 - 2^{-3/2})}{\pi^3} 2^{3n/2} - \frac{3}{\pi^2} 2^{2n/2} + \frac{3}{\pi} 2^{n/2} - 1 \right).$$

Multiplying both sides with the prefactor $2^{-3n/2}$, we have

$$2^{-3n/2} \mu(S_n) \geq \frac{2\pi^2}{3} \left(\frac{(1 - 2^{-3/2})}{\pi^3} - \frac{3}{\pi^2} 2^{-n/2} + \frac{3}{\pi} 2^{-2n/2} - 2^{-3n/2} \right),$$

where the expression on the right-hand side does not converge to zero in the limit as $n \rightarrow \infty$. Therefore the sequence s is not square-summable and consequently B is not 4-admissible.

3.2 Normal semigroups

Proposition 3.3 (Laplace–Carleson embeddings for normal semigroups). *Let A generate a strongly continuous semigroup of normal operators on a Hilbert space X . Denote the spectral measure of A by E and the spectral measure of the extension A_{-1} by E_{-1} . Consider a control operator $b \in X_{-1} \simeq \mathcal{L}(\mathbb{C}, X_{-1})$ and define the measure*

$$\mu(S) := \int_S (1 + |z|^2) d\langle E_{-1}(z)b, b \rangle_{X_{-1}} = (1 + |z|^2)(E_{-1})_{b,b}(S) \quad (9)$$

on Borel subsets S of $\mathbb{C}$.

Then b is L^p -admissible for T if and only if the Laplace transform is a bounded operator when considered as a map

$$\mathcal{L}: L^p([0, \infty)) \rightarrow L^2(\sigma(-A), \mu).$$

Proof. By the spectral theorem applied to the normal operator A_{-1} , we have the representations

$$(rI - A_{-1}) = \int_{\sigma(-A)} (r - z) dE_{-1}(z) \quad (r \in \mathbb{C}), \quad T(t) = \int_{\sigma(-A)} e^{-tz} dE_{-1}(z).$$

Let u be a locally integrable complex-valued function on $[0, \infty)$. To prove the claim, we show that

$$\|\mathcal{L}u\|_{L^2(\sigma(-A), \mu)} \simeq \left\| \int_0^\infty T(s)bu(s) ds \right\|_X,$$

where $a \simeq b$ means there exist $C_1, C_2 > 0$ such that $a \leq C_1 b$ and $b \leq C_2 a$.

To begin the proof, we expand the expression on the right-hand side and get

$$\begin{aligned} \left\| \int_0^\infty T(s)bu(s) ds \right\|_X^2 &= \left\| \int_{\sigma(-A)} \int_0^\infty e^{-sz} u(s) ds dE_{-1}(z)b \right\|_X^2 \\ &= \left\| (rI - A_{-1}) \int_{\sigma(-A)} \int_0^\infty e^{sz} u(s) ds dE_{-1}(z)b \right\|_{X_{-1}}^2 \\ &= \left\| \int_{\sigma(-A)} (r - z) \int_0^\infty e^{sz} u(s) ds dE_{-1}(z)b \right\|_{X_{-1}}^2 \\ &= \left\| \int_{\sigma(-A)} (r - z) \cdot (\mathcal{L}u)(z) dE_{-1}(z)b \right\|_{X_{-1}}^2 \end{aligned}$$

with $r \in \rho(A)$ being the element of $\rho(A)$ that was implicitly used to define the space X_{-1} .

Given a measurable function $f: \sigma(A) \rightarrow \mathbb{C}$, one has the following norm representation:

$$\left\| \left(\int_{\sigma(-A)} f dE_{-1} \right) b \right\|_{X_{-1}}^2 = \int_{\sigma(-A)} |f|^2 d(E_{-1})_{b,b},$$

see e.g. [6, Theorem 4.7]. Specifying the function $z \mapsto (r - z) \cdot (\mathcal{L}u)(z)$ as the function f gives

$$\begin{aligned}
\left\| \int_{\sigma(-A)} (r - z) \cdot (\mathcal{L}u)(z) dE_{-1}(z)b \right\|_{X_{-1}}^2 &= \int_{\sigma(-A)} |(r - z) \cdot (\mathcal{L}u)(z)|^2 d\langle E_{-1}(z)b, b \rangle_{X_{-1}} \\
&= \int_{\sigma(-A)} |(\mathcal{L}u)(z)|^2 |r - z|^2 d(E_{-1})_{b,b}(z) \\
&\simeq \int_{\sigma(-A)} |(\mathcal{L}u)(z)|^2 |r - z|^2 \frac{1 + |z|^2}{|r - z|^2} d(E_{-1})_{b,b}(z) \\
&= \|\mathcal{L}u\|_{L^2(\sigma(-A), \mu)}^2,
\end{aligned}$$

where we have used that $r \in \rho(A)$ implies that

$$\frac{1 + |z|^2}{|r - z|^2} \simeq C$$

on $\sigma(A)$ for some $C > 0$. Then we also have

$$\left\| \int_0^\infty T(s)bu(s) ds \right\|_X^2 \simeq \|\mathcal{L}u\|_{L^2(\sigma(-A), \mu)}^2.$$

Hence p -admissibility of b is equivalent to boundedness of the Laplace transform. $\square$

Remark 3.4. We note that in the literature the specific case of normal operators with discrete spectrum is treated in full detail, whereas the general case is often argued to follow similarly as for diagonal semigroups; cf. e.g. [21, 42, 52] with the exception of [48, Theorem 19], where an approximation argument is used.

More precisely, in these references the measure μ is chosen to be $\langle E(\cdot)b, b \rangle$ which is in fact only well-defined on bounded Borel sets for $b \in X_{-1}$ (see also the proof of Theorem 19 in [48]). In Proposition 3.3 we circumvent this technical subtlety by employing a slightly different definition of μ , which however coincides with the abovementioned choice for diagonal semigroups.

3.3 Conditions for multiplication semigroups

Comparable results to the ones in the previous subsection for normal semigroups on Hilbert spaces hold for semigroups generated by multiplication operators on L^q spaces.

Let (Ω, μ) be a σ -finite measure space. Given a measurable function $a: \Omega \rightarrow \mathbb{C}$, we let the multiplier M_a on $L^q(\Omega)$, $1 < q < \infty$, be defined by

$$g \mapsto M_a g = ag, \quad g \in \mathcal{D}(M_f) = \{g \in L^q(\Omega) : ag \in L^q(\Omega)\}.$$

For multiplication semigroups, the extrapolation and interpolation spaces have convenient forms, see e.g. [9, Example II.5.7]. Suppose that a is complex-valued with $\text{ess sup}_\Omega \text{Re } a < 0$. The semigroup generated by the multiplication operator M_a on X is given by $(T(t))_{t \geq 0}$, where $T(t)f = e^{ta}f$. Moreover, for $n \in \mathbb{Z}$, the abstract Sobolev spaces X_n are given by the weighted Lebesgue spaces $X_n = L^q(\Omega, |a|^{nq}\mu)$.

Lemma 3.5. *Let $1 < q < \infty$ and assume that the multiplier M_a generates a strongly continuous semigroup T on $L^q(\Omega, \mu)$ with $\operatorname{Re} a < 0$ μ -a.e. on Ω . Then we have that*

$$M_{|\operatorname{Re} a|^{-1/q}} b \in L^q(\Omega, \mu) \text{ implies that } b \text{ is } L^{q'}\text{-admissible for } T.$$

Proof. By the functional calculus for multiplication operators, the semigroup T is given by $(T(t)f)(s) = e^{sa(t)}f(s)$. Therefore we need to show that there exists a constant $M > 0$ such that

$$\left\| \int_0^\infty e^{sa} b u(s) \, ds \right\|_{L^q(\Omega, \mu)} \leq M \|u\|_{L^{q'}([0, \infty))}$$

holds for all $u \in L^{q'}([0, \infty))$.

Since

$$\left\| \int_0^\infty e^{sa} b u(s) \, ds \right\|_{L^q(\Omega, \mu)}^q = \int_\Omega \left| \int_0^\infty e^{sa(t)} b(t) u(s) \, ds \right|^q d\mu(t),$$

we can use Hölder's inequality to get

$$\begin{aligned} & \int_\Omega \left| \int_0^\infty e^{sa(t)} b(t) u(s) \, ds \right|^q d\mu(t) \\ & \leq \int_\Omega \left| \left(\int_0^\infty |e^{sa(t)} b(t)|^q \, ds \right)^{1/q} \left(\int_0^\infty |u(s)|^{q'} \, ds \right)^{1/q'} \right|^q d\mu(t) \\ & = \|u\|_{L^{q'}([0, \infty))}^q \int_\Omega |b(t)|^q \int_0^\infty |e^{sa(t)}|^q \, ds \, d\mu(t) \\ & = \|u\|_{L^{q'}([0, \infty))}^q \int_\Omega |b(t)|^q \int_0^\infty e^{qs \operatorname{Re}(a(t))} \, ds \, d\mu(t) \\ & = C(q) \|u\|_{L^{q'}([0, \infty))}^q \int_\Omega |b|^q \frac{1}{|\operatorname{Re} a|} \, d\mu. \end{aligned}$$

This implies that b is q' -admissible for T if $b \in L^q(\Omega, |\operatorname{Re} a|^{-1} \mu)$. $\square$

Lemma 3.6. *Fix $1 < q < \infty$. Given a measurable complex-valued function a on some σ -finite measure space (Ω, μ) such that $\operatorname{Re} a$ is essentially bounded from above, let M_a be the corresponding multiplier on $L^q(\Omega, \mu)$. Then $b \in \mathcal{L}(\mathbb{C}, X_{-1}) \simeq X_{-1} = L^q(\Omega, |a|^{-1} \mu)$ is $L^{q'}$ -admissible for T if and only if the Laplace transform induces a bounded map*

$$\mathcal{L}: L^{q'}([0, \infty)) \rightarrow L^q(\Omega, \nu), \quad u \mapsto \mathcal{L}u = \int_0^\infty e^{-t(\cdot)} u(t) \, dt,$$

where the measure ν is defined by $\nu := (-a)_* |b|^q \mu$.

Proof. We have

$$\begin{aligned} \left\| \int_0^\infty T(s) b u(s) \, ds \right\|_{L^q(\Omega, \mu)}^q &= \int_\Omega |b(t)|^q \left| \int_0^\infty e^{sa(t)} u(s) \, ds \right|^q d\mu(t) \\ &= \int_\Omega |b(t)|^q |(\mathcal{L}u)(-a(t))|^q d\mu(t) \\ &= \int_\Omega |(\mathcal{L}u) \circ (-a)|^q d(|b|^q \mu) \\ &= \|(\mathcal{L}u)\|_{L^q(\Omega, (-a)_* |b|^q \mu)}^q, \end{aligned}$$

thus b is $L^{q'}$ -admissible if and only if

$$\|(\mathcal{L}u)\|_{L^q(\Omega, \nu)} \leq \|u\|_{L^{q'}([0, \infty))},$$

which is what we wanted to show. $\square$

Due to invariance of admissibility under similarity transformations, this result extends to generators that are similar to a multiplier or unitarily equivalent to a multiplier, and thus applies in particular to normal operators on a Hilbert space by a version of the spectral theorem for unbounded operators. Furthermore, the diagonal case follows as a special case:

Example 3.7 (Diagonal semigroups). Let A generate a diagonal semigroup on the state space $X = \ell^q(\mathbb{N}) = L^q(\mathbb{N}, \xi)$, where $\xi = \sum_k \delta_k$ is the counting measure on $\mathbb{N}$. Componentwise, this means that $(Ax)_k = \lambda_k x_k$ holds for all $k \in \mathbb{N}$ and all $x \in X$. Interpreting A as a multiplication operator on X with symbol $a: \mathbb{N} \rightarrow \mathbb{C}$, $k \mapsto \lambda_k$, we also identify the control operator $b \in X_{-1}$ with a function $b: \mathbb{N} \rightarrow \mathbb{C}$, $k \mapsto b_k$. By Lemma 3.6, the input element b is admissible for T if and only if the Laplace transform is a bounded map $\mathcal{L}: L^p([0, \infty)) \rightarrow L^q(\Omega, \nu)$ with measure ν given by

$$\nu = (-a)_* |b|^q \xi = |b|^q \sum_k \delta_{-\lambda_k} = \sum_k |b_k|^q \delta_{-\lambda_k},$$

which is the same measure as μ in Theorems 3.2 and 3.5 in [31].

4 Examples with generators having non-discrete spectra

By considering generators with non-discrete spectrum, we can find examples to illustrate the usage of the Laplace–Carleson type condition in the non-diagonal setting. Candidates are the Dirichlet or Neumann Laplacians on a suitable domain, since their spectra need not be discrete. In passing, we note that the Neumann Laplacian on Ω has purely discrete spectrum if and only if the Rellich–Kondrachov compactness theorem holds for the domain Ω . Hence a possible example would be the Neumann Laplacian on a sufficiently rough domain that fails the Rellich–Kondrachov embedding theorem, see e.g. [23]. Another operator with non-discrete spectrum, which we will consider in the sequel, is the Laplacian on the whole space $\mathbb{R}^n$.

4.1 The Laplacian on the full space

Example 4.1. Let $n \leq 3$ and consider the state space $X = L^2(\mathbb{R}^n)$ and the Laplacian $A = \Delta$ on X with domain $\mathcal{D}(A) = H^2(\mathbb{R}^n)$, which generates a C_0 -semigroup on X .

We choose the Dirac delta distribution δ_0 as the control operator b and claim that $b \in X_{-1}$ if and only if $n \leq 3$.

Proof of the claim. By self-adjointness of A the domains $\mathcal{D}(A)$ and $\mathcal{D}(A^*)$ are identical. Therefore the extrapolation space X_{-1} is given by the dual of $\mathcal{D}(A) = H^2(\mathbb{R}^n)$ with respect to the pivot space $X = L^2(\mathbb{R}^n)$. As the spatial domain is the full space $\mathbb{R}^n$, we have $H^2(\mathbb{R}^n) = H_0^2(\mathbb{R}^n)$. Hence—by definition of Sobolev spaces with negative integer indices—it holds that $X_{-1} = H^{-2}(\mathbb{R}^n)$. In the Fourier domain the space $H^{-2}(\mathbb{R}^n)$ has the representation

$$H^{-2}(\mathbb{R}^n) = \{u \in \mathcal{S}'(\mathbb{R}^n) : (1 + |\cdot|^2)^{-1} \hat{u} \in L^2(\mathbb{R}^n)\},$$

where $\mathcal{S}'(\mathbb{R}^n)$ denotes the space of tempered distributions on $\mathbb{R}^n$. As the Fourier transform of δ_0 computes to $\hat{\delta}_0 \equiv 1$ in the sense of tempered distributions, we have $\delta_0 \in X_{-1}$ if and only if

$$\int_{\mathbb{R}^n} \frac{1}{(1 + |\xi|^2)^2} d\xi < \infty.$$

By transforming to spherical coordinates in $\mathbb{R}^n$, this is equivalent to finiteness of

$$\int_0^\infty \frac{r^{n-1}}{(1 + r^2)^2} dr,$$

which is the case if and only if $n \leq 3$. $\square$

The spectral measure $E_{g,g}$ corresponding to the operator A and some $g \in L^2(\mathbb{R}^n)$ is computed in e.g. [44, Equation (7.34)] using the representation of the Laplacian by its Fourier symbol $\xi \mapsto -|\xi|^2$ and is given by the Lebesgue measure λ^n weighted with the density

$$\rho_{g,g}(z) = \chi_{(-\infty, 0]}(z) |z|^{\frac{n}{2}-1} \int_{\partial B_1^n(0)} |\hat{g}(|z|^{1/2} w)|^2 d\mathcal{H}^{n-1}(w);$$

in particular the spectrum of A is purely absolutely continuous and $\sigma(A) = (-\infty, 0]$.

In the above formula $\partial B_1^n(0) := \{y \in \mathbb{R}^n : |y| = 1\}$ denotes the unit sphere in $\mathbb{R}^n$, $\hat{g}$ denotes the Fourier transform of $g \in L^2(\mathbb{R}^n)$ and $\mathcal{H}^{n-1}$ is the $(n-1)$ -dimensional Hausdorff measure on $\mathbb{R}^n$. Note that this representation of $\rho_{g,g}$ is also valid in dimension $n = 1$ using the 0-dimensional Hausdorff measure.

For our example of the control operator $b = \delta_0 \in X_{-1}$, we need to consider the extension of $E_{g,g}$ for $g \in H^{-2}(\mathbb{R}^n) = X_{-1}$. For $g \in X_{-1}$ the density $(\rho_{-1})_{g,g}$ is then given by

$$(\rho_{-1})_{g,g}(z) = (1 + |z|^2) \chi_{(-\infty, 0]}(z) |z|^{\frac{n}{2}-1} \int_{\partial B_1^n(0)} \left| (\mathcal{F}(R(1, A)g))(|z|^{1/2} w) \right|^2 d\mathcal{H}^{n-1}(w).$$

We use $\hat{\delta}_0 \equiv 1$ to get

$$\begin{aligned} (\rho_{-1})_{\delta_0, \delta_0}(z) &= (1 + |z|^2) \chi_{(-\infty, 0]}(z) |z|^{\frac{n}{2}-1} \int_{\partial B_1^n(0)} \left| \frac{1}{1 + ||z|^{1/2} w|^2} \right|^2 d\mathcal{H}^{n-1}(w) \\ &= (1 + |z|^2) \chi_{(-\infty, 0]}(z) |z|^{\frac{n}{2}-1} \int_{\partial B_1^n(0)} (1 + |z|)^{-2} d\mathcal{H}^{n-1}(w) \\ &= \frac{1 + |z|^2}{(1 + |z|)^2} \chi_{(-\infty, 0]}(z) |z|^{\frac{n}{2}-1} \cdot C(n) \end{aligned}$$

Since $\frac{1}{2} \leq \frac{1 + |z|^2}{(1 + |z|)^2} \leq 1$ holds for $z \in \mathbb{C}$, we can reduce to the density

$$(\rho_{-1})_{\delta_0, \delta_0}(z) = \chi_{(-\infty, 0]}(z) |z|^{\frac{n}{2}-1} \tag{10}$$

in applications of Proposition 3.3.

To make statements concerning finite-time p -admissibility of b , we shift the generator of the semigroup such that the resulting semigroup is exponentially stable and apply the test from Proposition 3.3 for infinite-time admissibility.

Concretely, since $0 \in \sigma(A)$, to apply Proposition 3.3 we consider the shifted semigroup generated by the operator $A_1 = A - I$, such that $\sigma(A_1) = \sigma(A) - 1 = (-\infty, -1]$. Then by Proposition 3.3, and using the expression for the density $(\rho_{-1})_{\delta_0, \delta_0}$ from (10), the proof of finite-time L^p -admissibility of b reduces to finding a Laplace transform bound of the form

$$\int_{-\infty}^{-1} \left| \int_0^{\infty} e^{ts} u(t) dt \right|^2 (-s)^{\frac{n}{2}-1} ds \leq C \|u\|_{L^p([0, \infty))}^2 \quad (11)$$

for some $C > 0$. Using the Hölder inequality, we have

$$\begin{aligned} \int_{-\infty}^{-1} \left| \int_0^{\infty} e^{ts} u(t) dt \right|^2 (-s)^{\frac{n}{2}-1} ds &= \int_1^{\infty} \left| \int_0^{\infty} e^{-ty} u(t) dt \right|^2 y^{\frac{n}{2}-1} dy \\ &\leq \int_1^{\infty} \|e^{-(\cdot)y}\|_{L^{p'}([0, \infty))}^2 \|u\|_{L^p([0, \infty))}^2 y^{\frac{n}{2}-1} dy. \end{aligned} \quad (12)$$

As $\|e^{-(\cdot)y}\|_{L^{p'}([0, \infty))}^2 = (yp')^{-2/p'}$, the integral above stays bounded if

$$-\frac{2}{p'} + \frac{n}{2} - 1 < -1 \Leftrightarrow p > \frac{4}{4-n}. \quad (13)$$

Since δ_0 only defines an element of X_{-1} for $n \leq 3$, one arrives at the condition $p > 4/3$ in the one-dimensional setting, $p > 2$ in two dimensions and $p > 4$ for $n = 3$.

4.2 The Laplacian on a half line

Example 4.2. We consider the Dirichlet controlled heat equation on the half axis $[0, \infty)$. Set $X = L^2([0, \infty))$ and consider the Laplacian $A = \frac{d^2}{dx^2}$ on the domain $\mathcal{D}(A) = H_0^1([0, \infty)) \cap H^2([0, \infty))$. We select the scalar input space $U = \mathbb{C}$ and choose the control operator $b = \delta'_0 \in X_{-1} \simeq \mathcal{L}(\mathbb{C}, X_{-1})$. Our aim is to investigate p -admissibility of b for this example system.

We want to compute the resolvent of A to aid us in finding the spectral measure of A . To that end, for $g \in L^1([0, \infty)) \cap L^2([0, \infty))$, define the odd extension $\tilde{g} \in L^1(\mathbb{R}) \cap L^2(\mathbb{R})$ by

$$\tilde{g}(x) = \begin{cases} g(x) & x \geq 0 \\ -g(-x) & x < 0. \end{cases}$$

Denoting the Fourier variable on the full real axis by η and employing the representation of the resolvent of the Laplacian by its symbol in the Fourier domain, we have

$$R(z, A)g = \mathcal{F}^{-1}((z + \eta^2)^{-1}(\mathcal{F}\tilde{g})(\eta))\Big|_{[0, \infty)},$$

which allows for an explicit integral representation of the resolvent. In fact, we have

$$\begin{aligned}
R(z, A)g(x) &= \frac{1}{\sqrt{2\pi}} \mathcal{F}^{-1} \left(\frac{1}{z + \eta^2} \int_{-\infty}^{\infty} e^{-iy\eta} \tilde{g}(y) dy \right) (x) \\
&= \frac{1}{\sqrt{2\pi}} \mathcal{F}^{-1} \left(\frac{1}{z + \eta^2} \int_0^{\infty} (e^{-iy\eta} - e^{iy\eta}) g(y) dy \right) (x) \\
&= \frac{2}{i\sqrt{2\pi}} \mathcal{F}^{-1} \left(\frac{1}{z + \eta^2} \int_0^{\infty} \sin(y\eta) g(y) dy \right) (x) \\
&= \frac{1}{\pi i} \int_{-\infty}^{\infty} \frac{e^{ix\eta}}{z + \eta^2} \int_0^{\infty} \sin(y\eta) g(y) dy d\eta \\
&= \frac{1}{\pi i} \int_0^{\infty} \frac{e^{ix\eta} - e^{-ix\eta}}{z + \eta^2} \int_0^{\infty} \sin(y\eta) g(y) dy d\eta \\
&= \frac{2}{\pi} \int_0^{\infty} \frac{\sin(x\eta)}{z + \eta^2} \int_0^{\infty} \sin(y\eta) g(y) dy d\eta.
\end{aligned}$$

For brevity, in the sequel we denote the one-sided sine transform by $\mathcal{S}$, i.e.

$$(\mathcal{S}f)(x) := \int_0^{\infty} f(\xi) \sin(x\xi) d\xi, \quad f \in L^1([0, \infty)) \cap L^2([0, \infty)).$$

Still considering $g \in L^1([0, \infty)) \cap L^2([0, \infty))$ we compute

$$\begin{aligned}
\langle g, R(z, A)g \rangle_{L^2([0, \infty))} &= \int_0^{\infty} g(t) \overline{(R(z, A)g)(t)} dt \\
&= \frac{2}{\pi} \int_0^{\infty} b(t) \int_0^{\infty} \frac{\sin(\tau t)}{\bar{z} + \tau^2} \int_0^{\infty} \sin(\tau s) \overline{g(s)} ds d\tau dt \\
&= \frac{2}{\pi} \int_0^{\infty} g(t) \int_0^{\infty} \frac{\sin(\tau t)}{\bar{z} + \tau^2} (\mathcal{S}\bar{g})(\tau) d\tau dt \\
&= \frac{2}{\pi} \int_0^{\infty} \frac{(\mathcal{S}\bar{g})(\tau)}{\bar{z} + \tau^2} \int_0^{\infty} \sin(\tau t) g(t) dt d\tau \\
&= \frac{2}{\pi} \int_0^{\infty} \frac{(\mathcal{S}\bar{g})(\tau)}{\bar{z} + \tau^2} (\mathcal{S}g)(\tau) d\tau \\
&= \frac{2}{\pi} \int_0^{\infty} \frac{|(\mathcal{S}g)(\tau)|^2}{\bar{z} + \tau^2} d\tau.
\end{aligned}$$

We set $\rho = \tau^2$ and get

$$\begin{aligned}
\langle g, R(z, A)g \rangle_{L^2([0, \infty))} &= \frac{1}{\pi} \int_0^{\infty} \frac{|(\mathcal{S}g)(\sqrt{\rho})|^2}{\bar{z} + \rho} \frac{d\rho}{\sqrt{\rho}} \\
&= \frac{1}{\pi} \int_{\mathbb{C}} \chi_{[0, \infty)}(\rho) \frac{|(\mathcal{S}g)(\sqrt{|\rho|})|^2}{\bar{z} + \rho} \frac{d\rho}{\sqrt{|\rho|}} \\
&= \frac{1}{\pi} \int_{\mathbb{C}} \chi_{(-\infty, 0]}(\rho) \frac{|(\mathcal{S}g)(\sqrt{|\rho|})|^2}{\bar{z} - \rho} \frac{d\rho}{\sqrt{|\rho|}} \\
&= \int_{\mathbb{C}} \frac{1}{\bar{z} - \rho} dE_{g, g}(\rho),
\end{aligned}$$

where $E_{g,g}$ is Lebesgue measure weighted with the function $\chi_{(-\infty,0]}(\cdot)|(\mathcal{S}g)(\sqrt{|\cdot|})|^2/(\pi\sqrt{|\cdot|})$. We note that—like the Fourier transform—the sine and cosine transforms extend to operators defined on L^2 functions by density; cf. e.g. [10].

Returning to the control operator $b = \delta'_0$, we note that

$$(\mathcal{S}\delta'_0)(t) = \langle \delta'_0, \sin((\cdot)t) \rangle = -\frac{d}{d\xi} \sin(t\xi)|_{\xi=0} = -t.$$

In the same manner as for the Laplacian on the full Euclidean space as in Section 4.1 above, we can determine the measure μ from Proposition 3.3. This measure μ is formally given by the expression “ $\langle E(\cdot)\delta'_0, \delta'_0 \rangle_X$ ”—which is not surprising regarding the extension procedure used by Weiss in [48] mentioned in Remark 3.4 above—giving a measure that is equivalent (in the sense of mutual absolute continuity) to the measure defined by

$$E_{\delta'_0, \delta'_0}(z) = \chi_{(-\infty,0]}(z) \frac{|(\mathcal{S}\delta'_0)(\sqrt{|z|})|^2}{\sqrt{|z|}} \lambda^1(z) = \chi_{(-\infty,0]}(z) |z|^{1/2} \lambda^1(z);$$

note also that the associated Radon–Nikodým densities are bounded away from 0 and ∞ .

In terms of p -admissibility of b , the computation in (11), (12) and (13) then yields finite-time admissibility of b for all $p > 4$.

Example 4.3. Proceeding in a similar manner as in the prior case of Dirichlet control, we can consider the Neumann controlled heat equation on the non-negative real axis. On the same state space $X = L^2([0, \infty))$ we now consider the the Laplace operator $A = \frac{d^2}{dt^2}$ on the domain $\mathcal{D}(A) = \{f \in H^2([0, \infty)) : f'(0) = 0\}$ and the control operator $b = \delta_0$ on the input space $U = \mathbb{C}$.

Analogously as in the Dirichlet setting, but using the even extension of functions defined on the half line, the resolvent of A computes to

$$(R(z, A)g)(x) = \frac{2}{\pi} \int_0^\infty \frac{\cos(x\eta)}{z + \eta^2} \int_0^\infty \cos(y\eta) g(y) dy d\eta$$

for $g \in L^1([0, \infty)) \cap L^2([0, \infty))$. For the spectral measure, an analogous calculation as in the Dirichlet case yields

$$E_{g,g}(z) = \chi_{(-\infty,0]}(z) \frac{|(\mathcal{C}g)(\sqrt{|z|})|^2}{\pi\sqrt{|z|}} \lambda^1(z), \quad g \in L^1([0, \infty)) \cap L^2([0, \infty)).$$

Here $\mathcal{C}$ denotes the one-sided cosine transform defined by $(\mathcal{C}f)(x) := \int_0^\infty f(\xi) \cos(x\xi) d\xi$ for $f \in L^1([0, \infty)) \cap L^2([0, \infty))$, and we also denote by $\mathcal{C}$ the extension of the transform to $L^2([0, \infty))$ and to tempered distributions.

As $\mathcal{C}\delta_0 \equiv 1$, the same argument as in the Dirichlet case above lets us determine the measure μ from Proposition 3.3 to be equivalent to

$$E_{\delta_0, \delta_0}(z) = \chi_{(-\infty,0]}(z) |z|^{-1/2} \lambda^1(z).$$

Since this is the same density as in the full space case with $n = 1$ in Example 4.1, the computation in (11), (12) and (13) there yields finite-time L^p -admissibility of $b = \delta_0$ for A for all $p > \frac{4}{3}$.

5 Applications to infinite-dimensional input spaces

Laplace–Carleson embeddings have been proven to be useful in the context of admissibility with finite-dimensional input spaces. In this section we propose a partial extension to infinite-dimensional input spaces. We note that the question of a full characterization of p -admissibility in terms of these criteria—without restrictive additional assumptions—seems to be open at this moment in time.

However, imposing additional assumptions on the system, we can extend admissibility results for finite-dimensional input spaces to infinite-dimensional ones. One of such conditions is that A has the p -Weiss property.

Definition 5.1 (p -Weiss property). Let $p \in [1, \infty]$ and A generate a C_0 -semigroup on the Banach space X . Then we say that the p -Weiss property for control operators holds for A if for any Banach space U and any $B \in \mathcal{L}(U, X_{-1})$, we have that

$$B \text{ is infinite-time } p\text{-admissible} \Leftrightarrow \sup_{\operatorname{Re} \lambda > 0} (\operatorname{Re} \lambda)^{\frac{1}{p}} \|R(\lambda, A)B\| < \infty.$$

The p -Weiss property for observation operators holds if for all $C \in \mathcal{L}(X_1, Y)$, Y Banach space,

$$C \text{ is infinite-time } p\text{-admissible} \Leftrightarrow \sup_{\operatorname{Re} \lambda > 0} (\operatorname{Re} \lambda)^{\frac{1}{p'}} \|CR(\lambda, A)\| < \infty.$$

For general semigroup generators A , the p -Weiss property does not hold, as indicated in the introduction; see e.g. [30, 54, 55] for $p = 2$. In special cases—such as the case of self-adjoint generators A —it does hold; see Proposition 5.4 below.

Theorem 5.2 (Uniform weak admissibility & Weiss property). *Let A be the generator of a C_0 -semigroup on the Banach space X . Fix an exponent $q \in (1, \infty)$ and assume that the q -Weiss property holds for A . Then the following are equivalent for $B \in \mathcal{L}(U, X_{-1})$:*

- (a) B is L^q -admissible for A .
- (b) For all $u \in U$ with $\|u\|_U = 1$ the input element $Bu \in X_{-1}$ is L^q -admissible and the admissibility constants, i.e. the numbers

$$C(u) := \inf \left\{ C > 0 : \forall v \in L^q([0, t]) \left\| \int_0^t T(t-s)Buv(s) \, ds \right\| \leq C \|v\|_{L^q([0, t])} \right\},$$

are uniformly bounded in u .

Proof. We note that for any $u \in U$ with norm 1 and any $v \in L^q([0, \infty))$ we have $uv \in L^q([0, \infty), U)$. To show “(a) $\Rightarrow$ (b)”, we assume that $B: U \rightarrow X_{-1}$ is q -admissible. Then by admissibility of B we infer

$$\left\| \int_0^t T(t-s)B(uv)(s) \, ds \right\|_X \leq C \|uv\|_{L^q([0, t], U)} = C \|v\|_{L^q([0, t])},$$

which implies that $Bu \in X_{-1}$ is q -admissible with admissibility constant that does not depend on u . Hence the admissibility constants are also uniformly bounded in u .

For the other implication, we assume that for all $u \in U$ with unit norm the input element Bu is q -admissible, i.e.

$$\left\| \int_0^t T(t-s)B(uv)(s) \, ds \right\|_X \leq C(u)\|v\|_{L^q([0,t])},$$

and that the associated constants are uniformly bounded in u , which is to say we have

$$\sup_{u \in U, \|u\|_U=1} C(u) < \infty.$$

Hence the mapping $\Phi_{t,Bu}: L^q([0, \infty)) \rightarrow X$, $v \mapsto \int_0^t T(s)Buv(s) \, ds$ allows for the bound $M_t := \sup_{u \in U, \|u\|=1} \|\Phi_{t,Bu}\| < \infty$. Using infinite-time q -admissibility of Bu , we also get a bound of the form $M := \limsup_{t \rightarrow \infty} M_t < \infty$. Furthermore, setting $\tilde{v}(s) := (q \operatorname{Re} \lambda)^{1/q} e^{-\lambda s}$ we get $\|\tilde{v}\|_{L^q([0, \infty))} = 1$. Thus $\|\Phi_{t,Bu}\tilde{v}\|_X \leq \|\Phi_{t,Bu}\| \|\tilde{v}\|_{L^q([0, \infty))} = \|\Phi_{t,Bu}\| \leq M$ and

$$\left\| \int_0^\infty T(s)Bu\tilde{v}(s) \, ds \right\|_X = \lim_{t \rightarrow \infty} \|\Phi_{t,Bu}\tilde{v}\|_X \leq M.$$

As the resolvent is the Laplace transform of the semigroup, it follows that

$$\begin{aligned} (\operatorname{Re} \lambda)^{1/q} \|R(\lambda, A)Bu\|_X &= (\operatorname{Re} \lambda)^{1/q} \left\| \int_0^\infty e^{-\lambda s} T(s)Bu \, ds \right\|_X \\ &= \|\Phi_{t,Bu}\tilde{v}\|_X \leq M. \end{aligned}$$

Due to our assumption, $\|\Phi_{t,Bu}\|$ is uniformly bounded, which implies

$$(\operatorname{Re} \lambda)^{1/q} \|R(\lambda, A)B\| = \sup_{u \in U, \|u\|_U=1} (\operatorname{Re} \lambda)^{1/q} \|R(\lambda, A)Bu\|_X \leq M;$$

taking the supremum among $\lambda \in \mathbb{C}$ with $\operatorname{Re} \lambda > 0$ then gives

$$\sup_{\operatorname{Re} \lambda > 0} (\operatorname{Re} \lambda)^{1/q} \|R(\lambda, A)B\|_X \leq M < \infty.$$

Hence B is infinite-time q -admissible by the assumed validity of the q -Weiss property for A . $\square$

Remark 5.3. Theorem 5.2 allows extending the Laplace–Carleson test for admissibility to infinite-dimensional input spaces. The practicality of this condition is yet to be seen, but this approach allows for assessing admissibility sharply in p . For example, a possible application to the heat equation could involve estimates for the norm of the trace of eigenfunctions on the boundary by means of eigenfunction expansions, as the eigenfunctions of the Laplacian with Dirichlet or Neumann boundary conditions are explicitly known for well-behaved geometries. Moreover, there is a wide array of work on norms of the appropriate traces of eigenfunctions on the boundary, see e.g. [2, 13, 22, 43, 53].

In this context we note that the p -Weiss property is strongly related to admissibility of $(-A)^{1/p}$ for A , which was shown by Haak in [14]. If A generates a semigroup of self-adjoint operators and $\sigma(A) \subset (-\infty, 0]$, this even leads to validity of the p -Weiss property for $p \leq 2$.

Proposition 5.4. *Let A be a self-adjoint operator on a Hilbert space X with spectrum contained in $(-\infty, a]$ for some $a \in \mathbb{R}$. Then A has the p -Weiss property (for control operators) if $p \leq 2$.*




Figure 2: The p -Weiss property for self-adjoint A , see [14].

Proof. The case $p = 2$ is contained in [35, Corollary 4.3] as the negative semidefinite self-adjoint operator A is the generator of a C_0 -semigroup of contractions on X .

Thus we fix $p < 2$. The claim is that $B \in \mathcal{L}(U, X_{-1})$ is infinite-time L^p -admissible if and only if the quantity $\sup_{\lambda > 0} \lambda^{1/p} \|R(\lambda, A)B\|_{\mathcal{L}(U, X)}$ is finite. To prove the claim we use results from Hosfeld, Jacob and Schwenninger [25] in conjunction with results from Haak [14].

First we note that due to the assumption of being bounded from above, the self-adjoint A generates a bounded analytic semigroup $(T(t))_{t \geq 0}$, see [9, Corollary II.4.7]. We observe that the spectral theorem for unbounded self-adjoint operators allows us to prove p' -admissibility of $(-A)^{1/p'}$ as an observation operator for $p' > 2$ more directly. Indeed, as $-A$ is self-adjoint, there exists a spectral measure E such that

$$-A = \int_{\sigma(-A)} z \, dE(z), \quad T(t) = \int_{\sigma(-A)} e^{-tz} \, dE(z), \quad t \geq 0,$$

holds. In particular, there is a representation of the norm, i.e.

$$\int_0^\infty \|(-A)^{1/p'} T(t)x\|_X^{p'} \, dt = \int_0^\infty \left(\int_{\sigma(-A)} |z^{1/p'} e^{-tz}|^2 \, dE_{x,x}(z) \right)^{p'/2} \, dt;$$

cf. [6, Theorem 4.7]. For σ -finite measure spaces (X, μ) and (Y, ν) , a measurable function f defined on $X \times Y$ and $1 \leq p < q < \infty$, Minkowski's integral inequality reads

$$\int_Y \left(\int_X |f(x, y)|^p \, d\mu(x) \right)^{q/p} \, d\nu(y) \leq \left(\int_X \left(\int_Y |f(x, y)|^q \, d\mu(y) \right)^{p/q} \, d\mu(x) \right)^{q/p},$$

see e.g. [3, Exercise 3.10.46] or [26, Proposition 1.2.22]. Applying Minkowski's inequality in (5) above yields

$$\begin{aligned} \int_0^\infty \left(\int_{\sigma(-A)} |z^{1/p'} e^{-tz}|^2 \, dE_{x,x}(z) \right)^{p'/2} \, dt &\leq \left(\int_{\sigma(-A)} \left(\int_0^\infty z e^{-tz p'} \, dt \right)^{2/p'} \, dE_{x,x}(z) \right)^{p'/2} \\ &= \left(\frac{1}{p'^{2/p'}} \int_{\sigma(-A)} dE_{x,x}(z) \right)^{p'/2} \\ &= \frac{1}{p'} \|x\|_X^{p'}. \end{aligned}$$

Therefore $(-A)^{1/p'}$ is p' -admissible as an observation operator, which is equivalent to requiring that the operator $-A$ satisfies $L^{p'}$ estimates in the sense of Haak, [14, page 29].

Because A was required to be self-adjoint, the same estimates hold for $(-A)^*$. Hence we can apply the generalization of [14, Satz 2.1.6] given in [14, page 29], which states that the p -Weiss property for control operators is valid for A . This ends the proof. $\square$

Remark 5.5. The last proposition leads us to the observation that there is a noticeable difference between the cases $p < 2$ and $p > 2$ in checking p -admissibility. In the control operator setting with a system on a Hilbert space X , the case $p < 2$ is simpler than $p > 2$. By the previous argument, once we transfer the reasoning from observation operators to control operators— A was self-adjoint—we would need to require $p'/2 \geq 1$ for the Minkowski inequality to hold, which means that we would need $p < 2$.

The complementary case does not allow for such arguments. Below we give an example of a self-adjoint and negative operator that does not have the 4-Weiss property. We note in passing that in the case $p = \infty$, the property of A^* being an L^1 -admissible observation operator is equivalent to A being bounded on X , see also [33].

Example 5.6. By means of the example of the one-dimensional heat equation from Section 2.2 above, we can show that a negative definite self-adjoint generator need not possess the p -Weiss property for $p > 2$.

As a continuation of our earlier computations, we prove that the resolvent condition is valid for $p = 4$. Since the control operator of system in question was not 4-admissible, this implies that the generator does not have the 4-Weiss property.

Recall that the Dirichlet Laplacian A on $L^2([0, 1])$ has the diagonal representation

$$Ae_n = -n^2\pi^2e_n, \quad n \in \mathbb{N},$$

where e_n are the unit vectors in $\ell^2(\mathbb{N})$. Moreover, the control operator b modeling the heat injection at the right endpoint of the interval was described by the sequence $b = \{b_n\}_{n \in \mathbb{N}}$, where $b_n = (-1)^n\sqrt{2}n\pi$.

The composition $R(\lambda, A)b$ of the resolvent and control operator is then given by the sequence

$$(R(\lambda, A)b)_n = \frac{(-1)^n\sqrt{2}n\pi}{\lambda + n^2\pi^2}, \quad n \in \mathbb{N}.$$

Validity of the resolvent condition in the 4-Weiss property is (by definition) equivalent to uniform boundedness of the quantity $(\operatorname{Re} \lambda)^{1/4} \|R(\lambda, A)b\|_{\mathcal{L}(U, X)}$ among complex numbers λ with positive real part.

In view of $U = \mathbb{C}$ and $X = \ell^2(\mathbb{N})$, we insert the expressions for the coefficients of $R(\lambda, A)b$; it then remains to show that

$$\sup_{\operatorname{Re} \lambda > 0} (\operatorname{Re} \lambda)^{1/4} \left(\sum_{n=1}^{\infty} \frac{2n^2\pi^2}{|\lambda + n^2\pi^2|^2} \right)^{1/2} < \infty. \quad (14)$$

Note that we can restrict to real λ . Moreover, it is convenient to consider $\mu = \sqrt{\lambda}$ in place of λ . Employing this substitution, after squaring the expression (14) becomes

$$\sup_{\mu > 0} \mu \sum_{n=1}^{\infty} \frac{2n^2\pi^2}{(\mu^2 + n^2\pi^2)^2} < \infty.$$

A closed form of the sum $\sum_{n=1}^{\infty} \frac{\pi^2 n^2}{(b + \pi^2 n^2)^2}$ exists in terms of hyperbolic functions. Indeed, using

$$\coth x := \frac{\cosh x}{\sinh x} = \frac{e^x + e^{-x}}{e^x - e^{-x}}, \quad \csc x := \frac{1}{\sin x}, \quad \operatorname{csch} x := \frac{1}{\sinh x} = \frac{2}{e^x - e^{-x}}$$

together with the series expansions

$$\coth(\pi x) = \frac{1}{\pi x} + \frac{2x}{\pi} \sum_{k=1}^{\infty} \frac{1}{k^2 + x^2}, \quad \csc^2(\pi x) = \frac{1}{\pi^2} \sum_{k=-\infty}^{\infty} \frac{1}{(x - k)^2},$$

see [12, page 36], and the identity $\operatorname{csch}^2(x) = -\csc^2(ix)$ for $x \in \mathbb{R}$, one can show that

$$\mu \sum_{n=1}^{\infty} \frac{2n^2\pi^2}{(\mu^2 + n^2\pi^2)^2} = \frac{1}{2}(\coth \mu - \mu \operatorname{csch}^2 \mu) = \frac{1}{2} + \frac{(2 - 4\mu)e^{2\mu} - 2}{2(e^{2\mu} - 1)^2}.$$

As $\frac{(2-4\mu)e^{2\mu}-2}{2(e^{2\mu}-1)^2} < 0$, we have

$$\sup_{\mu>0} \mu \sum_{n=1}^{\infty} \frac{2n^2\pi^2}{(\mu^2 + n^2\pi^2)^2} = \sup_{\mu>0} \left(\frac{1}{2} + \frac{(2 - 4\mu)e^{2\mu} - 2}{2(e^{2\mu} - 1)^2} \right) \leq \frac{1}{2};$$

validating the resolvent condition. Therefore the 4-Weiss property is violated for A .

The above example shows that Theorem 5.2 cannot be applied to show p -admissibility for systems arising from boundary control with infinite-dimensional input space if $p > 2$, even if A is self-adjoint. Consequently, determining admissibility with respect to sharp values of p remains difficult in this regime. Practically, the straight-forward tool to derive admissibility seems to be trace estimates in conjunction with the interpolation spaces as lined out in Section 2.1.

References

- [1] R. A. Adams and J. J. F. Fournier. *Sobolev Spaces*. 2nd edition. Pure and Applied Mathematics 140. Amsterdam, Boston: Academic Press, 2003. ISBN: 978-0-12-044143-3.
- [2] A. Bäcker, S. Fürstberger, R. Schubert, and F. Steiner. “Behaviour of Boundary Functions for Quantum Billiards”. In: *Journal of Physics A: Mathematical and General* 35.48 (Nov. 2002), p. 10293.
- [3] V. I. Bogachev. *Measure Theory*. Berlin, New York: Springer, 2007. ISBN: 978-3-540-34513-8.
- [4] H. Bounit, A. Driouich, and O. El-Mennaoui. “A Direct Approach to the Weiss Conjecture for Bounded Analytic Semigroups”. In: *Czechoslovak Mathematical Journal* 60.2 (June 2010), pp. 527–539.
- [5] C. I. Byrnes, D. S. Gilliam, V. I. Shubov, and G. Weiss. “Regular Linear Systems Governed by a Boundary Controlled Heat Equation”. In: *Journal of Dynamical and Control Systems* 8.3 (July 2002), pp. 341–370.
- [6] J. B. Conway. *A Course in Functional Analysis*. Vol. 96. Graduate Texts in Mathematics. New York, NY: Springer New York, 2007. ISBN: 978-1-4757-4383-8.
- [7] R. F. Curtain and A. J. Pritchard. *Infinite Dimensional Linear Systems Theory*. Ed. by A. V. Balakrishnan and M. Thoma. Vol. 8. Lecture Notes in Control and Information Sciences. Berlin, Heidelberg: Springer, 1978. ISBN: 978-3-540-08961-2.

- [8] K.-J. Engel. “On the Characterization of Admissible Control- and Observation Operators”. In: *Systems & Control Letters* 34.4 (July 1998), pp. 225–227.
- [9] K.-J. Engel and R. Nagel. *One-parameter semigroups for linear evolution equations*. Graduate texts in mathematics 194. New York, Heidelberg: Springer, 2000. ISBN: 978-0-387-22642-2.
- [10] R. R. Goldberg. “Certain Operators and Fourier Transforms on L^2 ”. In: *Proceedings of the American Mathematical Society* 10.3 (June 1959).
- [11] P. Grabowski and F. M. Callier. “Admissible Observation Operators. Semigroup Criteria of Admissibility”. In: *Integral Equations and Operator Theory* 25.2 (June 1996), pp. 182–198.
- [12] I. S. Gradshteyn, I. M. Ryzhik, and A. Jeffrey. *Table of integrals, series, and products*. Corrected and enlarged edition. New York: Academic Press, 1980. ISBN: 0-12-294760-6.
- [13] D. S. Grebenkov and B.-T. Nguyen. “Geometrical Structure of Laplacian Eigenfunctions”. In: *SIAM Review* 55.4 (Jan. 2013), pp. 601–667.
- [14] B. H. Haak. “Kontrolltheorie in Banachräumen und quadratische Abschätzungen”. Dissertation: University of Karlsruhe, 2005. 148 pp.
- [15] B. H. Haak. “On the Carleson Measure Criterion in Linear Systems Theory”. In: *Complex Analysis and Operator Theory* 4.2 (May 2010), pp. 281–299.
- [16] B. H. Haak. “The Weiss Conjecture and Weak Norms”. In: *Journal of Evolution Equations* 12.4 (Dec. 2012), pp. 855–861.
- [17] B. H. Haak and P. C. Kunstmann. “On Kato’s Method for Navier–Stokes Equations”. In: *Journal of Mathematical Fluid Mechanics* 11.4 (Dec. 2009), pp. 492–535.
- [18] B. H. Haak and P. C. Kunstmann. “Admissibility of Unbounded Operators and Well-posedness of Linear Systems in Banach Spaces”. In: *Integral Equations and Operator Theory* 55.4 (Aug. 2006), pp. 497–533.
- [19] B. H. Haak and P. C. Kunstmann. “Weighted Admissibility and Wellposedness of Linear Systems in Banach Spaces”. In: *SIAM Journal on Control and Optimization* 45.6 (Jan. 2007), pp. 2094–2118.
- [20] B. H. Haak and C. Le Merdy. “ α -Admissibility of Observation and Control Operators”. In: *Houston Journal of Mathematics* 31.4 (2005), pp. 1153–1167.
- [21] S. Hansen and G. Weiss. “The Operator Carleson Measure Criterion for Admissibility of Control Operators for Diagonal Semigroups on l^2 ”. In: *Systems & Control Letters* 16.3 (Mar. 1991), pp. 219–227.
- [22] A. Hassell and T. Tao. “Upper and Lower Bounds for Normal Derivatives of Dirichlet Eigenfunctions”. In: *Mathematical Research Letters* 9.3 (May 2002), pp. 289–305.
- [23] R. Hempel, L. A. Seco, and B. Simon. “The Essential Spectrum of Neumann Laplacians on Some Bounded Singular Domains”. In: *Journal of Functional Analysis* 102.2 (Dec. 1991), pp. 448–483.
- [24] L. F. Ho and D. L. Russell. “Admissible Input Elements for Systems in Hilbert Space and a Carleson Measure Criterion”. In: *SIAM Journal on Control and Optimization* 21.4 (July 1983), pp. 614–640.

- [25] R. Hoffeld, B. Jacob, and F. L. Schwenninger. “Characterization of Orlicz Admissibility”. In: *Semigroup Forum* 106.3 (June 2023), pp. 633–661.
- [26] T. Hytönen, J. Van Neerven, M. Veraar, and L. Weis. *Analysis in Banach Spaces*. Vol. I: Martingales and Littlewood-Paley Theory. Cham: Springer International Publishing, 2016. ISBN: 978-3-319-48519-5.
- [27] B. Jacob, R. Nabiullin, J. R. Partington, and F. L. Schwenninger. “Infinite-Dimensional Input-to-State Stability and Orlicz Spaces”. In: *SIAM Journal on Control and Optimization* 56.2 (Jan. 2018), pp. 868–889.
- [28] B. Jacob and J. R. Partington. “Admissibility of Control and Observation Operators for Semigroups: A Survey”. In: *Current Trends in Operator Theory and Its Applications*. Ed. by J. A. Ball, M. Klaus, J. W. Helton, and L. Rodman. Vol. 149. Operator Theory: Advances and Applications. Basel: Birkhäuser, 2004, pp. 199–221. ISBN: 978-3-0348-9608-5.
- [29] B. Jacob and J. R. Partington. “The Weiss Conjecture on Admissibility of Observation Operators for Contraction Semigroups”. In: *Integral Equations and Operator Theory* 40.2 (June 2001), pp. 231–243.
- [30] B. Jacob, J. R. Partington, and S. Pott. “Admissible and weakly admissible observation operators for the right shift semigroup”. In: *Proceedings of the Edinburgh Mathematical Society* 45.2 (June 2002), pp. 353–362.
- [31] B. Jacob, J. R. Partington, and S. Pott. “Applications of Laplace–Carleson Embeddings to Admissibility and Controllability”. In: *SIAM Journal on Control and Optimization* 52.2 (Jan. 2014), pp. 1299–1313.
- [32] B. Jacob, J. R. Partington, S. Pott, E. Rydhe, and F. L. Schwenninger. *Laplace-Carleson Embeddings and Infinity-Norm Admissibility*. Feb. 2, 2023. arXiv: 2109.11465.
- [33] B. Jacob, F. L. Schwenninger, and J. Wintermayr. “A refinement of Baillon’s theorem on maximal regularity”. In: *Studia Mathematica* 263 (2022), pp. 141–158.
- [34] I. Lasiecka and R. Triggiani. *Control theory for partial differential equations: continuous and approximation theories*. Vol. 1. Encyclopedia of mathematics and its applications 74-75. Cambridge, New York: Cambridge University Press, 2000. ISBN: 978-0-521-43408-9.
- [35] C. Le Merdy. “The Weiss Conjecture for Bounded Analytic Semigroups”. In: *Journal of the London Mathematical Society* 67.03 (June 2003), pp. 715–738.
- [36] F. Maragh, H. Bounit, A. Fadili, and H. Hammouri. “On the Admissible Control operators for Linear and Bilinear Systems and the Favard Spaces”. In: *Bulletin of the Belgian Mathematical Society - Simon Stevin* 21.4 (Oct. 2014).
- [37] A. Mironchenko and C. Prieur. “Input-to-State Stability of Infinite-Dimensional Systems: Recent Results and Open Questions”. In: *SIAM Review* 62.3 (Jan. 2020), pp. 529–614.
- [38] A. Pazy. *Semigroups of Linear Operators and Applications to Partial Differential Equations*. Vol. 44. Applied Mathematical Sciences. New York: Springer, 1983. ISBN: 0-387-90845-5.
- [39] D. Salamon. “Infinite Dimensional Linear Systems With Unbounded Control and Observation: A Functional Analytic Approach”. In: *Transactions of the American Mathematical Society* 300.2 (1987), pp. 383–431.

- [40] C. Schneider. *Beyond Sobolev and Besov: Regularity of Solutions of PDEs and Their Traces in Function Spaces*. Vol. 2291. Lecture Notes in Mathematics. Cham: Springer International Publishing, 2021. ISBN: 978-3-030-75139-5.
- [41] F.L. Schwenninger. “Input-to-state stability for parabolic boundary control: linear and semilinear systems”. In: *Control Theory of Infinite-Dimensional Systems*. Operator Theory: Advances and Applications. Cham: Springer, 2020, pp. 83–116. ISBN: 978-3-030-35898-3.
- [42] O.J. Staffans. *Well-posed linear systems*. Cambridge: Cambridge University Press, 2005. ISBN: 978-0-511-08208-5.
- [43] D. Tataru. “On the Regularity of Boundary Traces for the Wave Equation”. In: *Annali della Scuola Normale Superiore di Pisa - Classe di Scienze* 26.1 (1998), pp. 185–206.
- [44] G. Teschl. *Mathematical Methods in Quantum Mechanics: With Applications to Schrödinger Operators, Second Edition*. Vol. 157. Graduate Studies in Mathematics. American Mathematical Society, 2014. ISBN: 978-1-4704-1888-5.
- [45] H. Triebel. *Theory of Function Spaces II*. Basel: Springer, 1992. ISBN: 978-3-0346-0418-5.
- [46] M. Tucsnak and G. Weiss. *Observation and control for operator semigroups*. Birkhäuser advanced texts. Basel, Boston: Birkhäuser, 2009. ISBN: 978-3-764-38993-2.
- [47] M. Unteregge. “ p -Admissible Control Elements for Diagonal Semigroups on l^r -Spaces”. In: *Systems & Control Letters* 56.6 (June 2007), pp. 447–451.
- [48] G. Weiss. “A Powerful Generalization of the Carleson Measure Theorem?” In: *Open Problems in Mathematical Systems and Control Theory*. Ed. by V. Blondel, E.D. Sontag, M. Vidyasagar, and J.C. Willems. Communications and Control Engineering. London: Springer, 1999, pp. 267–272. ISBN: 978-1-4471-0807-8.
- [49] G. Weiss. “Admissibility of Input Elements for Diagonal Semigroups on l^2 ”. In: *Systems & Control Letters* 10.1 (Jan. 1988), pp. 79–82.
- [50] G. Weiss. “Admissibility of Unbounded Control Operators”. In: *SIAM Journal on Control and Optimization* 27.3 (May 1989), pp. 527–545.
- [51] G. Weiss. “Admissible Observation Operators for Linear Semigroups”. In: *Israel Journal of Mathematics* 65.1 (Feb. 1989), pp. 17–43.
- [52] G. Weiss. “Two Conjectures on the Admissibility of Control Operators”. In: *Estimation and Control of Distributed Parameter Systems: Proceedings of an International Conference on Control and Estimation of Distributed Parameter Systems, Vorau, July 8–14, 1990*. Ed. by W. Desch, F. Kappel, and K. Kunisch. International Series of Numerical Mathematics. Basel: Birkhäuser, 1991, pp. 367–378. ISBN: 978-3-0348-6418-3.
- [53] S. Zelditch. “Billiards and Boundary Traces of Eigenfunctions”. In: *Journées équations aux dérivées partielles* (2003), pp. 1–22.
- [54] H.J. Zwart and B. Jacob. *Disproof of an admissibility conjecture of Weiss*. Faculty of Mathematical Sciences, Memorandum 1539. Enschede, The Netherlands: University of Twente, 2000.
- [55] H.J. Zwart, B. Jacob, and O.J. Staffans. “Weak Admissibility Does Not Imply Admissibility for Analytic Semigroups”. In: *Systems & Control Letters*. Optimization and Control of Distributed Systems 48.3 (Mar. 2003), pp. 341–350.