## AN EFFECTIVE VERSION OF FEKETE'S THEOREM

NORM LEVENBERG AND MAYURESH LONDHE

ABSTRACT. A classical result of Fekete gives necessary conditions on a compact set in the complex plane so that it contains infinitely many sets of conjugate algebraic integers. We prove an effective version of Fekete's theorem in terms of a height function. As an application, we give a lower bound on the growth of the leading coefficient of certain polynomial sequences, generalizing a result by Schur. Lastly, we give an upper bound on minimal asymptotics of height over sequences of algebraic numbers.

### 1. INTRODUCTION

Let  $\Sigma$  be a compact set in the complex plane  $\mathbb{C}$ . Fekete studied which compact sets  $\Sigma$  contain only a finite number of sets of conjugate algebraic integers. This involves the notion of the *transfinite diameter of*  $\Sigma$ ; it is given by

$$d(\Sigma) = \lim_{n \to \infty} \left( \sup_{\xi_1, \dots, \xi_n \in \Sigma} \prod_{i < j} |\xi_i - \xi_j|^{\frac{2}{n(n-1)}} \right).$$

The transfinite diameter of a circle of radius r is equal to r, and for a line segment of length l it is equal to l/4. It turns out that the quantity  $d(\Sigma)$  coincides with the logarithmic capacity of  $\Sigma$ , denoted by  $c(\Sigma)$ ; see Subsection 2.1 for the definition. The product  $\prod_{i < j} |\xi_i - \xi_j|$  is related to the resultant of P and its derivative P', where  $P(z) = \prod_{j=1}^{n} (z - \xi_j)$ ; see (2.4). This motivates the use of potential-theoretic tools in studying sets of conjugate algebraic numbers. We now state a result by Fekete:

**Result 1.1** (Fekete, [Fek23]). Let  $\Sigma \subset \mathbb{C}$  be compact and symmetric about the real axis. If  $c(\Sigma) < 1$  then there exists a neighborhood U of  $\Sigma$  such that there are only a finite number of algebraic integers  $\alpha$  so that  $\alpha$  and all its conjugates lie in U.

The above result is not true if we replace algebraic integers by algebraic numbers in the statement, for example, if  $\Sigma = \{|z| = 1/2\}$  then all roots of  $2^n z^n - 1$  are in  $\Sigma$ . In [FS55], Fekete, together with Szegö showed that the converse of Result 1.1 is also true.

Our main results involve a notion of height  $h_{\Sigma}(\alpha)$  of an algebraic number  $\alpha$  relative to a compact set  $\Sigma$  as defined in [Rum99]. Let  $\overline{\mathbb{Q}}$  denote the set of algebraic numbers in  $\mathbb{C}$ . Given a nonzero  $\alpha \in \overline{\mathbb{Q}}$ , there is a polynomial  $P_{\alpha}$  with integer coefficients and leading coefficient  $b_{\alpha}$  and degree deg  $\alpha$  that is irreducible over the integers and satisfies  $P_{\alpha}(\alpha) = 0$  ( $P_{\alpha}$  is unique up to the sign of  $b_{\alpha}$ ). For an algebraic integer  $\alpha$ ,  $b_{\alpha} = 1$  or -1.

<sup>2020</sup> Mathematics Subject Classification. Primary: 11G50, 31A15; Secondary:11R06, 30C85.

Key words and phrases. height bound, capacity, conjugate algebraic integers, Mahler measure.

We denote by  $\operatorname{Conj}(\alpha)$  the set of all conjugates of  $\alpha$  which is same as the set of all roots of  $P_{\alpha}$  in  $\mathbb{C}$ . Recall that the (absolute logarithmic) Weil height of  $\alpha \in \overline{\mathbb{Q}}$  is given by

$$h(\alpha) = \frac{1}{\deg \alpha} \Big( \log |b_{\alpha}| + \sum_{x \in \operatorname{Conj}(\alpha)} \max\{0, \log |x|\} \Big).$$

We refer the reader to [BG06] for an introduction to the theory of heights. Note that  $\log^+ |z| := \max\{0, \log |z|\}$  is the Green's function of the unit circle centered at 0 with pole at  $\infty$ . For an arbitrary compact set  $\Sigma$  with positive capacity, the *height of*  $\alpha \in \overline{\mathbb{Q}}$  with respect to  $\Sigma$  is defined as

$$h_{\Sigma}(\alpha) = \frac{1}{\deg \alpha} \Big( \log |b_{\alpha}| + \sum_{x \in \operatorname{Conj}(\alpha)} g_{\Sigma}(x, \infty) \Big),$$
(1.1)

where  $g_{\Sigma}(z, \infty)$  denotes the Green's function of the set  $\Sigma$  with pole at  $\infty$ . Loosely speaking,  $h_{\Sigma}(\alpha)$  quantifies the distances of conjugates of  $\alpha$  from  $\Sigma$ . See Section 2 for more details about the Green's function and height  $h_{\Sigma}$ . Also, see [Rum99] and [Pri11] for an application of this height to limit distributions of sets of conjugate algebraic numbers.

In the case  $c(\Sigma) \geq 1$  and  $\Sigma$  is symmetric about the real axis, it follows from the Fekete-Szegö result in [FS55] that there exists a sequence  $\{\alpha_n\}$  in  $\overline{\mathbb{Q}}$  satisfying  $h_{\Sigma}(\alpha_n) \to 0$ . When  $0 < c(\Sigma) < 1$ , we show a nonzero uniform lower bound on  $\liminf_{n\to\infty} h_{\Sigma}(\alpha_n)$ .

**Theorem 1.2.** Let  $\Sigma \subset \mathbb{C}$  be compact and symmetric about the real axis. Assume that  $0 < c(\Sigma) < 1$ . Then for every sequence  $\{\alpha_n\}$  in  $\overline{\mathbb{Q}}$  such that deg  $\alpha_n \to \infty$ , we have

$$\liminf_{n \to \infty} h_{\Sigma}(\alpha_n) \ge \frac{1}{2} \log \frac{1}{c(\Sigma)}.$$

For any  $\Sigma$  with positive capacity,  $g_{\Sigma}(z, \infty) - \log^+ |z|$  is bounded, thus there exist a constant C such that  $|h(\alpha) - h_{\Sigma}(\alpha)| < C$  for every  $\alpha \in \overline{\mathbb{Q}}$ . Therefore, the height  $h_{\Sigma}$  also has the *Northcott property*: for every M > 0 and D > 0, the set  $\{\alpha \in \overline{\mathbb{Q}} : h_{\Sigma}(\alpha) \leq M, \deg \alpha \leq D\}$  is a finite set. Theorem 1.2 and the Northcott property of  $h_{\Sigma}$  immediately give us the following effective version of Result 1.1:

**Corollary 1.3.** Let  $\Sigma \subset \mathbb{C}$  be compact and symmetric about the real axis. Assume that  $0 < c(\Sigma) < 1$ . Then for any  $\epsilon > 0$  there are only a finite number of algebraic **numbers**  $\alpha$  such that

$$h_{\Sigma}(\alpha) \le \frac{1}{2} \log \frac{1}{c(\Sigma)} - \epsilon.$$

It follows from the properties of  $g_{\Sigma}(x, \infty)$  that if  $\alpha$  is an algebraic integer all of whose conjugates are in a small neighborhood of  $\Sigma$  then  $h_{\Sigma}(\alpha)$  is also small.

We next give an application of Theorem 1.2 to the growth of the leading coefficient of certain polynomial sequences. Let  $\mathbb{Z}[z]$  denote the set of polynomials with integer coefficients. Issai Schur showed that restricting the location of zeros of polynomials in  $\mathbb{Z}[z]$  with simple zeros to [-1, 1] leads to geometric growth of the leading coefficients of these polynomials. **Result 1.4** (Schur [Sch18], Satz VII). Let  $\{Q_n\}$  be a sequence of polynomials in  $\mathbb{Z}[z]$  with simple zeros such that deg  $Q_n \to \infty$  and all roots of  $Q_n$  are in [-1, 1]. Then

$$\liminf_{n \to \infty} |\text{lead}(Q_n)|^{1/\deg Q_n} \ge \sqrt{2},$$

where  $lead(Q_n)$  is the leading coefficient of  $Q_n$ .

For  $k \in \mathbb{N}$ , we denote by  $\mathbb{Z}^{k}[z]$  the set of polynomials in  $\mathbb{Z}[z]$  whose zeroes are repeated at most k times. A version of Theorem 1.2, Theorem 3.5, is valid in a more general setting of a sequence of polynomials in  $\mathbb{Z}^{k}[z]$ , using the notion of Mahler measure of a polynomial in place of the height of an algebraic number (see Subsection 2.2). As an application, we get a lower bound on the growth of the leading coefficients of a sequence of polynomials in  $\mathbb{Z}^{k}[z]$  whose roots are "near"  $\Sigma$ .

**Corollary 1.5.** Let  $\Sigma$  be as in the statement of Theorem 1.2. Let  $k \in \mathbb{N}$ . Assume that  $\{Q_n\}$  is a sequence of polynomials in  $\mathbb{Z}^k[z]$  such that deg  $Q_n \to \infty$  and

$$\frac{1}{\deg Q_n} \sum_{Q_n(x)=0} \mathbf{g}_{\Sigma}(x,\infty) \to 0$$

(e.g., if all roots of  $\{Q_n\}$  eventually belong to any neighborhood of  $\Sigma$ ). Then

$$\liminf_{n \to \infty} |\operatorname{lead}(Q_n)|^{1/\deg Q_n} \ge \frac{1}{\sqrt{c(\Sigma)}}$$

For the interval [-1, 1] we have c([-1, 1]) = 1/2; compare with Result 1.4. Pritsker (Proposition 1.7 in [Pri05]) proved a similar bound with the assumption that the roots lie in the set and with an additional hypothesis of irreducibility.

Having given a lower bound on heights in Theorem 1.2, we now turn our attention to an upper bound. By Result 1.1, if  $c(\Sigma) < 1$ , then there exists a neighborhood U of  $\Sigma$  such that there are only finitely many algebraic integers  $\alpha$  with the property that  $\alpha$ and all its conjugates lie in U. We then ask: can we find a sequence  $\{\alpha_n\}$  of algebraic integers such that "most" conjugates of  $\alpha_n$  are eventually near  $\Sigma$ ? In general, this can be achieved if the heights  $h_{\Sigma}(\alpha_n)$  go to  $\infty$ ; we show that this can also be achieved while keeping the heights bounded.

**Theorem 1.6.** Let  $\Sigma \subset \mathbb{C}$  be compact and symmetric about the real axis. If  $0 < c(\Sigma) < 1$ , then for every neighborhood U of  $\Sigma$ , there exists a sequence  $\{\alpha_n\}$  of distinct algebraic integers such that

$$\frac{\sharp\{\operatorname{Conj}(\alpha_n) \cap U\}}{\deg \alpha_n} \longrightarrow 1 \quad as \quad n \longrightarrow \infty.$$
(1.2)

Moreover, there exists a sequence  $\{\alpha_n\}$  of distinct algebraic integers such that  $\{\alpha_n\}$ satisfy (1.2) for any neighborhood U of  $\Sigma$  and  $h_{\Sigma}(\alpha_n) \to -\log c(\Sigma)$  as  $n \to \infty$ .

Owing to Result 1.1, deg  $\alpha_n \to \infty$  in the above theorem. In fact, we can find a sequence  $\{\alpha_n\}$  as in Theorem 1.6 such that the limiting distribution of the probability measures uniformly distributed on all conjugates of  $\alpha_n$  of is given by the equilibrium measure of  $\Sigma$  (see Subsection 2.1 for the definition; also see Remark 4.1). Theorem 1.6 together with

Theorem 1.2 immediately give us uniform bounds on minimal asymptotics of  $\{h_{\Sigma}(\alpha_n)\}$  over sequences of algebraic numbers.

**Corollary 1.7.** Let  $\Sigma$  be as in the statement of Theorem 1.2. Let L denote the infimum of  $\liminf_{n\to\infty} h_{\Sigma}(\alpha_n)$  over all sequences of algebraic numbers. Then

$$\frac{1}{2}\log\frac{1}{c(\Sigma)} \le L \le \log\frac{1}{c(\Sigma)}.$$

**Outine of proofs:** We briefly mention the idea of the proof of Theorem 1.2. Let  $\{\alpha_n\}$  be a sequence in  $\overline{\mathbb{Q}}$  for which  $\{h_{\Sigma}(\alpha_n)\}$  is convergent. Let  $\mu_n$  be the probability measure uniformly distributed on all conjugates of  $\alpha_n$ , i.e.,

$$\mu_n := \frac{1}{\deg \alpha_n} \sum_{x \in \operatorname{Conj}(\alpha_n)} \delta_x$$

It follows from Lemma 2.4 that the sequence  $\{\mu_n\}$  has a convergent subsequence. Although a limiting measure  $\mu$  need not be compactly supported, it turns out that such a  $\mu$  has finite logarithmic energy  $I(\mu)$  and satisfies  $\int \log^+ |z| d\mu < \infty$ . This allows us to use potential-theoretic methods. Refining the argument in Serre [Ser19], we further show that  $\mu$  satisfies the following arithmetic property: for any  $Q \in \mathbb{Z}[z]$ , we have

$$\frac{1}{\deg Q} \int \log |Q(z)| \, d\mu(z) \ge -\lim_{n \to \infty} \frac{\log |b_n|}{\deg \alpha_n} - \mathcal{E}(\Sigma; \{\alpha_n\}),$$

where  $b_n := b_{\alpha_n}$  and  $\mathcal{E}(\Sigma; \{\alpha_n\})$  is the escape rate of the sequence  $\alpha_n$  as defined in (3.1). We use this property to get a bound on the logarithmic energy  $I(\mu)$ :

$$I(\mu) \le 2 \left( \lim_{n \to \infty} \frac{\log |b_n|}{\deg \alpha_n} + \mathcal{E}(\Sigma; \{\alpha_n\}) \right)$$

Finally, we use a convexity property of the logarithmic energy to conclude our result.

To prove Theorem 1.6, we construct a sequence of auxiliary sets  $E_n$  so that  $c(\Sigma \cup E_n) = 1$ . The set  $E_n$  is chosen to be a subset of the level set  $g_{\Sigma}(z, \infty) = n$  and symmetric about the real axis. Let  $\nu_n$  be the equilibrium measure of  $\Sigma \cup E_n$ . We solve a Dirichlet problem to determine how much mass  $\nu_n$  puts on the set  $E_n$ . Using this information in the definition of height together with the Fekete-Szegö result in [FS55], we construct the appropriate sequences  $\{\alpha_n\}$ .

**Remarks on**  $c(\Sigma) \geq 1$  **case:** As noted earlier, in [FS55], Fekete–Szegö showed the converse of Result 1.1, i.e., if  $c(\Sigma) \geq 1$ , then every neighborhood U of  $\Sigma$  contains infinitely many algebraic integers  $\alpha$  so that  $\alpha$  and all its conjugates lie in U. By shrinking U, we can get a sequence  $\{\alpha_n\}$  of algebraic integers such that the sets of conjugates of  $\alpha_n$  are eventually contained in any neighborhood of  $\Sigma$ . Bilu [Bil97] studied limit distributions of the measures  $\mu_n$  associated to such sequences of algebraic integers, more generally, sequences  $\{\alpha_n\}$  in  $\overline{\mathbb{Q}}$  satisfying  $h(\alpha_n) \to 0$ , in the case of the unit circle centered at 0. Using potential-theoretic methods, limit distributions of  $\mu_n$  associated to sequences  $\{\alpha_n\}$  in  $\overline{\mathbb{Q}}$  satisfying  $h_{\Sigma}(\alpha_n) \to 0$  have been studied for general sets  $\Sigma$  of logarithmic capacity 1 by Rumely [Rum99] and Pritsker [Pri11]. For such sequences  $\{\alpha_n\}$  in  $\mathbb{Q}$ , they showed the limiting distribution of  $\mu_n$  is given by the equilibrium measure of  $\Sigma$ . Recently potential-theoretic methods have been employed when  $c(\Sigma) > 1$  in [Smi21] to study limit distributions in connection with the Schur–Siegel–Smyth trace problem (see also [OS23]).

We include the necessary potential-theoretic background in the next section, including a non-standard result on logarithmic potentials of certain measures with unbounded support. Sections 3 and 4 give the proofs of Theorems 1.2 and 1.6 respectively. We conclude in Section 5 with some further questions.

### 2. Preliminaries

2.1. **Potential theory.** We work exclusively in the complex plane  $\mathbb{C}$ . The term measure will refer to a positive Borel measure in  $\mathbb{C}$  unless otherwise stated. Given such a measure  $\mu$ , we can form its logarithmic potential function

$$U_{\mu}(z) = \int \log \frac{1}{|z-w|} d\mu(w)$$

provided this integral exists. In this case, if  $\mu$  has compact support, then

$$U_{\mu}(z) = \mu(\mathbb{C})(-\log|z| + O(1)) \text{ as } z \to \infty.$$

$$(2.1)$$

This is a lower semicontinuous, superharmonic function on  $\mathbb{C}$  which is harmonic off of the support of  $\mu$ . The logarithmic energy of  $\mu$  is

$$I(\mu) := \int \int \log \frac{1}{|z-w|} d\mu(w) d\mu(z);$$

this may be  $+\infty$ . If  $\mu$  is a positive measure with unbounded support such that

$$\int \log^+ |z| \, d\mu(z) < \infty,$$

then  $U_{\mu}$  is still locally integrable and superharmonic, and  $I(\mu) > -\infty$  (see [CKL98]).

Given a compact set  $K \subset \mathbb{C}$ , let  $\mathcal{M}(K)$  denote the cone of probability measures supported in K. The logarithmic capacity of K is

$$c(K) := \exp[-\inf\{I(\mu) : \mu \in \mathcal{M}(K)\}];$$

this coincides with the transfinite diameter d(K) in the introduction (see, for instance, [Ahl10, Ch 2] or [Ran95, Ch 5]). If c(K) > 0, the Frostman conditions (cf., [Ran95]) tell us that there is a unique measure  $\mu_K \in \mathcal{M}(K)$  with  $c(K) = \exp[-I(\mu_K)]$  and  $U_{\mu_K} = I(\mu_K)$  quasi-everywhere (abbreviated as q.e.) on K; i.e., on all of K except for a polar set. Here,  $E \subset \mathbb{C}$  is polar if there exists a subharmonic function  $u \not\equiv -\infty$  on a neighborhood of E with  $E \subset \{u = -\infty\}$ ; for E compact this is equivalent to c(E) = 0. If  $E \subset \mathbb{C}$  is compact and polar, for any  $K \subset \mathbb{C}$  compact we have  $c(K \cup E) = c(K)$ . The measure  $\mu_K$  is called the *equilibrium measure of* K.

For non-polar compact K, we will have use of the Green's function

$$g_K(z,\infty) := -U_{\mu_K}(z) + I(\mu_K)$$

for (the unbounded component of the complement of) K with pole at  $\infty$ ; note that  $g_K(z,\infty) = 0$  q.e. on K. This is an upper semicontinuous function which is harmonic outside of K; from (2.1), it follows that  $g_{\Sigma}(z,\infty) - \log^+ |z|$  is bounded as stated in the introduction. If  $U_{\mu_K}$  (equivalently,  $g_K$ ) is continuous, we say that K is regular. This is the same as  $U_{\mu_K} = I(\mu_K)$  ( $g_K = 0$ ) on K (cf. [Ran95]). The Green's function of the circle (or closed disk) of radius r centered at a is  $\log^+(|z-a|/r)$ . For the real interval K = [-1, 1], we have

$$g_K(z,\infty) = \log|z + \sqrt{z^2 - 1}|$$

More generally, the Green's function of  $K = [a, b] \subset \mathbb{R}$  is given by

$$g_K(z,\infty) = \log |f(z) + \sqrt{f(z)^2 - 1}|,$$

where  $f(z) = \frac{2}{b-a}z - \frac{a+b}{b-a}$ , and  $c(K) = \frac{b-a}{4}$ . If K is the Julia set of a polynomial P of degree  $n \ge 2$  then its Green's function is

$$g_K(z,\infty) = \lim_{j \to \infty} \frac{1}{n^j} \log^+ |P^j(z)|,$$

where  $P^{j}$  is the *j*-fold composition of P with itself and  $c(K) = |\text{lead}(P)|^{-1/(n-1)}$ .

It is a standard fact from potential theory that the logarithmic energy of compactly supported measures satisfies a convexity property (see [Ahl10, Lemma 2.1] or [ST97, Lemma I.1.8]). It turns out that this property also holds for certain measures that are not necessarily compactly supported; later we will need to consider measures  $\mu$  with unbounded support.

**Result 2.1** (Theorem 2.5 of [CKL98]). Let  $\tau$  be a signed measure with  $\tau(\mathbb{C}) = 0$  that satisfies  $\int \log^+ |z| \, d|\tau| < \infty$  and  $I(|\tau|) < \infty$ . Then the logarithmic energy  $I(\tau)$  of  $\tau$  is well-defined and  $I(\tau) \ge 0$ . In particular, if  $\tau_1$  and  $\tau_2$  are probability measures satisfying  $\int \log^+ |z| \, d\tau_i < \infty$  and  $I(\tau_i) < \infty$  for i = 1, 2, then, considering  $\tau = \tau_1 - \tau_2$ ,

$$2\int U_{\tau_1}(z) \, d\tau_2(z) \le I(\tau_1) + I(\tau_2).$$

In the case where we consider  $\tau = \tau_1 - \tau_2$ , the inequality follows because

$$I(\tau) = I(\tau_1) + I(\tau_2) - 2 \int U_{\tau_1}(z) \, d\tau_2(z).$$

2.2. Heights and Mahler measure. We recall the notion of height of  $\alpha \in \overline{\mathbb{Q}}$  with respect to a non-polar compact set  $\Sigma$  as defined in (1.1). Let  $P_{\alpha} \in \mathbb{Z}[z]$  be an irreducible polynomial with leading coefficient  $b_{\alpha}$  and degree deg  $\alpha$ . Then

$$h_{\Sigma}(\alpha) := \frac{1}{\deg \alpha} \Big( \log |b_{\alpha}| + \sum_{x \in \operatorname{Conj}(\alpha)} g_{\Sigma}(x, \infty) \Big).$$

The Mahler measure of a polynomial  $P_n(z) = a_n \prod_{j=1}^n (z - \alpha_j)$  of degree *n* (thus  $a_n \neq 0$ ) is

$$M(P_n) := \exp\left[\frac{1}{2\pi} \int_0^{2\pi} \log|P_n(e^{it})|dt\right] = |a_n| \prod_{j=1}^n \max\{1, |\alpha_j|\},$$

the latter equality following from Jensen's formula. Since the Green's function for the unit circle  $\Sigma = \{|z| = 1\}$  is  $g_{\Sigma}(z, \infty) = \log^+ |z| := \max\{0, \log |z|\}$ , we have

$$\frac{1}{n}\log M(P_n) = \frac{1}{n}\log|a_n| + \frac{1}{n}\sum_j g_{\Sigma}(\alpha_j, \infty).$$

This coincides, in the case where  $P_n \in \mathbb{Z}[z]$  is an irreducible polynomial with roots  $\alpha_1, ..., \alpha_n \in \overline{\mathbb{Q}}$ , to the Weil height  $h(\alpha_j)$ .

More generally, following Pritsker [Pri11], we define the Mahler measure of a polynomial  $P_n(z) = a_n \prod_{j=1}^n (z - \alpha_j)$  of degree n with respect to any non-polar compact set  $\Sigma$  as

$$M_{\Sigma}(P_n) := |a_n| \exp\left[\sum_{j=1}^n g_{\Sigma}(\alpha_j, \infty)\right]$$

so that, if  $P_n \in \mathbb{Z}[z]$  is an irreducible polynomial with roots  $\alpha_1, ..., \alpha_n \in \overline{\mathbb{Q}}$ , we again have the relation

$$\frac{1}{n}\log M_{\Sigma}(P_n) = h_{\Sigma}(\alpha_j).$$
(2.2)

Observe that if  $P = P_1 P_2 \cdots P_n$  with deg  $P_i > 0$  for every  $1 \le i \le n$ , then we have

$$M_{\Sigma}(P)^{\frac{1}{\deg P}} = \prod_{i=1}^{n} \left( M_{\Sigma}(P_i)^{\frac{1}{\deg P_i}} \right)^{\frac{\deg P_i}{\deg P}}$$
(2.3)

where  $\sum_{i=1}^{n} \frac{\deg P_i}{\deg P} = 1.$ 

Using this generalized notion of Mahler measure, Pritsker proved the following:

**Result 2.2** (Pritsker, [Pri11]). Let  $P_n(z) = a_n \prod_{j=1}^n (z - \alpha_{j,n})$  be a sequence of polynomials with integer coefficients and simple zeros. Let  $\Sigma \subset \mathbb{C}$  be compact with  $c(\Sigma) = 1$ . Then

$$\lim_{n \to \infty} (M_{\Sigma}(P_n))^{1/n} = 1$$

if and only if

(1)  $\lim_{n\to\infty} |a_n|^{1/n} = 1;$ (2)  $\lim_{R\to\infty} \lim_{n\to\infty} (\prod_{|\alpha_{j,n}|\geq R} |\alpha_{j,n}|)^{1/n} = 1;$  and (3)  $\mu_n := \frac{1}{n} \sum_{j=1}^n \delta_{\alpha_{j,n}} \to \mu_{\Sigma}$  in the weak\* topology as  $n \to \infty$ .

As an immediate corollary, we have the following result, which will be used later on.

**Lemma 2.3.** Let  $\Sigma \subset \mathbb{C}$  be compact and symmetric about the real axis. Suppose that  $c(\Sigma) = 1$ . Let  $\{\alpha_n\}$  be a sequence of distinct algebraic integers such that for every neighborhood U of  $\Sigma$  there exist  $N \in \mathbb{N}$  such that  $\operatorname{Conj}(\alpha_n) \subset U$  for all  $n \geq N$ . Then

$$\mu_n := \frac{1}{\deg \alpha_n} \sum_{x \in \operatorname{Conj}(\alpha_n)} \delta_x \longrightarrow \mu_{\Sigma} \quad as \quad n \longrightarrow \infty,$$

where  $\mu_{\Sigma}$  is the equilibrium measure of  $\Sigma$ .

Next, we recall that the *resultant* of two polynomials

$$P(z) = a_n z^n + \dots + a_0 = a_n \prod_{j=1}^n (z - \alpha_j)$$
  
and 
$$Q(z) = b_m z^m + \dots + b_0 = b_m \prod_{k=1}^m (z - \beta_k)$$

is given by

$$\operatorname{Res}(P,Q) := a_n b_m \prod_{j=1,\dots,n;\ k=1,\dots,m} (\alpha_j - \beta_k).$$

For  $P, Q \in \mathbb{Z}[z]$  it follows that  $\operatorname{Res}(P, Q) \in \mathbb{Z}$  (cf., [CLO92], page 153). Thus, if  $P, Q \in \mathbb{Z}[z]$  do not have any factor in common then  $|\operatorname{Res}(P, Q)| \ge 1$ . The discriminant of P is related to the resultant of P and P':

$$\operatorname{Disc}(P) = \frac{(-1)^{n(n-1)/2}}{a_n} \cdot \operatorname{Res}(P, P') = (-1)^{n(n-1)/2} a_n^{2n-2} \prod_{j \neq k} (\alpha_j - \alpha_k).$$
(2.4)

The reader will notice the resemblance to the quantity  $\prod_{i < j} |\xi_i - \xi_j|^{\frac{2}{n(n-1)}}$  in the definition of transfinite diameter from the introduction.

Finally, recall that a sequence of measures  $\{\mu_{\beta}\}_{\beta \in B}$  on  $\mathbb{C}$  is *tight* if for any  $\epsilon > 0$  there exists a compact set  $K_{\epsilon}$  such that  $\mu_{\beta}(\mathbb{C} \setminus K_{\epsilon}) < \epsilon$  for all  $\beta \in B$ . If  $\{\mu_{\beta}\}_{\beta \in B}$  is tight and uniformly bounded in the total variation norm, Prokhorov's theorem implies that it has a convergent subsequence. We will need the following application.

**Lemma 2.4.** Let  $\{\alpha_n\}$  be a sequence in  $\overline{\mathbb{Q}}$  such that corresponding sequence of heights  $\{h_{\Sigma}(\alpha_n)\}$  is bounded. If

$$\mu_n := \frac{1}{\deg \alpha_n} \sum_{x \in \operatorname{Conj}(\alpha_n)} \delta_x$$

is the probability measure uniformly distributed on all conjugates of  $\alpha_n$ , then the sequence  $\{\mu_n\}$  has a convergent subsequence in the weak\* topology.

Proof. We shall show that the sequence  $\{\mu_n\}$  is tight. Then the result follows from Prokhorov's theorem. Assume that the sequence  $\{\mu_n\}$  is not tight. By definition, there exists  $\rho > 0$  such that for every compact set  $K \subset \mathbb{C}$ , we have  $\mu_n(\mathbb{C} \setminus K) > \rho$  for some  $n \in \mathbb{N}$ . Since  $g_{\Sigma}(z, \infty) \to \infty$  as  $z \to \infty$ , by enlarging K, we get an  $\alpha \in \overline{\mathbb{Q}}$  from the sequence  $\{\alpha_n\}$  for which  $h_{\Sigma}(\alpha)$  is arbitrarily large. This contradicts the assumption that  $\{h_{\Sigma}(\alpha_n)\}$  is bounded. Thus the sequence  $\{\mu_n\}$  is tight, and there exists a convergent subsequence of  $\{\mu_n\}$ .

# 3. Proof of Theorem 1.2

Let  $\Sigma \subset \mathbb{C}$  be compact and symmetric about the real axis with  $0 < c(\Sigma) < 1$ . Given a sequence  $\{\alpha_n\}$  in  $\overline{\mathbb{Q}}$ , for simplicity of notation, we let  $b_n := b_{\alpha_n}$  and  $P_n := P_{\alpha_n}$ . We want a lower bound on  $\liminf_{n\to\infty} h_{\Sigma}(\alpha_n)$  when  $\deg \alpha_n \to \infty$ . If  $\liminf_{n\to\infty} h_{\Sigma}(\alpha_n) < \infty$ , then

there exists a subsequence  $\{\alpha_{n_k}\}$  for which  $\{h_{\Sigma}(\alpha_{n_k})\}$  and  $\{\log |b_{n_k}|/\deg \alpha_{n_k}\}$  are both convergent. Thus, without loss of generality, we make the following assumptions:

- (a) deg  $\alpha_n \to \infty$ ;
- (b) the sequence of heights  $\{h_{\Sigma}(\alpha_n)\}$  is convergent; and
- (c) the sequence  $\{\log |b_n|/\deg \alpha_n\}$  is convergent.

As in Lemma 2.4, we let

$$\mu_n := \frac{1}{\deg \alpha_n} \sum_{x \in \operatorname{Conj}(\alpha_n)} \delta_x.$$

This lemma ensures that there exists a convergent subsequence of  $\{\mu_n\}$ . In general, the limiting measure  $\mu$  of the subsequence need not be compactly supported. However, in the next result, we shall see that the limiting measure satisfies an interesting property that allows us to use potential-theoretic tools.

**Result 3.1** (Theorem 2.1 of [Pri15]). Let  $\{\alpha_n\}$  be a sequence in  $\overline{\mathbb{Q}}$  satisfying (a)–(c). Let  $\{\mu_n\}$  be as above. Assume that  $\mu_n$  converges to a probability measure  $\mu$  in the weak\* topology. Then,  $\mu$  has finite logarithmic energy  $I(\mu)$ . Furthermore, we have that

$$\int \log^+ |z| \, d\mu(z) < \infty.$$

This result has been stated in [Pri15] under the assumption that  $h(\alpha_n)$  is bounded, where h denotes the standard Weil height. However, as noted in the introduction, there exist a constant  $C = C(\Sigma)$  such that  $|h(\alpha) - h_{\Sigma}(\alpha)| < C$  for every  $\alpha \in \overline{\mathbb{Q}}$ . Thus the result holds with our assumptions as well.

By Result 3.1, if  $\{\mu_n\}$  converges to a probability measure  $\mu$ , then the logarithmic energy of  $\mu$  is finite. We shall now find a bound on its energy. We make a definition at this stage. Recall the notation:  $g_{\Sigma}(z, \infty)$  denotes the Green's function of  $\Sigma$  with pole at  $\infty$ .

**Definition 3.2.** Let  $\{\beta_n\}$  be a sequence in  $\overline{\mathbb{Q}}$ . The *escape rate* of the sequence  $\{\beta_n\}$  with respect to  $\Sigma$  is defined as

$$\mathcal{E}(\Sigma; \{\beta_n\}) = \lim_{r \to \infty} \left(\liminf_{n \to \infty} \frac{1}{\deg \beta_n} \sum_{\substack{x \in \operatorname{Conj}(\beta_n) \\ |x| \ge r}} g_{\Sigma}(x, \infty)\right).$$
(3.1)

Note that if  $\{h_{\Sigma}(\beta_n)\}$  is bounded, the escape rate  $\mathcal{E}(\Sigma; \{\beta_n\})$  is a finite positive number. We now show that the measures that give the limiting distribution of the conjugates of a sequence in  $\overline{\mathbb{Q}}$  satisfy an interesting arithmetic property. In the case of a sequence of algebraic integers with bounded conjugates lying near a compact, symmetric set  $\Sigma$  with  $c(\Sigma) \geq 1$ , the arithmetic property characterizes such measures (see Section 5; also see [Smi21], [OS23]).

**Lemma 3.3.** Let  $\{\alpha_n\}$  be a sequence in  $\overline{\mathbb{Q}}$  satisfying (a)-(c). If  $\{\mu_n\}$  converges to a probability measure  $\mu$  in the weak\* topology, then for any polynomial  $Q \in \mathbb{Z}[z]$ , we have

$$\frac{1}{\deg Q} \int \log |Q(z)| \, d\mu(z) \ge -\lim_{n \to \infty} \frac{\log |b_n|}{\deg \alpha_n} - \mathcal{E}(\Sigma; \{\alpha_n\}),$$

where  $\mathcal{E}(\Sigma; \{\alpha_n\})$  is the escape rate of the sequence  $\{\alpha_n\}$  as defined in (3.1).

*Proof.* Fix a polynomial  $Q \in \mathbb{Z}[z]$ . Let  $\phi$  be a non-negative compactly supported continuous function. Since  $\phi(z) \log |Q(z)|$  is an upper semicontinuous function that is bounded from above, we have

$$\frac{1}{\deg Q} \int \log |Q(z)| \, d\mu(z) \ge \limsup_{n \to \infty} \frac{1}{\deg Q} \int \phi(z) \log |Q(z)| \, d\mu_n(z)$$
$$\ge \liminf_{n \to \infty} \frac{1}{\deg Q} \int \log |Q(z)| \, d\mu_n(z)$$
$$-\liminf_{n \to \infty} \frac{1}{\deg Q} \int (1 - \phi(z)) \log |Q(z)| \, d\mu_n(z). \tag{3.2}$$

For the last inequality, we use the fact that  $\limsup_{n\to\infty} (k_n + l_n) \ge \liminf_{n\to\infty} k_n + \limsup_{n\to\infty} l_n$ . Note that, owing to Result 3.1,  $\int \log |Q(z)| d\mu(z)$  makes sense.

Since  $P_n$  and Q do not have roots in common for large n, we have  $|\text{Res}(Q, P_n)| \ge 1$ . Thus we get

$$\frac{1}{\deg Q} \int \log |Q(z)| \, d\mu_n(z) = \frac{\log |\operatorname{Res}(Q, P_n)| - \log |b_n^{\deg Q}|}{\deg Q \cdot \deg P_n} \ge -\frac{\log |b_n|}{\deg \alpha_n}.$$
(3.3)

Observe that, as  $z \to \infty$ ,  $g_{\Sigma}(z, \infty) = \log |z| + O(1)$  and  $(1/\deg Q) \log |Q(z)| = \log |z| + O(1)$ . We denote the disk of radius r centered at the origin by  $\mathbb{D}(0; r)$  and its complement by  $\mathbb{D}(0; r)^c$ . For every  $\delta > 0$  there exists r >> 1 such that  $\mu(\mathbb{D}(0; r)^c) < \delta$  and  $\mu_n(\mathbb{D}(0; r)^c) < \delta$  for sufficiently large n. Thus if  $0 \le \phi \le 1$  with  $\phi \equiv 1$  on  $\mathbb{D}(0; r)$  for sufficiently large r, we get

$$\liminf_{n \to \infty} \frac{1}{\deg Q} \int (1 - \phi(z)) \log |Q(z)| \, d\mu_n(z)$$
  
$$\leq \liminf_{n \to \infty} \frac{1}{\deg \alpha_n} \sum_{\substack{x \in \operatorname{Conj}(\alpha_n) \\ |x| \ge r}} g_{\Sigma}(x, \infty) + \delta \cdot O(1).$$
(3.4)

Since the choice of  $\phi$  in (3.2) is arbitrary, by using (3.3) and (3.4) in (3.2), it follows that

$$\frac{1}{\deg Q} \int \log |Q(z)| \, d\mu(z) \ge -\lim_{n \to \infty} \frac{\log |b_n|}{\deg \alpha_n} - \mathcal{E}(\Sigma; \{\alpha_n\}).$$

We now use the above lemma to get an upper bound on  $I(\mu)$ . Here we continue with the same notation as above.

**Proposition 3.4.** Let  $\{\alpha_n\}$  be a sequence in  $\overline{\mathbb{Q}}$  satisfying (a)–(c). If  $\{\mu_n\}$  converges to a probability measure  $\mu$  in the weak\* topology, then

$$I(\mu) \le 2 \left( \lim_{n \to \infty} \frac{\log |b_n|}{\deg \alpha_n} + \mathcal{E}(\Sigma; \{\alpha_n\}) \right).$$

where  $\mathcal{E}(\Sigma; \{\alpha_n\})$  is the escape rate of the sequence  $\{\alpha_n\}$  as defined in (3.1).

Proof. Let  $\phi$  be a cut-off function as in Lemma 3.3 so that  $0 \leq \phi \leq 1$  with  $\phi \equiv 1$  on  $\mathbb{D}(0;r)$  for sufficiently large r > 0. Since  $I(\mu) < \infty$  from Result 3.1, given  $\delta > 0$ , we can choose r >> 1 large so that  $\int \phi(z)U_{\mu}(z) d\mu(z) > I(\mu) - \delta$ ; in addition, as in the previous lemma, for such r we may assume  $\mu(\mathbb{D}(0;r)^c) < \delta$  and  $\mu_n(\mathbb{D}(0;r)^c) < \delta$ . Then since  $\phi(z)U_{\mu}(z)$  is a lower semicontinuous function that is bounded from below, we have

$$I(\mu) - \delta < \int \phi(z) U_{\mu}(z) \, d\mu(z) \le \liminf_{n \to \infty} \int \phi(z) U_{\mu}(z) \, d\mu_n(z)$$
$$\le \limsup_{n \to \infty} \int U_{\mu}(z) \, d\mu_n(z)$$
$$-\liminf_{n \to \infty} \int (1 - \phi(z)) U_{\mu}(z) \, d\mu_n(z). \tag{3.5}$$

For the last inequality, we use the fact that  $\liminf_{n\to\infty} (k_n + l_n) \leq \limsup_{n\to\infty} k_n + \limsup_{n\to\infty} l_n$ . By Fubini's theorem, we see that

$$\int U_{\mu}(z) \, d\mu_n(z) = \int U_{\mu_n}(z) \, d\mu(z) = -\frac{1}{\deg P_n} \int \log |P_n(z)| \, d\mu(z) + \frac{\log |b_n|}{\deg P_n}$$

Since Lemma 3.3 holds for an arbitrary polynomial Q with integer coefficients, in particular it holds for the polynomials  $P_n$ . Therefore, fixing  $Q = P_n$ , for all m sufficiently large,  $P_n$  and  $P_m$  have no common factors and we get

$$\limsup_{n \to \infty} \int U_{\mu}(z) \, d\mu_n(z) \le \lim_{n \to \infty} \frac{\log |b_n|}{\deg \alpha_n} + \mathcal{E}(\Sigma; \{\alpha_n\}) + \lim_{n \to \infty} \frac{\log |b_n|}{\deg P_n}.$$
 (3.6)

Now as  $z \to \infty$ ,

$$U_{\mu}(z) = \int_{\mathbb{D}(0;r)} \log \frac{1}{|z-w|} d\mu(w) + \int_{\mathbb{D}(0;r)^{c}} \log \frac{1}{|z-w|} d\mu(w)$$
  

$$\geq (1-\delta)(-\log|z| + O(1)) + \int_{\mathbb{D}(0;r)^{c}} \log \frac{1}{|z-w|} d\mu(w).$$

For any M > 0 and |z| > r,

$$\int_{\mathbb{D}(0;r)^c} \log \frac{1}{|z-w|} \, d\mu(w) \ge \int_{\mathbb{D}(0;r)^c} \min\left\{M, \log \frac{1}{|z-w|}\right\} \, d\mu(w)$$

which is finite; hence, as  $z \to \infty$ ,

$$U_{\mu}(z) \ge -\log|z| + O(1) = -g_{\Sigma}(z, \infty) + O(1).$$

Therefore, it follows that

$$\liminf_{n \to \infty} \int (1 - \phi(z)) U_{\mu}(z) \, d\mu_n(z) \ge -\liminf_{n \to \infty} \frac{1}{\deg \alpha_n} \sum_{\substack{x \in \operatorname{Conj}(\alpha_n) \\ |x| \ge r}} g_{\Sigma}(x, \infty) + \delta \cdot O(1).$$
(3.7)

Since  $\delta > 0$  is arbitrary, using (3.6) and (3.7) in (3.5), it follows that

$$I(\mu) \le 2 \Big( \lim_{n \to \infty} \frac{\log |b_n|}{\deg \alpha_n} + \mathcal{E}(\Sigma; \{\alpha_n\}) \Big).$$

We are now ready for

The proof of Theorem 1.2. We shall first prove the result when  $\Sigma$  is regular (see Section 2). Let  $\{\alpha_n\}$  be a sequence in  $\overline{\mathbb{Q}}$  satisfying (a)-(c). Observe that the result follows if we show that  $\lim_{n\to\infty} h_{\Sigma}(\alpha_n) \ge -(1/2) \log c(\Sigma)$ . With this notation, we have

$$h_{\Sigma}(\alpha_n) = \frac{\log |b_n|}{\deg \alpha_n} + \frac{1}{\deg \alpha_n} \sum_{x \in \operatorname{Conj}(\alpha_n)} g_{\Sigma}(x, \infty).$$

Recall  $\mu_n$  is the probability measure uniformly distributed on all conjugates of  $\alpha_n$ . By Lemma 2.4, there exists a convergent subsequence of  $\{\mu_n\}$ . We simply denote this subsequence by  $\{\alpha_n\}$  and the corresponding sequence of measures by  $\{\mu_n\}$ . Let  $\mu$  be the weak<sup>\*</sup> limit of  $\{\mu_n\}$ . Note that  $\mu$  need not be compactly supported.

Let  $\mu_{\Sigma}$  denote the equilibrium measure of  $\Sigma$ . We have  $U_{\mu_{\Sigma}}(z) = -g_{\Sigma}(z, \infty) - \log c(\Sigma)$ for  $z \in \mathbb{C}$ . Thus we get

$$h_{\Sigma}(\alpha_n) = \frac{\log |b_n|}{\deg \alpha_n} + \frac{1}{\deg \alpha_n} \sum_{x \in \operatorname{Conj}(\alpha_n)} \left( \log \frac{1}{c(\Sigma)} - U_{\mu_{\Sigma}}(x) \right)$$
$$= \log \frac{1}{c(\Sigma)} + \frac{\log |b_n|}{\deg \alpha_n} - \int U_{\mu_{\Sigma}}(z) \, d\mu_n(z).$$

Since  $\{h_{\Sigma}(\alpha_n)\}\$  and  $\{\log |b_n|/\deg \alpha_n\}\$  are both convergent, we get

$$\lim_{n \to \infty} h_{\Sigma}(\alpha_n) = \log \frac{1}{c(\Sigma)} + \lim_{n \to \infty} \frac{\log |b_n|}{\deg \alpha_n} - \lim_{n \to \infty} \int U_{\mu_{\Sigma}}(z) \, d\mu_n(z). \tag{3.8}$$

The last limit exists because all other limits in (3.8) exist. We now estimate the last limit in (3.8). By our assumption that  $\Sigma$  is regular,  $U_{\mu_{\Sigma}}$  is a continuous function. Let  $\phi$  be a non-negative compactly supported continuous function. Thus  $\phi(z)U_{\mu_{\Sigma}}(z)$  is bounded and continuous, and we get

$$\int \phi(z) U_{\mu_{\Sigma}} d\mu(z) = \lim_{n \to \infty} \int \phi(z) U_{\mu_{\Sigma}}(z) d\mu_n(z)$$
$$= \lim_{n \to \infty} \int U_{\mu_{\Sigma}}(z) d\mu_n(z) - \lim_{n \to \infty} \int (1 - \phi(z)) U_{\mu_{\Sigma}}(z) d\mu_n(z).$$
(3.9)

Fix  $\delta > 0$ . Since  $U_{\mu_{\Sigma}}(z) = -\log |z| + O(1)$  as  $z \to \infty$ , by Result 3.1, there exists r > 0 such that if  $\phi$  is a cut-off function as in Lemma 3.3 so that  $0 \le \phi \le 1$  with  $\phi \equiv 1$  on  $\mathbb{D}(0; r)$  for sufficiently large r, then

$$\int \phi(z) U_{\mu_{\Sigma}}(z) \, d\mu(z) \le (1+\delta) \int U_{\mu_{\Sigma}}(z) \, d\mu(z). \tag{3.10}$$

As in the proof of Lemma 3.3, increasing r if necessary, we have  $\mu(\mathbb{D}(0;r)^c) < \delta$  and  $\mu_n(\mathbb{D}(0;r)^c) < \delta$  for sufficiently large n. Since  $U_{\mu_{\Sigma}}(z) = -g_{\Sigma}(z,\infty) - \log c(\Sigma)$ , for large

r, as in (3.7), we also have

$$\lim_{n \to \infty} \int (1 - \phi(z)) U_{\mu_{\Sigma}}(z) \, d\mu_n(z) \leq -\liminf_{n \to \infty} \frac{1}{\deg \alpha_n} \sum_{\substack{x \in \operatorname{Conj}(\alpha_n) \\ |x| \geq r}} g_{\Sigma}(x, \infty) + \delta \cdot O(1).$$
(3.11)

Since  $\delta > 0$  is arbitrary, (3.10) and (3.11) together with (3.9) give us

$$\lim_{n \to \infty} \int U_{\mu_{\Sigma}}(z) \, d\mu_n(z) \leq \int U_{\mu_{\Sigma}}(z) \, d\mu(z) - \lim_{r \to \infty} \left(\liminf_{n \to \infty} \frac{1}{\deg \alpha_n} \sum_{\substack{x \in \operatorname{Conj}(\alpha_n) \\ |x| \ge r}} g_{\Sigma}(x, \infty)\right).$$

The limit on the right hand side of the last expression is the escape rate  $\mathcal{E}(\Sigma; \{\alpha_n\})$  of the sequence  $\{\alpha_n\}$  as in Proposition 3.4. By invoking Result 2.1 and Proposition 3.4, it follows that

$$\lim_{n \to \infty} \int U_{\mu_{\Sigma}}(z) \, d\mu_n(z) \leq \frac{I(\mu_{\Sigma}) + I(\mu)}{2} - \mathcal{E}(\Sigma; \{\alpha_n\})$$
$$\leq \frac{1}{2} \log \frac{1}{c(\Sigma)} + \lim_{n \to \infty} \frac{\log |b_n|}{\deg \alpha_n}.$$

Using this in (3.8), we get

$$\lim_{n \to \infty} h_{\Sigma}(\alpha_n) \ge \log \frac{1}{c(\Sigma)} + \lim_{n \to \infty} \frac{\log |b_n|}{\deg \alpha_n} - \frac{1}{2} \log \frac{1}{c(\Sigma)} - \lim_{n \to \infty} \frac{\log |b_n|}{\deg \alpha_n} \ge \frac{1}{2} \log \frac{1}{c(\Sigma)}.$$

Since the sequence  $\{\alpha_n\}$  is arbitrary, the result follows when  $\Sigma$  is regular.

If  $\Sigma$  is not regular, we consider a sequence  $\{\Sigma_m\}$  of regular compact sets symmetric about the real axis such that  $\Sigma_{m+1} \subset \Sigma_m$  for every  $m \in \mathbb{N}$  and  $\Sigma = \bigcap_m \Sigma_m$ . For example, we can take  $\Sigma_m = \{z \in \mathbb{C} : \operatorname{dist}(z, \Sigma) \leq 1/m\}$ . Note that  $c(\Sigma_m) \downarrow c(\Sigma)$ . Also, note that  $g_{\Sigma_m}(z, \infty) \leq g_{\Sigma}(z, \infty)$  for every  $m \in \mathbb{N}$  and  $z \in \mathbb{C}$ . Therefore, for any  $\alpha \in \overline{\mathbb{Q}}$ , we have  $h_{\Sigma_m}(\alpha) \leq h_{\Sigma}(\alpha)$  for every  $m \in \mathbb{N}$ . Let  $\{\alpha_n\}$  be a sequence in  $\overline{\mathbb{Q}}$  such that deg  $\alpha_n \to \infty$ . Since the sets  $\Sigma_m$  are regular, it follows that  $\liminf_{n\to\infty} h_{\Sigma}(\alpha_n) \geq$  $\liminf_{n\to\infty} h_{\Sigma_m}(\alpha_n) \geq -(1/2) \log c(\Sigma_m)$  for every  $m \in \mathbb{N}$ . Thus

$$\liminf_{n \to \infty} h_{\Sigma}(\alpha_n) \ge \frac{1}{2} \log \frac{1}{c(\Sigma)}.$$

We now prove a result alluded to in Section 1. Recall the definition of the Mahler measure of a polynomial from Subsection 2.2 and the relation (2.3). We shall use these in the proof of our next result. Recall that  $\mathbb{Z}^{k}[z]$  denotes the set of polynomials in  $\mathbb{Z}[z]$  whose zeroes are repeated at most k times.

**Theorem 3.5.** Let  $\Sigma$  be as in the statement of Theorem 1.2. Let  $k \in \mathbb{N}$ . Let  $\{Q_n\}$  be a sequence of polynomials in  $\mathbb{Z}^k[z]$  such that deg  $Q_n \to \infty$ . Then we have

$$\liminf_{n \to \infty} \frac{1}{\deg Q_n} \log M_{\Sigma}(Q_n) \ge \frac{1}{2} \log \frac{1}{c(\Sigma)}.$$

*Proof.* Fix  $\epsilon > 0$ . As observed in Section 1, Theorem 1.2 and the Northcott property of  $h_{\Sigma}$  imply that there are only finitely many  $\alpha \in \overline{\mathbb{Q}}$  such that  $h_{\Sigma}(\alpha) \leq -(1/2) \log c(\Sigma) - \epsilon$ . Thus there are only finitely many **irreducible** polynomials  $P \in \mathbb{Z}[z]$  satisfying

$$\frac{1}{\deg P}\log M_{\Sigma}(P) \le \frac{1}{2}\log \frac{1}{c(\Sigma)} - \epsilon.$$

Let  $\mathcal{P}$  denote the set of such polynomials.

Let  $\{Q_n\}$  be a sequence of polynomials in  $\mathbb{Z}^k[z]$  such that deg  $Q_n \to \infty$ . Write, for every  $n, Q_n = Q_n^1 Q_n^2$ , where  $Q_n^1$  is the product of factors of  $Q_n$  that are also in  $\mathcal{P}$  and  $Q_n^2$  is the product of the remaining factors of  $Q_n$ . Thus, from (2.3),

$$\frac{1}{\deg Q_n} \log M_{\Sigma}(Q_n) = \frac{\deg Q_n^1}{\deg Q_n} \left( \frac{\log M_{\Sigma}(Q_n^1)}{\deg Q_n^1} \right) + \frac{\deg Q_n^2}{\deg Q_n} \left( \frac{\log M_{\Sigma}(Q_n^2)}{\deg Q_n^2} \right).$$
(3.12)

Similar to (3.12),  $\log M_{\Sigma}(Q_n^1)/\deg Q_n^1$  can be further written as a convex combination using its irreducible sub-factors. Since these sub-factors are from the set  $\mathcal{P}$ , we have

$$\frac{\log M_{\Sigma}(Q_n^1)}{\deg Q_n^1} \le \frac{1}{2}\log \frac{1}{c(\Sigma)} - \epsilon.$$

Also,  $\log M_{\Sigma}(Q_n^2)/\deg Q_n^2$  can be further written as a convex combination using its irreducible sub-factors. Therefore,

$$\frac{\log M_{\Sigma}(Q_n^2)}{\deg Q_n^2} > \frac{1}{2}\log \frac{1}{c(\Sigma)} - \epsilon.$$

Since the zeroes of each  $Q_n$  are repeated at most k times,  $\{\deg Q_n^1\}$  is bounded. Therefore, by (3.12), there exists  $N \in \mathbb{N}$  such that for every  $n \geq N$ , we have

$$\frac{1}{\deg Q_n} \log M_{\Sigma}(Q_n) > \frac{1}{2} \log \frac{1}{c(\Sigma)} - \epsilon - \frac{\epsilon}{2}$$

Since  $\epsilon > 0$  is arbitrary, the result follows.

Corollary 1.5 follows from Theorem 3.5 (if  $\Sigma$  is not regular, for the case when all roots of  $\{Q_n\}$  eventually belong to any neighborhood of  $\Sigma$ , we use an argument as in the last part of the proof of Theorem 1.2).

### 4. Proof of Theorem 1.6

In this section, we prove Theorem 1.6. As in the last section, we assume that  $\Sigma \subset \mathbb{C}$  is compact and symmetric about the real axis with  $0 < c(\Sigma) < 1$ . Since  $\Sigma$  is symmetric about the real axis,  $g_{\Sigma}(z, \infty) = g_{\Sigma}(\bar{z}, \infty)$ ; thus level sets of  $g_{\Sigma}(z, \infty)$  are also symmetric about the real axis. For  $n \in \mathbb{N}$ , define

$$L_n := \{ z \in \mathbb{C} : g_{\Sigma}(z, \infty) = n \}$$

We choose a compact subset  $E_n$  of  $L_n$  that is symmetric about the real axis and satisfies  $c(\Sigma \cup E_n) = 1$ . For sufficiently large n, such a set always exists. To see this, we start with a compact subset E of  $L_n$  that is invariant under complex conjugation and which satisfies  $c(\Sigma \cup E) > 1$ . For example, if we choose E with c(E) > 1 then  $c(\Sigma \cup E) > 1$ .

Since capacity is a monotone function, by considering a decreasing family of subsets of  $L_n$ , we eventually get a compact subset  $E_n$  of  $L_n$  that satisfies  $c(\Sigma \cup E_n) = 1$ . Since  $c(\Sigma) < 1$ , the set  $E_n$  is non-polar.

We recall the Fekete-Szegö result in [FS55] discussed in the introduction, i.e., if  $c(K) \geq 1$  and K is symmetric about the real axis, then every neighborhood U of K contains infinitely many algebraic integers  $\alpha$  so that  $\alpha$  and all its conjugates lie in U. Let  $\nu_n$  be the equilibrium measure of  $\Sigma \cup E_n$ . Since  $c(\Sigma \cup E_n) = 1$ , we can get a sequence  $\{\alpha_{n,j}\}$  of algebraic integers such that  $\operatorname{Conj}(\alpha_{n,j})$  eventually belongs to any neighborhood of  $\Sigma \cup E_n$ . By Lemma 2.3,

$$\mu_{n,j} := \frac{1}{\deg \alpha_{n,j}} \sum_{x \in \operatorname{Conj}(\alpha_{n,j})} \delta_x \longrightarrow \nu_n \quad \text{as} \quad j \longrightarrow \infty.$$
(4.1)

Fix a neighborhood U of  $\Sigma$ . We want to determine what fraction of  $\operatorname{Conj}(\alpha_{n,j})$  is in U for sufficiently large j; thus we want to determine the value of  $\nu_n(\Sigma)$ . To do this, we solve a Dirichlet problem on the domain  $D_n$  in the Riemann sphere  $\widehat{\mathbb{C}}$ , where  $D_n$  is the component of  $\widehat{\mathbb{C}} \setminus (\Sigma \cup E_n)$  containing  $\infty$ .

Let  $\phi$  be the function on the boundary  $\partial D_n$  of  $D_n$  such that  $\phi \equiv 0$  on the exterior boundary of  $\Sigma$  and  $\phi \equiv 1$  on the exterior boundary of  $E_n$ . Since  $\phi$  is continuous on  $\partial D_n$ , there exists a unique bounded harmonic function f on  $D_n$  such that  $\lim_{z\to\xi} f(z) = \phi(\xi)$ for q.e.  $\xi \in \partial D_n$ . Then, by definition,  $f(\infty) = \nu_n(E_n)$ . We show

$$f(z) = \frac{g_{\Sigma}(z, \infty) - g_{\Sigma \cup E_n}(z, \infty)}{n}$$

Observe that, since  $g_{\Sigma}(z, \infty) \equiv n$  on  $E_n$ ,  $\lim_{z \to \xi} f(z) = \phi(\xi)$  for q.e.  $\xi \in \partial D_n$ . The function f is harmonic and bounded on  $D_n$  with

$$f(\infty) = \lim_{z \to \infty} \frac{g_{\Sigma}(z, \infty) - g_{\Sigma \cup E_n}(z, \infty)}{n}$$
$$= -\frac{1}{n} \log c(\Sigma).$$

Thus the function f defined above solves the Dirichlet problem for  $\phi$ . Therefore, we have  $\nu_n(\Sigma) = 1 + \frac{1}{n} \log c(\Sigma)$ .

Now, owing to the above observations, there exist  $j' \in \mathbb{N}$  such that  $\alpha_{n,j'} =: \alpha_n$  satisfies

$$\frac{\sharp\{\operatorname{Conj}(\alpha_n) \cap U\}}{\deg \alpha_n} \ge 1 + \frac{1}{n}\log c(\Sigma) - \frac{1}{n}.$$
(4.2)

Note that, since  $c(\Sigma) > 0$ , the right hand side of (4.2) converges to 1 as  $n \to \infty$ . Since n is arbitrary, we get a sequence  $\{\alpha_n\}$  of algebraic integers that satisfy (1.2).

We shall now construct a sequence  $\{\alpha_n\}$  of distinct algebraic integers that satisfy (1.2) for any neighborhood U of  $\Sigma$  and  $h_{\Sigma}(\alpha_n) \to -\log c(\Sigma)$  as  $n \to \infty$ . Let  $\{\alpha_{n,j}\}$  be a sequence of algebraic integers as above. There exists  $J \in \mathbb{N}$  such that for all  $j \geq J$ , we get  $g_{\Sigma}(z, \infty) \geq n - \frac{1}{n^2}$  for at least deg  $\alpha_{n,j} \cdot \left(\nu_n(E_n) - \frac{1}{n^2}\right)$  many  $z \in \operatorname{Conj}(\alpha_{n,j})$ .

Therefore, for all  $j \geq J$ , we get

$$h_{\Sigma}(\alpha_{n,j}) \ge \left(n - \frac{1}{n^2}\right) \left(-\frac{1}{n}\log c(\Sigma) - \frac{1}{n^2}\right).$$

$$(4.3)$$

Let r > 0 be such that  $\operatorname{Conj}(\alpha_{n,j}) \subset \mathbb{D}(0;r)$  for all j. Since  $g_{\Sigma}(z,\infty)$  is an upper semicontinuous function, by (4.1), we get

$$\limsup_{j \to \infty} h_{\Sigma}(\alpha_{n,j}) = \limsup_{j \to \infty} \int_{\{|z| \le r\}} g_{\Sigma}(z,\infty) \, d\mu_{n,j}(z) \le \int_{\{|z| \le r\}} g_{\Sigma}(z,\infty) \, d\nu_n(z).$$
(4.4)

Using the definition of  $g_{\Sigma}(z, \infty)$  and the fact that  $U_{\nu_n}(z) = 0$  q.e. on  $\Sigma \cup E_n$ , we get

$$\int g_{\Sigma}(z,\infty) d\nu_n(z) = \int (U_{\mu_{\Sigma}}(z) - \log c(\Sigma)) d\nu_n(z) = -\log c(\Sigma) + \int U_{\nu_n}(z) d\mu_{\Sigma}(z)$$
$$= -\log c(\Sigma). \tag{4.5}$$

By (4.3), (4.4) and (4.5), there exist  $j' \in \mathbb{N}$  such that  $\alpha_{n,j'}$  satisfies

$$|h_{\Sigma}(\alpha_{n,j'}) + \log c(\Sigma)| < \frac{1}{n}$$

and

$$\frac{\sharp\{\operatorname{Conj}(\alpha_{n,j'}) \cap U_n\}}{\deg \alpha_{n,j'}} \ge 1 + \frac{1}{n} \log c(\Sigma) - \frac{1}{n},$$

where  $U_n := \{z \in \mathbb{C} : \operatorname{dist}(x, z) < \frac{1}{n} \text{ for some } x \in \Sigma\}$ . Since *n* is arbitrary, we get a sequence  $\{\alpha_n := \alpha_{n,j'}\}$  of algebraic integers that satisfy the required properties. This finishes the proof of Theorem 1.6.

Remark 4.1. We can show that the sequence of measures  $\{\nu_n\}$  constructed above converge in the weak<sup>\*</sup> topology to the equilibrium measure  $\mu_{\Sigma}$  (and hence so does the appropriate subsequence of the measures  $\{\mu_{n,j}\}$ ). First, by construction,  $\{\nu_n\}$  is tight and uniformly bounded in the total variation norm, so by Lemma 2.4 there exists a weak<sup>\*</sup> convergent subsequence which we continue to denote by  $\{\nu_n\}$ . Let  $\mu$  be its weak<sup>\*</sup> limit. Writing  $\nu_n := \gamma_n + \zeta_n$  where  $\gamma_n := \nu_n|_{\Sigma}$  and  $\zeta_n := \nu_n|_{E_n}$ , we have  $\gamma_n(\Sigma) = 1 + \frac{1}{n} \log c(\Sigma) \to 1$  so that  $\mu$  is the weak<sup>\*</sup> limit of  $\{\gamma_n\}$  and  $\mu$  is a probability measure supported in  $\Sigma$ . Since the logarithmic potential  $U_{\mu}$  is lower semicontinuous, we have

$$\liminf_{n \to \infty} U_{\gamma_n}(z_n) \ge U_{\mu}(z) \quad \text{for } z \in \mathbb{C}$$

where  $z_n \to z$  (cf., Theorem I.6.8 of [ST97]). We will show that  $U_{\mu}(z) \leq -\log c(\Sigma)$  for q.e.  $z \in \Sigma$ . For then  $I(\mu) \leq -\log c(\Sigma) = I(\mu_{\Sigma})$  and the result follows.

To this end, since  $\nu_n$  is the equilibrium measure of a set of logarithmic capacity equal to 1, we have

$$U_{\nu_n}(z) = U_{\gamma_n}(z) + U_{\zeta_n}(z) = 0$$
 for q.e.  $z \in \Sigma \cup E_n$ 

In particular, the above holds for q.e.  $z \in \Sigma$ . Thus

$$-\limsup_{n \to \infty} U_{\zeta_n}(z) \ge U_{\mu}(z) \quad \text{for q.e. } z \in \Sigma.$$

16

Now for  $z \in \Sigma$ , since  $E_n \to \infty$  and  $\zeta_n(E_n) \to 0$ ,

$$\limsup_{n \to \infty} U_{\zeta_n}(z) = \limsup_{n \to \infty} \int_{E_n} \log \frac{1}{|z - w|} d\zeta_n(w) = \limsup_{n \to \infty} \int_{E_n} \log \frac{1}{|w|} d\zeta_n(w)$$
$$= \limsup_{n \to \infty} \left( -\int_{E_n} g_{\Sigma}(w, \infty) d\zeta_n(w) \right).$$

Since  $g_{\Sigma}(w, \infty) \equiv n$  on  $E_n$ , we get

$$\int_{E_n} g_{\Sigma}(w,\infty) \, d\zeta_n(w) = n\zeta_n(E_n) = n\left(-\frac{1}{n}\log c(\Sigma)\right) = -\log c(\Sigma).$$

Therefore, we have

$$U_{\mu}(z) \leq -\limsup_{n \to \infty} U_{\zeta_n}(z) = -\log c(\Sigma) \text{ for q.e. } z \in \Sigma.$$

## 5. FURTHER DISCUSSION

There remain several relevant questions related to Theorems 1.2 and 1.6.

- (1) Can the lower bound  $-(1/2) \log c(\Sigma)$  in Theorem 1.2 be improved?
- (2) Does there exist a sequence of distinct algebraic integers  $\{\alpha_n\}$  such that 1.2 holds for every neighborhood U of  $\Sigma$  and  $h_{\Sigma}(\alpha_n) \to L < -\log c(\Sigma)$  as  $n \to \infty$ ?
- (3) Let  $\{\alpha_n\}$  be a sequence of distinct algebraic integers such that  $\{\alpha_n\}$  satisfy (1.2) for any neighborhood U of  $\Sigma$  and  $h_{\Sigma}(\alpha_n) \to -\log c(\Sigma)$  as  $n \to \infty$ . Does the corresponding sequence of probability measures  $\{\mu_n\}$  converge to the equilibrium measure  $\mu_{\Sigma}$ ? If not, what other measures  $\mu$  can be realized in this fashion?

Note that, in the case where  $\Sigma$  is a compact set symmetric about the real axis with  $c(\Sigma) \geq 1$ , Orloski and Sardari [OS23] have recently characterized the so-called *arithmetic* probability measures on  $\Sigma$ , i.e., probability measures  $\mu$  on  $\Sigma$  which are weak\* limits of measures  $\{\mu_n\}$  coming from  $\{\alpha_n\}$  where all all conjugates of  $\alpha_n$  eventually lie inside any fixed neighborhood of  $\Sigma$ . In our setting when  $c(\Sigma) < 1$ , necessarily in the second question above we consider the less restrictive hypothesis (1.2).

### References

- [Ahl10] Lars Ahlfors, Conformal Invariants, AMS Chelsea Publishing, Providence, RI, 2010. xii+162 pp. Topics in geometric function theory. Reprint of the 1973 original. With a foreword by Peter Duren, F. W. Gehring and Brad Osgood.
- [Bil97] Yuri Bilu, Limit distribution of small points on algebraic tori, Duke Math. J. 89 (1997), no. 3, 465–476.
- [BG06] Enrico Bombieri and Walter Gubler, *Heights in Diophantine Geometry*, New Math. Monogr., 4 Cambridge University Press, Cambridge, 2006. xvi+652 pp.
- [CKL98] U. Cegrell, S. Kolodziej and N. Levenberg, Two problems on potential theory for unbounded sets, Math. Scand. 83 (1998), no. 2, 265–276.
- [CLO92] D. Cox, J. Little, D. O'Shea, Ideals, Varieties, and Algorithms, Springer-Verlag, New York, 1992.
- [Fek23] Michael Fekete, Über die Verteilung der Wurzeln bei gewissen algebraischen Gleichungen mit ganzzahligen Koeffizienten, Math. Z. 17 (1923), no. 1, 228–249.

- [FS55] Michael Fekete and Gábor Szegö, On algebraic equations with integral coefficients whose roots belong to a given point set, Math. Z. 63 (1955), 158–172.
- [OS23] Bryce Joseph Orloski and Naser Talebizadeh Sardari, *Limiting distributions of conjugate al*gebraic integers, arXiv reference: arXiv:2302.02872.
- [Pri05] Igor Pritsker, Small polynomials with integer coefficients, J. Anal. Math. 96 (2005), 151–190.
- [Pri11] Igor Pritsker, Distribution of algebraic numbers, J. reine angew. Math. 657 (2011), 57–80.
- [Pri15] Igor Pritsker, Asymptotic distribution and symmetric means of algebraic numbers, Acta Arith. 168 (2015), no. 2, 121–138.
- [Ran95] Thomas Ransford, Potential Theory in the Complex Plane, LMS Student Texts 25, Cambridge University Press, Cambridge (UK), 1995.
- [Rum99] Robert Rumely, On Bilu's equidistribution theorem, Spectral problems in geometry and arithmetic, Contemp. Math. 237, AMS (1999), 159–166.
- [ST97] Edward B. Saff and Vilmos Totik, Logarithmic Potentials with External Fields, Springer-Verlag, Berlin, 1997.
- [Sch18] I. Schur, Uber die Verteilung der Wurzeln bei gewissen algebraischen Gleichungen mit ganzzahligen Koeffizienten, Math. Z. 1 (1918), 377–402.
- [Ser19] Jean-Pierre Serre, Distribution asymptotique des valeurs propres des endomorphismes de Frobenius, Astérisque No. 414, Séminaire Bourbaki, Vol. 2017/2018. Exposés 1136–1150 (2019).
- [Smi21] Alexander Smith, Algebraic integers with conjugates in a prescribed distribution, to appear in Annals of Mathematics, arXiv reference: arXiv:2111.12660.

DEPARTMENT OF MATHEMATICS, INDIANA UNIVERSITY, BLOOMINGTON, INDIANA 47405, USA *Email address:* nlevenbe@indiana.edu, mmlondhe@iu.edu