A Novel Model-Independent Approach to Explore New Physics in Five-body Semileptonic Decays

Yao Yu,^{1,2,*} Han Zhang,^{3,†} and Bai-Cian Ke^{3,‡}

¹Chongqing University of Posts & Telecommunications, Chongqing, 400065, China

²Department of Physics and Chongqing Key Laboratory for Strongly Coupled Physics,

Chongqing University, Chongqing 401331, People's Republic of China

³School of Physics and Laboratory of Zhongyuan Light,

Zhengzhou University, Zhengzhou, Henan 450001, China

Abstract

We derive three parameters associated with the angular distribution of semileptonic five-body decays $D_{(s)}/B_{(s)} \rightarrow V\ell^+\nu_\ell$ ($\ell = e, \mu, \tau$) with $V \rightarrow PPP$, where V and P denote vector and pseudoscalar particles. These parameters, expected to be unity in the Standard Model, may deviate if new physics is involved. Our model-independent approach involves deriving the specific form of the angular distribution under the most general form of the decay matrix element $D_{(s)}/B_{(s)} \rightarrow V$. The outcomes have potential applications in precisely testing the Standard Model and exploring new physics. Relevant measurements can be carried out using data obtained from BESIII, Belle II, and LHCb.

^{*} Corresponding author: yuyao@cqupt.edu.cn

 $^{^\}dagger$ Corresponding author: zhanghanzzu@gs.zzu.edu.cn

[‡] Corresponding author: baiciank@ihep.ac.cn

I. INTRODUCTION

The hadronic and weak currents in semileptonic decays can be effectively disentangled, providing a clear environment for investigating the strong and weak interactions. With minimal impact from new physics interactions in the lepton pair (the weak current), semileptonic decays offer an effective method to explore new physics contributions to the hadronic form factors (the hadronic current). Furthermore, in the Standard Model (SM), left-handed chiral currents are exclusively involved, while in the context of new physics, right-handed currents may be present. The influence of these additional currents can be detected in the angular distribution of semileptonic decays.

Nowadays, extensive experimental studies of three- and four-body semileptonic decays of charmed mesons have been conducted [1]. The relevant hadronic form factors of D^1 transitioning to final-state hadronic are measured to test the SM by comparing the measured results with predictions by theoretical models, such as lattice QCD [2], the covariant confined quark model (CCQM) [3], the holographic QCD model [4], the QCD sum rule model [5], etc. However, none of these provide strong constraints on new physics.

In this study, we initially present the angular distribution form for five-body decays $D \rightarrow V \ell^+ \nu_\ell \ (\ell = e, \mu)$ with $V \rightarrow PPP$, where the decay matrix element takes the most general form. Five-body decays offer the advantage of introducing additional degrees of freedom compared to three- and four-body decays [6–8]. Subsequently, we propose three asymmetry parameters associated with the angular distributions by incorporating various forms of currents outside the framework of the SM, other than the left-handed current. These parameters can serve as sensitive measurement tools for searching new physics, without being reliant on any particular new physics model. Finally, a succinct summary will be presented.

II. DECAY RATE FORMALISM

The decay rate of a $D \to V \ell^+ \nu_\ell, V \to P_1 P_2 P_3$ decay can be parameterized using the following eight independent variables [9] as illustrated in Fig. 1: $Y^2/K^2/Z^2$ is the mass squared of the $\ell^+ \nu_\ell/P_1 P_2 P_3/P_2 P_3$ system; $\theta_1/\theta_2/\theta_3$ is the angle between the three-momenta

¹ We utilize the decay process of the D meson as a specific example to showcase the derivation of the pertinent equations. The calculation methodology remains identical for the D_s , B, and B_s mesons, with the only difference being the substitution of the corresponding physical parameters.



FIG. 1. Definition of angles in the cascade decay $D \to V\ell^+\nu_\ell, V \to P_1P_2P_3$. θ_1 (θ_2) is the angle between the three-momenta of ℓ^+ (P_1) and D in the $\ell^+\nu_\ell$ (V) rest frame; θ_3 is the angle between the three-momenta of P_3 and V in the P_2P_3 rest frame; ϕ_1 (ϕ_3) is the angle between the $\ell\nu_\ell$ (P_2P_3) and the VP_1 systems.

of $\ell^+/P_1/P_3$ in the $\ell^+\nu_\ell/V/P_2P_3$ rest frame and the line of the flight of the $\ell^+\nu_\ell/V/P_2P_3$ system in the D/D/V rest frame; ϕ_1/ϕ_3 is the angle between the planes formed in the D/Vrest frame by the momenta of the $\ell^+\nu_\ell/P_2P_3$ pair and the VP_1/DP_1 pair. The sense of ϕ_1/ϕ_3 is from the VP_1/DP_1 plane to the $\ell^+\nu_\ell/P_2P_3$ plane. The four-momenta of ℓ^+ , ν_ℓ , P_1 , P_2 , P_3 are denotes as q_ℓ , q_ν , p_1 , p_2 , and p_3 , and their masses as m_ℓ , m_ν , m_1 , m_2 , and m_3 , respectively. For convenience, we define $Q \equiv q_\ell + q_\nu + p_1 + p_2 + p_3$, $Y \equiv q_\ell + q_\nu$, $K \equiv p_1 + p_2 + p_3$, and $Z \equiv p_2 + p_3$, and denote the masses of D and V as m_A and m_V , respectively.

A. Amplitude

Given the assumption that neutrinos are left-handed, the most general effective Hamiltonian describing the $c \rightarrow de^+\nu^-$ decays, containing all possible parity-conserving four-fermion dimension-6 operators, can be written as [10, 11]

$$\mathcal{H}_{\mathcal{EFF}} = \frac{G_F}{\sqrt{2}} V_{cq} \left[(1+g_V) \bar{c} \gamma_\mu d\bar{u}(q_\ell) \gamma^\mu (1-\gamma_5) v(q_\nu) + (-1+g_A) \bar{c} \gamma_\mu \gamma_5 d\bar{u}(q_\ell) \gamma^\mu (1-\gamma_5) v(q_\nu) \right. \\ \left. + g_S \bar{c} d\bar{u}(q_\ell) (1-\gamma_5) v(q_\nu) + g_P \bar{c} \gamma_5 d\bar{u}(q_\ell) (1-\gamma_5) v(q_\nu) \right. \\ \left. + g_T \bar{c} \sigma_{\mu\nu} d\bar{u}(q_\ell) \sigma^{\mu\nu} (1-\gamma_5) v(q_\nu) + g_{T5} \bar{c} \sigma_{\mu\nu} \gamma_5 d\bar{u}(q_\ell) \sigma^{\mu\nu} (1-\gamma_5) v(q_\nu) \right] , \tag{1}$$

where G_F represents the Fermi constant and V_{cq} denotes the Cabibbo-Kobayashi-Maskawa matrix element. The coefficients g_V , g_A , g_S , g_P , g_T , and g_{T5} represent the modification from vector, axial-vector, scalar, pseudoscalar, tensor, axial-tensor currents to the SM, respectively, and $g_T = -g_{T5}$ due to the left-handed nature of neutrinos. It is important to note that g_V , g_A , g_S , g_P , g_T , and g_{T5} are zero in the SM. The leptonic part is written as

$$\mathcal{L}_{\lambda_W}^{\lambda_l} = \epsilon_{Y,\mu}(\lambda_W)\bar{u}_{\lambda_l}(q_l)\gamma^{\mu}(1-\gamma_5)v(q_{\nu}),$$

$$\mathcal{L}_{sp}^{\lambda_l} = \bar{u}_{\lambda_l}(q_l)(1-\gamma_5)v(q_{\nu}),$$

$$\mathcal{L}_{\lambda_1\lambda_2}^{\lambda_l} = -\mathcal{L}_{\lambda_2\lambda_1}^{\lambda_l} = -i\epsilon_{Y,\mu}(\lambda_1)\epsilon_{Y,\nu}(\lambda_2)\bar{u}_{\lambda_l}(q_l)\sigma^{\mu\nu}(1-\gamma_5)v(q_{\nu})$$
(2)

with the non-vanishing psuedo-scalar and vector type amplitudes [10]

$$\mathcal{L}^{+}_{+} = -\sqrt{2}m_{l}\beta_{l}\sin\theta_{1}e^{-2i\phi_{1}}, \\
\mathcal{L}^{+}_{-} = \sqrt{2}m_{l}\beta_{l}\cos\theta_{1}e^{-i\phi_{1}}, \\
\mathcal{L}^{+}_{0} = 2m_{l}\beta_{l}\cos\theta_{1}e^{-i\phi_{1}}, \\
\mathcal{L}^{-}_{\pm} = \sqrt{2Y^{2}}\beta_{l}(1\pm\cos\theta_{1})e^{\mp i\phi_{1}}, \\
\mathcal{L}^{-}_{0} = 2\sqrt{Y^{2}}\beta_{l}\sin\theta_{1}, \\
\mathcal{L}^{+}_{sp} = -2\sqrt{Y^{2}}\beta_{l}e^{-i\phi_{1}}.$$
(3)

and the non-vanishing tensor type amplitudes

$$\mathcal{L}_{+0}^{+} = -\sqrt{2Y^{2}}\beta_{l}\sin\theta_{1}e^{-2i\phi_{1}}, \\ \mathcal{L}_{-0}^{+} = -\mathcal{L}_{0t}^{+} = 2\sqrt{Y^{2}}\beta_{l}\cos\theta_{1}e^{-i\phi_{1}}, \\ \mathcal{L}_{+t}^{+} = -\mathcal{L}_{0t}^{+} = 2\sqrt{Y^{2}}\beta_{l}\cos\theta_{1}e^{-i\phi_{1}}, \\ \mathcal{L}_{-t}^{+} = -\sqrt{2Y^{2}}\beta_{l}\sin\theta_{1}, \\ \mathcal{L}_{\pm0}^{-} = \pm\sqrt{2}m_{l}\beta_{l}(1\pm\cos\theta_{1})e^{\mp i\phi_{1}}, \\ \mathcal{L}_{+-}^{-} = -\mathcal{L}_{0t}^{-} = 2m_{l}\beta_{l}\sin\theta_{1}, \\ \mathcal{L}_{\pm t}^{-} = -\sqrt{2}m_{l}\beta_{l}(1\pm\cos\theta_{1})e^{\mp i\phi_{1}}.$$
(4)

Here, $\beta_l = \sqrt{1 - \frac{m_l^2}{Y^2}}$. The hadronic part is given by

$$\mathcal{H}_{V(A),\lambda_{W}}^{\lambda_{m}} = \epsilon_{Q,\mu}^{\dagger}(\lambda_{W})\langle V(\lambda_{m}) | V^{\mu}(A^{\mu}) | D \rangle , ,$$

$$\mathcal{H}_{S(P)}^{\lambda_{m}} = \langle V(\lambda_{m}) | S(P) | D \rangle ,$$

$$\mathcal{H}_{T(T5),\lambda_{1}\lambda_{2}}^{\lambda_{m}} = i\epsilon_{Q,\mu}^{\dagger}(\lambda_{1})\epsilon_{Q,\nu}^{\dagger}(\lambda_{2})\langle V(\lambda_{m}) | \sigma^{\mu\nu}(\gamma_{5}) | D \rangle ,$$
(5)

with the non-vanishing terms [10, 11]

$$\begin{aligned}
\mathcal{H}_{V,\pm}^{\pm} &= \mp 2 \frac{m_A}{m_A + m_v} |\vec{K}| V_0, \mathcal{H}_{A,\pm}^{\pm} = -(m_A + m_v) A_1, \\
\mathcal{H}_{A,0}^0 &= 8 \frac{m_A m_v}{\sqrt{Y^2}} A_{12}, \mathcal{H}_{A,t}^0 = 2 \frac{m_A}{\sqrt{Y^2}} |\vec{K}| A_0, \\
\mathcal{H}_P^0 &= -2 \frac{m_A}{m_c + m_d} |\vec{K}| A_0, \\
\mathcal{H}_{T,+-}^0 &= -\mathcal{H}_{T5,0t}^0 = \frac{1}{2m_V} [(-m_A^2 + 3m_V^2 - Y^2) T_2 + 4 \frac{m_A^2}{m_A^2 - m_V^2} |\vec{K}|^2 T_3], \\
\mathcal{H}_{T,\pm0}^{\pm} &= \pm \mathcal{H}_{T,\pm t}^{\pm} = \pm \frac{m_A^2 - m_V^2}{\sqrt{Y^2}} T_2, \\
\mathcal{H}_{T5,\pm 0}^{\pm} &= \pm \mathcal{H}_{T5,\pm t}^{\pm} = -2 \frac{m_A}{\sqrt{Y^2}} |\vec{K}| T_1, \end{aligned}$$
(6)

with

$$A_{12} = \frac{1}{16m_A m_{\omega}^2} (m_A^2 - m_{\omega}^2 - Y^2) (m_A + m_{\omega}) A_1 - 4 \frac{m_A^2}{m_A + m_{\omega}} |\vec{K}|^2 A_2$$
(7)

where $V_0 A_{0,1,2}$, and, $T_{1,2,3}$ are the form factors of the vector meson.

Finally, the amplitude of $V \to P_1 P_2 P_3$ are given by [9]

$$\mathcal{A}(V \to P_1 P_2 P_3) = g \epsilon_{\mu\nu\alpha\beta} \epsilon_V^{\mu} p_1^{\nu} p_2^{\alpha} p_3^{\beta} = -\frac{1}{2} g \epsilon_{\mu\nu\alpha\beta} \epsilon_V^{\mu} K^{\nu} Z^{\alpha} p_{23}^{\beta}, \qquad (8)$$

with $p_{23} = p_2 - p_3$. The coupling incorporates information about the decay structure of V and can always be expressed as a function of Z^2 and θ_3 , specifically $g = g(Z^2, \theta_3)$. For the specific case of $\omega \to \pi^+ \pi^- \pi^0$,

$$g = g_{3\pi} + g_{\rho\pi}g_{\rho\to\pi\pi} (D^{-1}[\rho^0, p_{\pi^+} + p_{\pi^-}] + D^{-1}[\rho^+, p_{\pi^0} + p_{\pi^+}] + D^{-1}[\rho^-, p_{\pi^0} + p_{\pi^-}]), \quad (9)$$

where $D[\rho, P] = P^2 - m_{\rho}^2 + im_{\rho}\Gamma_{\rho}$. It is known that $(p_{\pi^+} + p_{\pi^-})^2$, $(p_{\pi^0} + p_{\pi^+})^2$ and $(p_{\pi^0} + p_{\pi^-})^2$ can be expressed as functions of Z^2 and θ_3 [12].

B. Decay rate

Combining the element of five-body phase space [9] and collecting amplitudes in Eqs. (2, 3, 4, 5, 6), one have the decay rate of $D \to V \ell^+ \nu_\ell, V \to P_1 P_2 P_3$ as

$$\Gamma_{5} = \frac{1}{2^{17}\pi^{9}m_{A}^{2}} \int \frac{|\vec{q_{1}}||\vec{q_{4}}||\vec{Z}||\vec{K}|}{\sqrt{K^{2}}\sqrt{Y^{2}}\sqrt{Z^{2}}} \sum_{\lambda_{l}} |\mathcal{A}_{p}^{\lambda_{l}}|^{2} d\Omega_{1} d\Omega_{3} d\cos\theta_{2} dK^{2} dY^{2} dZ^{2}.$$
(10)

Here, $\mathcal{A}_p^{\lambda_l}$ are expressed in terms of the form factors of the vector meson [9]:

$$\mathcal{A}_{p}^{+} = -\frac{g}{\sqrt{2}} |\vec{Z}| |\vec{q}_{4}| |\sqrt{K^{2}}| \sin \theta_{3} \frac{G_{F}}{\sqrt{2}} V_{cd}(F_{1}\alpha_{1} + F_{2}\alpha_{2} + F_{3}\alpha_{3} + F_{4}\alpha_{4}),$$

$$\mathcal{A}_{p}^{-} = -\frac{g}{\sqrt{2}} |\vec{Z}| |\vec{q}_{4}| |\sqrt{K^{2}}| \sin \theta_{3} \frac{G_{F}}{\sqrt{2}} V_{cd}(F_{1}'\beta_{1} + F_{2}'\beta_{2} + F_{3}'\beta_{3}).$$
(11)

Here, $F^{(\prime)}$ s are given by

$$F_{1} = -\frac{m_{l}}{\sqrt{Y^{2}}}f_{1} - \delta_{1}, F_{2} = -\frac{m_{2}}{\sqrt{Y^{2}}}f_{2} - \delta_{2},$$

$$F_{3} = \frac{m_{2}}{\sqrt{Y^{2}}}f_{3} + \delta_{3}, F_{4} = \frac{m_{2}}{\sqrt{Y^{2}}}f_{4},$$

$$F_{1}' = f_{1} + \frac{m_{l}}{\sqrt{Y^{2}}}\delta_{1}, F_{2}' = f_{2} + \frac{m_{l}}{\sqrt{Y^{2}}}\delta_{2}, F_{3}' = f_{3} + \frac{m_{l}}{\sqrt{Y^{2}}}\delta_{3},$$
(12)

with

$$f_{1} = -2\sqrt{2}\beta_{l}\frac{m_{A}}{m_{A}+m_{\omega}}|\vec{K}|(g_{V}+1)V_{0}, f_{2} = \sqrt{2}\beta_{l}(m_{A}+m_{\omega})(1-g_{A})A_{1},$$

$$f_{3} = 2\sqrt{2}\beta_{l}(1-g_{A})\mathcal{H}_{A,0}^{0}, f_{4} = 4\sqrt{2}\beta_{l}m_{A}|\vec{K}|A_{0}[(1-g_{A})+\frac{Y^{2}}{m_{l}(m_{c}+m_{d})}g_{P}],$$

$$\delta_{1} = 8\sqrt{2}\beta_{l}m_{A}|\vec{K}|g_{T5}T_{1}, \delta_{2} = -4\sqrt{2}\beta_{l}(m_{A}^{2}-m_{\omega}^{2})g_{T}T_{2},$$

$$\delta_{3} = 4\sqrt{2}\beta_{l}\sqrt{Y^{2}}(g_{T}-g_{5})H_{T,+-}^{0},$$
(13)

and

$$\begin{aligned}
\alpha_{1} &= i \sin \theta_{1} e^{-2i\phi_{1}} (\cos \phi_{3} + i \cos \theta_{2} \sin \phi_{3}) + i \sin \theta_{1} (\cos \phi_{3} - i \cos \theta_{2} \sin \phi_{3}), \\
\alpha_{2} &= i \sin \theta_{1} e^{-2i\phi_{1}} (\cos \phi_{3} + i \cos \theta_{2} \sin \phi_{3}) - i \sin \theta_{1} (\cos \phi_{3} - i \cos \theta_{2} \sin \phi_{3}), \\
\alpha_{3} &= \cos \theta_{1} e^{-i\phi_{1}} \sin \theta_{2} \sin \phi_{3}, \\
\alpha_{4} &= e^{-i\phi_{1}} \sin \theta_{2} \sin \phi_{3}, \\
\beta_{1} &= i (1 + \cos \theta_{1}) e^{-i\phi_{1}} (\cos \phi_{3} + i \cos \theta_{2} \sin \phi_{3}) - i (1 - \cos \theta_{1}) (\cos \phi_{3} - i \cos \theta_{2} \sin \phi_{3}), \\
\beta_{2} &= i (1 + \cos \theta_{1}) e^{i\phi_{1}} (\cos \phi_{3} + i \cos \theta_{2} \sin \phi_{3}) + i (1 - \cos \theta_{1}) (\cos \phi_{3} - i \cos \theta_{2} \sin \phi_{3}), \\
\beta_{3} &= \sin \theta_{1} \sin \theta_{2} \sin \phi_{3}.
\end{aligned}$$
(14)

With the additional degrees of freedom, the following unique asymmetries for five-body decays are considered:

$$\begin{split} \Upsilon_{1} &= \int_{S_{1}} d\phi_{3} \int_{S_{1}} d\phi_{1} \int_{D_{1}} d\cos\theta_{2} \int_{D_{1}} d\cos\theta_{1} d\Phi_{5} \\ &\propto \frac{16\pi^{2}}{9} (8F_{1}^{2} + 8F_{2}^{2} + 2F_{3}^{2} + 3F_{2}^{4} + 16F_{1}^{\prime 2} + 16F_{2}^{\prime 2} + 2F_{3}^{\prime 2}) \\ \Upsilon_{2} &= \int_{S_{3}} d\phi_{3} \int_{S_{2}} d\phi_{1} \int_{D_{1}} d\cos\theta_{2} \int_{D_{1}} d\cos\theta_{1} d\Phi_{5} \propto 8\pi^{2} (-F_{1}^{\prime}F_{3}^{\prime} + F_{2}F_{4}) \\ \Upsilon_{3} &= \int_{S_{3}} d\phi_{3} \int_{S_{3}} d\phi_{1} \int_{D_{2}} d\cos\theta_{2} \int_{D_{1}} d\cos\theta_{1} d\Phi_{5} \propto \frac{128}{3} (F_{1}^{2} - F_{2}^{2} + F_{1}^{\prime 2} - F_{2}^{\prime 2}) \\ \Upsilon_{4} &= \int_{S_{1}} d\phi_{3} \int_{S_{4}} d\phi_{1} \int_{D_{2}} d\cos\theta_{2} \int_{D_{1}} d\cos\theta_{1} d\Phi_{5} \propto -\frac{16\pi^{2}}{3} (F_{1}^{\prime}F_{3}^{\prime} + F_{2}F_{4}) \\ \Upsilon_{5} &= \int_{S_{2}} d\phi_{3} \int_{S_{1}} d\phi_{1} \int_{D_{1}} d\cos\theta_{2} \int_{D_{1}} d\cos\theta_{1} d\Phi_{5} \propto \frac{16\pi^{2}}{3} (8F_{1}^{\prime}F_{2}^{\prime} + F_{3}F_{4}) \\ \Upsilon_{6} &= \int_{S_{3}} d\phi_{3} \int_{S_{2}} d\phi_{1} \int_{D_{1}} d\cos\theta_{2} \int_{D_{2}} d\cos\theta_{1} d\Phi_{5} \propto \frac{32\pi}{3} (F_{2}F_{3} - F_{2}^{\prime}F_{3}^{\prime}) \\ \Upsilon_{7} &= \int_{S_{1}} d\phi_{3} \int_{S_{4}} d\phi_{1} \int_{D_{2}} d\cos\theta_{2} \int_{D_{2}} d\cos\theta_{1} d\Phi_{5} \propto -\frac{64\pi}{3} (F_{2}F_{3} + F_{2}F_{3}^{\prime}) \end{split}$$
(15)

where $\int_{D_{1(2)}} \equiv \int_{-1}^{0} \pm \int_{0}^{1}$, $\int_{S_{1(2)}} \equiv \int_{0}^{\pi} \pm \int_{\pi}^{2\pi}$, $\int_{S_{3(4)}} \equiv \int_{0}^{\pi/2} - \int_{\pi/2}^{\pi} \pm \int_{\pi}^{3\pi/2} \mp \int_{3\pi/2}^{2\pi}$, and $d\Phi_5 \equiv \frac{d\Gamma_5}{d\Omega_1 d\phi_3 d\cos\theta_2 dY^2}$.

Although these seven parameters $\Upsilon_{1,2,3,4,5,6,7}$ are independent on each other, there are only four independent form factors in the SM. Therefore, we introduce the following three parameters to describe the relationships among $\Upsilon_{1,2,3,4,5,6,7}$, which are constants within the framework of the SM.

$$\begin{aligned} \Theta_{1} &= \frac{2}{3} \frac{Y^{2} - m_{l}^{2}}{Y^{2} + m_{l}^{2}} \frac{\Upsilon_{6}}{\Upsilon_{7}} \\ \Theta_{2} &= \frac{\Delta}{\Upsilon_{3}} \left[(1 + \frac{m_{l}^{2}}{Y^{2}})^{2} \frac{2\Upsilon_{2} + 3\Upsilon_{4}}{18\pi^{3}\Upsilon_{6}} - \frac{\Upsilon_{6}}{2\pi(2\Upsilon_{2} + 3\Upsilon_{4})} \right] \\ \Theta_{3} &= \frac{1}{\Upsilon_{1}} \left\{ \frac{3\Upsilon_{4} + 2\Upsilon_{2}}{\Delta} \left[\frac{4\pi}{3} \frac{m_{l}^{2} + 2Y^{2}}{m_{l}^{2} + Y^{2}} \Upsilon_{6} + \frac{4}{9\pi} \frac{m_{l}^{2} + Y^{2}}{m_{l}^{2}} \frac{(2\Upsilon_{2} - 3\Upsilon_{4})^{2}}{\Upsilon_{6}} \right] \\ &+ \Delta \left(\frac{\pi}{6} \frac{m_{l}^{2} + 2Y^{2}}{m_{l}^{2} + Y^{2}} \frac{\Upsilon_{6}}{2\Upsilon_{2} + 3\Upsilon_{4}} + \frac{1}{54\pi} \frac{m_{l}^{2} + Y^{2}}{Y^{2}} \frac{m_{l}^{2} + 2Y^{2}}{Y^{2}} \frac{2\Upsilon_{2} + 3\Upsilon_{4}}{\Upsilon_{6}} \right) \right\}$$
(16)

where

$$\Delta = \begin{cases} 3\Upsilon_5 + \sqrt{-32\Upsilon_2^2 + 72\Upsilon_4^2 + 9\Upsilon_5^2}, & \text{if } Y^2 V_0 A_1 + 4m_A m_\omega \frac{m_1^2}{Y^2} A_0 A_{12} < 0\\ 3\Upsilon_5 - \sqrt{-32\Upsilon_2^2 + 72\Upsilon_4^2 + 9\Upsilon_5^2}, & \text{if } Y^2 V_0 A_1 + 4m_A m_\omega \frac{m_1^2}{Y^2} A_0 A_{12} > 0 \end{cases}$$

In the SM, Θ_1 , Θ_2 , and Θ_3 are strictly equal to unity for any Y^2 , but with effects from new physics, they may significantly deviate from unity for certain Y^2 . For convenience, we further assume that $\delta_{1(2,3)}$ is a small quantity and derive the first-order approximation of $\Theta_{1(2,3)}$:

$$\Theta_{1} \approx 1 + \frac{2t}{1+t^{2}} \left(\frac{\delta_{2}}{f_{2}} + \frac{\delta_{3}}{f_{3}}\right),
\Theta_{2} \approx 1 + \frac{t(1-t^{3})}{1+t^{2}} \left(\frac{\delta_{1}}{f_{1}} \frac{2f_{1}^{2}}{f_{1}^{2} - f_{2}^{2}} + \frac{\delta_{2}}{f_{2}} \frac{8f_{1}f_{2} + f_{3}f_{4}}{8f_{1}f_{2} - t^{2}f_{3}f_{4}} + \frac{\delta_{3}}{f_{3}}\varsigma\right),
\Theta_{3} \approx 1 + \frac{t(1-t^{2})}{(2+t^{2})(8f_{1}^{2} + 8f_{2}^{2} + f_{3}^{3}) + 3t^{2}f_{3}} \left[\frac{\delta_{1}}{f_{1}} 16f_{1}^{2} + \frac{\delta_{2}}{f_{2}} \varpi_{1} + \frac{\delta_{3}}{f_{3}} \varpi_{2}\right],$$
(17)

with

$$t = \frac{m_l}{\sqrt{Y^2}}, \varsigma = \frac{8f_1^3f_2 + 8f_1f_2^3 + f_2^2f_3f_4 - f_1^2f_3f_4(1+2t^2)}{(f_1^2 - f_2^2)(8f_1f_2 - t^2f_3f_4)},$$

$$\varpi_1 = -\frac{24f_1f_2f_4^2 - (8f_1f_2 + 2f_3f_4)(8f_1^2 + 8f_2^2 - f_3^2)}{8f_1f_2 - t^2f_3f_4} + \frac{8f_1^2 - 8f_2^2 - f_3^2 - 3f_4^2}{1+t^2},$$

$$\varpi_2 = \frac{24f_1f_2f_4^2 - (8f_1f_2 + 2f_3f_4)(8f_1^2 + 8f_2^2 - f_3^2)}{8f_1f_2 - t^2f_3f_4} + \frac{8f_1^2 - 8f_2^2 - f_3^2 - 3f_4^2}{1+t^2} + 16f_1^2.(18)$$

III. SUMMARY AND DISCUSSION

Five-body semileptonic decays have additional degrees of freedom compared to three- and four-body semileptonic decays, resulting in unique asymmetry correlations. Based on these correlations, we have introduced three parameters $\Theta_{1,2,3}$, which are strictly equal to one in the SM model for any Y^2 . However, according to Eqs. (13) and (17), these parameters may deviate when (axial-)tension current, i.e. $g_T(g_{T5})$, is present, and the magnitude of deviation is larger with larger $t = \frac{m_l}{\sqrt{Y^2}}$. As a result, experimentally studying five-body semileptonic decays to heavier leptons, such as $B_{(s)} \rightarrow V \tau^+ \nu_{\tau} / V \mu^+ \nu_{\mu}$ and $D_{(s)} \rightarrow V \mu^+ \nu_{\mu}$, could yield larger significance. Our investigation offers a new and model-independent approach to explore new physics and the parameters introduced can be measured with the dataset collected by BESIII [13], Belle II, and LHCb.

ACKNOWLEDGMENTS

YY was supported in part by National Natural Science Foundation of China (NSFC) under Contracts No. 11905023, No. 12047564 and No. 12147102, the Natural Science Foundation of Chongqing (CQCSTC) under Contract No. cstc2020jcyj-msxmX0555 and the Science and Technology Research Program of Chongqing Municipal Education Commission (STRPCMEC) under Contracts No. KJQN202200605 and No. KJQN202200621; HZ and BCK were supported in part by NSFC under Contracts No. 11875054 and No. 12192263 and Joint Large-Scale Scientific Facility Fund of the NSFC and the Chinese Academy of Sciences under Contract No. U2032104.

- B. C. Ke, J. Koponen, H. B. Li and Y. H. Zheng, Ann. Rev. Nucl. Part. Sci. 73, 285-314 (2023) [arXiv:2310.05228 [hep-ex]].
- [2] Y. Aoki *et al.* [Flavour Lattice Averaging Group (FLAG)], Eur. Phys. J. C 82, 869 (2022)
 [arXiv:2111.09849 [hep-lat]].
- [3] N. R. Soni, M. A. Ivanov, J. G. Körner, J. N. Pandya, P. Santorelli and C. T. Tran, Phys. Rev. D 98, no.11, 114031 (2018) [arXiv:1810.11907 [hep-ph]].
- [4] H. A. Ahmed, Y. Chen and M. Huang, Phys. Rev. D 109, no.2, 026008 (2024) [arXiv:2309.06156 [hep-ph]].
- [5] H. J. Tian, H. B. Fu, T. Zhong, X. Luo, D. D. Hu and Y. L. Yang, Phys. Rev. D 108, no.7, 076003 (2023) [arXiv:2306.07595 [hep-ph]].
- [6] R. Kumar, Phys. Rev. 185, 1865-1875 (1969).
- [7] T. Huber, Q. Qin and K. K. Vos, Eur. Phys. J. C 78, no.9, 748 (2018) [arXiv:1806.11521 [hep-ph]].
- [8] Q. Y. Hu, X. Q. Li, X. L. Mu, Y. D. Yang and D. H. Zheng, JHEP 06, 075 (2021)
 [arXiv:2104.04942 [hep-ph]].
- [9] H. Zhang, B. C. Ke, Y. Yu, Z. Xiong and J. W. Zhang, [arXiv:2404.04816 [hep-ph]].
- [10] R. Mandal, C. Murgui, A. Peñuelas and A. Pich, JHEP 08, no.08, 022 (2020) [arXiv:2004.06726 [hep-ph]].
- [11] D. Bečirević, M. Fedele, I. Nišandžić and A. Tayduganov, [arXiv:1907.02257 [hep-ph]].

- [12] D. G. Gudino and G. T. Sanchez, Int. J. Mod. Phys. A 27, 1250101 (2012) [arXiv:1106.1467
 [hep-ph]].
- [13] M. Ablikim et al. (BESIII Collaboration), Chin. Phys. C 44, 040001 (2020).