EXTREMES OF GENERALIZED INVERSIONS AND DESCENTS ON PERMUTATION GROUPS

PHILIP DÖRR

ABSTRACT. The classical inversion statistic on symmetric groups is the sum of all indicators $\mathbf{1}\{\pi(i) > \pi(j)\}$ for a random permutation $\pi = (\pi(1), \ldots, \pi(n))$ and the pairs (i, j) with $1 \le i < j \le n$. The descent statistic counts all $i \in \{1, \ldots, n-1\}$ with $\pi(i) > \pi(i+1)$. The number of inversions can be generalized by restricting the indicators to pairs (i, j) with i < j and $|i - j| \le d$ for some $d \in \{1, \ldots, n-1\}$. Likewise, the number of descents can be generalized by counting all $i \in \{1, \ldots, n-1\}$. Likewise, the number of descents can be generalized by counting all $i \in \{1, \ldots, n-d\}$ with $\pi(i) > \pi(i + d)$. These generalized statistics can be further extended to the signed and even-signed permutation groups. The bandwidth index d can be chosen in dependence of n, and the magnitude of d is significant for asymptotic considerations. It is known that each of these statistics is asymptotically normal for suitable choices of d. In this paper we prove the bivariate asymptotic normality and determine the extreme value asymptotics of generalized inversions and descents.

1. INTRODUCTION

The numbers of inversions and descents are two important quantities of permutations. For a permutation $\pi: \{1, \ldots, n\} \to \{1, \ldots, n\}$, an *inversion* is a pair (i, j) with i < j, but $\pi(i) > \pi(j)$. A *descent* is an index i with $\pi(i) > \pi(i+1)$, i.e., descents correspond to adjacent inversions. We now equip the symmetric group S_n with the discrete uniform probability measure induced by the point masses $\mathbb{P}(\{\pi\}) = 1/n! \ \forall \pi \in S_n$. In this context, we denote the random numbers of inversions and descents as X_{inv} and X_{des} . These random variables can also be represented with help of i.i.d. standard uniform variables $Z_1, Z_2, \ldots, \sim U(0, 1)$, namely,

(1)
$$X_{inv} = \sum_{1 \le i < j \le n} \mathbf{1} \{ Z_i > Z_j \},$$
$$X_{des} = \sum_{i=1}^{n-1} \mathbf{1} \{ Z_i > Z_{i+1} \}.$$

In (1), we have to sum the indicators $\mathbf{1}\{Z_i > Z_j\}$ over all pairs (i, j) with i < j. A class of generalized inversion statistics can be constructed by restricting the sum

Date: April 15, 2024.

²⁰¹⁰ Mathematics Subject Classification. Primary: 60G70, 05A16; Secondary: 20F55. Key words and phrases. extreme values, CLT, permutation statistics, Coxeter group. Philip Däm is supported by the DEC (214828170, CBK 2207, "MathCoRe")

in (1) to pairs (i, j) with $1 \le j - i \le d$, for some $d \in \{1, ..., n - 1\}$. Writing $\mathfrak{N}_{n,d} := \{(i, j) \in \{1, ..., n\}^2 \mid 1 < j - i \le d\}$, we can state that

(2)
$$X_{\text{inv}}^{(d)} := \sum_{(i,j)\in\mathfrak{N}_{n,d}} \mathbf{1}\{Z_i > Z_j\}$$

counts the so-called *d*-inversions. The term generalized inversions is an umbrella term for all *d*-inversions. By choosing d = n - 1 or d = 1, it is seen that this class includes common inversions and descents. Moreover, *d*-descents or generalized descents are given by

(3)
$$X_{\text{des}}^{(d)} := \sum_{i=1}^{n-d} \mathbf{1} \{ Z_i > Z_{i+d} \}.$$

Generalized inversions were first introduced by de Mari & Shayman [5] to solve a problem in algebraic geometry. The first stochastic considerations for the random variables $X_{inv}^{(d)}$ are due to Bona [2] and Pike [10], who computed the mean and variance and proved a central limit theorem (CLT).

Symmetric groups belong to the class of finite irreducible Coxeter groups, which have been classified by [4]. The concept of inversions and descents can be transferred to these groups, see [1, Section 1.4]. Two other important subfamilies of the finite irreducible Coxeter groups are the signed permutation groups B_n and the even-signed permutation groups D_n . The group B_n consists of all maps $\pi: \{1, \ldots, n\} \longrightarrow \{1, \ldots, n\} \cup$ $\{-1, \ldots, -n\}$ for which $|\pi|$ is a permutation. Such a map is called a signed permutation. The groups D_n are the subgroups of B_n containing all signed permutations with an even number of negative signs. The random number of inversions on these groups can be represented with help of i.i.d. variables $Z_1, \ldots, Z_n \sim U(-1, 1)$ as follows:

(4a)
$$X_{inv}^B = \sum_{1 \le i < j \le n} \mathbf{1}\{Z_i > Z_j\} + \sum_{1 \le i < j \le n} \mathbf{1}\{-Z_i > Z_j\} + \sum_{i=1}^n \mathbf{1}\{Z_i < 0\}$$

(4b)
$$X_{inv}^D = \sum_{1 \le i < j \le n} \mathbf{1} \{ Z_i > Z_j \} + \sum_{1 \le i < j \le n} \mathbf{1} \{ -Z_i > Z_j \}.$$

We refer to the three families S_n , B_n , and D_n as the *classical Weyl groups*. Recently, Meier & Stump [9] extended the concept of generalized inversions and descents to the groups B_n and D_n , again proving a CLT.

The investigation of extreme value asymptotics for common inversions and descents was initiated in [7, 6], so we aim to extend this knowledge to generalized inversions. We aim to prove Gumbel attraction for both the stand-alone statistics $X_{\text{inv}}^{(d_1)}, X_{\text{des}}^{(d_2)}$ and the joint statistic $(X_{\text{inv}}^{(d_1)}, X_{\text{des}}^{(d_2)})^{\top}$. Here, d_1 and d_2 can be either fixed or dependent on n. We will investigate the impact of the choice of d for these results.

This paper is structured as follows. Section 2 gives basic properties on generalized inversions and descents, and introduces Hájek projections that are used as independent sum approximations. Section 3 deduces the bivariate CLT and the extreme value

theorems for generalized inversions and descents. Section 4 gives the proofs of two auxiliary lemmas determining the choices of d for which the Hájek projection serves as a working approximation. We use typical Landau notation for positive sequences a_n, b_n as follows:

- a_n = O(b_n) means that lim sup_{n→∞} a_n/b_n < ∞.
 a_n = o(b_n) means that lim_{n→∞} a_n/b_n = 0. This is also written as a_n ≪ b_n or $b_n \gg a_n$.
- $a_n = \Theta(b_n)$ means that a_n and b_n have the same order of magnitude, i.e., both $a_n = O(b_n)$ and $b_n = O(a_n)$ hold.
- $a_n = b_n + o_{\mathbb{P}}(1)$ means that a_n, b_n are sequences of random variables with $a_n - b_n \xrightarrow{\mathbb{P}} 0.$

2. Basic definitions

Definition 2.1. Let S_n be a symmetric group and let $\pi \in S_n$. For $d \in \{1, \ldots, n-1\}$, *d*-inversions are all pairs (i, j) in $\mathfrak{N}_{n,d}$ with $\pi(i) > \pi(j)$. In this sense, common descents equal 1-inversions, and common inversions equal (n-1)-inversions. Moreover, *d*-descents are all numbers $i \in \{1, \ldots, n-d\}$ with $\pi(i) > \pi(i+d)$. We write $X_{inv}^{(d)}$ for the random number of *d*-inversions and $X_{des}^{(d)}$ for the number of *d*-descents. Probabilistic representations are given in (2), (3).¹

Remark 2.2. Obviously, every number $k \in \{1, \ldots, n\}$ can appear in at most 2dd-inversions. This bound is redundant if d > n/2. In fact, it is an important case distinction whether $d \leq n/2$ or d > n/2, e.g., when calculating the mean and variance of $X_{inv}^{(d)}$ and $X_{des}^{(d)}$. In the case of $d \leq n/2$, we can split $\{1, \ldots, n\}$ into the regions

$$K_1 := \{1, \dots, d\}, \qquad K_2 := \{d+1, \dots, n-d\}, \qquad K_3 := \{n-d+1, \dots, n\}.$$

For each $k \in K_2$, all larger indices $k+1, \ldots, k+d$ and all smaller indices $k-1, \ldots, k-d$ allow to form d-inversions. For any $k \notin K_2$, there are less than d indices available in one direction. If $k \in K_1$, then k-1 smaller indices are available for d-inversions, which is less than d numbers. If k > n - d is large, then there are only n - k < d larger indices available.

On the contrary, if d > n/2, then n - d < d and the above partition into three regions is now written as

$$K_1 := \{1, \dots, n-d\}, \qquad K_2 := \{n-d+1, \dots, d\}, \qquad K_3 := \{d+1, \dots, n\}.$$

Now, if $k \in K_2$, then there are only k-1 smaller indices and n-k larger indices available to form d-inversions.

¹In the literature, there are different terminologies, e.g., in [2, 10], d-inversions are called d-descents. However, we use the terms and notation provided in [9].

The mean and variance of $X_{inv}^{(d)}$ have been extensively computed by Pike [10]. It is easy to verify that the special cases d = 1 and d = n - 1 are consistent with [8, Corollaries 3.2 and 4.2]. The proof of [10, Theorem 1] is reviewed in [9, Theorem A.1], where the variance of *d*-descents is provided as well.

Theorem 2.3 (see [10], Theorem 1 and [9], Theorem A.1). For all d = 1, ..., n - 1, it holds that

$$\mathbb{E}\left(X_{\text{inv}}^{(d)}\right) = \frac{2nd - d^2 - d}{4}, \qquad \qquad \mathbb{E}\left(X_{\text{des}}^{(d)}\right) = \frac{n - d}{2}.$$

Moreover, if $d \leq n/2$, then

$$\operatorname{Var}\left(X_{\operatorname{inv}}^{(d)}\right) = \frac{6nd + 4d^3 + 3d^2 - d}{72}, \qquad \operatorname{Var}\left(X_{\operatorname{des}}^{(d)}\right) = \frac{n+d}{12}.$$

If d > n/2, then

$$\operatorname{Var}\left(X_{\mathrm{inv}}^{(d)}\right) = -\frac{1}{6}d^3 + \left(\frac{1}{3}n - \frac{7}{24}\right)d^2 - \left(\frac{1}{6}n^2 - \frac{5}{12}n + \frac{1}{8}\right)d + \frac{1}{36}n^3 - \frac{1}{12}n^2 + \frac{1}{18}n$$

and $\operatorname{Var}\left(X_{\mathrm{des}}^{(d)}\right) = (n-d)/4.$

Meier & Stump [9] introduced an extension of generalized inversions and descents from symmetric groups to the other classical Weyl groups B_n and D_n . This extension is based on the *root poset* of a classical Weyl group. We refer to [9, Section 2] for the details. On the symmetric group S_n , the ordered pairs of indices (i, j) correspond to the positive roots $[ij] := e_i - e_j$, where e_i, e_j are unit vectors in \mathbb{R}^n , and the height of [ij] within the root poset is ht([ij]) = j - i.

On the signed permutation group B_n , we also have to consider the positive roots $[\tilde{i}j] := e_i + e_j$ and $[i] := e_i$ for $1 \leq i < j \leq n$. The heights of these additional roots are $\operatorname{ht}([\tilde{i}j]) = i + j$ and $\operatorname{ht}([i]) = i$. The root $[\tilde{i}j]$ corresponds to the indicator $\mathbf{1}\{-\pi(i) > \pi(j)\}$ appearing in (4a), while [i] corresponds to $\mathbf{1}\{\pi(i) < 0\}$. See [9, Example 2.2] for an illustration of the root poset of B_n . On the even-signed permutation group D_n , the roots [i] are disregarded, and $[\tilde{i}j]$ has height i + j - 2.

Definition 2.4. For any classical Weyl group, *d*-inversions are determined by roots of height at most *d*, and *d*-descents are determined by roots of height exactly *d*, see [9, Definition 2.4]. For symmetric groups, this coincides with Definition 2.7. In addition to $\mathfrak{N}_{n,d}$, we introduce

$$\widetilde{\mathfrak{N}}_{n,d} := \{(i,j) \in \{1,\ldots,n\}^2 \mid i < j, j+i \le d\}.$$

Then, on the signed and even-signed permutation groups, $X_{inv}^{(d)}$ and $X_{des}^{(d)}$ can be expressed as follows:

(5a)
$$X_{\text{inv}}^{B,(d)} = \sum_{(i,j)\in\mathfrak{N}_{n,d}} \mathbf{1}\{Z_i > Z_j\} + \sum_{(i,j)\in\widetilde{\mathfrak{N}}_{n,d}} \mathbf{1}\{-Z_i > Z_j\} + \sum_{i=1}^d \mathbf{1}\{Z_i < 0\},$$

EXTREMES OF GENERALIZED INVERSIONS AND DESCENTS ON PERMUTATION GROUPS 5

(5b)
$$X_{\text{inv}}^{D,(d)} = \sum_{(i,j)\in\mathfrak{N}_{n,d}} \mathbf{1}\{Z_i > Z_j\} + \sum_{(i,j)\in\widetilde{\mathfrak{N}}_{n,d+2}} \mathbf{1}\{-Z_i > Z_j\},$$

(5c)
$$X_{\text{des}}^{B,(d)} = \sum_{i=1}^{n-d} \mathbf{1}\{Z_i > Z_{i+d}\} + \sum_{i=1}^{\lfloor d/2 \rfloor - 1} \mathbf{1}\{-Z_i > Z_{d-i}\} + \mathbf{1}\{\pi(i) > d\},$$

(5d)
$$X_{\text{des}}^{D,(d)} = \sum_{i=1}^{n-d} \mathbf{1}\{Z_i > Z_{i+d}\} + \sum_{i=1}^{\lceil d/2 \rceil} \mathbf{1}\{-Z_i > Z_{d+2-i}\}.$$

Remark 2.5. The largest possible choice of d equals the total height of the root poset, namely, $d_{\max} - 1$, where d_{\max} denotes the largest degree of the underlying classical Weyl group. In particular, $d_{\max} = n$ for S_n , $d_{\max} = 2n$ for B_n and $d_{\max} = 2n - 2$ for D_n . To precisely compute the variance of $X_{inv}^{(d)}$ and $X_{des}^{(d)}$ on the groups B_n and D_n , one needs to distinguish eight cases, as seen in [9, Theorems A.4 and A.13]. However, many of these cases give the same asymptotic quantification, which can be stated as follows:

Lemma 2.6 (cf. [9], Theorems A.4 and A.13). For the generalized inversions and descents on both the groups B_n and D_n , it holds that

$$\operatorname{Var}\left(X_{\mathrm{inv}}^{(d)}\right) = \begin{cases} \frac{1}{36}d^3 + \frac{1}{12}nd + O(d^2), & d \le n/2\\ \frac{1}{36}d^3 + O(d^2), & n/2 \le d < n\\ -\frac{1}{12}d^3 + \frac{1}{3}nd^2 - \frac{1}{3}n^2d + \frac{1}{9}n^3 + O(d^2), & d \ge n \end{cases}$$
$$\operatorname{Var}\left(X_{\mathrm{des}}^{(d)}\right) = \begin{cases} \frac{1}{24}d + \frac{1}{12}n + O(1), & d < n\\ -\frac{1}{8}d + \frac{1}{4}n + O(1), & d \ge n \end{cases}.$$

An evident issue seen in (2), (5a), and (5b) is the mutual dependence of the indicators $1\{Z_i > Z_j\}$ and $1\{-Z_i > Z_j\}$. Therefore, we aim to approximate $X_{inv}^{(d)}$ with a sum of independent variables. Hájek projections are a common tool for this.

Definition 2.7. Let Z_1, \ldots, Z_n be independent random variables, and let X be another random variable. Then, the *Hájek projection* of X with respect to Z_1, \ldots, Z_n is given by

$$\hat{X} := \sum_{k=1}^{n} \mathbb{E}(X \mid Z_k) - (n-1)\mathbb{E}(X).$$

As every $\mathbb{E}(X \mid Z_k)$ is a measurable function only in Z_k , the Hájek projection is a sum of independent random variables. If $(X_n)_{n \in \mathbb{N}}$ is a sequence of random variables with each X_n depending on Z_1, \ldots, Z_n , then the Hájek projections \hat{X}_n should give a sufficiently accurate approximation to X_n . To ensure this, the following criterion is useful.

Theorem 2.8 (cf. [11], Theorem 11.2). Consider a sequence $(X_n)_{n\geq 1}$ of random variables and their associated Hájek projections $(\hat{X}_n)_{n\geq 1}$. If $\operatorname{Var}(\hat{X}_n) \sim \operatorname{Var}(X_n)$ as

 $n \to \infty$, then

$$\frac{X_n - \mathbb{E}(X_n)}{\operatorname{Var}(X_n)^{1/2}} = \frac{\hat{X}_n - \mathbb{E}(\hat{X}_n)}{\operatorname{Var}(\hat{X}_n)^{1/2}} + o_{\mathbb{P}}(1).$$

3. Asymptotic results

Asymptotic normality. For the univariate statistics $X_{inv}^{(d)}$ and $X_{des}^{(d)}$, the asymptotic normality has been proven in [9]. For the joint statistic $\left(X_{inv}^{(d)}, X_{des}^{(d)}\right)^{\top}$, the simplest case is when d remains fixed. Then, both $X_{inv}^{(d)}$ and $X_{des}^{(d)}$ are m-dependent (m = d) and it is not necessary to apply the Hájek approximation. From the CLT for m-dependent random vectors, it follows that:

Theorem 3.1. For any two fixed numbers d_1, d_2 , the joint distribution $\left(X_{\text{inv}}^{(d_1)}, X_{\text{des}}^{(d_2)}\right)^{\top}$ satisfies the CLT.

An even stronger statement than the CLT is the uniform Gaussian approximation over all hyperrectangles for *m*-dependent random vectors given in [3, Theorem 2]. It also applies for random vectors with a sparse dependency graph. Let $(X_j^{(n)})_{j=1,\dots,n}$ be a triangular array of (w.l.o.g., centered) *p*-dimensional random vectors and let

$$X^{(n)} := \sum_{t=1}^{n} X_t^{(n)}, \qquad \Sigma^{(n)} := \operatorname{Var}(X^{(n)}), \qquad \mathcal{N} \sim \operatorname{N}_p(0, \Sigma^{(n)}).$$

Let $\mathcal{A}^{\mathrm{re}}$ be the system of all *p*-dimensional hyperrectangles, including infinite bounds, i.e., $\mathcal{A}^{\mathrm{re}} := \left\{ \{ \mathbf{w} \in \mathbb{R}^p : \mathbf{a} \leq \mathbf{w} \leq \mathbf{b} \} \mid \mathbf{a}, \mathbf{b} \in [-\infty, \infty]^p \right\}$. Let

$$r_n(\mathcal{A}^{\mathrm{re}}) := \sup_{A \in \mathcal{A}^{\mathrm{re}}} |\mathbb{P}(X^{(n)} \in A) - \mathbb{P}(\mathcal{N} \in A)|$$

In particular, $r_n(\mathcal{A}^{re})$ is an upper bound for

$$r_n(\mathcal{A}^{\mathrm{CLT}}) := \sup_{u \in \mathbb{R}^p} |\mathbb{P}(X^{(n)} \le u) - \mathbb{P}(\mathcal{N}_n \le u)|$$

which is relevant for the CLT, and for

$$r_n(\mathcal{A}^{\text{ext}}) := \sup_{u \in \mathbb{R}^p} |\mathbb{P}(X^{(n)} > u) - \mathbb{P}(\mathcal{N}_n > u)|$$

which is relevant for the asymptotics of extreme values. Now, let G_n be a dependency graph for $X_1^{(n)}, \ldots, X_n^{(n)}$, which consists of all edges (i, j) for which $X_i^{(n)}$ and $X_j^{(n)}$ are dependent. Let Δ_n be the maximum degree of G_n and let Δ_n^* be the maximum degree of the 2-reachability graph of G_n . If the graphs G_n are not too dense as $n \to \infty$, then $r_n(\mathcal{A}^{\text{re}})$ can be bounded as follows: **Theorem 3.2** (cf. [3], Theorem 2). Let $(X_j^{(n)})_{j=1,\dots,n}$, $r_n(\mathcal{A}^{re})$, Δ_n, Δ_n^* be as above. Under some conditions which are satisfied by bounded variables, it holds that

$$r_n(\mathcal{A}^{\rm re}) = O\left(\frac{(\Delta_n \Delta_n^*)^{1/3} \log(p)^{7/6}}{n^{1/6}}\right).$$

In particular, if $(X_j^{(n)})_{j=1,\dots,n}$ is m-dependent for some global constant $m \in \mathbb{N}$, then $r_n(\mathcal{A}^{\mathrm{re}}) = O(n^{-1/6}\log(p)^{7/6}).$

This theorem was stated by Chang *et al.* for high dimensions, but it also works with fixed p by artificially repeating the components of a random vector in fixed dimension.

From now, we consider the univariate statistics $X_{\text{inv}}^{(d_1)}, X_{\text{des}}^{(d_2)}$ and the joint statistic $\left(X_{\text{inv}}^{(d_1)}, X_{\text{des}}^{(d_2)}\right)^{\top}$ for two sequences $d_1 = d_1(n), d_2 = d_2(n)$ with $d_1(n), d_2(n) \leq d_{\max} - 1$ $\forall n \in \mathbb{N}$. Both $X_{\text{inv}}^{(d_1)}$ and $X_{\text{des}}^{(d_2)}$ are based on a sequence of classical Weyl groups $(W_n)_{n \in \mathbb{N}}$, where each W_n is one of S_n, B_n , or D_n .

If $d_2 = d_2(n)$ grows as $n \to \infty$, then there exists no constant $m \in \mathbb{N}$ for which all $X_{des}^{(d_2)}$ are *m*-dependent. However, the dependency structure of $X_{des}^{(d_2)}$ is still sparse. Recall that $X_{inv}^{(d_1)}$ and $X_{des}^{(d_2)}$ are constructed from i.i.d. random variables Z_1, \ldots, Z_n . According to (3), (5c), (5d), we can represent $X_{des}^{(d_2)}$ as a sum of indicator variables, each of which depends on at most three others. Therefore, the first-order and second-order maximum degrees Δ_n, Δ_n^* are bounded in the way of $\Delta_n \leq 3$ and $\Delta_n^* \leq 9$. In conclusion, there are no issues with applying Theorem 3.2 if d_2 grows.

If $d_1 = d_1(n)$ grows as well, then the maximum degrees Δ_n, Δ_n^* of $\left(X_{\text{inv}}^{(d_1)}, X_{\text{des}}^{(d_2)}\right)^{\top}$ are bounded in the way of $\Delta_n \leq 4d_1$ and $\Delta_n^* \leq 8d_1$. Moreover, we have to take into account that by (2), (5a), (5b), $X_{\text{inv}}^{(d_1)}$ is based on $\Theta(nd_1)$ summands, so we have to replace nwith nd in Theorem 3.2. Therefore, Theorem 3.2 gives an o(1) bound of $r_n(\mathcal{A}^{\text{re}})$ if and only if $(d_1)^{2/3} = o((nd_1)^{1/6}) \iff d_1 = o(n^{1/3})$. For any faster growth rate of d_1 , the dependency structure of $\left(X_{\text{inv}}^{(d_1)}, X_{\text{des}}^{(d_2)}\right)^{\top}$ is too complex to apply Theorem 3.2. In that case, we have to replace $X_{\text{inv}}^{(d_1)}$ with the Hájek projection $\hat{X}_{\text{inv}}^{(d_1)}$. The growth rate of d_1 determines whether the condition in Theorem 2.8 is fulfilled or not. To classify the validity of this condition, we distinguish the *short case* $d \leq d_{\text{max}}/2$ and the *long case* $d > d_{\text{max}}/2$.

Lemma 3.3. Given the short case, we have for symmetric groups that

$$\operatorname{Var}\left(\hat{X}_{\operatorname{inv}}^{(d)}\right) = \frac{1}{72}\left(4d^3 + 6d^2 + 2d\right)$$

and we have $\operatorname{Var}\left(\hat{X}_{\operatorname{inv}}^{(d)}\right) = d^3/36 + O(d^2)$ for the groups B_n and D_n . So, if $d_{\max}/2 > d \gg \sqrt{n}$, then $\operatorname{Var}\left(\hat{X}_{\operatorname{inv}}^{(d)}\right) \sim \operatorname{Var}\left(X_{\operatorname{inv}}^{(d)}\right)$.

Lemma 3.4. Given the long case, $\operatorname{Var}\left(\hat{X}_{\operatorname{inv}}^{(d)}\right) \sim \operatorname{Var}\left(X_{\operatorname{inv}}^{(d)}\right)$ holds on symmetric groups if and only if $d \sim d_{\max}$. On the other classical Weyl groups, $\operatorname{Var}\left(\hat{X}_{\operatorname{inv}}^{(d)}\right) \sim \operatorname{Var}\left(X_{\operatorname{inv}}^{(d)}\right)$ holds if and only if $d \sim d_{\max}$ or $d \sim d_{\max}/2$.

From Theorem 3.2 and Slutsky's lemma, we can conclude:

Corollary 3.5. $\left(X_{\text{inv}}^{(d_1)}, X_{\text{des}}^{(d_2)}\right)^{\top}$ satisfies the CLT if $d_1 = o(n^{1/3})$ or if d_1 satisfies the conditions of either Lemma 3.3 or Lemma 3.4.

Extreme values of the univariate statistics $X_{inv}^{(d)}$ and $X_{des}^{(d)}$. We now postulate the univariate and bivariate extreme value limit theorems (EVLTs) for $X_{inv}^{(d)}$ and $X_{des}^{(d)}$ with the help of the methods used in [6]. The proof of [6, Theorem 4.1] consists of two parts. The first part is to show the Gumbel attraction of the statistic that replaces $X_{inv}^{(d)}$ with $\hat{X}_{inv}^{(d)}$. This is done by an application of [3, Theorem 2], providing Gaussian approximation for *m*-dependent random vectors. The second part is to show that the error resulting from this replacement vanishes in probability. Recall that by use of

$$\alpha_n = \frac{1}{\sqrt{2\log n}}, \qquad \beta_n = \frac{1}{\alpha_n} - \frac{1}{2}\alpha_n \left(\log\log n + \log(4\pi)\right),$$

the maximum of n i.i.d. variables $N_1, \ldots, N_n \sim N(0, 1)$ is attracted to the Gumbel distribution, that is, $\forall x \in \mathbb{R}$:

$$\mathbb{P}\left(\max_{i=1,\dots,n} N_i \le \alpha_n x + \beta_n\right) \longrightarrow \exp(-\exp(-x)) =: \Lambda(x).$$

Likewise, the two-dimensional standard normal distribution is attracted to the twodimensional Gumbel distribution with independent marginals. Taking $\boldsymbol{\alpha}_n := (\alpha_n, \alpha_n)$, $\boldsymbol{\beta}_n := (\beta_n, \beta_n) \in \mathbb{R}^2$ and writing "*" for component-wise multiplication, we get that for i.i.d. bivariate standard normal $\mathbf{N}_1, \ldots, \mathbf{N}_n$ and $\mathbf{x} = (x_1, x_2) \in \mathbb{R}^2$:

$$\mathbb{P}\left(\max_{i=1,\dots,n}\mathbf{N}_{i} \leq \boldsymbol{\alpha}_{n} \ast \mathbf{x} + \boldsymbol{\beta}_{n}\right) \longrightarrow \exp\left(-\exp(-x_{1}) - \exp(-x_{2})\right) =: \Lambda_{2}(x).$$

Since the random numbers of generalized inversions and descents are discrete distributions, we are interested in the extremes of a triangular array $(X_{n1}, \ldots, X_{nk_n})$, where for each $n \in \mathbb{N}$, the block X_{n1}, \ldots, X_{nk_n} consists of i.i.d. samples taken from the *n*-th symmetric group S_n . Each of the following EVLTs imposes an upper bound on the number of samples k_n .

For a univariate triangular array consisting of generalized descents, it is not necessary to use the Hájek projection. It has already been argued in [6, Remark 4.2] that a subexponential bound of k_n can be obtained if the Hájek projection is not needed. So, in the univariate EVLT for generalized descents, we can use the bound of $k_n = \exp(o(n^{1/7}))$ stated in [6, Remark 4.2]: **Theorem 3.6.** Let $(X_{nj})_{j=1,\dots,k_n}$ be a row-wise i.i.d. triangular array with $X_{n1} \stackrel{\mathcal{D}}{=} X_{des}^{(d)}$ for any sequence d = d(n) with $1 \le d \le d_{max} - 1$, and let $M_n := \max\{X_{n1},\dots,X_{nk_n}\}$. Let $a_n := \sigma(X_{des}^{(d)})\alpha_{k_n}$ and let $b_n := \sigma(X_{des}^{(d)})\beta_{k_n} + \mu(X_{inv}^{(d)})$. If $k_n = \exp(o(n^{1/7}))$, then $\forall x \in \mathbb{R}$: $\mathbb{P}(M_n \le a_n x + b_n) \longrightarrow \Lambda(x)$.

A similar statement applies to $X_{inv}^{(d)}$ if d grows slow enough to permit the application of Theorem 3.2.

Theorem 3.7. Let $(X_{n1}, \ldots, X_{nk_n})$ be a row-wise i.i.d. triangular array with $X_{n1} \stackrel{D}{=} X_{inv}^{(d)}$, where $d = d(n) = o(n^{1/3})$. Let M_n, a_n, b_n be given in analogy to Theorem 3.6. If $k_n = \exp(o(n^{1/7}/d^{3/7}))$, then

$$\forall x \in \mathbb{R}: \quad \mathbb{P}(M_n \le a_n x + b_n) \longrightarrow \Lambda(x) \,.$$

Proof. According to the above considerations, the maximum degrees Δ_n, Δ_n^* in the dependency graph of the representations (2), (5a), and (5b) are bounded in the way of $\Delta_n \leq 4d$ and $\Delta_n^* \leq 8d$. Since these representations are based on $\Theta(nd)$ summands, we need to replace n with nd when applying Theorem 3.2. An application of Theorem 3.2 with $\max\{n, k_n\}$ i.i.d. iterations of $X_{inv}^{(d)}$ yields

$$|\mathbb{P}(M_n \le a_n x + b_n) - \mathbb{P}(\mathcal{M}_n \le \alpha_n x + \beta_n)| = O(n^{-1/6} d^{1/2}) \log(k_n)^{7/6} = o(1),$$

where \mathcal{M}_n is the maximum of n i.i.d. copies of the standard normal distribution. The claim follows.

For any other growth rate of d, we can state an EVLT only for the cases covered by Lemmas 3.3 and 3.4, and due to the use of Hájek's projection, we can only impose a strongly reduced asymptotic bound for k_n .

Theorem 3.8. Let $(X_{n1}, \ldots, X_{nk_n})$ be a row-wise i.i.d. triangular array with $X_{n1} \stackrel{\mathcal{D}}{=} X_{inv}^{(d)}$, where d = d(n) satisfies the conditions in either Lemma 3.3 or 3.4. Assume $k_n \log(k_n) = o(n^{-1}d^2)$ in the short case and $k_n \log(k_n) = o\left(\min\left\{n, \frac{n}{n-d}\right\}\right)$ in the long case. Let $M_n := \max\{X_{n1}, \ldots, X_{nk_n}\}$ be the row-wise maximum. Then,

$$\forall x \in \mathbb{R}: \quad \mathbb{P}(M_n \le a_n x + b_n) \longrightarrow \Lambda(x) \,,$$

with a_n, b_n given in analogy to Theorem 3.6.

Proof. The conditions on d ensure that $1 - \operatorname{Var}(X_{inv}^{(d)}) / \operatorname{Var}(\hat{X}_{inv}^{(d)}) = o(1)$, but the rate of convergence determines the bound for k_n by means of [6, Eq. (9), (10)]. We compute this rate for symmetric groups, since the same conclusions can be obtained on the other classical Weyl groups. In the short case, we have

$$1 - \frac{\operatorname{Var}(X_{\operatorname{inv}}^{(d)})}{\operatorname{Var}(\hat{X}_{\operatorname{inv}}^{(d)})} = 1 - \frac{4d^3 + 6nd + 3d^2 - d}{4d^3 + 6d^2 + 2d} = \frac{4d^2 + 6n + 3d - 1}{4d^2 + 6d + 2}$$

$$= 1 - \frac{4d^2}{4d^2 + 6d + 2} - \frac{3d + 1}{4d^2 + 6d + 2} - \frac{6n}{4d^2 + 6d + 2}$$
$$= \Theta\left(\frac{1}{d}\right) - \frac{6n}{4d^2 + 6d + 2}.$$

Apparently, $6n/(4d^2 + 6d + 2)$ always dominates 1/d. In conclusion,

$$1 - \frac{\operatorname{Var}(X_{\operatorname{inv}}^{(d)})}{\operatorname{Var}(\hat{X}_{\operatorname{inv}}^{(d)})} = \Theta\left(\frac{n}{d^2}\right),$$

giving the condition of $k_n \log(k_n) = o(d^2/n)$ according to the arguments in the proof of [6, Theorem 4.1]. The remaining steps of the proof are the same.

In the long case, we write $\delta := n - d$ and note that

$$\operatorname{Var}(\hat{X}_{inv}) = -\frac{1}{6}(n-\delta)^3 + \frac{1}{3}(n-\delta)^2n - \frac{5}{36}n^3 + \frac{16}{36}n^2\delta + O(n^2),$$
$$\operatorname{Var}(X_{inv}) = -\frac{1}{6}(n-\delta)^3 + \frac{1}{3}(n-\delta)^2n - \frac{5}{36}n^3 + \frac{6}{36}n^2\delta + O(n^2).$$

Therefore,

$$1 - \frac{\operatorname{Var}(X_{\operatorname{inv}})}{\operatorname{Var}(\hat{X}_{\operatorname{inv}})} = \frac{10n^2\delta + O(n^2)}{\Theta(n^3)} = \Theta\left(\max\left\{\frac{\delta}{n}, \frac{1}{n}\right\}\right),$$

giving the requirement of

$$k_n \log(k_n) \Theta\left(\max\left\{\frac{\delta}{n}, \frac{1}{n}\right\}\right) = o(1) \iff k_n \log(k_n) = o\left(\min\left\{n, \frac{n}{\delta}\right\}\right).$$

Again, the proof now follows the same steps as in [6, Theorem 4.1].

Remark 3.9. This result is only useful if $d \gg n^{3/4}$. Otherwise, we can obtain the EVLT with a bound of $k_n = o(n^{1/2})$ from the universal EVLT [7, Theorem 5.1].

Extreme values of the joint distribution of $X_{inv}^{(d)}$ and $X_{des}^{(d)}$. The bivariate EVLT for $\left(X_{inv}^{(d_1)}, X_{des}^{(d_2)}\right)^{\top}$ follows the same way, since the descent component does not interfere with any of the previously used arguments.

Theorem 3.10. Let $(X_{n1}, \ldots, X_{nk_n})$ be a row-wise i.i.d. triangular array with $X_{n1} \stackrel{\mathcal{D}}{=} (X_{inv}^{(d_1)}, X_{des}^{(d_2)})^{\top}$ on S_n .

- (a) If $d_1 = o(n^{1/3})$, then assume $k_n = \exp(o(n^{1/7}/d^{3/7}))$.
- (b) If d_1 satisfies the conditions in either Lemma 3.3 or 3.4, then assume $k_n \log(k_n) = o(n^{-1}d_1^2)$ in the short case and $k_n \log(k_n) = o\left(\min\left\{n, \frac{n}{n-d_1}\right\}\right)$ in the long case.

Let
$$M_n := \max\{X_{n1}, \ldots, X_{nk_n}\}$$
 be the row-wise maximum. Let $\mu_n := \mathbb{E}(X_{n1})$ and let $s_n := \left(\sigma(X_{inv}^{(d)}), \sigma(X_{des})\right)$. Let $a_n := s_n * \mathbf{\alpha}_{k_n}$ and let $b_n := s_n * \mathbf{\beta}_{k_n} + \mu_n$. Then,
 $\forall \mathbf{x} \in \mathbb{R}^2$: $\mathbb{P}(M_n \leq a_n * \mathbf{x} + b_n) \longrightarrow \Lambda_2(\mathbf{x})$.

4. Proofs of Lemmas 3.3 and 3.4

We first consider $\hat{X}_{inv}^{(d)}$ on symmetric groups and then provide the analogous observations on the other classical Weyl groups.

Proof of Lemma 3.3 for symmetric groups. By (2), we have

$$\mathbb{E}(X_{\text{inv}}^{(d)} \mid Z_k) = \sum_{(i,j)\in\mathfrak{N}_{n,d}} \mathbb{P}(Z_i > Z_j \mid Z_k) = \sum_{(i,j)\in\mathfrak{N}_{n,d}} \begin{cases} 1/2, & k \notin \{i,j\} \\ Z_k, & k=i \\ 1-Z_k, & k=j \end{cases}$$

Only the pairs (i, j) with $k \in \{i, j\}$ contribute to $\operatorname{Var}\left(\mathbb{E}\left(X_{\operatorname{inv}}^{(d)} \mid Z_k\right)\right)$. These contributions are called the *non-trivial parts* for simplicity. The number of these pairs depends on whether k belongs to K_1, K_2 , or K_3 , as already seen in Remark 2.2. Figure 1 visualizes this case distinction for the exemplary choice of n = 15 and d = 4. If $k \in K_2$, then the non-trivial parts are

$$\underbrace{Z_k + Z_k + \ldots + Z_k}_{d \text{ times}} + \underbrace{(1 - Z_k) + (1 - Z_k) + \ldots + (1 - Z_k)}_{d \text{ times}} = d$$

This means $\mathbb{E}\left(X_{\text{inv}}^{(d)} \mid Z_k\right)$ is constant due to cancellation, and vanishes when computing the variance. So, in the short case, $\operatorname{Var}\left(\mathbb{E}\left(X_{\text{inv}}^{(d)} \mid Z_k\right)\right)$ originates only from K_1 and K_3 . If $k \in K_1$, then $\mathbb{E}\left(X_{\text{inv}}^{(d)} \mid Z_k\right) = (k-1)(1-Z_k) + dZ_k + \operatorname{const} = (d+1-k)Z_k + \operatorname{const}$. If $k \in K_3$, then $\mathbb{E}\left(X_{\text{inv}}^{(d)} \mid Z_k\right) = (n-k)Z_k + d(1-Z_k) + \operatorname{const} = (n-d-k)Z_k + \operatorname{const}$. So, the overall representation of $\hat{X}_{\text{inv}}^{(d)}$ in the short case is

$$\hat{X}_{inv}^{(d)} = \sum_{k=1}^{n} \omega_d(k) Z_k + \text{const}, \quad \text{with} \quad \omega_d(k) := \begin{cases} d-k+1, & k \in K_1 \\ 0, & k \in K_2 \\ n-d-k, & k \in K_3 \end{cases}$$

Therefore,

$$\operatorname{Var}\left(\hat{X}_{inv}^{(d)}\right) = \operatorname{Var}\left(\sum_{k=n-d+1}^{n} (n-k-d)Z_k + \sum_{k=1}^{d} (d+1-k)Z_k\right)$$
$$= \sum_{k=n-d+1}^{n} \frac{1}{12}(n-k-d)^2 + \sum_{k=1}^{d} \frac{1}{12}(d+1-k)^2$$



FIGURE 1. Overview of relevant pairs (i, j) for computing the variance of $\mathbb{E}(X_{inv}^{(d)} | Z_k)$, where n = 15 and d = 4. For each of the regions K_1, K_2, K_3 , an exemplary index k is chosen, and the pairs that give a non-trivial contribution are highlighted in red for $k \in K_1$, in blue for $k \in K_2$, and in green for $k \in K_3$.

$$= \frac{1}{12} \sum_{k=1}^{d} k^2 + \frac{1}{12} \sum_{k=1}^{d} (-k)^2 = \frac{1}{6} \frac{d(d+1)(2d+1)}{6}$$
$$= \frac{1}{72} \left(4d^3 + 6d^2 + 2d \right).$$

So, the leading term is always $4d^3/72$. In light of Theorem 2.3, according to which $\operatorname{Var}\left(X_{\mathrm{inv}}^{(d)}\right)$ contains the monomials 6nd/72 and $4d^3/72$, it must be ensured that d^3 is dominant over nd. This means $d \gg \sqrt{n}$.

Now, we consider the long case. Due to d > n/2, we now have n - d < d, and the regions K_1, K_2, K_3 are redefined according to Remark 2.2. For the non-trivial parts, we note that:

• If $k \in K_1$ or $k \in K_3$, then the non-trivial parts are the same as in the short case.

• If $k \in K_2$, then the non-trivial parts yield $\mathbb{E}\left(X_{inv}^{(d)} \mid Z_k\right) = (n-2k+1)Z_k + \text{const.}$ So, we again obtain a representation in the way of

(6)
$$\hat{X}_{inv}^{(d)} = \sum_{k=1}^{n} \omega_d(k) Z_k + \text{const}, \quad \text{with} \quad \omega_d(k) := \begin{cases} d-k+1, & k \in K_1 \\ n-2k+1, & k \in K_2 \\ n-d-k, & k \in K_3 \end{cases}$$

The proof of the following lemma shows that if $d \not\sim n$, then the leading terms of $\operatorname{Var}\left(X_{\operatorname{inv}}^{(d)}\right)$ and $\operatorname{Var}\left(\hat{X}_{\operatorname{inv}}^{(d)}\right)$ do not match perfectly.

Proof of Lemma 3.4 for symmetric groups. From (6), we state that

(6a)
$$\operatorname{Var}\left(\hat{X}_{\mathrm{inv}}^{(d)}\right) = \frac{1}{12} \sum_{k=1}^{n-d} (d+1-k)^2$$

(6b)
$$+ \frac{1}{12} \sum_{k=n-d+1}^{d} (n-2k+1)^2$$

(6c)
$$+ \frac{1}{12} \sum_{k=d+1}^{n} (n-k-d)^2.$$

By appropriate index shifting, we calculate

$$(6a), (6c) = \frac{1}{12} \left(\sum_{k=1}^{d} k^2 - \sum_{k=1}^{2d-n} k^2 \right)$$

$$\implies (6a) + (6c) = \frac{1}{6} \left(\sum_{k=1}^{d} k^2 - \sum_{k=1}^{2d-n} k^2 \right)$$

$$= \frac{1}{36} (n-d) \left(14d^2 + d(9-10n) + 12n^2 - 3n + 1 \right),$$

$$(6b) = \frac{1}{36} (2d-n) (4d^2 - 4dn + n^2 - 1).$$

This gives the total result

$$\operatorname{Var}\left(\hat{X}_{\operatorname{inv}}^{(d)}\right) = \frac{1}{36} \left(-6d^3 + 12d^2n - 9d^2 - 16dn^2 + 12dn - 3d + 11n^3 - 3n^2 + 2n\right).$$

In contrast, by Theorem 2.3,

$$\operatorname{Var}\left(X_{\operatorname{inv}}^{(d)}\right) = \frac{1}{36}\left(-6d^3 + \left(12n - \frac{21}{2}\right)d^2 - \left(6n^2 - 15n + \frac{9}{2}\right)d + n^3 - 3n^2 + 2n\right).$$

Since the long case implies n/2 < d < n, all monomials of order 3 are leading terms. The monomials $-16dn^2 + 11n^3$ appearing in Var $(\hat{X}_{inv}^{(d)})$ and $-6n^2d + n^3$ appearing in

 $\operatorname{Var}\left(X_{\operatorname{inv}}^{(d)}\right)$ do not match in general. Using $\delta := n - d$, we can write

$$\operatorname{Var}(\hat{X}_{\mathrm{inv}}^{(d)}) = -\frac{1}{6}(n-\delta)^3 + \frac{1}{3}(n-\delta)^2n - \frac{5}{36}n^3 + \frac{16}{36}n^2\delta + O(n^2),$$
$$\operatorname{Var}(X_{\mathrm{inv}}^{(d)}) = -\frac{1}{6}(n-\delta)^3 + \frac{1}{3}(n-\delta)^2n - \frac{5}{36}n^3 + \frac{6}{36}n^2\delta + O(n^2).$$

This shows that we need $\delta = o(n)$, since otherwise, there are two different leading coefficients of n^3 .

We now derive the analogous statements for the other classical Weyl groups B_n and D_n . It is sufficient to prove the lemmas 3.3 and 3.4 for the groups B_n , since the difference between $X_{inv}^{B,(d)}$ and $X_{inv}^{D,(d)}$ is asymptotically negligible (cf. (5a) and (5b)). Recall the asymptotic quantification of $Var(X_{inv}^{(d)})$ given in Lemma 2.6.

To compute $\operatorname{Var}(\hat{X}_{\operatorname{inv}}^{(d)})$, we ignore all constant parts appearing in $\hat{X}_{\operatorname{inv}}^{(d)}$. By (5a), we can split $\mathbb{E}(X_{\operatorname{inv}}^{B,(d)} \mid Z_k) = \mathbb{E}(X_{\operatorname{inv}}^{B,(d)} \mid Z_k)^+ + \mathbb{E}(X_{\operatorname{inv}}^{B,(d)} \mid Z_k)^- + \mathbb{E}(X_{\operatorname{inv}}^{B,(d)} \mid Z_k)^\circ$, where $\mathbb{E}(X_{\operatorname{inv}}^{B,(d)} \mid Z_k)^+ := \sum_{(i,j)\in\mathfrak{N}_{n,d}} \mathbb{P}(Z_i > Z_j \mid Z_k)$, $\mathbb{E}(X_{\operatorname{inv}}^{B,(d)} \mid Z_k)^- := \sum_{(i,j)\in\mathfrak{\widetilde{N}}_{n,d}} \mathbb{P}(-Z_i > Z_j \mid Z_k)$, $\mathbb{E}(X_{\operatorname{inv}}^{B,(d)} \mid Z_k)^\circ := \sum_{i=1}^n \mathbb{P}(Z_i < 0 \mid Z_k)$.

The third sum $\mathbb{E}(X_{\text{inv}}^{B,(d)} \mid Z_k)^{\circ}$ is asymptotically negligible. So, to obtain a representation $\hat{X}_{\text{inv}}^{B,(d)} = \sum_{i=1}^{n} \omega_d(k) Z_k + \text{const}$, we write $\omega_d(k) = \omega_d(k)^+ + \omega_d(k)^-$, with $\omega_d(k)^+$ stemming from $\mathbb{E}(X_{\text{inv}}^{B,(d)} \mid Z_k)^+$ and $\omega_d(k)^-$ stemming from $\mathbb{E}(X_{\text{inv}}^{B,(d)} \mid Z_k)^-$, respectively.

For $\omega_d(k)^+$, we can use the previous counting method. However, we have to take into account that on B_n ,

$$\mathbb{P}(Z_k > Z_j \mid Z_k) = \frac{Z_k + 1}{2}, \qquad \mathbb{P}(Z_i > Z_k \mid Z_k) = \frac{1 - Z_k}{2},$$
$$\mathbb{P}(-Z_k > Z_j \mid Z_k) = \frac{1 - Z_k}{2}, \qquad \mathbb{P}(Z_i > Z_k \mid Z_k) = \frac{1 - Z_k}{2}.$$

In conclusion, the coefficients $\omega_d(k)^+$ on B_n are half of the coefficients $\omega_d(k)$ on S_n . Note that $\omega_d(k)^+ = (n - 2k + 1)/2$ for all $d \ge n$. Moreover, $\omega_d(k)^- = -\widetilde{N}_{n,d}^{(k)}/2$, where

$$\widetilde{N}_{n,d}^{(k)} := \left| \{ (i,j) \in \widetilde{\mathfrak{N}}_{n,d} \mid \text{either } i = k \text{ or } j = k \} \right|.$$

Recall that by Remark 2.5, the short case is now d < n and the long case is $d \ge n$. By counting the number of such (k, j) and (i, k), we can determine the asymptotic quantification of Var $(\hat{X}_{inv}^{B,(d)})$.

Proof of Lemma 3.3 for the groups B_n and D_n . If $d \leq n/2$, then all pairs in $\mathfrak{N}_{n,d}$ are located within K_1 , yielding

$$\omega_d(k) = \begin{cases} (d-k+1)/2 - (d-k-1)/2, & k \le d/2\\ (d-k+1)/2 - (d-k)/2, & d/2 < k \le d\\ 0, & d < k \le n-d\\ (n-d-k)/2, & n-d < k \le n \end{cases}$$
$$= \begin{cases} (n-d-k)/2, & n-d < k \le n\\ O(1), & \text{otherwise} \end{cases}.$$

In conclusion, if $d \leq n/2$, then

$$\operatorname{Var}\left(\hat{X}_{\operatorname{inv}}^{B,(d)}\right) = \frac{1}{3} \sum_{k=1}^{n} \omega_d(k)^2 + O(d^2) = \frac{1}{12} \sum_{k=n-d+1}^{n} (n-d-k)^2 + O(d^2)$$
$$= \frac{1}{12} \frac{d(d+1)(2d+1)}{6} + O(d^2) = \frac{1}{36} d^3 + O(d^2).$$

Due to Var $(X_{inv}^{B,(d)}) = d^3/36 + nd/12 + O(d^2)$ according to Lemma 2.6, we obtain the same condition as in Lemma 3.3, namely, $d \gg \sqrt{n}$.

If n/2 < d < n, then the pairs in $\mathfrak{N}_{n,d}$ also cover K_2 . For $k \in K_2$, there cannot be any pairs (k, j) if $n/2 < d \leq 2n/3$, while this is possible if d > 2n/3. However, the difference between these two subcases is only marginal. If $n/2 < d \leq 2n/3$, we obtain

$$\omega_d(k) = \begin{cases} O(1), & k \in K_1 \\ \left(n - 2k + 1 - (d - k)\right)/2, & k \in K_2 \\ (n - d - k)/2, & k \in K_3 \end{cases}$$

If d > 2n/3, then

$$\omega_d(k) = \begin{cases} O(1), & k \in K_1 \\ \left(n - 2k + 1 - (d - k - 1)\right)/2, & k \in K_2, k \le d/2 \\ \left(n - 2k + 1 - (d - k)\right)/2, & k \in K_2, k > d/2 \\ (n - d - k)/2, & k \in K_3 \end{cases}$$

In conclusion, if n/2 < d < n, then

$$\operatorname{Var}\left(\hat{X}_{\mathrm{inv}}^{B,(d)}\right) = \frac{1}{3} \sum_{k=1}^{n} \omega_d(k)^2 + O(d^2)$$
$$= \frac{1}{12} \sum_{k=n-d+1}^{d} (n-d-k+1)^2 + \frac{1}{12} \sum_{k=d+1}^{n} (n-d-k)^2 + O(d^2)$$

$$= \frac{1}{12} \sum_{k=n-d+1}^{d} (n-d-k+1)^2 + \frac{1}{12} \sum_{k=1}^{2d-n} k^2 + O(d^2)$$

$$= \frac{1}{12} \sum_{k=1}^{2d-n-1} k^2 + \frac{1}{12} \left(\sum_{k=1}^{d} k^2 - \sum_{k=1}^{2d-n} k^2 \right) + O(d^2)$$

$$= \frac{1}{36} d^3 + O(d^2).$$

Proof of Lemma 3.4 for the groups B_n and D_n . In the long case, the main focus is on counting $\widetilde{N}_{n,d}^{(k)}$. Figure 2 illustrates the positions of pairs $(i, j) \in \widetilde{\mathfrak{N}}_{n,d}$ in the long case for the exemplary choice of n = 12, d = 16.



FIGURE 2. Visualization of pairs (i, j) in $\widetilde{\mathfrak{N}}_{n,d}$ in the long case for n = 12and d = 16. The numbers d - n and d/2 are important case distinction thresholds for counting the pairs (k, j) and (i, k), respectively.

With help of Figure 2, it is straightforward to count

$$\widetilde{N}_{n,d}^{(k)} = \begin{cases} n-1, & 1 \le k \le d-n \\ d-k-1, & d-n < k \le d/2 \\ d-k, & d/2 < k \le n \end{cases}$$

This result is also illustrated in Figure 3, which displays the number of pairs (i, k) and the number of pairs (k, j).



FIGURE 3. Plots of the numbers of pairs (i, k) (red) and (k, j) (blue) in $\widetilde{\mathfrak{N}}_{n,d}$. The sum of these two numbers is $\widetilde{N}_{n,d}^{(k)}$, which is displayed by the black crossed points.

Therefore, in the long case, we have

$$\omega_d(k) = \begin{cases} 1-k, & 1 \le k \le d-n\\ (n+2-d-k)/2, & d-n < k \le d/2,\\ (n+1-d-k)/2, & d/2 < k \le n. \end{cases}$$

We compute

$$\sum_{k=1}^{n} \omega_d(k)^2 = \sum_{k=1}^{d-n} (k-1)^2 + \frac{1}{4} \sum_{k=d-n+1}^{n} (n-d+1-k)^2 + O(n^2)$$
$$= \sum_{k=1}^{d-n-1} k^2 + \frac{1}{4} \sum_{k=d-n+1}^{n} \left(k - (d-n-1)\right)^2 + O(n^2)$$

$$= \frac{1}{3}(d-n-1)^3 + \frac{1}{12}(2n-d-1)^3 + O(n^2)$$

= $\frac{d^3}{4} - \frac{d^2n}{2} - \frac{3d^2}{4} + dn + \frac{3d}{4} + \frac{n^3}{3} - \frac{n}{2} + \frac{1}{4} + O(n^2)$

Due to $\operatorname{Var}(Z_k) = 1/3$, we obtain

$$\operatorname{Var}(\hat{X}_{\operatorname{inv}}^{B,(d)}) = \frac{d^3}{12} - \frac{d^2n}{6} + \frac{n^3}{9} + O(d^2)$$

On the other hand, by Lemma 2.6,

$$\operatorname{Var}(X_{\operatorname{inv}}^{B,(d)}) = -\frac{1}{12}d^3 + \frac{1}{3}nd^2 - \frac{1}{3}n^2d + \frac{1}{9}n^3 + O(d^2)$$

i.e., the leading terms $d^3/12 - d^2n/6$ and $-d^3/12 + nd^2/3 - n^2d/3$ need to match. Writing d = cn(1 + o(1)) for $1 \le c \le 2$, this gives the condition

$$\frac{(cn)^3}{12} - \frac{c^2n^3}{6} = -\frac{(cn)^3}{12} + \frac{c^2n^3}{3} - \frac{cn^3}{3} \iff \frac{c^3}{6} - \frac{c^2}{2} + \frac{c}{3} = 0,$$

which holds precisely for c = 1 and c = 2.

References

- Anders Björner and Francesco Brenti, Combinatorics of Coxeter groups, vol. 231, Springer Science & Business Media, 2006.
- Miklós Bóna, Generalized descents and normality, Electronic Journal of Combinatorics 15 (2008), N21.
- Jinyuan Chang, Xiaohui Chen, and Mingcong Wu, Central limit theorems for high dimensional dependent data, arXiv preprint arXiv:2104.12929 (2021).
- 4. Harold S. M. Coxeter, The complete enumeration of finite groups of the form $r_i^2 = (r_i r_j)^{k_{ij}} = 1$, Journal of the London Mathematical Society 1 (1935), no. 1, 21–25.
- Filippo de Mari and Mark A. Shayman, Generalized eulerian numbers and the topology of the hessenberg variety of a matrix, Acta Applicandae Mathematica 12 (1988), 213–235.
- Philip Dörr and Johannes Heiny, Extremes of joint inversions and descents on finite coxeter groups, arXiv preprint arXiv:2309.17314 (2023).
- 7. Philip Dörr and Thomas Kahle, *Extreme values of permutation statistics*, arXiv preprint arXiv:2205.01426 (2022).
- Thomas Kahle and Christian Stump, Counting inversions and descents of random elements in finite Coxeter groups, Mathematics of Computation 89 (2020), no. 321, 437–464.
- Kathrin Meier and Christian Stump, Central limit theorems for generalized descents and generalized inversions in finite root systems, preprint, arXiv:2202.05580 (2022).
- 10. John Pike, *Convergence rates for generalized descents*, the electronic journal of combinatorics (2011), P236–P236.
- 11. Aad W. van der Vaart, Asymptotic statistics, vol. 3, Cambridge university press, 2000.

DEPARTMENT OF MATHEMATICS, MAGDEBURG... Email address: philip.doerr@ovgu.de

18