

Uniqueness to inverse acoustic and elastic medium scattering problems with hyper-singular source method

Chun Liu*, Guanghui Hu*, Jianli Xiang[†] and Jiayi Zhang[‡]

Abstract

This paper is concerned with inverse scattering problems of determining the support of an isotropic and homogeneous penetrable body from knowledge of multi-static far-field patterns in acoustics and in linear elasticity. The normal derivative of the total fields admits no jump on the interface of the scatterer in the trace sense. If the contrast function of the refractive index function or the density function has a positive lower bound near the boundary, we propose a hyper-singular source method to prove uniqueness of inverse scattering with all incoming plane waves at a fixed energy. It is based on subtle analysis on the leading part of the scattered field when hyper-singular sources caused by the first derivative of the fundamental solution approach to a boundary point. As a by-product, we show that this hyper-singular method can be also used to determine the boundary value of a Hölder continuous refractive index function in acoustics or a Hölder continuous density function in linear elasticity.

Keywords: Inverse medium scattering, Helmholtz equation, Navier equation, uniqueness.

1 Introduction

1.1 Acoustic medium scattering problem

Consider a time-harmonic acoustic plane wave incident onto a bounded penetrable obstacle $D \subset \mathbb{R}^3$ embedded in a homogeneous isotropic medium. The spatially-dependent incident field $u^{\text{in}}(x, d)$ takes the form $u^{\text{in}}(x, d) = e^{ikx \cdot d}$, $x \in \mathbb{R}^3$, where $d \in \mathbb{S}^2 := \{x \in \mathbb{R}^3 : |x| = 1\}$ is the incident direction and $k > 0$ is the wave number. We assume that the complement of D , $D^e := \mathbb{R}^3 \setminus \bar{D}$, is connected and D is of C^2 -smoothness. The acoustic properties of the scatterer can be described by the refractive index function $n(x)$ such that $n(x) = 1$ in D^e after some normalization. Hence the contrast function $1 - n(x)$ is compactly supported in D . The wave propagation is modeled by the Helmholtz equation

$$\Delta u + k^2 n u = 0 \text{ in } \mathbb{R}^3. \quad (1.1)$$

*School of Mathematical Sciences and LPMC, Nankai University, Tianjin, 300071, China (liuchun@nankai.edu.cn, ghhu@nankai.edu.cn)

[†]Three Gorges Mathematical Research Center, College of Science, China Three Gorges University, Yichang, 443002, China (xiangjianli@ctgu.edu.cn)

[‡]Corresponding author. School of Mathematical Sciences and LPMC, Nankai University, Tianjin, 300071, China (zhangjy97@mail.nankai.edu.cn)

In (1.1), $u = u^{\text{in}} + u^{\text{sc}}$ denotes the total field, where u^{sc} is the perturbed scattered field satisfying the Sommerfeld radiation condition

$$\lim_{|x| \rightarrow \infty} r \left\{ \frac{\partial u^{\text{sc}}}{\partial r} - iku^{\text{sc}} \right\} = 0, \quad r = |x|. \quad (1.2)$$

The Sommerfeld radiation condition (1.2) leads to the asymptotic expansion

$$u^{\text{sc}}(x) = \frac{e^{ik|x|}}{|x|} u^\infty(\hat{x}) + \mathcal{O}\left(\frac{1}{|x|^{3/2}}\right), \quad |x| \rightarrow +\infty,$$

uniformly in all directions $\hat{x} := x/|x|$, $x \in \mathbb{R}^3$. The function $u^\infty(\hat{x})$ is an analytic function defined on \mathbb{S}^2 and is referred to as the far-field pattern or the scattering amplitude. The vector $\hat{x} \in \mathbb{S}^2$ is called the observation direction of the far field. We also need transmission conditions on the interface ∂D . In this paper we assume the *continuity* of the total field and its normal derivative in the trace sense that,

$$u^+ = u^-, \quad \partial_\nu u^+ = \partial_\nu u^- \quad \text{on } \partial D. \quad (1.3)$$

Here $\nu \in \mathbb{S}^2$ is the unit normal vector on ∂D pointing into D^e , and the superscripts $(\cdot)^\pm$ stand for the limits taken from outside and inside, respectively.

The direct scattering problem for (1.1), (1.2) and (1.3) can be formulated as a task of finding the scattered field u^{sc} when the incident field u^{in} and n are given. If the refractive index $n \in C(\overline{D})$ satisfies $\text{Re } n > 0$ and $\Im n \geq 0$, it was shown in [3, Theorems 8.1 and 8.3] that the direct scattering problem admits a unique solution $u^{\text{sc}} \in H_{\text{loc}}^2(\mathbb{R}^3)$.

1.2 Elastic medium scattering problem

We also consider an isotropic and inhomogeneous elastic medium in three dimensions. The medium can be described by two Lamé coefficients λ and μ , and the mass density ρ . We simplify the discussion further by supposing that λ and μ are constants satisfying $\mu > 0$ and $2\mu + 3\lambda > 0$. We also assume that the density $\rho(x) \in L^\infty(\mathbb{R}^3)$ satisfies $\rho(x) = 1$ in $D^c := \mathbb{R}^3 \setminus \overline{D}$. Then the vector field $\mathbf{u} = (u_1, u_2, u_3)$ is governed by the following Lamé system

$$\Delta^* \mathbf{u} + \omega^2 \rho(x) \mathbf{u} = 0 \quad \text{in } \mathbb{R}^3, \quad \Delta^* \mathbf{u} := \mu \Delta \mathbf{u} + (\lambda + \mu) \text{grad div } \mathbf{u}. \quad (1.4)$$

Here $\omega > 0$ is a fixed frequency. Moreover, we assume the exterior D^c is connected. $\mathbf{u} = \mathbf{u}^{\text{in}} + \mathbf{u}^{\text{sc}}$ is the total field and \mathbf{u}^{sc} is the scattered field. \mathbf{u}^{sc} in D^c can be decomposed into the sum of the compressional (longitudinal) part \mathbf{u}_p^{sc} and shear (transversal) part \mathbf{u}_s^{sc} as follows

$$\mathbf{u}^{\text{sc}} = \mathbf{u}_p^{\text{sc}} + \mathbf{u}_s^{\text{sc}}, \quad \mathbf{u}_p^{\text{sc}} = \frac{1}{k_p^2} \text{grad div } \mathbf{u}^{\text{sc}}, \quad \mathbf{u}_s^{\text{sc}} = \frac{1}{k_s^2} \text{curl curl } \mathbf{u}^{\text{sc}}, \quad (1.5)$$

in which

$$k_s := \omega \sqrt{\frac{\rho}{\mu}}, \quad k_p := \omega \sqrt{\frac{\rho}{\lambda + 2\mu}}$$

denote the shear and compressional wave numbers, respectively. It follows from the decomposition in (1.5) that \mathbf{u}_p^{sc} and \mathbf{u}_s^{sc} satisfy the Helmholtz equations

$$(\Delta + k_\alpha^2) \mathbf{u}_\alpha^{\text{sc}} = 0 \quad \text{in } D^c, \quad \alpha = p, s.$$

The scattered field is required to satisfy the Kupradze radiation condition(see e.g. [2])

$$\lim_{r \rightarrow \infty} r \left(\frac{\partial \mathbf{u}_\alpha^{\text{sc}}}{\partial r} - ik_\alpha \mathbf{u}_\alpha^{\text{sc}} \right) = 0, \quad r = |x|, \quad \alpha = p, s, \quad (1.6)$$

uniformly with respect to all directions $\hat{x} = x/|x| \in \mathbb{S}^2 := \{x \in \mathbb{R}^3 : |x| = 1\}$. The radiation conditions in (1.6) lead to the P-part \mathbf{u}_p^∞ and S-part \mathbf{u}_s^∞ of the far-field pattern of u^{sc} , given by the asymptotic behavior

$$\mathbf{u}_s^{\text{sc}}(x) = \frac{\exp(ik_p|x|)}{|x|} \mathbf{u}_p^\infty(\hat{x}) + \frac{\exp(ik_s|x|)}{|x|} \mathbf{u}_s^\infty(\hat{x}) + \mathcal{O}\left(\frac{1}{|x|^2}\right), \quad |x| \rightarrow \infty,$$

where \mathbf{u}_p^∞ and \mathbf{u}_s^∞ are the far-field part of \mathbf{u}_p^{sc} and \mathbf{u}_s^{sc} , respectively. In this paper, we define the far-field pattern \mathbf{u}^∞ of the scattered field u^{sc} as the sum of \mathbf{u}_p^∞ and \mathbf{u}_s^∞ , which means $\mathbf{u}^\infty := \mathbf{u}_p^\infty + \mathbf{u}_s^\infty$. The transmission conditions on the scatterer's surface ∂D are supposed to be

$$\mathbf{u}^+ = \mathbf{u}^-, \quad T\mathbf{u}^+ = T\mathbf{u}^-, \quad \text{on } \partial D, \quad (1.7)$$

where

$$T\mathbf{u} := 2\mu\nu \cdot \nabla \mathbf{u} + \lambda\nu \nabla \cdot \mathbf{u} + \mu\nu \times (\nabla \times \mathbf{u}) \quad (1.8)$$

is the surface stress operator in terms of the outwardly directed unit normal vector ν . Again the superscripts $(\cdot)^\pm$ stand for the limits taken from outside and inside, respectively. The incident wave \mathbf{u}^{in} is supposed to an elastic plane wave of the form

$$\mathbf{u}^{\text{in}}(x, d, q) = \mathbf{u}_p^{\text{in}}(x, d) + \mathbf{u}_s^{\text{in}}(x, d, q)$$

where

$$\mathbf{u}_p^{\text{in}}(x, d) = d \exp(ik_p x \cdot d), \quad \mathbf{u}_s^{\text{in}}(x, d, q) = q \exp(ik_s x \cdot d)$$

with d being the unit incident direction vector and q being the unit polarization vector satisfying $d \cdot q = 0$.

The direct scattering problem for (1.4)-(1.6) is to find the scattered field \mathbf{u}^{sc} when the incident field \mathbf{u}^{in} and density ρ are given. For a real valued $\rho \in C^{1,\gamma}(\mathbb{R}^3)$ with $\text{supp}(1 - \rho) = \overline{D}$, it has been shown in [5, Theorem 5.10] that (1.4)-(1.7) admits a unique solution $\mathbf{u}^{\text{sc}}, \mathbf{u} \in C^2(\mathbb{R}^3 \setminus \overline{D})$.

Remark 1.1. *The well-posedness of the medium scattering problems (1.1)-(1.3) and (1.4)-(1.7) can be established by the integral equation method (see, e.g. [3, 5]). The key ingredients for proving uniqueness are Green's theorem, Rellich's lemma and the unique continuation for elliptic equations. Existence of a solution is derived with the help of the representation theorem which leads to a Fredholm integral equation of Lippmann-Schwinger type for the displacement u or \mathbf{u} . Under the weaker regularity assumption that $n, \rho \in L^\infty(\mathbb{R}^3)$, the well-posedness of the acoustic and elastic medium scattering problem can be established via variational approach together with properties of Dirichlet-to-Neumann operators for the Helmholtz and Navier equations, we refer to [3, Chapter 5.3] and [1] for the details.*

1.3 Inverse problems and literature review

Suppose that the contrast functions $|1 - n|$ and $|1 - \rho|$ is strictly positive in a neighborhood of ∂D in D . In this paper we study the following two *inverse medium scattering problems* with infinitely many plane waves at a fixed energy:

IP1: For a fixed k , determine the support D of the inhomogeneous medium together with $n|_{\partial D}$ from knowledge of $u^\infty(\hat{x}, d)$ for all $\hat{x}, d \in \mathbb{S}^2$.

IP2: For a fixed ω , determine the support D of the inhomogeneous medium together with $\rho|_{\partial D}$ from knowledge of $\mathbf{u}^\infty(\hat{x}, d, q)$ for all $\hat{x}, d, q \in \mathbb{S}^2, d \cdot q = 0$.

If D is an impenetrable obstacle, many uniqueness results have been obtained with one or many incident plane waves at a fixed energy. The first uniqueness result for sound-soft obstacles in acoustics was given in [3] based on the ideas of Schiffer. If two scatterers produce the same far field pattern for all incident directions, Isakov [9] derived a contradiction by considering a series of solutions with a singularity moving towards a point on the boundary of one obstacle but not contained in the other obstacle. Motivated by [9], Kirsch and Kress [3, 10] simplified Isakov's method by using a priori estimate of classical solutions to an integral equation system that is equivalent to the scattering problem. This method also works for sound-hard and Robin-type impenetrable obstacles as well as penetrable ones satisfying interface transmission conditions with a discontinuity condition on the normal derivative (see (1.9) below); see [10]. In linear elasticity it was applied in [6] to get uniqueness in inverse time-harmonic scattering of elastic waves.

If D is a penetrable body, Hähner in [8] introduced a new technique to prove the unique determination of a penetrable, inhomogeneous and anisotropic body from a knowledge of the scattered near fields for all incident plane waves. Hähner's method is based on the existence, uniqueness and regularity of solutions to an interior transmission problem in D . The work [11] uses complex geometrical optics solutions of the Helmholtz equation to prove that a penetrable obstacle with C^2 -smooth boundary in two dimensions can be uniquely reconstructed by near field measurements. The results of [11] was then extended to three dimensions in [12] and to the case that D is penetrable with Lipschitz continuous boundary [13]. Moreover, [14] also uses complex geometrical optics solution of the elastic wave equation to get uniqueness with far-field measurements. There are also other works dealing with the penetrable body with embedded obstacles. It was proved in [15] that D can be determined from knowledge of the acoustic far field pattern for incident plane waves when the refractive index function n is a known constant in D . All the above uniqueness results were obtained on the jump condition for the normal derivatives, i.e.,

$$\partial_\nu u^+ = \gamma \partial_\nu u^- \text{ on } \partial D, \quad \gamma \neq 1. \tag{1.9}$$

In this work we are interested in inverse medium scattering problems with the transmission coefficient $\gamma = 1$ (see (1.9)) both in acoustics and elastics. Physically, the *continuity* of the normal derivatives (in the sense of trace) leads to a weakly scattering effect by the penetrable obstacle, in comparing with the strongly scattering due to the transmission coefficient $\gamma \neq 1$. This perhaps explains why it is difficult to carry out the same a priori estimate as in [10] for singular solutions incited by the fundamental solution as an incoming wave. To overcome this difficulty, [16] proposed a new method to establish uniqueness for recovering the shape of a penetrable D in acoustic case. It is based on the L^p -estimate of hyper-singular solutions and also on the constructing a properly defined and well-posed interior transmission problem in a

small domain inside D . The selection of a small domain ensures that the lowest transmission eigenvalue is large, so that a given wave number k is not an eigenvalue of the constructed interior transmission problem. The same trick has been employed in [17] for inverse elastic scattering problems. The main contribution of this paper is to propose a more direct approach by avoiding a priori estimate as much as possible and without using interior transmission eigenvalue problems. Due to the weakly scattering, it still seems necessary to use hyper-singular solutions. If the refractive index function admits jumps across the boundary, we employ singular total field caused by the first derivative of the fundamental solution when $\gamma = 1$. This direct approach requires a more subtle analysis on the singular behavior of the scattered field at the boundary as the source position approaches to a boundary point from exterior, which has been used by Potthast [18] in acoustic and electromagnetic inhomogeneous medium scattering. In [18], the unknown domain D is assumed to be bounded, uniformly smooth and satisfy an exterior cone condition and the refractive index function is assumed to be Hölder continuous. Based on the lower and upper estimates of the scattered field generated by a multiple, Potthast [18] proved uniqueness for recovering the shape of D . The main contribution of this paper is to calculate the leading part of the asymptotics explicitly in terms of the medium parameters. Consequently, we show that the leading part yields information not only on the position and geometric shape of the scatterer but also on the value of the medium function at the boundary. Such an idea has been early used in [4] for inverse electromagnetic scattering problems in a multi-layered medium in the TM polarization case where $\gamma \neq 1$ is the ratio of the two wavenumbers on both sides of the boundary.

The rest of the paper is organized as follows. In Sections 2, we prove that far field patterns for incident plane waves with all incident directions at a fixed wave number uniquely determine the shape of the penetrable body D in acoustics. In Section 3, the hyper-singular source method will be carried over to inverse elastic scattering by penetrable bodies with the same uniqueness result.

2 Inverse acoustic scattering problem

We briefly introduce the Lippmann-Schwinger equation and the volume potential. We first present a mapping property for the volume potential in Sobolev spaces [3, Theorem 8.2].

Theorem 2.1. *Given two bounded domains D and G , the volume potential*

$$(V\varphi)(x) := \int_D \Phi(x, y)\varphi(y) \, dy, \quad x \in \mathbb{R}^3 \quad (2.1)$$

defines a bounded operator $V : L^2(D) \rightarrow H^2(G)$.

Then the Lippmann-Schwinger equation is defined as

$$u(x) = u^{\text{in}}(x) - k^2 \int_{\mathbb{R}^3} (1 - n(y))\Phi(x, y)u(y) \, dy, \quad x \in \mathbb{R}^3. \quad (2.2)$$

The medium scattering problem (1.1)–(1.3) is equivalent to the problem of solving the above integral equation; see [3, Theorem 8.3].

Next we prove the unique determination of the support D of the inhomogeneous medium from the far-field pattern $u^\infty(\hat{x}, d)$ for all $\hat{x}, d \in \mathbb{S}^2$ by using the Lippmann-Schwinger equation and first derivatives of the fundamental solution for the Helmholtz equation.

Theorem 2.2 (Uniqueness with infinitely many plane waves). *Let $n \in L^\infty(\mathbb{R}^3)$ be real valued. Suppose that $|1 - n(x)| \geq C_0 > 0$ a.e. for $x \in D$ lying in a neighboring area of ∂D . Then the measurement data $u^\infty(\hat{x}, d)$ for all $\hat{x}, d \in \mathbb{S}^2$ uniquely determine ∂D . Furthermore, if $n \in C^{0,\gamma}(\overline{D})$ for some $\gamma \in (0, 1)$, then the value of n on ∂D , that is, $n|_{\partial D}$, can also be uniquely determined by the same measurement data.*

Remark 2.3. *We remark that the determination of ∂D has been shown in [18, Chapter 2] for Hölder continuous refractive index functions.*

Proof. Assume that there are two penetrable obstacles D and D' such that

$$u^\infty(\hat{x}, d) = u'^\infty(\hat{x}, d) \text{ for all } \hat{x}, d \in \mathbb{S}^2. \quad (2.3)$$

If $D \neq D'$, we shall derive a contradiction from the above relation. By Rellich's lemma, $u^{\text{sc}}(x, d) = u'^{\text{sc}}(x, d)$ for all $x \in G, d \in \mathbb{S}^2$, where G denotes the unbounded component of $\mathbb{R}^3 \setminus \overline{D \cup D'}$. Using the mixed reciprocity relation [3, Theorem 3.24] one can deduce that

$$4\pi u^\infty(-d, x) = 4\pi u'^\infty(-d, x) \text{ for all } d \in \mathbb{S}^2, x \in G.$$

Rewriting the above equation with $-d = \hat{x}, x = z$ gives

$$u^\infty(\hat{x}, z) = u'^\infty(\hat{x}, z) \text{ for all } \hat{x} \in \mathbb{S}^2, z \in G.$$

Then we get

$$u^{\text{sc}}(x, z) = u'^{\text{sc}}(x, z) \text{ for all } x, z \in G,$$

due to the Rellich's lemma. This implies that

$$\nabla u^{\text{sc}}(x, z) = \nabla u'^{\text{sc}}(x, z) \text{ for all } x, z \in G. \quad (2.4)$$

Here $u^\infty(\hat{x}, z)$ and $u^{\text{sc}}(x, z)$ denote the far-field patterns and the scattered fields corresponding to a point source generated at $z \in \mathbb{R}^3$, respectively.

Choose $z^* \in \partial D \cap \partial G$ and $z^* \notin \overline{D'}$ (see Figure 1 for the choice of z^*) and let

$$z_j := z^* + \frac{\nu(z^*)}{j} \in \mathbb{R}^3 \setminus \overline{D}, \quad j = 1, 2, \dots,$$

where $\nu(z^*)$ is the unit outer normal at z^* . Define a sequence of incoming point source waves

$$u_j^{\text{in}}(x) = \Phi(x, z_j) := \frac{e^{ik|x-z_j|}}{4\pi|x-z_j|}, \quad x \neq z_j.$$

By straightforward computations, it holds that $u_j^{\text{in}} \in L_{\text{loc}}^2(\mathbb{R}^3)$. Recalling the Lippmann-Schwinger equation that is equivalent to the medium scattering problem, we get the integral equation

$$u(x, z_j) = u_j^{\text{in}}(x) - k^2 \int_{\mathbb{R}^3} (1 - n(y)) \Phi(x, y) u(y, z_j) dy, \quad x \in \mathbb{R}^3. \quad (2.5)$$

Then we define the volume operator $K: L_{\text{loc}}^2(\mathbb{R}^3) \rightarrow H_{\text{loc}}^2(\mathbb{R}^3)$ by

$$(K\varphi)(x) := -k^2 \int_{\mathbb{R}^3} (1 - n(y)) \Phi(x, y) \varphi(y) dy, \quad x \in \mathbb{R}^3 \quad (2.6)$$




Figure 1: Illustration of the location of z^* and the normal direction $\nu(z^*)$.

where $\text{supp}(1 - n) = \overline{D}$. With Theorem 2.1, it holds that $Ku_j^{\text{in}} \in H_{\text{loc}}^2(\mathbb{R}^3)$ if $n \in L^\infty(\mathbb{R}^3)$. By the definition of K , we can reformulate (2.5) as an abstract operator equation

$$(I - K)u(x, z_j) = u_j^{\text{in}}(x), \quad j = 1, 2, \dots.$$

Write $u_j(x) := u(x, z_j)$ for notational simplicity and introduce the functions $w_j := u_j - u_j^{\text{in}}$. It is easy to see that the functions w_j satisfy the integral equation

$$(I - K)w_j = Ku_j^{\text{in}}, \quad j \geq 1.$$

Let Ω be a subdomain of \mathbb{R}^3 containing D and z^* . The embedding of $H^2(\Omega)$ into $L^2(\Omega)$ implies that $I - K: L^2(\Omega) \rightarrow L^2(\Omega)$ is a Fredholm operator with the index zero. By the uniqueness of forward scattering, the integral equation $(I - K)w_j = 0$ has only the trivial solution $w_j = 0$. Then due to the Fredholm alternative, $(I - K)w_j = Ku_j^{\text{in}}$ has a unique solution in $L^2(\Omega)$ such that $\|w_j\|_{L^2(\Omega)} \leq C\|Ku_j^{\text{in}}\|_{L^2(\Omega)}$. Hence

$$\begin{aligned} \|w_j\|_{H^2(\Omega)} &= \|Kw_j + Ku_j^{\text{in}}\|_{H^2(\Omega)} \\ &\leq \|Kw_j\|_{H^2(\Omega)} + \|Ku_j^{\text{in}}\|_{H^2(\Omega)} \\ &\leq C\|w_j\|_{L^2(\Omega)} + \|Ku_j^{\text{in}}\|_{H^2(\Omega)} \\ &\leq C\|Ku_j^{\text{in}}\|_{H^2(\Omega)} + \|Ku_j^{\text{in}}\|_{H^2(\Omega)} \\ &\leq C\|Ku_j^{\text{in}}\|_{H^2(\Omega)}. \end{aligned}$$

Moreover, by the boundedness of operator K , we have

$$\|w_j\|_{C(\Omega)} \leq C\|w_j\|_{H^2(\Omega)} \leq C\|Ku_j^{\text{in}}\|_{H^2(\Omega)} \leq C\|u_j^{\text{in}}\|_{L^2(\Omega)} \leq C. \quad (2.7)$$

Note that the uniform boundedness also follows from the solvability of the integral equation in $C(\Omega)$.

It follows from the Lippmann-Schwinger equation that

$$w_j(x) = -k^2 \int_D (1 - n(y))\Phi(x, y)(w_j(y) + u_j^{\text{in}}(y)) \, dy, \quad x \in \mathbb{R}^3.$$

To prove uniqueness of the inverse problem we shall take the first derivative of w by setting $v_j(x) := \nabla w_j(x) \cdot \nu(z^*)$. Then we can compute $v_j(x)$ as

$$v_j(x) = -k^2 \int_D (1 - n(y)) (\nabla_x \Phi(x, y) \cdot \nu(z^*)) (w_j(y) + u_j^{\text{in}}(y)) \, dy.$$

Dividing $v_j(x)$ into three parts at $x = z_j$ yields

$$\begin{aligned} v_j(z_j) &= -k^2 \int_D (1 - n(y)) (\nabla_x \Phi(z_j, y) \cdot \nu(z^*)) w_j(y) \, dy \\ &\quad - k^2 \int_{D \cap B_\delta(z^*)} (1 - n(y)) (\nabla_x \Phi(z_j, y) \cdot \nu(z^*)) u_j^{\text{in}}(y) \, dy \\ &\quad - k^2 \int_{D \setminus B_\delta(z^*)} (1 - n(y)) (\nabla_x \Phi(z_j, y) \cdot \nu(z^*)) u_j^{\text{in}}(y) \, dy \\ &= I_1(n, w_j) + I_2(n) + I_3(n), \end{aligned}$$

where $\delta > 0$ is sufficiently small. Using (2.7) and the definition of $u_j^{\text{in}}(x)$, we can estimate $I_1(n, w_j)$, $I_2(n)$, $I_3(n)$ as follows:

$$|I_1(n, w_j)| + |I_3(n)| < M \quad \text{for all } j \in \mathbb{N}, \quad (2.8)$$

and

$$\begin{aligned} |I_2(n)| &= \left| -k^2 \int_{D \cap B_\delta(z^*)} (1 - n(y)) (\nabla_x \Phi(z_j, y) \cdot \nu(z^*)) u_j^{\text{in}}(y) \, dy \right| \\ &= \left| k^2 \int_{D \cap B_\delta(z^*)} (1 - n(y)) \frac{e^{ik|z_j - y|}}{4\pi|z_j - y|} \frac{e^{ik|z_j - y|}}{4\pi|z_j - y|^2} \left(\frac{z_j - y}{|z_j - y|} \cdot \nu(z^*) \right) \, dy \right| + \mathcal{O}(1) \\ &\geq k^2 \int_{D \cap B_\delta(z^*)} |1 - n(y)| \frac{\text{Re}(e^{i2k|z_j - y|})}{16\pi^2|z_j - y|^3} \left(\frac{z_j - y}{|z_j - y|} \cdot \nu(z^*) \right) \, dy + \mathcal{O}(1) \\ &\geq \frac{C_0}{2} \frac{k^2}{16\pi^2} \int_{D \cap B_\delta(z^*)} \frac{1}{|z_j - y|^3} \, dy + \mathcal{O}(1) \\ &= C_0 \frac{k^2}{16\pi} \ln(\delta j + 1) + \mathcal{O}(1) \end{aligned} \quad (2.9)$$

as $j \rightarrow \infty$. Note that the last step follows from Lemma 2.4 below and we have used the condition that $|1 - n| \geq C_0$ almost everywhere in $D \cap B_\delta(z^*)$. This together with (2.8) proves that

$$\lim_{j \rightarrow \infty} |v_j(z_j)| = \infty. \quad (2.10)$$

Analogously to the definition of v_j for D , we can define $v'_j(x) = \nabla w'_j(x) \cdot \nu(z^*)$, where $w'_j := u'_j - u_j^{\text{in}}$ denotes the scattered field corresponding to another penetrable obstacle D' . Recalling that $z^* \in \partial D$ and $z^* \notin \overline{D'}$, we obtain from the well-posedness of the forward scattering problem for D' that

$$|\nabla w'_j(x)| + |w'_j(x)| \leq C \quad \text{for all } |x - z_j| < \epsilon, \epsilon > 0.$$

This in particular implies that

$$|v'_j(z_j)| = |\nabla w'_j(z_j) \cdot \nu(z^*)| \leq C, \quad j \rightarrow \infty. \quad (2.11)$$

On the other hand, we have the relation

$$v_j(z_j) = v'_j(z_j) \text{ for all } j \geq 1, \quad (2.12)$$

which contradicts (2.10) and (2.11). This contradiction implies that $D = D'$.

Next, we are going to determine the value of $n(x) \in C^{0,\gamma}(\overline{D})$ on ∂D . Suppose there are two refractive index functions n and n' and assume on the contrary that $n(x) \neq n'(x)$ at some boundary point $x = z^* \in \partial D$. By the continuity, we assume that $|n(x) - n'(x)| \geq c_0 > 0$ for all $x \in D \cap B_\delta(z^*)$ for some $\delta > 0$. Then choose a sequence of incoming point source waves $u_j^{\text{in}}(x)$ with $z_j = z^* + \nu(z^*)/j \in \mathbb{R}^3 \setminus \overline{D}$ ($j = 1, 2, \dots$). Following the uniqueness proof in determining the boundary, we obtain

$$I_1(n, w_j) + I_2(n) + I_3(n) = I_1(n', w'_j) + I_2(n') + I_3(n'),$$

that is

$$I_2(n) - I_2(n') = I_1(n', w'_j) - I_1(n, w_j) + I_3(n') - I_3(n). \quad (2.13)$$

For all $j \in \mathbb{N}$, the right hand of (2.13) is bounded by

$$|I_1(n', w'_j) - I_1(n, w_j) + I_3(n') - I_3(n)| \leq |I_1(n', w'_j)| + |I_1(n, w_j)| + |I_3(n')| + |I_3(n)| < M.$$

While for the left hand of (2.13), the calculations in (2.9) show that

$$\begin{aligned} |I_2(n) - I_2(n')| &= \left| -k^2 \int_{D \cap B_\delta(z^*)} (n'(y) - n(y)) (\nabla_x \Phi(z_j, y) \cdot \nu(z^*)) u_j^{\text{in}}(y) dy \right| \\ &\geq k^2 \int_{D \cap B_\delta(z^*)} |n'(y) - n(y)| \frac{\text{Re}(e^{i2k|z_j - y|})}{16\pi^2 |z_j - y|^3} \left(\frac{z_j - y}{|z_j - y|} \cdot \nu(z^*) \right) dy + \mathcal{O}(1) \\ &\geq \frac{c_0}{2} \frac{k^2}{16\pi^2} \int_{D \cap B_\delta(z^*)} \frac{1}{|z_j - y|^3} dy + \mathcal{O}(1) \\ &= c_0 \frac{k^2}{16\pi} \ln(\delta j + 1) + \mathcal{O}(1) \end{aligned}$$

as $j \rightarrow \infty$, i.e. $\lim_{j \rightarrow \infty} |I_2(n) - I_2(n')| = \infty$. This is a contradiction which implies that $n(x) = n'(x)$ on ∂D . \square

Lemma 2.4. *Suppose that the boundary ∂D is C^2 -smooth and $\delta > 0$ is sufficiently small. A lower bound of the asymptotic behavior of $|I_2|$ can be estimated by*

$$|I_2| \geq C_0 \frac{k^2}{16\pi} \ln(\delta j + 1) + \mathcal{O}(1) \quad \text{as } j \rightarrow \infty,$$

where $C_0 > 0$ is the positive lower bound of $|1 - n|$ for $x \in D$ close to ∂D .

Proof. To compute the integral in $|I_2|$, we fix $z_j = z^* + \nu(z^*)/j$ for some $j \geq 1$ at the origin and assume $\nu(z^*) = (0, 0, -1)$. Introduce the cylindrical coordinate system (r, θ, z) for $y = (y^{(1)}, y^{(2)}, y^{(3)}) \in D \cap B_\delta(z^*)$:

$$\begin{cases} y^{(1)} &= r \cos \theta, \\ y^{(2)} &= r \sin \theta, \\ y^{(3)} &= z, \end{cases}$$

where $r \geq 0$, $0 < \theta \leq 2\pi$ and $z \in \mathbb{R}$. Then we know that

$$dy = dy^{(1)} dy^{(2)} dy^{(3)} = \left| \frac{\partial(y^{(1)}, y^{(2)}, y^{(3)})}{\partial(r, \theta, z)} \right| dr d\theta dz = r dr d\theta dz.$$

Since δ is sufficiently small, we approximate the integral by integrating $1/|z_j - y|^3$ on a half ball centered at z^* under the cylindrical coordinate system (see Figure 2). We observe that r should satisfy $0 < r < \sqrt{\delta^2 - (z - \frac{1}{j})^2}$ for fixed $z \in (\frac{1}{j}, \frac{1}{j} + \delta)$. Hence we have the following estimate




Figure 2: The cross section of domain $D \cap B_\delta(z^*)$ on the oy_2y_3 -plane.

$$\begin{aligned} \int_{D \cap B_\delta(z^*)} \frac{1}{|z_j - y|^3} dy &= \int_{\frac{1}{j}}^{\frac{1}{j} + \delta} \int_0^{2\pi} \int_0^{\sqrt{\delta^2 - (z - \frac{1}{j})^2}} (r^2 + z^2)^{-\frac{3}{2}} r dr d\theta dz + \mathcal{O}(1) \\ &= 2\pi \int_{\frac{1}{j}}^{\frac{1}{j} + \delta} \frac{1}{z} - \frac{1}{\sqrt{\delta^2 - \frac{1}{j^2} + \frac{2}{j}z}} dz + \mathcal{O}(1) \\ &= 2\pi \left(\ln z \Big|_{\frac{1}{j}}^{\frac{1}{j} + \delta} - j \sqrt{\delta^2 - \frac{1}{j^2} + \frac{2}{j}z} \Big|_{\frac{1}{j}}^{\frac{1}{j} + \delta} \right) + \mathcal{O}(1) \\ &= 2\pi \ln(\delta j + 1) - 4\pi\delta \frac{1}{\sqrt{\delta^2 + \frac{1}{j^2} + \frac{2\delta}{j}} + \sqrt{\delta^2 + \frac{1}{j^2}}} + \mathcal{O}(1) \\ &= 2\pi \ln(\delta j + 1) + \mathcal{O}(1). \end{aligned}$$

Now the desired estimate for the lower bound of $|I_2|$ follows from (2.9). \square

Remark 2.5. In Theorem 2.2, we have assumed that the contrast function $1 - n$ is real valued (that is, the medium is non-absorbing) and does not change sign near ∂D . In the case of an

absorbing medium (that is, $1 - n$ is complex valued) and $|\operatorname{Re}(1 - n)| \geq C > 0$ near ∂D , one can get the same uniqueness result. If the refractive index function is continuous at the interface but the absolute value of its higher order derivatives keep a positive distance from zero near ∂D , it is still possible to get uniqueness by using singular solutions incited by higher order derivatives of the fundamental solution. Moreover, we think that, through more delicate analysis, the hyper-singular source method developed here can be used to uniquely determine an analytic refractive index function defined on \overline{D} .

3 Inverse elastic scattering problem

In this section, we first introduce the Lippmann-Schwinger equation for the Navier equation and properties of the fundamental tensor to the elastic wave equation. Then we apply the technique of using hyper-singular sources to obtain similar uniqueness results for inverse elastic scattering from penetrable obstacles. Through out this paper, the symbol \cdot stands for either the product between a matrix and a vector or the inner product between two vectors in \mathbb{R}^3 . All vectors are supposed to be column vectors.

3.1 Fundamental solution and Lippmann-Schwinger equation

We begin with the fundamental solution (or Green's tensor) $\Pi(x, y)$ to the Navier equation $\Delta^* \mathbf{u} + \omega^2 \mathbf{u} = 0$:

$$\Pi(x, y) = \frac{1}{\mu} \Phi_{k_s}(x, y) \mathbf{I} + \frac{1}{\omega^2} \nabla_x \nabla_x^\top [\Phi_{k_s}(x, y) - \Phi_{k_p}(x, y)], \quad x \neq y \quad (3.1)$$

where $\Phi_k(x, y) = \frac{\exp(ik|x-y|)}{4\pi|x-y|}$ ($k = k_p, k_s$) is the fundamental solution to the Helmholtz equation. The symbol \mathbf{I} stands for the 3×3 identity matrix and $'\top'$ denotes the transpose of a matrix or a vector.

Next, we will present some properties of the fundamental solution $\Pi(x, y)$, especially its behavior for $|x - y| \rightarrow 0$. To this end we expand $\exp(ik|x|)/(4\pi|x|)$ to a power series to obtain

$$\begin{aligned} \frac{\exp(ik|x|)}{4\pi|x|} &= \frac{\cos(k|x|)}{4\pi|x|} + i \frac{\sin(k|x|)}{4\pi|x|} \\ &= \frac{1}{4\pi|x|} - \frac{k^2}{4\pi} |x| + k^4 |x|^3 f_1(k^2|x|^2) + ik f_2(k^2|x|^2) \end{aligned} \quad (3.2)$$

with two entire functions f_1 and f_2 .

Letting $r = |x - y| \rightarrow 0$ and inserting the above expression (3.2) into the definition of $\Pi(x, y)$, we obtain

$$\begin{aligned} \Pi(x, y) &= \frac{1}{\mu} \Phi_{k_s}(x, y) \mathbf{I} + \frac{1}{\omega^2} \nabla_x \nabla_x^\top [\Phi_{k_s}(x, y) - \Phi_{k_p}(x, y)] \\ &= \frac{1}{\mu} \frac{1}{4\pi r} \mathbf{I} + \frac{1}{\omega^2} \nabla_x \nabla_x^\top \left(\frac{1}{8\pi} (k_p^2 - k_s^2) |r| \right) + \mathcal{O}(1) \\ &= \frac{1}{\mu} \frac{1}{4\pi r} \mathbf{I} - \frac{\lambda + \mu}{8\pi\mu(\lambda + 2\mu)} \nabla_x \nabla_x^\top (|r|) + \mathcal{O}(1). \end{aligned} \quad (3.3)$$

We denote the primary part of $\Pi(x, y)$ as

$$\Pi^0(x, y) := \frac{1}{\mu} \frac{1}{4\pi r} \mathbf{I} - \frac{\lambda + \mu}{8\pi\mu(\lambda + 2\mu)} \nabla_x \nabla_x^\top (|r|) \quad (3.4)$$

which is known as the Kelvin's matrix [2]. Direct calculations show that

$$\nabla_x r = (x - y)/r, \quad \nabla_x \nabla_x^\top (|r|) = \frac{1}{r} \mathbf{I} - \frac{1}{r} (x \hat{-} y) \otimes (x \hat{-} y) \quad (3.5)$$

where the symbol ' $\hat{\otimes}$ ' represents the tensor product $a \hat{\otimes} a := aa^\top$ for the vector $a \in \mathbb{R}^3$, and $(x \hat{-} y) := (x - y)/|x - y|$. Set $\alpha := \frac{\lambda + 3\mu}{8\pi\mu(\lambda + 2\mu)}$ and $\beta := \frac{\lambda + \mu}{8\pi\mu(\lambda + 2\mu)}$. Then

$$\Pi^0(x, y) = \alpha \frac{1}{r} \mathbf{I} + \beta \frac{1}{r} (x \hat{-} y) \otimes (x \hat{-} y) \quad (3.6)$$

and

$$\Pi(x, y) = \Pi^0(x, y) + \mathcal{O}(1), \quad |x - y| \rightarrow 0 \quad (3.7)$$

which means that $\Pi(x, y)$ and $\Pi^0(x, y)$ have the same singularity of type $1/|x - y|$.

Lemma 3.1. *Let $b \in \mathbb{R}^3$ be a vector. Then it holds that*

(1)

$$\nabla_x (\Pi^0(x, y) \cdot b) = \frac{1}{r^2} [-\alpha (b \otimes \bar{x}) + \beta ((\bar{x} \cdot b) \mathbf{I} + \bar{x} \otimes \nu - 3(\bar{x} \cdot b) \bar{x} \otimes \bar{x})]. \quad (3.8)$$

(2)

$$\begin{aligned} & \nabla_x (\Pi^0(x, y) \cdot b) \cdot (\Pi^0(x, y) \cdot b) \\ &= \frac{1}{r^3} [-\alpha^2 (\bar{x} \cdot b) b + (\alpha\beta - (3\alpha\beta + \beta^2) (\bar{x} \cdot b)^2) \bar{x}]. \end{aligned} \quad (3.9)$$

Here $r = |x - y|$ and $\bar{x} := (x \hat{-} y)$.

(3) *In the special case that $b = x \hat{-} y$, we have*

$$\nabla_x (\Pi^0(x, y) \cdot b) \cdot (\Pi^0(x, y) \cdot b) = \frac{-1}{r^3} (\alpha + \beta)^2 \bar{x}. \quad (3.10)$$

Proof. (1) Set $z := x - y = (z^{(1)}, z^{(2)}, z^{(3)})^\top$ and $\mathbf{A} := \nabla_x \left(\frac{1}{r} \hat{z} \otimes \hat{z} \cdot b \right) \in \mathbb{R}^{3 \times 3}$. Since $\hat{z} \otimes \hat{z} \cdot b = (\hat{z} \cdot b) \hat{z}$, the j -th row of \mathbf{A} takes the form

$$\begin{aligned} \mathbf{A}_j &= \nabla_x \left(\frac{z^{(j)}}{r^3} (z \cdot b) \right) \\ &= (z \cdot b) \nabla_x \left(\frac{z^{(j)}}{r^3} \right) + \frac{z^{(j)}}{r^3} b^\top \\ &= (z \cdot b) \frac{1}{r^3} e_j^\top - (z \cdot b) \frac{3z^{(j)}}{r^5} z^\top + \frac{z^{(j)}}{r^3} b^\top. \end{aligned} \quad (3.11)$$

Here $\{e_j\}_{j=1}^3$ denote the unit vectors in \mathbb{R}^3 in the Cartesian coordinates. From the expression of \mathbf{A}_j we can derive

$$\mathbf{A} = \frac{1}{r^3} (z \cdot b) \mathbf{I} + \frac{1}{r^3} z \otimes b - \frac{3}{r^5} (z \cdot b) z \otimes z. \quad (3.12)$$

Substituting (3.12) into $\Pi^0(x, y)$ gives

$$\nabla_x (\Pi^0(x, y) \cdot b) = -\frac{\alpha}{r^3} b \otimes z + \beta \left[\frac{1}{r^3} (z \cdot b) \mathbf{I} + \frac{1}{r^3} z \otimes b - \frac{3}{r^5} (z \cdot b) z \otimes z \right]. \quad (3.13)$$

The proof is now completed by replacing z with $x - y$.

(2) Recalling (3.8), we know

$$\nabla_x (\Pi^0(x, y) \cdot b) = \frac{1}{r^2} [-\alpha (b \otimes \bar{x}) + \beta ((\bar{x} \cdot b) \mathbf{I} + \bar{x} \otimes b - 3(\bar{x} \cdot b) \bar{x} \otimes \bar{x})] \quad (3.14)$$

and

$$\Pi^0(x, y) \cdot b = \frac{1}{r} [\alpha \mathbf{I} + \beta \bar{x} \otimes \bar{x}] \cdot b \quad (3.15)$$

Now, taking the product between (3.14) and (3.15) yields

$$\begin{aligned} & \nabla_x (\Pi^0(x, y) \cdot b) \cdot (\Pi^0(x, y) \cdot b) \\ &= \frac{1}{r^3} [-\alpha(\alpha + \beta) (b \otimes \bar{x}) \cdot b - \beta^2 (\bar{x} \cdot b) (\bar{x} \otimes \bar{x}) \cdot b] \\ & \quad + \frac{\alpha\beta}{r^3} [(\bar{x} \cdot b) b + \bar{x} - 3(\bar{x} \cdot b) (\bar{x} \otimes \bar{x}) \cdot b]. \end{aligned} \quad (3.16)$$

The proof is finished by applying identities

$$\begin{aligned} (\nu \otimes \bar{x}) \cdot (\bar{x} \otimes \bar{x}) \cdot \nu &= \nu \cdot \bar{x}^\top \cdot \bar{x} \cdot \bar{x}^\top \cdot \nu = (\nu \otimes \bar{x}) \cdot \nu, \\ (\bar{x} \otimes \nu) \cdot (\bar{x} \otimes \bar{x}) \cdot \nu &= (\bar{x} \cdot \nu) (\bar{x} \otimes \bar{x}) \cdot \nu, \\ (\nu \otimes \bar{x}) \cdot \nu &= (\bar{x} \cdot \nu) \nu, \\ (\bar{x} \otimes \bar{x}) \cdot \nu &= (\bar{x} \cdot \nu) \bar{x}. \end{aligned}$$

(3) If $b = x \hat{=} y$, then using the results of the second assertion we have

$$\begin{aligned} & \nabla_x (\Pi^0(x, y) \cdot b) \cdot (\Pi^0(x, y) \cdot b) \\ &= \frac{1}{r^3} [-\alpha^2 (\bar{x} \cdot b) b + (\alpha\beta - (3\alpha\beta + \beta^2) (\bar{x} \cdot b)^2) \bar{x}] \\ &= \frac{1}{r^3} [-\alpha^2 b + (\alpha\beta - (3\alpha\beta + \beta^2)) \bar{x}] \\ &= \frac{-1}{r^3} (\alpha + \beta)^2 \bar{x}. \end{aligned} \quad (3.17)$$

□

At last, we introduce the Lippmann-Schwinger equation that is equivalent to the scattering problem (1.4)-(1.6). For this purpose we need to consider the volume potential

$$(V\varphi)(x) := \int_D \Pi(x, y) \varphi(y) \, dy, \quad x \in \mathbb{R}^3. \quad (3.18)$$

Below we show a classical mapping property property of V in Hölder spaces, the proof of which can be found in [5, Theorem 5.6].

Theorem 3.2. *Let $D \subset B_R$, where B_R is a ball centered at origin with radius R . If $\varphi \in [C^{0,\gamma}(D)]^3$ with $\gamma \in (0, 1)$, then the volume potential $V\varphi \in [C^2(B_R)]^3$.*

By using Lax's Theorem [3, Theorem 3.5], one can deduce a mapping property for the volume potential in Sobolev spaces based on the result of Theorem 3.2.

Lemma 3.3. *The volume potential defines a bounded operator $V : [L^2(D)]^3 \rightarrow [H^2(B_R)]^3$.*

The Lippmann-Schwinger equation is defined as

$$\mathbf{u}(x) = \mathbf{u}^{\text{in}}(x) - \omega^2 \int_{\mathbb{R}^3} (1 - \rho(y)) \Pi(x, y) \mathbf{u}(y) \, dy, \quad x \in \mathbb{R}^3. \quad (3.19)$$

It was proved in [5, Lemma 5.7] that a solution of the above Lippmann-Schwinger integral equation is a solution to the scattering problem (1.4)-(1.6) and vice versa. Then for $\rho \in [L^\infty(\mathbb{R}^3)]^3$ satisfying $\text{supp}(1 - \rho) = \overline{D}$, it's easy to show that the scattering problem (1.4)-(1.7) has a unique solution $\mathbf{u}^{\text{sc}} \in [H_{\text{loc}}^2(\mathbb{R}^3)]^3$.

3.2 Uniqueness for inverse elastic scattering

Given an incident plane wave $\mathbf{u}^{\text{in}}(x, d, q)$, we will use the notations $\mathbf{u}^{\text{sc}}(x, d, q)$, $u(x, d, q)$ and $\mathbf{u}^\infty(\hat{x}, d, q)$ to indicate dependence of the scattered field, of the total field and of the far field pattern on the incident direction d and polarization q , respectively. For an incoming point source $\mathbf{w}^{\text{in}}(x, y, a) = \Pi(x, y) \cdot a$ located at $y \in D^c$ with the unit vector $a \in \mathbb{C}^3$, we denote the scattered field by $\mathbf{w}^{\text{sc}}(x, y, a)$, the total field by $\mathbf{w}(x, y, a)$, and the far field pattern of the scattered field by $\mathbf{w}^\infty(\hat{x}, y, a)$. We refer to [17, Theorem 2.1] for a mixed reciprocity relation that is analogous to the acoustic case as follows.

Lemma 3.4. *For any $y \in D^c$ and a fixed unit vector a , we have*

$$q \cdot \mathbf{w}^\infty(-d, y, a) + d \cdot \mathbf{w}^\infty(-d, y, a) = a \cdot \mathbf{u}^{\text{sc}}(y, d, q) \quad (3.20)$$

for all $d, q \in \mathbb{S}^2$, $d \cdot q = 0$.

Now we state and prove the uniqueness result in recovering the shape of the inhomogeneous penetrable obstacle D in linear elasticity.

Theorem 3.5. *Suppose $\rho \in [L^\infty(\mathbb{R}^3)]^3$ is a real-valued function such that $|1 - \rho| \geq C$ for some constant $C > 0$ and for $x \in D$ lying in a neighboring area of ∂D . Then the measurement data $\mathbf{u}^\infty(\hat{x}, d, q)$ for all $\hat{x}, d, q \in \mathbb{S}^2$ can uniquely determine ∂D . Furthermore, if $\rho(x) \in [C^{0,\gamma}(\overline{D})]^3$ for some $\gamma \in (0, 1)$, then the value of $\rho(x)$ on ∂D can be also uniquely determined by the same measurement data.*

Proof. Assume that (D, ρ) and $(\tilde{D}, \tilde{\rho})$ generate the same far field patterns, that is,

$$\mathbf{u}^\infty(\hat{x}, d, q) = \tilde{\mathbf{u}}^\infty(\hat{x}, d, q) \quad \text{for all } \hat{x}, d, q \in \mathbb{S}^2, \quad d \cdot q = 0. \quad (3.21)$$

Then by using Rellich's lemma, we obtain

$$\mathbf{u}^{\text{sc}}(x, d, q) = \tilde{\mathbf{u}}^{\text{sc}}(x, d, q) \quad \text{for all } x \in G, \quad d, q \in \mathbb{S}^2, \quad d \cdot q = 0, \quad (3.22)$$

where G is denoted as the unbounded connected domain of $\mathbb{R}^3 \setminus \overline{D \cup \tilde{D}}$. Applying the above mixed reciprocity relation theorem, we can deduce that for any fixed vector a

$$\mathbf{w}^\infty(-d, x, a) = \tilde{\mathbf{w}}^\infty(-d, x, a) \quad \text{for all } x \in G, d \in \mathbb{S}^2. \quad (3.23)$$

Using Rellich's lemma again, we get

$$\mathbf{w}^{\text{sc}}(y, x, a) = \tilde{\mathbf{w}}^{\text{sc}}(y, x, a) \quad \text{for all } x, y \in G. \quad (3.24)$$

If $D \neq \tilde{D}$, we suppose without loss of generality that there exists a point z^* satisfying $z^* \in \partial D \cap \partial G$ and $z^* \notin \tilde{D}$. As done in the acoustic case, we set

$$z_j = z^* + \frac{\nu(z^*)}{j}, \quad j = 1, 2, \dots$$

where $\nu(z^*)$ is the unit normal at z^* directed into the exterior of D . Apparently, $z_j \in G$ for $j \geq j_0$. Next, we consider the scattering problems due to point sources $\mathbf{w}_j^{\text{in}}(x, z_j, \nu(z^*)) = \Pi(x, z_j) \cdot \nu(z^*)$. It's easy to obtain $w_j^{\text{in}} \in [L_{\text{loc}}^2(\mathbb{R}^3)]^3$ from the singularity of $\Pi(x, y)$. The total field $\mathbf{w}(x, z_j, \nu(z^*))$ can be given by the Lippmann-Schwinger equation [5], that is

$$\mathbf{w}(x, z_j, \nu(z^*)) = w_j^{\text{in}}(x, z_j, \nu(z^*)) - \omega^2 \int_{\mathbb{R}^3} (1 - \rho(y)) \Pi(x, y) \mathbf{w}(y, z_j, \nu(z^*)) \, dy, \quad x \in \mathbb{R}^3. \quad (3.25)$$

Then

$$\mathbf{w}_j^{\text{sc}}(x, z_j, \nu(z^*)) = -\omega^2 \int_{\mathbb{R}^3} (1 - \rho(y)) \Pi(x, y) (\mathbf{w}_j^{\text{in}}(x, z_j, \nu(z^*)) + \mathbf{w}_j^{\text{sc}}(x, z_j, \nu(z^*))) \, dy, \quad x \in \mathbb{R}^3. \quad (3.26)$$

Define the volume potential $V : [L_{\text{loc}}^2(\mathbb{R}^3)]^3 \rightarrow [H_{\text{loc}}^2(\mathbb{R}^3)]^3$

$$(V\varphi)(x) = -\omega^2 \int_{\mathbb{R}^3} (1 - \rho(y)) \Pi(x, y) \varphi(y) \, dy, \quad x \in \mathbb{R}^3. \quad (3.27)$$

We can rewrite the equation (3.26) as

$$(I - V)\mathbf{w}_j^{\text{sc}}(x, z_j, \nu(z^*)) = V\mathbf{w}_j^{\text{in}}(x, z_j, \nu(z^*)), \quad x \in \mathbb{R}^3, j = 1, 2, \dots. \quad (3.28)$$

Together with the facts $\text{supp}(1 - \rho) = \overline{D} \subset B_R$, $\rho(x) \in [L^\infty(\mathbb{R}^3)]^3$ and Lemma 3.3, we get a similar estimate to (2.7) as

$$\|\mathbf{w}_j^{\text{sc}}\|_{[C(B_R)]^3} \leq C_1 \|\mathbf{w}_j^{\text{sc}}\|_{[H^2(B_R)]^3} \leq C_2 \|V\mathbf{w}_j^{\text{in}}\|_{[H^2(B_R)]^3} \leq C_3 \|\mathbf{w}_j^{\text{in}}\|_{[L^2(B_R)]^3} \leq C_4 \quad (3.29)$$

where B_R is a ball with radius R containing D and z^* .

Define $\mathbf{v}_j(x) := \nabla_x [\mathbf{w}_j^{\text{sc}}(x, z_j, \nu(z^*)) \cdot \nu(z^*)]$. Then,

$$\mathbf{v}_j(x) = -\omega^2 \int_D (1 - \rho(y)) \nabla_x [\Pi(x, y) \cdot \nu(z^*)] (\mathbf{w}_j^{\text{in}}(y, z_j, \nu(z^*)) + \mathbf{w}_j^{\text{sc}}(y, z_j, \nu(z^*))) \, dy. \quad (3.30)$$

We continue to calculate $\mathbf{v}_j(x)$ at $x = z_j$ by

$$\begin{aligned} \mathbf{v}_j(z_j) &= -\omega^2 \int_D (1 - \rho(y)) \nabla_x [\Pi(x, y) \cdot \nu(z^*)] |_{x=z_j} \mathbf{w}_j^{\text{sc}}(y, z_j, \nu(z^*)) \, dy \\ &\quad - \omega^2 \int_{D \cap B_\delta(z^*)} (1 - \rho(y)) \nabla_x [\Pi(x, y) \cdot \nu(z^*)] |_{x=z_j} \mathbf{w}_j^{\text{in}}(y, z_j, \nu(z^*)) \, dy \\ &\quad - \omega^2 \int_{D \setminus B_\delta(z^*)} (1 - \rho(y)) \nabla_x [\Pi(x, y) \cdot \nu(z^*)] |_{x=z_j} \mathbf{w}_j^{\text{in}}(y, z_j, \nu(z^*)) \, dy \\ &:= I_1(\rho, \mathbf{w}_j^{\text{sc}}) + I_2(\rho) + I_3(\rho), \quad x \in \mathbb{R}^3, \end{aligned} \quad (3.31)$$

where $B_\delta(z^*)$ is a small ball centered at z^* with a small radius $\delta > 0$.

Combining equation (3.7) and Lemma 3.1 we conclude that $\nabla_x[\Pi(x, y) \cdot \nu(z^*)]$ has a singularity of type $1/|x - y|^2$. This together with the fact that $\mathbf{w}_j^{\text{in}} \in [L_{\text{loc}}^2(\mathbb{R}^3)]^3$ and (3.29) gives

$$|\nu(z^*) \cdot I_1(\rho, \mathbf{w}_j^{\text{sc}})| + |\nu(z^*) \cdot I_3(\rho)| \leq M, \quad (3.32)$$

where M is a positive constant independent of $j \in \mathbb{N}$.

From (3.7) it follows that $\mathbf{w}_j^{\text{in}}(y, z_j, \nu(z^*)) = \Pi^0(y, z_j) \cdot \nu(z^*) + \mathcal{O}(1)$. Hence,

$$I_2(\rho) = -\omega^2 \int_{D \cap B_\delta(z^*)} (1 - \rho(y)) \nabla_x[\Pi^0(x, y) \cdot \nu(z^*)]|_{x=z_j} \cdot (\Pi^0(y, z_j) \cdot \nu(z^*)) \, dy + \mathcal{O}(1). \quad (3.33)$$

Denote $\bar{z} = (z_j - y)/|z_j - y|$. Using Lemma 3.1 one can show

$$\begin{aligned} & |\nu(z^*) \cdot I_2(\rho)| \\ &= \left| -\omega^2 \int_{D \cap B_\delta(z^*)} (1 - \rho(y)) [\nabla_x[\Pi^0(x, y) \cdot \nu(z^*)]|_{x=z_j} \cdot (\Pi^0(y, z_j) \cdot \nu(z^*)) \cdot \nu(z^*) \, dy \right| + \mathcal{O}(1) \\ &= \omega^2 \left| \int_{D \cap B_\delta(z^*)} \frac{(1 - \rho(y))}{|z_j - y|^3} [-\alpha^2(\bar{z} \cdot \nu(z^*)) + (\alpha\beta - (3\alpha\beta + \beta^2)(\bar{z} \cdot \nu(z^*))^2) \bar{z} \cdot \nu(z^*)] \, dy \right| + \mathcal{O}(1). \end{aligned} \quad (3.34)$$

Since $\bar{z} \cdot \nu(z^*) \rightarrow 1$ as $j \rightarrow \infty$, it is easy to see

$$[-\alpha^2(\bar{z} \cdot \nu(z^*)) + (\alpha\beta - (3\alpha\beta + \beta^2)(\bar{z} \cdot \nu(z^*))^2) \bar{z} \cdot \nu(z^*)] \rightarrow -(\alpha + \beta)^2, \quad j \rightarrow \infty.$$

Recalling that $\alpha = \frac{\lambda + 3\mu}{8\pi\mu(\lambda + 2\mu)}$ and $\beta = \frac{\lambda + \mu}{8\pi\mu(\lambda + 2\mu)}$, we get $\alpha + \beta = \frac{1}{4\pi\mu}$. Then

$$\begin{aligned} & |\nu(z^*) \cdot I_2(\rho)| \\ & \geq C\omega^2 \left| \int_{D \cap B_\delta(z^*)} \frac{1}{|z_j - y|^3} [-(\alpha + \beta)^2] \, dy \right| + \mathcal{O}(1) \\ & \geq \frac{C\omega^2}{16\pi^2\mu^2} \left| \int_{D \cap B_\delta(z^*)} \frac{1}{|z_j - y|^3} \, dy \right| + \mathcal{O}(1) \\ & = \frac{C\omega^2}{8\pi\mu^2} \ln j + \mathcal{O}(1), \quad j \rightarrow \infty. \end{aligned} \quad (3.35)$$

Here C means the lower bound of the contrast function $|1 - \rho|$. This implies that $|\nu(z^*) \cdot \mathbf{v}_j(z_j)| \rightarrow \infty$ as $j \rightarrow \infty$.

On the other hand, we can define $\tilde{\mathbf{v}}_j(x) := \nabla_x[\tilde{\mathbf{w}}_j^{\text{sc}}(x, z_j, \nu(z^*)) \cdot \nu(z^*)]$, where $\tilde{\mathbf{w}}_j^{\text{sc}}(x, z_j, \nu(z^*))$ is the scattered field corresponding to \tilde{D} and the incident point source $\mathbf{w}_j^{\text{in}}(x, z_j, \nu(z^*))$. Since $z^* \in \partial D$ and $z^* \notin \tilde{D}$, it follows from the well-posedness of the direct scattering problem that

$$|\nu(z^*) \cdot \nabla_x[\tilde{\mathbf{w}}_j^{\text{sc}}(x, z_j, \nu(z^*)) \cdot \nu(z^*)]| + |\nu(z^*) \cdot \tilde{\mathbf{w}}_j^{\text{sc}}(x, z_j, \nu(z^*)) \cdot \nu(z^*)| \leq M, \quad x \in B_\delta(z_j). \quad (3.36)$$

Here M is a constant and $B_\delta(z_j)$ is a ball centered at z_j outside of \tilde{D} . The above inequality implies that $|\nu(z^*) \cdot \tilde{\mathbf{v}}_j(z_j)| \leq M$ as $j \rightarrow \infty$. Recalling from the relation (3.24) that $\mathbf{v}_j(z_j) = \tilde{\mathbf{v}}_j(z_j)$ for all $j \in \mathbb{N}$, we get a contradiction here, which finishes the proof of $D = \tilde{D}$.

Next, we are going to determine the value of $\rho(x) \in [C^{0,\gamma}(\overline{D})]^3$ on ∂D . Assume on the contrary that $\rho(z^*) \neq \tilde{\rho}(z^*)$ at some boundary point $z^* \in \partial D$. We assume further that $|\rho(z^*) - \tilde{\rho}(z^*)| \geq c > 0$ for $x \in D \cap B_\delta(z^*)$ and with some $\delta > 0$. Then choose a sequence of incoming point source waves $u_j^{\text{in}}(x)$ with $z_j = z^* + \nu(z^*)/j \in \mathbb{R}^3 \setminus \overline{D}$ ($j = 1, 2, \dots$). Following the above processes, we obtain that

$$I_1(\rho, \mathbf{w}_j^{\text{sc}}) + I_2(\rho) + I_3(\rho) = I_1(\tilde{\rho}, \tilde{\mathbf{w}}_j^{\text{sc}}) + I_2(\tilde{\rho}) + I_3(\tilde{\rho}),$$

that is,

$$\nu(z^*) \cdot [I_2(\rho) - I_2(\tilde{\rho})] = \nu(z^*) \cdot [I_1(\tilde{\rho}, \tilde{\mathbf{w}}_j^{\text{sc}}) - I_1(\rho, \mathbf{w}_j^{\text{sc}}) + I_3(\tilde{\rho}) - I_3(\rho)]. \quad (3.37)$$

The right hand of (3.37) can be bounded by

$$|\nu(z^*) \cdot I_1(\tilde{\rho}, \tilde{\mathbf{w}}_j^{\text{sc}})| + |\nu(z^*) \cdot I_1(\rho, \mathbf{w}_j^{\text{sc}})| + |\nu(z^*) \cdot I_3(\tilde{\rho})| + |\nu(z^*) \cdot I_3(\rho)| < M, \quad \forall j \in \mathbb{N}.$$

While for the left hand of (3.37), repeating the calculations in (3.33)–(3.35) one deduces that

$$\begin{aligned} & |\nu(z^*) \cdot I_2(\rho) - \nu(z^*) \cdot I_2(\tilde{\rho})| \\ &= \left| -\omega^2 \int_{D \cap B_\delta(z^*)} (\tilde{\rho}(y) - \rho(y)) \nabla_x [\Pi^0(x, y) \cdot \nu(z^*)] \Big|_{x=z_j} \cdot (\Pi^0(y, z_j) \cdot \nu(z^*)) \, dy \right| + \mathcal{O}(1) \\ &\geq \frac{c\omega^2}{16\pi^2\mu^2} \left| \int_{D \cap B_\delta(z^*)} \frac{1}{|z_j - y|^3} \, dy \right| + \mathcal{O}(1) = \frac{c\omega^2}{8\pi\mu^2} \ln j + \mathcal{O}(1), \end{aligned}$$

as $j \rightarrow \infty$, i.e. $\lim_{j \rightarrow \infty} |\nu(z^*) \cdot I_2(\rho) - \nu(z^*) \cdot I_2(\tilde{\rho})| = \infty$. This contradiction implies that $\rho(x) = \tilde{\rho}(x)$ on ∂D . \square

4 Acknowledgments

The work of G. Hu is partially supported by the National Natural Science Foundation of China (No. 12071236) and the Fundamental Research Funds for Central Universities in China (No. 63233071). The work of J. Xiang is supported by the Natural Science Foundation of China (No. 12301542), the Natural Science Foundation of Hubei (No. 2022CFB725) and the Natural Science Foundation of Yichang (No. A23-2-027).

References

- [1] G. Bao, G. Hu, J. Sun and T. Yin, Direct and inverse elastic scattering from anisotropic media, *J. Math. Pures Appl.* 117 (2018): 263-301.
- [2] V. D. Kupradze, Three-dimensional problems of the mathematical theory of elasticity and thermo elasticity, North-Holland, Amsterdam, (1979).
- [3] D. Colton and R. Kress, Inverse acoustic and electromagnetic scattering theory (4th Edition), Springer, Berlin, (2019).
- [4] J. Elschner and G. Hu, Uniqueness in inverse transmission scattering problems for multi-layered obstacles, *Inverse Problems and Imaging*, **5** (2011), 793-813.

- [5] P. Hähner, On acoustic, electromagnetic and elastic scattering problems in inhomogeneous media, Habilitation Thesis, (1998).
- [6] P. Hähner and G. C. Hsiao, Uniqueness theorems in inverse obstacle scattering of elastic waves, *Inverse Problems* **9** (1993), 525-534.
- [7] P. Hähner, A uniqueness theorem in inverse scattering of elastic waves, *IMA J. Appl. Math.* **51** (1993), 201-215.
- [8] P. Hähner, On the uniqueness of the shape of a penetrable anisotropic obstacle, *J. Comput. Appl. Math.* **116** (2000), 167-180.
- [9] V. Isakov, On uniqueness in the inverse transmission scattering problem, *Partial Differential Equations* **15** (1990), 1565-1587.
- [10] A. Kirsch and R. Kress, Uniqueness in inverse obstacle scattering (acoustics), *Inverse Problems* **9** (1993), 285-299.
- [11] S. Nagayasu, G. Uhlmann and J.N. Wang, Reconstruction of penetrable obstacles in acoustic scattering, *SIAM J. Math. Anal.* **43** (2011), 189-211.
- [12] K. Yoshida, Reconstruction of a penetrable obstacle by complex spherical waves, *J. Math. Anal. Appl.* **49** (2010), 645-C657.
- [13] M. Sini and K. Yoshida, On the reconstruction of interfaces using complex geometrical optics solutions for the acoustic case, *Inverse Probl.* **28** (2012), 055013.
- [14] M. Kar and M. Sini, Reconstruction of interfaces from the elastic farfield measurements using CGO solutions, *SIAM J. Math. Anal.* **46** (4) (2014), 2650-2691.
- [15] X. Liu, B. Zhang, Direct and inverse obstacle scattering problems in a piecewise homogeneous medium, *SIAM J. Appl. Math.* **70** (2010), 3105-3120.
- [16] J. Yang, B. Zhang, and H. Zhang, Uniqueness in inverse acoustic and electromagnetic scattering by penetrable obstacles with embedded objects, *J. Differ. Equ.* **265** (2017), 6352-6383.
- [17] J. Xiang and G. Yan, The uniqueness of the inverse elastic wave scattering problem based on the mixed reciprocity relation, *Inverse Problems and Imaging.* **15** (2021), 539-554.
- [18] R. Potthast, Point sources and multipoles in inverse scattering theory, Chapman and Hall/CRC, (2001).