Four-fifths laws in incompressible and magnetized fluids: Helicity, Energy and Cross-helicity

Yulin Ye^{*}, Yanqing Wang[†] and Otto Chkhetiani[‡]

Abstract

In this paper, we are concerned with the Kolmogorov's scaling laws of conserved quantities. By means of Eyink's longitudinal structure functions and the analysis of interaction of different physical quantities, we extend celebrated four-fifths laws from energy to helicity in incompressible fluid and, energy and cross-helicity in magneto-hydrodynamic flow. In contrast to pervious 4/5 laws of energy and cross-helicity in magnetized fluids obtained by Politano and Pouquet, they are in terms of the mixed three-order structure functions rather than the structure coupling correlation functions.

MSC(2020): 76F02, 35Q31, 76B03, 76W05

Keywords: 4/5 law; helicity; energy; cross-helicity; incompressible magnetohydrodynamic fluid

1 Introduction

The fluid contribution of Kolmogorov in [31–33] (the so-called K41 theory) is a milestone in the context of turbulence. In a seminal paper [33], under the homogeneity, isotropy and the finiteness of the energy dissipation, Kolmogorov presented the following four-fifths law in terms of third-order longitudinal structure function

$$\langle [\delta \boldsymbol{v}_L(\boldsymbol{r})]^3 \rangle = -\frac{4}{5} \epsilon \boldsymbol{r},$$
 (1.1)

where ϵ is the mean rate of kinetic energy dissipation per unit mass of the Navier-Stokes equations with sufficiently high Reynolds numbers. Here, $\delta \boldsymbol{v}_L(\boldsymbol{r}) = \delta \boldsymbol{v}(\boldsymbol{r}) \cdot \frac{\boldsymbol{r}}{|\boldsymbol{r}|} = [\boldsymbol{v}(x+\boldsymbol{r}) - \boldsymbol{v}(x)] \cdot \frac{\boldsymbol{r}}{|\boldsymbol{r}|}$ stands for the longitudinal velocity increment and $\langle \cdot \rangle$ denotes the mean value. The derivation of (1.1) in [33] relies on the famous Kármárth-Howarth equations in [30]. The Kolmogorov' 4/5 law (1.1) has attracted considerable attention in turbulence theory, experiment and numerical simulation (see e.g. [1, 2, 4, 11, 16, 20–22, 25, 28, 37, 38, 43, 44, 53]). The modern rigorous derivation of Kolmogorov' 4/5 law (1.1) can be found in [20–22].

^{*}School of Mathematics and Statistics, Henan University, Kaifeng, 475004, P. R. China. Email: ylye@vip.henu.edu.cn

[†]Corresponding author. College of Mathematics and Information Science, Zhengzhou University of Light Industry, Zhengzhou, Henan 450002, P. R. China Email: wangyanqing20056@gmail.com

[‡]A. M. Obukhov Institute of Atmospheric Physics, Russian Academy of Sciences, Pyzhevsky per. 3, Moscow, 119017, Russia. Email: ochkheti@ifaran.ru

In particular, Eyink derived the local version of Kolmogorov's 4/5 law (1.1) below without assumptions of homogeneity and isotropy,

$$S_L(\boldsymbol{v}) = -\frac{4}{5}D(\boldsymbol{v}), \qquad (1.2)$$

where

$$S_L(\boldsymbol{v}) = \lim_{\lambda \to 0} S_L(\boldsymbol{v}, \lambda) = \lim_{\lambda \to 0} \frac{1}{\lambda} \int_{\partial B} \boldsymbol{\ell} \cdot \delta \boldsymbol{v}(\lambda \boldsymbol{\ell}) [\delta \boldsymbol{v}_L(\lambda \boldsymbol{\ell})]^2 \frac{d\sigma(\boldsymbol{\ell})}{4\pi}$$

and

$$D(oldsymbol{v}) = \lim_{arepsilon
ightarrow 0} rac{1}{4} \int_{\mathbb{T}^3}
abla arphi_arepsilon(oldsymbol{\ell}) \cdot \delta oldsymbol{v}(oldsymbol{\ell}) [\delta oldsymbol{v}_L(oldsymbol{\ell})]^2 doldsymbol{\ell},$$

in [20]. Here, $\sigma(x)$ stands for the surface measure on the sphere $\partial B = \{x \in \mathbb{R}^3 : |x| = 1\}$ and φ is some smooth non-negative function supported in \mathbb{T}^3 with unit integral and $\varphi_{\varepsilon}(x) = \varepsilon^{-3}\varphi(\frac{x}{\varepsilon})$. To this end, Eyink defined the longitudinal and transverse velocity increments as follows

$$\delta \boldsymbol{v}_L(\boldsymbol{x},t,\boldsymbol{\ell}) = (\hat{\boldsymbol{\ell}} \otimes \hat{\boldsymbol{\ell}}) \delta \boldsymbol{v}(\boldsymbol{x},t,\boldsymbol{\ell}), \quad \delta \boldsymbol{v}_T(\boldsymbol{x},t,\boldsymbol{\ell}) = (\mathbf{1} - \hat{\boldsymbol{\ell}} \otimes \hat{\boldsymbol{\ell}}) \delta \boldsymbol{v}(\boldsymbol{x},t,\boldsymbol{\ell}), \quad (1.3)$$

and invoked the dissipation term D(v) introduced by Duchon and Robert in [18]. It is worth pointing out that Duchon and Robert also deduced the following 4/3 law

$$S(\boldsymbol{v}) = -\frac{4}{3}D(\boldsymbol{v})$$

for the energy of the Euler equations, which corresponds to

$$\langle \delta \boldsymbol{v}_L(\boldsymbol{r}) [\delta \boldsymbol{v}(\boldsymbol{r})]^2 \rangle = -\frac{4}{3} \epsilon \boldsymbol{r}.$$
 (1.4)

This kind of four-thirds law originated from Yaglom's work [43, 51] and was developed in [43]. The dissipation term D(v) was applied by Duchon and Robert to study the Onsager conjecture concerning the minimal regularity of weak solutions for keeping energy conservation of the Euler equations in [18]. In this direction, regarding the energy conservation of weak solutions for the turbulence models, we refer the readers to [3, 9, 13– 15, 18, 19, 23, 26, 29, 45, 50].

The character of turbulence rests on not only the non-linear cascades of energy, but also the conserved quantities in three dimensions (see [1]). As known to all, besides the energy $\frac{1}{2} \int_{\mathbb{T}^3} |\boldsymbol{v}|^2 dx$, the helicity $\frac{1}{2} \int_{\mathbb{T}^3} \boldsymbol{\omega} \cdot \boldsymbol{v} \, dx$ is another quadratic conserved quantity, where the vorticity $\boldsymbol{\omega} = \nabla \times \boldsymbol{v}$. The concept of helicity was first proposed by Moffatt in [42], which is important at a fundamental level in relation to flow kinematics because it admits topological interpretation in relation to the linkage or linkages of vortex lines of the flow. The interaction of the transfer of energy and helicity in incompressible flow was considered by Chen, Chen and Eyink in [7], and, Chen, Chen, Eyink and Holm in [8]. For this second invariant quantity, its first exact law was due to Chkhetiani in [10], where the derivation yielded the so-called 2/15 law

$$\langle \delta \boldsymbol{v}_L(\boldsymbol{x}+\boldsymbol{r})(\boldsymbol{v}_T(\boldsymbol{x}+\boldsymbol{r}) \times \boldsymbol{v}_T(\boldsymbol{x})) \rangle = \frac{2}{15}h\boldsymbol{r}^2 \epsilon_H, \quad \boldsymbol{v}_T = \boldsymbol{v} - \boldsymbol{v}_L,$$
 (1.5)

where ϵ_H represents the helicity dissipation rates. For the 2/15-law (1.5), see also [34, 35]. Subsequently, Gomez, Plitano and Pouquet [27] presented the following Yaglom type fourthirds law of helicity via the third-order structure function

$$\langle \delta \boldsymbol{v}_L(\mathbf{r}) [\delta \boldsymbol{v}(\boldsymbol{r}) \cdot \delta \boldsymbol{\omega}(\boldsymbol{r}) | \rangle - \frac{1}{2} \langle \delta \boldsymbol{\omega}_L(\boldsymbol{r}) [\delta \boldsymbol{v}(\boldsymbol{r})]^2 \rangle = -\frac{4}{3} \epsilon_H \boldsymbol{r}.$$
(1.6)

However, to the best of our knowledge, there is no analogues 4/5 law for helicity in Navier-Stokes or Euler equations. Hence, a natural question arises that weather or not the Kolmogorov's 4/5 law (1.1) for the energy can be extended to helicity in incompressible fluid. This is our main motivation of the current paper. Actually, compared with the 4/5 law for the energy based on the velocity only, it seems that this type law for the helicity requires the mixed moments of the velocity and vorticity. This destroys the symmetry in some sense, consequently, it becomes more complicated to derive 4/5 law for the velocity coupling other quantity (see (1.16)). It also occurs in magnetized fluid (see [5, p. 148]), where there exist the nonlinear terms based on the velocity field and the magnetic field. In what follows, we invoke Eyink's decomposition (1.3) that one splits the quantity into longitudinal and transverse parts. Firstly, we state our first result on scaling laws for helicity in incompressible fluids.

Theorem 1.1. Suppose that the pair (v, ω) satisfy the following 3D Euler equations

$$\begin{cases} \boldsymbol{v}_t + \operatorname{div}(\boldsymbol{v} \otimes \boldsymbol{v}) + \nabla P = 0, \\ \boldsymbol{\omega}_t + \boldsymbol{v} \cdot \nabla \boldsymbol{\omega} - \boldsymbol{\omega} \cdot \nabla \boldsymbol{v} = 0, \\ \operatorname{div} \boldsymbol{v} = \operatorname{div} \boldsymbol{\omega} = 0, \end{cases}$$
(1.7)

where v and P stand for the velocity and pressure of the fluid, respectively. Then, there hold the following local longitudinal and transverse Kármárth-Howarth equations for helicity, respectively

$$\partial_{t}(\boldsymbol{\omega}\cdot\boldsymbol{v}_{L}^{\varepsilon}+\boldsymbol{v}\cdot\boldsymbol{\omega}_{L}^{\varepsilon})+\operatorname{div}(\boldsymbol{\omega}P_{L}^{\varepsilon}+P\boldsymbol{\omega}_{L}^{\varepsilon})+\operatorname{div}\left[\boldsymbol{v}(\boldsymbol{\omega}\cdot\boldsymbol{v}_{L}^{\varepsilon})-\boldsymbol{\omega}(\boldsymbol{v}\cdot\boldsymbol{v}_{L}^{\varepsilon})+\boldsymbol{v}(\boldsymbol{v}\cdot\boldsymbol{\omega}_{L}^{\varepsilon})\right] \\ +\operatorname{div}\left[\left(\boldsymbol{v}(\boldsymbol{\omega}_{L}\cdot\boldsymbol{v}_{L})\right)^{\varepsilon}-\boldsymbol{v}(\boldsymbol{\omega}_{L}\cdot\boldsymbol{v}_{L})^{\varepsilon}\right]+\frac{1}{2}\operatorname{div}\left[\boldsymbol{\omega}(\boldsymbol{v}_{L}\cdot\boldsymbol{v}_{L})^{\varepsilon}-\left(\boldsymbol{\omega}(\boldsymbol{v}_{L}\cdot\boldsymbol{v}_{L})\right)^{\varepsilon}\right] \\ =-\frac{4}{3}D_{HL}^{\varepsilon}(\boldsymbol{v},\boldsymbol{\omega}), \tag{1.8}$$

and

$$\partial_{t}(\boldsymbol{\omega}\cdot\boldsymbol{v}_{T}^{\varepsilon}+\boldsymbol{v}\cdot\boldsymbol{\omega}_{T}^{\varepsilon})+\operatorname{div}(\boldsymbol{\omega}P_{T}^{\varepsilon}+P\boldsymbol{\omega}_{T}^{\varepsilon})+\operatorname{div}\left[\boldsymbol{v}(\boldsymbol{\omega}\cdot\boldsymbol{v}_{T}^{\varepsilon})+\boldsymbol{v}(\boldsymbol{v}\cdot\boldsymbol{\omega}_{T}^{\varepsilon})-\boldsymbol{\omega}(\boldsymbol{v}\cdot\boldsymbol{v}_{T}^{\varepsilon})\right] \\ +\operatorname{div}\left[\left(\boldsymbol{v}(\boldsymbol{v}_{T}\cdot\boldsymbol{\omega}_{T})\right)^{\varepsilon}-\boldsymbol{v}(\boldsymbol{v}_{T}\cdot\boldsymbol{\omega}_{T})^{\varepsilon}\right]+\frac{1}{2}\operatorname{div}\left[\boldsymbol{\omega}(\boldsymbol{v}_{T}\cdot\boldsymbol{v}_{T})^{\varepsilon}-\left(\boldsymbol{\omega}(\boldsymbol{v}_{T}\cdot\boldsymbol{v}_{T})\right)^{\varepsilon}\right] \quad (1.9) \\ =-\frac{8}{3}D_{HT}^{\varepsilon}(\boldsymbol{v},\boldsymbol{\omega}),$$

where the dissipation terms (Kármán-Howarth-Monin type relation) are defined by

$$D_{HL}^{\varepsilon}(\boldsymbol{v},\boldsymbol{\omega}) = \frac{3}{4} \int_{\mathbb{T}^3} \nabla \varphi^{\varepsilon}(\ell) \cdot \delta \boldsymbol{v} (\delta \boldsymbol{v}_L \cdot \delta \boldsymbol{\omega}_L) + \frac{2}{|\boldsymbol{\ell}|} \varphi^{\varepsilon}(\ell) \hat{\boldsymbol{\ell}} \cdot \left[\delta \boldsymbol{v} (\delta \boldsymbol{v}_T \cdot \boldsymbol{\omega}_T) + \delta \boldsymbol{v} \times (\delta \boldsymbol{\omega} \times \delta \boldsymbol{v}) \right] d^3 \boldsymbol{\ell}$$
(1.10)
$$- \frac{3}{8} \int_{\mathbb{T}^3} \nabla \varphi^{\varepsilon}(\ell) \cdot \delta \boldsymbol{\omega} [\delta \boldsymbol{v}_L]^2 + \frac{2}{|\boldsymbol{\ell}|} \varphi^{\varepsilon}(\ell) \hat{\boldsymbol{\ell}} \cdot \delta \boldsymbol{\omega} [\delta \boldsymbol{v}_T]^2 d^3 \boldsymbol{\ell},$$

$$D_{HT}^{\varepsilon}(\boldsymbol{v},\boldsymbol{\omega}) = \frac{3}{8} \int \nabla \varphi^{\varepsilon}(\ell) \cdot \delta \boldsymbol{v}(\delta \boldsymbol{v}_{T} \cdot \delta \boldsymbol{\omega}_{T}) - \frac{2}{|\boldsymbol{\ell}|} \varphi^{\varepsilon}(\ell) \hat{\boldsymbol{\ell}} \cdot \left[\delta \boldsymbol{v}(\delta \boldsymbol{v}_{T} \cdot \delta \boldsymbol{\omega}_{T}) + \delta \boldsymbol{v} \times (\delta \boldsymbol{\omega} \times \delta \boldsymbol{v}) \right] d^{3} \boldsymbol{\ell} \quad (1.11)$$
$$- \frac{3}{16} \int \nabla \varphi^{\varepsilon} \cdot \delta \boldsymbol{\omega} [\delta \boldsymbol{v}_{T}]^{2} - \frac{2}{|\boldsymbol{\ell}|} \varphi^{\varepsilon}(\ell) \hat{\boldsymbol{\ell}} \cdot \delta \boldsymbol{\omega} [\delta \boldsymbol{v}_{T}]^{2} d^{3} \boldsymbol{\ell}.$$

In addition, if suppose that for any $1 < m, n < \infty, 3 \le p, q < \infty$ with $\frac{2}{p} + \frac{1}{m} = 1$ and $\frac{2}{q} + \frac{1}{n} = 1$ such that $(\boldsymbol{v}, \boldsymbol{\omega})$ satisfies

$$\boldsymbol{v} \in L^p(0,T; L^q(\mathbb{T}^3)) \text{ and } \boldsymbol{\omega} \in L^m(0,T; L^n(\mathbb{T}^3)).$$
 (1.12)

Then the function $D_{HX}^{\varepsilon}(\boldsymbol{v},\boldsymbol{\omega})$ with X = L or T converges to a distribution $D_H(\boldsymbol{v},\boldsymbol{\omega})$ in the sense of distributions as $\varepsilon \to 0$, and $D_H(\boldsymbol{v},\boldsymbol{\omega})$ satisfies the local equation of helicity

$$\frac{1}{2}\partial_t(\boldsymbol{v}\cdot\boldsymbol{\omega}) + \frac{1}{2}\mathrm{div}\Big[\boldsymbol{v}(\boldsymbol{v}\cdot\boldsymbol{\omega}) - \frac{1}{2}\boldsymbol{\omega}(|\boldsymbol{v}|^2) + P\boldsymbol{\omega}\Big] = -D_H(\boldsymbol{v},\boldsymbol{\omega}), \quad (1.13)$$

Furthermore, we have the following 4/5 exact relation for the helicity

$$S_{HL}(\boldsymbol{v},\boldsymbol{\omega}) = -\frac{4}{5}D_H(\boldsymbol{v},\boldsymbol{\omega}), \qquad (1.14)$$

and 8/15 law

$$S_{HT}(\boldsymbol{v},\boldsymbol{\omega}) = -\frac{8}{15}D_H(\boldsymbol{v},\boldsymbol{\omega}),$$

where

$$S_{HX}(\boldsymbol{v},\boldsymbol{\omega}) = \lim_{\lambda \to 0} S_{HX}(\boldsymbol{v},\boldsymbol{\omega},\lambda), \text{ with } X = L,T,$$

and

$$\begin{split} S_{HL}(\boldsymbol{v},\boldsymbol{\omega},\lambda) &= \frac{1}{\lambda} \int_{\partial B} \hat{\boldsymbol{\ell}} \cdot \left[\delta \boldsymbol{v}(\lambda \boldsymbol{\ell}) \left[\delta \boldsymbol{v}_L(\lambda \boldsymbol{\ell}) \cdot \delta \boldsymbol{\omega}_L(\lambda \boldsymbol{\ell}) \right] - \frac{1}{2} \delta \boldsymbol{\omega}(\lambda \boldsymbol{\ell}) [\delta \boldsymbol{v}_L(\lambda \boldsymbol{\ell})]^2 \right] \frac{d\sigma(\boldsymbol{\ell})}{4\pi} \\ &\quad - \frac{2}{5} \frac{1}{\lambda} \int_{\partial B} \hat{\boldsymbol{\ell}} \cdot \left[\delta \boldsymbol{\omega}(\lambda \boldsymbol{\ell}) \left[\delta \boldsymbol{v}(\lambda \boldsymbol{\ell}) \cdot \delta \boldsymbol{v}(\lambda \boldsymbol{\ell}) \right] - \delta \boldsymbol{v}(\lambda \boldsymbol{\ell}) \left(\delta \boldsymbol{\omega}(\lambda \boldsymbol{\ell}) \cdot \delta \boldsymbol{v}(\lambda \boldsymbol{\ell}) \right) \right] \frac{d\sigma(\boldsymbol{\ell})}{4\pi}, \\ S_{HT}(\boldsymbol{v},\boldsymbol{\omega},\lambda) &= \frac{1}{\lambda} \int_{\partial B} \hat{\boldsymbol{\ell}} \cdot \left[\delta \boldsymbol{v}(\lambda \boldsymbol{\ell}) \left[\delta \boldsymbol{v}_T(\lambda \boldsymbol{\ell}) \cdot \delta \boldsymbol{\omega}_T(\lambda \boldsymbol{\ell}) \right] - \frac{1}{2} \delta \boldsymbol{\omega}(\lambda \boldsymbol{\ell}) [\delta \boldsymbol{v}_T(\lambda \boldsymbol{\ell})]^2 \right] \frac{d\sigma(\boldsymbol{\ell})}{4\pi} \\ &\quad + \frac{2}{5} \frac{1}{\lambda} \int_{\partial B} \hat{\boldsymbol{\ell}} \cdot \left[\delta \boldsymbol{\omega}(\lambda \boldsymbol{\ell}) \left(\delta \boldsymbol{v}(\lambda \boldsymbol{\ell}) \cdot \delta \boldsymbol{v}(\lambda \boldsymbol{\ell}) \right) - \delta \boldsymbol{v}(\lambda \boldsymbol{\ell}) \left(\delta \boldsymbol{\omega}(\lambda \boldsymbol{\ell}) \cdot \delta \boldsymbol{v}(\lambda \boldsymbol{\ell}) \right) \right] \frac{d\sigma(\boldsymbol{\ell})}{4\pi}. \end{split}$$

Remark 1.1. It is noted that the 4/5 law (1.14) corresponds to the following four-fifths law for longitudinal third-order moments

$$\langle \delta \boldsymbol{v}_L (\delta \boldsymbol{v}_L \cdot \delta \boldsymbol{\omega}_L) \rangle - \frac{1}{2} \langle \delta \boldsymbol{\omega}_L (\delta \boldsymbol{v}_L)^2 \rangle - \frac{2}{5} \langle \delta \boldsymbol{\omega}_L (\delta \boldsymbol{v})^2 \rangle + \frac{2}{5} \langle \delta \boldsymbol{v}_L (\delta \boldsymbol{v} \cdot \delta \boldsymbol{\omega}) \rangle = -\frac{4}{5} \epsilon_H \boldsymbol{r}. \quad (1.15)$$

Remark 1.2. The 4/5 law and 8/15 law for helicity in Theorem 1.1 rely on the assumption that the limits of $S_{HX}(\boldsymbol{v}, \boldsymbol{\omega}, \lambda)$ with X = L, T exist as $\lambda \to 0$. Actually, this assumption can be removed by following the strategy due to Novack [44], in which the author considered the validity of certain exact results from turbulence theory in the deterministic setting. We leave this to the interested reader.

Remark 1.3. A well known helicity criterion for the incompressible Euler equations obtained in [9] is that the helicity of weak solutions is conserved provided that $v \in L^3(0,T; B_{3,q^{\natural}}^{\frac{2}{3}})$ with $1 \leq q^{\natural} < \infty$. Hence, if $m \geq 3$ in (1.12), we require n < 9/4 in this theorem. Since a special case of (1.12) is p = m = 3, $q = \frac{9}{2}$ and $n = \frac{9}{5}$, the condition (1.12) is no empty. Remark 1.4. Thanks to the vector triple product formula

$$A \times (B \times C) = B(A \cdot C) - C(A \cdot B)$$
, for any vectors A, B and C ,

one may reformulate $S_{HX}(\boldsymbol{v},\boldsymbol{\omega})$ by $\delta \boldsymbol{v} \times (\delta \boldsymbol{\omega} \times \delta \boldsymbol{v})$ in this theorem.

The new ingredient in this theorem is two-fold. First, we prove new exact laws and generalize famous Kolmogorov's 4/5 law (1.1) from energy to helicity for the incompressible flow. Second, we present the effect of mixed moments in four-fifths laws in terms of longitudinal third order. The interaction of different quantities brings more difficulty to derive a scaling relation analogous to Kolmogorov's 4/5 law (see [5, p. 148]). In Eyink's framework, it seems that two identities (2.19) and (2.20) in [20] break down for the velocity coupling with other different quantities. To over this difficulty, we are trying to find the accurate relation between $\frac{\partial}{\partial \ell_k} (\hat{\ell}_i \hat{\ell}_j)$ and $(\frac{\partial \hat{\ell}_i}{\partial \ell_j} + \frac{\partial \hat{\ell}_j}{\partial \ell_i}) \hat{\ell}_k$ involving different variable quantities (see (2.27)), which are equivalent in deriving the 4/5 law of energy just involving the velocity field quantity in the incompressible fluids. To this end, we deduce two modified identities (2.26) and (2.46) allowing different quantities, which reduce to Eyink's identities when the different quantities are the same. The starting point is that the following new identity

$$\begin{bmatrix} \frac{\partial}{\partial \ell_k} \left(\hat{\ell}_i \hat{\ell}_j \right) - \left(\frac{\partial \hat{\ell}_i}{\partial \ell_j} + \frac{\partial \hat{\ell}_j}{\partial \ell_i} \right) \hat{\ell}_k \end{bmatrix} A_k B_i C_j$$

=
$$\frac{1}{|\ell|} \hat{\ell} \cdot \left[\mathbf{C} (\mathbf{A} \cdot \mathbf{B}) + \mathbf{B} (\mathbf{A} \cdot \mathbf{C}) - 2\mathbf{A} (\mathbf{B} \cdot \mathbf{C}) \right]$$
(1.16)

is observed, which immediately becomes zero when $\mathbf{A} = \mathbf{B} = \mathbf{C}$.

Furthermore, in light of the key observation (1.16), we can deal with other scaling laws involving different variable quantities. Next, we turn our attention to the incompressible magnetohydrodynamic fluid, in which there exist the nonlinear terms involving both the the velocity \boldsymbol{v} and the magnetic field \boldsymbol{h} . Two 4/5 laws for total energy and cross-helicity obtained by Politano and Pouquet in the magnetized fluids in [49] can be written as

$$\langle [\delta \boldsymbol{v}_L(\boldsymbol{r})]^3 \rangle - 6 \langle \boldsymbol{h}_L^2 \delta \boldsymbol{v}_L(\boldsymbol{r}) \rangle = -\frac{4}{5} \epsilon_N \boldsymbol{r},$$

$$\langle [\delta \boldsymbol{h}_L(\boldsymbol{r})]^3 \rangle - 6 \langle \boldsymbol{h}_L^2 \delta \boldsymbol{v}_T(\boldsymbol{r}) \rangle = -\frac{4}{5} \epsilon_C \boldsymbol{r},$$

$$(1.17)$$

where ϵ_N and ϵ_C stand for total energy and cross-helicity dissipation rates, respectively (see also [6]). In another contemporary paper [48], Politano and Pouquet established the Yaglom scaling laws for energy and cross-helicity. Indeed, it is very useful to invoke these scaling laws to model the energy flux in turbulent plasmas. We refer the reader to [40] for a detailed review of scaling laws for the energy transfer in space plasma turbulence recently provided by Marino and Sorriso-Valvo. However, notice that the second term in $(1.17)_1$ and $(1.17)_2$ are $\langle \mathbf{h}_L^2 \delta \mathbf{v}_L(\mathbf{r}) \rangle$ and $\langle \mathbf{h}_L^2 \delta \mathbf{v}_T(\mathbf{r}) \rangle$ based on the structure and correlation functions rather than $\langle \delta \mathbf{h}_L^2 \delta \mathbf{v}_L(\mathbf{r}) \rangle$ and $\langle \delta \mathbf{h}_L^2 \delta \mathbf{v}_T(\mathbf{r}) \rangle$ via the structure functions only. Hence, our second objective is to revisit Kolmogorov type scaling laws (1.17) in plasma turbulence. In the spirit of above theorem, we formulate 4/5 laws of total energy and cross-helicity in the inviscid magnetohydrodynamic flow.

Theorem 1.2. Suppose that the triplet $(\boldsymbol{v}, \boldsymbol{h}, \Pi)$ satisfy the following

$$\begin{cases} \partial_t \boldsymbol{v} + \operatorname{div}(\boldsymbol{v} \otimes \boldsymbol{v}) - \operatorname{div}(\boldsymbol{h} \otimes \boldsymbol{h}) + \nabla \Pi = 0, \\ \partial_t \boldsymbol{h} + \operatorname{div}(\boldsymbol{v} \otimes \boldsymbol{h}) - \operatorname{div}(\boldsymbol{h} \otimes \boldsymbol{u}) = 0, \\ \operatorname{div} \boldsymbol{v} = \operatorname{div} \boldsymbol{h} = 0, \end{cases}$$
(1.18)

where v, h and Π stand for the veolcity field, magnetic field and the pressure of the fluid,

respectively. Then, there hold the following local longitudinal and transverse Kármárth-Howarth equations for energy,

$$\begin{aligned} \partial_t (\boldsymbol{v}_L^{\varepsilon} \cdot \boldsymbol{v} + \boldsymbol{h} \cdot \boldsymbol{h}_L^{\varepsilon}) + \operatorname{div} \Big[\boldsymbol{v} (\boldsymbol{v} \cdot \boldsymbol{v}_L^{\varepsilon}) - \boldsymbol{h} (\boldsymbol{h} \cdot \boldsymbol{v}_L^{\varepsilon}) + \boldsymbol{v} (\boldsymbol{h} \cdot \boldsymbol{h}_L^{\varepsilon}) - \boldsymbol{h} (\boldsymbol{v} \cdot \boldsymbol{h}_L^{\varepsilon}) \Big] \\ &+ \operatorname{div} \Big[\Pi_L^{\varepsilon} \boldsymbol{v} + \Pi \boldsymbol{v}_L^{\varepsilon} \Big] - \operatorname{div} \Big[\big(\boldsymbol{h} (\boldsymbol{v}_L \cdot \boldsymbol{h}_L) \big)^{\varepsilon} - \boldsymbol{h} (\boldsymbol{v}_L \cdot \boldsymbol{h}_L)^{\varepsilon} \Big] \\ &+ \frac{1}{2} \operatorname{div} \Big[\big(\boldsymbol{v} (\boldsymbol{v}_L \cdot \boldsymbol{v}_L) \big)^{\varepsilon} - \boldsymbol{v} (\boldsymbol{v}_L \cdot \boldsymbol{v}_L)^{\varepsilon} \Big] + \frac{1}{2} \operatorname{div} \Big[\big(\boldsymbol{v} (\boldsymbol{h}_L \cdot \boldsymbol{h}_L) \big)^{\varepsilon} - \boldsymbol{v} (\boldsymbol{h}_L \cdot \boldsymbol{h}_L)^{\varepsilon} \Big] \\ &= -\frac{2}{3} D_{EL}^{\varepsilon} (\boldsymbol{v}, \boldsymbol{h}); \\ \partial_t (\boldsymbol{v}_T^{\varepsilon} \cdot \boldsymbol{v} + \boldsymbol{h} \cdot \boldsymbol{h}_T^{\varepsilon}) + \operatorname{div} \Big[\boldsymbol{v} (\boldsymbol{v} \cdot \boldsymbol{v}_T^{\varepsilon}) - \boldsymbol{h} (\boldsymbol{h} \cdot \boldsymbol{v}_T^{\varepsilon}) + \boldsymbol{v} (\boldsymbol{h} \cdot \boldsymbol{h}_T^{\varepsilon}) - \boldsymbol{h} (\boldsymbol{v} \cdot \boldsymbol{h}_T^{\varepsilon}) \Big] \\ &+ \operatorname{div} \Big[\Pi_T^{\varepsilon} \boldsymbol{v} + \Pi \boldsymbol{v}_T^{\varepsilon} \Big] - \operatorname{div} \Big[\big(\boldsymbol{h} (\boldsymbol{v}_T \cdot \boldsymbol{h}_T) \big)^{\varepsilon} - \boldsymbol{h} (\boldsymbol{v}_T \cdot \boldsymbol{h}_T)^{\varepsilon} \Big] \\ &+ \frac{1}{2} \operatorname{div} \Big[\big(\boldsymbol{v} (\boldsymbol{v}_T \cdot \boldsymbol{v}_T) \big)^{\varepsilon} - \boldsymbol{v} (\boldsymbol{v}_T \cdot \boldsymbol{v}_T)^{\varepsilon} \Big] + \frac{1}{2} \operatorname{div} \Big[\big(\boldsymbol{v} (\boldsymbol{h}_T \cdot \boldsymbol{h}_T) \big)^{\varepsilon} - \boldsymbol{v} (\boldsymbol{h}_T \cdot \boldsymbol{h}_T)^{\varepsilon} \Big] \\ &= -\frac{4}{3} D_{ET}^{\varepsilon} (\boldsymbol{v}, \boldsymbol{h}); \end{aligned}$$

and the following local longitudinal and transverse Kármárth-Howarth equations for cross-helicity

$$\begin{aligned} \partial_t (\boldsymbol{v}_L^{\varepsilon} \cdot \boldsymbol{h} + \boldsymbol{v} \cdot \boldsymbol{h}_L^{\varepsilon}) + \operatorname{div} \Big[\boldsymbol{v} (\boldsymbol{v} \cdot \boldsymbol{h}_L^{\varepsilon}) - \boldsymbol{h} (\boldsymbol{h} \cdot \boldsymbol{h}_L^{\varepsilon}) + \boldsymbol{v} (\boldsymbol{h} \cdot \boldsymbol{v}_L^{\varepsilon}) - \boldsymbol{h} (\boldsymbol{v} \cdot \boldsymbol{v}_L^{\varepsilon}) \Big] \\ &+ \operatorname{div} (\Pi_L^{\varepsilon} \boldsymbol{h} + \boldsymbol{h}_L^{\varepsilon} \Pi) + \operatorname{div} \Big[\big(\boldsymbol{v} (\boldsymbol{h}_L \cdot \boldsymbol{v}_L) \big)^{\varepsilon} - \boldsymbol{v} (\boldsymbol{h}_L \cdot \boldsymbol{v}_L)^{\varepsilon} \Big] \\ &- \frac{1}{2} \operatorname{div} \Big[\big(\boldsymbol{h} (\boldsymbol{h}_L \cdot \boldsymbol{h}_L) \big)^{\varepsilon} - \boldsymbol{h} (\boldsymbol{h}_L \cdot \boldsymbol{h}_L)^{\varepsilon} \Big] - \frac{1}{2} \operatorname{div} \Big[\big(\boldsymbol{h} (\boldsymbol{v}_L \cdot \boldsymbol{v}_L) \big)^{\varepsilon} - \boldsymbol{h} (\boldsymbol{v}_L \cdot \boldsymbol{v}_L)^{\varepsilon} \Big] \\ &= - \frac{2}{3} D_{CHL}^{\varepsilon} (\boldsymbol{v}, \boldsymbol{h}); \\ \partial_t (\boldsymbol{v}_T^{\varepsilon} \cdot \boldsymbol{h} + \boldsymbol{v} \cdot \boldsymbol{h}_T^{\varepsilon}) + \operatorname{div} \Big[\boldsymbol{v} (\boldsymbol{v} \cdot \boldsymbol{h}_T^{\varepsilon}) - \boldsymbol{h} (\boldsymbol{h} \cdot \boldsymbol{h}_T^{\varepsilon}) + \boldsymbol{v} (\boldsymbol{h} \cdot \boldsymbol{v}_T^{\varepsilon}) - \boldsymbol{h} (\boldsymbol{v} \cdot \boldsymbol{v}_T^{\varepsilon}) \Big] \\ &+ \operatorname{div} (\Pi_T^{\varepsilon} \boldsymbol{h} + \boldsymbol{h}_T^{\varepsilon} \Pi) + \operatorname{div} \Big[\big(\boldsymbol{v} (\boldsymbol{h}_T \cdot \boldsymbol{v}_T) \big)^{\varepsilon} - \boldsymbol{v} (\boldsymbol{h}_T \cdot \boldsymbol{v}_T)^{\varepsilon} \Big] \\ &- \frac{1}{2} \operatorname{div} \Big[\big(\boldsymbol{h} (\boldsymbol{h}_T \cdot \boldsymbol{h}_T) \big)^{\varepsilon} - \boldsymbol{h} (\boldsymbol{h}_T \cdot \boldsymbol{h}_T)^{\varepsilon} \Big] - \frac{1}{2} \operatorname{div} \Big[\big(\boldsymbol{h} (\boldsymbol{v}_T \cdot \boldsymbol{v}_T) \big)^{\varepsilon} - \boldsymbol{h} (\boldsymbol{v}_T \cdot \boldsymbol{v}_T)^{\varepsilon} \Big] \\ &= - \frac{4}{3} D_{CHT}^{\varepsilon} (\boldsymbol{v}, \boldsymbol{h}). \end{aligned}$$

Here,

$$\begin{split} D_{EL}^{\varepsilon}(\boldsymbol{v},\boldsymbol{h}) &= \frac{3}{4} \int_{\mathbb{T}^3} \nabla \varphi^{\varepsilon}(\ell) \cdot \delta \boldsymbol{v} \Big([\delta \boldsymbol{v}_L]^2 + [\delta \boldsymbol{h}_L]^2 \Big) + \frac{2}{|\boldsymbol{\ell}|} \varphi^{\varepsilon}(\ell) \hat{\boldsymbol{\ell}} \cdot \delta \boldsymbol{v} \Big([\delta \boldsymbol{v}_T]^2 + [\delta \boldsymbol{h}_T]^2 \Big) d^3 \boldsymbol{\ell} \\ &- \frac{3}{2} \int_{\mathbb{T}^3} \nabla \varphi^{\varepsilon}(\ell) \cdot \delta \boldsymbol{h} (\delta \boldsymbol{v}_L \cdot \delta \boldsymbol{h}_L) + \frac{2}{|\boldsymbol{\ell}|} \varphi^{\varepsilon}(\ell) \hat{\boldsymbol{\ell}} \cdot \delta \boldsymbol{h} (\delta \boldsymbol{v}_T \cdot \delta \boldsymbol{h}_T) d^3 \boldsymbol{\ell} \\ &- 3 \int_{\mathbb{T}^3} \frac{1}{|\boldsymbol{\ell}|} \varphi^{\varepsilon}(\ell) \hat{\boldsymbol{\ell}} \cdot \Big[\delta \boldsymbol{h} \times (\delta \boldsymbol{v} \times \delta \boldsymbol{h}) \Big] d^3 \boldsymbol{\ell}; \\ D_{ET}^{\varepsilon}(\boldsymbol{v},\boldsymbol{h}) &= \frac{3}{8} \int_{\mathbb{T}^3} \nabla \varphi^{\varepsilon}(\ell) \cdot \delta \boldsymbol{v} \Big([\delta \boldsymbol{v}_T]^2 + [\delta \boldsymbol{h}_T]^2 \Big) - \frac{2}{|\boldsymbol{\ell}|} \varphi^{\varepsilon} \hat{\boldsymbol{\ell}} \cdot \delta \boldsymbol{v} \Big([\delta \boldsymbol{v}_T]^2 + [\delta \boldsymbol{h}_T]^2 \Big) d^3 \boldsymbol{\ell} \\ &- \frac{3}{4} \int_{\mathbb{T}^3} \nabla \varphi^{\varepsilon}(\ell) \cdot \delta \boldsymbol{h} (\delta \boldsymbol{v}_T \cdot \delta \boldsymbol{h}_T) - \frac{2}{|\boldsymbol{\ell}|} \varphi^{\varepsilon}(\ell) \hat{\boldsymbol{\ell}} \cdot \delta \boldsymbol{h} (\delta \boldsymbol{v}_T \cdot \delta \boldsymbol{h}_T) d^3 \boldsymbol{\ell} \\ &+ \frac{3}{2} \int_{\mathbb{T}^3} \frac{1}{|\boldsymbol{\ell}|} \varphi^{\varepsilon}(\ell) \hat{\boldsymbol{\ell}} \cdot [\delta \boldsymbol{h} \times (\delta \boldsymbol{v} \times \delta \boldsymbol{h})] d^3 \boldsymbol{\ell}; \end{split}$$

$$\begin{split} D_{CHL}^{\varepsilon}(\boldsymbol{v},\boldsymbol{h}) &= \frac{3}{2} \int_{\mathbb{T}^3} \nabla \varphi^{\varepsilon}(\ell) \cdot \delta \boldsymbol{v} (\delta \boldsymbol{h}_L \cdot \delta \boldsymbol{v}_L) + \frac{2}{|\boldsymbol{\ell}|} \varphi^{\varepsilon}(\ell) \hat{\boldsymbol{\ell}} \cdot \delta \boldsymbol{v} (\delta \boldsymbol{h}_T \cdot \delta \boldsymbol{v}_T) d^3 \boldsymbol{\ell} \\ &\quad -\frac{3}{4} \int_{\mathbb{T}^3} \nabla \varphi^{\varepsilon}(\ell) \cdot \delta \boldsymbol{h} \Big([\delta \boldsymbol{h}_L]^2 + [\delta \boldsymbol{v}_L]^2 \Big) + \frac{2}{|\boldsymbol{\ell}|} \varphi^{\varepsilon}(\ell) \hat{\boldsymbol{\ell}} \cdot \delta \boldsymbol{h} \Big([\delta \boldsymbol{h}_T]^2 + [\delta \boldsymbol{v}_T]^2 \Big) d^3 \boldsymbol{\ell} \\ &\quad +\frac{3}{2} \int_{\mathbb{T}^3} \frac{2}{|\boldsymbol{\ell}|} \varphi^{\varepsilon}(\ell) \hat{\boldsymbol{\ell}} \cdot \Big[\delta \boldsymbol{v} \times (\delta \boldsymbol{h} \times \delta \boldsymbol{v}) \Big] d^3 \boldsymbol{\ell}; \\ D_{CHT}^{\varepsilon}(\boldsymbol{v},\boldsymbol{h}) &= \frac{3}{4} \int_{\mathbb{T}^3} \nabla \varphi^{\varepsilon}(\ell) \cdot \delta \boldsymbol{v} (\delta \boldsymbol{h}_T \cdot \delta \boldsymbol{v}_T) - \frac{2}{|\boldsymbol{\ell}|} \varphi^{\varepsilon} \hat{\boldsymbol{\ell}} \cdot \delta \boldsymbol{v} (\delta \boldsymbol{h}_T \cdot \delta \boldsymbol{v}_T) d^3 \boldsymbol{\ell} \\ &\quad -\frac{3}{8} \int_{\mathbb{T}^3} \nabla \varphi^{\varepsilon}(\ell) \cdot \delta \boldsymbol{h} \Big([\delta \boldsymbol{h}_T]^2 + [\delta \boldsymbol{v}_T]^2 \Big) - \frac{2}{|\boldsymbol{\ell}|} \varphi^{\varepsilon} \hat{\boldsymbol{\ell}} \cdot \delta \boldsymbol{h} \Big([\delta \boldsymbol{h}_T]^2 + [\delta \boldsymbol{v}_T]^2 \Big) d^3 \boldsymbol{\ell} \\ &\quad -\frac{3}{4} \int_{\mathbb{T}^3} \frac{2}{|\boldsymbol{\ell}|} \varphi^{\varepsilon}(\ell) \hat{\boldsymbol{\ell}} \cdot \Big[\delta \boldsymbol{v} \times (\delta \boldsymbol{h} \times \delta \boldsymbol{v}) \Big] d^3 \boldsymbol{\ell}. \end{split}$$

In addition, if suppose that $(\boldsymbol{v}, \boldsymbol{h})$ satisfies

$$\boldsymbol{v}, \boldsymbol{h} \in L^3(0, T; L^3(\mathbb{T}^3)).$$
 (1.19)

Then the function $D_{YX}^{\varepsilon}(\boldsymbol{v},\boldsymbol{h})$ with X = L,T and Y = E,CH converges to a distribution $D_Y(\boldsymbol{v},\boldsymbol{h})$ in the sense of distributions as $\varepsilon \to 0$, and $D_Y(\boldsymbol{v},\boldsymbol{h})$ satisfies the local equation of total energy and cross-helicity, respectively

$$\frac{1}{2}\partial_t(|\boldsymbol{v}|^2 + |\boldsymbol{h}|^2) + \operatorname{div}\left[\frac{1}{2}\boldsymbol{v}|\boldsymbol{v}|^2 + \frac{1}{2}\boldsymbol{v}|\boldsymbol{h}|^2 - \boldsymbol{h}(\boldsymbol{h}\cdot\boldsymbol{v}) + \boldsymbol{v}\Pi\right] = -D_E(\boldsymbol{v},\boldsymbol{h})$$
(1.20)

$$\partial_t (\boldsymbol{v} \cdot \boldsymbol{h}) + \operatorname{div} \left[\boldsymbol{v}(\boldsymbol{v} \cdot \boldsymbol{h}) - \frac{1}{2} (|\boldsymbol{h}|^2) \boldsymbol{h} - \frac{1}{2} \boldsymbol{h} |\boldsymbol{v}|^2 + \boldsymbol{h} \Pi \right] = -D_{CH}(\boldsymbol{v}, \boldsymbol{h}).$$
(1.21)

Furthermore, we have the following 4/5 and 8/15 laws

$$S_{EL}(\boldsymbol{v},\boldsymbol{h}) = -\frac{4}{5}D_E(\boldsymbol{v},\boldsymbol{h}), \ S_{ET}(\boldsymbol{v},\boldsymbol{h}) = -\frac{8}{15}D_E(\boldsymbol{v},\boldsymbol{h})$$
(1.22)

$$S_{CHL}(\boldsymbol{v},\boldsymbol{h}) = -\frac{4}{5}D_{CH}(\boldsymbol{v},\boldsymbol{h}), \ S_{CHT}(\boldsymbol{v},\boldsymbol{h}) = -\frac{8}{15}D_{CH}(\boldsymbol{v},\boldsymbol{h}),$$
(1.23)

where

$$S_{YX}(\boldsymbol{v},\boldsymbol{h}) = \lim_{\lambda \to 0} S_{YX}(\boldsymbol{v},\boldsymbol{h},\lambda), \text{ with } X = L, T, Y = E, CH,$$

$$\begin{split} S_{EL}(\boldsymbol{v},\boldsymbol{h},\lambda) \\ =& \frac{1}{\lambda} \int_{\partial B} \hat{\boldsymbol{\ell}} \cdot \left[\delta \boldsymbol{v}(\lambda \boldsymbol{\ell}) | \left(\delta \boldsymbol{v}_L(\lambda \boldsymbol{\ell}) \right]^2 + \left[\delta \boldsymbol{h}_L(\lambda \boldsymbol{\ell}) \right]^2 \right) - 2\delta \boldsymbol{h}(\boldsymbol{\ell}) \left(\delta \boldsymbol{v}_L(\lambda \boldsymbol{\ell}) \cdot \delta \boldsymbol{h}_L(\lambda \boldsymbol{\ell}) \right) \right] \frac{d\sigma(\boldsymbol{\ell})}{4\pi} \\ &- \frac{4}{5} \frac{1}{\lambda} \int_{\partial B} \hat{\boldsymbol{\ell}} \cdot \left[\delta \boldsymbol{h}(\lambda \boldsymbol{\ell}) \times \left(\delta \boldsymbol{v}(\lambda \boldsymbol{\ell}) \times \delta \boldsymbol{h}(\lambda \boldsymbol{\ell}) \right) \right] \frac{d\sigma(\boldsymbol{\ell})}{4\pi}; \\ S_{ET}(\boldsymbol{v},\boldsymbol{h},\lambda) \\ =& \frac{1}{\lambda} \int_{\partial B} \hat{\boldsymbol{\ell}} \cdot \left[\delta \boldsymbol{v}(\lambda \boldsymbol{\ell}) \left(\left[\delta \boldsymbol{v}_T(\lambda \boldsymbol{\ell}) \right]^2 + \left[\delta \boldsymbol{h}_T(\lambda \boldsymbol{\ell}) \right]^2 \right) - 2\delta \boldsymbol{h}(\lambda \boldsymbol{\ell}) \left(\delta \boldsymbol{v}_T(\lambda \boldsymbol{\ell}) \cdot \delta \boldsymbol{h}_T(\lambda \boldsymbol{\ell}) \right) \right] \frac{d\sigma(\boldsymbol{\ell})}{4\pi} \\ &+ \frac{4}{5} \frac{1}{\lambda} \int_{\partial B} \hat{\boldsymbol{\ell}} \cdot \left[\delta \boldsymbol{h}(\lambda \boldsymbol{\ell}) \times \left(\delta \boldsymbol{v}(\lambda \boldsymbol{\ell}) \times \delta \boldsymbol{h}(\lambda \boldsymbol{\ell}) \right) \right] \frac{d\sigma(\boldsymbol{\ell})}{4\pi}; \\ S_{CHL}(\boldsymbol{v},\boldsymbol{h},\lambda) \end{split}$$

$$\begin{split} &= \int_{\partial B} \hat{\boldsymbol{\ell}} \cdot \Big[2\delta \boldsymbol{v}(\lambda \boldsymbol{\ell}) \Big(\delta \boldsymbol{h}_L(\lambda \boldsymbol{\ell}) \cdot \delta \boldsymbol{v}_L(\lambda \boldsymbol{\ell}) \Big) - \delta \boldsymbol{h}(\lambda \boldsymbol{\ell}) \Big([\delta \boldsymbol{h}_L(\lambda \boldsymbol{\ell})]^2 + [\delta \boldsymbol{v}_L(\lambda \boldsymbol{\ell})]^2 \Big) \Big] \frac{d\sigma(\boldsymbol{\ell})}{4\pi} \\ &- \frac{4}{5} \frac{1}{\lambda} \int_{\partial B} \hat{\boldsymbol{\ell}} \cdot \Big[\delta \boldsymbol{v}(\lambda \boldsymbol{\ell}) \times \Big(\delta \boldsymbol{h}(\lambda \boldsymbol{\ell}) \times \delta \boldsymbol{v}(\lambda \boldsymbol{\ell}) \Big) \Big] \frac{d\sigma(\boldsymbol{\ell})}{4\pi}; \\ S_{CHT}(\boldsymbol{v}, \boldsymbol{h}, \lambda) \\ &= \frac{1}{\lambda} \int_{\partial B} \hat{\boldsymbol{\ell}} \cdot \Big[2\delta \boldsymbol{v}(\lambda \boldsymbol{\ell}) \Big(\delta \boldsymbol{h}_T(\lambda \boldsymbol{\ell}) \cdot \delta \boldsymbol{v}_T(\lambda \boldsymbol{\ell}) \Big) - \delta \boldsymbol{h}(\lambda \boldsymbol{\ell}) \Big([\delta \boldsymbol{v}_T(\lambda \boldsymbol{\ell})]^2 + [\delta \boldsymbol{h}_T(\lambda \boldsymbol{\ell})]^2 \Big) \Big] \frac{d\sigma(\boldsymbol{\ell})}{4\pi} \\ &+ \frac{4}{5} \frac{1}{\lambda} \int_{\partial B} \hat{\boldsymbol{\ell}} \cdot \Big[\delta \boldsymbol{v}(\lambda \boldsymbol{\ell}) \times \Big(\delta \boldsymbol{h}(\lambda \boldsymbol{\ell}) \times \delta \boldsymbol{v}(\lambda \boldsymbol{\ell}) \Big) \Big] \frac{d\sigma(\boldsymbol{\ell})}{4\pi}. \end{split}$$

Remark 1.5. According to the vector triple product formula in Remark 1.4, one could rewrite $S_{EL}(\boldsymbol{v}, \boldsymbol{h}, \lambda)$, $S_{ET}(\boldsymbol{v}, \boldsymbol{h}, \lambda)$, $S_{CHL}(\boldsymbol{v}, \boldsymbol{h}, \lambda)$ and $S_{CHT}(\boldsymbol{v}, \boldsymbol{h}, \lambda)$ slightly. Hence, the 4/5 laws (1.22) and (1.23) correspond to the following 4/5 laws for longitudinal third-order moments

$$\langle \delta \boldsymbol{v}_L (\delta \boldsymbol{v}_L)^2 \rangle + \langle \delta \boldsymbol{v}_L (\delta \boldsymbol{h}_L)^2 \rangle - 2 \langle \delta \boldsymbol{h}_L (\delta \boldsymbol{h}_L \cdot \delta \boldsymbol{v}_L) \rangle - \frac{4}{5} \langle \delta \boldsymbol{h}_L (\delta \boldsymbol{h} \cdot \delta \boldsymbol{v}) \rangle + \frac{4}{5} \langle \delta \boldsymbol{v}_L (\delta \boldsymbol{h})^2 \rangle = -\frac{4}{5} \epsilon_E \boldsymbol{r},$$

$$2 \langle \delta \boldsymbol{v}_L (\delta \boldsymbol{h}_L \cdot \delta \boldsymbol{v}_L) \rangle - \langle \delta \boldsymbol{h}_L (\delta \boldsymbol{h}_L)^2 \rangle - \langle \delta \boldsymbol{h}_L (\delta \boldsymbol{v}_L)^2 \rangle - \frac{4}{5} \langle \delta \boldsymbol{h}_L (\delta \boldsymbol{v})^2 \rangle + \frac{4}{5} \langle \delta \boldsymbol{v}_L (\delta \boldsymbol{v} \cdot \delta \boldsymbol{h}) \rangle = -\frac{4}{5} \epsilon_C \boldsymbol{r}.$$

$$(1.24)$$

Remark 1.6. The four-fifths law (1.22) concerning energy in the inviscid MHD system reduces to the celebrated Kolmogorov four-fifths law showed by Eyink in [20] when the magnetic field becomes constant.

Remark 1.7. We rewrite the inviscid MHD equations (1.18) in terms of the Elsässer variances $Z^+ = \frac{v+h}{2}$, $Z^- = \frac{v-h}{2}$ as follows

$$\begin{cases} \partial_t \mathbf{Z}^+ + \mathbf{Z}^- \cdot \mathbf{Z}^+ + \nabla \Pi = 0, \\ \partial_t \mathbf{Z}^- + \mathbf{Z}^+ \cdot \mathbf{Z}^- + \nabla \Pi = 0, \\ \operatorname{div} \mathbf{Z}^+ = \operatorname{div} \mathbf{Z}^- = 0. \end{cases}$$
(1.25)

Following the path of the above theorems, one could show that

$$\langle \delta \mathbf{Z}_{L}^{-} [\delta \mathbf{Z}_{L}^{+}|^{2} \rangle - \frac{2}{5} \langle \delta \mathbf{Z}_{L}^{+} (\mathbf{Z}_{L}^{+} \cdot \mathbf{Z}_{L}^{-}) \rangle + \frac{2}{5} \langle \delta \mathbf{Z}_{L}^{-} [\delta \mathbf{Z}^{+}|^{2} \rangle = -\frac{4}{5} \epsilon_{+} r,$$

$$\langle \delta \mathbf{Z}_{L}^{+} [\delta \mathbf{Z}_{L}^{-}|^{2} \rangle - \frac{2}{5} \langle \delta \mathbf{Z}_{L}^{-} (\mathbf{Z}_{L}^{-} \cdot \mathbf{Z}_{L}^{+}) \rangle + \frac{2}{5} \langle \delta \mathbf{Z}_{L}^{+} [\delta \mathbf{Z}^{-}|^{2} \rangle = -\frac{4}{5} \epsilon_{+} r.$$

$$(1.26)$$

This is left to the readers.

Remark 1.8. Compared with the application of the Kármárth-Howarth type equations in a statistical view, the filtering approach for derivation of the scaling laws developing in [18, 20] holds for individual realizations of fluids. We refer to [17, 20, 21, 36] for advantage of this method.

Remark 1.9. Just as the relationship between four-fifths law (1.1) and four-thirds law (1.4), the left-hand side of (1.6) also appears to the left of (1.15), so does that in (1.24) and (1.26). Remark 1.10. It is worth remarking that a four-fifths law of magnetic helicity in the electron magnetohydrodynamic flow was established in [11]. In a forthcoming paper, we will address four-fifths laws for energy and magnetic helicity in electron and Hall magnetohydrodynamic fluids.

The remainder of this paper is organized as follows. Section 2 is devoted to Kolmogorov type exact scaling relation of helicity in the incompressible fluids. In Section 3, we study dissipation rates of total energy and cross helicity in the magnetohydrodynamics flow, respectively. Finally, a summary is provided.

2 Four-fifths laws of helicity in incompressible fluids

In this section, we apply Eyink's decomposition (1.3) involving the longitudinal and transverse velocity increments to consider the dissipation rates of helicity in incompressible fluids. A new identity (2.27) is observed, which allows one to deal with the interaction of different physical quantities.

Proof of Theorem 1.1. The proof of Theorem 1.1 is divided into four steps.

<u>Step 1</u>: First, inspired by the work of Duchon-Robert [18] and Eyink [20], we estiblish the local longitudinal and transverse Kármárth-Howarth equations for the helicity of Euler equations (1.7). To do this, we need to consider the longitudinal and transverse balance equations for the Euler equations separately. We define the longitudinal and transverse velocity (vorticity) relative to a vector ℓ as follows:

$$\boldsymbol{v}_{L}(\boldsymbol{x},t,\boldsymbol{\ell}) = (\hat{\boldsymbol{\ell}} \otimes \hat{\boldsymbol{\ell}}) \cdot \boldsymbol{v}(\boldsymbol{x}+\boldsymbol{\ell},t), \quad \boldsymbol{v}_{T}(\boldsymbol{x},t,\boldsymbol{\ell}) = (\mathbf{1} - \hat{\boldsymbol{\ell}} \otimes \hat{\boldsymbol{\ell}}) \cdot \boldsymbol{v}(\boldsymbol{x}+\boldsymbol{\ell},t); \\ \boldsymbol{\omega}_{L}(\boldsymbol{x},t,\boldsymbol{\ell}) = (\hat{\boldsymbol{\ell}} \otimes \hat{\boldsymbol{\ell}}) \cdot \boldsymbol{\omega}(\boldsymbol{x}+\boldsymbol{\ell},t), \quad \boldsymbol{\omega}_{T}(\boldsymbol{x},t,\boldsymbol{\ell}) = (\mathbf{1} - \hat{\boldsymbol{\ell}} \otimes \hat{\boldsymbol{\ell}}) \cdot \boldsymbol{v}(\boldsymbol{x}+\boldsymbol{\ell},t);$$
(2.1)

which implies the fact that $\boldsymbol{v}_L + \boldsymbol{v}_T = \boldsymbol{v}$ and $\boldsymbol{\omega}_L + \boldsymbol{\omega}_T = \boldsymbol{\omega}$. Let $\varphi(\ell)$ be any C_0^{∞} function, nonnegative with unit integral, radially symmetric, and let $\varphi^{\varepsilon}(\ell) = \frac{1}{\varepsilon^3}\varphi(\frac{\ell}{\varepsilon})$. Then the mollified version of $\boldsymbol{v}_X, \boldsymbol{\omega}_X$ with X = L, T can be defined as follows

$$\boldsymbol{v}_X^{\varepsilon}(\boldsymbol{x},t) = \int_{\mathbb{T}^3} \varphi^{\varepsilon}(\boldsymbol{\ell}) \boldsymbol{v}_X(\boldsymbol{x},t;\boldsymbol{\ell}) d^3 \boldsymbol{\ell}, \quad X = L, T,$$
(2.2)

and

$$\boldsymbol{\omega}_{X}^{\varepsilon}(\boldsymbol{x},t) = \int_{\mathbb{T}^{3}} \varphi^{\varepsilon}(\boldsymbol{\ell}) \boldsymbol{\omega}_{X}(\boldsymbol{x},t;\boldsymbol{\ell}) d^{3}\boldsymbol{\ell}, \quad X = L,T,$$
(2.3)

respectively.

To obtain the longitudinal velocity (vorticity) equations relative to vector $\boldsymbol{\ell}$, multiplying $(1.7)_1$ and $(1.7)_2$ by $\varphi^{\varepsilon}(\boldsymbol{\ell})(\hat{\boldsymbol{\ell}} \otimes \hat{\boldsymbol{\ell}})$ and integrating over \mathbb{T}^3 , we have

$$\partial_t \boldsymbol{v}_L^{\varepsilon} + \operatorname{div}(\boldsymbol{v} \otimes \boldsymbol{v}_L)^{\varepsilon} + \operatorname{div}\boldsymbol{\Pi}_L^{\varepsilon} = 0, \partial_t \boldsymbol{\omega}_L^{\varepsilon} + \operatorname{div}(\boldsymbol{v} \otimes \boldsymbol{\omega}_L)^{\varepsilon} - \operatorname{div}(\boldsymbol{\omega} \otimes \boldsymbol{v}_L)^{\varepsilon} = 0,$$
(2.4)

where

$$\boldsymbol{\Pi}_{L}^{\varepsilon} = \int_{\mathbb{T}^{3}} \varphi^{\varepsilon}(\boldsymbol{\ell}) (\hat{\boldsymbol{\ell}} \otimes \hat{\boldsymbol{\ell}}) P(\boldsymbol{x} + \boldsymbol{\ell}, t) d^{3} \boldsymbol{\ell}.$$
(2.5)

By the same token, multiplying $(1.7)_1$ and $(1.7)_2$ by $\varphi^{\varepsilon}(\ell)(\mathbf{1} - \hat{\ell} \otimes \hat{\ell})$ and integrating over \mathbb{T}^3 , it follows that

$$\partial_t \boldsymbol{v}_T^{\varepsilon} + \operatorname{div}(\boldsymbol{v} \otimes \boldsymbol{v}_T)^{\varepsilon} + \operatorname{div} \boldsymbol{\Pi}_T^{\varepsilon} = 0, \partial_t \boldsymbol{\omega}_T^{\varepsilon} + \operatorname{div}(\boldsymbol{v} \otimes \boldsymbol{\omega}_T)^{\varepsilon} - \operatorname{div}(\boldsymbol{\omega} \otimes \boldsymbol{v}_T)^{\varepsilon} = 0,$$
(2.6)

where

$$\mathbf{\Pi}_{T}^{\varepsilon} = \int_{\mathbb{T}^{3}} \varphi^{\varepsilon}(\boldsymbol{\ell}) (\mathbf{1} - \hat{\boldsymbol{\ell}} \otimes \hat{\boldsymbol{\ell}}) P(\boldsymbol{x} + \boldsymbol{\ell}, t) d^{3} \boldsymbol{\ell}.$$
(2.7)

Before proceeding further, we claim that

$$\operatorname{div} \mathbf{\Pi}_{X}^{\varepsilon}(\boldsymbol{x},t) = \nabla P_{X}(\boldsymbol{x},t), \quad X = L,T,$$
(2.8)

holds in the sense of distributions, where $P_L(\boldsymbol{x},t), P_T(\boldsymbol{x},t)$ are scalar functions defined as follows

$$P_X^{\varepsilon}(\boldsymbol{x},t) = \int_{\mathbb{T}^3} \varphi_X^{\varepsilon}(\ell) P(\boldsymbol{x}+\boldsymbol{\ell},t) d^3 \boldsymbol{\ell}, \quad X = L, T,$$
(2.9)

with

$$\varphi_L(\ell) = \varphi(\ell) - \varphi_T(\ell), \quad \varphi_T(\ell) = 2 \int_{|\boldsymbol{\ell}|}^{\infty} \frac{\varphi(\ell')}{\ell'} d\ell'.$$
 (2.10)

With the help of the definition of $\varphi(\ell)$, it is easy to see that $\varphi_L(\ell)$ and $\varphi_T(\ell)$ defined here are compactly supported and C^{∞} everywhere except at 0, where they have a mild (logarithmic) singularity.

Now we are in a position to show the validity of (2.8). On the one hand, using a straightforward computation, we have the following basic relation

$$\partial_{\ell_k} \hat{\boldsymbol{\ell}}_i = \frac{\partial}{\partial \ell_k} \left(\frac{\ell_i}{|\boldsymbol{\ell}|} \right) = \frac{\delta^{ik} |\boldsymbol{\ell}| - \frac{\ell_i \ell_k}{|\boldsymbol{\ell}|}}{|\boldsymbol{\ell}|^2} = \frac{1}{|\boldsymbol{\ell}|} (\delta^{ik} - \hat{\boldsymbol{\ell}}_i \hat{\boldsymbol{\ell}}_k), \tag{2.11}$$

which together with the integration by parts yields

$$\operatorname{div} \mathbf{\Pi}_{L}^{\varepsilon} = \partial_{x_{i}} \int_{\mathbb{T}^{3}} \varphi^{\varepsilon}(\ell) (\hat{\boldsymbol{\ell}}_{i} \hat{\boldsymbol{\ell}}_{j}) P(\boldsymbol{x} + \boldsymbol{\ell}) d^{3} \boldsymbol{\ell}$$

$$= \int_{\mathbb{T}^{3}} \varphi^{\varepsilon}(\ell) (\hat{\boldsymbol{\ell}}_{i} \hat{\boldsymbol{\ell}}_{j}) \partial_{x_{i}} P(\boldsymbol{x} + \boldsymbol{\ell}) d^{3} \boldsymbol{\ell}$$

$$= -\int_{\mathbb{T}^{3}} \left[\frac{d\varphi^{\varepsilon}(\ell)}{d\ell} \hat{\boldsymbol{\ell}}_{j} + \varphi^{\varepsilon}(\ell) \frac{2}{|\boldsymbol{\ell}|} \hat{\boldsymbol{\ell}}_{j} \right] P(\boldsymbol{x} + \boldsymbol{\ell}) d^{3} \boldsymbol{\ell}.$$
(2.12)

On the other hand, in light of the definition of P_L^{ε} and (2.10), we have

$$\nabla \varphi_L^{\varepsilon}(\ell) = \left(\frac{d\varphi^{\varepsilon}}{d\ell}(\ell) + \frac{2}{|\ell|}\varphi^{\varepsilon}(\ell)\right)\hat{\ell}.$$
(2.13)

Combining (2.12), (2.13) and using the integration by parts again, we can deduce that

$$\operatorname{div} \mathbf{\Pi}_{L}^{\varepsilon} = -\int_{\mathbb{T}^{3}} \partial_{\ell_{j}} \varphi_{L}^{\varepsilon}(\ell) P(\boldsymbol{x}+\boldsymbol{\ell},t) d^{3}\ell = \partial_{x_{j}} \int_{\mathbb{T}^{3}} \varphi_{L}^{\varepsilon}(\ell) P(\boldsymbol{x}+\boldsymbol{\ell},t) d^{3}\ell = \nabla P_{L}^{\varepsilon}(\boldsymbol{x}). \quad (2.14)$$

Similarly, since $\mathbf{\Pi}_{L}^{\varepsilon} + \mathbf{\Pi}_{T}^{\varepsilon} = P^{\varepsilon}\mathbf{1}$ and $P_{L}^{\varepsilon} + P_{T}^{\varepsilon} = P^{\varepsilon}$, we can also obtain div $\mathbf{\Pi}_{T}^{\varepsilon} = \nabla P_{T}^{\varepsilon}$. Then we have proved the claim (2.8).

Now, thanks to (2.8), the equations (2.4) and (2.6) can be further simplified as follows

$$\partial_t \boldsymbol{v}_X^{\varepsilon} + \operatorname{div}(\boldsymbol{v} \otimes \boldsymbol{v}_X)^{\varepsilon} + \nabla P_X^{\varepsilon} = 0, \quad X = L, T, \\ \partial_t \boldsymbol{\omega}_X^{\varepsilon} + \operatorname{div}(\boldsymbol{v} \otimes \boldsymbol{\omega}_X)^{\varepsilon} - \operatorname{div}(\boldsymbol{\omega} \otimes \boldsymbol{v}_X)^{\varepsilon} = 0, \quad X = L, T.$$
(2.15)

Next, our task is to show v_L^{ε} , v_T^{ε} and ω_L^{ε} , ω_T^{ε} are divergence-free in the sense of distributions.

Actually, for any test function $\psi(\boldsymbol{x}) \in C_0^{\infty}(\mathbb{T}^3)$, it follows from a straitforward compu-

tation that

$$\int_{\mathbb{T}^{3}} \psi(\boldsymbol{x}) \operatorname{div} \boldsymbol{v}_{L}^{\varepsilon}(\boldsymbol{x}, t) d^{3}\boldsymbol{x} = \int_{\mathbb{T}^{3}} \psi(\boldsymbol{x}) \partial_{x_{i}} \int_{\mathbb{T}^{3}} \varphi^{\varepsilon}(\ell) \hat{\boldsymbol{\ell}}_{i} \hat{\boldsymbol{\ell}}_{j} \boldsymbol{v}(\boldsymbol{x} + \boldsymbol{\ell}) d^{3} \boldsymbol{\ell} d^{3} \boldsymbol{x}$$

$$= \int_{\mathbb{T}^{3}} \int_{\mathbb{T}^{3}} \psi(\boldsymbol{x}) \varphi^{\varepsilon}(\ell) \hat{\boldsymbol{\ell}}_{i} \hat{\boldsymbol{\ell}}_{j} \partial_{x_{i}} v_{j}(\boldsymbol{x} + \boldsymbol{\ell}) d^{3} \boldsymbol{\ell} d^{3} \boldsymbol{x}$$

$$= -\int_{\mathbb{T}^{3}} \int_{\mathbb{T}^{3}} \partial_{\ell_{i}} \left[\varphi^{\varepsilon}(\ell) \hat{\boldsymbol{\ell}}_{i} \hat{\boldsymbol{\ell}}_{j} \right] \psi(\boldsymbol{x}) v_{j}(\boldsymbol{x} + \boldsymbol{\ell}) d^{3} \boldsymbol{x} d^{3} \boldsymbol{\ell}$$

$$= \int_{\mathbb{T}^{3}} \varphi^{\varepsilon}_{L}(\ell) \operatorname{div} \left(\psi * \boldsymbol{v} \right) (\boldsymbol{\ell}) d^{3} \boldsymbol{\ell} = 0,$$

$$(2.16)$$

where in the last equality we have used the fact that \boldsymbol{v} is divergence-free in the sense of distributions. Thus, we have shown that $\operatorname{div} \boldsymbol{v}_L^{\varepsilon} = 0$ in the sense of distributions. Moreover, according to the fact $\boldsymbol{v}_L^{\varepsilon} + \boldsymbol{v}_T^{\varepsilon} = \boldsymbol{v}^{\varepsilon}$, it follows that $\operatorname{div} \boldsymbol{v}_T^{\varepsilon} = 0$. Using the same argument as above, we can easily carry out $\operatorname{div} \boldsymbol{\omega}_L^{\varepsilon} = \operatorname{div} \boldsymbol{\omega}_T^{\varepsilon} = 0$ in the sense of distributions.

Then we are going to establish the local longitudinal Kármárth-Howarth equations for the helicity of Euler equations. Multiplying $(2.15)_1, (1.7)_2, (2.15)_2, (1.7)_1$ with X = L by $\boldsymbol{\omega}, \boldsymbol{v}_L^{\varepsilon}, \boldsymbol{v}, \boldsymbol{\omega}_L^{\varepsilon}$, respectively, we have

$$\boldsymbol{\omega} \cdot \partial_t \boldsymbol{v}_L^{\varepsilon} + \boldsymbol{\omega} \cdot \operatorname{div}(\boldsymbol{v} \otimes \boldsymbol{v}_L)^{\varepsilon} + \boldsymbol{\omega} \cdot \nabla P_L^{\varepsilon} = 0, \qquad (2.17)$$

$$\boldsymbol{v}_{L}^{\varepsilon} \cdot \partial_{t} \boldsymbol{\omega} + \boldsymbol{v}_{L}^{\varepsilon} \cdot \operatorname{div}(\boldsymbol{v} \otimes \boldsymbol{\omega}) - \boldsymbol{v}_{L}^{\varepsilon} \cdot \operatorname{div}(\boldsymbol{\omega} \otimes \boldsymbol{v}) = 0, \qquad (2.18)$$

$$\boldsymbol{v} \cdot \partial_t \boldsymbol{\omega}_L^{\varepsilon} + \boldsymbol{v} \cdot \operatorname{div}(\boldsymbol{v} \otimes \boldsymbol{\omega}_L)^{\varepsilon} - \boldsymbol{v} \cdot \operatorname{div}(\boldsymbol{\omega} \otimes \boldsymbol{v}_L)^{\varepsilon} = 0, \qquad (2.19)$$

$$\boldsymbol{\omega}_{L}^{\varepsilon} \cdot \partial_{t} \boldsymbol{v} + \boldsymbol{\omega}_{L}^{\varepsilon} \cdot \operatorname{div}(\boldsymbol{v} \otimes \boldsymbol{v}) + \boldsymbol{\omega}_{L}^{\varepsilon} \cdot \nabla P = 0.$$
(2.20)

On account of the Leibniz formula and divergence-free condition, we infer that

$$\boldsymbol{\omega} \cdot \operatorname{div}(\boldsymbol{v} \otimes \boldsymbol{v}_L)^{\varepsilon} + \boldsymbol{v}_L^{\varepsilon} \cdot \operatorname{div}(\boldsymbol{v} \otimes \boldsymbol{\omega}) = \operatorname{div}[\boldsymbol{v}(\boldsymbol{\omega} \cdot \boldsymbol{v}_L^{\varepsilon})] + \omega_j \cdot \partial_i \Big[(v_i v_{L_j})^{\varepsilon} - (v_i v_{L_j}^{\varepsilon}) \Big], \\ - \boldsymbol{v}_L^{\varepsilon} \cdot \operatorname{div}(\boldsymbol{\omega} \otimes \boldsymbol{v}) - \boldsymbol{v} \cdot \operatorname{div}(\boldsymbol{\omega} \otimes \boldsymbol{v}_L)^{\varepsilon} = -\operatorname{div}[\boldsymbol{\omega}(\boldsymbol{v} \cdot \boldsymbol{v}_L^{\varepsilon})] - v_j \partial_i \Big[(\omega_i v_{L_j})^{\varepsilon} - \omega_i v_{L_j}^{\varepsilon} \Big], \\ \boldsymbol{v} \cdot \operatorname{div}(\boldsymbol{v} \otimes \boldsymbol{\omega}_L)^{\varepsilon} + \boldsymbol{\omega}_L^{\varepsilon} \cdot \operatorname{div}(\boldsymbol{v} \otimes \boldsymbol{v}) = \operatorname{div}[\boldsymbol{v}(\boldsymbol{v} \cdot \boldsymbol{\omega}_L^{\varepsilon})] + v_j \partial_i \Big[(v_i \omega_{L_j})^{\varepsilon} - (v_i \omega_{L_j}^{\varepsilon}) \Big].$$

With this in hand, after a few computations, we get

$$\partial_{t}(\boldsymbol{\omega}\cdot\boldsymbol{v}_{L}^{\varepsilon}+\boldsymbol{v}\cdot\boldsymbol{\omega}_{L}^{\varepsilon})+\operatorname{div}(\boldsymbol{\omega}P_{L}^{\varepsilon}+P\boldsymbol{\omega}_{L}^{\varepsilon})+\operatorname{div}\left[\boldsymbol{v}(\boldsymbol{\omega}\cdot\boldsymbol{v}_{L}^{\varepsilon})-\boldsymbol{\omega}(\boldsymbol{v}\cdot\boldsymbol{v}_{L}^{\varepsilon})+\boldsymbol{v}(\boldsymbol{v}\cdot\boldsymbol{\omega}_{L}^{\varepsilon})\right]$$
$$=\omega_{j}\partial_{i}\left[(v_{i}v_{L_{j}}^{\varepsilon})-(v_{i}v_{L_{j}})^{\varepsilon}\right]+v_{j}\partial_{i}\left[(v_{i}\boldsymbol{\omega}_{L_{j}}^{\varepsilon})-(v_{i}\boldsymbol{\omega}_{L_{j}})^{\varepsilon}\right]+v_{j}\partial_{i}\left[(\omega_{i}v_{L_{j}})^{\varepsilon}-(\omega_{i}v_{L_{j}}^{\varepsilon})\right].$$
(2.21)

To process further, we notice that

$$\delta \boldsymbol{v}_L(\ell) = (\hat{\boldsymbol{\ell}} \otimes \hat{\boldsymbol{\ell}}) \delta \boldsymbol{v} = \hat{\boldsymbol{\ell}}_i(\hat{\boldsymbol{\ell}}_j \delta v_j) = (\delta v_{L_1}, \delta v_{L_2}, \delta v_{L_3}), \text{ with } \delta v_{L_i} = \hat{\boldsymbol{\ell}}_i(\hat{\boldsymbol{\ell}}_j \delta v_j).$$
(2.22)

And for any vectors $\boldsymbol{E} = (e_1, e_2, e_3)$ and $\boldsymbol{F} = (f_1, f_2, f_3)$, it is easy to verify that

$$\delta \boldsymbol{E}_{L} \cdot \delta \boldsymbol{F}_{L} = \hat{\boldsymbol{\ell}}_{i}(\hat{\boldsymbol{\ell}}_{j}\delta e_{j}) \cdot \hat{\boldsymbol{\ell}}_{i}(\hat{\boldsymbol{\ell}}_{k}\delta f_{k}) = \hat{\boldsymbol{\ell}}_{j}\hat{\boldsymbol{\ell}}_{k}\delta e_{j}\delta f_{k} = \hat{\boldsymbol{\ell}}_{j}\hat{\boldsymbol{\ell}}_{i}\delta e_{j}\delta f_{i} = \hat{\boldsymbol{\ell}}_{i}\hat{\boldsymbol{\ell}}_{j}\delta e_{i}\delta f_{j}$$
(2.23)

$$\delta \boldsymbol{E}_T \cdot \delta \boldsymbol{F}_T = \delta \boldsymbol{e}_{T_i} \cdot \delta f_{T_i}$$

= $(\delta^{ij} - \hat{\boldsymbol{\ell}}_i \hat{\boldsymbol{\ell}}_j) \delta \boldsymbol{e}_j \cdot (\delta^{ik} - \hat{\boldsymbol{\ell}}_i \hat{\boldsymbol{\ell}}_k) \delta f_k$
= $(\delta^{ij} - \hat{\boldsymbol{\ell}}_i \hat{\boldsymbol{\ell}}_j) \delta \boldsymbol{e}_i \delta f_j.$ (2.24)

Combining (2.23) and (2.24), we can obtain

$$\int_{\mathbb{T}^3} \nabla \varphi^{\varepsilon}(\ell) \cdot \delta \boldsymbol{\omega}(\boldsymbol{\ell}) [\delta \boldsymbol{v}_L(\boldsymbol{\ell})]^2 + \frac{2}{|\boldsymbol{\ell}|} \varphi^{\varepsilon}(\ell) \hat{\boldsymbol{\ell}} \cdot \delta \boldsymbol{\omega}(\boldsymbol{\ell}) [\delta \boldsymbol{v}_T(\boldsymbol{\ell})]^2 d^3 \boldsymbol{\ell}$$

$$= \int_{\mathbb{T}^3} \frac{\partial \varphi^{\varepsilon}}{\partial_{\ell_k}} \cdot \delta \omega_k \hat{\boldsymbol{\ell}}_i \hat{\boldsymbol{\ell}}_j \delta v_i \delta v_j + \varphi^{\varepsilon} \hat{\boldsymbol{\ell}}_k \delta \omega_k (\frac{\partial \hat{\boldsymbol{\ell}}_i}{\partial \ell_j} + \frac{\partial \hat{\boldsymbol{\ell}}_j}{\partial \ell_i}) \delta v_i \delta v_j d^3 \boldsymbol{\ell},$$
(2.25)

where we have used (2.11).

Then it follows from the Leibniz formula that

$$\int_{\mathbb{T}^{3}} \nabla \varphi^{\varepsilon}(\ell) \cdot \delta \boldsymbol{\omega}(\boldsymbol{\ell}) [\delta \boldsymbol{v}_{L}(\boldsymbol{\ell})]^{2} + \frac{2}{|\boldsymbol{\ell}|} \varphi^{\varepsilon}(\ell) \hat{\boldsymbol{\ell}} \cdot \delta \boldsymbol{\omega}(\boldsymbol{\ell}) [\delta \boldsymbol{v}_{T}(\boldsymbol{\ell})]^{2} d^{3} \boldsymbol{\ell} \\
= \int_{\mathbb{T}^{3}} \frac{\partial}{\partial_{\ell_{k}}} \Big(\varphi^{\varepsilon} \hat{\boldsymbol{\ell}}_{i} \hat{\boldsymbol{\ell}}_{j} \Big) \delta \omega_{k} \delta v_{i} \delta v_{j} - \varphi^{\varepsilon} \Big[\frac{\partial}{\partial_{\ell_{k}}} (\hat{\boldsymbol{\ell}}_{i} \hat{\boldsymbol{\ell}}_{j}) - (\frac{\partial \hat{\boldsymbol{\ell}}_{i}}{\partial \ell_{j}} + \frac{\partial \hat{\boldsymbol{\ell}}_{j}}{\partial \ell_{i}}) \hat{\boldsymbol{\ell}}_{k} \Big] \delta \omega_{k} \delta v_{i} \delta v_{j} d^{3} \boldsymbol{\ell}.$$
(2.26)

Before going any further, assume we have proved that, for any vector fields $\mathbf{A} = (A_1, A_2, A_3), \mathbf{B} = (B_1, B_2, B_3)$ and $\mathbf{C} = (C_1, C_2, C_3)$, there holds

$$\begin{bmatrix} \frac{\partial}{\partial \ell_k} (\hat{\ell}_i \hat{\ell}_j) - (\frac{\partial \hat{\ell}_i}{\partial \ell_j} + \frac{\partial \hat{\ell}_j}{\partial \ell_i}) \hat{\ell}_k \end{bmatrix} A_k B_i C_j$$

= $\frac{1}{|\ell|} \hat{\ell} \cdot \left[\mathbf{C} (\mathbf{A} \cdot \mathbf{B}) + \mathbf{B} (\mathbf{A} \cdot \mathbf{C}) - 2\mathbf{A} (\mathbf{B} \cdot \mathbf{C}) \right].$ (2.27)

As a consequent, by choosing $\mathbf{A} = \delta \boldsymbol{\omega}, \mathbf{B} = \mathbf{C} = \delta \boldsymbol{v}$ in (2.27), we have

$$\begin{bmatrix} \frac{\partial}{\partial \ell_{k}} (\hat{\boldsymbol{\ell}}_{i} \hat{\boldsymbol{\ell}}_{j}) - (\frac{\partial \hat{\boldsymbol{\ell}}_{i}}{\partial \ell_{j}} + \frac{\partial \hat{\boldsymbol{\ell}}_{j}}{\partial \ell_{i}}) \hat{\boldsymbol{\ell}}_{k} \end{bmatrix} \delta \omega_{k} \delta v_{i} \delta v_{j} \\
= \frac{2}{|\boldsymbol{\ell}|} \hat{\boldsymbol{\ell}} \cdot \left[\delta \boldsymbol{v} (\delta \boldsymbol{\omega} \cdot \delta \boldsymbol{v}) - \delta \boldsymbol{\omega} (\delta \boldsymbol{v} \cdot \delta \boldsymbol{v}) \right].$$
(2.28)

Substituting (2.28) into (2.26), it yields that

$$\int_{\mathbb{T}^{3}} \nabla \varphi^{\varepsilon}(\ell) \cdot \delta \boldsymbol{\omega}(\ell) [\delta \boldsymbol{v}_{L}(\ell)]^{2} + \frac{2}{|\boldsymbol{\ell}|} \varphi^{\varepsilon}(\ell) \hat{\boldsymbol{\ell}} \cdot \delta \boldsymbol{\omega}(\ell) [\delta \boldsymbol{v}_{T}(\boldsymbol{\ell})]^{2} d^{3}\boldsymbol{\ell} \\
= \int_{\mathbb{T}^{3}} \frac{\partial}{\partial_{\ell_{k}}} \Big(\varphi^{\varepsilon} \hat{\boldsymbol{\ell}}_{i} \hat{\boldsymbol{\ell}}_{j} \Big) \delta \omega_{k} \delta v_{i} \delta v_{j} - \frac{2}{|\boldsymbol{\ell}|} \varphi^{\varepsilon} \hat{\boldsymbol{\ell}} \cdot \Big[\delta \boldsymbol{v}(\delta \boldsymbol{\omega} \cdot \delta \boldsymbol{v}) - \delta \boldsymbol{\omega}(\delta \boldsymbol{v} \cdot \delta \boldsymbol{v}) \Big] d^{3}\boldsymbol{\ell},$$
(2.29)

Next, for simplicity of presentation, we denote

$$\boldsymbol{v}(\boldsymbol{x}+\boldsymbol{\ell}) = \bar{\boldsymbol{v}} = (\bar{v}_1, \bar{v}_2, \bar{v}_3), \boldsymbol{\omega}(\boldsymbol{x}+\boldsymbol{\ell}) = \bar{\boldsymbol{\omega}} = (\bar{\omega}_1, \bar{\omega}_2, \bar{\omega}_3).$$
(2.30)

A routine computation gives rise to

$$\int_{\mathbb{T}^{3}} \frac{\partial}{\partial_{\ell_{k}}} \left(\varphi^{\varepsilon} \hat{\ell}_{i} \hat{\ell}_{j}\right) \delta\omega_{k} \delta v_{i} \delta v_{j} d^{3} \ell$$

$$= \int_{\mathbb{T}^{3}} \frac{\partial}{\partial_{\ell_{k}}} \left(\varphi^{\varepsilon} \hat{\ell}_{i} \hat{\ell}_{j}\right) \left[\omega_{k}(\boldsymbol{x}+\boldsymbol{\ell}) - \omega_{k}(\boldsymbol{x})\right] \left[v_{i}(\boldsymbol{x}+\boldsymbol{\ell}) - v_{i}(\boldsymbol{x})\right] \left[v_{j}(\boldsymbol{x}+\boldsymbol{\ell}) - v_{j}(\boldsymbol{x})\right] d^{3} \ell$$

$$= \int_{\mathbb{T}^{3}} \frac{\partial}{\partial_{\ell_{k}}} \left(\varphi^{\varepsilon} \hat{\ell}_{i} \hat{\ell}_{j}\right)$$

$$\times \left(\bar{\omega}_{k} \bar{v}_{i} \bar{v}_{j} - \bar{\omega}_{k} \bar{v}_{i} v_{j} - \bar{\omega}_{k} v_{i} \bar{v}_{j} + \bar{\omega}_{k} v_{i} v_{j} - \omega_{k} \bar{v}_{i} \bar{v}_{j} + \omega_{k} \bar{v}_{i} \bar{v}_{j} + \omega_{k} \bar{v}_{i} v_{j} - \omega_{k} v_{i} v_{j}\right) d^{3} \ell.$$
(2.31)

Then we will deal with the terms on the right-hand side of (2.31) one by one. For example, according to integration by parts and (2.23), we get

$$\int_{\mathbb{T}^{3}} \frac{\partial}{\partial_{\ell_{k}}} \left(\varphi^{\varepsilon} \hat{\ell}_{i} \hat{\ell}_{j} \right) \bar{\omega}_{k} \bar{v}_{i} \bar{v}_{j} d^{3} \ell$$

$$= -\int_{\mathbb{T}^{3}} \varphi^{\varepsilon} \hat{\ell}_{i} \hat{\ell}_{j} \frac{\partial}{\partial_{\ell_{k}}} \left(\bar{\omega}_{k} \bar{v}_{i} \bar{v}_{j} \right) d^{3} \ell$$

$$= -\frac{\partial}{\partial_{x_{k}}} \int_{\mathbb{T}^{3}} \varphi^{\varepsilon} \hat{\ell}_{i} \hat{\ell}_{j} \bar{\omega}_{k} \bar{v}_{i} \bar{v}_{j} d^{3} \ell$$

$$= -\operatorname{div} \left(\boldsymbol{\omega} (\boldsymbol{v}_{L} \cdot \boldsymbol{v}_{L}) \right)^{\varepsilon}.$$
(2.32)

Performing the same procedure for the rest terms on the right-hand side of (2.31), we have

$$-\int_{\mathbb{T}^{3}} \frac{\partial}{\partial_{\ell_{k}}} \Big[\varphi^{\varepsilon} \hat{\ell}_{i} \hat{\ell}_{j} \Big] \bar{\omega}_{k} \bar{v}_{i} v_{j} d^{3} \ell = v_{j} \partial_{k} (\omega_{k} v_{L_{j}})^{\varepsilon},$$

$$-\int_{\mathbb{T}^{3}} \frac{\partial}{\partial_{\ell_{k}}} \Big[\varphi^{\varepsilon} \hat{\ell}_{i} \hat{\ell}_{j} \Big] \bar{\omega}_{k} v_{i} \bar{v}_{j} d^{3} \ell = v_{i} \partial_{k} (\omega_{k} v_{L_{i}})^{\varepsilon},$$

$$\int_{\mathbb{T}^{3}} \frac{\partial}{\partial_{\ell_{k}}} \Big[\varphi^{\varepsilon} \hat{\ell}_{i} \hat{\ell}_{j} \Big] \bar{\omega}_{k} v_{i} v_{j} d^{3} \ell = -(\boldsymbol{v}_{L} \cdot \boldsymbol{v}_{L}) \partial_{k} (\omega_{k})^{\varepsilon} = 0,$$

$$-\int_{\mathbb{T}^{3}} \frac{\partial}{\partial_{\ell_{k}}} \Big[\varphi^{\varepsilon} \hat{\ell}_{i} \hat{\ell}_{j} \Big] \omega_{k} \bar{v}_{i} \bar{v}_{j} d^{3} \ell = \omega_{k} \partial_{k} (\boldsymbol{v}_{L} \cdot \boldsymbol{v}_{L})^{\varepsilon} = \operatorname{div} \Big[\boldsymbol{\omega} (\boldsymbol{v}_{L} \cdot \boldsymbol{v}_{L})^{\varepsilon} \Big], \qquad (2.33)$$

$$\int_{\mathbb{T}^{3}} \frac{\partial}{\partial_{\ell_{k}}} \Big[\varphi^{\varepsilon} \hat{\ell}_{i} \hat{\ell}_{j} \Big] \omega_{k} \bar{v}_{i} v_{j} d^{3} \ell = -\omega_{k} v_{j} \partial_{k} (v_{L_{j}})^{\varepsilon} = -v_{j} \partial_{k} (\omega_{k} v_{L_{j}}^{\varepsilon}),$$

$$\int_{\mathbb{T}^{3}} \frac{\partial}{\partial_{\ell_{k}}} \Big[\varphi^{\varepsilon} \hat{\ell}_{i} \hat{\ell}_{j} \Big] \omega_{k} v_{i} \bar{v}_{j} d^{3} \ell = -\omega_{k} v_{i} \partial_{k} v_{L_{i}}^{\varepsilon} = -v_{i} \partial_{k} (\omega_{k} v_{L_{j}}^{\varepsilon}),$$

$$-\int_{\mathbb{T}^{3}} \frac{\partial}{\partial_{\ell_{k}}} \Big[\varphi^{\varepsilon} \hat{\ell}_{i} \hat{\ell}_{j} \Big] \omega_{k} v_{i} \bar{v}_{j} d^{3} \ell = 0,$$

where the divergence-free condition of the velocity and the vorticity was used many times. Then plugging (2.32) and (2.33) into (2.31), we can obtain

$$\int_{\mathbb{T}^3} \frac{\partial}{\partial_{\ell_k}} \Big[\varphi^{\varepsilon} \hat{\boldsymbol{\ell}}_i \hat{\boldsymbol{\ell}}_j \Big] \delta\omega_k \delta v_i \delta v_j d^3 \boldsymbol{\ell}$$

$$= -\operatorname{div} \Big[\Big(\boldsymbol{\omega} (\boldsymbol{v}_L \cdot \boldsymbol{v}_L) \Big)^{\varepsilon} - \boldsymbol{\omega} (\boldsymbol{v}_L \cdot \boldsymbol{v}_L)^{\varepsilon} \Big] + 2v_j \partial_k \Big[(\omega_k v_{L_j})^{\varepsilon} - (\omega_k v_{L_j}^{\varepsilon}) \Big].$$
(2.34)

Furthermore, by inserting (2.34) into (2.29), we observe that

$$\frac{1}{2} \int_{\mathbb{T}^3} \nabla \varphi^{\varepsilon}(\ell) \cdot \delta \boldsymbol{\omega} [\delta \boldsymbol{v}_L]^2 + \frac{2}{|\boldsymbol{\ell}|} \varphi^{\varepsilon}(\ell) \hat{\boldsymbol{\ell}} \cdot \left[\delta \boldsymbol{\omega} [\delta \boldsymbol{v}_T]^2 + \delta \boldsymbol{v} (\delta \boldsymbol{\omega} \cdot \delta \boldsymbol{v}) - \delta \boldsymbol{\omega} (\delta \boldsymbol{v} \cdot \delta \boldsymbol{v}) \right] d^3 \boldsymbol{\ell}
+ \frac{1}{2} \operatorname{div} \left[\left(\boldsymbol{\omega} (\boldsymbol{v}_L \cdot \boldsymbol{v}_L) \right)^{\varepsilon} - \boldsymbol{\omega} (\boldsymbol{v}_L \cdot \boldsymbol{v}_L)^{\varepsilon} \right]
= v_j \partial_k \left[\left(\omega_k v_{L_j} \right)^{\varepsilon} - \left(\omega_k v_{L_j}^{\varepsilon} \right) \right].$$
(2.35)

Along the same lines as (2.26), we find

$$\int_{\mathbb{T}^{3}} \nabla \varphi^{\varepsilon}(\ell) \cdot \delta \boldsymbol{v}(\ell) (\delta \boldsymbol{v}_{L} \cdot \delta \boldsymbol{\omega}_{L}) + \frac{2}{|\boldsymbol{\ell}|} \varphi^{\varepsilon}(\ell) \hat{\boldsymbol{\ell}} \cdot \delta \boldsymbol{v}(\boldsymbol{\ell}) (\delta \boldsymbol{v}_{T} \cdot \delta \boldsymbol{\omega}_{T}) d^{3} \boldsymbol{\ell}$$

$$= \int_{\mathbb{T}^{3}} \frac{\partial}{\partial_{\ell_{k}}} \Big(\varphi^{\varepsilon} \hat{\boldsymbol{\ell}}_{i} \hat{\boldsymbol{\ell}}_{j} \Big) \delta v_{k} \delta v_{i} \delta \omega_{j} - \varphi^{\varepsilon} \Big[\frac{\partial}{\partial_{\ell_{k}}} (\hat{\boldsymbol{\ell}}_{i} \hat{\boldsymbol{\ell}}_{j}) - (\frac{\partial \hat{\boldsymbol{\ell}}_{i}}{\partial \ell_{j}} + \frac{\partial \hat{\boldsymbol{\ell}}_{j}}{\partial \ell_{i}}) \hat{\boldsymbol{\ell}}_{k} \Big] \delta v_{k} \delta v_{i} \delta \omega_{j} d^{3} \boldsymbol{\ell}$$

$$(2.36)$$

From (2.27), we reformulate the above equation as

$$\int_{\mathbb{T}^3} \nabla \varphi^{\varepsilon}(\ell) \cdot \delta \boldsymbol{v}(\boldsymbol{\ell}) (\delta \boldsymbol{v}_L \cdot \delta \boldsymbol{\omega}_L) + \frac{2}{|\boldsymbol{\ell}|} \varphi^{\varepsilon}(\ell) \hat{\boldsymbol{\ell}} \cdot \delta \boldsymbol{v}(\boldsymbol{\ell}) (\delta \boldsymbol{v}_T \cdot \delta \boldsymbol{\omega}_T) d^3 \boldsymbol{\ell}$$

$$= \int_{\mathbb{T}^3} \frac{\partial}{\partial_{\ell_k}} \Big(\varphi^{\varepsilon} \hat{\boldsymbol{\ell}}_i \hat{\boldsymbol{\ell}}_j \Big) \delta v_k \delta v_i \delta \omega_j + \frac{1}{|\boldsymbol{\ell}|} \varphi^{\varepsilon} \hat{\boldsymbol{\ell}} \cdot \Big[\delta \boldsymbol{v}(\delta \boldsymbol{\omega} \cdot \delta \boldsymbol{v}) - \delta \boldsymbol{\omega}(\delta \boldsymbol{v} \cdot \delta \boldsymbol{v}) \Big] d^3 \boldsymbol{\ell}.$$
(2.37)

Repeating the deduction process of (2.34), we arrive at

$$\int_{\mathbb{T}^3} \frac{\partial}{\partial_{\ell_k}} \Big(\varphi^{\varepsilon} \hat{\ell}_i \hat{\ell}_j \Big) \delta v_k \delta v_i \delta \omega_j d^3 \ell$$

= $- \operatorname{div} \Big[\Big(\boldsymbol{v} (\boldsymbol{v}_L \cdot \boldsymbol{\omega}_L) \Big)^{\varepsilon} - \boldsymbol{v} (\boldsymbol{v}_L \cdot \boldsymbol{\omega}_L)^{\varepsilon} \Big] + \omega_j \partial_k \Big[(v_k v_{L_j})^{\varepsilon} - (v_k v_{L_j}^{\varepsilon}) \Big] + v_i \partial_k \Big[(v_k \omega_{L_i})^{\varepsilon} - (v_k \omega_{L_i}^{\varepsilon}) \Big]$ (2.38)

Then, it follows from (2.37) and (2.38) that

$$\int_{\mathbb{T}^{3}} \nabla \varphi^{\varepsilon}(\ell) \cdot \delta \boldsymbol{v}(\delta \boldsymbol{v}_{L} \cdot \delta \boldsymbol{\omega}_{L}) + \frac{2}{|\boldsymbol{\ell}|} \varphi^{\varepsilon}(\ell) \hat{\boldsymbol{\ell}} \cdot \left[\delta \boldsymbol{v}(\delta \boldsymbol{v}_{T} \cdot \delta \boldsymbol{\omega}_{T}) - \frac{1}{2} \left(\delta \boldsymbol{v}(\delta \boldsymbol{\omega} \cdot \delta \boldsymbol{v}) - \delta \boldsymbol{\omega}(\delta \boldsymbol{v} \cdot \delta \boldsymbol{v}) \right) \right] d^{3} \boldsymbol{\ell} + \operatorname{div} \left[\left(\boldsymbol{v}(\boldsymbol{v}_{L} \cdot \boldsymbol{\omega}_{L}) \right)^{\varepsilon} - \boldsymbol{v}(\boldsymbol{v}_{L} \cdot \boldsymbol{\omega}_{L})^{\varepsilon} \right] = \omega_{j} \partial_{k} \left[\left(v_{k} v_{L_{j}} \right)^{\varepsilon} - \left(v_{k} v_{L_{j}}^{\varepsilon} \right) \right] + v_{i} \partial_{k} \left[\left(v_{k} \omega_{L_{i}} \right)^{\varepsilon} - \left(v_{k} \omega_{L_{i}}^{\varepsilon} \right) \right].$$

$$(2.39)$$

Finally, substituting (2.35) and (2.39) into (2.21), we conclude that

$$\partial_{t}(\boldsymbol{\omega}\cdot\boldsymbol{v}_{L}^{\varepsilon}+\boldsymbol{v}\cdot\boldsymbol{\omega}_{L}^{\varepsilon})+\operatorname{div}(\boldsymbol{\omega}P_{L}^{\varepsilon}+P\boldsymbol{\omega}_{L}^{\varepsilon})+\operatorname{div}\left[\boldsymbol{v}(\boldsymbol{\omega}\cdot\boldsymbol{v}_{L}^{\varepsilon})-\boldsymbol{\omega}(\boldsymbol{v}\cdot\boldsymbol{v}_{L}^{\varepsilon})+\boldsymbol{v}(\boldsymbol{v}\cdot\boldsymbol{\omega}_{L}^{\varepsilon})\right] \\ +\operatorname{div}\left[\left(\boldsymbol{v}(\boldsymbol{v}_{L}\cdot\boldsymbol{\omega}_{L})\right)^{\varepsilon}-\boldsymbol{v}(\boldsymbol{v}_{L}\cdot\boldsymbol{\omega}_{L})^{\varepsilon}\right]-\frac{1}{2}\operatorname{div}\left[\left(\boldsymbol{\omega}(\boldsymbol{v}_{L}\cdot\boldsymbol{v}_{L})\right)^{\varepsilon}-\boldsymbol{\omega}(\boldsymbol{v}_{L}\cdot\boldsymbol{v}_{L})^{\varepsilon}\right] \\ =-\frac{4}{3}D_{HL}^{\varepsilon}(\boldsymbol{v},\boldsymbol{\omega}), \tag{2.40}$$

where

$$D_{HL}^{\varepsilon}(\boldsymbol{v},\boldsymbol{\omega}) = \frac{3}{4} \int \nabla \varphi^{\varepsilon}(\ell) \cdot \delta \boldsymbol{v}(\delta \boldsymbol{v}_{L} \cdot \delta \boldsymbol{\omega}_{L}) + \frac{2}{|\boldsymbol{\ell}|} \varphi^{\varepsilon}(\ell) \hat{\boldsymbol{\ell}} \cdot \left[\delta \boldsymbol{v}(\delta \boldsymbol{v}_{T} \cdot \delta \boldsymbol{\omega}_{T}) + \delta \boldsymbol{v} \times (\delta \boldsymbol{\omega} \times \delta \boldsymbol{v}) \right] d^{3} \boldsymbol{\ell} \quad (2.41)$$
$$- \frac{3}{8} \int_{\mathbb{T}^{3}} \nabla \varphi^{\varepsilon}(\ell) \cdot \delta \boldsymbol{\omega} [\delta \boldsymbol{v}_{L}]^{2} + \frac{2}{|\boldsymbol{\ell}|} \varphi^{\varepsilon}(\ell) \hat{\boldsymbol{\ell}} \cdot \delta \boldsymbol{\omega} [\delta \boldsymbol{v}_{T}]^{2} d^{3} \boldsymbol{\ell},$$

and in the last equality we have used the following vector identity

$$\delta \boldsymbol{v} \times (\delta \boldsymbol{\omega} \times \delta \boldsymbol{v}) = \delta \boldsymbol{\omega} (\delta \boldsymbol{v} \cdot \delta \boldsymbol{v}) - \delta \boldsymbol{v} (\delta \boldsymbol{\omega} \cdot \delta \boldsymbol{v}).$$
(2.42)

Then we have proved the local longitudinal Kármárth-Howarth equations for helicity. Next, we are in a position to get the transverse Kármárth-Howarth equations for helicity. Actually, arguing in the same manner as in the derivation of (2.21), we have

$$\partial_{t}(\boldsymbol{\omega}\cdot\boldsymbol{v}_{T}^{\varepsilon}+\boldsymbol{v}\cdot\boldsymbol{\omega}_{T}^{\varepsilon})+\operatorname{div}(\boldsymbol{\omega}P_{T}^{\varepsilon}+P\boldsymbol{\omega}_{T}^{\varepsilon})+\operatorname{div}\left[\boldsymbol{v}(\boldsymbol{\omega}\cdot\boldsymbol{v}_{T}^{\varepsilon})+\boldsymbol{v}(\boldsymbol{v}\cdot\boldsymbol{\omega}_{T}^{\varepsilon})-\boldsymbol{\omega}(\boldsymbol{v}\cdot\boldsymbol{v}_{T}^{\varepsilon})\right]$$
$$=\omega_{j}\partial_{i}\left[(v_{i}v_{T_{j}}^{\varepsilon})-(v_{i}v_{T_{j}})^{\varepsilon}\right]+v_{j}\partial_{i}\left[(v_{i}\omega_{T_{j}}^{\varepsilon})-(v_{i}\omega_{T_{j}})^{\varepsilon}\right]+v_{j}\partial_{i}\left[(\omega_{i}v_{T_{j}})^{\varepsilon}-(\omega_{i}v_{T_{j}}^{\varepsilon})\right].$$
(2.43)

In view of (2.11), (2.24) and the Leibniz formula, we see that

$$\int_{\mathbb{T}^{3}} \nabla \varphi^{\varepsilon}(\ell) \cdot \delta \boldsymbol{\omega} [\delta \boldsymbol{v}_{T}]^{2} - \frac{2}{|\boldsymbol{\ell}|} \varphi^{\varepsilon} \hat{\boldsymbol{\ell}} \cdot \delta \boldsymbol{\omega} [\delta \boldsymbol{v}_{T}]^{2} d^{3} \boldsymbol{\ell} \\
= \int_{\mathbb{T}^{3}} \partial_{\ell_{k}} \Big[\varphi^{\varepsilon} (\delta^{ij} - \hat{\boldsymbol{\ell}}_{i} \hat{\boldsymbol{\ell}}_{j}) \Big] \delta \omega_{k} \delta v_{i} \delta v_{j} + \varphi^{\varepsilon} \Big[\partial_{\ell_{k}} (\hat{\boldsymbol{\ell}}_{i} \hat{\boldsymbol{\ell}}_{j}) - (\frac{\partial \hat{\boldsymbol{\ell}}_{i}}{\partial_{\ell_{j}}} + \frac{\partial \hat{\boldsymbol{\ell}}_{j}}{\partial_{\ell_{i}}}) \hat{\boldsymbol{\ell}}_{k} \Big] \delta \omega_{k} \delta v_{i} \delta v_{j} d^{3} \boldsymbol{\ell},$$
(2.44)

which together with (2.28) and (2.42) yields

$$\int_{\mathbb{T}^{3}} \nabla \varphi^{\varepsilon}(\ell) \cdot \delta \boldsymbol{\omega} [\delta \boldsymbol{v}_{T}]^{2} - \frac{2}{|\boldsymbol{\ell}|} \varphi^{\varepsilon} \hat{\boldsymbol{\ell}} \cdot \delta \boldsymbol{\omega} [\delta \boldsymbol{v}_{T}]^{2} d^{3} \boldsymbol{\ell}$$

$$= \int_{\mathbb{T}^{3}} \partial_{\ell_{k}} \Big[\varphi^{\varepsilon} (\delta^{ij} - \hat{\boldsymbol{\ell}}_{i} \hat{\boldsymbol{\ell}}_{j}) \Big] \delta \omega_{k} \delta v_{i} \delta v_{j} - \frac{2}{|\boldsymbol{\ell}|} \varphi^{\varepsilon} \hat{\boldsymbol{\ell}} \cdot \big[\delta \boldsymbol{v} \times (\delta \boldsymbol{\omega} \times \delta \boldsymbol{v}) \big] d^{3} \boldsymbol{\ell}$$

$$(2.45)$$

Likewise, combining (2.27) and (2.42), we know that

$$\int_{\mathbb{T}^{3}} \nabla \varphi^{\varepsilon}(\ell) \cdot \delta \boldsymbol{v} (\delta \boldsymbol{\omega}_{T} \cdot \delta \boldsymbol{v}_{T}) - \frac{2}{|\boldsymbol{\ell}|} \varphi^{\varepsilon} \hat{\boldsymbol{\ell}} \cdot \delta \boldsymbol{v} (\delta \boldsymbol{\omega}_{T} \cdot \delta \boldsymbol{v}_{T}) d^{3} \boldsymbol{\ell}$$

$$= \int_{\mathbb{T}^{3}} \partial_{\ell_{k}} \varphi^{\varepsilon} \cdot \delta v_{k} (\delta^{ij} - \hat{\boldsymbol{\ell}}_{i} \hat{\boldsymbol{\ell}}_{j}) \delta \omega_{i} \delta v_{j} - \frac{2}{|\boldsymbol{\ell}|} \varphi^{\varepsilon} \hat{\boldsymbol{\ell}}_{k} \cdot \delta v_{k} (\delta^{ij} - \hat{\boldsymbol{\ell}}_{i} \hat{\boldsymbol{\ell}}_{j}) \delta \omega_{i} \delta v_{j} d^{3} \boldsymbol{\ell}$$

$$= \int_{\mathbb{T}^{3}} \partial_{\ell_{k}} \Big[\varphi^{\varepsilon} (\delta^{ij} - \hat{\boldsymbol{\ell}}_{i} \hat{\boldsymbol{\ell}}_{j}) \Big] \delta v_{k} \delta \omega_{i} \delta v_{j} + \varphi^{\varepsilon} \Big[\partial_{\ell_{k}} (\hat{\boldsymbol{\ell}}_{i} \hat{\boldsymbol{\ell}}_{j}) - (\frac{\partial \hat{\boldsymbol{\ell}}_{i}}{\partial_{\ell_{j}}} + \frac{\partial \hat{\boldsymbol{\ell}}_{j}}{\partial_{\ell_{i}}}) \hat{\boldsymbol{\ell}}_{k} \Big] \delta v_{k} \delta \omega_{i} \delta v_{j} d^{3} \boldsymbol{\ell}$$

$$= \int_{\mathbb{T}^{3}} \partial_{\ell_{k}} \Big[\varphi^{\varepsilon} (\delta^{ij} - \hat{\boldsymbol{\ell}}_{i} \hat{\boldsymbol{\ell}}_{j}) \Big] \delta v_{k} \delta \omega_{i} \delta v_{j} + \frac{1}{|\boldsymbol{\ell}|} \varphi^{\varepsilon} \hat{\boldsymbol{\ell}} \cdot \big[\delta \boldsymbol{v} \times (\delta \boldsymbol{\omega} \times \delta \boldsymbol{v}) \big] d^{3} \boldsymbol{\ell}$$
(2.46)

To control the first term on the right-hand side of (2.46), it follows from a straightforward calculation that

$$\begin{split} &\int_{\mathbb{T}^3} \partial_{\ell_k} \Big[\varphi^{\varepsilon} (\delta^{ij} - \hat{\ell}_i \hat{\ell}_j) \Big] \delta v_k \delta \omega_i \delta v_j d^3 \ell \\ &= \int_{\mathbb{T}^3} \partial_{\ell_k} \Big[\varphi^{\varepsilon} (\delta^{ij} - \hat{\ell}_i \hat{\ell}_j) \Big] \\ &\times \big[\overline{v}_k \overline{\omega}_i \overline{v}_j - \overline{v}_k \overline{\omega}_i v_j - \overline{v}_k \omega_i \overline{v}_j + \overline{v}_k \omega_i v_j - v_k \overline{\omega}_i \overline{v}_j + v_k \overline{\omega}_i v_j - v_k \omega_i v_j \big] d^3 \ell , \end{split}$$

where we have used the notation (2.30). For example, by virtue of integration by parts and (2.24), we obtain

$$\int_{\mathbb{T}^{3}} \frac{\partial}{\partial \ell_{k}} \Big[\varphi^{\varepsilon} (\delta^{ij} - \hat{\ell}_{i} \hat{\ell}_{j}) \Big] \bar{v}_{k} \bar{\omega}_{i} \bar{v}_{j} d^{3} \ell = - \int_{\mathbb{T}^{3}} \Big[\varphi^{\varepsilon} (\delta^{ij} - \hat{\ell}_{i} \hat{\ell}_{j}) \Big] \frac{\partial}{\partial \ell_{k}} \Big(\bar{v}_{k} \bar{\omega}_{i} \bar{v}_{j} \Big) d^{3} \ell
= - \frac{\partial}{\partial x_{k}} \int_{\mathbb{T}^{3}} \Big[\varphi^{\varepsilon} (\delta^{ij} - \hat{\ell}_{i} \hat{\ell}_{j}) \Big] \bar{v}_{k} \bar{\omega}_{i} \bar{v}_{j} d^{3} \ell$$

$$= - \operatorname{div} \Big[\boldsymbol{v} (\boldsymbol{\omega}_{T} \cdot \boldsymbol{v}_{T}) \Big]^{\varepsilon}.$$
(2.47)

Similar to the above derivation, it follows that

$$-\int_{\mathbb{T}^{3}} \frac{\partial}{\partial_{\ell_{k}}} \Big[\varphi^{\varepsilon} (\delta^{ij} - \hat{\ell}_{i} \hat{\ell}_{j}) \Big] \bar{v}_{k} \bar{\omega}_{i} v_{j} d^{3} \ell = v_{j} \partial_{k} (v_{k} \omega_{T_{j}})^{\varepsilon}, -\int_{\mathbb{T}^{3}} \frac{\partial}{\partial_{\ell_{k}}} \Big[\varphi^{\varepsilon} (\delta^{ij} - \hat{\ell}_{i} \hat{\ell}_{j}) \Big] \bar{v}_{k} \omega_{i} \bar{v}_{j} d^{3} \ell = \omega_{i} \partial_{k} (v_{k} v_{T_{i}})^{\varepsilon}, \int_{\mathbb{T}^{3}} \frac{\partial}{\partial_{\ell_{k}}} \Big[\varphi^{\varepsilon} (\delta^{ij} - \hat{\ell}_{i} \hat{\ell}_{j}) \Big] \bar{v}_{k} \omega_{i} v_{j} d^{3} \ell = -\omega_{T} \cdot v_{T} \int_{\mathbb{T}^{3}} \varphi^{\varepsilon} \frac{\partial}{\partial_{\ell_{k}}} \bar{v}_{k} d^{3} \ell = 0, -\int_{\mathbb{T}^{3}} \frac{\partial}{\partial_{\ell_{k}}} \Big[\varphi^{\varepsilon} (\delta^{ij} - \hat{\ell}_{i} \hat{\ell}_{j}) \Big] v_{k} \bar{\omega}_{i} \bar{v}_{j} d^{3} \ell = v_{k} \partial_{k} (\omega_{T} \cdot v_{T})^{\varepsilon} = \operatorname{div} [\boldsymbol{v} (\omega_{T} \cdot v_{T})^{\varepsilon}],$$
(2.48)
$$\int_{\mathbb{T}^{3}} \frac{\partial}{\partial_{\ell_{k}}} \Big[\varphi^{\varepsilon} (\delta^{ij} - \hat{\ell}_{i} \hat{\ell}_{j}) \Big] v_{k} \bar{\omega}_{i} v_{j} d^{3} \ell = -v_{k} v_{j} \partial_{k} (\omega_{T_{j}})^{\varepsilon} = -v_{j} \partial_{k} (v_{k} \omega_{T_{j}}^{\varepsilon}), \int_{\mathbb{T}^{3}} \frac{\partial}{\partial_{\ell_{k}}} \Big[\varphi^{\varepsilon} (\delta^{ij} - \hat{\ell}_{i} \hat{\ell}_{j}) \Big] v_{k} \omega_{i} \bar{v}_{j} d^{3} \ell = -v_{k} \omega_{i} \partial_{k} v_{T_{i}}^{\varepsilon} = -\omega_{i} \partial_{k} (v_{k} v_{T_{i}}^{\varepsilon}), -\int_{\mathbb{T}^{3}} \frac{\partial}{\partial_{\ell_{k}}} \Big[\varphi^{\varepsilon} (\delta^{ij} - \hat{\ell}_{i} \hat{\ell}_{j}) \Big] v_{k} \omega_{i} v_{j} d^{3} \ell = 0.$$

Consequently, combining (2.46) through (2.48), we get

$$\int_{\mathbb{T}^3} \nabla \varphi^{\varepsilon}(\ell) \cdot \delta \boldsymbol{v} (\delta \boldsymbol{\omega}_T \cdot \delta \boldsymbol{v}_T) - \frac{2}{|\boldsymbol{\ell}|} \varphi^{\varepsilon} \hat{\boldsymbol{\ell}} \cdot \delta \boldsymbol{v} (\delta \boldsymbol{\omega}_T \cdot \delta \boldsymbol{v}_T) - \frac{1}{|\boldsymbol{\ell}|} \varphi^{\varepsilon} \hat{\boldsymbol{\ell}} \cdot \left[\delta \boldsymbol{v} \times (\delta \boldsymbol{\omega} \times \delta \boldsymbol{v}) \right] d^3 \boldsymbol{\ell}$$

$$= \int_{\mathbb{T}^3} \partial_{\ell_k} \left[\varphi^{\varepsilon} (\delta^{ij} - \hat{\boldsymbol{\ell}}_i \hat{\boldsymbol{\ell}}_j) \right] \delta v_k \delta \omega_i \delta v_j d^3 \boldsymbol{\ell}$$

$$= - \operatorname{div} \left[\left(\boldsymbol{v} (\boldsymbol{\omega}_T \cdot \boldsymbol{v}_T) \right)^{\varepsilon} - \boldsymbol{v} (\boldsymbol{\omega}_T \cdot \boldsymbol{v}_T)^{\varepsilon} \right] + v_j \partial_k \left[(v_k \omega_{T_j})^{\varepsilon} - (v_k \omega_{T_j}^{\varepsilon}) \right] + \omega_i \partial_k \left[(v_k v_{T_i})^{\varepsilon} - (v_k v_{T_i}^{\varepsilon}) \right],$$
(2.49)

which implies

$$\int_{\mathbb{T}^{3}} \nabla \varphi^{\varepsilon}(\ell) \cdot \delta \boldsymbol{v} (\delta \boldsymbol{\omega}_{T} \cdot \delta \boldsymbol{v}_{T}) - \frac{2}{|\boldsymbol{\ell}|} \varphi^{\varepsilon} \hat{\boldsymbol{\ell}} \cdot \delta \boldsymbol{v} (\delta \boldsymbol{\omega}_{T} \cdot \delta \boldsymbol{v}_{T}) - \frac{1}{|\boldsymbol{\ell}|} \varphi^{\varepsilon} \hat{\boldsymbol{\ell}} \cdot \left[\delta \boldsymbol{v} \times (\delta \boldsymbol{\omega} \times \delta \boldsymbol{v}) \right] d^{3} \boldsymbol{\ell} + \operatorname{div} \left[\left(\boldsymbol{v}(\boldsymbol{\omega}_{T} \cdot \boldsymbol{v}_{T}) \right)^{\varepsilon} - \boldsymbol{v}(\boldsymbol{\omega}_{T} \cdot \boldsymbol{v}_{T})^{\varepsilon} \right] = v_{j} \partial_{k} \left[\left(v_{k} \omega_{T_{j}} \right)^{\varepsilon} - \left(v_{k} \omega_{T_{j}}^{\varepsilon} \right) \right] + \omega_{i} \partial_{k} \left[\left(v_{k} v_{T_{i}} \right)^{\varepsilon} - \left(v_{k} v_{T_{i}}^{\varepsilon} \right) \right].$$

$$(2.50)$$

On the other hand, by a suitable modification to the derivation of (2.49), we infer that

$$\int_{\mathbb{T}^3} \nabla \varphi^{\varepsilon}(\ell) \cdot \delta \boldsymbol{\omega} [\delta \boldsymbol{v}_T]^2 - \frac{2}{|\boldsymbol{\ell}|} \varphi^{\varepsilon} \hat{\boldsymbol{\ell}} \cdot \delta \boldsymbol{\omega} [\delta \boldsymbol{v}_T]^2 + \frac{2\varphi^{\varepsilon}}{|\boldsymbol{\ell}|} \hat{\boldsymbol{\ell}} \cdot \left[\delta \boldsymbol{v} \times (\delta \boldsymbol{\omega} \times \delta \boldsymbol{v}) \right] d^3 \boldsymbol{\ell}$$

$$= \int_{\mathbb{T}^3} \partial_{\ell_k} \left[\varphi^{\varepsilon} (\delta^{ij} - \hat{\boldsymbol{\ell}}_i \hat{\boldsymbol{\ell}}_j) \right] \delta \omega_k \delta v_i \delta v_j d^3 \boldsymbol{\ell}$$

$$= -\operatorname{div} \left[\left(\boldsymbol{\omega} (\boldsymbol{v}_T \cdot \boldsymbol{v}_T) \right)^{\varepsilon} - \boldsymbol{\omega} (\boldsymbol{v}_T \cdot \boldsymbol{v}_T)^{\varepsilon} \right] + 2v_j \partial_k \left[(\omega_k v_{T_j})^{\varepsilon} - (\omega_k v_{T_j}^{\varepsilon}) \right], \qquad (2.51)$$

which means

$$\frac{1}{2} \int_{\mathbb{T}^{3}} \nabla \varphi^{\varepsilon}(\ell) \cdot \delta \boldsymbol{\omega} [\delta \boldsymbol{v}_{T}]^{2} - \frac{2}{|\boldsymbol{\ell}|} \varphi^{\varepsilon} \hat{\boldsymbol{\ell}} \cdot \delta \boldsymbol{\omega} [\delta \boldsymbol{v}_{T}]^{2} + \frac{2}{|\boldsymbol{\ell}|} \varphi^{\varepsilon} \hat{\boldsymbol{\ell}} \cdot \left[\delta \boldsymbol{v} \times (\delta \boldsymbol{\omega} \times \delta \boldsymbol{v}) \right] d^{3} \boldsymbol{\ell}
+ \frac{1}{2} \operatorname{div} \left[\left(\boldsymbol{\omega} (\boldsymbol{v}_{T} \cdot \boldsymbol{v}_{T}) \right)^{\varepsilon} - \boldsymbol{\omega} (\boldsymbol{v}_{T} \cdot \boldsymbol{v}_{T})^{\varepsilon} \right]
= v_{j} \partial_{k} \left[(\omega_{k} v_{T_{j}})^{\varepsilon} - (\omega_{k} v_{T_{j}}^{\varepsilon}) \right].$$
(2.52)

Thus, we plug (2.50) and (2.52) into (2.43) to deduce that

$$\partial_{t}(\boldsymbol{\omega}\cdot\boldsymbol{v}_{T}^{\varepsilon}+\boldsymbol{v}\cdot\boldsymbol{\omega}_{T}^{\varepsilon})+\operatorname{div}(\boldsymbol{\omega}P_{T}^{\varepsilon}+P\boldsymbol{\omega}_{T}^{\varepsilon})+\operatorname{div}\left[\boldsymbol{v}(\boldsymbol{\omega}\cdot\boldsymbol{v}_{T}^{\varepsilon})+\boldsymbol{v}(\boldsymbol{v}\cdot\boldsymbol{\omega}_{T}^{\varepsilon})-\boldsymbol{\omega}(\boldsymbol{v}\cdot\boldsymbol{v}_{T}^{\varepsilon})\right]$$
$$+\operatorname{div}\left[\left(\boldsymbol{v}(\boldsymbol{\omega}_{T}\cdot\boldsymbol{v}_{T})\right)^{\varepsilon}-\boldsymbol{v}(\boldsymbol{\omega}_{T}\cdot\boldsymbol{v}_{T})^{\varepsilon}\right]-\frac{1}{2}\operatorname{div}\left[\left(\boldsymbol{\omega}(\boldsymbol{v}_{T}\cdot\boldsymbol{v}_{T})\right)^{\varepsilon}-\boldsymbol{\omega}(\boldsymbol{v}_{T}\cdot\boldsymbol{v}_{T})^{\varepsilon}\right]$$
$$=-\frac{8}{3}D_{HT}^{\varepsilon}(\boldsymbol{v},\boldsymbol{\omega}),$$
$$(2.53)$$

where

$$D_{HT}^{\varepsilon}(\boldsymbol{v},\boldsymbol{\omega}) = \frac{3}{8} \int_{\mathbb{T}^3} \nabla \varphi^{\varepsilon}(\ell) \cdot \delta \boldsymbol{v} (\delta \boldsymbol{v}_T \cdot \delta \boldsymbol{\omega}_T) - \frac{2}{|\boldsymbol{\ell}|} \varphi^{\varepsilon}(\ell) \hat{\boldsymbol{\ell}} \cdot \left[\delta \boldsymbol{v} (\delta \boldsymbol{v}_T \cdot \delta \boldsymbol{\omega}_T) + \delta \boldsymbol{v} \times (\delta \boldsymbol{\omega} \times \delta \boldsymbol{v}) \right] d^3 \boldsymbol{\ell}$$

$$- \frac{3}{16} \int_{\mathbb{T}^3} \nabla \varphi^{\varepsilon}(\ell) \cdot \delta \boldsymbol{\omega} [\delta \boldsymbol{v}_T]^2 - \frac{2}{|\boldsymbol{\ell}|} \varphi^{\varepsilon}(\ell) \hat{\boldsymbol{\ell}} \cdot \delta \boldsymbol{\omega} [\delta \boldsymbol{v}_T]^2 d^3 \boldsymbol{\ell}.$$
(2.54)

Then the validity of transverse Kármárth-Howarth equations for helicity has been proved.

Step 2: The next objective is passing to the limit of ε on the left hand side of (2.40) and (2.53). Firstly, due to the definition of φ^{ε} , we have the following elementary fact

$$\int_{\mathbb{T}^3} \varphi^{\varepsilon}(\ell) (\hat{\ell} \otimes \hat{\ell}) d^3 \ell = \frac{1}{3} \mathbf{1}.$$
(2.55)

Together with the definition of v_L^{ε} and using the Minkowski inequality, we can obtain

$$\begin{aligned} \left\| \boldsymbol{v}_{L}^{\varepsilon}(\boldsymbol{x},t) - \frac{1}{3} \boldsymbol{v}(\boldsymbol{x},t) \right\|_{L^{q}(\mathbb{T}^{3})} &= \left\| \int_{\mathbb{T}^{3}} \varphi^{\varepsilon}(\ell) (\hat{\boldsymbol{\ell}} \otimes \hat{\boldsymbol{\ell}}) \cdot \left[\boldsymbol{v}(\boldsymbol{x}+\boldsymbol{\ell},t) - \boldsymbol{v}(\boldsymbol{x},t) \right] d^{3}\boldsymbol{\ell} \right\|_{L^{q}(\mathbb{T}^{3})} \\ &\leq \int_{\mathbb{T}^{3}} \varphi^{\varepsilon}(\ell) \| \boldsymbol{v}(\boldsymbol{x}+\boldsymbol{\ell},t) - \boldsymbol{v}(\boldsymbol{x},t) \|_{L^{q}(\mathbb{T}^{3})} d^{3}\boldsymbol{\ell}, \end{aligned}$$
(2.56)

which implies $\|\boldsymbol{v}_{L}^{\varepsilon}(\boldsymbol{x},t) - \frac{1}{3}\boldsymbol{v}(\boldsymbol{x},t)\|_{L^{q}(\mathbb{T}^{3})} \to 0$ as $\varepsilon \to 0$ according to the fact $\boldsymbol{v} \in L^{q}(\mathbb{T}^{3})$. Since $\boldsymbol{v}_{T}^{\varepsilon} = \boldsymbol{v}^{\varepsilon} - \boldsymbol{v}_{L}^{\varepsilon}$, we can deduce that $\|\boldsymbol{v}_{T}^{\varepsilon}(\boldsymbol{x},t) - \frac{2}{3}\boldsymbol{v}(\boldsymbol{x},t)\|_{L^{q}(\mathbb{T}^{3})} \to 0$ as $\varepsilon \to 0$. In the same manner, since $\boldsymbol{\omega} \in L^{n}(\mathbb{T}^{3})$, we can get that $\|\boldsymbol{\omega}_{L}^{\varepsilon}(\boldsymbol{x},t) - \frac{1}{3}\boldsymbol{\omega}(\boldsymbol{x},t)\|_{L^{n}(\mathbb{T}^{3})} \to 0$ and $\|\boldsymbol{\omega}_{T}^{\varepsilon}(\boldsymbol{x},t) - \frac{2}{3}\boldsymbol{\omega}(\boldsymbol{x},t)\|_{L^{n}(\mathbb{T}^{3})} \to 0$. Moreover, in light of Calderón-Zygmund theorem and $\boldsymbol{v} \in L^{q}(\mathbb{T}^{3})$, we know that $P \in L^{\frac{q}{2}}(\mathbb{T}^{3})$. Consequently, we also deduce that $\|P_{L}^{\varepsilon}(\boldsymbol{x},t) - \frac{1}{3}P(\boldsymbol{x},t)\|_{L^{\frac{q}{2}}(\mathbb{T}^{3})} \to 0$ and $\|P_{T}^{\varepsilon}(\boldsymbol{x},t) - \frac{2}{3}P(\boldsymbol{x},t)\|_{L^{\frac{q}{2}}(\mathbb{T}^{3})} \to 0$ as $\varepsilon \to 0$. Then using the standard properties of mollification, we have $\left[\left(\boldsymbol{v}(\boldsymbol{\omega}_{L}\cdot\boldsymbol{v}_{L})\right)^{\varepsilon} - \boldsymbol{v}(\boldsymbol{\omega}_{L}\cdot\boldsymbol{v}_{L})^{\varepsilon}\right] \to 0$ and $\left[\boldsymbol{\omega}(\boldsymbol{v}_{L}\cdot\boldsymbol{v}_{L})^{\varepsilon} - \left(\boldsymbol{\omega}(\boldsymbol{v}_{L}\cdot\boldsymbol{v}_{L})\right)^{\varepsilon}\right] \to 0$, the left hand side of (2.40) converges to

$$\frac{4}{3}\partial_t(\frac{1}{2}\boldsymbol{\omega}\cdot\boldsymbol{v}) + \frac{4}{3}\operatorname{div}(\frac{1}{2}\boldsymbol{\omega}P) + \frac{4}{3}\operatorname{div}\left[\frac{1}{2}\boldsymbol{v}(\boldsymbol{\omega}\cdot\boldsymbol{v}) - \frac{1}{4}\boldsymbol{\omega}(\boldsymbol{v}\cdot\boldsymbol{v})\right] =: -\frac{4}{3}D_H(\boldsymbol{v},\boldsymbol{\omega})$$
(2.57)

in the sense of distributions. Similarly, we can also prove the left hand side of (2.53) converges to $-\frac{8}{3}D_H(\boldsymbol{v},\boldsymbol{\omega})$, as $\varepsilon \to 0$, when X = T. In conclusion, we have

$$D_{HX}^{\varepsilon}(\boldsymbol{v},\boldsymbol{\omega}) \to D_H(\boldsymbol{v},\boldsymbol{\omega}),$$
 (2.58)

for both X = L, T, as $\varepsilon \to 0$, in the sense of distributions.

<u>Step 3</u>: Next, we have to show the 4/5 law for the helicity in the incompressible fluids based on the previous the local longitudinal and transverse Kármárth-Howarth equations for the helicity of Euler equations (1.7) we have just proved. To this end, we let

$$\begin{split} \overline{S}_{HL}(\boldsymbol{v},\boldsymbol{\omega},\lambda) &= \frac{1}{\lambda} \int_{\partial B} \hat{\boldsymbol{\ell}} \cdot \left[\delta \boldsymbol{v}(\lambda \boldsymbol{\ell}) (\delta \boldsymbol{v}_L(\lambda \boldsymbol{\ell}) \cdot \delta \boldsymbol{\omega}_L(\lambda \boldsymbol{\ell})) - \frac{1}{2} \delta \boldsymbol{\omega}(\lambda \boldsymbol{\ell}) [\delta \boldsymbol{v}_L(\lambda \boldsymbol{\ell})]^2 \right] \frac{d\sigma(\boldsymbol{\ell})}{4\pi}, \\ \overline{S}_{HT}(\boldsymbol{v},\boldsymbol{\omega},\lambda) &= \frac{1}{\lambda} \int_{\partial B} \hat{\boldsymbol{\ell}} \cdot \left[\delta \boldsymbol{v}(\lambda \boldsymbol{\ell}) (\delta \boldsymbol{v}_T(\lambda \boldsymbol{\ell}) \cdot \delta \boldsymbol{\omega}_T(\lambda \boldsymbol{\ell})) - \frac{1}{2} \delta \boldsymbol{\omega}(\lambda \boldsymbol{\ell}) [\delta \boldsymbol{v}_T(\lambda \boldsymbol{\ell})]^2 \right] \frac{d\sigma(\boldsymbol{\ell})}{4\pi}, \\ \overline{S}_H(\boldsymbol{v},\boldsymbol{\omega},\lambda) &= \frac{1}{\lambda} \int_{\partial B} \hat{\boldsymbol{\ell}} \cdot \left[\delta \boldsymbol{v}(\lambda \boldsymbol{\ell}) \times (\delta \boldsymbol{\omega}(\lambda \boldsymbol{\ell}) \times \delta \boldsymbol{v}(\lambda \boldsymbol{\ell})) \right] \frac{d\sigma(\boldsymbol{\ell})}{4\pi}. \end{split}$$

By taking a direct computation and using the radially symmetric of the test function $\varphi^{\varepsilon}(\ell)$, we conclude by coarea formula and the change of variable that

$$D_{H}(\boldsymbol{v},\boldsymbol{\omega}) = \lim_{\varepsilon \to 0} D_{HT}^{\varepsilon}(\boldsymbol{v},\boldsymbol{\omega})$$

$$= \lim_{\varepsilon \to 0} \frac{3}{8} \int_{\mathbb{T}^{3}} \left[\nabla \varphi^{\varepsilon}(\ell) - \frac{2}{|\ell|} \varphi^{\varepsilon} \hat{\ell} \right] \left[\delta \boldsymbol{v} (\delta \boldsymbol{v}_{T} \cdot \delta \boldsymbol{\omega}_{T}) - \frac{1}{2} \delta \boldsymbol{\omega} [\delta \boldsymbol{v}_{T}]^{2} \right] d^{3} \ell$$

$$- \lim_{\varepsilon \to 0} \frac{3}{4} \int_{\mathbb{T}^{3}} \frac{1}{|\ell|} \varphi^{\varepsilon} \hat{\ell} \cdot \delta \boldsymbol{v} \times (\delta \boldsymbol{\omega} \times \delta \boldsymbol{v}) d^{3} \ell$$

$$= \lim_{\varepsilon \to 0} \frac{3}{8} \left(\int_{0}^{\infty} r^{3} \varphi'(r) - 2r^{2} \varphi(r) dr 4\pi \bar{S}_{HT}(\boldsymbol{v},\boldsymbol{\omega},\varepsilon \ell) \right) - \lim_{\varepsilon \to 0} \frac{3}{4} \left(\int_{0}^{\infty} r^{2} \varphi(r) dr 4\pi \bar{S}_{H}(\boldsymbol{v},\boldsymbol{\omega},\varepsilon \ell) \right)$$

$$= -\frac{15}{8} \left(\bar{S}_{HT}(\boldsymbol{v},\boldsymbol{\omega}) + \frac{2}{5} \bar{S}_{H}(\boldsymbol{v},\boldsymbol{\omega}) \right)$$

$$= -\frac{15}{8} S_{HT}(\boldsymbol{v},\boldsymbol{\omega}), \qquad (2.59)$$

which leads to

$$S_{HT}(\boldsymbol{v}, \boldsymbol{\omega}) = -\frac{8}{15} D_H(\boldsymbol{v}, \boldsymbol{\omega}),$$

where

$$S_{HT}(\boldsymbol{v},\boldsymbol{\omega}) = \lim_{\lambda \to 0} \frac{1}{\lambda} \int_{\partial B} \hat{\boldsymbol{\ell}} \cdot \left[\delta \boldsymbol{v}(\lambda \boldsymbol{\ell}) (\delta \boldsymbol{v}_T(\lambda \boldsymbol{\ell}) \cdot \delta \boldsymbol{\omega}_T(\lambda \boldsymbol{\ell})) - \frac{1}{2} \delta \boldsymbol{\omega}(\lambda \boldsymbol{\ell}) [\delta \boldsymbol{v}_T(\lambda \boldsymbol{\ell})]^2 \right] \frac{d\sigma(\boldsymbol{\ell})}{4\pi} + \frac{2}{5} \lim_{\lambda \to 0} \frac{1}{\lambda} \int_{\partial B} \hat{\boldsymbol{\ell}} \cdot \left[\delta \boldsymbol{v}(\lambda \boldsymbol{\ell}) \times (\delta \boldsymbol{\omega}(\lambda \boldsymbol{\ell}) \times \delta \boldsymbol{v}(\lambda \boldsymbol{\ell})) \right] \frac{d\sigma(\boldsymbol{\ell})}{4\pi}.$$
(2.60)

Following exactly the lines of reasoning which led to (2.59), we find that

$$D_{H}(\boldsymbol{v},\boldsymbol{\omega}) = \lim_{\varepsilon \to 0} D_{HL}^{\varepsilon}(\boldsymbol{v},\boldsymbol{\omega})$$

$$= \lim_{\varepsilon \to 0} \frac{3}{4} \int \nabla \varphi^{\varepsilon}(\ell) \cdot \left[\delta \boldsymbol{v}(\delta \boldsymbol{v}_{L} \cdot \delta \boldsymbol{\omega}_{L}) - \frac{1}{2} \delta \boldsymbol{\omega}[\delta \boldsymbol{v}_{L}]^{2} \right] d^{3}\ell$$

$$+ \lim_{\varepsilon \to 0} \frac{3}{2} \int \frac{1}{|\ell|} \varphi^{\varepsilon} \hat{\ell} \cdot \left[\delta \boldsymbol{v}(\delta \boldsymbol{v}_{T} \cdot \delta \boldsymbol{\omega}_{T}) - \frac{1}{2} \delta \boldsymbol{\omega}[\delta \boldsymbol{v}_{T}]^{2} \right] d^{3}\ell$$

$$+ \lim_{\varepsilon \to 0} \frac{3}{2} \int \frac{1}{|\ell|} \varphi^{\varepsilon} \hat{\ell} \cdot \left[\delta \boldsymbol{v} \times (\delta \boldsymbol{\omega} \times \delta \boldsymbol{v}) \right] d^{3}\ell$$

$$= \lim_{\varepsilon \to 0} \left(\frac{3}{4} \int_{0}^{\infty} r^{3} \varphi'(r) dr 4\pi \bar{S}_{HL}(\boldsymbol{v}, \boldsymbol{\omega}, \varepsilon \ell) \right) + \lim_{\varepsilon \to 0} \left(\frac{3}{2} \int_{0}^{\infty} \varphi(r) r^{2} dr 4\pi \bar{S}_{HT}(\boldsymbol{v}, \boldsymbol{\omega}, \varepsilon \ell) \right)$$

$$+ \lim_{\varepsilon \to 0} \left(\frac{3}{2} \int_{0}^{\infty} r^{2} \varphi(r) dr 4\pi \bar{S}_{H}(\boldsymbol{v}, \boldsymbol{\omega}, \varepsilon \ell) \right)$$

$$= -\frac{9}{4} \bar{S}_{HL}(\boldsymbol{v}, \boldsymbol{\omega}) + \frac{3}{2} \bar{S}_{HT}(\boldsymbol{v}, \boldsymbol{\omega}) + \frac{3}{2} \bar{S}_{H}(\boldsymbol{v}, \boldsymbol{\omega}).$$

$$(2.61)$$

A combination of (2.59) and (2.61), gives, as $\varepsilon \to 0$,

$$\begin{cases} D_H(\boldsymbol{v},\boldsymbol{\omega}) = -\frac{15}{8}\bar{S}_{HT}(\boldsymbol{v},\boldsymbol{\omega}) - \frac{3}{4}\bar{S}_H(\boldsymbol{v},\boldsymbol{\omega}), \\ D_H(\boldsymbol{v},\boldsymbol{\omega}) = -\frac{9}{4}\bar{S}_{HL}(\boldsymbol{v},\boldsymbol{\omega}) + \frac{3}{2}\bar{S}_{HT}(\boldsymbol{v},\boldsymbol{\omega}) + \frac{3}{2}\bar{S}_H(\boldsymbol{v},\boldsymbol{\omega}), \end{cases}$$
(2.62)

which implies that

$$D_H(\boldsymbol{v},\boldsymbol{\omega}) = -\frac{5}{4}\bar{S}_{HL}(\boldsymbol{v},\boldsymbol{\omega}) + \frac{1}{2}\bar{S}_H(\boldsymbol{v},\boldsymbol{\omega})$$

$$= -\frac{5}{4}S_{HL}(\boldsymbol{v},\boldsymbol{\omega}),$$
 (2.63)

where

$$S_{HL}(\boldsymbol{v},\boldsymbol{\omega}) = \lim_{\lambda \to 0} \frac{1}{\lambda} \int_{\partial B} \hat{\boldsymbol{\ell}} \cdot \left[\delta \boldsymbol{v}(\lambda \boldsymbol{\ell}) (\delta \boldsymbol{v}_L(\lambda \boldsymbol{\ell}) \cdot \delta \boldsymbol{\omega}_L(\lambda \boldsymbol{\ell})) - \frac{1}{2} \delta \boldsymbol{\omega}(\lambda \boldsymbol{\ell}) [\delta \boldsymbol{v}_L(\lambda \boldsymbol{\ell})]^2 \right] \frac{d\sigma(\boldsymbol{\ell})}{4\pi} - \frac{2}{5} \lim_{\lambda \to 0} \frac{1}{\lambda} \int_{\partial B} \hat{\boldsymbol{\ell}} \cdot \left[\delta \boldsymbol{v}(\lambda \boldsymbol{\ell}) \times (\delta \boldsymbol{\omega}(\lambda \boldsymbol{\ell}) \times \delta \boldsymbol{v}(\lambda \boldsymbol{\ell})) \right] \frac{d\sigma(\boldsymbol{\ell})}{4\pi}.$$
(2.64)

Then, we have proved the 4/5 law and 8/15 law for the helicity of Euler equations.

Step 4: To complete the proof, it remains to to prove the equality (2.27) we have assumed. Indeed, for any vectors $\mathbf{A} = (A_1, A_2, A_3), \mathbf{B} = (B_1, B_2, B_3)$ and $\mathbf{C} = (C_1, C_2, C_3)$, we conclude by (2.11) that

$$\begin{split} & \left[\frac{\partial}{\partial \ell_{k}}(\hat{\ell}_{i}\hat{\ell}_{j}) - (\frac{\partial\hat{\ell}_{i}}{\partial \ell_{j}} + \frac{\partial\hat{\ell}_{j}}{\partial \ell_{i}})\hat{\ell}_{k}\right]A_{k}B_{i}C_{j} \\ &= \left[\left(\frac{\partial\hat{\ell}_{i}}{\partial \ell_{k}}\hat{\ell}_{j} + \frac{\partial\hat{\ell}_{j}}{\partial \ell_{k}}\hat{\ell}_{i}\right) - \frac{2}{|\ell|}(\delta^{ij} - \hat{\ell}_{i}\hat{\ell}_{j})\hat{\ell}_{k}\right]A_{k}B_{i}C_{j} \\ &= \frac{1}{|\ell|} \left[(\delta^{ik} - \hat{\ell}_{i}\hat{\ell}_{k})\hat{\ell}_{j} + (\delta^{jk} - \hat{\ell}_{j}\hat{\ell}_{k})\hat{\ell}_{i} - 2(\delta^{ij} - \hat{\ell}_{i}\hat{\ell}_{j})\hat{\ell}_{k}\right]A_{k}B_{i}C_{j} \\ &= \frac{1}{|\ell|} \left[(\delta^{ik}\hat{\ell}_{j} + \delta^{jk}\hat{\ell}_{i} - 2\delta^{ij}\hat{\ell}_{k}]A_{k}B_{i}C_{j}. \end{split}$$
(2.65)

For the sake of convience, we denote the left-hand side of above equality by I. Then, we will discuss term I into five cases.

Case 1: i = k = j. For this case, we see that

$$I = \frac{1}{|\boldsymbol{\ell}|} (\hat{\boldsymbol{\ell}}_j + \hat{\boldsymbol{\ell}}_i - 2\hat{\boldsymbol{\ell}}_k) A_k B_i C_j = \mathbf{0}.$$
 (2.66)

Case 2: i = k but $j \neq k$. We notice that

$$I = \frac{1}{|\boldsymbol{\ell}|} \hat{\boldsymbol{\ell}}_{j} A_{k} B_{i} C_{j}$$

$$= \frac{1}{|\boldsymbol{\ell}|} \Big[\hat{\boldsymbol{\ell}}_{1} C_{1} (A_{2} B_{2} + A_{3} B_{3}) + \hat{\boldsymbol{\ell}}_{2} C_{2} (A_{1} B_{1} + A_{3} B_{3}) + \hat{\boldsymbol{\ell}}_{3} C_{3} (A_{1} B_{1} + A_{2} B_{2}) \Big]$$

$$= \frac{1}{|\boldsymbol{\ell}|} \Big[\hat{\boldsymbol{\ell}}_{1} C_{1} (\mathbf{A} \cdot \mathbf{B}) - \hat{\boldsymbol{\ell}}_{1} A_{1} B_{1} C_{1} + \hat{\boldsymbol{\ell}}_{2} C_{2} (\mathbf{A} \cdot \mathbf{B}) - \hat{\boldsymbol{\ell}}_{2} A_{2} B_{2} C_{2} + \hat{\boldsymbol{\ell}}_{3} C_{3} (\mathbf{A} \cdot \mathbf{B}) - \hat{\boldsymbol{\ell}}_{3} A_{3} B_{3} C_{3} \Big]$$

$$= \frac{1}{|\boldsymbol{\ell}|} \Big[\hat{\boldsymbol{\ell}} \cdot \mathbf{C} (\mathbf{A} \cdot \mathbf{B}) - \hat{\boldsymbol{\ell}}_{1} A_{1} B_{1} C_{1} - \hat{\boldsymbol{\ell}}_{2} A_{2} B_{2} C_{2} - \hat{\boldsymbol{\ell}}_{3} A_{3} B_{3} C_{3} \Big].$$
(2.67)

Case 3: $i \neq k$ but i = j. We compute

$$I = \frac{1}{|\boldsymbol{\ell}|} (-2\hat{\boldsymbol{\ell}}_k) A_k B_i C_j$$

= $-\frac{2}{|\boldsymbol{\ell}|} \Big[\hat{\boldsymbol{\ell}}_1 A_1 (B_2 C_2 + B_3 C_3) + \hat{\boldsymbol{\ell}}_2 A_2 (B_1 C_1 + B_3 C_3) + \hat{\boldsymbol{\ell}}_3 A_3 (B_1 C_1 + B_2 C_2) \Big]$ (2.68)
= $-\frac{2}{|\boldsymbol{\ell}|} \Big[\hat{\boldsymbol{\ell}} \cdot \mathbf{A} (\mathbf{B} \cdot \mathbf{C}) - \hat{\boldsymbol{\ell}}_1 A_1 B_1 C_1 - \hat{\boldsymbol{\ell}}_2 A_2 B_2 C_2 - \hat{\boldsymbol{\ell}}_3 A_3 B_3 C_3 \Big].$

Case 4: $i \neq k$ but j = k. A straightforward computation leads to

$$I = \frac{1}{|\ell|} \hat{\ell}_i A_k B_i C_j$$

= $\frac{1}{|\ell|} \Big[\hat{\ell}_1 B_1 (A_2 C_2 + A_3 C_3) + \hat{\ell}_2 B_2 (A_1 C_1 + A_3 C_3) + \hat{\ell}_3 B_3 (A_1 C_1 + A_2 C_2) \Big]$ (2.69)
= $\frac{1}{|\ell|} \Big[\hat{\ell} \cdot \mathbf{B} (\mathbf{A} \cdot \mathbf{C}) - \hat{\ell}_1 A_1 B_1 C_1 - \hat{\ell}_2 A_2 B_2 C_2 - \hat{\ell}_3 A_3 B_3 C_3 \Big].$

Case 5: $i \neq k, j \neq k$ and $j \neq i$. It is clear that

$$I = \mathbf{0}.\tag{2.70}$$

Thus, combining all above cases together, we have

$$\begin{bmatrix} \frac{\partial}{\partial \ell_k} (\hat{\ell}_i \hat{\ell}_j) - (\frac{\partial \hat{\ell}_i}{\partial \ell_j} + \frac{\partial \hat{\ell}_j}{\partial \ell_i}) \hat{\ell}_k \end{bmatrix} A_k B_i C_j$$

=
$$\frac{1}{|\ell|} \hat{\ell} \cdot \Big[\mathbf{C} (\mathbf{A} \cdot \mathbf{B}) + \mathbf{B} (\mathbf{A} \cdot \mathbf{C}) - 2\mathbf{A} (\mathbf{B} \cdot \mathbf{C}) \Big].$$
 (2.71)

The proof of this theorem is now completed.

3 Four-fifths laws of total energy and cross-helicity in magnetized fluids

This section is to prove Theorem 1.2. We will consider dissipation rates of total energy and cross-helicity in magnetohydrodynamics flow, respectively.

3.1 Total energy

In this subsection, we present 4/5 law of total energy for the MHD equations.

Proof of Theorem 1.2: Energy. On the one hand, following the same path of (2.17)-(2.20), we can obtain

$$\begin{cases} \partial_t \boldsymbol{v}_L^{\varepsilon} \cdot \boldsymbol{v} + \operatorname{div}(\boldsymbol{v} \otimes \boldsymbol{v}_L)^{\varepsilon} \cdot \boldsymbol{v} - \operatorname{div}(\boldsymbol{h} \otimes \boldsymbol{h}_L)^{\varepsilon} \cdot \boldsymbol{v} + \nabla \Pi_L^{\varepsilon} \cdot \boldsymbol{v} = 0, \\ \partial_t \boldsymbol{v} \cdot \boldsymbol{v}_L^{\varepsilon} + \operatorname{div}(\boldsymbol{v} \otimes \boldsymbol{v}) \cdot \boldsymbol{v}_L^{\varepsilon} - \operatorname{div}(\boldsymbol{h} \otimes \boldsymbol{h}) \cdot \boldsymbol{v}_L^{\varepsilon} + \nabla \Pi \cdot \boldsymbol{v}_L^{\varepsilon} = 0, \\ \partial_t \boldsymbol{h}_L^{\varepsilon} \cdot \boldsymbol{h} + \operatorname{div}(\boldsymbol{v} \otimes \boldsymbol{h}_L)^{\varepsilon} \cdot \boldsymbol{h} - \operatorname{div}(\boldsymbol{h} \otimes \boldsymbol{v}_L)^{\varepsilon} \cdot \boldsymbol{h} = 0, \\ \partial_t \boldsymbol{h} \cdot \boldsymbol{h}_L^{\varepsilon} + \operatorname{div}(\boldsymbol{v} \otimes \boldsymbol{h}) \cdot \boldsymbol{h}_L^{\varepsilon} - \operatorname{div}(\boldsymbol{h} \otimes \boldsymbol{v}) \cdot \boldsymbol{h}_L^{\varepsilon} = 0, \\ \operatorname{div} \boldsymbol{v}_L^{\varepsilon} = \operatorname{div} \boldsymbol{h}_L^{\varepsilon} = 0. \end{cases}$$

Then by using the Leibniz formula, it shows that

$$\begin{aligned} \partial_i (v_i v_{L_j})^{\varepsilon} v_j + \partial_i (v_i v_j) v_{L_j}^{\varepsilon} &= \partial_i [(v_i v_j) v_{L_j}^{\varepsilon}] + v_j \partial_i \Big((v_i v_{L_j})^{\varepsilon} - (v_i v_{L_j}^{\varepsilon}) \Big), \\ &- \Big[\partial_i (h_i h_{L_j})^{\varepsilon} v_j + \partial_i (h_i h_j) v_{L_j}^{\varepsilon} \Big] = - \partial_i [(h_i h_j) v_{L_j}^{\varepsilon}] - \partial_i (h_i h_{L_j})^{\varepsilon} v_j + h_j \partial_i (h_i v_{L_j}^{\varepsilon}), \\ &\partial_i (v_i h_{L_j})^{\varepsilon} h_j + \partial_i (v_i h_j) h_{L_j}^{\varepsilon} = \partial_i [(v_i h_j) h_{L_j}^{\varepsilon}] + h_j \partial_i \Big((v_i h_{L_j})^{\varepsilon} - (v_i h_{L_j}^{\varepsilon}) \Big), \\ &- \Big[\partial_i (h_i v_{L_j})^{\varepsilon} h_j + \partial_i (h_i v_j) h_{L_j}^{\varepsilon} \Big] = - \partial_i [(h_i v_j) h_{L_j}^{\varepsilon}] - \partial_i (h_i v_{L_j})^{\varepsilon} h_j + v_j \partial_i (h_i h_{L_j}^{\varepsilon}). \end{aligned}$$

Hence, we arrive at

$$\partial_{t}(\boldsymbol{v}_{L}^{\varepsilon}\cdot\boldsymbol{v}+\boldsymbol{h}\cdot\boldsymbol{h}_{L}^{\varepsilon}) + \operatorname{div}\left[\Pi_{L}^{\varepsilon}\boldsymbol{v}+\Pi\boldsymbol{v}_{L}^{\varepsilon}\right] + \operatorname{div}\left[\boldsymbol{v}(\boldsymbol{v}\cdot\boldsymbol{v}_{L}^{\varepsilon})-\boldsymbol{h}(\boldsymbol{h}\cdot\boldsymbol{v}_{L}^{\varepsilon})+\boldsymbol{v}(\boldsymbol{h}\cdot\boldsymbol{h}_{L}^{\varepsilon})-\boldsymbol{h}(\boldsymbol{v}\cdot\boldsymbol{h}_{L}^{\varepsilon})\right]$$
$$= -v_{j}\partial_{i}\left[(v_{i}v_{L_{j}})^{\varepsilon}-(v_{i}v_{L_{j}}^{\varepsilon})\right] + v_{j}\partial_{i}\left[(h_{i}h_{L_{j}})^{\varepsilon}-(h_{i}h_{L_{j}}^{\varepsilon})\right]$$
$$-h_{j}\partial_{i}\left[(v_{i}h_{L_{j}})^{\varepsilon}-(v_{i}h_{L_{j}}^{\varepsilon})\right] + h_{j}\partial_{i}\left[(h_{i}v_{L_{j}})^{\varepsilon}-(h_{i}v_{L_{j}}^{\varepsilon})\right].$$
(3.1)

On the other hand, we derive from (2.35) that

$$\frac{1}{2} \int_{\mathbb{T}^3} \nabla \varphi^{\varepsilon}(\ell) \cdot \delta \boldsymbol{v}[\delta \boldsymbol{v}_L]^2 + \frac{2}{|\boldsymbol{\ell}|} \varphi^{\varepsilon}(\ell) \hat{\boldsymbol{\ell}} \cdot \delta \boldsymbol{v}[\delta \boldsymbol{v}_T]^2 d^3 \boldsymbol{\ell}
+ \frac{1}{2} \operatorname{div} \Big[\big(\boldsymbol{v}(\boldsymbol{v}_L \cdot \boldsymbol{v}_L) \big)^{\varepsilon} - \boldsymbol{v}(\boldsymbol{v}_L \cdot \boldsymbol{v}_L)^{\varepsilon} \Big] = v_j \partial_k \Big[(v_k v_{L_j})^{\varepsilon} - (v_k v_{L_j}^{\varepsilon}) \Big],$$
(3.2)

and

$$\frac{1}{2} \int_{\mathbb{T}^{3}} \nabla \varphi^{\varepsilon}(\ell) \cdot \delta \boldsymbol{v} [\delta \boldsymbol{h}_{L}]^{2} + \frac{2}{|\boldsymbol{\ell}|} \varphi^{\varepsilon}(\ell) \hat{\boldsymbol{\ell}} \cdot \left[\delta \boldsymbol{v} [\delta \boldsymbol{h}_{T}]^{2} + \delta \boldsymbol{h} (\delta \boldsymbol{v} \cdot \delta \boldsymbol{h}) - \delta \boldsymbol{v} (\delta \boldsymbol{h} \cdot \delta \boldsymbol{h}) \right] d^{3} \boldsymbol{\ell}
+ \frac{1}{2} \operatorname{div} \left[\left(\boldsymbol{v} (\boldsymbol{h}_{L} \cdot \boldsymbol{h}_{L}) \right)^{\varepsilon} - \boldsymbol{v} (\boldsymbol{h}_{L} \cdot \boldsymbol{h}_{L})^{\varepsilon} \right]
= h_{j} \partial_{k} \left[(v_{k} h_{L_{j}})^{\varepsilon} - (v_{k} h_{L_{j}}^{\varepsilon}) \right].$$
(3.3)

Similarly, making use of (2.39), we discover that

$$\int_{\mathbb{T}^{3}} \nabla \varphi^{\varepsilon}(\ell) \cdot \delta \boldsymbol{h}(\delta \boldsymbol{v}_{L} \cdot \delta \boldsymbol{h}_{L}) + \frac{2}{|\boldsymbol{\ell}|} \varphi^{\varepsilon}(\ell) \hat{\boldsymbol{\ell}} \cdot \delta \boldsymbol{h}(\delta \boldsymbol{v}_{T} \cdot \delta \boldsymbol{h}_{T})
- \frac{1}{|\boldsymbol{\ell}|} \varphi^{\varepsilon} \hat{\boldsymbol{\ell}} \cdot \left[\delta \boldsymbol{h}(\delta \boldsymbol{h} \cdot \delta \boldsymbol{v}) - \delta \boldsymbol{v}(\delta \boldsymbol{h} \cdot \delta \boldsymbol{h}) \right] d^{3} \boldsymbol{\ell} + \operatorname{div} \left[\left(\boldsymbol{h}(\boldsymbol{v}_{L} \cdot \boldsymbol{h}_{L}) \right)^{\varepsilon} - \boldsymbol{h}(\boldsymbol{v}_{L} \cdot \boldsymbol{h}_{L})^{\varepsilon} \right] \qquad (3.4)$$

$$= h_{j} \partial_{k} \left[(h_{k} v_{L_{j}})^{\varepsilon} - (h_{k} v_{L_{j}}^{\varepsilon}) \right] + v_{i} \partial_{k} \left[(h_{k} h_{L_{i}})^{\varepsilon} - (h_{k} h_{L_{i}}^{\varepsilon}) \right].$$

Plugging (3.2)-(3.4) into (3.1), we end up with

$$\partial_{t}(\boldsymbol{v}_{L}^{\varepsilon}\cdot\boldsymbol{v}+\boldsymbol{h}\cdot\boldsymbol{h}_{L}^{\varepsilon}) + \operatorname{div}\left[\boldsymbol{v}(\boldsymbol{v}\cdot\boldsymbol{v}_{L}^{\varepsilon}) - \boldsymbol{h}(\boldsymbol{h}\cdot\boldsymbol{v}_{L}^{\varepsilon}) + \boldsymbol{v}(\boldsymbol{h}\cdot\boldsymbol{h}_{L}^{\varepsilon}) - \boldsymbol{h}(\boldsymbol{v}\cdot\boldsymbol{h}_{L}^{\varepsilon})\right] \\ + \operatorname{div}\left[\Pi_{L}^{\varepsilon}\boldsymbol{v}+\Pi\boldsymbol{v}_{L}^{\varepsilon}\right] - \operatorname{div}\left[\left(\boldsymbol{h}(\boldsymbol{v}_{L}\cdot\boldsymbol{h}_{L})\right)^{\varepsilon} - \boldsymbol{h}(\boldsymbol{v}_{L}\cdot\boldsymbol{h}_{L})^{\varepsilon}\right] \\ + \frac{1}{2}\operatorname{div}\left[\left(\boldsymbol{v}(\boldsymbol{v}_{L}\cdot\boldsymbol{v}_{L})\right)^{\varepsilon} - \boldsymbol{v}(\boldsymbol{v}_{L}\cdot\boldsymbol{v}_{L})^{\varepsilon}\right] + \frac{1}{2}\operatorname{div}\left[\left(\boldsymbol{v}(\boldsymbol{h}_{L}\cdot\boldsymbol{h}_{L})\right)^{\varepsilon} - \boldsymbol{v}(\boldsymbol{h}_{L}\cdot\boldsymbol{h}_{L})^{\varepsilon}\right] \\ = -\frac{2}{3}D_{EL}^{\varepsilon}(\boldsymbol{v},\boldsymbol{h}), \tag{3.5}$$

where

$$\begin{split} D_{EL}^{\varepsilon}(\boldsymbol{v},\boldsymbol{h}) \\ = &\frac{3}{4} \int_{\mathbb{T}^3} \nabla \varphi^{\varepsilon}(\ell) \cdot \delta \boldsymbol{v} \Big([\delta \boldsymbol{v}_L]^2 + [\delta \boldsymbol{h}_L]^2 \Big) + \frac{2}{|\boldsymbol{\ell}|} \varphi^{\varepsilon} \hat{\boldsymbol{\ell}} \cdot \delta \boldsymbol{v} \Big([\delta \boldsymbol{v}_T]^2 + [\delta \boldsymbol{h}_T]^2 \Big) d^3 \boldsymbol{\ell} \\ &- \frac{3}{2} \int_{\mathbb{T}^3} \nabla \varphi^{\varepsilon}(\ell) \cdot \delta \boldsymbol{h} (\delta \boldsymbol{v}_L \cdot \delta \boldsymbol{h}_L) + \frac{2}{|\boldsymbol{\ell}|} \varphi^{\varepsilon} \hat{\boldsymbol{\ell}} \cdot \delta \boldsymbol{h} (\delta \boldsymbol{v}_T \cdot \delta \boldsymbol{h}_T) d^3 \boldsymbol{\ell} \\ &- 3 \int_{\mathbb{T}^3} \frac{1}{|\boldsymbol{\ell}|} \varphi^{\varepsilon} \hat{\boldsymbol{\ell}} \cdot \big(\delta \boldsymbol{h} \times (\delta \boldsymbol{v} \times \delta \boldsymbol{h}) \big) d^3 \boldsymbol{\ell}, \end{split}$$

and the indentity $\delta \boldsymbol{h} \times (\delta \boldsymbol{v} \times \delta \boldsymbol{h}) = \delta \boldsymbol{v} (\delta \boldsymbol{h} \cdot \delta \boldsymbol{h}) - \delta \boldsymbol{h} (\delta \boldsymbol{h} \cdot \delta \boldsymbol{v})$ has been used.

Proceeding similarly as the derivation of (3.1), we have

$$\partial_{t}(\boldsymbol{v}_{T}^{\varepsilon}\cdot\boldsymbol{v}+\boldsymbol{h}\cdot\boldsymbol{h}_{T}^{\varepsilon}) + \operatorname{div}\left[\Pi_{T}^{\varepsilon}\boldsymbol{v}+\Pi\boldsymbol{v}_{T}^{\varepsilon}\right] + \operatorname{div}\left[\boldsymbol{v}(\boldsymbol{v}\cdot\boldsymbol{v}_{T}^{\varepsilon}) - \boldsymbol{h}(\boldsymbol{h}\cdot\boldsymbol{v}_{T}^{\varepsilon}) + \boldsymbol{v}(\boldsymbol{h}\cdot\boldsymbol{h}_{T}^{\varepsilon}) - \boldsymbol{h}(\boldsymbol{v}\cdot\boldsymbol{h}_{T}^{\varepsilon})\right] \\ = -v_{j}\partial_{i}\left[(v_{i}v_{T_{j}})^{\varepsilon} - (v_{i}v_{T_{j}}^{\varepsilon})\right] + v_{j}\partial_{i}\left[(h_{i}h_{T_{j}})^{\varepsilon} - (h_{i}h_{T_{j}}^{\varepsilon})\right] \\ -h_{j}\partial_{i}\left[(v_{i}h_{T_{j}})^{\varepsilon} - (v_{i}h_{T_{j}}^{\varepsilon})\right] + h_{j}\partial_{i}\left[(h_{i}v_{T_{j}})^{\varepsilon} - (h_{i}v_{T_{j}}^{\varepsilon})\right].$$

$$(3.6)$$

In light of (2.50), it gives that

$$\int_{\mathbb{T}^{3}} \nabla \varphi^{\varepsilon}(\ell) \cdot \delta \boldsymbol{h}(\delta \boldsymbol{v}_{T} \cdot \delta \boldsymbol{h}_{T}) - \frac{2}{|\boldsymbol{\ell}|} \varphi^{\varepsilon} \hat{\boldsymbol{\ell}} \cdot \delta \boldsymbol{h}(\delta \boldsymbol{v}_{T} \cdot \delta \boldsymbol{h}_{T}) - \frac{1}{|\boldsymbol{\ell}|} \hat{\boldsymbol{\ell}} \cdot \left[\delta \boldsymbol{h} \times (\delta \boldsymbol{v} \times \delta \boldsymbol{h}) \right] d^{3} \boldsymbol{\ell} + \operatorname{div} \left[\left(\boldsymbol{h}(\boldsymbol{v}_{T} \cdot \boldsymbol{h}_{T}) \right)^{\varepsilon} - \boldsymbol{h}(\boldsymbol{v}_{T} \cdot \boldsymbol{h}_{T})^{\varepsilon} \right]$$

$$= h_{j} \partial_{k} \left[(h_{k} v_{T_{j}})^{\varepsilon} - (h_{k} v_{T_{j}}^{\varepsilon}) \right] + v_{i} \partial_{k} \left[(h_{k} h_{T_{i}})^{\varepsilon} - (h_{k} h_{T_{i}}^{\varepsilon}) \right].$$

$$(3.7)$$

It follows from (2.52) that

$$\frac{1}{2} \int_{\mathbb{T}^3} \nabla \varphi^{\varepsilon}(\ell) \cdot \delta \boldsymbol{v} [\delta \boldsymbol{v}_T]^2 - \frac{2}{|\boldsymbol{\ell}|} \varphi^{\varepsilon} \hat{\boldsymbol{\ell}} \cdot \delta \boldsymbol{v} [\delta \boldsymbol{v}_T]^2 d^3 \boldsymbol{\ell} + \frac{1}{2} \operatorname{div} \Big[\big(\boldsymbol{v}(\boldsymbol{v}_T \cdot \boldsymbol{v}_T) \big)^{\varepsilon} - \boldsymbol{v}(\boldsymbol{v}_T \cdot \boldsymbol{v}_T)^{\varepsilon} \Big] \\
= v_j \partial_k \Big[(v_k v_{T_j})^{\varepsilon} - (v_k v_{T_j}^{\varepsilon}) \Big],$$
(3.8)

$$\frac{1}{2} \int_{\mathbb{T}^3} \nabla \varphi^{\varepsilon}(\ell) \cdot \delta \boldsymbol{v} [\delta \boldsymbol{h}_T]^2 - \frac{2}{|\boldsymbol{\ell}|} \varphi^{\varepsilon} \hat{\boldsymbol{\ell}} \cdot \delta \boldsymbol{v} [\delta \boldsymbol{h}_T]^2 + \frac{2}{|\boldsymbol{\ell}|} \hat{\boldsymbol{\ell}} \cdot \left[\delta \boldsymbol{h} \times (\delta \boldsymbol{v} \times \delta \boldsymbol{h}) \right] d^3 \boldsymbol{\ell}
+ \frac{1}{2} \operatorname{div} \left[\left(\boldsymbol{v} (\boldsymbol{h}_T \cdot \boldsymbol{h}_T) \right)^{\varepsilon} - \boldsymbol{v} (\boldsymbol{h}_T \cdot \boldsymbol{h}_T)^{\varepsilon} \right]
= h_j \partial_k \left[(v_k h_{T_j})^{\varepsilon} - (v_k h_{T_j}^{\varepsilon}) \right].$$
(3.9)

Inserting (3.7)-(3.9) into (3.6), we arrive at

$$\partial_{t}(\boldsymbol{v}_{T}^{\varepsilon}\cdot\boldsymbol{v}+\boldsymbol{h}\cdot\boldsymbol{h}_{T}^{\varepsilon}) + \operatorname{div}\left[\boldsymbol{v}(\boldsymbol{v}\cdot\boldsymbol{v}_{T}^{\varepsilon}) - \boldsymbol{h}(\boldsymbol{h}\cdot\boldsymbol{v}_{T}^{\varepsilon}) + \boldsymbol{v}(\boldsymbol{h}\cdot\boldsymbol{h}_{T}^{\varepsilon}) - \boldsymbol{h}(\boldsymbol{v}\cdot\boldsymbol{h}_{T}^{\varepsilon})\right] + \operatorname{div}\left[\Pi_{T}^{\varepsilon}\boldsymbol{v} + \Pi\boldsymbol{v}_{T}^{\varepsilon}\right] - \operatorname{div}\left[\left(\boldsymbol{h}(\boldsymbol{v}_{T}\cdot\boldsymbol{h}_{T})\right)^{\varepsilon} - \boldsymbol{h}(\boldsymbol{v}_{T}\cdot\boldsymbol{h}_{T})^{\varepsilon}\right] + \frac{1}{2}\operatorname{div}\left[\left(\boldsymbol{v}(\boldsymbol{v}_{T}\cdot\boldsymbol{v}_{T})\right)^{\varepsilon} - \boldsymbol{v}(\boldsymbol{v}_{T}\cdot\boldsymbol{v}_{T})^{\varepsilon}\right] + \frac{1}{2}\operatorname{div}\left[\left(\boldsymbol{v}(\boldsymbol{h}_{T}\cdot\boldsymbol{h}_{T})\right)^{\varepsilon} - \boldsymbol{v}(\boldsymbol{h}_{T}\cdot\boldsymbol{h}_{T})^{\varepsilon}\right] = -\frac{4}{3}D_{ET}^{\varepsilon}(\boldsymbol{v},\boldsymbol{h}),$$

$$(3.10)$$

where

$$\begin{split} D_{ET}^{\varepsilon}(\boldsymbol{v},\boldsymbol{h}) = &\frac{3}{8} \int_{\mathbb{T}^3} \nabla \varphi^{\varepsilon}(\ell) \cdot \delta \boldsymbol{v} \Big[(\delta \boldsymbol{v}_T)^2 + (\delta \boldsymbol{h}_T)^2 \Big] - \frac{2}{|\boldsymbol{\ell}|} \varphi^{\varepsilon} \hat{\boldsymbol{\ell}} \cdot \delta \boldsymbol{v} \Big[(\delta \boldsymbol{v}_T)^2 + (\delta \boldsymbol{h}_T)^2 \Big] d^3 \boldsymbol{\ell} \\ &- \frac{3}{4} \int_{\mathbb{T}^3} \nabla \varphi^{\varepsilon}(\ell) \cdot \delta \boldsymbol{h} (\delta \boldsymbol{v}_T \cdot \delta \boldsymbol{h}_T) - \frac{2}{|\boldsymbol{\ell}|} \varphi^{\varepsilon} \hat{\boldsymbol{\ell}} \cdot \delta \boldsymbol{h} (\delta \boldsymbol{v}_T \cdot \delta \boldsymbol{h}_T) d^3 \boldsymbol{\ell} \\ &+ \frac{3}{2} \int_{\mathbb{T}^3} \frac{1}{|\boldsymbol{\ell}|} \hat{\boldsymbol{\ell}} \cdot \big[\delta \boldsymbol{h} \times (\delta \boldsymbol{v} \times \delta \boldsymbol{h}) \big] d^3 \boldsymbol{\ell}. \end{split}$$

An argument similar to the proof of Theorem 1.1 in Step 2 shows that the left-hand side of (3.5) and (3.10) converge to

$$\frac{1}{3} \Big[\partial_t ([\boldsymbol{v}]^2 + [\boldsymbol{h}]^2) + \operatorname{div}(\boldsymbol{\Pi}\boldsymbol{v}) + \operatorname{div}\left(\boldsymbol{v}[\boldsymbol{v}]^2 - 2\boldsymbol{h}(\boldsymbol{h}\cdot\boldsymbol{v}) + \boldsymbol{v}[\boldsymbol{h}]^2\right) \Big] =: -\frac{2}{3} D_E(\boldsymbol{v}, \boldsymbol{h}), \quad (3.11)$$

and

$$\frac{2}{3} \Big[\partial_t ([\boldsymbol{v}]^2 + [\boldsymbol{h}]^2) + \operatorname{div}(\boldsymbol{\Pi}\boldsymbol{v}) + \operatorname{div}\left(\boldsymbol{v}[\boldsymbol{v}]^2 - 2\boldsymbol{h}(\boldsymbol{h}\cdot\boldsymbol{v}) + \boldsymbol{v}[\boldsymbol{h}]^2\right) \Big] =: -\frac{4}{3} D_E(\boldsymbol{v},\boldsymbol{h}), \quad (3.12)$$

respectively, in the sense of distributions. Namely, we have

$$D_{EX}^{\varepsilon}(\boldsymbol{v},\boldsymbol{h}) \to D_E(\boldsymbol{v},\boldsymbol{h}),$$
 (3.13)

for both X = L, T, as $\varepsilon \to 0$, in the sense of distributions.

To proceed further, we set

$$\overline{S}_{EL}(\boldsymbol{v},\boldsymbol{h},\lambda) = \frac{1}{\lambda} \int_{\partial B} \hat{\boldsymbol{\ell}} \cdot \left[\delta \boldsymbol{v}(\lambda \boldsymbol{\ell}) \left([\delta \boldsymbol{v}_L(\lambda \boldsymbol{\ell})]^2 + [\delta \boldsymbol{h}_L(\lambda \boldsymbol{\ell})]^2 \right) - 2\delta \boldsymbol{h}(\lambda \boldsymbol{\ell}) \left(\delta \boldsymbol{v}_L(\lambda \boldsymbol{\ell}) \cdot \delta \boldsymbol{h}_L(\lambda \boldsymbol{\ell}) \right) \right] \frac{d\sigma(\boldsymbol{\ell})}{4\pi},$$

$$\overline{S}_{ET}(\boldsymbol{v},\boldsymbol{h},\lambda) = \frac{1}{\lambda} \int_{\partial B} \hat{\boldsymbol{\ell}} \cdot \left[\delta \boldsymbol{v}(\lambda \boldsymbol{\ell}) \left([\delta \boldsymbol{v}_T(\lambda \boldsymbol{\ell})]^2 + [\delta \boldsymbol{h}_T(\lambda \boldsymbol{\ell})]^2 \right) - 2\delta \boldsymbol{h}(\lambda \boldsymbol{\ell}) \left(\delta \boldsymbol{v}_T(\lambda \boldsymbol{\ell}) \cdot \delta \boldsymbol{h}_T(\lambda \boldsymbol{\ell}) \right) \right] \frac{d\sigma(\boldsymbol{\ell})}{4\pi},$$

$$\overline{S}_E(\boldsymbol{v},\boldsymbol{h},\lambda) = \frac{1}{\lambda} \int_{\partial B} \hat{\boldsymbol{\ell}} \cdot \left[\delta \boldsymbol{h}(\lambda \boldsymbol{\ell}) \times \left(\delta \boldsymbol{v}(\lambda \boldsymbol{\ell}) \times \delta \boldsymbol{h}(\lambda \boldsymbol{\ell}) \right) \right] \frac{d\sigma(\boldsymbol{\ell})}{4\pi}.$$

A simple computation leads to

$$D_{E}(\boldsymbol{v},\boldsymbol{h}) = \lim_{\varepsilon \to 0} D_{ET}^{\varepsilon}(\boldsymbol{v},\boldsymbol{h})$$

$$= \lim_{\varepsilon \to 0} \frac{3}{8} \int_{\mathbb{T}^{3}} \left[\nabla \varphi^{\varepsilon}(\ell) - \frac{2}{|\ell|} \varphi^{\varepsilon} \hat{\ell} \right] \cdot \left[\delta \boldsymbol{v} [\delta \boldsymbol{v}_{T}]^{2} + \delta \boldsymbol{v} [\delta \boldsymbol{h}_{T}]^{2} - 2\delta \boldsymbol{h} (\delta \boldsymbol{v}_{T} \cdot \delta \boldsymbol{h}_{T}) \right] d^{3} \ell$$

$$+ \frac{3}{2} \lim_{\varepsilon \to 0} \int_{\mathbb{T}^{3}} \frac{1}{|\ell|} \hat{\ell} \cdot [\delta \boldsymbol{h} \times (\delta \boldsymbol{v} \times \delta \boldsymbol{h})] d^{3} \ell$$

$$= -\frac{15}{8} \left(\bar{S}_{ET}(\boldsymbol{v},\boldsymbol{h}) - \frac{4}{5} \bar{S}_{E}(\boldsymbol{v},\boldsymbol{h}) \right)$$

$$= -\frac{15}{8} S_{ET}(\boldsymbol{v},\boldsymbol{h}), \qquad (3.14)$$

and

$$\begin{split} D_{E}(\boldsymbol{v},\boldsymbol{h}) &= \lim_{\varepsilon \to 0} D_{EL}^{\varepsilon}(\boldsymbol{v},\boldsymbol{h}) \\ &= \frac{3}{4} \lim_{\varepsilon \to 0} \int_{\mathbb{T}^{3}} \nabla \varphi^{\varepsilon}(\ell) \cdot \left[\delta \boldsymbol{v} \left([\delta \boldsymbol{v}_{L}]^{2} + [\delta \boldsymbol{h}_{L}]^{2} \right) - 2\delta \boldsymbol{h} (\delta \boldsymbol{v}_{L} \cdot \delta \boldsymbol{h}_{L}) \right] d^{3}\ell \\ &\quad + \frac{3}{4} \lim_{\varepsilon \to 0} \int_{\mathbb{T}^{3}} \frac{2}{|\ell|} \varphi^{\varepsilon} \hat{\ell} \cdot \left[\delta \boldsymbol{v} \left([\delta \boldsymbol{v}_{T}]^{2} + [\delta \boldsymbol{h}_{T}]^{2} \right) - 2\delta \boldsymbol{h} (\delta \boldsymbol{v}_{T} \cdot \boldsymbol{h}_{T}) \right] d^{3}\ell \\ &\quad - 3 \lim_{\varepsilon \to 0} \int_{\mathbb{T}^{3}} \frac{1}{|\ell|} \varphi^{\varepsilon} \hat{\ell} \cdot \left[\delta \boldsymbol{h} \times (\delta \boldsymbol{v} \times \delta \boldsymbol{h}) \right] d^{3}\ell \\ &= \frac{3}{4} \int_{0}^{\infty} r^{3} \varphi'(r) dr 4\pi \bar{S}_{EL}(\boldsymbol{v},\boldsymbol{h}) + \frac{3}{4} \lim_{\varepsilon \to 0} \int_{0}^{\infty} 2\varphi(r) r^{2} dr 4\pi \bar{S}_{ET}(\boldsymbol{v},\boldsymbol{h}) \\ &\quad - 3 \int_{0}^{\infty} r^{2} \varphi(r) dr 4\pi \bar{S}_{E}(\boldsymbol{v},\boldsymbol{h}) \\ &= -\frac{9}{4} \bar{S}_{EL}(\boldsymbol{v},\boldsymbol{h}) + \frac{3}{2} \bar{S}_{ET}(\boldsymbol{v},\boldsymbol{h}) - 3\bar{S}_{E}(\boldsymbol{v},\boldsymbol{h}). \end{split}$$

This together with (3.14) yields that

$$\begin{cases} D_E(\boldsymbol{v}, \boldsymbol{h}) = -\frac{15}{8} \bar{S}_{ET}(\boldsymbol{v}, \boldsymbol{h}) + \frac{3}{2} \bar{S}_E(\boldsymbol{v}, \boldsymbol{h}), \\ D_E(\boldsymbol{v}, \boldsymbol{h}) = -\frac{9}{4} \bar{S}_{EL}(\boldsymbol{v}, \boldsymbol{h}) + \frac{3}{2} \bar{S}_{ET}(\boldsymbol{v}, \boldsymbol{h}) - 3\bar{S}_E(\boldsymbol{v}, \boldsymbol{h}), \end{cases}$$
(3.15)

which turns out that

$$D_E(\boldsymbol{v}, \boldsymbol{h}) = -\frac{5}{4}\bar{S}_{EL}(\boldsymbol{v}, \boldsymbol{h}) - \bar{S}_E(\boldsymbol{v}, \boldsymbol{h}) = -\frac{5}{4}\Big(\bar{S}_{EL}(\boldsymbol{v}, \boldsymbol{h}) + \frac{4}{5}\bar{S}_E(\boldsymbol{v}, \boldsymbol{h})\Big) = -\frac{5}{4}S_{EL}.$$
 (3.16)

We derive from (3.14) and (3.16) that

$$S_{ET}(\boldsymbol{v},\boldsymbol{h}) = -\frac{8}{15}D_E(\boldsymbol{v},\boldsymbol{h}), \quad S_{EL}(\boldsymbol{v},\boldsymbol{h}) = -\frac{4}{5}D_E(\boldsymbol{v},\boldsymbol{h}),$$

where

$$S_{EL} = \lim_{\lambda \to 0} \frac{1}{\lambda} \int_{\partial B} \hat{\boldsymbol{\ell}} \cdot \left[\delta \boldsymbol{v}(\lambda \boldsymbol{\ell}) \left([\delta \boldsymbol{v}_L(\lambda \boldsymbol{\ell})]^2 + [\delta \boldsymbol{h}_L(\lambda \boldsymbol{\ell})]^2 \right) - 2\delta \boldsymbol{h}(\lambda \boldsymbol{\ell}) \left(\delta \boldsymbol{v}_L(\lambda \boldsymbol{\ell}) \cdot \delta \boldsymbol{h}_L(\lambda \boldsymbol{\ell}) \right) \right] \frac{d\sigma(\boldsymbol{\ell})}{4\pi} \\ + \frac{4}{5} \lim_{\lambda \to 0} \frac{1}{\lambda} \int_{\partial B} \hat{\boldsymbol{\ell}} \cdot \left[\delta \boldsymbol{h}(\lambda \boldsymbol{\ell}) \times \left(\delta \boldsymbol{v}(\lambda \boldsymbol{\ell}) \times \delta \boldsymbol{h}(\lambda \boldsymbol{\ell}) \right) \right] \frac{d\sigma(\boldsymbol{\ell})}{4\pi},$$

and

$$S_{ET} = \lim_{\lambda \to 0} \frac{1}{\lambda} \int_{\partial B} \hat{\boldsymbol{\ell}} \cdot \left[\delta \boldsymbol{v}(\lambda \boldsymbol{\ell}) \left([\delta \boldsymbol{v}_T(\lambda \boldsymbol{\ell})]^2 + [\delta \boldsymbol{h}_T(\lambda \boldsymbol{\ell})]^2 \right) - 2\delta \boldsymbol{h}(\lambda \boldsymbol{\ell}) \left(\delta \boldsymbol{v}_T(\lambda \boldsymbol{\ell}) \cdot \delta \boldsymbol{h}_T(\lambda \boldsymbol{\ell}) \right) \right] \frac{d\sigma(\boldsymbol{\ell})}{4\pi} - \frac{4}{5} \lim_{\lambda \to 0} \frac{1}{\lambda} \int_{\partial B} \hat{\boldsymbol{\ell}} \cdot \left[\delta \boldsymbol{h}(\lambda \boldsymbol{\ell}) \times \left(\delta \boldsymbol{v}(\lambda \boldsymbol{\ell}) \times \delta \boldsymbol{h}(\lambda \boldsymbol{\ell}) \right) \right] \frac{d\sigma(\boldsymbol{\ell})}{4\pi}.$$

Thus, the proof of four-fifths laws of total energy in inviscid MHD equations is completed. $\hfill \Box$

3.2 Cross-helicity

In this subsection, we consider Kolmogorov' 4/5 law of cross-helicity in terms of mixed velocity field and magnetic field in MHD turbulence.

Theorem 1.2: Cross-helicity. A slight modification of the proof in (2.17)-(2.20) gives that

$$\begin{cases} \partial_t \boldsymbol{v}_L^{\varepsilon} \cdot \boldsymbol{h} + \operatorname{div}(\boldsymbol{v} \otimes \boldsymbol{v}_L)^{\varepsilon} \cdot \boldsymbol{h} - \operatorname{div}(\boldsymbol{h} \otimes \boldsymbol{h}_L)^{\varepsilon} \cdot \boldsymbol{h} + \nabla \Pi_L^{\varepsilon} \cdot \boldsymbol{h} = 0, \\ \partial_t \boldsymbol{h} \cdot \boldsymbol{v}_L^{\varepsilon} + \operatorname{div}(\boldsymbol{v} \otimes \boldsymbol{h}) \cdot \boldsymbol{v}_L^{\varepsilon} - \operatorname{div}(\boldsymbol{h} \otimes \boldsymbol{v}) \cdot \boldsymbol{v}_L^{\varepsilon} = 0, \\ \partial_t \boldsymbol{h}_L^{\varepsilon} \cdot \boldsymbol{v} + \operatorname{div}(\boldsymbol{v} \otimes \boldsymbol{h}_L)^{\varepsilon} \cdot \boldsymbol{v} - \operatorname{div}(\boldsymbol{h} \otimes \boldsymbol{v}_L)^{\varepsilon} \cdot \boldsymbol{v} = 0, \\ \partial_t \boldsymbol{v} \cdot \boldsymbol{h}_L^{\varepsilon} + \operatorname{div}(\boldsymbol{v} \otimes \boldsymbol{v}) \cdot \boldsymbol{h}_L^{\varepsilon} - \operatorname{div}(\boldsymbol{h} \otimes \boldsymbol{h}) \cdot \boldsymbol{h}_L^{\varepsilon} + \nabla \Pi \cdot \boldsymbol{h}_L^{\varepsilon} = 0, \\ \operatorname{div} \boldsymbol{v}_L^{\varepsilon} = \operatorname{div} \boldsymbol{h}_L^{\varepsilon} = 0. \end{cases}$$

It is easy to check that

$$\begin{aligned} \partial_i (v_i v_{L_j})^{\varepsilon} h_j &+ \partial_i (v_i h_j) v_{L_j}^{\varepsilon} = \partial_i (v_i h_j v_{L_j}^{\varepsilon}) + h_j \partial_i \Big[(v_i v_{L_j})^{\varepsilon} - (v_i v_{L_j}^{\varepsilon}) \Big], \\ &- \Big[\partial_i (h_i h_{L_j})^{\varepsilon} h_j + \partial_i (h_i v_j) v_{L_j}^{\varepsilon} \Big] = -\partial_i (h_i v_j v_{L_j}^{\varepsilon}) + v_j \partial_i (h_i v_{L_j}^{\varepsilon}) - h_j \partial_i (h_i h_{L_j})^{\varepsilon}, \\ &\partial_i (v_i h_{L_j})^{\varepsilon} v_j + \partial_i (v_i v_j) h_{L_j}^{\varepsilon} = \partial_i (v_i v_j h_{L_j}^{\varepsilon}) + v_j \partial_i \Big[(v_i h_{L_j})^{\varepsilon} - (v_i h_{L_j}^{\varepsilon}) \Big], \\ &- \Big[\partial_i (h_i v_{L_j})^{\varepsilon} v_j + \partial_i (h_i h_j) h_{L_j}^{\varepsilon} \Big] = -\partial_i (h_i h_j h_{L_j}^{\varepsilon}) + h_j \partial_i (h_i h_{L_j}^{\varepsilon}) - v_j \partial_i (h_i v_{L_j})^{\varepsilon}. \end{aligned}$$

This in turn means that

$$\partial_{t}(\boldsymbol{v}_{L}^{\varepsilon}\cdot\boldsymbol{h}+\boldsymbol{v}\cdot\boldsymbol{h}_{L}^{\varepsilon}) + \operatorname{div}(\Pi_{L}^{\varepsilon}\boldsymbol{h}+\boldsymbol{h}_{L}^{\varepsilon}\Pi) + \operatorname{div}\left[\boldsymbol{v}(\boldsymbol{h}\cdot\boldsymbol{v}_{L}^{\varepsilon})+\boldsymbol{v}(\boldsymbol{v}\cdot\boldsymbol{h}_{L}^{\varepsilon})-\boldsymbol{h}(\boldsymbol{v}\cdot\boldsymbol{v}_{L}^{\varepsilon})-\boldsymbol{h}(\boldsymbol{h}\cdot\boldsymbol{h}_{L}^{\varepsilon})\right]$$
$$= -h_{j}\partial_{i}\left[(v_{i}v_{L_{j}})^{\varepsilon}-(v_{i}v_{L_{j}}^{\varepsilon})\right] - v_{j}\partial_{i}\left[(v_{i}h_{L_{j}})^{\varepsilon}-(v_{i}h_{L_{j}}^{\varepsilon})\right] + h_{j}\partial_{i}\left[(h_{i}h_{L_{j}})^{\varepsilon}-(h_{i}h_{L_{j}}^{\varepsilon})\right]$$
$$+ v_{j}\partial_{i}\left[(h_{i}v_{L_{j}})^{\varepsilon}-(h_{i}v_{L_{j}}^{\varepsilon})\right]. \tag{3.17}$$

Repeating the derivation of (2.35), we have

$$\frac{1}{2} \int_{\mathbb{T}^3} \nabla \varphi^{\varepsilon}(\ell) \cdot \delta \boldsymbol{h} [\delta \boldsymbol{h}_L]^2 + \frac{2}{|\boldsymbol{\ell}|} \varphi^{\varepsilon} \hat{\boldsymbol{\ell}} \cdot \delta \boldsymbol{h} [\delta \boldsymbol{h}_T]^2 d^3 \boldsymbol{\ell} \\
+ \frac{1}{2} \operatorname{div} \Big[\left(\boldsymbol{h} (\boldsymbol{h}_L \cdot \boldsymbol{h}_L) \right)^{\varepsilon} - \boldsymbol{h} (\boldsymbol{h}_L \cdot \boldsymbol{h}_L)^{\varepsilon} \Big] \\
= h_j \partial_k \Big[(h_k h_{L_j})^{\varepsilon} - (h_k h_{L_j}^{\varepsilon}) \Big],$$
(3.18)

$$\frac{1}{2} \int_{\mathbb{T}^{3}} \nabla \varphi^{\varepsilon}(\ell) \cdot \delta \boldsymbol{h} [\delta \boldsymbol{v}_{L}]^{2} + \frac{2}{|\boldsymbol{\ell}|} \varphi^{\varepsilon} \hat{\boldsymbol{\ell}} \cdot \left[\delta \boldsymbol{h} [\delta \boldsymbol{v}_{T}]^{2} + \delta \boldsymbol{v} (\delta \boldsymbol{h} \cdot \delta \boldsymbol{v}) - \delta \boldsymbol{h} (\delta \boldsymbol{v} \cdot \delta \boldsymbol{v}) \right] d^{3} \boldsymbol{\ell}
+ \frac{1}{2} \operatorname{div} \left[\left(\boldsymbol{h} (\boldsymbol{v}_{L} \cdot \boldsymbol{v}_{L}) \right)^{\varepsilon} - \boldsymbol{h} (\boldsymbol{v}_{L} \cdot \boldsymbol{v}_{L})^{\varepsilon} \right]
= v_{j} \partial_{k} \left[(h_{k} v_{L_{j}})^{\varepsilon} - (h_{k} v_{L_{j}}^{\varepsilon}) \right].$$
(3.19)

Moreover, along the same line of (2.39), we get

$$\int_{\mathbb{T}^{3}} \nabla \varphi^{\varepsilon}(\ell) \cdot \delta \boldsymbol{v} (\delta \boldsymbol{h}_{L} \cdot \delta \boldsymbol{v}_{L}) + \frac{2}{|\boldsymbol{\ell}|} \varphi^{\varepsilon} \hat{\boldsymbol{\ell}} \cdot \delta \boldsymbol{v} (\delta \boldsymbol{h}_{T} \cdot \delta \boldsymbol{v}_{T}) - \frac{1}{|\boldsymbol{\ell}|} \varphi^{\varepsilon} \hat{\boldsymbol{\ell}} \cdot \left[\delta \boldsymbol{v} (\delta \boldsymbol{v} \cdot \delta \boldsymbol{h}) - \delta \boldsymbol{h} (\delta \boldsymbol{v} \cdot \delta \boldsymbol{v}) \right] d^{3} \boldsymbol{\ell} + \operatorname{div} \left[\left(\boldsymbol{v}(\boldsymbol{h}_{L} \cdot \boldsymbol{v}_{L}) \right)^{\varepsilon} - \boldsymbol{v}(\boldsymbol{h}_{L} \cdot \boldsymbol{v}_{L})^{\varepsilon} \right]$$
(3.20)
$$= h_{j} \partial_{k} \left[(v_{k} v_{L_{j}})^{\varepsilon} - (v_{k} v_{L_{j}}^{\varepsilon}) \right] + v_{i} \partial_{k} \left[(v_{k} h_{L_{i}})^{\varepsilon} - (v_{k} h_{L_{i}}^{\varepsilon}) \right].$$

Inserting (3.18)-(3.20) into (3.17), we conclude that

$$\partial_{t}(\boldsymbol{v}_{L}^{\varepsilon}\cdot\boldsymbol{h}+\boldsymbol{v}\cdot\boldsymbol{h}_{L}^{\varepsilon}) + \operatorname{div}\left(\boldsymbol{v}(\boldsymbol{h}\cdot\boldsymbol{v}_{L}^{\varepsilon})+\boldsymbol{v}(\boldsymbol{v}\cdot\boldsymbol{h}_{L}^{\varepsilon})-\boldsymbol{h}(\boldsymbol{v}\cdot\boldsymbol{v}_{L}^{\varepsilon})-\boldsymbol{h}(\boldsymbol{h}\cdot\boldsymbol{h}_{L}^{\varepsilon})\right) + \operatorname{div}(\Pi_{L}^{\varepsilon}\boldsymbol{h}+\boldsymbol{h}_{L}^{\varepsilon}\Pi) + \operatorname{div}\left[\left(\boldsymbol{v}(\boldsymbol{h}_{L}\cdot\boldsymbol{v}_{L})\right)^{\varepsilon}-\boldsymbol{v}(\boldsymbol{h}_{L}\cdot\boldsymbol{v}_{L})^{\varepsilon}\right] - \frac{1}{2}\operatorname{div}\left[\left(\boldsymbol{h}(\boldsymbol{h}_{L}\cdot\boldsymbol{h}_{L})\right)^{\varepsilon}-\boldsymbol{h}(\boldsymbol{h}_{L}\cdot\boldsymbol{h}_{L})^{\varepsilon}\right] - \frac{1}{2}\operatorname{div}\left[\left(\boldsymbol{h}(\boldsymbol{v}_{L}\cdot\boldsymbol{v}_{L})\right)^{\varepsilon}-\boldsymbol{h}(\boldsymbol{v}_{L}\cdot\boldsymbol{v}_{L})^{\varepsilon}\right] = -\frac{2}{3}D_{CHL}^{\varepsilon}(\boldsymbol{v},\boldsymbol{h})$$
(3.21)

where

$$\begin{split} &D_{CHL}^{\varepsilon}(\boldsymbol{v},\boldsymbol{h}) \\ = &\frac{3}{2} \int_{\mathbb{T}^3} \nabla \varphi^{\varepsilon}(\ell) \cdot \delta \boldsymbol{v}(\delta \boldsymbol{h}_L \cdot \delta \boldsymbol{v}_L) + \frac{2}{|\boldsymbol{\ell}|} \varphi^{\varepsilon} \hat{\boldsymbol{\ell}} \cdot \left[\delta \boldsymbol{v}(\delta \boldsymbol{h}_T \cdot \delta \boldsymbol{v}_T) + \delta \boldsymbol{v} \times (\delta \boldsymbol{h} \times \delta \boldsymbol{v}) \right] d^3 \boldsymbol{\ell} \\ &- \frac{3}{4} \int_{\mathbb{T}^3} \nabla \varphi^{\varepsilon}(\ell) \cdot \delta \boldsymbol{h} \Big([\delta \boldsymbol{h}_L]^2 + [\delta \boldsymbol{v}_L]^2 \Big) + \frac{2}{|\boldsymbol{\ell}|} \varphi^{\varepsilon} \hat{\boldsymbol{\ell}} \cdot \delta \boldsymbol{h} \Big([\delta \boldsymbol{h}_T]^2 + [\delta \boldsymbol{b}_T]^2 \Big) d^3 \boldsymbol{\ell}, \end{split}$$

and the indentity $\delta \boldsymbol{v} \times (\delta \boldsymbol{h} \times \delta \boldsymbol{v}) = \delta \boldsymbol{h} (\delta \boldsymbol{v} \cdot \delta \boldsymbol{v}) - \delta \boldsymbol{v} (\delta \boldsymbol{h} \cdot \delta \boldsymbol{v})$ has been used.

On the other hand, following the path of (3.17), we infer that

$$\partial_{t} (\boldsymbol{v}_{T}^{\varepsilon} \cdot \boldsymbol{h} + \boldsymbol{v} \cdot \boldsymbol{h}_{T}^{\varepsilon}) + \operatorname{div}(\Pi_{T}^{\varepsilon}\boldsymbol{h} + \boldsymbol{h}_{T}^{\varepsilon}\Pi) + \operatorname{div}\left[\boldsymbol{v}(\boldsymbol{h} \cdot \boldsymbol{v}_{T}^{\varepsilon}) + \boldsymbol{v}(\boldsymbol{v} \cdot \boldsymbol{h}_{T}^{\varepsilon}) - \boldsymbol{h}(\boldsymbol{v} \cdot \boldsymbol{v}_{T}^{\varepsilon}) - \boldsymbol{h}(\boldsymbol{h} \cdot \boldsymbol{h}_{T}^{\varepsilon})\right]$$

$$= -h_{j}\partial_{i}\left[(v_{i}v_{T_{j}})^{\varepsilon} - (v_{i}v_{T_{j}}^{\varepsilon})\right] - v_{j}\partial_{i}\left[(v_{i}h_{T_{j}})^{\varepsilon} - (v_{i}h_{T_{j}}^{\varepsilon})\right] + h_{j}\partial_{i}\left[(h_{i}h_{T_{j}})^{\varepsilon} - (h_{i}h_{T_{j}}^{\varepsilon})\right]$$

$$+ v_{j}\partial_{i}\left[(h_{i}v_{T_{j}})^{\varepsilon} - (h_{i}v_{T_{j}}^{\varepsilon})\right].$$
(3.22)

Arguing as (2.52), we remark that

$$\frac{1}{2} \int_{\mathbb{T}^{3}} \nabla \varphi^{\varepsilon}(\ell) \cdot \delta \boldsymbol{h} [\delta \boldsymbol{h}_{T}]^{2} - \frac{2}{|\boldsymbol{\ell}|} \varphi^{\varepsilon} \hat{\boldsymbol{\ell}} \cdot \delta \boldsymbol{h} [\delta \boldsymbol{h}_{T}]^{2} d^{3} \boldsymbol{\ell}
+ \frac{1}{2} \operatorname{div} \Big[\big(\boldsymbol{h} (\boldsymbol{h}_{T} \cdot \boldsymbol{h}_{T}) \big)^{\varepsilon} - \boldsymbol{h} (\boldsymbol{h}_{T} \cdot \boldsymbol{h}_{T})^{\varepsilon} \Big]
= h_{j} \partial_{k} \Big[(h_{k} h_{T_{j}})^{\varepsilon} - (h_{k} h_{T_{j}}^{\varepsilon}) \Big],$$
(3.23)

$$\frac{1}{2} \int_{\mathbb{T}^{3}} \nabla \varphi^{\varepsilon}(\ell) \cdot \delta \boldsymbol{h}[\delta \boldsymbol{v}_{T}]^{2} - \frac{2}{|\boldsymbol{\ell}|} \varphi^{\varepsilon} \hat{\boldsymbol{\ell}} \cdot \delta \boldsymbol{h}[\delta \boldsymbol{v}_{T}]^{2} + \frac{2}{|\boldsymbol{\ell}|} \varphi^{\varepsilon} \hat{\boldsymbol{\ell}} \cdot \left[\delta \boldsymbol{v} \times (\delta \boldsymbol{h} \times \delta \boldsymbol{v}) \right] d^{3} \boldsymbol{\ell}
+ \frac{1}{2} \operatorname{div} \left[\left(\boldsymbol{h}(\boldsymbol{v}_{T} \cdot \boldsymbol{v}_{T}) \right)^{\varepsilon} - \boldsymbol{h}(\boldsymbol{v}_{T} \cdot \boldsymbol{v}_{T})^{\varepsilon} \right]
= v_{j} \partial_{k} \left[(h_{k} v_{T_{j}})^{\varepsilon} - (h_{k} v_{T_{j}}^{\varepsilon}) \right].$$
(3.24)

By suitable modification of (2.50), one arrives at

$$\int_{\mathbb{T}^{3}} \nabla \varphi^{\varepsilon}(\ell) \cdot \delta \boldsymbol{v} (\delta \boldsymbol{h}_{T} \cdot \delta \boldsymbol{v}_{T}) - \frac{2}{|\boldsymbol{\ell}|} \varphi^{\varepsilon} \hat{\boldsymbol{\ell}} \cdot \delta \boldsymbol{v} (\delta \boldsymbol{h}_{T} \cdot \delta \boldsymbol{v}_{T}) - \frac{1}{|\boldsymbol{\ell}|} \varphi^{\varepsilon} \hat{\boldsymbol{\ell}} \cdot \left[\delta \boldsymbol{v} \times (\delta \boldsymbol{h} \times \delta \boldsymbol{v}) \right] d^{3} \boldsymbol{\ell} + \operatorname{div} \left[\left(\boldsymbol{v}(\boldsymbol{h}_{T} \cdot \boldsymbol{v}_{T}) \right)^{\varepsilon} - \boldsymbol{v}(\boldsymbol{h}_{T} \cdot \boldsymbol{v}_{T})^{\varepsilon} \right] = v_{j} \partial_{k} \left[\left(v_{k} h_{T_{j}} \right)^{\varepsilon} - \left(v_{k} h_{T_{j}}^{\varepsilon} \right) \right] + h_{i} \partial_{k} \left[\left(v_{k} v_{T_{i}} \right)^{\varepsilon} - \left(v_{k} v_{T_{i}}^{\varepsilon} \right) \right].$$

$$(3.25)$$

Substituting (3.23)-(3.25) into (3.22), it follows that

$$\partial_{t}(\boldsymbol{v}_{T}^{\varepsilon}\cdot\boldsymbol{h}+\boldsymbol{v}\cdot\boldsymbol{h}_{T}^{\varepsilon})+\operatorname{div}\left[\boldsymbol{v}(\boldsymbol{h}\cdot\boldsymbol{v}_{T}^{\varepsilon})+\boldsymbol{v}(\boldsymbol{v}\cdot\boldsymbol{h}_{T}^{\varepsilon})-\boldsymbol{h}(\boldsymbol{v}\cdot\boldsymbol{v}_{T}^{\varepsilon})-\boldsymbol{h}(\boldsymbol{h}\cdot\boldsymbol{h}_{T}^{\varepsilon})\right] +\operatorname{div}(\Pi_{T}^{\varepsilon}\boldsymbol{h}+\boldsymbol{h}_{T}^{\varepsilon}\Pi)+\operatorname{div}\left[\left(\boldsymbol{v}(\boldsymbol{h}_{T}\cdot\boldsymbol{v}_{T})\right)^{\varepsilon}-\boldsymbol{v}(\boldsymbol{h}_{T}\cdot\boldsymbol{v}_{T})^{\varepsilon}\right] -\frac{1}{2}\operatorname{div}\left[\left(\boldsymbol{h}(\boldsymbol{h}_{T}\cdot\boldsymbol{h}_{T})\right)^{\varepsilon}-\boldsymbol{h}(\boldsymbol{h}_{T}\cdot\boldsymbol{h}_{T})^{\varepsilon}\right]-\frac{1}{2}\operatorname{div}\left[\left(\boldsymbol{h}(\boldsymbol{v}_{T}\cdot\boldsymbol{v}_{T})\right)^{\varepsilon}-\boldsymbol{h}(\boldsymbol{v}_{T}\cdot\boldsymbol{v}_{T})^{\varepsilon}\right] =-\frac{4}{3}D_{CHT}^{\varepsilon}(\boldsymbol{v},\boldsymbol{h}),$$

$$(3.26)$$

where

$$\begin{split} &D_{CHT}^{\varepsilon}(\boldsymbol{v},\boldsymbol{h}) \\ = &\frac{3}{4} \int_{\mathbb{T}^3} \nabla \varphi^{\varepsilon}(\ell) \cdot \delta \boldsymbol{v}(\boldsymbol{\ell}) (\delta \boldsymbol{h}_T \cdot \delta \boldsymbol{v}_T) - \frac{2}{|\boldsymbol{\ell}|} \varphi^{\varepsilon} \hat{\boldsymbol{\ell}} \cdot \left[\delta \boldsymbol{v} (\delta \boldsymbol{h}_T \cdot \delta \boldsymbol{v}_T) + \delta \boldsymbol{v} \times (\delta \boldsymbol{h} \times \delta \boldsymbol{v}) \right] d^3 \boldsymbol{\ell} \\ &- \frac{3}{8} \int_{\mathbb{T}^3} \nabla \varphi^{\varepsilon}(\ell) \cdot \delta \boldsymbol{h} \Big([\delta \boldsymbol{h}_T]^2 + [\delta \boldsymbol{v}_T]^2 \Big) - \frac{2}{|\boldsymbol{\ell}|} \varphi^{\varepsilon} \hat{\boldsymbol{\ell}} \cdot \delta \boldsymbol{h} \Big([\delta \boldsymbol{h}_T]^2 + [\delta \boldsymbol{v}_T]^2 \Big) d^3 \boldsymbol{\ell}. \end{split}$$

Next, proceeding similarly as (2.55)-(2.58) in Section 2 shows that the left-hand side of (3.21) and (3.26) converge to

$$\frac{2}{3}\left[\partial_t(\boldsymbol{v}\cdot\boldsymbol{h}) + \operatorname{div}(\Pi\boldsymbol{v}) + \operatorname{div}\left(\boldsymbol{v}(\boldsymbol{v}\cdot\boldsymbol{h}) - \frac{1}{2}\boldsymbol{h}|\boldsymbol{v}|^2 - \frac{1}{2}\boldsymbol{h}|\boldsymbol{h}|^2\right)\right] =: -\frac{2}{3}D_{CH}(\boldsymbol{v},\boldsymbol{h}), \quad (3.27)$$

and

$$\frac{4}{3}\left[\partial_t(\boldsymbol{v}\cdot\boldsymbol{h}) + \operatorname{div}(\boldsymbol{\Pi}\boldsymbol{v}) + \operatorname{div}\left(\boldsymbol{v}(\boldsymbol{v}\cdot\boldsymbol{h}) - \frac{1}{2}\boldsymbol{h}|\boldsymbol{v}|^2 - \frac{1}{2}\boldsymbol{h}|\boldsymbol{h}|^2\right)\right] =: -\frac{4}{3}D_{CH}(\boldsymbol{v},\boldsymbol{h}) \quad (3.28)$$

respectively, in the sense of distributions. In other words, we have

$$D_{CHX}^{\varepsilon}(\boldsymbol{v},\boldsymbol{h}) \to D_{CH}(\boldsymbol{v},\boldsymbol{h}),$$
 (3.29)

for both X = L, T, as $\varepsilon \to 0$, in the sense of distributions.

Finally, to show the scaling law of cross-helicity, we let

$$\begin{split} \overline{S}_{CHL}(\boldsymbol{v},\boldsymbol{h}) \\ &= \lim_{\lambda \to 0} \frac{1}{\lambda} \int_{\partial B} \hat{\boldsymbol{\ell}} \cdot \left[2\delta \boldsymbol{v}(\lambda \boldsymbol{\ell}) (\delta \boldsymbol{h}_L(\lambda \boldsymbol{\ell}) \cdot \delta \boldsymbol{v}_L(\lambda \boldsymbol{\ell})) - \delta \boldsymbol{h}(\lambda \boldsymbol{\ell}) [\delta \boldsymbol{v}_L(\lambda \boldsymbol{\ell})]^2 - \delta \boldsymbol{h}(\lambda \boldsymbol{\ell}) [\delta \boldsymbol{h}_L(\lambda \boldsymbol{\ell})]^2 \right] \frac{d\sigma(\boldsymbol{\ell})}{4\pi}, \\ \overline{S}_{CHT}(\boldsymbol{v},\boldsymbol{h}) \\ &= \lim_{\lambda \to 0} \frac{1}{\lambda} \int_{\partial B} \hat{\boldsymbol{\ell}} \cdot \left[2\delta \boldsymbol{v}(\lambda \boldsymbol{\ell}) (\delta \boldsymbol{h}_T(\lambda \boldsymbol{\ell}) \cdot \delta \boldsymbol{v}_T(\lambda \boldsymbol{\ell})) - \delta \boldsymbol{h}(\lambda \boldsymbol{\ell}) [\delta \boldsymbol{v}_T(\lambda \boldsymbol{\ell})]^2 - \delta \boldsymbol{h}(\lambda \boldsymbol{\ell}) [\delta \boldsymbol{h}_T(\lambda \boldsymbol{\ell})]^2 \right] \frac{d\sigma(\boldsymbol{\ell})}{4\pi}, \\ \overline{S}_{CH}(\boldsymbol{v},\boldsymbol{h}) \\ &= \lim_{\lambda \to 0} \frac{1}{\lambda} \int_{\partial B} \hat{\boldsymbol{\ell}} \cdot \left[\delta \boldsymbol{v}(\lambda \boldsymbol{\ell}) \times (\delta \boldsymbol{h}(\lambda \boldsymbol{\ell}) \times \delta \boldsymbol{v}(\lambda \boldsymbol{\ell})) \right] \frac{d\sigma(\boldsymbol{\ell})}{4\pi}. \end{split}$$

By a straightforward computation, it follows that

$$D_{CH}(\boldsymbol{v},\boldsymbol{h}) = \lim_{\varepsilon \to 0} D_{CHT}^{\varepsilon}(\boldsymbol{v},\boldsymbol{h})$$

$$= \frac{3}{8} \lim_{\varepsilon \to 0} \int_{\mathbb{T}^3} \left[\nabla \varphi^{\varepsilon}(\ell) - \frac{2}{|\boldsymbol{\ell}|} \varphi^{\varepsilon} \hat{\boldsymbol{\ell}} \right] \cdot \left[2\delta \boldsymbol{v} (\delta \boldsymbol{h}_T \cdot \delta \boldsymbol{v}_T) - \delta \boldsymbol{h} [\delta \boldsymbol{v}_T]^2 - \delta \boldsymbol{h} [\delta \boldsymbol{h}_T]^2 \right] d^3 \boldsymbol{\ell}$$

$$- \frac{3}{2} \lim_{\varepsilon \to 0} \int_{\mathbb{T}^3} \frac{1}{|\boldsymbol{\ell}|} \varphi^{\varepsilon} \hat{\boldsymbol{\ell}} \cdot \left[\delta \boldsymbol{v} \times (\delta \boldsymbol{h} \times \delta \boldsymbol{v}) \right] d^3 \boldsymbol{\ell}$$

$$= \frac{3}{8} \left(\int_0^{\infty} r^3 \varphi'(r) - 2r^2 \varphi(r) dr \right) 4\pi \bar{S}_{CHT}(\boldsymbol{v},\boldsymbol{h}) - \frac{3}{2} \int_0^{\infty} r^2 \varphi(r) dr 4\pi \bar{S}_{CH}(\boldsymbol{v},\boldsymbol{h})$$

$$= -\frac{15}{8} \left(\bar{S}_{CHT}(\boldsymbol{v},\boldsymbol{h}) + \frac{4}{5} \bar{S}_{CH}(\boldsymbol{v},\boldsymbol{h}) \right)$$

$$= -\frac{15}{8} S_{CHT}(\boldsymbol{v},\boldsymbol{h}),$$
(3.30)

and

$$D_{CH}(\boldsymbol{v},\boldsymbol{h}) = \lim_{\varepsilon \to 0} D_{CHL}^{\varepsilon}(\boldsymbol{v},\boldsymbol{h})$$

$$= \frac{3}{4} \lim_{\varepsilon \to 0} \int_{\mathbb{T}^{3}} \nabla \varphi^{\varepsilon}(\ell) \cdot \left[2\delta \boldsymbol{v}(\delta \boldsymbol{h}_{L} \cdot \delta \boldsymbol{v}_{L}) - \delta \boldsymbol{h}[\delta \boldsymbol{h}_{L}]^{2} - \delta \boldsymbol{h}[\delta \boldsymbol{v}_{L}]^{2} \right] d^{3}\ell$$

$$+ \frac{3}{4} \lim_{\varepsilon \to 0} \int_{\mathbb{T}^{3}} \frac{2}{|\ell|} \varphi^{\varepsilon}(\ell) \hat{\ell} \cdot \left[2\delta \boldsymbol{v}(\delta \boldsymbol{h}_{T} \cdot \delta \boldsymbol{v}_{T}) - \delta \boldsymbol{h}[\delta \boldsymbol{h}_{T}]^{2} - \delta \boldsymbol{h}[\delta \boldsymbol{v}_{T}(\ell)]^{2} \right] d^{3}\ell$$

$$+ 3 \lim_{\varepsilon \to 0} \int_{\mathbb{T}^{3}} \frac{1}{|\ell|} \varphi^{\varepsilon} \hat{\ell} \cdot \left[\delta \boldsymbol{v} \times (\delta \boldsymbol{h} \times \delta \boldsymbol{v}) \right] d^{3}\ell$$

$$= \frac{3}{4} \int_{0}^{\infty} r^{3} \varphi'(r) dr 4\pi \bar{S}_{CHL}(\boldsymbol{v},\boldsymbol{h}) + \frac{3}{4} \int_{0}^{\infty} 2\varphi(r) r^{2} dr 4\pi \bar{S}_{CHT}(\boldsymbol{v},\boldsymbol{h})$$

$$+ 3 \int_{0}^{\infty} r^{2} \varphi(r) dr 4\pi \bar{S}_{CH}(\boldsymbol{v},\boldsymbol{h}) + 3\bar{S}_{CH}(\boldsymbol{v},\boldsymbol{h}).$$
(3.31)

Combining this and (3.30), we infer that, as $\varepsilon \to 0$,

$$\begin{cases} D_{CH}(\boldsymbol{v}, \boldsymbol{h}) = -\frac{15}{8} \bar{S}_{CHT}(\boldsymbol{v}, \boldsymbol{h}) - \frac{3}{2} \bar{S}_{CH}(\boldsymbol{v}, \boldsymbol{h}), \\ D_{CH}(\boldsymbol{v}, \boldsymbol{h}) = -\frac{9}{4} \bar{S}_{CHL}(\boldsymbol{v}, \boldsymbol{h}) + \frac{3}{2} \bar{S}_{CHT}(\boldsymbol{v}, \boldsymbol{h}) + 3\bar{S}_{CH}(\boldsymbol{v}, \boldsymbol{h}), \end{cases}$$
(3.32)

which imples

$$D_{CH}(\boldsymbol{v},\boldsymbol{h}) = -\frac{5}{4}\bar{S}_{CHL}(\boldsymbol{v},\boldsymbol{h}) + \bar{S}_{CH}(\boldsymbol{v},\boldsymbol{h})$$

$$= -\frac{5}{4}\left[\bar{S}_{CHL}(\boldsymbol{v},\boldsymbol{h}) - \frac{4}{5}\bar{S}_{CH}(\boldsymbol{v},\boldsymbol{h})\right]$$

$$= -\frac{5}{4}S_{CHL}(\boldsymbol{v},\boldsymbol{h}).$$

(3.33)

Hence, according to (3.30) and (3.33), we deduce the 4/5 law and 8/15 law of cross-helicity in magnetied fluids

$$S_{CHL}(\boldsymbol{v},\boldsymbol{h}) = -\frac{4}{5}D_{CH}(\boldsymbol{v},\boldsymbol{h}), \text{ and } S_{CHT}(\boldsymbol{v},\boldsymbol{h}) = -\frac{8}{15}D_{CH}(\boldsymbol{v},\boldsymbol{h}),$$

where

$$S_{CHL}(\boldsymbol{v},\boldsymbol{h}) = \lim_{\lambda \to 0} \frac{1}{\lambda} \int_{\partial B} \hat{\boldsymbol{\ell}} \cdot \left[2\delta \boldsymbol{v}(\lambda \boldsymbol{\ell}) \left(\delta \boldsymbol{h}_L(\lambda \boldsymbol{\ell}) \cdot \delta \boldsymbol{v}_L(\lambda \boldsymbol{\ell}) \right) - \delta \boldsymbol{h}(\lambda \boldsymbol{\ell}) [\delta \boldsymbol{v}_L(\lambda \boldsymbol{\ell})]^2 - \delta \boldsymbol{h}(\lambda \boldsymbol{\ell}) [\delta \boldsymbol{h}_L(\lambda \boldsymbol{\ell})]^2 \right] \frac{d\sigma(\boldsymbol{\ell})}{4\pi} - \frac{4}{5} \lim_{\lambda \to 0} \frac{1}{\lambda} \int_{\partial B} \hat{\boldsymbol{\ell}} \cdot \left[\delta \boldsymbol{v}(\lambda \boldsymbol{\ell}) \times \left(\delta \boldsymbol{h}(\lambda \boldsymbol{\ell}) \times \delta \boldsymbol{v}(\lambda \boldsymbol{\ell}) \right) \right] \frac{d\sigma(\boldsymbol{\ell})}{4\pi},$$

$$(3.34)$$

and

$$S_{CHT}(\boldsymbol{v}, \boldsymbol{h}) = \lim_{\lambda \to 0} \frac{1}{\lambda} \int_{\partial B} \hat{\boldsymbol{\ell}} \cdot \left[2\delta \boldsymbol{v}(\lambda \boldsymbol{\ell}) \left(\delta \boldsymbol{h}_T(\lambda \boldsymbol{\ell}) \cdot \delta \boldsymbol{v}_T(\lambda \boldsymbol{\ell}) \right) - \delta \boldsymbol{h}(\lambda \boldsymbol{\ell}) [\delta \boldsymbol{v}_T(\lambda \boldsymbol{\ell})]^2 - \delta \boldsymbol{h}(\lambda \boldsymbol{\ell}) [\delta \boldsymbol{h}_T(\lambda \boldsymbol{\ell})]^2 \right] \frac{d\sigma(\boldsymbol{\ell})}{4\pi} + \frac{4}{5} \lim_{\lambda \to 0} \frac{1}{\lambda} \int_{\partial B} \hat{\boldsymbol{\ell}} \cdot \left[\delta \boldsymbol{v}(\lambda \boldsymbol{\ell}) \times (\delta \boldsymbol{h}(\lambda \boldsymbol{\ell}) \times \delta \boldsymbol{v}(\lambda \boldsymbol{\ell})) \right] \frac{d\sigma(\boldsymbol{\ell})}{4\pi}.$$
(3.35)

The proof of Theorem 1.2 is finished.

4 Conclusion

One accurate result of the well-known K41 theory is four-fifths law of energy

$$\langle [\delta \boldsymbol{v}_L(\boldsymbol{r})]^3
angle = -rac{4}{5} \epsilon \boldsymbol{r},$$

for incompressible flow in turbulence. Another famous scaling law of energy is Yaglom four-thirds type

$$\langle \delta oldsymbol{v}_L(oldsymbol{r}) [\delta oldsymbol{v}(oldsymbol{r})]^2
angle = -rac{4}{3} \epsilon oldsymbol{r}$$

in incompressible fluid. Helicity is the second quadratic conserved quantity in the Euler equations. Its 4/3 law is governed by

$$\langle \delta \boldsymbol{v}_L(\boldsymbol{r}) [\delta \boldsymbol{v}(\boldsymbol{r}) \cdot \delta \boldsymbol{\omega}(\boldsymbol{r})]
angle - rac{1}{2} \langle \delta \boldsymbol{\omega}_L(\boldsymbol{r}) [\delta \boldsymbol{v}(\boldsymbol{r})]^2
angle = -rac{4}{3} \epsilon_H \boldsymbol{r},$$

which can be found in [27]. Moreover, there also exists 2/15 law of helicity (see [10]). A natural question is that weather Kolmogorov type law is valid for the helicity. We consider this issue and invoke Eyink's velocity decomposition in [20] to establish the local 4/5 law and 8/15 law for the helicity. In contrast to classical Kolmogorov law and Eyink's work [20], we have to handle with the interaction between the velocity and vorticity. To overcome this difficulty, we first try to find the gap between $\frac{\partial}{\partial_{\ell_k}}(\hat{\ell}_i\hat{\ell}_j)$ and $(\frac{\partial\hat{\ell}_i}{\partial_{\ell_j}} + \frac{\partial\hat{\ell}_j}{\partial_{\ell_i}})\hat{\ell}_k$ involving different variable quantities (see (2.27)). More precisely, a new identity (2.65) is observed, which allows us to deal with the interaction of different physical quantities. Finally, we deduce the exact law

$$S_{HL}(\boldsymbol{v},\boldsymbol{\omega}) = -\frac{4}{5}D_H(\boldsymbol{v},\boldsymbol{\omega}),$$

which corresponds to

$$\langle \delta \boldsymbol{v}_L (\delta \boldsymbol{v}_L \cdot \delta \boldsymbol{\omega}_L) \rangle - \frac{1}{2} \langle \delta \boldsymbol{\omega}_L (\delta \boldsymbol{v}_L)^2 \rangle - \frac{2}{5} \langle \delta \boldsymbol{\omega}_L (\delta \boldsymbol{v})^2 \rangle + \frac{2}{5} \langle \delta \boldsymbol{v}_L (\delta \boldsymbol{v} \cdot \delta \boldsymbol{\omega}) \rangle = -\frac{4}{5} \epsilon_H \boldsymbol{r}.$$

The observation (2.65) helps us to revisit the 4/5 laws due to Politano and Pouquet in [49]

$$\langle [\delta \boldsymbol{v}_L(\boldsymbol{r})]^3 \rangle - 6 \langle \boldsymbol{h}_L^2 \delta \boldsymbol{v}_L(\boldsymbol{r}) \rangle = -\frac{4}{5} \epsilon_E \boldsymbol{r}, \quad \langle [\delta \boldsymbol{h}_L(\boldsymbol{r})]^3 \rangle - 6 \langle \boldsymbol{h}_L^2 \delta \boldsymbol{v}_T(\boldsymbol{r}) \rangle = -\frac{4}{5} \epsilon_C \boldsymbol{r},$$

for the total energy and cross-helicity in magnetized fluids, where the velocity field couples the magnetic field. The scaling law plays an important role in plasma turbulence. We refer the reader to [1, 40] for recent review in this direction. Here, we obtain that

$$S_{EL}(\boldsymbol{v},\boldsymbol{h}) = -\frac{4}{5}D_E(\boldsymbol{v},\boldsymbol{h}), \quad S_{CL}(\boldsymbol{v},\boldsymbol{h}) = -\frac{4}{5}D_{CH}(\boldsymbol{v},\boldsymbol{h}),$$

which correspond to

$$\langle \delta \boldsymbol{v}_L (\delta \boldsymbol{v}_L)^2 \rangle + \langle \delta \boldsymbol{v}_L (\delta \boldsymbol{h}_L)^2 \rangle - 2 \langle \delta \boldsymbol{h}_L (\delta \boldsymbol{h}_L \cdot \delta \boldsymbol{v}_L) \rangle - \frac{4}{5} \langle \delta \boldsymbol{h}_L (\delta \boldsymbol{h} \cdot \delta \boldsymbol{v}) \rangle + \frac{4}{5} \langle \delta \boldsymbol{v}_L (\delta \boldsymbol{h})^2 \rangle = -\frac{4}{5} \epsilon_E \boldsymbol{r},$$

$$2 \langle \delta \boldsymbol{v}_L (\delta \boldsymbol{h}_L \cdot \delta \boldsymbol{v}_L) \rangle - \langle \delta \boldsymbol{h}_L (\delta \boldsymbol{h}_L)^2 \rangle - \langle \delta \boldsymbol{h}_L (\delta \boldsymbol{v}_L)^2 \rangle - \frac{4}{5} \langle \delta \boldsymbol{h}_L (\delta \boldsymbol{v})^2 \rangle + \frac{4}{5} \langle \delta \boldsymbol{v}_L (\delta \boldsymbol{v} \cdot \delta \boldsymbol{h}) \rangle = -\frac{4}{5} \epsilon_C \boldsymbol{r}.$$

It should be pointed out that an analogue of the above four-fifths laws also hold for the inviscid MHD equations based on the Elsässer variances. It is an interesting question to deduce these exact laws via the corresponding Kármán-Howarth type equations directly.

In summary, three new four-fifths laws for the helicity and, total energy, cross-helicity in the incompressible Euler and inviscid MHD equations are deduced, respectively.

Acknowledgement

Ye was partially sponsored by National Natural Science Foundation of China (No. 11701145) and Natural Science Foundation of Henan (No. 232300420111). Wang was partially supported by the National Natural Science Foundation of China under grant (No. 11971446 and No. 12071113) and sponsored by Natural Science Foundation of Henan (No. 232300421077).

References

- A. Alexakis and L. Biferale. Cascades and transitions in turbulent flows. Physics Reports, 767 (2018), 1–101.
- [2] R. Antonia, M. Ould-Rouis, F. Anselmet and Y. Zhu. Analogy between predictions of Kolmogorov and Yaglom. J. Fluid Mech., 332 (1997), 395–409.
- [3] C. Bardos, P. Gwiazda, A. Świerczewska-Gwiazda, E. S. Titi and E. Wiedemann. Onsager's conjecture in bounded domains for the conservation of entropy and other companion laws. Proc. R. Soc. A, 475 (2019), 18 pp.
- [4] J. Bedrossian, M. Coti Zelati, S. Punshon-Smith and F. Weber. A Sufficient Condition for the Kolmogorov 4/5 Law for Stationary Martingale Solutions to the 3D Navier-Stokes Equations. Commun. Math. Phys. 367 (2019), 1045–1075.
- [5] D. Biskamp. Magnetohydrodynamic turbulence. Cambridge Univ. Press. 2003.

- [6] S. Chandrasekhar. The invariant theory of isotropic turbulence in magnetohydrodynamics. Proc. R. Soc. Lond. A 204 (1951), 435–449.
- [7] Q. Chen, S. Chen and G. L. Eyink. The joint cascade of energy and helicity in threedimensional turbulence. Phys. Fluids 15 (2003), 361–374.
- [8] Q. Chen, S. Chen, G. L. Eyink and D. Holm. Intermittency in the joint cascade of energy and helicity. Phys. Rev. Lett. 90 (2003), 214503.
- [9] A. Cheskidov, P. Constantin, S. Friedlander and R. Shvydkoy. Energy conservation and Onsager's conjecture for the Euler equations. Nonlinearity, 21 (2008), 1233–52.
- [10] O. Chkhetiani. On the third moments in helical turbulence. JETP Lett. 63 (1996), 808–812.
- [11] O. Chkhetiani, On triple correlations in isotropic electronic magnetohydrodynamic turbulence. JETP Lett., 69 (1999), 664–668.
- [12] O. Chkhetiani. On the local structure of helical turbulence. Doklady Physics. 53 (2008), 513-516.
- [13] P. Constantin, W. E, and E. Titi. Onsager's conjecture on the energy conservation for solutions of Euler's equation. Commun. Math. Phys. 165 (1994), 207–209.
- [14] C. De Lellis and L J. Székelyhidi. The Euler equations as a differential inclusion, Ann. of Math., 170 (2009), 1417–1436.
- [15] C. De Lellis and L J. Székelyhidi. Dissipative continuous Euler flows. Invent Math. 193 (2013), 377–407.
- [16] T. D. Drivas. Self-regularization in turbulence from the Kolmogorov 4/5-law and alignment. Phil. Trans. R. Soc. A 380 (2022), 20210033, 1–15.
- [17] B. Dubrulle. Beyond Kolmogorov cascades. J. Fluid Mech., 867 (2019), 1–63.
- [18] J. Duchon and R. Robert. Inertial Energy Dissipation for Weak Solutions of Incompressible Euler and Navier-Stokes Equations. Nonlinearity. 13 (2000), 249–255.
- [19] G. Eyink. Energy dissipation without viscosity in ideal hydrodynamics I. Fourier analysis and local energy transfer. Phys. D 78(1994), 222–240.
- [20] G. Eyink. Local 4/5-law and energy dissipation anomaly in turbulence. Nonlinearity 16(2003), 137–145.
- [21] G. Eyink. Turbulence theory, course notes, 2007-2008, available at http://www.ams.jhu.edu/~eyink/Turbulence/notes.html.
- [22] U. Frisch. Turbulence. Cambridge Univ. Press. 1995
- [23] U. S. Fjordholm and E. Wiedemann. Statistical solutions and Onsager's conjecture, Phys. D 376-377 (2018), 259–265.
- [24] S. Galtier. Introduction to Modern Magnetohydrodynamics. Cambridge Univ. Press. 2016.

- [25] S. Galtier. Physics of Wave Turbulence. Cambridge Univ. Press. 2022.
- [26] V. Giri, H. Kwon and M. Novack. The L^3 -based strong Onsager theorem. arxiv: 2305. 18509.
- [27] T. Gomez, H. Politano and A. Pouquet. Exact relationship for third-order structure functions in helical flows. Phys. Rev. E 61 (2000), 5321–5325.
- [28] M. Hofmanova, U. Pappalettera, R. Zhu and X. Zhu. Kolmogorov 4/5 law for the forced 3D Navier-Stokes equations. arXiv: 2304.14470.
- [29] P. Isett. A proof of Onsager's conjecture. Ann. of Math., 188(3):871, 2018.
- [30] T. von Kármán and L. Howarth. On the statistical theory of isotropic turbulence, Proc. R. Soc. Lond. A 164 (1938), 192–215.
- [31] A. N. Kolmogorov. The local structure of turbulence in incompressible viscous fluid for very large Reynolds number, Dokl. Akad. Nauk SSSR 30 (1941), 9-13 (reprinted in Proc. R. Soc. Lond. A 434, 9-13 (1991)).
- [32] A. N. Kolmogorov. On degeneration (decay) of isotropic turbulence in an incompressible viscous liquid, Dokl. Akad. Nauk SSSR 31 (1941), 538–540.
- [33] A. N. Kolmogorov. Dissipation of energy in the locally isotropic turbulence. Dokl. Adad. Nauk SSSR 32(1941). English transl. Proc. R. Soc. Lond. A 434 (1991), 15–17.
- [34] S. Kurien. The reflection-antisymmetric counterpart of the Kármán-Howarth dynamical equation. Phys. D 175 (2003), 167–176.
- [35] S. Kurien, M. A. Taylor and T. Matsumoto. Isotropic third-order statistics in turbulence with helicity: the 2/15-law. J. Fluid Mech. 515 (2004), 87–97.
- [36] D. Kuzzay, O. Alexandrova and L. Matteini. Local approach to the study of energy transfers in incompressible magnetohydrodynamic turbulence. Physical Review E., 99 (2019), 053202, 1–20.
- [37] D. Kuzzay, D. Faranda and B. Dubrulle. Global vs local energy dissipation: The energy cycle of the turbulent von Kármán flow. Physics of Fluids. 27 (2015), 075105, 1–21.
- [38] E. Lindborg, A note on Kolmogorov's third-order structure-function law, the local isotropy hypothesis and the pressure-velocity correlation, J. Fluid Mech. 326 (1996), 343–356.
- [39] V. S. L'vov, E. Podivilov and I. Procaccia. Exact result for the 3rd order correlations of velocity in turbulence with helicity. arXiv:chao-dyn/9705016v2.
- [40] R. Marino and L. Sorriso-Valvo. Scaling laws for the energy transfer in space plasma turbulence. Physics Reports, 1006 (2023), 1–144.
- [41] H. K. Moffatt and A. Tsinober. Helicity in Laminar and Turbulent Flow. Ann. Rev. Fluid Mech. 24 (1992), 281–312.
- [42] H. K. Moffatt. The degree of knottedness of tangled vortex lines. J. Fluid Mech., 35 (1969), 117–129.

- [43] A. S. Monin and A. M. Yaglom. Statistical Fluid Mechanics: Mechanics of Turbulence, Vol. 2. The MIT Press. 1975.
- [44] M. Novack. Scaling laws and exact results in deterministic flows. arXiv:2310.01375.
- [45] M. Novack and V. Vicol. An Intermittent Onsager Theorem. Invent Math., 233, 2023.
- [46] J. Podesta. Laws for third-order moments in homogeneous anisotropic incompressible magnetohydrodynamic turbulence, J. Fluid Mech., 609 (2008), 171–194.
- [47] J. Podesta, M. Forman and C. Smith. Anisotropic form of third-order moments and relationship to the cascade rate in axisymmetric magnetohydrodynamic turbulence. Physics of Plasmas, 14 (2007), 092305.
- [48] H. Politano and A. Pouquet. Dynamical length scales for turbulent magnetized flows. Geophys. Res. Lett., 25 (1998), 273–276.
- [49] H. Politano and A. Pouquet, von Kármán-Howarth equation for magnetohydrodynamics and its consequences on third-order longitudinal structure and correlation functions. Phys. Rev. E 57 (1998), R21.
- [50] Y. Wang, W. Wei, G. Wu and Y. Ye. On the energy and helicity conservation of the incompressible Euler equations. arXiv:2307.08322.
- [51] A. M. Yaglom. Local Structure of the Temperature Field in a Turbulent Flow. Dokl. Akad. Nauk SSSR, 69 (1949), 743–746.
- [52] K. Yoshimatsu. Examination of the four-fifths law for longitudinal third-order moments in incompressible magnetohydrodynamic turbulence in a periodic box. Phys. Rev. E 85 (2012), 066313.
- [53] T. A. Yousef, F. Rincon, and A. A. Schekochihin. Exact scaling laws and the local structure of isotropic magnetohydrodynamic turbulence. J. Fluid Mech. (2007), vol. 575, 111–120.