FIRST EIGENVALUE ESTIMATES FOR ASYMPTOTICALLY HYPERBOLIC MANIFOLDS AND THEIR SUBMANIFOLDS

SAMUEL PÉREZ-AYALA AND AARON J. TYRRELL

ABSTRACT. We derive a sharp upper bound for the first eigenvalue $\lambda_{1,p}$ of the p-Laplacian on asymptotically hyperbolic manifolds for 1 . We then prove that a particular class of conformally compact submanifolds within asymptotically hyperbolic manifolds are themselves asymptotically hyperbolic. As a corollary, we show that for any minimal conformally compact submanifold Y^{k+1} within $\mathbb{H}^{n+1}(-1)$, $\lambda_{1,p}(Y) = \left(\frac{k}{p}\right)^p$. We then obtain lower bounds on $\lambda_{1,2}(Y)$ in the case where minimality is replaced with a bounded mean curvature assumption and where the ambient space is a general Poincaré-Einstein space whose conformal infinity is of non-negative Yamabe type. In the process, we introduce an invariant $\hat{\beta}^{Y}$ for each such submanifold, enabling us to generalize a result due to Cheung-Leung in [7].

CONTENTS

1.	Introduction	1
2.	Preliminaries for Estimates on Asymptotically Hyperbolic Manifolds	7
3.	Upper Bounds for $\lambda_{1,p}(X)$ Proof of Theorem 1.2 and of Corollary 1.3	8
4.	Estimates on Submanifolds - Proofs	12
5.	Appendix: One-to-one correspondence between $\mathcal{L}(g_+)$ and $[g_+]_{\infty}^{Sc\geq 0}$	16
References		17

1. INTRODUCTION

Let \overline{X}^{n+1} be a compact manifold with boundary and interior X. A complete Riemannian metric g_+ on X is said to be conformally compact of order $C^{k,\alpha}$ if $\overline{g} := r^2 g_+$ extends as a metric to \overline{X} which is $C^{k,\alpha}(\overline{X})$, where r is a defining function for $M := \partial \overline{X}$. We will assume this extension is $C^{3,\alpha}(\overline{X})$, however some of our results hold under weaker assumptions. By a defining function for M we mean

$$r > 0 \text{ on } X,$$

$$r = 0 \text{ on } M,$$

$$lr \neq 0 \text{ on } M.$$

If ρ is another defining function for M, then the induced boundary metrics $(r^2g_+)|_{TM}$ and $(\rho^2g_+)|_{TM}$ are conformal to each other, giving rise to an invariant $[g_+]_{\infty} := [\overline{g}|_{TM}]$ of g_+ known as the conformal infinity of (X^{n+1}, g_+) . Here $[\overline{g}|_{TM}]$ denotes the conformal class of $\overline{g}|_{TM}$. As observed by Mazzeo in [22], conformally transforming the Riemann tensor of $g_+ =: g$ yields

(1.1)
$$R_{ijkl}^{g_+} = -|dr|_{\overline{g}}^2(g_{ik}g_{jl} - g_{il}g_{jk}) + O_{ijkl}(r^{-3}),$$

where $O_{ijkl}(r^{-3})$ is the component function of a covariant 4-tensor which is $O(r^{-3})$ as $r \to 0$. Notice that since $(g_+)^{ij} = r^2 \overline{g}^{ij}$, this implies that the tensor corresponding to those components vanishes with respect to the g_+ - norm as $r \to 0^+$. Moreover, (1.1) also gives us that the sectional curvatures of (X^{n+1}, g_+) approach $-|dr|^{\frac{2}{q}}$ at the boundary M. Hence, $|dr|^{\frac{2}{r^2g_+}}$ restricted to M is another invariant of the metric g_+ (see Section 2 in [12]).

Asymptotically hyperbolic (AH) manifolds are conformally compact manifolds that have asymptotic sectional curvatures equal to a constant $-\kappa^2$, where $\kappa > 0$; unless otherwise stated, we will assume this constant

Key words and phrases. Asymptotically hyperbolic, Dirichlet eigenvalue, minimal submanifold.

to be -1. We denote them by $AH(-\kappa^2)$, or just AH if $-\kappa^2 = -1$. Note that, based on (1.1), this constant $-\kappa^2$ equals $-|dr|^2_{\overline{\sigma}|M}$.

A special defining function for M is a defining function r such that $|dr|_g^2 \equiv 1$ in a neighborhood of M rather than just on M itself. It is a lemma in Graham-Lee's work [13] (see also Lemma 2.1 in [12]) that for every metric g_{∞} in the conformal infinity of an asymptotically hyperbolic manifold (X^{n+1}, g_+) , there exists a special defining function r such that $(r^2g_+)|_{TM} = g_{\infty}$. A conformally compact manifold that also satisfies the Einstein condition $\operatorname{Ric}(g_+) = -ng_+$ is called a Poincaré-Einstein manifold or conformally compact Einstein manifold. For such a manifold, it can be verified by contracting (1.1) that $(|dr|_g^2)|_M = 1$, thus implying that Poincaré-Einstein manifolds. In summary,

 ${Poincaré-Einstein} \subseteq {Asymptotically Hyperbolic} \subseteq {Conformally Compact}.$

The model case of a conformally compact Riemannian manifold is the hyperbolic space $\mathbb{H}^{n+1}(-1)$. We use the Poincaré ball model for hyperbolic space:

(1.2)
$$(\mathbb{H}^{n+1}, g_H) = \left(\mathbb{B}^{n+1}, \left(\frac{2}{1-|y|^2}\right)^2 \cdot \sum_{i=1}^{n+1} (dy_i)^2\right),$$

where $\mathbb{B}^{n+1} \subseteq \mathbb{R}^{n+1}$ denotes the standard open unit ball in Euclidean space. The functions $r_1, r_2 : \mathbb{B}^{n+1} \to \mathbb{R}^{n+1}$, defined as $r_1(y) = 1 - |y|$ and $r_2(y) = \frac{1-|y|}{1+|y|}$, where $|\cdot|$ denotes the standard euclidean norm, are examples of defining functions for $\partial \mathbb{B}^{n+1} = \mathbb{S}^n$ with both $r_1^2 g_H$ and $r_2^2 g_H$ extending to metrics on $\overline{\mathbb{B}}^{n+1}$. As discussed in [12], r_2 is an example of a special defining function. Moreover, $[g_H]_{\infty} = [(r_2^2 g_H)|_{T\mathbb{S}^n}] = [(\text{standard metric on } \mathbb{S}^n)]$ is the conformal infinity of (\mathbb{H}^{n+1}, g_H) . Finally, g_H has constant sectional curvatures equal to -1, and so $\operatorname{Ric}(g_H) = -ng_H$, which means that $(\mathbb{H}^{n+1}(-1), g_H)$ is also an example of a Poincaré-Einstein manifold.

We now proceed to introduce our main results. We begin with a discussion of upper bounds for Dirichlet eigenvalues on asymptotically hyperbolic manifolds, followed by an examination of general eigenvalue estimates on their submanifolds. All submanifolds $\iota: \overline{Y} \to \overline{X}$ are assumed to be immersed and we will often identify \overline{Y} with $\iota(\overline{Y})$ when there is no ambiguity.

1.1. Eigenvalue Estimates in Complete, non-Compact Manifolds. Let (X^{n+1}, g_+) be a complete and non-compact manifold. For any bounded domain Ω with smooth boundary, we consider the Dirichlet eigenvalue problem

(1.3)
$$\begin{cases} \Delta_p f + \lambda |f|^{p-2} f = 0 & \text{in } \Omega, \\ f = 0 & \text{on } \partial\Omega, \end{cases}$$

where $\Delta_p f = \operatorname{div}(|\nabla_{g_+} f|^{p-2}\nabla_{g_+} f)$ is the so-called *p*-Laplace operator and 1 . Notice that when <math>p = 2, we recover the standard Laplace-Beltrami operator. Any $\lambda \in \mathbb{R}$ for which a nontrivial solution f of (1.3) exists is called a Dirichlet eigenvalue of Δ_p or simply a *p*-Dirichlet eigenvalue. Solutions to (1.3) are not smooth, in general, and thus by "solutions" we always mean a solution in the sense of distributions; see [18] for a precise formulation. Our focus is on the first *p*-Dirichlet eigenvalue, whose variational characterization is given by

(1.4)
$$\lambda_{1,p}(\Omega) = \inf_{f \neq 0} \frac{\int_{\Omega} |\nabla_{g_+} f|^p \, dv_{g_+}}{\int_{\Omega} |f|^p \, dv_{g_+}}$$

and where the infimum is taken over the space $W^{1,p}_o(\Omega)$ - the completion of $C^{\infty}_o(\Omega)$ under the Sobolev norm $\|\cdot\|_{L^{1,p}(g_+)}$. The reader can consult [15, 19], and references therein, for the definition of the first *p*-Laplace Dirichlet eigenvalue (1.4) and basic properties of the associated eigenfunction.

Due to domain monotonicity (Lemma 1.1 in [10]), it makes sense to define the first *p*-Dirichlet eigenvalue of (X^{n+1}, g_+) as follows:

Definition 1.1 (First *p*-Dirichlet Eigenvalue).

(1.5)
$$\lambda_{1,p}(X) := \inf_{\Omega} \lambda_{1,p}(\Omega),$$

where the infimum is being taken over all bounded domains of X with smooth boundary.

We emphasize that the definition of $\lambda_{1,p}(X)$ also depends on g_+ , and so a better notation would be $\lambda_{1,p}(X, g_+)$. However, since g_+ will be fixed throughout the paper, we omit any reference to it and simply write $\lambda_{1,p}(X)$. In the case where $X = \mathbb{H}^{n+1}$, we will denote $g_+ = g_H$. When dealing with immersed submanifolds $\iota: Y^{k+1} \to X^{n+1}$, $\lambda_{1,p}(Y)$ is meant to be with respect to the induced metric $h_+ = \iota^* g_+$.

The first *p*-Dirichlet eigenvalue is an invariant of (X^{n+1}, g_+) which is often difficult to compute. A lot of attention has been given to the case when p = 2, that is, to the first Dirichlet eigenvalue of the Laplace-Beltrami operator, when (X^{n+1}, g_+) is assumed to be asymptotically hyperbolic. In [21], Mazzeo showed that the essential spectrum of $\Delta_2 = \Delta_{g_+}$ consists of the ray $[\left(\frac{n}{2}\right)^2, \infty)$. As a consequence, he obtained the following upper bound: $\lambda_{1,2}(X) \leq \left(\frac{n}{2}\right)^2$. For a general $p \in (1, \infty)$, upper bounds are known but under global assumptions on the Ricci curvature. In fact, on complete, (n + 1)-dimensional manifolds (X^{n+1}, g_+) with Ricci curvature bounded from below, $\operatorname{Ric}(g_+) \geq -ng_+$, classical techniques due to Cheng [6] yield $\left(\frac{n}{p}\right)^p$ as an upper bound for $\lambda_{1,p}(X)$; see [14, 27] for discussion on lower bounds under the same conditons on the Ricci curvature. This upper bound is sharp since $\lambda_{1,p}(\mathbb{H}^{n+1}(-1)) = \left(\frac{n}{p}\right)^p$; this equality will become clear in the discussion that follows.

On an AH manifold (X^{n+1}, g_+) , we argue that upper bounds on $\lambda_{1,p}(X)$ should only be influenced by the asymptotic behavior of the manifold at infinity, and that no global information should be needed to derive sharp estimates. Indeed, if Ω is a bounded domain with a smooth boundary in a collar neighborhood of $M = \partial \overline{X}$, then $\lambda_{1,p}(X) \leq \lambda_{1,p}(\Omega)$. On the other hand, the sectional curvatures are uniformly approaching -1 near the boundary, suggesting that the geometry of (X^{n+1}, g_+) should reflect some properties of \mathbb{H}^{n+1} near M. In particular, we expect $\lambda_{1,p}(\Omega)$ to be close to $\left(\frac{n}{p}\right)^p = \lambda_{1,p}(\mathbb{H}^{n+1}(-1))$. In our first theorem, we prove that this intuition is indeed correct by generalizing Mazzeo's upper estimate for any $p \in (1, \infty)$.

Theorem 1.2. Let (X^{n+1}, g_+) be an asymptotically hyperbolic manifold. Then $\lambda_{1,p}(X) \leq \left(\frac{n}{p}\right)^p$.

A few remarks are in order. To our knowledge, this result is novel for $p \neq 2$. The methods and techniques are also novel, even for the case when p = 2, involving only the construction of a family of test functions "near infinity" to capture the behavior of the manifold near its boundary. Additionally, we would like to point out that a slight generalization of Theorem 1.2 is available for manifolds which are $AH(-\kappa^2)$. Indeed, if a manifold (X^{n+1}, g_+) is $AH(-\kappa^2)$, then $(X^{n+1}, \kappa^2 g_+)$ would be AH(-1) thanks to the scaling properties of the Riemann tensor and (1.1). Since $\lambda_{1,p}(X, \kappa^2 g_+) = \kappa^{-p} \lambda_{1,p}(X^{n+1}, g_+)$ by the variational characterization (1.4), we derive the following consequence to Theorem 1.2:

Corollary 1.3. Let (X^{n+1}, g_+) be $AH(-\kappa^2)$. Then $\lambda_{1,p}(X) \leq \left(\frac{n\kappa}{p}\right)^p$.

Remark 1.4. Recall that $\kappa = (|dr|_{r^2g_+})|_M$. So, the upper bound can be re-written as

(1.6)
$$\lambda_{1,p}(X) \le \left(\frac{n(|dr|_{r^2g_+}^2)|_M}{p}\right)^p.$$

Different bounds have been derived for $\lambda_{1,p}(X)$ under various geometric assumptions and through a variety of methods. In [9], Carvalho-Cavalcante generalized a classical result due to McKean [23] and showed that for simply connected manifolds with negative sectional curvature bounded above by $-\kappa^2$, the first *p*-Dirichlet eigenvalue is at least $\left(\frac{n\kappa}{p}\right)^p$ - this bound is sharp as it is precisely what you obtain in $\mathbb{H}^{n+1}(-\kappa^2)$. Therefore, together with Corollary 1.3, we derive

Corollary 1.5. Let (X^{n+1}, g_+) be simply connected and $AH(-\kappa^2)$ with sectional curvature bounded above by $-\kappa^2$. Then $\lambda_{1,p}(X) = \left(\frac{n\kappa}{p}\right)^p$.

There is limited literature on upper bounds for $\lambda_{1,p}(X)$ when X is asymptotically hyperbolic or under other geometric assumptions. In [18], Lima-Montenegro-Santos obtained an upper bound for what they call the essential *p*-first eigenvalue in terms of the exponential volume growth O(X) of the manifold. In particular, they proved that for manifolds with infinite volume, the essential *p*-first eigenvalue is bounded above by $\left(\frac{O(X)}{p}\right)^p$ (see Theorem 1.4 in [18]). Their results, as well as their techniques, are a generalization of the corresponding result by Brooks [1] when p = 2. Bounds of the same type as in Theorem 1.2 have been found in [17], but there they do not work with Dirichlet boundary conditions.

1.2. Eigenvalue Estimates on Submanifolds of AH Spaces. We proceed with a discussion of eigenvalue estimates on submanifolds of asymptotically hyperbolic spaces. Let us start with some consequences of Theorem 1.2 for a generic $p \in (1, \infty)$.

In [10], Du and Mao, using techniques developed by Cheung-Leung in [7], showed that if Y is a (k + 1)-dimensional, complete and noncompact immersed submanifold in $\mathbb{H}^{n+1}(-1)$ with norm of the mean curvature satisfying $||H|| \leq \alpha < k$, then

(1.7)
$$0 < \left(\frac{k-\alpha}{p}\right)^p \le \lambda_{1,p}(Y)$$

For complete, (k+1)-dimensional minimal submanifolds in $\mathbb{H}^{n+1}(-1)$, the lower bound becomes $\left(\frac{k}{p}\right)^p$, which is sharp as it is the value on totally geodesic planes. We demonstrate that the upper bound is also sharp on a broader class of minimal submanifolds of AH spaces that we call conformally compact submanifolds. More generally, we define:

Definition 1.6 (Conformally Compact Submanifold & Asymptotically CMC Submanifold). Let (X^{n+1}, g_+) be a conformally compact manifold. Let \overline{Y}^{k+1} be a compact manifold with boundary and interior Y. Let $\iota: \overline{Y} \to \overline{X}^{n+1}$ be a $C^{2,\alpha}$ immersion. If $(Y, \iota^* g_+)$ is conformally compact with the property that $\iota|_{\partial \overline{Y}}: \partial \overline{Y} \to M$ is an immersion such that $\iota(\overline{Y})$ meets M transversely, then we say that $(Y^{k+1}, \iota^* g_+)$ is a conformally compact submanifold of (X^{n+1}, g_+) . Furthermore, if its mean curvature vector H satisfies $g_+(H, H) = C^2 + O(r)$ for some $C \in [0, \infty)$ and where r is any defining function M, then we say that $(Y, \iota^* g_+)$ is asymptotically CMC with asymptotic mean curvature equal to C^2 . In the case where C = 0, we say that Y is asymptotically minimal.

Remark 1.7. Given any pair of defining functions r and ρ for M, $r = O(\rho)$. It follows that the asymptotic mean curvature as described in the previous definition is well-defined.

Conformally compact submanifolds are complete and non-compact. Some examples of conformally compact hypersurfaces in hyperbolic space are the totally geodesic planes and the spherical catenoids; a nonexample is given by the horospheres. Horospheres are a non-example since although they are complete and non-compact, they do not have a boundary under compactification. Another class of conformally compact submanifolds are the type of submanifolds introduced by Graham and Witten in [25]. These submanifolds are also called polyhomogeneously immersed submanifolds; for more information on these, the reader may consult [20] and [3].

We are now in a position to state our first result related to this class of submanifolds. We prove that any asymptotically CMC submanifold with asymptotic mean curvature equal to C^2 will be $AH\left(-\left(1-\frac{C^2}{(k+1)^2}\right)\right)$ itself, assuming that it lives within an AH space. Thus allowing us to apply Corollary 1.3.

Proposition 1.8. Let Y^{k+1} be an asymptotically CMC submanifold of an asymptotically hyperbolic space (X^{n+1}, g_+) with asymptotic mean curvature C^2 . Then Y is asymptotically hyperbolic with asymptotic sectional curvatures equal to $-\left(1-\frac{C^2}{(k+1)^2}\right)$. In particular, if Y is asymptotically minimal, then Y is asymptotically hyperbolic.

As a consequence of Proposition 1.8 and Corollary 1.3, we obtain:

Corollary 1.9. Let the setup be the same as that in Proposition 1.8. Then

(1.8)
$$\lambda_{1,p}(Y^{k+1}) \le \left(\frac{k}{p}\right)^p \left(1 - \frac{C^2}{(k+1)^2}\right)^{\frac{p}{2}}.$$

In particular, if Y is asymptotically minimal, then the upper bound becomes $\left(\frac{k}{p}\right)^p$.

Remark 1.10. It will be shown in Section 4.1 that in our context C < k + 1 always holds.

Together with (1.7) for the minimal case ($\alpha = 0$), we conclude that the lower bound is attained for minimal conformally compact submanifolds of $\mathbb{H}^{n+1}(-1)$ for any $p \in (1, \infty)$. In other words, the bound $\left(\frac{k}{n}\right)^p$ is optimal for these submanifolds. We summarize our discussion in the next corollary:

Corollary 1.11. Let Y^{k+1} be a minimal conformally compact submanifold of $\mathbb{H}^{n+1}(-1)$. Then $\lambda_{1,p}(Y) = \left(\frac{k}{p}\right)^p$.

We turn our attention to acquiring bounds on the first Dirichlet eigenvalue (p = 2) of submanifolds of asymptotically hyperbolic spaces. First, let us recall some previous work on eigenvalue estimates for submanifolds of hyperbolic space. As previously mentioned, for complete, simply connected, (n+1)-dimensional manifolds X with sectional curvature bounded above by $-\kappa^2$, $\kappa \neq 0$, we have

(1.9)
$$\left(\frac{n\kappa}{2}\right)^2 \le \lambda_{1,2}(X);$$

see [9,23]. In [7], Cheung-Leung extended inequality (1.9) to complete non-compact submanifolds Y^{k+1} of hyperbolic space $\mathbb{H}^{n+1}(-1)$ with sectional curvature satisfying $||H|| \leq \alpha < k$ by showing that

(1.10)
$$\left(\frac{k-\alpha}{2}\right)^2 \le \lambda_{1,2}(Y).$$

Their result follows from the following Poincaré-type inequality (Proposition 1 in [7]): for all $f \in C_0^{\infty}(Y)$, we have

(1.11)
$$\left(\frac{k-\alpha}{2}\right)^2 \int_Y f^2 \le \int_Y |\nabla f|^2.$$

A key element in their work is the following. Let $p \in \mathbb{H}^{n+1}(-1) \setminus Y$, and set $x(y) := d_{g_H}(y, p)$, where g_H is the hyperbolic metric introduced in (1.2). Then $u : \mathbb{H}^{n+1}(-1) \to \mathbb{R}$, $u(y) := \cosh(x(y))$, satisfies $\Delta_{g_H} u = (n+1)u$ on $\mathbb{H}^{n+1}(-1)$. In fact u is an example of what we call a Lee-eigenfunction (see Definition 4.1). By noticing this fact, we are able to give a simpler argument in the spirit of Lee's [16] that also generalizes their result to any such submanifold living in a general Poincaré-Einstein manifold (X^{n+1}, g_+) with conformal infinity having non-negative Yamabe invariant.

Theorem 1.12. Let $\iota: Y^{k+1} \to X^{n+1}$ be a complete and noncompact immersion into a Poincaré-Einstein space (X^{n+1}, g_+) whose conformal infinity $(M, [g_+]_{\infty})$ has non-negative Yamabe invariant, and let u be a Lee-eigenfunction. Denote by $b(u) := \nabla_{g_+}^2 u - \frac{1}{n+1}(\Delta_{g_+}u)g_+ = \nabla_{g_+}^2 u - ug_+$ the trace-free hessian of u. If the mean curvature vector of Y has norm satisfying $||H|| \leq \alpha$ for some constant α , and if $\beta^Y(u) := \sup_Y \left(u^{-1} \cdot tr\left(b(u)|_{(TY^{\perp})^2}\right)\right)$ satisfies $\alpha + \beta^Y(u) < k$, then it follows that

(1.12)
$$0 < \left(\frac{k - \alpha - \beta^Y(u)}{2}\right)^2 \le \lambda_{1,2}(Y).$$

 $Note: \ 0 \leq \beta^{Y}(u) \ since \ u = O(r^{-1}) \ and \ tr\big(b(u)|_{(TY^{\perp})^{2}}\big) = O(1), \ as \ r \to 0.$

Let $\mathcal{L}(g_+)$ be the collection of all Lee-eigenfunctions (see Definition 4.1). As explained in Proposition 5.1 (see Appendix), there is a one-to-one correspondence between $\mathcal{L}(g_+)$ and those metrics in the conformal infinity of (X^{n+1}, g_+) whose scalar curvature is nowhere negative on M. By taking the infimum of $\beta^Y(u)$ over $u \in \mathcal{L}(g_+)$, an improved version of (1.12) is obtained:

Corollary 1.13. Let the setup be the same as that in Theorem 1.12, and define $\hat{\beta}^Y := \inf_{u \in \mathcal{L}(g_+)} \beta^Y(u)$. Then

(1.13)
$$\left(\frac{k-\alpha-\hat{\beta}^Y}{2}\right)^2 \le \lambda_{1,2}(Y).$$

Note that $\hat{\beta}^{Y}$ is an invariant of $\iota^{*}g_{+}$.

Estimate (1.13) can be thought of as a generalization of Cheung-Leung's estimate (1.10) to submanifolds living in a general Poincaré-Einstein space (X^{n+1}, g_+) whose conformal infinity has non-negative Yamabe invariant. When the ambient space is $(\mathbb{H}^{n+1}(-1), g_H)$, we observe that $\hat{\beta}^Y = 0$ for every Y and recover Cheung-Leung's result: **Corollary 1.14.** Let Y^{k+1} be a complete and noncompact submanifold in hyperbolic space $\mathbb{H}^{n+1}(-1)$. If the mean curvature vector of Y has norm satisfying $||H|| \leq \alpha < k$, then it follows that $\left(\frac{k-\alpha}{2}\right)^2 \leq \lambda_{1,2}(Y)$.

In the minimal case, Corollary 1.14 and Corollary 1.9 together give:

Corollary 1.15. If Y^{k+1} is a minimal conformally compact submanifold of $\mathbb{H}^{n+1}(-1)$, then $\left(\frac{k}{2}\right)^2 = \lambda_{1,2}(Y)$.

As another consequence of Theorem 4.2, we derive a stability result for complete and non-compact minimal hypersurfaces in Poincaré-Einstein spaces whose conformal infinity has non-negative Yamabe invariant. Indeed, recall that a minimal hypersurface Y^n living in a Poincaré-Einstein space (i.e. $\operatorname{Ric}(g_+) = -ng_+$) is stable if

(1.14)
$$\int_{Y} (|\nabla_{h_{+}} f|^{2} - (|B|^{2} - n))f^{2} dv_{h_{+}} \ge 0$$

for every compactly supported function f, where B denotes the second fundamental form of Y. It is expected then that lower bounds on $\lambda_{1,2}(Y)$ together with upper bounds on $|B|^2$ will yield stability results; for instance, see section 5 in [2]. This generalizes a result of Seo in [26]¹.

Corollary 1.16. Let Y^n be a complete and non-compact minimal hypersurface in a Poincaré-Einstein space (X^{n+1}, g_+) where the conformal infinity of g_+ is of non-negative Yamabe type. If $|B|^2 \leq \frac{(n-1-\hat{\beta}^Y)^2}{4} + n$ at every point of Y and $\hat{\beta}^Y \leq n-1$, then Y is stable. If $(X^{n+1}, g_+) = (\mathbb{H}^{n+1}(-1), g_H)$, then $|B|^2 \leq \frac{(n+1)^2}{4}$ everywhere implies stability.

We now provide a geometric interpretation of the invariant $\hat{\beta}^{Y}$ introduced in Corollary 1.13. On any AH space (X^{n+1}, g_+) , for any defining function r, a unique u solving $\Delta_{g_+} u = (n+1)u$ and satisfying $u - r^{-1} = O(1)$ exists; see discussion in Section 4.2. Therefore, it makes sense to compactify the metric g_+ using u^{-1} , that is, we consider the compact space $(\overline{X}^{n+1}, u^{-2}g_+)$. It was shown by Qing in [24] that this compactification satisfies nice properties, allowing the author to obtain an important rigidity result for Poincaré-Einstein spaces whose conformal infinity is the round sphere. Inspired by the same approach, a simple calculation leads to the following:

Proposition 1.17. Let (X^{n+1}, g_+) be a Poincaré-Einstein space and let u be as described above. Set $g_u = u^{-2}g_+$. If E_{g_u} denotes the trace-free Ricci tensor of g_u and $b(u) = \nabla^2_{g_+}u - ug_+$ is the trace-free hessian of u, then $E_{g_u} = (n-1)\frac{b(u)}{u}$. Moreover, b(u) = 0 if and only if R_{g_u} is constant.

Recall that in the context of Theorem 1.12, for a given $u \in \mathcal{L}(g_+)$, in order to obtain $\beta^Y(u)$ one first need to restrict $u^{-1}b(u) = (n-1)^{-1}E_{g_u}$ to the normal bundle of Y, take the trace with respect to $(g_+)|_{TY^{\perp}}$, and then take the supremum over Y. The invariant $\hat{\beta}^Y$ is then $\hat{\beta}^Y = (n-1)^{-1} \inf_{u \in \mathcal{L}(g_+)} \operatorname{tr}_{g_+}(E_{g_u}|_{TY^{\perp}})$. An

interesting problem is to investigate for which submanifolds Y is $\hat{\beta}^Y = 0$. For instance, we could ask if there are submanifolds for which $\hat{\beta}^Y > 0$ and whether or not (1.13) is optimal in such cases.

We close our introduction with a few remarks about the estimate (1.8) in Corollary 1.9, particularly in the case where p = 2, while reviewing some previous work. In [26], the author works with complete non-compact minimal hypersurfaces Y^n in $\mathbb{H}^{n+1}(-1)$ which are stable, that is, for which the second variation of the area functional with respect to compactly supported normal variations satisfy a sign condition (see condition (1.2) in [26]). Under a finite assumption on the L^2 -norm of the second fundamental form of such submanifolds, Seo managed to prove that

(1.15)
$$\lambda_{1,2}(Y) \le n^2$$

The key element in their proof is the construction of a test function for the Rayleigh quotient in an arbitrary geodesic ball involving the norm of the second fundamental form - this is how the finite assumption and the stability condition come into play.

In Fu-Tao's work [11], they managed to improve Seo's upper bound (1.15). In fact, under similar integrability and stability assumptions, they show that for a complete and non-compact hypersurface Y^n in

¹See remark 4.2

 $\mathbb{H}^{n+1}(-1)$ it holds that

(1.16)
$$\lambda_{1,2}(Y) \le \left(\frac{n-1}{2}\right)^2 \left(1 - \frac{|H|^2}{(k+1)^2}\right),$$

where H is the scalar mean curvature of Y. In particular, if it is minimal, then it gives the sharp upper bound. They also obtain the same upper bound for higher codimensional submanifolds Y^{k+1} under the assumption that H is parallel and that $\int_{Y} |\mathring{B}|^p dv_h < \infty$ for $p \ge k+1$, where \mathring{B} is the traceless second fundamental form of Y and $h = \iota^* g_H$ is the induced metric from (\mathbb{H}^{n+1}, g_H) ; see Theorem 1 in [11]. This integrability assumption always holds: indeed, let r be any defining function for (\mathbb{H}^{n+1}, g_H) , and denote by \bar{h} the metric on Y induced from the compactified metric $\overline{g}_H = r^2 g_H$. Then, denoting by $\overset{\circ}{B}$ the traceless second fundamental form with respect to \bar{h} and using conformal covariance, we deduce

(1.17)
$$\int_{Y} |\mathring{B}|^{p} dv_{h} = \int_{Y} r^{p} \left| \overset{\circ}{\overline{B}} \right|^{p} dv_{h} = \int_{Y} r^{p-(k+1)} \left| \overset{\circ}{\overline{B}} \right|^{p} dv_{\bar{h}} < \infty,$$

where the inequality follows from the fact that \bar{h} extends smoothly to the compact set \overline{Y} and the integrand is continuous there.

Finally, we would like to reiterate that upper bounds for Dirichlet eigenvalues should only be influenced by the asymptotic behavior of the manifold at infinity; see the discussion prior to the statement of Theorem 1.2. In particular, no global integrability assumptions on the second fundamental form or stability assumptions should be necessary if the manifold satisfies certain asymptotics at infinity. This is precisely the case when the submanifolds are asymptotically CMC and the ambient manifold is AH.

1.3. Organization of the paper. In section 2, we explain the asymptotic properties of an AH manifold in terms of a special defining function r for its boundary M. In particular, we describe how the metric can be written in normal form in a neighborhood of the boundary and the specific asymptotics of the volume form there. In section 3, we provide the proof of one of our main results, Theorem 1.2. The techniques herein are novel and self-contained. Finally, in section 4, we furnish the proofs of results concerning eigenvalue estimates on submanifolds of asymptotically hyperbolic manifolds, as elucidated in section 1.2. From this section, there are two aspects that we would like to highlight. In subsection 4.1, while proving Proposition 1.8, we noticed that the asymptotic norm of the mean curvature determines both the angles at which the submanifold meets the boundary at infinity and the asymptotic sectional curvature, which could be of independent interest. On the other hand, in section 4.2, we explain what we refer to as a Lee-eigenfunction. The reader will notice that we employ two type of test functions throughout our work: u^{-s} and $r^{s'}$ for particular $s, s' \in \mathbb{R}$. For the upper bounds, we utilize $r^{s'}$ as we only need to know what happens at infinity. For lower bounds, we employ $(\Delta_{g_+} u = (n+1)u$ everywhere on X).

1.4. Acknowledgements. This work was initiated during the International Doctoral Summer School In Conformal Geometry and Non-local Operators at the "Instituto de Matemáticas de la Universidad de Granada" (IMAG). We thank the organizers, Azahara DelaTorre Pedraza and María del Mar González, for their kind invitation. The first author was supported by the NSF grant RTG-DMS-1502424.

2. Preliminaries for Estimates on Asymptotically Hyperbolic Manifolds

Recall that given a compact manifold \overline{X}^{n+1} with boundary M and interior X, a complete Riemannian metric g_+ on X is said conformally compact if $\overline{g} := r^2 g_+$ extends to a metric on \overline{X} . Here r is a defining function for M, that is, a positive function on X which vanishes on M and with non-vanishing gradient on M. Furthermore, if the asymptotic sectional curvatures approach -1 at the boundary, then we say that (X^{n+1}, g_+) is asymptotically hyperbolic.

A defining function r determines for some $r_0 > 0$ an identification of $M \times [0, r_0)$ with a neighborhood of M in X, as follows: any $(p, t) \in M \times [0, r_0)$ corresponds to $\phi(p, t)$, where ϕ is the flow of $\nabla_{\overline{g}} r$. If r is a special defining function, that is, if r is such that $|\nabla_{g_+} r|^2 = r^2$ in a neighborhood of M, then $r(\phi(p, t)) = t$. This means that we can think about the t coordinate as just being r and $\nabla_{\overline{g}} r$ is orthogonal to the slices $M \times \{t\}$. The metric \overline{g} then takes the form

(2.1)
$$\overline{g} = g_r + dr^2$$

where g_r is a 1-parameter family of metrics on M. It can be shown that

(2.2)
$$g_r = g_0 - 2Lr + O(r^2),$$

where L is the second fundamental form of M with respect to \overline{g} . In $M \times [0, r_0)_r$, we can write the volume form of g_+ as

(2.3)
$$dv_{g_{+}} = r^{-n-1} \left(\frac{\det g_{r}}{\det g_{0}}\right)^{1/2} dv_{g_{0}} dr$$

Then using Jacobi's formula

(2.4)
$$\frac{d}{dt}(\det A(t)) = (\det A(t)) * \operatorname{trace}(A^{-1}(t)A'(t)),$$

we deduce

(2.5)
$$\left(\frac{\det g_r}{\det g_0}\right)^{1/2} = 1 + \nu^{(1)}r + O(r^2)$$

where $\nu^{(1)} = -H^M$ and H^M is the mean curvature of M with respect to \overline{g} .

3. Upper Bounds for $\lambda_{1,p}(X)$ Proof of Theorem 1.2 and of Corollary 1.3

3.1. **Proof of Theorem 1.2.** Our strategy to prove Theorem 1.2 is quite standard. Using the variational characterization of *p*-Dirichlet eigenvalues, given in (1.4), we compute the Rayleigh quotient for a suitable test function that will yield the sharp bound. Let us be more precise. For $\gamma \in (0, r_0)$, consider the bounded domains given by

(3.1)
$$X_{\gamma} := \{ x \in X^{n+1} : r(x) \ge \gamma \}.$$

Notice that the boundary of each X_{γ} , the γ -level set of r, is a smooth hypersurface. By the domain monotonicity property of p-Dirichlet eigenvalues (Lemma 1.1 in [15]), we have that $\lambda_{1,p}(X_{\gamma})$ decreases monotonically to $\lambda_{1,p}(X)$ as $\gamma \to 0^+$. Moreover,

(3.2)
$$\lambda_{1,p}(X) \le \lambda_{1,p}(X_{\gamma}) = \inf_{f \in W_0^{1,p}(X_{\gamma}) \setminus \{0\}} \frac{\int_{X_{\gamma}} |\nabla_{g_+}|f||^p \, dv_{g_+}}{\int_{X_{\gamma}} |f|^p \, dv_{g_+}}.$$

We want to use $r^{\frac{n}{p}}$ as a test function because a special defining function should capture the behavior of X at infinity. The issue is that r is positive everywhere in the interior of \overline{X} and so it cannot be used as a test function. However, we can multiply it by a suitable cut-off ϕ in such a way that we capture the same behavior near M. To this end, we introduce

Definition 3.1. For $\epsilon \in (0, r_0)$ and $\delta > 0$ small enough, define $\phi_{\epsilon\delta} \colon \overline{X} \to \mathbb{R}$ by

(3.3)
$$\phi_{\epsilon\delta}(p) := \begin{cases} 1, & \epsilon \le r(p) \\ \frac{1+\delta}{\epsilon}(r(p) - \frac{\epsilon\delta}{1+\delta}), & \frac{\epsilon\delta}{1+\delta} \le r(p) \le \epsilon \\ 0, & 0 \le r(p) \le \frac{\epsilon\delta}{1+\delta} \end{cases}$$

We also need to introduce some big O notation that will make some of the computations clearer. We say that a function w is $O(a, b) = O(a(\epsilon), b(\delta))$ if there exist positive constants A, B, ϵ_o and δ_o such that $|w(\epsilon, \delta)| \leq A|a(\epsilon)|$ for all $0 \leq \epsilon < \epsilon_o$ and $|w(\epsilon, \delta)| \leq B|b(\delta)|$ for all $0 \leq \delta < \delta_o$. For instance, a function $w(\epsilon, \delta)$ being $O(\epsilon^2, 1)$ means that, in particular, the function remains bounded as $\delta \to 0^+$ while for any fixed $\delta > 0$ it is $O(\epsilon^2)$ as $\epsilon \to 0^+$. A function is O(1, 1) if it remains bounded as both $\epsilon \to 0^+$ and $\delta \to 0^+$. As in other sections of our work, we employ classical big O notation as well. For instance, a function t(r) is O(r) if there exist positive constants R and r_o such that $|t(r)| \leq R \cdot r$ for $0 \leq r \leq r_o$.

Proof of Theorem 1.2. Set

(3.4)
$$q_p(\epsilon, \delta, s) := \frac{\int_X |\nabla_{g_+}(r^s \phi_{\epsilon \delta})|^p \, dv_{g_+}}{\int_X r^{ps} \phi_{\epsilon \delta}^p \, dv_{g_+}} = \frac{N_p(\epsilon, \delta, s)}{D_p(\epsilon, \delta, s)},$$

where $\frac{n-p}{p} < s < \frac{n}{p}$, and notice that thanks to (3.2) we have

(3.5)
$$\lambda_{1,p}(X) \le \lim_{\epsilon \to 0^+} \lim_{\delta \to 0^+} q_p(\epsilon, \delta, s).$$

Also, combining (2.3) and (2.5), we get

(3.6)
$$dv_{g_{+}} = r^{-n-1}(1 - H^{M}r + O(r^{2})) \ dv_{g_{0}}dr = r^{-n-1}[1 + O(r)] \ dv_{g_{0}}dr$$

The range in the values of s will be explained as we proceed with the proof.

We begin by studying the denominator in $q_p(\epsilon, \delta, s)$:

$$D_{p}(\epsilon, \delta, s) = \int_{\{r \ge r_{0}\}} r^{ps} dv_{g_{+}} + \int_{\epsilon}^{r_{0}} \int_{M} r^{ps-n-1} [1+O(r)] dv_{g_{0}} dr$$

$$+ \left(\frac{1+\delta}{\epsilon}\right)^{p} \int_{\frac{\epsilon\delta}{1+\delta}}^{\epsilon} \int_{M} \left(r - \frac{\epsilon\delta}{1+\delta}\right)^{p} \cdot r^{ps-n-1} [1+O(r)] dv_{g_{0}} dr$$

$$= O(1,1) + \frac{\operatorname{Vol}_{g_{0}}(M)}{ps-n} \cdot [r^{ps-n}|_{\epsilon}^{r_{o}} + O(\epsilon^{ps-n+1},1)$$

$$+ \left(\frac{1+\delta}{\epsilon}\right)^{p} \int_{\frac{\epsilon\delta}{1+\delta}}^{\epsilon} \int_{M} \left(r - \frac{\epsilon\delta}{1+\delta}\right)^{p} \cdot r^{ps-n-1} [1+O(r)] dv_{g_{0}} dr$$

For $\frac{\epsilon\delta}{1+\delta} < r < \epsilon$, we can expand

(3.8)
$$\left(r - \frac{\epsilon\delta}{1+\delta}\right)^p = r^p - \left(pr^{p-1}\epsilon\left(\frac{\delta}{1+\delta}\right) - \frac{p(p-1)}{2}r^{p-2}\epsilon^2\left(\frac{\delta}{1+\delta}\right)^2 + \cdots\right).$$

Therefore,

$$\begin{split} D_p(\epsilon,\delta,s) &= O(1,1) - \frac{\operatorname{Vol}_{g_0}(M)}{ps-n} \epsilon^{ps-n} + O(\epsilon^{ps-n+1},1) \\ &+ \left(\frac{1+\delta}{\epsilon}\right)^p \int_{\frac{\epsilon}{\epsilon+\delta}}^{\epsilon} \int_M \left\{ r^{p+ps-n-1} - pr^{p+ps-n-2}\epsilon \left(\frac{\delta}{1+\delta}\right) + \cdots \right\} [1+O(r)] \, dv_{g_0} dr \\ &= O(1,1) - \frac{\operatorname{Vol}_{g_0}(M)}{ps-n} \epsilon^{ps-n} + O(\epsilon^{ps-n+1},1) \\ &+ \operatorname{Vol}_{g_0}(M) \left(\frac{1+\delta}{\epsilon}\right)^p \cdot \left[\frac{r^{p+ps-n}}{p+ps-n} - p\epsilon \left(\frac{\delta}{1+\delta}\right) \frac{r^{p+ps-n-1}}{p+ps-n-1} + \cdots \right]_{\frac{\epsilon}{1+\delta}}^{\epsilon} + O(\epsilon^{ps-n+1},\delta) \\ &= O(1,1) - \frac{\operatorname{Vol}_{g_0}(M)}{ps-n} \epsilon^{ps-n} + O(\epsilon^{ps-n+1},1) \\ &+ \operatorname{Vol}_{g_0}(M) \left(\frac{1+\delta}{\epsilon}\right)^p \frac{1}{p+ps-n} \left(\epsilon^{p+ps-n} - \left(\frac{\epsilon\delta}{1+\delta}\right)^{p+ps-n}\right) \\ &- \operatorname{Vol}_{g_0}(M) \left(\frac{1+\delta}{\epsilon}\right)^p \frac{p\epsilon}{p+ps-n-1} \left(\frac{\delta}{1+\delta}\right) \left(\epsilon^{p+ps-n-1} - \left(\frac{\epsilon\delta}{1+\delta}\right)^{p+ps-n-1}\right) \\ &+ O(\epsilon^{ps-n}, \delta^{p+ps-n}) \\ &= O(1,1) - \frac{\operatorname{Vol}_{g_0}(M)}{ps-n} \epsilon^{ps-n} + O(\epsilon^{ps-n+1},1) + \operatorname{Vol}_{g_0}(M) \frac{(1+\delta)^p}{p+ps-n} e^{ps-n} \\ &- \operatorname{Vol}_{g_0}(M) \frac{\delta^{p+ps-n}}{p+ps-n} \left(\frac{\epsilon}{1+\delta}\right)^{ps-n} - \operatorname{Vol}_{g_0}(M)(1+\delta)^{p-1} \frac{p\epsilon^{ps-n}\delta}{p-ps-n-1} \\ &+ \operatorname{Vol}_{g_0}(M) \frac{p\delta^{p+ps-n}}{p+ps-n-1} \left(\frac{\epsilon}{1+\delta}\right)^{ps-n} \end{split}$$

That is,

(3.9)
$$D_p(\epsilon, \delta, s) = O(1, 1) + \operatorname{Vol}_{g_0}(M)\epsilon^{ps-n} \left(\frac{(1+\delta)^p}{p+ps-n} - \frac{1}{ps-n}\right) + O(\epsilon^{ps-n+1}, 1) + O(\epsilon^{ps-n}, \delta^{p+ps-n}).$$

Notice that some of the powers in δ are p + ps - n. Later we will take the limit as $\delta \to 0^+$, and so we need p + ps - n > 0, that is, we need $s > \frac{n-p}{p}$. We turn our focus to the numerator in $q_p(\epsilon, \delta, s)$. Recall that r is a special defining function, and thus $|\nabla_{g_+}r|^2 = r^2$ in a collar neighborhood of M. Then

$$\begin{split} N_{p}(\epsilon,\delta,s) &= \int_{X} |\nabla_{g_{+}}(r^{s}\phi_{\epsilon\delta})|^{p} dv_{g_{+}} = \int_{\{r \geq r_{0}\}} |\nabla_{g_{+}}(r^{s})|^{p} dv_{g_{+}} + \int_{X \setminus \{r \geq r_{0}\}} |r^{s}\nabla_{g_{+}}\phi_{\epsilon\delta} + sr^{s-1}\phi_{\epsilon\delta}\nabla_{g_{+}}r|^{p} dv_{g_{+}} \\ &= O(1,1) + \int_{X \setminus \{r \geq r_{0}\}} (r^{2s}|\nabla_{g_{+}}\phi_{\epsilon\delta}|^{2} + s^{2}r^{2(s-1)}\phi_{\epsilon\delta}^{2}|\nabla_{g_{+}}r|^{2} + 2sr^{2s-1}\phi_{\epsilon\delta}\langle\nabla_{g_{+}}r, \nabla_{g_{+}}\phi_{\epsilon\delta}\rangle_{g_{+}})^{\frac{p}{2}} dv_{g_{+}} \\ &= O(1,1) + \int_{\epsilon}^{r_{o}} \int_{M} s^{p}r^{p(s-1)}r^{p}r^{-n-1}[1+O(r)] dv_{g_{o}}dr \\ &+ \int_{\frac{\epsilon\delta}{1+\delta}}^{\epsilon} \int_{M} (r^{2s}|\nabla_{g_{+}}\phi_{\epsilon\delta}|^{2} + s^{2}r^{2(s-1)}\phi_{\epsilon\delta}^{2}r^{2} + 2sr^{2s-1}\phi_{\epsilon\delta}\langle\nabla_{g_{+}}r, \nabla_{g_{+}}\phi_{\epsilon\delta}\rangle_{g_{+}})^{\frac{p}{2}}r^{-n-1}[1+O(r)] dv_{g_{0}}dr \\ &= O(1,1) - \frac{\operatorname{Vol}_{g_{0}}(M)}{ps-n}s^{p}\epsilon^{ps-n} + O(\epsilon^{ps-n+1},1) \\ &+ \int_{\frac{\epsilon\delta}{1+\delta}}^{\epsilon} \int_{M} (r^{2s}|\nabla_{g_{+}}\phi_{\epsilon\delta}|^{2} + s^{2}r^{2(s-1)}\phi_{\epsilon\delta}^{2}r^{2} + 2sr^{2s-1}\phi_{\epsilon\delta}\langle\nabla_{g_{+}}r, \nabla_{g_{+}}\phi_{\epsilon\delta}\rangle_{g_{+}})^{\frac{p}{2}}r^{-n-1}[1+O(r)] dv_{g_{0}}dr \end{split}$$

Using $\nabla_{g_+}\phi_{\epsilon\delta} = \frac{1+\delta}{\epsilon}\nabla_{g_+}r$ on $\frac{\epsilon\delta}{1+\delta} < r < \epsilon$, we can write

$$N_{p}(\epsilon, \delta, s) = O(1, 1) - \frac{\operatorname{Vol}_{g_{0}}(M)}{ps - n} s^{p} \epsilon^{ps - n} + O(\epsilon^{ps - n + 1}, 1) + \int_{\frac{\epsilon\delta}{1+\delta}}^{\epsilon} \int_{M} \left(r^{2s+2} \left(\frac{1+\delta}{\epsilon} \right)^{2} + s^{2} r^{2s} \left(\frac{1+\delta}{\epsilon} \right)^{2} \left(r - \frac{\epsilon\delta}{1+\delta} \right)^{2} + 2s r^{2s+1} \left(\frac{1+\delta}{\epsilon} \right)^{2} \left(r - \frac{\epsilon\delta}{1+\delta} \right) \right)^{\frac{p}{2}} r^{-n-1} [1+O(r)] \, dv_{g_{0}} dr$$

We now proceed to simplify the main factor in the integrand:

$$r^{2s+2} \left(\frac{1+\delta}{\epsilon}\right)^2 + s^2 r^{2s} \left(\frac{1+\delta}{\epsilon}\right)^2 \left(r - \frac{\epsilon\delta}{1+\delta}\right)^2 + 2sr^{2s+1} \left(\frac{1+\delta}{\epsilon}\right)^2 \left(r - \frac{\epsilon\delta}{1+\delta}\right)$$

$$= \left(\frac{1+\delta}{\epsilon}\right)^2 \left\{r^{2s+2} + s^2 r^{2s} \left(r^2 - \frac{2r\epsilon\delta}{1+\delta} + \left(\frac{\epsilon\delta}{1+\delta}\right)^2\right) + 2sr^{2s+1} \left(r - \frac{\epsilon\delta}{1+\delta}\right)\right\}$$

$$(3.11) \qquad = \left(\frac{1+\delta}{\epsilon}\right)^2 \left\{r^{2s+2} + s^2 r^{2s+2} - \frac{2s^2 r^{2s+1}\epsilon\delta}{1+\delta} + \frac{s^2 r^{2s} \epsilon^2 \delta^2}{(1+\delta)^2} + 2sr^{2s+2} - \frac{2sr^{2s+1}\epsilon\delta}{1+\delta}\right\}$$

$$= \left(\frac{1+\delta}{\epsilon}\right)^2 \left\{r^{2s+2} (s^2 + 2s + 1) - \frac{2r^{2s+1}\epsilon\delta}{1+\delta} (s^2 + s) + \frac{s^2 r^{2s} \epsilon^2 \delta^2}{(1+\delta)^2}\right\}$$

$$= \left(\frac{1+\delta}{\epsilon}\right)^2 r^{2s+2} (s^2 + 2s + 1) + \epsilon^{-1} r^{2s+1} \cdot O(1,\delta) + r^{2s} \cdot O(1,\delta^2)$$

Recall that we will first take the limit $\delta \to 0^+$. Therefore, we rewrite (3.11) as

(3.12)
$$\left(\frac{1+\delta}{\epsilon}\right)^2 r^{2s+2}(s^2+2s+1) + r^{2s+1}O(\delta) + r^{2s}O(\delta^2).$$

After raising to the power of $\frac{p}{2}$, we obtain

$$(3.13) \qquad \qquad \left(\frac{1+\delta}{\epsilon}\right)^{p} r^{ps+p} (s^{2}+2s+1)^{\frac{p}{2}} + \frac{p}{2} r^{(2s+2)(\frac{p}{2}-1)} (r^{2s+1}O(\delta) + r^{2s}O(\delta^{2})) \\ \qquad \qquad + \frac{1}{2} \cdot \frac{p}{2} \left(\frac{p}{2}-1\right) r^{(2s+2)(\frac{p}{2}-2)} (r^{2s+1}O(\delta) + r^{2s}O(\delta^{2}))^{2} + \cdots \\ \qquad \qquad = \left(\frac{1+\delta}{\epsilon}\right)^{p} r^{ps+p} (s^{2}+2s+1)^{\frac{p}{2}} + \frac{p}{2} r^{(2s+2)(\frac{p}{2}-1)} (r^{2s+1}O(\delta) + r^{2s}O(\delta^{2})) \\ \qquad \qquad + \frac{1}{2} \cdot \frac{p}{2} \left(\frac{p}{2}-1\right) r^{(2s+2)(\frac{p}{2}-2)} (r^{2s+1}O(\delta) + r^{2s}O(\delta^{2}))^{2} + \cdots \\ \qquad \qquad = \left(\frac{1+\delta}{\epsilon}\right)^{p} r^{ps+p} (s^{2}+2s+1)^{\frac{p}{2}} + \frac{p}{2} (r^{ps+p-1}O(\delta) + r^{ps+p-2}O(\delta^{2})) + \cdots,$$

and notice that, starting from the second term above, we have a Laurent expansion in both r and δ such that the exponent of r and that of δ in each term always adds up to ps + p. Moreover, the power of δ is positive. Now, multiplying by $r^{-n-1}[1 + O(r)]$, gives

(3.14)
$$\begin{pmatrix} \frac{1+\delta}{\epsilon} \end{pmatrix}^p r^{ps+p-n-1} (s^2+2s+1)^{\frac{p}{2}} + \frac{p}{2} (r^{ps+p-n-2}O(\delta) + r^{ps+p-n-3}O(\delta^2)) + F_1(r,\delta) \\ + \epsilon^{-p}O(r^{ps+p-n}) + O(r^{ps+p-n-1},\delta) + O(r^{ps+p-n-2},\delta^2) + F_2(r,\delta),$$

where $F_1(r, \delta)$ is a Laurent expansion in both r and δ such that in each term the exponents of r and δ add up to ps + p - n - 1, while $F_2(r, \delta)$ is the same but the powers add up to ps + p - n. Once again, the power of δ is positive in every term of the expansions F_1 and F_2 .

Recall that ps + p - n > 0. Therefore we have the following two cases to consider. First, let us assume that $ps + p - n - k \neq -1$ for all $k \in \mathbb{N} \setminus \{1\}$. Performing the double integral in (3.10) gives

$$(3.15) \qquad \frac{\operatorname{Vol}_{g_0}(M)}{ps+p-n} (1+\delta)^p (s^2+2s+1)^{\frac{p}{2}} \epsilon^{ps-n} \left(1 - \left(\frac{\delta}{1+\delta}\right)^{ps+p-n}\right) \\ + [r^{ps+p-n-1}\Big|_{\frac{\epsilon\delta}{1+\delta}}^{\epsilon} \cdot O(\delta) + [r^{ps+p-n-2}\Big|_{\frac{\epsilon\delta}{1+\delta}}^{\epsilon} \cdot O(\delta^2) + O(\delta^{\alpha}) \\ + O(\epsilon^{ps-n+1}, 1) + [r^{ps+p-n}\Big|_{\frac{\epsilon\delta}{1+\delta}}^{\epsilon} \cdot O(\delta) + [r^{ps+p-n-1}\Big|_{\frac{\epsilon\delta}{1+\delta}}^{\epsilon} \cdot O(\delta^2),$$

where $\alpha > 0$. The terms coming from integrating F_1 and F_2 have both been absorbed into the term $O(\delta^{\alpha})$. Taking the limit as $\delta \to 0^+$, we obtain

(3.16)
$$\frac{\operatorname{Vol}_{g_0}(M)}{ps+p-n}(s^2+2s+1)^{\frac{p}{2}}\epsilon^{ps-n}+O(\epsilon^{ps-n+1},1).$$

On the other hand, if ps + p - n - k = -1 for some $k \in \mathbb{N} \setminus \{1\}$, then log terms will appear somewhere after the first term. However, in those cases, the contribution will be of the form $\log(\delta) \cdot O(\delta^{\alpha})$, where $\alpha > 0$. Therefore, after taking $\delta \to 0^+$, we still get (3.16).

Combine (3.10) and (3.16) to get

(3.17)
$$\lim_{\delta \to 0^+} N_p(\epsilon, \delta, s) = O(1, 1) - \frac{\operatorname{Vol}_{g_0}(M)}{ps - n} s^p \epsilon^{ps - n} + O(\epsilon^{ps - n + 1}, 1) + \frac{\operatorname{Vol}_{g_0}(M)}{ps + p - n} (s^2 + 2s + 1)^{\frac{p}{2}} \epsilon^{ps - n}.$$

Putting (3.9) and (3.17) together gives

(3.18)
$$\lim_{\delta \to 0^+} q_p(\epsilon, \delta, s) \cdot \left\{ O(1, 1) + \operatorname{Vol}_{g_0}(M) \epsilon^{ps-n} \left(\frac{1}{p+ps-n} - \frac{1}{ps-n} \right) + O(\epsilon^{ps-n+1}, 1) \right\} \\ = O(1, 1) - \frac{\operatorname{Vol}_{g_0}(M)}{ps-n} s^p \epsilon^{ps-n} + O(\epsilon^{ps-n+1}, 1) + \frac{\operatorname{Vol}_{g_0}(M)}{ps+p-n} (s^2 + 2s + 1)^{\frac{p}{2}} \epsilon^{ps-n}.$$

Let us recall that n - ps > 0. Multiply across by ϵ^{n-ps} , let $\epsilon \to 0^+$ and divide by $\operatorname{Vol}_{g_0}(M)$ to obtain

(3.19)
$$\lim_{\epsilon \to 0^+} \lim_{\delta \to 0^+} q_p(\epsilon, \delta, s) \cdot \left\{ \frac{1}{p+ps-n} - \frac{1}{ps-n} \right\} = \frac{(s^2+2s+1)^{\frac{p}{2}}}{p+ps-n} - \frac{s^p}{ps-n}$$

Going back to (3.5), we conclude that

(3.20)
$$\lambda_{1,p}(X) \le s^p \left(s+1-\frac{n}{p}\right) + (1+2s+s^2)^{\frac{p}{2}} \left(\frac{n}{p}-s\right)$$

for all $\frac{n-p}{p} < s < \frac{n}{p}$. Letting $s \to \frac{n-p}{p}^+$ or $s \to \frac{n}{p}^-$ yields the desired upper bound.

3.2. Proof of Corollary 1.3.

Proof of Corollary 1.3. Let r be a defining function for M, and recall that \bar{g} denotes the compactified metric r^2g_+ . Using the Transformation law (1.1) for $\kappa^2g_+ = r^{-2}(\kappa^2\bar{g})$, we obtain

(3.21)
$$R_{ijkl}^{\kappa^{2}g_{+}} = -|dr|_{\kappa^{2}\overline{g}}^{2}((\kappa^{2}g)_{ik}(\kappa^{2}g)_{jl} - (\kappa^{2}g)_{il}(\kappa^{2}g)_{jk}) + O_{ijkl}(r^{-3})$$
$$= -(\kappa^{-2}|dr|_{\overline{g}}^{2})((\kappa^{2}g)_{ik}(\kappa^{2}g)_{jl} - (\kappa^{2}g)_{il}(\kappa^{2}g)_{jk}) + O_{ijkl}(r^{-3})$$

By assumption the manifold (X^{n+1}, g_+) is $\operatorname{AH}(-\kappa^2)$. As explained in the introduction, this means that $|dr|_{\overline{g}}^2|_M = \kappa^2 \iff \kappa^{-2}|dr|_{\overline{g}}^2|_M = 1$. This allow us to conclude that the sectional curvatures of (X^{n+1}, g_+) approach -1 at M, that is, $(X^{n+1}, \kappa^2 g_+)$ is asymptotically hyperbolic. Therefore, it follows from Theorem 1.2 that $\lambda_{1,p}(X, \kappa^2 g_+) \leq \left(\frac{n}{p}\right)^p$. Finally, by the variational characterization (1.4) of the first *p*-Dirichlet eigenvalue, the scaling property

(3.22)
$$\lambda_{1,p}(X,\kappa^2 g_+) = \inf_{\Omega} \lambda_{1,p}(\Omega,\kappa^2 g_+) = \kappa^{-p} \inf_{\Omega} \lambda_{1,p}(\Omega,g_+) = \kappa^{-p} \lambda_{1,p}(X,g_+)$$

holds and the result follows.

4. Estimates on Submanifolds - Proofs

4.1. **Proof of Proposition 1.8.** We start with some preliminaries. Let (x^{α}) be a coordinate system on Y, and let $(\mu_{\alpha'})$ be a unit-normal frame for g_+ to Y that is adapted in the following sense: if $g(\nabla_{\overline{g}}r, \mu_{\alpha'}) < 0$, then replace $\mu_{\alpha'}$ with $-\mu_{\alpha'}$. Let $\overline{\mu}_{\alpha'} = r^{-1}\mu_{\alpha'}$; this is a unit-normal vector for \overline{g} . Then the vector-valued second fundamental form is given by

(4.1)
$$II(X,Z) = B^{\alpha'}(X,Z)\mu_{\alpha'},$$

where

(4.2)
$$g_{\alpha'\gamma'}B^{\gamma'}(X,Z) = g_+(\nabla_X^{g_+}Z,\mu_{\alpha'})$$

For $q \in Y$, let $P_q: T_q X \to T_q Y^{\perp}$ be the canonical normal-projection map. More explicitly, with respect to our chosen frame, this canonical projection takes the form $[v \mapsto \sum_{\alpha'} g(v, \mu_{\alpha'})\mu_{\alpha'}]$. We point out that this projection is independent of our choice of frame. Notice that $\sum_{\alpha'} g(v, \mu_{\alpha'})\mu_{\alpha'} = \sum_{\alpha'} \overline{g}(v, \overline{\mu}_{\alpha'})\overline{\mu}_{\alpha'}$, and observe that

(4.3)
$$\sum_{\alpha'} \overline{g}(\overline{\mu}_{\alpha'}, \nabla_{\overline{g}}r)^2 = |P(\nabla_{\overline{g}}r)|_{\overline{g}}^2 \le |\nabla_{\overline{g}}r|_{\overline{g}}^2 = 1.$$

We define $\Theta_{\alpha'}$ to be the angle between $\nabla_{\overline{q}}r$ and $\mu_{\alpha'}$,

(4.4)
$$\cos \Theta_{\alpha'} := \overline{\mu}_{\alpha'}(r) = \overline{g}(\overline{\mu}_{\alpha'}, \nabla_{\overline{g}}r) = g(\mu_{\alpha'}, r\nabla_{\overline{g}}r).$$

Note that $\Theta_{\alpha'}$ is manifestly a conformal invariant. Using this, we can now rewrite $P(\nabla_{\bar{q}}r)$ as

(4.5)
$$P(\nabla_{\overline{g}}r) = \sum_{\alpha'} \cos \Theta_{\alpha'} \overline{\mu}_{\alpha'}.$$

This gives us an interpretation of $P(\nabla_{\overline{g}}r)$ as being a vector field which contains all the information about the angles formed between $\nabla_{\overline{g}}r$ and the orthonormal frame $\overline{\mu}_{\alpha'}$. Note that if we choose a different orthonormal frame then the individual angles $\Theta_{\alpha'}$ may change but $P(\nabla_{\overline{g}}r)$ is invariant.

Recall the conformal transformation law

(4.6)
$$B_{\alpha\beta}^{\alpha'} = \frac{\overline{B}_{\alpha\beta}^{\alpha'}}{r} + \frac{\overline{\mu}_{\alpha'}(r)\overline{h}_{\alpha\beta}}{r^2}.$$

If we impose our assumption, $g_+(H, H) = C^2 + O(r)$, then

(4.7)
$$H^{\alpha'} = h^{\alpha\beta} B^{\alpha'}_{\alpha\beta}$$

(4.8)
$$= r\overline{h}^{\alpha\beta}\overline{B}^{\alpha}_{\alpha\beta} + \overline{\mu}_{\alpha'}(r)(k+1)$$

Therefore,

(4.9)
$$\cos \Theta_{\alpha'} = \overline{\mu}_{\alpha'}(r) = \frac{H^{\alpha'}}{k+1} + O(r),$$

thus it follows that

(4.10)
$$|P(\nabla_{\overline{g}}r)|_{\overline{g}}^2 = \frac{C^2}{(k+1)^2} + O(r).$$

After recalling (4.3) and taking $r \to 0$, we conclude

(4.11)
$$C^2 \le (k+1)^2.$$

Before we start with the proof of Proposition 1.8, we explain some interesting geometric consequences of the previous formulas. First, (4.10) gives us that the multi-angle vector $P(\nabla_{\overline{g}}r)$ has constant length at infinity given by $\frac{C}{(k+1)}$. If we use (4.9) and think in terms of our frame $\overline{\mu}_{\alpha'}$, then we deduce that the angle between $\overline{\mu}_{\alpha'}$ and $\nabla_{\overline{g}}r$ at infinity is given by $\arccos\left(\frac{H^{\alpha'}}{(k+1)}\right)$. Given $q \in \partial \overline{Y}$, we say that $\Theta := \arccos(|P(\nabla r)(q)|)$ is the **generalized non-oblique angle** at which Y meets M at q. In the hypersurface case, this gives us the non-oblique angle between Y and M at q in the plane determined by any choice of normal vector to Y and $\nabla_{\overline{q}}r$. We will sometimes just refer to this as the generalized angle.

In particular, asymptotically minimal submanifolds (C = 0) meet M at right angles in the sense that all $\Theta_{\alpha'} = \frac{\pi}{2}$ no matter which normal frame we choose, and that asymptotically CMC submanifolds meet M at a constant generalized angle. Furthermore, this gives an upper bound on the possible value of the asymptotic mean curvature for asymptotically CMC submanifolds; we have seen $C^2 \leq (k + 1)^2$. Note that CMC conformally compact submanifolds are special cases of these, so this bound applies to them globally. This shows that the bound is determined purely by the asymptotics. If C = k + 1, then $P(\nabla_{\overline{g}} r)|_{r=0} = \nabla_{\overline{g}} r|_{r=0}$, since $\nabla_{\overline{g}} r|_{r=0}$ is normal to M at r = 0, and, therefore, it follows that $T_q Y \subset T_q M$ for all $q \in \partial \overline{Y}$. This violates the assumption that Y meets M transversely, hence it must be true that C < (k + 1).

Now we proceed with the proof.

Proof of Proposition 1.8. Consider the Gauss Equation

$$\begin{split} R^{X}_{\alpha\beta\gamma\delta} &= R^{Y}_{\alpha\beta\gamma\delta} + g_{\alpha'\beta'} (B^{\alpha'}_{\alpha\delta} B^{\beta'}_{\beta\gamma} - B^{\alpha'}_{\alpha\gamma} B^{\beta'}_{\beta\delta}) \\ &= R^{Y}_{\alpha\beta\gamma\delta} + \sum_{\alpha'} \left(\left[\frac{\overline{B}^{\alpha'}_{\alpha\delta}}{r} + \frac{\overline{\mu}_{\alpha'}(r)\overline{h}_{\alpha\delta}}{r^2} \right] \left[\frac{\overline{B}^{\alpha'}_{\beta\gamma}}{r} + \frac{\overline{\mu}_{\alpha'}(r)\overline{h}_{\beta\gamma}}{r^2} \right] - \left[\frac{\overline{B}^{\alpha'}_{\alpha\gamma}}{r} + \frac{\overline{\mu}_{\alpha'}(r)\overline{h}_{\alpha\gamma}}{r^2} \right] \left[\frac{\overline{B}^{\alpha'}_{\beta\delta}}{r} + \frac{\overline{\mu}_{\alpha'}(r)\overline{h}_{\beta\delta}}{r^2} \right] \right) \\ &= R^{Y}_{\alpha\beta\gamma\delta} + \frac{|P(\nabla_{\overline{g}}r)|^2}{r^4} (\overline{h}_{\alpha\delta}\overline{h}_{\beta\gamma} - \overline{h}_{\alpha\gamma}\overline{h}_{\beta\delta}) + O_{\alpha\beta\gamma\delta}(r^{-3}) \\ &= R^{Y}_{\alpha\beta\gamma\delta} + |P(\nabla_{\overline{g}}r)|^2 (h_{\alpha\delta}h_{\beta\gamma} - h_{\alpha\gamma}h_{\beta\delta}) + O_{\alpha\beta\gamma\delta}(r^{-3}). \end{split}$$

It now follows from (1.1) and (4.10) that

$$R^{Y}_{\alpha\beta\gamma\delta} = \left[\frac{C^2}{(k+1)^2} - 1\right] (h_{\alpha\gamma}h_{\beta\delta} - h_{\alpha\delta}h_{\beta\gamma}) + O_{\alpha\beta\gamma\delta}(r^{-3})$$
$$= -\left[1 - \frac{C^2}{(k+1)^2}\right] (h_{\alpha\gamma}h_{\beta\delta} - h_{\alpha\delta}h_{\beta\gamma}) + O_{\alpha\beta\gamma\delta}(r^{-3})$$

It follows that

(4.12)
$$\sec^{Y}(X,Z) = -\left[1 - \frac{C^{2}}{(k+1)^{2}}\right] + O(r),$$

where (X, Z) is any pair of vector fields on Y that always span a plane and are defined in some collar neighborhood. Since Y is asymptotically CMC, we know by the discussion prior to the proof that C < k+1, it now follows that Y is $\operatorname{AH}\left(-\left(1-\frac{C^2}{(k+1)^2}\right)\right)$.

4.2. **Proof of Theorem 1.12.** In [21], Mazzeo showed that the essential spectrum of an (n+1)-dimensional asymptotically hyperbolic manifold (X, g_+) is $\left[\frac{n^2}{4}, \infty\right)$ with no embedded eigenvalues, leaving the possibility of the existence of finitely many eigenvalues in $(0, \frac{n^2}{4})$. A few years after, Lee proved that if the manifold is, in addition, Poincaré-Einstein and its conformal infinity has nonnegative Yamabe invariant, then the first Dirichlet eigenvalue is exactly $\frac{n^2}{4}$. In particular, such manifolds have no L^2 - eigenvalues.

Since it will be relevant to our work, in what follows we explain Lee's approach in detail. Here is where the assumption that g_+ is conformally compact of order $C^{3,\alpha}$ becomes necessary. Thanks to a result of Barta (see Chapter 3 in [4]), the first Dirichlet eigenvalue $\lambda_{1,2}(\Omega)$ on any bounded domain with smooth boundary satisfies

(4.13)
$$\inf_{x \in \Omega} \left(\frac{-\Delta_{g_+} f}{f} \right) \le \lambda_{1,2}(\Omega),$$

where f is a smooth positive function vanishing on $\partial \Omega$. In [5] (see Section 3), the authors extended Barta's result to complete, non-compact manifolds and showed that

(4.14)
$$\inf_{x \in X} \left(\frac{-\Delta_{g_+} f}{f} \right) \le \lambda_{1,2}(X),$$

where f is a smooth positive function on X.

If $|\nabla_{g_+} u|^2 < 1$ then

With this result at hand, Lee's idea is to construct a positive smooth test function φ on X such that $\frac{-\Delta \varphi}{\varphi} \geq \frac{n^2}{4}$. The test function is constructed as follows. On any asymptotically hyperbolic manifold X and for any smooth defining function r, there exists a unique, smooth and strictly positive function u on X satisfying

(4.15)
$$\begin{cases} \Delta_{g_+} u &= (n+1)u \\ u - r^{-1} &= O(1); \end{cases}$$

see Proposition 4.1 in [16]². We will call any such solution u a Lee-eigenfunction. Notice that if s > 0 and we take $\varphi = u^{-s}$, then

(4.16)
$$\frac{-\Delta_{g_+}\varphi}{\varphi} = -\operatorname{div}_{g_+}(-su^{-s-1}\nabla_{g_+}u) = s(n+1) - s(s+1)\frac{|\nabla_{g_+}u|^2}{u^2}.$$

(4.17)
$$\frac{-\Delta_{g_+}\varphi}{\varphi} \ge s(n+1) - s(s+1) = s(n-s).$$

Then the result follows after observing that s(n-s), as a function of s > 0, has a maximum at $s = \frac{n}{2}$. The main difficulty is proving that

$$\frac{|\nabla_{g_+}u|^2}{u^2} \le 1$$

holds globally on X. It turns out that if (X^{n+1}, g_+) is Poincaré-Einstein and the conformal infinity is of non-negative Yamabe type, then we can select a smooth defining function r such that $(r^2g_+)|_{TM}$ has nonnegative scalar curvature. For the corresponding solution u, (4.18) holds and the result follows; see Theorem A in [16]. This motivates our following definition:

Definition 4.1 (Lee-eigenfunction). Let (X^{n+1}, g_+) be a PE space whose conformal infinity $(M, [g_+]_{\infty})$ is of non-negative Yamabe type. As explained, for a smooth defining function r, there exists a unique, smooth positive solution u solving (4.15). If $(r^2g_+)|_{TM}$ has non-negative scalar curvature, then we say that u is a Lee-eigenfunction.

 $^{^{2}}$ The sign discrepancy is due to different conventions in the definition of the Laplace operator.

Our submanifolds are not necessarily Poincaré-Einstein, and we do not impose any assumptions on its conformal boundary. However, they live inside a Poincaré-Einstein manifold whose conformal infinity has non-negative Yamabe invariant. That means that a Lee-eigenfunction exists on the ambient manifold, as it is AH in particular, and it globally satisfies (4.18). We utilize Barta's inequality (4.14) and Lee's trick applied to the restriction of u^{-s} to Y to derive the lower bound.

Proof of Theorem 1.12. Let us set $\hat{u} := u|_Y$. Also, we denote by h_+ the induced metric on Y, and we use $g^{\alpha'\beta'}, \alpha', \beta' = k + 2, \cdots, n + 1$, to denote the components of g_+^{-1} with respect to a local orthonormal frame for TY^{\perp} . Recall that (see Lemma 2 in [8], for instance)

(4.19)
$$\Delta_{h_{+}}\hat{u} = (\Delta_{g_{+}}u)|_{Y} + H^{\alpha'}u_{\alpha'} - (g^{\alpha'\beta'}\nabla^{2}_{\alpha'\beta'}u)|_{Y},$$

and that, for a Lee-eigenfunction u, its trace-free hessian (w.r.t. g_+) equals $b(u) = \nabla_{g_+}^2 u - ug_+$. Setting $T := \operatorname{tr}_{g_+}(b(u)|_{(TY^{\perp})^2})$, we can therefore write

(4.20)
$$\Delta_{h_{+}}\hat{u} = (\Delta_{g_{+}}u)|_{Y} + H^{\alpha'}u_{\alpha'} - T - (n-k)\hat{u} \\ = (k+1)\hat{u} + H^{\alpha'}u_{\alpha'} - T.$$

Next, for s > 0, consider the test function $\varphi := \hat{u}^{-s}$, and compute

(4.21)
$$\frac{-\Delta_{h+}\varphi}{\varphi} = s\frac{\Delta_{h+}\hat{u}}{\hat{u}} - s(s+1)\frac{|d\hat{u}|_{h+}^2}{\hat{u}^2} = s\left(k+1+H^{\alpha'}\log(u)_{\alpha'}-\frac{T}{\hat{u}}\right) - s(s+1)\frac{|\nabla_{h+}\hat{u}|^2}{\hat{u}^2}.$$

Using that $\frac{|\nabla_{h_{\pm}}\hat{u}|^2}{\hat{u}^2} \leq \frac{|\nabla_{g_{\pm}}\hat{u}|^2}{\hat{u}^2} \leq 1$, we derive

(4.22)
$$\frac{-\Delta_{h_+}\varphi}{\varphi} \ge s(k+1) - s(s+1) + sH_{\alpha'}\log(u)_{\alpha'} - s\frac{T}{\hat{u}} = s(k-s) + sH^{\alpha'}\log(u)_{\alpha'} - s\frac{T}{\hat{u}}.$$

Now the Cauchy-Schwarz inequality gives $H^{\alpha'}\log(u)_{\alpha'} = H^{\alpha'}\frac{u_{\alpha'}}{\hat{u}} \ge -\alpha \frac{|\nabla_{g_+}u|}{\hat{u}} \ge -\alpha$. Therefore,

(4.23)
$$\frac{-\Delta_{h+}\varphi}{\varphi} \ge s(k-s-\alpha-\beta^Y(u))$$

Observe that $s(k - s - \alpha - \beta^Y(u))$, as a function of s > 0, has a maximum at $s = \frac{k - \alpha - \beta^Y(u)}{2}$. The result now follows from (4.14).

4.3. Proof of Corollary 1.13.

Proof of Corollary 1.13. Notice that for any Lee-eigenfunction u, that is, for any $u \in \mathcal{L}(g_+)$, inequality (1.12) holds. The result follows after taking the supremum over $\mathcal{L}(g_+)$ on both sides.

4.4. Proof of Corollary 1.14.

Proof of Corollary 1.14. Without loss of generality assume $0 \notin Y$. Let $p \in \mathbb{H}^{n+1} \setminus Y$. Define $x \colon \mathbb{H}^{n+1} \to [0, \infty)$ by $x(y) := d_{g_H}(p, y)$ where g_H is the hyperbolic metric. Then (Lemma 3 in [8])

(4.24)
$$\nabla_{g_H}^2 \cosh(x(y)) = \cosh(x(y))g_H,$$

and so $\Delta_{g_H} \cosh(x(y)) = (n+1) \cosh(x(y))$. Let r(y) be a smooth defining function for \mathbb{S}^n that agrees with 1 - |y| outside of some compact set containing 0. Then $\cosh(x(y)) - r^{-1}(y)$ remains bounded as $|y| \to 1$, therefore $u(y) = \cosh(x(y))$ is a Lee-eigenfunction for which $\beta^Y(u) = 0$ thanks to (4.24).

4.5. Stability - Proof of Corollary 1.16.

Proof of Corollary 1.16. Let f be a compactly supported Lipschitz function, and recall from Corollary 1.13 in the minimal case ($\alpha = 0$) that

r

(4.25)
$$\frac{(n-1-\hat{\beta}^Y)^2}{4} \le \lambda_{1,2}(Y) \le \frac{\int_Y |\nabla f|^2 \, dv_{h_+}}{\int_Y f^2 \, dv_{h_+}}.$$

Therefore,

(4.26)
$$\int_{Y} (|\nabla f|^{2} - (|B|^{2} - n)f^{2}) \, dv_{h_{+}} \ge \int_{Y} \left(\frac{(n - 1 - \hat{\beta}^{Y})^{2}}{4} + n - |B|^{2} \right) f^{2} \, dv_{h_{+}} \ge 0,$$

and the result follows. The second part of the statement follows from the fact that when the ambient manifold is \mathbb{H}^{n+1} , then $\hat{\beta}^Y = 0$ for any such submanifold; see proof Corollary 1.14.

Remark 4.2. In Seo's paper in [26], the work of Cheung-Leung is quoted incorrectly. For instance, Cheung-Leung's lower bound estimate does not require stability; see Theorem 2 in [7]. Also, in the statement of Theorem 3.1 in [26], the norm of the second fundamental form should be squared.

4.6. Proof of Corollary 1.17.

Proof. Recall that $\operatorname{Ric}(g_+) = -ng_+$, $b(u) = \nabla_{g_+}^2 u - ug_+$ and $g_u = u^{-2}g_+$. Using the conformal transformation law for the Ricci tensor, we obtain

(4.27)
$$\operatorname{Ric}_{g_u} = g_+ + (n-1)u^{-1}\nabla^2_{g_+}u - nu^{-2}|\nabla_{g_+}u|^2g_+$$
$$= ng_+ + (n-1)u^{-1}b(u) - n|\nabla_{g_+}u|^2g_u.$$

Since $\operatorname{tr}_{g_u}(\operatorname{Ric}_{g_u}) = R_{g_u} = n(n+1)(u^2 - |\nabla_{g_+}^2 u|^2)$ and $E_{g_u} = \operatorname{Ric}_{g_u} - \frac{R_{g_u}}{n+1}g_u$, we get

(4.28)
$$E_{g_u} = ng_+ + (n-1)u^{-1}b(u) - n|\nabla_{g_+}u|^2g_u - ng_+ + n|\nabla_{g_+}u|^2g_u = (n-1)u^{-1}b(u),$$

as desired.

For the second part of the statement, notice that now we can conclude that b(u) = 0 if and only if $E_{g_u} = 0$, thus either condition implies that R_{g_u} is constant. On the other hand, if R_{g_u} is constant, then it is harmonic and using Bochner's formula we conclude that $|b(u)|_{g_+}^2 = 0$. This concludes the proof.

5. Appendix: One-to-one correspondence between $\mathcal{L}(q_+)$ and $[q_+]_{\infty}^{Sc\geq 0}$

We refer to section 4.2 for some of the terminology employed here.

Proposition 5.1. Let $\mathcal{L}(g_+)$ be the set of all Lee-eigenfunctions on a Poincaré-Einstein space (X^{n+1}, g_+) and let $[g_+]_{\infty}^{Sc\geq 0}$ be the subset of $[g_+]_{\infty}$ whose elements have non-negative scalar curvature. Then there is a 1-1 correspondence between $\mathcal{L}(g_+)$ and $[g_+]_{\infty}^{Sc\geq 0}$.

Proof. Define the map $\Omega: [g_+]_{\infty}^{Sc\geq 0} \to \mathcal{L}(g_+)$ as follows. For $\hat{g} \in [g_+]_{\infty}^{Sc\geq 0}$, let r be a defining function such that $(r^2g_+)|_{TM} = \hat{g}$. Set $\Omega(\hat{g}) := u$, where u is the Lee-eigenfunction determined by r. We claim that this is independent of the choice of r consider. Indeed, let ρ be another defining function such that $(\rho^2g_+)|_{TM} = \hat{g} = (r^2g_+)|_{TM}$. Then $\rho = r + O(r^2)$, and so $u - r^{-1} = u - (\rho + O(r^2))^{-1} = u - \rho^{-1} + O(\rho^2)$. Since $u - r^{-1}$ is bounded as $r \to 0$, we conclude that $u - \rho^{-1}$ is also bounded as $\rho \to 0$. It follows by uniqueness that u is the eigenfunction which is induced by ρ , showing the claim.

The map Ω is surjective by construction. We proceed to showing that Ω is injective. To this end, suppose $\Omega(\hat{g}) = \Omega(\hat{g}') = u$, and let r and r_0 be defining functions which correspond to \hat{g} and \hat{g}' , respectively. Then $u - r^{-1}$ and $u - r_0^{-1}$ are both bounded as $r, r_0 \to 0$. Consequently, $(u - r^{-1}) - (u - r_0^{-1})$ is bounded as $r \to 0$, which implies $r^{-1} - r_0^{-1}$ is bounded too as $r \to 0$. We know there exists $c \in \mathbb{R}$ such that $r = cr_0 + O(r_0^2)$. It follows that $c^{-1}r_0^{-1} - r_0^{-1}$ is bounded as $r \to 0$, therefore $(c^{-1} - 1)r_0^{-1}$ is bounded as $r \to 0$. It follows that c = 1 and therefore

(5.1)
$$(r^2g_+)|_{TM} = ([r_0 + O(r_0^2))]^2g_+)|_{TM}$$
$$= (r_0^2g_+)|_{TM}.$$

References

- R. Brooks, A relation between growth and the spectrum of the laplacian, Mathematische Zeitschrift 178 (1981), no. 4, 501–508 (eng).
- [2] M. P. D. Carmo and M. Dajczer, Rotation hypersurfaces in spaces of constant curvature, Springer (2012) (eng).
- [3] J. S. Case, C. R. Graham, T.-M. Kuo, A. J. Tyrrell, and A. Waldron, A gauss-bonnet formula for the renormalized area of minimal submanifolds of poincaré-einstein manifolds, arXiv preprint arXiv:2403.16710 (2024).
- [4] I. Chavel, Eigenvalues in riemannian geometry, Pure and applied mathematics (Academic Press); 115, Academic Press, Orlando, 1984 (eng).
- [5] S. Y. Cheng and S. T. Yau, Differential equations on riemannian manifolds and their geometric applications, Communications on pure and applied mathematics 28 (1975), no. 3, 333–354 (eng).
- [6] S.-Y. Cheng, Eigenvalue comparison theorems and its geometric applications, Mathematische Zeitschrift 143 (1975), no. 3, 289–297 (eng).
- [7] L.-F. Cheung and P.-F. Leung, Eigenvalue estimates for submanifolds with bounded mean curvature in the hyperbolic space, Mathematische Zeitschrift 236 (2001), no. 3, 525–530 (eng).
- [8] J. Choe and R. Gulliver, Isoperimetric inequalities on minimal submanifolds of space forms, Manuscripta mathematica 77 (1992), no. 1, 169–189 (eng).
- [9] F. G. de S. Carvalho and M. P. de A. Cavalcante, On the fundamental tone of the p-laplacian on riemannian manifolds and applications, Journal of mathematical analysis and applications 506 (2022), no. 2, 125703 (eng).
- [10] F. Du and J. Mao, Estimates for the first eigenvalue of the drifting laplace and the p-laplace operators on submanifolds with bounded mean curvature in the hyperbolic space, Journal of mathematical analysis and applications 456 (2017), no. 2, 787–795 (eng).
- [11] H. Fu and Y. Tao, Eigenvalue estimates for complete submanifolds in the hyperbolic spaces, Journal of Mathematical Research with Applications 33 (2013), no. 5, 598–606 (eng).
- [12] C R. Graham, Volume and area renormalizations for conformally compact einstein metrics, arXiv.org (1999) (eng).
- [13] C. R. Graham and J. M Lee, Einstein metrics with prescribed conformal infinity on the ball, Advances in mathematics (New York. 1965) 87 (1991), no. 2, 186–225 (eng).
- [14] O. Hijazi, S. Montiel, and S. Raulot, The cheeger constant of an asymptotically locally hyperbolic manifold and the yamabe type of its conformal infinity, Communications in mathematical physics 374 (2020), no. 2, 873–890 (eng).
- [15] A. LE, Eigenvalue problems for the p-laplacian, Nonlinear analysis 64 (2006), no. 5, 1057–1099 (eng).
- [16] J. M. Lee, The spectrum of an asymptotically hyperbolic einstein manifold, Communications in analysis and geometry 3 (1995), no. 2, 253–271 (eng).
- [17] Z. Li and G. Huang, Upper bounds on the first eigenvalue for the p-laplacian, Mediterranean journal of mathematics 17 (2020), no. 4 (eng).
- [18] B. P. Lima, J. F. B. Montenegro, and N. L. Santos, Eigenvalue estimates for the p-laplace operator on manifolds, Nonlinear analysis 72 (2010), no. 2, 771–781 (eng).
- [19] P. Lindqvist, On the equation $div(|\nabla u|^{p-2}\nabla u) + \lambda |u|^{p-2}u = 0$, Proceedings of the American Mathematical Society 109 (1990), no. 1, 157–164 (eng).
- [20] J. Marx-Kuo, Variations of renormalized volume for minimal submanifolds of poincare-einstein manifolds, arXiv preprint arXiv:2111.04294 (2021).
- [21] R. Mazzeo, Unique continuation at infinity and embedded eigenvalues for asymptotically hyperbolic manifolds, American journal of mathematics **113** (1991), no. 1, 25–45 (eng).
- [22] R. R. Mazzeo, Hodge cohomology of negatively curved manifolds, Ph.D. Thesis, 1986.
- [23] H. P. McKean, An upper bound to the spectrum of Δ on a manifold of negative curvature, Journal of differential geometry **4** (1970), no. 3, 359–366 (eng).
- [24] J. Qing, On the rigidity for conformally compact einstein manifolds, International Mathematics Research Notices 2003 (2003), no. 21, 1141–1153 (eng).
- [25] C. Robin Graham and E. Witten, Conformal anomaly of submanifold observables in ads/cft correspondence, Nuclear physics. B 546 (1999), no. 1, 52–64 (eng).
- [26] K.-K. Seo, Stable minimal hypersurfaces in the hyperbolic space, Journal of the Korean Mathematical Society 48 (2011), no. 2, 253–266 (eng).
- [27] C.-J. A. Sung and J. Wang, Sharp gradient estimate and spectral rigidity for p-laplacian, Mathematical research letters 21 (2014), no. 4, 885–904 (eng).

FINE HALL 101, PRINCETON UNIVERSITY, PRINCETON, NEW JERSEY, NJ 08544, USA *Email address:* sperezayala@princeton.edu

18A DEPARTMENT OF MATHEMATICS AND STATISTICS, TEXAS TECH UNIVERSITY, LUBBOCK, TX 79409, USA *Email address:* aatyrrel@ttu.edu