THE IMMERSION POSET ON PARTITIONS

LISA JOHNSTON, DAVID KENEPP, EVUILYNN NGUYEN, DIGJOY PAUL, ANNE SCHILLING, MARY CLAIRE SIMONE, AND REGINA ZHOU

ABSTRACT. We introduce the immersion poset $(\mathcal{P}(n), \leq_I)$ on partitions, defined by $\lambda \leq_I \mu$ if and only if $s_\mu(x_1, \ldots, x_N) - s_\lambda(x_1, \ldots, x_N)$ is monomial-positive. Relations in the immersion poset determine when irreducible polynomial representations of $GL_N(\mathbb{C})$ form an immersion pair, as defined by Prasad and Raghunathan [PR22]. We develop injections $SSYT(\lambda, \nu) \hookrightarrow SSYT(\mu, \nu)$ on semistandard Young tableaux given constraints on the shape of λ , and present results on immersion relations among hook and two column partitions. The standard immersion poset $(\mathcal{P}(n), \leq_{std})$ is a refinement of the immersion poset, defined by $\lambda \leq_{std} \mu$ if and only if $\lambda \leq_D \mu$ in dominance order and $f^\lambda \leq f^{\mu}$, where f^{ν} is the number of standard Young tableaux of shape ν . We classify maximal elements of certain shapes in the standard immersion poset using the hook length formula. Finally, we prove Schur-positivity of power sum symmetric functions p_{A_μ} on conjectured lower intervals in the immersion poset, addressing questions posed by Sundaram [Sun18].

Contents

1. 1	ntroduction	l
1.1.	Immersion of representations	1
1.2.	Polynomial representation theory of $GL_N(\mathbb{C})$ and symmetric polynomials	2
1.3.	Monomial positivity	2
1.4.	Immersion of Weyl modules: the immersion poset	2
1.5.	Representation theory of symmetric groups	2
1.6.	Schur-positivity	9
1.7.	Power sum symmetric polynomials and restricted row sums of character table	e e
1.8.	Results	9
Ackn	owledgements	4
2. S	tandard immersion poset	4
2.1.	Definition of the standard immersion poset	4
2.2.	Properties of the standard immersion poset	E.
2.3.	Classifying maximal elements	6
3. II	mmersion poset	11
3.1.	Properties of the immersion poset	12
3.2.	Explicit injections	12
3.3.	Immersion poset on hook partitions	22
3.4.	Immersion poset on two column partitions	24
3.5.	Lower intervals and Schur-positivity of interval power sums	27
4. L	Discussion	32
Refer	rences	33

1. INTRODUCTION

1.1. Immersion of representations. Given two finite-dimensional representations $\pi_1: G \to GL(W_1)$ and $\pi_2: G \to GL(W_2)$ of a group G, we say that the representation π_1 is *immersed* in the representation π_2 if the eigenvalues of $\pi_1(g)$, counting multiplicities, are contained in the eigenvalues of $\pi_2(g)$ for all

²⁰²⁰ Mathematics Subject Classification. 06A07, 06A11, 05E05, 05A17, 20C05.

Key words and phrases. immersion poset, monomial-positivity, Schur-positivity, characters of symmetric group.

 $g \in G$. In this case, we call (W_1, W_2) an *immersion pair* denoted by $W_1 \leq W_2$. Note that, if π_1 is a subrepresentation of π_2 , then $W_1 \leq W_2$, but the converse is not true.

Question 1.1 (Prasad and Raghunathan [PR22]). Classify immersion of representations $W_1 \leq_I W_2$ for a given group.

Recently, some progress was made on the above problem for symmetric groups [PPS23] and alternating groups [PPS24]. In this paper, we study immersion pairs for finite-dimensional irreducible polynomial representations of the general linear group $GL_N(\mathbb{C})$.

1.2. Polynomial representation theory of $GL_N(\mathbb{C})$ and symmetric polynomials. The polynomial representation theory of $GL_N(\mathbb{C})$ was developed by Schur [Sch07] and later popularized by Weyl [Wey39] in his expository book on the representation theory of the classical groups. Briefly, the homogeneous irreducible polynomial representations (of degree n) of $GL_N(\mathbb{C})$, also known as Weyl modules $W_\lambda(\mathbb{C}^N)$, are indexed by integer partitions λ (of size n) with at most N non-zero parts. The corresponding irreducible characters, known as Schur polynomials $s_\lambda(x_1, \ldots, x_N)$, are homogeneous symmetric polynomials (of degree n) in N variables x_1, \ldots, x_N .

1.3. Monomial positivity. Given a partition λ of n with at most N parts, the monomial symmetric polynomials is $m_{\lambda}(x_1, \ldots, x_N) \coloneqq \sum_{\alpha} x_1^{\alpha_1} \cdots x_N^{\alpha_N}$, where the sum is over all distinct permutations α of the parts of λ . For example, $m_{(2,1)}(x_1, x_2, x_3) = x_1^2 x_2 + x_1^2 x_3 + x_2^2 x_1 + x_2^2 x_3 + x_3^2 x_1 + x_3^2 x_2$.

The Schur polynomials $\{s_{\lambda} \mid \lambda \vdash n\}$ as well as the monomial symmetric polynomials $\{m_{\lambda} \mid \lambda \vdash n\}$ form a basis for the vector space of symmetric polynomials of degree n. A symmetric polynomial $f(x_1, \ldots, x_N)$ is called *monomial-positive* if

$$f(x_1,\ldots,x_N) = \sum_{\lambda} c_{\lambda} m_{\lambda}(x_1,\ldots,x_N),$$

where the coefficients c_{λ} are non-negative integers.

1.4. Immersion of Weyl modules: the immersion poset. For a partition λ of n with length $\ell(\lambda) \leq N$, let

$$\rho_{\lambda}: GL_N(\mathbb{C}) \longrightarrow GL\left(W_{\lambda}(\mathbb{C}^N)\right)$$

be the irreducible polynomial representation of degree n of highest weight λ . It is a known fact that (for example, see [Sta99, Chapter 7]) if $g \in GL_N(\mathbb{C})$ has the eigenvalues x_1, \ldots, x_N , then the eigenvalues of $\rho_{\lambda}(g)$ are the monomials appearing in the Schur polynomial $s_{\lambda}(x_1, \ldots, x_N)$.

Thus, given two partitions λ, μ of n with $\ell(\lambda), \ell(\mu) \leq N$, the Weyl module $W_{\lambda}(\mathbb{C}^N)$ is immersed in $W_{\mu}(\mathbb{C}^N)$ if and only if $s_{\mu}(x_1, \ldots, x_N) - s_{\lambda}(x_1, \ldots, x_N)$ is monomial-positive. Hence studying the immersion of Weyl modules is equivalent to studying monomial positivity of the difference of Schur polynomials.

Let $\mathcal{P}(n)$ denote the set of integer partitions of n.

Definition 1.2. We define a partial order on $\mathcal{P}(n)$ as follows. For $\lambda, \mu \in \mathcal{P}(n)$, we define $\lambda \leq_I \mu$ if $s_{\mu}(x_1, \ldots, x_N) - s_{\lambda}(x_1, \ldots, x_N)$ is monomial-positive. We call the poset $(\mathcal{P}(n), \leq_I)$ the *immersion poset*.

1.5. Representation theory of symmetric groups. The irreducible representations as well as the conjugacy classes of the symmetric group S_n are indexed by partitions of n. Let $\chi^{\lambda}(\mu)$ denote the character value of the irreducible character χ^{λ} evaluated at an element of cycle type μ . The character table of S_n is a square matrix encoding character values, whose rows are indexed by irreducible characters χ^{λ} and whose columns are indexed by conjugacy classes C_{μ} . The character values of S_n are all integers. Solomon [Sol61] proved that all row sums of the character table of S_n are non-negative integers. Finding a combinatorial interpretation of the row sums is still an open problem (see [Sta99, Exercise 7.71]).

1.6. Schur-positivity. A symmetric polynomial $f(x_1, \ldots, x_N)$ of degree n is called *Schur-positive* if

$$f(x_1,\ldots,x_N) = \sum_{\lambda \vdash n} c_\lambda s_\lambda(x_1,\ldots,x_N),$$

where the coefficients c_{λ} are non-negative integers. Schur-positivity is intimately tied to representation theory. Namely, the symmetric function f is Schur-positive if it is the character of a representation W of $GL_N(\mathbb{C})$ which admits the decomposition into irreducibles $W \cong \bigoplus W_{\lambda}(\mathbb{C}^N)^{\oplus c_{\lambda}}$.

The *Frobenius characteristic map* is a bridge between characters of the symmetric group and symmetric polynomials. The irreducible character χ^{λ} maps to s_{λ} under the Frobenius characteristic map. Via the Frobenius map, Schur-positivity of f implies that there exists a representation V of S_n such that $V \cong \bigoplus_{\lambda \vdash n} V_{\lambda}^{\oplus c_{\lambda}}$, where V_{λ} is the irreducible representation of S_n indexed by λ .

1.7. Power sum symmetric polynomials and restricted row sums of character table. Define the *r*-th power sum symmetric polynomial as

$$p_r(x_1,\ldots,x_N) \coloneqq \sum_{i=1}^N x_i^r.$$

For a partition $\mu = (\mu_1, \mu_2, ...) \vdash n$, define the *power sum symmetric polynomial* as $p_{\mu} \coloneqq p_{\mu_1} p_{\mu_2} \cdots$.

Given a subset A_n of partitions of n, consider the sum of power sum symmetric polynomials

(1.1)
$$p_{A_n} \coloneqq \sum_{\mu \in A_n} p_{\mu}.$$

By the Murnaghan–Nakayama rule [Sta99, Corollary 7.17.4], p_{μ} can be expressed in the basis of Schur polynomials as

$$p_{\mu} = \sum_{\lambda \vdash n} \chi^{\lambda}(\mu) s_{\lambda}.$$

Observe that the coefficient of s_{λ} in the expansion of p_{A_n} is $\sum_{\mu \in A_n} \chi^{\lambda}(\mu)$. This is precisely the restricted row sum (ignoring the columns not in A_n) of the character table of S_n . These values need not always be non-negative integers, that is, p_{A_n} need not be Schur-positive. For example, if $A_4 = \{(1^4), (2, 1, 1), (4)\}$, then one can deduce from the character table of S_4 that $p_{A_4} = 3s_{(4)} + 3s_{(3,1)} + 2s_{(2,2)} + 3s_{(2,1,1)} - s_{(1^4)}$ is not Schur-positive.

Question 1.3 (Sundaram [Sun18]). For which choices of A_n is the symmetric polynomial p_{A_n} Schurpositive? In other words, which subsets A_n of columns in the character table of S_n result in non-negative row sums?

In pursuit of Sundaram's question, we explore the immersion poset in detail, hence understanding the immersion of polynomial representations for $GL_N(\mathbb{C})$. Given a partition μ of n, consider the interval in the immersion poset $[(1^n), \mu] := \{\lambda \mid (1^n) \leq_I \lambda \leq_I \mu\}$. One may ask for what choices of μ , the symmetric polynomial $p_{[(1^n),\mu]}$ defined in Equation (1.1) is Schur-positive. Assuming Conjectures 3.40 and 3.43, we prove that:

- (1) $p_{[(1^n),(n-2,1,1)]}$ is Schur-positive;
- (2) $p_{\lceil (1^n), (n-2,2)\rceil}$ is Schur-positive for $n \neq 7$.

One natural question which arises from the Schur-positivity of the above symmetric functions is to explore the representation theory behind it. It would be interesting to construct a natural representation V of the symmetric group such that its character maps to the symmetric polynomial $p_{[(1^n),\mu]}$ under the Frobenius map, when $\mu = (n-2, 1, 1)$ or (n-2, 2).

1.8. **Results.** In this paper we analyze various properties of the immersion poset. We begin in Section 2 by defining the standard immersion poset. The relation $\lambda \leq_I \mu$ in the immersion poset for $\lambda, \mu \in \mathcal{P}(n)$ holds if the Kostka numbers $K_{\lambda,\alpha} \leq K_{\mu,\alpha}$ for all $\alpha \in \mathcal{P}(n)$. In the standard immersion poset, one only compares the number of standard tableaux of shape λ and μ (instead of semistandard tableaux of all content). Relations in the immersion poset imply relations for the standard immersion poset, but not vice versa. We study properties and maximal elements of the standard immersion. In particular, maximal elements in the standard immersion poset are also maximal elements in the immersion poset. In Section 3, we study properties of the immersion poset. In particular, in Section 3.2 we study relations and covers in the immersion poset using explicit injections between sets of semistandard tableaux. In Section 3.3, we analyze the immersion poset restricted to partitions of hook shape. In Section 3.4, we analyze the immersion relations on partitions with at most two columns. Finally, in Section 3.5 we conjecture the structure of certain lower intervals in the immersion poset and prove that $p_{[(1^n),(n-2,1,1)]}$ and $p_{[(1^n),(n-2,2)]}$ $(n \neq 7)$ are Schur-positive. We conclude in Section 4 with a discussion of open problems.

Acknowledgements

We thank Amritanshu Prasad and Dipendra Prasad for discussions. In particular, we thank Amritanshu Prasad for his probabilistic questions posed in Section 4.

AS was partially supported by NSF grant DMS-2053350. AS thanks IPAM for hospitality at the program "Geometry, Statistical Mechanics, and Integrability" in Spring 2024, where this work was completed.

2. Standard immersion poset

In this section, we introduce the standard immersion poset, which is a refinement of the immersion poset. The definition is given in Section 2.1. Basic properties of the standard immersion poset are proved in Section 2.2. In Section 2.3, the maximal elements of the standard immersion poset are studied. We follow the notational conventions in [Sta99, Chapter 6,7].

2.1. Definition of the standard immersion poset. The Schur polynomial s_{λ} for $\lambda \vdash n$ is defined as (2.1) $s_{\lambda}(x_1, \dots, x_N) = \sum_{\mu \vdash n} K_{\lambda,\mu} m_{\mu}(x_1, \dots, x_N),$

where $K_{\lambda,\mu}$ are the *Kostka numbers* which count the number of semistandard Young tableaux of shape λ and content μ . Note that with this definition the Schur polynomials are zero unless $N \ge \ell(\lambda)$, that is, the number of variables needs to be at least as large as the number of parts in λ .

Lemma 2.1. For $\lambda, \mu \in \mathcal{P}(n)$, $\lambda \leq_I \mu$ if $K_{\lambda,\alpha} \leq K_{\mu,\alpha}$ for all $\alpha \in \mathcal{P}(n)$.

Proof. By Definition 1.2, two partitions $\lambda, \mu \in \mathcal{P}(n)$ are comparable in the immersion poset $\lambda \leq \mu$ if

$$s_{\mu}(x_1,\ldots,x_N) - s_{\lambda}(x_1,\ldots,x_N)$$

is monomial-positive. Using (2.1), this can be restated as saying $\lambda \leq I \mu$ if $K_{\lambda,\alpha} \leq K_{\mu,\alpha}$ for all $\alpha \in \mathcal{P}(n)$. \Box

In particular, Lemma 2.1 implies that a necessary condition for $\lambda \leq_I \mu$ is that $K_{\lambda,(1^n)} \leq K_{\mu,(1^n)}$, which count the standard Young tableaux of shape λ and μ , respectively. Note that $f^{\lambda} \coloneqq K_{\lambda,(1^n)}$ is also the dimension of the Specht module V_{λ} (the irreducible representation of S_n) indexed by λ .

Let $\lambda, \mu \in \mathcal{P}(n)$. Define $\lambda \leq_D \mu$ in *dominance order* on partitions by requiring that

$$\sum_{i=1}^{k} \lambda_i \leq \sum_{i=1}^{k} \mu_i \quad \text{for all } k \geq 1.$$

The Kostka matrix $(K_{\lambda,\alpha})_{\lambda,\alpha\in\mathcal{P}(n)}$ is unit upper-triangular with respect to dominance order, that is, $K_{\lambda,\lambda} = 1$ and $K_{\lambda,\alpha} = 0$ unless $\alpha \leq_D \lambda$. This implies another necessary condition for $\lambda \leq_I \mu$, namely $\lambda \leq_D \mu$. This motivates the definition of the standard immersion poset.

Definition 2.2. On $\mathcal{P}(n)$, define $\lambda \leq_{std} \mu$ if $\lambda \leq_D \mu$ in dominance order and $f^{\lambda} \leq f^{\mu}$. We call this poset the *standard immersion poset*.

As argued above, the standard immersion poset is a refinement of the immersion poset, that is, $\lambda \leq_I \mu$ implies that $\lambda \leq_{std} \mu$. The converse is not always true. For $n \geq 12$, there are examples of $\lambda \leq_{std} \mu$, which do not satisfy $\lambda \leq_I \mu$. For example (5,3,1,1,1,1) covers (4,2,2,2,1,1) in the standard immersion poset for n = 12, but not in the immersion poset.



FIGURE 1. The (standard) immersion poset for n = 8.

Example 2.3. The immersion poset for n = 8 is given in Figure 1. It is equal to the standard immersion poset.

2.2. Properties of the standard immersion poset. We now state and prove properties of the standard immersion poset. Our main tool is the *hook length formula* for $\lambda \in \mathcal{P}(n)$

(2.2)
$$f^{\lambda} = \frac{n!}{\prod_{u \in \lambda} h(u)}$$

where h(u) is the hook length of the cell u in λ which counts the cells weakly to the right of u and strictly below u (in English notation for partitions).

We write $\lambda \leq_{std} \mu$ if μ covers λ in the standard immersion poset. More precisely, $\lambda \leq_{std} \mu$ if $\lambda \leq_{std} \mu$ and there does not exist any ν such that $\lambda \leq_{std} \nu \leq_{std} \mu$.

Lemma 2.4. The partition (1^n) is the unique minimal element in the standard immersion poset.

Proof. The partition (1^n) is the unique minimal element in dominance order. Furthermore, $f^{(1^n)} = 1 \leq f^{\lambda}$ for all $\lambda \in \mathcal{P}(n)$. This proves the claim.

Lemma 2.5. We have

- (1) $(1^n) \leq_{std} (n)$ for all n and
- (2) $(2, 1^{n-2}) \leq_{std} (n-1, 1)$ for all $n \ge 3$.

Proof. We have $(1^n) <_D (n)$ and $f^{(1^n)} = f^{(n)} = 1$. There is no other partition λ with $f^{\lambda} = 1$. This implies $(1^n) \leq_{std} (n)$. Similarly, $(2, 1^{n-2}) <_D (n-1, 1)$ and $f^{(2, 1^{n-2})} = f^{(n-1, 1)} = n-1$. There is no other partition λ with $f^{\lambda} = n - 1$. This implies $(2, 1^{n-2}) \leq_{std} (n - 1, 1)$.

Remark 2.6.

- (1) Let $\lambda <_{std} \mu$. If μ covers λ in dominance order, then μ covers λ with respect to $<_{std}$. The converse is not true. Take $\lambda = (1^n)$ and $\mu = (n)$.
- (2) For a given partition λ with transpose λ^t , if $\lambda <_D \lambda^t$, then $\lambda <_{std} \lambda^t$ as both representations have the same dimension, that is, $f^{\lambda} = f^{\lambda^{t}}$. In general, λ^{t} does not cover λ .

Given a partition λ such that $\lambda <_D \lambda^t$, it would be interesting to find all partitions $\lambda <_D \nu <_D \lambda^t$ satisfying $f^{\lambda} = f^{\nu}$. This would help to understand when the transpose of λ covers λ in the immersion poset.

Lemma 2.7. Let
$$\lambda = (2^a, 1^b)$$
 and $\mu = (2^{a+1}, 1^{b-2})$. Then $\lambda \leq_{std} \mu$ if and only if $\frac{b(b-1)}{2} > a$.

Proof. We have $\lambda \leq_D \mu$. Hence by Remark 2.6(1), it suffices to show that $\lambda \leq_{std} \mu$. By the hook length formula, this is true if $\frac{f^{\lambda}}{f^{\mu}} = \frac{(b+1)(a+1)}{(b-1)(a+b+1)} \leq 1$, which is equivalent to the condition $\frac{b(b-1)}{2} > a$.

2.3. Classifying maximal elements. In this section, we study the maximal elements of the standard immersion poset. Recall that the standard immersion poset is a refinement of the immersion poset. This implies that if a partition is maximal in the standard immersion poset, then it is also maximal in the immersion poset.

Proposition 2.8. The partition (a + b, a) is a maximal element in the standard immersion poset if and only if $\frac{b(b+3)}{2} \ge a$.

Proof. Let $\lambda = (a + b, a)$. Any partition ν which dominates λ , that is, $\nu >_D \lambda$, must have the form $\nu = \nu^{(i)} = (a + b + i, a - i)$ for some $i \ge 1$. Note that $\frac{f^{\nu^{(1)}}}{f^{\lambda}} = \frac{a(b+3)}{(b+1)(a+b+2)}$. Hence $f^{\nu^{(1)}} < f^{\lambda}$ if and only if $\frac{b(b+3)}{a-1} > 2 \text{ (which is equivalent to } \frac{b(b+3)}{2} \ge a). \text{ Thus, the condition is necessary.}$ To prove that the condition $\frac{b(b+3)}{a-1} > 2$ is sufficient, note that

$$\frac{f^{\nu^{(i+1)}}}{f^{\nu^{(i)}}} = \frac{(a-i)(b+3+2i)}{(b+1+2i)(a+b+i+2)} < \frac{a(b+3)}{(b+1)(a+b+2)} = \frac{f^{\nu^{(1)}}}{f^{\lambda}}$$

Since $f^{\nu^{(1)}} < f^{\lambda}$ when $\frac{b(b+3)}{a-1} > 2$, we must have $\frac{f^{\nu^{(i+1)}}}{f^{\nu^{(i)}}} < 1$. This is true for each *i*. Hence λ is a maximal element.

Proposition 2.9. Let $\lambda = (a + b, a, 1)$ where $a \ge 2$. Then λ is maximal in the standard immersion poset if and only if $a \leq \frac{(b+1)(b+2)}{2}$.

Proof. We first prove the reverse direction by inducting on a. For our base case, let a = 2 and $2 \leq \frac{(b+1)(b+2)}{2}$. To prove that $\lambda = (2 + b, 2, 1)$ is maximal, we show that there exists no partition ν such that $\lambda <_D \nu$ and $f^{\lambda} < f^{\nu}$. We start by classifying all partitions ν such that $\lambda <_D \nu$. It is known that $\lambda <_D \nu$ if and only if the Young diagram of ν can be obtained from the Young diagram of λ by moving a single box in row k to row k-1 or by moving a single box in column k to column k+1. This means that the partition

(2+b,2,1) has exactly two covers: (2+b,3) and $(3+b,1^2)$. The former is obtained by moving the box in row 3 to row 2, and the latter is obtained by moving the box at the end of row 2 to row 1. Furthermore, (2+b,3) and $(3+b,1^2)$ are only covered by (3+b,2). Below is the Hasse diagram in dominance order summarizing the specific covering relations:



Let ν be any partition such that $\lambda <_D \nu$. By our covering relations, we have that either $\nu = (2 + b, 3)$ or ν is contained in some chain $\lambda <_D (3 + b, 1^2) <_D \cdots <_D \nu$.

Now, we will show that $f^{\lambda} > f^{\nu}$ for all ν such that $\lambda <_D \nu$.

By Proposition 2.12, we know that $(3 + b, 1^2)$ is maximal in the standard immersion poset. That is, if $(3 + b, 1^2) <_D \nu$ then $f^{(3+b,1^2)} > f^{\nu}$. Note that our assumption that $2 \leq \frac{(b+1)(b+2)}{2}$ implies $b \geq 1$. By this fact and the hook length formula,

$$\frac{f^{(2+b,3)}}{f^{\lambda}} = \frac{b(b+4)}{2(b+1)(b+3)} < 1 \qquad \text{and} \qquad \frac{f^{(3+b,1^2)}}{f^{\lambda}} = \frac{3(b+4)}{2(b+1)(b+5)} < 1$$

Since $f^{\lambda} > f^{(2+b,3)}$ and $f^{\lambda} > f^{(3+b,1^2)} > f^{\nu}$, we have shown that $f^{\lambda} > f^{\nu}$ for all ν such that $\lambda <_D \nu$. Now, let $\lambda = (a+b,a,1)$ where $a \leq \frac{(b+1)(b+2)}{2}$ and suppose that for some $a \geq 2$, the partition (c+b,c,1)

Now, let $\lambda = (a + b, a, 1)$ where $a \leq \frac{(b+1)(b+2)}{2}$ and suppose that for some $a \geq 2$, the partition (c + b, c, 1) is maximal when $c < a \leq \frac{(b+1)(b+2)}{2}$. We follow a similar argument as the base case and show that $f^{\lambda} > f^{\nu}$ for $\lambda \leq_D \nu$. Observe that the Hasse diagram in dominance order around λ looks as follows:



We first consider the partition (a + b, a + 1). Then by the hook length formula,

$$\frac{f^{(a+b,a+1)}}{f^{\lambda}} = \frac{(a+b+2)(b)}{(a+b+1)(a)(b+1)} < \frac{(a+b+2)}{(a+b+1)(a)} < 1$$

where the last inequality follows since $a \ge 2$.

Next, consider the partition (a + b + 1, a - 1, 1). By our inductive hypothesis, (a + b + 1, a - 1, 1) = ((a - 1) + (b + 2), a - 1, 1) is maximal since $a - 1 < a \le \frac{(b+1)(b+2)}{2} < \frac{(b+3)(b+4)}{2}$. Suppose that a is the upper

bound of our inequality $a \leq \frac{(b+1)(b+2)}{2}$, that is, $a = \frac{(b+1)(b+2)}{2}$. Then by the hook length formula,

$$(2.3) \qquad \frac{f^{(a+b+1,a-1,1)}}{f^{\lambda}} = \frac{(a+b+2)(a+1)(a-1)(b+3)}{(a+b+3)(a+b+1)(a)(b+1)} \\ = \frac{((b+1)(b+2)+2b+4)((b+1)(b+2)+2)((b+1)(b+1-2))(2b+6)}{((b+1)(b+2)+2b+6)((b+1)(b+2)+2b+2)((b+1)(b+2))(2b+2)} \\ = \frac{b^7 + 14b^6 + 82b^5 + 260b^4 + 477b^3 + 486b^2 + 216b}{b^7 + 14b^6 + 82b^5 + 260b^4 + 477b^3 + 502b^2 + 280b + 64} < 1.$$

It follows that if $a < \frac{(b+1)(b+2)}{2}$ then $f^{(a+b+1,a-1,1)} < f^{\lambda}$ because for fixed b, Equation (2.3) decreases as a decreases. To see this, we examine the effect of decreasing a on $\frac{a+b+2}{a+b+3}$, $\frac{a+1}{a+b+1}$, and $\frac{a-1}{a}$ individually. Each of these factors is of the form $\frac{x}{x+d}$ for fixed d > 0. Notice that $g(x) = \frac{x}{x+d}$ is a strictly increasing function for x > 0. Therefore, each of the above factors decreases as a decreases. Thus, we have shown that for all ν such that $\lambda <_D \nu$, $f^{\lambda} > f^{(a+b,a+1)}$ and $f^{\lambda} > f^{(a+b+1,a-1,1)} > f^{\nu}$. Hence, (a+b,a,1) is maximal whenever $a \leq \frac{(b+1)(b+2)}{2}$.

Now in the reverse direction, if $a > \frac{(b+1)(b+2)}{2}$, then $f^{(a+b+1,a-1,1)} > f^{\lambda}$. To see this it suffices to consider $a = \frac{(b+1)(b+2)}{2} + 1$ since Equation (2.3) increases as a increases for the same reason as above. If $a = \frac{(b+1)(b+2)}{2} + 1$, then

$$\frac{f^{(a+b+1,a-1,1)}}{f^{\lambda}} = \frac{b^7 + 14b^6 + 88b^5 + 322b^4 + 739b^3 + 1056b^2 + 852b + 288}{b^7 + 14b^6 + 88b^5 + 318b^4 + 707b^3 + 964b^2 + 740b + 240} > 1.$$

Therefore, λ is maximal if only if $a \leq \frac{(b+1)(b+2)}{2}$.

Proposition 2.10. Let $\lambda = (a + b, a, 2)$ where $a \ge 3$. Then λ is maximal in the standard immersion poset if and only if $a \le \frac{(b+1)(b+2)}{2}$.

Proof. We first prove the reverse direction by inducting on a. For our base case, let a = 3 and $3 \leq \frac{(b+1)(b+2)}{2}$. To prove that $\lambda = (3+b,3,2)$ is maximal, we follow a similar argument to Proposition 2.9. We first classify all partitions ν such that $\lambda <_D \nu$ and then show that $f^{\lambda} > f^{\nu}$ for all such ν by finding chains in the dominance order that contain maximal elements from the standard immersion poset. Our assumption that $3 \leq \frac{(b+1)(b+2)}{2}$ implies that $b \geq 1$. Hence it suffices to show that $\lambda = (3+b,3,2)$ is maximal for all $b \geq 1$. We consider the cases b = 1, b = 2, and $b \geq 3$ separately. It can be checked explicitly (for example using SAGEMATH [The24]) that (4,3,2) and (5,3,2) are maximal in the standard immersion poset.

For $b \ge 3$, the Hasse diagram in dominance order around $\lambda = (3 + b, 3, 2)$ looks as follows:



If $\lambda <_D \nu$ then $\nu = (3 + b, 4, 1), (4 + b, 2, 2)$, or ν is contained in some chain $\lambda <_D (3 + b, 4, 1) <_D \nu$. By Proposition 2.9, (3+b, 4, 1) is maximal in the standard immersion poset so it suffices to show that $f^{\lambda} > f^{\nu}$ for $\nu = (3 + b, 4, 1)$ and (4 + b, 2, 2). By the hook length formula,

$$\frac{f^{(3+b,4,1)}}{f^{\lambda}} = \frac{4(b)(b+4)}{5(b+1)(b+3)} = \frac{4(b^2+4b)}{5(b^2+4b+3)} < 1 \qquad \text{and} \qquad \frac{f^{(4+b,2,2)}}{f^{\lambda}} = \frac{4(b+4)}{2(b+1)(b+6)} < 1.$$

Hence, for $b \ge 3$, (3+b,3,1) is maximal in the standard immersion poset, so we have shown that (3+b,3,1)is maximal for all $b \ge 1$.

Now, let $\lambda = (a + b, a, 2)$ where $a \leq \frac{(b+1)(b+2)}{2}$ and suppose that for some $a \geq 3$, the partition (c + b, c, 2)is maximal when $c < a \leq \frac{(b+1)(b+2)}{2}$. Again, we show that $f^{\lambda} > f^{\nu}$ for $\lambda <_D \nu$. Observe that the Hasse diagram in dominance order around λ looks as follows:



By the Hasse diagram, if ν is a partition such that $\lambda <_D \nu$, then $\nu = (a+b,a+1,1), (a+b,a+2), (a+b+2), (a$ 1, a-1, 2, or ν is contained in some chain $\lambda <_D (a+b+1, a-1, 2) <_D \nu$. Observe that (a+b+1, a-1, 2) = 0((a-1) + (b+2), a-1, 2) is maximal by our inductive hypothesis since $a-1 < a \leq \frac{(b+1)(b+2)}{2} \leq \frac{(b+3)(b+4)}{2}$. Therefore, it suffices to check that $f^{\lambda} > f^{\nu}$ for $\nu = (a+b,a+1,1), (a+b,a+2)$, and (a+b+1,a-1,2). For $\nu = (a + b, a + 1, 1)$, we have that

(2.4)
$$\frac{f^{(a+b,a+1,1)}}{f^{\lambda}} = \frac{2b(a+b+1)(a+1)}{(a+b)(a-1)(a+2)(b+1)}$$

Since

$$\frac{d}{da}\frac{f^{(a+b,a+1,1)}}{f^{\lambda}} = -\frac{\left(2b(a^4+2a^3b+4a^3+a^2b^2+5a^2b+7a^2+2ab^2+8ab+2a+3b^2+3b-2)\right)}{\left((a-1)^2(a+2)^2(b+1)(a+b)^2\right)}$$

we have that Equation (2.4) decreases as a increases. Therefore, it suffices to consider a = 3 which we have done in our base case. Hence, $f^{(a+b,a+1,1)} < f^{\lambda}$.

For $\nu = (a + b, a + 2)$, we have that

$$\frac{f^{(a+b,a+2)}}{f^{\lambda}} = \frac{2(b-1)(a+b+2)}{(a-1)(b+1)(a+b)(a+2)} \leqslant \frac{a+b+2}{(a+b)(a+2)}$$

since $a \ge 3$. As $(a+b)(a+2) = a^2 + 2a + ab + 2b \ge a + b + 2$, we have that $f^{(a+b,a+2)} < f^{\lambda}$. Lastly, for $\nu = (a+b+1, a-1, 2)$, we first consider when $a = \frac{(b+1)(b+2)}{2}$. By the hook length formula, we have

$$(2.5) \qquad \frac{f^{(a+b+1,a-1,2)}}{f^{\lambda}} = \frac{(b+3)(a+b+1)(a-2)(a+1)}{(b+1)(a+b+3)(a-1)(a+b)} \\ = \frac{(2b+6)((b+1)(b+2)+2b+2)((b+1)(b+2)-4)((b+1)(b+2)+2)}{((b+1)(b+2)+2b)(2b+2)((b+1)(b+2)+2b+6)((b+1)(b+2)-2)} \\ = \frac{b^7+14b^6+78b^5+220b^4+321b^3+182b^2-80b-96}{b^7+14b^6+78b^5+220b^4+321b^3+214b^2+48b} \\ < 1.$$

Following a similar argument as in Proposition 2.9 for Equation (2.3), we can see that for fixed b, Equation (2.5) decreases as a decreases by considering $\frac{a+b+1}{a+b+3}$, $\frac{a-2}{a-1}$, and $\frac{a+1}{a+b}$

Value for α	$\lambda = (\alpha, \beta)$	$\lambda = (\alpha, \beta, 1)$	$\lambda = (\alpha, \beta, 2)$
$\alpha \geqslant 2$	$(\alpha, 1)$		
$\alpha \geqslant 3$	$(\alpha, 2)$	$(\alpha, 2, 1)$	
$\alpha \geqslant 4$		$(\alpha, 3, 1)$	(lpha,3,2)
$\alpha \geqslant 5$	$(\alpha,3)$		
$\alpha \geqslant 6$	$(\alpha, 4)$	$(\alpha, 4, 1)$	(lpha,4,2)
$\alpha \geqslant 7$	$(\alpha, 5)$	$(\alpha, 5, 1)$	$(\alpha, 5, 2)$
$\alpha \geqslant 8$		$(\alpha, 6, 1)$	(lpha,6,2)
$\alpha \geqslant 9$	$(\alpha, 6)$		
$\alpha \geqslant 10$	$(\alpha,7)$	$(\alpha, 7, 1)$	(lpha,7,2)
$\alpha \geqslant 11$	$(\alpha, 8)$	$(\alpha, 8, 1)$	(lpha, 8, 2)
$\alpha \geqslant 12$	$(\alpha,9)$	(lpha,9,1)	(lpha,9,2)
$\alpha \geqslant 13$		$(\alpha, 10, 1)$	$(\alpha, 10, 2)$
$\alpha \geqslant 14$	$(\alpha, 10)$		
$\alpha \geqslant 15$	$(\alpha, 11)$	$(\alpha, 11, 1)$	$(\alpha, 11, 2)$
$\alpha \geqslant 16$	$(\alpha, 12)$	$(\alpha, 12, 1)$	$(\alpha, 12, 2)$
$\alpha \geqslant 17$	$(\alpha, 13)$	$(\alpha, 13, 1)$	$(\alpha, 13, 2)$
$\alpha \geqslant 18$	$(\alpha, 14)$	$(\alpha, 14, 1)$	$(\alpha, 14, 2)$
$\alpha \geqslant 19$		$(\alpha, 15, 1)$	$(\alpha, 15, 2)$
$\alpha \geqslant 20$	$(\alpha, 15)$		
÷	÷	÷	:

TABLE 1. Necessary and sufficient conditions for maximality of a partition λ .

We have thus shown that when $a \ge 3$, $\lambda = (a + b, a, 2)$ is maximal if $a \le \frac{(b+1)(b+2)}{2}$. For the reverse direction, consider Equation (2.5) when $a = \frac{(b+1)(b+2)}{2} + 1$. We have that

$$\frac{f^{(a+b+1,a-1,2)}}{f^{\lambda}} = \frac{b^6 + 12b^5 + 60b^4 + 162b^3 + 243b^2 + 162b}{b^6 + 12b^5 + 60b^4 + 158b^3 + 219b^2 + 150b + 40} > 1.$$

Since Equation (2.5) increases as a increases, $\lambda \leq_{std} (a+b+1, a-1, 2)$ when $a > \frac{(b+1)(b+2)}{2}$. Therefore, λ is maximal if and only if $a \leq \frac{(b+1)(b+2)}{2}$.

Remark 2.11. We may translate the results of Propositions 2.8, 2.9, and 2.10 into statements about partitions of the form (α, β) , $(\alpha, \beta, 1)$, and $(\alpha, \beta, 2)$. Table 1 summarizes our maximality conditions.

We next classify all maximal hook shape partitions. As noted in Lemma 2.5, $(1^n) <_{std} (n)$ and so the single column shape is only maximal when n = 1. By Lemma 2.7, $(2, 1^b) <_{std} (2^2, 1^{b-2})$ whenever $b \ge 3$. Since $(2, 1, 1) <_{std} (3, 1)$, the only maximal hook shape with arm length 2 is (2, 1). In the following proposition, we investigate all hook shape partitions with arm length greater than 2.

Proposition 2.12. Let $\lambda = (a, 1^b)$ be a hook shape partition such that a > 2. Then λ is a maximal element in the standard immersion poset if and only if $b \leq 2$.

Proof. When b = 1, the only partition that dominates (a, 1) is (a + 1) and $f^{(a,1)} = (a + 1) - 1 > 1 = f^{(a+1)}$. Thus, (a, 1) is maximal. When b = 2, the only partitions that dominate $(a, 1^2)$ are (a + 2), (a + 1, 1), and (a,2). By the hook length formula,

$$\frac{f^{(a+1,1)}}{f^{(a,1^2)}} = \frac{2}{a} < 1 \qquad \text{and} \qquad \frac{f^{(a,2)}}{f^{(a,1^2)}} = \frac{(a+2)(a-1)}{(a+1)a} = \frac{a^2+a-2}{a^2+a} < 1.$$

Therefore, no partition dominates $(a, 1^2)$ and has more standard Young tableaux, so $(a, 1^2)$ is maximal. When $b \ge 3$, $(a, 1^b) <_{std} (a, 2, 1^{b-2})$, by the hook length formula:

$$\frac{f^{(a,1^b)}}{f^{(a,2,1^{b-2})}} = \frac{a(a+b-1)}{(a+b)(a-1)(b-1)} \leqslant 1$$

since

$$(a+b)(a-1)(b-1) \ge 2(a-1)(a+b) \ge a(a+b-1).$$

Therefore $f^{(a,1^b)} \leq f^{(a,2,1^{b-2})}$ and $(a,1^b) <_D (a,2,1^{b-2})$, so we have $(a,1^b) <_{std} (a,2,1^{b-2})$ whenever $b \geq 3$.

Proposition 2.13. If λ is a maximal element in the standard immersion poset, then $\lambda_1 > \lambda_2$.

Proof. Suppose by contradiction that $\lambda = (a^b, \lambda_{b+1}, ...)$ with $a > \lambda_{b+1}$ and $b \ge 2$. Let $\mu = (a + 1, a^{b-2}, a - 1, \lambda_{b+1}, ...)$ and denote by $\mathsf{SYT}(\lambda)$ the set of all standard Young tableaux of shape λ . The map

$$\varphi$$
:SYT $(\lambda) \rightarrow$ SYT (μ) ,

where $\varphi(T)$ is the standard Young tableau obtained from T by moving the box in position (b,a) to position (1, a + 1), is an injection. Therefore, $\lambda <_D \mu$ and $f^{\lambda} \leq f^{\mu}$, which implies $\lambda \leq_I \mu$ and thus demonstrates that λ is not a maximal element in the standard immersion poset.

In fact, the injection φ used in the proof of Proposition 2.13 remains an injection when the domain and codomain are extended to semistandard Young tableaux of content ν , for any $\nu \vdash |\lambda|$. Injection arguments between sets of semistandard Young tableaux are expanded on in Section 3.2. In particular, this result is extended to the immersion poset in Corollary 3.7.

We conclude this section with a conjecture about more general maximal elements in the standard immersion poset.

Conjecture 2.14. Suppose $\lambda = (\sum_{i=1}^{\ell} a_i, \sum_{i=1}^{\ell-1} a_i, \dots, a_2 + a_1, a_1)$ for $\ell > 2$. If

$$\binom{a_j+2}{2} \geqslant \sum_{i=1}^{j-1} a_i + j - 2$$

is satisfied for all $2 \leq j \leq \ell$, then λ is maximal in the standard immersion poset.

This conjecture has been verified with SAGEMATH [The24] for $|\lambda| \leq 30$.

Remark 2.15. Proposition 2.8 addresses the case $\ell = 2$ associated to Conjecture 2.14. Note that for $\ell = 2$ the condition stated in Conjecture 2.14 reads

$$\binom{a_2+2}{2} \ge a_1$$
, whereas the condition from Proposition 2.8 is $\binom{a_2+2}{2} \ge a_1$.

This discrepancy comes from the fact that for $\ell > 2$, there are more factors contributing to the inequality in $\frac{f^{\mu}}{f^{\lambda}} < 1$.

3. Immersion poset

In this section we turn to the immersion poset. In Section 3.1, we study basic properties of the immersion poset. In Section 3.2, we provide explicit injections between certain sets of semistandard Young tableaux, which are used to determine statements about maximal elements and cover relations in the immersion poset. In Sections 3.3 and 3.4, we study the immersion poset restricted to hook partitions and two column partitions, respectively. We conclude in Section 3.5 with conjectures about certain lower intervals in the immersion poset and prove that the conjectured intervals give Schur-positive sums of power sum symmetric functions.

3.1. Properties of the immersion poset. We begin by specifying the minimal element.

Lemma 3.1. The partition (1^n) is the unique minimal element in the immersion poset $(\mathcal{P}(n), \leq_I)$.

Proof. We have $f^{(1^n)} = 1 \leq f^{\lambda}$ for all $\lambda \in \mathcal{P}(n)$. Furthermore $K_{(1^n),\alpha} = 0 \leq K_{\lambda,\alpha}$ for all $\alpha \neq (1^n)$ and $\lambda \in \mathcal{P}(n)$. By Lemma 2.1 this proves the claim.

Analogously to Lemma 2.5, we prove the following result.

Lemma 3.2. We have

- (1) $(1^n) \leq_I (n)$ for all n and
- (2) $(2, 1^{n-2}) \leq_I (n-1, 1)$ for all $n \geq 3$.

Proof. By Lemma 3.1, we have $(1^n) <_I (n)$. By Lemma 2.5, $(1^n) <_{std} (n)$. Since in the immersion poset there are fewer order relations than in the standard immersion poset, the first part of the lemma follows. We have $(2, 1^{n-2}) <_I (n-1, 1)$ since

$$s_{(2,1^{n-2})} = (n-1)m_{(1^n)} + m_{(2,1^{n-2})} \text{ and } s_{(n-1,1)} = (n-1)m_{(1^n)} + (n-2)m_{(2,1^{n-2})} + \sum_{\mu \neq (1^n), (2,1^{n-2})} K_{(n-1,1),\mu}m_{\mu}.$$

Again, since by Lemma 2.5 we have $(2, 1^{n-2}) \leq_{std} (n-1, 1)$, the second part of the lemma follows.

Unlike in the standard immersion poset, where λ and λ^t are always comparable as long as they are comparable in dominance order (see Remark 2.6), this is not always true in the immersion poset. For example $\lambda = (4, 4, 2, 1, 1)$ and λ^t are not comparable in the immersion poset since $K_{(4,4,2,1,1),(4,4,1,1,1,1)} > K_{(5,3,2,2),(4,4,1,1,1,1)}$. For hook partitions, it is however true that $\lambda <_I \lambda^t$ if $\lambda <_D \lambda^t$ (see Corollary 3.30).

We prove the analog of Lemma 2.7 in the next section using injections on semistandard Young tableaux. See Corollaries 3.6, 3.13, and 3.21.

3.2. Explicit injections. Recall from Lemma 2.1 that $\lambda \leq_I \mu$ if and only if $K_{\lambda,\nu} \leq K_{\mu,\nu}$ for all $\nu \in \mathcal{P}(n)$. The Kostka number $K_{\lambda,\nu}$ is the cardinality of the set of semistandard Young tableaux SSYT (λ,ν) of shape λ and content ν . Hence we can analyze the order relations $\lambda \leq_I \mu$ by constructing explicit injections

(3.1)
$$\varphi: \mathsf{SSYT}(\lambda, \nu) \to \mathsf{SSYT}(\mu, \nu)$$

for all $\nu \in \mathcal{P}(n)$.

To this end, we present one such injection, where μ differs from λ by moving a single cell from the *c*-th column to the (c+1)-th column, and λ has a bound on the relative size of the two columns. Upon establishing this first injection, we refine it to obtain more precise bounds on the relative size of the columns. We partially characterize what elements cannot be maximal in the immersion poset, similar to those given in Section 2.3 for the standard immersion poset.

Let

(3.2)
$$\lambda = (\lambda_1, \dots, \lambda_{\alpha}, c^{\beta}, \lambda_{\alpha+\beta+1}, \dots),$$
$$\mu = (\lambda_1, \dots, \lambda_{\alpha}, c+1, c^{\beta-2}, c-1, \lambda_{\alpha+\beta+1}, \dots).$$

such that either $\alpha > 0$ and $\lambda_{\beta+\alpha+1} < c < \lambda_{\alpha}$, or $\alpha = 0$ and $\lambda_{\beta+\alpha+1} < c$. In particular, $\lambda_{\beta+\alpha+1}$ can be 0. We define a map

$$\varphi_0$$
:SSYT $(\lambda, \nu) \rightarrow$ YT $(\mu, \nu),$

where $\mathsf{YT}(\mu,\nu)$ is the set of all tableaux of shape μ and content ν , not necessarily semistandard. We will show in Proposition 3.5 that when $\beta \ge \alpha + 2$, the image of φ_0 will be contained in $\mathsf{SSYT}(\mu,\nu)$, so φ_0 will be as in (3.1).

For $T \in SSYT(\lambda, \nu)$, we define $\varphi_0(T)$ as follows. Suppose the entries in the *c*-th column of *T* in increasing order are $x_{\beta+\alpha}, x_{\beta+\alpha-1}, \ldots, x_1$ and the entries in the (c+1)-th column of *T* in increasing order are $y_{\alpha}, y_{\alpha-1}, \ldots, y_1$. Let *i* be the smallest index such that $x_i > y_i$. If no such index exists, let $i = \alpha + 1$. Then $\varphi_0(T)$ is the tableau such that the entries in the *c*-th column of $\varphi_0(T)$ are

$$x_{\beta+\alpha}, x_{\beta+\alpha-1}, \ldots, x_{i+1}, y_{i-1}, y_{i-2}, \ldots, y_1,$$

the entries in the (c+1)-th column of $\varphi_0(T)$ are

(3.3)

$$y_{\alpha}, y_{\alpha-1}, \ldots, y_i, x_i, x_{i-1}, \ldots, x_1,$$

and all other entries are the same as those in T. In other words, φ_0 moves the cell containing x_1 to the $(\alpha + 1)$ -th row of the (c + 1)-th column, and swaps each x_j with y_{j-1} for all $2 \leq j \leq i$.

More concretely, the c-th and (c+1)-th column in T and $\varphi_0(T)$ look as follows:



The cells marked in green contain the entries that move from the c-th column to the (c + 1)-th column, and the cells marked in yellow are the entries that move from (c + 1)-th column to the c-th column. We continue to use this convention for all subsequent examples of φ_0 .

Remark 3.3. Observe that by our choice of *i*, both $x_{i+1} < x_{i-1} \leq y_{i-1}$ and $y_i < x_i$, so the columns of $\varphi_0(T)$ are strictly increasing by construction.

Example 3.4. For $\lambda = (3, 2, 1^4)$ and $\mu = (3, 2, 2, 1^2, 0)$, we have c = 1, $\alpha = 2$ and $\beta = 4$. Here are some examples of the injection φ_0 on various tableaux of shape λ :



Proposition 3.5. Let λ and μ be as in (3.2) with $\beta \ge \alpha + 2$. Then φ_0 as defined above is an injection

 φ_0 :SSYT $(\lambda, \nu) \rightarrow$ SSYT (μ, ν) .

Proof. Let $T \in SSYT(\lambda, \nu)$. Note that the content does not change under φ_0 . We need to check that the *c*-th and (c + 1)-th columns of $\varphi_0(T)$ are strictly increasing, and that the $(\alpha - i + 2)$ -th through $(\alpha + \beta - 1)$ -th rows of $\varphi_0(T)$ are weakly increasing, since all other entries are identical to those in T. (It may be helpful to consult (3.3).) The columns are strictly increasing by Remark 3.3.

For rows, we first consider the $(\alpha - i + 2)$ -th through $(\alpha + 1)$ -th rows. Due to the bound $\beta \ge \alpha + 2$, in the *c*-th column, these rows contain $x_{\beta+i-1}, \ldots, x_{\beta}$. In the (c+1)-th column, irrespective of the bound on α and β , these rows contain x_i, \ldots, x_1 . In particular, the bound $\beta \ge \alpha + 2$ makes it so that there is no y_j entry in these rows, so there is no "overlap" of y_j and x_k for $1 \le k \le i$. The rows are thus strictly increasing because $x_j > x_k$ for all j < k, so the x_j entries in the (c+1)-th column are greater than the entries to their left in the *c*-th column; and $x_j < x_{j-1} \le y_{j-1}$ for $2 \le j \le i$, so the x_j entries in the (c+1)-th column are less than any entries to their right, originally from *T*. (Such entries on the right do not necessarily exist. In particular, x_1 never has any cell to its right.) Now consider the $(\alpha + 2)$ -th through $(\alpha + \beta - 1)$ -th rows. In the *c*-th column, these rows contain

$$x_{\beta-1}, \ldots, x_{i+1}, y_{i-1}, \ldots, y_1,$$

and in the (c+1)-th column, these rows contain no cells. They are weakly increasing because $y_{j-1} \ge x_{j-1} > x_j$ for $2 \le j \le i$, so the y_{j-1} entries in the *c*-th column are greater than the entries to their left, originally from *T*, and they have no cells to their right.

To show injectivity, we define an explicit inverse ψ_0 . Let $T' \in \varphi_0(\mathsf{SSYT}(\lambda,\nu))$. Suppose the entries in the *c*-th column of T' in increasing order are $x'_{\beta+\alpha-1}, x'_{\beta+\alpha-2}, \ldots, x'_1$, and the entries in the (c+1)-th column of T' in increasing order are $y'_{\alpha+1}, y'_{\alpha}, \ldots, y'_1$. Let *i'* be the smallest index such that $y'_{i'} > x'_{i'}$. This *i'* will be equal to the *i* from the definition of φ_0 , because $x_i > x_{i+1}$ and $x_j \leq y_j$ for $1 \leq j \leq i-1$ in T.

Then $\psi_0(T')$ is the tableau of shape λ such that the entries in the *c*-th column of $\psi_0(T')$ are

$$x'_{\beta+\alpha-1}, x'_{\beta+\alpha-2}, \dots, x'_{\beta-1}, x'_{\beta-2}, x'_{\beta-3}, \dots, x'_{i'}, y'_{i'}, y'_{i'-1}, \dots, y'_{1},$$

the entries in the (c+1)-th column of $\psi_0(T')$ are

$$y'_{\alpha+1}, y'_{\alpha}, \dots, y'_{i'+1}, x'_{i'-1}, x'_{i'-2}, \dots, x'_1,$$

and all other entries are the same as those in T'. In other words, ψ_0 moves the cell containing y'_1 to the $(\alpha + \beta)$ -th position in the *c*-th column, and swaps each $y'_{j'}$ with $x'_{j'-1}$ for all $2 \leq j' \leq i'$. Concretely:

(3.4)



Since i' = i, ψ_0 moves back exactly the entries in T' that were originally moved by φ_0 in T, so ψ_0 is the inverse of φ_0 .

As a corollary, the injection describes a class of cover relations in the immersion poset. As a specific example, it can partially address the two column case, which was completely addressed by Lemma 2.7 for the standard immersion poset.

Corollary 3.6. The partitions λ and μ as in (3.2) with $\beta \ge \alpha + 2$ form a cover in the immersion poset. In particular, $\lambda = (2^{\alpha}, 1^{\beta})$ and $\mu = (2^{\alpha+1}, 1^{\beta-2})$ form a cover.

Proof. The partition μ covers λ in dominance order, and the injection shows that μ is greater than λ in the immersion poset, so μ must also cover λ in the immersion poset.

The injection also gives a few conditions on which partitions cannot be maximal.

Corollary 3.7. If $\lambda = (a^{\beta}, b, ...)$, where a > b, and $\beta \ge 2$, then λ is not maximal.

Proof. We have $\beta \ge 2$ with $\alpha = 0$, so we can apply the injection.

Corollary 3.8. If $\lambda = (a, b^{\beta}, c, ...)$, where a > b > c, and $\beta \ge 3$, then λ is not maximal. In particular, $\lambda = (a, 1^{\beta})$ is not maximal for $a \ge 2$, $\beta \ge 3$.

Proof. We have $\beta \ge 3$ with $\alpha = 1$, so we can apply the injection.

Note that Corollary 3.7 and Corollary 3.8 repeat the results from Proposition 2.13 and the forward direction of Proposition 2.12 concerning nonmaximal elements in the standard immersion poset.

Corollary 3.9. If $\lambda = (a, b, c, d)$ is maximal in the immersion poset, then it has no more than two identical non-zero parts.

Proof. If λ has three or more identical parts, then λ is one of (a^4) , (a^3, d) , or (a, b^3) , so we can apply the injection.

As stated in the proof of Proposition 3.5, the bound $\beta \ge \alpha + 2$ is necessary for $\varphi_0(T)$ to be semistandard for T semistandard. When $\beta < \alpha + 2$, φ_0 can cause an "overlapping" row, where for certain $1 \le j \le \alpha - \beta + 2$, $y_{\beta-2+j}$ is to the left of x_j , yet $y_{\beta-2+j} > x_j$.

Example 3.10. For $\lambda = (2^2, 1^3)$, so $\alpha = 2$ and $\beta = 3 = \alpha + 1$, φ_0 can give:

1	6	\rightarrow	1	3
2	7		2	4
3			6	5
4			7	
5				

One natural modification to restore weakly increasing rows is to swap $y_{\beta-2+j}$ and x_j whenever the problem occurs. Unfortunately, doing so on its own would not maintain injectivity. If we try to swap the 5 and 6 in the previous example, our final tableau is the same as the following tableau obtained from φ_0 with no switches:

1	5	₽	1	3
2	7		2	4
3			5	6
4			7	
6				•

However, if we are able to implement subsequent modifications in a way such that the resulting tableau is semistandard, yet cannot be obtained from φ_0 alone, then we can restore injectivity.

We now define the modification of our original φ_0 injection for the case when $\beta = \alpha + 1$, and $\alpha \ge 2$, which we call

$$\varphi_1$$
:SSYT $(\lambda) \rightarrow$ SSYT (μ) .

From now on, we drop the content ν as all maps in this subsection preserve the content.

Let $T \in \mathsf{SSYT}(\lambda)$. As before, suppose that the entries in the *c*-th column of *T* in increasing order are $x_{\beta+\alpha}, x_{\beta+\alpha-1}, \ldots, x_1$, and the entries in the (c+1)-th column of *T* in increasing order are $y_{\alpha}, y_{\alpha-1}, \ldots, y_1$. We define $\varphi_1(T)$ to be the same as $\varphi_0(T)$ if $\varphi_0(T) \in \mathsf{SSYT}(\mu)$.

If $\varphi_0(T) \notin \mathsf{SSYT}(\mu)$, then necessarily $i = \alpha + 1$ as defined for φ_0 and x_1 is to the right of $y_{\beta-1} = y_\alpha$, with $y_\alpha > x_1$. Then $\varphi_1(T)$ is the same as $\varphi_0(T)$, except we swap y_α with x_1 , as well as $x_{\beta+1}$ with x_β .

Concretely, when $\varphi_1(T) \neq \varphi_0(T)$, the *c*-th and (c+1)-th columns of T, $\varphi_0(T)$, and $\varphi_1(T)$ look as follows:



We indicate the additional swaps φ_1 adds to φ_0 using boldface on the relevant entries, $x_{\beta+1}$ and x_{β} . We will continue to use this convention for any subsequent modifications to φ_0 .

Observe that for the $x_{\beta+1}$ with x_{β} swap to be between cells in different rows, we must have $\alpha \ge 2$. This property is necessary for the tableau to remain semistandard after the swap.

The intuition behind the swaps is that the swap of y_{α} with x_1 makes the tableau semistandard, and the swap of $x_{\beta+1}$ with x_{β} prevents the new tableau from being in the image of φ_0 .

Example 3.11. For T in Example 3.10, φ_1 maps:





Proposition 3.12. Let λ and μ be as in (3.2) with $\beta = \alpha + 1 \ge 3$. Then φ_1 as defined above is an injection $\varphi_1: SSYT(\lambda) \to SSYT(\mu).$

Proof. Let $T \in \mathsf{SSYT}(\lambda)$. We need to check that $\varphi_1(T)$ is semistandard. It suffices to do so for the case when $\varphi_1(T) \neq \varphi_0(T)$, where there is an overlap in $\varphi_0(T)$ consisting of a single decreasing pair of cells in a row, $y_{\alpha} > x_1$. In particular, $\varphi_0(T)$ would be semistandard if it were not for this single pair by the proof of Proposition 3.5, so it suffices to check that swapping the x_1 with y_{α} makes the tableau semistandard, and swapping the $x_{\beta+1}$ with x_{β} keeps it semistandard, by examining the changed entries.

Swapping the entries x_1 with y_{α} makes the $(\alpha+1)$ -th row weakly increasing since $x_1 < y_{\alpha}$ by assumption. The *c*-th column remains strictly increasing since $x_{\beta+1} < x_1 < y_{\alpha} < y_{\alpha-1}$, and the (c+1)-th column remains strictly increasing since $x_2 < x_1 < y_{\alpha}$.

Swapping the entries x_{β} with $x_{\beta+1}$ keeps the relevant rows, namely the 1st row and α -th row, weakly increasing and their columns strictly increasing since $x_j > x_k$ for all j < k.

Specifically, the c-th column remains strictly increasing since $x_{\beta+2} < x_{\beta} < x_1$, and (c+1)-th column remains strictly increasing since $x_{\beta+1} < x_{\beta-1}$. The 1st row remains weakly increasing because $x_{\beta+1} < x_{\beta}$, so $x_{\beta+1}$ is also less than all entries to its right, which are originally right of x_{β} . The α -th row remains weakly increasing because $x_{\beta+1} < x_{\beta}$, so x_{β} is greater than all entries to its left, which are originally left of $x_{\beta+1}$.

To show injectivity, it suffices to check that the modified tableau cannot be in the image of φ_0 , allowing us to define an explicit inverse ψ_1 by likewise modifying ψ_0 . Namely, we must verify that $\varphi_1(T)$ is not equal to $\varphi_0(S)$ for any $S \in SSYT(\lambda)$.

Indeed, consider $\varphi_0(S)$ for any $S \in \mathsf{SSYT}(\lambda)$. Let *i* be as in the definition of φ_0 . Then the *i*-th entry from the bottom of the (c+1)-th column in $\varphi_0(S)$ must have originally been below and hence greater than the *i*-th entry from the bottom of the *c*-th column in $\varphi_0(S)$, which stays in the same place in $\varphi_0(S)$.

That is, if the *j*-th entry from the bottom of the (c + 1)-th column in $\varphi_1(T)$ is less than or equal to the *j*-th entry from the bottom of the *c*-th column in $\varphi_1(T)$ for all $1 \leq j \leq \alpha + 1$, then $\varphi_1(T) \neq \varphi_0(S)$ for all $S \in SSYT(\lambda)$. If we check these corresponding pairs of entries in $\varphi_1(T)$, we have $y_{\alpha} < y_1, x_j < y_j$ for $2 \leq j \leq \alpha - 1, x_{\beta-1} < x_1$, and $x_{\beta+1} < x_\beta$. Thus, we do not have a requisite pair of entries, and no *S* satisfies $\varphi_1(T) = \varphi_0(S)$.

We can now define our explicit inverse ψ_1 . Let $T' \in \varphi_1(\mathsf{SSYT}(\lambda))$. As before, the entries in the *c*-th column of T' in increasing order are $x'_{\beta+\alpha-1}, x'_{\beta+\alpha-2}, \ldots, x'_1$, and the entries in the (c+1)-th column of T' in increasing order are $y'_{\alpha+1}, y'_{\alpha}, \ldots, y'_1$. Let $\psi_1(T') = \psi_0(T')$ when $T' \in \varphi_0(\mathsf{SSYT}(\lambda))$, the domain of ψ_0 . This occurs when there exists an i'

Let $\psi_1(T') = \psi_0(T')$ when $T' \in \varphi_0(SSYT(\lambda))$, the domain of ψ_0 . This occurs when there exists an i' such that $y'_{i'} > x'_{i'}$, so we can take the smallest such i' as in the definition of ψ_0 .

If such an i' does not exist, then ψ_1 first swaps x'_{β} with $y'_{\alpha+1}$, and $x'_{\beta-1}$ with y'_1 , undoing the modifications. Relabelling the new tableau obtained after these swaps T'', we now let $\psi_1(T') = \psi_0(T'')$.

Concretely, when ψ_1 differs from ψ_0 , the *c*-th and (c+1)-th columns look as follows:

(3.7)								
T':	$x'_{\beta+\alpha-1}$	y'_{lpha+1}	T'':	$x'_{\beta+\alpha-1}$	x'_eta	$\psi_1(T'):$	$x'_{\beta+\alpha-1}$	y'_1
	$x'_{\beta+\alpha-2}$	y'_{lpha}		$x'_{\beta+\alpha-2}$	y'_{lpha}		$x'_{\beta+\alpha-2}$	x'_{eta-2}
	:	:		•••	:		:	÷
	x'_eta	y_2'		y'_{lpha+1}	y_2'		$y'_{\alpha+1}$	x'_1
	$x'_{\beta-1}$	y_1'		y_1'	x'_{eta-1}		x'_{eta}	
	x'_{eta-2}			x'_{eta-2}			y'_{lpha}	
	÷						÷	
	x'_1			x'_1			y_2'	
		-			-		r'	

It is straightforward to check that for $T' = \varphi_1(T)$, ψ_1 exactly reverses all the swaps done by φ_1 .

We now obtain stronger versions of the corollaries obtained from the previous injection, in particular Corollary 3.6.

Corollary 3.13. The partitions λ and μ as in (3.2) with $\beta \ge \alpha + 1 \ge 3$ form a cover in the immersion poset.

Corollary 3.14. If $\lambda = (a^2, b^\beta, c, ...)$, where a > b > c, and $\beta \ge 3$, then λ is not maximal.

Corollary 3.15. If $\lambda = (a, b, c, d, e)$ is maximal in the immersion poset, then it has no more than two identical non-zero parts.

In order to further improve the bound for the injection, we must continue to apply modifications to resolve decreasing pairs in "overlapping" rows, and then apply further modifications to establish injectivity. However, there are now multiple cases to consider.

Firstly, any combination of the overlapping rows containing both $y_{\beta-2+j}$ and x_j can be decreasing.

Example 3.16. For $\lambda = (2^4, 1^4)$, so $\alpha = \beta = 4$, following φ_0 can give two overlapping rows. We have each possible combination of rows with decreasing pairs as follows:

(3.8)	1	7	↦	1	4	1	8	\rightarrow	1	4	1	9	\rightarrow	1	4
()	2	10		2	5	2	9		2	5	2	10		2	5
	3	11		3	6	3	11		3	6	3	11		3	6
	4	12		7	8	4	12		8	7	4	12		9	7
	5			10	9	5			9	10	5			10	8
	6			11		6			11		6			11	
	8			12		7			12		7			12	
	9					10					8				

While all these previous tableaux give 2 rows of overlap, it is also possible for a tableau of the same shape to give 0 or 1 rows of overlap instead. More generally, $\varphi_0(T)$ for T of shape λ as in (3.2) can have anywhere between 0 and max $\{0, \alpha - \beta + 2\}$ rows of overlap.

Example 3.17. For the same $\lambda = (2^4, 1^4)$ as in Example 3.16, φ_0 can give a single overlapping row, which contains a decreasing pair:

(3.9)

1	5	\rightarrow	1	5
2	10		2	6
3	11		3	7
4	12		4	8
6			10	9
7			11	
8			12	
9				

Hence, in our next modifications of φ_0 , we must encode the information of every possible case in a way that both distinguishes the cases from φ_0 with no modifications, and distinguishes the cases from each other. To achieve this, our modifications will involve cyclically rotating certain entries in the *c*-th and (c+1)-th columns. These rotations will be analogous to the x_β with $x_{\beta+1}$ swap in φ_1 , which can be thought of as a rotation of 2 elements.

We now define a second set of modifications of our original φ_0 injection for the case when $\beta = \alpha$, and $\alpha \ge 4$, which we call

$$\varphi_2$$
:SSYT $(\lambda) \rightarrow$ SSYT (μ) .

Let $T \in \mathsf{SSYT}(\lambda)$. As before, suppose that the entries in the *c*-th column of *T* in increasing order are $x_{\beta+\alpha}, x_{\beta+\alpha-1}, \ldots, x_1$, and the entries in the (c+1)-th column of *T* in increasing order are $y_{\alpha}, y_{\alpha-1}, \ldots, y_1$. We define $\varphi_2(T)$ to be the same as $\varphi_0(T)$ if $\varphi_0(T) \in \mathsf{SSYT}(\mu)$. If $\varphi_0(T) \notin SSYT(\mu)$, then we have several cases. If there are two rows of overlap, then *i* as defined for φ_0 is $\alpha + 1$, x_1 is to the right of $y_{\alpha-1}$, and x_2 is to the right of $y_{\alpha} = y_{\beta}$:

(3.10)	T:	$x_{\beta+\alpha}$	y_{lpha}	$\varphi_0(T)$:	$x_{\beta+\alpha}$	$x_{\beta+1}$
		$x_{\beta+\alpha-1}$	y_{lpha-1}	10()	$x_{\beta+\alpha-1}$	x_eta
		$x_{\beta+\alpha-2}$	y_{lpha-2}		$x_{\beta+\alpha-2}$	$x_{\beta-1}$
		:	:		:	÷
		$x_{\beta+3}$	y_3		$x_{\beta+3}$	x_4
		$x_{\beta+2}$	y_2		$x_{\beta+2}$	x_3
		$x_{\beta+1}$	y_1		y_{lpha}	x_2
		x_eta			y_{lpha-1}	x_1
		$x_{\beta-1}$			y_{lpha-2}	
		:			:	
		x_2			y_1	
		\overline{x}_1				

If $y_{\alpha-1} > x_1$ and $y_{\alpha} \le x_2$, then we swap $y_{\alpha-1}$ with x_1 . We also "clockwise rotate" the entries $x_{\beta+2}$ and $x_{\beta+3}$ in the *c*-th column, and $x_{\beta+1}$ in the (c+1)-th column, as shown in our next diagram.

In our definition of φ_2 , a clockwise rotation of a set of entries in the *c*-th and (c+1)-th columns moves all entries in the *c*-th column up one cell except the topmost entry, which moves to the topmost cell in the (c+1)-th column containing an entry being rotated. The rotation moves all entries in the (c+1)-th column down one cell except the bottommost entry, which moves to the bottommost cell in the *c*-th column containing an entry being rotated. As another example, a rotation of a single entry in the *c*-th column and a single entry in the (c+1)-th column is a swap of those entries. We will continue to describe all cases of φ_2 with rotations of different sets of entries.

If $y_{\alpha-1} \leq x_1$ and $y_{\alpha} > x_2$, then we swap y_{α} with x_2 . We also clockwise rotate $x_{\beta+2}$ in the *c*-th column, and $x_{\beta+1}$ and x_{β} in the (c+1)-th column.

If $y_{\alpha-1} > x_1$ and $y_{\alpha} > x_2$, then we swap both $y_{\alpha-1}$ with x_1 and y_{α} with x_2 . We also clockwise rotate $x_{\beta+2}$ and $x_{\beta+3}$ in the *c*-th column, and $x_{\beta+1}$ and x_{β} in the (*c*+1)-th column.

Concretely, the two row overlap cases are as follows:



Consider the entries involved in the clockwise rotation in each case. For the topmost entry in the *c*-th column to move strictly up to the (c + 1)-th column, and the bottommost entry in the (c + 1)-th column move strictly down to the *c*-th column, we must have $\alpha \ge 3$. This property is necessary for the tableau to remain semistandard after the rotation, which partially necessitates the $\alpha \ge 4$ assumption, which is analogous to the $\alpha \ge 2$ assumption for φ_1 .

If there is one row of overlap, then i as defined for φ_0 is α , and x_1 is to the right of $y_{\alpha-1}$ with $x_1 < y_{\alpha-1}$:



In this case, we only have the single pair of decreasing entries, $y_{\alpha-1} < x_1$, so we swap $y_{\alpha-1}$ with x_1 . However, for the additional modifications after and in addition to this swap, we have different subcases.

If $y_{\alpha} < x_{\beta+2}$, we clockwise rotate $x_{\beta+1}$ and $x_{\beta+2}$ in the *c*-th column, and x_{β} in the (c+1)-th column. If $x_{\beta+2} \leq y_{\alpha} < x_{\beta+1}$, we swap $x_{\beta+2}$ with y_{α} , and $x_{\beta+1}$ with x_{β} . Observe in particular that this subcase

is two separate swaps, and not a rotation.

If $x_{\beta+1} \leq y_{\alpha}$, we clockwise rotate $x_{\beta+1}$ and $x_{\beta+2}$ in the *c*-th column, and y_{α} , x_{β} , and $x_{\beta-1}$ in the (*c*+1)-th column. Concretely, the one row overlap cases are as follows:



Again, consider the entries involved in the modifications in each case, either the rotations when $y_{\alpha} < x_{\beta+2}$ or $x_{\beta+1} \leq y_{\alpha}$, or the swaps when $x_{\beta+2} \leq y_{\alpha} < x_{\beta+1}$. For the topmost entry in the *c*-th column to move strictly up to the (c+1)-th column, and the bottommost entry in the (c+1)-th column to move strictly down to the *c*-th column, we must have $\alpha \geq 4$. This property is necessary for the tableau to remain semistandard after the rotation, as we will see in Lemma 3.19, which fully necessitates the $\alpha \geq 4$ assumption.

Example 3.18. For $\lambda = (2^4, 1^4)$ and T from Example 3.16, we get all the two row overlap cases of φ_2 :

(3.14)

1	7	₽	1	2	1	8	→	1	3	1	9	Ŷ	1	2
2	10		3	5	2	9		2	4	2	10		4	3
3	11		4	6	3	11		5	6	3	11		5	6
4	12		7	8	4	12		7	8	4	12		7	9
5			9	10	5			9	10	5			8	10
6			11		6			11		6			11	
8			12		7			12		7			12	
9				•	10				-	8				

For the same λ , we have all the one row overlap cases of φ_2 as follows, including the T from Example 3.17:

(3.15)



The proof that φ_2 is an injection of semistandard tableaux relies on the following lemma regarding the clockwise rotation.

Lemma 3.19. Suppose S is a semistandard tableau. Suppose that we perform a clockwise rotation on elements in the c-th and (c+1)-th columns of S to obtain S', such that the following are true:

- (1) The bottommost rotated entry in the c-th column in S is less than or equal to the topmost rotated entry in the (c+1)-th column in S.
- (2) The entry moving from the c-th column in S to the (c+1)-th column in S' moves strictly upwards.
- (3) The entry moving from the (c+1)-th column in S to the c-th column in S' moves strictly downward.
- (4) The entry moving from the c-th column in S to the (c + 1)-th column in S' is greater than the entry above it in the (c + 1)-th column in S', if such an entry exists.
- (5) The entry moving from the (c+1)-th column to the c-th column is less than the entry below it in the c-th column in S'.

Then S' is semistandard.

Proposition 3.20. Let λ and μ be as in (3.2) with $\beta = \alpha \ge 4$. Then φ_2 as defined above is an injection

$$\varphi_2$$
: SSYT $(\lambda) \rightarrow$ SSYT (μ) .

The proof of Proposition 3.20 is technical and omitted here. It follows similar ideas to the proof of Proposition 3.12. We can now further improve upon Corollary 3.6 and Corollary 3.13.

Corollary 3.21. The partitions λ and μ as in (3.2) with $\beta \ge \alpha \ge 4$ form a cover in the immersion poset.

We summarize the bounds on α and β needed for each map to be an injection:

Map	α	β
$arphi_0$	$\alpha \ge 0$	$\beta \geqslant \alpha + 2$
φ_1	$\alpha \geqslant 2$	β = α + 1
φ_2	$\alpha \geqslant 4$	β = α

3.3. Immersion poset on hook partitions. For this section, set $\lambda^i = (i, 1^{n-i}) \vdash n$ and let $S = \{\lambda^i \mid 1 \leq i \leq n\}$ be the set of all hook partitions. We study the immersion poset restricted to S.

Proposition 3.22. Let $1 \leq i \leq n$ and $\alpha = (\alpha_1, \ldots, \alpha_k) \vdash n$ such that $\alpha \leq_D \lambda^i$. Then

$$K_{\lambda^i,\alpha} = \binom{k-1}{n-i}.$$

Proof. Since λ^i dominates α , we know that $K_{\lambda^i,\alpha} \ge 1$. To form a semistandard Young tableau of shape λ^i and content α , the α_1 entries 1 must be placed leftmost in the first row of λ^i . The remaining n - i positions in the first column of λ^i can be filled with distinct values from the set $\{2, 3, \ldots, k\}$. This gives $\binom{k-1}{n-i}$ choices. Once these are placed, there is only one way to fill the remainder of the first row so that the resulting tableau is semistandard.

Recall from Lemma 2.1 that $\mu \leq I \lambda$ if and only if $K_{\mu,\alpha} \leq K_{\lambda,\alpha}$ for all $\alpha \vdash n$. Hence with Proposition 3.22, we are now ready to describe all the relations between hook partitions $\lambda^i \in S$ in the immersion poset. To illustrate what the proposition implies, we form a matrix of values in the following way:

- The j-th column is indexed by the content α^{j} , where α^{j} is any content that has j parts.
- The *i*-th row is indexed by the shape λ^i .
- The (i, j) entry of this matrix is the value $T_{i,j} := K_{\lambda^i, \alpha^j} = {j-1 \choose n-i}$ for $1 \le i, j \le n$.

Example 3.23. We give the matrix for n = 7:

# of parts Partition	1	2	3	4	5	6	7
(1 ⁷)	0	0	0	0	0	0	1
$(2,1^5)$	0	0	0	0	0	1	6
$(3,1^4)$	0	0	0	0	1	5	15
$(4,1^3)$	0	0	0	1	4	10	20
$(5,1^2)$	0	0	1	3	6	10	15
(6,1)	0	1	2	3	4	5	6
(7)	1	1	1	1	1	1	1

Remark 3.24. In this context, $\lambda^i \ge_I \lambda^j$ if and only if $T_{i,m} \ge T_{j,m}$ for all m. Equivalently, since $T_{j,m} = 0$ when $n - j \ge m$ and λ^i dominates λ^j when i > j, we need only show $T_{i,m} \ge T_{j,m}$ for all m > n - j when i > j.

The following lemma is used to prove the structure of the immersion poset restricted to hook partitions. **Lemma 3.25.** Suppose $\binom{n-1}{n-i} \ge \binom{n-1}{n-j}$ and i > j (note that this implies $j \le \frac{n}{2}$). Then for all $0 \le p \le j-1$, we have

$$\binom{n-1-p}{n-i} \ge \binom{n-1-p}{n-j}$$

The proof follows from basic properties of binomial coefficients, and is omitted here.

Corollary 3.26. If $T_{i,n} \ge T_{j,n}$ for i > j, then $\lambda^i \ge_I \lambda^j$.

Proof. By Proposition 3.22, $T_{i,n-p} = K_{\lambda^i,\alpha^{n-p}} = \binom{(n-p)-1}{n-i}$. Hence if $T_{i,n} \ge T_{j,n}$, by Lemma 3.25, we also have $T_{i,n-p} \ge T_{j,n-p}$ for $0 \le p \le j-1$. By Remark 3.24, this implies $\lambda^i \ge_I \lambda^j$.

Example 3.27. Take the rows corresponding to the partitions $(5, 1^2)$ and $(3, 1^4)$ in Example 3.23. Since the last column entries give $T_{5,7} = 15 \ge 15 = T_{3,7}$, then by Corollary 3.26 we also have $T_{5,7-p} \ge T_{3,7-p}$ for $1 \le p \le 2$: $T_{5,6} = 10 \ge 5 = T_{3,6}$, $T_{5,5} = 6 \ge 1 = T_{3,5}$.

We now describe the relations in the immersion poset on S depending upon whether n is even or odd.



FIGURE 2. Immersion poset restricted to hook partitions for n = 2k + 1 (left) and n = 2k(right).

Proposition 3.28. Let n = 2k + 1 be odd, then:

- $\begin{array}{ll} (1) \ \lambda^{\ell+1} \geq_I \lambda^{\ell} \ for \ all \ 1 \leq \ell \leq k. \\ (2) \ (\lambda^{k+1-\ell})^t = \lambda^{k+1+\ell} \geq_I \lambda^{k+1-\ell} \ for \ all \ 1 \leq \ell \leq k. \end{array}$
- (3) For any $1 < i \le k+1$, λ^i is incomparable to λ^j for all j > n-i+1.
- (4) For any $k + 2 \leq i < n$, λ^i is incomparable to λ^j for all j > i.

These describe all relations in the immersion poset restricted to hook partitions S.

Proof. Let us first prove (1). Fix an ℓ with $1 \leq \ell \leq k$. Then by Corollary 3.26, $\lambda^{\ell+1} \geq_I \lambda^{\ell}$ if and only if $T_{\ell+1,n} \ge T_{\ell,n}$. Note that $T_{\ell+1,n} = \binom{n-1}{n-\ell-1} = \binom{n-1}{\ell}$ and $T_{\ell,n} = \binom{n-1}{n-\ell} = \binom{n-1}{\ell-1}$. Since $1 \le \ell \le k$, we have $\binom{n-1}{\ell} \ge \binom{n-1}{\ell-1}$ and the result follows.

To prove (2), note that $\lambda^{k+1+\ell} \ge_I \lambda^{k+1-\ell}$ if and only if $T_{k+1+\ell,n} \ge T_{k+1-\ell,n}$. Since $T_{k+1+\ell,n} = \binom{n-1}{k-\ell} = \binom{n-1}{k-\ell}$ $\binom{2k}{k-\ell} = \binom{2k}{k+\ell} = \binom{n-1}{k+\ell} = T_{k+1-\ell,n}$, the result follows.

To prove (3) we show for any $1 < i \le k+1$ that λ^i is incomparable to λ^j for all j > n-i+1. Since λ^j dominates λ^i , we need only show there exists some α such that $K_{\lambda^i,\alpha} > K_{\lambda^j,\alpha}$. Choose $\alpha = (1^n)$. Then $K_{\lambda^{i},\alpha} = \binom{n-1}{n-i} = \binom{n-1}{i-1} > \binom{n-1}{n-j} = K_{\lambda^{j},\alpha} \text{ because } i-1 > n-j \text{ and } 1 < i \leq k+1.$

Lastly, to prove (4) we follow the same strategy as (3). Since λ^j dominates λ^i , we can let $\alpha = (1^n)$, and since k+1 < i < j we get $K_{\lambda^i,\alpha} = \binom{n-1}{n-i} > \binom{n-1}{n-j} = K_{\lambda^j,\alpha}$, and the result follows. \square

Proposition 3.29. Let n = 2k be even, then:

- (1) $\lambda^{\ell+1} \geq_I \lambda^{\ell}$ for all $1 \leq \ell < k$. (2) $(\lambda^{k-\ell})^t = \lambda^{k+1+\ell} \geq_I \lambda^{k-\ell}$ for all $0 \leq \ell \leq k-1$.
- (3) For any $1 < i \le k$, λ^i is incomparable to λ^j for all j > n i + 1.
- (4) For any $k + 1 \leq i < n$, λ^i is incomparable to λ^j for all j > i.

These describe all relations in the immersion poset restricted to hook partitions S.

The proof of the even case is similar to the odd case.

The Hasse diagram of the immersion poset restricted to hook partitions is given in Figure 2. Notice that item (1) in Propositions 3.28 and 3.29 proves the string of covers on the left going up each Hasse diagram, while (2) proves the covers going up the right side which are the transposes.

Corollary 3.30. Let $\lambda \in S$ be a hook partition such that $\lambda \leq \lambda^t$. Then $\lambda \leq \lambda^t$.

The *rank* of a poset is the length of the longest chain of elements of the poset.

Corollary 3.31. The rank of the immersion poset $(\mathcal{P}(n), \leq_I)$ is at least |n/2|.

3.4. Immersion poset on two column partitions. Now let S be the set partitions with at most two columns, that is, $S = \{\lambda \mid \lambda_1 \leq 2\}$. If n = 2k, then for this section we define $\lambda^j = (2^{k-j}, 1^{2j})$ for $0 \leq j \leq k$. Similarly, if n = 2k + 1, then $\lambda^j = (2^{k-j}, 1^{2j+1})$ for $0 \leq j \leq k$. In this section, we study the immersion poset restricted to S.

Remark 3.32. Note that $K_{\lambda,\mu} = 0$ if $\lambda \in S$ and $\mu \notin S$. Hence there does not exist an immersion pair $\mu \leq_I \lambda$ with $\lambda \in S$ and $\mu \notin S$. This implies that if λ^i is a cover for λ^j in the subposet restricted to S, then λ^i is a cover for λ^j in the immersion poset.

This remark implies that we only need to consider $K_{\lambda,\mu}$ for $\lambda, \mu \in S$ when determining the immersion relations for this subset. Recall that f^{λ} is the number of standard Young tableaux of shape λ .

Proposition 3.33.

- (1) Let $\lambda^j = (2^{k-j}, 1^{2j}) \vdash 2k$ for $0 \leq j \leq k$. Then $K_{\lambda^i, \lambda^j} = f^{(j+i,j-i)}$ when $i \leq j$ and $K_{\lambda^i, \lambda^j} = 0$ when i > j.
- (2) Let $\lambda^j = (2^{k-j}, 1^{2j+1}) \vdash 2k+1$ for $0 \leq j \leq k$. Then $K_{\lambda^i, \lambda^j} = f^{(j+i+1,j-i)}$ when $i \leq j$ and $K_{\lambda^i, \lambda^j} = 0$ when i > j.

Proof. For (1), if i > j, λ^j dominates λ^i and hence $K_{\lambda^i,\lambda^j} = 0$. If i = j, clearly $K_{\lambda^i,\lambda^i} = f^{(2i)} = 1$. Suppose j > i. Then the first k - j of the k - i two length rows of any tableau $T \in \mathsf{SSYT}(\lambda^i, \lambda^j)$ are fixed by the content. Hence, there is a bijection $\mathsf{SSYT}(\lambda^i, \lambda^j) \to \mathsf{SSYT}((2^{j-i}, 1^{2i}), (1^{2j}))$ by removing the first k - j rows. Note that $K_{(2^{j-i}, 1^{2i}), (1^{2j})} = f^{(2^{j-i}, 1^{2i})}$, which is also equal to the number of standard tableaux of the transpose of $(2^{j-i}, 1^{2i})$. The result follows.

The proof of part (2) is similar.

Using the hook length formula with Proposition 3.33, we can describe K_{λ^i,λ^j} . We present this in the form of a matrix. More explicitly, suppose n = 2k or n = 2k + 1. Then for all $0 \le i, j \le k$, the *i*-th row and *j*-th column entry of the matrix $T = (T_{i,j})$ is $T_{i,j} = K_{\lambda^i,\lambda^j}$. Note that the indexing starts with 0.

Example 3.34. Below are the matrices in tabular form for cases n = 14, 15.

The case when $n = 14$:													
Content Shape	(2^7)	$(2^6, 1^2)$	$(2^5, 1^4)$	$(2^4, 1^6)$	$(2^3, 1^8)$	$(2^2, 1^{10})$	$(2, 1^{12})$	(1^{14})					
(2^7)	1	1	2	5	14	42	132	429					
$(2^6, 1^2)$	0	1	3	9	28	90	297	1001					
$(2^5, 1^4)$	0	0	1	5	20	75	275	1001					
$(2^4, 1^6)$	0	0	0	1	7	35	154	637					
$(2^3, 1^8)$	0	0	0	0	1	9	54	273					
$(2^2, 1^{10})$	0	0	0	0	0	1	11	77					
$(2,1^{12})$	0	0	0	0	0	0	1	13					
(1^{14})	0	0	0	0	0	0	0	1					

Content Shape	$(2^{7}, 1)$	$(2^6, 1^3)$	$(2^5, 1^5)$	$(2^4, 1^7)$	$(2^3, 1^9)$	$(2^2, 1^{11})$	$(2, 1^{13})$	(1^{15})
$(2^7, 1)$	1	2	5	14	42	132	429	1430
$(2^6, 1^3)$	0	1	4	14	48	165	572	2002
$(2^5, 1^5)$	0	0	1	6	27	110	429	1638
$(2^4, 1^7)$	0	0	0	1	8	44	208	910
$(2^3, 1^9)$	0	0	0	0	1	10	65	350
$(2^2, 1^{11})$	0	0	0	0	0	1	12	90
$(2,1^{13})$	0	0	0	0	0	0	1	14
(1^{15})	0	0	0	0	0	0	0	1

The case when n = 15:

Since the columns and rows are decreasing in dominance order, for any i < j we have $\lambda^i \ge_I \lambda^j$ if $T_{i,m} \ge T_{j,m}$ for all $0 \le m \le k$. In the following lemma, we prove some properties of the matrix T that will show that this statement is equivalent to only comparing values in the last column of the matrix. That is, if i < j and $T_{i,k} \ge T_{j,k}$, then $\lambda^i \ge_I \lambda^j$. The reader can verify this in Example 3.34.

Lemma 3.35. The matrix $(T_{i,j}) = (K_{\lambda^i,\lambda^j})$ defined above with $0 \le i, j \le k$ has the following properties:

- (1) The entries weakly increase within each row.
- (2) The entries within each column are unimodal.
- (3) The rate of change of entries within a row increases as the row number increases. In particular, for any fixed i and j with $0 \le i < j \le k$, we have for all $j \le r < k$:

$$\frac{T_{i,r+1}}{T_{i,r}} < \frac{T_{j,r+1}}{T_{j,r}}$$

(4) For any fixed i and j with i < j, if $T_{i,k} \ge T_{j,k}$, then $T_{i,m} \ge T_{j,m}$ for all $0 \le m \le k$.

Proof. We begin by proving (1). Let n = 2k be even. Then for a fixed row *i*, given any $i \leq j < k$, we need to show that $T_{i,j+1} \geq T_{i,j}$. Using Proposition 3.33, we have:

$$\frac{T_{i,j+1}}{T_{i,j}} = \frac{f^{(j+1+i,j+1-i)}}{f^{(j+i,j-i)}} = \frac{(2j+2)(2j+1)}{(j+i+2)(j+1-i)} \ge 1$$

because $j \ge i$ implies

$$2j+2 \ge j+i+2$$
 and $2j+1 \ge j+1-i$.

Now let n = 2k + 1 be odd. Using the same strategy as the even case we have:

$$\frac{T_{i,j+1}}{T_{i,j}} = \frac{f^{(j+2+i,j+1-i)}}{f^{(j+i+1,j-i)}} = \frac{(2j+3)(2j+2)}{(j+i+3)(j+1-i)} \ge 1$$

because $j \ge i$ implies

 $2j+3 \ge j+i+3$ and $2j+2 \ge j+1-i$.

Next we prove statement (2). Let n = 2k be even. Since statement (2) holds trivially if there is only one non-zero entry in the column, we focus on columns with more than one non-zero entry. Fix a $2 \le j \le k$. To determine when the column is increasing and decreasing we consider the fraction:

$$\frac{T_{i+1,j}}{T_{i,j}} = \frac{f^{(j+i+1,j-i-1)}}{f^{(j+i,j-i)}} = \frac{\frac{(2j)!(2i+3)}{(j+i+2)!(j-i-1)!}}{\frac{(2j)!(2i+1)!}{(j+i+1)!(j-i)!}} = \frac{(2i+3)(j-i)}{(2i+1)(j+i+2)}.$$

Analyzing the following inequalities gives:

(3.16)
$$\begin{aligned} \frac{T_{i+1,j}}{T_{i,j}} > 1 &\iff 2i^2 + 4i + 1 < j, \\ \frac{T_{i+1,j}}{T_{i,j}} = 1 &\iff 2i^2 + 4i + 1 = j, \\ \frac{T_{i+1,j}}{T_{i,j}} < 1 &\iff 2i^2 + 4i + 1 > j. \end{aligned}$$

Thus, for values of *i* such that $2i^2 + 4i + 1 < j$ the column entries are increasing, and when the values of *i* satisfy $2i^2 + 4i + 1 > j$ the column entries are decreasing. This proves (2) for the even case.

Now let n = 2k + 1 be odd. Fix a $2 \le j \le k$. Similar to the even case we have:

$$\frac{T_{i+1,j}}{T_{i,j}} = \frac{f^{(j+i+2,j-i-1)}}{f^{(j+i+1,j-i)}} = \frac{\frac{(2j+1)!(2i+4)}{(j+i+3)!(j-i-1)!}}{\frac{(2j+1)!(2i+2)}{(j+i+2)!(j-i)!}} = \frac{(2i+4)(j-i)}{(2i+2)(j+i+3)}$$

Analyzing the following inequalities gives:

(3.17)
$$\begin{aligned} \frac{T_{i+1,j}}{T_{i,j}} > 1 &\iff 2i^2 + 6i + 3 < j, \\ \frac{T_{i+1,j}}{T_{i,j}} = 1 &\iff 2i^2 + 6i + 3 = j, \\ \frac{T_{i+1,j}}{T_{i,j}} < 1 &\iff 2i^2 + 6i + 3 > j. \end{aligned}$$

Again, we notice that for values of *i* such that $2i^2 + 6i + 3 < j$ the column entries are increasing, and when the values of *i* satisfy $2i^2 + 6i + 3 > j$ the column entries are decreasing. This concludes the proof of (2).

To prove statement (3), we use Proposition 3.33 and the hook length formula to get the following equivalences:

$$\frac{T_{i,r+1}}{T_{i,r}} < \frac{T_{j,r+1}}{T_{j,r}} \quad \Longleftrightarrow \quad \frac{K_{\lambda^i,\lambda^{r+1}}}{K_{\lambda^i,\lambda^r}} < \frac{K_{\lambda^j,\lambda^{r+1}}}{K_{\lambda^j,\lambda^r}} \quad \Longleftrightarrow \quad \frac{(r+j+2)(r+1-j)}{(r+i+2)(r+1-i)} < 1 \quad \Longleftrightarrow \quad i^2+i < j^2+j.$$

The last inequality is always true since $0 \le i < j$, thus proving (3).

To prove (4), fix *i* and *j* with i < j where $T_{i,k} \ge T_{j,k}$. Then by statement (3) it directly follows that $T_{i,m} \ge T_{j,m}$ for all $j \le m \le k$. Because $T_{j,m} = 0$ for all $0 \le m < j$, it trivially follows that $T_{i,m} \ge T_{j,m}$ for these values of *m*, this finishes the proof of (4).

The beauty of Lemma 3.35, in particular statement (4), is that we can now reduce much of the work in determining the immersion relations between partitions in S to just comparing the numbers of standard Young tableaux, as is done in the next proposition.

Proposition 3.36. For n = 2k even or n = 2k + 1 odd, the last (k-th) column of T can be used to completely determine relations in the immersion poset restricted to the subset S. In particular:

- (1) $\lambda^i \geq_I \lambda^j$ if and only if i < j and $T_{i,k} \geq T_{j,k}$,
- (2) For i < j, λ^i and λ^j are incomparable if and only if $T_{j,k} > T_{i,k}$.

Proof. To prove (1), by definition $\lambda^i \ge_I \lambda^j$ if and only if $\lambda^i >_D \lambda^j$ and $T_{i,\ell} \ge T_{j,\ell}$ for all $0 \le \ell \le k$. But λ^i dominates λ^j if and only if i < j, and by (4) of Lemma 3.35, $T_{i,\ell} \ge T_{j,\ell}$ for all $0 \le \ell \le k$ if and only if $T_{i,k} \ge T_{j,k}$ (when i < j).

To prove (2), let i < j. If λ^i and λ^j are incomparable, then there exists some ℓ such that $T_{j,\ell} > T_{i,\ell}$. By (3) of Lemma 3.35, we have:

$$\frac{T_{i,r+1}}{T_{i,r}} \leqslant \frac{T_{j,r+1}}{T_{j,r}}$$

for all $\ell \leq r < k$, which guarantees that $T_{i,k} > T_{i,k}$.

As a consequence we obtain the following immediate corollary.

Corollary 3.37. The cover relations for the immersion poset of the set S are the exact same as those in the standard immersion poset.

We can now explain the cover relations of the immersion poset restricted to the set S.

Proposition 3.38. Let n = 2k be even or n = 2k + 1 be odd, then:

- (1) $\lambda^i >_I \lambda^{i+1}$ when $2i^2 + 4i + 2 > k$ for n even and $2i^2 + 6i + 4 > k$ for n odd. This also coincides with Lemma 2.7, taking a = k i 1 and b = 2i + 2 (n even) or b = 2i + 3 (n odd).
- (2) λ^i and λ^j are incomparable in the immersion poset for all $0 \le i, j \le i_{max}$ with $i \ne j$ and i_{max} being the largest i value not satisfying (1).
- (3) Fix i with $0 \le i \le i_{max}$ and let $m > i_{max} i$ be smallest such that $T_{i,k} \ge T_{i+m,k}$. Then $\lambda^i \ge_I \lambda^{i+m}$.

Proof. If $T_{i,k} \ge T_{i+1,k}$, then by Proposition 3.36 and Remark 3.32, we have that $\lambda^i \ge_I \lambda^{i+1}$ is a cover. We determine the values for *i* such that $T_{i,k} \ge T_{i+1,k}$ by using the middle equation and bottom inequality of (3.16) (for *n* even) and (3.17) (for *n* odd), where we replace *j* with *k*. Specifically, for *n* even:

$$2i^2 + 4i + 1 \ge k \implies 2i^2 + 4i + 2 > k,$$

and for n odd:

$$2i^2 + 6i + 3 \ge k \implies 2i^2 + 6i + 4 > k.$$

To prove (2), notice that since i_{max} is the number of the row containing the first maximum, by the increasing nature of the column up to the maximum value given by (2) of Lemma 3.35, then for any $0 \le i < j \le i_{max}$ we have $T_{i,k} < T_{j,k}$. Hence by Proposition 3.36 (2), λ^i and λ^j are incomparable.

To prove (3) notice that by Proposition 3.36 statement (1), since m is the smallest value it must be a cover.

Example 3.39. Suppose n = 14, so that k = 7. By (1) of Proposition 3.38, the inequality holds for $1 \le i \le k = 7$ so we obtain:

$$\lambda^7 \lessdot_I \lambda^6 \lessdot_I \lambda^5 \lessdot_I \lambda^4 \lessdot_I \lambda^3 \lessdot_I \lambda^2 \lessdot_I \lambda^1$$

Applying (3) of Proposition 3.38 to λ^0 , with i = 0 we find that m = 4:

$$T_{0.7} = 429 \ge 273 = T_{4.7}$$
.

Notice that m = 3 does not satisfy the inequality:

$$T_{0,7} = 429 \ge 637 = T_{3,7}.$$

So our final cover relation for the poset is $\lambda^4 \leq_I \lambda^0$.

3.5. Lower intervals and Schur-positivity of interval power sums. In this section, we make conjectures about certain lower intervals $A_{\mu} := \{\lambda | (1^n) \leq_I \lambda \leq_I \mu\}$ in the immersion poset. Determining intervals will

- (1) enhance our understanding of the immersion of polynomial representations for $GL_N(\mathbb{C})$ and
- (2) allow us to investigate when $p_{A_{\mu}}$ of Equation (1.1) is Schur-positive, as asked in Question 1.3. We call $p_{A_{\mu}}$ an interval power sum. It also helps towards constructing a natural corresponding representation of the symmetric group.

In this section, we prove that $p_{A_{\mu}}$ is Schur-positive for the conjectured intervals.

Conjecture 3.40. For n = 5 and $n \ge 9$, the interval $A_{(n-2,2)} = \{\lambda \mid (1^n) \le I \lambda \le I (n-2,2)\}$ is exactly

$$(1^n) \leq_I (2, 1^{n-2}) \leq_I (2, 2, 1^{n-4}) \leq_I (n-2, 2)$$

Remark 3.41. The first two covers are consequences of Proposition 3.38 (1). The map

 $\mathsf{SSYT}((2,2,1^{n-4}),\nu) \longrightarrow \mathsf{SSYT}((n-2,2),\nu),$



FIGURE 3. Subposet of the immersion poset only containing partitions in $A_{(n-2,2)}$ for n = 8 (left), n = 9 (middle), and n = 15 (right).

which is the transpose if $\nu_1 = 1$ and which moves the boxes in positions $(3, 1), \ldots, (n-2, 1)$ to positions $(1,3), \ldots, (1, n-2)$ if $\nu_1 = 2$ is an injection. This shows that $(2,2,1^{n-4}) <_I (n-2,2)$. Therefore, we have

$$\{(1^n), (2, 1^{n-2}), (2, 2, 1^{n-4}), (n-2, 2)\} \subseteq A_{(n-2,2)}$$

However, we have not proven the cover relation $(2, 2, 1^{n-4}) \leq_I (n-2, 2)$. One strategy to show the reverse containment is to argue that for all partitions λ such that λ and λ^t are not included in the above list, we have $f^{\lambda} > \frac{n(n-3)}{2} = f^{(n-2,2)}$. This would prove that $\lambda \notin_I (n-2,2)$, hence $\lambda \notin A_{(n-2,2)}$. We have confirmed the conjecture up to n = 18. See Figure 3.

Proposition 3.42.

- (1) For n < 7 and n = 8, $p_{A_{(n-2,2)}}$ is Schur-positive.
- (2) For $n \ge 9$, $p_{(1^n)} + p_{(2,1^{n-2})} + p_{(2,2,1^{n-4})} + p_{(n-2,2)}$ is Schur-positive.

Proof. Part (1) can be checked explicitly by SAGEMATH. For part (2), let

(3.18)
$$p_{(1^n)} + p_{(2,1^{n-2})} + p_{(2,2,1^{n-4})} + p_{(n-2,2)} = \sum_{\lambda \vdash n} c_\lambda s_\lambda$$

We prove that $c_{\lambda} \ge 0$ for all $\lambda \vdash n$ by proving that all partial sums $p_{(1^n)}$, $p_{(1^n)} + p_{(2,1^{n-2})}$, $p_{(1^n)} + p_{(2,1^{n-2})} + p_{(2,2,1^{n-4})} + p_{(n-2,2)}$ are Schur-positive. We employ the combinatorial Murnaghan–Nakayama rule involving ribbon tableaux (see for example [Sta99, Chapter 7.17])

$$p_{\mu} = \sum_{\lambda \vdash n} \chi^{\lambda}(\mu) s_{\lambda}$$
 where $\chi^{\lambda}(\mu) = \sum_{T \in \mathsf{R}(\lambda,\mu)} (-1)^{\operatorname{ht}(T)}$

and $\mathsf{R}(\lambda,\mu)$ is the set of all ribbon tableaux of shape λ and type μ and $\mathsf{ht}(T)$ is equal to the sum of the heights of all ribbons in T. We show that each subset of ribbon tableaux that contributes a negative term to c_{λ} is in bijection with a distinct subset of ribbon tableaux that contributes a positive number to c_{λ} , ensuring that $c_{\lambda} \ge 0$. We examine each partial sum of power sum symmetric functions, and demonstrate Schur-positivity at each step through these bijections.

(1) It is well-known that $p_{(1^n)} = \sum_{\lambda \vdash n} f^{\lambda} s_{\lambda}$ since $\mathsf{R}(\lambda, (1^n))$ is the set of all standard Young tableaux of shape λ .

(2) For $T \in \mathbb{R}(\lambda, (2, 1^{n-2}))$, T has either a horizontal or a vertical 2-ribbon and the remaining are single box ribbons. If T has a horizontal 2-ribbon, then $\operatorname{ht}(T) = 0$ and T contributes +1 to $\chi^{\lambda}((2, 1^{n-2}))$. There are $f^{\lambda/(2)}$ such ribbon tableaux in $\mathbb{R}(\lambda, (2, 1^{n-2}))$, where $f^{\lambda/\mu}$ is the cardinality of $\mathsf{SYT}(\lambda/\mu)$, the set of standard Young tableaux of skew shape λ/μ . If T has a vertical 2-ribbon, then $\operatorname{ht}(T) = 1$ and Tcontributes -1 to $\chi^{\lambda}((2, 1^{n-2}))$. There are $f^{\lambda/(1,1)}$ such ribbon tableaux in $\mathbb{R}(\lambda, (2, 1^{n-2}))$. Therefore, the coefficient of s_{λ} in $p_{(2,1^{n-2})}$ is $f^{\lambda/(2)} - f^{\lambda/(1,1)}$. If $(1,1) \subseteq \lambda$, then c_{λ} includes $-f^{\lambda/(1,1)}$. The natural bijection

$$\mathsf{SYT}(\lambda/(1,1)) \rightarrow \{T \in \mathsf{SYT}(\lambda) \mid T_{1,1} = 1 \text{ and } T_{2,1} = 2\}$$

demonstrates that $f^{\lambda} - f^{\lambda/(1,1)} \ge 0$. Hence $p_{(1^n)} + p_{(2,1^{n-2})}$ is Schur-positive.

(3) For any $T \in \mathsf{R}(\lambda, (2, 2, 1^{n-4}))$, there are six possible ways to arrange two 2-ribbons.



ht(T) = 0 ht(T) = 2 ht(T) = 0 ht(T) = 2 ht(T) = 1 ht(T) = 1

The remaining n-4 ribbons in T are single boxes. Therefore, the coefficient of s_{λ} in $p_{(2,2,1^{n-4})}$ is

$$2f^{\lambda/(2,2)} + f^{\lambda/(4)} + f^{\lambda/(1^4)} - f^{\lambda/(3,1)} - f^{\lambda/(2,1,1)}.$$

If $(3,1) \subseteq \lambda$, then c_{λ} includes $-f^{\lambda/(3,1)}$. Consider the bijection

$$\mathsf{SYT}(\lambda/(3,1)) \to \{T \in \mathsf{SYT}(\lambda/(2)) \mid T_{1,3} = 1, \text{ and } T_{2,1} = 2\}.$$

If $(2,1,1) \subseteq \lambda$, then c_{λ} includes $-f^{\lambda/(2,1,1)}$. Consider the bijection

$$\text{SYT}(\lambda/(2,1,1)) \to \{T \in \text{SYT}(\lambda/(2)) \mid T_{2,1} = 1 \text{ and } T_{3,1} = 2\}.$$

Hence $f^{\lambda/(2)}$ from (2) and the terms $-f^{\lambda/(3,1)} - f^{\lambda/(2,1,1)}$ from (3) satisfy $f^{\lambda/(2)} - f^{\lambda/(3,1)} - f^{\lambda/(2,1,1)} \ge 0$. So far, we have shown that $p_{(1^n)} + p_{(2,1^{n-2})} + p_{(2,2,1^{n-4})}$ is Schur-positive.

(4) For $T \in \mathsf{R}(\lambda, (n-2,2))$, the possible ways of arranging one (n-2)-ribbon and one 2-ribbon in T are the following. Note that 1^0 appearing in λ means that there are no parts of size 1 in λ .



 $ht(T) = b \qquad ht(T) = b - 2 + 1 \qquad ht(T) = b + 1 \qquad ht(T) = b + 2 + 1 \qquad ht(T) = n - 3 + 1 \qquad ht(T) = 0$

If $\lambda = (a, 1^b)$ with $2 \leq b \leq n-3$, then cases $\lambda^{(1)}$ and $\lambda^{(2)}$ both apply. Since $\operatorname{ht}(\lambda^{(1)})$ and $\operatorname{ht}(\lambda^{(2)})$ have opposite parity, $\chi^{(a,1^b)}((n-2,2)) = 0$. For $\lambda = (2, 1^{n-2})$, only $\lambda^{(2)}$ applies and $\chi^{(2,1^{n-2})}((n-2,2)) = (-1)^{n-3}$. For $\lambda = (1^n)$, only $\lambda^{(2)}$ applies and $\chi^{(1^n)}((n-2,2)) = (-1)^{n-1}$. Since (1^4) is contained in both $\lambda = (2, 1^{n-2}), (1^n), c_{\lambda}$ also includes $f^{\lambda/(1^4)} \geq 1$. For $\lambda = (n-1,1)$, only $\lambda^{(1)}$ applies and $\chi^{(n-1,1)}((n-2,2)) = -1$. In this case, $(4) \subseteq \lambda$, and thus c_{λ} also includes $f^{\lambda/(4)} \geq 1$. For $\lambda = (n)$, the height of any ribbon tableaux is 0, so there are no negatives to worry about.

If λ is of the form $\lambda^{(3)}$, $\lambda^{(4)}$, or $\lambda^{(5)}$, then it is possible that $c_{\lambda^{(i)}}$ includes -1 from the unique $T \in \mathbb{R}(\lambda^{(i)}, (n-2,2))$ for i = 3, 4, 5. In any of these disjoint cases, $(2,2) \subseteq \lambda^{(i)}$, which means $c_{\lambda^{(i)}}$ also includes $f^{\lambda^{(i)}/(2,2)} \ge 1$. Hence $f^{\lambda/(1^4)}$, $f^{\lambda/(4)}$, and $f^{\lambda/(2,2)}$ from (3) and $\chi^{\lambda}((n-2,2))$ from (4) satisfy $f^{\lambda/(1^4)} + f^{\lambda/(4)} + f^{\lambda/(2,2)} + \chi^{\lambda}((n-2,2)) \ge 0$. We have shown that $p_{(1^n)} + p_{(2,1^{n-2})} + p_{(2,2,1^{n-4})} + p_{(n-2,2)}$ is Schur-positive.

Conjecture 3.43. For $n \ge 9$, the interval $A_{(n-2,1,1)} = \{\lambda \mid (1^n) \le_I \lambda \le_I (n-2,1,1)\}$ is exactly

$$(1^n) \leq_I (2, 1^{n-2}) \leq_I (2, 2, 1^{n-4}) \leq_I (3, 1^{n-3}) \leq_I (n-2, 1, 1)$$

Remark 3.44. The first two covers are consequences of Corollary 3.6. By Proposition 3.5,

$$\varphi_0: SSYT((2,2,1^{n-4}),\nu) \to SSYT((3,1^{n-3}),\nu)$$

is an injection (with $\alpha = 0, \beta = 2$). Since $(2, 2, 1^{n-4}) <_D (3, 1^{n-3})$, this implies $(2, 2, 1^{n-4}) <_I (3, 1^{n-3})$. By Corollary 3.30, we know $(3, 1^{n-3}) <_I (n-2, 1, 1)$ because $(3, 1^{n-3}) <_D (3, 1^{n-3})^t = (n-2, 1, 1)$. This implies

$$\{(1^n), (2, 1^{n-2}), (2, 2, 1^{n-4}), (3, 1^{n-3}), (n-2, 1, 1)\} \subseteq A_{(n-2, 1, 1)}$$

However, we have not proven the cover relation $(3, 1^{n-3}) \leq_I (n-2, 1, 1)$. To show that $A_{(n-2,1,1)}$ is contained in the above set, one could show that for all partitions λ such that λ and λ^t are not included in the above list, we have $f^{\lambda} > \frac{(n-1)(n-2)}{2} = f^{(n-2,1,1)}$. This would prove that $\lambda \notin A_{(n-2,1,1)}$. We have confirmed the conjecture up to n = 18.

Proposition 3.45.

- (1) For n < 9, $p_{A_{(n-2,1,1)}}$ is Schur-positive.
- (2) For $n \ge 9$, $p_{(1^n)} + p_{(2,1^{n-2})} + p_{(2,2,1^{n-4})} + p_{(3,1^{n-3})} + p_{(n-2,1,1)}$ is Schur-positive.

Proof. Part (1) can be checked explicitly using SAGEMATH. For part (2), as shown in the proof of Proposition 3.42, $p_{(1^n)} + p_{(2,1^{n-2})} + p_{(2,2,1^{n-4})}$ is Schur-positive. We next show $p_{(1^n)} + p_{(2,1^{n-2})} + p_{(2,2,1^{n-4})} + p_{(3,1^{n-3})}$ is Schur-positive. For $T \in \mathsf{R}(\lambda, (3, 1^{n-3}))$, there are three possible ways of arranging one 3-ribbon in T.



ht(T) = 0 ht(T) = 1 ht(T) = 2

Therefore, the coefficient of s_{λ} in $p_{(3,1^{n-3})}$ is $f^{\lambda/(3)} - f^{\lambda/(2,1)} + f^{\lambda/(1,1,1)}$.

If $(2,1) \subseteq \lambda$, then c_{λ} includes $-f^{\lambda/(2,1)}$. Consider the bijection

$$SYT(\lambda/(2,1)) \rightarrow \{T \in SYT(\lambda) \mid T_{1,1} = 1, T_{1,2} = 2, \text{ and } T_{2,1} = 3\}.$$

Note, the above subset of standard Young tableaux of shape λ in the term f^{λ} in $p_{(1^n)}$ was not used in the bijections in the proof of Proposition 3.42. This shows that $p_{(1^n)} + p_{(2,1^{n-2})} + p_{(2,2,1^{n-4})} + p_{(3,1^{n-3})}$ is Schur-positive.

We now examine the Schur expansion of $p_{(n-2,1,1)}$. There are a few specific shapes λ where $\mathsf{R}(\lambda, (n-2,1,1))$ is nonempty. Note that for a ribbon tableau T of type (n-2,1,1), $\mathsf{ht}(T) = \mathsf{ht}(R_1)$, where R_1 is the (n-2)-ribbon of 1's in T. In Case 1, we examine all hook shapes $\lambda = (a, 1^b)$. In Case 2, we examine all shapes $\lambda = (a, 2, 1^b)$. In Case 3, we examine the remaining two shapes $(a, 3, 1^b)$ and $(a, 2, 2, 1^b)$.

Case 1a: $\lambda = (a, 1^b)$ with $4 \le a \le n-3$ and $3 \le b \le n-4$. These conditions on a, b require that the (n-2)-ribbon forms a hook with nontrivial arm and nontrivial leg.



ht(T) = b ht(T) = b - 2 ht(T) = b - 1 ht(T) = b - 1

Since b, b-2 have opposite parity to b-1, $\chi^{\lambda}((n-2,1,1)) = 0$ for $\lambda = (a, 1^b)$ with $4 \le a \le n-3$ and $3 \le b \le n-4$.

Case 1b: $\lambda = (3, 1^{n-3}).$



ht(T) = n - 3 ht(T) = n - 4 ht(T) = n - 4 ht(T) = n - 5

Since n - 3, n - 5 have opposite parity to $n - 4, \chi^{\lambda}((n - 2, 1, 1)) = 0$ for $\lambda = (3, 1^{n-3})$. Case 1c: $\lambda = (2, 1^{n-2})$.



ht(T) = n - 3 ht(T) = n - 3 ht(T) = n - 4

Since n-3 and n-4 have opposite parity, $\chi^{\lambda}((n-2,1,1)) = (-1)^{n-3}$ for $\lambda = (2,1^{n-2})$. Since $n \ge 5$, $(1^4) \subseteq \lambda$ and $f^{\lambda/(1^4)} \ge 1$, the coefficient of s_{λ} in $p_{(2,2,1^{n-4})}$ will cancel this potential negative.

Case 1d: $\lambda = (1^n)$. The unique ribbon tableau $T \in \mathsf{R}((1^n), (n-2, 1, 1))$ has height n-3, hence $\chi^{\lambda}((n-2, 1, 1)) = (-1)^{n-3}$ for $\lambda = (1^n)$. Since $n \ge 5$, $(1^4) \subseteq \lambda$ and $f^{\lambda/(1^4)} \ge 1$, the coefficient of s_{λ} in $p_{(2,2,1^{n-4})}$ will cancel this potential negative.

Case 1e: $\lambda = (n - 2, 1, 1).$



Thus, $\chi^{\lambda}((n-2,1,1)) = 0$ for $\lambda = (n-2,1,1)$.

Case 1f: $\lambda = (n - 1, 1)$.



Case 1g: $\lambda = (n)$. The unique ribbon tableau $T \in \mathsf{R}((n), (n-2, 1, 1))$ has height 0, hence $\chi^{\lambda}((n-2, 1, 1)) = 1$ for $\lambda = (n)$.

Case 2a: $\lambda = (a, 2, 1^b)$ with $3 \le a \le n - 3$ and $1 \le b \le n - 5$.



 $ht(T) = b + 1 \quad ht(T) = b + 1 \quad ht(T) = b \quad ht(T) = b$

Since b and b+1 have opposite parity, $\chi^{\lambda}((n-2,1,1)) = 0$ for $\lambda = (a,2,1^b)$ with $3 \le a \le n-3$ and $1 \le b \le n-5$.

Case 2b: $\lambda = (n - 2, 2)$.



Thus, $\chi^{\lambda}((n-2,1,1)) = -1$ for $\lambda = (n-2,2)$. Since $(2,2) \subseteq \lambda$ and $f^{\lambda/(2,2)} \ge 1$, the coefficient of s_{λ} in $p_{(2,2,1^{n-4})}$ will cancel this negative.

Case 2c: $\lambda = (2, 2, 1^{n-4}).$



ht(T) = n - 3 ht(T) = n - 4 ht(T) = n - 4

Since n-3 and n-4 have opposite parity, $\chi^{\lambda}((n-2,1,1)) = (-1)^{n-4}$ for $\lambda = (2,2,1^{n-4})$. Since $(2,2) \subseteq \lambda$ and $f^{\lambda/(2,2)} \ge 1$, the coefficient of s_{λ} in $p_{(2,2,1^{n-4})}$ will cancel this potential negative.

Case 3a: $\lambda = (n - (3 + b), 3, 1^b)$ for $0 \le b \le n - 6$. The unique ribbon tableau $T \in \mathsf{R}((n - (3 + b), 3, 1^b), (n - 2, 1, 1))$ has height b + 1, hence $\chi^{\lambda}((n - 2, 1, 1)) = (-1)^{b+1}$ for $\lambda = (n - (3 + b), 3, 1^b)$. Since $(2, 2) \subseteq \lambda$ and $f^{\lambda/(2,2)} \ge 1$, the coefficient of s_{λ} in $p_{(2,2,1^{n-4})}$ will cancel this potential negative.

Case 3b: $\lambda = (n - (4+b), 2, 2, 1^b)$ for $0 \le b \le n-6$. The unique ribbon tableau $T \in \mathsf{R}((n - (4+b), 2, 2, 1^b), (n - 2, 1, 1))$ has height b + 2, hence $\chi^{\lambda}((n - 2, 1, 1)) = (-1)^{b+2}$ for $\lambda = (n - (4+b), 2, 2, 1^b)$. Since $(2, 2) \subseteq \lambda$ and $f^{\lambda/(2,2)} \ge 1$, the coefficient of s_{λ} in $p_{(2,2,1^{n-4})}$ will cancel this potential negative.

4. DISCUSSION

In this paper, we studied various properties of the immersion and standard immersion poset, which are tightly linked to finite-dimensional irreducible polynomial representations of $GL_N(\mathbb{C})$ through their immersion pairs.

There are still many open questions to pursue in this line of research. It would be interesting to characterize all maximal elements in the immersion and standard immersion poset. In particular, a proof of Conjecture 2.14 seems in reach with the methods developed in this paper. In Corollary 3.30, we showed that for hook shapes λ and λ^t form an immersion pair. The same seems true for two column partitions. It would be interesting to classify when λ and its transpose form an immersion pair. In Corollary 3.31, we showed that the rank of the immersion poset is at least $\lfloor n/2 \rfloor$. It would be desirable to find better bounds for the rank.

33

Furthermore, it would be interesting to classify all intervals and chains in the immersion poset, in particular to obtain proofs of Conjectures 3.40 and 3.43. In view of the results of Section 3.5, the following question is natural.

Question 4.1. Which intervals $A_{\mu} := \{\lambda | (1^n) \leq_I \lambda \leq_I \mu\}$ in the immersion poset give rise to Schurpositivity of $p_{A_{\mu}}$?

Sundaram conjectured that all intervals $[(1^n), \mu]$ in reverse lexicographic order make (1.1) Schurpositive [Sun18, Conjecture 1], and has proven the conjecture for certain intervals [Sun19]. When $n \ge 5$, it appears that the immersion poset always contains some interval(s) which do not give rise to Schurpositivity. For example, $p_{A_{(n-1,1)}} = p_{(1^n)} + p_{(2,1^{n-2})} + p_{(n-1,1)}$ contains $-s_{(1^n)}$ when n is odd. This observation shows that the analog of [Sun18, Conjecture 1] is false for the immersion poset order. However, it does seem true that a large percentage of intervals A_{μ} in the immersion poset yield Schur-positivity. Using SAGEMATH [The24], we observe that when $6 \le n \le 9$ at least 91% of the intervals in the immersion poset make (1.1) Schur-positive. When n = 10, 11 the percentage of Schur-positive intervals drops to at least 81%, and when n = 18, the percentage is approximately 73.5%.

We conclude with some probabilistic and asymptotic questions.

Question 4.2. For randomly chosen partition $\lambda <_D \mu$, what is the probability that $\lambda \leq_I \mu$?

Based on computer evidence, we conjecture that the probability is near 0.5. For a partition λ of any size, consider the padded partition $\lambda[N] \coloneqq (N - |\lambda|, \lambda_1, \lambda_2, ...)$ of size N, where $N \ge |\lambda|$. For any two partitions $\lambda \le_D \mu$ (of any size), what can we say about $\lambda[N] \le_I \mu[N]$ for $N \gg 1$? Furthermore, it would be interesting to study the asymptotical behaviors of the (standard) immersion poset.

References

- [PPS23] Amrutha P, Amritanshu Prasad, and Velmurugan S. On the existence of elementwise invariant vectors in representations of symmetric groups. preprint, arXiv:2308.08146, 2023.
- [PPS24] Amrutha P, Amritanshu Prasad, and Velmurugan S. Cyclic characters of alternating groups. preprint, arXiv:2403.05109, 2024.
- [PR22] Dipendra Prasad and Ravi Raghunathan. Relations between cusp forms sharing Hecke eigenvalues. Represent. Theory, 26:1063–1079, 2022.
- [Sch07] J. Schur. Untersuchungen über die Darstellung der endlichen Gruppen durch gebrochene lineare Substitutionen. J. Reine Angew. Math., 132:85–137, 1907.
- [Sol61] Louis Solomon. On the sum of the elements in the character table of a finite group. Proc. Amer. Math. Soc., 12:962–963, 1961.
- [Sta99] Richard P. Stanley. Enumerative combinatorics. Vol. 2, volume 62 of Cambridge Studies in Advanced Mathematics. Cambridge University Press, Cambridge, 1999. With a foreword by Gian-Carlo Rota and appendix 1 by Sergey Fomin.
- [Sun18] Sheila Sundaram. The conjugacy action of S_n and modules induced from centralisers. J. Algebraic Combin., 48(2):179–225, 2018.
- [Sun19] Sheila Sundaram. On a positivity conjecture in the character table of S_n . Electronic J. Algebraic Combin., 26(1), 2019.
- [The24] The Sage Developers. SageMath, the Sage Mathematics Software System (Version 10.3), 2024. https://www.sagemath.org.
- [Wey39] Hermann Weyl. The Classical Groups. Their Invariants and Representations. Princeton University Press, Princeton, NJ, 1939.

(L. Johnston) DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA, ONE SHIELDS AVENUE, DAVIS, CA 95616-8633, U.S.A.

 $Email \ address: \verb"lisjohnston@ucdavis.edu"$

(D. Kenepp) DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA, ONE SHIELDS AVENUE, DAVIS, CA 95616-8633, U.S.A.

Email address: dskenepp@ucdavis.edu

(E. Nguyen) DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA, ONE SHIELDS AVENUE, DAVIS, CA 95616-8633, U.S.A.

 $Email \ address: \ {\tt evtnguyen@ucdavis.edu}$

(D. Paul) DEPARTMENT OF MATHEMATICS, INDIAN INSTITUTE OF SCIENCE, BANGALORE 560012, INDIA *Email address*: digjoypaul@iisc.ac.in

(A. Schilling) DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA, ONE SHIELDS AVENUE, DAVIS, CA 95616-8633, U.S.A.

Email address: anne@math.ucdavis.edu URL: http://www.math.ucdavis.edu/~anne

(M. C. Simone) DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA, ONE SHIELDS AVENUE, DAVIS, CA 95616-8633, U.S.A.

 $Email \ address: \ \tt mcsimone@ucdavis.edu$

(R. Zhou) DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA, ONE SHIELDS AVENUE, DAVIS, CA 95616-8633, U.S.A.

Email address: rgazhou@ucdavis.edu