

Gas velocity in the merging cluster Abell 2256 ¹

1 Merging clusters X-ray spectroscopy

1.1 Cluster dynamics and gas velocity

Galaxy clusters form hierarchically through collisions and mergers of smaller systems. Spatial and radial velocity distributions of member galaxies and X-ray observations of the intracluster medium (ICM) have revealed that some systems are still forming and un-relaxed. Sharp X-ray images revealed shocks and density discontinuities (“cold fronts”) suggesting supersonic or transonic gas motions, respectively. These heating processes would develop gas turbulence, cascading down from the driving scales to dissipative ones, which are determined by unknown viscosity. These are assumed to accelerate particles, reorder and amplify the magnetic fields, and generate diffuse radio halos and relics. To understand these energy flows, the gas motions in a range of scales should be measured.

1.1.1 X-ray gas velocity CCD measurements

The gas bulk motion can be measured most directly using X-ray line emission Doppler shift. These measurements have been challenging due to the limited energy resolutions of current X-ray instruments. Typical CCD energy resolution is about 130 eV (FWHM), compared with a possible energy shift of ~ 20 eV corresponding to a radial velocity of 1000 km s^{-1} , all at the Fe-K line energy here and hereafter.

[Tamura et al.(2014)] found a hint of gas bulk motion (v_{bulk}) at < 100 kpc west of the Perseus cluster center, where a cold front is seen in X-ray images. [Ota & Yoshida(2016)] reported *Suzaku* measurements of gas v_{bulk} in 8 clusters without significant signs of velocity variations. We also used *XMM* PN spectra to measure gas v_{bulk} maps in the Perseus and coma [Sanders et al.(2020)], A 3667 [Omiya et al.(2024)], and others.

1.1.2 X-ray imaging and *XMM* RGS limits

X-ray brightness (ICM density) fluctuation analyses are used to study gas dynamics (e.g., Zhuravleva et al. 2014). Furthermore, the *XMM* RGS are used to limit gas dynamics based on line broadening or resonant scattering.

1.2 High energy resolution spectroscopy

Hitomi Perseus results ([Hitomi Collaboration et al.(2018)]) : We used the Hitomi ~ 5 eV resolution spectra to resolve emission lines and measure the gas Doppler shifts and width. Within the central 100 kpc core, we found a line of sight (LOS) velocity gradient of $100 - 200 \text{ km s}^{-1}$. This could be due to the buoyant AGN bubble or central gas sloshing. A velocity dispersion (σ_{dis}) is also revealed. These velocities correspond to a galaxy scale potential and $< 10\%$ of the ICM thermal pressure. We also used resolved X-ray lines of various elements and ions to limit the non-Gaussianity of the line shape, test ion-electron thermal equilibrium, and resonant scattering. These observations are limited to the Perseus core.

¹submitted to the XRISM GO1 proposal. Takayuki TAMURA et al. April 15, 2024

From the *HITOMI* analysis, we learned that high resolution spectra include a wealth of information but are complex. For example, the line profiles depend not only on small to large-scale motions but also on the atomic and thermal nature of the ICM projected onto the telescope beam.

X-ray imaging studies are useful but require specific collision configurations to be applied to observations. In contrast, X-ray Doppler shift measurements are more direct and commonly applicable to merging systems. This method is sensitive to the motion parallel to the LOS. These two measurements are complementary and could provide a direct measurement of three-dimensional motion if applied to a merging system simultaneously.

1.3 Target selection and Abell 2256

General motivations: In almost all cases we found that X-ray gas redshift is consistent with those of optical galaxies around the corresponding region. However, previous measurements are limited mostly to X-ray bright and relaxed regions. In systems under a violent phase, we expect to find segregation in the location and velocity of the gas and galaxies. Such detachments in the sky position have been observed in some merging systems such as the Bullet cluster.

To understand cluster formation, which is dominated by non-linear processes, systematic measurements in a sample of clusters are required. In [Kitayama et al.(2014)], we examined and proposed various types of bright targets for ASTRO-H studies of clusters. Five merging clusters were selected, including *XRISM* PV targets, Coma, and A 2319.

1.3.1 The Abell 2256 cluster

From targets in [Kitayama et al.(2014)] and recent literature to study gas dynamics in a different environment, we select an X-ray bright cluster A 2256 ($z=0.056$). The past observations revealed that this system has recently experienced a strong merger shock and is at a more violent merger stage than Coma. This is the most suitable for the *XRISM* spectral mapping of gas motions and related physics, as explained below. **Double peaked structures in X-ray image and galaxy dynamics:** At the center, there are two substructures in the X-ray surface brightness separating by $3'.5$ in the sky. The two are separated by $\sim 2000 \text{ km s}^{-1}$ in radial velocity peaks of member galaxies as well. These angular and velocity separations are unique in this system. Almost all other X-ray bright mergers show LOS velocities smaller than this (e.g., 700 km s^{-1} in the Coma, 100 km s^{-1} in A3667).

1.3.2 *Suzaku* discovery of gas bulk motion

Using *Suzaku* CCD, we found a radial velocity difference between the main and subsystems to be 1500 ± 300 (statistical) ± 300 (systematic) km s^{-1} . See Fig.3. This is the first detection of gas bulk motion in a cluster. Except for this, Coma cluster [Sanders et al.(2020)], and *HITOMI* Perseus given above, no clear detection of the gas motion is found.

We found that the X-ray determined absolute redshifts of and hence the difference between the main and sub components are consistent with those of member galaxies in optical. In fact, the X-ray redshift of the main component, is the same as the galaxy redshift within the error. These suggests that the X-ray emitting gas is moving together with galaxies as a substructure. In addition to these, we found a hint of X-ray gas velocity shift with respect to the galaxy (G2) around the sub-component at about 500 km s^{-1} . Note that this difference is within the combined errors from X-ray statistical (150 km s^{-1}), systematic (300 km s^{-1}), and optical fitting (160 km s^{-1}).

1.3.3 Recent deep X-ray and radio observations

Chandra observations revealed detailed gas structures in and around the main (P1 = CORE), second (P2 = NW-SUB), and fainter (P3) peaks. [Ge et al.(2020)] analyzed the X-ray temperature, metallicity, and density map, showing clear evidence of a series of merging activities. They examined these X-ray and radio features and discussed a scenario including an infalling gas from NW to the central primary. We aim to probe LOS motions associated with this collision. Radio observations revealed an NW relic (the 2nd brightest one among all known systems) and a faint halo. Several head tail galaxies (B, C, I) also indicates the dynamic interaction between gas and galaxies. We are fascinated by galaxy C, which has a tail extending > 500 kpc, but not completely aligned with the merger direction. [Miller et al.(2003)] suggested that since the tail emission is not disrupted the region would have low turbulence. To test this, we measure the X-ray line width (§ 2.1).

2 Science Goals and Observation Plan

2.1 Resolve Measurement goals

2.1.1 line shifts

We measure a map of X-ray line redshift and search for gas v_{bulk} structures associated with cluster formation. By proposed *XRISM* observations, we aim to improve and advance previous CCD results given above by not only reducing velocity errors (§ 2.4.2) but also resolving velocity structure spatially.

Does the gas move together with galaxies in a merging cluster? We compare the obtained X-ray velocities with those of galaxies and examine if the gas components around the merging core are moving together with galaxies.

We use the gas v_{bulk} map around central galaxies to directly constrain critical merging parameters, including the direction of merger axis and the angular moment. There are radial velocity differences among bright galaxies, G1, G2, and G3, indicating the merger axis to be not in the sky plane. Radio relic shape and polarization fraction also indicate a tilted direction (van Weeren et al. 2012). This cluster has the largest shift in optical radial velocities (§ 1.3.1). Therefore we expect to discover the largest angular moment in the cluster plasma.

2.1.2 Line profiles and turbulent motions

We resolve line profiles to measure the velocity dispersion (σ_{dis}), which is essentially the 1D LOS component of the characteristic velocity of isotropic turbulence.

Along with gas bulk, turbulent motion is assumed to play an important role in the galactic and larger scales (§ 1.1). Nevertheless, turbulence has been more challenging to measure and hence to model theoretically. *HITOMI* Perseus spectroscopy revealed a velocity dispersion (σ_{dis}) peaked toward the central AGN at 200 km s^{-1} and being uniform at 100 km s^{-1} around the core. Given the gas super-sonic motion found with *Suzaku* and other active gas features given above, do we expect a large σ_{dis} around merging systems? To answer this, we search for gas bulk and turbulent motions simultaneously across the violent collision site. We compare A 2256 motions with those in other places such as completely relaxed ones, and systematically characterize these motions.

2.2 Pointing plan

We aim to cover a central $6'$ (400 kpc) region. We estimated spectral parameters from the *Suzaku* Fe-line map and *Chandra* spectral fit [Sun et al.(2002)]. We expect the NW-SUB region to provide Fe line flux comparable to the core one.

2.3 EXTEND analysis

We use the Extend data to map thermal structure in a large area and wide energy range. We search for a hotter (> 10 keV) thermal and non-thermal emission components. These spectra are used to constrain the inverse-Compton emission and hence limit the magnetic field in a cluster scale along with radio non-thermal (radio relic) imaging. [Nakazawa et al.(2009)] demonstrated these analysis in a merging cluster A 3667.

2.4 Visibility and technical feasibility

2.4.1 Visibility and roll angle

The target is visible over the GO phase with a large freedom of roll angle. We do not constrain the roll angle. For a given observation date, we will align the Resolve angle to the cluster merger axis to cover the largest scale as much as possible.

2.4.2 Statistical limit and exposure time request

The statistical accuracy of the line shift and broadening depends on intrinsic cluster line width (σ_{dis}). Ota et al. (2018) showed that > 200 Fe line counts are required to achieve an accuracy better than 20% in the gas σ_{dis} , when $\sigma_{\text{dis}} < 200 \text{ km s}^{-1}$. We checked these errors using the *XRISM* response files. Based on these and the spectral parameters of the proposed regions from the *Suzaku* spectra, we simulated *XRISM* observations, examined feasibility, and planned the observations.

Table 1: Expected accuracy of X-ray gas velocities (1σ statistical) from [Ota et al.(2018)]. We assume thermal 5–8 keV CIE emission.

Fe Line (counts/100ks)	σ_{dis} (km/s)	$\Delta\sigma_{\text{dis}}$ (km/s)	Δv_{bulk} (km/s)
200	0	30	30
200	200	30	75
50	0	50	30
50	200	50	100

2.5 Systematic limits

2.5.1 Instrumental and astronomical errors

In [Hitomi Collaboration et al.(2018)] we evaluated instrumental calibration effects on the velocity measurements. We expect the Resolve energy scale accuracy to be reduced down to 1 eV (goal, $\sim 50 \text{ km s}^{-1}$). These levels of systematic can be comparable when the cluster σ_{dis} is significantly larger than the instrumental energy resolution ($\sigma \sim 2 \text{ eV}$). Note that energy scale variations among

observations (integrated over the FOV) and pixels can be important. Note also that (1) absolute energy is critical to compare with other instruments, and (2) relative one is to measure astronomical variations.

Based on the *HITOMI* analysis, we expect other errors to be considered. Those include line spread function (< 0.5 eV), spectral and atomic modeling errors (< 1 eV).

2.5.2 Spatial-spectral mixing

Within the proposed region, we aim to measure v_{bulk} and σ_{dis} structures in $0'.5 - 1'.5$ and $3'$ (Resolve FOV) spacial resolutions, respectively, considering the pixel size of $0'.5$ and angular resolution of $\sim 1'.3$ HPD. Because of a relatively flat brightness distribution in this cluster, the PSF scattering is less severe than that of centrally-peaked cool core clusters. We model spacial mixing using existing high spatial resolution X-ray data. We will use a simultaneous fitting method that considers PSF mixing to reconstruct velocity maps as we demonstrated for the *HITOMI* Perseus analysis.

2.6 Dark matter and cosmology

In clusters gas, galaxies, and associated interactions are dynamically controlled by dark matter gravity. Based on this assumption, we have studied these structure formations. In this proposal, we focus on dark matter in an unrelated merging system and compare it with those in a relaxed one. Is the gas trapped at the bottom of the cluster potential or at those of the central galaxy (cD) one? There is a redshift difference at about 400 km s^{-1} . between the cluster mean and cD (A 2256 G1), which is within the feasibility above.

Accurate measurements of dark matter distribution are required for precise cosmology. Energies in gas motions could be the key uncertainty to calibrating total mass distribution, as we discussed in [Kitayama et al.(2014)]. From the Perseus core, *HITOMI* revealed little ($< 10\%$) deviation from hydrostatic equilibrium. We aim to test this basic assumption in a merging cluster.

References

- [Berrington et al.(2002)] Berrington, R. C., Lugger, P. M., & Cohn, H. N. 2002, *Astron. J.*, 123, 2261. doi:10.1086/339840
- [Briel et al.(1991)] Briel, U. G., et al. 1991, *Astron. Astrophys.*, 246, L10
- [Clarke & Ensslin(2006)] Clarke, T. E. & Ensslin, T. A. 2006, *Astron. J.*, 131, 2900. doi:10.1086/504076
- [Gatuzz et al.(2022)] Gatuzz, E., Sanders, J. S., Dennerl, K., et al. 2022, *Monthly Notices Roy. Astron. Soc.*, 511, 4511. doi:10.1093/mnras/stab2661
- [Gatuzz et al.(2022)] Gatuzz, E., Sanders, J. S., Canning, R., et al. 2022, *Monthly Notices Roy. Astron. Soc.*, 513, 1932. doi:10.1093/mnras/stac846cen
- [Ge et al.(2020)] Ge, C., Liu, R.-Y., Sun, M., et al. 2020, *Monthly Notices Roy. Astron. Soc.*, 497, 4704. doi:10.1093/mnras/staa2320
- [Hitomi Collaboration et al.(2018)] Hitomi Collaboration, Aharonian, F., Akamatsu, H., et al. 2018, *Publ. Astron. Soc. Japan*, 70, 11. doi:10.1093/pasj/psy004

- [Hitomi Collaboration et al.(2018)] Hitomi Collaboration, Aharonian, F., Akamatsu, H., et al. 2018, *Publ. Astron. Soc. Japan*, 70, 12. doi:10.1093/pasj/psx156
- [Hitomi Collaboration et al.(2018)] Hitomi Collaboration, Aharonian, F., Akamatsu, H., et al. 2018, *Publ. Astron. Soc. Japan*, 70, 9. doi:10.1093/pasj/psx138
- [Kitayama et al.(2014)] Kitayama, T., Bautz, M., Markevitch, M., et al. 2014, arXiv:1412.1176. doi:10.48550/arXiv.1412.1176
- [Ota et al.(2018)] Ota, N., Nagai, D., & Lau, E. T. 2018, *Publ. Astron. Soc. Japan*, 70, 51. doi:10.1093/pasj/psy040
- [Ota & Yoshida(2016)] Ota, N. & Yoshida, H. 2016, *Publ. Astron. Soc. Japan*, 68, S19. doi:10.1093/pasj/psv128
- [Omiya et al.(2024)] Omiya, Y., Nakazawa, K., Tamura, T., et al. 2024, arXiv:2403.10150. doi:10.48550/arXiv.2403.10150
- [Markevitch & Vikhlinin(2007)] Markevitch, M. & Vikhlinin, A. 2007, *Physics Report*, 443, 1. doi:10.1016/j.physrep.2007.01.001
- [Miller et al.(2003)] Miller, N. A., Owen, F. N., & Hill, J. M. 2003, *Astron. J.*, 125, 2393. doi:10.1086/374628
- [Nakazawa et al.(2009)] Nakazawa, K., Sarazin, C. L., Kawaharada, M., et al. 2009, *Publ. Astron. Soc. Japan*, 61, 339. doi:10.1093/pasj/61.2.339
- [Sun et al.(2002)] Sun, M., Murray, S. S., Markevitch, M., et al. 2002, *Astrophys. J.*, 565, 867. doi:10.1086/324721
- [Sanders et al.(2020)] Sanders, J. S., Dennerl, K., Russell, H. R., et al. 2020, *Astron. Astrophys.*, 633, A42. doi:10.1051/0004-6361/201936468
- [Tamura et al.(2014)] Tamura, T., Yamasaki, N. Y., Iizuka, R., et al. 2014, *Astrophys. J.*, 782, 38. doi:10.1088/0004-637X/782/1/38
- [Tamura et al.(2011)] Tamura, T., Hayashida, K., Ueda, S., et al. 2011, *Publ. Astron. Soc. Japan*, 63, S1009. doi:10.1093/pasj/63.sp3.S1009

Figure 1: Gas and galaxies velocities from observations and XRISM/Resolve simulations. Resolve errors bars are statistical one ($\pm 80 \text{ km s}^{-1}$).

Figure 2: Simulated *XRISM* spectra of the A2256 core component. Input σ_{dis} are 0 km s^{-1} (red histogram) and 200 km s^{-1} (black points).