STRONGLY ROBUSTNESS OF TORIC IDEALS OF WEIGHTED ORIENTED CYCLES SHARING A SINGLE VERTEX

RAMAKRISHNA NANDURI* AND TAPAS KUMAR ROY[†]

ABSTRACT. In this article, we study the strongly robust property of toric ideals of weighted oriented graphs. Let D be a weighted oriented graph consists of weighted oriented cycles (balanced or unbalanced) sharing a single vertex v and D' be a weighted oriented graph consists of D and a finite number of disjoint cycles such that each of these cycles is connected by a path at the sharing vertex v of D. Then we show that the toric ideals $I_D, I_{D'}$ of D and D' respectively, are strongly robust and hence robust. That is, for the toric ideal I_D , of D, its Graver basis is a minimal generating set of I_D . We explicitly give a unique minimal generating set of primitive binomials of I_D . We explicitly describe the circuit binomials in any weighted oriented graph.

1. INTRODUCTION

Toric ideals are important in the modern theory of Algebra and impacted within a variety of research areas including Commutative Algebra, Algebraic Geometry, Combinatorial Optimization and Algebraic Coding Theory (see [23], [28, Chapter 8], [21]). For any $n \times m$ matrix $A = [\mathbf{a_1} \dots \mathbf{a_m}]$ with non-negative integers, let $M = (\mathbf{x^{a_1}}, \dots, \mathbf{x^{a_m}})$ be the monomial ideal in $R = K[x_1, \dots, x_n]$, where K is a field. Let $S = K[e_1, \dots, e_m]$ be a polynomial ring the variables e_1, \dots, e_m . Define a ring homomorphism $\varphi : S \to R$, as $\varphi(e_i) = \mathbf{x^{a_i}}$. Then the kernel of φ is called the *toric ideal* associated to A or M and we denote by I_A . Note that S/I_A is the affine semigroup ring corresponding to the semigroup $\mathbb{Z}_+\mathbf{a_1} + \cdots + \mathbb{Z}_+\mathbf{a_m}$. It is well known that I_A is a prime ideal generated by binomials $\mathbf{e^u} - \mathbf{e^v}$ such that $\deg_A(\mathbf{e^u}) = \deg_A(\mathbf{e^v})$, where $\deg_A(\mathbf{e^u}) = u_1\mathbf{a_1} + \cdots + u_n\mathbf{a_n}$.

A binomial $\mathbf{e}^{\mathbf{u}} - \mathbf{e}^{\mathbf{v}}$ in I_A is called a *primitive* binomial if there exists no other binomial $\mathbf{e}^{\mathbf{u}'} - \mathbf{e}^{\mathbf{v}'}$ in I_A such that $\mathbf{e}^{\mathbf{u}'}|\mathbf{e}^{\mathbf{u}}$ and $\mathbf{e}^{\mathbf{v}'}|\mathbf{e}^{\mathbf{v}}$. The set of primitive binomials in I_A is called the *Graver basis* of I_A and denoted by Gr_A . Minimal system of binomial generators of

[†] Supported by PMRF research fellowship, India.

^{*} Corresponding author and supported by SERB grant No: CRG/2021/000465, India.

AMS Classification 2020: 13F65, 13A70, 05E40, 05C38, 05C22.

toric ideals have been studied by many authors (see [23], [4]). It is a demanding problem in Algebraic Statistics, when toric ideals have unique minimal system of binomial generators (see [26]). To study this problem many authors defined various notions of binomial generators of toric ideals. In [4], an explicit combinatorial formula for the number of minimal systems of binomial generators of toric ideals was given. The notion of robustness of toric ideals has been studied in several papers. A toric ideal I_A is said to be *robust* if it is minimally generated by its universal Gröbner basis. The universal Gröbner basis of I_A is denoted by \mathcal{U}_A and is defined as the union of all reduced Gröbner bases of I_A with respect to all possible term orders. The robust toric ideals have been studied in [1] for simple graphs, in [3] for toric ideals generated by quadratic binomials, and in [7] for toric graph ideals generated by binomials. Nevertheless, robustness is a rare property and in literature we know few nontrivial examples of robust ideals, see [3, 7]. A weaker version of robust ideals known as generalized robust ideals have been studied (see [25]). A toric ideal I_A is said to be generalized robust if its universal Gröbner basis \mathcal{U}_A is equal to its universal Markov basis. A Markov basis of I_A is a minimal binomial generating set of I_A and the universal Markov basis of I_A is the union of all its minimal sets of binomial generators and we denote by \mathcal{M}_A . A toric ideal is said to be strongly robust if its Graver basis equals the union of all minimal sets of binomial generators \mathcal{M}_A of I_A , equivalently, Gr_A equals the set of all indispensable binomials of I_A . A binomial $f = \mathbf{e}^{\mathbf{u}} - \mathbf{e}^{\mathbf{v}}$ in I_A is called indispensable if every system of binomial generators of I_A contains f or -f. Generalized robust toric ideals have been studied in [24] for codimension 2 toric ideals, and in [25] for toric ideals of simple graphs. In fact, the notions robust and strongly robust are coincide for toric ideals of simple graphs (see [25]). See [22, 24, 11] for some family of strongly robust toric ideals. In this work we study the minimal systems of binomial generators of toric ideals weighted oriented graphs. In particular, we study when such toric ideals have unique minimal set of binomial generators. Nevertheless, the notions of robust, generalized robust and strongly robust properties of toric ideals of weighted oriented graphs are completely unknown. However the toric ideals of edge ideals of simple graphs (simply we call as toric graph ideals) are broadly studied and many algebraic and combinatorial properties and characterizations of such ideal are explored enough in the literature (see [17, 18, 27, 15, 12]).

A weighted oriented graph D (shortly we write as WOG) is a multiple object D = (V(D), E(D), w), where $V(D) = \{x_1, \ldots, x_n\}$ is the (finite) set of vertices of D, E(D) is

the set of edges of D (without multiple edges nor loops),

$$E(D) = \{(x_i, x_j) : \text{there is an edge from } x_i \text{ to } x_j\}$$

and w is a weight function $w : V(D) \to \mathbb{N}$, which puts a weight to each vertex of D. That is, each edge is specified a orientation and each vertex is given a weight. WOGs are significant because of their uses in various research fields (see for example [13, 5, 8, 19, 29, 9, 10]). The edge ideal of D is defined as the ideal in R, $I(D) = (x_i x_j^{w_j} : (x_i, x_j) \in E(D))$. The toric ideal of I(D), simply we write as I_D , is the toric ideal corresponding to the incidence matrix A(D) (see Definition 2.1) of D. We call I_D , the toric ideal of D. The toric ideal I_D has been studied in [2] and showed when it is principal and also they summarized the complexity about unhampered number of generators (or their exponents) for weighted oriented cycles (balanced or unbalanced or odd) sharing a vertex. Recently, in [16], the authors give an explicit formulas for the generators of toric ideals of any number of balanced cycles sharing a vertex and toric ideals of any two weighted oriented cycles (balanced or odd) sharing a vertex. Let $\mathcal{M}_D, \mathcal{U}_D, \mathcal{G}_{T_D}$ denote the universal Markov basis, universal Gröbner basis and Graver basis of I_D respectively. A binomial $\mathbf{e^u} - \mathbf{e^v}$ in I_A is called a *circuit* of I_D if it has minimal support with respect to set inclusion. The set of circuits of I_D is denoted by \mathcal{C}_D . Moreover it is known that

$$\mathcal{C}_D \subseteq \mathcal{U}_D \subseteq Gr_D$$
 and $\mathcal{M}_D \subseteq Gr_D$.

If D is a WOG of any number of balanced cycles sharing a vertex, then by [16, Theorem 3.1], we have that I_D is a circuit ideal, that is I_D is generated by C_D and hence it is strongly robust and robust as well. Nevertheless the toric ideal of any two oriented cycles (balanced or unbalanced or odd) is principal and whose generator is explicitly computed (see [16]). Thus the first nontrivial class of D to study the robustness of I_D is three oriented cycles sharing a vertex.

In this work, our aim is to study strongly robustness of toric ideals of WOGs. We show that the toric ideals of a WOG D of finite oriented cycles sharing a vertex, and a WOG D' consists of above D and a finite number of disjoint cycles such that each of these cycles is connected by a path at the sharing vertex of cycles of D, are strongly robust (Theorems 3.11, 3.12). We explicitly compute the unique minimal generating set M_D consists of primitive binomials, of the toric ideal I_D of D consists of three oriented cycles (balanced or unbalanced or odd) sharing a common vertex (see Theorems 3.13, 3.15, 3.16, 3.17). That is, $M_D = \mathcal{U}_D = Gr_D$. We characterize the circuit binomials in any WOG (see Proposition 4.2). If a WOG D consists of finitely many oriented cycles sharing a vertex such that at most two among these cycles are unbalanced, then we show that \mathcal{C}_D is a generating set of I_D and hence $\mathcal{C}_D = \mathcal{M}_D = \mathcal{U}_D = Gr_D$ (see Propositions 4.3, 4.4, 4.5, 4.6). We characterize primitive binomials of a WOG consists of n unbalanced cycles sharing a vertex. Further if D is a WOG of finitely many oriented cycles sharing a common vertex, then we describe primitive binomials in I_D (Theorem 4.8). Unlike the case of two oriented cycles sharing a vertex, I_D can have many minimal binomial generators whose support is E(D). For robust toric ideals I_D , we have that the multigraded Betti numbers of I_D and $\operatorname{in}_<(I_D)$ are equal, for any term order <. By using our main results, we give some interesting applications to the case that V^+ are sink vertices. If D consists of n unbalanced (even or odd) cycles, then we show that $I_D = I_G$, if V^+ are sink vertices (Proposition 4.14) or each vertex of D is either a source or a sink vertex (Proposition 4.7). We observe some class of non-trivial graphs where $\mathcal{C}_D = \mathcal{M}_D = \mathcal{U}_D = Gr_D$ (see 4.9, 4.11).

We constructed the paper as follows. In the section 2, we recall the definitions and some results needed for the rest of the paper. In the section 3, first we characterized WOGs Dwhose toric ideal I_D is generic (Theorem 3.2). We show that I_D is strongly robust, if Dis a WOG of three oriented cycles sharing a vertex by explicitly giving a unique minimal generating set of primitive binomials of I_D . Let D be a WOG consists of weighted oriented cycles (balanced or unbalanced) sharing a single vertex v and D' be a weighted oriented graph consists of D and a finite number of disjoint cycles such that each of these cycles is connected by a path at the sharing vertex v of D. Then we show that the toric ideals $I_D, I_{D'}$ of D and D' respectively, are strongly robust. In the section 4, we characterize circuit binomials in any WOG and we showed some WOGs which are generated by circuit binomials. Finally, we describe primitive binomials in a WOG consists of finitely many unbalanced cycles.

2. Preliminaries

Let $R = K[x_1, \ldots, x_n]$, where K is a field. Recall that the *support* of a binomial $f = \prod_{k=1}^{m} e_k^{p_k} - \prod_{k=1}^{m} e_k^{q_k}$, denoted as supp(f), is the set of variables e_k which appear in f with non-zero exponent. Also, support of a vector $\mathbf{a} \in \mathbb{Z}_+^n$, is defined as $supp(\mathbf{a}) := \{i : a_i \neq 0\}$.

A (vertex) weighted oriented graph is a triplet $D = (V(D), E(D), \mathbf{w})$, where $V(D) = \{x_1, \ldots, x_n\}$ is the vertex set of D,

$$E(D) = \{(x_i, x_j) : \text{there is an edge from } x_i \text{ to } x_j\}$$

is the edge set of D and the weight function $\mathbf{w} : V(D) \to \mathbb{N}$. We simply denote the weight function \mathbf{w} by the vector $\mathbf{w} = (w_1, \dots, w_n)$, where $w_j = w(x_j)$ for each j. If $e = (x_i, x_j)$ is an edge of D, then we call x_j as *head* of e and x_i is called *tail* of e. The *indegree* of v_i is defined as $|\{v_j \in V(D) : (v_j, v_i) \in E(D)\}|$ and the *outdegree* of v_i is defined as $|\{v_j \in V(D) : (v_i, v_j) \in E(D)\}|$. The *edge ideal* of D is defined as the ideal

$$I(D) = \left(x_i x_j^{w_j} : (x_i, x_j) \in E(D)\right)$$

in R. Let $V(D) = \{x_1, \dots, x_n\}$ and $E(D) = \{e_1, \dots, e_m\}.$

Definition 2.1. The incidence matrix of D is an $n \times m$ matrix whose $(i, j)^{th}$ entry $a_{i,j}$ is defined by

$$a_{i,j} = \begin{cases} 1 & \text{if } e_j = (x_i, x_l) \in E(D) \text{ for some } 1 \le l \le n, \\ w_i & \text{if } e_j = (x_l, x_i) \in E(D) \text{ for some } 1 \le l \le n, \\ 0 & \text{otherwise}, \end{cases}$$

and we denote by A(D).

The toric ideal of D is defined as the toric ideal of the incidence matrix A(D) or the edge ideal I(D) and we shortly denote as I_D . A weighted oriented r-cycle denoted by C_r is a weighted oriented graph whose underlying graph is a cycle of length r and the edges oriented in any direction (need not be naturally oriented). Recall that a weighted oriented even cycle C_m on m vertices is said to be balanced if $\prod_{k=1}^m a_{k,k} = a_{1,m} \prod_{k=1}^m a_{k+1,k}$, where $A(C_m) = [a_{i,j}]_{m \times m}$. We denote Null(A(D)), the null space of A(D) over \mathbb{Q} .

A weighted oriented cycle is called naturally oriented if edges are either oriented in clockwise direction or in anti-clockwise direction.

Definition 2.2. For a vector $\mathbf{b} = ((-1)^{p_1}b_1, (-1)^{p_2}b_2, \dots, (-1)^{p_l}b_l) \in \mathbb{Z}^l$, with $p_i \ge 1, b_i \ge 0$ integers, define the corresponding binomial in the variables e_1, \dots, e_l as $f_{\mathbf{b}} := f_{\mathbf{b}}^+ - f_{\mathbf{b}}^-$, where

$$f_{\mathbf{b}}^{+} = \prod_{i=1 \ (p_i \ even)}^{l} e_i^{b_i}, \ and \ f_{\mathbf{b}}^{-} = \prod_{i=1 \ (p_i \ odd)}^{l} e_i^{b_i}$$

For any vector $\mathbf{b} \in \mathbb{Z}^l$, we can always write $\mathbf{b} = \mathbf{b}_+ - \mathbf{b}_-$ uniquely, where $\mathbf{b}_+, \mathbf{b}_- \in \mathbb{Z}^l_+$. Note that $f_{\mathbf{m}}^+ = \mathbf{e}^{\mathbf{m}_+}$ and $f_{\mathbf{m}}^- = \mathbf{e}^{\mathbf{m}_-}$.

Definition 2.3. For $\mathbf{a}, \mathbf{b} \in \mathbb{Z}^l$, we define $\mathbf{a} \prec \mathbf{b}$, if $supp(\mathbf{a}_+) \subseteq supp(\mathbf{b}_+)$ and $supp(\mathbf{a}_-) \subseteq supp(\mathbf{b}_-)$. For binomials $f_{\mathbf{a}}, f_{\mathbf{b}}$, we define $f_{\mathbf{a}} \prec f_{\mathbf{b}}$. if $\mathbf{a} \prec \mathbf{b}$, equivalently, $supp(f_{\mathbf{a}}^+) \subseteq supp(f_{\mathbf{b}}^+)$ and $supp(f_{\mathbf{a}}^-) \subseteq supp(f_{\mathbf{b}}^-)$.

Let $[\mathbf{b}]_i$ denotes the *i*-th entry of **b**. Also, $f_{\mathbf{m}} \in I_D$ implies that $\mathbf{m} \in \text{Null}(A(D))$ and $[\mathbf{m}]_i$ denotes the *i*-th entry of **m** corresponding to the edge $e_i \in E(D)$. For any $S \subseteq \{1, \ldots, l\}$ and $\mathbf{a} \in \mathbb{Z}^l$, we denote $\mathbf{a}|_S$ be the vector in $\mathbb{Z}^{|S|}$ such that $[\mathbf{a}|_S]_i = [\mathbf{a}]_i$ for all $i \in S$. Throughout we use minimal element in \mathbb{Z}^2 , with respect to the following partial order: $(a, b) \leq (c, d)$, if $a \leq c$ and $b \leq d$.

Theorem 2.4. [2, Theorem 5.1]. Let D be weighted oriented graph consists of two unbalanced cycles share a vertex. Then I_D is a principal ideal.

Lemma 2.5. Let $\mathbf{a} = ((-1)^{p_i} a_i)_{i=1}^l, \mathbf{b} = ((-1)^{q_i} b_i)_{i=1}^l \in \mathbb{Z}^l$, where $p_i, q_i \in \mathbb{N}, a_i, b_i \in \mathbb{N} \cup \{0\}$ such that $supp(\mathbf{a}) \cap supp(\mathbf{b}) = \{z_1, z_2, \dots, z_{l_1}\} = S$. Let $\lambda_1, \lambda_2 \in \mathbb{N}$.

(i) $f_{\lambda_1 \mathbf{a}} \in (f_{\mathbf{a}})$.

(ii) Suppose $(-1)^{p_{z_i}} = (-1)^{q_{z_i}}$ for each $i \in \{1, 2, \dots, l_1\}$. Then $f_{\lambda_1 \mathbf{a} + \lambda_2 \mathbf{b}} \in (f_{\mathbf{a}}, f_{\mathbf{b}})$.

Proof. (i) $f_{\lambda_1 \mathbf{a}} = (f_{\mathbf{a}}^+)^{\lambda_1} - (f_{\mathbf{a}}^-)^{\lambda_1} = (f_{\mathbf{a}}^+ - f_{\mathbf{a}}^-)h_1 = f_{\mathbf{a}}h_1$ for some polynomial h_1 . Thus $f_{\lambda_1 \mathbf{a}} \in (f_{\mathbf{a}})$.

(ii) We have $f_{\lambda_1 \mathbf{a} + \lambda_2 \mathbf{b}} = f_{\lambda_1 \mathbf{a}} f_{\lambda_2 \mathbf{b}}^+ + f_{\lambda_1 \mathbf{a}}^- f_{\lambda_2 \mathbf{b}} \in (f_{\mathbf{a}}, f_{\mathbf{b}})$ as $f_{\lambda_1 \mathbf{a}}, f_{\lambda_2 \mathbf{b}}$ are multiples of $f_{\mathbf{a}}, f_{\mathbf{b}}$ respectively.

Below proposition is graph version of [23, Proposition 4.13]. Note that A(H) is a matrix obtained from A(D) by deleting some columns.

Proposition 2.6. Let H be a oriented subgraph of a weighted oriented graph D such that V(D) = V(H). Then

- (i) $I_H = I_D \cap K[e_i : e_i \in E(H)],$
- (ii) $\mathcal{C}_H = \mathcal{C}_D \cap K[e_i : e_i \in E(H)],$
- (iii) $\mathcal{U}_H = \mathcal{U}_D \cap K[e_i : e_i \in E(H)],$
- (iv) $Gr_H = Gr_D \cap K[e_i : e_i \in E(H)].$

Below we prove a lemma which we use in many proofs.

Lemma 2.7. Let D be any weighted oriented graph and $f_{\mathbf{m}} \neq 0 \in I_D$. Let $v \in V(D)$ of degree n. If (n-1) edges of D incident with v are not in $supp(f_{\mathbf{m}})$, then the other edge incident with v is not in $supp(f_{\mathbf{m}})$. Moreover if the edge e_i incident with v belongs to $supp(f_{\mathbf{m}}^+)($ or $supp(f_{\mathbf{m}}^-))$, then there exists an edge e_j incident with v belongs to $supp(f_{\mathbf{m}}^-)($ or $supp(f_{\mathbf{m}}^+))$.

Proof. Note that $\mathbf{m} \in \text{Null}(A(D))$. Let e_1, e_2, \ldots, e_n be the edges precisely incident with v. Suppose $e_1, e_2, \ldots, e_{i-1}, e_{i+1}, \ldots, e_n \notin \text{supp}(f_{\mathbf{m}})$. Then $[\mathbf{m}]_k = 0$ for $k = 1, 2, \ldots, i - 1, i + 1, \ldots, n$. Then from the equation $A(D)\mathbf{m} = \mathbf{0}$, we get that $[\mathbf{m}]_i = 0$. This implies that $e_i \notin \text{supp}(f_{\mathbf{m}})$, as required.

Suppose $e_i \in \operatorname{supp}(f_{\mathbf{m}}^+)(\operatorname{or} \in \operatorname{supp}(f_{\mathbf{m}}^-))$. Then $[\mathbf{m}]_i > 0(\operatorname{or} [\mathbf{m}]_i < 0)$. From the equation $A(D)\mathbf{m} = \mathbf{0}$, we get $\sum_{k=1}^{i-1} [\mathbf{m}]_k x_k + [\mathbf{m}]_i x_i + \sum_{k=i+1}^n [\mathbf{m}]_k x_k = 0$ for some positive integers x_k 's. This implies that there exists j such that $[\mathbf{m}]_j < 0(\operatorname{or} [\mathbf{m}]_j > 0)$. Thus $e_j \in \operatorname{supp}(f_{\mathbf{m}}^-)(\operatorname{or} \operatorname{supp}(f_{\mathbf{m}}^+))$.

3. Strongly robustness of toric ideal of three unbalanced cycles sharing a vertex

In this section, first we characterize WOGs D those toric ideal I_D is generic. We show that the toric ideals of a WOG D of finite oriented cycles sharing a vertex, and a WOG D'consists of above D and a finite number of disjoint cycles such that each of these cycles is connected by a path at the sharing vertex of cycles of D, are strongly robust and the toric ideals have unique minimal generating sets (by [25, Theorem 5.10]) consists of primitive binomials. We explicitly compute this minimal generating set of I_D if D is a WOG of three oriented cycles (even or odd or balanced or unbalanced) sharing a vertex.

Definition 3.1. [20] Let D be weighted oriented graph and E(D) be its edge set. Then the toric ideal I_D is called generic if supp(f) = E(D), for all $f \in M_D$.

Theorem 3.2. The toric ideal I_D of a weighted oriented graph D is generic if and only if D is one of below type graphs:

- (i) D is a balanced cycle.
- (ii) D consists of precisely two unbalanced cycles share a vertex.
- (iii) D consists of precisely two unbalanced cycles C_1, C_2 share a path P such that the induced cycle $D \setminus (P \setminus end points of P)$ is unbalanced.

RAMAKRISHNA NANDURI AND TAPAS KUMAR ROY

(iv) D consists of precisely two unbalanced cycles connected by a path.

Proof. Suppose I_D is generic. Then D cannot have any leaf vertices because edge incident with leaf cannot appear in any element of minimal generating set of I_D . Since $I_D \neq (0)$, D is not an unbalanced cycle. Suppose D is none of above type required graphs. Then either D has an proper balanced cycle or D has no proper balanced cycles.

<u>Case-1</u>: Suppose D has an proper balanced cycle C. If $f_{\mathbf{c}}$ is not indispensable, then there exists a balanced cycle C_1 in D such that $f_{\mathbf{c}_1}^+|f_{\mathbf{c}}^+$ and there is another cycle C_2 in D where C_1, C_2 share an edge e and $E(C) = (E(C_1) \cup E(C_2)) \setminus \{e\}$. If $f_{\mathbf{c}_1}$ is not indispensable, we continue as above. As the graph D has finite number of edges, continuing in this way, after finite number of steps, we see that there is proper induced balanced cycle C' in Dsuch that $f_{\mathbf{c}'}$ is indispensable. This implies that $f_{\mathbf{c}'} \in M_D$ where $|\operatorname{supp}(f_{\mathbf{c}'})| < |E(D)|$ which contradicts the fact that I_D is generic.

<u>Case-2</u>: Suppose D has no proper balanced cycles. Then there is proper subgraph H of D such that H consists of two unbalanced cycles share a vertex or two unbalanced cycles connected by a path or two unbalanced cycles share a path and for any $f \in M_D$ with $\operatorname{supp}(f) = E(D)$, $\operatorname{supp}(f^+)$, $\operatorname{supp}(f^-)$ both contains at least one edge not belonging to E(H). By using [2, Theorem 5.1], I_H is a principal ideal, say $I_H = (f_m)$. We see that $|\operatorname{supp}(f_m)| < |E(D)|$. Then as I_D is generic, there exists $f_n \in M_D$, $\operatorname{supp}(f_n) = E(D)$ such that $f_n^+|f_m^+$ or $f_n^-|f_m^+$ which is contradiction as from above argument $\operatorname{supp}(f_n^+)$, $\operatorname{supp}(f_n^-)$ both contains at least one edge not belonging to E(H). Thus D is one of above type required graphs.

Suppose D is one of above type required graphs. Then we know that I_D is principal and f is the generator of I_D , $\operatorname{supp}(f) = E(D)$. Hence, I_D is generic.

Remark 3.3. By the Theorem 3.2, if I_D is generic, then I_D is a principal ideal. But, the converse is need not be true. For example if D consists of a balanced cycle and certain number of trees, then I_D is principal ideal but I_D is not generic.

Lemma 3.4. Let D be weighted oriented graph consists of one unbalanced cycle C_{m_1} and n-1 (balanced or unbalanced) cycles $C_{m_2}, C_{m_3}, \dots, C_{m_n}$ sharing a single vertex. Then $\dim_{\mathbb{Q}} Null(A(D)) = n-1$.

Proof. Note that $|V(D)| = m_1 + \cdots + m_n - (n-1)$. Let $v_1 \in V(D)$ be the common vertex of all the cycles $\mathcal{C}_{m_1}, \mathcal{C}_{m_2}, \mathcal{C}_{m_3}$. Let $V(\mathcal{C}_{m_i}) = \{v_{\sum_{j=1}^{i-1} m_j+1}, \dots, v_{\sum_{j=1}^{i} m_j}\}$, with the

convention that $v_1 = v_{\sum_{j=1}^{i} m_j + 1}$, for $1 \le i \le n$. Let the edges e_i of D be labelled in the following fashion:

for $1 \leq i \leq n$, each e_k is the edge incident to the vertices v_k and v_{k+1} , for $\sum_{j=1}^{i-1} m_j + 1 \leq i$

 $k \leq \sum_{j=1}^{r} m_j - 1. \text{ Let } A(D) \text{ be the incidence matrix of } D \text{ where the rows are ordered by } v_1, v_2, \ldots, v_{m_1}, v_{m_1+2}, v_{m_1+3}, \ldots, v_{m_1+m_2}, \ldots, v_{m_1+m_2+\ldots+m_{n-1}}, v_{m_1+m_2+\ldots+m_{n-1}+2}, \ldots, v_{m_1+m_2+\ldots+m_n}, \text{ and the columns are ordered by } e_1, e_2, \ldots, e_{m_1+m_2+\ldots+m_n}. We define A(D) \\ \text{to be the matrix obtained from the incidence matrix } A(D) \text{ by deleting } (\sum_{i=1}^{r} m_i)^{\text{th column}} \\ \text{for } r = 2, 3, \ldots, n. \text{ Then } A(D) \text{ is a block triangular matrix of the form } \left[\frac{A \mid B}{O \mid C}\right] \text{ where } A \\ \text{is an } m_1 \times m_1 \text{ matrix and } C \text{ is an } m_2 + m_3 + \ldots + m_n - (n-1) \times m_2 + m_3 + \ldots + m_n - (n-1) \\ \text{upper triangular matrix. Again } A \text{ is the incidence matrix of } \mathcal{C}_{m_1}. \text{ Since toric ideal of unbalanced cycle is zero, } \det(A) \neq 0. \text{ On the other hand, } C \text{ is an upper triangular matrix having each diagonal entry positive. This implies that <math>\det(C) \neq 0$. These two arguments show that $\det(\widetilde{A(D)}) \neq 0$. So, rank of the matrix $\widetilde{A(D)}$ is $m_1 + m_2 + \ldots + m_n - (n-1)$. Therefore rank of the matrix $A(D) \geq m_1 + m_2 + \ldots + m_n - (n-1)$. This implies rank of the matrix $A(D) \leq m_1 + m_2 + \ldots + m_n - (n-1)$. This implies rank of the matrix $A(D) = m_1 + m_2 + \ldots + m_n - (n-1)$. This implies rank of the matrix $A(D) = m_1 + m_2 + \ldots + m_n - (n-1)$.

Lemma 3.5. Let D be a weighted oriented graph consists of n cycles sharing a common vertex. Let D' be a weighted oriented graph consists D and disjoint cycles C_1, C_2, \ldots, C_k such that C_1 is unbalanced and each C_i is connected by a path P_i with D. Then

$$\dim_{\mathbb{Q}} Null(A(D')) = n + k - 1.$$

Proof. Let C'_1, C'_2, \ldots, C'_n be the cycles in D sharing a vertex. Let C_1 be connected by the path P_1 with the cycle C'_1 in D. We label vertices and edges of D' in the following order: first we start labelling C_1 (of vertices and edges) followed by labelling of P_1 , then label C'_1, C'_2, \ldots, C'_n , and then label P_2 followed by C_2 , P_3 followed by C_3 , and so on lastly P_k followed by C_k . Let A(D') be the incidence matrix of D' with respect to above vertex and edge ordering. We define $\widetilde{A(D')}$ is the matrix obtained from A(D') by deleting columns corresponding to last labelling edge of $E(C'_j)$ and $E(C_l)$, for $j = 1, 2, \ldots, n$, $l = 2, \ldots, k$. Then $\widetilde{A(D')}$ is a block triangular matrix of the form $\left[\begin{array}{c} A & B \\ O & C \end{array} \right]$ where A is

an $|V(\mathcal{C}_1)| \times |E(\mathcal{C}_1)|$ matrix and C is an $(|V(D')| - |V(\mathcal{C}_1)|) \times (|E(D')| - (n+k-1) - |E(\mathcal{C}_1)|)$ upper triangular matrix. Note that A is the incidence matrix of \mathcal{C}_1 . Since the toric ideal of unbalanced cycle is zero, $\det(A) \neq 0$. On the other hand, C is an upper triangular matrix having each diagonal entry positive. This implies that $\det(C) \neq 0$. These two arguments show that $\det(\widetilde{A(D')}) \neq 0$. So, rank of the matrix $\widetilde{A(D')}$ is |V(D')| = |E(D')| -(n+k-1). Therefore $\operatorname{rank}(A(D')) \geq |V(D')| = |E(D')| - (n+k-1)$. This implies that $\dim_{\mathbb{Q}}\operatorname{Null}(A(D')) \leq n+k-1$. Let $f_{\mathbf{a_j}}$ be the generator of toric ideal of unbalanced cycle \mathcal{C}_1 and cycle \mathcal{C}_j connected by path and $f_{\mathbf{a'_i}}$ be the generator of toric ideal of unbalanced cycle \mathcal{C}_1 and cycle \mathcal{C}'_l connected by path for $j = 2, 3, \ldots, k, \ l = 1, 2, \ldots, n$. Taking embedding of $\mathbf{a_j}, \mathbf{a'_i}$ in $\operatorname{Null}(A(D'))$, we see that $\{\mathbf{a_j}, \mathbf{a'_i} : j = 2, 3, \ldots, k, \ l = 1, 2, \ldots, n\}$ is linearly independent. Thus $\dim_{\mathbb{Q}}\operatorname{Null}(A(D')) \geq n+k-1$. Hence $\dim_{\mathbb{Q}}\operatorname{Null}(A(D')) = n+k-1$. \Box

Proposition 3.6. Let *D* be weighted oriented graph and $A(D) = [\mathbf{a_1} \ \mathbf{a_2} \ \dots \ \mathbf{a_{|E(D)|}}]$ be the incidence matrix of *D*. Let *S* be an affine semigroup in $\mathbb{Z}_+^{|V(D)|}$ minimally generated by $\mathbf{a_1}, \mathbf{a_2}, \dots, \mathbf{a_{|E(D)|}}$. Then the Krull dimension of K[S] is equal to rank of A(D).

Proof. Let $\phi : \mathbb{Z}^{|E(D)|} \to \mathbb{Z}^{|V(D)|}$ be a group homomorphism defined by $e_i \to \mathbf{a_i}$ where $\{\mathbf{e_1}, \mathbf{e_2}, \dots, \mathbf{e_{|E(D)|}}\}$ is the canonical basis of $\mathbb{Z}^{|E(D)|}$. Then by [14, Proposition 7.5], the Krull dimension of K[S] is equal to $|E(D)| - \operatorname{rank}(\ker(\phi))$. Since $\ker(\phi)$ is a submodule of a free module, we have $\ker\phi \cong \mathbb{Z}^r$, where $r \leq |E(D)|$. Then by Rank-Nullity theorem, we have $\operatorname{rank}(A(D)) = |E(D)| - r$, as required.

Corollary 3.7. Let D be weighted oriented graph as in Lemma 3.4. Then the Krull dimension of $K[e : e \in E(D)]/I_D$ is equal to $m_1 + m_2 + \cdots + m_m - (n-1)$.

Proof. Let $A(D) = [\mathbf{a_1} \ \mathbf{a_2} \ \dots \ \mathbf{a_{|E(D)|}}]$ be the incidence matrix of D. Let A be the affine semigroup in $\mathbb{Z}_+^{|V(D)|}$ minimally generated by $\mathbf{a_1}, \mathbf{a_2}, \dots, \mathbf{a_{|E(D)|}}$. Then $K[A] \cong \frac{K[e:e \in E(D)]}{I_D}$. Using Proposition 3.6, we have the Krull dimension of K[A] is equal to the rank of A(D). Then using Lemma 3.4, this corollary follows.

In the rest of the section, we show that the toric ideal of three weighted oriented unbalanced cycles sharing a vertex, is strongly robust. We achieve this by giving explicitly the unique minimal binomial generating set of the toric ideal. Below we give the figure and notation that we are going to use rest of this section.



Notation 3.8. Let D be weighted oriented graph consists of three unbalanced cycles $C_{m_1}, C_{m_2}, C_{m_3}$ such that these cycles share only a single vertex v_1 as given in above figure. Let $V(C_{m_i}) = \{v_{\sum_{j=1}^{i-1} m_j+1}, \dots, v_{\sum_{j=1}^{i} m_j}\}$, with the convention that $v_1 = v_{\sum_{j=1}^{i} m_j+1}$, for $1 \leq i \leq 3$. and for $1 \leq i \leq 3$, each e_k is the edge incident to the vertices v_k and v_{k+1} , for $\sum_{j=1}^{i-1} m_j + 1 \leq k \leq \sum_{j=1}^{i} m_j - 1$, and the edge $e_{\sum_{j=1}^{i} m_j}$ is incident to the vertices $v_{\sum_{j=1}^{i} m_j}$ and v_1 . Let D_i be the induced subgraph of D that contains precisely $C_{m_i}, C_{m_{i+1}}$ for $1 \leq i \leq 3$ under convention $C_{m_4} = C_{m_1}$. Below we draw picture of D.

Lemma 3.9. For any matrix A with entries of non-negative integers, the corresponding toric ideal I_A is generated by the set of primitive binomials belonging to I_A . That is, I_A is generated by Gr_A .

Proof. Let $f_{\mathbf{m}} \in I_A$ be a part of a generating set of binomials of I_A . Suppose $f_{\mathbf{m}}$ is not a primitive binomial. Then there exists $f_{\mathbf{x}} \in I_A$ such that $f_{\mathbf{x}}^+|f_{\mathbf{m}}^+$ and $f_{\mathbf{x}}^-|f_{\mathbf{m}}^-$. This implies that $\mathbf{e}^{\mathbf{x}_+}|\mathbf{e}^{\mathbf{m}_+}$ and $\mathbf{e}^{\mathbf{x}_-}|\mathbf{e}^{\mathbf{m}_-}$. This gives that $\mathbf{y} := \mathbf{m} - \mathbf{x} \in \text{Null}(A)$ and $\mathbf{y}_+ = \mathbf{m}_+ - \mathbf{x}_+ \ge 0$ and $\mathbf{y}_- = \mathbf{m}_- - \mathbf{x}_- \ge 0$. Thus $f_{\mathbf{y}} \in I_A$ and $f_{\mathbf{m}} \in (f_{\mathbf{x}}, f_{\mathbf{y}})$ (using Lemma 2.5(ii)). If $f_{\mathbf{x}}, f_{\mathbf{y}}$ are not primitive binomials, then we proceed as above until we end with primitive binomials. Continuing in this way after finite number of steps we can show that $f_{\mathbf{m}}$ is a polynomial combination of primitive binomials belonging to I_A . Thus I_A is generated by primitive binomials in it.

Lemma 3.10. Let D be weighted oriented graph and C_1 be a balanced cycle such that both these are connected by a path of length $k \ge 0$ (k = 0 means C_1 shares only one vertex with D). If $f_{\mathbf{a}} \in I_D$ such that $E(\mathcal{C}_1) \subseteq supp(f)$, then $f_{\mathbf{c_1}}^+|f_{\mathbf{a}}^+$ and $f_{\mathbf{c_1}}^-|f_{\mathbf{a}}^-$, for some primitive binomial $f_{\mathbf{c_1}} \in I_{\mathcal{C}_1}$ and $f_{\mathbf{c_1}} \prec f_{\mathbf{a}}$.

Proof. Let $V(\mathcal{C}_1) = \{v_1, v_2, \ldots, v_{m_1}\}$, $E(\mathcal{C}_1) = \{e_1, e_2, \ldots, e_{m_1}\}$, where v_1 is vertex belonging to the connecting path, and e_i is incident with v_i and v_{i+1} , for $i = 1, 2, \ldots, m_1$, under convention that $v_{m_1+1} = v_1$. Let $A(\mathcal{C}_1) = [a_{kl}]_{m_1 \times m_1}$. Assume $f_{\mathbf{a}} \in I_D$ and $E(\mathcal{C}_1) \subseteq \operatorname{supp}(f)$. Then we have $A(D)\mathbf{a} = 0$ which gives that $a_{k(k-1)}[\mathbf{a}]_{k-1} = a_{kk}[\mathbf{a}]_k$, for $2 \leq k \leq m_1$. This implies that

$$a_{11}[\mathbf{a}]_{1} - a_{1m_{1}}[\mathbf{a}]_{m_{1}} = \frac{a_{22}a_{33}\dots a_{m_{1}m_{1}}}{a_{21}a_{32}\dots a_{m_{1}m_{1}-1}}[\mathbf{a}]_{m_{1}} - a_{1m_{1}}[\mathbf{a}]_{m_{1}}$$

$$= \frac{a_{11}a_{22}a_{33}\dots a_{m_{1}m_{1}} - a_{21}a_{32}\dots a_{m_{1}(m_{1}-1)}a_{1m_{1}}}{a_{21}a_{32}\dots a_{m_{1}(m_{1}-1)}}[\mathbf{a}]_{m_{1}}$$

$$= \frac{\det(A(\mathcal{C}_{1}))}{a_{21}a_{32}\dots a_{m_{1}(m_{1}-1)}}[\mathbf{a}]_{m_{1}}$$

$$= 0 \text{ (because } \det(A(\mathcal{C}_{1})) = 0 \text{ as } \mathcal{C}_{1} \text{ is balanced}).$$

Thus the above equalities gives that $\mathbf{a}|_{E(\mathcal{C}_1)} \in \operatorname{Null}(A(\mathcal{C}_1))$. Since $\operatorname{Null}(A(\mathcal{C}_1))$ has dimension 1, let $\{\mathbf{a}_1\}$ be a basis of $\operatorname{Null}(A(\mathcal{C}_1))$, where \mathbf{a}_1 with integer entries. Then $\mathbf{a}|_{E(\mathcal{C}_1)} = k\mathbf{c}_1$ for some positive integer k. This implies that $f_{\mathbf{c}_1}^+|f_{\mathbf{a}}^+$ and $f_{\mathbf{c}_1}^-|f_{\mathbf{a}}^-$. This completes the proof.

Below we show that a WOG of finite oriented cycles sharing a common vertex is strongly robust.

Theorem 3.11. Let D be a weighted oriented graph of finite number of cycles share a single vertex. Then I_D is strongly robust.

Proof. Note that $\mathcal{M}_D \subseteq Gr_D$. Suppose $f_{\mathbf{n}} \in Gr_D \setminus \mathcal{M}_D$. Then $f_{\mathbf{n}}$ is not a part of a system of minimal generators of I_D . This implies that $f_{\mathbf{n}}$ is a combination of remaining elements of that minimal system of generators of I_D . Then by comparing the monomials both sides of this equality we get that there exists $f_{\mathbf{m}} \in Gr_D \setminus \{f_{\mathbf{n}}\}$ such that $f_{\mathbf{m}}^+|f_{\mathbf{n}}^+$.

For any $e_i \in \operatorname{supp}(f_{\mathbf{m}}^-)$, there exists an edge $e_j \in \operatorname{supp}(f_{\mathbf{m}}^+)$ such that e_i, e_j are incident with a vertex v of degree 2. Since $e_j \in \operatorname{supp}(f_{\mathbf{m}}^+) \subseteq \operatorname{supp}(f_{\mathbf{n}}^+)$ and v is incident with only e_i, e_j , then $e_i \in \operatorname{supp}(f_{\mathbf{n}}^-)$. If a_i, a_j are exponents of e_i, e_j respectively in $f_{\mathbf{m}}$, then from $A(D)\mathbf{m} = \mathbf{0}$ (by comparing the row corresponding to v), we get $a_jy - a_ix = 0$ for some positive integers x, y. If b_i, b_j are exponents of e_i, e_j respectively in $f_{\mathbf{n}}$, then from $A(D)\mathbf{n} = \mathbf{0}$ (by comparing the row corresponding to v), we get $b_j y - b_i x = 0$ for the same above x, y. Since $f_{\mathbf{m}}^+|f_{\mathbf{n}}^+$, this implies that $a_j \leq b_j$ which yields that $a_i \leq b_i$ and hence $e_i^{a_i}|e_i^{b_i}$. Thus $f_{\mathbf{m}}^-|f_{\mathbf{n}}^-$. This is a contradiction because $f_{\mathbf{n}}$ is a primitive binomial. Hence $f_{\mathbf{n}}$ is a part of minimal generators of I_D . Thus $Gr_D \subseteq \mathcal{M}_D$. The reverse inclusion is always true. Therefore $Gr_D = \mathcal{M}_D$. Hence I_D is strongly robust. \Box

Theorem 3.12. Let D be the weighted oriented graph consists of certain number of cycles share a vertex v. Let D' be the graph consists of D and certain number of cycles such that each of these cycles is connected by a path at v with D. Then $I_{D'}$ is strongly robust.

Proof. Note that $\mathcal{M}_{D'} \subseteq Gr_{D'}$. Suppose $f_{\mathbf{n}} \in Gr_{D'} \setminus \mathcal{M}_{D'}$. Then there exists $f_{\mathbf{m}} \in Gr_{D'} \setminus \{f_{\mathbf{n}}\}$ such that $f_{\mathbf{m}}^+|f_{\mathbf{n}}^+$. Let $e_i \in \operatorname{supp}(f_{\mathbf{m}}^-)$. We prove that $e_i \in \operatorname{supp}(f_{\mathbf{n}}^-)$ and the exponent of e_i in $f_{\mathbf{m}}^-$ is less than or equal to the exponent of e_i in $f_{\mathbf{n}}^-$.

Step 1: If e_i belongs to cycle or connecting path where the length of this path is greater than 1, then there exists an edge e_j such that e_i, e_j are incident with same vertex of degree 2. Then using arguments as in Theorem 3.11, $e_i \in \text{supp}(f_n^-)$ and exponent of e_i in $\text{supp}(f_m^-)$ is less than or equal to exponent of e_i in $\text{supp}(f_n^-)$.

Step 2: Now assume e_i belongs to connecting path of length 1. Let C_s be the cycle connected by e_i at v with D. Let v_1 be the vertex belonging to $V(\mathcal{C}_s)$ such that e_i is incident with v_1 . Let other two edges in $E(\mathcal{C}_s)$ incident with v_1 be e_x, e_y . Using Lemma 2.7 and $\operatorname{supp}(f_{\mathbf{m}}^+) \subseteq \operatorname{supp}(f_{\mathbf{n}}^+)$, we get that $E(\mathcal{C}_s) \subseteq \operatorname{supp}(f_{\mathbf{m}}) \cap \operatorname{supp}(f_{\mathbf{n}})$ and $\mathbf{m}|_{E(\mathcal{C}_s)} \prec$ $\mathbf{n}|_{E(\mathcal{C}_s)}$. Now we show that \mathcal{C}_s is unbalanced. If possible, suppose \mathcal{C}_s is balanced. Then by the Lemma 3.10, we get that $\mathbf{m}|_{E(\mathcal{C}_s)} \in \operatorname{Null}(A(\mathcal{C}_s))$. This implies that $f_{\mathbf{m}|_{E(\mathcal{C}_s)}} \in I_{D'}$. Note that $f_{\mathbf{m}|_{E(\mathcal{C}_s)}} \neq f_{\mathbf{m}}$, because $e_i \notin \operatorname{supp}(f_{\mathbf{m}|_{E(\mathcal{C}_s)})$ and $e_i \in \operatorname{supp}(f_{\mathbf{m}})$. We see that $f_{\mathbf{m}|_{E(\mathcal{C}_s)}}^+|f_{\mathbf{m}}^+, f_{\mathbf{m}|_{E(\mathcal{C}_s)}}^-|f_{\mathbf{m}}^-$ which is contradiction as $f_{\mathbf{m}}$ is primitive. Thus \mathcal{C}_s is unbalanced. Using similar arguments as above, $\operatorname{supp}(f_{\mathbf{m}}) \cap \operatorname{supp}(f_{\mathbf{n}})$ can not contain edge set of any balanced cycle. Since $E(\mathcal{C}_s) \subseteq \operatorname{supp}(f_{\mathbf{m}}) \cap \operatorname{supp}(f_{\mathbf{n}})$ and $f_{\mathbf{m}} \neq 0, f_{\mathbf{n}} \neq 0$, this implies that there exists an unbalanced cycle $\mathcal{C}_t(t \neq s)$ in D' such that $E(\mathcal{C}_t) \subseteq \operatorname{supp}(f_{\mathbf{m}}) \cap \operatorname{supp}(f_{\mathbf{n}})$.

Let $\operatorname{supp}(f_{\mathbf{n}}) = E(D_1)$ where D_1 is the subgraph of D' consists of all unbalanced cycles and their connecting paths in $\operatorname{supp}(f_{\mathbf{n}})$. Denote all the unbalanced cycles of D_1 by $\mathcal{C}_1, \mathcal{C}_2 \ldots, \mathcal{C}_l$. As $\operatorname{supp}(f_{\mathbf{m}}^+) \subseteq \operatorname{supp}(f_{\mathbf{n}}^+)$ and using Lemma 2.7, Proposition 2.6(i) and using the argument that toric ideal of unbalanced cycle is zero, it follows that $\operatorname{supp}(f_{\mathbf{m}}) \subseteq E(D_1)$ and $f_{\mathbf{m}} \in I_{D_1}$. For any $j \in \{1, 2, \ldots, l\}, j \neq t$, by [2, Theorem 5.1], the toric ideal of unbalanced cycles \mathcal{C}_t and \mathcal{C}_j connected by a path of length $k \geq 0$ (k = 0 means share a vertex) is principal, say, generated by the primitive binomial $f_{\mathbf{a}_j}$ for some vector \mathbf{a}_j . Let $A = {\mathbf{a}_s, \mathbf{a}_j : j = 1, 2, ..., l, j \notin {s, t}}$. For each $k \in {1, 2, ..., l}, k \neq t$ and $e_j \in E(\mathcal{C}_k)$, we have $j \in \operatorname{supp}(\mathbf{a}_k), j \notin \operatorname{supp}(\mathbf{a}_r)$ for each $r \in {1, 2, ..., l} \setminus {t, k}$. Thus A is linearly independent. By Lemma 3.5, we get that $\dim_{\mathbb{Q}} \operatorname{Null}(A(D_1)) = l - 1 = |A|$. Hence A is basis of $\operatorname{Null}(A(D_1))$. As $E(\mathcal{C}_s) \subseteq \operatorname{supp}(f_m)$, without loss of any generality, assume that $\mathbf{a}_s|_{E(\mathcal{C}_s)} \prec \mathbf{m}|_{E(\mathcal{C}_s)}$. Since $\mathbf{m}|_{E(\mathcal{C}_s)} \prec \mathbf{n}|_{E(\mathcal{C}_s)} \prec \mathbf{n}$.

Since A is a basis of Null(A(D₁)) and $\mathbf{m}|_{E(D_1)}, \mathbf{n}|_{E(D_1)} \in \text{Null}(A(D_1))$, then there exist $\mu_k, \lambda_k \in \mathbb{Q}$ such that $\mathbf{m}|_{E(D_1)} = \sum_{k=1,k\neq t}^l \mu_k \mathbf{a}_k$ and $\mathbf{n}|_{E(D_1)} = \sum_{k=1,k\neq t}^l \lambda_k \mathbf{a}_k$. Now compare the k^{th} component both sides, this gives that $[\mathbf{m}]_k = \mu_s[\mathbf{a}_s]_k$ and $[\mathbf{n}]_k = \lambda_s[\mathbf{a}_s]_k$ for all $e_k \in E(\mathcal{C}_s)$. Since for any $e_k \in E(\mathcal{C}_s)$, we have either $k \in \text{supp}(\mathbf{m}_+) \cap \text{supp}(\mathbf{n}_+) \cap \text{supp}((\mathbf{a}_s)_+)$ or $k \in \text{supp}(\mathbf{m}_-) \cap \text{supp}(\mathbf{n}_-) \cap \text{supp}((\mathbf{a}_s)_-)$, this implies that $\mu_s > 0, \lambda_s > 0$. Now compare the i^{th} component both sides of expressions of \mathbf{m}, \mathbf{n} , we get $[\mathbf{m}]_i = \mu_s[\mathbf{a}_s]_i$ and $[\mathbf{n}]_i = \lambda_s[\mathbf{a}_s]_i$. Since $\mu_s > 0, \lambda_s > 0$, and $e_i \in \text{supp}(f_{\mathbf{m}}^-)$, this implies that $e_i \in \text{supp}(f_{\mathbf{n}}^-)$.

Since $e_i \in \operatorname{supp}(f_{\mathbf{m}}^-)$, and v_1 is incident to the edges e_i, e_x, e_y , then the Lemma 2.7, we get either e_x or e_y belongs to $\operatorname{supp}(f_{\mathbf{m}}^+)$, say $e_x \in \operatorname{supp}(f_{\mathbf{m}}^+) \subseteq \operatorname{supp}(f_{\mathbf{n}}^+)$. Now compare the x^{th} component both sides of expressions of \mathbf{m}, \mathbf{n} , we get $0 < [\mathbf{m}]_x = \mu_s[\mathbf{a}_s]_x$ and $0 < [\mathbf{n}]_x = \lambda_s[\mathbf{a}_s]_x$. As $f_{\mathbf{m}}^+|f_{\mathbf{n}}^+$, this implies that $[\mathbf{m}]_x \leq [\mathbf{n}]_x$ which gives that $\mu_s \leq \lambda_s$. Since $e_i \in \operatorname{supp}(f_{\mathbf{m}}^-) \subseteq \operatorname{supp}(f_{\mathbf{n}}^-)$, we have $[\mathbf{m}]_i < 0$ and $[\mathbf{n}]_i < 0$. Since $[\mathbf{m}]_i = \mu_s[\mathbf{a}_s]_i$ and $0 < \mu_s \leq \lambda_s$, we get that $[\mathbf{m}]_i \geq [\mathbf{n}]_i$. This implies that the exponent of e_i in $f_{\mathbf{m}}^-$ is less than or equal to the exponent of e_i in $f_{\mathbf{n}}^-$. Hence in both the Steps 1 and 2, we get that $f_{\mathbf{m}}^-|f_{\mathbf{n}}^-$. This is a contradiction because $f_{\mathbf{n}}$ is primitive. Thus $Gr_{D'} = \mathcal{M}_{D'}$.

Below we explicitly compute unique minimal generating set of I_D , where D consists of three odd cycles sharing a vertex.

Theorem 3.13. Let D be the weighted oriented graph consists of three odd cycles $C_{m_1}, C_{m_2}, C_{m_3}$ such that these cycles share only a single vertex as in Notation 3.8. Let $I_{D_1}, I_{D_2}, I_{D_3}$ are generated by $\{f_{\mathbf{a}}\}, \{f_{\mathbf{b}}\}, \{f_{\mathbf{c}}\}$ respectively, where $\mathbf{a} = ((-1)^{i+1}a_i)_{i=1}^{m_1+m_2}, \mathbf{b} = ((-1)^{i}b_i)_{i=m_1+1}^{m_1+m_2+m_3}$ $\mathbf{c} = (((-1)^{i+1}c_i)_{i=1}^{m_1}, ((-1)^{i}c_i))_{i=m_1+m_2+1}^{m_1+m_2+m_3}$. Then

$$\mathcal{U}_D = Gr_D = M_D = \{f_{\mathbf{a}}, f_{\mathbf{b}}, f_{\mathbf{c}}, f_{\mathbf{x}}, f_{\mathbf{y}}, f_{\mathbf{z}} : \mathbf{x} \in S_1, \mathbf{y} \in S_2, \mathbf{z} \in S_3\},\$$

where $d_1 = gcd(a_1, a_2, \dots, a_{m_1}), d_2 = gcd(b_{m_1+m_2+1}, \dots, b_{m_1+m_2+m_3}),$ $d_3 = gcd(a_{m_1+1}, \dots, a_{m_1+m_2}), d_4 = gcd(c_{m_1+m_2+1}, \dots, c_{m_1+m_2+m_3}),$ $d_5 = gcd(b_{m_1+1}, \dots, b_{m_1+m_2}), d_6 = gcd(c_1, c_2, \dots, c_{m_1}),$

$$\begin{split} S_1 &= \left\{ (((-1)^{i+1} \frac{p_1}{d_1} a_i)_{i=1}^{m_1}, ((-1)^{i+1} (\frac{p_1}{d_1} a_i + \frac{p_2}{d_2} b_i))_{i=m_1+1}^{m_1+m_2}, ((-1)^{i+1} \frac{p_2}{d_2} b_i)_{i=m_1+m_2+1}^{m_1+m_2+m_3}) : \\ (p_1, p_2) &\in E_1 \text{ is a minimal element in } E_1 \right\}, \\ S_2 &= \left\{ (((-1)^{i+1} \frac{p_3}{d_3} a_i + \frac{p_4}{d_4} c_i)_{i=1}^{m_1}, ((-1)^{i+1} \frac{p_3}{d_3} a_i)_{i=m_1+1}^{m_1+m_2}, ((-1)^i \frac{p_4}{d_4} c_i)_{i=m_1+m_2+1}^{m_1+m_2+m_3}) : \\ (p_3, p_4) &\in E_2 \text{ is a minimal element in } E_2 \right\}, \\ S_3 &= \left\{ (((-1)^{i+1} \frac{p_6}{d_6} c_i)_{i=1}^{m_1}, ((-1)^i \frac{p_5}{d_5} b_i)_{i=m_1+1}^{m_1+m_2}, ((-1)^i (\frac{p_5}{d_5} b_i + \frac{p_6}{d_6} c_i))_{i=m_1+m_2+1}^{m_1+m_2+m_3}) : \\ (p_5, p_6) &\in E_3 \text{ is a minimal element in } E_3 \right\}, \\ E_1 &= \left\{ (p_1, p_2) \in [d_1 - 1] \times [d_2 - 1] : \frac{p_1}{d_1} a_j + \frac{p_2}{d_2} b_j \in \mathbb{N}, m_1 + 1 \le j \le m_1 + m_2 \right\}, \\ E_2 &= \left\{ (p_3, p_4) \in [d_3 - 1] \times [d_4 - 1] : \frac{p_3}{d_3} a_j + \frac{p_4}{d_4} c_j \in \mathbb{N}, 1 \le j \le m_1 \right\}, \\ E_3 &= \left\{ (p_5, p_6) \in [d_5 - 1] \times [d_6 - 1] : \frac{p_5}{d_5} b_j + \frac{p_6}{d_6} c_j \in \mathbb{N}, m_1 + m_2 + 1 \le j \le m_1 + m_2 + m_3 \right\} \\ \text{Before previous this theorem first up prove the below proposition:} \end{split}$$

Before proving this theorem, first we prove the below proposition:

Proposition 3.14. Assume all the notations of the Theorem 3.13. Then the set of primitive binomials in I_D is precisely M_D .

Proof. Let $f_{\mathbf{n}} \in I_D$ be a pure binomial such that $\operatorname{supp}(f_{\mathbf{n}}) = E(D)$. Without loss of any generality, assume that $1 \in \operatorname{supp}((\mathbf{n}|_{E(\mathcal{C}_1)})_+), m_1 + 1 \in \operatorname{supp}((\mathbf{n}|_{E(\mathcal{C}_2)})_+), m_1 + m_2 + 1 \in \operatorname{supp}((\mathbf{n}|_{E(\mathcal{C}_2)})_+)$. Then by using the Lemma 2.7 applied to each vertex of degree 2 in D, we have the following possible forms for \mathbf{n} :

$$\begin{split} \mathbf{n} &= (\mathbf{n}|_{\mathbf{E}(\mathcal{C}_{1})}, -\mathbf{n}|_{\mathbf{E}(\mathcal{C}_{2})}, \mathbf{n}|_{\mathbf{E}(\mathcal{C}_{3})}) \text{ or } \mathbf{n} = (\mathbf{n}|_{\mathbf{E}(\mathcal{C}_{1})}, \mathbf{n}|_{\mathbf{E}(\mathcal{C}_{2})}, -\mathbf{n}|_{\mathbf{E}(\mathcal{C}_{3})}) \text{ or } \\ \mathbf{n} &= (\mathbf{n}|_{\mathbf{E}(\mathcal{C}_{1})}, -\mathbf{n}|_{\mathbf{E}(\mathcal{C}_{2})}, -\mathbf{n}|_{\mathbf{E}(\mathcal{C}_{3})}). \text{ This implies that} \end{split}$$

(1)
$$\mathbf{n} = ((-1)^{i+1}n_i)_{i=1}^{m_1+m_2+m_3} \quad \text{or}$$

(2)
$$\mathbf{n} = ((-1)^{i+1}n_i)_{i=1}^{m_1+m_2}, ((-1)^i n_i)_{i=m_1+m_2+1}^{m_1+m_2+m_3} \text{ or }$$

(3)
$$\mathbf{n} = ((-1)^{i+1}n_i)_{i=1}^{m_1}, ((-1)^i n_i)_{i=m_1+1}^{m_1+m_2+m_3}$$

where $n_i \in \mathbb{N}$.

Assume that **n** is of the form as in (1). Since $\{\mathbf{a}, -\mathbf{b}\}$ is a basis of Null(A(D)) and we get that $\mathbf{n} = \frac{p'_1}{d'_1}\mathbf{a} + \frac{p'_2}{d'_2}(-\mathbf{b})$, for some $\frac{p'_1}{d'_1}, \frac{p'_2}{d'_2} \in \mathbb{Q}_+$ with $\operatorname{gcd}(p'_1, d'_1) = 1$, $\operatorname{gcd}(p'_2, d'_2) = 1$. This implies that

(4)
$$n_i = \frac{p'_1}{d'_1} a_i + \frac{p'_2}{d'_2} b_i, \ n_j = \frac{p'_1}{d'_1} a_j, \quad n_k = \frac{p'_2}{d'_2} b_k$$

for $i = m_1 + 1, \ldots, m_1 + m_2, \ j = 1, 2, \ldots, m_1, \ k = m_1 + m_2 + 1, \ldots, m_1 + m_2 + m_3.$ From (4), it follows that $d'_1|a_i, \ d'_2|b_j$, for $i = 1, 2, \ldots, m_1, \ j = m_1 + m_2 + 1, \ldots, m_1 + m_2 + m_3$. Then $d'_1|d_1, \ d'_2|d_2$, by definition of d_1, d_2 . Then we can write $\frac{p'_1}{d_1} = \frac{p''_1}{d_1}, \frac{p'_2}{d_2} = \frac{p''_2}{d_2}$ for some $p''_1 \in \mathbb{N}, p''_2 \in \mathbb{N}$. By division algorithm, we have $p''_1 = p'''_1 + r_1d_1, \ p''_2 = p'''_2 + r_2d_2$, for some $0 \le p'''_1 \le d_1 - 1, 0 \le p'''_2 \le d_2 - 1$, and $r_1, r_2 \in \mathbb{N} \cup \{0\}$. Thus we have

(5)
$$\mathbf{n} = \left(\frac{p_1''}{d_1} + r_1\right)\mathbf{a} + \left(\frac{p_2''}{d_2} + r_2\right)(-\mathbf{b}), \ p_i'' < d_i, \ i = 1, 2.$$

First we show that each binomial in M_D is primitive. Since $f_{\mathbf{a}}$, $f_{\mathbf{b}}$, $f_{\mathbf{c}}$ are circuit binomials, we have that $f_{\mathbf{a}}$, $f_{\mathbf{b}}$, $f_{\mathbf{c}}$ are primitive binomials. For $\mathbf{x} = \frac{p_1}{d_1}\mathbf{a} + \frac{p_2}{d_2}(-\mathbf{b}) \in S_1$, we will show that $f_{\mathbf{x}}$ is primitive binomial. On the contrary suppose $f_{\mathbf{x}}$ is not primitive. Then there exists a pure binomial $f_{\mathbf{n}} \in I_D$ such that $f_{\mathbf{n}}^+|f_{\mathbf{x}}^+$ and $f_{\mathbf{n}}^-|f_{\mathbf{x}}^-$ and $\operatorname{supp}(\mathbf{n}_+) \subseteq \operatorname{supp}(\mathbf{x}_+)$, $\operatorname{supp}(\mathbf{n}_-) \subseteq \operatorname{supp}(\mathbf{x}_-)$. If $\operatorname{supp}(f_{\mathbf{n}}) \subset E(D)$, then by the Proposition 2.6, we get $\operatorname{supp}(f_{\mathbf{n}}) \subseteq E(D_i)$, for i = 1, 2, 3. In fact $\operatorname{supp}(f_{\mathbf{n}}) = E(D_i)$, i = 1, 2, 3. This implies that $f_{\mathbf{n}} \in I_{D_i}$. This gives that $f_{\mathbf{n}} \in (f_{\mathbf{a}})$ or $(f_{-\mathbf{b}})$ or $(f_{\mathbf{c}})$. Thus $f_{\mathbf{y}}|f_{\mathbf{n}}$ for $\mathbf{y} \in \{\mathbf{a}, -\mathbf{b}, \mathbf{c}\}$. This yields that $f_{\mathbf{y}}^+|f_{\mathbf{n}}^+$ or $f_{\mathbf{y}}^-|f_{\mathbf{n}}^+$. Thus $f_{\mathbf{y}}^+|f_{\mathbf{x}}^+$ or $f_{\mathbf{y}}^-|f_{\mathbf{x}}^+$, for $\mathbf{y} \in \{\mathbf{a}, -\mathbf{b}, \mathbf{c}\}$.

This leads to a contradiction because of the following reasons:

if $f_{\mathbf{a}}^+|f_{\mathbf{x}}^+$, then by comparing exponents of e_1 we get a contradiction. If $f_{\mathbf{a}}^-|f_{\mathbf{x}}^+$, then by comparing the second components of \mathbf{a} and \mathbf{x} , we get a contradiction. In a similar way, one can drive contradictions for the other two cases.

Therefore $\operatorname{supp}(f_{\mathbf{n}}) = E(D)$. Now we show that **n** must be in the form (1). If **n** is in the form (2) (or (3)), then by comparing $m_1 + m_2 + 2$ and $m_1 + m_2 + 3$ components (or

 $m_1 + 1$ and $m_1 + 2$ components in case of form (3)) of \mathbf{n} and \mathbf{x} , we get a contradiction to the above support inclusions. Thus \mathbf{n} must be in the form (1). This implies that $\mathbf{e}^{\mathbf{n}_+}|\mathbf{e}^{\mathbf{x}_+}$ and $\mathbf{e}^{\mathbf{n}_-}|\mathbf{e}^{\mathbf{x}_-}$ which yields that $\mathbf{n}_+ \leq \mathbf{x}_+$ and $\mathbf{n}_- \leq \mathbf{x}_-$. Since $\mathrm{supp}(\mathbf{a})$ is neither a subset nor a superset of $\mathrm{supp}(\mathbf{b})$, then there exist indices i_0, j_0 such that $[\mathbf{a}]_{i_0} \neq 0$, $[\mathbf{b}]_{i_0} = 0$ and $[\mathbf{a}]_{j_0} = 0$, $[\mathbf{b}]_{j_0} \neq 0$. Since $\mathbf{n}_+ \leq \mathbf{x}_+$ and $\mathbf{n}_- \leq \mathbf{x}_-$, this implies that $\frac{p_i''}{d_i} + r_i \leq \frac{p_i}{d_i}$ for i = 1, 2. Since $p_i < d_i$, this forces that $r_1 = 0 = r_2$ and $(p_1'', p_2'') \leq (p_1, p_2)$. By the minimality of \mathbf{x} in E_1 , this gives that $(p_1'', p_2'') = (p_1, p_2)$. Therefore $f_{\mathbf{x}}$ must be primitive. In a similar way, if \mathbf{n} is in the form (2) or (3), then one can prove that $f_{\mathbf{y}}, f_{\mathbf{z}}$ are primitive for $\mathbf{y} \in S_2$ and $\mathbf{z} \in S_3$ respectively.

Now we show that if $f_{\mathbf{n}} \in I_D$ is primitive, then $f_{\mathbf{n}} \in M_D$. Let $f_{\mathbf{n}} \in I_D$ be a primitive binomial. Suppose $|\operatorname{supp}(f_{\mathbf{n}})| < |E(D)|$. Then $\operatorname{supp}(f_{\mathbf{n}}) \subseteq E(D_1)$ or $E(D_2)$ or $E(D_3)$. If $\operatorname{supp}(f_{\mathbf{n}}) \subseteq E(D_1)$, then by the Proposition 2.6, we get that $f_{\mathbf{n}} \in I_{D_1} = (f_{\mathbf{a}})$. This implies that $f_{\mathbf{a}}^+|f_{\mathbf{n}}^+$ and $f_{\mathbf{a}}^-|f_{\mathbf{n}}^-$ which implies that $f_{\mathbf{n}} = f_{\mathbf{a}}$ because $f_{\mathbf{n}}$ is primitive. Similarly one can prove for D_2 and D_3 . Assume $\operatorname{supp}(f_{\mathbf{n}}) = E(D)$.

Suppose $r_1 \neq 0$. This implies that $\mathbf{n}_+ > \mathbf{a}_+$ and $\mathbf{n}_- > \mathbf{a}_-$. This gives that $\mathbf{e}^{\mathbf{a}_+} | \mathbf{e}^{\mathbf{n}_+}$ and $\mathbf{e}^{\mathbf{a}_-} | \mathbf{e}^{\mathbf{n}_-}$ which gives that $f_{\mathbf{a}}^+ | f_{\mathbf{n}}^+$ and $f_{\mathbf{a}}^- | f_{\mathbf{n}}^-$. This is a contradiction to the fact that $f_{\mathbf{n}}$ is primitive. Thus $r_1 = 0$ and similarly $r_2 = 0$. Thus $\mathbf{n} = \frac{p_1''}{d_1} \mathbf{a} + \frac{p_2''}{d_2}(-\mathbf{b})$ and $(p_1'', p_2'') \in E_1$. If (p_1'', p_2'') is not minimal, then there exists $(p_1''', p_2'') \in E_1$ such that $p_1''' < p_1''$ or $p_2''' < p_2''$. Let $\mathbf{x} := \frac{p_1'''}{d_1} \mathbf{a} + \frac{p_2''}{d_2}(-\mathbf{b})$. Then $\mathbf{x} \in \text{Null}(A(D))$ such that $\mathbf{n}_+ > \mathbf{x}_+$ and $\mathbf{n}_- > \mathbf{x}_-$. This gives that $\mathbf{e}^{\mathbf{x}_+} | \mathbf{e}^{\mathbf{n}_+}$ and $\mathbf{e}^{\mathbf{x}_-} | \mathbf{e}^{\mathbf{n}_-}$ which gives that $f_{\mathbf{x}}^+ | f_{\mathbf{n}}^+$ and $f_{\mathbf{x}}^- | f_{\mathbf{n}}^-$. This is a contradiction to the fact that $f_{\mathbf{n}}$ is primitive. Therefore (p_1'', p_2'') is minimal. Thus $\mathbf{n} \in S_1$. One can prove in a similar way, that if \mathbf{n} is in the form (2) or (3), then $\mathbf{n} \in S_2$ or S_3 by using the similar arguments as above.

Proof of Theorem 3.13: From Theorem 3.11 and Proposition 3.14, we get that I_D is strongly robust and $\mathcal{U}_D = Gr_D = M_D$.

Below we explicitly compute the unique minimal generating set of I_D , where D consists of two odd cycles and an unbalanced even cycle sharing a vertex.

Theorem 3.15. Let *D* be weighted oriented graph consists of two odd cycles C_{m_1}, C_{m_2} and an unbalanced even cycle C_{m_3} such that these cycles share only a single vertex as in Notation 3.8. Let $I_{D_1}, I_{D_2}, I_{D_3}$ are generated by $\{f_{\mathbf{a}}\}, \{f_{\mathbf{b}}\}, \{f_{\mathbf{c}}\}$ respectively, where $\mathbf{a} = ((-1)^{i+1}a_i)_{i=1}^{m_1+m_2}, \mathbf{b} = (((-1)^ib_i)_{i=m_1+1}^{m_1+m_2}, (l_ib_i)_{i=m_1+m_2+1}^{m_1+m_2+m_3}),$

$$\mathbf{c} = (((-1)^{i+1}c_i)_{i=1}^{m_1}, (l_ic_i)_{i=m_1+m_2+1}^{m_1+m_2+m_3}), \ l_i = \begin{cases} (-1)^{i+1}, & \text{if } r > 0\\ (-1)^i, & \text{if } r < 0 \end{cases}$$

Then

$$\mathcal{U}_D = Gr_D = M_D = \{ f_{\mathbf{a}}, f_{\mathbf{b}}, f_{\mathbf{c}}, f_{\mathbf{x}}, f_{\mathbf{y}}, f_{\mathbf{z}} : \mathbf{x} \in S_1, \mathbf{y} \in S_2, \mathbf{z} \in S_3 \},\$$

where

$$S_{1} = \left\{ (((-1)^{i+1} \frac{p_{1}}{d_{1}} a_{i})^{m_{1}}_{i=1}, ((-1)^{i+1} (\frac{p_{1}}{d_{1}} a_{i} + \frac{p_{2}}{d_{2}} b_{i}))^{m_{1}+m_{2}}_{i=m_{1}+1}, (l_{i}^{*} \frac{p_{2}}{d_{2}} b_{i})^{m_{1}+m_{2}+m_{3}}_{i=m_{1}+m_{2}+1}) : (p_{1}, p_{2}) \in E_{1} \text{ is a minimal element in } E_{1} \right\},$$

$$S_{2} = \left\{ (((-1)^{i+1} \frac{p_{3}}{d_{3}} a_{i} + \frac{p_{4}}{d_{4}} c_{i})^{m_{1}}_{i=1}, ((-1)^{i+1} \frac{p_{3}}{d_{3}} a_{i})^{m_{1}+m_{2}}_{i=m_{1}+1}, (l_{i}^{**} \frac{p_{4}}{d_{4}} c_{i})^{m_{1}+m_{2}+m_{3}}_{i=m_{1}+m_{2}+1}) : (p_{3}, p_{4}) \in E_{2} \text{ is a minimal element in } E_{2} \right\},$$

$$S_{3} = \left\{ \left(((-1)^{i+1} \frac{p_{6}}{d_{6}} c_{i})_{i=1}^{m_{1}}, ((-1)^{i} \frac{p_{5}}{d_{5}} b_{i})_{i=m_{1}+1}^{m_{1}+m_{2}}, (l_{i}^{**} (\frac{p_{5}}{d_{5}} b_{i} + \frac{p_{6}}{d_{6}} c_{i}))_{i=m_{1}+m_{2}+1}^{m_{1}+m_{2}+m_{3}} \right) :$$
$$(p_{5}, p_{6}) \in E_{3} \text{ is a minimal element in } E_{3} \right\},$$

 $l_i^* = \begin{cases} (-1)^{i+1}, & if \ r > 0 \\ (-1)^i, & if \ r < 0 \end{cases}, \quad l_i^{**} = \begin{cases} (-1)^i, & if \ r > 0 \\ (-1)^{i+1}, & if \ r < 0 \end{cases}, E_1, E_2, E_3, d_1, d_2, d_3, d_4, d_5, d_6, are same as mentioned in Theorem 3.13 and <math>r = \det(A(\mathcal{C}_{m_3})). \end{cases}$

Proof. Let $f_{\mathbf{n}} \in I_D$ be a pure binomial such that $\operatorname{supp}(f_{\mathbf{n}}) = E(D)$. Since $\mathbf{n} \in \operatorname{Null}(A(D))$, using Lemma 2.7, we get that \mathbf{n} is of the following possible forms (with appropriate signs)

(6)
$$((-1)^{i+1}n_i)_{i=1}^{m_1}, (-1)^i n_i)_{i=m_1+1}^{m_1+m_2}, ((-1)^{i+1}n_i)_{i=m_1+m_2+1}^{m_1+m_2+m_3}) \text{ or }$$

(7)
$$((-1)^{i+1}n_i)_{i=1}^{m_1}, (-1)^i n_i)_{i=m_1+1}^{m_1+m_2}, ((-1)^i n_i)_{i=m_1+m_2+1}^{m_1+m_2+m_3})$$
 or

(8)
$$((-1)^{i+1}n_i)_{i=1}^{m_1+m_2+m_3}$$
 or

(9)
$$((-1)^{i+1}n_i)_{i=1}^{m_1+m_2}, (-1)^i n_i)_{i=m_1+m_2+1}^{m_1+m_2+m_3}$$

where $n_i \in \mathbb{N}$. The element mentioned in (6) can not belong to $\operatorname{Null}(A(D))$ if r > 0 because if it belongs to $\operatorname{Null}(A(D))$, then from $A(D)\mathbf{n} = 0$, we get $\frac{\det(A(\mathcal{C}_{m_1}))}{a_1} + \frac{\det(A(\mathcal{C}_{m_2}))}{b_1} + \frac{\det(A(\mathcal{C}_{m_3}))}{c_1} = 0$ where $a_1, b_1, c_1 \in \mathbb{N}$ which is a contradiction as $\det(A(\mathcal{C}_{m_i})) > 0$ for i = 1, 2, 3. Similarly, the element mentioned in (7) can not belong to $\operatorname{Null}(A(D))$ if r < 0 because if it belongs to $\operatorname{Null}(A(D))$, then from $A(D)\mathbf{n} = 0$, we get $\frac{\det(A(\mathcal{C}_{m_1}))}{a_2} + \frac{\det(A(\mathcal{C}_{m_2}))}{b_2} - \frac{\det(A(\mathcal{C}_{m_3}))}{c_2} = 0$ where $a_2, b_2, c_2 \in \mathbb{N}$ which is a contradiction as $\det(A(\mathcal{C}_{m_i})) > 0$ for i = 1, 2 and $\det(A(\mathcal{C}_{m_3})) < 0$.

Since $\dim_{\mathbb{Q}}(\operatorname{Null}(A(D))) = 2$, then for any $\mathbf{n} \in \operatorname{Null}(A(D))$, we have that \mathbf{n} is a \mathbb{Q} -linear combination of any subset of two linearly independent elements of the set $\{\mathbf{a}, -\mathbf{a}, \mathbf{b}, -\mathbf{b}, \mathbf{c}, -\mathbf{c}\}$ satisfying sign conditions, are described in below table.

Form of n Case	(6)	(7)	(8)	(9)
r > 0	×	$\checkmark \{\mathbf{b},\!\mathbf{c}\}$	$\checkmark \{a, -b \}$	$\checkmark \{a,c\}$
r < 0	$\checkmark \{b,c\}$	×	$\checkmark \{a, c \}$	$\checkmark \{a, \text{-}b\}$

For each case with the possible basis as mentioned in the above table, we proceed as in the proofs of the Proposition 3.14 and the Theorem 3.13, one can show that the primitive binomials in I_D are in the required forms mentioned in the statement and this set is a minimal generating set.

Below we explicitly compute the unique minimal generating set of I_D , where D consists of an odd cycle and two unbalanced even cycles sharing a vertex.

 $\begin{array}{l} \textbf{Theorem 3.16. Let } D \ be \ weighted \ oriented \ graph \ consists \ of \ odd \ cycle \ \mathcal{C}_{m_1} \ and \ two \ unbalanced \ even \ cycles \ \mathcal{C}_{m_2}, \ \mathcal{C}_{m_3} \ such \ that \ these \ cycles \ share \ only \ a \ single \ vertex \ as \ in \ Notation \ 3.8. \ Let \ I_{D_1}, I_{D_2}, I_{D_3} \ are \ generated \ by \ f_{\mathbf{a}}, f_{\mathbf{b}}, f_{\mathbf{c}} \ respectively, \ where \ \mathbf{a} = (((-1)^{i+1}a_i)_{i=1}^{m_1}, (l_ia_i)_{i=m_1+1}^{m_1+m_2}), \ \mathbf{b} = (((-1)^{ib})_{i=m_1+1}^{m_1+m_2}, (l_i'a_i)_{i=m_1+m_2+1}^{m_1+m_2+m_3}), \ \mathbf{c} = (((-1)^{i+1}b_i)_{i=1}^{m_1}, (l_i''a_i)_{i=m_1+m_2+1}^{m_1+m_2+m_3}), \ l_i = \begin{cases} (-1)^{i+1}, \ if \ q > 0 \\ (-1)^i, \ if \ q < 0 \end{cases}, \ l_i'' = \begin{cases} (-1)^{i+1}, \ if \ qr > 0 \\ (-1)^i, \ if \ qr < 0 \end{cases}, \ l_i'' = \begin{cases} (-1)^{i+1}, \ if \ r > 0 \\ (-1)^i, \ if \ r < 0 \end{cases}. \ Then \end{aligned}$

$$\mathcal{U}_D = Gr_D = M_D = \{f_{\mathbf{a}}, f_{\mathbf{b}}, f_{\mathbf{c}}, f_{\mathbf{x}}, f_{\mathbf{y}}, f_{\mathbf{z}} : \mathbf{x} \in S_1, \mathbf{y} \in S_2, \mathbf{z} \in S_3\},\$$

where

$$\begin{split} S_1 &= \begin{cases} (((-1)^{i+1}\frac{p_1}{d_1}a_i)_{i=1}^{m_1}, (k_i^*(\frac{p_1}{d_1}a_i + \frac{p_2}{d_2}b_i))_{i=m_1+1}^{m_1+m_2}, (l_i^*\frac{p_2}{d_2}b_i)_{i=m_1+m_2+1}^{m_1+m_2+m_3}) :\\ (p_1, p_2) \in E_1 \text{ is a minimal element in } E_1 \\ \\ S_2 &= \begin{cases} (((-1)^{i+1}\frac{p_3}{d_3}a_i + \frac{p_4}{d_4}c_i)_{i=1}^{m_1}, (k_i^{**}\frac{p_3}{d_3}a_i)_{i=m_1+1}^{m_1+m_2}, (l_i^{**}\frac{p_4}{d_4}c_i)_{i=m_1+m_2+1}^{m_1+m_2+m_3}) :\\ (p_3, p_4) \in E_2 \text{ is a minimal element in } E_2 \\ \\ \\ S_3 &= \begin{cases} (((-1)^{i+1}\frac{p_6}{d_6}c_i)_{i=1}^{m_1}, (k_i^{***}\frac{p_5}{d_5}b_i)_{i=m_1+1}^{m_1+m_2}, (l_i^{***}(\frac{p_5}{d_5}b_i + \frac{p_6}{d_6}c_i))_{i=m_1+m_2+1}^{m_1+m_2+m_3}) :\\ (p_5, p_6) \in E_3 \text{ is a minimal element in } E_3 \\ \\ \\ k_i^* &= k_i^{**} = \begin{cases} (-1)^i, & \text{if } q < 0 \\ (-1)^{i+1}, & \text{if } q < 0 \end{cases}, \quad k_i^{***} &= \begin{cases} (-1)^i, & \text{if } q > 0 \\ (-1)^{i+1}, & \text{if } q < 0 \end{cases},\\ \\ \\ l_i^* &= \begin{cases} (-1)^{i+1}, & \text{if } r < 0 \\ (-1)^{i+1}, & \text{if } r < 0 \end{cases}, \quad l_i^{**} &= l_i^{***} &= \begin{cases} (-1)^i, & \text{if } r < 0 \\ (-1)^{i+1}, & \text{if } r > 0 \end{cases},\\ \\ \\ E_1, E_2, E_3, d_1, d_2, d_3, d_4, d_5, d_6, \text{ are same as mentioned in Theorem 3.13 and q &= det(A(C_{m_2})), \\ \\ \end{array} \right), \end{split}$$

Proof. Let $f_{\mathbf{n}} \in I_D$ be a pure binomial such that $\operatorname{supp}(f_{\mathbf{n}}) = E(D)$. Since $\mathbf{n} \in \operatorname{Null}(A(D))$, using Lemma 2.7, we get that **n** is of the following possible forms (with appropriate signs)

(10)
$$(((-1)^{i+1}n_i)_{i=1}^{m_1}, ((-1)^i n_i))_{i=m_1+1}^{m_1+m_2+m_3}) \text{ or }$$

(11)
$$(((-1)^{i+1}n_i)_{i=1}^{m_1}, ((-1)^i n_i)_{i=m_1+1}^{m_1+m_2}, ((-1)^{i+1}n_i)_{i=m_1+m_2+1}^{m_1+m_2+m_3}) \text{ or }$$

(12)
$$(((-1)^{i+1}n_i)_{i=1}^{m_1+m_2}, ((-1)^i n_i)_{i=m_1+m_2+1}^{m_1+m_2+m_3}) \text{ or }$$

(13)
$$((-1)^{i+1}n_i)_{i=1}^{m_1+m_2+m_3}$$

where $n_i \in \mathbb{N}$. The element mentioned in (10) can not belong to $\operatorname{Null}(A(D))$ if q > 0, r > 0because if it belongs to $\operatorname{Null}(A(D))$, then we get $\frac{\det(A(\mathcal{C}_{m_1}))}{a_1} + \frac{\det(A(\mathcal{C}_{m_2}))}{b_1} + \frac{\det(A(\mathcal{C}_{m_3}))}{c_1} = 0$ where $a_1, b_1, c_1 \in \mathbb{N}$ which is a contradiction as $\det(A(\mathcal{C}_{m_i})) > 0$ for i = 1, 2, 3. The element

20

 E_1

mentioned in (11) can not belong to $\operatorname{Null}(A(D))$ if q > 0, r < 0 because if it belongs to $\operatorname{Null}(A(D))$, then we get $\frac{\det(A(\mathcal{C}_{m_1}))}{a_2} + \frac{\det(A(\mathcal{C}_{m_2}))}{b_2} - \frac{\det(A(\mathcal{C}_{m_3}))}{c_2} = 0$ where $a_2, b_2, c_2 \in \mathbb{N}$ which is a contradiction as $\det(A(\mathcal{C}_{m_i})) > 0$ for i = 1, 2 and $\det(A(\mathcal{C}_{m_3})) < 0$. The element mentioned in (12) can not belong to $\operatorname{Null}(A(D))$ if q < 0, r > 0 because if it belongs to $\operatorname{Null}(A(D))$, then we get $\frac{\det(A(\mathcal{C}_{m_1}))}{a_3} - \frac{\det(A(\mathcal{C}_{m_2}))}{b_3} + \frac{\det(A(\mathcal{C}_{m_3}))}{c_3} = 0$ where $a_3, b_3, c_3 \in \mathbb{N}$ which is a contradiction as $\det(A(\mathcal{C}_{m_i})) > 0$ for i = 1, 3 and $\det(A(\mathcal{C}_{m_2})) < 0$. The element mentioned in (13) can not belong to $\operatorname{Null}(A(D))$ if q < 0, r < 0 because if it belongs to $\operatorname{Null}(A(D))$, then we get $\frac{\det(A(\mathcal{C}_{m_1}))}{a_3} - \frac{\det(A(\mathcal{C}_{m_2}))}{b_3} - \frac{\det(A(\mathcal{C}_{m_3}))}{c_3} = 0$ where $a_3, b_3, c_3 \in \mathbb{N}$ which is a contradiction as $\det(A(\mathcal{C}_{m_1})) < 0$ for i = 2, 3 and $\det(A(\mathcal{C}_{m_1})) > 0$.

Since $\dim_{\mathbb{Q}}(\operatorname{Null}(A(D))) = 2$, then for any $\mathbf{n} \in \operatorname{Null}(A(D))$, we have that \mathbf{n} is a \mathbb{Q} -linear combination of any subset of two linearly independent elements of the set $\{\mathbf{a}, -\mathbf{a}, \mathbf{b}, -\mathbf{b}, \mathbf{c}, -\mathbf{c}\}$ satisfying sign conditions, are described in the below table.

Form of n Case	(10)	(11)	(12)	(13)
q > 0, r > 0	×	$\checkmark \{\mathbf{b},\!\mathbf{c}\}$	$\checkmark \{a, \ -b \ \}$	$\checkmark \{a,\!c\}$
q < 0, r < 0	$\checkmark \{a,c\}$	$\checkmark \{a,\!b\}$	√{-b, c}	×
q > 0, r < 0	$\checkmark \{b, c\}$	×	$\checkmark \{a, c\}$	$\checkmark \{a, -b \}$
q < 0, r > 0	$\checkmark \{a,b\}$	$\checkmark \{a,\!c\}$	×	$\checkmark \{\text{-b,c}\}$

For each case with the possible basis as mentioned in the above table, we proceed as in the proofs of the Proposition 3.14 and the Theorem 3.13, one can show that the primitive binomials in I_D are in the required forms mentioned in the statement and this set is a minimal generating set.

Below we explicitly compute the unique minimal generating set of I_D , where D consists of three unbalanced even cycles sharing a vertex.

Theorem 3.17. Let *D* be weighted oriented graph consists of three unbalanced even cycles C_{m_1} , C_{m_2} , C_{m_3} such that these cycles share only a single vertex. Let I_{D_1} , I_{D_2} , I_{D_3} are generated by $f_{\mathbf{a}}$, $f_{\mathbf{b}}$, $f_{\mathbf{c}}$ respectively, where $\mathbf{a} = (((-1)^{i+1}a_i)_{i=1}^{m_1}, (l_ia_i)_{i=m_1+1}^{m_1+m_2}), \mathbf{b} = (((-1)^{i+1}b_i)_{i=m_1+1}^{m_1+m_2}, (l'_ia_i)_{i=m_1+m_2+1}^{m_1+m_2+m_3}),$ $\mathbf{c} = (((-1)^{i+1}c_i)_{i=1}^{m_1}, (l''_ia_i)_{i=m_1+m_2+1}^{m_1+m_2+m_3}),$

$$l_i = \begin{cases} (-1)^i, & if \ pq > 0\\ (-1)^{i+1}, & if \ pq < 0 \end{cases}, \quad l'_i = \begin{cases} (-1)^i, & if \ qr > 0\\ (-1)^{i+1}, & if \ qr < 0 \end{cases}, \quad l''_i = \begin{cases} (-1)^i, & if \ pr > 0\\ (-1)^{i+1}, & if \ pr < 0 \end{cases}$$

Then

$$\mathcal{U}_{D} = Gr_{D} = M_{D} = \{f_{\mathbf{a}}, f_{\mathbf{b}}, f_{\mathbf{c}}, f_{\mathbf{x}}, f_{\mathbf{y}}, f_{\mathbf{z}} : \mathbf{x} \in S_{1}, \mathbf{y} \in S_{2}, \mathbf{z} \in S_{3}\},\$$
where $k_{i}^{*} = k_{i}^{**} = \begin{cases} (-1)^{i}, & \text{if } pq > 0\\ (-1)^{i+1}, & \text{if } pq < 0 \end{cases}, \quad k_{i}^{***} = \begin{cases} (-1)^{i+1}, & \text{if } pq > 0\\ (-1)^{i}, & \text{if } pq < 0 \end{cases},$

$$l_{i}^{*} = \begin{cases} (-1)^{i+1}, & \text{if } pr > 0\\ (-1)^{i}, & \text{if } pr < 0 \end{cases}, \quad l_{i}^{**} = l_{i}^{***} = \begin{cases} (-1)^{i}, & \text{if } pr > 0\\ (-1)^{i+1}, & \text{if } pr < 0 \end{cases},$$

$$S_{1}, S_{2}, S_{3}, E_{1}, E_{2}, E_{3}, d_{1}, d_{2}, d_{3}, d_{4}, d_{5}, d_{6} \text{ are same as mentioned in Theorem 3.13 and } p = det(A(\mathcal{C}_{m_{1}})), q = det(A(\mathcal{C}_{m_{2}})), r = det(A(\mathcal{C}_{m_{3}})).$$

Proof. Let $f_{\mathbf{n}} \in I_D$ be a pure binomial such that $\operatorname{supp}(f_{\mathbf{n}}) = E(D)$. Since $\mathbf{n} \in \operatorname{Null}(A(D))$, using Lemma 2.7, we get that \mathbf{n} is of the following possible forms (with appropriate signs)

(14)
$$(((-1)^{i+1}n_i)_{i=1}^{m_1+m_2+m_3} \text{ or }$$

(15)
$$(((-1)^{i+1}n_i)_{i=1}^{m_1+m_2}, ((-1)^i n_i))_{i=m_1+m_2+1}^{m_1+m_2+m_3}) \text{ or }$$

(16)
$$(((-1)^{i+1}n_i)_{i=1}^{m_1}, ((-1)^i n_i))_{i=m_1+1}^{m_1+m_2}, ((-1)^{i+1}n_i))_{i=m_1+m_2+1}^{m_1+m_2+m_3}) \text{ or }$$

(17)
$$(((-1)^{i+1}n_i)_{i=1}^{m_1}, ((-1)^i n_i)_{i=m_1+1}^{m_1+m_2+m_3}) \text{ or }$$

where $n_i \in \mathbb{N}$. The element mentioned in (14) can not belong to $\operatorname{Null}(A(D))$ if p > 0, q > 0, r > 0 as well as if p < 0, q < 0, r < 0 because if it belongs to $\operatorname{Null}(A(D))$, then we get $\frac{\det(A(\mathcal{C}_{m_1}))}{a_1} + \frac{\det(A(\mathcal{C}_{m_2}))}{b_1} + \frac{\det(A(\mathcal{C}_{m_3}))}{c_1} = 0$ where $a_1, b_1, c_1 \in \mathbb{N}$ which is a contradiction in this case. The element mentioned in (15) can not belong to $\operatorname{Null}(A(D))$ if p > 0, q > 0, r < 0 as well as if p < 0, q < 0, r > 0 because if it belongs to $\operatorname{Null}(A(D))$, then we get $\frac{\det(A(\mathcal{C}_{m_1}))}{a_2} + \frac{\det(A(\mathcal{C}_{m_2}))}{b_2} - \frac{\det(A(\mathcal{C}_{m_3}))}{c_2} = 0$ where $a_2, b_2, c_2 \in \mathbb{N}$ which is a contradiction in this case. The element mentioned in (16) can not belong to $\operatorname{Null}(A(D))$ if p > 0, q < 0, r > 0 as well as if p < 0, q > 0, r < 0 because if it belongs to $\operatorname{Null}(A(D))$ if p > 0, q < 0, r > 0 as well as if p < 0, q > 0, r < 0 because if it belongs to $\operatorname{Null}(A(D))$ if p > 0, q < 0, r > 0 as well as if p < 0, q > 0, r < 0 because if it belongs to $\operatorname{Null}(A(D))$ if p > 0, q < 0, r > 0 as well as if p < 0, q > 0, r < 0 because if it belongs to $\operatorname{Null}(A(D))$ if p > 0, q < 0, r < 0 because if $\operatorname{Log}(A(\mathcal{C}_{m_1})) = 0$ of r = 1, 3 and $\operatorname{Log}(A(\mathcal{C}_{m_2})) < 0$. The element mentioned in (17) can not belong to $\operatorname{Null}(A(D))$ if p > 0, q < 0, r < 0 as well as if p < 0, q < 0, r < 0 as well as if p < 0, q < 0, r < 0 as well as if p < 0, q < 0, r < 0 as well as if p < 0, q < 0, r < 0 because if it belongs to $\operatorname{Null}(A(D))$.

22

it belongs to Null(A(D)), then we get $\frac{\det(A(\mathcal{C}_{m_1}))}{a_4} - \frac{\det(A(\mathcal{C}_{m_2}))}{b_4} - \frac{\det(A(\mathcal{C}_{m_3}))}{c_4} = 0$ where $a_4, b_4, c_4 \in \mathbb{N}$ which is a contradiction in this case.

Since $\dim_{\mathbb{Q}}(\operatorname{Null}(A(D))) = 2$, then for any $\mathbf{n} \in \operatorname{Null}(A(D))$, we have that \mathbf{n} is a \mathbb{Q} -linear combination of any subset of two linearly independent elements of the set $\{\mathbf{a}, -\mathbf{a}, \mathbf{b}, -\mathbf{b}, \mathbf{c}, -\mathbf{c}\}$ satisfying sign conditions, are described in the below table.

Form of n Case	(14)	(15)	(16)	(17)
p > 0, q > 0, r > 0				
p < 0, q < 0, r < 0	×	$\checkmark \{\mathbf{b},\!\mathbf{c}\}$	$\checkmark \{a, -b \}$	$\checkmark \{a,\!c\}$
p > 0, q > 0, r < 0				
p < 0, q < 0, r > 0	$\checkmark \{\mathbf{b}, \mathbf{c}\}$	×	$\checkmark \{a,c\}$	$\checkmark \{a, \text{-}b \ \}$
p > 0, q < 0, r > 0				
p < 0, q > 0, r < 0	$\checkmark \{a,b\}$	$\checkmark \{a,\!c\}$	×	$\checkmark\{\text{-}\mathbf{b},\mathbf{c}\}$
p > 0, q < 0, r < 0				
p < 0, q > 0, r > 0	$\checkmark \{a,c\}$	$\checkmark \{a,b\}$	\checkmark {-b,c}	×

For each case with the possible basis as mentioned in the above table, we proceed as in the proofs of the Proposition 3.14 and the Theorem 3.13, one can show that the primitive binomials in I_D are in the required forms mentioned in the statement and this set is a minimal generating set.

Example 3.18. Let D be a weighted oriented graph as shown in the below diagram consists of three odd cycles $C_{m_1}, C_{m_2}, C_{m_3}$ that share a vertex as in Notation 3.8. Let w_i be the weight of vertex v_i . Assume that $w_1 = 2, w_2 = 2, w_3 = 3, w_5 = 3, w_6 = 4, w_8 = 2, w_9 = 4$. Also,



$$I_{D_1} = (f_{\mathbf{a}}), I_{D_2} = (f_{\mathbf{b}}), I_{D_3} = (f_{\mathbf{c}}), where$$

$$\mathbf{a} = (25, 50, 150, 13, 39, 156, 0, 0, 0),$$

$$\mathbf{b} = (0, 0, 0, 17, 51, 204, 25, 50, 200),$$

$$\mathbf{c} = (17, 34, 102, 0, 0, 0, 13, 26, 104)$$

Then according to the notation of the Theorem 3.13, we have $d_1 = d_2 = 25, d_3 = d_4 = 13, d_5 = d_6 = 17$ and

$$\begin{split} E_1 &= \{(1,11), (3,8), (5,5), (7,2), (16,1), (2,22), (4,19), (6,16), (8,13), (9,24), (10,10), \\ &\quad (11,21), (12,7), (13,18), (14,4), (15,15), (17,12), (18,23), (19,9), (20,20), (21,6), \\ &\quad (22,17), (23,3), (24,14)\}, \end{split} \\ E_2 &= \{(1,10), (2,7), (3,4), (4,1), (5,11), (6,8), (7,5), (8,2), (9,12), (10,9), (11,6), (12,3)\}, \end{split}$$

$$\begin{split} E_3 &= \{(1,2),(9,1),(2,4),(3,6),(4,8),(5,10),(6,12),(7,14),(8,16),(10,3),(11,5),(12,7),\\ &\quad (13,9),(14,11),(15,13),(16,15)\}. \end{split}$$

The sets of minimal elements in E_1 , E_2 , E_3 are

 $\{(1,11), (3,8), (5,5), (7,2), (16,1)\}, \{(1,10), (2,7), (3,4), (4,1)\}, \{(1,2), (9,1)\}$ respectively. Thus $|Gr_D| = |\mathcal{U}_D| = |M_D| = 3 + |S_1| + |S_2| + |S_3| = 3 + 5 + 4 + 2 = 14.$

If D is weighted oriented graph consists of two balanced cycles share a path, then I_D need not be generalized robust. Below we provide an example.

Example 3.19. Consider the following weighted oriented graph consists of two even cycles C_m, C_n that share a path of length 1. Let w_i be the weight of vertex v_i .



Here m = n = 4. If $w_1 = w_2 = w_3 = w_7 = 2$, then above cycles C_m, C_n are balanced. Also, $I_{C_m} = (f_{\mathbf{a}}), I_{C_n} = (f_{\mathbf{b}}), I_{\mathcal{C}} = (f_{\mathbf{c}})$ where $\mathbf{a} = (2, 2, 1, 1, 0, 0, 0), \mathbf{b} = (1, 0, 0, 0, 2, 2, 1), \mathbf{c} = (0, 2, 1, 1, 4, 4, 1)$. Using Macaulay-2, we compute $I_D = (f_{\mathbf{a}}, f_{\mathbf{b}})$.

24

Let A be any minimal generating set of I_D . Then there exists $f \in A$ such that $f^+|f_{\mathbf{a}}^+$. If $supp(f) = E(\mathcal{C}_n)$ or $E(D) \setminus \{e_1\}$ or E(D), then we arrive at a contradiction as in this case $supp(f^+) \notin supp(f_{\mathbf{a}}^+)$. Thus $supp(f) = E(\mathcal{C}_m)$. Then using arguments as in Theorem 3.11, we get $f^-|f_{\mathbf{a}}^-$. This implies that f and $f_{\mathbf{a}}$ are associates as $f_{\mathbf{a}}$ is primitive binomial. This proves that f_a is indispensable binomial. Similarly, we can show that f_b is indispensable binomial. As $\{f_{\mathbf{a}}, f_{\mathbf{b}}\}$ is generating set of I_D , it is unique minimal generating set of I_D i.e $\mathcal{M}_D = \{f_{\mathbf{a}}, f_{\mathbf{b}}\}$. Using Macaulay-2 [6], we get $B = \{f_{\mathbf{a}}, f_{\mathbf{b}}, e_2^2 e_4 e_6^2 - e_1 e_3 e_5^2 e_7\}$ is Gröbner basis of I_D . We see that $B \neq \mathcal{M}_D$. Thus $\mathcal{U}_D \neq \mathcal{M}_D$. Hence I_D is not generalized robust. This implies that I_D is not robust and not strongly robust.

Notation 3.20. Let D be weighted oriented graph consists of two cycles C_m, C_n share a vertex. Let $V(C_m) = \{v_1, \ldots, v_m\}, V(C_n) = \{v_1, v_{m+2}, \ldots, v_{m+n}\}$ under convention $v_{m+1} = v_{m+n+1} = v_1, e_i$ be the edge incident with v_i and v_{i+1} for $i = 1, 2, \ldots, m+n$.

Below we recall the results from [16] which are used in the proofs of main results.

Theorem 3.21. [16, Theorem 4.7] Let D be weighted oriented graph consists of two odd cycles C_m, C_n as in above notation 3.20. Then the toric ideal I_D is generated by

$$\prod_{i=1,odd}^{m+n} e_i^{\frac{a_i}{d}} - \prod_{i=1,even}^{m+n} e_i^{\frac{a_i}{d}}, \text{ where }$$

$$\begin{aligned} a_{1} &= \alpha_{2}\alpha_{3}\dots\alpha_{m}|p|, \\ a_{i} &= \alpha'_{2}\alpha'_{3}\dots\alpha'_{i}\alpha_{i+1}\dots\alpha_{m}|p|, \ i=2,3,\dots,m-1, \\ a_{m} &= \alpha'_{2}\alpha'_{3}\dots\alpha'_{m}|p|, \\ a_{m+1} &= \alpha_{m+2}\alpha_{m+3}\dots\alpha_{m+n}|q|, \\ a_{m+i} &= \alpha'_{m+2}\dots\alpha'_{m+i}\alpha_{m+i+1}\dots\alpha_{m+n}|q|, \ i=2,\dots,n-1, \\ a_{m+n} &= \alpha'_{m+2}\alpha'_{m+3}\dots\alpha'_{m+n}|q|, \\ p &= det(A(\mathcal{C}_{n})), \\ q &= det(A(\mathcal{C}_{m})), \\ d &= gcd(a_{i})_{i=1}^{m+n}, \\ det(A(\mathcal{C}_{n})) &= \alpha_{m+2}\alpha_{m+3}\dots\alpha_{m+n}\alpha_{1,m+1} + (-1)^{n+1}\alpha'_{m+2}\alpha'_{m+3}\alpha'_{m+4}\dots\alpha'_{m+n}\alpha_{1,m+n}, \\ det(A(\mathcal{C}_{m})) &= \alpha_{1}\alpha_{2}\alpha_{3}\dots\alpha_{m} + (-1)^{m+1}\alpha'_{2}\alpha'_{3}\alpha'_{4}\dots\alpha'_{m}\alpha_{1,m} \end{aligned}$$

$$\alpha'_{k} = \begin{cases} 1, & \text{if } v_{k} \text{ is tail of } e_{k-1}, \\ w_{k}, & \text{if } v_{k} \text{ is head of } e_{k-1}, \end{cases}, \qquad \alpha_{j} = \begin{cases} 1, & \text{if } v_{j} \text{ is tail of } e_{j}, \\ w_{j}, & \text{if } v_{j} \text{ is head of } e_{j}, \end{cases}$$
$$\alpha_{1,k} = \begin{cases} 1, & \text{if } v_{j} \text{ is tail of } e_{k}, \\ w_{1}, & \text{if } v_{1} \text{ is head of } e_{k}, \end{cases}.$$

Theorem 3.22. [16, Theorem 4.9] Let D be weighted oriented graph consists of two even unbalanced cycles C_m, C_n as in above Notation 3.20. Then the toric ideal I_D is generated by

$$\begin{array}{l} (1) & \prod_{i=1,odd}^{m} e_{i}^{\frac{a_{i}}{d}} \prod_{i=m+1,even}^{m+n} e_{i}^{\frac{a_{i}}{d}} - \prod_{i=1,even}^{m} e_{i}^{\frac{a_{i}}{d}} \prod_{i=m+1,odd}^{m+n} e_{i}^{\frac{a_{i}}{d}} & \text{if } pq > 0 \\ (2) & \prod_{i=1,odd}^{m+n} e_{i}^{\frac{a_{i}}{d}} - \prod_{i=1,even}^{m+n} e_{i}^{\frac{a_{i}}{d}} & \text{if } pq < 0 \end{array}$$

where a_i, d, p are same as in Theorem 3.13.

Lemma 3.23. Let C_m be a weighted oriented even cycle. If V^+ are sinks, then C_m is balanced and exponent of each variable into irreducible generator of I_{C_m} is 1.

Proof. Since main-diagonal and off-diagonal entries of each row of incidence matrix corresponding to source vertex as well as non-source non-sink vertex are 1 and main-diagonal and off-diagonal entries of each row of incidence matrix corresponding to sink are its weight, determinant of its incidence matrix is zero and $\mathbf{x} = (1, -1, 1, -1, \dots, 1, -1)^T \in$ Null $(A(\mathcal{C}_m)$. This implies that the corresponding binomial of \mathbf{x} , is the required generator of $I_{\mathcal{C}_m}$.

Lemma 3.24. Let D be weighted oriented graph consists of unbalanced cycles C_m, C_n share a vertex such that V^+ are sinks. Then C_m, C_n are odd cycles and exponent of each variable in $f_{\mathbf{c}_m \cup \mathbf{c}_n}$ is 1.

Proof. By using Lemma 3.23, we get that C_m, C_n are odd cycles. We use notation as in Theorem 3.21. Let v_1 be the sharing vertex and w_i denotes weight corresponding to vertex v_i . Then $\alpha_j = \alpha'_j = \begin{cases} 1, & \text{if } v_j \text{ is not sink of } D, \\ w_j, & \text{if } v_j \text{ is sink of } D, \end{cases}$ for $j = 2, 3, \ldots, m, m + 2, \ldots, m + n$. Also, $p = 2 \prod_{v_i \text{ is sink of } c_n} w(v_i)$ and $q = 2 \prod_{v_j \text{ is sink of } c_m} w(v_j)$. Using Theorem 3.13,

26

$$a_{i} = \begin{cases} 2 \prod_{v_{i} \text{ is sink of } c_{1}} w_{i} \prod_{v_{j} \text{ is sink of } c_{2}} w_{j} & \text{if } v_{1} \text{ is not sink of } D \\ 2w_{1} \prod_{v_{i} \text{ is sink of } c_{1}} w_{i} \prod_{v_{j} \text{ is sink of } c_{2}} w_{j} & \text{if } v_{1} \text{ is sink of } D \\ \text{for each } i = 1, 2, \dots, m+n. \end{cases}$$

Hence, exponent of each variable in $f_{\mathbf{c_m}\cup\mathbf{c_n}}$ is 1.

Corollary 3.25. Let D be a weighted oriented graph consists of unbalanced cycles C_1, C_2, C_3 sharing a vertex such that V^+ are sinks. Then $I_D = (f_{\mathbf{c}_1 \cup \mathbf{c}_2}, f_{\mathbf{c}_1 \cup \mathbf{c}_3}, f_{\mathbf{c}_2 \cup \mathbf{c}_3})$.

Proof. Using the Lemma 3.24, Theorem 3.13 and according to the notation of this theorem, $d_1 = d_2 = d_3 = d_4 = d_5 = d_6 = 1$ and hence $E_1 = E_2 = E_3 = \emptyset$. This implies the result follows.

Lemma 3.26. Let D be a weighted oriented graph consists of two naturally oriented odd cycles C_m , C_n (or unbalanced even cycles C_m , C_n) sharing only a single vertex and $\prod_{v \in V(C_m)} w(v) = \prod_{v \in V(C_n)} w(v)$. Then the irreducible generator of I_D has exponents of at least one variable from the edge set $E(C_m)$ and at least one variable from the edge set $E(C_n)$ are equal to 1.

Proof. We use notation as in Theorems 3.21 or 3.22. Since cycles are naturally oriented, without loss of any generality, we can assume that $e_1 = (v_1, v_2)$, $e_{m+1} = (v_1, v_{m+2})$.

In this case, $\alpha_i = 1$ for i = 2, 3, ..., m, m + 2, m + 3, ..., m + n,

$$p = \begin{cases} 1 + \prod_{v \in V(\mathcal{C}_n)} w(v), & \text{if } n \text{ odd,} \\ 1 - \prod_{v \in V(\mathcal{C}_n)} w(v), & \text{if } n \text{ even,} \end{cases}, \qquad q = \begin{cases} 1 + \prod_{v \in V(\mathcal{C}_m)} w(v), & \text{if } n \text{ odd,} \\ 1 - \prod_{v \in V(\mathcal{C}_m)} w(v), & \text{if } n \text{ even,} \end{cases}, a_1 = a_1 + a_2 + a_2 + a_3 + a_4 + a_$$

 $a_{m+1} = |p| = |q|$ and $|p||a_i$ for any i = 1, 2, ..., m + n. Thus d = |p| = |q|. Therefore using Theorem 3.13 or 3.16, exponents of e_1, e_{m+1} are 1.

Corollary 3.27. Let D be weighted oriented graph consists of three naturally oriented odd cycles $C_{m_1}, C_{m_2}, C_{m_3}$ (or unbalanced even cycles $C_{m_1}, C_{m_2}, C_{m_3}$) sharing only a single vertex as in Notation 3.8. Suppose $\prod_{v \in V(C_{m_1})} w(v) = \prod_{v \in V(C_{m_2})} w(v) = \prod_{v \in V(C_{m_3})} w(v)$. Then mini-

mum generating set of I_D is precisely the set of three irreducible generators corresponding to I_{D_1} , I_{D_2} , I_{D_3} respectively.

Proof. Let $f_{\mathbf{a}}, f_{\mathbf{b}}, f_{\mathbf{c}}$ be irreducible generators of $I_{D_1}, I_{D_2}, I_{D_3}$ respectively. Then using Lemma 3.26 and using Theorems 3.13 or 3.17(notation as in these theorems), $d_1 = d_2 = d_3 = d_4 = d_5 = d_6 = 1$. Hence, $E_1 = E_2 = E_3 = \emptyset$ and this completes the proof.

Lemma 3.28. Let D be a weighted oriented graph consists of unbalanced cycles C_m, C_n share a vertex v_1 such that C_m has only one non-source non-sink vertex x and C_n has only one non-source non-sink vertex y such that w(x) = w(y) and the sharing vertex is either sink of D or is not sink of any cycle or the weight of this vertex is 1. Then both cycles are odd and the exponent of at least one variable corresponding to edge set of each cycle in irreducible generator of I_D is 1.

Proof. As in a cycle graph, the number of sources is equal to the number of sinks and each of the given cycles has only one non-source non-sink vertex, each of them is odd cycle.

We use notation as in Theorem 3.21. If v_i is non-source non-sink vertex of cycles C_m or C_n incident with edges e_{j-1} and e_j , then without loss of any generality, we can assume that v_i is head of e_{j-1} and v_i is tail of e_j . Let w_i denotes weight corresponding to vertex v_i and w(x) = w(y) = w. We get $p = (1 + w) \prod_{v_i \text{ is sink of } C_n} w(v_i)$ and $q = (1 + w) \prod_{v_j \text{ is sink of } c_m} w(v_j)$, . We use Theorem 3.21. Now, we have the following casess :

<u>**Case 1:**</u> Suppose $v_i (\neq v_1)$ is non-source non-sink vertex of \mathcal{C}_m and v_{m+1} is non-source nonsink vertex of \mathcal{C}_n . Then $\alpha'_i = w$, $\alpha_i = 1$ and $a_{i-1} = \prod_{\substack{v_i (\neq v_1) \text{ sink of } \mathcal{C}_m \\ v_i \neq v_1 \text{ sink of } \mathcal{C}_n \\ v_i \neq v_i \text{ sink of } \mathcal{C}_n \\ w$), $a_j = a_{i-1}$, $a_k = wa_{i-1}$ for $1 \leq j < i-1$, $k \geq i$, $a_{m+j} = a_{i-1}$ for $j = 1, 2, \ldots, n$. Thus $d = a_{i-1}$ and exponents of e_i in irreducible generator of I_D is 1 for $i = 1, 2, \ldots, i-1, m+1, \ldots, m+n$.

<u>**Case 2:**</u> Suppose $v_i \neq v_1$ is non-source non-sink vertex of C_m and $v_j \neq v_{m+1}$ is non-source non-sink vertex of C_n . Using arguments like in Case 1, exponents of e_i in irreducible generator of I_D is 1 for i = 1, 2, ..., i - 1, j - 1, ..., m + n.

<u>**Case 3:**</u> Suppose v_1 is non-source non-sink vertex of \mathcal{C}_m and v_{m+1} is non-source non-sink vertex of \mathcal{C}_n . Using arguments like in Case 1, exponents of e_i in irreducible generator of I_D is 1 for $i = 1, 2, \ldots, m + n$.

Corollary 3.29. Let D be a weighted oriented graph consists of three unbalanced cycles share a vertex as in Notation 3.8 such that each cycle has only one non-source non-sink vertex with same weight and the sharing vertex is either sink of D or is not sink of any cycle or the weight of this vertex is 1. Then minimum generating set of I_D is precisely the set of three irreducible generators corresponding to I_{D_1} , I_{D_2} , I_{D_3} .

Proof. Let $f_{\mathbf{a}}, f_{\mathbf{b}}, f_{\mathbf{c}}$ be irreducible generators of $I_{D_1}, I_{D_2}, I_{D_3}$ respectively. Then using Lemma 3.28 and using Theorem 3.13 (notation as in this theorem), $d_1 = d_2 = d_3 = d_4 = d_5 = d_6 = 1$. Hence, $E_1 = E_2 = E_3 = \emptyset$ and this completes the proof.

4. CIRCUIT BINOMIALS AND STRONGLY ROBUSTNESS OF TORIC IDEALS

In this section, first we characterize the circuit binomials in any WOG. If D consists of n oriented cycles sharing a vertex such that at most two among these cycles are unbalanced, then we show that C_D is a generating set of I_D and hence $C_D = \mathcal{M}_D = \mathcal{U}_D = Gr_D$. We characterize primitive binomials of a WOG consists of n unbalanced cycles sharing a vertex. First we fix the below notation.

Notation 4.1. For a balanced cycle C_i , we know that its toric ideal I_{C_i} is generated by a single primitive binomial, say $f_{\mathbf{c}_i}$, where $\mathbf{c}_i \in Null(A(C_i))$. For two unbalanced cycles C_i, C_j sharing a vertex, the toric ideal of $C_i \cup C_j$ is principal and generated by a primitive binomial, say $f_{\mathbf{c}_i \cup \mathbf{c}_j}$, where $\mathbf{c}_i \cup \mathbf{c}_j$ is a vector in $Null(A(C_i \cup C_j))$. Let $f_{\mathbf{c}_i\mathbf{P}_k\mathbf{c}_j}$ denotes the primitive binomial which is the generator of toric ideal of unbalanced cycles C_i and C_j share a path P_k such that the cycle whose edge set is $E(C_i \cup C_j) \setminus E(P_k)$ is unbalanced. Also, $f_{\mathbf{c}_i \cup \mathbf{P}_k \cup \mathbf{c}_j}$ denotes the primitive binomial which is the generator of toric ideal of the WOG of unbalanced cycles C_i and C_j that are connected by a path P_k (such a toric ideal is principal by [2, Theorem 5.1]).

Proposition 4.2. For any weighted oriented graph D, the set of all circuit binomials in I_D is given by

 $\mathcal{C}_D = \{ f_{\mathbf{c}}, f_{\mathbf{c}_i \cup \mathbf{c}_i}, f_{\mathbf{c}_i \cup \mathbf{p}_k \cup \mathbf{c}_i}, f_{\mathbf{c}_i \mathbf{p}_k \mathbf{c}_i} : \mathcal{C} \text{ is balanced, } P_k \text{ is path }, \mathcal{C}_i, \mathcal{C}_j \text{ are unbalanced in } D \}.$

Proof. Let $A = \{f_{\mathbf{c}}, f_{\mathbf{c}_{\mathbf{i}}\cup\mathbf{c}_{\mathbf{j}}}, f_{\mathbf{c}_{\mathbf{i}}\cup\mathbf{p}_{\mathbf{k}}\cup\mathbf{c}_{\mathbf{j}}}, f_{\mathbf{c}_{\mathbf{i}}\mathbf{p}_{\mathbf{k}}\mathbf{c}_{\mathbf{j}}} : \mathcal{C} \text{ is balanced}, P_k \text{ is path }, \mathcal{C}_i, \mathcal{C}_j \text{ are unbalanced} \text{ in } D\}$. It is easy to see that $A \subseteq \mathcal{C}_D$. Let $f \in \mathcal{C}_D$ such that $\operatorname{supp}(f)$ contains more than three cycles. Then either $\operatorname{supp}(f_{\mathbf{c}}) \subsetneqq \operatorname{supp}(f)$ or $\operatorname{supp}(f_{\mathbf{c}_{\mathbf{i}}\cup\mathbf{p}_{\mathbf{k}}\cup\mathbf{c}_{\mathbf{j}}}) \subsetneqq \operatorname{supp}(f)$ or $\operatorname{supp}(f_{\mathbf{c}_{\mathbf{i}}\mathbf{p}_{\mathbf{k}}\mathbf{c}_{\mathbf{j}}}) \subsetneqq \operatorname{supp}(f)$, where $\operatorname{supp}(f)$ strictly contains edge set of a balanced cycle or two unbalanced cycles or three unbalanced cycles. This is a contradiction because f is a circuit binomial. Thus $\operatorname{supp}(f)$ contains precisely edge set of one balanced

cycle or two unbalanced cycles or three unbalanced cycles. Hence $f \in \mathcal{C}_D$. This completes the proof.

Proposition 4.3. Let D be a weighted oriented graph such that I_D is strongly robust. If C_D is a generating set of I_D , then $C_D = \mathcal{M}_D = \mathcal{U}_D = Gr_D$.

Proof. As I_D is strongly robust and $\mathcal{C}_D \subseteq Gr_D$, then $\mathcal{C}_D \subseteq M_D$. Thus if \mathcal{C}_D is generating set of I_D , then \mathcal{C}_D is a minimal generating set of I_D , that is, $\mathcal{C}_D = \mathcal{M}_D$. Since I_D is strongly robust, $\mathcal{C}_D = \mathcal{M}_D = \mathcal{U}_D = Gr_D$.

Proposition 4.4. Let D be weighted oriented graph consists of certain number of balanced cycles share a vertex. Then $C_D = \mathcal{M}_D = \mathcal{U}_D = Gr_D$.

Proof. Using [16, Theorem 3.1], we have that C_D is a generating set of I_D . Thus the result follows from the Theorem 3.11 and Proposition 4.3.

Proposition 4.5. Let $D = C_1 \cup \ldots \cup C_n$ be a weighted oriented graph such that C_1, \ldots, C_n are oriented cycles share a vertex, C_1 is unbalanced and C_2, \ldots, C_n are balanced. Then $C_D = \mathcal{M}_D = \mathcal{U}_D = Gr_D$.

Proof. Let $f \in Gr_D \setminus \mathcal{C}_D$. Then using Lemma 3.10, $\operatorname{supp}(f)$ can not contain edge set of any balanced cycle. This implies that $\operatorname{supp}(f) \subseteq E(\mathcal{C}_1)$. Therefore $f \in I_{\mathcal{C}_1} = (0)$ and hence f = 0. This is a contradiction. Thus $Gr_D = \mathcal{C}_D$ and as I_D is strongly robust, $\mathcal{C}_D = \mathcal{M}_D = \mathcal{U}_D = Gr_D$.

Proposition 4.6. Let *D* be a weighted oriented graph consists of balanced cycles C_1, C_2, \ldots , C_{n-2} and unbalanced cycles C_{n-1}, C_n share a vertex. Then $C_D = \mathcal{M}_D = \mathcal{U}_D = Gr_D$.

Proof. By the Lemma 3.9, we have I_D is generated by Gr_D . To show that $\mathcal{C}_D = Gr_D$. Suppose there exists $f \in Gr_D \setminus \mathcal{C}_D$. If $\operatorname{supp}(f)$ contains edge set of one balanced cycle \mathcal{C} , then using Lemma 3.10, $f_{\mathbf{c}}^+|f^+$ and $f_{\mathbf{c}}^-|f^-$ which is a contradiction as $f \in Gr_D \setminus \mathcal{C}_D$. Thus $\operatorname{supp}(f)$ contains precisely edge set of unbalanced cycles $\mathcal{C}_{n-1}, \mathcal{C}_n$. This implies that $f_{\mathbf{c}_{n-1}\cup\mathbf{c}_n}^+|f^+$ and $f_{\mathbf{c}_{n-1}\cup\mathbf{c}_n}^-|f^-$ which is a contradiction as $f \in Gr_D \setminus \mathcal{C}_D$. Thus $Gr_D = \mathcal{C}_D$. As I_D is strongly robust, $M_D = \mathcal{U}_D = Gr_D = \mathcal{C}_D$.

Proposition 4.7. Let D be a weighted oriented graph consists of n cycles share a vertex such that every vertex of D is either source or sink. Then every cycle in D is even balanced cycle and $\mathcal{M}_D = \mathcal{U}_D = Gr_D = \mathcal{C}_D$.

Proof. As every vertex of D is either source or sink, if D has any odd cycle, then that odd cycle has non-source non-sink vertex. Thus D can not have any odd cycles. Then every cycle in D is even and balanced. Hence using Lemma 3.10 and as I_D is strongly robust, it follows.

Theorem 4.8. Let D be weighted oriented graph consists of n unbalanced cycles C_1, \ldots, C_n share a common vertex. Let $I_{\mathcal{C}_i \cup \mathcal{C}_j} = (f_{\mathbf{c}_i \cup \mathbf{c}_j})$, for some vector $\mathbf{c}_i \cup \mathbf{c}_j \in Null(A(D))$ with integer entries, for all i, j. Let $\mathbf{m} \in Null(A(D)) \cap \mathbb{Z}^{|E(D)|}$. Suppose $f_{\mathbf{m}}$ is a primitive binomial in I_D such that $supp(f_{\mathbf{m}}) = E(D)$. Then there exists a cycle \mathcal{C}_k depending on \mathbf{m} such that \mathbf{m} is of the following form

$$\mathbf{m} = \sum_{r \in X} (\frac{p_r}{d_r} + k_r) (\mathbf{c_k} \cup \mathbf{c_r}) + \sum_{r \notin X} (\frac{p_r}{d_r} - k_r) (\mathbf{c_k} \cup \mathbf{c_r}),$$

where $X = \{r : (\mathbf{c_k} \cup \mathbf{c_r}) \prec \mathbf{m}\}$, such that $(\mathbf{c_k} \cup \mathbf{c_r})|_{E(\mathcal{C}_k)} \prec \mathbf{m}$, $\sum_{r \in X} (\frac{p_r}{d_r} + k_r) [\mathbf{c_k} \cup \mathbf{c_r}]_j + \sum_{r \notin X} (\frac{p_r}{d_r} - k_r) [\mathbf{c_k} \cup \mathbf{c_r}]_j \in \mathbb{Z}, \text{ for } e_j \in E(\mathcal{C}_k), \text{ where } d_r = gcd([\mathbf{c_k} \cup \mathbf{c_r}]_j : e_j \in E(\mathcal{C}_r)), \ 0 \le p_r < d_r, \ k_r \in \mathbb{N} \cup \{0\}.$

Proof. Fix an arbitrary cycle C_k from D. Let $A = {\mathbf{c_k} \cup \mathbf{c_l} : C_l \text{ is cycle in } D}$. Since $\dim_{\mathbb{Q}} \operatorname{Null}(A(D)) = n - 1$ and A is linearly independent set containing n - 1 elements, therefore A is a basis of $\operatorname{Null}(A(D))$. Let $\mathbf{m} = \sum_{C_i \in D} \mu_i(\mathbf{c_k} \cup \mathbf{c_l})$, for some $\mu_i \in \mathbb{Q}$. Without loss of generality, we can assume that $(\mathbf{c_k} \cup \mathbf{c_l})|_{E(C_k)} \prec \mathbf{m}$, for each cycle C_l in D.

Now we prove that there exists a cycle C_j such that $(\mathbf{c_k} \cup \mathbf{c_j})|_{E(C_j)} \prec \mathbf{m}$. Suppose not, that is, $(\mathbf{c_k} \cup \mathbf{c_l})|_{E(\mathcal{C}_l)} \not\prec \mathbf{m}$ for each cycle C_l in D. This implies that $t \in \operatorname{supp}(\mathbf{m}_+) \cap$ $\operatorname{supp}((\mathbf{c_k} \cup \mathbf{c_l})_-)$ or $t \in \operatorname{supp}(\mathbf{m}_-) \cap \operatorname{supp}((\mathbf{c_k} \cup \mathbf{c_l})_+)$ for $e_t \in E(C_l)$. Since $[\mathbf{m}]_t =$ $\sum_l \mu_l[(\mathbf{c_k} \cup \mathbf{c_l})]_t = \mu_l[(\mathbf{c_k} \cup \mathbf{c_l})]_t$, for all $t \in E(\mathcal{C}_l)$, we get that $\mu_l < 0$. Thus we have $\mu_l < 0$ for all $l \neq k$. This implies that $t \in \operatorname{supp}(\mathbf{m}_+) \cap \operatorname{supp}(\mathbf{m}_-)$ for $e_t \in E(\mathcal{C}_k)$ which is a contradiction because $\operatorname{supp}(\mathbf{m}_+) \cap \operatorname{supp}(\mathbf{m}_-) = \emptyset$. Thus there exists a cycle C_j such that $(\mathbf{c_k} \cup \mathbf{c_j}) \prec \mathbf{m}$. Let $X = \{r : (\mathbf{c_k} \cup \mathbf{c_r}) \prec \mathbf{m}\}$. Then

(18)
$$\mathbf{m} = \sum_{\mathcal{C}_r \text{ in } D} \mu_r(\mathbf{c}_{\mathbf{k}} \cup \mathbf{c}_{\mathbf{r}}) = \sum_{r \in X} \mu_r(\mathbf{c}_{\mathbf{k}} \cup \mathbf{c}_{\mathbf{r}}) + \sum_{r \notin X} \mu_r(\mathbf{c}_{\mathbf{k}} \cup \mathbf{c}_{\mathbf{r}})$$

Now we show that $\mu_r > 0$ for $r \in X$ and $\mu_r < 0$ for $r \notin X$. Let $r \in X$. Then for $e_t \in E(\mathcal{C}_r)$, and from (18), we get that $[\mathbf{m}]_t = \mu_r[\mathbf{c_k} \cup \mathbf{c_r}]_t$. Since $(\mathbf{c_k} \cup \mathbf{c_r}) \prec \mathbf{m}$, we get that $t \in \operatorname{supp}(\mathbf{m}_+) \cap \operatorname{supp}((\mathbf{c_k} \cup \mathbf{c_r})_+)$ or $t \in \operatorname{supp}(\mathbf{m}_-) \cap \operatorname{supp}((\mathbf{c_k} \cup \mathbf{c_r})_-)$ for $e_t \in E(\mathcal{C}_r)$.

This implies that $\mu_r > 0$ for $r \in X$ because of the equality $[\mathbf{m}]_t = \mu_r [\mathbf{c}_{\mathbf{k}} \cup \mathbf{c}_{\mathbf{r}}]_t$. In a similar way one can show that $\mu_r < 0$ for $r \notin X$.

Let $\mu_r = \frac{p'_r}{d'_r}$ such that $gcd(p'_r, d'_r) = 1$, $p'_r \in \mathbb{Z}$, $d'_r \in \mathbb{N}$. Then $d'_r |[\mathbf{c_k} \cup \mathbf{c_r}]_j$ for $e_j \in E(\mathcal{C}_r)$. Then by definition of d_r , $d'_r | d_r$ and this implies that $\mu_r = \frac{p'_r}{d'_r} = \frac{p''_r}{d_r}$, $p''_r \in \mathbb{Z}$. For $r \in X$, $p''_r > 0$ and let $p''_r = k_r d_r + p_r$, with $0 \le p_r < d_r$, $p_r \in \mathbb{N}$, $k_r \in \mathbb{N} \cup \{0\}$. For $r \notin X$, $p''_r < 0$ and let $p''_r = -k_r d_r + p_r$, with $0 \le p_r < d_r$, $p_r \in \mathbb{N}$, $k_r \in \mathbb{N}$. Thus **m** is of the required form.

Corollary 4.9. Let D be a weighted oriented graph consists of certain number of naturally oriented odd cycles(or unbalanced even cycles) and certain number of balanced cycles share a vertex such that $\prod_{v \in V(\mathcal{C})} w(v)$ has same value for each unbalanced cycle \mathcal{C} in D. Then $\mathcal{M}_D = \mathcal{U}_D = Gr_D = \mathcal{C}_D$.

Proof. Let $f \in Gr_D \setminus \mathcal{C}_D$. Then using Lemma 3.10, $\operatorname{supp}(f_{\mathbf{m}}) \subseteq \bigcup_{\mathcal{C}} E(\mathcal{C})$, where union is taken over all unbalanced cycles \mathcal{C} in D and $\operatorname{supp}(f_{\mathbf{m}})$ contains edges of at least three unbalanced cycles. Assume that $\operatorname{supp}(f_{\mathbf{m}}) = E(D_1)$ where D_1 consists of precisely nunbalanced cycles. Then using Theorem 4.8 and Lemma 3.26, \mathbf{m} is of the form :

$$\sum_{r \in X} k_r(\mathbf{c}_{\mathbf{k}} \cup \mathbf{c}_{\mathbf{r}}) + \sum_{r \notin X} -k_r(\mathbf{c}_{\mathbf{k}} \cup \mathbf{c}_{\mathbf{r}})$$

where $X = \{r : (\mathbf{c_k} \cup \mathbf{c_r}) \prec \mathbf{m}\}, k_r \in \mathbb{N}, C_1, C_2, \dots, C_n \text{ are cycles of } D_1.$ Note that $X \neq \emptyset$. Let $r \in X$. Then $(\mathbf{c_k} \cup \mathbf{c_r}) \prec \mathbf{m}$. Then Using Theorems 3.13 or 3.16 and arguments of proof of Lemma 3.26, we get $[\mathbf{c_k} \cup \mathbf{c_r}]_j = [\mathbf{c_k} \cup \mathbf{c_l}]_j$ for $e_j \in E(\mathcal{C}_k)$. Then $f_{\mathbf{c_k} \cup \mathbf{c_r}}^+ | f_{\mathbf{m}}^+$ and $f_{\mathbf{c_k} \cup \mathbf{c_r}}^- | f_{\mathbf{m}}^-$ which is a contradiction as $f_{\mathbf{m}}$ is primitive. Thus $Gr_D = \mathcal{C}_D$. As I_D is strongly robust, $\mathcal{M}_D = \mathcal{U}_D = Gr_D = \mathcal{C}_D$.

Remark 4.10. In the Corollary 4.9, I_D is independent of the weight of that sharing vertex, that is, if we change weight of this vertex, generators remains same. Thus it is possible that there exists a weighted oriented graph D such that the generators of I_D are independent of the weight of a non-source non-sink vertex.

Corollary 4.11. Let D be a weighted oriented graph consists of certain number of balanced cycles and certain number of unbalanced cycles share a vertex such that each unbalanced cycle has only one non-source non-sink vertex such that all these non-source non-sink vertices have equal weights and the sharing vertex is one of the below type :

- (i) sink of each unbalanced cycle.
- (ii) not a sink of any unbalanced cycle.
- (iii) weight of this vertex is 1.

Then $\mathcal{M}_D = \mathcal{U}_D = Gr_D = \mathcal{C}_D$.

Proof. If C_1, C_2, C_3 are cycles in D, then according to given condition, $[\mathbf{c_1} \cup \mathbf{c_2}]_i = [\mathbf{c_1} \cup \mathbf{c_3}]_i$ for $e_i \in E(C_1)$ using Theorems 3.13 or 3.16 and arguments of proof of Lemma 3.28. Using Theorem 4.8, Lemma 3.28, Corollary 3.29, the proof of arguments is same as in Corollary 4.9.

Remark 4.12. Let *D* be weighted oriented graph as in Corollary 4.11. If *f* is a part of minimal generators of I_D , then exponent of each variable in *f* is 1 or *w*, where *w* is the weight of any non-source non-sink vertex. For explicitly providing form of *f*, take $f_{\mathbf{c}_1 \cup \mathbf{c}_2}$ be a part of minimal generators of I_D corresponding to cycles C_1 , C_2 in *D* and $V(C_1) = \{v_1, v_2, \ldots, v_{m_1}\}$, e_i be the edge incident with v_i and v_{i+1} for $i = 1, 2, \ldots, m_1$ under convention $v_{m_1+1} = v_1$, v_1 is sharing vertex. If v_j is non-source non-sink vertex, $v_j \neq v_1$, then exponent of e_i is 1 for $1 \leq i < j$ and exponent of e_i is *w* for $j \leq i \leq m_1$ and if the sharing vertex is non-source non-sink vertex of both C_1 and C_2 , then exponent of each variable in $f_{\mathbf{c}_1 \cup \mathbf{c}_2}$ is 1. Also I_D is independent of weight of sink vertices.

Corollary 4.13. Let D be weighted oriented graph consists of cycles share a vertex such that V^+ are sinks. Then $\mathcal{M}_D = \mathcal{U}_D = Gr_D = \mathcal{C}_D$.

Proof. Using Lemma 3.10, Theorem 4.8, Lemma 3.24 and using proof of arguments of Corollary 4.9, it follows. \Box

Proposition 4.14. Let D be weighted oriented graph consists of cycles share a vertex such that V^+ are sinks. Let G be the underlying simple graph of D. Then $I_D = I_G$.

Proof. As V^+ are sinks, so every even cycle is balanced. Thus every unbalanced cycle is odd. Let $f_{\mathbf{m}} \in I_D$ be a primitive binomial such that $f_{\mathbf{m}} \neq f_{\mathbf{c}}$ for any even cycle C in D. Then using Lemma 3.10, $\operatorname{supp}(f_{\mathbf{m}}) \subseteq \cup_{\mathcal{C}} E(\mathcal{C})$, where union is taken over all unbalanced cycles \mathcal{C} in D. We will show that $f_{\mathbf{m}} \in (f_{\mathbf{c}_i \cup \mathbf{c}_j} : \mathcal{C}_i, \mathcal{C}_j$ are unbalanced in D). Suppose the number of unbalanced cycles in D is n. We will prove the proposition by induction on n. Basic step : If n = 1, then $I_D = (0)$. If n = 2 and D consists of two unbalanced cycles \mathcal{C}_1 , \mathcal{C}_2 , then $I_D = (f_{\mathbf{c}_1 \cup \mathbf{c}_2})$ by the Theorem 3.13. If n = 3, the result is true by using Corollary 3.25.

Induction hypothesis : Suppose the result is true for any above type graph which consists of precisely n_1 unbalanced cycles, $n_1 < n$.

Inductive step : Let $f_{\mathbf{m}} \in I_D$ be primitive binomial such that $\operatorname{supp}(f_{\mathbf{m}}) \subset \{E(\mathcal{C}) : \mathcal{C} \text{ is unbalanced in } D\}$. Then there is an subgraph D_1 of D such that $\operatorname{supp}(f_{\mathbf{m}}) = E(D_1)$, D_1 consists of precisely t unbalanced cycles sharing a vertex. If t < n, then we are done by induction hypothesis. Assume that t = n. Since $\dim_{\mathbb{Q}}\operatorname{Null}(A(D_1)) = n - 1$ (by the Lemma 3.4), there exists a cycle \mathcal{C}_1 such that \mathbf{m} is \mathbb{Q} -linear combination of n - 1 elements from the set $\{\mathbf{c_1} \cup \mathbf{c_i} : \mathcal{C}_i \text{ is cycle in } D_1\}$. As $\{\mathbf{c_1} \cup \mathbf{c_2}, \mathbf{c_1} \cup \mathbf{c_3}, \ldots, \mathbf{c_1} \cup \mathbf{c_n}\}$ is a linearly independent set, therefore this set is a basis of $\operatorname{Null}(A(D_1))$. Let $\mathbf{m} = \sum_{i=2}^n \lambda_{i-1}(\mathbf{c_1} \cup \mathbf{c_i})$, for some $\lambda_1, \lambda_2, \ldots, \lambda_{n-1} \in \mathbb{Q}$. Without loss of generality we can assume that $(\mathbf{c_1} \cup \mathbf{c_i})|_{E(\mathcal{C}_1)} \prec \mathbf{m}$ for $i = 2, 3, \ldots, n$. Assume that $(\mathbf{c_1} \cup \mathbf{c_i})|_{E(\mathcal{C}_i)} \not\prec \mathbf{m}$ for $i = 2, 3, \ldots, n$. Then $\lambda_i < 0$ for $i = 1, 2, \ldots, n-1$. This implies that $j \in \operatorname{supp}(\mathbf{m}_+) \cap \operatorname{supp}(\mathbf{m}_-)$ for $e_j \in E(\mathcal{C}_1)$ which is a contradiction as $\operatorname{supp}(\mathbf{m}_+) \cap \operatorname{supp}(\mathbf{m}_-) = \emptyset$. Thus there exists a cycle $\mathcal{C}_j(\operatorname{say})$ such that $(\mathbf{c_1} \cup \mathbf{c_j}) \prec \mathbf{m}$.

Since V^+ are sinks, $|[\mathbf{m}]_i| = |[\mathbf{m}]_j|$ for all $e_i, e_j \in E(\mathcal{C}_1)$ and similarly true for \mathcal{C}_j . Then using the Lemma 3.24, we can write $f_{\mathbf{m}} = h_1 f_{\mathbf{c}_1 \cup \mathbf{c}_j} + h_2 f$, for some polynomials h_1, h_2 and $f \in I_{D'}, E(D') = E(D_1) \setminus E(\mathcal{C}_1)$ or $E(D') = E(D_1) \setminus E(\mathcal{C}_j)$. Then by induction hypothesis, $f \in (f_{\mathbf{c}_i \cup \mathbf{c}_j} : \mathcal{C}_i, \mathcal{C}_j \text{ are cycles in } D_1)$. Thus $\{f_{\mathbf{c}}, f_{\mathbf{c}_i \cup \mathbf{c}_j} : \mathcal{C} \text{ is even cycle}, \mathcal{C}_i \text{ and } \mathcal{C}_j \text{ are odd}$ cycles in $D\}$ generates I_D where exponent of each variable in $f_{\mathbf{c}}$ and $f_{\mathbf{c}_i \cup \mathbf{c}_j}$ is 1 i.e. $I_D = I_G$.

Data availability statement: Data sharing is not applicable to this article as no datasets were generated or analyzed during the current study.

References

- A. Boocher, B.C. Brown, T. Duff, L. Lyman, T. Murayama, A. Nesky and K. Schaefer, *Robust graph ideals*, Ann. Comb. **19** (2015), no.4, 641–660.
- J. Biermann, S. Kara, K-N. Lin and A. B. O'Keefe, *Toric ideals of weighted oriented graphs*, Internat. J. Algebra Comput. **32** (2022), no. 2, 307–325.
- [3] A. Boocher, and E. Robeva, Robust toric ideals, Symbolic Comput. 68 (2015), 254–264.
- [4] H. Charalambous, A. Katsabekis and A. Thoma, Minimal systems of binomial generators and the indispensable complex of a toric ideal, Proc. Amer. Math. Soc. 135 (2007), no.11, 3443–3451.

- [5] P. Gimenez, J. Martínez-Bernal, A. Simis, R. H. Villarreal, and C. Vivares, Symbolic powers of monomial ideals and Cohen-Macaulay vertex-weighted digraphs, Singularities, algebraic geometry, commutative algebra, and related topics, 491–510, Springer, Cham, 2018.
- [6] D. R. Grayson and M. E. Stillman, Macaulay2, a software system for research in algebraic geometry. Available at http://www.math.uiuc.edu/Macaulay2/.
- [7] I. Garcia-Marco, and C. Tatakis, On robustness and related properties on toric ideals, J. Algebraic Combin. 57 (2023), no.1, 21–52.
- [8] H. T. Há, K. N. Lin, S. Morey, E. Reyes, and R. H. Villarreal, *Edge ideals of oriented graphs*. Internat. J. Algebra Comput., **29** (2019), no. 3, 535–559.
- [9] S. Kara and J. Biermann, K. N. Lin, and A. O'Keefe, Algebraic invariants of weighted oriented graphs, J. Algebraic. Comb. 55 (2022), no. 2, 461–491.
- [10] M. Kumar and R. Nanduri, Regularity of powers of edge ideals of Cohen-Macaulay weighted oriented forests, J. Algebraic Comb., 58 (2023), no. 3, 867-893.
- [11] D. Kosta, A. Thoma, and M. Vladoiu, On the strongly robust property of toric ideals, J. Algebra 616 (2023), 1–25.
- [12] K. Mori, H. Ohsugi, and A. Tsuchiya, Edge rings with q-linear resolutions, J. Algebra 593(2022), 550–567.
- [13] J. Martínez-Bernal, Y. Pitones and R. H. Villarreal, Minimum distance functions of graded ideals and Reed-Muller-type codes, J. Pure Appl. Algebra 221 (2017), no. 2, 251–275.
- [14] E. N. Miller and B. Sturmfels, Combinatorial commutative algebra, Graduate Texts in Mathematics, 227, Springer, New York, 2005.
- [15] R. Nandi, and R. Nanduri, On regularity bounds and linear resolutions of toric algebras of graphs, J. Commut. Algebra 14(2022), no. 2, 285–296.
- [16] R. Nanduri and T. K. Roy, On combinatorial characterization of generators of toric ideals of weighted oriented graphs, http://arxiv.org/abs/2312.16841v1.
- [17] H. Ohsugi and T. Hibi, Toric ideals generated by quadratic binomials, J. Algebra 218 (1999), no. 2, 509–527.
- [18] H. Ohsugi, and T. Hibi, Indispensable binomials of finite graphs, J. Algebra Appl. 4(2005), no. 4, 421–434.
- [19] Y. Pitones, E. Reyes, and J. Toledo, Monomial ideals of weighted oriented graphs, Electron. J. Combin. 26(2019), no. 3, Paper No. 3.44, 18 pp.
- [20] I. V. Peeva and B. Sturmfels, Generic lattice ideals, J. Amer. Math. Soc. 11 (1998), no. 2, 363–373.
- [21] S. Petrović and D. Stasi, Toric algebra of hypergraphs, J. Algebraic Combin. 39 (2014), no. 1, 187– 208.
- [22] S. Petrović, A. Thoma and M. Vladoiu, Hypergraph encodings of arbitrary toric ideals, J. Combin. Theory Ser. A 166 (2019), 11–41.
- [23] B. Sturmfels, Gröbner bases and convex polytopes, University Lecture Series, 8, Amer. Math. Soc., Providence, RI, (1995).

RAMAKRISHNA NANDURI AND TAPAS KUMAR ROY

- [24] S. Sullivant, Strongly robust toric ideals in codimension 2, J. Algebr. Stat. 10 (2019), no.1, 128–136.
- [25] C. Tatakis, Generalized robust toric ideals, J. Pure Appl. Algebra 220 (2016), no.1, 263–277.
- [26] A. Takemura and S. Aoki, Some characterizations of minimal Markov basis for sampling from discrete conditional distributions, Ann. Inst. Statist. Math. 56 (2004), no.1, 1–17.
- [27] R. H. Villarreal, Rees algebras of edge ideals, Comm. Algebra 23 (1995), no. 9, 3513–3524.
- [28] R. H. Villarreal, Monomial algebras, Monographs and Textbooks in Pure and Applied Mathematics, 238, Marcel Dekker, Inc., New York, 2001.
- [29] G. Zhu, L. Xu, H. Wang, and Z. Tang, Projective dimension and regularity of edge ideals of some weighted oriented graphs, Rocky Mountain J. Math. 49 (2019), no. 4, 1391–1406.

DEPARTMENT OF MATHEMATICS, INDIAN INSTITUTE OF TECHNOLOGY KHARAGPUR, WEST BEN-GAL, INDIA - 721302.

Email address: nanduri@maths.iitkgp.ac.in

DEPARTMENT OF MATHEMATICS, INDIAN INSTITUTE OF TECHNOLOGY KHARAGPUR, WEST BEN-GAL, INDIA - 721302.

Email address: tapasroy147@kgpian.iitkgp.ac.in