

TIT/HEP-700

April 2024

Brane expansions for anti-symmetric line operator index

Yosuke IMAMURA*and Masato INOUE†

*Department of Physics, Tokyo Institute of Technology,
Tokyo 152-8551, Japan*

Abstract

Based on the D5-brane realization of Wilson line operators in anti-symmetric representations, we propose brane expansion formulas for $I_{N,k}$, the Schur index of $\mathcal{N} = 4$ $U(N)$ SYM decorated by line operators in the anti-symmetric representation of rank k . For the large N index $I_{\infty,k}$ we propose a double-sum expansion, and for finite N index $I_{N,k}$ we propose a quadruple-sum expansion. Objects causing finite k and finite N corrections are disk D3-branes ending on the D5-brane.

*E-mail: imamura@phys.titech.ac.jp

†E-mail: masatoinoue@phys.titech.ac.jp

Contents

1	Introduction	1
2	Finite k corrections	3
2.1	Large N and large k limit	3
2.2	Full expansion	5
2.3	Reduced expansion	7
2.4	Half BPS limit	8
3	Finite N corrections	8
3.1	Full expansion	8
3.2	Reduced expansion	10
3.3	Half BPS limit	12
4	Conclusions and discussion	12
A	Character expansion method	14

Contents

1 Introduction

Wilson lines are the most basic and important observables in gauge theories. They serve as order parameters for detecting phase transitions. They are also useful in gaining deeper insights into phenomena such as dualities. In this work, we study Wilson line operators of $\mathcal{N} = 4$ $U(N)$ SYM within the framework of the AdS/CFT correspondence [1, 2, 3].

Wilson line operators are labeled by representations of the gauge group. We consider $U(N)$ gauge theories, and irreducible representations are specified by partitions μ ($\ell(\mu) \leq N$), which are depicted as Young diagrams. The number of boxes $k = |\mu|$ in μ is referred to as the rank of the representation.

Different holographic realizations exist for Wilson lines. Wilson line operators in the fundamental representation ($\mu = \{1\}$) are realized by fundamental strings in the AdS space [4, 5]. If the rank of the representation is of order N , the coincident strings are more naturally described by D-branes. A Wilson line in the symmetric representation with $\mu = \{k\}$ is related to a tubular D3-brane with a cross section $S^2 \subset AdS_5$ [6] (See also a discussion in Section 4). For the anti-symmetric representation with $\mu = \{1^k\}$, the corresponding object is a tubular D5-brane with a cross section $S^4 \subset S^5$ [7]. In both cases, the charge of k fundamental strings is carried by the electric flux on the D-brane. If the rank is of order N^2 , the strings or D-branes deform the background spacetime significantly, and the line operators are naturally described by bubbling geometries [8].

Among these realizations of line operators, we specifically focus on the anti-symmetric line operators realized by D5-branes. We investigate them by computing the Schur index [9], which is a specialization of the superconformal index [10, 11]. See [12, 13, 14] for analytic formulas of the Schur index without line operator insertion. These indices are regarded as the partition function in $S^3 \times S^1$, and line operator indices [15, 16, 17, 18, 19, 20, 21] are defined as the partition functions with insertion of BPS line operators. We insert $1/2$ BPS Wilson line operators at the poles of S^3 , thereby breaking the rotational symmetry $so(4)$ of S^3 to $so(3)$. The R-symmetry $so(6)$ is also broken to $so(5)$. Let J_1 be the Cartan generators of the $so(3)$, and R_x and R_y be the Cartan generators of the $so(5)$. The Schur index of $\mathcal{N} = 4$ SYM is defined using these generators by

$$\text{tr}[(-1)^F q^{J_1} x^{R_x} y^{R_y}], \quad q = xy. \quad (1)$$

We can regard the index with line operator insertion as a correlator of the line operators in the spacetime $S^3 \times S^1$, and we use the notation $\langle \dots \rangle^{U(N)}$. We can calculate the index with line operators in a representation μ and the conjugate of a representation ν inserted at the north and the south poles, respectively, by the localization formula

$$\langle s_\mu(U) s_\nu(U^{-1}) \rangle^{U(N)} = \int_{U(N)} dU \text{Pexp}(i_{\text{vec}} \chi_{\text{adj}}^{U(N)}(U)) s_\mu(U) s_\nu(U^{-1}), \quad (2)$$

where i_{vec} is the letter index of the $\mathcal{N} = 4$ vector multiplet

$$i_{\text{vec}} = 1 - \frac{(1-x)(1-y)}{1-q}, \quad (3)$$

and $\chi_{\text{adj}}^{U(N)}(U)$ is the $U(N)$ adjoint character

$$\chi_{\text{adj}}^{U(N)}(U) = \text{tr}(U) \text{tr}(U^{-1}). \quad (4)$$

Pexp is the plethystic exponential, and $\int_{U(N)} dU$ is the integral over the $U(N)$ gauge group manifold with the Haar measure. $s_\mu(U)$ is the $U(N)$ character of an irreducible representation μ . For a diagonalized matrix $U = \text{diag}(z_1, \dots, z_N)$ $s_\mu(U)$ are symmetric polynomials of the N gauge fugacities z_a ($a = 1, \dots, N$), which are called Schur polynomials. We will abuse this term to refer to $s_\mu(U)$, too. On the left hand side in (2) we express the line operators by the corresponding Schur polynomials.

The line operator index we are interested in is

$$I_{N,k} = \langle s_{\{1^k\}}(U) s_{\{1^k\}}(U^{-1}) \rangle^{U(N)} \quad (0 \leq k \leq N). \quad (5)$$

It is well known that the large N index without insertion, $I_{\infty,0}$, is reproduced as the supergravity index on the AdS side [11];

$$I_{\infty,0} = I_{\text{sugra}} = \text{Pexp} \left(\frac{x}{1-x} + \frac{y}{1-y} - \frac{q}{1-q} \right). \quad (6)$$

For the insertion of fundamental line operators, it was confirmed in [16] that the index $I_{\infty,1}$ is reproduced by including the contribution from the fluctuation modes on a fundamental string extended along AdS_2 analyzed in [22].

$$\frac{I_{\infty,1}}{I_{\text{sugra}}} = I_{\text{F1}} = \text{Pexp}(x + y - q) = 1 + \frac{x}{1-x} + \frac{y}{1-y}. \quad (7)$$

For the finite N index $I_{N,0}$ without line operator insertion, the finite N corrections are reproduced by D3-branes wrapped around topologically trivial three-cycles in S^5 , called giant gravitons [23, 24]. The finite N index can be given in the form of expansions with respect to the number of giant gravitons, and such expansions are called giant graviton expansions [25, 26, 27, 28]. See also [29, 30, 31, 32].

Such expansions were generalized to the line operator index of the fundamental representation $I_{N,1}$ in [33]. (See also [34].)

The purpose of this paper is to propose similar holographic relations for $I_{N,k}$ with finite N and finite k based on the D5-brane realization of anti-symmetric line operators.

We will heavily rely on numerical analysis in which we treat indices as Taylor series. We use two different Taylor series with respect to two different variables. One is the y -series in the form

$$\sum_{i=0}^{\infty} f_i(x) y^i, \quad (8)$$

and the other is the t -series in the form

$$\sum_{i=0}^{\infty} f_i(u) t^i, \quad (9)$$

where t and u are defined by the relations

$$x = tu, \quad y = tu^{-1}. \quad (10)$$

In the following we mainly use the variables x and y . When we consider t -expansion of a function, we implicitly regard it as a function of t and u , even if it is written in terms of x and y .

2 Finite k corrections

2.1 Large N and large k limit

Let us first consider the large N index $I_{\infty,k}$ given by the formula [18]

$$\frac{I_{\infty,k}}{I_{\text{sugra}}} = \sum_{\lambda \vdash k} \frac{1}{z_\lambda} [I_{\text{F1}}]_\lambda. \quad (11)$$

(See Appendix for a derivation.) $\sum_{\lambda \vdash k}$ is the summation over partitions of an integer k , and $[\cdots]_\lambda$ is defined by

$$[f(x, y)]_\lambda = \prod_{i=1}^{\ell(\lambda)} f(x^{\lambda_i}, y^{\lambda_i}) \quad (12)$$

for a function $f(x, y)$ of fugacities x and y . z_λ is defined for a partition $\lambda = \{\lambda_1, \lambda_2, \dots, \lambda_{\ell(\lambda)}\} = \{\dots, 3^{r_3}, 2^{r_2}, 1^{r_1}\}$ by $z_\lambda = \prod_{m=1}^{\infty} m^{r_m} r_m!$. The t -series expansions of $I_{\infty, k}/I_{\text{sugra}}$ for small k are

$$\begin{aligned} I_{\infty, 0}/I_{\text{sugra}} &= 1, \\ I_{\infty, 1}/I_{\text{sugra}} &= 1 + 2t + 2t^2 + 2t^3 + 2t^4 + 2t^5 + 2t^6 + 2t^7 + \mathcal{O}(t^8), \\ I_{\infty, 2}/I_{\text{sugra}} &= 1 + 2t + 5t^2 + 6t^3 + 9t^4 + 10t^5 + 13t^6 + 14t^7 + \mathcal{O}(t^8), \\ I_{\infty, 3}/I_{\text{sugra}} &= 1 + 2t + 5t^2 + 10t^3 + 15t^4 + 22t^5 + 31t^6 + 40t^7 + \mathcal{O}(t^8), \\ I_{\infty, 4}/I_{\text{sugra}} &= 1 + 2t + 5t^2 + 10t^3 + 20t^4 + 30t^5 + 48t^6 + 68t^7 + \mathcal{O}(t^8), \\ I_{\infty, 5}/I_{\text{sugra}} &= 1 + 2t + 5t^2 + 10t^3 + 20t^4 + 36t^5 + 58t^6 + 90t^7 + \mathcal{O}(t^8), \\ I_{\infty, 6}/I_{\text{sugra}} &= 1 + 2t + 5t^2 + 10t^3 + 20t^4 + 36t^5 + 65t^6 + 102t^7 + \mathcal{O}(t^8). \end{aligned} \quad (13)$$

We showed unrefined index with $x, y \rightarrow t$ to save the space. We can clearly see the convergence in the limit $k \rightarrow \infty$.

The analytic expression for the large k limit is obtained as follows. Notice that we can regard the right hand side of (11) as the partition function of k identical particles. If we take the large k limit with fixed energy, almost all particles are in the ground state (Bose-Einstein condensate), and the number of excited particles is not constrained. The ground state is given by the term “1” in (7), and particles in the ground state do not contribute to the index. Therefore, the large k limit of the index becomes [16]

$$\begin{aligned} \lim_{k \rightarrow \infty} \frac{I_{\infty, k}}{I_{\text{sugra}}} &= \text{Pexp}(I_{\text{F1}} - 1) \\ &= 1 + 2t + 5t^2 + 10t^3 + 20t^4 + 36t^5 + 65t^6 + 110t^7 + \mathcal{O}(t^8). \end{aligned} \quad (14)$$

In the second line we again showed the unrefined index.

This result agrees with the index of the fluctuation modes on the D5-brane. The insertion of large k anti-symmetric line operators is realized on the AdS side by introducing a D5-brane extended along $AdS_2 \times S^4$ [7]. The radius of S^4 is given by $L_{\text{AdS}} \sin \theta_{\text{D5}}$ with the angle θ_{D5} determined by the relation

$$\frac{k}{N} = \frac{1}{\pi} (\theta_{\text{D5}} - \sin \theta_{\text{D5}} \cos \theta_{\text{D5}}). \quad (15)$$

If k is of order N , the radius of S^4 is comparable to the AdS radius L_{AdS} , and we can treat the brane as a semi-classical object. Mode analysis of the fields on the D5-brane was performed in [22, 35], and the corresponding index

$$I_{\text{D5}} = \text{Pexp} \left(\frac{x}{1-x} + \frac{y}{1-y} \right) \quad (16)$$

agrees with (14).

2.2 Full expansion

In the large k limit, the index is given by (14). We are interested in the finite k correction of $\mathcal{O}(t^{k+1})$ in the index $I_{\infty,k}$.

$$\frac{I_{\infty,k}}{I_{\text{sugra}} I_{\text{D5}}} = 1 + \mathcal{O}(t^{k+1}). \quad (17)$$

What are objects causing the finite k correction on the AdS side?

Let \mathcal{C}_x and \mathcal{C}_y be the R_x -fixed locus and R_y -fixed locus in S^5 , respectively. They are homologically trivial S^3 in S^5 . In the giant graviton expansion of the Schur index without line operator insertion [26], D3-branes wrapped around \mathcal{C}_x and \mathcal{C}_y give finite N corrections to the index. Let us define the latitude θ ($0 \leq \theta \leq \pi$) in S^5 so that two poles $\theta = 0$ and $\theta = \pi$ are located on the intersection $\mathcal{C}_x \cap \mathcal{C}_y$. The metric of S^5 is

$$ds_{S^5}^2 = L_{\text{AdS}}^2 (d\theta^2 + \sin^2 \theta ds_{S^4}^2). \quad (18)$$

A cross section of the tubular D5-brane is $\theta = \theta_{\text{D5}}$, and the three-cycle \mathcal{C}_x is divided by the D5-brane into two parts $0 \leq \theta \leq \theta_{\text{D5}}$ and $\theta_{\text{D5}} \leq \theta \leq \pi$, which we denote by \mathcal{D}_x and \mathcal{D}'_x . Similarly, \mathcal{C}_y is divided into two parts \mathcal{D}_y and \mathcal{D}'_y . (See Figure 1.)

A suggestive fact is that the right hand side of (15) is the ratio of the volumes

$$\frac{k}{N} = \frac{\text{vol}(\mathcal{D}_x)}{\text{vol}(\mathcal{C}_x)} = \frac{\text{vol}(\mathcal{D}_y)}{\text{vol}(\mathcal{C}_y)}. \quad (19)$$

It is essentially important in the giant graviton expansion that the mass of a D3-brane wrapped around \mathcal{C}_x and \mathcal{C}_y is N/L_{AdS} , and the corresponding operator dimension is N . (19) means that the mass of a D3-brane on \mathcal{D}_x and \mathcal{D}_y is k/L_{AdS} , and this fact suggests that such D3-branes give the finite k corrections to the index.

Based on analogy with giant graviton expansions for finite N corrections to $I_{N,0}$, we propose the expansion

$$\frac{I_{\infty,k}}{I_{\text{sugra}} I_{\text{D5}}} = \sum_{m_x, m_y=0}^{\infty} x^{km_x} y^{km_y} \mathcal{F}_{m_x, m_y}, \quad (20)$$




Figure 1: The small circle is a cross-section (S^4) of the tubular D5-brane. The R_x -fixed locus \mathcal{C}_x shown as the horizontal circle is divided into two parts \mathcal{D}_x and \mathcal{D}'_x by the D5-brane. Similarly, the R_y -fixed locus \mathcal{C}_y shown as the vertical circle is divided into \mathcal{D}_y and \mathcal{D}'_y . Some important features are not shown correctly. For example, \mathcal{D}_x and \mathcal{D}_y intersect not at a point but along a segment, and \mathcal{D}_x and \mathcal{D}'_y are not separated as shown but in contact at two points.

where m_x and m_y are the numbers of disk D3-branes on \mathcal{D}_x and \mathcal{D}_y , respectively. In this section we consider the large N limit with finite k , and then $k' = N - k$ is infinite and D3-branes on \mathcal{D}'_x and \mathcal{D}'_y are decoupled. k -independent functions \mathcal{F}_{m_x, m_y} are the indices of the theories realized on the system consisting of m_x D3-branes on \mathcal{D}_x and m_y D3-branes on \mathcal{D}_y (and the tubular D5-brane).

Because the branes causing the finite k correction are not giant gravitons but disk D3-branes, we call (20) (and similar expansions we will see below) “a brane expansion” rather than a giant graviton expansion.

Let us first consider the functions $\mathcal{F}_{m,0}$, which are the index of the theories realized on \mathcal{D}_x . Such an index of theories living on a three-dimensional disk is known as the half-index [15], and for $\mathcal{N} = 4$ $U(m)$ SYM it is given by

$$\Pi_m = \int_{U(m)} dU \text{Pexp}(i_{\text{half}} \chi_{\text{adj}}^{U(m)}(U)), \quad i_{\text{half}} = \frac{x - q}{1 - q}. \quad (21)$$

Analytic form of the integral is¹

$$\Pi_m = \text{Pexp} \left(\frac{x^m - q}{1 - q} + \frac{x - x^m}{1 - x} \right). \quad (22)$$

¹We did not analytically derive this analytic form, but only checked the agreement of Taylor expansions up to a very high order.

We need to apply the variable change

$$\sigma_x : (x, y) \rightarrow (x^{-1}, xy) \quad (23)$$

to translate the index of the boundary theory to the index on the theory on the R_x -fixed locus [25, 26, 28]. Namely, $\mathcal{F}_{m,0}$ should be given by $\mathcal{F}_{m,0} = \sigma_x \Pi_m$. With the analytic expression (22) for Π_m it is easy to calculate $\sigma_x \Pi_m$ explicitly. For small m the explicit forms of t -expansion of $\mathcal{F}_{m,0}$ are

$$\begin{aligned} \mathcal{F}_{1,0} &= \frac{-x}{1 - \frac{y}{x}} + \frac{x^2(-1 + \frac{y}{x} - \frac{y^2}{x^2})}{1 - \frac{y}{x}} + \frac{x^3(-1 + \frac{y}{x} - \frac{y^2}{x^2})}{1 - \frac{y}{x}} + \frac{x^4(-1 + \frac{y}{x} - \frac{y^2}{x^2})}{1 - \frac{y}{x}} + \mathcal{O}(t^5), \\ \mathcal{F}_{2,0} &= \frac{-\frac{x^5}{y}}{1 - \frac{y^2}{x^2}} + \frac{\frac{x^7}{y^2}(-1 + \frac{y^2}{x^2} - \frac{y^3}{x^3})}{1 - \frac{y}{x}} + \frac{\frac{x^9}{y^3}(-1 - \frac{y}{x} - \frac{y^2}{x^2} + \frac{y^3}{x^3} - \frac{y^5}{x^5} - \frac{y^8}{x^8})}{1 - \frac{y^2}{x^2}} + \mathcal{O}(t^7), \\ \mathcal{F}_{3,0} &= \frac{-\frac{x^{12}}{y^3}}{1 - \frac{y^3}{x^3}} + \frac{\frac{x^{15}}{y^5}(-1 - \frac{y^2}{x^2} + \frac{y^3}{x^3} - \frac{y^6}{x^6})}{1 - \frac{y^3}{x^3}} + \frac{\frac{x^{18}}{y^7}(-1 - \frac{y^2}{x^2} - 2\frac{y^4}{x^4} + \frac{y^5}{x^5} - \frac{y^8}{x^8} - \frac{y^{12}}{x^{12}})}{1 - \frac{y^3}{x^3}} + \mathcal{O}(t^{12}) \end{aligned} \quad (24)$$

$\mathcal{F}_{0,m}$ are obtained from $\mathcal{F}_{m,0}$ by exchanging x and y ; $\mathcal{F}_{0,m} = \mathcal{F}_{m,0}|_{x \leftrightarrow y}$.

If both m_x and m_y are positive, we naively expect that \mathcal{F}_{m_x, m_y} is the index of a $U(m_x) \times U(m_y)$ gauge theory with bi-fundamental fields living on the intersection $\mathcal{D}_x \cap \mathcal{D}_y$. The intersection is a segment, and we need to carefully determine the boundary conditions at the ends of the segment to calculate the letter index of the bi-fundamental fields. We will not examine the boundary conditions here. In fact, numerical analysis shows that only $\mathcal{F}_{m_x,0}$ and \mathcal{F}_{0,m_y} are sufficient to reproduce the finite k index in (11), and we do not have to include the contributions from intersecting branes. This means $\mathcal{F}_{m_x, m_y} = 0$ for $m_x, m_y \geq 1$, and strongly suggests that the supersymmetry is spontaneously broken in the theory realized on $\mathcal{D}_x \cup \mathcal{D}_y$.

In summary, the brane expansion of $I_{\infty,k}$ is

$$\frac{I_{\infty,k}}{I_{\text{sugra}} I_{\text{D5}}} = 1 + \sum_{m=1}^{\infty} x^{mk} \mathcal{F}_{m,0} + \sum_{m=1}^{\infty} y^{mk} \mathcal{F}_{0,m}. \quad (25)$$

We numerically confirmed that this relation holds for $k \leq 20$ up to $\mathcal{O}(t^{20})$.

2.3 Reduced expansion

For the Schur index without line insertion the double sum expansion in [26] is reduced to simple-sum expansion if we treat the index as a y -series [28]. See also [36, 37] for the mechanism of the reduction. This reduction mechanism works for the brane expansion of the line-operator index (25). We can easily confirm that y -series expansion of $\mathcal{F}_{0,m} = (\sigma_x \Pi_m)|_{x \leftrightarrow y}$ ($m \geq 1$) is zero, and

(25) reduces to the simple-sum expansion

$$\frac{I_{\infty,k}}{I_{\text{sugra}}I_{\text{D5}}} = \sum_{m=0}^{\infty} x^{km} \mathcal{F}_{m,0}. \quad (26)$$

$\mathcal{F}_{m,0}$ expressed as y -series are shown for small m below.

$$\begin{aligned} \mathcal{F}_{1,0} &= \frac{-x}{1-x} - y + \left(-\frac{1}{x} - 1\right)y^2 + \left(-\frac{1}{x^2} - \frac{1}{x}\right)y^3 + \mathcal{O}(y^4), \\ \mathcal{F}_{2,0} &= \frac{x^3}{(1-x)(1-x^2)} + \frac{x}{1-x}y + \frac{\frac{1}{x}+x}{1-x}y^2 + \frac{\frac{1}{x^3}+\frac{1}{x}}{1-x}y^3 + \mathcal{O}(y^4), \\ \mathcal{F}_{3,0} &= \frac{-x^6}{(1-x)(1-x^2)(1-x^3)} + \frac{-x^3}{(1-x)(1-x^2)}y + \frac{1+x-x^2}{(1-x)^2}y^2 + \mathcal{O}(y^3). \end{aligned} \quad (27)$$

We numerically confirmed that (26) holds for $k \leq 20$ up to $\mathcal{O}(y^{10})$ and $\mathcal{O}(x^{20})$.

2.4 Half BPS limit

We can simplify the problem by taking the half-BPS limit $y \rightarrow 0$. Then, functions appearing in (26) take simple forms

$$\frac{I_{\infty,n}}{I_{\text{sugra}}} = \Pi_n = \text{Pexp}(x + \cdots + x^n), \quad I_{\text{D5}} = I_{\text{sugra}} = \text{Pexp} \frac{x}{1-x}, \quad (28)$$

and we can analytically confirm (26). In fact, the simple-sum brane expansion (26) is identical with the simple-sum giant graviton expansion of the Schur index without line insertion [28] in the half BPS limit.

3 Finite N corrections

3.1 Full expansion

If N is finite, in addition to disk D3-branes on \mathcal{D}_x and \mathcal{D}_y , ones on \mathcal{D}'_x and \mathcal{D}'_y also contribute to the index. We expect the quadruple-sum brane expansion

$$\frac{I_{N,k}}{I_{\text{sugra}}I_{\text{D5}}} = \sum_{m_x, m_y, m'_x, m'_y=0}^{\infty} x^{km_x+k'm'_x} y^{km_y+k'm'_y} \mathcal{F}_{m_x, m_y, m'_x, m'_y}, \quad (29)$$

where $k' = N - k$. Functions $\mathcal{F}_{m_x, m_y, m'_x, m'_y}$ are not all independent but satisfy

$$\mathcal{F}_{m_x, m_y, m'_x, m'_y} = \mathcal{F}_{m'_x, m'_y, m_x, m_y} = \mathcal{F}_{m_y, m_x, m'_y, m'_x}|_{x \leftrightarrow y} \quad (30)$$

and by definition

$$\mathcal{F}_{m_x, m_y, 0, 0} = \mathcal{F}_{m_x, m_y}. \quad (31)$$

We have not yet fully understood how to calculate the functions $\mathcal{F}_{m_x, m_y, m'_x, m'_y}$ for general wrapping numbers. In this subsection, we will calculate the t -series expansion of $I_{N, k}$ on the gauge theory side for different values of N and k , and we extract functions $\mathcal{F}_{m_x, m_y, m'_x, m'_y}$ for small wrapping numbers by assuming the expansion (29).

Let m_{tot} be the total wrapping number; $m_{\text{tot}} = m_x + m_y + m'_x + m'_y$. There are four functions for $m_{\text{tot}} = 1$, and they are all related to $\mathcal{F}_{1, 0, 0, 0} = \mathcal{F}_{1, 0} = \sigma_x \Pi_1$ by (30).

For $m_{\text{tot}} = 2$, there are four essentially different functions

$$\mathcal{F}_{2, 0, 0, 0}, \quad \mathcal{F}_{1, 1, 0, 0}, \quad \mathcal{F}_{1, 0, 1, 0}, \quad \mathcal{F}_{1, 0, 0, 1}. \quad (32)$$

The first two are the same as the functions given in the previous section

$$\begin{aligned} \mathcal{F}_{2, 0, 0, 0} &= \mathcal{F}_{2, 0} = \sigma_x \Pi_2, \\ \mathcal{F}_{1, 1, 0, 0} &= \mathcal{F}_{1, 1} = 0, \end{aligned} \quad (33)$$

and the other two extracted from the comparison to $I_{N, k}$ are

$$\begin{aligned} \mathcal{F}_{1, 0, 1, 0} &= \frac{x^2(1 + \frac{y}{x})}{(1 - \frac{y}{x})^2} + \frac{x^3(2 - 2\frac{y}{x} + 2\frac{y^2}{x^2} + 2\frac{y^3}{x^3})}{(1 - \frac{y}{x})^2} \\ &\quad + \frac{x^4(3 - 4\frac{y}{x} + 4\frac{y^2}{x^2} - 6\frac{y^3}{x^3} + 6\frac{y^4}{x^4} + 3\frac{y^5}{x^5})}{(1 - \frac{y}{x})^2} + \mathcal{O}(t^5), \\ \mathcal{F}_{1, 0, 0, 1} &= \frac{2xy}{(1 - \frac{y}{x})(1 - \frac{x}{y})} + \frac{2(x^3 + y^3)}{(1 - \frac{y}{x})(1 - \frac{x}{y})} \\ &\quad + \frac{2(\frac{x^5}{y} + x^4 - x^3y + x^2y^2 - xy^3 + y^4 + \frac{y^5}{x})}{(1 - \frac{y}{x})(1 - \frac{x}{y})} + \mathcal{O}(t^5). \end{aligned} \quad (34)$$

These are expected to be the indices of $U(1) \times U(1)$ gauge theories with bi-fundamental fields.

The brane system corresponding to $\mathcal{F}_{1, 0, 1, 0}$ consists of two disks \mathcal{D}_x and \mathcal{D}'_x , and they are adjacent on $\mathcal{D}_x \cap \mathcal{D}'_x = S^2$. There exists a bi-fundamental hypermultiplet on S^2 . Indeed, we can confirm that $\mathcal{F}_{1, 0, 1, 0}$ in (34) is given by

$$\mathcal{F}_{1, 0, 1, 0} = \mathcal{F}_{1, 0}^2 \sigma_x I_{3d}, \quad (35)$$

where I_{3d} is the 3d index [38, 39] of a charged hypermultiplet

$$I_{3d} = \int \frac{dz}{2\pi i z} \text{Pexp} \left[i_{3d} \left(z + \frac{1}{z} \right) \right], \quad i_{3d} = \frac{y^{\frac{1}{2}}(1-x)}{1-q}. \quad (36)$$

$\mathcal{F}_{1,0,0,1}$ is the index of the theory realized on $\mathcal{D}_x \cup \mathcal{D}'_y$. Two disks \mathcal{D}_x and \mathcal{D}'_y are touching each other at two points, and bi-fundamental fields live on the points. We find the relation

$$\mathcal{F}_{1,0,0,1} = 2\mathcal{F}_{1,0}\mathcal{F}_{0,1} \quad (37)$$

holds, and identify the factor 2 as the contribution from the bi-fundamental fields. Indeed, a pair of bi-fundamental fermion with the letter index $-(z + \frac{1}{z})$ gives this factor

$$2 = \int \frac{dz}{2\pi iz} \text{Pexp} \left[- \left(z + \frac{1}{z} \right) \right]. \quad (38)$$

Using these functions $\mathcal{F}_{m_x, m_y, m'_x, m'_y}$ for $m_{\text{tot}} \leq 2$, the quadruple-sum expansion (29) correctly reproduces $I_{N,k}$ up to order $\mathcal{O}(t^n)$ with $n = \min(2k + k', k + 2k')$.

For $m_{\text{tot}} = 3$, there are five essentially different functions:

$$\mathcal{F}_{3,0,0,0}, \quad \mathcal{F}_{2,1,0,0}, \quad \mathcal{F}_{2,0,1,0}, \quad \mathcal{F}_{2,0,0,1}, \quad \mathcal{F}_{1,1,1,0}. \quad (39)$$

The first two are given by $\mathcal{F}_{3,0,0,0} = \mathcal{F}_{3,0} = \sigma_x \Pi_3$ and $\mathcal{F}_{2,1,0,0} = \mathcal{F}_{2,1} = 0$, and we can extract the other three by assuming the quadruple-sum expansion (29). The leading terms in the t -expansions are

$$\begin{aligned} \mathcal{F}_{2,0,1,0} &= \frac{x^4}{(1 - \frac{y}{x})^2 (1 - \frac{y^2}{x^2})} + \mathcal{O}(t^5), \\ \mathcal{F}_{2,0,0,1} &= \frac{-x^3 y}{(1 - \frac{y}{x})^2 (1 - \frac{y^2}{x^2})} + \mathcal{O}(t^5), \\ \mathcal{F}_{1,1,1,0} &= \frac{-xy^3}{(1 - \frac{y}{x})^3} + \mathcal{O}(t^5). \end{aligned} \quad (40)$$

Although it would also be possible to obtain higher order terms in (40) and functions for $m_{\text{tot}} \geq 4$ by continuing a similar analysis, we stop here the analysis of the quadruple-sum expansion (29).

3.2 Reduced expansion

We can reduce the quadruple-sum expansion (29) to the double-sum expansion

$$\frac{I_{N,k}}{I_{\text{sugra}} I_{\text{D5}}} = \sum_{m, m'=0}^{\infty} x^{km+k'm'} \mathcal{F}_{m,0,m',0}, \quad (41)$$

by treating the index as a y -series. In this expansion only the R_x -fixed locus \mathcal{C}_x contributes to the index, and the summation is taken over the two wrapping numbers m and m' associated with \mathcal{D}_x and \mathcal{D}'_x , respectively.

$\mathcal{F}_{m,0,m',0}$ is the index of the theory on \mathcal{C}_x divided by the D5-brane to \mathcal{D}_x and \mathcal{D}'_x . We take the S-duality frame in which the tubular D5-brane becomes an NS5-brane. Then, the gauge theory on \mathcal{C}_x is the quiver gauge theory with gauge group $U(m) \times U(m')$. The vector multiplets live in two disks \mathcal{D}_x and \mathcal{D}'_x , and the bi-fundamental hypermultiplet lives on the wall dividing \mathcal{C}_x into the two parts. The functions $\mathcal{F}_{m,0,m',0}$ are given by the localization formula

$$\mathcal{F}_{m,0,m',0} = \sigma_x \left[\int_{U(m)} dU \int_{U(m')} dU' \text{Pexp} \left(i_{\text{half}} \chi_{\text{adj}}^{U(m)}(U) + i_{\text{half}} \chi_{\text{adj}}^{U(m')}(U') + i_{3d} \chi_{\text{bifund}}^{U(m) \times U(m')}(U, U') \right) \right], \quad (42)$$

where the bi-fundamental character is defined by

$$\chi_{\text{bifund}}^{U(m) \times U(m')}(U, U') = \text{tr}(U) \text{tr}(U'^{-1}) + \text{tr}(U^{-1}) \text{tr}(U'). \quad (43)$$

The first few terms in the y -series expansions of the functions for small m_{tot} are

$$\begin{aligned} \mathcal{F}_{1,0,1,0} &= \frac{x^2}{(1-x)^2} + \frac{x(3-x)}{1-x}y + \frac{5-x+x^2-x^3}{1-x}y^2 + \mathcal{O}(y^3), \\ \mathcal{F}_{2,0,1,0} &= \frac{-x^4}{(1-x)^2(1-x^2)} + \frac{x^2(-2-2x+x^2)}{(1-x)(1-x^2)}y \\ &\quad + \frac{-2-2x+2x^2-2x^3+x^4}{(1-x)^2}y^2 + \mathcal{O}(y^3), \\ \mathcal{F}_{3,0,1,0} &= \frac{x^7}{(1-x)^2(1-x^2)(1-x^3)} + \frac{x^4(2+x+2x^2-x^3)}{(1-x)(1-x^2)(1-x^3)}y \\ &\quad + \frac{x(2+2x+3x^2-x^3+2x^4+x^6-x^7)}{(1-x)(1-x^2)(1-x^3)}y^2 + \mathcal{O}(y^3), \\ \mathcal{F}_{2,0,2,0} &= \frac{x^6}{(1-x)^2(1-x^2)^2} + \frac{x^3(1+2x-x^2)}{(1-x)^2(1-x^2)}y \\ &\quad + \frac{2x(1+3x-3x^2+2x^4-x^5)}{(1-x)^2(1-x^2)}y^2 + \mathcal{O}(y^3). \end{aligned} \quad (44)$$

See (27) for $\mathcal{F}_{m,0,0,0} = \mathcal{F}_{m,0} = \sigma_x \Pi_m$.

We numerically confirmed the expansion (41) works well. For example, for $N = 2$ and $k = 1$, if we sum up the contributions from $m_{\text{tot}} \leq 4$, the discrepancy is

$$\frac{I_{2,1}}{I_{\text{sugra}} I_{\text{D5}}} - \sum_{m+m' \leq 4} (\dots) = \mathcal{O}(x^{14})y^0 + \mathcal{O}(x^{10})y^1 + \mathcal{O}(x^7)y^2 + \mathcal{O}(x^4)y^3 + \mathcal{O}(y^4), \quad (45)$$

and a lot of terms are correctly reproduced.

3.3 Half BPS limit

The half-BPS limit $y \rightarrow 0$ of $I_{N,k}$ for general values of N and k is given by [20]

$$I_{N,k} = \prod_{m=1}^k \frac{1}{1-x^m} \prod_{m=1}^{N-k} \frac{1}{1-x^m}. \quad (46)$$

The expansion (41) reproduces this factorized form. In the half-BPS limit the letter index $\sigma_x i_{3d}$ of the hypermultiplet on the wall vanishes, and the wall contribution becomes trivial. As the result, the functions $\mathcal{F}_{m,0,m',0}$ are factorized to the product of $\mathcal{F}_{m,0}$ and $\mathcal{F}_{m',0}$;

$$\mathcal{F}_{m,0,m',0} = \mathcal{F}_{m,0} \mathcal{F}_{m',0} = \sigma_x (\Pi_m \Pi_{m'}). \quad (47)$$

Then, the double-sum expansion (41) factorizes into two copies of the simple-sum expansion (26) for the large N index

$$\frac{I_{N,k}}{I_{\text{sugra}} I_{\text{D5}}} = \left(\sum_{m=0}^{\infty} x^{km} \sigma_x \Pi_m \right) \left(\sum_{m'=0}^{\infty} x^{k'm'} \sigma_x \Pi_{m'} \right) = \frac{I_{\infty,k}}{I_{\text{sugra}} I_{\text{D5}}} \frac{I_{\infty,k'}}{I_{\text{sugra}} I_{\text{D5}}}. \quad (48)$$

This correctly reproduces the factorized form (46).

4 Conclusions and discussion

In this paper we investigated the brane expansions for the index $I_{N,k}$ of the $\mathcal{N} = 4$ $U(N)$ SYM with the line operators of the rank k anti-symmetric representations. Our analysis was based on the D5-brane realization of the line-operators.

The large N index $I_{\infty,k}$ with finite k is given by the double-sum brane expansion (20), and the finite N index $I_{N,k}$ is given by the quadruple-sum brane expansion (29). By taking an appropriate expansion scheme they reduce to the simple-sum expansion (26) and the double-sum expansion (41), respectively. The objects causing the finite k and finite N corrections are disk D3-branes ending on the D5-brane.

For the quadruple-sum expansion (29) for finite N , the brane indices $\mathcal{F}_{m_x, m_y, m'_x, m'_y}$ are labeled by the four wrapping numbers, and the corresponding brane systems consist of branes extended on the four disks \mathcal{D}_x , \mathcal{D}_y , \mathcal{D}'_x and \mathcal{D}'_y . Due to the complication of the systems, we did not give a general formula to calculate such contributions. One interesting fact we found is the contributions $\mathcal{F}_{m_x, m_y} = \mathcal{F}_{m_x, m_y, 0, 0}$ for $m_x, m_y \geq 1$ are vanishing. This indicates the supersymmetry is spontaneously broken in the theory realized on $\mathcal{D}_x \cup \mathcal{D}_y$. In connection with this, another interesting fact

we found is $\mathcal{F}_{1,1,1,0} \neq 0$. This means that the corresponding brane system $\mathcal{D}_x \cup \mathcal{D}_y \cup \mathcal{D}'_x$ preserves supersymmetry even though it includes the supersymmetry breaking configuration $\mathcal{D}_x \cup \mathcal{D}_y$ as a subsystem. This indicates that whether the supersymmetry is broken depends on the global structure of the brane system. This situation reminds us of the s -rules in Hanany-Witten type brane systems [40]. It would be interesting to investigate the supersymmetry breaking/preserving mechanism on the brane systems with general wrapping numbers.

There are different directions of generalization of the results in this work.

One is generalization to multiple tubular D5-branes. In this work we focused on the anti-symmetric Wilson lines with $\mu = \{1^k\}$. We can consider a more general line labeled by $\mu = \{n^{k_1-k_2}, (n-1)^{k_2-k_3}, \dots, 1^{k_n}\}$. The Young diagram has n columns, and the line corresponds to concentric n D5-branes. The n D5-branes divide each of \mathcal{C}_x and \mathcal{C}_y into $n+1$ zones. If we treat the index as y -series, only branes on \mathcal{C}_x will contribute to the index, and we will have a gauge theory depicted as a linear quiver with $n+1$ nodes. The full expansion will be more complicated and the theory on a brane system is in general a quiver gauge theory with $2(n+1)$ nodes.

It is also important to find precise line operators corresponding to tubular D3-branes, and investigate the finite k and finite N corrections to the corresponding line operator index. Although it is often claimed in the literature that the line operator in a symmetric representation is holographically dual to a tubular D3-brane configuration found in [6], fluctuation modes on the D3-brane analyzed in [22, 35] do not reproduce the index of the symmetric line operators, which is identical in the large N limit to the index (11) of the anti-symmetric line operator with the same rank [18]. This means some modification is necessary for line operators dual to the tubular D3-branes. Regardless of whatever correspond to tubular D3-branes, it would be interesting to analyze the line-operator index for the tubular D3-brane insertion by using brane expansions. We point out that D-strings are strong candidates for causing finite k corrections. Let us introduce a coordinate u in AdS_5 so that the metric of AdS_5 is given by

$$ds_{AdS_5}^2 = L_{AdS}^2 (\cosh^2 u ds_{AdS_2}^2 + du^2 + \sinh^2 u ds_{S^2}^2). \quad (49)$$

For a tubular D3-brane with string charge k , its worldvolume is given by $u = u_{D3}$ [6] where u_{D3} is given by

$$\sinh u_{D3} = \frac{k\sqrt{4\pi g_s N}}{4N}. \quad (50)$$

The mass of a D-string stretched between the two poles of S^2 is

$$T_{D1} \int_{-u_{D3}}^{u_{D3}} L_{AdS} \cosh u du = 2T_{D1} L_{AdS} \sinh u_{D3} = \frac{k}{L_{AdS}}, \quad (51)$$

and the dimension of the corresponding operator is k . This fact strongly suggests that D-strings cause finite k corrections to the line operator index. In addition, D3 giants will give finite N corrections. It would be interesting to study brane expansions including the contribution from D-strings and D3 giants.

It is expected that when the numbers of columns and rows of Young diagrams become of order N the holographic correspondent should be bubbling geometry [8]. Recently, relation between the Schur indices of line operators with such large representations and bubbling geometries were discussed in [21]. It is interesting to investigate the finite N corrections in such background geometries by using brane expansions.

We hope to revisit these issues in future works.

Acknowledgments

The authors thank A. Sei and D. Yokoyama for valuable discussions and comments. The work of Y. I. was supported by JSPS KAKENHI Grant Number JP21K03569.

A Character expansion method

In this appendix we derive some useful formulas using character expansion method [41, 42]. Let us introduce the power sum symmetric polynomials

$$p_\lambda(U) = \prod_{i=1}^{\ell(\lambda)} \text{tr } U^{\lambda_i} = [\text{tr}(U)]_\lambda \quad (52)$$

labeled by partitions λ , and define the index with insertion of line operators associated with these polynomials. We start from the localization formula (2) with s_μ and s_ν replaced by p_λ and $p_{\lambda'}$:

$$\begin{aligned} I_{N,\lambda,\lambda'} &= \langle p_\lambda(U) p_{\lambda'}(U^{-1}) \rangle^{U(N)} \\ &= \int_{U(N)} dU \text{Pexp}(i_{\text{vec}} \text{tr}(U) \text{tr}(U^{-1})) p_\lambda(U) p_{\lambda'}(U^{-1}). \end{aligned} \quad (53)$$

By using the definition of the plethystic exponential we can rewrite this as

$$I_{N,\lambda,\lambda'} = \int_{U(N)} dU \sum_{\lambda''} \frac{1}{z_{\lambda''}} [i_{\text{vec}}]_{\lambda''} p_{\lambda+\lambda''}(U) p_{\lambda'+\lambda''}(U^{-1}), \quad (54)$$

where $\lambda + \lambda'$ is the union of two partitions λ and λ' , which is given by arranging $\{\lambda_1, \lambda_2, \dots, \lambda_{\ell(\lambda)}, \lambda'_1, \lambda'_2, \dots, \lambda'_{\ell(\lambda')}\}$ in descending order. The polynomials $p_\lambda(U)$ are expressed as linear combinations of the Schur polynomials

$s_\mu(U)$ by the formula

$$p_\lambda(U) = \sum_{\mu, \ell(\mu) \leq N} \chi_\mu(\lambda) s_\mu(U) \quad (55)$$

given by Frobenius. $\chi_\mu(\lambda)$ is the character of an irreducible representation of the symmetric group S_n ($n = |\lambda|$) corresponding to the partition μ evaluated at a permutation of cycle type λ . With this formula and the orthonormal relation of the Schur polynomials

$$\int_{U(N)} dU s_\mu(U) s_\nu(U) = \delta_{\mu, \nu}, \quad (56)$$

we obtain the expression

$$I_{N, \lambda, \lambda'} = \sum_{\lambda''} \frac{1}{z_{\lambda''}} [i_{\text{vec}}]_{\lambda''} \sum_{\mu, \ell(\mu) \leq N} \chi_\mu(\lambda + \lambda'') \chi_\mu(\lambda' + \lambda''). \quad (57)$$

In the large N limit, the bound $\ell(\mu) \leq N$ in the μ -sum in (57) disappears, and we can use the orthogonal relation of the characters $\chi_\mu(\lambda)$:

$$\sum_{\mu} \chi_\mu(\lambda) \chi_\mu(\lambda') = z_\lambda \delta_{\lambda, \lambda'}. \quad (58)$$

Then, we obtain

$$I_{\infty, \lambda, \lambda'} = \delta_{\lambda, \lambda'} \sum_{\lambda''} \frac{z_\lambda}{z_{\lambda''}} [i_{\text{vec}}]_{\lambda''} = z_\lambda \delta_{\lambda, \lambda'} [I_{\text{F1}}]_\lambda I_{\text{sugra}}. \quad (59)$$

For $\lambda = \lambda' = \{1\}$ this gives (7).

By using the inverse relation to (55)

$$s_\mu(U) = \sum_{\lambda \vdash |\mu|} \frac{1}{z_\lambda} \chi_\mu(\lambda) p_\lambda(U), \quad (60)$$

We can express $\langle s_\mu(U) s_\nu(U^{-1}) \rangle^{U(N)}$ as a linear combination of $I_{N, \lambda, \lambda'}$. For $N = \infty$, combining (59) and (60) we obtain

$$\langle s_\mu(U) s_\nu(U^{-1}) \rangle^{U(\infty)} = I_{\text{sugra}} \sum_{\lambda \vdash |\mu|} \frac{1}{z_\lambda} [I_{\text{F1}}]_\lambda \chi_\mu(\lambda) \chi_\nu(\lambda). \quad (61)$$

This is non-vanishing only when $|\mu| = |\nu|$.

The line operator indices for the symmetric and anti-symmetric representations can be obtained by substituting $\sigma_{\{1^k\}}(\lambda) = \pm 1$ and $\sigma_{\{k\}}(\lambda) = 1$ for $\lambda \vdash k$ into (61). The common result is shown in (11).

References

- [1] J. M. Maldacena, “The Large N limit of superconformal field theories and supergravity,” *Adv. Theor. Math. Phys.* **2**, 231-252 (1998) doi:10.4310/ATMP.1998.v2.n2.a1 [arXiv:hep-th/9711200 [hep-th]].
- [2] S. S. Gubser, I. R. Klebanov and A. M. Polyakov, “Gauge theory correlators from noncritical string theory,” *Phys. Lett. B* **428**, 105-114 (1998) doi:10.1016/S0370-2693(98)00377-3 [arXiv:hep-th/9802109 [hep-th]].
- [3] E. Witten, “Anti-de Sitter space and holography,” *Adv. Theor. Math. Phys.* **2**, 253-291 (1998) doi:10.4310/ATMP.1998.v2.n2.a2 [arXiv:hep-th/9802150 [hep-th]].
- [4] S. J. Rey and J. T. Yee, “Macroscopic strings as heavy quarks in large N gauge theory and anti-de Sitter supergravity,” *Eur. Phys. J. C* **22**, 379-394 (2001) doi:10.1007/s100520100799 [arXiv:hep-th/9803001 [hep-th]].
- [5] J. M. Maldacena, “Wilson loops in large N field theories,” *Phys. Rev. Lett.* **80**, 4859-4862 (1998) doi:10.1103/PhysRevLett.80.4859 [arXiv:hep-th/9803002 [hep-th]].
- [6] N. Drukker and B. Fiol, “All-genus calculation of Wilson loops using D-branes,” *JHEP* **02**, 010 (2005) doi:10.1088/1126-6708/2005/02/010 [arXiv:hep-th/0501109 [hep-th]].
- [7] S. Yamaguchi, “Wilson loops of anti-symmetric representation and D5-branes,” *JHEP* **05**, 037 (2006) doi:10.1088/1126-6708/2006/05/037 [arXiv:hep-th/0603208 [hep-th]].
- [8] E. D’Hoker, J. Estes and M. Gutperle, “Gravity duals of half-BPS Wilson loops,” *JHEP* **06**, 063 (2007) doi:10.1088/1126-6708/2007/06/063 [arXiv:0705.1004 [hep-th]].
- [9] A. Gadde, L. Rastelli, S. S. Razamat and W. Yan, “Gauge Theories and Macdonald Polynomials,” *Commun. Math. Phys.* **319**, 147-193 (2013) doi:10.1007/s00220-012-1607-8 [arXiv:1110.3740 [hep-th]].
- [10] C. Romelsberger, “Counting chiral primaries in $N = 1$, $d=4$ superconformal field theories,” *Nucl. Phys. B* **747**, 329-353 (2006) doi:10.1016/j.nuclphysb.2006.03.037 [arXiv:hep-th/0510060 [hep-th]].
- [11] J. Kinney, J. M. Maldacena, S. Minwalla and S. Raju, “An Index for 4 dimensional super conformal theories,” *Commun. Math. Phys.* **275**, 209-254 (2007) doi:10.1007/s00220-007-0258-7 [arXiv:hep-th/0510251 [hep-th]].

- [12] J. Bourdier, N. Drukker and J. Felix, “The exact Schur index of $\mathcal{N} = 4$ SYM,” JHEP **11**, 210 (2015) doi:10.1007/JHEP11(2015)210 [arXiv:1507.08659 [hep-th]].
- [13] Y. Pan and W. Peelaers, “Exact Schur index in closed form,” Phys. Rev. D **106**, no.4, 045017 (2022) doi:10.1103/PhysRevD.106.045017 [arXiv:2112.09705 [hep-th]].
- [14] Y. Hatsuda and T. Okazaki, “ $\mathcal{N} = 2^*$ Schur indices,” JHEP **01**, 029 (2023) doi:10.1007/JHEP01(2023)029 [arXiv:2208.01426 [hep-th]].
- [15] T. Dimofte, D. Gaiotto and S. Gukov, “3-Manifolds and 3d Indices,” Adv. Theor. Math. Phys. **17**, no.5, 975-1076 (2013) doi:10.4310/ATMP.2013.v17.n5.a3 [arXiv:1112.5179 [hep-th]].
- [16] D. Gang, E. Koh and K. Lee, “Line Operator Index on $S^1 \times S^3$,” JHEP **05**, 007 (2012) doi:10.1007/JHEP05(2012)007 [arXiv:1201.5539 [hep-th]].
- [17] N. Drukker, “The $\mathcal{N} = 4$ Schur index with Polyakov loops,” JHEP **12**, 012 (2015) doi:10.1007/JHEP12(2015)012 [arXiv:1510.02480 [hep-th]].
- [18] Y. Hatsuda and T. Okazaki, “Exact $\mathcal{N} = 2^*$ Schur line defect correlators,” JHEP **06**, 169 (2023) doi:10.1007/JHEP06(2023)169 [arXiv:2303.14887 [hep-th]].
- [19] Z. Guo, Y. Li, Y. Pan and Y. Wang, “ $\mathcal{N}=2$ Schur index and line operators,” Phys. Rev. D **108**, no.10, 106002 (2023) doi:10.1103/PhysRevD.108.106002 [arXiv:2307.15650 [hep-th]].
- [20] Y. Hatsuda and T. Okazaki, “Large N and large representations of Schur line defect correlators,” JHEP **01**, 096 (2024) doi:10.1007/JHEP01(2024)096 [arXiv:2309.11712 [hep-th]].
- [21] Y. Hatsuda and T. Okazaki, “Excitations of bubbling geometries for line defects,” Phys. Rev. D **109**, no.6, 066013 (2024) doi:10.1103/PhysRevD.109.066013 [arXiv:2311.13740 [hep-th]].
- [22] A. Faraggi and L. A. Pando Zayas, “The Spectrum of Excitations of Holographic Wilson Loops,” JHEP **05**, 018 (2011) doi:10.1007/JHEP05(2011)018 [arXiv:1101.5145 [hep-th]].
- [23] J. McGreevy, L. Susskind and N. Toumbas, “Invasion of the giant gravitons from Anti-de Sitter space,” JHEP **0006**, 008 (2000) doi:10.1088/1126-6708/2000/06/008 [hep-th/0003075].
- [24] A. Mikhailov, “Giant gravitons from holomorphic surfaces,” JHEP **0011**, 027 (2000) doi:10.1088/1126-6708/2000/11/027 [hep-th/0010206].

- [25] R. Arai and Y. Imamura, “Finite N Corrections to the Superconformal Index of S-fold Theories,” PTEP **2019**, no.8, 083B04 (2019) doi:10.1093/ptep/ptz088 [arXiv:1904.09776 [hep-th]].
- [26] R. Arai, S. Fujiwara, Y. Imamura and T. Mori, “Schur index of the $\mathcal{N} = 4$ $U(N)$ supersymmetric Yang-Mills theory via the AdS/CFT correspondence,” Phys. Rev. D **101**, no.8, 086017 (2020) doi:10.1103/PhysRevD.101.086017 [arXiv:2001.11667 [hep-th]].
- [27] Y. Imamura, “Finite- N superconformal index via the AdS/CFT correspondence,” PTEP **2021**, no.12, 123B05 (2021) doi:10.1093/ptep/ptab141 [arXiv:2108.12090 [hep-th]].
- [28] D. Gaiotto and J. H. Lee, “The Giant Graviton Expansion,” [arXiv:2109.02545 [hep-th]].
- [29] S. Murthy, “Unitary matrix models, free fermions, and the giant graviton expansion,” Pure Appl. Math. Quart. **19**, no.1, 299-340 (2023) doi:10.4310/PAMQ.2023.v19.n1.a12 [arXiv:2202.06897 [hep-th]].
- [30] J. H. Lee, “Exact stringy microstates from gauge theories,” JHEP **11**, 137 (2022) doi:10.1007/JHEP11(2022)137 [arXiv:2204.09286 [hep-th]].
- [31] M. Beccaria and A. Cabo-Bizet, “On the brane expansion of the Schur index,” JHEP **08**, 073 (2023) doi:10.1007/JHEP08(2023)073 [arXiv:2305.17730 [hep-th]].
- [32] M. Beccaria and A. Cabo-Bizet, “Giant graviton expansion of Schur index and quasimodular forms,” [arXiv:2403.06509 [hep-th]].
- [33] Y. Imamura, “Giant graviton expansions for line operator index,” [arXiv:2403.11543 [hep-th]].
- [34] M. Beccaria, “Schur line defect correlators and giant graviton expansion,” [arXiv:2403.14553 [hep-th]].
- [35] A. Faraggi, W. Mueck and L. A. Pando Zayas, “One-loop Effective Action of the Holographic Antisymmetric Wilson Loop,” Phys. Rev. D **85**, 106015 (2012) doi:10.1103/PhysRevD.85.106015 [arXiv:1112.5028 [hep-th]].
- [36] Y. Imamura, “Analytic continuation for giant gravitons,” PTEP **2022**, no.10, 103B02 (2022) doi:10.1093/ptep/ptac127 [arXiv:2205.14615 [hep-th]].
- [37] S. Fujiwara, Y. Imamura, T. Mori, S. Murayama and D. Yokoyama, “Simple-Sum Giant Graviton Expansions for Orbifolds and Orientifolds,” PTEP **2024**, no.2, 023B02 (2024) doi:10.1093/ptep/ptae006 [arXiv:2310.03332 [hep-th]].

- [38] S. Kim, “The Complete superconformal index for N=6 Chern-Simons theory,” Nucl. Phys. B **821**, 241-284 (2009) [erratum: Nucl. Phys. B **864**, 884 (2012)] doi:10.1016/j.nuclphysb.2009.06.025 [arXiv:0903.4172 [hep-th]].
- [39] Y. Imamura and S. Yokoyama, “Index for three dimensional superconformal field theories with general R-charge assignments,” JHEP **04**, 007 (2011) doi:10.1007/JHEP04(2011)007 [arXiv:1101.0557 [hep-th]].
- [40] A. Hanany and E. Witten, “Type IIB superstrings, BPS monopoles, and three-dimensional gauge dynamics,” Nucl. Phys. B **492**, 152-190 (1997) doi:10.1016/S0550-3213(97)00157-0 [arXiv:hep-th/9611230 [hep-th]].
- [41] F. A. Dolan, “Counting BPS operators in N=4 SYM,” Nucl. Phys. B **790**, 432-464 (2008) doi:10.1016/j.nuclphysb.2007.07.026 [arXiv:0704.1038 [hep-th]].
- [42] S. Dutta and R. Gopakumar, “Free fermions and thermal AdS/CFT,” JHEP **03**, 011 (2008) doi:10.1088/1126-6708/2008/03/011 [arXiv:0711.0133 [hep-th]].