

New instanton on a warped Kerr spacetime

Reinoud Jan Slagter^{1,*}

Abstract

We find an exact time-dependent instanton solution on a vacuum Kerr-like warped spacetime in conformal dilaton gravity. Remarkably, the metric solution results from a first-order PDE, allowing the connection with self-duality. The antipodal boundary condition on the hypersurface of a Klein bottle $\sim \mathbb{C}^1 \times \mathbb{C}^1$ is applied to describe the Hawking particles. We used the Hopf fibration to get S^2 as the black hole horizon, where the centrix is not in a torus but in the Klein bottle. The twist fits very well with the antipodal identification of the point on the horizon. No "cut and past" is necessary, so the Hawking particles remain pure without instantaneous information transport. A local observer passing the horizon will not notice a central singularity in suitable coordinates. The black hole paradoxes are also revisited in our new black hole model. A connection is made with the geometric quantization of $\mathbb{C}^1 \times \mathbb{C}^1 \sim S^3$, by considering the symplectic 2-form. The model can be easily extended to the non-vacuum situation by including a scalar field. Both the dilaton and the scalar field can be treated as quantum fields when approaching the Planck area.

Keywords: Kerr black hole, conformal invariance, brane-world models, Klein surface, antipodality, Hawking radiation, instantons
PACS: 04.20.-q, 04.50.-h, 04.62.+v, 02.40.-k, 04.70.Dy, 02.40.Dr, 03.65.Ud, 03.65.Ta, 03.65.Vf, 02.30.Jr

1. Introduction

Following the pioneering work of Hawking (1975) on the evaporation of black holes, many scientists began to study the quantum-gravitational properties of black holes. However, there are some profound unsolved problems as one approaches the Planck scale. Due to the extremely high curvature, quantum effects can no longer be denied. It is thought that knowledge of the quantum effects near the horizon of a black hole will eventually lead to a consistent quantum gravity model. The properties of the black hole, i.e. its entropy, mass and emission, need to be revised because there are some serious paradoxes. When fields are quantized on a manifold, one needs a one-to-one map between the entire asymptotic domain of the maximally extended Penrose diagram. We will follow here the approach of 't Hooft (2016), i.e. an antipodal topological boundary condition, already proposed by Schrödinger (1957). This approach differs significantly from other approaches to attack the paradoxes, i.e. the information paradox, the firewall paradox, the complementarity paradox and the unitarity of the S-matrix. The Randall-Sundrum (RS) warped brane world model (Randall, et al. (1999), Randall, et al. (1999a)) is a simplification of the full string model. The extra-dimensional bulk space has a negative cosmological constant. The dilution of gravity by the presence of the bulk effectively weakens gravity on the brane. The result is that the true higher dimensional Planck scale can be much lower than the effective 4D Planck scale, possibly down to the

electroweak scale. It also extends the range of graviton modes that can be felt on the brane, the so-called Kaluza-Klein (KK) modes. The model does not rely on compactification to localize gravity on the brane, but on the curvature of the bulk. The notion of an instanton, or pseudoparticle, was introduced by 't Hooft and represents a localized finite action topological invariant in field theory. It was first developed in Yang-Mills theory, in particular in $SU(n)$ gauge theories. The classical field equations with self-dual field strength then gave rise to a new non-perturbative treatment. The Minkowski spacetime was replaced by a Wick rotation $t \rightarrow i\tau$, where τ is a special parameterization of the favoured tunneling path through the configuration space. Feynman's path-integral approach to quantization is then unambiguous. Solutions are concentrated around a point in four-dimensional Euclidean space, and at one instant there is a "bump" in the field strength, which quickly dies away. There are striking similarities between these instantons and their gravitational counterparts. They are obtained in general relativity by a Wick rotation on Riemannian spacetime. The field equations then lead to first order differential equations. For an overview see Gibbons, et al. (1993). The structure of this letter is as follows. In section two we present the model under consideration. In section three we treat the Hopf fibration needed to describe the embedding of the Klein surface in the 5D model. In section four we make the connection with the instanton and in section five a possible treatment of the Hawking particles.

2. The model

In an earlier study Slagter (2023) we studied the conformally invariant warped spacetime.

$$ds^2 = \omega(t, r, y_5)^{4/3} y_0 \left[-N(t, r)^2 dt^2 + \frac{1}{N(t, r)^2} dr^2 + dz^2 \right]$$

*Former: University of Amsterdam

Email address: info@asfyon.com/info@asfyon.org (Reinoud Jan Slagter)

URL: www.asfyon.com (Reinoud Jan Slagter)

¹Asfyon, Astronomisch Fysisch Onderzoek Nederland

$$+r^2(d\phi + N^\phi(t, r)dt)^2 + dy_5^2 \quad (1)$$

where y_5 is the fifth coordinate, y_0 the bulk dimension and ω is a warp factor, reinterpreted as a dilaton field. We write $^{(5)}g_{\mu\nu} = \omega^{4/(d-2)}\tilde{g}_{\mu\nu}$ and $\tilde{g}_{\mu\nu} = ^{(4)}\tilde{g}_{\mu\nu} + n_\mu n_\nu$. A "unphysical" spacetime is thus split off. Here n_μ is the unit normal to the brane. Again we write $^{(4)}\tilde{g}_{\mu\nu} = \tilde{\omega}^{2(d-2)}\bar{g}_{\mu\nu}$. The Lagrangian under consideration is

$$S = \int d^d x \sqrt{-\bar{g}} \left[-\frac{1}{2}\xi(\Phi\Phi^* + \omega^2)\tilde{R} - \frac{1}{2}\tilde{g}^{\mu\nu} \left(\mathcal{D}_\mu\Phi(\mathcal{D}_\mu\Phi)^* + \partial_\mu\omega\partial_\nu\omega \right) - \frac{1}{4}F_{\alpha\beta}F^{\mu\nu} - V(\Phi, \omega) - \Lambda\kappa^{\frac{4}{d-2}}\xi^{\frac{d}{d-2}}\omega^{\frac{2d}{d-2}} \right] \quad (2)$$

which is invariant under 't Hooft (2015))

$$\tilde{g}_{\mu\nu} \rightarrow \Omega^{\frac{4}{d-2}}\tilde{g}_{\mu\nu}, \quad \omega \rightarrow \omega^{-\frac{d-2}{2}}, \quad \Phi \rightarrow \omega^{-\frac{d-2}{2}}\Phi \quad (3)$$

where \mathcal{D} is the gauge covariant derivative and F the Abelian field strength. Furthermore, $\xi = (d-2)/(4(d-1))$ and we applied the Wick rotations $\omega^2 \rightarrow -6\frac{\omega^2}{\kappa^2}$, with $\kappa = 8\pi G_N$, to ensure that the field ω has the same unitarity and positivity properties as the scalar field. The ω freedom is used by the in-falling observer to describe his experience of the vacuum. Since we are working in a RS warped brane world model, one solves the 5D and the effective 4D equations simultaneously, with the contribution of the projected bulk Weyl tensor appearing in the latter. We are mainly interested in the vacuum case. For the inclusion of the scalar field we refer to Slagter (2022, 2023, 2024). It turns out that one can isolate the field equations for N, ω and N^ϕ . The exact solution of N , without the scalar field, becomes

$$N^2 = \frac{C_2}{r^2(t-b)^4 + C_3} \left[\frac{(r-a)^{k+1}(r(k+1)+a)}{k+2} + C_1 \right] \quad (4)$$

where C_i are constant. The solution is valid for both the 5D and the effective field equations (except for the constants). The constant $k \in \mathbb{N}$. For $k=3$ we get our metric solution, a quintic equation with in general five zero's. The solution for N^ϕ becomes

$$N^\phi = \int \frac{dr}{r^3 \omega^{\frac{n-1}{n-3}}} + F_n(t) \quad (5)$$

where $F_n(t)$ is a arbitrary function determined by a constraint equation. Since the solution for N is separable, we write $N^2 = N_1(r)^2 N_2(t)^2$. The solution for ω is²

$$\omega = \left(\frac{c}{(r-a)(t-b)} \right)^{\frac{1}{2}n-1} \quad (6)$$

with $n=4, 5$ for the effective and 5D situation respectively. The r -dependent part, N_1 of equation (4) is determined by a first-order differential equation

$$rN_1 \frac{\partial N_1}{\partial r} + N_1^2 = \frac{k+1}{2}(r-a)^k \quad (7)$$

²On a FLRW spacetime, the dilaton field is then considered as a warp factor, which can also be expressed exactly in (r, t, y_5) , where the bulk part becomes $e^{\sqrt{\frac{\Lambda_5}{6}}(y_5 - y_0)}$ (Slagter, et.al. (2016)). You will need these solutions when the Hawking particles become hard.

For $C_1 = 0$ the zeros are at $r_H = a_2$ with multiplicity $(k+1)$ and at $r_H = -\frac{a_2}{k+1}$. For $k=3$ and $C_1 = -a^5/5$ there is one real solution, $r_H \approx 1.606a$, while the others are complex. For $k=2$ and $C_1 < a^4/4$ there are 4 zeros, as in the BTZ solution, two of them negative. For $k=1$ and $C_1 < -a^3/3$ there is still a real solution. For $k=0$ we surprisingly get two horizons $r_H = \pm a_2$. The ergo region can be determined by solving $g_{tt} = 0$, with (for $k=3$)

$$\bar{g}_{tt} = \frac{(t-b)^6}{4r^2} \left(6ar^2 \log r - a^2 + 6a^2r - 2r^3 \right)^2 - \frac{C_2(r-a)^4(4r+a)}{5r^2((t-b)^4 + C_3)} \quad (8)$$

One can make the spacetime free of an ergo region by suitable values of the parameters and $F_k(t)$. The effective 4d spacetime in suitable coordinates becomes

$$ds_4^2 = \omega^{4/3}\tilde{\omega}^2 \left[\frac{N_1^2}{N_2^2}(-dt^{*2} + dr^{*2}) + dz^2 + r^2 d\phi^{*2} \right] \quad (9)$$

with

$$r^* = \frac{1}{4} \sum_{r_i^H} \frac{r_i^H \log(r - r_i^H)}{(r_i^H + a_2)^3}, \quad t^* = \frac{1}{4C_2} \sum_{t_i^H} \frac{\log(t - t_i^H)}{(t_i^H + a_3)^3}. \quad (10)$$

The sum is taken over the roots of $(10a_2^3r^2 + 20a_2^2r^3 + 15a_2r^4 + 4r^5 + C_1)$ and $C_2(t+a_3)^4 + C_3$, i. e. r_i^H and t_i^H . We also defined the azimuthal angle coordinate $d\phi^* = d\phi + (N^\phi/N_2^2)dt^*$, which can be used when an incoming null geodesic falls on the horizon, i.e. in a coordinate system that rotates about the z -axis relative to the Boyer-Lindquist coordinates. In Kruskal-Szekeres lightcone coordinates (U, V) we have

$$ds^2 = \omega^{4/3}\tilde{\omega}^2 \left[\frac{N_1^2}{N_2^2} \frac{dUdV}{\varepsilon^2 UV} + dz^2 + r^2 d\phi^{*2} \right] \quad (11)$$

with ε a constant related to the surface gravity at the horizon. If we omit the contribution of the projected Weyl




Figure 1: Penrose diagram. The antipodal points $P(X)$ and $P(\bar{X})$ are identified. Particles entering the black hole will create waves that approach the horizon from the outside, and those that pass the horizon will emerge from the "other side" of the black hole. Note that $t^* = \log(U/V) \sim \log(t - t_H)$. In this approach, I and II are CPT invariant. This means that time runs backwards in II.

tensor in the effective 4D Einstein equations, we obtain

the Bañados-Teitelboim-Zanelli 3D solution (Bañados, et al. (1992), Compère (2018)). This is equivalent to the substitution $k = 2$ in Eq. (4). We also found a remarkable connection with the icosahedral Möbius group A_5 when inscribed in S^2 . The isometry group is of order 5 and has three orbits which are invariant under the antipodal map. Our general polynomial of N^2 is a quintic, which cannot be a coincident (Slagter (2023)). If we include a complex scalar-gauge (Φ, A_μ) , we find from the superfluous dilaton equation, that the scalar-dilaton potential $V(\omega, \Phi)$ satisfies

$$V(\omega, \Phi) = \beta_1 \Phi^{\beta_2} \omega^{\frac{2}{3} - \frac{\eta \beta_2}{y_0}} \quad (12)$$

So we get a quartic conformal invariant matter coupling $\sim \Phi^2 \omega^2$ for $|y_0| = 3/2\eta$. Where η is the vacuum expectation and we have used a constant gauge field.

3. Hopf fibrating $\mathbb{C}^1 \otimes \mathbb{C}^1$ and the Klein bottle

We have used the embedding of the Klein bottle in the warped 5D spacetime in our model. To get back to our black hole in the effective 4D spacetime, we can use the Hopf fibration of $S^3 \sim \mathbb{C}^1 \otimes \mathbb{C}^1$ and apply the antipodal map ('t Hooft (2016)) to get the S^2 of our black hole horizon. The Hawking particle created near the horizon at $P(U, V, z, \varphi, y_5)$, leaves the horizon at the antipode $\tilde{P}(-U, -V, -z, \varphi + \pi, -y_5)$ after a finite proper time. In our situation we define $W = x + iy, Z = z + iy_5^3$.




Figure 2: *Fibering the S^3 .*

We write the 3-sphere $S^3 = \{x, y, z, y_5 \in \mathbb{R}^4 | x^2 + y^2 + z^2 + y_5^2 = r^2 + R^2\} = \{(Z, W) \in \mathbb{C}^2 | Z\bar{Z} + W\bar{W} = r^2 + R^2\}$. See figure (2)

³or W in polar coordinates $re^{i\varphi}$.

and the diagram of Eq. (13).

$$\begin{array}{ccc} (Z, W) \in \mathbb{C}^2 / \{0\} & \longleftarrow & S^3 \\ \downarrow H & & \downarrow \mathbb{C}P^1 \quad \downarrow P \\ u \equiv \frac{Z}{W} \in \mathbb{C} \cup \{\infty\} & = & S^2 \end{array} \quad (13)$$

There are two basic compact self-dual manifolds, the round metric on S^4 and the Fubini-Study metric on $\mathbb{C}P^2$. In our situation we are mainly interested in the quantization on the latter. It is possible to construct a complex two-dimensional Hilbert space \mathcal{H} . If we write on \mathbb{C}^2 , $|\mathcal{U}\rangle = (Z, W)$ with density matrix $\rho = |\mathcal{U}\rangle\langle\mathcal{U}|$, we can construct a quantization of $\mathbb{C}P^2$ (Urbanke (1990); Slagter (2023)). On the non-orientable Klein surface, meromorphic functions remain constant and our quintic is meromorphic. It is also homeomorphic to the connected sum of two projected planes. It is also possible to construct geodesics




Figure 3: *Geodesics on the torus and the Klein bottle. Note the difference: because of the \mathbb{Z}_2 symmetry of our warped spacetime, we need a non-closed geodesic for the torus.*

and a minimum surface needed to follow the particles. You can use the torus, on which the orbits are easy to find. Note that the orbit is still in 4D space. The geodesics on the Klein surface will never intersect each other because there is no "inside".

4. The instanton

It turns out that self-duality in the Yang-Mills model, leads to first-order differential equations. They played a fundamental role in non-abelian gauge theories. See, for example, the textbook of Felsager (1998). It is now known that there are




Figure 4: *Plot of the instanton for $\tilde{g}_{\mu\nu}$ for $a = \pm 1$ respectively. We have chosen $C_1 = 0$.*

Euclidean self-dual instanton solutions in gravity theory. The self-duality is then expressed in terms of the Riemann curvature tensor. They are thought to play a role in understanding

quantum gravitational effects near the horizon of BH's, such as the evaporation process and the formation of Planck-sized BH's⁴. Well-known examples are the Eguchi-Hanson solution (Eguchi, et al. (1978)) and the Fubini study, which can be described on a $(\mathbb{C}^1 \times \mathbb{C}^1)$ Kähler manifold (Joyce (1995)).

If we switch to Riemannian space with $t \rightarrow i\tau$, we get the same solution as in Eq.(3) - (5). The line element can be written in the stationary case and for $C_1 = 0, k = 3$ as

$$ds^2 = \omega^{4/3} \bar{\omega}^2 \left[\frac{(a+4r)(r-a)^4}{5r^2} d\tau^2 + dr^{*2} + dz^2 + r^2 d\varphi^{*2} + dy_5^2 \right] \quad (14)$$

with




Figure 5: Left: Plot of r^* against r and the inverse for the case $a = 1$, with range $-\frac{a}{4} < r < a$. If $r \rightarrow a$, then $r^* \rightarrow \pm\infty$. Right: Surface of revolution around the r axis. A possible geodesic is drawn. The surface is not orientable, but there is a cusp singularity at $r = -1/4a$.

$$r^* = \frac{1}{5\sqrt{a}} \left[6a \operatorname{arctanh} \left(\sqrt{\frac{a+4r}{5a}} \right) + \frac{\sqrt{5a(a+4r)}}{r-a} \right] \quad (15)$$

with $-\frac{a}{4} < r < a$. It describes the interior of the black hole. In figure (6) we have plotted r^* , valid only inside the horizon $r_h = 1$, for $a = 1$. We have also plotted the surface of revolution (y_5 suppressed). Note the difference from the wormhole




Figure 6: Left: Plot of $z(r)$ of the Euclidean 3-space for our model. Right: The surface of revolution over φ . A particle arriving at the horizon $r = a$ "jumps" to the antipode $\varphi \rightarrow \varphi + \pi$.

case⁵. The region $r < \frac{a}{4}$ is inaccessible. It would be challenging to consider large values of k . We will then get a horizon and the singularity for $\rightarrow 0$ will never be reached. It became quite clear that Euclidean self-dual solutions in Einstein's theory play

a role in quantum gravity. These solutions have vanishing classical action and non-trivial topological invariants. Moreover, there is a similarity with the YM Euclidean self-dual solutions Eguchi, et al. (1978). Despite their Euclidean origin, instantons, like solitons, play a fundamental role in quantum theories. Their importance arises, for example, in *tunneling* effects using the WKB approximation. Instanton-like solutions are expected to be all self-dual, localized in Euclidean spacetimes and free of singularities. Asymptotically locally Euclidean self-dual metrics were also important for the path integral calculations. They give a dominant contribution to the path integral that is as important as that of the flat metric itself. They also have time-parity inversion. However, gravitational instantons do not contribute to the zero-frequency modes of a spin-1/2 Dirac particle, while they produce zero-frequency solutions of spin-3/2 particles. It is conjectured (Gibbons, et al. (1993), Atiyah, et al. (1978)) that there is a deep connection between the gravitational instanton and the alternating discrete symmetry group \mathcal{A}_5 which serves as a discrete subgroup of $SO(4)$ of its boundary. A gravitational instanton is a four-dimensional regular spacetime M , self-dual Ricci curvature and asymptotically local Euclidean, whose boundary has a $\mathcal{A}_k, D_k, T, O, I$ symmetry⁶. In fact, every finite subgroup of $S^3 \sim SU(2)$, is either cyclic or conjugate to these binary subgroups. For our 5D warped spacetime, our boundary is the Klein surface, embedded in $\mathbb{R}^4 \sim \mathbb{C}^1 \times \mathbb{C}^1$ and the symmetry the double cover group I of the icosahedron. An interesting study has also been done by Gibbons, et al. (1993). They studied, on $\mathbb{C}P^2$, a gravitational instanton surrounded by an event horizon and compared the solution with the $SU(2)$ Yang-Mills instanton. This study is based on the Riemannian solution of the Einstein equations with a cosmological constant on $\mathbb{C}P^2$ by applying the Hopf-fiberings. The Weyl tensor is then anti-self dual and they conclude that there is no Lorentzian counterpart connection, i. e. it cannot be related to a real 4-dimensional manifold. Our model does.

5. Treatment of the Hawking particles

We have taken a slightly different approach to the antipodal map used by 't Hooft (2015). To revise the information, firewall and complementarity paradoxes, 't Hooft uses the antipodal map on S^2 , where the non-orientability is provided by the Möbius strip. His method restores quantum purity and CPT invariance. The particles remain entangled. The incoming particles return at the antipode and the particles remain entangled. There is no inside of the black hole, i.e. region II is not "really there" for the observer inside by the cut-and-paste procedure. In our model we use the Klein surface \mathbb{K} . Our conjecture is that our solution can be seen as a dynamical solution embedded in our $(4+1)$ dimensional spacetime. More precisely, if \mathcal{M} is the sub-manifold with metric g , i.e. the metric on our effective 4D spacetime, then \mathcal{M} is diffeomorphic to the hyperbolic 5D spacetime: $\chi : \mathbb{R}^5 \rightarrow \mathcal{N}$. The Riemannian manifold (\mathcal{M}, g)

⁴Recently, primordial BHs have been observed, which could not be formed by contracting matter.

⁵In our case we have no explicit expression for the surface of revolution.

⁶These are the dihedral, Möbius, tetrahedral, octahedral and icosahedral groups respectively.




Figure 7: Treatment of the Hawking particles on the Klein bottle.

must be conformally flat, which it is. Now physicists are interested in the topology of moduli spaces of self-dual connections on vector bundles over Riemannian manifolds. One reason was that on these spaces the instanton approximation to the Green functions of Euclidean quantum gravity, Yang-Mills theory, can be expressed in terms of integrals over moduli spaces. One then needs the metric and volume form of the moduli spaces. We suspect that we can consider \mathbb{K} as a 4-sphere in our hyperbolic pseudo-Riemannian spacetime. A note must be made about the \mathbb{Z}_2 symmetry in the original description of the antipodal mapping ('t Hooft (2021)). At the boundary of regions I and II in the Penrose diagram, the antipodes were glued together on a 3-sphere, and the transverse (θ, φ) part is a projected 2-sphere. In our model it is replaced by the projected 3-sphere (z, φ, y_5) using the \mathbb{Z}_2 symmetry of the bulk space (U, V, z, φ, y_5) . We "blow up" the 4-manifold to 5D to handle the singularities in the curvature and to apply the antipodal map. One can mathematically formulate the topology of a 4-manifold using self-dual connections over de Riemannian S^4 (Freed (1984)). It depends only on the conformal class of the Riemannian metric. This self-dual connection can be interpreted by the conformal map: $\mathbb{R}^4 \rightarrow S^4/\{0\}$ as a self-dual connection or "instanton". We interpret the Riemannian 5D warped spacetime, an open 5-ball, as an instanton on \mathbb{R}^4 . In the pseudo-Riemannian spacetime, the boundary is the non-orientable Klein surface, which we used for the antipodality to preserve the pure states of the Hawking particles. It is known that a static solution in m space dimensions is completely equivalent to an instanton in m space-time dimensions. An outside observer will register the Hawking radiation as thermal, i.e. in a mixed state, while a local observer will doubt the vacuum state. Furthermore, the evolution of the wave function of the incoming particle must be unitary, i.e. it must satisfy the Schrödinger equation $|\chi(t_1)\rangle = U(t_1, t_2)|\chi(t_2)\rangle$ and be bijective. It is thought that information about the quantum state is preserved during black hole evaporation. The loss of information is incompatible with unitarity. Thus, 't Hooft's method resolves the controversy between the pure quantum state of the incoming particles and the mixed state property of the Hawking radiation. The information is retrieved by the outgoing particles. They carry the information of the incoming particles. In quantum mechanical terms, the incoming posi-

tion of the wave packet is imprinted on the outgoing momentum. The positions are then highly de-localised. This is how long-wavelength Hawking particles are produced from short-wavelength waves. In our model, the Hawking particles don't jump infinitely fast to the antipode, but stay on the Klein surface for a while. Moreover, the complementarity in the different notion of the dilaton ω used by the inside and outside observer is formulated by the invariance of Eq.(3). Then we have the problem of handing over the "hard" gravitons. 't Hooft uses the Shapiro delay method to describe the gravitational interaction near the horizon. Hard particles become "soft" by taking into account the equations in the transverse spherical harmonics $\mathbf{Y}_{l,m}(\theta, \varphi)$. The dynamics is factorized in (l, m) . The dilaton is written as

$$\omega = \mathbf{J}(r, t)e^{im\varphi}\mathbf{Y}_m(\varphi) \quad (16)$$

where \mathbf{J} is the elliptic Jacobi function and \mathbf{Y}_m is the cylindrical harmonic. The firewall decays due to the identification $(\theta, \varphi) \rightarrow (\pi - \theta, \varphi + \pi)$. In our model, the shape of the probability function makes the gravitons hard in the bulk (Randall, et al. (1999a)) and soft on the brane. The result is that the presence of the bulk causes the KK contribution of the Weyl tensor on the brane, leading to our new solution. The spherical harmonics are replaced by the cylindrical $\mathbf{Y}_m(\varphi)$. The second quantum number is provided by the "winding number", because the dilaton behaves as an ordinary quantum field with axial symmetry, just like the scalar field, when it is included. Furthermore, we obtained PDE's in (r, t) with exact solution. There will be a time delay between the incoming and outgoing waves or particles depending on y_0 , which can be expressed in the parameters of our model. Thus, experimental data could provide an estimate of the bulk dimension. Furthermore, 't Hooft's method requires high wave numbers when approaching the Planck region, and the cut-off is then determined by the number of microstates. In our model we don't need a cut-off. We have the elapsed time on the Klein surface in the bulk, which gives the degree of "hardness". Furthermore, in our model the wavelength increases with time. This would imply that the particles gain momentum and the quanta become blue-shifted. However, our process takes place in the bulk, so it is not visible. One can also apply Schrödinger's original treatment of the Klein-Gordon equation on a hypersurface (Schrödinger (1957)). He finds conditions for the values of the quantum numbers (l, m) ⁷ when the lines of constant phase on the hypersurface will close. In our case we can have the situation where the phase lines spiral around the Klein surface many times without the possibility of intersecting. It should be noted that the antipodal map is obligatory in order to make the mapping of the coordinates in regions I and II one-to-one, and yet they are distinct because they are spatially separated. There is no summation over "invisible" quantum states to form the density matrix, we still have pure states. If you sum over the other states, which are far away in the antipodal region and unobservable to the local observer, you would conclude that these particles have the Hawking temperature. But the state is not thermal at all. The Hawking particles on one hemisphere

⁷Schrödinger applies the case of an expanding universe.

are entangled with the antipode. To the outside observer, the Hartle-Hawking wave function is no longer a thermally mixed state, but a single pure state. In general, the antipodes are completely entangled. From the outside, the black hole is not in a stationary Hartle-Hawking quantum state. In other words, there is no complete heat bath. Finally, we can study the surface gravity of our model. In the φ^* coordinate it becomes

$$\kappa = \frac{2}{5} \left(\frac{10C_2(t-b)^3 + 6r^2 + 10a^2 - 15ar - \frac{5C+a^5}{r^3}}{C_2(t-b)^4 + C_3} \right) \quad (17)$$

Our solution becomes an extremal black hole, i.e. the surface gravity is zero, for

$$t = b + \frac{1}{a} \sqrt[3]{\frac{C}{2C_2}} \quad (18)$$

In general, the black hole never becomes extremal in (t, r, φ) coordinates.

6. Conclusions

We formulate our results in the following proposition:

1. The vacuum solution found on the conformal invariant 5D warped spacetime, where the metric is determined by a quintic polynomial, can be described by a Laplace-type equation on a Klein surface. The zeros of this quintic polynomial depend on the constraints on the parameters of the model under consideration. In a special case, the zeros can be expressed analytically as $\sim (r - r_H)^{k+1} [r(k+1) + r_H] + C(k+2)$ where k is an integer and C a constant. They are related to the eigenvalues of the Laplacian on the Klein surface. The minimal area is determined by an elliptic integral of the second kind.
2. The antipodal mapping of regions I and II in the Penrose diagram is accompanied by the continuous transition in the extra bulk dimension by means of the non-orientable Klein surface embedded in $\mathbb{R}^4 \sim \mathbb{C}^1 \times \mathbb{C}^1$ by means of a Hopf fibration. Measuring the elapsed time that the Hawking particles spend on the Klein surface could give an estimate of the dimension of the bulk.
3. In the conformal dilaton-scalar field model on a warped spacetime, the fields equations for the dilaton ω , which is part of the gravitational metric, and the scalar field obey comparable Klein-Gordon-type PDE's. The interaction of these quantum fields on the "un-physical" conformally flat spacetime also describes the time evolution of the horizon.
4. The information problem could be solved by the antipodal boundary condition, i.e. by removing the inside of the black hole. The result was that the black hole is not thermal. The reason is that there is no "hidden sector" anymore to produce the thermodynamically mixed state of the Hawking radiation. All states remain pure for the outside observer. In our model, the antipodal map on the brane is accompanied by the \mathbb{Z}_2 symmetry in the bulk in the Randall-Sundrum model. No "cut-and-past" is required. This means that there is no infinitely fast transport to the other side of the hemisphere. The Hawking particles remain on the Klein surface for a while, as seen by the outside

observer. A local observer will not notice the $r = 0$ singularity in a suitably chosen coordinate system for the S^3 , which is conformal flat. The vacuum solution described an instanton in Riemannian space. In suitable coordinates, the inside of the black hole shows no central singularity.

5. The geometric quantization of our Klein surface can be constructed using a two-level Hilbert space on the complex hypersurface $\mathbb{C}^1 \times \mathbb{C}^1$.

References

- M. F. Atiyah, N. J. Hitchin and I. M. Singer, Proc. R. Soc. Lond. A 362 (1978) 425.
M. Bañados, M. C. Teitelboim and T. Zanelli, Phys. Rev. Lett. 69, (1992) 1849.
G. Compère, Advanced Lectures on General Relativity, Lecture notes in Physics, Springer, 2019.
T. Eguchi and A. J. Hanson, Ann. of Phys. 120 (1979) 82.
B. Felsager, Geometry, Particles and Fields, Springer, New York, 1998.
D. S. Freed and K. K. Uhlenbeck, Instantons and four-manifolds, Springer-Verlag: New York, 1984.
G. W. Gibbons and S. W. Hawking, Euclidean Quantum Gravity, World Scientific, Singapore, 1993.
S. Hawking, Comm. Math. Phys. 43 (1975) 199.
G. 't Hooft, Found. Phys., 46 (2016) 1185.
G. 't Hooft, Int. J. of Mod. Phys. D, 24 (2015) 1543001.
G. 't Hooft, Universe, 7 (2021) 298.
D. D. Joyce, Duke Math. J. 77 (1995) 519.
L. Randall, R. Sundrum, Phys. Rev Lett 83 (1999) 3370.
L. Randall, R. Sundrum, Phys. Rev Lett 83 (1999) 4690.
E. Schrödinger, Expanding Universe, Cambridge University. Press, Cambridge U.K. 1957.
R. J. Slagter and S. Pan, Found. Phys. 46 (2016) 1075.
R. J. Slagter, Int. J. Mod. Phys. A, 37 (2022) 2250176.
R. J. Slagter, Universe, 9 (2023) 383.
R. J. Slagter, Black holes and Cosmic Strings Revisited, (2024). To appear.
H. Urbantke, Am. J. Phys. 59 (1990) 503.