Concentration properties of fractional posterior in 1-bit matrix completion

The Tien Mai

Department of Mathematical Sciences, Norwegian University of Science and Technology, Trondheim, 7034, Norway.

Corresponding author(s). E-mail(s): the.t.mai@ntnu.no;

Abstract

The problem of estimating a matrix based on a set of its observed entries is commonly referred to as the matrix completion problem. In this work, we specifically address the scenario of binary observations, often termed as 1-bit matrix completion. While numerous studies have explored Bayesian and frequentist methods for real-value matrix completion, there has been a lack of theoretical exploration regarding Bayesian approaches in 1-bit matrix completion. We tackle this gap by considering a general, non-uniform sampling scheme and providing theoretical assurances on the efficacy of the fractional posterior. Our contributions include obtaining concentration results for the fractional posterior and demonstrating its effectiveness in recovering the underlying parameter matrix. We accomplish this using two distinct types of prior distributions: low-rank factorization priors and a spectral scaled Student prior, with the latter requiring fewer assumptions. Importantly, our results exhibit an adaptive nature by not mandating prior knowledge of the rank of the parameter matrix. Our findings are comparable to those found in the frequentist literature, yet demand fewer restrictive assumptions.

Keywords: matrix completion, binary response, logistic regression, fractional posterior, concentration rate, low-rank priors

1 Introduction

Matrix completion has been extensively explored in the fields of machine learning and statistics, attracting considerable attention in recent years due to its relevance to various contemporary applications such as recommendation systems (Bobadilla et al.,

2013; Koren et al., 2009), including the notable Netflix challenge (Bennett and Lanning, 2007), image processing (Ji et al., 2010; Han et al., 2014), genotype imputation (Chi et al., 2013; Jiang et al., 2016), and quantum statistics (Gross, 2011). Although completing a matrix in general is often deemed infeasible, seminal works by Candès and Tao (2010); Candes and Plan (2010); Candès and Recht (2009) have demonstrated its potential feasibility under the assumption of a low-rank structure. This assumption aligns naturally with practical scenarios, particularly in recommendation systems, where it implies the presence of a limited number of latent features that capture user preferences. Various theoretical and computational approaches to matrix completion have been proposed and investigated, as in Tsybakov et al. (2011); Lim and Teh (2007); Salakhutdinov and Mnih (2008); Recht and Ré (2013); Chatterjee (2015); Mai and Alquier (2015); Alquier and Ridgway (2020); Chen et al. (2019).

The previously mentioned studies primarily focused on matrices with realnumbered elements. However, in many practical situations, the observed elements are often binary, taking values from the set $\{-1, 1\}$. This type of data is prevalent in diverse contexts, such as voting or rating data, where responses typically involve binary distinctions like "yes/no", "like/dislike", or "true/false". Tackling the challenge of reconstructing a matrix from incomplete binary observations, known as 1-bit matrix completion, was initially investigated in Davenport et al. (2014). Subsequent studies in this field have been conducted by various researchers (Cai and Zhou, 2013; Klopp et al., 2015; Hsieh et al., 2015; Cottet and Alquier, 2018; Herbster et al., 2016; Alquier et al., 2019), most of whom have taken a frequentist approach. However, there remains a gap in the literature concerning the theoretical assessment of Bayesian methodologies in this domain.

In this study, we aim to address this gap by focusing on a generalized Bayesian approach, where we utilize a fractional power of the likelihood. This leads to what is commonly referred to as fractional posteriors or tempered posteriors, as elucidated in Bhattacharya et al. (2019); Alquier and Ridgway (2020). It is noteworthy to emphasize that generalized Bayesian methods, where the likelihood is substituted by its fractional power or by a concept of risk, has garnered increased attention in recent years, as demonstrated by various works such as Hammer et al. (2023); Jewson and Rossell (2022); Yonekura and Sugasawa (2023); Mai and Alquier (2017); Matsubara et al. (2022); Medina et al. (2022); Grünwald and Van Ommen (2017); Bissiri et al. (2016); Yang et al. (2020); Lyddon et al. (2019); Syring and Martin (2019); Knoblauch et al. (2022); Mai (2023b); Hong and Martin (2020). Additionally, we tackle this problem by considering a general, non-uniform sampling scheme. While a general sampling scheme for 1-bit matrix completion has been examined in Klopp et al. (2015), our requirements are less stringent than theirs.

Initially, we present results concerning the employment of a widely used low-rank factorized prior distribution. Such priors have demonstrated practical efficacy, as evidenced in works such as Cottet and Alquier (2018); Lim and Teh (2007); Salakhutdinov and Mnih (2008). However, due to the typically large dimensionalities of matrix completion problems, employing low-rank factorized priors necessitates intricate Markov Chain Monte Carlo (MCMC) adaptations, which can be computationally expensive and lack scalability. Consequently, in practical applications, variational inference is

often favored for such priors, as discussed in works like Cottet and Alquier (2018); Lim and Teh (2007); Babacan et al. (2012). We present novel results regarding the consistency and concentration properties of the fractional posterior. Specifically, we establish concentration results for the recovering distribution within the α -Rényi divergence framework. Consequently, as particular instances, we derive concentration outcomes relative to metrics such as the Hellinger metric. Furthermore, we broaden our investigation to establish concentration rates for parameter estimation utilizing specific distance measures such as the Frobenius norm. Our findings are comparable to those in the frequentist literature as documented in Davenport et al. (2014); Cai and Zhou (2013), and Klopp et al. (2015).

In addition to the aforementioned type of prior, we also undertake theoretical examination utilizing a spectral scaled Student prior. This prior, introduced by Dalalyan (2020), shares conceptual similarities with a hierarchical prior discussed in Yang et al. (2018). The spectral scaled Student prior enables posterior sampling through Langevin Monte Carlo, a gradient-based sampling technique that has recently garnered considerable attention in various high-dimensional problems, as observed in Durmus and Moulines (2017, 2019); Dalalyan (2017); Dalalyan and Karagulyan (2019). We demonstrate that by employing this prior, it is possible to achieve concentration results for the fractional posterior without necessitating a boundedness assumption, as is typically required for low-rank factorization priors.

The remainder of the paper is structured as follows. In Section 2, we introduce the notations essential for our work and discuss the problem of 1-bit matrix completion. We also present the fractional posterior along with the low-rank factorization prior in this section. Section 3 presents the results pertaining to the low-rank factorization prior, while Section 4 is dedicated to the outcomes obtained using the spectral scaled Student prior. All technical proofs are consolidated in Section 5.

2 Notations and method

2.1 Notations

For any integer m, let $[m] = \{1, \ldots, m\}$. Given integers m and k, and a matrix $M \in \mathbb{R}^{m \times k}$, we write $||M||_{\infty} := \max_{(i,j) \in [m] \times [k]} |M_{ij}|$. For a matrix M, its spectral norm is denoted by ||M||, its Fobenius norm is denoted by $||M||_F = \sqrt{\sum_{ij} M_{ij}^2}$, and its nuclear norm is denoted by $||M||_*$ (the sum of the singular values).

Let $\alpha \in (0, 1)$, the α -Rényi divergence between two probability distributions Q and R is defined by

$$D_{\alpha}(Q,R) = \frac{1}{\alpha - 1} \log \int \left(\frac{\mathrm{d}Q}{\mathrm{d}\mu}\right)^{\alpha} \left(\frac{\mathrm{d}R}{\mathrm{d}\mu}\right)^{1 - \alpha} \mathrm{d}\mu,$$

where μ is any measure such that $Q \ll \mu$ and $R \ll \mu$. The Kullback-Leibler (KL) divergence is defined by

$$\mathcal{K}(Q,R) = \int \log\left(\frac{\mathrm{d}Q}{\mathrm{d}R}\right) \mathrm{d}Q$$
 if $Q \ll R, +\infty$ otherwise

2.2 1-bit matrix completion

We assume that the observed data $(w_1, Y_1), \ldots, (w_n, Y_n)$ are i.i.d. (independent and identically distributed) random variables drawn from a joint distribution characterized by a matrix $M^* \in \mathbb{R}^{d_1 \times d_2}$, denoted by P_{M^*} . Additionally, we assume that $(\omega_s)_{s=1}^n \in ([d_1] \times [d_2])^n$ are i.i.d. and denoted by Π its marginal distribution. These indices correspond to observations, denoted by $(Y_s)_{s=1}^n \in (-1, +1)^n$, distributed accordingly:

$$Y|\omega = \begin{cases} 1 & \text{with probability } f(M_{\omega}^*), \\ -1 & \text{with probability } 1 - f(M_{\omega}^*), \end{cases}$$
(1)

where f is the logistic link function $f(x) = \frac{\exp(x)}{1+\exp(x)}$. This model is similar to Klopp et al. (2015). In this model, we have the likelihood of the observations as $L_n(M) := \prod_{s=1}^n f(M_{\omega_i})^{1_{[Y_i=1]}} (1-f(M_{\omega_i}))^{1_{[Y_i=-1]}}$.

In this study, we operate under the assumption that the rank of M^* , denoted as r, is substantially smaller than its dimensions, specifically $r \ll \min(d_1, d_2)$. This is a prevalent assumption in 1-bit matrix completion research, (Davenport et al., 2014; Cai and Zhou, 2013; Cottet and Alquier, 2018; Klopp et al., 2015; Alquier et al., 2019).

We concentrate on the fractional posterior for $\alpha \in (0,1)$, as discussed in Bhattacharya et al. (2019); Alquier and Ridgway (2020), which is formulated as follows:

$$\pi_{n,\alpha}(M) \propto L_n^{\alpha}(M)\pi(M).$$

In the case $\alpha = 1$, one recovers the traditional posterior distribution.

We define the mean estimator as

$$\hat{M} := \int M \pi_{n,\alpha}(\mathrm{d}M). \tag{2}$$

Low-rank factorization prior

In Bayesian matrix completion methodologies (Babacan et al., 2012; Salakhutdinov and Mnih, 2008; Lim and Teh, 2007), a prevalent concept involves decomposing a matrix into two matrices in order to establish a prior distribution on low-rank matrices. It is commonly acknowledged that any matrix with a rank of r can be decomposed as follows: $M = LR^{\top}, L \in \mathbb{R}^{d_1 \times r}, R \in \mathbb{R}^{d_2 \times r}$. This approach is grounded in the assumption that the underlying matrix M^* exhibits a low rank, or is at least well approximated by a low-rank matrix.

However, in practical scenarios, the rank of the matrix is typically unknown. Thus, for a fixed $K \in \{1, \ldots, \min(d_1, d_2)\}$, one can express $M = LR^{\top}$ with $L \in \mathbb{R}^{d_1 \times K}$, $R \in \mathbb{R}^{d_2 \times K}$. Subsequently, potential ranks $r \in [K]$ are adjusted by diminishing certain columns of L and R to zero. To address this, the reference Cottet and Alquier (2018) considers the following hierarchical model:

$$\gamma_k \overset{iid}{\sim} \pi^{\gamma}, \forall k \in [K],$$
$$L_{i,\cdot}, R_{j,\cdot} | \gamma \overset{iid}{\sim} \mathcal{N}(0, \operatorname{diag}(\gamma)), \forall (i, j) \in [d_1] \times [d_2].$$

The prior distribution on the variances π^{γ} plays a crucial role in controlling the shrinkage of the columns of L and R towards zero. It is common for π^{γ} to follow an inverse-Gamma distribution (Salakhutdinov and Mnih, 2008). This hierarchical prior distribution bears resemblance to the Bayesian Lasso proposed in Park and Casella (2008), and particularly resembles the Bayesian Group Lasso (Kyung et al., 2010), where the variance term follows a Gamma distribution.

The paper Cottet and Alquier (2018) shows that the Gamma distribution is also a possible alternative in matrix completion, both for theoretical results and practical considerations. Thus all the results in this paper are stated under the assumption that π^{γ} is either the Gamma or the inverse-Gamma distribution: $\pi^{\gamma} = \Gamma(a, b)$, or $\pi^{\gamma} = \Gamma^{-1}(a, b)$. In this study, we regard *a* as a fixed constant, while *b* is seen as a small parameter requiring adjustment.

3 Main results

For $r \geq 1$ and B > 0, we define $\mathcal{M}(r, B)$ as the set of pairs of matrices (\bar{U}, \bar{V}) , with dimensions $d_1 \times K$ and $d_2 \times K$ respectively, satisfying that: $\|\bar{U}\|_{\infty} \leq B$, $\|\bar{V}\|_{\infty} \leq B$ and $\bar{U}_{i,\ell} = 0$ for i > r and $\bar{V}_{j,\ell} = 0$ for j > r. Similar to Cottet and Alquier (2018); Alquier and Ridgway (2020), we make the following assumption on the true parameter matrix.

Assumption 3.1. We assume that $M^* = \overline{U}\overline{V}^t$ for $(\overline{U}, \overline{V}) \in \mathcal{M}(r, B)$.

The following theorem presents the first consistency result for the fractional posterior in 1-bit matrix completion with Gaussian priors which frequently employed in practical applications.

Theorem 1. Assume that Assumption 3.1 holds. Then, there is a small enough b > 0 such that

$$\mathbb{E}\left[\int D_{\alpha}(P_M, P_{M^*})\pi_{n,\alpha}(\mathrm{d}M)\right] \leq \frac{1+\alpha}{1-\alpha}\varepsilon_n.$$

where

$$\varepsilon_n = C_{a,B} \frac{r(d_1 + d_2)\log(nd_1d_2)}{n},$$

for some universal constant $C_{a,B}$ depending only on a, B. Specifically, the result remains valid for the selection $b = B^2/[512(nd_1d_2)^4K^2\max^2(d_1, d_2)]$.

It is reminded that all technical proofs are postponed to Section 5. The main argument is based on a general scheme for fractional posteriors derived in Bhattacharya et al. (2019); Alquier and Ridgway (2020).

In practical applications, it is noted that $b = B^2/[512(nd_1d_2)^4K^2\max^2(d_1,d_2)]$ may not the best choice; rather, Alquier et al. (2014); Alquier and Ridgway (2020) suggests employing cross-validation to select b. Ensuring a small b is crucial in practical situations to guarantee a reliable approximation of low-rank matrices (Alquier et al., 2014; Alquier and Ridgway, 2020).

The following theorem introduces the first concentration results for the fractional posterior in 1-bit matrix completion when utilizing commonly employed Gaussian priors in practical applications.

Theorem 2. Assume that Assumption 3.1 holds. Then, for a sufficiently small b > 0, such as $b = \frac{B^2}{512(nd_1d_2)^4K^2\max^2(d_1,d_2)}$, it holds that

$$\mathbb{P}\left[\int \mathcal{D}_{\alpha}(P_{M}, P_{M^{*}})\pi_{n,\alpha}(\mathrm{d}M) \leq \frac{2(\alpha+1)}{1-\alpha}\varepsilon_{n}\right] \geq 1 - \frac{2}{n\varepsilon_{n}}$$

where,

$$\varepsilon_n = C_{a,B} \frac{r(d_1 + d_2)\log(nd_1d_2)}{n},$$

for some universal constant $C_{a,B}$ depending only on a and B. Remark 1. It is important to note that our results are formulated without prior knowledge of r, the rank of the true underlying parameter matrix. This aspect highlights the adaptive nature of our results, indicating their ability to adjust and perform effectively regardless of the specific rank of the true underlying parameter matrix. F

$$c_{\alpha} = \begin{cases} \frac{2(\alpha+1)}{1-\alpha}, \alpha \in [0.5, 1), \\ \frac{2(\alpha+1)}{\alpha}, \alpha \in (0, 0.5). \end{cases}$$

Corollary 3. As a special case, Theorem 1 leads to a concentration result in terms of the classical Hellinger distance

$$\mathbb{P}\left[\int H^2(P_M, P_{M^*})\pi_{n,\alpha}(\mathrm{d}M) \le c_\alpha \varepsilon_n\right] \ge 1 - \frac{2}{n\varepsilon_n}.$$
(3)

Remark 2. The rate specified in (3), of the order $r(d_1 + d_2)\log(n)/n$, bears resemblance to that observed in prior studies in frequentist literature like Klopp et al. (2015) when examining a general sampling framework. To elaborate further, Theorem 1 and Lemma 9 in Klopp et al. (2015) delve into the recovery of the distribution f(M); however, they necessitate stricter assumptions. In comparison, our findings in (3)demonstrate a faster rate than those outlined in Davenport et al. (2014) where their results is of order $\sqrt{r(d_1+d_2)\log(\max(d_1,d_2))/n}$.

To derive results directly concerning the parameter matrix, we must make use of the following assumption.

Assumption 3.2. We assume that there exist a constant $C_1 > 0$, such that,

$$\min_{i \in [d_1], j \in [d_2]} \mathbb{P}(\omega_i = (i, j)) \ge C_1.$$

$\mathbf{6}$

This assumption guarantees that each coefficient has a non-zero probability of being observed. For instance, with the uniform distribution, we can express it as $C_1 = 1/(d_1d_2)$. This assumption was initially introduced in Klopp (2014) within the classical unquantized (continuous) matrix completion setting. It is also used for 1-bit matrix completion under a general sampling distribution, as demonstrated in Klopp et al. (2015).

Assumption 3.3. We assume that $||M^*||_{\infty} \leq \kappa < \infty$ and there exist a constant $C_{\kappa} > 0$ such that

$$C_{\kappa} = \inf_{|x| \le \kappa} \frac{f'(x)^2}{8f(x)(1 - f(x))},$$

where $f(x) = e^x/(1 + e^x)$.

Assumption 3.3 stands as a cornerstone requirement essential for deriving insights into estimation errors. It was first introduced in Davenport et al. (2014) and has since served as a fundamental premise in various prior works, including Cai and Zhou (2013) and Klopp et al. (2015).

We are ready to state our main results regarding the recovering of the parameter matrix.

Theorem 4. Under the same assumption as in Theorem 2 and additionally assuming that Assumption 3.2 and Assumption 3.3 hold. We have that

$$\mathbb{P}\left[\int \frac{\|M - M^*\|_F^2}{d_1 d_2} \pi_{n,\alpha}(\mathrm{d}M) \le \frac{c_\alpha}{C_1 C_\kappa} \varepsilon_n\right] \ge 1 - \frac{2}{n\varepsilon_n}.$$
(4)

and

$$\mathbb{P}\left[\frac{\|\hat{M} - M^*\|_F^2}{d_1 d_2} \le \frac{c_\alpha}{C_1 C_\kappa} \varepsilon_n\right] \ge 1 - \frac{2}{n\varepsilon_n}.$$
(5)

Remark 3. Up to a logarithmic factor, the error rate for the mean estimator in the squared Frobenius norm, given in (5), is of order $r(d_1 + d_2)/n$ which is minimax-optimal according to Theorem 3 in Klopp et al. (2015). The result stated in (5) is achieved by applying Jensen's inequality to the mean. By employing similar methods, one can readily derive outcomes for other estimator derived from the fractional posterior, such as the median, drawing upon insights provided in Merkle (2005).

Remark 4. Under the uniform sampling assumption and that $||X||_{\infty} \leq \gamma$, Theorem 1 in Davenport et al. (2014) presented results of order $\sqrt{r(d_1 + d_2)/n}$. A similar result using max-norm minimization was also obtained in Cai and Zhou (2013). The paper Klopp et al. (2015) proves a faster estimation error rate as $r(d_1 + d_2)\log(d_1 + d_2)/n$. A comparable result to Klopp et al. (2015) is also established in Alquier et al. (2019). Subsequently, this rate has been recently enhanced to $r(d_1+d_2)/n$, without the presence of a logarithmic term, in Alaya and Klopp (2019) (refer to Theorem 7). Consequently, the work presented in Alaya and Klopp (2019) attains the precise minimax estimation rate of convergence for 1-bit matrix completion.

Remark 5. It is noteworthy that our findings are established within a general sampling framework. In contrast to the requirements set forth in Klopp et al. (2015), our approach necessitates only that the probability of observing any entries is strictly

positive, without imposing additional assumptions. This aspect further enhances the robustness of employing a fractional posterior.

4 Results with a spectral scaled Student prior

We have opted to initially present results in Section 3 with factorization-type priors, as they are widely favored in the matrix completion literature for utilization with MCMC or Variational Bayes (VB) methods. However, another spectral scaled Student prior has garnered particular interest due to its promising outcomes, whether employed with VB (Yang et al., 2018) or Langevin Monte Carlo, a gradient-based sampling method (Dalalyan, 2020). This prior has previously been applied in different problems involving matrix parameters (Mai, 2023a,b).

With $\tau > 0$, we consider the following spectral scaled Student prior, given as

$$\pi_{st}(M) \propto \det(\tau^2 \mathbf{I}_{d_1} + M M^{\mathsf{T}})^{-(d_1+d_2+2)/2}.$$
 (6)

This prior possesses the capability to introduce approximate low-rankness in matrices M. This is evident from the fact that $\pi_{st}(M) \propto \prod_{j=1}^{d_1} (\tau^2 + s_j(M)^2)^{-(d_1+d_2+2)/2}$, where $s_j(M)$ represents the j^{th} largest singular value of M. Consequently, the distribution follows a scaled Student's t-distribution evaluated at $s_j(M)$, which induces approximate sparsity on $s_j(M)$, as discussed in Dalalyan and Tsybakov (2012b,a). Thus, under this prior distribution, the majority of $s_j(M)$ tend to be close to 0, suggesting that M is approximately low-rank.

We are now present a consistency result using the spectral scaled Student prior. Theorem 5. For $\tau = 1/n$, we have that

$$\mathbb{E}\left[\int \mathcal{D}_{\alpha}(P_M, P_{M^*})\pi_{n,\alpha}(\mathrm{d}M)\right] \leq \frac{1+\alpha}{1-\alpha}\varepsilon_n$$

where

$$=\frac{2r(d_1+d_2+2)\log\left(1+\frac{n\|M^*\|_F}{\sqrt{2r}}\right)}{n}$$

 ε_n

The proofs of this section can be found in Section 5.2. It is noted that in the rate ε_n outlined in Theorem 5 and Theorem 6 below, the condition $r = \operatorname{rank}(M^*) \neq 0$ is not necessary. This is because we interpret $0 \log(1 + 0/0)$ as 0 for the scenario where $r^* = 0$ and $M^* = 0$.

The next Theorem presents a concentration result for the fractional posterior. **Theorem 6.** For $\tau = 1/n$, we have that

$$\mathbb{P}\left[\int \mathcal{D}_{\alpha}(P_M, P_{M^*})\pi_{n,\alpha}(\mathrm{d}M) \leq \frac{2(\alpha+1)}{1-\alpha}\varepsilon_n\right] \geq 1 - \frac{2}{n\varepsilon_n}$$

where

$$\varepsilon_n = \frac{2r(d_1 + d_2 + 2)\log\left(1 + \frac{n\|M^*\|_F}{\sqrt{2r}}\right)}{n}$$

Remark 6. We do not assert that $\tau = 1/n$, in both Theorem 5 and 6, represents the optimal selection. In practical applications, users can utilize cross-validation to fine-tune the value of τ .

Remark 7. It is interesting to observe that by utilizing the spectral scaled Student prior described in (6), we are not required to impose a boundedness assumption on M^* , as was necessary in the previous section with low-rank factorized priors. Furthermore, the additional logarithmic factor in Theorem 5 and Theorem 6 can be further simplified. This can be achieved by employing the inequality $||M^*||_F \leq ||M^*||\sqrt{r}$, resulting in $\log(1 + n||M^*||)$.

Similar to Theorem 4, with the inclusion of additional assumptions, we can derive concentration results for recovering the underlying matrix parameter as well as results for the mean estimator defined in (2).

Theorem 7. Under the same assumption as in Theorem 6 and additional assume that Assumption 3.2 and Assumption 3.3 hold. We have that

$$\mathbb{P}\left[\int \frac{\|M - M^*\|_F^2}{d_1 d_2} \pi_{n,\alpha}(\mathrm{d}M) \le \frac{c_\alpha}{C_1 C_\kappa} \varepsilon_n\right] \ge 1 - \frac{2}{n\varepsilon_n}.$$
(7)

and

$$\mathbb{P}\left[\frac{\|\hat{M} - M^*\|_F^2}{d_1 d_2} \le \frac{c_\alpha}{C_1 C_\kappa} \varepsilon_n\right] \ge 1 - \frac{2}{n\varepsilon_n}.$$
(8)

Remark 8. Similar to the outcomes detailed in Section 3, the results presented in this section for the spectral scaled Student prior do not necessitate prior knowledge of r, the rank of the true underlying parameter matrix. This underscores the adaptive nature of our results, demonstrating their capacity to adjust and perform effectively, regardless of the rank of the true underlying parameter matrix.

5 Proofs

5.1 Proofs for Section 3

Proof of Theorem 1. As the logistic loss is 1-Lipschitz, the log-likelihood satisfies that

$$\left|\log f(x) - \log f(y)\right| \le |x - y|.$$

One has that

$$\begin{aligned} \mathcal{K}(P_{M^*}, P_M) &\leq \frac{1}{d_1 d_2} \sum_{i \in [d_1]} \sum_{j \in [d_2]} \Pi_{ij} |M_{ij}^* - M_{ij}| \\ &\leq \frac{1}{d_1 d_2} \sum_{i \in [d_1]} \sum_{j \in [d_2]} |M_{ij}^* - M_{ij}| \\ &\leq \frac{\|M^* - M\|_F}{\sqrt{d_1 d_2}}, \end{aligned}$$

where $\Pi_{ij} \leq 1$ is the probability to observe the (i, j)-th entry. For any (U, V) in the support of ρ_n , given in (13), one has that

$$\begin{split} \|M^* - UV^t\|_F &= \|\bar{U}\bar{V}^t - \bar{U}V^t + \bar{U}V^t - UV^t\|_F \\ &\leq \|\bar{U}(\bar{V}^t - V^t)\|_F + \|(\bar{U} - U)V^t\|_F \\ &\leq \|\bar{U}\|_F \|\bar{V} - V\|_F + \|\bar{U} - U\|_F \|V^t\|_F \\ &\leq d_1 d_2 \|\bar{U}\|_{\infty}^{1/2} \|\bar{V} - V\|_{\infty}^{1/2} + d_1 d_2 \|V\|_{\infty}^{1/2} \|\bar{U} - U\|_{\infty}^{1/2} \\ &\leq d_1 d_2 \delta^{1/2} [B^{1/2} + (B + \delta)^{1/2}] \\ &\leq 2d_1 d_2 \delta^{1/2} (B + \delta)^{1/2} \\ &\leq 2^{3/2} d_1 d_2 \delta^{1/2} B^{1/2}. \end{split}$$

Therefore,

$$\mathcal{K}(P_{M^*}, P_M) \le \frac{\|M^* - M\|_F}{\sqrt{d_1 d_2}} \le \sqrt{\delta 2^3 d_1 d_2 B}.$$
(9)

For $\delta = B/[8(nd_1d_2)^2]$ that satisfies $0 < \delta < B$, we have that

$$\|M^* - UV^t\|_F \le B/n \tag{10}$$

and

$$\int \mathcal{K}(P_{M^*}, P_M)\rho_n(dM) \le \frac{B}{n\sqrt{d_1d_2}}$$

Now, from Lemma 8, we have that

$$\frac{1}{n}\mathcal{K}(\rho_n,\pi) \le \frac{2(1+2a)r(d_1+d_2)\left[\log(nd_1d_2) + C_a\right]}{n}.$$

We now can apply Theorem 2.6 in Alquier and Ridgway (2020) with ρ_n in (13) and

$$\varepsilon_n = C_{a,B} \frac{r(d_1 + d_2)\log(nd_1d_2)}{n}$$

to obtain the result. The proof is completed.

Proof of Theorem 2. As the logistic loss is 1-Lipschitz, the log-likelihood satisfies that $|\log f(x) - \log f(y)| \le |x - y|$. Thus, we can deduce that

$$\mathbb{E}\left[\log\left(\frac{p_M}{p_{M^*}}\right)^2\right] \le \frac{1}{d_1 d_2} \sum_{i \in [d_1]} \sum_{j \in [d_2]} \prod_{ij} \left(\log f(M_{ij}^*) - \log f(M_{ij})\right)^2$$

$$\leq \frac{1}{d_1 d_2} \sum_{i \in [d_1]} \sum_{j \in [d_2]} \left(\log f(M_{ij}^*) - \log f(M_{ij}) \right)^2$$

$$\leq \frac{1}{d_1 d_2} \sum_{i \in [d_1]} \sum_{j \in [d_2]} \left(M_{ij}^* - M_{ij} \right)^2$$

$$\leq \frac{1}{d_1 d_2} \| M^* - M \|_F^2, \tag{11}$$

where $\Pi_{ij} \leq 1$ is the probability to observe the (i, j)-th entry. From (9), we have that

$$\mathcal{K}(P_{M^*}, P_M) \le \frac{\|M^* - M\|_F}{\sqrt{d_1 d_2}} \le \sqrt{\delta 2^3 d_1 d_2 B},$$

and

$$\mathbb{E}\left[\log\left(\frac{p_M}{p_{M^*}}\right)^2\right] \le \frac{\|M^* - M\|_F^2}{d_1 d_2} \le \delta 2^3 d_1 d_2 B.$$

For any (U, V) in the support of ρ_n , given in (13), we observe that for $\delta = \frac{B}{8(nd_1d_2)^2}$, where δ satisfies $0 < \delta < B$, and from equation (10) we can deduce that

$$\int \mathcal{K}(P_{M^*}, P_M)\rho_n(dM) \le \frac{B}{n\sqrt{d_1d_2}},$$

and

$$\int \mathbb{E}\left[\log^2\left(\frac{p_M}{p_{M^*}}\right)\right] \rho_n(dM) \le \frac{B^2}{n^2 d_1 d_2}.$$

Now, from Lemma 8, we have that

$$\frac{1}{n}\mathcal{K}(\rho_n,\pi) \le \frac{2(1+2a)r(d_1+d_2)\left[\log(nd_1d_2) + C_a\right]}{n}.$$

We now can apply Corollary 2.5 and Theorem 2.4 in Alquier and Ridgway (2020) with ρ_n in (13) and

$$\varepsilon_n = C_{a,B} \frac{r(d_1 + d_2)\log(nd_1d_2)}{n}$$

to obtain the result. The proof is completed.

Proof of Corollary 3. From Van Erven and Harremos (2014), we have that

$$H^{2}(P,Q) \le D_{1/2}(P,Q) \le D_{\alpha}(P,Q)$$

for $\alpha \in [0.5, 1)$. In addition, we also have that

$$D_{1/2}(P,Q) \le \frac{(1-\alpha)1/2}{\alpha(1-1/2)} D_{\alpha}(P,Q) = \frac{(1-\alpha)}{\alpha} D_{\alpha}(P,Q),$$

for $\alpha \in (0, 0.5)$.

Thus, using definition of c_{α} and Theorem 2, we obtain the results.

Proof of Theorem 4. From (3), we have that

$$\mathbb{P}\left[\int H^2(P_M, P_{M^*})\pi_{n,\alpha}(\mathrm{d}M) \le c_\alpha \varepsilon_n\right] \ge 1 - \frac{2}{n\varepsilon_n},$$

from Lemma 9, one has that

$$\mathbb{P}\left[\int \frac{C_1 C_{\kappa} \|M - M^*\|_F^2}{d_1 d_2} \pi_{n,\alpha}(\mathrm{d}M) \le c_{\alpha} \varepsilon_n\right] \ge 1 - \frac{2}{n \varepsilon_n},$$

thus, we obtain (4). To obtain (5), one can apple Jensen's inequality for a convex function, that

$$\|\hat{M} - M^*\|_F^2 = \left\|\int M\pi_{n,\alpha}(\mathrm{d}M) - M^*\right\|_F^2 \le \int \|M - M^*\|_F^2\pi_{n,\alpha}(\mathrm{d}M).$$

This completes the proof.

Proof of Theorem 5. From (9), we have that

$$\mathcal{K}(P_{M^*}, P_M) \le \frac{\|M^* - M\|_F}{\sqrt{d_1 d_2}},$$

When integrating with respect to $\rho_n := \rho_0$ given in (14), we have that

$$\int \mathcal{K}(P_{M^*}, P_M) \rho_n(dM) \leq \int \frac{\|M^* - M\|_F}{\sqrt{d_1 d_2}} \rho_0(dM)$$

= $\int \frac{\|M^* - M\|_F}{\sqrt{d_1 d_2}} \pi_{st}(M - M^*) dM$
= $\frac{1}{\sqrt{d_1 d_2}} \int \|M\|_F \pi_{st}(M) dM$
 $\leq \frac{1}{\sqrt{d_1 d_2}} \left(\int \|M\|_F^2 \pi_{st}(M) dM\right)^{1/2}$

12

L		

$$\leq \frac{1}{\sqrt{d_1 d_2}} \sqrt{d_1 d_2 \tau^2} = \tau,$$
 (12)

where we have used Holder's inequality and Lemma 10 to obtain the result. Now, from Lemma 11, we have that

$$\frac{1}{n}\mathcal{K}(\rho_n, \pi_{st}) \le \frac{2r(d_1+d_2+2)\log\left(1+\frac{\|M^*\|_F}{\tau\sqrt{2r}}\right)}{n}$$

Taking $\tau = 1/n$, we obtain that

$$\int \mathcal{K}(P_{M^*}, P_M)\rho_n(dM) \le \frac{1}{n},$$

$$\frac{1}{n}\mathcal{K}(\rho_n, \pi_{st}) \le \frac{2r(d_1 + d_2 + 2)\log\left(1 + \frac{n\|M^*\|_F}{\sqrt{2r}}\right)}{n}.$$

We now can apply Theorem 2.6 in Alquier and Ridgway (2020) with

$$\varepsilon_n = \frac{2r(d_1 + d_2 + 2)\log\left(1 + \frac{n\|M^*\|_F}{\sqrt{2r}}\right)}{n}$$

to obtain the result. The proof is completed.

Proof of Theorem 6. From (9), we have that

$$\mathcal{K}(P_{M^*}, P_M) \le \frac{\|M^* - M\|_F}{\sqrt{d_1 d_2}},$$

When integrating with respect to $\rho_n := \rho_0$ given in (14), and from (12), we have that

$$\int \mathcal{K}(P_{M^*}, P_M)\rho_n(dM) \le \tau.$$

Now, from Lemma 11, we have that

$$\frac{1}{n}\mathcal{K}(\rho_n, \pi_{st}) \le \frac{2r(d_1 + d_2 + 2)\log\left(1 + \frac{\|M^*\|_F}{\tau\sqrt{2r}}\right)}{n}.$$

Moreover, from (11), one has that

$$\mathbb{E}\left[\log\left(\frac{p_M}{p_{M^*}}\right)^2\right] \le \frac{\|M - M^*\|_F^2}{d_1 d_2}$$

1	0
Т	.5
	~

and when integrating with respect to $\rho_n := \rho_0$ given in (14), it leads to

$$\int \mathbb{E} \left[\log \left(\frac{p_M}{p_{M^*}} \right)^2 \right] \rho_n(dM) \le \int \frac{\|M - M^*\|_F^2}{d_1 d_2} \rho_n(dM)$$

= $\int \frac{\|M - M^*\|_F^2}{d_1 d_2} \pi_{st}(M - M^*) dM$
= $\frac{1}{d_1 d_2} \int \|M\|_F^2 \pi_{st}(M) dM$
 $\le \frac{d_1 d_2 \tau^2}{d_1 d_2} = \tau^2,$

where we have used a change of variable and Lemma 10 to obtain the result. Now, by taking $\tau = 1/n$, we obtain that

$$\int \mathcal{K}(P_{M^*}, P_M)\rho_n(dM) \leq \frac{1}{n},$$

$$\int \mathbb{E} \left[\log \left(\frac{p_M}{p_{M^*}} \right)^2 \right] \rho_n(dM) \leq \frac{1}{n^2},$$

$$\frac{1}{n} \mathcal{K}(\rho_n, \pi_{st}) \leq \frac{2r(d_1 + d_2 + 2)\log\left(1 + \frac{n ||M^*||_F}{\sqrt{2r}}\right)}{n}.$$

We now can apply Theorem 2.4 and Corollary 2.5 in Alquier and Ridgway (2020) with

$$\varepsilon_n = \frac{2r(d_1 + d_2 + 2)\log\left(1 + \frac{n\|M^*\|_F}{\sqrt{2r}}\right)}{n}$$

to obtain the result. The proof is completed.

Proof of Theorem 7. From Theorem 6, using a bound for Hellinger distance as in Corollary 3, we have that

$$\mathbb{P}\left[\int H^2(P_M, P_{M^*})\pi_{n,\alpha}(\mathrm{d}M) \le c_\alpha \varepsilon_n\right] \ge 1 - \frac{2}{n\varepsilon_n},$$

from Lemma 9, it yields that

$$\mathbb{P}\left[\int \frac{C_1 C_{\kappa} \|M - M^*\|_F^2}{d_1 d_2} \pi_{n,\alpha}(\mathrm{d}M) \le c_{\alpha} \varepsilon_n\right] \ge 1 - \frac{2}{n \varepsilon_n},$$

thus, we obtain (7). To obtain (8), one can apple Jensen's inequality for a convex function, that

$$\|\hat{M} - M^*\|_F^2 = \left\|\int M\pi_{n,\alpha}(\mathrm{d}M) - M^*\right\|_F^2 \le \int \|M - M^*\|_F^2\pi_{n,\alpha}(\mathrm{d}M),$$

and combine with result in (7). This completes the proof.

5.3 Lemma

Definition 1. Fix B > 0, $r \ge 1$. For any pair $(\overline{U}, \overline{V}) \in \mathcal{M}(r, B)$, we define for $\delta \in (0, B)$ that will be chosen later,

$$\rho_n(\mathrm{d}U,\mathrm{d}V,\mathrm{d}\gamma) \propto \mathbf{1}_{(\|U-\bar{U}\|_{\infty} \le \delta, \|U-\bar{U}\|_{\infty} \le \delta)} \pi(\mathrm{d}U,\mathrm{d}V,\mathrm{d}\gamma).$$
(13)

Lemma 8. Put $C_a := \log(8\sqrt{\pi}\Gamma(a)2^{10a+1}) + 3$ and with $\delta = B/[8(nd_1d_2)^2]$ that satisfies $0 < \delta < B$, we have for ρ_n in (13) that

$$\mathcal{K}(\rho_n, \pi) \le 2(1+2a)r(d_1+d_2)\left[\log(nd_1d_2) + C_a\right].$$

Proof of Lemma 8. This result can found, for example, in the proof of Theorem 4.1 in Alquier and Ridgway (2020).

Lemma 9. For any matrix $A \in \mathbb{R}^{d_1 \times d_2}$ and $B \in \mathbb{R}^{d_1 \times d_2}$ satisfying that $||A||_{\infty} \leq \kappa$ and $||B||_{\infty} \leq \kappa$, under Assumption 3.3 and Assumption 3.2, one has that

$$\frac{\|A - B\|_F^2}{d_1 d_2} \le \frac{H^2(P_A, P_B)}{C_1 C_{\kappa}}.$$

Proof of Lemma 9. This is Lemma A.2 in Davenport et al. (2014). With $d_H^2(p,q) := (\sqrt{p} - \sqrt{q})^2 + (\sqrt{1-p} - \sqrt{1-q})^2$ for two number $p, q \in [0, 1]$, it is noting under Assumption 3.2 that

$$H^{2}(P_{A}, P_{B}) = \frac{1}{d_{1}d_{2}} \sum_{i \in [d_{1}]} \sum_{j \in [d_{2}]} \Pi_{ij} d_{H}^{2}(f(A_{ij}), f(B_{ij}))$$

$$\geq \frac{C_{1}}{d_{1}d_{2}} \sum_{i \in [d_{1}]} \sum_{j \in [d_{2}]} d_{H}^{2}(f(A_{ij}), f(B_{ij}))$$

$$\geq C_{1}H^{2}(f(A), f(B)).$$

where Π_{ij} is the probability to observe the (i, j)-th entry. Now from Lemma A.2 in Davenport et al. (2014), under Assumption 3.3, one has that

$$H^2(f(A), f(B)) \ge C_{\kappa} \frac{\|A - B\|_F^2}{d_1 d_2}.$$

1	5
т	J

The argument is also similar to Lemma 9 and Lemma 11 in Klopp et al. (2015). This completes the proof.

Finally, we will use quite often the following distribution that will be defined as translations of the prior π_{st} in (6). We introduce the following notation. **Definition 2.** Let's define

$$\rho_0(M) = \pi_{st}(M - M^*). \tag{14}$$

The following technical lemmas will be useful in the proofs. Lemma 10 (Lemma 1 in Dalalyan (2020)). We have

$$\int \|M\|_F^2 \pi_{st}(M) \mathrm{d}M \le d_1 d_2 \tau^2.$$

Lemma 11 (Lemma 2 in Dalalyan (2020)). We have

$$KL(\rho_0, \pi_{st}) \le 2r(d_1 + d_2 + 2)\log\left(1 + \frac{\|M^*\|_F}{\tau\sqrt{2r}}\right),$$

with the convention $0\log(1+0/0) = 0$.

Acknowledgements. The author was supported by the Norwegian Research Council, grant number 309960, through the Centre for Geophysical Forecasting at NTNU.

Conflict of interest/Competing interests:

The author declares no potential conflict of interests.

References

- Alquier, P., Cottet, V., Chopin, N., Rousseau, J.: Bayesian matrix completion: prior specification and consistency. arXiv preprint arXiv:1406.1440 (2014)
- Alquier, P., Cottet, V., Lecué, G.: Estimation bounds and sharp oracle inequalities of regularized procedures with lipschitz loss functions. The Annals of Statistics 47(4), 2117–2144 (2019)
- Alaya, M.Z., Klopp, O.: Collective matrix completion. J. Mach. Learn. Res. 20, 148–1 (2019)
- Alquier, P., Ridgway, J.: Concentration of tempered posteriors and of their variational approximations. The Annals of Statistics 48(3), 1475–1497 (2020)
- Bissiri, P.G., Holmes, C.C., Walker, S.G.: A general framework for updating belief distributions. Journal of the Royal Statistical Society: Series B (Statistical Methodology) 78(5), 1103–1130 (2016)

- Bennett, J., Lanning, S.: The netflix prize. In: Proceedings of KDD Cup and Workshop, vol. 2007, p. 35 (2007)
- Babacan, S.D., Luessi, M., Molina, R., Katsaggelos, A.K.: Sparse Bayesian methods for low-rank matrix estimation. IEEE Transactions on Signal Processing 60(8), 3964–3977 (2012)
- Bobadilla, J., Ortega, F., Hernando, A., Gutiérrez, A.: Recommender systems survey. Knowledge-based systems 46, 109–132 (2013)
- Bhattacharya, A., Pati, D., Yang, Y.: Bayesian fractional posteriors. Annals of Statistics 47(1), 39–66 (2019)
- Cottet, V., Alquier, P.: 1-bit matrix completion: Pac-bayesian analysis of a variational approximation. Machine Learning **107**(3), 579–603 (2018)
- Chen, Y., Fan, J., Ma, C., Yan, Y.: Inference and uncertainty quantification for noisy matrix completion. Proceedings of the National Academy of Sciences 116(46), 22931–22937 (2019)
- Chatterjee, S.: Matrix estimation by universal singular value thresholding. The Annals of Statistics **43**(1), 177–214 (2015)
- Candes, E.J., Plan, Y.: Matrix completion with noise. Proceedings of the IEEE **98**(6), 925–936 (2010)
- Candès, E.J., Recht, B.: Exact matrix completion via convex optimization. Found. Comput. Math. 9(6), 717–772 (2009) https://doi.org/10.1007/s10208-009-9045-5
- Candès, E.J., Tao, T.: The power of convex relaxation: near-optimal matrix completion. IEEE Trans. Inform. Theory 56(5), 2053–2080 (2010) https://doi.org/10. 1109/TIT.2010.2044061
- Cai, T., Zhou, W.-X.: A max-norm constrained minimization approach to 1-bit matrix completion. J. Mach. Learn. Res. 14(1), 3619–3647 (2013)
- Chi, E.C., Zhou, H., Chen, G.K., Del Vecchyo, D.O., Lange, K.: Genotype imputation via matrix completion. Genome research **23**(3), 509–518 (2013)
- Dalalyan, A.S.: Theoretical guarantees for approximate sampling from smooth and log-concave densities. Journal of the Royal Statistical Society: Series B Statistical Methodology 79(3), 651–676 (2017)
- Dalalyan, A.S.: Exponential weights in multivariate regression and a low-rankness favoring prior. Annales de l'Institut Henri Poincaré, Probabilités et Statistiques **56**(2), 1465–1483 (2020)
- Dalalyan, A.S., Karagulyan, A.: User-friendly guarantees for the langevin monte carlo

with inaccurate gradient. Stochastic Processes and their Applications 129(12), 5278-5311 (2019)

- Durmus, A., Moulines, E.: Nonasymptotic convergence analysis for the unadjusted langevin algorithm. The Annals of Applied Probability **27**(3), 1551–1587 (2017)
- Durmus, A., Moulines, E.: High-dimensional Bayesian inference via the unadjusted langevin algorithm. Bernoulli 25(4A), 2854–2882 (2019)
- Davenport, M.A., Plan, Y., Van Den Berg, E., Wootters, M.: 1-bit matrix completion. Information and Inference: A Journal of the IMA 3(3), 189–223 (2014)
- Dalalyan, A.S., Tsybakov, A.: Mirror averaging with sparsity priors. Bernoulli 18(3), 914–944 (2012)
- Dalalyan, A.S., Tsybakov, A.B.: Sparse regression learning by aggregation and langevin monte-carlo. Journal of Computer and System Sciences **78**(5), 1423–1443 (2012)
- Gross, D.: Recovering low-rank matrices from few coefficients in any basis. IEEE Transactions on Information Theory **57**(3), 1548–1566 (2011)
- Grünwald, P., Van Ommen, T.: Inconsistency of Bayesian inference for misspecified linear models, and a proposal for repairing it. Bayesian Analysis 12(4), 1069–1103 (2017)
- Hong, L., Martin, R.: Model misspecification, Bayesian versus credibility estimation, and Gibbs posteriors. Scandinavian Actuarial Journal 2020(7), 634–649 (2020)
- Hsieh, C.-J., Natarajan, N., Dhillon, I.: Pu learning for matrix completion. In: International Conference on Machine Learning, pp. 2445–2453 (2015). PMLR
- Herbster, M., Pasteris, S., Pontil, M.: Mistake bounds for binary matrix completion. Advances in Neural Information Processing Systems **29** (2016)
- Hammer, H.L., Riegler, M.A., Tjelmeland, H.: Approximate bayesian inference based on expected evaluation. Bayesian Analysis, 1–22 (2023)
- Han, X., Wu, J., Wang, L., Chen, Y., Senhadji, L., Shu, H.: Linear total variation approximate regularized nuclear norm optimization for matrix completion. In: Abstract and Applied Analysis, vol. 2014 (2014). Hindawi
- Ji, H., Liu, C., Shen, Z., Xu, Y.: Robust video denoising using low rank matrix completion. In: 2010 IEEE Computer Society Conference on Computer Vision and Pattern Recognition, pp. 1791–1798 (2010). IEEE
- Jiang, B., Ma, S., Causey, J., Qiao, L., Hardin, M.P., Bitts, I., Johnson, D., Zhang, S., Huang, X.: Sparrec: An effective matrix completion framework of missing data

imputation for gwas. Scientific reports 6(1), 35534 (2016)

- Jewson, J., Rossell, D.: General bayesian loss function selection and the use of improper models. Journal of the Royal Statistical Society Series B: Statistical Methodology 84(5), 1640–1665 (2022)
- Koren, Y., Bell, R., Volinsky, C.: Matrix factorization techniques for recommender systems. Computer 42(8), 30–37 (2009)
- Kyung, M., Gill, J., Ghosh, M., Casella, G.: Penalized regression, standard errors, and bayesian lassos. Bayesian Analysis 5(2), 369–412 (2010)
- Knoblauch, J., Jewson, J., Damoulas, T.: An optimization-centric view on bayes' rule: Reviewing and generalizing variational inference. Journal of Machine Learning Research 23(132), 1–109 (2022)
- Klopp, O., Lafond, J., Moulines, É., Salmon, J.: Adaptive multinomial matrix completion. Electronic Journal of Statistics 9, 2950–2975 (2015)
- Klopp, O.: Noisy low-rank matrix completion with general sampling distribution. Bernoulli **20**(1), 282–303 (2014) https://doi.org/10.3150/12-BEJ486
- Lyddon, S.P., Holmes, C., Walker, S.: General bayesian updating and the losslikelihood bootstrap. Biometrika **106**(2), 465–478 (2019)
- Lim, Y.J., Teh, Y.W.: Variational bayesian approach to movie rating prediction. Proceedings of KDD cup and workshop 7, 15–21 (2007)
- Mai, T.T., Alquier, P.: A bayesian approach for noisy matrix completion: Optimal rate under general sampling distribution. Electron. J. Statist. 9(1), 823–841 (2015) https://doi.org/10.1214/15-EJS1020
- Mai, T.T., Alquier, P.: Pseudo-Bayesian quantum tomography with rank-adaptation. Journal of Statistical Planning and Inference 184, 62–76 (2017)
- Mai, T.T.: From bilinear regression to inductive matrix completion: a quasi-bayesian analysis. Entropy **25**(2), 333 (2023)
- Mai, T.T.: A reduced-rank approach to predicting multiple binary responses through machine learning. Statistics and Computing **33**(6), 136 (2023)
- Merkle, M.: Jensen's inequality for medians. Statistics & probability letters 71(3), 277–281 (2005)
- Matsubara, T., Knoblauch, J., Briol, F.-X., Oates, C.J.: Robust generalised bayesian inference for intractable likelihoods. Journal of the Royal Statistical Society Series B: Statistical Methodology 84(3), 997–1022 (2022)

- Medina, M.A., Olea, J.L.M., Rush, C., Velez, A.: On the robustness to misspecification of α -posteriors and their variational approximations. Journal of Machine Learning Research **23**(147), 1–51 (2022)
- Park, T., Casella, G.: The bayesian lasso. Journal of the american statistical association **103**(482), 681–686 (2008)
- Recht, B., Ré, C.: Parallel stochastic gradient algorithms for large-scale matrix completion. Math. Program. Comput. 5(2), 201–226 (2013) https://doi.org/10.1007/s12532-013-0053-8
- Salakhutdinov, R., Mnih, A.: Bayesian probabilistic matrix factorization using markov chain monte carlo. In: Proceedings of the 25th International Conference on Machine Learning, pp. 880–887 (2008). ACM
- Syring, N., Martin, R.: Calibrating general posterior credible regions. Biometrika 106(2), 479–486 (2019)
- Tsybakov, A.B., Koltchinskii, V., Lounici, K.: Nuclear-norm penalization and optimal rates for noisy low-rank matrix completion. Annals of Statistics **39**(5), 2302–2329 (2011)
- Van Erven, T., Harremos, P.: Rényi divergence and kullback-leibler divergence. IEEE Transactions on Information Theory 60(7), 3797–3820 (2014)
- Yang, L., Fang, J., Duan, H., Li, H., Zeng, B.: Fast low-rank Bayesian matrix completion with hierarchical gaussian prior models. IEEE Transactions on Signal Processing 66(11), 2804–2817 (2018)
- Yang, Y., Pati, D., Bhattacharya, A.: α-variational inference with statistical guarantees. Annals of Statistics 48(2), 886–905 (2020)
- Yonekura, S., Sugasawa, S.: Adaptation of the tuning parameter in general bayesian inference with robust divergence. Statistics and Computing **33**(2), 39 (2023)