

00178. The evolutionary divergence of Mars, Venus and Earth

Keiko Hamano^a, Cedric Gillmann^b, Gregor J. Golabek^c, Diogo Lourenço^b,
Frances Westall^d

^a*Earth-Life Science Institute, Tokyo Institute of Technology, 2-12-1-IE-1
Ookayama, Meguro-ku, 152-8550, Tokyo, Japan*

^b*ETH Zürich, Institute of Geophysics, Sonneggstrasse 5, 8092, Zürich, Switzerland*

^c*Bayerisches Geoinstitut, University of Bayreuth, Universitätsstrasse
30, 95440, Bayreuth, Germany*

^d*CNRS-Centre de Biophysique Moléculaire, Rue Charles Sadron CS
80054, 45071, Orléans, France*

Abstract

Mars, Venus and Earth are expected to have started in a hot molten state. Here, we discuss how these three terrestrial planets diverged in their evolution and what mechanisms could be the cause. We discuss that early-on after magma ocean crystallization the mantle/surface redox state and water inventory may already differ considerably, depending on planetary mass and orbital distance from the Sun. During the subsequent internal evolution, the three planets also diverged in terms of their tectonic regime, affecting the long-term planetary evolution via heat flux and outgassing rate, and possibly the physical state of their respective core and the onset and end of a planetary dynamo. We discuss how, throughout the evolution of these rocky planets, the dominant process for atmospheric loss would shift from hydrodynamic escape and impact erosion to non-thermal escape, where small terrestrial planets like Mars are here more vulnerable to volatile loss.

Keywords: atmosphere evolution, comparative planetology, Earth, habitability, life, magma ocean, mantle dynamics, Mars, planetary evolution, Solar System, terrestrial planets, Venus, volatile cycles, volcanism

Key points/objectives box

- We describe characteristics and processes that played a role for the divergence among the three planets Mars, Venus and Earth.
- We discuss how the primitive magma ocean phase could cause the divergence early on.
- We investigate long-term interior, surface and atmosphere processes that can enhance those early established differences.

1. Introduction

Earth, Mars and Venus are all classified as “terrestrial planets”, as opposed to gaseous planets. They have solid surfaces and are composed mostly of silicate rocks and metals with a relatively thin outer atmospheric layer of liquids and gas trapped by gravity. Their building blocks are generally considered to be of similar origin (Smrekar et al., 2018; Lammer et al., 2018), coming from the planetary accretion disc, but their respective present-day states are remarkably different.

The characteristics in the past and present on each planet are described in detail in other chapters, thus here, we briefly compare the three planets. Venus and Earth share similarities in their sizes, masses, and densities, and their orbits are rather close to each other (Table 1). These characteristics make it possible for Venus to have a comparable structure to Earth, although measurements are currently not precise enough to fully resolve the question (e.g. Dumoulin et al., 2017). Comparatively, Mars is smaller and less dense than the two other planets and orbits further away from the Sun.

The most obvious differences between the three planets are readily observable in their atmospheres and surface conditions. Earth has a mild climate with an average temperature of 288 K, and harbors stable liquid water at its surface. It currently has a 1 bar atmosphere mainly composed of nitrogen and oxygen with minor species including noble gases and carbon dioxide. By contrast, Venus and Mars have similar atmosphere compositions dominated by CO₂ and N₂. SO₂ can be observed on Venus (150 ppmv), but very little water (20 ppmv). Mars offers a wide range of species with O₂ (0.16%), CO (0.06%), H₂O and NO. Compared to Earth’s atmosphere, Venus has an atmosphere that is about 100 times more massive ($\sim 4.8 \times 10^{20}$ kg), while the Martian atmosphere is about 100 times less massive ($\sim 2.5 \times 10^{16}$ kg). Surface temperatures also cover a wide range, between Mars’ chilly 210 K average (with large variations) and Venus’ hellish 737 K. It is interesting to

note that, due to its thick atmosphere, Venus boasts a larger atmospheric mass of N_2 than the Earth, despite it making up a much smaller percentage of the total surface pressure. Additionally, much of Earth's superficial carbon inventory is stored in carbonates rather than in the planet's atmosphere. This inventory is estimated to be similar to the amount of CO_2 , present in the atmosphere of Venus (Donahue and Pollack, 1983; Wedepohl, 1995; Lécuyer et al., 2000; Hartmann, 2012).

Table 1: Main characteristics of the present Earth, Venus and Mars

	unit	Earth	Venus	Mars
Radius	(Earth=1)	1	0.949	0.532
Mass	(Earth=1)	1	0.815	0.107
Mean density	(Earth=1)	1	0.951	0.714
Semi-major axis	au	1	0.723	1.52
Mean surface temperature	K	288	737	210
Average surface pressure	bar	1	92	6.36×10^{-3}
Atmospheric composition	vol. %	N ₂ (78.08), O ₂ (20.95), with noble gases and CO ₂	CO ₂ (96.5), N ₂ (3.5), and SO ₂ (0.015)	CO ₂ (95.1), N ₂ (2.59), O ₂ (0.16), CO(0.06) with H ₂ O and NO
H ₂ O inventory				
surface+atmosphere		1 ocean	$< 2 \times 10^{-4}$ ocean	20-30 mGEL (polar caps and permafrost)
interior		0-6 oceans	-	100-1000 ppm
Redox state in the upper mantle		IW+1-5	-	IW+0-1
Magnetic field				
Surface field	T	5×10^{-5}	$< 10^{-8}$	$10^{-9} - 10^{-4}$ (patchy but locally strong)
Interpretation		Core dynamo	No dynamo	Remanent crustal field
Tectonic mode		Plate tectonics	Single plate/ Stagnant lid	Stagnant lid

Data sources: Radius, mass, mean density, semi-major axis, mean surface temperature, surface pressure and atmospheric composition are from Planetary Fact Sheet (https://nssdc.gsfc.nasa.gov/planetary/factsheet/planet_table_ratio.html). H₂O inventory from Peslier et al. (2017) for Earth and Scheller et al. (2021) for Mars. H₂O inventory in Venus’s atmosphere was estimated from concentration of water vapor obtained by Venus Express (~ 30 ppmv, Marcq et al., 2017). The unit “ocean” represents the Earth’s ocean mass, 1.4×10^{21} kg, and “GEL” represents global equivalent layer. Redox state in the upper mantle from Righter et al. (2016). For Earth, the redox state shows a large variation, depending on the rock samples (xenoliths, peridotites, MORB, and Kilauea lava samples). Magnetic field informations from Stevenson (2003).

Going deeper than surface observation and delving into the interior characteristics of the planets is hampered by the relative paucity of data available beyond Earth. Samples from Mars and in-situ measurements provide some constraints (i.e. water inventory, redox state, isotopic measurements, core size - see Table 1) that are still unavailable for Venus, which implies that a large part of our understanding of Venus relies on educated guesses based on Earth observations and the consequences of common mechanisms. The present Earth is known for having an oxidized upper mantle, reflected by its oxygen fugacity (a measure of the oxidation state of the mantle, from the ratio of the relative abundance of iron (III) ions to total iron, and depending on pressure and temperature; see also Section 2). In Earth’s present-day upper mantle, oxygen fugacity relative to the iron-wüstite (IW) buffer is estimated to be about IW+1~IW+5 (Richter et al., 2016; Hirschmann, 2022, and references therein), that is between 1 and 5 logarithmic units above the IW buffer. Higher values (up to IW+8) also have been measured in oxidized samples possibly due to processes in the crust or metasomatism. These values are several orders of magnitude higher than the value estimated from FeO content in the upper mantle based on the assumption that mantle material was in equilibrium with Fe metal during core formation ($\sim$ IW-2)(e.g. Frost et al., 2008). Earth’s upper mantle oxygen fugacity appears to have been stable over the last $\approx$ 2 Gyr (Hirschmann, 2023; Stagno and Aulbach, 2021). Some heterogeneity is expected depending on the source region of the samples and their evolution (Hirschmann, 2023). Based on Martian meteorites, the Martian upper mantle seems to have an oxygen fugacity of IW+0~IW+1(Richter et al., 2016, and references therein). This value is also larger than the level set by metal-silicate equilibrium ($\sim$ IW-1) based on FeO content in the Martian mantle, but the difference is much smaller than Earth’s. The redox state of the Venusian mantle remains unknown. Calculations based on the equilibration during core formation yield a range of values for oxygen fugacity similar to Earth’s (Richter et al., 2016), that would then have probably evolved toward more oxidized values. Venus’s surface may be highly oxidized as suggested by thermodynamic calculations and spectroscopic data (Fegley et al., 1997).

It is known that Venus shows no signs of a self-generated magnetic field at present-day, even though it has been speculated that remanent magnetisation may still exist in the crust (O’Rourke et al., 2018). On the contrary, Earth sustains an intrinsic magnetic field at present-day and possibly since the Hadean (Tarduno et al., 2015), while Mars shows signs of a past magnetic

dynamo in patches of remanently magnetized ancient crustal material (Acuña et al., 1999; Plattner and Simons, 2015).

Observation of the three planets has also led to hypotheses regarding their internal dynamics. Earth’s mantle convection regime is known as plate tectonics (i.e. Wessel and Müller, 2015), where the strong lithosphere is broken into distinct mobile plates with deformation mostly localized at the plate boundaries and subduction of oceanic plates being the dominant mechanism for driving plate tectonics (e.g. Forsyth and Uyeda, 1975). As a result, very little ancient crust still remains at the surface of the Earth. In contrast, Mars exhibits a so-called stagnant-lid regime, where a thick rigid lithosphere is decoupled from the convective mantle and mostly immobile on top of the mantle (e.g. Solomatov, 1995). Most of Mars’ surface is thus ancient (billions of years old), but shows a specific pattern: a dichotomy between the southern highlands and the northern lowlands, with a ~ 5 km difference in elevation. Venus is located somewhere in the middle. Its surface shows no sign of (i) global plate tectonics, (ii) localized subduction zones and (iii) a clear topographic dichotomy between oceanic and continental crust as observed on Earth. However, its surface is on average relatively young and signs of deformation have been noted. Several explanations have been proposed, such as episodic activity (Turcotte, 1993) or lithospheric weakening by magma intrusion (the so-called plutonic squishy lid; Rozel et al., 2017; Lourenço et al., 2018). The offset between the center of mass and the center of figure of a planet (CM-CF offset) indicates differences in interior dynamics, since they would emerge from substantial large scale density anomalies in the solid planet, possibly due to interior temperature or topography (King, 2018). The CM-CF offset has been estimated to be smaller on Venus (280 m, Bindshadler et al. (1994)) than for Earth (2.1 km, Yoder and Ahrens (1995)) and Mars (3.3 km, Wieczorek (2015)). The martian offset has been suggested to be caused by the crustal dichotomy and the Tharsis province (Wieczorek, 2015). The difference between Earth and Venus is more difficult to explain, with no uniquely identified mechanism that could act as a cause. It has been suggested that the small CM-CF on Venus could be indicative of the lack of substantial downwellings during the last tens to hundreds of million years of the planet evolution and possible lack of a mobile lid regime (King, 2018).

In this chapter, we attempt to provide an overview of recent efforts to clarify what factors and processes make the current state of the three planets so diverse. We focus on physical and chemical processes that occurred after


their formation stage, while it should be noted that diversification of terrestrial planets may also originate during their accretion stage (e.g. Jacobson et al., 2017; Rubie et al., 2015a). Section 2 describes what differences are expected to arise as a result of the formation of deep magma oceans. In Section 3, we address the diversity in tectonic regime of the three planets during the subsequent internal evolution and its consequences on the long-term planetary evolution. How the dominant process for atmospheric loss would shift throughout the evolution of these rocky planets is discussed in Section 4. In Section 5, we review the unique features in the history of Mars, Venus and Earth that arose from their diverging evolutionary pathways. Finally, we conclude this chapter, with some perspectives in Section 6.

2. Diversity originating during a magma ocean stage

Terrestrial planets, especially Earth, likely undergo a single or multiple global melting by giant impacts (e.g. Canup, 2012; Tucker and Mukhopadhyay, 2014; Nakajima and Stevenson, 2015; Lock et al., 2018). The energy imparted into the planet during a succession of major collision events can be large enough to melt a thick silicate outer layer, possibly down to the core-mantle boundary, producing a deep magma ocean (e.g. Wetherill, 1976; Drake, 2000; Nakajima et al., 2021). The planetary characteristics originated during crystallization of a magma ocean would set initial conditions for the subsequent long-term evolution of planetary interior and atmosphere. Here, we focus on two processes that could occur during a magma ocean phase, iron redox disproportionation and appearance of threshold flux of hot steam atmospheres. Recent studies on these two processes highlight the importance of the planetary mass and the orbital distance from the Sun as potential causes of planetary diversity originated during their magma ocean stages, as described in details in this section. We address the proposed mechanisms and predicted characteristics of planets that evolved from a hot molten state.


2.1. Iron disproportionation in silicate liquid

Early planetary mantles could have a redox state which reflects O partitioning among interior reservoirs, affecting the composition of subsequently outgassed volcanic gases and thus long-term climate. One of the mechanisms proposed to affect the redox state in a planetary mantle is iron redox disproportionation,




The original idea is that crystallization of Fe^{3+} -rich solid phases such as bridgmanite (the dominant mineral in the Earth’s lower mantle) produces Fe metal by this reaction. Segregation of the produced Fe metal into the core enriches the lower mantle in Fe^{3+} . Then, the later mantle convection homogenizes the Fe^{3+} -rich lower mantle with the upper mantle, and raises the Fe^{3+} content in the upper mantle gradually (e.g. Frost et al., 2008). If the same iron disproportionation occurs and produces Fe metal directly in liquid silicate, segregation of Fe metal and homogenization of the remaining Fe^{3+} melt can occur very quickly, possibly increasing the Fe^{3+} abundance in the whole part of the magma ocean (Armstrong et al., 2019).

Under metal-saturated conditions, the partitioning of O between metal and silicate melt determines the oxygen fugacity (f_{O_2}).



where the superscripts ‘met’ and ‘sil’ indicate a molten metallic phase and a silicate liquid phase, respectively. Ferric iron in silicate melt is also set by the reaction.



The molar ratio of Fe^{3+} to total Fe ($\text{Fe}^{3+}/\Sigma\text{Fe}$) in silicate melt increases with the oxygen fugacity, and also varies with pressure, because of the difference in partial molar volumes Δv between the ferric and ferrous oxide components (e.g. Kress and Carmichael, 1991; O’Neill et al., 2006; Hirschmann, 2012)¹. This suggests that the $\text{Fe}^{3+}/\Sigma\text{Fe}$ ratio in a magma ocean may depend on the planetary mass, since pressures in a magma ocean increase with the planetary mass.

The pressure dependence of Δv and $\text{Fe}^{3+}/\Sigma\text{Fe}$ ratios was explored recently by high-pressure experiments (Zhang et al., 2017; Armstrong et al., 2019; Kuwahara et al., 2023) and first principle calculations (Deng et al., 2020). $\text{Fe}^{3+}/\Sigma\text{Fe}$ decreases with pressure up to about 10 GPa owing to the large volume of $\text{FeO}_{1.5}$ compared to that of FeO (Kress and Carmichael, 1991;

¹At an oxygen fugacity fixed relative to a specific oxygen buffer, the pressure effect depends on Δv relative to a change in partial molar volumes associated with the buffer reaction (Hirschmann, 2022).

O'Neill et al., 2006; Zhang et al., 2017). However, this trend reverses at higher pressures because $\text{FeO}_{1.5}$ has a larger compressibility than FeO (Armstrong et al., 2019; Deng et al., 2020). Accordingly, Fe^{3+} is more stable at pressures higher than around 10 GPa, possibly up to 90 GPa. These studies show that silicate melt with a higher $\text{Fe}^{3+}/\Sigma\text{Fe}$ ratio can be in equilibrium with Fe metal at higher pressure, suggesting that the disproportionation could occur in silicate melt. The increasing stability of Fe^{3+} at high pressures is also reported for basaltic and peridotitic melts based on high pressure experiments by Kuwahara et al. (2023). Pressures in a magma ocean would be higher for larger planets. Therefore, the enhanced stability of Fe^{3+} suggests that larger planets have a higher abundance of Fe^{3+} in their magma ocean under the same oxygen fugacity and the presence of Fe metal.

2.2. Potential correlation between planetary mass and atmospheric composition

Fe metal produced by the disproportionation could sink quickly in the magma ocean (e.g. Ichikawa et al., 2010). If the segregated Fe metal ponds and equilibrates with the magma ocean at its base, then reaction (2) sets the f_{O_2} at the base of the magma ocean. Due to the pressure dependence of Δv as described above, the f_{O_2} in the silicate melt can vary as it ascends along a geotherm in the metal-free magma ocean (Hirschmann, 2012), when the $\text{Fe}^{3+}/\Sigma\text{Fe}$ ratio is assumed to be constant in a highly turbulent magma ocean. The f_{O_2} at the top of a magma ocean is likely higher than that at its base, when the base pressure exceeds roughly 10 GPa (Armstrong et al., 2019; Deng et al., 2020; Kuwahara et al., 2023), though some differences exist among these studies. In short, on small planets like the Moon and Mercury, the oxygen fugacity would monotonically decrease from the base of the magma ocean towards the surface (Zhang et al., 2017), while relatively large planets on Earth and Mars the shallow part would be more oxidized than its base (Armstrong et al., 2019; Deng et al., 2020), when the $\text{FeO-FeO}_{1.5}$ reaction (3) buffers the oxygen fugacity in the magma ocean.

Based on the assumption that metal-silicate equilibration occurred at the base of the magma ocean, these recent findings suggest that a larger planet features more Fe^{3+} in the silicate melt and a more oxidized surface during its deep magma ocean stage. Deng et al. (2020) proposed a trend between the Fe^{3+} abundance, the surface redox state, the speciation of atmospheres, and the planetary mass (Figure 1). As the authors used FeO content in the present upper mantle of Earth, Mars and Moon, the f_{O_2} at the base of the




Figure 1: The proposed trend in atmospheric compositions formed above a deep magma ocean. The atmospheric composition reflects the surface redox state and the partitioning of C-H-O-N-S elements between a magma ocean and its atmosphere. The oxygen fugacity at the surface is estimated to be IW-2~-1.5 for the Moon, IW-1~-+1 for Mars, and IW+3~-+5 for Earth based on the assumption of metal-silicate equilibrium at the base of the magma ocean (Deng et al., 2020). The difference in oxygen fugacity affects the gas speciation and the volatile partitioning. H_2O is highly soluble in silicate melt so that its majority is dissolved in the magma ocean. Modified from Deng et al. (2020) under CC by 4.0 license (<http://creativecommons.org/licenses/by/4.0/>), based on numerical results by Gaillard et al. (2022).

magma ocean also differs among the planets, but the overall trend would be similar even if the same f_{O_2} is imposed at the base: a larger planet has an oxidizing surface above a Fe^{3+} -rich magma ocean.

The oxygen fugacity at the surface controls the speciation of volatile elements in the gas phase (Schaefer and Fegley, 2017; Gaillard et al., 2021). Because volatile solubility in silicate melt strongly depends on gas speciation (see Lichtenberg et al., 2021, and references therein), it also affects the partitioning behaviour of volatile elements between a magma ocean and its atmosphere. A gas species having a lower solubility is preferentially out-gassed into the atmosphere. The composition of atmospheres formed above a magma ocean also depends on the pressure and temperature, relative abundance of volatile elements, as well as the depth of a magma ocean. Gaillard et al. (2022) modeled the partitioning of C-H-O-N-S elements between a deep magma ocean and its atmosphere at 1,500 °C using volatile inventories on

the present Earth. According to the scenario proposed by Deng et al. (2020) and the numerical results by Gaillard et al. (2022), large planets like Earth could have an oxidizing atmosphere consisting of CO_2 , SO_2 , and N_2 during its deep magma ocean stage (Figure 1). Although we have little information about the Venusian mantle, Venus could be expected to have had an oxidizing atmosphere by analogy, if Venus experienced metal-silicate equilibration at high pressures as well. Intermediate-sized planets like Mars might have a weakly reducing atmosphere consisting of CO , CO_2 , and N_2 . And if planets as small as the Moon formed with C-H-O-N-S whose relative abundance is as the same as that of the Earth, the shallow magma ocean might produce a highly reducing atmosphere consisting of CO and H_2 .

The oxidation mechanism involving the high stability of Fe^{3+} in silicate melt at high pressures seems a promising way to oxidize a large part of the silicate portion of planets rapidly. On the other hand, the Fe^{3+} abundance in the magma ocean estimated for each planet relies on the simplified assumption of perfect equilibration between silicate melt and metal at the base of the magma ocean. For Earth, the estimated $\text{Fe}^{3+}/\Sigma\text{Fe}$ ratios show significant differences among studies and do not necessarily coincide with the observations (Section 5.3). The cause for the differences between the predicted and observed $\text{Fe}^{3+}/\Sigma\text{Fe}$ ratios in the mantle requires further investigation.

The solubility of volatile elements also depends on the chemical composition of silicate melts and temperature, as well as cross-interaction among dissolved volatiles, and much remains to be determined especially for extreme conditions relevant to deep magma oceans (e.g. ultramafic compositions and high temperatures). Under more reducing conditions, the formation and flotation of graphite also would be an important factor on the C budget by controlling the oxygen fugacity and limiting the amount of outgassed C (Keppler and Golabek, 2019). On planets like Earth, another C-bearing phase, namely diamonds, possibly forms due to the reducing condition at greater depths (Armstrong et al., 2019). Keeping C as diamonds in the interior could cause the atmosphere and surface to be C-depleted (Dasgupta, 2013). The precipitation and stability of diamonds in a deep magma ocean remains to be investigated.

Note that the atmospheric compositions predicted for a deep magma ocean stage in Figure 1 would be different from those finally remaining at the end of the magma ocean stage. Later outgassing processes along with magma ocean crystallization can alter the mass and composition of atmospheres (see the next section). The redox state in a magma ocean can also vary due to

changes in $\text{Fe}^{3+}/\Sigma\text{Fe}$ as crystallization proceeds (e.g. Boujibar et al., 2016; Davis and Cottrell, 2021). According to recent modeling results by Maurice et al. (2023), the $\text{Fe}^{3+}/\Sigma\text{Fe}$ ratio and the oxygen fugacity in the crystallizing magma ocean can increase by several orders of magnitude throughout magma ocean crystallization. Although this would not change the total amount of Fe^{3+} in the mantle, the significant oxidation in the shallow part could cause a transition from a reducing atmosphere towards an oxidizing atmosphere during the solidification process.

2.3. Runaway threshold as a lower limit of thermal radiation from hot steam atmospheres

Following core formation, a magma ocean crystallizes by convective heat transport. At high degrees of melting, ultramafic melts are known to have a very low viscosity (Liebske et al., 2005; Posner et al., 2018). Rayleigh numbers of a planetary-scale magma ocean are extremely high, ranging from 10^{20} to 10^{30} (e.g. Solomatov, 2007), thus the convection in a deep magma ocean is supposed to be highly turbulent. At the early stage of solidification, volatile species close to the surface are expected to dissolve in the surface magma and to be transported downwards and homogenized by the vigorous convection. As solidification proceeds, dissolved volatiles would be released towards the surface, forming an overlying atmosphere. The outgassed atmosphere insulates the surface and slows the cooling of a magma ocean. Recently, such an interplay between a magma ocean and planetary atmosphere has been modeled and investigated numerically by several groups (e.g. Elkins-Tanton, 2008; Hamano et al., 2013; Hamano et al., 2015; Lebrun et al., 2013; Schaefer et al., 2016; Salvador et al., 2017; Katyal et al., 2019; Bower et al., 2019, 2022; Lichtenberg et al., 2021). According to theoretical studies, the effectiveness of thermal blanketing depends on the mass and composition of overlying atmospheres. In particular, if a steam-dominated atmosphere forms during magma ocean crystallization, terrestrial planets could experience different evolutionary paths despite the same initial state, where the path is being controlled by the orbital distance from the Sun (Hamano et al., 2013; Lebrun et al., 2013; Salvador et al., 2017).

Why is the formation of a steam-dominated atmosphere important for the possible divergent evolution of terrestrial planets? This is because it can limit the thermal radiation from hot molten planets to a certain value. Thermal radiation from steam atmospheres has been extensively explored especially regarding conditions of the formation of liquid water ocean (Komabayasi, 1967;




Figure 2: Thermal radiation from steam atmospheres. For these radiative transfer calculations, the atmospheric pressure (P_{atm}) is given by the saturation vapor pressure of water at surface temperature below the critical point of water, and set to 260 bar above it. As the surface temperature increases, the thermal radiation increases and then levels off at about 280 W/m² (radiation limit or runaway threshold), thus the runaway threshold is an upper limit in thermal radiation from water-saturated steam atmospheres. Above the critical point of water, the thermal radiation gradually starts to increase from the radiation limit with increasing surface temperature, owing to thermal emissions from the hot surface and the hot and dry (unsaturated) lower atmosphere. Under these hot surface conditions, the threshold flux corresponds to a lower limit in the thermal radiation from steam atmospheres. Modified from Hamano et al. (2015).

Ingersoll, 1969; Rasool and De Bergh, 1970; Pollack, 1971; Abe and Matsui, 1988; Kasting, 1988; Goldblatt et al., 2013; Kopparapu et al., 2013). Radiative transfer calculations show that moist (water-saturated) atmospheres have an upper limit for the outgoing infrared flux that can be emitted from its top. If any energy input exceeds this limit, the surface temperature increases until a new radiative equilibrium state is reached and all liquid water completely evaporates from the surface. This is called a runaway greenhouse state.

The physical mechanism behind the radiation limit was elucidated by

Nakajima et al. (1992) as follows. The thermal radiation from the top of the atmosphere F_{pl} is given by

$$F_{\text{pl}} = \int F_{\lambda} d\lambda \quad (4)$$

and

$$F_{\lambda} = \int_0^{\tau_{\lambda,s}} \pi B_{\lambda}(\tau') \exp[-\tau'] d\tau' + \pi B_{\lambda}(\tau_{\lambda,s}) \exp[-\tau_{\lambda,s}] \quad (5)$$

where τ is the effective optical depth at the top of the atmosphere, and F_{λ} and B_{λ} is the spectral irradiance and the Planck function emitted at wavelength λ . The subscript s represents the planetary surface. The first term at the right hand side of eq.(5) represents contributions of thermal emission from atmospheric layers and the second term is the emission from the planetary surface. The thermal radiation is most sensitive to the temperature around $\tau \sim 1$.

At surface temperatures lower than the critical point of water, the amount of H_2O is roughly governed by the saturation water vapor pressure. At low surface temperatures, the atmosphere is optically thin so the planetary radiation comes mainly from the surface. As the surface temperature increases, more water evaporates from the surface so the atmosphere becomes dominated by water vapor. The emission from the surface attenuates within the opaque atmosphere, while the infrared thermal radiation at the top of the atmosphere mainly comes from a moist convective region (troposphere), in which the optical depth becomes ~ 1 .

Assuming that the steam-rich atmosphere is saturated with water vapor, its thermal structure (a pressure-temperature relation) in the moist convective region approaches asymptotically the saturation vapor pressure curve of water vapor. The pressure level at which the optical depth is equal to unity also lies on the saturated vapor pressure curve. Consequently, the thermal radiation from the top of the atmosphere also approaches a certain value (radiation limit or runaway threshold) as the surface temperature increases (Figure 2).

If the absorbed solar flux exceeds the critical flux, the surface temperature increases until the outgoing thermal flux balances with the net solar insolation. Above the critical point of water, the thermal radiation starts to increase again as surface temperature increases (Figure 2). This is because the thermal emissions from the hot surface and the hot and water-unsaturated lower atmosphere escape through atmospheric windows, thus

increasing both the first and second terms at the right hand side of Eq.(5) (Goldblatt et al., 2013; Hamano et al., 2015). When the mass of the steam atmosphere increases, the thermal radiation starts to increase above the radiation limit, but never falls below the runaway threshold. In short, under hot surface conditions, like those during magma ocean crystallization, the radiation limit corresponds to a lower limit for the outgoing thermal radiation of steam atmospheres. It can be speculated that the presence of such a limit affects the energy budget of solidifying terrestrial planets, depending on whether the net incoming flux from the young Sun exceeds the lower limit or not. In a runaway greenhouse state, the surface temperature could exceed the solidus temperature of rocks (e.g. Kasting, 1988). A 30 bar steam atmosphere is sufficient for surface temperature to exceed the solidus temperature of peridotite rocks with the Earth’s gravity (e.g. Hamano, 2020).

2.4. Two distinctive cooling paths depending on the distance from the Sun

The effect of the radiation limit was first incorporated into a coupled model of a magma ocean and its overlying atmosphere by Hamano et al. (2013) and Lebrun et al. (2013). In both studies, the authors calculated the thermal evolution of molten terrestrial planets along with the formation of outgassed atmospheres. Hamano et al. (2013) considered a simple gray model of a pure H_2O atmosphere overlying a magma ocean and also incorporated loss of water associated with hydrodynamic escape of hydrogen in their model. In the Lebrun et al. (2013) model, the escape process of atmospheres was neglected, while radiative transfer of $\text{H}_2\text{O}\text{--CO}_2\text{(--N}_2\text{)}$ atmospheres was treated in a more realistic way with non-gray gas opacities.

Hamano et al. (2013) and Lebrun et al. (2013) assumed that volatiles, which are initially dissolved in a deep magma ocean are exsolved at the surface as solidification proceeds. They evaluated the thermal blanketing effect by an overlying atmosphere, whose mass and composition also evolve by volatile outgassing from the interior and atmospheric escape to space. The results for a pure steam atmosphere are shown in Figure 3. The evolutionary paths of solidifying Earth-sized planets formed at 1 au (Type I) and 0.7 au (Type II) are imposed on results of radiative transfer calculations. Based on the assumption of dissolution equilibrium between the magma ocean and the overlying atmosphere, both planets start out with an about 40 bar steam atmosphere with a total H_2O inventory of 5 Earth’s ocean masses. At a surface temperature of 3,000 K, the initial thermal radiation at the top of the atmosphere amounts to approximately $1,500 \text{ W/m}^2$.




Figure 3: Two types of evolutionary paths during magma ocean solidification. Both paths start at the same initial condition at 3,000 K. On a planet at 1 au (“Type I”), the atmosphere monotonically grows during cooling. In contrast, a planet at 0.7 au (“Type II”) loses its atmosphere after the planetary radiation reaches the net stellar radiation at the orbit. Calculation results from Hamano et al. (2013).

On the Type I planet, the absorbed solar flux is lower than the radiation limit. As the thermal radiation drops due to a decrease in surface temperature, the planet moves in this diagram from right to left until it reaches the threshold flux. The subsequent cooling heat flux is given by a flux difference between the radiation limit and the absorbed solar flux, and is high enough for the outgassing of water to overwhelm the escape. Therefore, the curve for the Type I planet bends upward along the radiation limit due to a net growth of the steam atmosphere. The resulting crystallization timescale is on the order of 1 Myr or less. Due to the short duration, most water remains at the end of the crystallization of the magma ocean (Figure 4). The remaining dense steam atmosphere condenses and forms an ocean immediately after the end of the magma ocean phase (Lebrun et al., 2013).

In contrast to the Type I planet, the solar radiation absorbed by the Type II planet exceeds the radiation limit due to the proximity to the young Sun. In this diagram, the Type II planet starts at the same position as the Type




Figure 4: Evolutionary dichotomy for terrestrial planets. Beyond the critical distance (see text for its definition), an Earth-sized planet solidifies on the order of 10^6 years or less, keeping its water (Type I). In contrast, inside the critical distance, a magma ocean could last longer for a larger inventory of water, and moreover, its solidification time is well approximated by the time required for total loss of the primordial water (Type II), indicated with thin dotted curves in the top panel. In short, a Type II planet is desiccated during the slow solidification process. Modified from Hamano et al. (2013)

I and moves from right to left as well. However, it reaches the absorbed solar flux, prior to the threshold flux. The Type II planets cannot move further to the left, otherwise the absorbed flux exceeds the outgoing thermal radiation, thus leading to an increase in the surface temperature and the thermal radiation. As all surface H_2O is present as water vapor that can be rapidly lost by photodissociation and followed by hydrogen escape to space. The net loss of the steam atmosphere allows the planet to lose its heat by making the atmosphere optically thin enough to emit the thermal radiation slightly larger than the absorbed solar flux. This is why the curve for the Type II planet sharply bends downward approximately along the value of the insolation. As a result, the Type II planet is desiccated and much less water is left on the surface when it completely solidifies (Figure 4). In the case that internal heat flux from a magma ocean is extremely low, surface temperature may also decrease due to structural changes in steam atmospheres, possibly leading to a shorter-lived magma ocean (Selsis et al., 2023). This possibility remains to be examined further, with a coupled magma-ocean-atmosphere model, which includes outgassing processes.

A similar dichotomy was also reported for $\text{CO}_2\text{-H}_2\text{O}$ atmospheres by Lebrun et al. (2013), Salvador et al. (2017) and Massol et al. (2023). A critical condition that separates these two evolutionary paths is given by the balance of absorbed solar flux and the critical threshold for a runaway greenhouse. This condition can be expressed as a critical orbital distance a_{cr} from the Sun² (Hamano et al., 2013),

$$a_{\text{cr}} \approx 0.77 \left(\frac{F_{\text{lim}}}{280 \text{ W/m}^2} \right)^{-1/2} \left(\frac{S}{0.7 S_{\text{sun}}} \right)^{1/2} \left(\frac{1 - \alpha_{\text{pl}}}{1 - 0.3} \right) [\text{au}] \quad (6)$$

where S and S_{sun} are the past and present solar constants, α_{pl} is the planetary albedo, and F_{lim} is the radiation limit, which depends on the planetary gravity. The threshold value obtained from non-gray cloud-free radiative transfer calculations is about 280 W/m^2 for the Earth's gravity and about 250 W/m^2 for the Mars' gravity (Goldblatt et al., 2013; Kopparapu et al., 2014). Therefore, the critical distance for Mars-sized planets is about 6 % larger than that for Earth-sized planets. According to the value of the critical distance (Eq. 6), Earth and Mars are categorized as Type I. If they had

²The expression for a_{cr} is re-scaled with the critical flux for Earth-sized planets obtained by Goldblatt et al. (2013) and Kopparapu et al. (2013) (about 280 W/m^2).

acquired abundant water during formation, both planets would have formed oceans soon after their solidification period on timescales of $\sim 10^6$ years or less. By contrast, it is not obvious which planetary type Venus belongs to. Three different scenarios that Venus might have followed are discussed in Section 5.2.

The critical distance is identical to the inner edge of the habitable zone, which is set by the onset of a runaway greenhouse state (the so-called runaway greenhouse limit). According to recent 3D simulations, the onset condition for a runaway greenhouse also depends on the initial state of the planetary atmosphere due to different cloud feedbacks (Way and Del Genio, 2020; Turbet et al., 2021). If Venus started from an extremely hot condition (magma ocean) like Earth, the Type II scenario seems more probable (Turbet et al., 2021). In our solar system a_{cr} lies in the middle of the formation region of the terrestrial planets. This is simply because the critical flux of steam atmospheres is comparable to the typical net insolation at these orbits. The concept of a runaway greenhouse itself is applicable to atmospheres dominated by other condensable gas species such as CO_2 and CH_4 . However, the critical flux estimated for these gas species is much smaller than that for water vapor (Pierrehumbert, 2010; McKay et al., 1999), making the location of the runaway greenhouse limit for other gas species irrelevant for the terrestrial planet region.

In reality, planets would acquire not only water vapor, but also other gas species during accretion. Many modeling studies have treated C-O-H atmospheres, especially oxidizing atmospheres consisting of $\text{CO}_2\text{--H}_2\text{O}$, in a more general case (Elkins-Tanton, 2008; Lebrun et al., 2013; Salvador et al., 2017; Bower et al., 2019, 2022). Sossi et al. (2020) found that early atmospheres are depleted in H-bearing components because most H is dissolved in the magma ocean as H_2O and the main component for the Earth’s earliest atmosphere could be CO. The similar feature of the H-depleted atmosphere and the hydrous magma ocean is seen in calculations of the C-H-O partitioning (Bower et al., 2022) and the C-H-O-N-S partitioning (Gaillard et al., 2022) between a deep magma ocean and its atmosphere in a wide range of surface oxygen fugacity. Gaillard et al. (2022) also found that, under highly reducing conditions with oxygen fugacity less than IW-4, most H is present in the atmosphere as H_2 and the atmosphere is C-depleted owing to the formation of graphite at the surface. At a fixed f_{O_2} , Bower et al. (2022) also calculated the change in the mass and composition of the atmospheres by outgassing processes. They showed that later outgassing of H_2O and CO_2 , as well as a

decrease in surface temperature, increases the amount of H-bearing species such as H_2O and H_2 , and the CO_2/CO ratio in the atmosphere.

The formation of H_2O -dominated atmospheres would depend on the outgassing efficiency at the late stage of solidification, as well as the volatile inventories and the evolution of the surface redox state. As long as the melt fraction in the surface magma is high enough for it to behave like a liquid, water dissolved in the magma ocean likely outgases fast enough to keep the system nearly at dissolution equilibrium (Pahlevan et al., 2019; Hamano et al., 2013). The efficient outgassing would occur by coalescence and ascent of bubbles in a low viscosity magma flow (Massol et al., 2016) or the diffusion of water molecules through a surface thermal boundary layer, which can be as thin as on the order of centimeters with a low liquid-like magma viscosity (Solomatov, 2007). On the other hand, when the melt fraction drops along with crystallization so that rheological transition from liquid-like to solid-like behavior occurs, the outgassing rate of water may decrease. At this stage, a highly viscous boundary layer would start to grow at the top of the magma ocean. Although melt percolation seems to transport volatiles efficiently upwards (e.g. Lebrun et al., 2013), its velocity decrease as the melt fraction decreases. Some of the volatiles including water might be trapped in the highly viscous crystal mush under the thick rigid surface layer. Bower et al. (2022) suggest that the outgassing efficiency at late stage of a magma ocean period would depend on the style of crystallization, i.e. fractional crystallization or equilibrium crystallization. The outgassing efficiency may also depend on the vigor of convection and the water abundance in the magma ocean (Salvador and Samuel, 2023). Even if the oxygen fugacity is extremely low at an early stage so that most of H is present as H_2 in the atmosphere, subsequent fractionation of Fe^{2+} and Fe^{3+} during the crystallization process may gradually oxidize the surface magma ocean, converting H_2 into H_2O , possibly leading to the transition to a steam-dominated atmosphere (Maurice et al., 2023).

3. Post-magma ocean internal evolution

The long-term evolution of a planet’s interior profoundly impacts surface conditions through the release of volatile species from the mantle into the atmosphere. Here, we examine how terrestrial planets in the Solar System could diverge during their long-term post-magma ocean internal evolution. Furthermore, we discuss the current knowledge regarding the differences in

interior evolution for Mars, Earth and Venus. The interior evolution of terrestrial planets is governed by heat transport in the mantle and its loss at the surface. Heat is generated by the decay of radioactive elements incorporated into the planet during formation, plus any residual accretion/differentiation heat. This determines the dynamics of the planet’s mantle and temperature field, but also partial melting of mantle material and magmatism/volcanism, mantle structure, topography, and even possible magnetic field generation (e.g. Schubert et al., 2001). Heat can be transported through convection, which is the more efficient mechanism, or through conduction.

3.1. Convection regimes

A convection (or tectonic) regime is the surface expression of interior dynamics. In general, various primary convection dynamic regimes are considered: mobile lid (with plate tectonics being a specific subset), stagnant lid, heat pipe, and plutonic-squishy lid (see Figure 5).

The mobile lid regime is characterized by its surface mobility, as the lithosphere is part of the convecting cell and surface velocities are equal to or greater than those of the underlying mantle. In the specific case of plate tectonics, the surface is broken into a small number of plates exhibiting differing relative motion, with deformation highly concentrated along the narrow boundaries between these plates. Plate tectonics is further characterized by long-term continuous subduction processes (e.g. Tackley, 2000; Stein et al., 2004; Rolf et al., 2022).

In a stagnant lid, a rigid lithosphere shows no lateral mobility and little deformation (Nataf and Richter, 1982; Christensen, 1984; Solomatov, 1995). The lid sits on top of a convective mantle; the hot mantle and cold lithosphere are decoupled. Stagnant lid naturally occurs when planetary lithospheres cannot be broken or mobilized by mantle convection, either because the lithosphere is too strong or the convective stresses are too low. Therefore, planets with a thick lithosphere or sluggish convection, for example, would fall into this regime (e.g. Sleep, 2000a; O’Neill et al., 2016; Stern et al., 2018; Rolf et al., 2022). As a result, heat flux and volcanic production are expected to be generally lower on stagnant lid than on mobile lid planets. For small or old planets, stagnant lid appears to be the final regime as geological activity dies down and the end-member toward which all terrestrial planets ultimately evolve as they cool down.

However, for larger, younger planets, the stagnant lid can insulate the mantle from the surface because heat transport by conduction through the lid is inefficient. Reducing the heat loss from the mantle leads to increasing interior temperatures that can cause substantial partial melting. A subset of the stagnant lid is a direct result of this scenario and is called the heat pipe regime, where heat is transported to the surface through erupting silicate melt (Moore and Webb, 2013; Moore et al., 2017). This subset of the stagnant lid regime has been suggested for young terrestrial planets and, at present-day, for Io, one of Jupiter’s moons (O’Reilly and Davies, 1981). In such cases, volcanic production and heat transport can be as high as in the plate tectonics regime.

Plutonic-squishy lid has been suggested recently as an alternative to the previous scenarios, where most of the melt produced by the planet is intruded, and remains within the lithosphere rather than reaching the surface. This leads to a warm ductile lithosphere that prevents plate tectonics (Rozel et al., 2017; Lourenço et al., 2018, 2020). However, the surface can be moderately mobile (more so than in stagnant lid but less than in mobile lid regime), but surface velocities are lower than the convective mantle’s. Deformation is more localized than in stagnant lid but less than in a plate tectonics, with generally small tectonic units. Heat is extracted from the mantle more efficiently than in the stagnant lid regime. Isolated subduction by lithospheric dripping is possible. Volcanic production would be relatively low due to the high fraction of intruded melt, but the surface heat flux could be close to that of the mobile lid regime (Lourenço et al., 2018).

The regimes discussed above are possible end-members for what is likely closer to a continuum of interior dynamics. Real-life is more complicated than these ideal configurations and spatial and temporal variations are expected for each given planet. It is also likely that terrestrial planets can switch from one regime to the other during their evolution (Sleep, 2000b). An episodic lid regime, for example, could correspond to a scenario with short occurrences of mobile lid separated by long periods of stagnant lid (e.g. Armann and Tackley, 2012) or the other way around (Lenardic and Crowley, 2012). Finally, it is likely that transition periods between regimes could last hundreds of millions to billions of years (e.g. Weller and Kiefer, 2020; Rolf et al., 2022).

The interior of present-day Earth is convecting, and heat is lost efficiently by plate tectonics. Evidence suggests that modern-day plate tectonics has




Figure 5: The three main end-member convection regimes: Plate tectonics, Stagnant Lid and Plutonic-squishy lid. Two additional specific cases are also highlighted: episodic lid, which is the transition between plate tectonics/mobile lid and the stagnant lid end member, and heat pipe, a stagnant lid with a large volcanic production (i.e., eruption). Schematic views of the interior dynamics of the three main regimes are indicated as cutaway illustrations of planetary interior structures, next to their respective domains. Regimes are distinguished by possibly observable characteristics of the planet. These axes should not be considered as the causes of the regimes depicted and are not fully independent from each other. Surface heat flux, the most readily available observable in the solar system, depends on many factors, such as size, mass, age and composition of the planet. The intrusion to extrusion ratio defines how much of the melt produced in the upper mantle reaches the surface. It is not a direct proxy of total volcanism and is likely to vary spatially. It could be a function of the surface temperature, which is a more direct observable, but can change over time. Surface yield stress is the convective stress caused by mantle convection required to break the lithosphere. It varies with the age of the planet and probably surface temperature. The presence of surface water and other fluids, and processes such as erosion, are also known to facilitate subduction and therefore lower the surface yield stress, which could introduce additional points of divergence between Mars, Venus and Earth. Lithospheric thickness could be considered as a first order observable for this parameter, although it is also influenced, to a lesser extent, by the intrusion to extrusion ratio. Convection regime figures are modified from Lourenço and Rozel (2023).

existed in its current form for at least 1 Gyr (Stern, 2005), with suggested onsets at least 3-4 Gyr ago (e.g. Dhuime et al., 2015; Cawood et al., 2006; Nutman et al., 2002). The clear topographic dichotomy between oceanic and continental crust, with the majority of the volcanic activity localized at the plate boundaries, is specific to plate tectonics. Earth’s tectonic regime is also characterized by the subduction of oceanic lithosphere that enables the large-scale recycling of surface material into the mantle. Earth’s tectonic processes and erosion combined are also responsible for the small number of impact craters to be found on Earth (see Figure 6, top panel).

This convection regime has not been observed anywhere else on present-day terrestrial planets. Mars is currently in a stagnant-lid regime. On Mars, the lack of tectonic activity and limited erosion due the thin atmosphere result in a heavily cratered, mostly old surface (see Figure 6, middle panel). It should be noted that stagnant lid does not mean geologically dead, since Mars appears to have been operating in the stagnant-lid regime for billions of years and still shows recent signs of activity (e.g. Hartmann, 2005; Hauber et al., 2011). Other planets, like Mercury, could also be in stagnant lid, but closer to being geologically dead bodies (see Grott et al., 2011).

The current convection regime active on Venus is uncertain: while Venus does not display global plate tectonics, the planet shows signs of deformation, volcanic and tectonic activity, and lateral variation (e.g. Basilevsky and Head, 2003; Rolf et al., 2022; Herrick et al., 2023; Hanmer, 2023), which would be out of place in a fully stagnant-lid regime, such as observed on Mars. As a result, the term “single plate planet” rather than “stagnant lid planet” has been used to describe Venus. It has been suggested that Venus could be an episodic lid planet, with brief overturns of the lithosphere into the mantle, separated by long stagnant periods (Turcotte, 1993; Armann and Tackley, 2012). The “plutonic-squishy lid” regime has also been proposed, and could possibly explain the relatively young surface age, deformation and heat flux measurements better than relying on a “well timed” overturn (Lourenço and Rozel, 2023; Rolf et al., 2022). However, no definitive evidence has been found yet to settle the question. Venus’ stronger tectonic activity compared to Mars can be also seen in the relatively small number of impact craters on its surface (see Figure 6, bottom panel) as erosion processes are negligible on this planet.

Evidence for Mars’ or Venus’ ancient convection regimes is even more difficult to identify. Mars’s surface is old and indicates that its most active phase occurred during the first billion year of its evolution (Carr, 2007),

however the available evidence does not shed much light on what the planet’s early dynamics looked like. Venus’ relatively young surface does not allow any direct insight into its distant past, as older surfaces have been covered by more recent basaltic material. Indirect evidence from the atmosphere of the planet indicate that large-scale outgassing may have been required to generate the current volatile inventory (Weller et al., 2022; Gillmann et al., 2022), which could indicate an active regime like a mobile lid in the past. Such an active phase may not be required when considering other sources of volatiles like impact delivery (Sakuraba et al., 2019; Gillmann et al., 2020) or could contradict ^{40}Ar measurements unless outgassing occurs very early (Kaula, 1999; Gillmann et al., 2022).

3.2. Volcanic production

While it is obvious that Earth is volcanically active at present-day, the same cannot be said about Mars and Venus. However, dating of lava flows on Mars indicates volcanic activity in specific areas in the last few hundred million years (Neukum et al., 2004; Werner, 2009; Carr and Head, 2010; Niles et al., 2010) and marsquakes in the Cerberus Fossa region indicate the presence of magma in the martian underground (Stähler et al., 2022). On Venus, transient IR anomalies localized close to identified lava flows and surface changes in volcanic regions have been interpreted as current volcanic activity (Smrekar et al., 2010; Herrick and Hensley, 2023). Global volcanic production rates are estimated to be around 20-30 km^3/yr for Earth and the extrusive volcanic production to be about 4 km^3/yr (Crisp, 1984; Morgan, 1998). In the case of Venus, it has been estimated to range from 0.3 to 10 km^3/yr (Fegley Jr and Prinn, 1989; Bullock and Grinspoon, 2001) to ≤ 2 km^3/yr (Head et al., 1991). Estimates for Mars indicate lower production rates around 0.3 km^3/yr (Greeley and Schneid, 1991).

Past volcanic production is even more difficult to constrain, but all three planets show signs of an active past. Mars boasts some of the largest volcanoes in the solar system and multiple volcanic provinces. Venus’ surface is mainly covered by basaltic plains, its mostly young surface indicates active resurfacing, and extremely long lava flows have been observed (Fegley Jr, 2014; Conrad and Nimmo, 2023).

It is likely that the three planets were more active during the first billion years as they lost accretion heat. Furthermore, heating by short-lived ^{26}Al might have played a role in the early evolution of the growing planetary embryos (Lichtenberg et al., 2019). Mars is considered to have completed




Figure 6: (top) Map of Earth impact craters larger than 5 km diameter together with their respective ages coded by color. Filled yellow circles indicate an uncertain age. (middle) Map of Mars impact craters larger than 20 km diameter. Smaller impact craters are not shown for visibility. (bottom) Map of Venus impact craters larger than 5 km diameter. Venus impact craters are thought to be relatively young (less than 1 Ga) but relative and absolute ages are uncertain. Earth topography credit: Copyright:©2013 National Geographic Society, i-cubed, published by Esri for use in ArcGIS. Earth crater database adapted from (Mazrouei et al., 2019) and (Spray, 2018). Mars MOLA topography map: WMS server used for JMARS client by Arizona State University for Jmars (Christensen et al., 2009), credit: ASU, NASA, Goddard. Mars crater database from Robbins and Hynes (2012). Venus map Credit: Credit: USGS/NASA/JPL Magellan Mission, http://planetarymaps.usgs.gov/cgi-bin/mapserv?map=/maps/venus/venus_simp_cyl.map. Venus crater database is from (Herrick et al., 1997), and available at <https://www.lpi.usra.edu/resources/vc/>.

accretion and core formation already within ~ 10 Myrs (Dauphas and Pour-

mand, 2011; Mezger et al., 2013) and magma ocean crystallisation occurred within ~ 20 Myrs after the start of the solar system (Bouvier et al., 2018). In comparison to Earth-like planets, smaller bodies such as Mars would have lost heat more rapidly during their early stages of evolution. This could explain differences in estimated volcanic production rates. However, once a stagnant lid is in place, the planet loses heat inefficiently, which may be why Mars’ core appears to still be mostly liquid (Irving et al., 2023). It could also cause the mantle to heat up and cause widespread partial melting (which is in turn countered by increasing thick lids; (O’Neill et al., 2014)). Differences between Venus and Earth may originate from other characteristics, since their sizes are similar.

3.3. Outgassing as evidence for planetary history

The volcanic history of a terrestrial planet is critical to the evolution of its atmosphere. Volcanic outgassing is one of the major sources of volatiles for the fluid envelopes of a rocky planet. In general, a higher production of volcanic material is expected to increase global volatile extraction from the mantle, but transport from the interior to the atmosphere is modulated by interior and surface conditions (see Gillmann et al., 2022). For example, the specific composition and mass of outgassed species depends on interior composition, redox state (Hirschmann and Withers, 2008; Ortenzi et al., 2020) and P - T conditions, both in the mantle (during melting) and at the surface (upon release, Gaillard and Scaillet (2014)).

As a rule, due to their size and heat budget, Venus and Earth are expected to produce more melt than Mars and possibly outgas more. Differences between Earth and Venus would be caused by past history and mantle dynamics differences: at present Venus is expected to experience less volcanism, but precise estimates are difficult to make. Based on their atmosphere composition all three planet would release CO_2 . Section 3.6 discusses the consequences for the volatile inventory. Mars could outgas CO_2 at low volatile concentrations due to its composition and reduced interior (Hirschmann and Withers, 2008). It is unknown what the composition of Venus’ outgassing is, but it is thought that water may not be released into the atmosphere because the planet’s mantle is dry and/or the surface pressure is high (Gaillard and Scaillet, 2014).

The concentration of radioactive Ar in the atmosphere has been suggested to be a proxy for the volcanic history of a terrestrial planet. As potassium decays into ^{40}Ar , argon partitions into the liquid phase during partial melting

and is outgassed into the atmosphere. Based on assumed initial compositions for the planet (i.e. K/U ratio), one can compare the relative outgassing of the planets. Measurements for ^{40}Ar suggest that Earth could be $\approx 50\%$ degassed, and Venus would have outgassed $\approx 25\%$ of its volatiles, making its atmosphere more primordial (less affected by outgassed volatiles, at least after the end of the magma ocean phase, Kaula, 1999). The uncertainties in initial compositions however, make this number poorly constrained (Gillmann et al., 2022). Additionally, the timing of the outgassing is important, as an early massive outgassing, when little ^{40}Ar has been produced, could possibly release large amounts of volatiles without being mirrored by ^{40}Ar measurements. Finally, especially in the case of Mars, losses of Argon through atmospheric escape could be difficult to quantify (i.e. Hutchins and Jakosky, 1996; Gillmann et al., 2011; Leblanc et al., 2012). However, models that track the evolution of outgassing and ^{40}Ar production under a continuous stagnant lid regime throughout the simulated history of Mars (O'Neill et al., 2007; Gillmann et al., 2011) produce volcanic fluxes that are consistent with present-day ^{40}Ar measurements (Pollack and Black, 1982). In the end, though qualitative, differences in the records of Earth and Venus point at a different style of outgassing, which could be explained if Earth operated mostly under mobile lid/plate tectonics (associated with larger outgassing), while Venus has had an episodic regime (O'Neill et al., 2014), a change in its regime (Gillmann and Tackley, 2014), or if its mobile-lid/plate-tectonics regime is limited to its early evolution (Arkani-Hamed, 1993; Gillmann and Tackley, 2014; Weller et al., 2022).

The generation (or not) of a magnetic field in the planetary core is also a clue pointing towards different evolutionary paths. Earth shows a self-sustained magnetic field with a strength similar to present-day since at least 3.2 Ga, possibly 3.45 Ga (Tarduno et al., 2014). Mars, however, sustains no dynamo at present-day but traces of ancient remanent magnetisation of its crust exist and date back to the first 600 Myr of its evolution (Steele et al., 2023). Finally, Venus shows no sign of a global self-sustained magnetic field at present-day (Phillips and Russell, 1987) and no remanent magnetisation has been observed so far. A magnetic dynamo is generated by convection in the liquid layer of the metallic core. It can be thermal (sufficient heat is lost at the core mantle boundary to drive core convection) and/or chemical (iron solidifies and sinks towards the solid inner core, thus stirring the liquid core), but it is always related to heat extraction by mantle convection and core cooling. The lack of a dynamo is thus indicative of either a completely

frozen core, a stably stratified core (Jacobson et al., 2017) or a too low heat flux out of the core (insufficient to cause core convection, thus no thermal dynamo).

The lack of a dynamo on Venus is especially puzzling if one assumes that Earth and Venus share a similar core composition and characteristics, as simulations would predict that slow rotating present-day Venus could generate a detectable magnetic field (i.e. O’Rourke et al., 2018). The current estimates for the tidal Love number on Venus are not precise enough to discriminate between a solid and liquid core (Dumoulin et al., 2017). It has been suggested that Venus did not experience giant collisions such as Earth’s moon-forming impact, and therefore retained a stratified core (Jacobson et al., 2017), thus explaining the lack of magnetic field. In that case, the slow retrograde rotation observed for Venus would be caused by tidal interaction (Dobrovolskis and Ingersoll, 1980; Correia and Laskar, 2003; Leconte et al., 2015), not impact despinning. However, it has also been proposed that impacts were the cause of the peculiar Venus rotation (i.e. McCord, 1968; Alemi and Stevenson, 2006; Raymond et al., 2013). If the core is unstratified, the lack of a present-day magnetic field would suggest near-complete solidification or that the thermal conductivity of Venus’ core material is higher than generally assumed (O’Rourke et al., 2018).

3.4. The role of impacts in interior evolution

Impacts have also been suggested to affect the interior dynamics of terrestrial planets. During a collision, the kinetic energy of the impactor can be transferred to the target planet’s mantle as thermal energy. If the impact is energetic enough and large enough (so that the heated volume extends beyond the lithosphere and into the mantle, typically for impactors above 100 km radius), then the mantle dynamics can be affected by the collision (Gillmann et al., 2016).

Such large impacts have been shown to cause temporary subduction events in numerical simulations of Venus and Earth (Gillmann et al., 2016; O’Neill et al., 2017; Borgeat and Tackley, 2022) and to increase lithosphere mobility. However, their consequences in simulations are relatively short-term (a few hundred million years at most) and any mobile lid regime they cause would have naturally emerged within the simulation parameters (Borgeat and Tackley, 2022).

Simulations of Mars’ largest recorded impacts have also failed to result in any lasting change for both climate or internal state (Turbet et al., 2020;

Ruedas and Breuer, 2019). Only early-on (≥ 4 Ga) impactors are likely to be large enough (above a few hundred kilometer radius, in order to significantly penetrate into the mantle) to have affected the interior and climate significantly on the long-term. Finally, it is unclear how different the impact flux would have been on the three terrestrial planets and what marker would allow us to distinguish between possible scenarios (e.g. Gillmann et al., 2020).

3.5. Modelling efforts and governing parameters

While numerical models are now capable of simulating Earth-like single sided plate tectonics, where only one of the plates sinks back into the mantle even when both are oceanic plates (i.e. Cramer and Tackley, 2014, 2015), the actual conditions for a planet to develop this regime are more difficult to assess. In essence, it requires the lithosphere to be brittle enough so it can break rather than flow in a ductile way, and not so strong that convective stress can never break it. In terms of modelling, this means that the yield strength should be low (a few tens of MPa - see for example Lourenço and Rozel, 2023). However, these values do not fit with laboratory measurements for Earth materials, which are reportedly much higher, potentially because these experiments also do not account for localized weak zones, faults and lateral material variations occurring on larger length scales.

Temperature is a major parameter that affects rheology. It has been shown that surface temperature variations, due to the evolution of the climate, could have direct consequences on the convection regime. For example, Lenardic et al. (2008) have shown that a surface temperature increase of about 100 K could shut down existing plate tectonics in an Earth-like model by reducing the viscosity of the convective mantle below the lithosphere. Because of the exponential dependence of the viscosity upon temperature, increasing temperature by 100 K reduces the viscosity by about one order of magnitude (Kohlstedt et al., 1995). In turn, lower viscosities would reduce convective stresses upon the lithosphere (Moresi and Solomatov, 1998). If convective stress drops below the yield strength of the lithosphere, plate tectonics stops (Moresi and Solomatov, 1998). Models of Venus (Gillmann and Tackley, 2014) indicate that water abundance variations in the atmosphere of Venus could change surface temperatures by several 10s to 100s K and cause a switch from a mobile to a episodic lid regime as partial pressure of water evolves towards its present-day value. Noack et al. (2012) highlight that particularly high surface temperatures (higher than at present-day) could reduce lithosphere viscosity enough to allow it to be dragged by the mantle

and become mobile. Such high temperatures (and possibly present-day Venus temperatures) could also lead to a relatively ductile lithosphere and prevent a plate-like behaviour by allowing faults to “heal” more easily (Bercovici and Ricard, 2014). It is still unclear whether any of those scenarios occurred during the evolution of Venus.

The temperature within the lithosphere can also be affected by intrusive volcanism, in turn possibly favored by high surface temperatures (Lourenço et al., 2020). The phenomenon can be local and generate weak zones or be more widespread and lead to a warm and soft lithosphere.

Water has also been suggested to play an important role in the rheology of the mantle and lithosphere of terrestrial planet due to its effect on the friction mechanisms and pore pressure (Korenaga, 2013). Through the dependence of viscosity on water, the water cycle affects mantle dynamics. The correlation between plate tectonics and liquid surface water in the solar system has motivated this line of thought. If water is needed for plate tectonics (i.e. Solomatov, 2004), a dry early state, as described before, would favor the single plate/episodic lid regime as suggested for present-day Venus. On the other hand, low water abundances may lead to a higher mantle viscosity (Xie and Tackley, 2004), which could, for example, improve the coupling between the mantle and lithosphere. Because the abundance of water in the mantle is also dependent on mantle dynamics, a feedback mechanism would exist between thermal evolution and the water cycle (Crowley et al., 2011).

In the end, despite important advances in our understanding of different regimes and mechanisms at work in planetary mantles, the lack of actual knowledge regarding the state and conditions inside solar system terrestrial planets prevents us from proposing a definitive explanation for the differences we observe. This is in part because we have little direct knowledge of the rheology and water content of both Mars and Venus. In recent years, partly in preparation for future exoplanet atmosphere characterization, efforts have been made to use atmosphere composition to investigate interior evolution, by using the consequences of outgassing integrated over the planets’ history e.g. (Gillmann et al., 2020; Krissansen-Totton et al., 2021a; Ortenzi et al., 2020; Weller et al., 2022; Warren and Kite, 2023).

3.6. Consequences on volatile inventory

The convection regime of all three terrestrial planets had important consequences for their whole evolution and surface conditions. On Earth, the composition of the atmosphere is the result of many processes, collectively

referred to as volatile cycles (see Dasgupta and Hirschmann, 2010). Over geological timescales, surface solid-gas reactions, the effect of life and the recycling flux at the subduction zones governs, for example, the abundance of CO_2 in the atmosphere, with complex and uncertain feedbacks (e.g. Friedlingstein et al., 2006), while on short timescales, ocean-atmosphere-biosphere interaction would dominate.

Comparatively, volatile exchanges are likely to be much simpler on Mars and Venus (e.g. Van Thienen et al., 2007). The long-term stagnant lid on Mars makes any recycling of gaseous species into the mantle unlikely. Same applies for Venus, at least since the time of the formation of the present-day surface and atmosphere. Additionally, at present-day Venus surface conditions, carbonates are not stable and, due to the lack of surface water, can not form (Fegley Jr, 2014). A limited form of volatile storage can act as a temporary sink even on stagnant lid planets. However, with no volatile cycle and no pathway back into the mantle, it can be easily destabilized. Höning et al. (2021) have described such an evolution and suggest that volatiles would be released back into the atmosphere during the first billion year as temperature increases either at the surface or at the depth where the material is buried.

In the absence of a stabilizing feedback, volatiles would accumulate in the atmosphere when not lost by escape processes, which would be the situation we observe for Mars and Venus relative to Earth (Bean et al., 2017; Turbet et al., 2019). The nature and amount of gases produced by partial melting and volcanism, then released into the atmosphere, are primarily dependent on the composition and redox state of the mantle, which both remain uncertain, even for Earth. The redox state of the mantle can be expressed using the oxygen fugacity measured as the ratio between the abundances of the different iron ions relative to a buffer, like the IW buffer (see also Section 2). Earth’s upper mantle is, despite some heterogeneities, rather oxidized at present-day, at approximately IW+1 to IW+5 (e.g. Ballhaus et al., 1990; Hirschmann, 2022), while Mars is more reduced, around IW+0 to IW+1 (Herd et al., 2002; Shearer et al., 2006; Karner et al., 2007; Hirschmann and Withers, 2008), and Venus remains unconstrained, possibly close to Earth (Schaefer and Fegley, 2017). As a result, outgassing on Mars may be characterized by lower CO_2 concentrations in the magma than on Earth and, possibly, Venus (Hirschmann and Withers, 2008).

However, it has also been suggested that surface conditions, such as surface pressure, may affect the composition of the outgassing into the atmosphere. For example, surface pressure affects the gas-gas and gas-melt equi-

libria, which modifies volatile speciation during release (i.e. Gaillard and Scaillet, 2014). As a result, high surface pressures (above ≈ 10 bar) may inhibit part of water outgassing into the atmosphere, increasing its solubility in the lava. Gaillard and Scaillet (2014) suggest $\approx 90\%$ of the initial water content of lava could be trapped. Instead, more Earth-like surface pressures would lead to water being the dominant outgassed species. On the other hand, CO_2 can outgas even at pressures of ≈ 100 bar. Venus-like surface pressures would prevent most H_2O from reaching the atmosphere under present-day Venus conditions, and possibly in the past, as long as surface pressures remain high (above a few bar). Earth may thus occupy a niche where the mantle is oxidized enough and the surface pressure low enough to allow for the outgassing of both CO_2 and water. Under Mars- or Venus-like conditions, either the mantle or at the surface, respectively, could stifle the release of one or both of those species (Gillmann et al., 2022).

Episodic volcanic outgassing is also often suggested for Venus. Based on the most recent resurfacing event that covered most of the surface with basalt (or the equivalent production under the equilibrium hypothesis; Phillips et al. (1992); see also Smrekar et al. (2018), Lopez et al. (1998) estimate that 9-10 such events would be required to build up the present-day CO_2 atmosphere, assuming an Earth-like lava composition. This corresponds to a total amount of extrusive lava of about $5 \times 10^{10} \text{ km}^3$ (based on possibly generous estimates from (Bullock and Grinspoon, 2001)), which would translate into a total volume of melt of $2.5\text{-}5 \times 10^{11} \text{ km}^3$. As a reference, the volume of Venus' mantle is about $7.8 \times 10^{11} \text{ km}^3$. Therefore, episodic production of the atmosphere requires massive outgassing, which could be in conflict with ^{40}Ar measurements, unless it occurred early or a large fraction of the atmosphere was already acquired during accretion or from other sources (impacts, destabilization of surface material).

Based on a detailed survey of the surface, Head et al. (2022) suggest that 80-100 (likely smaller scale) resurfacing events would be required and that the atmosphere of Venus is therefore ancient, which seems to be supported by ^{40}Ar measurements. Simulations that reproduce episodic evolution (Armann and Tackley, 2012; Gillmann and Tackley, 2014; Gillmann et al., 2020; Weller et al., 2022) require multiple billions of years to generate enough volcanic episodes to produce Venus' 90 bar CO_2 atmosphere, or fail to generate enough large events, due to the large amount of heat required for such large melting episodes - heat that is extracted from the mantle during the process.

Additionally, if CO_2 would outgas easily (provided the mantle is oxidized

enough), water may have been unable to reach the atmosphere for a large part of the Venus evolution due to high surface pressures (Gaillard and Scaillet, 2014; Gaillard et al., 2022). Way et al. (2022a) propose that such large-scale volcanism (possibly akin to Earth’s Large Igneous Provinces) may have contributed to the present harsh venusian climate. However, little is known about the ancient volcanic production and outgassing on Venus, and modelling based on existing constraints, like ^{40}Ar , noble gases, D/H ratio is an active and promising avenue of research (Noack et al., 2012; Gillmann et al., 2020; Krissansen-Totton et al., 2021a).

4. Atmospheric loss

Looking at the different atmosphere masses on Earth, Mars and Venus, signs of volatile losses can for instance be inferred from the fractionation of non-radiogenic gases and the deuterium to hydrogen ratio (D/H). Indeed, Mars has been observed to be more enriched in deuterium by a factor of 5 than Earth (Owen et al., 1988) and Venus by a factor of 150 (Donahue et al., 1997). The preferential loss of the lighter isotope compared to the heavier one affects the ratio between the two isotopes - the more a sample deviates from the primordial ratio, the stronger the loss. One should however remember that this is not true for all loss mechanisms (impact erosion of the atmosphere or intense hydrodynamic escape would not fractionate H, for example), and that quantifying that loss (Pahlevan et al., 2019; Kurokawa et al., 2018; Zahnle and Kasting, 2023) or how the ratio is affected by other processes is still uncertain (Grinspoon, 1993; Avicé et al., 2022).

Generally speaking, three main categories of escape are considered and discussed below: hydrodynamic escape (or thermal escape, as it is due to heating of the atmosphere), non-thermal escape (chemical reactions and collisions with particles from the solar wind) and impact erosion (caused by meteoritic collisions).

Atmospheric escape fluxes have been directly measured or calculated for the three planets. Additionally, Mars and Venus appear drier than the Earth, and water plays an important role in the long-term terrestrial planet evolution. Therefore, it is no surprise that atmospheric loss processes have been proposed as a possible cause of the diverging evolutionary pathways for terrestrial planets in the Solar System.

Noble gases in the Venus atmosphere have been analyzed by mass spectrometers and gas chromatographs on board the Pioneer Venus and Venera

probes. Although a similar pattern is observed for the three terrestrial planets, the $^{20}\text{Ne}/^{36}\text{Ar}$ ratio is depleted by a factor of ≈ 3 in the Venus atmosphere relative to Earth's. Ne and Ar are more abundant in Venus' atmosphere than in Earth's atmosphere, by at least factors of 10 and 50, respectively. ^{20}Ne and ^{36}Ar abundances are close to chondritic for Venus, but depleted for Mars and Earth, as are ^{84}Kr and ^{130}Xe for those two planets. In addition, $^{130}\text{Xe}/^{84}\text{Kr}$ is measured at one order of magnitude below the chondritic ratio, indicating further loss (Pepin, 1991). In the case of Venus, however, Kr and Xe are still poorly constrained, thus so far it is not possible to assess whether or not Venus follows the Earth-like pattern (Lammer et al., 2018). With the caveat of Venus Xe measurements, all three planets are depleted in noble gases compared to the solar composition. In general though, Venus' noble gas elemental spectrum is suggested to be more solar-like than those of Earth and Mars.

On Mars, the isotopic components of atmospheric Kr are close to those of the solar wind and remained mainly unfractionated (e.g. Pepin, 1991; Becker et al., 2003; Conrad et al., 2016). Xe isotopes in Earth's and Mars' atmosphere are enriched in heavier isotopes. The case of Xe is still actively debated, with the fractionation possibly attributed to long-term (4.5-2 Ga; Avice and Marty (2020)) escape of ionized hydrogen dragging of Xe (Zahnle et al., 2019).

Neon is less fractionated (Wieler, 2002) in Venus atmosphere ($^{20}\text{Ne}/^{22}\text{Ne} = 11.9 \pm 0.7$) than in Earth's atmosphere ($^{20}\text{Ne}/^{22}\text{Ne} = 9.8$) with respect to solar composition ($^{20}\text{Ne}/^{22}\text{Ne} = 13.5$). Because Ne is only slightly heavier than O, hydrodynamic escape of Ne implies hydrodynamic escape of oxygen, and Ne may be used as a tracer of oxygen escape at primitive stages (e.g. Gillmann et al., 2009a). There is also no evidence of changes in Ne, Ar and Kr isotopic compositions (non-radiogenic) during the past 3.5 Ga on Earth (Avice et al., 2017). On all three planets it has generally been suggested that their lighter noble gases were fractionated during an early hydrodynamic escape phase (the first few hundred million years).

4.1. *Hydrodynamic escape*

Hydrodynamic (or thermal) escape occurs due to heating of the atmosphere by intense EUV radiation from the Sun or by radiation from a primitive hot rocky planet (Ginzburg et al., 2018, "core-powered"). It causes atoms in the upper atmosphere to exceed the escape velocity, leading to very efficient escape of the bulk gas of the upper atmosphere where the atmosphere

remains a collisional fluid as it passes through the transonic point (Chamberlain, 1963; Chassefière, 1996; Fossati et al., 2017). Light species (especially H) are also the most efficient at escaping via this process, as they require less energy to reach bulk escape velocity.

Heavier species (such as helium, carbon or oxygen atoms) can undergo escape via so-called hydrodynamic drag where they are carried along with the stream of lighter hydrogen atoms (e.g. Hunten, 1973). Due to solar evolution (the decrease in EUV flux with time (Claire et al., 2012), this implies the most favourable conditions for such a process occur during the first few hundred million years. At lower energy levels, a transition occurs from hydrodynamic escape to the much less efficient Jeans escape, causing the loss of atoms populating the high energy tail of a Maxwell distribution (Jeans, 1925; Chamberlain, 1963), and still occurring at present-day on Earth, Mars and Venus.

Hydrodynamic escape would have efficiently removed the primordial H/He atmosphere that all three planets obtained from the solar nebula during their formation (Lammer et al., 2018, 2020). Earth’s D/H seems to be incompatible with oceans originating from the solar nebula (Marty, 2012). Then, during the magma ocean phase, the ongoing thermal loss would have affected the water vapor/CO₂ atmosphere of Earth, Venus and Mars to various degrees, whether it came from interaction of the atmosphere with the magma ocean or from catastrophic outgassing during magma ocean solidification. Finally, if a planet, such as Venus, entered a runaway greenhouse state, then hydrodynamic escape would also have been a dominant factor for water loss. The loss of water would have remained efficient but the presence of more abundant heavier species would have led to diminished loss rates. Still, modelling efforts have shown that a Mars-like body orbiting at Earth-like distance from the Sun would lose its atmosphere (85 bar H₂O and 11 bar CO₂) in about 15 Myr (Odert et al., 2018), and that Venus would only take 50 Myr to lose about 2 Earth ocean equivalents (Gillmann et al., 2009b). Gravity, the composition of the atmosphere and the proximity to the star all affect the loss rates, with smaller or closer planets being more affected. Thus, Mars could have lost a large part of its initial water inventory and even CO₂ (Tian et al., 2009; Odert et al., 2018). Mars being drier than Earth due to early escape (≥ 4 Ga) is consistent with martian meteorites measurements (Cassata, 2017).

The planets are efficiently dried out, as long as a supply of water is available in the atmosphere. Kasting and Pollack (1983) calculated that hydrodynamic escape was the dominant process as long as the water mass

mixing ratio of water in the lower atmosphere exceeded 0.1, even during the later evolution.

In short, factors that favor a strong hydrodynamic escape are (in order of importance): (i) an H-rich atmosphere (water vapour for instance), (ii) a strong EUV flux (for example during the first few hundred million years of the evolution) and (iii) a small planet with low gravity (see also Catling and Zahnle, 2009). Because of (i) and (ii) thermal escape is likely the dominant process during the first few hundred million years of planetary evolution. The escape of lighter species (H) is favored, but a strong enough flux can cause the loss of heavier components. Conversely a CO₂ atmosphere, low EUV flux and stronger gravity would reduce escape rates, while magnetic fields do not impede hydrodynamic escape.

4.2. Non-thermal escape

Non-thermal escape covers a few mechanisms that involve the interaction of the upper atmosphere molecules (above the exobase) and ions with energy and particles brought by the solar flux, such as: charge exchange, photochemical dissociation, sputtering, polar wind, plasma clouds. In each case, the interaction leads specific molecules to reach high enough velocities to escape (Catling and Zahnle, 2009).

Over the last decades, a large number of studies have attempted to quantify the escape rates of heavy ions (such as O⁺, O₂⁺ and CO₂⁺) on Mars, Earth and Venus based on observations. (Ramstad and Barabash, 2021) have summarized these results (their figure 5) and found that estimates for Earth, Mars and Venus tend to be roughly similar, despite the large variations in planetary size, atmosphere composition, presence (or absence) of a global magnetic field. Mars' heavy ion escape rates range between $2 \times 10^{23} \text{ s}^{-1}$ and 10^{25} s^{-1} ; for Venus reported rates cover $1\text{-}6 \times 10^{24} \text{ s}^{-1}$ (with an outlier one order of magnitude above), while Earth's escape fluxes are higher: 7×10^{24} to 10^{26} s^{-1} . For O⁺ alone, rates ranges from $2\text{-}4 \times 10^{24} \text{ s}^{-1}$ for Mars, to $1\text{-}4 \times 10^{24} \text{ s}^{-1}$ for Venus and $5 \times 10^{24} \text{ s}^{-1}$ to $5 \times 10^{25} \text{ s}^{-1}$ for Earth (see Ramstad and Barabash, 2021, and references therein).

These reported values also cast doubt on the previously accepted notion that magnetic fields provide an efficient shield against atmospheric escape, which in turn was used to explain why Earth was so different compared to Mars and Venus, and retained its water. To explain the similar measured escape rates reported above, it has been suggested that Earth's magnetic field extends much further into space than its atmosphere (a few Earth radii) and

therefore intercepts more energy from the Sun than both Mars or Venus (Brain et al., 2012), leading to increased atmospheric escape.

It has also been suggested that magnetic fields could actually enhance atmospheric escape (e.g. Gunell et al., 2018). This idea has been disputed by Tarduno et al. (2014), for example for underestimating the return fluxes (ionized molecules escaping following magnetic lines that reconnect to the planet rather than open lines in the magnetotail). The current state of the research indicates that the magnetic field affects escape mechanisms (a magnetized planet will lose molecules at the cusps of the field rather than in the tail of the induced field) and affect which mechanism can be dominant. However, so far we are not able to explain exactly what the resulting escape flux will be (Gronoff et al., 2020). As a consequence, it remains uncertain by how much the presence of a past magnetic field could have affected the respective evolution of Mars, Venus and Earth, although as discussed above there may be hints that the presence of a magnetic field does not result in a radically different volatile loss history.

4.3. Past escape reconstruction

Attempts have been made to reconstruct past escape rates (Hunten et al., 1989) based on the present-day measurements for Mars (Chassefière and Leblanc, 2004; Gillmann et al., 2009b), with the latest contributions being the MAVEN results (Jakosky, 2021), and Venus (Gillmann and Tackley, 2014; Persson et al., 2020). These are obtained by extrapolating the present-day escape rates obtained at high and low solar EUV fluxes during a solar cycle. However, they ignore changes in atmosphere composition and structure, or exosphere temperature and remain dependent on other factors, such as the existence of a magnetic field (Gronoff et al., 2020; Way et al., 2022b). Depending on the specific conditions at a given time, escape rates caused by a given mechanism are likely to vary and the dominant escape process would certainly change: water vapour atmospheres would be defined by strong H loss from hydrodynamic escape, while dry atmospheres would see non-thermal processes gain importance. Generally speaking, the two types of atmospheres are often used to describe, respectively, (1) the primitive phase of planetary evolution during magma evolution or runaway greenhouse, and (2) the last ≈ 4 Gyr and present-day.

For example, CO₂ in the thermosphere radiates energy back to space, leading to infrared cooling of the upper atmosphere and affecting the structure of the atmosphere (Roble and Dickinson, 1989). This is why Venus'

exobase is both cooler and occurs at a lower altitude than Earth's. A hot, expanded atmosphere, resulting from a switch from a CO₂-rich atmosphere to a N₂ atmosphere for example, could possibly lead to escape rates orders of magnitude higher (Lammer et al., 2006; Way et al., 2022b, even though estimates are very uncertain;). Past reconstructed escape rates should thus be considered with caution, only as a first order approximation. They are currently, however, our best shot at ancient loss rates, since we are not yet able to predict a total escape rate from a given set of conditions.

Non-thermal escape is modulated by (i) a planet's gravity, with weaker gravity leaving planets more vulnerable to loss. Non-thermal escape is also a function of (ii) the composition of the atmosphere, since most mechanisms depend on chemistry. However the precise effects of different atmosphere composition on loss rates is still uncertain. Finally, as a large part of the escape involves ions, (iii) magnetic fields existence and configuration will affect non-thermal escape. Non-thermal escape is dominant during late evolution and for species such as O (all planets) or C, N, Ar (for Mars).

4.4. Loss of water and the fate of remnant oxygen

From observation, for non-thermal processes, the ratio of the escape of H and O appears to be close to or below the stoichiometry of water (between 1.1 and 2.6 Persson et al., 2018, for instance, in the case of Venus). Such a ratio leads to the full escape of water or a moderate loss of O relative to the source water. On the other hand, thermal escape is expected to lead to a relative accumulation of oxygen in the atmosphere due to the strong differences in loss efficiencies for species of different masses. Oxygen, being heavier, is lost at a much reduced rate compared to H. The ratio of the loss of H to O would thus be much higher than that of their respective abundance in the atmosphere (2 H for 1 O, assuming they come from the photodissociation of water; Kasting and Pollack, 1983; Gillmann et al., 2009a). As a result the atmosphere becomes more oxidized unless the excess O can be removed through other processes such as oxidizing reducing gas species and atmosphere-surface reactions. The fate of the remnant O has been considered to be a constraint for the evolution of the Venusian atmosphere, and is discussed in detail in Section 5.2. On Earth, reconstructed atmosphere compositions dating back from before the Great Oxygenation Event (Lyons et al., 2014; Catling and Zahnle, 2020) indicate that a long-lived O₂-rich atmosphere resulting from the process mentioned above is unlikely. Banded iron formations (Canfield, 2005, and references therein), for example are on

Earth indicators of oxidation of iron in oceans under an atmosphere with low O_2 concentrations and low seawater sulfates (and provide a link to the consequences of life for planetary evolution, see below, Cloud Jr, 1968).

As mentioned above (see Section 4), signs for long-term water loss are evidenced by the D/H measurements on Earth, Mars and Venus. Alsaed and Jakosky (2019), for example, reconstruct the evolution of water reservoirs based on atmospheric escape on Mars and suggest that at 3.3 Ga, a global equivalent layer of 40-120 m of water could have existed to explain the present-day D/H ratio. Further, Mars' atmosphere is enriched in ^{15}N over ^{14}N , which has been suggested to result from non-thermal loss processes (Shizgal and Arkos, 1996; Gillmann et al., 2011). On the other hand, non-thermal escape on Mars is unable to account for the loss of a water equivalent above a few 100 mbar over 3-4 Gyr (Gillmann et al., 2009b, 2011). At the same time, the cumulated water escape on Venus over the last 3.8 Gyr was recently estimated to amount to a few centimeters to tens of centimeters only (Persson et al., 2020).

It seems unlikely that non-thermal escape could, by itself, cause the divergences observed in the solar system. However, it can critically influence already thin atmospheres of low mass terrestrial planets like Mars'. Additionally it remains a significant evolutionary process that shaped the observed characteristics of thicker atmospheres like Earth's and Venus'.

4.5. Atmosphere erosion by impacts

Impact-related erosion of the atmosphere can lead to volatile loss (Cameron, 1983; Melosh and Vickery, 1989). Four main mechanisms can cause escape (e.g. Dehant et al., 2019): (i) direct ejection of the atmosphere by the shock wave, (ii) the rising of the vaporized projectile and sediments in a vapor plume, (iii) heating of the atmosphere by ejected particles and (iv) in the case of very large collisions, ground motion of the planet can accelerate particles of the atmosphere above escape velocities. That last process is deemed to be the most efficient one. Genda and Abe (2003) estimate that such an impact ($R_i \sim 1000$ km) could remove about 20% of a preexisting atmosphere and the presence of a water ocean could even enhance the atmosphere loss (Genda and Abe, 2005). More recent work (Schlichting et al., 2015), derives estimates depending on relative mass of the impactor to the planet and its velocity. Significant loss (above 10% of a preexisting atmosphere) requires the impactor to be traveling at several times the planet's escape velocity and/or

to be of a comparable mass to the target, which is more easily achieved for Mars than for larger planets.

A key feature of impact erosion is that it generates bulk ejection of the atmosphere and does not isotopically fractionate individual elements in the planetary atmosphere, unlike other processes described above. It could, however fractionate the volatile budget of two elements, in case those elements are not distributed equally within the planet, for example if one resides preferentially within the atmosphere while the other one is trapped in the interior (e.g. Massol et al., 2016). Impact erosion of the atmosphere is probably the main mean of removing Kr from the atmospheres of terrestrial planets, since Kr (generally non-fractionated) is too heavy and difficult to ionize to be significantly lost through the previously discussed thermal and non-thermal processes (Pepin, 1994; Catling and Zahnle, 2009; Cassata et al., 2022).

Smaller impactors ($R_i \leq 1000$ km) individually remove less atmospheric volatiles through processes (i)-(iii) than larger ones, however, they are more efficient at atmospheric erosion per unit of mass. Schlichting and Mukhopadhyay (2018) highlight that the much more numerous population of smaller impactors ($1 \text{ km} \leq R_i \leq 20 \text{ km}$), as a whole, can remove more volatiles than the few larger collisions ($25 \text{ km} \leq R_i \leq 1000 \text{ km}$). They argue that such a flux of small impactors would be able to erode an Earth-like atmosphere within a few billion years. While this can affect decisively thin atmospheres (such as Mars'), this escape rate makes little difference, by itself, on a planet with a multi-bar atmosphere such as present-day Venus.

In short, volatile loss is very dependent on the planetary mass and impact velocity: less massive planets such as Mars are more vulnerable to erosion than Earth or Venus (Pham et al., 2011) and faster impactors can drive a greater loss. Furthermore, massive atmospheres would be only marginally affected by erosion, relative to their total masses, in contrast to planets with thinner atmospheres. Finally, impacts can also act as sources of volatiles depending on their exact composition (Gillmann et al., 2020, for example) and affect atmosphere chemistry (Schaefer and Fegley, 2017). While erosion, from a relatively predictable population of impactors, averages out, volatile deposition by collisions is much more susceptible to stochastic events (such as a single large volatile-rich impact), which may cause drastic changes in atmosphere bulk composition and chemistry.

5. Unique features of the history of Mars, Venus and Earth

5.1. *Mars*

As Mars is less massive than Earth, magma ocean pressures on Mars would have been lower, compared to the Earth’s magma ocean pressures. Geochemical studies on martian meteorites support core formation pressure of about 14 GPa to explain the depletion in siderophile and chalcophile elements in the martian mantle (e.g. Righter and Chabot, 2011). Therefore, the iron disproportionation would have had a limited effect on Fe^{3+} abundance and a $f\text{O}_2$ gradient in the martian magma ocean. The estimated surface $f\text{O}_2$ above the magma ocean is about $\text{IW}-1 \sim \text{IW}+1$ (Deng et al., 2020). This $f\text{O}_2$ is high enough that a large fraction of H would be dissolved in the magma ocean as H_2O (Gaillard et al., 2022). As the orbit of Mars is far enough to escape from the runaway greenhouse state, Mars is expected to have crystallized rapidly, though the predicted duration of magma ocean crystallization shows a large discrepancy between thermal models ($< \sim 1$ Myr, Elkins-Tanton, 2008; Lebrun et al., 2013) and geochemical constraints (several to tens Myr after core formation, Borg et al., 2016; Kruijer et al., 2017, 2020). Several processes such as late differentiation caused by cumulate overturn (e.g. Elkins-Tanton et al., 2005) and heating by late accretion (e.g. Marchi et al., 2018) have been proposed as potential causes of the prolonged Martian magma ocean. Depending on its outgassing efficiency, its water would remain at the surface or in its interior (Odert et al., 2018).

Mars’ ancient surface shows a dichotomy between the southern highlands and the northern lowlands. The dichotomy is considered to be the oldest geological feature on Mars and was created within the first 400-500 Ma after the planet’s formation (Nimmo and Tanaka, 2005; Carr and Head, 2010). Different processes have been suggested as the origin of this topographic feature like (i) impact excavation of the northern hemisphere (Wilhelms and Squyres, 1984; Frey and Schultz, 1988; Andrews-Hanna et al., 2008; Marinova et al., 2008; Nimmo et al., 2008), (ii) low degree convection (Zhong and Zuber, 2001; Roberts and Zhong, 2006; Zhong, 2009; Keller and Tackley, 2009; Šrámek and Zhong, 2010) and (iii) an impact in the southern hemisphere forming a so-called impact megadome (Reese and Solomatov, 2006, 2010; Reese et al., 2010; Golabek et al., 2011; Ballantyne et al., 2023; Cheng et al., 2024). Also it has been suggested that Mars could have developed short-term plate tectonics during its first few hundred million years (Frey, 2006; Sautter et al., 2015).

In short, due to its relatively small mass, and possibly limited water inventory, Mars would have transitioned from a possibly active first billion year into a stagnant lid regime. At about the same time, it lost a large part of its initial water inventory and its magnetic field. Details regarding the climate on ancient Mars beyond increasing desiccation remain debated, and both "warm and wet" and "cold and wet/snowy" origins have been proposed (Wordsworth et al., 2015), as well as episodic warming by volcanism and reducing greenhouse gases (Halevy and Head III, 2014).

Observations of the martian surface give an idea of the consequences that hydrodynamic escape could have as Mars' terrain age allows studying its distant past. Fluvial features indicate a decline in water's role in shaping the surface of the planet, evolving from valley networks in the Noachian to episodic outflow channels in the Hesperian towards the more arid planet since the Amazonian (Carr, 2007). Based on mineral alteration in the presence of diminishing amounts of water the subsurface mineralogy analysis suggests a similar evolution from Phyllosian to Theikian to Siderikian (Bibring et al., 2006).

Long-term evolution models suggest outgassing would be relatively limited, possibly due to interior redox conditions inherited from its primitive history (Hirschmann and Withers, 2008; Deng et al., 2020). A large portion of the outgassing would have occurred during the first billion years of its evolution (Lammer et al., 2018), before most of its interior heat was lost. However, even such low volatile release rates could be sufficient to affect surface conditions and the composition of the atmosphere. Based on the estimated lava composition and simulated volcanic production over the evolution of Mars, outgassing histories have been modelled (i.e. Gillmann et al., 2009b, 2011; Grott et al., 2011; Leblanc et al., 2012; Morschhauser et al., 2011; Sandu and Kiefer, 2012; Hu et al., 2015).

It is likely that water loss on Mars was stronger early in the evolution of the planet, both for hydrodynamic and non-thermal escape (Lammer et al., 2018), and that for most of the evolution, escape was limited to a fraction of a bar. Some simulations indicate that ancient surface pressures on Mars, around 3.5-4 Ga, may have been low (a few to a few hundred millibar) and that much of present-day CO₂ atmosphere could originate from late outgassing (Gillmann et al., 2009b, 2011; Leblanc et al., 2012; Lammer et al., 2013). Volcanic outgassing could also account for most of present-day water surface inventory (up to a 100 m global equivalent layer, Alsaeed and Jakosky (2019)). This is in line with work by Kite et al. (2014) suggest-

ing an upper limit of ~ 1 bar atmospheric pressure ~ 3.6 Ga ago. This thin atmosphere was more prone to atmosphere loss than Earth's and Venus', accordingly over time favourable conditions for the presence of liquid water at the martian surface ceased.

5.2. Venus

5.2.1. Potential evolution paths

Compared to Earth and Mars, there is little information on the history of Venus. This is, in a large part, due to the overall young apparent age of most of the planetary surface, which limits observations regarding most of its evolution. No definitive evidence for surface water exists in the case of Venus, due to the lack of surface features dating back more than 300-1000 Ma. Surface mineralogy of tesserae (8% of the present-day venusian crust) has been proposed to indicate the presence of felsic material in unknown quantities (Shellnutt, 2019). This would suggest that an uncertain amount of liquid water could have existed on Venus for an unknown period of time. Most of the constraints on Venus' past are derived from its atmosphere that retains the accumulated effects of every process affecting volatile inventories. Needless to say, unraveling plausible scenarios is a complicated endeavour. Current research has not yet managed to converge toward a single agreed succession of events that would lead to present-day Venus.

If Venus and Earth formed in the same way from the same building blocks, the apparent dryness of the Venus could have been caused by its proximity to the Sun. Venus receives about 1.9 times more solar flux than Earth, and so it can relatively easily enter a runaway greenhouse state (Zahnle et al., 2007). However, while the atmosphere and surface of Venus are dry compared to Earth, the presence of water inside Venus has not been ruled out. Three different scenarios have been proposed to explain the lack of water at the surface of Venus (Kasting, 1988; Hamano et al., 2013; Gillmann et al., 2022): (i) Venus was once an ocean planet like Earth, and then, its climate was destabilized by internal factors, and/or, as the Sun got brighter over time, the surface water was evaporated and lost by photolysis and escape of hydrogen. (ii) Venus evolved into a desiccated planet as it is today during its magma ocean crystallization process. (iii) Water was never outgassed from the interior of the planet and was trapped in its interior.

One-dimensional cloud-free climate calculations yield the radiation limit of about 280 W/m^2 considering a planetary albedo of about 0.2 for thick steam-dominated atmospheres (Kopparapu et al., 2013; Goldblatt et al.,

2013). Simply substituting these values into equation (6) gives the critical orbital distance of 0.82 au, seemingly suggesting that Venus was well inside the critical distance and thus followed scenario (ii). However, planetary albedo plays a critical role in defining the radiative balance of rocky planets. 1D climate models cannot fully handle the effects of atmospheric circulation, such as heat transport by winds and the distribution of relative humidity and clouds, which may favor the formation of an ocean. Recent 3D simulations have shown that the stability of water oceans on early Venus would depend on starting conditions in its atmosphere. If all water on early Venus was initially present as a liquid ocean in a moderate climate, clouds formed on the dayside would have reflected sunlight efficiently and maintained the surface water ocean considering a slow rotation rate of Venus (Yang et al., 2013; Way et al., 2016). By contrast, if early Venus started out with a hot steam atmosphere, it would have remained in a runaway greenhouse state due to thermal blanketing by clouds formed on the nightside and strong absorption of sunlight by abundant water vapor (Turbet et al., 2021). If Venus experienced a giant impact at the end of its accretion, like Earth, the latter case seems more plausible. On the other hand, there is a possibility that Venus had a different initial thermal state when it formed in a much different way from Earth (e.g. Jacobson et al., 2017). This topic is out of scope of this chapter.

In scenarios (i) and (ii), water vapor is considered to be dissociated and lost by extensive escape of hydrogen to space. If Venus started with a water equivalent of one Earth’ ocean mass, a preferential escape of hydrogen might have left about 240 bars of oxygen on Venus. Some fraction of oxygen may also have escaped directly if the energy input is high enough (possible roughly before 4 Ga). However, based on recent numerical results (Johnstone, 2020; Johnstone et al., 2021; Saxena et al., 2019), the activity of the young Sun was not strong enough to drive rapid oxygen escape, thus some oxygen would have remained in the Venusian atmosphere.

The fate of the remnant oxygen has been considered to be a constraint for the recent evolution of the Venusian atmosphere. If oxygen builds up, the mixing ratio of water vapor drops so that oxygen could become less prone to escape (Kasting and Pollack, 1983; Wordsworth and Pierrehumbert, 2014). It had long been suggested that a mechanism that could explain the loss of oxygen from the atmosphere of Venus (and other terrestrial planets) was long-term ongoing loss of volatiles by non-thermal escape processes (see for example Gronoff et al. (2020)). However, total loss of O through non-thermal

escape processes is estimated to be far less than those needed to explain the loss of an amount of water comparable to one Earth ocean (Gillmann et al., 2020; Persson et al., 2020). Rapid water loss requires additional oxygen sink(s) other than thermal and non-thermal escape processes, which can efficiently dispose of the remaining oxygen.

In scenario (i), some of the oxygen may have reacted with, for example, CO to form CO₂ (or reduced species outgassed at the time) and some of the oxygen may have been consumed by oxidation of Venusian crustal material (experiments by Berger et al. (2019) and Filiberto et al. (2020), models by Gillmann et al. (2020); Warren and Kite (2023), and review by Gillmann et al. (2022)). However, it is unclear whether a combination of all these processes could have removed tens to hundreds of bars of O₂ immediately without preventing hydrogen from escaping. Generally speaking, current results suggest that only a moderate amount of oxygen can be extracted from the atmosphere by solid-gas reactions. Experiments and models currently suggest the total removal of O₂ may be limited to amounts similar to those caused by non-thermal escape (about a fraction of a bar to 1 bar at most). This does not outright make scenario (i) implausible but rather limits the possible range of water inventories (perhaps to a fraction of an Earth ocean, or arid conditions) and the time/length of the "wet" period (during the first few hundred million years). Recent studies suggest that a runaway greenhouse transition with enough water in the atmosphere (a minimum of a few tens of bars) may have caused global melting of the surface, which may become a volatile and oxygen sink (Gillmann et al., 2022; Warren and Kite, 2023). However, no evidence of such an event has been recorded due to the young age of the surface of Venus. This remains an active research topic, but the lack of constraints and multiple processes involved still make quantitative estimates unreliable.

In scenario (ii) the remnant oxygen would have been immediately removed from the atmosphere by oxidizing ferrous iron in the surface magma e.g. (Gillmann et al., 2009a; Lammer et al., 2018). Compared with the solid rocky surface, the surface magma over the vigorously convective magma ocean is expected to react with the overlying atmosphere more efficiently, due to its higher mobility and resurfacing rate. Furthermore, modeling results suggest that, in scenario (ii), the planetary surface can remain molten until most of its water is lost (Hamano et al., 2013). This is because the surface tem-

perature could exceed the solidus temperature ($\sim 1,340$ K for peridotites) as far as a few tens of bars of steam atmosphere remains (Hamano, 2020). On the other hand, a recent numerical study (Selsis et al., 2023) suggest that, under some conditions, a water-vapor atmosphere may see its structure dominated by radiative rather than convective layers, leading to cooler surface temperatures than for an adiabatic temperature profile. As a result, the occurrence and/or lifetime of a magma ocean may be greatly reduced. The magma ocean could solidify earlier than previously assumed, before the steam atmosphere escapes, which would reduce its efficiency as a trap for volatile species, including oxygen.

Nonetheless, if the magma ocean acts as a volatile sink, the oxygen buildup associated with hydrodynamic escape would not have occurred when Venus followed scenario (ii). One of the implications expected from this scenario is that the Venus’ magma ocean may have a redox evolution different than Earth’s, depending on the amount of the remnant oxygen reacted. The oxidation of the magma ocean associated with extensive loss of water may affect the speciation and partitioning of volatile elements. It may also explain an elevated atmospheric N inventory on Venus, compared with Earth (Wordsworth, 2016). Due to the lack of direct samples, the redox state of the present Venusian mantle remains poorly constrained. As deduced from FeO contents of surfacial basalts, the overall oxygen budget may be similar to Earth’s (e.g. Richter et al., 2016), whereas thermodynamic calculations and spectroscopic data suggests that the oxygen fugacity of Venus’ surface may be as high as the magnetite–hematite phase boundary (Fegley et al., 1997).

Accumulation of oxygen is not an issue in scenario (iii) either, as water would have only be a minor component of early outgassing in the first place. Venus surface and atmosphere would be dry, but its interior may not be. The potentially different dryness of the interior among the three scenarios might affect the subsequent evolution of the interior. In such an evolution, inefficient outgassing of the magma ocean (Ikoma et al., 2018; Solomatova and Caracas, 2021; Salvador and Samuel, 2023) combined with high surface pressures (increasing solubility of water in the magma ocean Gaillard and Scaillet, 2014), would have limited outgassing of water. Early loss of hydrogen could have favored the oxidation of the magma ocean (Gaillard et al., 2022), therefore an increased loss of H on Venus may help explain the divergent evolution path of Venus. Later water outgassing related to volcanism would

be prevented by high surface pressures due to the thick CO₂ atmosphere (Gaillard and Scaillet, 2014).

5.2.2. *The interior of Venus*

Our understanding of the interior of Venus is very limited. Observations are generally consistent with an approximately Earth-like internal structure and composition, but error bars are large and accommodate a wide range of possible configurations. In short, no direct observations strongly imply any radical structural differences compared to Earth, despite the surface expressions of interior dynamics diverging widely from what we can observe on Earth.

It is consensus that Venus displays no present-day global plate tectonics, but is possibly (1) in a plutonic squishy lid regime because of its high surface temperature or (2) in a quiet period of an episodic lid regime, or (3) in stagnant-lid after a transition from mobile-lid regime. Specific sequences of regime transitions have been discussed by Rolf et al. (2022) and Gillmann et al. (2022) (and references therein), but so far the question remains unsolved. The absence of a global self-generated magnetic field on Venus (Nimmo, 2002; Smrekar et al., 2018), possibly caused by insufficient heat extraction out of the core, is likely tied to the question of the mantle dynamics evolution through time (O’Rourke et al., 2018). Further investigation into past evidence of magnetic field generation may provide new constraints for the evolution of Venus’ mantle thermal state and convection regime (O’Rourke et al., 2019).

Differences in the dynamics of the interior of Earth and Venus and convection regimes may be related to the suspected dryness of the interior of Venus (Kiefer et al., 1986; Grinspoon, 1993). Because water affects melting and the mantle viscosity, it also affects mantle dynamics: for example, a dry mantle produces less melt and has a higher viscosity; Green et al. (2014); Karato (2015). However, the extent to which water can be a critical factor still remains to be fully analysed and quantified. Additionally, some works suggesting dry Venus (Kiefer et al., 1986; Grinspoon, 1993) made strong assumptions on the processes at work on Venus. Kiefer et al. (1986), for example, linked the correlation between Venus’ geoid and topography to the absence of an asthenosphere, which could be due to dry conditions (and is now questioned; Armann and Tackley (2012); Maia and Wieczorek (2022)). The dryness of the interior inferred by Grinspoon (1993) relied on assumptions on the rate of volcanic production and outgassing. However, volcanic

production is poorly constrained and highly debated (Herrick et al., 2023), while outgassing itself would be affected by Venus’ high surface pressure and reduced water release (see Section 3.6). As a result, whether Venus’ interior is dry or wet remains uncertain.

So far, no measurement has allowed us to rule out a specific set of evolution scenarios with any degree of certainty. We have no sample of volcanic rocks or volatiles from Venus, to estimate its interior composition and conditions. Data on the state of the core are consistent with a solid, liquid or stably stratified core (Dumoulin et al., 2017). Models (O’Rourke et al., 2018) suggest that if Venus had an Earth-like structure and composition, it would likely generate a similar global magnetic field. Therefore, without a better understanding of the interior of the planet, we are unable to decide whether Venus lost its global magnetic field or never generated one in the first place.

This need for more observations is driving the current Venus exploration projects with multiple missions planned or selected: DaVinci (USA), VERITAS (USA), EnVision (Europe), Venus Orbiter Mission (also known as Shukryaan-1) (India), Venera-D (Russia) and VOICE (China). Those missions are set to improve our knowledge of the interior structure of the planet, state of its core, its subsurface structure, surface topography, geology and composition and atmosphere structure and composition (both close to the ground and higher up, with a special interest in volcanic production), with a focus on traces of activity through time.

5.3. *Earth*

5.3.1. *Oxidation of Earth’s magma ocean*

Compared with other planetary bodies such as Mars and the Moon, Earth has a highly oxidized mantle characterized by high Fe^{3+} abundance (e.g. Frost et al., 2008). The evolution of the redox state of the Earth’s mantle is one of the research fields actively explored (chapter 00008). Geochemical studies of Hadean zircons suggest that the oxidation of the Earth’s mantle might have occurred very early (Trail et al., 2011), possibly during its magma ocean stage.

One of the mechanisms proposed to explain the oxidation of the terrestrial magma ocean is hydrogen escape accompanying dissociation of water (Hamano et al., 2013; Sharp et al., 2013). The present Earth’s upper mantle has a $\text{Fe}^{3+}/\Sigma\text{Fe}$ of 0.02–0.06, which corresponds to the Fe_2O_3 content of 0.16–0.48 wt% (Hirschmann, 2022, and references therein). Using the simple assumption of oxidation of FeO to Fe_2O_3 , the amount of water required to

raise the Fe_2O_3 content from zero to the present level is equivalent to 0.5-1.5 Earth's ocean masses for the whole mantle, and about 1/3 of this value (i.e. 0.2-0.5 ocean masses) for the upper mantle only. Earth is sufficiently far from the critical orbital distance (6) discussed in Section 2.4, thus Earth's magma ocean would have crystallized rapidly on the order of 1 Myr or less (Hamano et al., 2013; Lebrun et al., 2013; Salvador et al., 2017; Bower et al., 2022). Hamano et al. (2013) estimated the total loss of water throughout the Earth's magma ocean stage to be about 0.2 ocean masses at most for a pure steam atmosphere. Due to the low mixing ratio of H-bearing species in the upper atmosphere, the loss of water would become probably lower if the early atmosphere included large amounts of other gaseous species such as CO_2 . Thus, water loss during the Earth's magma ocean stage might have a partial contribution to oxidation of the upper mantle, but the effect could be marginal for the oxidation of the whole mantle.

An alternative mechanism is the redox disproportionation of Fe^{2+} to Fe^{3+} and Fe^0 . This oxidation mechanism was originally proposed to be driven by crystallization of a solid phase with affinity for Fe^{3+} , such as perovskite. However, it is unclear how quickly Fe metal formed by iron disproportionation could sink to the core through the solid mantle. As described in section 2.1, recent experimental and theoretical studies reported that redox disproportionation could occur directly in a magma ocean owing to a high stability of Fe^{3+} at high pressure (Armstrong et al., 2019; Deng et al., 2020; Kuwahara et al., 2023). Metal silicate equilibration at the base of the magma ocean can produce a high $\text{Fe}^{3+}/\Sigma\text{Fe}$ in the melt even at an oxygen fugacity as low as IW-2. This oxidation mechanism requires that the base of the magma ocean attains pressures as high as tens of GPa. This is in line with estimations of terrestrial magma ocean depths caused by giant impacts during the late stage of Earth's formation (Nakajima et al., 2021). The Earth's large planetary mass and the accretion history involving giant impacts during the late stage of formation may be key factors that set the different initial condition for the redox evolution of its mantle.

This oxidation scenario seems promising for Earth, although the estimated $\text{Fe}^{3+}/\Sigma\text{Fe}$ in the silicate melt shows significant differences among studies. Based on the assumption of metal-silicate equilibration at 25-90 GPa, Deng et al. (2020) computed the $\text{Fe}^{3+}/\Sigma\text{Fe}$ of about 0.01-0.02, which is comparable to the value observed in the Earth's upper mantle. On the other hand, the $\text{Fe}^{3+}/\Sigma\text{Fe}$ estimated by Armstrong et al. (2019) and Kuwahara et al. (2023) are much higher. Using a relatively low base pressure of 25-

33 GPa, Kuwahara et al. (2023) obtained a $\text{Fe}^{3+}/\Sigma\text{Fe}$ value of 0.2-0.5, which is one order of magnitude higher than the $\text{Fe}^{3+}/\Sigma\text{Fe}$ in the upper mantle. The cause of such differences remains unclear. If the excess in the $\text{Fe}^{3+}/\Sigma\text{Fe}$ is confirmed, then it may suggest reduction processes such as the late accretion of reducing material (Dauphas, 2017; Fischer-Gödde and Kleine, 2017; Genda et al., 2017), which could form a transient reducing surface environment, or may reflect incomplete metal-silicate equilibration, rather than a single equilibration event at the base of the magma ocean (e.g. Rubie et al., 2015b).

5.3.2. *Earth’s most distinctive feature: Life affects planetary evolution and volatile cycles*

The discussion above has made abundantly clear the fact that the Earth today is very different to present-day Venus and Mars. The present-day Earth is characterised by an equable climate (from the point of view of life), stable water at its surface, an oxidised atmosphere, and a robust plate tectonic regime that links surface processes to those in the mantle, in the process buffering the whole system Earth.

At present, Venus’ surface is dry and hot with a less oxidized atmosphere (and H_2SO_4 clouds), and no plate tectonics. Mars is a dry cold planet with a CO_2 atmosphere and a stagnant lid. Neither has life at its surface and the question of whether Venus could host some form of life in its clouds, or Mars in its subsurface, remains debated.

However, as the above discussions point out, the Earth did not start out as it is today. Indeed, life could not have emerged on an oxidised planet since abiogenesis (the emergence of life, as we know it) requires reduced organic building bricks (e.g. Hudson et al. (2020)). On the other hand, the oxidation that is so characteristic of its surface is a product of biological processes. Thus, while the early planets, Earth, Venus and Mars may have shared certain characteristics that permitted life to emerge, their evolution paths diverged significantly. As we understand the emergence of life, it was conceivable on Mars but remains uncertain for Venus, due to the remaining uncertainties regarding early surface conditions (see section 5.2.1). Since life is an integral component of the Earth system, it is pertinent here to consider, in the context of terrestrial geology and geophysics, how it appeared and evolved through geological time. Since reconstructions of the past Earth’s atmosphere composition do not highlight significant changes in bulk composition related purely to long-term escape (Gronoff et al., 2020), the intimate

interaction between the effects of life at the surface, on internal processes, as well as on the atmospheric envelope and vice versa, is of equal importance.

There are numerous theories about how and where life emerged, in deep ocean hydrothermal vents (Baross and Hoffman, 1985; Russell and Hall, 1997; Martin and Russell, 2003), shallow water hydrothermal vents (Westall et al., 2018), subaerial hydrothermal vents (Damer and Deamer, 2020; Van Kranendonk et al., 2021), in impact craters (Sasselov et al., 2020), in deep-seated fault systems (Schreiber et al., 2012), in pumice rafts (Brasier et al., 2011) or in nuclear sands (Maruyama and Ebisuzaki, 2023) to name but a few hypotheses (for a review see (Westall et al., 2018)). Westall et al. (2023) briefly reviewed the conditions that existed during the Hadean (4.5-4.0 Ga), and that were apparently compatible with prebiotic processes and the emergence of life. They included an anoxic atmosphere, (i) a high EUV flux reaching the surface of the Earth that may or may not have been beneficial for early prebiotic processes, depending upon the scenario preferred, (ii) a highly volcanic environment including abundant hydrothermal activity, and (iii) a largely ocean-covered planet with little exposed land mass. Modern-style plate tectonics did not exist and there is still much debate as to what style of tectonic regime was active during the Hadean (Capitanio et al., 2020; Gerya, 2022). A Hadean initiation of plate tectonics has been hypothesised on the basis of an andesitic source for zircons dating back to the Hadean (Turner et al., 2020), while evidence for both vertical and horizontal deformation styles has been documented in Paleo-Meso Archean formations (Heubeck et al., 2023), suggesting some form of tectonic activity. Recent studies suggest that modern style plate tectonics did not arise until much later in Earth’s history, towards the end of the Proterozoic, about 800 Ma (see Figure 7, (Hawkesworth et al., 2020; Stern, 2018), therefore the plutonic squishy lid might offer an alternative explanation (Rozel et al., 2017; Lourenço et al., 2018).

As soon as it appeared, maybe multiple times before really gaining a foothold, early life forms were intimately linked to the volcanic substrates from which they emerged. Note that all the proposed scenarios for the the emergence of life referred to above were located in volcanic settings. They were chemolithotrophic microorganisms obtaining their energy from oxidation of inorganic materials, such as H_2 , or through redox alteration of reactive minerals at the surface of the volcanic rocks (e.g. olivine, pyroxene, metal sulphides), or chemoorganotrophs oxidising organic substrates, such as dead organisms or bio-available abiotic organics (Weiss et al., 2016). At this stage, life was completely dependent upon the mineral substrates. At some stage,




Figure 7: Indicators of crustal evolution and the gradual rise of modern style plate tectonics, using zircon crystallization age distribution (lower serrated yellow curves), throughout geological history. Vertical grey bars mark the development of supercontinents/supercratons (see references in Hawkesworth et al. (2020)). The crustal growth curve (brown) is adapted from modelling by Dhuime et al. (2012). The crustal reworking curve is based on the Hf isotope ratios of zircons in different time slices (Dhuime et al., 2012). Variations in the thickness of continental crust at the sites of crust generation through time is inferred from Rb/Sr measurements. The variation of $\delta^{18}\text{O}$ values in zircons also marks the shifts in the composition of juvenile crust away from mafic compositions (Hawkesworth et al., 2020) (adapted from Hawkesworth et al. (2020)).

through mutation, life developed the ability to capture energy from sunlight (phototrophs, (Martin et al., 2018)), thus freeing it of the mineral substrates and providing it with access to an unlimited source of energy. The oldest, well-preserved terrestrial rocks dating back to about 3.5 Ga testify to the already widespread presence of anoxygenic phototrophs in shallow water environments.

While access to unlimited sunlight gave the phototrophs an energy-rich, metabolic advantage over the chemotrophs (litho- and organ-) enabling them to grow faster and create greater biomass, it was the mutation that led to oxygenic photosynthesis that would change the Earth, both physically, chemically, and biologically. The timing of this biological invention is debated with certain geochemical studies suggesting that “whiffs of oxygen”, i.e. biological




Figure 8: Relative concentrations of atmospheric oxygen throughout Earth’s history, including the classical two-step evolution (orange line, Kump, 2008) and the more recent model including unstable climate and O_2 concentrations after the first rise of O_2 concentrations (Lyons et al., 2014). Possible “whiffs of oxygen” (blue arrows) during the Neoarchean – and even before according to (Anbar et al., 2007) (adapted from (Lyons et al., 2014))

oxygen, may have occurred as early as the Paleo-Mesoarchean (~ 3.5 - 3.0 Ga) and during the next 500 Myr (see Fig. 8; Anbar et al. (2007)). Certainly by 2.9-3.0 Ga there is geochemical evidence for free oxygen and bioconstructions, such as stromatolites, start becoming common. It has been suggested that changes in the redox state of the Earth’s surface during the NeoArchean-Proterozoic transition may have been influenced not only by the production of oxygen by photosynthesisers, but also by changes in the redox state of the mantle, that affected the redox of outgassing species from volcanic eruptions (Wang et al. (2020)), although others have suggested that there was little change in the oxygen fugacity of the Archean mantle (e.g. Nicklas et al. (2018)). Whatever the major influences during the Neoarchean, during this period, huge formations of stromatolites produced by oxygenic phototrophs occur in Neoarchean formations across the world and, indeed, dominated the paleontological record for most of the Proterozoic.

The turning point was when the oxygen contents in the atmosphere started slowly to rise between about 2.4–2.3 Ga (the so-called “Great Oxygenation Event” (GOE), Holland (2002), see Fig. 8). In fact, the gradual accumulation of oxygen depended upon a delicate interplay of sinks and sources that probably started as early as about 3.0 Ga (Lyons et al., 2014) and may have continued until about 800 Ma (Lyons et al., 2021). As noted above, we keep in mind, however, that there may not have been major changes to the redox state of the mantle (Nicklas et al., 2018). The production of oxygen was balanced by its consumption by the initially totally reduced surface of the Earth, including reduced organic carbon, sulphide minerals, other ferrous minerals, and reduced volcanic gases, such as H_2S , as well as bio-

genic methane and its photodissociated product ethane. Concurrent with the gradual oxidation of the Earth's surface envelope, including its atmosphere, gradual changes in mantle buffering resulted in changes in the composition of volcanic gases, more reduced during the Archean, but more oxidised after the GOE.

While the presence of oxygen in the atmosphere led to elimination of methane as greenhouse gas, there is still the problem of reducing CO₂ in the Neoarchean atmosphere. The isotopic ratios of carbon in the Neoarchean are relatively constant, suggesting that there was not a major change in the burial flux of the total carbon inventory. However, one solution to the conundrum could be that the flux of buried organic carbon during the Archean may have been far higher than previously believed (Krissansen-Totton et al., 2021b). Whatever the processes involved, oxygenation of the atmosphere had the catastrophic consequence of lowering surface temperatures because the Sun's luminosity was still about 20-30% lower than today (as was the case during the Hadean and Archean, Sagan and Mullen (1972)), which led to the first series of global glaciations, the Huronian glaciations between 2.45 and 2.22 Ga, when low latitude continental terranes were covered with glaciers (Bekker and Holland, 2012). This effectively cut off phototrophic production of oxygen, increased temperatures due to the accumulation of volcanic CO₂ that led to melting of the glaciers and a sudden massive influx of nutrients that fed the rapid re-development of the oxygen producing cyanobacteria and the precipitation of biomediated carbonate deposits, the so-called cap carbonates, Hoffman et al. (1998). The sudden explosion of biomass production led to a very large rise in oxygen levels between 2.3 and 2.06 Ga, called the Lomagundi-Jatuli event, during which oxygen levels almost reached present-day levels (Melezhik et al., 2013), before falling to very low atmospheric partial pressure levels of O₂ < 0.1% of present atmospheric levels (Lyons et al., 2014) that reigned for the rest of the Proterozoic until about 0.8 Ga (Canfield, 1998), as illustrated by Figure 8. It is thought that this long period of stasis during the Meso-Neoproterozoic was due to the fact that only the upper part of the oceans were oxygenated while the deep ocean remained reduced.

However, it was during the Proterozoic that the extra energy available via oxygenic photosynthesis fuelled the mutations giving rise to eukaryotes, i.e. organisms with a membrane bound nucleus (prokaryotes, including Bacteria and Archaea do not have membrane bound nuclei). Indeed, it was only after complete oxygenation of the entire water column and the stabilisation of nu-

trient cycles and environmental conditions could life evolve into the diverse biota that characterised the terminal stages of the Neoproterozoic (1-0.54 Ga) and the dawn of the Cambrian period (Planavsky et al., 2015). The Neoproterozoic period was characterized by extreme biogeochemical and climatic volatility that led to a series of very severe, low latitude glaciations (Hoffman et al., 1998; Canfield, 2005; Lyons et al., 2014) resulting in a highly unstable carbon cycle. This period of multiple “snowball Earths” could have been triggered or at least influenced by the onset of modern plate tectonics (e.g. Stern et al., 2018). The global distribution of continents and their positions relative to the equator likely also had an effect on albedo and therefore temperatures (Endal and Schatten, 1982; Schrag et al., 2002). At the same time, changes in oceanic ecosystems led to the rise of eukaryotic algal primary producers as opposed to the prokaryotic cyanobacteria so prevalent from the Neoarchean (3.0-2.5 Ga) through the Mesoproterozoic (2.0-1.0 Ga), and the eventual emergence of metazoan life. Add to this the rise in oxygen globally and the conditions for significant evolutionary changes to life were all present (Planavsky et al., 2015).

Today, there appears to be a delicate balance between atmospheric composition, especially the greenhouse gases CO_2 and CH_4 , the presence of water at Earth’s surface, subduction of organic matter and carbonates, and subsequent release of volcanic CO_2 back into the Earth’s atmosphere (see Charnay et al. (2020) and Catling and Zahnle (2020) for recent reviews). It was likely established gradually throughout the geological evolution of the Earth, possibly as far back as the Paleoarchean or even before during the Hadean. The exact process for the recycling of this early carbon cycle is still unknown, but it was most likely different from modern-day plate tectonics. Initially, the higher CO_2 concentration in the Hadean/Archean atmosphere would have offset the lower luminosity from the Sun (i.e. Kasting, 1988). As solar luminosity increased, CO_2 was gradually removed from the atmosphere by the silicate/carbonate weathering cycle and sequestration of carbonates and organic carbon into the crust. In this way, throughout time, climatic stability has been ensured – as has been the presence of liquid water, one of the prime necessities of life on Earth.

6. Conclusions

The range of possible terrestrial planet evolution is difficult to explore with only three very particular examples, however, even in our Solar System,

the diversity among the rocky planets is striking. The exact causes for the clear differences between present-day Mars, Venus and Earth are still uncertain. It is now suspected that both their early (first few 100 Myr) and long-term evolution had a role to play. Atmospheric and interior conditions are the result of billions of years of accumulated and interacting processes that affect the physical and chemical properties of every planetary layer.

In recent years, our understanding of how magma ocean processes affect the early evolution of terrestrial planets has grown considerably. Studies show that in case the terrestrial planets started similarly in a hot molten state, the iron disproportionation at high pressures in a magma ocean can set a different initial condition for their subsequent redox evolution. Based on the assumption that metal-silicate equilibration occurred at the base of the magma ocean, due to their higher mass, Earth and Venus could have had a deep magma ocean with a higher abundance of Fe^{3+} and an oxidizing atmosphere consisting of CO_2 , N_2 and SO_2 , while a shallow magma ocean on Mars would be relatively Fe^{3+} -poor and overlaid by a weakly reducing atmosphere consisting of CO , CO_2 , and N_2 . On all the three planets, the majority of H would be dissolved in the magma ocean as H_2O at an early stage, and then would have outgassed at the surface as solidification proceeded.

In the case that a steam-dominated atmosphere forms, the subsequent evolution of the terrestrial planets depends on their orbital distance from the Sun. Earth and Mars could be sufficiently far from the Sun to keep their water intact against hydrodynamic escape owing to their short crystallization period. On the other hand, for Venus, three main different scenarios are possible: (i) rapid crystallization where water is retained or (ii) desiccation during a long-lived magma ocean period or (iii) that a high pressure CO_2 dominated atmosphere suppresses water outgassing. In the second scenario, the Venusian magma ocean can be oxidized by remnant oxygen produced by photodissociation of water, followed by preferential escape of hydrogen. At this point, the redox evolution of Earth and Venus may also diverge. If H_2O outgassing is insufficient to form a steam-dominated atmosphere, such an evolutionary dichotomy would not appear, and early planetary mantles would contain high abundance of water as discussed in scenario (iii). Planetary evolution during a magma ocean phase would set the initial conditions in terms of redox state in the mantle and atmosphere. This can have consequences for the crystallization history, the total and interior inventory of water and the climate throughout the entire planetary evolution.

Tectonic regimes can be divided into three main types of convection

regimes: plate tectonics, plutonic-squishy lid, and stagnant lid. Present-day Earth lies in the plate tectonics regime and features high heat flow and volcanic production rates. The onset time of modern plate tectonics on Earth is still under debate and might be related to the presence of life on Earth. The onset of plate tectonics depends on the rheology of the planetary lithosphere and the stresses in the convective mantle. Although surface temperature and water abundance are considered to be important factors affecting the rheology, whether they favor plate tectonics or not is still debated due to complicated feedbacks. The presence of subduction zones favors volatile exchanges between the atmosphere and the mantle. Mars has probably been in the stagnant lid regime for billion of years with limited geological activity in the last billion years. The heat flux in the stagnant lid regime is expected to be low, and this could explain why its core is still in a mostly liquid state, while not featuring a present-day core dynamo. Its smaller mass and more reduced interior is expected to limit total outgassing from its interior.

In contrast to Earth and Mars, the tectonic regime of past and present Venus remains unclear. Present-day Venus does not display signs of Earth-like plate tectonics, but seems to be more active than Mars and its stagnant lid regime. Its relatively young surface obscures most signs of prior tectonic and volcanic activity. A major constraint on Venus' interior is that at present it does not feature a global magnetic field. As Venus and Earth are comparable in size and mass, their differences may originate early on during their respective accretion and magma ocean stage.

In addition to volatile exchanges between surface and the interior, atmospheric escape is also a crucial process shaping and driving the evolution of planetary atmospheres. During the early evolution, hydrodynamic escape and impact erosion are important processes. Hydrodynamic escape is driven by strong EUV flux from the young Sun or by heating from a hot surface. Intense hydrodynamic escape would occur when the upper atmosphere is rich in H-bearing gaseous species such as H_2 and H_2O . This process would efficiently remove any primordial H_2 -He atmosphere the planetary embryos obtained from the solar nebula during their formation. Also this process would control the crystallization rate of a magma ocean located under a runaway greenhouse steam atmosphere. Impact erosion removes portions of planetary atmospheres. The most efficient loss occurs on small planets like Mars and/or in the presence of a surface water ocean. Impacts can also supply material, including volatiles. The net effect largely depends on the volatile content of the impactors, but also on the specific impact conditions.

As both impact and EUV flux decrease with time, non-thermal escape processes gain importance. Non-thermal escape involves photochemistry and charged particle interactions with the solar wind and among particles. Particles that obtained sufficient energy from the interaction can escape. Small planets like Mars are again more vulnerable to loss of heavier volatile species, such as O, C, N, and noble gases. Water loss, through escape of H and O ions is possible on all three planets at a comparable, but relatively slow rate. Based on recent measurements the previous idea that magnetic fields would provide an efficient shield against non-thermal escape is now being questioned as escape rates of heavy ions are comparable on Earth (featuring a global magnetic field), Venus, and Mars (no present-day core dynamo).

Upcoming missions to Venus specifically aim to shed light onto its past evolution compared to those of Mars and Earth. Likewise, sample return missions from Mars should lead to a dramatically better understanding of both Mars interior and surface conditions, as well as whether Mars might be currently hosting life in the subsurface. The future characterization of a large number of terrestrial exoplanets will provide further reference points for the possible evolution paths of rocky planets at all stages of their histories, including the critical magma ocean phase. The combination of terrestrial planet studies and exoplanet observations will provide both a detailed and statistical view of terrestrial evolution pathways and will help us to solve remaining mysteries that we have touched upon within the framework of this chapter.

Acknowledgements

The authors thank the anonymous reviewer for his constructive comments that helped to improve the manuscript. K.H. acknowledges financial support by JSPS KAKENHI grant No. JP18K13603 and MEXT KAKENHI grant No. JP22H05150. CG acknowledges that this work has been carried out within the framework of the NCCR PlanetS supported by the Swiss National Science Foundation under grants 51NF40_182901 and 51NF40_205606.

References

- Abe, Y., Matsui, T., 1988. Evolution of an impact-generated H₂O–CO₂ atmosphere and formation of a hot proto-ocean on earth. *Journal of Atmospheric Sciences* 45, 3081 – 3101. URL: https://journals.ametsoc.org/view/journals/atsc/45/21/1520-0469_1988_045_3081_eoaigh_2_0_co_2.xml, doi:10.1175/1520-0469(1988)045<3081:EOAIGH>2.0.CO;2.

- Acuña, M.H., Connerney, J.E.P., Ness, N.F., Lin, R.P., Mitchell, D., Carlson, C.W., McFadden, J., Anderson, K.A., Rème, H., Mazelle, C., Vignes, D., Wasilewski, P., Cloutier, P., 1999. Global distribution of crustal magnetization discovered by the mars global surveyor mag/er experiment. *Science* 284, 790–793.
- Alemi, A., Stevenson, D., 2006. Why Venus has No Moon, in: AAS/Division for Planetary Sciences Meeting Abstracts #38, p. 07.03.
- Alsaeed, N., Jakosky, B., 2019. Mars water and d/h evolution from 3.3 ga to present. *Journal of Geophysical Research: Planets* 124, 3344–3353.
- Anbar, A.D., Duan, Y., Lyons, T.W., Arnold, G.L., Kendall, B., Creaser, R.A., Kaufman, A.J., Gordon, G.W., Scott, C., Garvin, J., et al., 2007. A whiff of oxygen before the great oxidation event? *Science* 317, 1903–1906.
- Andrews-Hanna, J.C., Zuber, M.T., Banerdt, W.B., 2008. The Borealis basin and the origin of the martian crustal dichotomy. *Nature* 453, 1212 – 1215.
- Arkani-Hamed, J., 1993. On the tectonics of venus. *Physics of the earth and planetary interiors* 76, 75–96.
- Armann, M., Tackley, P.J., 2012. Simulating the thermochemical magmatic and tectonic evolution of venus’s mantle and lithosphere: Two-dimensional models. *Journal of Geophysical Research: Planets* 117.
- Armstrong, K., Frost, D.J., McCammon, C.A., Rubie, D.C., Ballaran, T.B., 2019. Deep magma ocean formation set the oxidation state of Earth’s mantle. *Science* 365, 903 – 906.
- Avicé, G., Marty, B., 2020. Perspectives on atmospheric evolution from noble gas and nitrogen isotopes on earth, mars & venus. *Space Science Reviews* 216, 36.
- Avicé, G., Marty, B., Burgess, R., 2017. The origin and degassing history of the earth’s atmosphere revealed by archean xenon. *Nature communications* 8, 15455.
- Avicé, G., Parai, R., Jacobson, S., Labidi, J., Trainer, M.G., Petkov, M.P., 2022. Noble Gases and Stable Isotopes Track the Origin and Early Evolution of the Venus Atmosphere. *Space Science Reviews* 218,

60. URL: <https://link.springer.com/10.1007/s11214-022-00929-9>, doi:10.1007/s11214-022-00929-9.
- Ballantyne, H.A., Jutzi, M., Golabek, G.J., Mishra, L., Cheng, K.W., Rozel, A.B., Tackley, P.J., 2023. Investigating the feasibility of an impact-induced martian dichotomy. *Icarus* 392, 115395. URL: <https://www.sciencedirect.com/science/article/pii/S0019103522004870>, doi:<https://doi.org/10.1016/j.icarus.2022.115395>.
- Ballhaus, C., Berry, R., Green, D., 1990. Oxygen fugacity controls in the earth's upper mantle. *Nature* 348, 437–440.
- Baross, J.A., Hoffman, S.E., 1985. Submarine hydrothermal vents and associated gradient environments as sites for the origin and evolution of life. *Origins of Life and Evolution of the Biosphere* 15, 327–345.
- Basilevsky, A.T., Head, J.W., 2003. The surface of venus. *Reports on Progress in Physics* 66, 1699.
- Bean, J.L., Abbot, D.S., Kempton, E.M.R., 2017. A statistical comparative planetology approach to the hunt for habitable exoplanets and life beyond the solar system. *The Astrophysical Journal Letters* 841, L24.
- Becker, R., Clayton, R., Galimov, E., Lammer, H., Marty, B., Pepin, R., Wieler, R., 2003. Isotopic signatures of volatiles in terrestrial planets: Working group report, in: *Solar System History from Isotopic Signatures of Volatile Elements: Volume Resulting from an ISSI Workshop 14–18 January 2002*, Bern, Switzerland, Springer. pp. 377–410.
- Bekker, A., Holland, H., 2012. Oxygen overshoot and recovery during the early paleoproterozoic. *Earth and Planetary Science Letters* 317, 295–304.
- Bercovici, D., Ricard, Y., 2014. Plate tectonics, damage and inheritance. *Nature* 508, 513–516.
- Berger, G., Cathala, A., Fabre, S., Borisova, A.Y., Pages, A., Aigouy, T., Esvan, J., Pinet, P., 2019. Experimental exploration of volcanic rocks-atmosphere interaction under venus surface conditions. *Icarus* 329, 8–23. URL: <https://www.sciencedirect.com/science/article/pii/S0019103518306134>, doi:<https://doi.org/10.1016/j.icarus.2019.03.033>.

- Bibring, J.P., Langevin, Y., Mustard, J.F., Poulet, F., Arvidson, R., Gendrin, A., Gondet, B., Mangold, N., Pinet, P., Forget, F., et al., 2006. Global mineralogical and aqueous mars history derived from omega/mars express data. *Science* 312, 400–404.
- Bindschadler, D.L., Schubert, G., Ford, P.G., 1994. Venus’ center of figure-center of mass offset. *Icarus* 111, 417–432. URL: <https://www.sciencedirect.com/science/article/pii/S0019103584711535>, doi:<https://doi.org/10.1006/icar.1994.1153>.
- Borg, L.E., Brennecka, G.A., Symes, S.J., 2016. Accretion timescale and impact history of mars deduced from the isotopic systematics of martian meteorites. *Geochimica et Cosmochimica Acta* 175, 150–167. URL: <https://www.sciencedirect.com/science/article/pii/S0016703715006948>, doi:<https://doi.org/10.1016/j.gca.2015.12.002>.
- Borgeat, X., Tackley, P., 2022. Hadean/eoarchean tectonics and mantle mixing induced by impacts: a three-dimensional study. *Prog Earth Planet Sci* 9, 38.
- Boujibar, A., Bolfan-Casanova, N., Andrault, D., Bouhifd, M.A., Trcera, N., 2016. Incorporation of Fe²⁺ and Fe³⁺ in bridgmanite during magma ocean crystallization. *American Mineralogist* 101, 1560–1570. URL: <https://doi.org/10.2138/am-2016-5561>, doi:10.2138/am-2016-5561.
- Bouvier, L.C., Costa, M.M., Connelly, J.N., Jensen, N.K., Wielandt, D., Storey, M., Nemchin, A.A., Whitehouse, M.J., Snape, J.F., Bellucci, J.J., Moynier, F., Agranier, A., Gueguen, B., Schönbächler, M., Martin Bizzarro, M., 2018. Evidence for extremely rapid magma ocean crystallization and crust formation on Mars. *Nature* 558, 586–589.
- Bower, D.J., Hakim, K., Sossi, P.A., Sanan, P., 2022. Retention of water in terrestrial magma oceans and carbon-rich early atmospheres. *The Planetary Science Journal* 3, 93. URL: <https://doi.org/10.3847/psj/ac5fb1>, doi:10.3847/psj/ac5fb1.
- Bower, D.J., Kitzmann, D., Wolf, A.S., Sanan, P., Dorn, C., Oza, A.V., 2019. Linking the evolution of terrestrial interiors and an early outgassed

- atmosphere to astrophysical observations. *Astronomy & Astrophysics* 631, A103.
- Brain, D., Leblanc, F., Luhmann, J., Moore, T.E., Tian, F., 2012. Planetary magnetic fields and climate evolution. *Comparative climatology of terrestrial planets* 1675, 8049.
- Brasier, M.D., Matthewman, R., McMahon, S., Wacey, D., 2011. Pumice as a remarkable substrate for the origin of life. *Astrobiology* 11, 725–735.
- Bullock, M.A., Grinspoon, D.H., 2001. The recent evolution of climate on venus. *Icarus* 150, 19–37.
- Šrámek, O., Zhong, S., 2010. Long-wavelength stagnant lid convection with hemispheric variation in lithospheric thickness: Link between martian crustal dichotomy and tharsis? *Journal of Geophysical Research: Planets* 115. URL: <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2010JE003597>, doi:<https://doi.org/10.1029/2010JE003597>, arXiv:<https://agupubs.onlinelibrary.wiley.com/doi/pdf/10.1029/2010JE003597>.
- Cameron, A., 1983. Origin of the atmospheres of the terrestrial planets. *Icarus* 56, 195–201. URL: <https://www.sciencedirect.com/science/article/pii/0019103583900325>, doi:[https://doi.org/10.1016/0019-1035\(83\)90032-5](https://doi.org/10.1016/0019-1035(83)90032-5).
- Canfield, D.E., 1998. A new model for proterozoic ocean chemistry. *Nature* 396, 450–453.
- Canfield, D.E., 2005. The early history of atmospheric oxygen: homage to robert m. garrels. *Annu. Rev. Earth Planet. Sci.* 33, 1–36.
- Canup, R.M., 2012. Forming a moon with an earth-like composition via a giant impact. *Science* 338, 1052–1055. URL: <https://www.science.org/doi/abs/10.1126/science.1226073>, doi:10.1126/science.1226073, arXiv:<https://www.science.org/doi/pdf/10.1126/science.1226073>.
- Capitanio, F.A., Nebel, O., Cawood, P.A., 2020. Thermochemical lithosphere differentiation and the origin of cratonic mantle. *Nature* 588, 89–94.
- Carr, M.H., 2007. *The surface of Mars. volume 6.* Cambridge University Press.

- Carr, M.H., Head, J.W., 2010. Geologic history of mars. *Earth and Planetary Science Letters* 294, 185–203. URL: <https://www.sciencedirect.com/science/article/pii/S0012821X09003847>, doi:<https://doi.org/10.1016/j.epsl.2009.06.042>. mars Express after 6 Years in Orbit: Mars Geology from Three-Dimensional Mapping by the High Resolution Stereo Camera (HRSC) Experiment.
- Cassata, W.S., 2017. Meteorite constraints on martian atmospheric loss and paleoclimate. *Earth and Planetary Science Letters* 479, 322–329.
- Cassata, W.S., Zahnle, K.J., Samperton, K.M., Stephenson, P.C., Wimpenny, J., 2022. Xenon isotope constraints on ancient martian atmospheric escape. *Earth and Planetary Science Letters* 580, 117349.
- Catling, D.C., Zahnle, K.J., 2009. The planetary air leak. *Scientific American* 300, 36–43.
- Catling, D.C., Zahnle, K.J., 2020. The archean atmosphere. *Science advances* 6, eaax1420.
- Cawood, P.A., Kroner, A., Pisarevsky, S., 2006. Precambrian plate tectonics: criteria and evidence. *GSA today* 16, 4.
- Chamberlain, J.W., 1963. Planetary coronae and atmospheric evaporation. *Planetary and Space Science* 11, 901–960.
- Charnay, B., Wolf, E.T., Marty, B., Forget, F., 2020. Is the faint young sun problem for earth solved? *Space Science Reviews* 216, 1–29.
- Chassefière, E., 1996. Hydrodynamic escape of oxygen from primitive atmospheres: applications to the cases of venus and mars. *Icarus* 124, 537–552.
- Chassefière, E., Leblanc, F., 2004. Mars atmospheric escape and evolution; interaction with the solar wind. *Planetary and Space Science* 52, 1039–1058.
- Cheng, K.W., Rozel, A.B., Golabek, G.J., Ballantyne, H.A., Jutzi, M., Tackley, P.J., 2024. Mars’s crustal and volcanic structure explained by southern giant impact and resulting mantle depletion. *Geophysical Research Letters* 51, e2023GL105910. doi:<https://doi.org/10.1029/2023GL105910>.

- Christensen, P., Engle, E., Anwar, S., Dickenshied, S., Noss, D., Gorelick, N., Weiss-Malik, M., 2009. Jmars-a planetary gis, in: AGU fall meeting Abstracts, pp. IN22A–06.
- Christensen, U.R., 1984. Heat transport by variable viscosity convection and implications for the Earth’s thermal evolution. *Physics of the Earth and Planetary Interiors* 35, 264–282.
- Claire, M.W., Sheets, J., Cohen, M., Ribas, I., Meadows, V.S., Catling, D.C., 2012. The evolution of solar flux from 0.1 nm to 160 μm : quantitative estimates for planetary studies. *The Astrophysical Journal* 757, 95.
- Cloud Jr, P.E., 1968. Atmospheric and hydrospheric evolution on the primitive earth: Both secular accretion and biological and geochemical processes have affected earth’s volatile envelope. *Science* 160, 729–736.
- Conrad, J., Nimmo, F., 2023. Constraining characteristic morphological wavelengths for venus using baltis vallis. *Geophysical Research Letters* 50, e2022GL101268.
- Conrad, P.G., Malespin, C.A., Franz, H.B., Pepin, R.O., Trainer, M.G., Schwenzer, S.P., Atreya, S., Freissinet, C., Jones, J., Manning, H., et al., 2016. In situ measurement of atmospheric krypton and xenon on mars with mars science laboratory. *Earth and Planetary Science Letters* 454, 1–9.
- Correia, A.C., Laskar, J., 2003. Long-term evolution of the spin of venus: II. numerical simulations. *Icarus* 163, 24–45. URL: <https://www.sciencedirect.com/science/article/pii/S0019103503000435>, doi:[https://doi.org/10.1016/S0019-1035\(03\)00043-5](https://doi.org/10.1016/S0019-1035(03)00043-5).
- Crameri, F., Tackley, P.J., 2014. Spontaneous development of arcuate single-sided subduction in global 3-d mantle convection models with a free surface. *Journal of Geophysical Research: Solid Earth* 119, 5921–5942. URL: <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1002/2014JB010939>, doi:<https://doi.org/10.1002/2014JB010939>, arXiv:<https://agupubs.onlinelibrary.wiley.com/doi/pdf/10.1002/2014JB010939>.
- Crameri, F., Tackley, P.J., 2015. Parameters controlling dynamically self-consistent plate tectonics and single-sided subduction in global models of mantle convection. *Journal of*

- Geophysical Research: Solid Earth 120, 3680–3706. URL: <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1002/2014JB011664>, doi:<https://doi.org/10.1002/2014JB011664>, arXiv:<https://agupubs.onlinelibrary.wiley.com/doi/pdf/10.1002/2014JB011664>.
- Crisp, J.A., 1984. Rates of magma emplacement and volcanic output. *Journal of Volcanology and Geothermal Research* 20, 177–211. URL: <https://www.sciencedirect.com/science/article/pii/0377027384900398>, doi:[https://doi.org/10.1016/0377-0273\(84\)90039-8](https://doi.org/10.1016/0377-0273(84)90039-8).
- Crowley, J.W., Gérard, M., O’Connell, R.J., 2011. On the relative influence of heat and water transport on planetary dynamics. *Earth and Planetary Science Letters* 310, 380–388.
- Damer, B., Deamer, D., 2020. The hot spring hypothesis for an origin of life. *Astrobiology* 20, 429–452.
- Dasgupta, R., 2013. Ingassing, Storage, and Outgassing of Terrestrial Carbon through Geologic Time. *Reviews in Mineralogy and Geochemistry* 75, 183–229. URL: <https://doi.org/10.2138/rmg.2013.75.7>, doi:10.2138/rmg.2013.75.7, arXiv:<https://pubs.geoscienceworld.org/msa/rimg/article-pdf/75/1/183/2952840/183>
- Dasgupta, R., Hirschmann, M.M., 2010. The deep carbon cycle and melting in earth’s interior. *Earth and Planetary Science Letters* 298, 1–13.
- Dauphas, N., 2017. The isotopic nature of the Earth’s accreting material through time. *Nature* 541, 521–524. doi:10.1038/nature20830.
- Dauphas, N., Pourmand, A., 2011. Hf–W–Th evidence for rapid growth of Mars and its status as a planetary embryo. *Nature* 473, 489–492.
- Davis, F.A., Cottrell, E., 2021. Partitioning of Fe_2O_3 in peridotite partial melting experiments over a range of oxygen fugacities elucidates ferric iron systematics in mid-ocean ridge basalts and ferric iron content of the upper mantle. *Contributions to Mineralogy and Petrology* 176, 67. URL: <https://doi.org/10.1007/s00410-021-01823-3>, doi:10.1007/s00410-021-01823-3.

- Dehant, V., Debaille, V., Dobos, V., Gaillard, F., Gillmann, C., Goderis, S., Grenfell, J.L., Höning, D., Javaux, E.J., Karatekin, Ö., et al., 2019. Geoscience for understanding habitability in the solar system and beyond. *Space Science Reviews* 215, 1–48.
- Deng, J., Du, Z., Karki, B.B., Ghosh, D.B., Lee, K.K.M., 2020. A magma ocean origin to divergent redox evolutions of rocky planetary bodies and early atmospheres. *Nature Communications* 11, 2007. URL: <https://doi.org/10.1038/s41467-020-15757-0>, doi:10.1038/s41467-020-15757-0.
- Dhuime, B., Hawkesworth, C.J., Cawood, P.A., Storey, C.D., 2012. A change in the geodynamics of continental growth 3 billion years ago. *Science* 335, 1334–1336.
- Dhuime, B., Wuestefeld, A., Hawkesworth, C.J., 2015. Emergence of modern continental crust about 3 billion years ago. *Nature Geoscience* 8, 552–555.
- Dobrovolskis, A.R., Ingersoll, A.P., 1980. Atmospheric tides and the rotation of venus i. tidal theory and the balance of torques. *Icarus* 41, 1–17.
- Donahue, T., Grinspoon, D., Hartle, R., Hodges Jr, R., 1997. Ion/neutral escape of hydrogen and deuterium: Evolution of water. *Venus II: Geology, geophysics, atmosphere, and solar wind environment* 385.
- Donahue, T., Pollack, J., 1983. Origin and evolution of the atmosphere of venus, in: *Venus*. University of Arizona Press, Tucson, p. 1003–1036.
- Drake, M.J., 2000. Accretion and primary differentiation of the earth: a personal journey. *Geochimica et Cosmochimica Acta* 64, 2363–2369. URL: <https://www.sciencedirect.com/science/article/pii/S0016703700003720>, doi:[https://doi.org/10.1016/S0016-7037\(00\)00372-0](https://doi.org/10.1016/S0016-7037(00)00372-0).
- Dumoulin, C., Tobie, G., Verhoeven, O., Rosenblatt, P., Rambaux, N., 2017. Tidal constraints on the interior of venus. *J. Geophys. Res. Planets* 122, 1338–1352.
- Elkins-Tanton, L., 2008. Linked magma ocean solidification and atmospheric growth for earth and mars. *Earth and Planetary Science Letters* 271, 181–191. URL: <https://www.sciencedirect.com/science/article/>

- pii/S0012821X08002306, doi:<https://doi.org/10.1016/j.epsl.2008.03.062>.
- Elkins-Tanton, L., Zaranek, S., Parmentier, E., Hess, P., 2005. Early magnetic field and magmatic activity on mars from magma ocean cumulate overturn. *Earth and Planetary Science Letters* 236, 1–12. URL: <https://www.sciencedirect.com/science/article/pii/S0012821X05002761>, doi:<https://doi.org/10.1016/j.epsl.2005.04.044>.
- Endal, A.S., Schatten, K.H., 1982. The faint young sun-climate paradox: Continental influences. *Journal of Geophysical Research: Oceans* 87, 7295–7302.
- Fegley, B., Zolotov, M.Y., Lodders, K., 1997. The oxidation state of the lower atmosphere and surface of venus. *Icarus* 125, 416–439. URL: <https://www.sciencedirect.com/science/article/pii/S0019103596956286>, doi:<https://doi.org/10.1006/icar.1996.5628>.
- Fegley Jr, B., 2014. Venus. *Planets, Asteroids, Comets and The Solar System* 2, 127–148.
- Fegley Jr, B., Prinn, R.G., 1989. Estimation of the rate of volcanism on venus from reaction rate measurements. *Nature* 337, 55–58.
- Filiberto, J., Trang, D., Treiman, A.H., Gilmore, M.S., 2020. Present-day volcanism on venus as evidenced from weathering rates of olivine. *Science Advances* 6, eaax7445.
- Fischer-Gödde, M., Kleine, T., 2017. Ruthenium isotopic evidence for an inner Solar System origin of the late veneer. *Nature* 541, 525–527. doi:10.1038/nature21045.
- Forsyth, D., Uyeda, S., 1975. On the Relative Importance of the Driving Forces of Plate Motion*. *Geophysical Journal International* 43, 163–200. URL: <https://doi.org/10.1111/j.1365-246X.1975.tb00631.x>, doi:10.1111/j.1365-246X.1975.tb00631.x, arXiv:<https://academic.oup.com/gji/article-pdf/43/1/163/1576393/43-1-163.pdf>.
- Fossati, L., Erkaev, N., Lammer, H., Cubillos, P., Odert, P., Juvan, I., Kislyakova, K., Lendl, M., Kubyshkina, D., Bauer, S., 2017. Aeronomical

- constraints to the minimum mass and maximum radius of hot low-mass planets. *Astronomy & Astrophysics* 598, A90.
- Frey, H., Schultz, R.A., 1988. Large impact basins and the mega-impact origin for the crustal dichotomy on mars. *Geophysical Research Letters* 15, 229–232. URL: <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/GL015i003p00229>, doi:<https://doi.org/10.1029/GL015i003p00229>, arXiv:<https://agupubs.onlinelibrary.wiley.com/doi/pdf/10.1029/GL015i003p00229>.
- Frey, H.V., 2006. Impact constraints on the age and origin of the lowlands of mars. *Geophysical Research Letters* 33. URL: <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2005GL024484>, doi:<https://doi.org/10.1029/2005GL024484>, arXiv:<https://agupubs.onlinelibrary.wiley.com/doi/pdf/10.1029/2005GL024484>.
- Friedlingstein, P., Cox, P., Betts, R., Bopp, L., von Bloh, W., Brovkin, V., Cadule, P., Doney, S., Eby, M., Fung, I., et al., 2006. Climate–carbon cycle feedback analysis: results from the c4mip model intercomparison. *Journal of climate* 19, 3337–3353.
- Frost, D., Mann, U., Asahara, Y., Rubie, D., 2008. The redox state of the mantle during and just after core formation. *Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences* 366, 4315–4337. URL: <https://royalsocietypublishing.org/doi/abs/10.1098/rsta.2008.0147>, doi:10.1098/rsta.2008.0147, arXiv:<https://royalsocietypublishing.org/doi/pdf/10.1098/rsta.2008.0147>.
- Gaillard, F., Bernadou, F., Roskosz, M., Bouhifd, M.A., Marrocchi, Y., Iacono-Marziano, G., Moreira, M., Scaillet, B., Rogerie, G., 2022. Redox controls during magma ocean degassing. *Earth and Planetary Science Letters* 577, 117255. URL: <https://www.sciencedirect.com/science/article/pii/S0012821X21005112>, doi:<https://doi.org/10.1016/j.epsl.2021.117255>.
- Gaillard, F., Bouhifd, M.A., Füre, E., Malavergne, V., Marrocchi, Y., Noack, L., Ortenzi, G., Roskosz, M., Vulpius, S., 2021. The diverse planetary ingassing/outgassing paths produced over billions of years of magmatic activity. *Space Science Reviews* 217, 22. URL: <https://doi.org/10.1007/s11214-021-00802-1>, doi:10.1007/s11214-021-00802-1.

- Gaillard, F., Scaillet, B., 2014. A theoretical framework for volcanic degassing chemistry in a comparative planetology perspective and implications for planetary atmospheres. *Earth and Planetary Science Letters* 403, 307–316.
- Genda, H., Abe, Y., 2003. Survival of a proto-atmosphere through the stage of giant impacts: the mechanical aspects. *Icarus* 164, 149–162. URL: <https://www.sciencedirect.com/science/article/pii/S0019103503001015>, doi:[https://doi.org/10.1016/S0019-1035\(03\)00101-5](https://doi.org/10.1016/S0019-1035(03)00101-5).
- Genda, H., Abe, Y., 2005. Enhanced atmospheric loss on protoplanets at the giant impact phase in the presence of oceans. *Nature* 433, 842–844.
- Genda, H., Iizuka, T., Sasaki, T., Ueno, Y., Ikoma, M., 2017. Ejection of iron-bearing giant-impact fragments and the dynamical and geochemical influence of the fragment re-accretion. *Earth and Planetary Science Letters* 470, 87–95. doi:10.1016/j.epsl.2017.04.035.
- Gerya, T., 2022. Numerical modeling of subduction: State of the art and future directions. *Geosphere* 18, 503–561.
- Gillmann, C., Chassefière, E., Lagnonné, P., 2009a. A consistent picture of early hydrodynamic escape of venus atmosphere explaining present ne and ar isotopic ratios and low oxygen atmospheric content. *Earth and Planetary Science Letters* 286, 503–513.
- Gillmann, C., Golabek, G., Raymond, S., Schönbächler, M., Tackley, P., Dehant, V., Vinciane, D., 2020. Dry late accretion inferred from venus’s coupled atmosphere and internal evolution. *Nature Geoscience* 13, 1–5. doi:10.1038/s41561-020-0561-x.
- Gillmann, C., Golabek, G.J., Tackley, P.J., 2016. Effect of a single large impact on the coupled atmosphere-interior evolution of venus. *Icarus* 268, 295–312. URL: <https://www.sciencedirect.com/science/article/pii/S0019103515005795>, doi:<https://doi.org/10.1016/j.icarus.2015.12.024>.
- Gillmann, C., Lagnonné, P., Chassefière, E., Moreira, M., 2009b. The present-day atmosphere of mars: Where does it come from? *Earth and Planetary Science Letters* 277, 384–393. URL: <https://www>.

sciencedirect.com/science/article/pii/S0012821X08007097,
doi:<https://doi.org/10.1016/j.epsl.2008.10.033>.

- Gillmann, C., Lognonné, P., Moreira, M., 2011. Volatiles in the atmosphere of mars: The effects of volcanism and escape constrained by isotopic data. *Earth and Planetary Science Letters* 303, 299–309. URL: <https://www.sciencedirect.com/science/article/pii/S0012821X11000239>, doi:<https://doi.org/10.1016/j.epsl.2011.01.009>.
- Gillmann, C., Tackley, P., 2014. Atmosphere/mantle coupling and feedbacks on venus. *Journal of Geophysical Research: Planets* 119, 1189–1217. URL: <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1002/2013JE004505>, doi:<https://doi.org/10.1002/2013JE004505>, arXiv:<https://agupubs.onlinelibrary.wiley.com/doi/pdf/10.1002/2013JE004505>.
- Gillmann, C., Way, M.J., Avice, G., Breuer, D., Golabek, G.J., Höning, D., Krissansen-Totton, J., Lammer, H., O'Rourke, J.G., Persson, M., Plesa, A.C., Salvador, A., Scherf, M., Zolotov, M.Y., 2022. The long-term evolution of the atmosphere of venus: Processes and feedback mechanisms. *Space Science Reviews* 218, 56.
- Ginzburg, S., Schlichting, H.E., Sari, R., 2018. Core-powered mass-loss and the radius distribution of small exoplanets. *Monthly Notices of the Royal Astronomical Society* 476, 759–765.
- Golabek, G.J., Keller, T., Gerya, T.V., Zhu, G., Tackley, P.J., Connolly, J.A., 2011. Origin of the martian dichotomy and tharsis from a giant impact causing massive magmatism. *Icarus* 215, 346–357. URL: <https://www.sciencedirect.com/science/article/pii/S0019103511002211>, doi:<https://doi.org/10.1016/j.icarus.2011.06.012>.
- Goldblatt, C., Robinson, T.D., Zahnle, K.J., Crisp, D., 2013. Low simulated radiation limit for runaway greenhouse climates. *Nature Geoscience* 6, 661–667. URL: <https://doi.org/10.1038/ngeo1892>, doi:10.1038/ngeo1892.
- Greeley, R., Schneid, B.D., 1991. Magma generation on mars: Amounts, rates, and comparisons with earth, moon, and venus. *Science* 254, 996–998.

- Green, D.H., Hibberson, W.O., Rosenthal, A., Kovács, I., Yaxley, G.M., Falloon, T.J., Brink, F., 2014. Experimental study of the influence of water on melting and phase assemblages in the upper mantle. *Journal of Petrology* 55, 2067–2096.
- Grinspoon, D.H., 1993. Implications of the high d/h ratio for the sources of water in venus’ atmosphere. *Nature* 363, 428–431.
- Gronoff, G., Arras, P., Baraka, S., Bell, J.M., Cessateur, G., Cohen, O., Curry, S.M., Drake, J.J., Elrod, M., Erwin, J., et al., 2020. Atmospheric escape processes and planetary atmospheric evolution. *Journal of Geophysical Research: Space Physics* 125, e2019JA027639.
- Grott, M., Breuer, D., Laneuville, M., 2011. Thermo-chemical evolution and global contraction of mercury. *Earth and Planetary Science Letters* 307, 135–146.
- Gunell, H., Maggiolo, R., Nilsson, H., Wieser, G.S., Slapak, R., Lindkvist, J., Hamrin, M., De Keyser, J., 2018. Why an intrinsic magnetic field does not protect a planet against atmospheric escape. *Astronomy & Astrophysics* 614, L3.
- Halevy, I., Head III, J.W., 2014. Episodic warming of early mars by punctuated volcanism. *Nature Geoscience* 7, 865–868.
- Hamano, K., 2020. Steam atmospheres and magma oceans on planets. *Oxford Research Encyclopedia of Planetary Science*. URL: <https://oxfordre.com/planetaryscience/view/10.1093/acrefore/9780190647926.001.0001/acrefore-9780190647926-e-203>, doi:10.1093/acrefore/9780190647926.013.203.
- Hamano, K., Abe, Y., Genda, H., 2013. Emergence of two types of terrestrial planet on solidification of magma ocean. *Nature* 497, 607–610. URL: <https://doi.org/10.1038/nature12163>, doi:10.1038/nature12163.
- Hamano, K., Kawahara, H., Abe, Y., Onishi, M., Hashimoto, G.L., 2015. Lifetime and Spectral Evolution of a Magma Ocean with a Steam Atmosphere: Its Detectability by Future Direct Imaging. *The Astrophysical Journal* 806, 216. doi:10.1088/0004-637X/806/2/216.

- Hanmer, S., 2023. Basic structural geology of venus: A review of the gaps and how to bridge them. *Earth-Science Reviews* , 104331.
- Hartmann, W.K., 2005. Martian cratering 8: Isochron refinement and the chronology of mars. *Icarus* 174, 294–320.
- Hartmann, W.K., 2012. The terrestrial planets at the dawn of the. *The Century of Space Science* , 257.
- Hauber, E., Brož, P., Jagert, F., Jodłowski, P., Platz, T., 2011. Very recent and wide-spread basaltic volcanism on mars. *Geophysical Research Letters* 38.
- Hawkesworth, C.J., Cawood, P.A., Dhuime, B., 2020. The evolution of the continental crust and the onset of plate tectonics. *Frontiers in earth science* 8, 326.
- Höning, D., Baumeister, P., Grenfell, J.L., Tosi, N., Way, M.J., 2021. Early habitability and crustal decarbonation of a stagnant-lid venus. *Journal of Geophysical Research: Planets* 126, e2021JE006895. URL: <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2021JE006895>, doi:<https://doi.org/10.1029/2021JE006895>, arXiv:<https://agupubs.onlinelibrary.wiley.com/doi/pdf/10.1029/2021JE006895>. e2021JE006895 2021JE006895.
- Head, J., Wilson, L., Ivanov, M., Wordsworth, R., Kane, S., 2022. Detecting fossil atmospheres in venus-like exoplanets: Lessons from the current venus atmosphere and preserved geologic record. *LPI Contributions* 2687, 3026.
- Head, J.W., Campbell, D.B., Elachi, C., Guest, J.E., McKenzie, D.P., Saunders, R.S., Schaber, G.G., Schubert, G., 1991. Venus volcanism: Initial analysis from magellan data. *Science* 252, 276–288.
- Herd, C.D., Borg, L.E., Jones, J.H., Papike, J.J., 2002. Oxygen fugacity and geochemical variations in the martian basalts: Implications for martian basalt petrogenesis and the oxidation state of the upper mantle of mars. *Geochimica et Cosmochimica Acta* 66, 2025–2036.
- Herrick, R., Sharpton, V., Malin, M., Lyons, S., Feely, K., 1997. Morphology and morphometry of impact craters. *Venus II: Geology, Geophysics, Atmosphere, and Solar Wind Environment* , 1015.

- Herrick, R.R., Bjonnes, E.T., Carter, L.M., Gerya, T., Ghail, R.C., Gillmann, C., Gilmore, M., Hensley, S., Ivanov, M.A., Izenberg, N.R., et al., 2023. Resurfacing history and volcanic activity of venus. *Space Science Reviews* 219, 29.
- Herrick, R.R., Hensley, S., 2023. Surface changes observed on a venusian volcano during the magellan mission. *Science* 379, 1205–1208. URL: <https://www.science.org/doi/abs/10.1126/science.abm7735>, doi:10.1126/science.abm7735, arXiv:<https://www.science.org/doi/pdf/10.1126/science.abm7735>.
- Heubeck, C., Thomsen, T.B., Heredia, B.D., Zeh, A., Balling, P., 2023. The malolotsha klippe: Large-scale subhorizontal tectonics along the southern margin of the archean barberton greenstone belt, eswatini. *Tectonics* 42, e2022TC007359.
- Hirschmann, M.M., 2012. Magma ocean influence on early atmosphere mass and composition. *Earth and Planetary Science Letters* 341–344, 48–57. URL: <https://www.sciencedirect.com/science/article/pii/S0012821X12002993>, doi:<https://doi.org/10.1016/j.epsl.2012.06.015>.
- Hirschmann, M.M., 2022. Magma oceans, iron and chromium redox, and the origin of comparatively oxidized planetary mantles. *Geochimica et Cosmochimica Acta* 328, 221–241. URL: <https://www.sciencedirect.com/science/article/pii/S0016703722001715>, doi:<https://doi.org/10.1016/j.gca.2022.04.005>.
- Hirschmann, M.M., 2023. The deep earth oxygen cycle: mass balance considerations on the origin and evolution of mantle and surface oxidative reservoirs. *Earth and Planetary Science Letters* 619, 118311.
- Hirschmann, M.M., Withers, A.C., 2008. Ventilation of co₂ from a reduced mantle and consequences for the early martian greenhouse. *Earth and Planetary Science Letters* 270, 147–155.
- Hoffman, P.F., Kaufman, A.J., Halverson, G.P., Schrag, D.P., 1998. A neo-proterozoic snowball earth. *Science* 281, 1342–1346.
- Holland, H.D., 2002. Volcanic gases, black smokers, and the great oxidation event. *Geochimica et Cosmochimica acta* 66, 3811–3826.

- Hu, R., Kass, D.M., Ehlmann, B.L., Yung, Y.L., 2015. Tracing the fate of carbon and the atmospheric evolution of mars. *Nature communications* 6, 10003.
- Hudson, R., de Graaf, R., Strandoo Rodin, M., Ohno, A., Lane, N., McGlynn, S.E., Yamada, Y.M., Nakamura, R., Barge, L.M., Braun, D., et al., 2020. Co₂ reduction driven by a pH gradient. *Proceedings of the National Academy of Sciences* 117, 22873–22879.
- Hunten, D., Donahue, T., Walker, J., Kasting, J., 1989. Escape of atmospheres and loss of water. *Origin and evolution of planetary and satellite atmospheres*, 386–422.
- Hunten, D.M., 1973. The escape of light gases from planetary atmospheres. *Journal of Atmospheric Sciences* 30, 1481–1494.
- Hutchins, K.S., Jakosky, B.M., 1996. Evolution of martian atmospheric argon: Implications for sources of volatiles. *Journal of Geophysical Research: Planets* 101, 14933–14949.
- Ichikawa, H., Labrosse, S., Kurita, K., 2010. Direct numerical simulation of an iron rain in the magma ocean. *Journal of Geophysical Research: Solid Earth* 115. URL: <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2009JB006427>, doi:<https://doi.org/10.1029/2009JB006427>, arXiv:<https://agupubs.onlinelibrary.wiley.com/doi/pdf/10.1029/2009JB006427>.
- Ikoma, M., Elkins-Tanton, L., Hamano, K., Suckale, J., 2018. Water partitioning in planetary embryos and protoplanets with magma oceans. *Space Science Reviews* 214, 1–28.
- Ingersoll, A.P., 1969. The Runaway Greenhouse: A History of Water on Venus. *Journal of Atmospheric Sciences* 26, 1191 – 1198. URL: https://journals.ametsoc.org/view/journals/atsc/26/6/1520-0469_1969_026_1191_trgaho_2_0_co_2.xml, doi:10.1175/1520-0469(1969)026<1191:TRGAH0>2.0.CO;2.
- Irving, J.C., Lekić, V., Durán, C., Drilleau, M., Kim, D., Rivoldini, A., Khan, A., Samuel, H., Antonangeli, D., Banerdt, W.B., et al., 2023. First observations of core-transiting seismic phases on mars. *Proceedings of the National Academy of Sciences* 120, e2217090120.

- Jacobson, S.A., Rubie, D.C., Hernlund, J., Morbidelli, A., Nakajima, M., 2017. Formation, stratification, and mixing of the cores of earth and venus. *Earth and Planetary Science Letters* 474, 375–386. URL: <https://www.sciencedirect.com/science/article/pii/S0012821X17303333>, doi:<https://doi.org/10.1016/j.epsl.2017.06.023>.
- Jakosky, B.M., 2021. Atmospheric loss to space and the history of water on mars. *Annual Review of Earth and Planetary Sciences* 49, 71–93.
- Jeans, J., 1925. *The dynamical theory of gases*. University Press.
- Johnstone, C.P., 2020. Hydrodynamic escape of water vapor atmospheres near very active stars. *The Astrophysical Journal* 890, 79. URL: <https://doi.org/10.3847/1538-4357/ab6224>, doi:10.3847/1538-4357/ab6224.
- Johnstone, C.P., Lammer, H., Kislyakova, K.G., Scherf, M., Güdel, M., 2021. The young Sun’s XUV-activity as a constraint for lower CO₂-limits in the Earth’s Archean atmosphere. *Earth and Planetary Science Letters* 576, 117197. URL: <https://www.sciencedirect.com/science/article/pii/S0012821X21004520>, doi:<https://doi.org/10.1016/j.epsl.2021.117197>.
- Karato, S., 2015. Water in the evolution of the earth and other terrestrial planets. *Treatise on Geophysics* 9, 105–144.
- Karner, J., Papike, J., Shearer, C., McKay, G., Le, L., Burger, P., 2007. Valence state partitioning of cr and v between pyroxene-melt: Estimates of oxygen fugacity for martian basalt que 94201. *American Mineralogist* 92, 1238–1241.
- Kasting, J.F., 1988. Runaway and moist greenhouse atmospheres and the evolution of Earth and Venus. *Icarus* 74, 472–494. doi:10.1016/0019-1035(88)90116-9.
- Kasting, J.F., Pollack, J.B., 1983. Loss of water from Venus. I. Hydrodynamic escape of hydrogen. *Icarus* 53, 479–508. URL: <https://www.sciencedirect.com/science/article/pii/0019103583902129>, doi:[https://doi.org/10.1016/0019-1035\(83\)90212-9](https://doi.org/10.1016/0019-1035(83)90212-9).

- Katyal, N., Nikolaou, A., Godolt, M., Grenfell, J.L., Tosi, N., Schreier, F., Rauer, H., 2019. Evolution and spectral response of a steam atmosphere for early earth with a coupled climate–interior model. *The Astrophysical Journal* 875, 31. URL: <https://dx.doi.org/10.3847/1538-4357/ab0d85>, doi:10.3847/1538-4357/ab0d85.
- Kaula, W.M., 1999. Constraints on venus evolution from radiogenic argon. *Icarus* 139, 32–39.
- Keller, T., Tackley, P.J., 2009. Towards self-consistent modeling of the martian dichotomy: The influence of one-ridge convection on crustal thickness distribution. *Icarus* 202, 429–443. URL: <https://www.sciencedirect.com/science/article/pii/S0019103509001377>, doi:<https://doi.org/10.1016/j.icarus.2009.03.029>.
- Keppler, H., Golabek, G., 2019. Graphite floatation on a magma ocean and the fate of carbon during core formation. *Geochemical Perspectives Letters* 11, 12–17. URL: <https://www.geochemicalperspectivesletters.org/article1918>.
- Kiefer, W.S., Richards, M.A., Hager, B.H., 1986. A dynamic model of venus’s gravity field. *Geophys. Res. Lett.* 13, 14–17. doi:<https://doi.org/10.1029/GL013i001p00014>.
- King, S.D., 2018. Venus resurfacing constrained by geoid and topography. *Journal of Geophysical Research: Planets* 123, 1041–1060. URL: <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1002/2017JE005475>, doi:<https://doi.org/10.1002/2017JE005475>, arXiv:<https://agupubs.onlinelibrary.wiley.com/doi/pdf/10.1002/2017JE005475>.
- Kite, E.S., Williams, J.P., Lucas, A., Aharonson, O., 2014. Low palaeo-pressure of the martian atmosphere estimated from the size distribution of ancient craters. *Nature Geoscience* 7, 335–339.
- Kohlstedt, D., Evans, B., Mackwell, S., 1995. Strength of the lithosphere: Constraints imposed by laboratory experiments. *Journal of Geophysical Research: Solid Earth* 100, 17587–17602.
- Komabayasi, M., 1967. Discrete equilibrium temperatures of a hypothetical planet with the atmosphere and the hydrosphere of one component-two

- phase system under constant solar radiation. *Journal of the Meteorological Society of Japan. Ser. II* 45, 137–139. doi:10.2151/jmsj1965.45.1_137.
- Kopparapu, R.K., Ramirez, R., Kasting, J.F., Eymet, V., Robinson, T.D., Mahadevan, S., Terrien, R.C., Domagal-Goldman, S., Meadows, V., Deshpande, R., 2013. Habitable zones around main sequence stars: New estimates. *The Astrophysical Journal* 765, 131. URL: <https://doi.org/10.1088/0004-637x/765/2/131>, doi:10.1088/0004-637x/765/2/131.
- Kopparapu, R.K., Ramirez, R.M., Schottelkotte, J., Kasting, J.F., Domagal-Goldman, S., Eymet, V., 2014. Habitable zones around main sequence stars: Dependence of planetary mass. *The Astrophysical Journal* 787, L29. URL: <https://doi.org/10.1088/2041-8205/787/2/129>, doi:10.1088/2041-8205/787/2/129.
- Korenaga, J., 2013. Initiation and evolution of plate tectonics on earth: theories and observations. *Annual review of earth and planetary sciences* 41, 117–151.
- Kress, V.C., Carmichael, I.S.E., 1991. The compressibility of silicate liquids containing Fe_2O_3 and the effect of composition, temperature, oxygen fugacity and pressure on their redox states. *Contributions to Mineralogy and Petrology* 108, 82–92. URL: <https://doi.org/10.1007/BF00307328>, doi:10.1007/BF00307328.
- Krissansen-Totton, J., Fortney, J.J., Nimmo, F., 2021a. Was venus ever habitable? constraints from a coupled interior–atmosphere–redox evolution model. *The Planetary Science Journal* 2, 216.
- Krissansen-Totton, J., Kipp, M.A., Catling, D.C., 2021b. Carbon cycle inverse modeling suggests large changes in fractional organic burial are consistent with the carbon isotope record and may have contributed to the rise of oxygen. *Geobiology* 19, 342–363.
- Kruijer, T.S., Borg, L.E., Wimpenny, J., Sio, C.K., 2020. Onset of magma ocean solidification on mars inferred from mn-cr chronometry. *Earth and Planetary Science Letters* 542, 116315. URL: <https://www.sciencedirect.com/science/article/pii/S0012821X20302594>, doi:<https://doi.org/10.1016/j.epsl.2020.116315>.

- Kruijer, T.S., Kleine, T., Borg, L.E., Brennecka, G.A., Irving, A.J., Bischoff, A., Agee, C.B., 2017. The early differentiation of mars inferred from hf-w chronometry. *Earth and Planetary Science Letters* 474, 345–354. URL: <https://www.sciencedirect.com/science/article/pii/S0012821X17303710>, doi:<https://doi.org/10.1016/j.epsl.2017.06.047>.
- Kurokawa, H., Kurosawa, K., Usui, T., 2018. A lower limit of atmospheric pressure on early mars inferred from nitrogen and argon isotopic compositions. *Icarus* 299, 443–459.
- Kuwahara, H., Nakada, R., Kadoya, S., Yoshino, T., Irifune, T., 2023. Hadean mantle oxidation inferred from melting of peridotite under lower-mantle conditions. *Nature Geoscience* 16, 461–465. doi:10.1038/s41561-023-01169-4.
- Lammer, H., Chassefière, E., Karatekin, Ö., Morschhauser, A., Niles, P.B., Mousis, O., Odert, P., Möstl, U.V., Breuer, D., Dehant, V., et al., 2013. Outgassing history and escape of the martian atmosphere and water inventory. *Space Science Reviews* 174, 113–154.
- Lammer, H., Kulikov, Y.N., Lichtenegger, H.I., 2006. Thermospheric x-ray and euv heating by the young sun on early venus and mars. *Space Science Reviews* 122, 189–196.
- Lammer, H., Scherf, M., Kurokawa, H., Ueno, Y., Burger, C., Maindl, T., Johnstone, C.P., Leizinger, M., Benedikt, M., Fossati, L., et al., 2020. Loss and fractionation of noble gas isotopes and moderately volatile elements from planetary embryos and early venus, earth and mars. *Space Science Reviews* 216, 1–50.
- Lammer, H., Zerkle, A.L., Gebauer, S., Tosi, N., Noack, L., Scherf, M., Pilat-Lohinger, E., Güdel, M., Grenfell, J.L., Godolt, M., et al., 2018. Origin and evolution of the atmospheres of early venus, earth and mars. *The Astronomy and Astrophysics Review* 26, 1–72.
- Lécuyer, C., Simon, L., Guyot, F., 2000. Comparison of carbon, nitrogen and water budgets on venus and the earth. *Earth. Earth Planet Sci Lett* 181, 33–40.

- Leblanc, F., Chassefière, E., Gillmann, C., Breuer, D., 2012. Mars' atmospheric 40Ar: A tracer for past crustal erosion. *Icarus* 218, 561–570.
- Lebrun, T., Massol, H., Chassefière, E., Davaille, A., Marcq, E., Sarda, P., Leblanc, F., Brandeis, G., 2013. Thermal evolution of an early magma ocean in interaction with the atmosphere. *Journal of Geophysical Research: Planets* 118, 1155–1176. URL: <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1002/jgre.20068>, doi:<https://doi.org/10.1002/jgre.20068>, arXiv:<https://agupubs.onlinelibrary.wiley.com/doi/pdf/10.1002/jgre.20068>.
- Leconte, J., Wu, H., Menou, K., Murray, N., 2015. Asynchronous rotation of earth-mass planets in the habitable zone of lower-mass stars. *Science* 347, 632–635.
- Lenardic, A., Crowley, J.W., 2012. On the notion of well-defined tectonic regimes for terrestrial planets in this solar system and others. *The Astrophysical Journal* 755, 132.
- Lenardic, A., Jellinek, A., Moresi, L.N., 2008. A climate induced transition in the tectonic style of a terrestrial planet. *Earth and Planetary Science Letters* 271, 34–42. URL: <https://www.sciencedirect.com/science/article/pii/S0012821X08001878>, doi:<https://doi.org/10.1016/j.epsl.2008.03.031>.
- Lichtenberg, T., Bower, D.J., Hammond, M., Boukrouche, R., Sanan, P., Tsai, S.M., Pierrehumbert, R.T., 2021. Vertically resolved magma ocean–protoatmosphere evolution: H₂, h₂o, co₂, ch₄, co, o₂, and n₂ as primary absorbers. *Journal of Geophysical Research: Planets* 126, e2020JE006711. URL: <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2020JE006711>, doi:<https://doi.org/10.1029/2020JE006711>, arXiv:<https://agupubs.onlinelibrary.wiley.com/doi/pdf/10.1029/2020JE006711>. e2020JE006711 2020JE006711.
- Lichtenberg, T., Golabek, G.J., Burn, R., Meyer, M.R., Alibert, Y., Gerya, T.V., Mordasini, C., 2019. A water budget dichotomy of rocky proto-planets from ²⁶Al-heating. *Nature Astronomy* 3, 307–313. doi:[10.1038/s41550-018-0688-5](https://doi.org/10.1038/s41550-018-0688-5), arXiv:1902.04026.

- Liebske, C., Schmickler, B., Terasaki, H., Poe, B.T., Suzuki, A., Ichi Funakoshi, K., Ando, R., Rubie, D.C., 2005. Viscosity of peridotite liquid up to 13 gpa: Implications for magma ocean viscosities. *Earth and Planetary Science Letters* 240, 589–604. URL: <https://www.sciencedirect.com/science/article/pii/S0012821X05006825>, doi:<https://doi.org/10.1016/j.epsl.2005.10.004>.
- Lock, S.J., Stewart, S.T., Petaev, M.I., Leinhardt, Z., Mace, M.T., Jacobsen, S.B., Cuk, M., 2018. The origin of the moon within a terrestrial synestia. *Journal of Geophysical Research: Planets* 123, 910–951. URL: <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1002/2017JE005333>, doi:<https://doi.org/10.1002/2017JE005333>, arXiv:<https://agupubs.onlinelibrary.wiley.com/doi/pdf/10.1002/2017JE005333>.
- Lopez, I., Oyarzun, R., Marquez, A., Doblas-Reyes, F., Laurrieta, A., 1998. Progressive build up of CO₂ in the atmosphere of Venus through multiple volcanic resurfacing events. *Earth, Moon, and Planets* 81, 187–192.
- Lourenço, D.L., Rozel, A.B., 2023. Chapter 9 - the past and the future of plate tectonics and other tectonic regimes, in: Duarte, J.C. (Ed.), *Dynamics of Plate Tectonics and Mantle Convection*. Elsevier, pp. 181–196. URL: <https://www.sciencedirect.com/science/article/pii/B9780323857338000044>, doi:<https://doi.org/10.1016/B978-0-323-85733-8.00004-4>.
- Lourenço, D.L., Rozel, A.B., Ballmer, M.D., Tackley, P.J., 2020. Plutonic-squishy lid: A new global tectonic regime generated by intrusive magmatism on earth-like planets. *Geochemistry, Geophysics, Geosystems* 21, e2019GC008756. URL: <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2019GC008756>, doi:<https://doi.org/10.1029/2019GC008756>, arXiv:<https://agupubs.onlinelibrary.wiley.com/doi/pdf/10.1029/2019GC008756>.
- Lourenço, D.L., Rozel, A.B., Gerya, T., Tackley, P.J., 2018. Efficient cooling of rocky planets by intrusive magmatism. *Nature Geoscience* 11, 322–327.
- Lyons, T.W., Diamond, C.W., Planavsky, N.J., Reinhard, C.T., Li, C., 2021. Oxygenation, life, and the planetary system during earth’s middle history: An overview. *Astrobiology* 21, 906–923.

- Lyons, T.W., Reinhard, C.T., Planavsky, N.J., 2014. The rise of oxygen in earth's early ocean and atmosphere. *Nature* 506, 307–315.
- Maia, J.S., Wieczorek, M.A., 2022. Lithospheric structure of venusian crustal plateaus. *Journal of Geophysical Research: Planets* 127, e2021JE007004.
- Marchi, S., Canup, R.M., Walker, R.J., 2018. Heterogeneous delivery of silicate and metal to the earth by large planetesimals. *Nature Geoscience* 11, 77–81. doi:10.1038/s41561-017-0022-3.
- Marcq, E., Mills, F.P., Parkinson, C.D., Vandaele, A.C., 2017. Composition and chemistry of the neutral atmosphere of venus. *Space Science Reviews* 214, 10. URL: <https://doi.org/10.1007/s11214-017-0438-5>, doi:10.1007/s11214-017-0438-5.
- Marinova, M.M., Aharonson, O., Asphaug, E., 2008. Mega-impact formation of the Mars hemispheric dichotomy. *Nature* 453, 1216 – 1219.
- Martin, W., Russell, M.J., 2003. On the origins of cells: a hypothesis for the evolutionary transitions from abiotic geochemistry to chemoautotrophic prokaryotes, and from prokaryotes to nucleated cells. *Philosophical Transactions of the Royal Society of London. Series B: Biological Sciences* 358, 59–85.
- Martin, W.F., Bryant, D.A., Beatty, J.T., 2018. A physiological perspective on the origin and evolution of photosynthesis. *FEMS microbiology reviews* 42, 205–231.
- Marty, B., 2012. The origins and concentrations of water, carbon, nitrogen and noble gases on earth. *Earth and Planetary Science Letters* 313, 56–66.
- Maruyama, S., Ebisuzaki, T., 2023. A nuclear geyser origin of life: Life assembly plant–three-step model for the emergence of the first life on earth and cell dynamics for the coevolution of life's functions. *Conflicting Models for the Origin of Life* , 327–359.
- Massol, H., Davaille, A., Sarda, P., 2023. Early Formation of a Water Ocean as a Function of Initial CO₂ and H₂O Contents in a Solidifying Rocky Planet. *Journal of Geophysical Research: Planets* 128, e2023JE007848. URL: <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2023JE007848>, doi:<https://doi.org/10.1029/2023JE007848>,

arXiv:<https://agupubs.onlinelibrary.wiley.com/doi/pdf/10.1029/2023JE007848>.
e2023JE007848 2023JE007848.

- Massol, H., Hamano, K., Tian, F., Ikoma, M., Abe, Y., Chassefière, E., Davaille, A., Genda, H., Güdel, M., Hori, Y., Leblanc, F., Marcq, E., Sarda, P., Shematovich, V.I., Stökl, A., Lammer, H., 2016. Formation and evolution of protoatmospheres. *Space Science Reviews* 205, 153–211. URL: <https://doi.org/10.1007/s11214-016-0280-1>, doi:10.1007/s11214-016-0280-1.
- Maurice, M., Dasgupta, R., Hassanzadeh, P., 2023. Redox Evolution of the Crystallizing Terrestrial Magma Ocean and Its Influence on the Outgassed Atmosphere. *The Planetary Science Journal* 4, 31. doi:10.3847/PSJ/acb2ca.
- Mazrouei, S., Ghent, R.R., Bottke, W.F., Parker, A.H., Gernon, T.M., 2019. Earth and moon impact flux increased at the end of the paleozoic. *Science* 363, 253–257.
- McCord, T.B., 1968. The loss of retrograde satellites in the solar system. *Journal of Geophysical Research* 73, 1497–1500.
- McKay, C.P., Lorenz, R.D., Lunine, J.I., 1999. Analytic solutions for the antigreenhouse effect: Titan and the early earth. *Icarus* 137, 56–61. URL: <https://www.sciencedirect.com/science/article/pii/S0019103598960390>, doi:<https://doi.org/10.1006/icar.1998.6039>.
- Melezhik, V.A., Fallick, A.E., Martin, A.P., Condon, D.J., Kump, L.R., Brasier, A.T., Salminen, P.E., 2013. 7.3 the palaeoproterozoic perturbation of the global carbon cycle: the lomagundi-jatuli isotopic event. *Reading the Archive of Earth's Oxygenation: Volume 3: Global Events and the Fennoscandian Arctic Russia-Drilling Early Earth Project*, 1111–1150.
- Melosh, H., Vickery, A., 1989. Impact erosion of the primordial atmosphere of mars. *Nature* 338, 487–489.
- Mezger, K., Debaille, V., T., K., 2013. Core Formation and Mantle Differentiation on Mars. *Space Science Reviews* 174, 27–48.
- Moore, W., Webb, A., 2013. Heat-pipe earth. *Nature* 501, 501–505. doi:10.1038/nature12473.

- Moore, W.B., Simon, J.I., Webb, A.A.G., 2017. Heat-pipe planets. *Earth and Planetary Science Letters* 474, 13–19.
- Moresi, L., Solomatov, V., 1998. Mantle convection with a brittle lithosphere: thoughts on the global tectonic styles of the earth and venus. *Geophysical Journal International* 133, 669–682.
- Morgan, J.P., 1998. Thermal and rare gas evolution of the mantle. *Chemical Geology* 145, 431–445.
- Morschhauser, A., Grott, M., Breuer, D., 2011. Crustal recycling, mantle dehydration, and the thermal evolution of mars. *Icarus* 212, 541–558.
- Nakajima, M., Golabek, G.J., Wünnemann, K., Rubie, D.C., Burger, C., Melosh, H.J., Jacobson, S.A., Manske, L., Hull, S.D., 2021. Scaling laws for the geometry of an impact-induced magma ocean. *Earth and Planetary Science Letters* 568, 116983. URL: <https://www.sciencedirect.com/science/article/pii/S0012821X21002429>, doi:<https://doi.org/10.1016/j.epsl.2021.116983>.
- Nakajima, M., Stevenson, D.J., 2015. Melting and mixing states of the earth’s mantle after the moon-forming impact. *Earth and Planetary Science Letters* 427, 286–295. doi:<https://doi.org/10.1016/j.epsl.2015.06.023>.
- Nakajima, S., Hayashi, Y.Y., Abe, Y., 1992. A study on the “run-away greenhouse effect” with a one-dimensional radiative-convective equilibrium model. *Journal of Atmospheric Sciences* 49, 2256 – 2266. URL: https://journals.ametsoc.org/view/journals/atsc/49/23/1520-0469_1992_049_2256_asotge_2_0_co_2.xml, doi:[https://doi.org/10.1175/1520-0469\(1992\)049<2256:ASOTGE>2.0.CO;2](https://doi.org/10.1175/1520-0469(1992)049<2256:ASOTGE>2.0.CO;2).
- Nataf, H.C., Richter, F.M., 1982. Convection experiments in fluids with highly temperature-dependent viscosity and the thermal evolution of the planets. *Physics of the Earth and Planetary Interiors* 29, 320–329.
- Neukum, G., Jaumann, R., Hoffmann, H., Hauber, E., Head, J.W., Basilevsky, A.T., Ivanov, B.A., Werner, S.C., van Gasselt, S., Murray, J.B., McCord, T., Team, T.H.C.I., 2004. Recent and episodic volcanic and glacial activity on mars revealed by the high resolution stereo camera. *Nature* 432, 971–979.

- Nicklas, R.W., Puchtel, I.S., Ash, R.D., 2018. Redox state of the archean mantle: evidence from v partitioning in 3.5–2.4 ga komatiites. *Geochimica et Cosmochimica Acta* 222, 447–466.
- Niles, P.B., Boynton, W.V., Hoffman, J.H., Ming, D.W., Hamara, D., 2010. Stable isotope measurements of martian atmospheric CO_2 at the phoenix landing site. *Science* 329, 1334–1337. URL: <https://www.science.org/doi/abs/10.1126/science.1192863>, doi:10.1126/science.1192863, arXiv:<https://www.science.org/doi/pdf/10.1126/science.1192863>.
- Nimmo, F., 2002. Why does venus lack a magnetic field? *Geology* 30, 987–990.
- Nimmo, F., Hart, S.D., Korycansky, D.G., Agnor, C.B., 2008. Implications of an impact origin for the martian hemispheric dichotomy. *Nature* 453, 1220 – 1223.
- Nimmo, F., Tanaka, K., 2005. Early crustal evolution of mars. *Annual Review of Earth and Planetary Sciences* 33, 133–161. URL: <https://doi.org/10.1146/annurev.earth.33.092203.122637>, doi:10.1146/annurev.earth.33.092203.122637, arXiv:<https://doi.org/10.1146/annurev.earth.33.092203.122637>.
- Noack, L., Breuer, D., Spohn, T., 2012. Coupling the atmosphere with interior dynamics: Implications for the resurfacing of venus. *Icarus* 217, 484–498. URL: <https://www.sciencedirect.com/science/article/pii/S0019103511003435>, doi:<https://doi.org/10.1016/j.icarus.2011.08.026>. advances in Venus Science.
- Nutman, A.P., Friend, C.R., Bennett, V.C., 2002. Evidence for 3650–3600 ma assembly of the northern end of the itsaq gneiss complex, greenland: implication for early archaean tectonics. *Tectonics* 21, 5–1.
- Odert, P., Lammer, H., Erkaev, N., Nikolaou, A., Lichtenegger, H., Johnstone, C., Kislyakova, K., Leitzinger, M., Tosi, N., 2018. Escape and fractionation of volatiles and noble gases from mars-sized planetary embryos and growing protoplanets. *Icarus* 307, 327–346. URL: <https://www.sciencedirect.com/>

science/article/pii/S0019103517301677, doi:<https://doi.org/10.1016/j.icarus.2017.10.031>.

- O'Neill, C., Lenardic, A., Jellinek, A.M., Kiefer, W.S., 2007. Melt propagation and volcanism in mantle convection simulations, with applications for martian volcanic and atmospheric evolution. *Journal of Geophysical Research: Planets* 112. URL: <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2006JE002799>, doi:<https://doi.org/10.1029/2006JE002799>, arXiv:<https://agupubs.onlinelibrary.wiley.com/doi/pdf/10.1029/2006JE002799>.
- O'Neill, H.S., Berry, A.J., McCammon, C.C., Jayasuriya, K.D., Campbell, S.J., Foran, G., 2006. An experimental determination of the effect of pressure on the $\text{Fe}^{3+}/\Sigma\text{Fe}$ ratio of an anhydrous silicate melt to 3.0 GPa. *American Mineralogist* 91, 404–412. URL: <https://doi.org/10.2138/am.2005.1929>, doi:[doi:10.2138/am.2005.1929](https://doi.org/10.2138/am.2005.1929).
- O'Reilly, T.C., Davies, G.F., 1981. Magma transport of heat on io: A mechanism allowing a thick lithosphere. *Geophys. Res. Lett.* 8, 313–316.
- O'Rourke, J., Buz, J., Fu, R., Lillis, R., 2019. Detectability of remanent magnetism in the crust of venus. *Geophysical Research Letters* 46, 5768–5777.
- Ortenzi, G., Noack, L., Sohl, F., Guimond, C., Grenfell, J., Dorn, C., Schmidt, J., Vulpius, S., Katyal, N., Kitzmann, D., et al., 2020. Mantle redox state drives outgassing chemistry and atmospheric composition of rocky planets. *Scientific reports* 10, 10907.
- Owen, T., Maillard, J.P., De Bergh, C., Lutz, B.L., 1988. Deuterium on mars: The abundance of hdo and the value of d/h. *Science* 240, 1767–1767.
- O'Neill, C., Lenardic, A., Höink, T., Coltice, N., 2014. Mantle convection and outgassing on terrestrial planets. *Comparative climatology of terrestrial planets* 473.
- O'Neill, C., Lenardic, A., Weller, M., Moresi, L., Quenette, S., Zhang, S., 2016. A window for plate tectonics in terrestrial planet evolution? *Physics of the Earth and Planetary Interiors* 255, 80–92.

- O'Neill, C., Marchi, S., Zhang, S., Bottke, W., 2017. Impact-driven subduction on the hadean earth. *Nature Geoscience* 10, 793–797.
- O'Rourke, J., C., G., Tackley, P., 2018. Prospects for an ancient dynamo and modern crustal remanent magnetism on venus. *Earth Planet Sci Lett* 502, 46–56.
- Pahlevan, K., Schaefer, L., Hirschmann, M.M., 2019. Hydrogen isotopic evidence for early oxidation of silicate Earth. *Earth and Planetary Science Letters* 526, 115770. URL: <https://www.sciencedirect.com/science/article/pii/S0012821X19304625>, doi:<https://doi.org/10.1016/j.epsl.2019.115770>.
- Pepin, R.O., 1991. On the origin and early evolution of terrestrial planet atmospheres and meteoritic volatiles. *Icarus* 92, 2–79.
- Pepin, R.O., 1994. Evolution of the martian atmosphere. *Icarus* 111, 289–304.
- Persson, M., Futaana, Y., Fedorov, A., Nilsson, H., Hamrin, M., Barabash, S., 2018. H⁺/O⁺ escape rate ratio in the venus magnetotail and its dependence on the solar cycle. *Geophysical Research Letters* 45, 10–805.
- Persson, M., Futaana, Y., Ramstad, R., Masunaga, K., Nilsson, H., Hamrin, M., Fedorov, A., Barabash, S., 2020. The Venusian Atmospheric Oxygen Ion Escape: Extrapolation to the Early Solar System. *Journal of Geophysical Research: Planets* 125, e2019JE006336. URL: <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2019JE006336>, doi:<https://doi.org/10.1029/2019JE006336>, arXiv:<https://agupubs.onlinelibrary.wiley.com/doi/pdf/10.1029/2019JE006336>. e2019JE006336 2019JE006336.
- Peslier, A.H., Schönbächler, M., Busemann, H., Karato, S.I., 2017. Water in the earth's interior: Distribution and origin. *Space Science Reviews* 212, 743–810. URL: <https://doi.org/10.1007/s11214-017-0387-z>, doi:10.1007/s11214-017-0387-z.
- Pham, L., Karatekin, Ö., Dehant, V., 2011. Effects of impacts on the atmospheric evolution: Comparison between mars, earth, and venus. *Planetary and Space Science* 59, 1087–1092.

- Phillips, J.L., Russell, C.T., 1987. Upper limit on the intrinsic magnetic field of venus. *Journal of Geophysical Research: Space Physics* 92, 2253–2263. URL: <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/JA092iA03p02253>, doi:<https://doi.org/10.1029/JA092iA03p02253>, arXiv:<https://agupubs.onlinelibrary.wiley.com/doi/pdf/10.1029/JA092iA03p02253>.
- Phillips, R.J., Raubertas, R.F., Arvidson, R.E., Sarkar, I.C., Herrick, R.R., Izenberg, N., Grimm, R.E., 1992. Impact craters and venus resurfacing history. *Journal of Geophysical Research: Planets* 97, 15923–15948.
- Pierrehumbert, R.T., 2010. *Principles of Planetary Climate*. Cambridge University Press. doi:10.1017/CB09780511780783.
- Planavsky, N.J., Tarhan, L.G., Bellefroid, E.J., Evans, D.A., Reinhard, C.T., Love, G.D., Lyons, T.W., Polly, P., Head, J., Fox, D., 2015. Late proterozoic transitions in climate, oxygen, and tectonics, and the rise of complex life. *Earth-life transitions: paleobiology in the context of earth system evolution* 21, 1–36.
- Plattner, A., Simons, F.J., 2015. High-resolution local magnetic field models for the martian south pole from mars global surveyor data. *Journal of Geophysical Research: Planets* 120, 1543–1566. URL: <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1002/2015JE004869>, doi:<https://doi.org/10.1002/2015JE004869>, arXiv:<https://agupubs.onlinelibrary.wiley.com/doi/pdf/10.1002/2015JE004869>.
- Pollack, J.B., 1971. A nongrey calculation of the runaway greenhouse: Implications for Venus’ past and present. *Icarus* 14, 295–306. URL: <https://www.sciencedirect.com/science/article/pii/0019103571900017>, doi:[https://doi.org/10.1016/0019-1035\(71\)90001-7](https://doi.org/10.1016/0019-1035(71)90001-7).
- Pollack, J.B., Black, D.C., 1982. Noble gases in planetary atmospheres: Implications for the origin and evolution of atmospheres. *Icarus* 51, 169–198.
- Posner, E.S., Schmickler, B., Rubie, D.C., 2018. Self-diffusion and chemical diffusion in peridotite melt at high pressure and implications for magma ocean viscosities. *Chemical Geology* 502, 66–75. URL: <https://www.sciencedirect.com/science/article/pii/>

- S0009254118305175, doi:<https://doi.org/10.1016/j.chemgeo.2018.10.014>.
- Ramstad, R., Barabash, S., 2021. Do intrinsic magnetic fields protect planetary atmospheres from stellar winds? lessons from ion measurements at mars, venus, and earth. *Space Science Reviews* 217, 1–39.
- Rasool, S.I., De Bergh, C., 1970. The Runaway Greenhouse and the Accumulation of CO₂ in the Venus Atmosphere. *Nature* 226, 1037–1039. URL: <https://doi.org/10.1038/2261037a0>, doi:10.1038/2261037a0.
- Raymond, S.N., Schlichting, H.E., Hersant, F., Selsis, F., 2013. Dynamical and collisional constraints on a stochastic late veneer on the terrestrial planets. *Icarus* 226, 671–681. URL: <https://www.sciencedirect.com/science/article/pii/S0019103513002753>, doi:<https://doi.org/10.1016/j.icarus.2013.06.019>.
- Reese, C., Solomatov, V., 2006. Fluid dynamics of local martian magma oceans. *Icarus* 184, 102–120. URL: <https://www.sciencedirect.com/science/article/pii/S0019103506001308>, doi:<https://doi.org/10.1016/j.icarus.2006.04.008>.
- Reese, C., Solomatov, V., 2010. Early martian dynamo generation due to giant impacts. *Icarus* 207, 82–97. URL: <https://www.sciencedirect.com/science/article/pii/S0019103509004321>, doi:<https://doi.org/10.1016/j.icarus.2009.10.016>.
- Reese, C.C., Orth, C.P., Solomatov, V.S., 2010. Impact origin for the martian crustal dichotomy: Half emptied or half filled? *Journal of Geophysical Research: Planets* 115. URL: <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2009JE003506>, doi:<https://doi.org/10.1029/2009JE003506>, arXiv:<https://agupubs.onlinelibrary.wiley.com/doi/pdf/10.1029/2009JE003506>.
- Righter, K., Chabot, N.L., 2011. Moderately and slightly siderophile element constraints on the depth and extent of melting in early mars. *Meteoritics & Planetary Science* 46, 157–176. URL: <https://onlinelibrary.wiley.com/doi/abs/10.1111/j.1945-5100.2010.01140.x>, doi:<https://doi.org/10.1111/j.1945-5100.2010.01140.x>, arXiv:<https://onlinelibrary.wiley.com/doi/pdf/10.1111/j.1945-5100.2010.01140.x>.

- Righter, K., Sutton, S.R., Danielson, L., Pando, K., Newville, M., 2016. Redox variations in the inner solar system with new constraints from vanadium XANES in spinels. *American Mineralogist* 101, 1928–1942. URL: <https://doi.org/10.2138/am-2016-5638>, doi:10.2138/am-2016-5638, arXiv:<https://pubs.geoscienceworld.org/msa/ammin/article-pdf/101/9/1928/3597833/>
- Robbins, S.J., Hynek, B.M., 2012. A new global database of mars impact craters > 1 km: 1. database creation, properties, and parameters. *Journal of Geophysical Research: Planets* 117.
- Roberts, J.H., Zhong, S., 2006. Degree-1 convection in the martian mantle and the origin of the hemispheric dichotomy. *Journal of Geophysical Research: Planets* 111. URL: <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2005JE002668>, doi:<https://doi.org/10.1029/2005JE002668>, arXiv:<https://agupubs.onlinelibrary.wiley.com/doi/pdf/10.1029/2005JE002668>.
- Roble, R., Dickinson, R., 1989. How will changes in carbon dioxide and methane modify the mean structure of the mesosphere and thermosphere? *Geophysical Research Letters* 16, 1441–1444.
- Rolf, T., Weller, M., Gülcher, A., Byrne, P., O’Rourke, J.G., Herrick, R., Bjornes, E., Davaille, A., Ghail, R., Gillmann, C., Plesa, A.C., Smrekar, S., 2022. Dynamics and evolution of venus’ mantle through time. *Space Science Reviews* 218, 70.
- Rozel, A., Golabek, G.J., Jain, C., Tackley, P.J., T., G., 2017. Continental crust formation on early earth controlled by intrusive magmatism. *Nature* 545, 332–335.
- Rubie, D., Jacobson, S., Morbidelli, A., O’Brien, D., Young, E., de Vries, J., Nimmo, F., Palme, H., Frost, D., 2015a. Accretion and differentiation of the terrestrial planets with implications for the compositions of early-formed solar system bodies and accretion of water. *Icarus* 248, 89–108. URL: <https://www.sciencedirect.com/science/article/pii/S0019103514005545>, doi:<https://doi.org/10.1016/j.icarus.2014.10.015>.
- Rubie, D., Nimmo, F., Melosh, H., 2015b. Formation of the Earth’s Core, in: Schubert, G. (Ed.), *Treatise on Geophysics* (Second Edition). second edition ed.. Elsevier, Oxford, pp. 43–79. URL: <https://www.sciencedirect.com>

- com/science/article/pii/B9780444538024001548, doi:<https://doi.org/10.1016/B978-0-444-53802-4.00154-8>.
- Ruedas, T., Breuer, D., 2019. Dynamical effects of multiple impacts: Large impacts on a mars-like planet. *Physics of the Earth and Planetary Interiors* 287, 76–92.
- Russell, M.J., Hall, A., 1997. The emergence of life from iron monosulphide bubbles at a submarine hydrothermal redox and ph front. *Journal of the Geological Society* 154, 377–402.
- Sagan, C., Mullen, G., 1972. Earth and mars: Evolution of atmospheres and surface temperatures. *Science* 177, 52–56.
- Sakuraba, H., Kurokawa, H., Genda, H., 2019. Impact degassing and atmospheric erosion on venus, earth, and mars during the late accretion. *Icarus* 317, 48–58.
- Salvador, A., Massol, H., Davaille, A., Marcq, E., Sarda, P., Chassefière, E., 2017. The relative influence of H₂O and CO₂ on the primitive surface conditions and evolution of rocky planets. *Journal of Geophysical Research: Planets* 122, 1458–1486. URL: <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1002/2017JE005286>, doi:<https://doi.org/10.1002/2017JE005286>, arXiv:<https://agupubs.onlinelibrary.wiley.com/doi/pdf/10.1002/2017JE005286>.
- Salvador, A., Samuel, H., 2023. Convective outgassing efficiency in planetary magma oceans: Insights from computational fluid dynamics. *Icarus* 390, 115265. URL: <https://www.sciencedirect.com/science/article/pii/S0019103522003578>, doi:<https://doi.org/10.1016/j.icarus.2022.115265>.
- Sandu, C., Kiefer, W.S., 2012. Degassing history of mars and the lifespan of its magnetic dynamo. *Geophysical Research Letters* 39.
- Sasselov, D.D., Grotzinger, J.P., Sutherland, J.D., 2020. The origin of life as a planetary phenomenon. *Science Advances* 6, eaax3419.
- Sautter, V., Toplis, M.J., Wiens, R.C., Cousin, A., Fabre, C., Gasnault, O., Maurice, S., Forni, O., Lasue, J., Ollila, A., Bridges, J.C., Mangold, N., Mouélic, S.L., Fisk, M., Meslin, P.Y., Beck, P., Pinet, P., Deit, L.L., Rapin,

- W., Stolper, E.M., Newsom, H., Dyar, D., Lanza, N., Vaniman, D., Clegg, S., Wray, J.J., 2015. In situ evidence for continental crust on early mars. *Nature Geoscience* 8, 605–609.
- Saxena, P., Killen, R.M., Airapetian, V., Petro, N.E., Curran, N.M., Mandell, A.M., 2019. Was the sun a slow rotator? sodium and potassium constraints from the lunar regolith. *The Astrophysical Journal Letters* 876, L16.
- Schaefer, L., Fegley, B., 2017. Redox states of initial atmospheres outgassed on rocky planets and planetesimals. *The Astrophysical Journal* 843, 120.
- Schaefer, L., Wordsworth, R.D., Berta-Thompson, Z., Sasselov, D., 2016. Predictions of the atmospheric composition of gj 1132b. *The Astrophysical Journal* 829, 63. URL: <https://dx.doi.org/10.3847/0004-637X/829/2/63>, doi:10.3847/0004-637X/829/2/63.
- Scheller, E.L., Ehlmann, B.L., Hu, R., Adams, D.J., Yung, Y.L., 2021. Long-term drying of mars by sequestration of ocean-scale volumes of water in the crust. *Science* 372, 56–62. URL: <https://www.science.org/doi/abs/10.1126/science.abc7717>, doi:10.1126/science.abc7717, arXiv:<https://www.science.org/doi/pdf/10.1126/science.abc7717>.
- Schlichting, H.E., Mukhopadhyay, S., 2018. Atmosphere impact losses. *Space Science Reviews* 214, 1–31.
- Schlichting, H.E., Sari, R., Yalinewich, A., 2015. Atmospheric mass loss during planet formation: The importance of planetesimal impacts. *Icarus* 247, 81–94. URL: <https://www.sciencedirect.com/science/article/pii/S0019103514005284>, doi:<https://doi.org/10.1016/j.icarus.2014.09.053>.
- Schrag, D.P., Berner, R.A., Hoffman, P.F., Halverson, G.P., 2002. On the initiation of a snowball earth. *Geochemistry, Geophysics, Geosystems* 3, 1–21.
- Schreiber, U., Locker-Grütjen, O., Mayer, C., 2012. Hypothesis: Origin of life in deep-reaching tectonic faults. *Origins of Life and Evolution of Biospheres* 42, 47–54.
- Schubert, G., Turcotte, D.L., Olson, P., 2001. Mantle convection in the Earth and planets. Cambridge University Press.

- Selsis, F., Leconte, J., Turbet, M., Chaverot, G., Bolmont, E., 2023. A cool runaway greenhouse without surface magma ocean. *Nature* 620, 287–291. doi:<https://doi.org/10.1038/s41586-023-06258-3>.
- Sharp, Z.D., McCubbin, F.M., Shearer, C.K., 2013. A hydrogen-based oxidation mechanism relevant to planetary formation. *Earth and Planetary Science Letters* 380, 88–97. URL: <https://www.sciencedirect.com/science/article/pii/S0012821X1300441X>, doi:<https://doi.org/10.1016/j.epsl.2013.08.015>.
- Shearer, C., McKay, G., Papike, J., Karner, J., 2006. Valence state partitioning of vanadium between olivine-liquid: Estimates of the oxygen fugacity of y980459 and application to other olivine-phyric martian basalts. *American Mineralogist* 91, 1657–1663.
- Shellnutt, J.G., 2019. The curious case of the rock at venera 8. *Icarus* 321, 50–61.
- Shizgal, B.D., Arkos, G.G., 1996. Nonthermal escape of the atmospheres of venus, earth, and mars. *Reviews of Geophysics* 34, 483–505.
- Sleep, N.H., 2000a. Evolution of the mode of convection within terrestrial planets. *Journal of Geophysical Research: Planets* 105, 17563–17578. URL: <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2000JE001240>, doi:<https://doi.org/10.1029/2000JE001240>, arXiv:<https://agupubs.onlinelibrary.wiley.com/doi/pdf/10.1029/2000JE001240>.
- Sleep, N.H., 2000b. Evolution of the mode of convection within terrestrial planets. *Journal of Geophysical Research: Planets* 105, 17563–17578.
- Smrekar, S.E., Davaille, A., Sotin, C., 2018. Venus interior structure and dynamics. *Space Science Reviews* 214, 1–34.
- Smrekar, S.E., Stofan, E.R., Mueller, N., Treiman, A., Elkins-Tanton, L., Helbert, J., Piccioni, G., Drossart, P., 2010. Recent hotspot volcanism on venus from virtis emissivity data. *Science* 328, 605–608. URL: <https://www.science.org/doi/abs/10.1126/science.1186785>, doi:10.1126/science.1186785, arXiv:<https://www.science.org/doi/pdf/10.1126/science.1186785>.

Solomatov, V., 2007. Magma Oceans and Primordial Mantle Differentiation, in: Schubert, G. (Ed.), *Evolution of the Earth*. volume 9, pp. 91–119. doi:10.1016/B978-044452748-6.00141-3.

Solomatov, V.S., 1995. Scaling of temperature- and stress-dependent viscosity convection. *Physics of Fluids* 7, 266–274. URL: <https://doi.org/10.1063/1.868624>, doi:10.1063/1.868624, arXiv:https://pubs.aip.org/aip/pof/article-pdf/7/2/266/12739046/266_1_online.pdf

Solomatov, V.S., 2004. Initiation of subduction by small-scale convection. *Journal of Geophysical Research: Solid Earth* 109. URL: <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2003JB002628>, doi:<https://doi.org/10.1029/2003JB002628>, arXiv:<https://agupubs.onlinelibrary.wiley.com/doi/pdf/10.1029/2003JB002628>.

Solomatova, N.V., Caracas, R., 2021. Genesis of a co₂-rich and h₂o-depleted atmosphere from earth’s early global magma ocean. *Science advances* 7, eabj0406.

Sossi, P.A., Burnham, A.D., Badro, J., Lanzirotti, A., Newville, M., O’Neill, H.S., 2020. Redox state of earth’s magma ocean and its venus-like early atmosphere. *Science Advances* 6, eabd1387. URL: <https://www.science.org/doi/abs/10.1126/sciadv.abd1387>, doi:10.1126/sciadv.abd1387, arXiv:<https://www.science.org/doi/pdf/10.1126/sciadv.abd1387>.

Spray, J., 2018. Earth impact database: Planetary and space science centre, university of new brunswick.

Stagno, V., Aulbach, S., 2021. Redox processes before, during, and after earth’s accretion affecting the deep carbon cycle. *Magma Redox Geochemistry* , 19–32.

Stähler, S.C., Mittelholz, A., Perrin, C., Kawamura, T., Kim, D., Knapmeyer, M., Zenhäusern, G., Clinton, J., Giardini, D., Lognonné, P., et al., 2022. Tectonics of cerberus fossae unveiled by marsquakes. *Nature Astronomy* 6, 1376–1386.

Steele, S.C., Fu, R.R., Volk, M.W., North, T.L., Brenner, A.R., Muxworthy, A.R., Collins, G.S., Davison, T.M., 2023. Paleomagnetic evidence for a long-lived, potentially reversing martian dynamo at 3.9

- ga. *Science Advances* 9, eade9071. URL: <https://www.science.org/doi/abs/10.1126/sciadv.ade9071>, doi:10.1126/sciadv.ade9071, arXiv:<https://www.science.org/doi/pdf/10.1126/sciadv.ade9071>.
- Stein, C., Schmalzl, J., Hansen, U., 2004. The effect of rheological parameters on plate behaviour in a self-consistent model of mantle convection. *Phys. Earth. Plan. Int.* 142, 225–255.
- Stern, R.J., 2005. Evidence from ophiolites, blueschists, and ultrahigh-pressure metamorphic terranes that the modern episode of subduction tectonics began in neoproterozoic time. *Geology* 33, 557–560.
- Stern, R.J., 2018. The evolution of plate tectonics. *Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences* 376, 20170406.
- Stern, R.J., Gerya, T., Tackley, P.J., 2018. Stagnant lid tectonics: Perspectives from silicate planets, dwarf planets, large moons, and large asteroids. *Geoscience Frontiers* 9, 103–119.
- Stevenson, D.J., 2003. Planetary magnetic fields. *Earth and Planetary Science Letters* 208, 1–11. URL: <https://www.sciencedirect.com/science/article/pii/S0012821X02011263>, doi:[https://doi.org/10.1016/S0012-821X\(02\)01126-3](https://doi.org/10.1016/S0012-821X(02)01126-3).
- Tackley, P.J., 2000. Self consistent generation of tectonic plates in time-dependent, three dimensional mantle convection simulations, part 2: Strain weakening and asthenosphere. *G3* 1.
- Tarduno, J.A., Blackman, E.G., Mamajek, E.E., 2014. Detecting the oldest geodynamo and attendant shielding from the solar wind: Implications for habitability. *Physics of the Earth and Planetary Interiors* 233, 68–87.
- Tarduno, J.A., Cottrell, R.D., Davis, W.J., Nimmo, F., Bono, R.K., 2015. A hadean to paleoarchean geodynamo recorded by single zircon crystals. *Science* 349, 521–524.
- Tian, F., Kasting, J.F., Solomon, S.C., 2009. Thermal escape of carbon from the early martian atmosphere. *Geophysical Research Letters* 36. URL: <https://agupubs.onlinelibrary.wiley.com/doi/abs/>

10.1029/2008GL036513, doi:<https://doi.org/10.1029/2008GL036513>,
arXiv:<https://agupubs.onlinelibrary.wiley.com/doi/pdf/10.1029/2008GL036513>.

Trail, D., Watson, E.B., Tailby, N.D., 2011. The oxidation state of hadean magmas and implications for early earth's atmosphere. *Nature* 480, 79–82. URL: <https://doi.org/10.1038/nature10655>, doi:10.1038/nature10655.

Tucker, J.M., Mukhopadhyay, S., 2014. Evidence for multiple magma ocean outgassing and atmospheric loss episodes from mantle noble gases. *Earth and Planetary Science Letters* 393, 254–265. URL: <https://www.sciencedirect.com/science/article/pii/S0012821X1400137X>, doi:<https://doi.org/10.1016/j.epsl.2014.02.050>.

Turbet, M., Bolmont, E., Chaverot, G., Ehrenreich, D., Leconte, J., Marcq, E., 2021. Day–night cloud asymmetry prevents early oceans on Venus but not on Earth. *Nature* 598, 276–280. URL: <https://doi.org/10.1038/s41586-021-03873-w>, doi:10.1038/s41586-021-03873-w.

Turbet, M., Ehrenreich, D., Lovis, C., Bolmont, E., Fauchez, T., 2019. The runaway greenhouse radius inflation effect-an observational diagnostic to probe water on earth-sized planets and test the habitable zone concept. *Astronomy & Astrophysics* 628, A12.

Turbet, M., Gillmann, C., Forget, F., Baudin, B., Palumbo, A., Head, J., Karatekin, O., 2020. The environmental effects of very large bolide impacts on early mars explored with a hierarchy of numerical models. *Icarus* 335, 113419.

Turcotte, D.L., 1993. An episodic hypothesis for venusian tectonics. *Journal of Geophysical Research: Planets* 98, 17061–17068.

Turner, S., Wilde, S., Wörner, G., Schaefer, B., Lai, Y.J., 2020. An andesitic source for jack hills zircon supports onset of plate tectonics in the hadean. *Nature Communications* 11, 1241.

Van Kranendonk, M.J., Baumgartner, R., Djokic, T., Ota, T., Steller, L., Garbe, U., Nakamura, E., 2021. Elements for the origin of life on land: a deep-time perspective from the pilbara craton of western australia. *Astrobiology* 21, 39–59.

- Van Thienen, P., Benzerara, K., Breuer, D., Gillmann, C., Labrosse, S., Lognonné, P., Spohn, T., 2007. Water, life, and planetary geodynamical evolution. *Geology and Habitability of Terrestrial Planets*, 167–203.
- Wang, Z., Liu, Y., Zong, K., Lin, J., Kusky, T.M., 2020. Mantle degassing related to changing redox and thermal conditions during the precambrian supercontinent cycle. *Precambrian Research* 350, 105895.
- Warren, A.O., Kite, E.S., 2023. Narrow range of early habitable venus scenarios permitted by modeling of oxygen loss and radiogenic argon degassing. *Proceedings of the National Academy of Sciences* 120, e2209751120.
- Way, M., Ernst, R.E., Scargle, J.D., 2022a. Large-scale volcanism and the heat death of terrestrial worlds. *The Planetary Science Journal* 3, 92.
- Way, M., Ostberg, C.M., Foley, B.J., Gillmann, C., Höning, D., Lammer, H., O'Rourke, J.G., Persson, M., Plesa, A.C., Salvador, A., et al., 2022b. Searching for synergies between venus and exoplanetary worlds, in: *AGU Fall Meeting Abstracts*, pp. P56B–01.
- Way, M.J., Del Genio, A.D., 2020. Venusian habitable climate scenarios: Modeling venus through time and applications to slowly rotating venus-like exoplanets. *Journal of Geophysical Research: Planets* 125, e2019JE006276. URL: <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2019JE006276>, doi:<https://doi.org/10.1029/2019JE006276>, arXiv:<https://agupubs.onlinelibrary.wiley.com/doi/pdf/10.1029/2019JE006276>. e2019JE006276 10.1029/2019JE006276.
- Way, M.J., Del Genio, A.D., Kiang, N.Y., Sohl, L.E., Grinspoon, D.H., Aleinov, I., Kelley, M., Clune, T., 2016. Was venus the first habitable world of our solar system? *Geophysical research letters* 43, 8376–8383.
- Wedepohl, K.H., 1995. The composition of the continental crust. *Geochimica et Cosmochimica Acta* 59, 1217–1232. URL: <https://www.sciencedirect.com/science/article/pii/0016703795000382>, doi:[https://doi.org/10.1016/0016-7037\(95\)00038-2](https://doi.org/10.1016/0016-7037(95)00038-2).
- Weiss, M.C., Sousa, F.L., Mrnjavac, N., Neukirchen, S., Roettger, M., Nelson-Sathi, S., Martin, W.F., 2016. The physiology and habitat of the last universal common ancestor. *Nature microbiology* 1, 1–8.

- Weller, M., Evans, A., Ibarra, D., Johnson, A., Kukla, T., 2022. Atmospheric evidence of early plate tectonics on venus. *LPI Contrib* 2678, 2328.
- Weller, M.B., Kiefer, W.S., 2020. The physics of changing tectonic regimes: Implications for the temporal evolution of mantle convection and the thermal history of venus. *Journal of Geophysical Research: Planets* 125, e2019JE005960.
- Werner, S.C., 2009. The global martian volcanic evolutionary history. *Icarus* 201, 44–68. URL: <https://www.sciencedirect.com/science/article/pii/S0019103508004478>, doi:<https://doi.org/10.1016/j.icarus.2008.12.019>.
- Wessel, P., Müller, R., 2015. 6.02 - plate tectonics, in: Schubert, G. (Ed.), *Treatise on Geophysics* (Second Edition). second edition ed.. Elsevier, Oxford, pp. 45–93. URL: <https://www.sciencedirect.com/science/article/pii/B9780444538024001111>, doi:<https://doi.org/10.1016/B978-0-444-53802-4.00111-1>.
- Westall, F., Hickman-Lewis, K., Hinman, N., Gautret, P., Campbell, K.A., Bréhéret, J.G., Foucher, F., Hubert, A., Sorieul, S., Dass, A.V., et al., 2018. A hydrothermal-sedimentary context for the origin of life. *Astrobiology* 18, 259–293.
- Westall, F., Höning, D., Avice, G., Gentry, D., Gerya, T., Gillmann, C., Izenberg, N., Way, M.J., Wilson, C., 2023. The habitability of venus. *Space Science Reviews* 219, 17.
- Wetherill, G.W., 1976. The role of large bodies in the formation of the earth and moon. *Lunar and Planetary Science Conference Proceedings* 3, 3245–3257.
- Wieczorek, M.A., 2015. Gravity and topography of the terrestrial planets. *Treatise on geophysics* 10, 165–206.
- Wieler, R., 2002. Noble gases in the solar system. *Reviews in Mineralogy and geochemistry* 47, 21–70.
- Wilhelms, D.E., Squyres, S.W., 1984. The martian hemispheric dichotomy may be due to a giant impact. *Nature* 309, 138–140.

- Wordsworth, R., 2016. Atmospheric nitrogen evolution on earth and venus. *Earth and Planetary Science Letters* 447, 103–111. URL: <https://www.sciencedirect.com/science/article/pii/S0012821X16301583>, doi:<https://doi.org/10.1016/j.epsl.2016.04.002>.
- Wordsworth, R., Pierrehumbert, R., 2014. Abiotic oxygen-dominated atmospheres on terrestrial habitable zone planets. *The Astrophysical Journal* 785, L20. URL: <https://doi.org/10.1088/2041-8205/785/2/L20>, doi:10.1088/2041-8205/785/2/L20.
- Wordsworth, R.D., Kerber, L., Pierrehumbert, R.T., Forget, F., Head, J.W., 2015. Comparison of “warm and wet” and “cold and icy” scenarios for early mars in a 3-d climate model. *Journal of Geophysical Research: Planets* 120, 1201–1219.
- Xie, S., Tackley, P.J., 2004. Evolution of helium and argon isotopes in a convecting mantle. *Physics of the Earth and Planetary Interiors* 146, 417–439.
- Yang, J., Cowan, N.B., Abbot, D.S., 2013. Stabilizing cloud feedback dramatically expands the habitable zone of tidally locked planets. *The Astrophysical Journal Letters* 771, L45.
- Yoder, C., Ahrens, T., 1995. *Global earth physics: A handbook of physical constants*. Washington DC: American Geophysical Union .
- Zahnle, K., Arndt, N., Cockell, C., Halliday, A., Nisbet, E., Selsis, F., Sleep, N.H., 2007. Emergence of a habitable planet. *Space Science Reviews* 129, 35–78.
- Zahnle, K.J., Gacesa, M., Catling, D.C., 2019. Strange messenger: A new history of hydrogen on earth, as told by xenon. *Geochimica et Cosmochimica Acta* 244, 56–85.
- Zahnle, K.J., Kasting, J.F., 2023. Elemental and isotopic fractionation as fossils of water escape from venus. *Geochimica et Cosmochimica Acta* 361, 228–244.
- Zhang, H., Hirschmann, M., Cottrell, E., Withers, A., 2017. Effect of pressure on $\text{Fe}^{3+}/\Sigma\text{Fe}$ ratio in a mafic magma and consequences for

magma ocean redox gradients. *Geochimica et Cosmochimica Acta* 204, 83–103. URL: <https://www.sciencedirect.com/science/article/pii/S0016703717300376>, doi:<https://doi.org/10.1016/j.gca.2017.01.023>.

Zhong, S., 2009. Migration of tharsis volcanism on mars caused by differential rotation of the lithosphere. *Nature Geoscience* 2, 19–23.

Zhong, S., Zuber, M.T., 2001. Degree-1 mantle convection and the crustal dichotomy on mars. *Earth and Planetary Science Letters* 189, 75–84. URL: <https://www.sciencedirect.com/science/article/pii/S0012821X01003454>, doi:[https://doi.org/10.1016/S0012-821X\(01\)00345-4](https://doi.org/10.1016/S0012-821X(01)00345-4).