# A q-ANALOG OF THE STIRLING-EULERIAN POLYNOMIALS

YAO DONG, ZHICONG LIN, AND QIONGQIONG PAN

ABSTRACT. In 1974, Carlitz and Scoville introduced the Stirling-Eulerian polynomial  $A_n(x, y | \alpha, \beta)$  as the enumerator of permutations by descents, ascents, left-to-right maxima and right-to-left maxima. Recently, Ji considered a refinement of  $A_n(x, y | \alpha, \beta)$ , denoted  $P_n(u_1, u_2, u_3, u_4 | \alpha, \beta)$ , which is the enumerator of permutations by valleys, peaks, double ascents, double descents, left-to-right maxima and right-to-left maxima. Using Chen's context-free grammar calculus, Ji proved a formula for the generating function of  $P_n(u_1, u_2, u_3, u_4 | \alpha, \beta)$ , generalizing the work of Carlitz and Scoville. Ji's formula has many nice consequences, one of which is an intriguing  $\gamma$ -positivity expansion for  $A_n(x, y | \alpha, \beta)$ . In this paper, we prove a q-analog of Ji's formula by using Gessel's q-compositional formula and provide a combinatorial approach to her  $\gamma$ -positivity expansion of  $A_n(x, y | \alpha, \beta)$ .

### 1. INTRODUCTION

Using Chen's context-free grammar [5], Ji [10] recently proved a formula for the generating function of permutations by valleys, peaks, double ascents, double descents, left-to-right maxima and right-to-left maxima. The main objective of this paper is to prove a q-analog of Ji's formula by using Gessel's q-compositional formula [9]. We need some further notations and definitions before we can state our q-analog of Ji's formula.

Let  $\mathfrak{S}_n$  be the set of permutations of  $[n] := \{1, \ldots, n\}$ . An index  $i \in [n-1]$  is a descent (resp., ascent) of  $\sigma = \sigma_1 \cdots \sigma_n \in \mathfrak{S}_n$  if  $\sigma_i > \sigma_{i+1}$  (resp.,  $\sigma_i < \sigma_{i+1}$ ). The Eulerian polynomials  $A_n(x)$  can be described as the enumerator of  $\mathfrak{S}_n$  by descents or ascents, that is

$$A_n(x) := \sum_{\sigma \in \mathfrak{S}_n} x^{\operatorname{des}(\sigma)} = \sum_{\sigma \in \mathfrak{S}_n} x^{\operatorname{asc}(\sigma)}, \qquad (1.1)$$

where des( $\sigma$ ) (resp., asc( $\sigma$ )) denotes the number of descents (resp., ascents) of  $\sigma$ . There is a well-known formula (see [14, p. 13]) for the exponential generating function for  $A_n(x)$ :

$$1 + \sum_{n \ge 1} x A_n(x) \frac{t^n}{n!} = \frac{1 - x}{1 - xe^{(1 - x)t}}.$$
(1.2)

A permutation statistic whose generating function equals (1.1) is called *Eulerian*. Another classical Eulerian statistic (see [17, p. 23]) is the number of excedances "exc", where

$$exc(\sigma) := \#\{i \in [n-1] : \sigma_i > i\}.$$

Another class of fundamental permutation statistics is the so-called Stirling statistics. A *left-to-right maximum* (resp., *left-to-right minimum*) of  $\sigma \in \mathfrak{S}_n$  is an element  $\sigma_i$  such that

Date: April 16, 2024.

Key words and phrases. Stirling-Eulerian polynomials;  $\gamma$ -positivity; inversions; descents; left-to-right minima; right-to-left minima.

#### Y. DONG, Z. LIN, AND Q.Q. PAN

 $\sigma_j < \sigma_i$  (resp.,  $\sigma_j > \sigma_i$ ) for every j < i. Let  $\operatorname{Ima}(\sigma)$  and  $\operatorname{Imi}(\sigma)$  denote the numbers of leftto-right maxima and left-to-right minima of  $\sigma$ , respectively. Similarly, we can define the *right-to-left maximum* (resp., *right-to-left minimum*) of  $\sigma$  and denote  $\operatorname{rma}(\sigma)$  (resp.,  $\operatorname{rmi}(\sigma)$ ) the number of right-to-left maxima (resp., right-to-left minima) of  $\sigma$ . It is well known (see [17, p. 20]) that the unsigned *Stirling numbers of the first kind* count permutations by their number of left-to-right maxima, namely

$$(x)_n := x(x+1)(x+2)\cdots(x+n-1) = \sum_{\sigma \in \mathfrak{S}_n} x^{\operatorname{Ima}(\sigma)}.$$

Therefore, each statistic whose distribution over  $\mathfrak{S}_n$  gives  $(x)_n$  is called *Stirling*. So "lmi", "rma" and "rmi" are all Stirling statistics by the symmetry of permutations and according to *Foata's first fundamental transformation* (see [17, Prop. 1.3.1]), the statistic "cyc" is also Stirling, where  $\operatorname{cyc}(\sigma)$  denotes the number of cycles of  $\sigma$ .

Carlitz and Scoville [4] considered the following polynomials involving Stirling-Eulerian Statistics, which we refer to as the *Stirling-Eulerian polynomials*<sup>1</sup>:

$$A_n(x, y | \alpha, \beta) = \sum_{\sigma \in \mathfrak{S}_n} x^{\operatorname{asc}(\sigma)} y^{\operatorname{des}(\sigma)} \alpha^{\operatorname{lma}(\sigma) - 1} \beta^{\operatorname{rma}(\sigma) - 1}$$
$$= \sum_{\sigma \in \mathfrak{S}_n} x^{\operatorname{des}(\sigma)} y^{\operatorname{asc}(\sigma)} \alpha^{\operatorname{lmi}(\sigma) - 1} \beta^{\operatorname{rmi}(\sigma) - 1}.$$

Here the second interpretation follows from the complement

$$\sigma = \sigma_1 \sigma_2 \cdots \sigma_n \mapsto \sigma^c = (n+1-\sigma_1)(n+1-\sigma_2) \cdots (n+1-\sigma_n)$$

of permutations. In [4, Theorem 9], they obtained the following generating function formula for  $A_n(x, y | \alpha, \beta)$ :

$$\sum_{n \ge 0} A_{n+1}(x, y | \alpha, \beta) \frac{t^n}{n!} = (1 + xF(x, y; t))^{\alpha} (1 + yF(x, y; t))^{\beta},$$
(1.3)

where

$$F(x,y;t) := \frac{e^{xt} - e^{yt}}{xe^{yt} - ye^{xt}}.$$
(1.4)

Recently, Ji [10] refined the Carlitz–Scoville formula (1.3) by using the following refinements of descents and ascents.

**Definition 1.** For a permutation  $\sigma \in \mathfrak{S}_n$  with some specified boundary condition (i.e., specified values of  $\sigma_0$  and  $\sigma_{n+1}$ ), the index  $i \in [n]$  is

- a **peak** if  $\sigma_{i-1} < \sigma_i > \sigma_{i+1}$ ;
- *a* valley if  $\sigma_{i-1} > \sigma_i < \sigma_{i+1}$ ;
- a double ascent if  $\sigma_{i-1} < \sigma_i < \sigma_{i+1}$ ;
- a double descent if  $\sigma_{i-1} > \sigma_i > \sigma_{i+1}$ .

<sup>&</sup>lt;sup>1</sup>These polynomials are called the  $(\alpha, \beta)$ -Eulerian polynomials by Ji [10].

Let  $M(\sigma)$ ,  $V(\sigma)$ ,  $da(\sigma)$  and  $dd(\sigma)$  (resp.,  $\widetilde{M}(\sigma)$ ,  $\widetilde{V}(\sigma)$ ,  $d\widetilde{a}(\sigma)$  and  $d\widetilde{d}(\sigma)$ ) denote respectively the number of peaks, valleys, double ascents and double descents in  $\sigma$  with boundary condition  $\sigma_0 = \sigma_{n+1} = 0$  (resp.,  $\sigma_0 = \sigma_{n+1} = +\infty$ ). Note that  $M(\sigma) = V(\sigma) + 1$  and

$$\sum_{\sigma \in \mathfrak{S}_n} u_1^{\mathrm{V}(\sigma)} u_2^{\mathrm{M}(\sigma)} u_3^{\mathrm{da}(\sigma)} u_4^{\mathrm{dd}(\sigma)} = \sum_{\sigma \in \mathfrak{S}_n} u_1^{\widetilde{\mathrm{M}}(\sigma)} u_2^{\widetilde{\mathrm{V}}(\sigma)} u_3^{\widetilde{\mathrm{dd}}(\sigma)} u_4^{\widetilde{\mathrm{da}}(\sigma)}$$

Ji [10] introduced a refinement of  $A_n(x, y | \alpha, \beta)$  as

$$P_n(u_1, u_2, u_3, u_4 | \alpha, \beta) := \sum_{\sigma \in \mathfrak{S}_n} u_1^{\mathcal{V}(\sigma)} u_2^{\mathcal{M}(\sigma) - 1} u_3^{\mathrm{da}(\sigma)} u_4^{\mathrm{dd}(\sigma)} \alpha^{\mathrm{lma}(\sigma) - 1} \beta^{\mathrm{rma}(\sigma) - 1}$$

and proved the following generalization of (1.3) using Chen's context-free grammar.

**Theorem 1.1** (Ji [10, Thm. 1.4]). We have

$$\sum_{n\geq 0} P_{n+1}(u_1, u_2, u_3, u_4 | \alpha, \beta) \frac{t^n}{n!} = (1 + yF(x, y; t))^{\frac{\alpha+\beta}{2}} (1 + xF(x, y; t))^{\frac{\alpha+\beta}{2}} e^{\frac{1}{2}(\beta-\alpha)(u_4-u_3)t},$$

where  $x + y = u_3 + u_4$  and  $xy = u_1u_2$ .

When  $\alpha = \beta = 1$ , Theorem 1.1 reduces to another generating function formula of Carlitz and Scoville [4], which is equivalent to

$$\sum_{n \ge 1} \sum_{\sigma \in \mathfrak{S}_n} u_1^{\mathcal{V}(\sigma)} u_2^{\mathcal{M}(\sigma)-1} u_3^{\mathrm{da}(\sigma)} u_4^{\mathrm{dd}(\sigma)} \frac{t^n}{n!} = F(x, y; t),$$
(1.5)

where  $x + y = u_3 + u_4$  and  $xy = u_1u_2$ . A grammatical proof of (1.5) was provided by Fu [7].

Let us enter the world of q-series. For an integer  $n \ge 0$ , let  $[n]_q! := \prod_{i=1}^n (1+q+\cdots+q^{i-1})$  be the q-analog of n!. Introduce a classical q-analog of the exponential function by

$$\exp_q(t) := \sum_{n \ge 0} \frac{t^n}{[n]_q!}.$$

Stanley (see [14, Sec. 6.4]) proved the following q-analog of (1.2)

$$1 + \sum_{n \ge 1} \sum_{\sigma \in \mathfrak{S}_n} x^{\operatorname{des}(\sigma) + 1} q^{\operatorname{inv}(\sigma)} \frac{t^n}{[n]_q!} = \frac{1 - x}{1 - x \exp_q(t(1 - x))},$$
(1.6)

where  $\operatorname{inv}(\sigma) := |\{(i,j) \in [n]^2 : i < j, \sigma_i > \sigma_j\}|$  is the number of inversions of  $\sigma$ . Pan and Zeng [13] proved (see also [6,19] for alternative proofs) the following q-analog of (1.5), which refines Stanley's formula (1.6).

Theorem 1.2 (Pan and Zeng). We have

$$\sum_{n\geq 1}\sum_{\sigma\in\mathfrak{S}_n}u_1^{\widetilde{\mathcal{V}}(\sigma)}u_2^{\widetilde{\mathcal{M}}(\sigma)}u_3^{\widetilde{\mathcal{d}}(\sigma)}u_4^{\widetilde{\mathrm{dd}}(\sigma)}q^{\mathrm{inv}(\sigma)}\frac{t^n}{[n]_q!}=F(x,y,u_4,q;t),$$

where

$$F(x, y, u, q; t) := u_1 \frac{\exp_q((x - u)t) - \exp_q((y - u)t)}{x \exp_q((y - u)t) - y \exp_q((x - u)t)},$$
(1.7)

 $x + y = u_3 + u_4$  and  $xy = u_1u_2$ .

Inspired by these work mentioned above, it is natural to ask for a q-analog of Theorem 1.1. For this purpose, we introduce the q-analog of  $P_n(u_1, u_2, u_3, u_4 | \alpha, \beta)$  by

$$P_n(u_1, u_2, u_3, u_4 | \alpha, \beta; q) := \sum_{\sigma \in \mathfrak{S}_n} u_1^{\mathcal{V}(\sigma)} u_2^{\mathcal{M}(\sigma)-1} u_3^{\operatorname{da}(\sigma)} u_4^{\operatorname{dd}(\sigma)} \alpha^{\operatorname{Ima}(\sigma)-1} \beta^{\operatorname{rma}(\sigma)-1} q^{\operatorname{inv}(\sigma)}.$$

Our main result is the following q-analog of Theorem 1.1, which will be proved via Gessel's q-compositional formula [9].

**Theorem 1.3.** Let F(x, y, u, q; t) be defined in (1.7). Then

$$\sum_{n \ge 0} P_{n+1}(u_1, u_2, u_3, u_4 | \alpha, \beta; q) \frac{t^n}{[n]_q!} = \prod_{k=0}^{\infty} G_k(x, y, u_1, u_2, u_3, u_4; t),$$
(1.8)

where  $x + y = u_3 + u_4$ ,  $xy = u_1u_2$  and

$$G_k(x, y, u_1, u_2, u_3, u_4; t) := \left[ 1 - t\alpha q^k (1 - q)(u_3 + u_2 F(x, y, u_4, q; tq^{k+1})) \right]^{-1} \\ \times \left[ 1 - \frac{t\beta}{q^k} (q - 1)(u_4 + u_2 F(x, y, u_3, \frac{1}{q}; \frac{t}{q^k})) \right]^{-1}.$$

We will show that Theorem 1.3 reduces to Theorem 1.1 when  $q \to 1$ . As shown by Ji [10], Theorem 1.1 has many nice applications, one of which is the  $\gamma$ -positivity of the Stirling-Eulerian polynomials  $A_n(x, y | \alpha, \beta)$  that we recall below.

If h(x, y) is symmetry and homogeneous of degree n, then it can be expanded as

$$h(x,y) = \sum_{k=0}^{\lfloor n/2 \rfloor} \gamma_k (xy)^k (x+y)^{n-2k}.$$

Moreover, if  $\gamma_k \geq 0$  for all k, then h(x, y) is said to be  $\gamma$ -positive. One of the typical examples arising from permutation statistics due to Foata and Schüzenberger is the bivariate Eulerian polynomials  $A_n(x, y) := \sum_{\sigma \in \mathfrak{S}_n} x^{\operatorname{asc}(\sigma)} y^{\operatorname{des}(\sigma)}$ ; see the survey of Athanasiadis [3] for more information.

Using Theorem 1.1, Ji proved the following refined  $\gamma$ -positivity expansion.

**Theorem 1.4** (Ji [10, Theorem 1.9]). Let  $\mathcal{P}_{n,k}$  be the set of  $\sigma \in \mathfrak{S}_n$  such that  $\widetilde{M}(\sigma) = k$ . Then

$$A_n(x,y|\frac{\alpha+\beta}{2},\frac{\alpha+\beta}{2}) = \sum_{k=0}^{\lfloor (n-1)/2 \rfloor} \gamma_{n,k}(\alpha,\beta)(xy)^k (x+y)^{n-2k-1},$$
 (1.9)

where

$$\gamma_{n,k}(\alpha,\beta) = 2^{2k+1-n} \sum_{\sigma \in \mathcal{P}_{n,k}} \alpha^{\operatorname{lmi}(\sigma)-1} \beta^{\operatorname{rmi}(\sigma)-1}.$$

On the other hand, Ji and Lin [11] proved the  $\alpha = \beta$  case combinatorially via introducing a group action on permutations, namely,

$$A_n(x,y|\alpha,\alpha) = \sum_{k=0}^{\lfloor (n-1)/2 \rfloor} \gamma_{n,k}(\alpha) (xy)^k (x+y)^{n-2k-1},$$
(1.10)

where

$$\gamma_{n,k}(\alpha) = 2^{2k+1-n} \sum_{\sigma \in \mathcal{P}_{n,k}} \alpha^{\operatorname{lmi}(\sigma) + \operatorname{rmi}(\sigma) - 2}.$$

Comparing (1.10) with (1.9) we get the following identity.

**Theorem 1.5.** For  $n \ge 1$ , we have

$$\sum_{\sigma \in \mathcal{P}_{n,k}} (2\alpha)^{\operatorname{lmi}(\sigma)-1} (2\beta)^{\operatorname{rmi}(\sigma)-1} = \sum_{\sigma \in \mathcal{P}_{n,k}} (\alpha + \beta)^{\operatorname{lmi}(\sigma) + \operatorname{rmi}(\sigma)-2}.$$
 (1.11)

We will introduce a group action on permutations to prove Theorem 1.5 combinatorially, which provides a combinatorial approach to (1.9) in view of the group action proof of (1.10) in [11].

The rest of this paper is organized as follows. In Section 2, we recall a q-analog of the exponential formula from [9] and use it to prove Theorem 1.3. In Section 3, we show that Theorem 1.3 can be described as an integral formula which reduces to Theorem 1.1 when  $q \rightarrow 1$ . The combinatorial proof of Theorem 1.5 is proved in Section 4 via introducing a new group action on permutations.

#### 2. Gessel's q-compositional formula and proof of Theorem 1.3

2.1. Gessel's *q*-compositional formula. Throughout this paper, let *R* be a ring of polynomials in some finite indeterminates with rational coefficients. The *Eulerian differential* operator  $\delta_t$  (see [2]) is defined by

$$\delta_t(f(t)) = \frac{f(qt) - f(t)}{(q-1)t}$$

for any function  $f(t) \in R[[t]]$  in the ring of formal power series in t over R. Note that for  $q = 1, \delta_t$  reduces to the ordinary derivative. For example, we have

$$\delta_t(t^n/[n]_q!) = t^{n-1}/[n-1]_q! \quad \text{for } n \ge 1.$$
(2.1)

For the sake of convenience, we shall often write f' for  $\delta_t(f)$ .

Following Gessel [9], we define the following q-analog of the map  $f \mapsto f^k/k!$  for exponential generating functions.

**Definition 2.** Suppose that 
$$f(0) = 0$$
. Define  $f^{[k]}$  by  $f^{[0]} = 1$  and for  $k \ge 1$ ,  
 $\delta_t(f^{[k]}) = f' \cdot f^{[k-1]}$  with  $f^{[k]}(0) = 0$ . (2.2)

If we write  $f^{[k]}(t) = \sum_{n \ge 0} f_n^{(k)} \frac{t^n}{[n]_q!}$  for  $k \ge 1$ , then Eq. (2.2) is equivalent to the recursion

$$f_{n+1}^{(k)} = \sum_{i=0}^{n} {n \brack i}_{q} f_{n-i+1}^{(1)} f_{i}^{(k-1)},$$

where  $\begin{bmatrix} n \\ i \end{bmatrix}_q := \frac{[n]_q!}{[i]_q![n-i]_q!}$  are the *q*-binomial coefficients.

**Definition 3.** Suppose that f(0) = 0 and  $g(t) = \sum_{n \ge 0} g_n \frac{t^n}{[n]_q!}$ . Define the q-composition g[f] by

$$g[f] := \sum_{n \ge 0} g_n f^{[n]}.$$

Note that  $t^{[k]} = t^k / [k]_q!$  and so g[t] = g(t). Gessel [9] proved the following q-compositional formula, which express  $\exp_q[f]$  as an infinite product.

**Theorem 2.1** (Gessel [9]). Suppose that f(0) = 0. Then

$$\exp_q[f] = \prod_{k=0}^{\infty} \left( 1 - tq^k (1-q)f'(q^k t) \right)^{-1}.$$
(2.3)

A permutation is *basic* if it begins with its greatest element. Denote by  $\mathfrak{B}_n$  the set of all basic permutations in  $\mathfrak{S}_n$ . For a permutation  $\sigma$  with  $\operatorname{Ima}(\sigma) = k$ , there exists a unique decomposition of  $\sigma$  into basic components as

$$\sigma = \beta_1 \beta_2 \cdots \beta_k, \tag{2.4}$$

where

- each  $\beta_i$  is basic;
- and the first letter of each  $\beta_i$  is a left-to-right maximum of  $\sigma$ .

Such a decomposition of  $\sigma$  is called the *basic decomposition* of  $\sigma$  in [9]. For example,  $\sigma = 2164573$  has the basic decomposition  $\beta_1 = 21$ ,  $\beta_2 = 645$  and  $\beta_3 = 73$ .

Given any permutation  $\sigma$  with n distinct letters, let  $red(\sigma)$  be the permutation in  $\mathfrak{S}_n$  obtained from  $\sigma$  by replacing its *i*-th smallest element by *i* for  $1 \leq i \leq n$ . For example, red(8425) = 4213. A function  $\omega$  from the set of all permutations to a commutative algebra over the rationals is called *multiplicative* if for all permutations  $\sigma$ :

(i)  $\omega(\sigma) = \omega(\operatorname{red}(\sigma));$ 

(ii) If  $\beta_1 \beta_2 \cdots \beta_k$  is the basic decomposition of  $\sigma$ , then  $\omega(\sigma) = \omega(\beta_1)\omega(\beta_2)\cdots\omega(\beta_k)$ . For instance, the function  $\omega(\sigma) = x^{\operatorname{Ima}(\sigma)}$  is multiplicative.

**Theorem 2.2** (Gessel [9]). Let  $\omega$  be a multiplicative function. Suppose  $g_n = \sum_{\sigma \in \mathfrak{S}_n} \omega(\sigma) q^{\mathrm{inv}(\sigma)}$ and  $f = \sum_{\sigma \in \mathfrak{S}_n} \omega(\sigma) q^{\mathrm{inv}(\sigma)}$ . Then

and 
$$f_n = \sum_{\beta \in \mathfrak{B}_n} \omega(\beta) q^{\mathrm{inv}(\beta)}$$
. Then

$$\sum_{n\geq 0} g_n \frac{t^n}{[n]_q!} = \exp_q \left[ \sum_{n\geq 1} f_n \frac{t^n}{[n]_q!} \right].$$
(2.5)

# 2.2. Proof of Theorem 1.3.

**Definition 4.** For  $\sigma \in \mathfrak{S}_n$ , let  $M_0(\sigma)$ ,  $V_0(\sigma)$ ,  $da_0(\sigma)$  and  $dd_0(\sigma)$  (resp.,  $M_{\infty}(\sigma)$ ),  $V_{\infty}(\sigma)$ ,  $da_{\infty}(\sigma)$  and  $dd_{\infty}(\sigma)$ ) denote respectively the numbers of peaks, valleys, double ascents and double descents in  $\sigma$  with boundary conditions  $\sigma_0 = 0$  and  $\sigma_{n+1} = +\infty$  (resp.,  $\sigma_0 = +\infty$  and  $\sigma_{n+1} = 0$ ). Introduce two weighted functions by

$$\omega_0(\sigma) = u_1^{\mathcal{V}_0(\sigma)} u_2^{\mathcal{M}_0(\sigma)} u_3^{\mathrm{da}_0(\sigma)} u_4^{\mathrm{dd}_0(\sigma)} \alpha^{\mathrm{lma}(\sigma)}$$

and

$$\omega_{\infty}(\sigma) = u_1^{\mathcal{V}_{\infty}(\sigma)} u_2^{\mathcal{M}_{\infty}(\sigma)} u_3^{\mathrm{da}_{\infty}(\sigma)} u_4^{\mathrm{dd}_{\infty}(\sigma)} \beta^{\mathrm{rma}(\sigma)}$$

The following observation can be checked routinely.

# **Lemma 2.3.** The function $\omega_0$ is multiplicative.

We are now in position to prove Theorem 1.3.

**Proof of Theorem 1.3.** Throughout the proof, we assume that  $x + y = u_3 + u_4$  and  $xy = u_1u_2$ . Let

$$L_n(u_3, u_4, \alpha, q) := \sum_{\sigma \in \mathfrak{S}_n} \omega_0(\sigma) q^{\mathrm{inv}(\sigma)} \quad \text{and} \quad B_n(\alpha, q) := \sum_{\sigma \in \mathfrak{B}_n} \omega_0(\sigma) q^{\mathrm{inv}(\sigma)}.$$

It follows from Lemma 2.3 and Theorem 2.2 that

$$\sum_{n\geq 0} L_n(u_3, u_4, \alpha, q) \frac{t^n}{[n]_q!} = \exp_q \left[ \sum_{n\geq 1} B_n(\alpha, q) \frac{t^n}{[n]_q!} \right].$$
(2.6)

As  $B_1(\alpha, q) = u_3 \alpha$  and for  $n \ge 2$ ,

$$B_n(\alpha,q) = u_2 \alpha q^{n-1} \sum_{\sigma \in \mathfrak{S}_{n-1}} u_1^{\widetilde{\mathcal{V}}(\sigma)} u_2^{\widetilde{\mathcal{M}}(\sigma)} u_3^{\widetilde{\mathcal{d}}(\sigma)} u_4^{\widetilde{\mathcal{d}}(\sigma)} q^{\operatorname{inv}(\sigma)},$$

by (2.1) we have

$$\delta_t \left( \sum_{n \ge 1} B_n(\alpha, q) \frac{t^n}{[n]_q!} \right) = \sum_{n \ge 1} B_n(\alpha, q) \frac{t^{n-1}}{[n-1]_q!}$$
$$= u_3 \alpha + u_2 \alpha \sum_{n \ge 1} \sum_{\sigma \in \mathfrak{S}_n} u_1^{\widetilde{\mathcal{V}}(\sigma)} u_2^{\widetilde{\mathcal{M}}(\sigma)} u_3^{\widetilde{\mathcal{d}}(\sigma)} u_4^{\widetilde{\mathcal{d}}(\sigma)} q^{\operatorname{inv}(\sigma)} \frac{(tq)^n}{[n]_q!}$$
$$= u_3 \alpha + u_2 \alpha F(x, y, u_4, q; tq)$$

in view of Theorem 1.2. It then follows from Theorem 2.1 and Eq. (2.6) that

$$\sum_{n\geq 0} L_n(u_3, u_4, \alpha, q) \frac{t^n}{[n]_q!} = \prod_{k\geq 0} \left[ 1 - t\alpha q^k (1-q)(u_3 + u_2 F(x, y, u_4, q; tq^{k+1})) \right]^{-1}.$$
 (2.7)

On the other hand, we need to compute the exponential generating function for

$$R_n(u_3, u_4, \beta, q) := \sum_{\sigma \in \mathfrak{S}_n} \omega_{\infty}(\sigma) q^{\operatorname{inv}(\sigma)}.$$

For  $\sigma \in \mathfrak{S}_n$ , let  $\sigma^r := \sigma_n \sigma_{n-1} \cdots \sigma_1$  be the *reversal* of  $\sigma$ . As

$$V_{\infty}(\sigma^r) = V_0(\sigma), \ M_{\infty}(\sigma^r) = M(\sigma), \ da_{\infty}(\sigma^r) = dd_0(\sigma), \ dd_{\infty}(\sigma^r) = da_0(\sigma)$$

and

$$\operatorname{rma}(\sigma^r) = \operatorname{lma}(\sigma), \quad \operatorname{inv}(\sigma^r) = \binom{n}{2} - \operatorname{inv}(\sigma),$$

we have

$$\sum_{n\geq 0} R_n(u_3, u_4, \beta, q) \frac{t^n}{[n]_q!} = \sum_{n\geq 0} q^{\binom{n}{2}} L_n(u_4, u_3, \beta, q^{-1}) \frac{t^n}{[n]_q!}$$
$$= \sum_{n\geq 0} L_n(u_4, u_3, \beta, q^{-1}) \frac{t^n}{[n]_{q^{-1}}!}.$$

It then follows from (2.7) that

$$\sum_{n\geq 0} R_n(u_3, u_4, \beta, q) \frac{t^n}{[n]_q!} = \prod_{k\geq 0} \left[ 1 - \frac{t\beta}{q^k} (1 - \frac{1}{q}) (u_4 + u_2 F(x, y, u_3, \frac{1}{q}; \frac{t}{q^{k+1}})) \right]^{-1}.$$
 (2.8)

It is known [17, Prop 1.3.17] that  $\binom{n}{k}_q$  has the following interpretation

$$\begin{bmatrix} n \\ k \end{bmatrix}_q = \sum_{(\mathcal{A},\mathcal{B})} q^{\mathrm{inv}(\mathcal{A},\mathcal{B})}$$

summed over all ordered partitions  $(\mathcal{A}, \mathcal{B})$  of [n] such that  $|\mathcal{A}| = k$ . Let

$$\mathfrak{S}_{n+1,k+1} := \{ \sigma \in \mathfrak{S}_{n+1} : \sigma_{k+1} = n+1 \}.$$

As any permutation  $\sigma \in \mathfrak{S}_{n+1,k+1}$  can be written as  $\sigma = \sigma_L(n+1)\sigma_R$  with  $\sigma_L = \sigma_1 \cdots \sigma_k$ and  $\sigma_R = \sigma_{k+2} \cdots \sigma_{n+1}$ , we have

$$\sum_{\sigma \in \mathfrak{S}_{n+1,k+1}} u_1^{\mathcal{V}(\sigma)} u_2^{\mathcal{M}(\sigma)-1} u_3^{\operatorname{da}(\sigma)} u_4^{\operatorname{dd}(\sigma)} \alpha^{\operatorname{Ima}(\sigma)-1} \beta^{\operatorname{rma}(\sigma)-1} q^{\operatorname{inv}(\sigma)}$$
$$= \begin{bmatrix} n \\ k \end{bmatrix}_q q^{n-k} L_k(u_3, u_4, \alpha, q) R_{n-k}(u_3, u_4, \beta, q)$$

in view of the above interpretation of  $\begin{bmatrix} n \\ k \end{bmatrix}_q$ . Thus,

$$P_{n+1}(u_1, u_2, u_3, u_4 | \alpha, \beta; q) = \sum_{k=0}^n {n \brack k}_q q^{n-k} L_k(u_3, u_4, \alpha, q) R_{n-k}(u_3, u_4, \beta, q),$$

which in terms of exponential generating function gives

$$\sum_{n\geq 0} P_{n+1}(u_1, u_2, u_3, u_4 | \alpha, \beta; q) \frac{t^n}{[n]_q!} = \left(\sum_{n\geq 0} L_n(u_3, u_4, \alpha, q) \frac{t^n}{[n]_q!}\right) \left(\sum_{n\geq 0} R_n(u_3, u_4, \beta, q) \frac{(tq)^n}{[n]_q!}\right)$$
  
Substituting (2.7) and (2.8) into the above identity proves Theorem 1.3.

Substituting (2.7) and (2.8) into the above identity proves Theorem 1.3.

# 3. A NEW PROOF OF THEOREM 1.1

For any  $f(t) \in R[[t]]$ , define the *q*-integral (see [8, Section 1.11]) of f by

$$\int_0^t f(x) d_q x := t(1-q) \sum_{n \ge 0} f(q^n t) q^n.$$
(3.1)

As  $\lim_{q \to 1} \int_0^t x^k d_q x = t^{k+1}/(k+1)$  for  $k \ge 0$ , we have

$$\lim_{q \to 1} \int_0^t f(x) d_q x = \int_0^t f(x) dx.$$
 (3.2)

Our main result Theorem 1.3 can also be described as an integral formula.

## Theorem 3.1. We have

$$\sum_{n\geq 0} P_{n+1}(u_1, u_2, u_3, u_4 | \alpha, \beta; q) \frac{t^n}{[n]_q!}$$
  
=  $\exp\left(\int_0^t \frac{\ln\left[1 - (1 - q)z\alpha(u_3 + u_2F(zq))\right]^{-1}}{(1 - q)z} d_q z + \int_0^t \frac{\ln\left[1 - (q - 1)z\beta(u_4 + u_2\tilde{F}(z))\right]^{-1}}{(1 - \frac{1}{q})z} d_{\frac{1}{q}}z\right)$ 

with  $x + y = u_3 + u_4$ ,  $xy = u_1u_2$ ,

$$F(t) := F(x, y, u_4, q; t) \quad and \quad \tilde{F}(t) := F(x, y, u_3, \frac{1}{q}; t).$$
(3.3)

 $\begin{aligned} Proof. \ \text{As } \ln(1-t)^{-1} &= \sum_{n\geq 1} \frac{t^n}{n}, \text{ we have} \\ &\prod_{k\geq 0} \left[ 1 - t\alpha q^k (1-q)(u_3 + u_2F(tq^{k+1})) \right]^{-1} \\ &= \exp\left( \ln\left(\prod_{k\geq 0} [1 - t\alpha q^k (1-q)(u_3 + u_2F(tq^{k+1}))]^{-1} \right) \right) \\ &= \exp\left( \sum_{k\geq 0} \ln[1 - t\alpha q^k (1-q)(u_3 + u_2F(tq^{k+1}))]^n \right) \\ &= \exp\left( \sum_{k\geq 0} \sum_{n\geq 1} \frac{[t\alpha q^k (1-q)(u_3 + u_2F(tq^{k+1}))]^n}{n} \right) \\ &= \exp\left( \sum_{n\geq 1} \frac{(1-q)^{n-1}}{n} (1-q)t \sum_{k\geq 0} q^k (tq^k)^{n-1} (u_3\alpha + u_2\alpha F(tq^{k+1}))^n \right) \\ &= \exp\left( \sum_{n\geq 1} \frac{(1-q)^{n-1}}{n} \int_0^t z^{n-1} (u_3\alpha + u_2\alpha F(zq))^n d_q z \right) \\ &= \exp\left( \int_0^t \sum_{n\geq 1} \frac{[(1-q)z(u_3\alpha + u_2\alpha F(zq))]^n}{n(1-q)z} d_q z \right) \\ &= \exp\left( \int_0^t \frac{\ln[1 - (1-q)z\alpha(u_3 + u_2F(zq))]^{-1}}{(1-q)z} d_q z \right). \end{aligned}$ 

Similarly, we have

$$\prod_{k\geq 0} \left[ 1 - \frac{t\beta}{q^k} (q-1)(u_4 + u_2\tilde{F}(tq^{-k})) \right]^{-1} = \exp\left(\int_0^t \frac{\ln\left[1 - (q-1)z\beta\left(u_4 + u_2\tilde{F}(z)\right)\right]^{-1}}{(1 - \frac{1}{q})z} d_{\frac{1}{q}}z \right).$$

The desired integral formula then follows from Theorem 1.3.

Next, we show that our integral formula in Theorem 3.1 reduces to Theorem 1.1 when  $q \rightarrow 1$ . We need the following simple lemma.

**Lemma 3.2.** Let F(x, y; t) be defined in (1.4). Then,

$$\int_{0}^{t} xy F(x, y; z) dz = \ln\left(\frac{x - y}{xe^{yt} - ye^{xt}}\right).$$
 (3.4)

*Proof.* We have

$$\frac{d\left(-\ln[xe^{yt}-ye^{xt}]\right)}{dt} = \frac{xy(e^{xt}-e^{yt})}{xe^{yt}-ye^{xt}} = xyF(x,y;t)$$

from which the result then follows.

We are now ready for a new proof of Theorem 1.1.

A new proof of Theorem 1.1. When  $q \to 1$ , the two infinitesimal quantities

 $\ln[1 - (1 - q)t\alpha(u_3 + u_2F(tq))]$  and  $(1 - q)t\alpha(u_3 + u_2F(tq))$ 

are equivalent, and so are

$$\ln\left[1-(q-1)t\beta\left(u_4+u_2\tilde{F}(t)\right)\right] \quad \text{and} \quad (q-1)t\beta\left(u_4+u_2\tilde{F}(t)\right),$$

where F(t) and  $\tilde{F}(t)$  are defined in (3.3). Thus, if we let  $q \to 1$  in Theorem 3.1, then by (3.2) we have

$$\sum_{n\geq 0} P_{n+1}(u_1, u_2, u_3, u_4 | \alpha, \beta) \frac{t^n}{n!}$$
  
=  $\exp\left(\int_0^t \alpha (u_3 + u_2 F(x, y, u_4, 1; z)) dz + \int_0^t \beta (u_4 + u_2 F(x, y, u_3, 1; z)) dz\right)$   
=  $\exp\left(\int_0^t \alpha (u_3 + xy F(x, y; z)) dz + \int_0^t \beta (u_4 + xy F(x, y; z)) dz\right)$ 

with  $xy = u_1u_2$  and  $x + y = u_3 + u_4$ . Using (3.4) we get

$$\sum_{n\geq 0} P_{n+1}(u_1, u_2, u_3, u_4 | \alpha, \beta) \frac{t^n}{n!} = e^{(u_3\alpha + u_4\beta)t} \left(\frac{x - y}{xe^{yt} - ye^{xt}}\right)^{\alpha + \beta},$$

completing the proof.

### 4. Combinatorics of the $\gamma$ -positivity of Stirling-Eulerian polynomials

This section aims to prove Theorem 1.5 combinatorially by introducing a new group action on permutations.

For  $\sigma \in \mathfrak{S}_n$  with  $\operatorname{lmi}(\sigma) - 1 = k$  and  $\operatorname{rmi}(\sigma) - 1 = l$ , there exists a unique decomposition of  $\sigma$  as

$$\sigma = \alpha_1 \alpha_2 \cdots \alpha_k 1 \beta_1 \beta_2 \cdots \beta_l,$$

where

- the first letter of each  $\alpha_i$   $(1 \le i \le k)$ , which is smallest inside  $\alpha_i$ , is a left-to-right minimum of  $\sigma$ ;
- and the last letter of each  $\beta_i$   $(1 \le i \le l)$ , which is smallest inside  $\beta_i$ , is a right-to-left minimum of  $\sigma$ .

We call this decomposition the *bi-basic decomposition* of  $\sigma$  and each  $\alpha_i$  or  $\beta_i$  a block of  $\sigma$ . An example of the bi-basic decomposition is

$$\sigma = 5\,10|\mathbf{2}\,12\,4\,13\,6|\mathbf{1}|11\,\mathbf{3}|9\,8\,15\,\mathbf{7}|\mathbf{14},\tag{4.1}$$

where blocks are separated by bars. For each word  $w = w_1 w_2 \cdots w_d$ , let  $w^r := w_d w_{d-1} \cdots w_1$ be the reversal of w. For each  $x \in [n]$ , we define  $\psi_x(\sigma)$  according to the following cases

(1) if x is the first letter of  $\alpha_i$  for some i, then  $\psi_x(\sigma)$  is obtained from  $\sigma$  by deleting the block  $\alpha_i$  and then inserting  $\alpha_i^r$  in some proper gap between blocks so that x becomes an additional right-to-left minimum;

- (2) if x is the first letter of  $\beta_i$  for some i, then  $\psi_x(\sigma)$  is obtained from  $\sigma$  by deleting the block  $\beta_i$  and then inserting  $\beta_i^r$  in some proper gap between blocks so that x becomes an additional left-to-right minimum;
- (3) otherwise, set  $\psi_x(\sigma) = \sigma$ .

Taking  $\sigma$  as in (4.1), then

$$\psi_2(\sigma) = \mathbf{5}\,10|\mathbf{1}|6\,13\,4\,12\,\mathbf{2}|11\,\mathbf{3}|9\,8\,15\,\mathbf{7}|\mathbf{14},$$

and

$$\psi_3(\sigma) = 5\,10|3\,11|2\,12\,4\,13\,6|1|9\,8\,15\,7|14.$$

It is clear that the map  $\psi_x$  is an involution acting on  $\mathfrak{S}_n$ , and for all  $x, y \in [n]$ ,  $\psi_x$  and  $\psi_y$  commute. The following lemma is evident from the construction of  $\psi_x$ .

**Lemma 4.1.** If x is a left-to-right (resp., right-to-left) minimum of  $\sigma$ , then x becomes a right-to-left (resp., left-to-right) minimum of  $\psi_x(\sigma)$ . Moreover,  $\psi_x$  preserves the peak statistic " $\widetilde{M}$ ".

We are now ready for the combinatorial proof of Theorem 1.5.

**Proof of Theorem 1.5.** Denote by  $LMI(\sigma)$  and  $RMI(\sigma)$  the set of left-to-right minima and the set of right-to-left minima of  $\sigma$  other than 1, respectively. In order to interpret the left-hand side of (1.11), we introduce the set

$$\mathcal{P}_{n,k} = \{(\sigma, S) : \sigma \in \mathcal{P}_{n,k}, S \subseteq \text{LMI}(\sigma) \cup \text{RMI}(\sigma)\}$$

of marked permutations, where  $\mathcal{P}_{n,k} := \{ \sigma \in \mathfrak{S}_n | \widetilde{M}(\sigma) = k \}$ . Thus,

$$\sum_{\sigma \in \mathcal{P}_{n,k}} (2\alpha)^{\operatorname{lmi}(\sigma)-1} (2\beta)^{\operatorname{rmi}(\sigma)-1} = \sum_{(\sigma,S) \in \widetilde{\mathcal{P}}_{n,k}} \alpha^{\operatorname{lmi}(\sigma)-1} \beta^{\operatorname{rmi}(\sigma)-1}.$$
(4.2)

For  $x \in [n]$  and  $(\sigma, S) \in \widetilde{\mathcal{P}}_{n,k}$ , we define the action  $\Psi_x$  on  $\widetilde{\mathcal{P}}_{n,k}$  as

$$\Psi_x(\sigma, S) = \begin{cases} (\psi_x(\sigma), S \triangle \{x\}), & \text{if } x \in \text{LMI}(\sigma) \cup \text{RMI}(\sigma); \\ (\sigma, S), & \text{otherwise.} \end{cases}$$

It is routine to check that all  $\Psi_x$ 's are involutions and commute. Therefore, for any  $X \subseteq [n]$ we can define the function  $\Psi_X : \widetilde{\mathcal{P}}_{n,k} \to \widetilde{\mathcal{P}}_{n,k}$  by  $\Psi_X = \prod_{x \in X} \Psi_x$ , where multiplication is the composition of functions. Hence the group  $\mathbb{Z}_2^n$  acts on  $\widetilde{\mathcal{P}}_{n,k}$  via the function  $\Psi_X$ . Let  $\operatorname{Orb}(\sigma, S)$  be the orbit of  $(\sigma, S)$  under this action. Inside each orbit  $\operatorname{Orb}(\sigma, S)$  there exists a unique element  $(\bar{\sigma}, \emptyset)$ , where  $\bar{\sigma} = \Psi_S(\sigma)$ . It then follows from Lemma 4.1 that

$$\sum_{(\pi,T)\in \operatorname{Orb}(\sigma,S)} \alpha^{\operatorname{lmi}(\pi)-1} \beta^{\operatorname{rmi}(\pi)-1} = (\alpha+\beta)^{\operatorname{lmi}(\bar{\sigma})+\operatorname{rmi}(\bar{\sigma})-2}.$$

Summing over all orbits of  $\widetilde{\mathcal{P}}_{n,k}$  then results in

$$\sum_{(\sigma,S)\in\widetilde{\mathcal{P}}_{n,k}} \alpha^{\operatorname{lmi}(\sigma)-1} \beta^{\operatorname{rmi}(\sigma)-1} = \sum_{\sigma\in\mathcal{P}_{n,k}} (\alpha+\beta)^{\operatorname{lmi}(\sigma)+\operatorname{rmi}(\sigma)-2}$$

which proves (1.11) in view of (4.2).

As a byproduct of the action  $\psi_x$ , we can also prove the following neat expression for  $\gamma_{n,k}(\alpha,\beta)$ .

**Theorem 4.2.** For  $n \ge 1$ , we have

$$\sum_{\sigma \in \mathcal{P}_{n,k}} \alpha^{\operatorname{lmi}(\sigma)-1} \beta^{\operatorname{rmi}(\sigma)-1} = \sum_{\sigma \in \mathcal{L}_{n-1,k}} (\alpha + \beta)^{\operatorname{rmi}(\sigma)},$$
(4.3)

where  $\mathcal{L}_{n-1,k} = \{ \sigma \in \mathfrak{S}_{n-1} : \mathrm{M}_0(\sigma) = k \}$ . Consequently,

$$\gamma_{n,k}(\alpha,\beta) = 2^{2k+1-n} \sum_{\sigma \in \mathcal{L}_{n-1,k}} (\alpha+\beta)^{\operatorname{rmi}(\sigma)}.$$

*Proof.* We consider the function  $\psi_X : \mathcal{P}_{n,k} \to \mathcal{P}_{n,k}$  defined by  $\psi_X = \prod_{x \in X} \psi_x$ . Then, the group  $\mathbb{Z}_2^n$  acts on  $\mathcal{P}_{n,k}$  via the function  $\psi_X$ . Let  $\operatorname{Orb}(\sigma)$  be the orbit of  $\sigma$  under this action. Inside each orbit  $\operatorname{Orb}(\sigma)$  there exists a unique element  $\bar{\sigma}$  whose first letter is 1. It then follows from Lemma 4.1 that

$$\sum_{\pi \in \operatorname{Orb}(\sigma)} \alpha^{\operatorname{lmi}(\pi) - 1} \beta^{\operatorname{rmi}(\pi) - 1} = (\alpha + \beta)^{\operatorname{rmi}(\bar{\sigma}) - 1}.$$

Summing over all orbits of  $\mathcal{P}_{n,k}$  then results in

$$\sum_{\sigma \in \mathcal{P}_{n,k}} \alpha^{\operatorname{Imi}(\sigma)-1} \beta^{\operatorname{rmi}(\sigma)-1} = \sum_{\substack{\sigma \in \mathcal{P}_{n,k} \\ \sigma_1 = 1}} (\alpha + \beta)^{\operatorname{rmi}(\bar{\sigma})-1}$$
$$= \sum_{\sigma \in \mathcal{L}_{n-1,k}} (\alpha + \beta)^{\operatorname{rmi}(\sigma)},$$

where the last equality follows from the simple one-to-one correspondence

$$\sigma \mapsto (\sigma_2 - 1)(\sigma_3 - 1) \cdots (\sigma_n - 1)$$

between  $\{\sigma \in \mathcal{P}_{n,k} : \sigma_1 = 1\}$  and  $\mathcal{L}_{n-1,k}$ .

The Euler numbers  $\{E_n\}_{n\geq 0}$  can be defined by the Taylor expansion

ŝ

$$\sec(t) + \tan(t) = \sum_{n \ge 0} E_n \frac{t^n}{n!}.$$

The numbers  $E_n$  with even indices and odd indices are known as *secant numbers* and *tangent numbers*, respectively. A permutation  $\sigma \in \mathfrak{S}_n$  is *alternating* if

$$\sigma_1 > \sigma_2 < \sigma_3 > \sigma_4 < \sigma_5 > \cdots$$

André [1] proved that  $|\mathcal{A}_n| = E_n$ , where  $\mathcal{A}_n$  denotes the set of all alternating permutations of length n. An immediate consequence of Theorem 4.2 is the following interesting interpretation of the  $\alpha$ -extension of secant numbers

$$A_{2n+1}(-1,1|\alpha/2,\alpha/2) = (-1)^n \sum_{\sigma \in \mathcal{A}_{2n}} \alpha^{\mathrm{rmi}(\sigma)},$$

which was proved by Ji [10] via different approach.

#### 5. Concluding remarks

The main achievement of this paper is the computation of the generating function for a qanalog of  $P_n(u_1, u_2, u_3, u_4 | \alpha, \beta)$  by using Gessel's q-compositional formula. In particular, Ji's formula for  $P_n(u_1, u_2, u_3, u_4 | \alpha, \beta)$  in Theorem 1.1 can be proved directly from Carlitz and Scoville's formula (1.3) by using the classical compositional formula (see [18, Thm. 5.1.4]). However, it should be noted that this approach via the compositional formula does not work for the variation of  $P_n(u_1, u_2, u_3, u_4 | \alpha, \beta)$ :

$$\tilde{P}_n(u_1, u_2, u_3, u_4 | \alpha, \beta) := \sum_{\sigma \in \mathfrak{S}_n} u_1^{\tilde{\mathcal{V}}(\sigma) - 1} u_2^{\tilde{\mathcal{M}}(\sigma)} u_3^{\tilde{\mathrm{da}}(\sigma)} u_4^{\tilde{\mathrm{dd}}(\sigma)} \alpha^{\mathrm{lma}(\sigma) - 1} \beta^{\mathrm{rma}(\sigma) - 1}$$

It remains open to compute the generating function for  $\tilde{P}_n(u_1, u_2, u_3, u_4 | \alpha, \beta)$ .

A statistic whose distribution over  $\mathfrak{S}_n$  gives  $[n]_q!$  is called a *Mahonian* statistic. It is well known that 'inv' is a Mahonian statistic on permutations. Another classical Mahonian statistic is the *major index* (see [17, Sec. 1.3])

$$\operatorname{maj}(\sigma) = \sum_{\sigma_i > \sigma_{i+1}} i$$

A third Mahonian statistic arising from poset topology was introduced by Shareshian and Wachs [15] as 'ai + des', where ai( $\sigma$ ) is the number of *admissible inversions* of  $\sigma$  (see also [12]). The generating functions for the four pairs of Euler–Mahonian statistics in {exc, des} × {inv, maj} have already been calculated in the literature; see [15]. In view of Theorem 1.3, it would be interesting to investigate systematically the generating functions for the Stirling–Euler–Mahonian statistics on permutations in

 $\{\rm{lmi},\rm{rmi},\rm{lma},\rm{rma},\rm{cyc}\}\times\{\rm{des},\rm{exc},\rm{M}\}\times\{\rm{inv},\rm{maj},\rm{ai}\}$ 

and their possible refinements. Some related results in this direction have been obtained by Sokal and Zeng [16].

# STATEMENTS AND DECLARATIONS

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

## Acknowledgement

This work was supported by the National Science Foundation of China grants 12271301, 12322115 and 12201468. Part of this work was done while the first author was visiting the third author at Wenzhou University in the winter of 2023.

### References

- [1] D. André, Développement de sec x and tan x, C. R. Math. Acad. Sci. Paris, 88 (1879), 965–979. 12
- [2] G.E. Andrews, On the foundations of combinatorial theory V, Eulerian differential operators, Stud. Appl. Math., 50 (1971), 345–375. 5
- [3] C.A. Athanasiadis, Gamma-positivity in combinatorics and geometry, Sém. Lothar. Combin. 77 (2018), Article B77i, 64pp (electronic). 4
- [4] L. Carlitz and R. Scoville, Generalized Eulerian numbers: combinatorial applications, J. Reine Angew. Math., 265 (1974), 110–137. 2, 3

- [5] W.Y.C. Chen, Context-free grammars, differential operators and formal power series, Theor. Comput. Sci., 117 (1993), 113–129. 1
- [6] Y. Dong and Z. Lin, Counting and signed counting permutations by descent-based statistics, preprint.
- [7] A.M. Fu, A context-free grammar for peaks and double descents of permutations, Adv. in Appl. Math., 100 (2018), 179–196. 3
- [8] G. Gasper and M. Rahman, Basic Hypergeometric Series, second ed., Cambridge University Press, Cambridge, 2004. 8
- [9] I.M. Gessel, A q-analog of the exponential formula, Discrete Math., 40 (1982), 69–80. 1, 4, 5, 6
- [10] K.Q. Ji, The  $(\alpha, \beta)$ -Eulerian Polynomials and Descent-Stirling Statistics on Permutations, arXiv:2310.01053. 1, 2, 3, 4, 12
- [11] K.Q. Ji and Z. Lin, The Binomial-Stirling-Eulerian Polynomials, arXiv:2310.04969.4, 5
- [12] Z. Lin and J. Zeng, The  $\gamma$ -positivity of basic Eulerian polynomials via group actions, J. Combin. Theory, Ser. A, **135** (2015), 112–129. 13
- [13] Q. Pan and J. Zeng, A q-analogue of generalized Eulerian polynomials with applications, Adv. in Appl. Math., 104 (2019), 85–99. 3
- [14] T.K. Petersen, Eulerian numbers, with a foreword by Richard Stanley, Birkhäuser Advanced Texts: Basler Lehrbücher. Birkhäuser/Springer, New York, 2015. 1, 3
- [15] J. Shareshian and M.L. Wachs, q-Eulerian polynomials: excedance number and major index, Electron. Res. Announc. Amer. Math. Soc., 13 (2007), 33–45. 13
- [16] A.D. Sokal and J. Zeng, Some multivariate master polynomials for permutations, set partitions, and perfect matchings, and their continued fractions, Adv. in Appl. Math., 138 (2022), Article 102341. 13
- [17] R.P. Stanley, *Enumerative combinatorics, Vol. 1*, Cambridge University Press, Cambriage, 1997. 1, 2, 8, 13
- [18] R.P. Stanley, Enumerative Combinatorics, Vol. 2, Cambridge University Press, Cambridge, 1999 13
- [19] Y. Zhuang, Eulerian polynomials and descent statistics, Adv. in Appl. Math., 90 (2017), 86–144. 3

(Yao Dong) RESEARCH CENTER FOR MATHEMATICS AND INTERDISCIPLINARY SCIENCES, SHANDONG UNIVERSITY & FRONTIERS SCIENCE CENTER FOR NONLINEAR EXPECTATIONS, MINISTRY OF EDUCA-TION, QINGDAO 266237, P.R. CHINA

Email address: y.dong@mail.sdu.edu.cn

(Zhicong Lin) Research Center for Mathematics and Interdisciplinary Sciences, Shandong University & Frontiers Science Center for Nonlinear Expectations, Ministry of Education, Qingdao 266237, P.R. China

Email address: linz@sdu.edu.cn

(Qiongqiong Pan) College of Mathematics and Physics, Wenzhou University, Wenzhou 325035, P.R. China

Email address: qpan@wzu.edu.cn