

# GLOBAL SOLUTIONS FOR CUBIC QUASILINEAR SCHRÖDINGER FLOWS IN TWO AND HIGHER DIMENSIONS

MIHAELA IFRIM AND DANIEL TATARU

**ABSTRACT.** In recent work the authors proposed a broad global well-posedness conjecture for cubic defocusing dispersive equations in one space dimension, and then proved this conjecture in two cases, namely for one dimensional semilinear and quasilinear Schrödinger flows.

Inspired by the circle of ideas developed in the proof of the above conjecture, in this paper we expand the reach of these methods to higher dimensional quasilinear cubic Schrödinger flows. The study of this class of problems, in all dimensions, was initiated in pioneering work of Kenig-Ponce-Vega for localized initial data, and then continued by Marzuola-Metcalf-Tataru (MMT) for initial data in Sobolev spaces.

The outcomes of this work are (i) a new, potentially sharp local well-posedness result in low regularity Sobolev spaces, one derivative below MMT and just one half derivative above scaling, (ii) a small data global well-posedness and scattering result at the same regularity level, the first result of its kind at least in two space dimensions, and (iii) a new way to think about this class of problems, which, we believe, will become the standard approach in the future.

## CONTENTS

1. Introduction	1
2. Notations and preliminaries	12
3. A frequency envelope formulation of the results	18
4. A paradifferential/resonant expansion of the equation	22
5. Interaction Morawetz estimates: the constant coefficient case	24
6. Density-flux relations for mass and momentum	26
7. Interaction Morawetz bounds for the linear paradifferential flow	28
8. Low regularity Strichartz bounds for the paradifferential flow	41
9. The local/global well-posedness result	44
10. The 2D global result at low regularity	50
References	54

## 1. INTRODUCTION

This article is devoted to the study of both local and global well-posedness for quasilinear cubic Schrödinger flows in two and higher space dimensions, following the authors' recent work in [36] on the similar problem in one space dimension.

---

2020 *Mathematics Subject Classification.* Primary: 35Q55 Secondary: 35B40 .

*Key words and phrases.* NLS problems, quasilinear, defocusing, scattering, interaction Morawetz.

The starting point for the work in [36] was in turn the *global well-posedness conjecture for 1D cubic defocusing dispersive flows* which was introduced also by the authors one year earlier in [33]. This conjecture was first proved in [33] for semilinear Schrödinger flows, and then in [36] for quasilinear Schrödinger flows. Further, the approach in [36] also led to a major improvement to the local well-posedness theory, up to the optimal threshold.

The cubic higher dimensional problem we consider here has received considerable interest over the years, beginning with the first local well-posedness results of Kenig-Ponce-Vega [45], for regular and localized initial data, which was expanded to a full well-posedness theory for small data in lower regularity Sobolev spaces by Marzuola-Metcalfe-Tataru [51], and later for large data in [52]. One key element of the latter paper was the introduction of the local energy spaces, which was instrumental in allowing the study of the problem for nonlocalized data, i.e. in standard Sobolev spaces.

On the other hand, the approach introduced in [36] for the one dimensional problem departs radically from [51], and is instead based on bilinear estimates via an interaction Morawetz analysis, and normal forms ideas implemented via modified energies. Our aim here is to extend the reach of these ideas to higher dimensional problems, leading to a drastic expansion of both local and global results. Precisely, our goals in this paper are fourfold:

- i) to produce a novel, better approach for the study of both local and global dynamics in higher dimensional problems.
- ii) to propose a higher dimensional version of our global well-posedness conjecture, particularly in the most interesting case of two dimensional problems.
- iii) to sharply improve the local well-posedness theory for cubic quasilinear Schrödinger flows in higher dimension, one full derivative below the earlier results of [51] and just one half derivative above scaling, which may well be optimal depending on the dimension.
- iv) to prove the global well-posedness conjecture for cubic quasilinear Schrödinger flows with nonlocalized data, at the same Sobolev regularity as the local well-posedness result. This is the first result of its kind in the most interesting case of 2D quasilinear flows.

**1.1. The global well-posedness conjectures.** As noted earlier, the present work is in part motivated by a broad conjecture formulated by the authors in a recent paper [33]. Our global well-posedness (GWP) conjecture, which applies to both semilinear and quasilinear 1D problems, is as follows:

**Conjecture 1** (Non-localized data defocusing GWP conjecture). *One dimensional dispersive flows on the real line with cubic defocusing nonlinearities and small initial data have global in time, scattering solutions.*

The main result of [33] and then [36] asserts that this conjecture is true first in the semilinear 1D Schrödinger setting. This was the first global in time well-posedness results of this type.

Next, in [36], the authors established the above conjecture for 1D quasilinear Schrödinger flows, which represented the first validation of the conjecture in a quasilinear setting. Precisely, we proved that if the problem has *phase rotation symmetry* and is *conservative* and *defocusing*, then small data in Sobolev spaces yields global, scattering solutions. Notably,

scattering here was interpreted in a weak sense, to mean that the solution satisfies global  $L^6$  Strichartz estimates and bilinear  $L^2$  bounds. This is because of the strong nonlinear effects, which preclude any classical scattering for the problem.

The defocusing condition for the nonlinearity is essential for the global result in 1D. In the focusing case, the existence of small solitons generally prevents global, scattering solutions. Nevertheless, in another recent paper the authors have conjectured that long time solutions can be obtained on a potentially optimal time scale:

**Conjecture 2** (Non-localized data long time well-posedness conjecture). *One dimensional dispersive flows on the real line with cubic conservative nonlinearities and initial data of size  $\epsilon \ll 1$  have long time solutions on the  $\epsilon^{-8}$  time scale.*

The conservative assumption here is very natural, and heuristically aims to prevent nonlinear ode blow-up of solutions with wave packet localization.

In this article we are interested in the counterpart of this conjecture in two and higher space dimensions. Compared to our earlier one dimensional work, the difference is that here we have more dispersion, which is reflected in the fact that the  $L^4$  space-time norm of the solutions, rather than the  $L^6$  one, plays the leading role. In particular, in the semilinear case the small data problem may be approached directly using Strichartz estimates, and global well-posedness and scattering follows. The focusing/defocusing character of the problem only plays a role in the large data case, see e.g. Dodson's results in [12, 13].

Thus, the interesting class of open problems remains the quasilinear one, where Strichartz estimate are no longer readily available. For such problems, we formulate here the following conjecture:

**Conjecture 3** (Non-localized data GWP conjecture). *Two and higher dimensional quasilinear dispersive flows on the real line with cubic nonlinearities and small initial data have global in time, scattering solutions.*

Here scattering is interpreted in the classical fashion, with the only proviso that, due to the nature of quasilinear well-posedness, convergence to a linear evolution can only be expected in a weaker topology.

While for convenience we have stated this conjecture in all dimensions, the most interesting case by far is the two dimensional case, where, at this point, no small data global well-posedness result exists. This is in part due to the fact that in two space dimensions  $L^4$  is a sharp Strichartz norm, whereas in higher dimensions there is some room. Indeed, in dimension three and higher global well-posedness has been proved for very specific models, using a combination of energy estimates and constant coefficient Strichartz estimates, properly interpolated, see e.g. [22].

In this article we will prove this conjecture for all cubic quasilinear Schrödinger flows. This requires no additional structural assumptions in dimension three and higher, and also in dimension two at higher regularity. However, in order to reach the local well-posedness Sobolev threshold in dimension two, we require the flow to be conservative.

**1.2. Quasilinear Schrödinger flows: local solutions.** The most general form for a quasilinear Schrödinger flow in one space dimension is

$$(DQNLS) \quad \begin{cases} iu_t + g(u, \partial_x u) \partial_x^2 u = N(u, \partial_x u), & u : \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{C} \\ u(0, x) = u_0(x), \end{cases}$$

with a metric  $g$  which is a real valued function and a source term  $N$  which is a complex valued smooth function of its arguments. Here smoothness is interpreted in the real sense. But if  $g$  and  $N$  are analytic, then they can also be thought of as separately (complex) analytic functions of  $u$  and  $\bar{u}$ . This is the interpretation we use in the present paper.

In parallel we consider the simpler problem

$$(QNLS) \quad \begin{cases} iu_t + g(u) \partial_x^2 u = N(u, \partial_x u), & u : \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{C} \\ u(0, x) = u_0(x), \end{cases}$$

where  $N$  is at most quadratic in  $\partial u$ . This can be seen as the differentiated form of (DQNLS). We will state our results for both of these flows, but, in order to keep the exposition focused on the important issues, we will provide complete arguments only for (QNLS).

In the present work we make the key assumption that

$$(1.1) \quad g(0) = I_n,$$

which can be easily generalized to any positive definite matrix. But there has also been considerable interest in the ultra-hyperbolic case, where  $g(0)$  is a symmetric nondegenerate matrix of indefinite case.

The first question we consider here is the local well-posedness question in Sobolev spaces. This is not at all a straightforward question, and is in particular much harder than the corresponding question for nonlinear wave equations. One primary reason for this is the infinite speed of propagation, which introduces short time growth effects which are only visible over long time scales for hyperbolic problems. This goes back to the work of Takeuchi [59], Mizohata [53] and Ichinose [27].

One effect of this is that in problems with quadratic nonlinearities  $N$ , one in general does not have local well-posedness in any  $H^s$  space, at least not without making additional decay assumptions at infinity. This happens even for semilinear equations where the metric  $g$  is simply the euclidean one.

Historically, the first problems of this type which were studied were the semilinear ones, starting with the work of Kenig-Ponce-Vega [43] for small data, Hayashi-Ozawa [20] and then Chihara [6] for large data, also using ideas of Doi [14]. Finally, Kenig-Ponce-Vega [44] also considered the ultrahyperbolic case, where the Laplacian is replaced by a nondegenerate second order operator with indefinite signature. All the results above are for localized data, and the first result in translation invariant Sobolev spaces was due to Bejenaru-Tataru [2].

Moving on to the full quasilinear case, the first problems that were considered there were the one results ones, for which we refer the reader to due to de Bouard-Hayashi-Naumkin-Saut [11], Colin [7], Poppenberg [56] and Lin-Ponce [47], leading up to the authors' recent article [36].

The first local well-posedness result for the quasilinear problem in higher dimension goes back to work of Kenig-Ponce-Vega [45], where these equations were studied for regular and

localized initial data. Similar results in the ultra-hyperbolic case were also obtained by Kenig-Ponce-Vega-Rovlung [41, 42]. The next step was to study the local well-posedness question in translation invariant Sobolev spaces and at lower regularity; this was accomplished by Marzuola-Metcalf-Tataru, first for small data in [50, 51], and then for large data in [52]; the latter article considers only the elliptic case, while the ultra-hyperbolic case was only recently studied by Pineau-Taylor [54], also related to the work of Jeong-Oh [38]. Another very recent work of Shao-Zhou [57] provides a nice alternate approach for the results of [45, 51].

One key distinction made in these last papers was between quadratic and cubic nonlinearities. It is only in the latter case that the local well-posedness is proved in classical  $H^s$  Sobolev spaces. Another distinction is between small and large data; in the latter case, large energy growth may occur on arbitrarily short time scales, and a nontrapping condition is also required.

For work on other related models we refer the reader to Colin [8], Chemin-Salort [5], as well as the work of Huang-Tataru [23, 24] as well as the small data global result of Huang-Li-Tataru [22]; the last three articles are concerned with the skew mean curvature flow, which can be formulated as a quasilinear Schrödinger evolution when represented in a suitable gauge.

In order to work in  $H^s$  Sobolev spaces, in the present article we restrict our attention to the cubic case. By this, we mean

**Definition 1.1.** *We say that the equation (QNLS)/(DQNLS) is cubic if  $g$  is at least quadratic and  $N$  is at least cubic at zero.*

For reference and easy comparison, we state the earlier results here, specialized to cubic nonlinearities:

**Theorem 1** (cubic nonlinearities [51, 52]). *The  $nD$  cubic problem (QNLS) is locally well-posed for nontrapping data in  $H^s$  for  $s > \frac{n+3}{2}$ , and the cubic problem (DQNLS) is locally well-posed in  $H^s$  for  $s > \frac{n+5}{2}$ .*

The nontrapping condition above is needed for large data only, as it is automatically satisfied for small data. In the 1D case, this result was improved in [36] up to the natural, optimal Sobolev threshold:

**Theorem 2** (1D cubic nonlinearities [36]). *The 1D cubic problem (QNLS) is locally well-posed for small initial data in  $H^s$  for  $s > 1$ , and the 1D cubic problem (DQNLS) is locally well-posed in  $H^s$  for  $s > 2$ .*

This is not only a major improvement over Theorem 1, it is also sharp for generic problems of this type, see the heuristic arguments in [37].

We now turn our attention to the first objective of this paper, which is to study the same local well-posedness problem in two and higher space dimensions. Our results are as follows:

**Theorem 3.** *Let  $n = 2$ . The 2D cubic problem (QNLS) is locally well-posed for small data in  $H^s$  for  $s > \frac{3}{2}$ , and the 2D cubic problem (DQNLS) is locally well-posed for small data in  $H^s$  for  $s > \frac{5}{2}$ .*

**Theorem 4.** *Let  $n \geq 3$ . The  $nD$  cubic problem (QNLS) is locally well-posed for small data in  $H^s$  for  $s > \frac{n+1}{2}$ , and the  $nD$  cubic problem (DQNLS) is locally well-posed for small data in  $H^s$  for  $s > \frac{n+3}{2}$ .*

Here well-posedness is interpreted in the Hadamard sense, and in effect in an enhanced Hadamard form, which includes :

- existence of solutions in  $C([0, T]; H^s)$ .
- uniqueness of regular solutions.
- uniqueness of rough solutions as uniform limits of regular solutions.
- continuous dependence on the initial data.
- Lipschitz dependence of the solutions on the initial data in a weaker topology ( $L^2$ )
- higher regularity: more regular data yields more regular solutions.

These results do not require any structure for the nonlinearity, other than that it is cubic. While the formulation is identical in two and higher dimensions, we have stated the two results separately in order to emphasize the fact the proofs are different, with the two dimensional case being the more difficult one.

To better understand their significance, one may compare both of these results with the leading order scaling for these problems, which corresponds to the exponents  $s_c = \frac{n}{2}$  for (QNLS), respectively  $s_c = \frac{n+2}{2}$  for (QNLS). So our results are only 1/2 derivative above scaling, which is similar to the earlier one dimensional result. However, this similarity at the level of the results does not translate into a similarity at the level of the proofs.

Concerning the optimality of the local well-posedness results in the above theorems, we propose the following

**Conjecture 4.** *The range of  $s$  in Theorems 3,4 is generically sharp, except possibly for the endpoint.*

This is however due to a different reason than in the one dimensional case, where the main obstruction appears to come from the self-interaction of a single wave-packet, see [37]. Instead, in the higher dimensional setting the obstruction seem related to trapping, which generically should fail for  $s$  below our thresholds even in the small data case.

In a related vein, we also propose

**Conjecture 5.** *The results in Theorems 3,4 also hold for large data, under an additional nontrapping assumption.*

One main difference between various dimensions arises at the level of the Strichartz estimates. In one dimension we have a better toolbox for short time Strichartz, see [36]. On the other hand in dimension three and higher the  $L^4$  Strichartz bound is obtained directly from the bilinear analysis. The most difficult case is the two dimensional one, for which the 2D toolbox is much smaller than in the 1D case, while the bilinear estimates are weaker than in higher dimension.

The starting point for the proof of our results is provided by the earlier results stated in Theorem 1. The lifespan of these solutions, say for (QNLS), apriori depends on the  $H^s$  size of the initial data, which is not necessarily small. However, once we prove suitable a-priori bounds on these solutions, we can extend the lifespan so that it depends only on the  $H^s$  size of the data.

Once we have regular solutions on uniform time scales, we produce the rough,  $H^s$  solutions as unique uniform limits of regular solutions. This requires careful estimates for the linearized flow, which we do at the  $L^2$  level.

**1.3. Quasilinear Schrödinger flows: global solutions.** Our objective here is to consider the global well-posedness for the 2D problem for initial data which is small in  $H^s$ , in the spirit of Conjecture 3.

This 1D counterpart of this conjecture was proved in [36], and we recall this first. By contrast to the local well-posedness result, refined long time and global in time results in 1D require some natural structural assumptions on the equations, which will be described next.

**Definition 1.2.** *We say that the equation (QNLS)/(DQNLS) has phase rotation symmetry if it is invariant with respect to the transformation  $u \rightarrow ue^{i\theta}$  for  $\theta \in \mathbb{R}$ .*

One consequence of this assumption is that the Taylor series of the nonlinearity only has odd terms, which are multilinear expressions in  $u$  and  $\bar{u}$  (and their derivatives) with one more  $u$  factor. It suffices in effect to assume that this holds for the cubic terms in the equation.

To state the next assumptions it is convenient to write the equations in a semilinear way. Assuming the phase rotation symmetry, and also harmlessly assuming that  $g(0) = 1$ , this semilinear form is

$$(1.2) \quad iu_t + \partial_x^2 u = C(u, \bar{u}, u) + \text{higher order terms},$$

where  $C$  is a translation invariant trilinear form with symbol  $c(\xi_1, \xi_2, \xi_3)$ . We refer the reader to Section 2 for a description of our notations.

**Definition 1.3.** *We say that the 1D equation (QNLS)/(DQNLS) is conservative if the cubic component of the nonlinearity satisfies*

$$c(\xi, \xi, \xi), \partial_{\xi_j} c(\xi, \xi, \xi) \in \mathbb{R}.$$

It is also immediately satisfied if the problem admits a coercive conservation law, but, conversely, it does not imply the existence of such a conservation law.

One final necessary assumption in 1D is that the problem is defocusing:

**Definition 1.4.** *We say that the equation (QNLS)/(DQNLS) is defocusing if the cubic component of the nonlinearity satisfies*

$$c(\xi, \xi, \xi) \gtrsim 1 + \xi^2.$$

Now we can state the one dimensional result:

**Theorem 5.** *Consider a 1D (QNLS)/(DQNLS) problem, which we assume to be cubic, phase rotation invariant and defocusing. Assume that the initial data  $u_0$  is small in  $H^s$ , where  $s > 1$  for (QNLS) and  $s > 2$  for (DQNLS). Then the solutions are global in time, and satisfy*

$$(1.3) \quad \|u\|_{L_t^\infty H_x^s} \lesssim \|u_0\|_{H_x^s}.$$

Now we turn our attention to the higher dimensional problem, which differs from the 1D case in two key ways:

- (1) The dispersive decay is stronger in nD, where cubic terms are even perturbative in the semilinear case; this makes the problem easier, and in particular it allows us to discard the defocusing assumption.
- (2) There are many more cubic resonant interactions in nD; this makes the problem harder, and potentially it mostly disables normal form/modified energy type arguments.

The first feature above allows us to weaken the hypothesis of the theorem, compared to the  $1D$  case, dispensing with both the phase rotation invariance, the conservative assumption and the defocusing condition. Without phase rotation the expansion in (1.2) acquires extra terms,

$$(1.4) \quad iu_t + \partial_x^2 u = C(u, \bar{u}, u) + \text{non rotation invariant cubic terms} + \text{higher order terms},$$

where all wave-packet self-interactions are captured by the  $C$  term. This term does not play any role in higher dimension  $n \geq 3$ , but we do need to consider its properties in dimension  $n = 2$ . There we introduce the following more relaxed conservative condition:

**Definition 1.5.** *We say that the 2D equation (QNLS)/(DQNLS) is conservative if the cubic component of the nonlinearity satisfies*

$$c(\xi, \xi, \xi) \in \mathbb{R}, \quad \xi \in \mathbb{R}.$$

After these preliminaries we are ready to state our global results. We begin with the higher dimensional case  $n \geq 3$ , where we work at the same regularity level as in the local well-posedness result in Theorem 4:

**Theorem 6.** *Let  $n \geq 3$ . a) Assume that the equation (QNLS) is cubic, and that the initial data  $u_0$  is small in  $H^s$ , with  $s > \frac{n+1}{2}$ ,*

$$(1.5) \quad \|u_0\|_{H^s} \leq \epsilon \ll 1.$$

*Then the solutions are global in time, satisfy the uniform bound*

$$(1.6) \quad \|u\|_{L_t^\infty H_x^s} \lesssim \epsilon,$$

*and scatter at infinity, in the sense that*

$$(1.7) \quad u_\infty = \lim_{t \rightarrow \infty} e^{-it\Delta} u(t) \quad \text{in } H^s$$

*exists and has a continuous dependence on the initial data in the same topology.*

*b) The same result holds for the equation (DQNLS) for  $s > \frac{n+3}{2}$ .*

Next we have a similar result in two space dimensions, but for a more restrictive class of exponents, namely  $1/4$  derivative above the local well-posedness threshold in Theorem 3:

**Theorem 7.** *Let  $n = 2$ . a) Assume that the equation (QNLS) is cubic, and that the initial data  $u_0$  is small in  $H^s$ , with  $s \geq \frac{n+1}{2} + \frac{1}{4}$ ,*

$$(1.8) \quad \|u_0\|_{H^s} \leq \epsilon \ll 1.$$

*Then the solutions are global in time, satisfy the uniform bound*

$$(1.9) \quad \|u\|_{L_t^\infty H_x^s} \lesssim \epsilon,$$

*and scatter at infinity, in the sense that*

$$(1.10) \quad u_\infty = \lim_{t \rightarrow \infty} e^{-it\Delta} u(t) \quad \text{in } H^s$$

*exists and has a continuous dependence on the initial data in the same topology.*

*b) The same result holds for the equation (DQNLS) for  $s \geq \frac{n+3}{2} + \frac{1}{4}$ .*

Finally, in the low regularity case in dimension  $n = 2$ , under an additional conservative assumption, we are able to bring the global well-posedness result to the same regularity level as the local well-posedness result:



**Theorem 8.** *Let  $n = 2$ . a) Assume that the equation (QNLS) is cubic and conservative, and that the initial data  $u_0$  is small in  $H^s$ , with  $s > \frac{n+1}{2}$ ,*

$$(1.11) \quad \|u_0\|_{H^s} \leq \epsilon \ll 1.$$

*Then the solutions are global in time, satisfy the uniform bound*

$$(1.12) \quad \|u\|_{L_t^\infty H_x^s} \lesssim \epsilon,$$

*and scatter at infinity, in the sense that*

$$(1.13) \quad u_\infty = \lim_{t \rightarrow \infty} e^{-it\Delta} u(t) \quad \text{in } H^s$$

*exists and has a continuous dependence on the initial data in the same topology.*

*b) The same result holds for the equation (DQNLS) for  $s > \frac{n+3}{2}$ .*

As noted earlier, these results represent the first global well-posedness results for general cubic quasilinear Schrödinger equations with small initial data in Sobolev spaces in dimension  $n \geq 2$ . Indeed, all prior global results are of following two types:

- (1) Problems with initial data which is not only small but also *smooth and localized*, see for instance Ginibre-Hayashi [16], Hayashi-Naumkin [18] and Bernal-Vilchis-Hayashi-Naumkin. [3].
- (2) Specific problems with additional structure in higher dimension  $n \geq 3$ , see e.g. Huang-Li-Tataru [22].

The contrast is even more stark in dimension  $n = 2$ , where as far as we are aware there is no global result known for any cubic quasilinear dispersive problem with small but non-localized initial data.

We continue with two remarks about the scattering result:

**Remark 1.6.** *One should compare the results on scattering with our earlier 1D results in [36]. In the 1D case, the cubic nonlinear effects are too strong to prevent any standard scattering; even for localized data, the best one could hope for is some form of modified scattering. By contrast, in two and higher dimension we have full classical scattering.*

**Remark 1.7.** *A natural question here is to study the regularity of the wave operator  $u_0 \rightarrow u_\infty$ . Our analysis implies that it has properties akin to the (quasilinear) Hadamard well-posedness:*

- *it is continuous in  $H^s$*
- *it is Lipschitz continuous<sup>1</sup> in a weaker topology (e.g.  $L^2$ ), and in effect close to the identity in the Lipschitz topology.*

We note that the above results are stated here in a short form, but the proofs shows in effect that the solutions satisfy several types of estimates:

- (i) Bilinear  $L^2$  bounds, which can be stated in a balanced form

$$(1.14) \quad \| |D|^{\frac{3-n}{2}} |\langle D \rangle^{s-\frac{1}{4}} u|^2 \|_{L_{t,x}^2} \lesssim \epsilon^2,$$

and in an imbalanced form, using standard paraproduct notation,

$$(1.15) \quad \|T_{\partial u} \bar{u}\|_{L_t^2 H_x^{s+\frac{1}{2}}} \lesssim \epsilon^2.$$

---

<sup>1</sup>When restricted to small  $H^s$  data.

(ii) Lossless  $L^4$  Strichartz bounds for  $n \geq 3$ :

$$(1.16) \quad \|\langle D \rangle^{s-\frac{2-n}{4}} u\|_{L_{t,x}^4} \lesssim \epsilon.$$

(iii) Lossless  $L^4 L^8$  Strichartz bounds for  $n = 2$

$$(1.17) \quad \|\langle D \rangle^{s-\frac{1}{4}} u\|_{L_t^4 L_x^8} \lesssim \epsilon.$$

(iv) Full Strichartz bounds with derivative loss:

$$(1.18) \quad \|\langle D \rangle^{s-\frac{2}{p}} u\|_{L_t^p L_x^q} \lesssim \epsilon$$

for any pair  $(p, q)$  of sharp Strichartz exponents.

The reader is referred to Theorem 9 and Theorem 10 for a more complete, frequency envelope based description of the results. One may also compare the  $L_{t,x}^4$  bounds above with the  $L_{t,x}^6$  bounds which play the leading role in the 1D case, where 6 is the sharp Strichartz exponent.

One may also frame our results with the broader question of obtaining long time solutions for one or two dimensional dispersive flows with quadratic/cubic nonlinearities, which has attracted a lot of attention in recent years. One can distinguish two different but closely related types of results that have emerged, as well as several successful approaches.

On one hand, *normal form methods* have been developed in order to extend the lifespan of solutions, beginning with [58] in the late '80's. Somewhat later, around 2000, the *I-method*, introduced in [9] brought forth the idea of constructing better almost conserved quantities. These two ideas serve well in the study of semilinear flows, where it was later understood that they are connected [4].

Neither of these techniques can be directly applied to quasilinear problems. Addressing this problem, it was discovered in the work of the authors and collaborators [26], [29] that one can adapt the normal form method to quasilinear problems by constructing energies which simultaneously capture both the quasilinear and the normal form structures. This idea was called the *modified energy method*, and can also be seen in some way as a quasilinear adaptation of the I-method. Other alternate approaches, also in the quasilinear setting, are provided by the *flow method* of Hunter-Ifrim [25], where a better nonlinear normal form transformation is constructed using a well chosen auxiliary flow, and by the paradiagonalization method of Alazard and-Delort [1], where a paradiifferential symmetrization is applied instead.

Going further, the question of obtaining scattering, global in time solutions for one or two dimensional dispersive flows with quadratic/cubic nonlinearities has also been extensively studied in the last two decades for a number of models, under the assumption that the initial data is both *small* and *localized*; for a few examples out of many, see for instance [17, 19, 49, 39, 28] in one dimension, and [15] in two dimensions. The nonlinearities in the one dimensional models are primarily cubic, though the analysis has also been extended via normal form and modified energy methods to problems which also have nonresonant quadratic interactions; several such examples are [1, 29, 12, 30, 48], see also further references therein, as well as the authors' expository paper [35]. On the other hand some quadratic nonlinearities can also be handled, see for instance [58], based on the idea of *space-time* resonances.

Comparing the above class of results where the initial data is both small and localized with the present results, without any localization assumption, it is clear that in the latter case the problem becomes much more difficult, because the absence of localization allows for far stronger nonlinear interactions over long time-scales. Another twist in the one dimensional setting was that one also needs to distinguish between the focusing and defocusing case, as was made clear in our earlier semilinear work in [33] and [34], and then quasilinear work in [36].

**1.4. Outline of the paper.** While the short form of the results provided in the introduction represents a good starting point, from the perspective of both proving these results and of understanding the global in time dispersive properties of the solutions it becomes essential to have an appropriate family of quantitative bounds for the solutions. As briefly indicated with the bounds (1.14)-(1.18), these bounds are roughly of two types:

**Strichartz estimates:** these represent the more classical side of it, and are well-known in the constant coefficient case, though proving such bounds in quasilinear settings becomes anything but straightforward.

**Bilinear  $L^2$  estimates:** these play the leading role here, and should be seen as transversality bounds, i.e. which in essence capture the bilinear interaction of two transversal waves. Again these are easily proved in linear, constant coefficient settings but not at all easy in any quasilinear context.

As it is frequently the case in the study of quasilinear problems, proving such estimates requires having a considerable amount of information on the solutions to start with, which is why the entire proof of the results is wrapped within a carefully designed bootstrap argument, where, to start with, one assumes that a weaker version of such estimates already holds, not only at the linear level but also at the bilinear level.

While the bootstrap requirements make it a necessity that the main line of our argument cannot close until the last section, we are still able to organize the proofs in a modular fashion, with the only proviso that the bootstrap assumptions are imposed at each step, but not fully recovered until the very end. These modules are as follows:

**I. Littlewood-Paley theory and frequency envelopes.** While the estimates (1.14)-(1.18) serve give the reader an idea of our overall set-up, for our proofs it is far better to have a more precise accounting of these estimates relative to the size of the frequency. This is achieved first by using a Littlewood-Paley decomposition of the solutions, and secondly by using the language of frequency envelopes in order to track the size of the dyadic pieces in both linear and bilinear bounds. This is described in Section 3, where we also provide sharper, frequency envelope versions of our estimates for the solutions, as well as the set-up of the global bootstrap argument.

**II. The paradifferential formalism.** As it is standard in the study of nonlinear evolutions, in order to prove local well-posedness we need good estimates not only for the solution itself, but also for the corresponding linearized equation. As it turns out, rather than study each of these two equations separately, it is better to seek a common denominator; this turns out to be the associated linear paradifferential flow, which will indeed play the leading role. In Section 4 we introduce all these related flows, and recast both the full equation and the linearized equation as paradifferential flows with source terms. These source terms should ideally play a perturbative role, and indeed they do with two provisos: (i) that we have

a suitable notion of “perturbative” which is sufficiently broad, and (ii) that in the most difficult case of two dimensional flows at low regularity we give special consideration to the cubic balanced terms.

**III. Conservation laws in density-flux form.** Efficient energy estimates must necessarily play an important role in our analysis, but for the proof of the bilinear estimates we need to consider these estimates in a frequency localized manner, and, more importantly, as local conservation laws in density-flux form. This is discussed primarily in Section 6, and refined further in the last section for 2D flows.

**IV. Interaction Morawetz bounds.** These represent the primary tool in the proof of the crucial bilinear  $L^2$  bounds, and originate in work of the I-team in [10]. However our starting point is provided by the work of Planchon-Vega [55] for the constant coefficient case. We modify and adapt this to our purposes in Section 7, following the ideas developed in our prior one dimensional work in [36]. However, the higher dimensional case turns out to be more difficult due to a certain lack of coercivity, which considerably increases the burden of estimating “perturbative” contributions.

**V. Strichartz estimates in the paradifferential world.** Directly proving Strichartz estimates for the paradifferential was possible in [36] in the 1D case locally in time, but it becomes a daunting task in higher dimension. This is partially rectified by the stronger bilinear bounds in the balanced case, which in particular yields a critical scale invariant  $L^4$  bound in dimension three and higher. But in 2D this is no longer the case, so in Section 8 we develop instead a strategy to prove the Strichartz estimates for a full range of exponents, globally in time, but with a loss of derivatives.

**VI. Rough solutions as limits of smooth solutions.** This is by now a standard step in obtaining rough solutions for quasilinear flows, which is best carried out under the umbrella of frequency envelopes. We do this in Section 9, following the path described in our earlier expository paper [32].

**VII. The 2D global result at low regularity.** While cubic balanced terms can be seen as perturbative in dimension three and higher, this is no longer the case in two space dimensions, particularly at low regularity. This is where our “conservative” assumption plays an important role, which is investigated in the last section of the paper.

**1.5. Acknowledgements.** The first author was supported by the Sloan Foundation, and by an NSF CAREER grant DMS-1845037. The second author was supported by the NSF grant DMS-2054975 as well as by a Simons Investigator grant from the Simons Foundation.

## 2. NOTATIONS AND PRELIMINARIES

Here we introduce the usual Littlewood-Paley decomposition, and discuss some standard facts for constant coefficient linear Schrödinger flows, including the Strichartz and bilinear  $L^2$  estimates, as well as the notion of cubic resonant interactions. Finally, we describe our notations for multilinear forms and their symbols.

**2.1. The Littlewood-Paley decomposition.** We use a standard Littlewood-Paley decomposition in frequency: let  $\psi$  be a bump function adapted to  $[-2, 2]$  and equal to 1 on  $[-1, 1]$ . We define the Littlewood-Paley operators  $P_k$ , as well as  $P_{\leq} = P_{\leq k+1}$  for  $k \in \mathbb{N}$  by defining

$$\widehat{P_{\leq k} f}(\xi) := \psi(2^{-k}\xi) \hat{f}(\xi)$$

with  $P_1 := P_{\leq 1}$  and  $P_k := P_{\leq k} - P_{\leq k-1}$  for  $k \geq 2$ . All the operators  $P_k$ ,  $P_{\leq k}$  are bounded on all translation-invariant Banach spaces, thanks to Minkowski's inequality. We also define  $P_{>k} := P_{\geq k+1} := 1 - P_{\leq k}$ .

Thus

$$1 = \sum_{k \in \mathbb{N}} P_k,$$

where the multipliers  $P_k$  have smooth symbols localized at frequency  $2^k$ . Correspondingly, our solution  $u$  will be decomposed as

$$u = \sum_{k \in \mathbb{N}} u_k, \quad u_k = P_k u.$$

The main estimates we will establish for our solution  $u$  will be linear and bilinear estimates for the functions  $u_k$ .

In the proof of the local well-posedness result in Section 9 it will be convenient to switch from a discrete to a continuous Littlewood-Paley decomposition. There we think of  $k$  as a real nonnegative parameter and define

$$P_k u := \frac{d}{dk} P_{<k} u,$$

and our Littlewood-Paley decomposition for  $u$  becomes

$$u = u_0 + \int_0^\infty u_k dk.$$

To shorten some calculations we will also use Greek indices for Littlewood-Paley projectors, e.g.  $P_\lambda$ ,  $P_\mu$  where we take  $\lambda, \mu \in 2^\mathbb{N}$ . Correspondingly, our Littlewood-Paley decomposition of a function  $u$  will read

$$u = \sum_{\lambda \in 2^\mathbb{N}} u_\lambda, \quad u_\lambda = P_\lambda u.$$

**2.2. Strichartz and bilinear  $L^2$  bounds.** Here we begin by recalling the Strichartz inequalities, which apply to solutions to the inhomogeneous linear Schrödinger equation:

$$(2.1) \quad (\partial_t + \Delta)u = f, \quad u(0) = u_0.$$

These are  $L_t^p L_x^q$  bounds which are used as measures of the dispersive effect. The allowed exponents  $(p, q)$  depend on the dimension:

**Definition 2.1.** *The pair  $(p, q)$  is a standard Strichartz exponent in  $n$  space dimensions if*

$$(2.2) \quad \frac{2}{p} + \frac{n}{q} = \frac{2}{n}, \quad 2 \leq p, q \leq \infty$$

*with the only exception of the forbidden endpoint  $(2, \infty)$  in dimension  $n = 2$ .*

With this definition, the Strichartz estimates in the  $L^2$  setting are summarized in the following

**Lemma 2.2.** *Assume that  $u$  solves (2.1) in  $[0, T] \times \mathbb{R}$ . Then the following estimate holds for all sharp Strichartz pairs  $(p, q), (p_1, q_1)$ :*

$$(2.3) \quad \|u\|_{L_t^p L_x^q} \lesssim \|u_0\|_{L^2} + \|f\|_{L_t^{p'_1} L_x^{q'_1}}.$$

For further details and references we direct the reader to [40], where the final endpoint  $p = 2$  was proved.

For practical purposes it is useful to place all these estimates under a single umbrella. In dimension three and higher we have access to both end-points, and it is most efficient to define the Strichartz space  $S$  associated to the  $L^2$  flow by

$$S = L_t^\infty L_x^2 \cap L_t^2 L_x^{\frac{2n}{n-2}},$$

as well as its dual

$$S' = L_t^1 L_x^2 + L_t^2 L_x^{\frac{2n}{n+2}}.$$

Then the Strichartz estimates can be summarized all as

$$(2.4) \quad \|u\|_S \lesssim \|u_0\|_{L^2} + \|f\|_{S'}.$$

In two space dimensions we lose access to the  $(2, \infty)$  endpoint so the above definition of the Strichartz space  $S$  no longer applies. Instead, it is convenient to use the  $U_\Delta^p$  and  $V_\Delta^p$  spaces. These were introduced in unpublished work of the second author [62], see also [46] and [21] for some of the first applications of these spaces. In this setting the Strichartz estimates can be summarized as

$$(2.5) \quad \|u\|_{V_\Delta^2} \lesssim \|u_0\|_{L^2} + \|f\|_{DV_\Delta^2},$$

where the transition to the Strichartz bounds is provided by the embeddings

$$(2.6) \quad V_\Delta^2 \subset U_\Delta^p \subset L_t^p L_x^q, \quad L_t^{p'} L_x^{q'} \subset DV_\Delta^{p'} \subset DV_\Delta^2.$$

The last property of the linear Schrödinger equation we will use here is the bilinear  $L^2$  estimate, which is as follows:

**Lemma 2.3.** *Let  $u^1, u^2$  be two solutions to the inhomogeneous Schrödinger equation with data  $u_0^1, u_0^2$  and inhomogeneous terms  $f^1$  and  $f^2$ . Assume that  $u^1$  and  $u^2$  are frequency localized in balls  $B_1, B_2$  with radius  $\lambda_1 \lesssim \lambda_2$  so that*

$$d(B_1, B_2) \gtrsim \lambda_2.$$

*Then we have*

$$(2.7) \quad \|u^1 u^2\|_{L_{t,x}^2} \lesssim \lambda_1^{\frac{n-1}{2}} \lambda_2^{-\frac{1}{2}} (\|u_0^1\|_{L_x^2} + \|f^1\|_{S'}) (\|u_0^2\|_{L_x^2} + \|f^2\|_{S'}).$$

**2.3. Resonant analysis.** In this subsection we aim to classify the trilinear interactions associated to the linear constant coefficient flow; for simplicity, the reader may think of the context of an equation of the form

$$(i\partial_t + \Delta)u = C(\mathbf{u}, \mathbf{u}, \mathbf{u}),$$

where  $C$  is a translation invariant trilinear form and we have used the notation  $\mathbf{u} = \{u, \bar{u}\}$  in order to allow for both  $u$  and  $\bar{u}$  in each of the arguments of  $C$ . For clarity, we note that this includes expressions of the form

$$C(u, \bar{u}, u), \quad C(u, u, u), \quad C(u, \bar{u}, \bar{u}), \quad C(\bar{u}, \bar{u}, \bar{u}).$$

Within these choices we distinguish the first one, namely  $C(u, \bar{u}, u)$ , as the only one with *phase rotation symmetry*.

Given three input frequencies  $\xi^1, \xi^2, \xi^3$  for our cubic nonlinearity, the output will be at frequency

$$(2.8) \quad \xi^4 = \pm \xi^1 \pm \xi^2 \pm \xi^3,$$

where the signs are chosen depending on whether we use the input  $u$  or  $\bar{u}$ . We call such an interaction balanced if all  $\xi^j$ 's are of comparable size, and *unbalanced* otherwise. This is a *resonant interaction* if and only if we have a similar relation for the associated time frequencies, namely

$$|\xi^4|^2 = \pm |\xi^1|^2 \pm |\xi^2|^2 \pm |\xi^3|^2.$$

On the other hand from the perspective of bilinear estimates we seek to distinguish the *transversal interactions*, which are defined as those for which no more than two of the frequencies  $\xi^j$  coincide. We seek to classify interactions from this perspective:

**(i) With phase rotation symmetry:** Here the relation (2.8) can be expressed in a more symmetric fashion as

$$\Delta^4 \xi = 0, \quad \Delta^4 \xi = \xi^1 - \xi^2 + \xi^3 - \xi^4.$$

This is a resonant interaction if and only if we have a similar relation for the associated time frequencies, namely

$$\Delta^4 \xi^2 = 0, \quad \Delta^4 \xi^2 = |\xi^1|^2 - |\xi^2|^2 + |\xi^3|^2 - |\xi^4|^2.$$

Hence, we define the resonant set in a symmetric fashion as

$$\mathcal{R} := \{\Delta^4 \xi = 0, \Delta^4 \xi^2 = 0\}.$$

In 1D it is easily seen that this set may be characterized as

$$\mathcal{R} = \{\{\xi^1, \xi^3\} = \{\xi^2, \xi^4\}\}.$$

However, in higher dimension this is no longer the case, and the resonant set  $\mathcal{R}$  consists of all quadruples  $\xi^1, \xi^2, \xi^3$  and  $\xi^4$  which are the vertices of a rectangle, in this order. Of these, the only non-transversal interactions are given by the diagonal set

$$\mathcal{R}_2 = \{\xi^1 = \xi^2 = \xi^3 = \xi^4\}.$$

We will refer to these interactions as the *doubly resonant interactions*.

**(ii) Without phase rotation symmetry:** Here we concentrate our attention on describing the worst case scenario, which corresponds to interactions which are both resonant and non-transversal. The first requirement corresponds to frequencies where

$$\xi^1 \pm \xi^2 \pm \xi^3 \pm \xi^4 = 0,$$

and

$$|\xi^1|^2 \pm |\xi^2|^2 \pm |\xi^3|^2 \pm |\xi^4|^2 = 0.$$

The second requirement is that three of the frequencies are equal. Without any loss in generality assume that  $\xi^1 = \xi^2 = \xi^3 := \xi$ . Depending on the choice of signs and excluding the case of phase rotation symmetry, we obtain one of the following systems:

$$\xi^4 = \pm 3\xi, \quad |\xi^4|^2 = \pm 3|\xi|^2,$$

with matched signs, or

$$\xi^4 = -\xi, \quad |\xi^4|^2 = -|\xi|^2.$$

In both cases the only solution is  $\xi = 0$ . Hence we have established the following

**Lemma 2.4.** *Except for the case  $(0,0,0,0)$ , all cubic interactions without phase rotation symmetry are either nonresonant or transversal.*

We will use the above classification to separate the cubic interactions using smooth frequency cut-offs as

$$C(\mathbf{u}, \mathbf{u}, \mathbf{u}) = C^{res}(u, \bar{u}, u) + C^0(\mathbf{u}, \mathbf{u}, \mathbf{u}) + C^{mr}(\mathbf{u}, \mathbf{u}, \mathbf{u}) + C^{tr}(\mathbf{u}, \mathbf{u}, \mathbf{u}),$$

where

- $C^{res}(u, \bar{u}, u)$  contains balanced interactions with phase rotation symmetry
- $C^0(\mathbf{u}, \mathbf{u}, \mathbf{u})$  contains only low frequency interactions
- $C^{mr}(\mathbf{u}, \mathbf{u}, \mathbf{u})$  contains only balanced nonresonant interactions
- $C^{tr}(\mathbf{u}, \mathbf{u}, \mathbf{u})$  contains only transversal interactions, balanced or unbalanced.

**2.4. Translation invariant multilinear forms.** As our problem and many of our estimates are invariant with respect to translations, it is natural that translation invariant multilinear forms play an important role in the analysis.

The simplest examples of multilinear forms are multipliers, which are convolution operators,

$$m(D)u(x) = \int K(y)u(x-y)dy, \quad \hat{K} = m.$$

More generally, we will denote by  $L$  any convolution operator

$$Lu(x) = \int K(y)u^y(x)dy,$$

where  $K$  is an integrable kernel, or a bounded measure, with a universal bound and where

$$u^y(x) := u(x-y),$$

is just a shorthand notation for translations of  $u$  by  $y$ .

Throughout the paper we will use multilinear operators, and denote by  $L$  any multilinear form

$$L(u_1, \dots, u_k)(x) = \int K(y_1, \dots, y_k)u_1^{y_1}(x) \cdots u_k^{y_k}(x)dy,$$

where, again  $K$  is assumed to have a kernel which is integrable, or more generally, a bounded measure (we allow for the latter in order to be able to include products here). Multilinear forms satisfy the same bounds as corresponding products do, as long as we work in translation invariant Sobolev spaces. Such a form may be equivalently described via its symbol  $a(\xi^1, \dots, \xi^k)$ , as

$$L(\widehat{u_1, \dots, u_k})(\xi) = (2\pi)^{-\frac{n(k-1)}{2}} \int_{\xi^1 + \dots + \xi^k = \xi} a(\xi^1, \dots, \xi^k) \hat{u}_1(\xi^1) \cdots \hat{u}_k(\xi^k) d\xi^1 \cdots d\xi^{k-1},$$

where the symbol  $a$  is the inverse Fourier transform of the kernel  $K$ .



A special role in our analysis is played by multilinear forms which exhibit phase rotation symmetry, where the arguments  $u$  and  $\bar{u}$  are alternating. For such forms we borrow the notations from our earlier paper [33].

Precisely, for an integer  $k \geq 2$ , we will use translation invariant  $k$ -linear forms

$$(\mathcal{D}(\mathbb{R}^n))^k \ni (u_1, \dots, u_k) \rightarrow Q(u_1, \bar{u}_2, \dots) \in \mathcal{D}'(\mathbb{R}^n),$$

where the nonconjugated and conjugated entries are alternating.

Such a form is uniquely described by its symbol  $q(\xi_1, \xi_2, \dots, \xi_k)$  via

$$Q(u_1, \bar{u}_2, \dots)(x) = (2\pi)^{-nk} \int e^{i(x-x_1)\xi^1} e^{-i(x-x_2)\xi^2} \dots q(\xi^1, \dots, \xi^k) \\ u_1(x_1) \bar{u}_2(x_2) \dots dx_1 \dots dx_k d\xi^1 \dots d\xi^k,$$

or equivalently on the Fourier side

$$\mathcal{F}Q(u_1, \bar{u}_2, \dots)(\xi) = (2\pi)^{-\frac{n(k-1)}{2}} \int_D q(\xi^1, \dots, \xi^k) \hat{u}_1(\xi^1) \bar{\hat{u}}_2(\xi^2) \dots d\xi^1 \dots d\xi^{k-1},$$

where, with alternating signs,

$$D := \{\xi = \xi^1 - \xi^2 + \dots\}.$$

They can also be described via their kernel

$$Q(u_1, \bar{u}_2, \dots)(x) = \int K(x - x_1, \dots, x - x_k) u_1(x_1) \bar{u}_2(x_2) \dots dx_1 \dots dx_k,$$

where  $K$  is defined in terms of the Fourier transform of  $q$

$$K(x_1, x_2, \dots, x_k) = (2\pi)^{-\frac{kn}{2}} \hat{q}(-x_1, x_2, \dots, (-1)^k x_k).$$

These notations are convenient but slightly nonstandard because of the alternation of complex conjugates. Another important remark is that, for  $k$ -linear forms, the cases of odd  $k$ , respectively even  $k$  play different roles here, as follows:

i) The  $2k + 1$  multilinear forms will be thought of as functions, e.g. those which appear in some of our evolution equations.

ii) The  $2k$  multilinear forms will be thought of as densities, e.g. which appear in some of our density-flux pairs.

Correspondingly, to each  $2k$ -linear form  $Q$  we will associate a  $2k$ -linear functional  $\mathbf{Q}$  defined by

$$\mathbf{Q}(u_1, \dots, u_{2k}) := \int_{\mathbb{R}} Q(u_1, \dots, \bar{u}_{2k})(x) dx,$$

which takes real or complex values. This may be alternatively expressed on the Fourier side as

$$\mathbf{Q}(u_1, \dots, u_{2k}) = (2\pi)^{n(1-k)} \int_D q(\xi^1, \dots, \xi^{2k}) \hat{u}_1(\xi^1) \bar{\hat{u}}_2(\xi^2) \dots \bar{\hat{u}}_{2k}(\xi^{2k}) d\xi^1 \dots d\xi^{2k-1},$$

where, with alternating signs, the diagonal  $D_0$  is given by

$$D_0 = \{0 = \xi^1 - \xi^2 + \dots\}.$$

Note that in order to define the multilinear functional  $\mathbf{Q}$  we only need to know the symbol  $q$  on  $D_0$ .

### 3. A FREQUENCY ENVELOPE FORMULATION OF THE RESULTS

For expository purposes, the main results of this paper, namely Theorem 4, Theorem 3, Theorem 6, Theorem 7 and Theorem 8 are stated in a simplified form in the introduction. However, the full results that we prove provide a much more detailed picture, which gives a full family of bilinear  $L^2$  bounds for the solutions. Furthermore, the proofs of the results are complex bootstrap arguments relative to all these bounds, both linear and bilinear. For these reasons, it is important to have a good setup for both the results and for the bootstrap assumptions. An elegant way to do this is to use the language of frequency envelopes. Our goals in this section are

- (i) to define frequency envelopes,
- (ii) to provide a more accurate, frequency envelope version of our main results,
- (iii) to provide the bootstrap assumptions in the proofs of each of the theorems,
- (iv) to outline the continuity argument allowing us to use these bootstrap assumptions.

**3.1. Frequency envelopes.** Before stating one of the main theorems of this paper, we revisit the *frequency envelope* notion. This elegant and useful tool will streamline the exposition of our results, and one should think at it as a bookkeeping device that is meant to efficiently track the evolution of the energy of the solutions between dyadic energy shells.

Following Tao's paper [61] we say that a sequence  $c_k \in \ell^2$  is an  $L^2$  frequency envelope for  $\phi \in L^2$  if

i)  $\sum_{k=0}^{\infty} c_k^2 < \infty$ ;

ii) it is slowly varying,

$$c_j/c_k \leq 2^{\delta|j-k|}, \quad j, k \in \mathbb{N}$$

with a small universal constant  $\delta$ ;

iii) it bounds the dyadic norms of  $\phi$ , namely  $\|P_k \phi\|_{L^2} \leq c_k$ .

Given a frequency envelope  $c_k$  we define

$$c_{\leq k} := \left( \sum_{j \leq k} c_j^2 \right)^{\frac{1}{2}}, \quad c_{\geq k} := \left( \sum_{j \geq k} c_j^2 \right)^{\frac{1}{2}}.$$

In practice we may choose our envelopes in a minimal fashion so that we have the equivalence

$$\sum_{k=0}^{\infty} c_k^2 \approx \|u\|_{L^2}^2$$

and so that we also have

$$c_0 \approx \|u\|_{L^2}.$$

The same applies with respect to any  $H^s$  norm.

**Remark 3.1.** *Another useful variation is to weaken the slowly varying assumption to*

$$2^{-\delta|j-k|} \leq c_j/c_k \leq 2^{C|j-k|}, \quad j < k,$$

*where  $C$  is a fixed but possibly large constant. All the results in this paper are compatible with this choice. This offers the extra flexibility of providing higher regularity results by the same arguments.*

**3.2. The frequency envelope form of the results.** The goal of this section is twofold: (i) restate the bounds needed for our main results in Theorem 4, Theorem 3, Theorem 6 and Theorem 7 in the frequency envelope setting, and (ii) to set up the bootstrap argument for the proof of these bounds.

The set-up for the bootstrap is most conveniently described using the language of frequency envelopes. This was originally introduced in the context of dyadic Littlewood-Paley decompositions in work of Tao, see e.g. [60]. For another interesting general frequency envelope setup one can see [31], which was the first place to use it to bootstrap bilinear bounds.

To start with, we assume that the initial data has small size in  $H^s$ ,

$$\|u_0\|_{H^s} \lesssim \epsilon.$$

We consider a dyadic frequency decomposition for the initial data,

$$u_0 = \sum_{\lambda \in 2^{\mathbb{N}}} u_{0,\lambda}.$$

Then we place the initial data components under an admissible frequency envelope,

$$\|u_{0,\lambda}\|_{H^s} \leq \epsilon c_\lambda, \quad c \in \ell^2,$$

where the envelope  $\{c_\lambda\}$  is not too large,

$$\|c\|_{\ell^2} \approx 1.$$

Our goal will be to establish similar frequency envelope bounds for the solution. We now state the frequency envelope bounds that will be the subject of the next theorems. These are as follows:

(i) Uniform frequency envelope bound:

$$(3.1) \quad \|u_\lambda\|_{L_t^\infty L_x^2} \lesssim \epsilon c_\lambda \lambda^{-s}$$

(ii) Unbalanced bilinear  $L^2$  bounds:

$$(3.2) \quad \|u_\lambda \bar{u}_\mu^{x_0}\|_{L_{t,x}^2} \lesssim \epsilon^2 c_\lambda c_\mu \lambda^{-s-\frac{1}{2}} \mu^{-s+\frac{n-1}{2}}, \quad \mu \ll \lambda$$

(iii) Balanced bilinear  $L^2$  bound

$$(3.3) \quad \||D|^{\frac{3-n}{2}}(u_\lambda \bar{u}_\mu^{x_0})\|_{L_{t,x}^2} \lesssim \epsilon^2 c_\lambda c_\mu \lambda^{-2s+\frac{1}{2}}(1 + \lambda|x_0|), \quad \lambda \approx \mu$$

(iv) Strichartz bounds with a loss, for any sharp Strichartz exponents  $(p, q)$ :

$$(3.4) \quad \|u_\lambda\|_{L_t^p L_x^q} \lesssim \epsilon c_\lambda \lambda^{-s+\frac{2}{p}}.$$

We remark that, as a consequence of the balanced bilinear bound (3.3), one also obtains the linear bounds

$$(3.5) \quad \|u_\lambda\|_{L_{t,x}^4} \lesssim \epsilon c_\lambda \lambda^{-s+\frac{n-2}{4}}, \quad n \geq 3,$$

respectively

$$(3.6) \quad \|u_\lambda\|_{L_t^4 L_x^8} \lesssim \epsilon c_\lambda \lambda^{-s+\frac{1}{4}}, \quad n = 2.$$

We also remark on a special case of the Strichartz bound (3.4) which has a  $1/2$  derivative loss and will play a leading role in the sequel:

$$(3.7) \quad \|u_\lambda\|_{L^4_{t,x}} \lesssim \epsilon c_\lambda \lambda^{-s+\frac{1}{2}}, \quad n = 2.$$

Here we distinguish between the 2D case and higher dimensions. The 2D bounds bear a closer resemblance to the 1D case studied in [36]; this reflects the fact that there is less dispersion in 2D, and in particular the  $L^4$  Strichartz bound lies on the sharp Strichartz line and cannot be obtained directly using interaction Morawetz based tools. By contrast, in three and higher dimension we are able to prove the  $L^4$  bound directly from interaction Morawetz.

We also remark on the need to add translations to the bilinear  $L^2$  estimates. This is because, unlike the linear bounds (3.1) and (3.4) which are inherently invariant with respect to translations, bilinear estimates are not invariant with respect to separate translations for the two factors. Hence, in particular, they cannot be directly transferred to nonlocal bilinear forms. However, if we allow translations, then one immediate corollary of (3.3) is that for any bilinear form  $L$  with smooth and bounded symbol we have

$$(3.8) \quad \| |D|^{\frac{3-n}{2}} L(u_\lambda \bar{u}_\lambda) \|_{L^2_{t,x}} \lesssim \epsilon^2 c_{\lambda^2} \lambda^{-2s-\frac{1}{2}}.$$

This is essentially the only way we will use this translation invariance in our proofs. We also remark on the  $(1+\lambda|x_0|)$  factor, which appears in the balanced bilinear bounds but not in the unbalanced ones. Heuristically this is because in the unbalanced case transversality remains valid even when one of the metrics gets translated; on the other hand, in the balanced case we also estimate interactions of near parallel waves, and there transversality is very sensitive to changes of the metric, so there is a price to pay for translations.

We are now ready to state our local results, beginning with the case of three and higher space dimensions:

**Theorem 9.** *a) Let  $n \geq 3$ ,  $s > \frac{n+1}{2}$ ,  $\epsilon \ll 1$  and  $T > 0$ . Consider the equation (QNLS) with cubic nonlinearity and let  $u \in C[0, T; H^s]$  be a smooth solution with initial data  $u_0$  which has  $H^s$  size at most  $\epsilon$ . Let  $\{\epsilon c_\lambda\}$  be a frequency envelope for the initial data in  $H^s$ . Then the solution  $u$  satisfies the bounds (3.1), (3.2), (3.3) and (3.4) uniformly with respect to  $x_0 \in \mathbb{R}$ .*

*b) The same result holds for (DQNLS) but with  $s > \frac{n+3}{2}$ .*

We continue with the case of two space dimensions, where we provide several variants of the result:

**Theorem 10.** *a) Let  $n = 2$ ,  $s > \frac{3}{2}$ ,  $\epsilon \ll 1$  and  $T > 0$ . Consider the equation (QNLS) with cubic nonlinearity, and for which one of the following three conditions holds:*

- (i) either  $s \geq \frac{7}{4}$  and  $T$  is arbitrary large,*
- (ii) or  $s > \frac{3}{2}$  and  $T \ll \epsilon^{-6}$ ,*
- (iii) or  $s > \frac{3}{2}$ , the problem is conservative and  $T$  is arbitrarily large.*

*Let  $u \in C[0, T; H^s]$  be a smooth solution with initial data  $u_0$  which has  $H^s$  size at most  $\epsilon$ . Let  $\{\epsilon c_\lambda\}$  be a frequency envelope for the initial data in  $H^s$ . Then the solution  $u$  satisfies the bounds (3.1), (3.2), (3.3) and (3.4) uniformly with respect to  $x_0 \in \mathbb{R}$ .*

*b) The same result holds for (DQNLS) but with  $s$  increased by one.*

The results in Theorem 9 and Theorem 10 [(i), (iii)] apply independently of the size of  $T$ , so they yield the global in time solutions in Theorems 6, 7, 8, given an appropriate local well-posedness result.

We continue with a theorem that applies to the linearized equation, which will be essential in the proof of all of our well-posedness results:

**Theorem 11.** *Consider the cubic equation (QNLS) for  $n \geq 2$ . Let  $u$  be a solution as in Theorem 10 or Theorem 9. Let  $v$  be a solution to the linearized equation around  $u$  with  $L^2$  initial data  $v_0$  and with frequency envelope  $d_\lambda$ . Then  $v$  satisfies the following bounds:*

(i) *Uniform frequency envelope bound:*

$$(3.9) \quad \|v_\lambda\|_{L_t^\infty L_x^2} \lesssim \epsilon d_\lambda$$

(ii) *Balanced bilinear  $(v, v)$ - $L^2$  bound:*

$$(3.10) \quad \| |D|^{\frac{1}{2}}(v_\lambda \bar{v}_\mu^{x_0}) \|_{L_{t,x}^2} \lesssim d_\lambda d_\mu (1 + \lambda |x_0|), \quad \lambda \approx \mu$$

(iii) *Unbalanced bilinear  $(v, v)$ - $L^2$  bound:*

$$(3.11) \quad \|v_\lambda \bar{v}_\mu^{x_0}\|_{L_{t,x}^2} \lesssim d_\lambda d_\mu \mu^{\frac{1}{2}} \lambda^{-\frac{1}{2}}, \quad \mu \ll \lambda$$

(iv) *Balanced bilinear  $(u, v)$ - $L^2$  bound:*

$$(3.12) \quad \| |D|^{\frac{1}{2}}(u_\lambda \bar{v}_\mu^{x_0}) \|_{L_{t,x}^2} \lesssim c_\lambda d_\mu \lambda^{-s} (1 + \lambda |x_0|), \quad \lambda \approx \mu$$

(v) *Unbalanced bilinear  $(u, v)$ - $L^2$  bound:*

$$(3.13) \quad \|u_\lambda \bar{v}_\mu^{x_0}\|_{L_{t,x}^2} \lesssim c_\lambda d_\mu \lambda^{-s} \frac{\min\{\lambda, \mu\}^{\frac{1}{2}}}{(\lambda + \mu)^{\frac{1}{2}}}, \quad \mu \not\approx \lambda.$$

This theorem in particular yields  $L^2$  well-posedness for the linearized equation.

**3.3. The bootstrap hypotheses.** To prove the above theorems, we make a bootstrap assumption where we assume the same bounds but with a worse constant  $C$ , as follows:

(i) *Uniform frequency envelope bound:*

$$(3.14) \quad \|u_\lambda\|_{L_t^\infty L_x^2} \leq C \epsilon c_\lambda \lambda^{-s}$$

(ii) *Unbalanced bilinear  $L^2$  bounds:*

$$(3.15) \quad \|u_\lambda \bar{u}_\mu^{x_0}\|_{L_{t,x}^2} \leq C^2 \epsilon^2 c_\lambda c_\mu \lambda^{-s-\frac{1}{2}} \mu^{-s+\frac{n-1}{2}}, \quad \mu \ll \lambda$$

(iii) *Balanced bilinear  $L^2$  bound:*

$$(3.16) \quad \| |D_x|^{\frac{3-n}{2}}(u_\lambda \bar{u}_\mu^{x_0}) \|_{L_{t,x}^2} \leq C^2 \epsilon^2 c_\lambda c_\mu \lambda^{-2s+\frac{n-1}{2}} \mu^{-s} \lambda^{\frac{1}{2}} (1 + \lambda |x_0|), \quad \lambda \approx \mu.$$

We note that in dimension  $n \geq 3$  the estimate (3.16) implies an  $L^4$  bound for  $u_\lambda$ , but that is no longer the case in dimension  $n = 2$ . For later use, we will add to the list above an  $L^4$  bootstrap bound for  $u_\lambda$ , which corresponds to the estimate (3.4):

(4)  $L^4$  Strichartz bound for  $n = 2$ :

$$(3.17) \quad \|u_\lambda\|_{L_{t,x}^4} \lesssim \epsilon c_\lambda \lambda^{-s+\frac{1}{2}}, \quad n = 2.$$

Then we seek to improve the constant in these bounds. The gain will come from the fact that the  $C$ 's will always come paired with extra  $\epsilon$ 's. Precisely, a continuity argument shows that

**Proposition 3.2.** *It suffices to prove Theorems 9, 10 under the bootstrap assumptions (3.14), (3.16) and (3.15).*

We remark that the Strichartz bounds (3.4) are not part of this bootstrap loop. Similar bootstrap assumptions are made for  $v$  in the proof of Theorem 11 in Section 4.

#### 4. A PARADIFFERENTIAL/RESONANT EXPANSION OF THE EQUATION

In this section we consider the equation for the frequency localized portions of the solution. We expand this equation in a dual fashion, separating two principal components:

- (1) The paradifferential part, which accounts for the quasilinear character of the problem. This is essential for the proofs of all the results in this paper.
- (2) The doubly resonant part, which accounts for the nonperturbative semilinear part of the nonlinearity. This is only needed for the global results in Theorem 8 in two dimensions.

In a first approximation, one may view the paradifferential part as simply obtained by truncating the coefficients in the principal part to lower frequencies,

$$(4.1) \quad i\partial_t v + \partial_j T_{g^{jk}(u)} \partial_k v = f.$$

While this might suffice for the local well-posedness results stated in Theorems 3 and 4, it is not precise enough for all the long time results, as the low frequencies of  $g(u)$  may also include some doubly resonant *high*  $\times$  *high* quadratic contributions. This leads us to make a better choice in the  $f$  term.

Algebraically it will be easier to deal with the corresponding frequency  $\lambda$  evolution, which may be taken as

$$(4.2) \quad i\partial_t v_\lambda + \partial_x g_{[\leq \lambda]} \partial_x v_\lambda = f_\lambda, \quad v_\lambda(0) = v_{0,\lambda},$$

where the truncated metric is defined as

$$(4.3) \quad g_{[\leq \lambda]} := P_{\leq \lambda} g(u_{\leq \lambda}).$$

It is important to note that we carefully localize  $u$  to lower frequencies, rather than  $g(u)$  directly; this is essential later on in order to exclude the doubly resonant interactions from the paradifferential flow. On the other hand, using the second order elliptic operator in divergence form is more a convenience, as the corresponding commutator terms play a perturbative role.

One may express both the full equation and the linearized equation in terms of the paradifferential flow, in the form

$$(4.4) \quad i\partial_t u_\lambda + \partial_x g_{[\leq \lambda]} \partial_x u_\lambda = N_\lambda(u), \quad u_\lambda(0) = u_{0,\lambda},$$

respectively

$$(4.5) \quad i\partial_t v_\lambda + \partial_x g_{[\leq \lambda]} \partial_x v_\lambda = N_\lambda^{lin} v, \quad v_\lambda(0) = v_{0,\lambda},$$

The source terms  $N_\lambda(u)$  and  $N_\lambda^{lin}$  will be written explicitly later when needed. They will be expected to play a perturbative role for the short time results.

For the 2D global results there is one additional portion of the nonlinearity which plays a nonperturbative role, namely the balanced cubic part, which we separate from  $N_\lambda$ , writing instead the frequency localized evolution (4.4) in the form

$$(4.6) \quad i\partial_t u_\lambda + \partial_x g_{[\leq \lambda]} \partial_x u_\lambda = C_\lambda(\mathbf{u}, \mathbf{u}, \mathbf{u}) + F_\lambda(u),$$

where we recall that the notation  $\mathbf{u}$  stands for  $\mathbf{u} = \{u, \bar{u}\}$ . Following the discussion in Section 2.3, if  $\lambda \gg 1$  then the cubic part is further decomposed into

$$C_\lambda(\mathbf{u}, \mathbf{u}, \mathbf{u}) = C_\lambda^{res}(u, \bar{u}, u) + C_\lambda^{nr}(\mathbf{u}, \mathbf{u}, \mathbf{u}) + C_\lambda^{tr}(\mathbf{u}, \mathbf{u}, \mathbf{u}).$$

Here the last term contains only transversal interactions and may be harmlessly included in  $F_\lambda$ . Correspondingly, we arrive at the final expansion

$$(4.7) \quad i\partial_t u_\lambda + \partial_x g_{[\leq \lambda]} \partial_x u_\lambda = C_\lambda^{res}(u, \bar{u}, u) + C_\lambda^{nr}(\mathbf{u}, \mathbf{u}, \mathbf{u}) + F_\lambda(u),$$

where the first two components on the right represent terms as follows:

- $C_\lambda^{res}(u, \bar{u}, u)$  contains balanced cubic terms with phase rotation symmetry, including the doubly resonant interactions
- $C_\lambda^{nr}(\mathbf{u}, \mathbf{u}, \mathbf{u})$  contains balanced nonresonant terms.

To explicitly make a choice for  $C_\lambda^{res}$  we introduce a symmetric symbol  $c_{diag}$ , smooth on the corresponding dyadic scale so that

$$c_{diag}(\xi_1, \xi_2, \xi_3) := \begin{cases} 1 & \sum |\xi_i - \xi_j| \ll \langle \xi_1 \rangle + \langle \xi_2 \rangle + \langle \xi_3 \rangle \\ 0 & \sum |\xi_i - \xi_j| \gtrsim \langle \xi_1 \rangle + \langle \xi_2 \rangle + \langle \xi_3 \rangle. \end{cases}$$

Then cubic doubly resonant part of the nonlinearity is defined in terms of the symbol  $c(\cdot, \cdot, \cdot)$  of the form in (1.2) as

$$(4.8) \quad c_\lambda(\xi_1, \xi_2, \xi_3) = p_\lambda(\xi_1 - \xi_2 + \xi_3) c_{diag}(\xi_1, \xi_2, \xi_3) c(\xi_1, \xi_2, \xi_3).$$

In particular, one sees that the conservative condition implies that this symbol is real on the diagonal,

$$\Im c_\lambda^{res}(\xi, \xi, \xi) = 0, \quad \xi \in \mathbb{R}.$$

Finally, we also briefly discuss  $F_\lambda(u)$ , containing the remaining source terms in the paradiifferential equation for  $u_\lambda$ , which can be classified as follows:

- (i) cubic *high*  $\times$  *high* terms of the form

$$u_{\leq \mu} \bar{u}_\mu \partial_x^2 u_\mu, \quad \mu \gtrsim \lambda,$$

- (ii) cubic *low*  $\times$  *high* commutator terms of the form

$$\partial_x(u_{< \lambda} \bar{u}_{< \lambda}) \partial_x u_\lambda,$$

- (iii) cubic terms with only transversal interactions of the form

$$\lambda^2 L(u_\lambda, u_\lambda, u_\lambda), \quad \lambda^2 L(u_\lambda, \bar{u}_\lambda, \bar{u}_\lambda), \quad \lambda^2 L(\bar{u}_\lambda, \bar{u}_\lambda, \bar{u}_\lambda),$$

- (iv) all cubic terms at frequency  $\lambda = 1$ ,

- (v) quartic and higher terms of the form

$$\mu^2 u_{\leq \mu}^2 \bar{u}_\mu u_\mu, \quad \mu \gtrsim \lambda.$$

In our analysis the  $F_\lambda$  terms will play a perturbative role, though establishing that is not at all immediate, and is instead part of the challenge.

## 5. INTERACTION MORAWETZ ESTIMATES: THE CONSTANT COEFFICIENT CASE

Our main tool in the proof of the bilinear  $L^2$  bounds is provided by the interaction Morawetz estimates, which in turn are based on density-flux identities for the mass and momentum. In this section we first review the constant coefficient setting, which is broadly based on [55]. However, our choices for the fluxes are different, customized to fit the nonlinear setting, akin to the work in the one dimensional case in [33], [36].

**5.1. Conservation laws in the flat case.** For clarity, we begin our discussion with the linear Schrödinger equation

$$(5.1) \quad iu_t + \Delta u = 0, \quad u(0) = u_0.$$

For this we consider the following conserved quantities, the mass

$$\mathbf{M}(u) = \int |u|^2 dx,$$

and the momentum

$$\mathbf{P}_j(u) = 2 \int \Im(\bar{u} \partial_j u) dx.$$

To these quantities we associate corresponding densities

$$M(u) = |u|^2, \quad P_j(u) = i(\bar{u} \partial_j u - u \partial_j \bar{u}).$$

The densities here are not uniquely determined, and our choices are motivated by the conservation law computations

$$(5.2) \quad \partial_t M(u) = \partial_j P_j(u),$$

$$(5.3) \quad \partial_t P_j(u) = \partial_m E_{jm}(u),$$

where we choose

$$E_{jm}(u) := \partial_m u \partial_j \bar{u} + \partial_j u \partial_m \bar{u} - v \partial_j \partial_m \bar{v} - \bar{v} \partial_j \partial_m v.$$

The reader may compare this with [55], where a different choice is made.

The symbols of these densities viewed as bilinear forms are

$$m(\xi, \eta) = 1, \quad p_j(\xi, \eta) = \xi_j + \eta_j,$$

respectively

$$e_{jm}(\xi, \eta) = (\xi_j + \eta_j)(\xi_m + \eta_m).$$

**5.2. Interaction Morawetz identities in the flat case.** We define the interaction Morawetz functional as

$$(5.4) \quad \mathbf{I}(u, v) := \iint a_j(x - y) (M(u)(x) P_j(v)(y) - P_j(u)(x) M(v)(y)) dx dy,$$

with the bounded weights  $a_j$  to be chosen later. Assuming that  $u, v$  solve the equation (5.1), the time derivative of  $\mathbf{I}(u, v)$  is

$$(5.5) \quad \frac{d}{dt} \mathbf{I}(u, v) = \mathbf{J}^4(u, v),$$

where

$$\mathbf{J}^4(u, v) = \int \partial_m a_j(x - y) (M(u) E_{jm}(v) - P_j(u) P_m(v) + M(v) E_{jm}(u) - P_j(v) P_m(u)) dx dy.$$



Now we consider the choice for  $a_j$ . To get a good sign and symmetry for  $\mathbf{J}^4$ , it is convenient to take

$$a_j(x) = \partial_j a, \quad \partial_m a_j = a_{mj},$$

where  $a$  is a well chosen convex function.

For the quadrilinear form  $J_{mj}^4$  which is the coefficient of  $a_{jm}$  we write the symbol

$$\begin{aligned} J_{mj}^4 &= (\xi_m^1 + \xi_m^2)(\xi_j^1 + \xi_j^2) + (\xi_m^3 + \xi_m^4)(\xi_j^3 + \xi_j^4) - (\xi_j^1 + \xi_j^2)(\xi_m^3 + \xi_m^4) - (\xi_m^1 + \xi_m^2)(\xi_j^3 + \xi_j^4) \\ &= (\xi_j^1 + \xi_j^2 - \xi_j^3 - \xi_j^4)(\xi_m^1 + \xi_m^2 - \xi_m^3 - \xi_m^4) \\ &= 2(\xi_j^1 - \xi_j^4)(\xi_m^3 - \xi_m^2) + 2(\xi_m^1 - \xi_m^4)(\xi_j^3 - \xi_j^2) \quad (\text{mod } \Delta^4 \xi), \end{aligned}$$

where terms with  $\Delta^4 \xi$  factors do not contribute to the integral  $\mathbf{J}_{mj}^4$ .

So for instance if  $a(x) = x^2$  then we get

$$J^4(u, v) = 8(\xi^1 - \xi^4) \cdot (\xi^3 - \xi^2) \quad (\text{mod } \Delta^4 \xi),$$

which yields the positive form

$$\mathbf{J}^4(u, v) = 8 \|\partial(u\bar{v})\|_{L^2}^2.$$

Unfortunately we cannot use this because the  $a_j$ 's are not bounded.

More generally, the above formula for the symbol of  $J_{mj}^4$  directly yields

$$(5.6) \quad \mathbf{J}^4(u, v) = 2 \int a_{jm}(x - y) F_j \bar{F}_m dx dy,$$

where

$$(5.7) \quad F_j(x, y) = u(x) \partial_j \bar{v}(y) + \partial_j u(x) \bar{v}(y).$$

This is clearly positive definite if  $a$  is convex.

The special case when  $u = v$  is also interesting. Using the identity

$$|u(x) \partial_j \bar{v}(y) + \partial_j u(x) \bar{v}(y)|^2 = |u(x) \partial_j v(y) - \partial_j u(x) v(y)|^2 + \partial_j |u(x)|^2 \partial_j |v(y)|^2,$$

and its bilinear version, setting  $u = v$  we obtain the inequality

$$(5.8) \quad \mathbf{J}^4(u, u) \geq \int a_{jm}(x - y) \partial_j |u(x)|^2 \partial_m |u(y)|^2 dx dy.$$

For the function  $a$  we will use  $a(x) = |x|$ , in which case we have the Hessian

$$D^2 a(x) = \frac{1}{|x|} (I - \frac{x}{|x|} \otimes \frac{x}{|x|}).$$

To complete our computations we use the identity

$$(5.9) \quad \int a_{jm}(x - y) \partial_j |u(x)|^2 \partial_m |u(y)|^2 dx dy = c_n \| |D|^{\frac{3-n}{2}} |u|^2 \|_{L^2}^2,$$

which is easily verified by interpreting  $a_{jm}$  in (5.8) as kernels for appropriate multipliers. This finally leads to

$$(5.10) \quad \mathbf{J}^4(u, u) \gtrsim \| |D|^{\frac{3-n}{2}} |u|^2 \|_{L^2}^2.$$

We remark, however, that the two expressions above are not equivalent, which is harmless here but will generate considerable difficulties in the nonlinear case.

We are also interested in frequency localized versions of these objects. Given a dyadic frequency  $\lambda$ , we start with a symbol  $a_\lambda(\xi, \eta)$  which is symmetric, in the sense that

$$a_\lambda(\eta, \xi) = \overline{a_\lambda(\xi, \eta)},$$

and localized at frequency  $\lambda$ , and then define an associated weighted mass density by

$$M_\lambda(u) = A_\lambda(u, \bar{u}).$$

We also define corresponding momentum symbols  $p_{j,\lambda}$  by

$$p_\lambda(\xi, \eta) = (\xi_j + \eta_j) a(\xi, \eta).$$

Then a direct computation yields the density flux relations

$$\frac{d}{dt} M_\lambda(u, \bar{u}) = \partial_j P_{j,\lambda}(u, \bar{u}), \quad \frac{d}{dt} P_{j,\lambda}(u, \bar{u}) = \partial_m E_{mj,\lambda}(u, \bar{u}).$$

## 6. DENSITY-FLUX RELATIONS FOR MASS AND MOMENTUM

In this section we develop the density-flux identities for the paradifferential and the non-linear flows, leaving the full interaction Morawetz analysis for the next section.

**6.1. Density-flux identities for the paradifferential problem.** Now that we have the flat case as a model, we turn our attention to the paradifferential flow (4.2) for which we consider solutions  $v_\lambda$ . For these we have

$$\begin{aligned} \frac{d}{dt} M(v_\lambda) &= 2\Re(\partial_t v_\lambda \cdot \bar{v}_\lambda) \\ &= 2\Im(\partial_j g_{[\leq \lambda]}^{jl} \partial_l v_\lambda \cdot \bar{v}_\lambda) + 2\Im(f_\lambda \bar{v}_\lambda) \\ &= 2\partial_j [g_{[\leq \lambda]}^{jl} \Im(\partial_l v_\lambda \cdot \bar{v}_\lambda)] + 2\Im(f_\lambda \bar{v}_\lambda). \end{aligned}$$

Defining the covariant momenta as

$$P^j(v_\lambda) = g_{[\leq \lambda]}^{jl} P_l(v_\lambda),$$

we rewrite this in the form

$$(6.1) \quad \frac{d}{dt} M(v_\lambda) = \partial_j [P^j(v_\lambda)] + F_{\lambda,m}^{para}, \quad F_{\lambda,m}^{para} = 2\Im(f_\lambda \bar{v}_\lambda).$$

Similarly, one computes, with  $g := g_{[\leq \lambda]}$  and  $v := v_\lambda$ ,

$$\begin{aligned} \frac{d}{dt} P_j(v_\lambda) &= 2\Re(\partial_k g^{kl} \partial_l v \partial_j \bar{v}) - 2\Re(v \partial_j \partial_k g^{kl} \partial_l \bar{v}) - 2\Re(f \partial_j \bar{v}) + 2\Re(v \partial_j \bar{f}) \\ &= 2\partial_k \Re(g^{kl} \partial_l v \partial_j \bar{v} - v \partial_j g^{kl} \partial_l \bar{v}) - 2\Re(g^{kl} \partial_l v \partial_j \partial_k \bar{v}) + 2\Re(\partial_k v \partial_j g^{kl} \partial_l \bar{v}) \\ &\quad - 2\Re(f \partial_j \bar{v}) + 2\Re(v \partial_j \bar{f}) \\ &= 2\partial_k \Re(g^{kl} \partial_l v \partial_j \bar{v} - v \partial_j g^{kl} \partial_l \bar{v}) + 2\Re(\partial_k v (\partial_j g^{kl}) \partial_l \bar{v}) - 2\Re(f \partial_j \bar{v}) + 2\Re(v \partial_j \bar{f}) \\ &= \partial_k [E_j^k(v_\lambda)] + F_{j,\lambda,p}^{para}, \end{aligned}$$

where

$$(6.2) \quad E_j^k(v_\lambda) := 2\Re(g^{kl} \partial_l v \partial_j \bar{v} - v \partial_j g^{kl} \partial_l \bar{v}),$$

and

$$(6.3) \quad F_{j,\lambda,p}^{para} := 2\Re(\partial_k v (\partial_j g^{kl}) \partial_l \bar{v}) - 2\Re(f \partial_j \bar{v}) + 2\Re(v \partial_j \bar{f}).$$

Now we switch to the covariant versions,

$$P^j(v_\lambda) = g_{[\prec\lambda]}^{jk} P_k(v_\lambda),$$

for which we similarly write

$$(6.4) \quad \frac{d}{dt} P^j(v_\lambda) = \partial_k [E^{kj}(v_\lambda)] + F_{\lambda,p}^{para,j},$$

where for  $E^{kj}$  we have simply raised indices tensorially,

$$(6.5) \quad E^{kj}(v_\lambda) = 2\Re(\partial^k v_\lambda \partial^j \bar{v}_\lambda - v_\lambda \partial^j \partial^k \bar{v}_\lambda),$$

but for the source term we get additional commutator terms,

$$(6.6) \quad F_{\lambda,p}^{j,para} = G_{\lambda,p}^{j,para} - 2\Re(f_\lambda \partial^j \bar{v}_\lambda) + 2\Re(v_\lambda \partial^j \bar{f}_\lambda),$$

where the quadratic term  $G_{\lambda,p}^{j,para}$  is given by

$$(6.7) \quad G_{\lambda,p}^{j,para} = \partial_k v_\lambda (\partial^j g_{[\prec\lambda]}^{kl}) \partial_l \bar{v}_\lambda + (\partial_t g_{[\prec\lambda]}^{jk}) P_k(v_\lambda) - (\partial_k g_{[\prec\lambda]}^{kl}) E_l^j(v_\lambda).$$

Fortunately its exact form is less important, as it will be perturbatively estimated in  $L^1$  later on. Here we simply note that we can schematically write it in the form

$$(6.8) \quad G_{\lambda,p}^{j,para} = h(\mathbf{u}_{<\lambda}) \mathbf{u}_{<\lambda} \partial \mathbf{u}_{<\lambda} \partial v_\lambda \partial \bar{v}_\lambda,$$

which is all that will be needed.

**6.2. Nonlinear density flux identities for the frequency localized mass and momentum.** The interaction Morawetz identities for the paradifferential equation suffice for the proof of our results in dimension  $n \geq 3$ . However, a finer analysis is needed in the most interesting case  $n = 2$ , and particularly for the proof of the global result in Theorem 8. This is the aim of this subsection. We note that the analysis below is only needed at large frequencies, where the  $L^4$  Strichartz bound loses too much to allow us to treat cubic terms perturbatively.

We begin with a simpler computation for the mass density of  $u_\lambda$ , using the equation (4.7). We have

$$\begin{aligned} \frac{d}{dt} M_\lambda(u) &= 2\Re(\partial_t u \cdot \bar{u}) \\ &= 2\Im(\partial_j g_{[\prec\lambda]}^{jk} \partial_k u_\lambda \cdot \bar{u}_\lambda) + 2\Im((C_\lambda^{res}(u, \bar{u}, u) + C_\lambda^{nr}(\mathbf{u}, \mathbf{u}, \mathbf{u}) + F_\lambda(\mathbf{u})) \bar{u}_\lambda) \\ &= 2\partial_j [g_{[\prec\lambda]}^{jk} \Im(\partial_k u_\lambda \cdot \bar{u}_\lambda)] + 2\Im((C_\lambda^{res}(u, \bar{u}, u) + C_\lambda^{nr}(\mathbf{u}, \mathbf{u}, \mathbf{u}) + F_\lambda(\mathbf{u})) \bar{u}_\lambda), \end{aligned}$$

which we rewrite in the form

$$(6.9) \quad \frac{d}{dt} M_\lambda(u) = 2\partial_j [P_\lambda^j(u)] + C_{\lambda,m}^{4,res}(u, \bar{u}, u, \bar{u}) + C_{\lambda,m}^{4,nr}(\mathbf{u}, \mathbf{u}, \mathbf{u}, \mathbf{u}) + 2F_{\lambda,m}^4(\mathbf{u}),$$

where

$$C_{\lambda,m}^{4,res}(u, \bar{u}, u, \bar{u}) := 2\Im((C_\lambda^{res}(u, \bar{u}, u) \bar{u}_\lambda), \quad C_{\lambda,m}^{4,nr}(\mathbf{u}, \mathbf{u}, \mathbf{u}, \mathbf{u}) := 2\Im(C_\lambda^{nr}(\mathbf{u}, \mathbf{u}, \mathbf{u}) \bar{u}_\lambda),$$

$$F_{\lambda,m}^4(\mathbf{u}) := 2\Im((F_\lambda(\mathbf{u}) \bar{u}_\lambda).$$

The three source terms in (6.9) will be treated differently in the last section.

The term  $C_{\lambda,m}^{4,res}$  contains only interactions which are localized at frequency  $\lambda$ , but include the parallel interactions where all frequencies are equal. Under the conservative assumption in Definition 1.5, we obtain the key symbol property

$$(6.10) \quad c_{\lambda,m}^4(\xi, \xi, \xi, \xi) = 0.$$

One should compare this with the 1D case in [33], [36], where a stronger conservative condition guaranteed a second order vanishing for  $c_{\lambda,m}^4$  on the diagonal. This in turn allowed us to remove this term via a quartic correction to the mass density, and a quartic correction to the mass flux, modulo perturbative errors. In the 2D case there are too many resonant interactions in order to allow such a strategy. Instead, we will interpret the above vanishing condition as allowing us to get a representation

$$C_{\lambda,m}^{4,res}(u, \bar{u}, u, \bar{u}) \approx \lambda L(\partial|u|^2, u, \bar{u}) + \lambda \partial R_{\lambda,m}^4(u, \bar{u}, \bar{u}, \bar{u}),$$

which in turn will be used in order to treat it as a perturbative term, by separating it into an  $L_{t,x}^1$  component and a flux correction.

The term  $C_{\lambda,m}^{4,res}$  contains a larger range of interactions which are also localized at frequency  $\lambda$ , and are nonresonant. This will allow us to remove it using both density and flux corrections

$$C_{\lambda,m}^{4,nr}(\mathbf{u}, \mathbf{u}, \mathbf{u}, \mathbf{u}) \approx \partial_t L(\mathbf{u}, \mathbf{u}, \mathbf{u}, \mathbf{u}) + \lambda \partial_x L(\mathbf{u}, \mathbf{u}, \mathbf{u}, \mathbf{u}) + F^{4,nr}(\mathbf{u})$$

with a perturbative source term  $F^{4,nr}(\mathbf{u})$ .

Finally,  $F_{\lambda,m}^4$  is at least quartic and only involves double transversal interactions, and will be estimated in  $L^1$  and treated perturbatively.

Similarly, we have a corresponding density-flux relation for the momentum

$$(6.11) \quad \frac{d}{dt} P_{\lambda}^j(u) = \partial_k E_{\lambda}^{jk}(u) + C_{\lambda,p}^{j,4,res}(u, \bar{u}, u, \bar{u}) + C_{\lambda,p}^{4,nr}(\mathbf{u}, \mathbf{u}, \mathbf{u}, \mathbf{u}) + F_{\lambda,p}^{j,4}(\mathbf{u})$$

with a cancellation relation akin to (6.10) and a similar set of density and flux corrections for the middle terms on the right.

## 7. INTERACTION MORAWETZ BOUNDS FOR THE LINEAR PARADIFFERENTIAL FLOW

In this section as well as in the next one we study the linear paradifferential equation associated to a solution  $u$  to the quasilinear Schrödinger flow (QNLS). Here we prove bilinear  $L^2$  bounds and  $L^2$  well-posedness. On one hand, this is a key step in the proof of both the local well-posedness results in Theorem 3, Theorem 4 and the global results in Theorem 7, Theorem 6. On the other hand, an enhanced version of these arguments will be used later in order to prove estimates for the full equation, and in particular to prove our final global result in Theorem 8. In the next section we prove Strichartz estimates for the paradifferential equation.

To start with, we consider a one parameter family of functions  $v_{\lambda}$  which solve the equations (4.2). For now we assume no connection between these functions. Later we will apply these bounds in the case when  $v_{\lambda} = P_{\lambda}u$  or  $v_{\lambda} = P_{\lambda}v$  where  $v$  solves the linearized equation.

For these functions we have the conservation laws

$$(7.1) \quad \partial_t M(v_{\lambda}) = \partial_k P^k(v_{\lambda}) + F_{\lambda,m}^{para},$$

respectively

$$(7.2) \quad \partial_t P^j(v_{\lambda}) = \partial_k E^{kj}(v_{\lambda}) + F_{p,\lambda}^{j,para},$$

where the source terms are

$$(7.3) \quad F_{\lambda,m}^4 := 2\Im M(f_\lambda, \bar{v}_\lambda) = 2\Im(f_\lambda \bar{v}_\lambda),$$

respectively

$$(7.4) \quad F_{p,\lambda}^{j,para} := G_{p,\lambda}^{j,para} - 2\Re(f_\lambda \partial^j \bar{v}_\lambda) + 2\Re(v_\lambda \partial^j \bar{f}_\lambda)$$

with  $G_{p,\lambda}^{j,para}$  given by (6.7).

We will measure each  $v_\lambda$  based on the size of the initial data and of the source term  $f_\lambda$ . However, following an idea introduced in [36], rather than measure  $f_\lambda$  directly, we will instead measure its interaction with  $v_\lambda$  and its translates. So we denote

$$(7.5) \quad d_\lambda^2 := \sup_{\mu, \nu \approx \lambda} \|v_\mu(0)\|_{L^2}^2 + \sup_{x_0 \in \mathbb{R}} \|v_\mu f_\nu^{x_0}\|_{L_{t,x}^1}.$$

Using the more implicit parameter  $d_\lambda$  allows for a larger range of source terms, which is absolutely essential in the present work.

Our main result here asserts that we can obtain energy and bilinear  $L^2$  bounds for  $v_\lambda$ :

**Theorem 12.** *Assume that  $u$  solves (QNLS) in a time interval  $[0, T]$ , and satisfies the bounds (3.1), (3.2) and (3.3) in the same time interval for some  $s > \frac{n+1}{2}$ . If  $n = 2$ , we also assume that  $T \lesssim \epsilon^{-6}$ . Assume  $v_\lambda$  solve (4.2), and let  $d_\lambda$  be as in (7.5). Then the following bounds hold for the functions  $v_\lambda$  uniformly in  $x_0 \in \mathbb{R}$ :*

(1) *Uniform energy bounds:*

$$(7.6) \quad \|v_\lambda\|_{L_t^\infty L_x^2} \lesssim d_\lambda$$

(2) *Balanced bilinear  $(u, v)$ - $L^2$  bound:*

$$(7.7) \quad \| |D|^{\frac{3-n}{2}}(v_\lambda \bar{u}_\mu^{x_0}) \|_{L_{t,x}^2} \lesssim \epsilon d_\lambda c_\mu \lambda^{-s+\frac{1}{2}}(1 + \lambda|x_0|), \quad \mu \approx \lambda$$

(3) *Unbalanced bilinear  $(u, v)$ - $L^2$  bounds:*

$$(7.8) \quad \|v_\lambda \bar{u}_\mu^{x_0}\|_{L_{t,x}^2} \lesssim \epsilon d_\lambda c_\mu \mu^{-s} \frac{\min\{\lambda, \mu\}^{\frac{n-1}{2}}}{(\lambda + \mu)^{\frac{1}{2}}} \quad \mu \not\approx \lambda$$

(4) *Balanced bilinear  $(v, v)$ - $L^2$  bound:*

$$(7.9) \quad \| |D|^{\frac{3-n}{2}}(v_\lambda \bar{v}_\mu^{x_0}) \|_{L_{t,x}^2} \lesssim d_\lambda d_\mu \lambda^{\frac{1}{2}}(1 + \lambda|x_0|), \quad \mu \approx \lambda$$

(5) *Unbalanced bilinear  $(v, v)$ - $L^2$  bound:*

$$(7.10) \quad \|v_\lambda \bar{v}_\mu^{x_0}\|_{L_{t,x}^2} \lesssim d_\lambda d_\mu \frac{\min\{\lambda, \mu\}^{\frac{n-1}{2}}}{(\lambda + \mu)^{\frac{1}{2}}} \quad \mu \not\approx \lambda.$$

Before proceeding to the proof, we remark that the unbalanced bounds are more robust, and do not require any relation between the metric in the  $v_\lambda$  equation and the one in the  $v_\mu$  equation:

**Corollary 7.1.** *The unbalanced bounds (7.8) and (7.10) are still valid if the equation (4.2) for  $v_\lambda$  is replaced by the linear Schrödinger equation*

$$(i\partial_t + \Delta)v_\lambda = f_\lambda.$$

This should be seen as a corollary of the proof of the theorem, and not of the theorem itself. Indeed, the unbalanced case of the proof of the theorem applies in this case without any change. We now return to the proof of the theorem.

*Proof of Theorem 12.* In order to treat the dyadic bilinear  $(u, v)$  and  $(v, v)$  bounds at the same time, we write the equation for  $u_\lambda$  in the form (4.4) where we estimate favourably the source term  $N_\lambda$ , and prove the following estimate:

**Proposition 7.2.** *a) Let  $n \geq 3$  and  $s > \frac{n+1}{2}$ . Assume that the function  $u$  satisfies the bounds (3.1), (3.2) and (3.3) in a time interval  $[0, T]$ . Then for  $\epsilon$  small enough, the functions  $N_\lambda(u)$  in (4.4) satisfy*

$$(7.11) \quad \|N_\lambda(\mathbf{u})\bar{u}_\lambda^{x_0}\|_{L_{t,x}^1} \lesssim \epsilon^4 c_\lambda^2 \lambda^{-2s}.$$

*b) Let  $n = 2$  and  $s > \frac{n+1}{2}$ . Then (7.11) holds under the additional assumption  $T \leq \epsilon^{-6}$ . Furthermore, for any  $T$  we have the partial bound*

$$(7.12) \quad \|F_\lambda(\mathbf{u})\bar{u}_\lambda^{x_0}\|_{L_{t,x}^1} \lesssim \epsilon^4 c_\lambda^2 \lambda^{-2s}.$$

*Proof.* We start the proof without discriminating between the spatial dimensions  $n = 2$  and  $n \geq 3$ ; we will do so when the estimates require it. The bound in (7.11) relies on a careful analysis of the frequency localized source term  $N_\lambda(\mathbf{u})$ . The key features of the multilinear terms contained in  $N_\lambda(u)$  are :

- they are at least cubic
- they have exactly two derivatives
- the highest frequency is at least  $\lambda$ .
- the two highest frequencies have to be comparable when either
  - the two derivatives apply to the highest frequency or
  - the highest frequency is  $\gg \lambda$ .

This leads to a large number of cases which are also informed by the comments in Section 4 which discuss a potential partition of  $N_\lambda$ , and which we next organize efficiently. For this, recall that

$$N_\lambda(\mathbf{u}) = C_\lambda(\mathbf{u}, \mathbf{u}, \mathbf{u}) + F_\lambda(\mathbf{u}),$$

where the entries of  $C_\lambda$  are localized around frequency  $\lambda$  (but not strictly localized), and where  $F_\lambda$  contain not only the remaining cubic source terms but also higher order terms. The cubic terms in  $F_\lambda$  are either unbalanced, or balanced transversal, and do not necessarily have the phase rotation symmetry property. We begin discussing all cases, one at a time:

**A.** Highest frequency is  $\mu \gg \lambda$ . In this case we must have at least two frequency  $\mu$  factors. We separate this case further.

A1. Terms which have at least one lower frequency  $\mu_1 \ll \mu$ . For brevity we consider a typical term of the form

$$N_\lambda^{A1} = \mathbf{u}_{\mu_1} \partial_x^2 \mathbf{u}_\mu^2,$$

which can be expressed as a multilinear expression in the form

$$N_\lambda^{A1} = \mu^2 L(\mathbf{u}_{\mu_1}, \mathbf{u}_\mu, \mathbf{u}_\mu).$$

Here the estimate goes as follows

$$\|N_\lambda^{A1} \bar{u}_\lambda^{x_0}\|_{L^1} \lesssim \mu^2 \|L(\mathbf{u}_\mu, \mathbf{u}_\lambda^{x_0})\|_{L^2} \|L(\mathbf{u}_{\mu_1}, \mathbf{u}_\mu)\|_{L^2} \lesssim \epsilon^4 c_\lambda c_{\mu_1} c_\mu^2 \mu^{1-2s} \lambda^{-s+\frac{n-1}{2}} \mu_1^{-s+\frac{n-1}{2}}.$$

Here we have arranged the factors to obtain two unbalanced pairs to which we then applied the unbalanced bilinear  $L^2$  bounds in (3.2). The summation with respect to  $\mu_1$  and  $\mu$  is straightforward. One last observation is regarding the complex conjugates which do not matter in obtaining this bound.

A2. Terms with exactly three  $\mu$  frequencies. Here the three  $\mu$  frequencies must add up to a frequency  $\ll \mu$ , so at least two must be  $\mu$  separated, and the above argument still applies.

A3. Terms with at least four  $\mu$  frequencies, e.g.

$$N_\lambda^{A3} = \mathbf{u}_\mu^3 \partial_\mu^2 \mathbf{u}_\mu,$$

which we can re-express as

$$N_\lambda^{A3} = \mu^2 L(\mathbf{u}_\mu, \mathbf{u}_\mu, \mathbf{u}_\mu, \mathbf{u}_\mu).$$

To get the desired estimate we group the two unbalanced terms and use (3.2),

$$\begin{aligned} \|N_\lambda^{A3} \bar{u}_\lambda^{x_0}\|_{L_{t,x}^1} &\lesssim \mu^2 \|L(\mathbf{u}_\mu, u_\lambda^{x_0})\|_{L_{t,x}^2} \|L(\mathbf{u}_\mu, \mathbf{u}_\mu, \mathbf{u}_\mu)\|_{L_{t,x}^2} \\ &\lesssim \epsilon^2 c_\lambda c_\mu \mu^{-s+\frac{3}{2}} \lambda^{-s+\frac{n-1}{2}} \|L(\mathbf{u}_\mu, \mathbf{u}_\mu, \mathbf{u}_\mu)\|_{L_{t,x}^2}. \end{aligned}$$

Then for the remaining factor we need to discuss separate ways of obtaining the bounds when  $n = 2$ , respectively  $n \geq 3$ . We begin with the seemingly easier case  $n \geq 3$ , and bound the remaining cubic terms using the  $L_{t,x}^4$  Strichartz bound (3.5) for two copies of  $u_\mu$  and Bernstein's inequality for the last  $u_\mu$

$$\|L(\mathbf{u}_\mu, \mathbf{u}_\mu, \mathbf{u}_\mu)\|_{L_{t,x}^2} \lesssim \|u_\mu\|_{L_{t,x}^4}^2 \|u_\mu\|_{L_{t,x}^\infty} \lesssim \epsilon^2 c_\mu^2 \mu^{-s+\frac{n-2}{2}} \mu^{\frac{n}{2}} \|u_\mu\|_{L_t^\infty L_x^2} \lesssim \epsilon^3 c_\mu^3 \mu^{-2s+n-1}.$$

We summarize the bound we obtained in the  $n \geq 3$  case below

$$\|N_\lambda^{A3} \bar{u}_\lambda^{x_0}\|_{L_{t,x}^1} \lesssim \epsilon^5 c_\lambda c_\mu^4 \mu^{-3s+n+\frac{1}{2}} \lambda^{-s+\frac{n-1}{2}}.$$

It remains to bound the same expression when  $n = 2$ :

$$\|N_\lambda^{A3} \bar{u}_\lambda^{x_0}\|_{L_{t,x}^1} \lesssim \mu^2 \|L(\mathbf{u}_\mu, u_\lambda^{x_0})\|_{L_{t,x}^2} \|L(\mathbf{u}_\mu, \mathbf{u}_\mu, \mathbf{u}_\mu)\|_{L_{t,x}^2} \lesssim \epsilon^2 c_\lambda c_\mu \mu^{-s+\frac{3}{2}} \lambda^{-s+\frac{1}{2}} \|L(\mathbf{u}_\mu, \mathbf{u}_\mu, \mathbf{u}_\mu)\|_{L_{t,x}^2}.$$

For the last trilinear term we rely on the Strichartz bound (3.6) as well on the Bernstein's inequality

$$\|L(\mathbf{u}_\mu, \mathbf{u}_\mu, \mathbf{u}_\mu)\|_{L_{t,x}^2} \lesssim \|u_\mu\|_{L_t^4 L_x^8}^2 \|u_\mu\|_{L_t^\infty L_x^4} \lesssim \epsilon^3 c_\mu^3 \mu^{-3s+1}.$$

Hence, when we are in spatial dimension  $n = 2$  we get

$$\|N_\lambda^{A3} \bar{u}_\lambda^{x_0}\|_{L_{t,x}^1} \lesssim \mu^2 \|L(\mathbf{u}_\mu, u_\lambda^{x_0})\|_{L_{t,x}^2} \|L(\mathbf{u}_\mu, \mathbf{u}_\mu, \mathbf{u}_\mu)\|_{L_{t,x}^2} \lesssim \epsilon^5 c_\lambda c_\mu^4 \mu^{-4s+2} \lambda^{-s+\frac{1}{2}},$$

which still suffices.

**B.** The highest frequency is comparable to  $\lambda$ . We also split this case further.

B1. Terms in  $F_\lambda(\mathbf{u})$  which have exactly one frequency  $\lambda$  factor. Such terms arise only from commuting the metric  $g_{[\cdot < \lambda]}$  with a derivative or a frequency  $\lambda$  projector (see the similar computation in [36]), so we have at most one derivative on the frequency  $\lambda$  factor. Thus we are led to consider expressions of the form

$$N_\lambda^{B1} = \partial_x(\mathbf{u}_{\lambda_1} \mathbf{u}_{\lambda_2}) \partial_x u_\lambda + \partial_x \mathbf{u}_{\lambda_1} \partial_x \mathbf{u}_{\lambda_2} u_\lambda, \quad \lambda_2 \leq \lambda_1 \ll \lambda,$$

which will be useful to write as

$$N_\lambda^{B1} = \lambda \lambda_1 L(\mathbf{u}_\lambda, \mathbf{u}_{\lambda_1}, \mathbf{u}_{\lambda_2}) + \lambda_1 \lambda_2 L(\mathbf{u}_\lambda, \mathbf{u}_{\lambda_1}, \mathbf{u}_{\lambda_2}).$$

We again take advantage of the unbalanced bilinear Strichartz bound (3.2)

$$\begin{aligned}
\|N_\lambda^{B1} \bar{u}_\lambda^{x_0}\|_{L_{t,x}^1} &\lesssim (\lambda_1 \lambda + \lambda_1 \lambda_2) \|u_{\lambda_1} u_\lambda^{x_0}\|_{L_{t,x}^2} \|u_{\lambda_2} u_\lambda\|_{L_{t,x}^2} \\
&\lesssim (\lambda_1 \lambda + \lambda_1 \lambda_2) \epsilon^4 c_\lambda^2 c_{\lambda_1} c_{\lambda_2} \lambda^{-2s-1} \lambda_1^{-s+\frac{n-1}{2}} \lambda_2^{-s+\frac{n-1}{2}} \\
&\lesssim \epsilon^4 c_\lambda^2 c_{\lambda_1} c_{\lambda_2} \lambda^{-2s} \lambda_1^{1-s+\frac{n-1}{2}} \lambda_2^{-s+\frac{n-1}{2}}.
\end{aligned}$$

B2. We have exactly two frequency  $\lambda$  factors. Here the two  $\lambda$  frequencies must add up to  $O(\lambda)$  so they must be  $\lambda$  separated, and the above argument still applies.

B3. We have at least three factors at frequency  $\lambda$ . This is the most difficult case, which includes in particular the cubic balanced terms in 2D. We first dispense with everything else.

B3a.  $n \geq 3$ . Here we can simply use four  $L^4$  bounds:

$$\|N_\lambda^{B3} \bar{u}_\lambda^{x_0}\|_{L_{t,x}^1} \lesssim \lambda^2 \|u_\lambda\|_{L_{t,x}^4}^4 \lesssim \epsilon^4 c_\lambda^4 \lambda^{-4s+n}.$$

B3b.  $n = 2$ . Here we separate into several cases:

B3b(i). We have at least four factors at frequency  $\lambda$ . This includes quartic and higher order terms in  $F_\lambda(\mathbf{u})$ . Such terms are given, for example, by

$$N_\lambda^{B3} = \partial_x^2 u_\lambda \mathbf{u}_\lambda^3.$$

We reinterpret this term as a multilinear form

$$N_\lambda^{B3} = \lambda^2 L(\mathbf{u}_\lambda, \mathbf{u}_\lambda, \mathbf{u}_\lambda, \mathbf{u}_\lambda).$$

Since we are in the case  $n = 2$ , this leads to

$$\begin{aligned}
\|N_\lambda^{B3} \bar{u}_\lambda^{x_0}\|_{L_{t,x}^1} &\lesssim \lambda^2 \|u_\lambda\|_{L_t^4 L_x^8}^4 \|u_\lambda\|_{L_t^\infty L_x^2} \\
&\lesssim \epsilon^5 c_\lambda^5 \lambda^{-5s+3}.
\end{aligned}$$

B3b(ii). We have quartic or higher order source terms with exactly three factors at frequency  $\lambda$  and at least one smaller frequency, e.g.

$$N_\lambda^{B3} = \lambda^2 L(\mathbf{u}_\lambda, \mathbf{u}_\lambda, \mathbf{u}_\lambda, \mathbf{u}_{\lambda_1}), \quad \lambda_1 < \lambda.$$

We bound its contribution as follows

$$\begin{aligned}
\|N_\lambda^{B3} \bar{u}_\lambda^{x_0}\|_{L_{t,x}^1} &\lesssim \lambda^2 \|u_\lambda u_{\lambda_1}\|_{L_{t,x}^2} \|u_\lambda\|_{L_t^4 L_x^8}^2 \|u_\lambda\|_{L_t^\infty L_x^4} \\
&\lesssim \lambda^2 \epsilon^5 c_\lambda^4 c_{\lambda_1} \lambda^{-s-\frac{1}{2}} \lambda_1^{-s+\frac{1}{2}} \lambda^{-2s+\frac{1}{2}} \lambda^{-s+\frac{1}{2}} \\
&\lesssim \epsilon^5 c_\lambda^4 c_{\lambda_1} \lambda^{-4s+\frac{5}{2}} \lambda_1^{-s+\frac{1}{2}}.
\end{aligned}$$

B3b(iii) We have exactly a cubic balanced form at frequency  $\lambda$  which further satisfies the transversality condition. Here we simply group the four entries into two groups with  $\lambda$ -frequency separated pairs, and use two bilinear  $L^2$  bounds.

B3b(iv) We have exactly three balanced terms at frequency  $\lambda$ , possibly resonant. In this case we know in addition that  $T < \epsilon^{-6}$ . We take

$$N_\lambda^{B3} = \lambda^2 L(\mathbf{u}_\lambda, \mathbf{u}_\lambda, \mathbf{u}_\lambda)$$



and its bound is given by

$$\|N_\lambda^{B3} \bar{u}_\lambda^{x_0}\|_{L_{t,x}^1} \lesssim \lambda^2 \|u_\lambda\|_{L_{x,t}^4}^4 \lesssim \lambda^2 T^{\frac{1}{3}} \|u_\lambda\|_{L_t^6 L_x^4}^4 \lesssim T^{\frac{1}{3}} \epsilon^4 c_\lambda^4 \lambda^{-4s+\frac{8}{3}}.$$

□

Given the estimate (7.11), the  $(u, v)$  bilinear bounds and the  $(v, v)$  bilinear bounds are absolutely identical. Hence in what follows we will just consider the  $(v, v)$  bounds. To prove the theorem it is convenient to make the following bootstrap assumptions:

(1) Uniform energy bounds:

$$(7.13) \quad \|v_\lambda\|_{L_t^\infty L_x^2} \leq C d_\lambda$$

(2) Unbalanced bilinear  $(u, v)$ - $L^2$  bound:

$$(7.14) \quad \|v_\lambda \bar{u}_\mu^{x_0}\|_{L_{t,x}^2} \leq C \epsilon d_\lambda c_\mu \mu^{-s+\frac{n-1}{2}} \lambda^{-\frac{1}{2}} \quad \mu < \lambda$$

(3) Balanced bilinear  $(v, v)$ - $L^2$  bound:

$$(7.15) \quad \| |D_x|^{\frac{3-n}{2}} (v_\lambda \bar{v}_\mu^{x_0}) \|_{L_{t,x}^2} \leq C^2 d_\lambda d_\mu \lambda^{\frac{1}{2}} (1 + \lambda |x_0|), \quad \mu \approx \lambda$$

(4) Unbalanced bilinear  $(v, v)$ - $L^2$  bound:

$$(7.16) \quad \|v_\lambda \bar{v}_\mu^{x_0}\|_{L_{t,x}^2} \leq C^2 \epsilon d_\lambda d_\mu \mu^{\frac{n-1}{2}} \lambda^{-\frac{1}{2}} \quad \mu < \lambda.$$

These are assumed to hold with a large universal constant  $C$ , and then will be proved to hold without it. The way  $C$  will be handled is by always pairing it with  $\epsilon$  factors in the estimates, so that it can be eliminated simply by assuming that  $\epsilon$  is sufficiently small.

Here we note that to prove (7.6), (7.7) and (7.8) it suffices to work with a fixed  $v_\lambda$ , while for the bilinear  $v$  bounds we need to work with exactly two  $v_\lambda$ 's. Since the functions  $v_\lambda$  are completely independent, without any restriction in generality we could assume that  $d_\lambda = 1$  in what follows.

We first observe that the bound (7.6) follows directly via a energy estimate. Next we use the bootstrap assumptions to estimate the sources  $F^{para}$  in the density flux energy identities for  $v_\lambda$  in  $L^1$ ,

**Lemma 7.3.** *Assume that (7.14) holds. Then we have*

$$(7.17) \quad \|F_{\lambda,m}^{para}\|_{L_{t,x}^1} \lesssim C^2 \epsilon^2 d_\lambda^2, \quad \|F_{\lambda,p}^{para}\|_{L_{t,x}^1} \lesssim C^2 \epsilon^2 \lambda d_\lambda^2.$$

*Proof.* The only nontrivial part is the first term in  $F_{\lambda,p}^{para}$ , namely  $G_{\lambda,p}^{para}$  given by the formula (6.7), which schematically has the form

$$\partial g_{[\prec \lambda]} \partial v_\lambda \partial \bar{v}_\lambda,$$

where we can use two bilinear  $L^2$  bounds. Precisely, since  $g$  is at least quadratic in  $u$ , it suffices to show that

$$(7.18) \quad \|\partial u_{\prec \lambda} v_\lambda\|_{L_{t,x}^2} \lesssim C \epsilon d_\lambda \lambda^{-\frac{1}{2}},$$

as well as its easier version

$$(7.19) \quad \|u_{\prec \lambda} v_\lambda\|_{L_{t,x}^2} \lesssim C \epsilon d_\lambda \lambda^{-\frac{1}{2}},$$

where the  $\epsilon$  factor serves to absorb the constant  $C$ . But these are easily obtained from our bootstrap assumption (7.14) after dyadic summation.

□

Now we use interaction Morawetz identities to prove the bilinear bounds in the theorem. For each pair  $v_\lambda, v_\mu$  we define the interaction Morawetz functional

$$(7.20) \quad \mathbf{I}^{x_0}(v_\lambda, v_\mu) := \iint a_j(x-y) (M(v_\lambda)(x) P^j(v_\mu^{x_0})(y) - P^j(v_\lambda)(x) M(v_\mu^{x_0})(y)) dx dy,$$

with  $a_j(x) = \partial_j |x|$ . The time derivative of  $\mathbf{I}^{x_0}(v_\lambda, v_\mu)$  is

$$(7.21) \quad \frac{d}{dt} \mathbf{I}^{x_0}(v_\lambda, v_\mu) = \mathbf{J}^4(v_\lambda, v_\mu^{x_0}) + \mathbf{K}(v_\lambda, v_\mu^{x_0}),$$

where

$$(7.22) \quad \begin{aligned} \mathbf{J}^4(v_\lambda, v_\mu^{x_0}) := & \iint a_{jk}(x-y) (M(v_\lambda) E^{jk}(v_\mu^{x_0}) - P^j(v_\lambda) P^k(v_\mu^{x_0}) \\ & + M(v_\lambda) E^{jk}(v_\mu^{x_0}) - P^j(v_\lambda) P^k(v_\mu^{x_0})) dx dy, \end{aligned}$$

respectively

$$(7.23) \quad \begin{aligned} \mathbf{K}(v_\lambda, v_\mu^{x_0}) := & \iint a_j(x-y) M(v_\lambda)(x) F_{j,\lambda,p}^{4,x_0}(y) + P_j(v_\lambda)(y) F_{\lambda,m}^{4,x_0}(x) \\ & - M(v_\mu^{x_0})(y) F_{j,\lambda,p}^4(x) - P_j(v_\mu^{x_0})(x) F_{\lambda,m}^4(y) dx dy. \end{aligned}$$

We will use the (integrated) interaction Morawetz identity (7.21) to produce  $L^2$  bilinear bounds as follows:

- The spacetime term  $\mathbf{J}^4$  contains at leading order the squared  $L^2$  norm we aim to bound.
- The fixed time expression  $\mathbf{I}^{x_0}$  gives the primary bound for  $\mathbf{J}^4$ .
- The space-time term  $\mathbf{K}$  will be estimated perturbatively.

We successively consider each of these three terms.

**I. The bound for  $\mathbf{I}^{x_0}(v_\lambda, v_\mu)$ .** For this we observe that

$$(7.24) \quad |\mathbf{I}^{x_0}(v_\lambda, v_\mu)| \lesssim (\lambda + \mu) d_\lambda^2 d_\mu^2,$$

which is a direct consequence of the straightforward fixed time density bounds

$$(7.25) \quad \|M(v_\lambda)\|_{L_{t,x}^1} \lesssim d_\lambda^2, \quad \|P^j(v_\lambda)\|_{L_{t,x}^1} \lesssim \lambda d_\lambda^2, \quad \|E^{jk}(v_\lambda)\|_{L_{t,x}^1} \lesssim \lambda^2 d_\lambda^2.$$

These in turn follow from the uniform  $L^2$  bound from  $v_\lambda$  and its frequency localization; the source terms  $f_\lambda$  play no role.

**II. The bound for  $\mathbf{K}$ .** This has the form

$$(7.26) \quad \left| \int_0^T \mathbf{K}(v_\lambda, v_\mu) dt \right| \lesssim (\lambda + \mu) d_\lambda^2 d_\mu^2,$$

and is an immediate consequence of the bounds (7.17) and (7.25).

**III. The contribution of  $\mathbf{J}^4$ .** Here we begin with an algebraic computation for  $\mathbf{J}^4$ , after which we specialize to the balanced, respectively the unbalanced case.

The expression  $\mathbf{J}^4$  can be further written as  
(7.27)

$$\mathbf{J}^4(v_\lambda, v_\mu^{x_0}) = \iint a_{jk}(x-y) (|v_\lambda|^2 E^{jk}(v_\mu^{x_0}) + |u_\mu^{x_0}|^2 E^{jk}(v_\lambda) - 8\Im(\bar{v}_\lambda \partial^j v_\lambda) \Im(\bar{v}_\mu^{x_0} \partial^j v_\mu^{x_0})) \, dx dy,$$

where

$$\begin{aligned} E^{jk}(v_\lambda) &= \partial^j v_\lambda \partial^k \bar{v}_\lambda + \partial^k v_\lambda \partial^j \bar{v}_\lambda - v \partial^j \partial^k \bar{v} - \bar{v} \partial^j \partial^k v \\ &= 2\partial^j v_\lambda \partial^k \bar{v}_\lambda + 2\partial^k v_\lambda \partial^j \bar{v}_\lambda - \partial^j(v \partial^k \bar{v}) - \partial^j(\bar{v} \partial^k v). \end{aligned}$$

At this point we integrate by parts the contributions of the last two terms in  $E^{jk}$ , noting that we have two types of error terms, arising from

- the fact that  $\partial^j$  is not skew-adjoint, and thus we have derivatives applied to the metric
- the fact that the metric  $g$  in  $\partial^j$  may be  $g_{[\leq \lambda]}(x)$  or  $g_{[\leq \mu]}^{x_0}$ , depending on whether this operator is applied to  $v_\lambda$  or to  $v_\mu^{x_0}$ .

We obtain

$$\begin{aligned} \mathbf{J}^4(v_\lambda, v_\mu^{x_0}) &= 4 \iint a_{jk}(x-y) (|v_\lambda|^2 \Re(\partial^j v_\mu^{x_0} \partial^k \bar{v}_\mu^{x_0}) + |u_\mu^{x_0}|^2 \Re(\partial^j v_\lambda \partial^k \bar{v}_\lambda) \\ &\quad - 2\Im(\bar{v}_\lambda \partial^j v_\lambda) \Im(\bar{v}_\mu^{x_0} \partial^j v_\mu^{x_0})) \, dx dy \\ &\quad + 4 \iint a_{jk}(x-y) (\partial^j |v_\lambda|^2 \partial^k |\bar{v}_\mu^{x_0}|^2 + \partial^k |u_\mu^{x_0}|^2 \partial^j |v_\lambda|^2) \, dx dy \\ &\quad - \iint a_{jk}(x-y) |v_\lambda|^2 (g_{[\leq \mu]}^{lj}(y+x_0) - g_{[\leq \lambda]}^{lj}(x)) \partial_l \partial^j |v_\mu^{x_0}|^2 \, dx dy \\ &\quad + 2 \iint a_{jk}(x-y) (\partial_l g_{[\leq \lambda]}^{lj}(x)) (|v_\lambda|^2 \partial^k |v_\mu^{x_0}|^2 + |u_\mu^{x_0}|^2 \partial^k |v_\lambda|^2) \, dx dy \\ &:= \mathbf{J}_{main}^4 + \mathbf{R}_1^4 + \mathbf{R}_2^4. \end{aligned} \tag{7.28}$$

Here the leading term  $\mathbf{J}_{main}^4$  contains the contribution of the first two lines, and the last two lines represent the error terms  $\mathbf{R}_1^4$ , respectively  $\mathbf{R}_2^4$ . Just as in the constant coefficient case,  $\mathbf{J}_{main}^4$  can be rewritten as

$$\mathbf{J}_{main}^4 = 2 \iint a_{jk}(x-y) F^j \bar{F}^k \, dx dy, \tag{7.29}$$

where

$$F^j = \partial^j v_\lambda \bar{v}_\mu^{x_0} + v_\lambda \partial^j v_\mu^{x_0}. \tag{7.30}$$

At this point we separate the analysis into the balanced and the unbalanced case.

*III A. The balanced symmetric case,  $\lambda = \mu$ ,  $x_0 = 0$ .* We use the interaction Morawetz identity (7.21) with  $a(x) = |x|$ . Then we will show that

**Lemma 7.4.** *Under our bootstrap assumptions on  $u$  and our bilinear bootstrap assumptions (7.13)-(7.14) on  $v_\lambda$  we have*

$$\int_0^T \mathbf{J}^4(v_\lambda, v_\lambda) \, dt \gtrsim \| |D_x|^{\frac{3-n}{2}} |v_\lambda|^2 \|_{L^2}^2 + O(C^2 \epsilon^2 \lambda d^4). \tag{7.31}$$

Combined with (7.24) and (7.26), this implies the bound (7.9) in this case.

*Proof.* We begin with the leading term  $J^{4,main}$ , for which we have the algebraic inequality

$$(7.32) \quad \mathbf{J}_{main}^4(v_\lambda, v_\lambda) \geq 2 \iint a_{jk}(x-y) \partial^j |v_\lambda(x)|^2 \partial^k |v_\lambda(y)|^2 dx dy.$$

Here the derivatives are covariant, so we cannot directly use the identity (5.9). We would like to replace  $g$  by the identity matrix,

$$\partial^j |v_\lambda(x)|^2 = \partial_j |v_\lambda(x)|^2 + \partial_m [(g_{[\leq \lambda]}^{jm} - I) |v_\lambda(x)|^2] - (\partial_m g_{[\leq \lambda]}^{jm}) |v_\lambda(x)|^2.$$

By (5.9), the first term on the right applied to both entries in the right hand side of (7.32) yields exactly the desired expression on the right in (7.31).

The contributions of the third term on the right, involving a derivative of the metric, may be included in  $R_2^4$  and will be estimated perturbatively there. It remains to consider the contributions of the second term on the right, which, after integrating by parts and excluding again the terms with a differentiated metric, have the form

$$\begin{aligned} & \iint a_{jk}(x-y) \partial((g_{[\leq \lambda]} - I) |v_\lambda(x)|^2) \partial^k (g |v_\lambda(y)|^2) dx dy dt \\ &= \iint \partial^2 a_{jk}(x-y) ((g_{[\leq \lambda]} - I) |v_\lambda(x)|^2) (g_{[\leq \lambda]} |v_\lambda(y)|^2) dx dy dt. \end{aligned}$$

Here  $\partial^2 a_{jk}$  are smooth homogeneous distributions of order  $-3$ , so they act like  $|D|^{3-n}$ . Hence the above integral is estimated by

$$\lesssim \|(g_{[\leq \lambda]} - I) |v_\lambda(x)|^2\|_{L_t^2 \dot{H}_x^{\frac{3-n}{2}}} \|g_{[\leq \lambda]} |v_\lambda(y)|^2\|_{L_t^2 \dot{H}_x^{\frac{3-n}{2}}}.$$

Here we know that  $\|g_{[\leq \lambda]} - I\|_{H_x^s} \lesssim C^2 \epsilon^2$  so we can use a fixed time multiplicative bound of the form

$$\|(g_{[\leq \lambda]} - I)w\|_{\dot{H}^{\frac{3-n}{2}}} \lesssim \|(g_{[\leq \lambda]} - I)\|_{H^2} \|w\|_{\dot{H}^{\frac{3-n}{2}}},$$

in order to estimate the above expression by

$$\lesssim C^2 \epsilon^2 \| |D_x|^{\frac{3-n}{2}} |v_\lambda|^2 \|_{L_x^2}^2.$$

But this can be absorbed into the leading term on the right in (7.31). Hence we arrive at the bound (7.31) for  $J^{4,main}$ , modulo additional  $\mathbf{R}_2^4$  type terms. It remains to estimate the expressions  $\mathbf{R}_1^4$  and  $\mathbf{R}_2^4$  perturbatively.

To bound  $\mathbf{R}_2^4$  we can start by estimating one of the factors by using two bilinear  $L^2$  bounds to obtain

$$\|\partial g_{[\leq \lambda]} |v_{< \lambda}|^2\|_{L_{t,x}^1} \lesssim C^2 \epsilon^2 d_\lambda^2 \lambda^{-1}.$$

This allows us to bound the entire  $\mathbf{R}_2^4$  as follows

$$(7.33) \quad \left| \int_0^T \mathbf{R}_2^4 dt \right| \lesssim \lambda^2 \| |v_\lambda|^2 \|_{L_t^\infty L_x^1} \|\partial g_{[\leq \lambda]} |v_{< \lambda}|^2\|_{L_{t,x}^1} \lesssim C^2 \epsilon^2 d_\lambda^2 \lambda$$

as needed.

Finally, we turn our attention to  $\mathbf{R}_1^4$ , where we seek to prove a similar bound

$$(7.34) \quad \left| \int_0^T \mathbf{R}_1^4 dt \right| \lesssim C^2 \epsilon^2 d_\lambda^2 \lambda.$$

We recall that the integrand has the form

$$\mathbf{R}_1^4 = a_{jk}(x-y)\lambda^2(g_{[\prec\lambda]}(x) - g_{[\prec\lambda]}(y))|v_\lambda(x)|^2|v_\lambda(y)|^2.$$

Here we begin with a paradifferential decomposition of  $g_{\prec\lambda}$ , based on the highest entry frequency  $\mu \ll \lambda$ ,

$$g_{\prec\lambda}(u) = \sum_{\mu \ll \lambda} g^1(u_{\leq \mu}) u_\mu := \sum_{\mu \ll \lambda} g_{[\mu]},$$

and estimate separately the contribution of each summand. For each  $\mu$ , we split the kernel  $a_{jk}$  on the dual scale, depending on the relative size of  $x-y$  and  $\mu^{-1}$ . For large  $x-y$  we treat the two terms in the  $g_{[\mu]}$  difference as separate, and estimate an expression of the form

$$\iint 1_{|x-y|>\mu^{-1}} a_{jk}(x-y)|v_\lambda(x)|^2 g_{[\mu]}^{lj}(x) \partial_t \partial^j |v_\lambda(y)|^2 dx dy dt.$$

Here we can use two bilinear  $L^2$  bounds to write

$$\| |v_\lambda(x)|^2 g_{[\mu]}^{lj}(x) \|_{L_{t,x}^1} \lesssim C^2 \epsilon^2 d_\lambda^2 \lambda^{-1} \mu^{\frac{n-1}{2}-s}$$

and combine this with the energy bound for the second  $v_\lambda$  factor and the uniform bound

$$|1_{|x-y|>\mu^{-1}} a_{jk}(x-y)| \lesssim \mu,$$

to arrive at the final bound

$$\leq C^2 \epsilon^2 \lambda d_\lambda^2 \mu^{\frac{n+1}{2}-s},$$

which has a trivial  $\mu$  summation for our choice of  $s$ .

For small  $x-y$  we need to exploit the difference instead, but the delicate matter is that we still want to be able to use bilinear  $L^2$  estimates. To achieve this we use the following

**Lemma 7.5.** *There exists a (matrix valued) kernel  $K(w, z)$  in  $B(0, 1) \times \mathbb{R}^n$ , uniformly smooth in  $w$  and uniformly Schwartz in  $h$ , so that for any function  $u$  in  $\mathbb{R}^n$  localized at frequency  $\leq 1$  we have the representation*

$$(7.35) \quad u(x) - u(y) = (x-y) \int K(x-y, z) \nabla u(x+z) dh, \quad |x-y| \leq 1.$$

To apply this to functions which are localized at frequency  $\leq \mu$  we simply rescale and use the kernel

$$K_\mu(z, h) = \mu^n K(\mu z, \mu h).$$

Then we need the constraint  $|x-y| \lesssim \mu^{-1}$ .

*Proof.* We expand the difference on the left using a smooth projector  $P_{\leq 1}$  to frequencies  $\leq 1$  and its associated Schwartz kernel  $K_{\leq 1}$  as

$$\begin{aligned} u(x) - u(y) &= \int_0^1 (x-y) \cdot \nabla u(x+h(y-x)) dh \\ &= \int_0^1 (x-y) \cdot \nabla P_{\leq 1} u(x+h(y-x)) dh \\ &= \int_0^1 \int (x-y) \cdot \nabla u(x+h(y-x)-x_1) K_{\leq 1}(x_1) dh dx_1. \end{aligned}$$

Changing variables to  $h(x - y) - x_1 = z$  we arrive at the desired formula (7.35) with

$$K(x - y, z) = \int K_{\leq 1}(h(y - x) - z) dh,$$

which is uniformly smooth in  $x - y$  and uniformly Schwartz in  $z$  for  $x - y$  inside the unit ball.  $\square$

Now we split the difference

$$g_{[\mu]}(y) - g_{[\mu]}(x) = g^1(u_{\leq \mu}(y))(u_{\mu}(y) - u_{\mu}(x)) + (g^1(u_{\leq \mu})(y) - g^1(u_{\leq \mu})(x))u_{< \mu}(x),$$

and apply the above lemma to the corresponding difference. For the first term above, for instance, we need to estimate the integral

$$\iint 1_{|x-y| < \mu^{-1}} a_{jk}(x-y)(x-y) K_{\mu}(x-y, z) g^1(u_{\leq \mu}(y)) \nabla u_{\mu}(y+z) |v_{\lambda}(x)|^2 \partial_l \partial^j |v_{\lambda}(y)|^2 dx dy dt dh,$$

where we can use two bilinear  $L^2$  estimates in  $y$ , an energy bound in  $x$  and a trivial  $h$  integration to arrive at the same bound as in (7.34), concluding the proof of the Lemma.  $\square$

*III B. The balanced non-symmetric case,  $\mu \approx \lambda$ ,  $x_0 \in \mathbb{R}$ .* Here we do not repeat the computation in the previous, symmetric case, but rather use the result there. Precisely, we apply the symmetric diagonal bound to the functions

$$w_{\lambda} = av_{\lambda} + bv_{\mu}^{x_0}, \quad |a|, |b| \leq 1.$$

To be able to do this, we need to estimate the source term in the paradifferential  $w_{\lambda}$  equation

$$(i\partial_t + \partial_j g_{[\leq \lambda]}^{jk} \partial_k) w_{\lambda} = g_{\lambda},$$

where

$$g_{\lambda} = af_{\lambda} + bf_{\mu}^{x_0} + \partial_j (g_{[\leq \mu]}^{x_0} - g_{[\leq \lambda]}^{jk} \partial_k) v_{\mu}^{x_0}.$$

For  $g_{\lambda}$  we claim that

$$(7.36) \quad \|g_{\lambda} w_{\lambda}^{y_0}\|_{L_{t,x}^1} \lesssim C^2 \epsilon^2 d_{\lambda}^2 (1 + \lambda |x_0|)$$

Here we split the  $g$  difference as

$$g_{[\leq \mu]}^{x_0} - g_{[\leq \lambda]} = (g_{[\leq \mu]}^{x_0} - g_{[\leq \mu]}) + (g_{[\leq \mu]} - g_{[\leq \lambda]}).$$

The contribution of the second difference requires the bound

$$\|\partial(g_{[\leq \mu]} - g_{[\leq \lambda]}) \partial v_{\mu}^{x_0} w_{\lambda}^{y_0}\|_{L_{t,x}^1} \lesssim C^2 \epsilon^2 d_{\lambda}^2,$$

which is obtained (with an extra  $C^2 \epsilon^2$  factor) by applying twice the bootstrap assumption (7.14), since the  $g$  difference is at least quadratic and contains at least one entry at frequency comparable to  $\lambda$ , as  $\mu \approx \lambda$ .

The contribution of the first difference requires the bound

$$\|\partial(g_{[\leq \mu]} - g_{[\leq \mu]}^{x_0}) \partial v_{\mu}^{x_0} w_{\lambda}^{y_0}\|_{L_{t,x}^1} \lesssim C^2 \epsilon^2 \lambda |x_0| d_{\lambda}^2,$$

which is estimated similarly with two bilinear  $L^2$  bounds after expanding

$$g_{[\leq \mu]} - g_{[\leq \mu]}^{x_0} = \int_0^1 x_0 \cdot \nabla g_{< \mu}^{hx_0} dh.$$

Applying the symmetric version of (7.9) to  $w_\lambda$  we obtain the estimate

$$\| |D|^{\frac{3-n}{2}} \Re(abv_\lambda v_\mu^{x_0}) \|_{L_{t,x}^2}^2 \lesssim C^2 \epsilon^2 \lambda d_\lambda^2 (1 + \lambda |x_0|),$$

from which the full bound (7.9) follows by making appropriate choices for  $a$  and  $b$ .

*III C. The unbalanced case  $\mu \ll \lambda$ .* The argument here is similar, but simpler because we will be able to directly treat  $(g - I)$  perturbatively.

The advantage we have is that the dyadic annuli where  $v_\mu$  and  $v_\lambda$  are supported are separate. Here we are allowed to freely localize  $v_\lambda$  with respect to angular directions by applying suitable multipliers:

**Lemma 7.6.** *Let  $A(D)$  be a smooth, bounded multiplier at frequency  $\lambda$ . If  $v_\lambda$  satisfies (7.5), then so does  $A(D)v_\lambda$ .*

*Proof.* The source term in the paradifferential equation for  $A(D)v_\lambda$  is

$$g_{[\lambda]} = A(D)f_\lambda + \partial_j[A(D), g_{[\lambda]}^{jk}] \partial_k v_\lambda.$$

The  $d_\lambda$  bound corresponding to the first term on the right is trivial, while the second term can be written in the form

$$\lambda L(\partial g_{[\lambda]}, v_\lambda).$$

Then its contribution to  $d_\lambda$  can also be easily estimated using two bilinear unbalanced  $L^2$  bounds,

$$\| \lambda L(\partial g_{[\lambda]}, v_\lambda) v_\lambda^{x_0} \|_{L_{t,x}^1} \lesssim \lambda \| L(\partial \mathbf{u}_{<\lambda} v_\lambda) \|_{L_{t,x}^2} \| L(\mathbf{u}_{<\lambda} v_\lambda) \|_{L_{t,x}^2} \lesssim C^2 \epsilon^2 d_\lambda^2,$$

as needed.  $\square$

Frequency localizing  $v_\lambda$  to smaller sets, we may assume that  $v_\mu$  and  $v_\lambda$  are  $O(\lambda)$  uniformly separated in frequency in a well chosen open set  $\Theta$  of angular directions. We still use the weight  $a(x) = |x|$  in our interaction Morawetz identity.

Here we start with  $\mathbf{J}^4$  as in (7.22), and seek to show that

$$(7.37) \quad \int_0^T \mathbf{J}^4(v_\lambda, v_\mu^{x_0}) dt \gtrsim \lambda^2 \mu^{1-n} \| v_\lambda \bar{v}_\mu^{x_0} \|_{L_{t,x}^2}^2 + O(C^2 \epsilon^2 \lambda d_\lambda^2 d_\mu^2).$$

Combined with (7.24) and (7.26), this would imply the unbalanced bilinear bound (7.10).

For this we use the earlier computation in (7.28), and begin by estimating perturbatively the terms  $\mathbf{R}_2^4$  and  $\mathbf{R}_1^4$ . The bound for  $\mathbf{R}_2^4$  is identical to the earlier one in the proof of Lemma 7.4, and is omitted. However, the  $\mathbf{R}_1^4$  bound is more difficult now, because we want our estimate to be uniform in  $x_0$  and so we have no control over the difference of the metrics, other than the naive comparison with the identity matrix. We will argue differently depending on the size of  $x - y$ .

a) If  $x - y$  is small,  $|x - y| \lesssim \mu^{-1}$ , we rewrite the integral as

$$\iint \chi_{|z| < \mu^{-1}} a_{jk}(z) |v_\lambda(x)|^2 (g_{[\mu]}^{lj}(x + z + x_0) - g_{[\lambda]}^{lj}(x)) \partial_l \partial^j |v_\mu(x + x_0 + z)|^2 dx dz.$$

Here we bound the metric difference by  $C^2 \epsilon^2$ , and use twice the bound (3.15) for fixed  $z$ . Finally the  $z$  integral yields another  $\mu^{1-n}$  factor. So the above integral is estimated by

$$\lesssim C^2 \epsilon^2 \mu^2 \lambda^{-1} \mu^{n-1} \mu^{1-n} = C^2 \epsilon^2 \mu^2 \lambda^{-1},$$

which suffices.

a) If  $x - y$  is large,  $|x - y| \gtrsim \mu^{-1}$ , then we integrate by parts to move the two derivatives on  $a$ . If one derivative falls on the metric we get an  $\mathbf{R}_2^4$  type term which is good. Else, we obtain an integral of the form

$$\iint \partial^2(\chi_{>\mu^{-1}} a_{jk})(x - y) |v_\lambda(x)|^2 (g_{[\leq \mu]}^{lj}(y + x_0) - g_{[\leq \lambda]}^{lj}(x)) g_{<\mu}(y + x_0) |v_\mu(y + x_0)|^2 dx dy.$$

We separate the  $(g - I)$  factors and regroup with either  $|v_\lambda|^2$  or  $|v_\mu|^2$ , The Fourier transform of  $\partial^2(\chi_{>\mu^{-1}} a_{jk})$  is bounded by  $\lesssim |\xi|^{3-n}$ , so we estimate the above integral by

$$\lesssim C^2 \epsilon^2 \| |D|^{\frac{3-n}{2}} |v_\lambda|^2 \|_{L_{t,x}^2} \| |D|^{\frac{3-n}{2}} |v_\mu|^2 \|_{L_{t,x}^2},$$

where the  $C^2 \epsilon^2$  factor comes from  $(g - I)$ . This we can estimate using (7.15) twice.

We are now left with the leading term  $J_{main}^4$ , where we have the chief advantage that the integrand is nonnegative pointwise. This allows us to localize it on the  $\mu^{-1}$  scale around the diagonal, using a nonnegative Schwartz cutoff  $\phi(\mu x)$  to be chosen later:

$$\mathbf{J}_{main}^4 \gtrsim \iint \phi(\mu(x - y)) a_{jk}(x - y) F^j \bar{F}^k d\theta.$$

At this point we can perturbatively replace the metric  $g$  with the identity in  $F^j$ , with the same argument as in the case of  $\mathbf{R}_1^4$ , arriving at

$$\mathbf{J}_{main}^4 \gtrsim \mathbf{J}_{flat}^4 + O(C^2 \epsilon^2 d_\lambda^2 d_\mu^2), \quad \mathbf{J}_{flat}^4 := \iint \phi(\mu(x - y)) a_{jk}(x - y) F_j \bar{F}_k d\theta.$$

It remains to show that we have the constant coefficient bound

$$(7.38) \quad \int_0^T \mathbf{J}_{flat}^4 dt \gtrsim \lambda^2 \mu^{1-n} \|v_\lambda \bar{v}_\mu^{x_0}\|_{L_{t,x}^2}^2.$$

To simplify the analysis we decompose our weight  $a$  as

$$a(x) = |x| = c_n \int_{\mathbf{S}^n} a^\theta(x) d\theta, \quad a^\theta(x) = |x \cdot \theta|.$$

This leads to a corresponding nonnegative foliation of the Hessian

$$D^2 a = c_n \int_{\mathbf{S}^n} D^2 a^\theta(x) d\theta,$$

and further

$$\mathbf{J}_{flat}^4 = \int_{\mathbf{S}^n} \iint \phi(\mu(x - y)) a_{jk}^\theta(x - y) F_j \bar{F}_k dx dy d\theta := \int_{\mathbf{S}^n} \mathbf{J}_\theta^4 d\theta.$$

It suffices now to prove that the bound (7.38) holds for  $\mathbf{J}_\theta^4$  for an open set of directions  $\theta$ . Here we choose exactly the set of directions in which  $v_\mu$  and  $v_\lambda$  are separated in frequency. Without any loss of generality assume that  $\theta = e_1$ . Then we are looking at

$$\mathbf{J}_{flat}^4(v_\lambda, v_\mu^{x_0}) = \int_{x_1=y_1} \phi(\mu(x' - y')) |F_1|^2 dx dy',$$

where

$$F_1(x, y) := \partial_1 v_\lambda(x) \bar{v}_\mu^{x_0}(y) + v_\lambda \partial_1 \bar{v}_\mu^{x_0}(y).$$



The last form of the right hand side allows us to interpret  $F$  as a bilinear form for a fixed translation parameter  $z$ , and the cutoff  $\chi_\mu$  allows us to estimate  $z'$  integrals.

Given our angular frequency localizations, within the support of  $v_\lambda$  we have  $\xi_1 \approx \lambda$ , while in the support of  $v_\mu$  we have  $|\xi_1| \lesssim \mu$ . Hence the first term in  $F_1$  is the leading one, while the second is smaller. Precisely, using the bootstrap bilinear bound we can write

$$\int \int_{z_1=0} \chi_\mu(z) |F_1(x, y+z)|^2 dx dz' dt = \int \int_{z_1=0} \chi_\mu(z) |\partial_1 v_\lambda(x)|^2 |v_\mu^{x_0}(x+z)|^2 dx dz' dt + O(C^2 \mu d_\lambda^2 d_\mu^2).$$

For a well chosen bump function  $\chi$ , the frequency localization of  $v_\mu$  allows us to collapse the integral on the right to

$$\mu^{1-n} \int |\partial_1 v_\lambda(x)|^2 |u_\mu^{x_0}(x)|^2 dx dt \gtrsim \|\partial_1(v_\lambda v_\mu^{x_0})\|_{L_{t,x}^2}^2,$$

where the  $x_1$  derivative yields a  $\lambda$  factor due to the frequency separation between  $v_\lambda$  and  $v_\mu$  in the  $x_1$  direction, and where the derivative applied to  $v_\mu$  yields a smaller, perturbative  $\mu$  factor. This implies (5.6) and thus (7.37), concluding the proof of the theorem in the final unbalanced case. □

## 8. LOW REGULARITY STRICHARTZ BOUNDS FOR THE PARADIFFERENTIAL FLOW

While the bilinear  $L^2$  bounds play the leading role in estimating unbalanced interactions, the balanced interactions are instead controlled using Strichartz bounds. Since here we are considering cubic problems, the  $L^4$  bound is the single most important one. The way we handle the  $L^4$  bounds depends somewhat on the dimension, where we distinguish two cases:

- (1) high dimension,  $n \geq 3$ . There, on one hand,  $L^4$  is not a sharp Strichartz norm, and on the other hand the  $L^4$  bound is obtained directly from the interaction Morawetz analysis, so this section is not playing any role in obtaining it.
- (2) low dimension,  $n = 2$ . Here  $L^4$  is a sharp Strichartz norm, but the  $L^4$  bound cannot be obtained directly from the interaction Morawetz analysis. This is why this section is needed.

Further elaborating on the case  $n = 2$ , we can make a further distinction [in this case](#) there depending on whether we seek local or global in time estimates:

- (1)  $n = 2$ , local in time bounds (needed for local well-posedness). The interaction Morawetz analysis yields a lossless  $L_t^4 L_x^8$  bound, which interpolated with the  $L_t^\infty L_x^2$  energy estimate gives a lossless  $L_t^6 L_x^4$  bound. Applying Holder's inequality in time yields an  $L_{t,x}^4$  bound with a  $1/6$  derivative loss, which suffices for the local result.
- (2)  $n = 2$ , global in time bounds (generically needed for global well-posedness). This is the single most interesting case, and also, the one case where the analysis here is absolutely necessary, providing an  $L_{t,x}^4$  bound with a  $1/2$  derivative loss.

Comparing the above discussion with our prior work [36] in one space dimension, we note that there the  $L_{t,x}^6$  bound was of the main one. For the local in time analysis, we proved a sharp  $L_{t,x}^6$  bound, which by interpolation and Holder's inequality gave an  $L_{t,x}^4$  bound with a  $1/4$  derivative loss, which sufficed. For the global in time analysis on the other hand, we used the interaction Morawetz analysis to prove a global  $L_{t,x}^6$  bound with a loss of derivatives;

this in turn required the conservative and defocusing assumption, but avoided any need for an  $L^4_{t,x}$  bound.

Instead, in the  $n = 2$  case, no defocusing assumption is necessary. The conservative assumption does play a role though, in reducing the Sobolev regularity at which the global result holds; this would have otherwise required an  $L^4_{t,x}$  bound with a smaller  $1/4$  derivative loss.

The result in this section provides a full range of Strichartz estimates in all dimensions, globally in time, but with a loss of derivatives: precisely one full derivative at the Pecher endpoint. Strictly speaking we need it only in two space dimensions, but the proof is new and dimension independent, and we believe that the bounds it provides are of independent interest also in higher dimensions, and that the ideas here will also be useful for other problems.

To review the set-up here, we seek  $L^p_t L^q_x$  Strichartz bounds for the paradifferential problem

$$(8.1) \quad i\partial_t v_\lambda + \partial_x g_{[\prec \lambda]} \partial_x v_\lambda = f_\lambda, \quad v_\lambda(0) = v_{0,\lambda},$$

where the pair of indices  $(p, q)$  lies on the sharp Strichartz line

$$(8.2) \quad \frac{2}{p} + \frac{n}{q} = \frac{n}{2}, \quad p \geq 2.$$

We will refer to the case  $p = 2$  as the Pecher endpoint, which is forbidden in dimension  $n = 2$ .

Classically one places the source term in dual Strichartz spaces. While we certainly want to allow such sources, in our analysis we would like to allow for a larger class of functions which is consistent with our needs for the quasilinear problem. One such class will be the space  $DV^2_\Delta$  associated to the flat Schrödinger flow. Another class may be defined by testing against solutions to the adjoint constant coefficient problem

$$(8.3) \quad (i\partial_t + \Delta)w_\lambda = h_\lambda, \quad w(T) = w_T,$$

defining a norm as follows:

$$(8.4) \quad \|f_\lambda\|_{N^\sharp} := \sup_{\|h_\lambda\|_{S'} + \|w_T\|_{L^2_{t,x}} \leq 1} \left| \int f_\lambda w_\lambda dx dt \right|.$$

For technical reasons we will use both  $DV^2_\Delta$  and  $N^\sharp$ ; the first choice turns out to be better in the low dimension  $n = 2$ , and the second is better in the higher dimension  $n \geq 3$ . Their common property is provided by the embeddings

$$(8.5) \quad N \subset N^\sharp, \quad N \subset DV^2,$$

which hold regardless of the dimension, where  $N$  represents any of the admissible sharp dual Strichartz norms.

We will also need to measure  $f_\lambda$  relative to  $v_\lambda$ , via the quantity

$$I(v_\lambda, f_\lambda^2) = \sup_{x_0} \|v_\lambda f_\lambda^{2,x_0}\|_{L^1_{t,x}}.$$

One consequence of allowing such terms is that we can commute the  $x$  derivatives as we like in the equation, placing errors in the perturbative source term  $f_\lambda$ .

**Theorem 13.** Assume that  $u$  solves (QNLS) in a time interval  $[0, T]$ , and satisfies the bootstrap bounds (3.14), (3.15), (3.16) and (3.17) for some  $s > \frac{n+1}{2}$ . Then the following Strichartz estimates hold for the linear paradifferential equation (4.5) for any pair of sharp Strichartz exponents  $(p, q)$  and  $(p_1, q_1)$ :

$$(8.6) \quad \|v_\lambda\|_{L_t^\infty L_x^2} + \lambda^{-1} \|v_\lambda\|_{V_\Delta^2} \lesssim \|v_{\lambda,0}\|_{L_x^2} + \|f_\lambda\|_{DV_\Delta^2} + I(v_\lambda, f_\lambda)^{\frac{1}{2}}, \quad n = 2,$$

$$(8.7) \quad \|v_\lambda\|_{L_t^\infty L_x^2} + \lambda^{-1} \|v_\lambda\|_{L^2 L^{\frac{2n}{n-2}}} \lesssim \|v_{\lambda,0}\|_{L_x^2} + \|f_\lambda\|_{N^\#} + I(v_\lambda, f_\lambda)^{\frac{1}{2}}, \quad n \geq 3.$$

In particular we have

$$(8.8) \quad \lambda^{-\frac{2}{p}} \|v_\lambda\|_{L_t^p L_x^q} \lesssim \|v_{\lambda,0}\|_{L^2} + \lambda^{\frac{2}{p_1}} \|f_\lambda\|_{L_t^{p_1'} L_x^{q_1'}}.$$

*Proof.* We first argue that (8.8) follows from (8.6), respectively (8.7). On the left, we have a direct interpolation in the case of (8.7), while in the two dimensional case the interpolation is slightly more subtle,

$$\lambda^{-\frac{2}{p}} \|v_\lambda\|_{L_t^p L_x^q} \lesssim \lambda^{-\frac{2}{p}} \|v_\lambda\|_{U_\Delta^p} \lesssim \|v_\lambda\|_{L_t^\infty L_x^2} + \lambda^{-1} \|v_\lambda\|_{V_\Delta^2}.$$

For the second term on the right in the case of (8.7) we use the embedding  $N \subset N^\#$ . Similarly, in the case of (8.6) we have the embedding  $L_t^{p_1'} L_x^{q_1'} \subset DV_\Delta^{p_1'}$ . Finally, the expression  $I(v_\lambda, f_\lambda)$  can be estimated directly using Hölder's inequality.

Now we turn our attention to the proof of (8.6) and (8.7). We trivially have energy estimates

$$\|v_\lambda\|_{L_t^\infty L_x^2} \lesssim \|v_\lambda(0)\|_{L_x^2} + I(v_\lambda, f_\lambda)^{\frac{1}{2}},$$

so it remains to prove that

$$(8.9) \quad \lambda^{-1} \|v_\lambda\|_{V_\Delta^2} \lesssim \|v_{\lambda,0}\|_{L_x^2} + \|f_\lambda\|_{DV_\Delta^2} + I(v_\lambda, f_\lambda)^{\frac{1}{2}}, \quad n = 2$$

respectively

$$(8.10) \quad \lambda^{-1} \|v_\lambda\|_{L^2 L^{\frac{2n}{n-2}}} \lesssim \|v_{\lambda,0}\|_{L_x^2} + \|f_\lambda\|_{N^\#} + I(v_\lambda, f_\lambda)^{\frac{1}{2}}, \quad n \geq 3.$$

For this we reinterpret our equation as a flat Schrödinger equation by moving the  $(g - I)$  contribution to the right, and then viewing it as a source term, which can be estimated as in  $d_\lambda$  using two bilinear  $L^2$  bounds. Precisely, we have

$$(8.11) \quad (i\partial_t + \Delta)v_\lambda = f_\lambda + f_\lambda^1, \quad v_\lambda(0) = v_{0,\lambda},$$

where

$$(8.12) \quad f_\lambda^1 := \partial_x(g_{[\cdot, \lambda]} - I)\partial_x v_\lambda.$$

For (8.9) we estimate directly

$$\|v_\lambda\|_{V_\Delta^2} \lesssim \|v_\lambda(0)\|_{L_x^2} + \|f_\lambda\|_{DV_\Delta^2} + \|f_\lambda^1\|_{DV_\Delta^2},$$

where only the last term on the right still needs to be controlled,

$$(8.13) \quad \|f_\lambda^1\|_{DV_\Delta^2}^2 \lesssim \lambda(\|v_\lambda(0)\|_{L_x^2}^2 + I(v_\lambda, f_\lambda)).$$

We postpone the proof of the last bound, and turn our attention to (8.10). There the argument goes by duality. Given  $w_\lambda$  a solution to the constant coefficient adjoint Schrödinger equation (which is the same) with source term  $h_\lambda \in S'$  we write the duality relation

$$\langle v_\lambda, w_\lambda \rangle_0^T = \int_0^T \int v_\lambda g_\lambda - w_\lambda (f_\lambda^1 + f_\lambda^2 + f_\lambda^3) dx dt,$$

where we maximize the first term on the right over  $\|h_\lambda\|_{S'} \leq 1$ , which in turn gives  $\|v\|_S \lesssim 1$ . This yields

$$\|v_\lambda\|_S \lesssim \|v_\lambda(0)\|_{L^2} + \|f_\lambda\|_{N^\#} + \|f_\lambda^1\|_{N^\#}.$$

It remains to control the last term,

$$(8.14) \quad \|f_\lambda^1\|_{N^\#}^2 \lesssim \lambda(\|v_\lambda(0)\|_{L_x^2}^2 + I(v_\lambda, f_\lambda)).$$

This requires an  $L^1$  bound for  $w_\lambda f_\lambda^1$ , for which we write

$$\|w_\lambda f_\lambda^1\|_{L_{t,x}^1} \lesssim \lambda^2 \|v_\lambda u_{<\lambda}\|_{L_{t,x}^2} \|w_\lambda u_{<\lambda}\|_{L_{t,x}^2}.$$

Here we observe that for  $w_\lambda$  we can control the associated  $d_\lambda$  as in (7.5), so we can use two unbalanced bilinear  $L^2$  estimates from Theorem 12, respectively Corollary 7.1 to bound this by

$$\lesssim \lambda(\|v_\lambda(0)\|_{L_x^2} + I(v_\lambda, f_\lambda)^{\frac{1}{2}}),$$

as needed.

The proof of (8.13) is somewhat similar. By duality it suffices to estimate in  $L^1$  the product  $f_\lambda^1 w_\lambda$  but this time for  $w_\lambda \in U_\Delta^2$ . It suffices to do this for an  $U_\Delta^2$  atom,

$$w_\lambda = \sum 1_{I_k} w_{\lambda,k},$$

where  $I_k$  are arbitrary disjoint time intervals and  $w_{\lambda,k}$  are  $L^2$  homogeneous Schrödinger waves. Using Cauchy-Schwarz and two bilinear  $L^2$  bounds for each of the time intervals  $I_k$  we have

$$\begin{aligned} \|w_\lambda f_\lambda^1\|_{L_{t,x}^1} &\lesssim \lambda^2 \|v_\lambda u_{<\lambda}\|_{L^2} \left( \sum_k \|w_{\lambda,k} u_{<\lambda}\|_{L_{t,x}^2(I_k)}^2 \right)^{\frac{1}{2}} \\ &\lesssim \lambda(\|v_\lambda(0)\|_{L_x^2} + I(v_\lambda, f_\lambda)^{\frac{1}{2}}) \left( \sum_k \|w_{\lambda,k}(0)\|_{L_x^2}^2 \right)^{\frac{1}{2}}, \end{aligned}$$

as needed. □

## 9. THE LOCAL/GLOBAL WELL-POSEDNESS RESULT

In this section we prove our main local well-posedness result in Theorems 3,4. The same proof also yields the global well-posedness result in Theorem 6 in high dimensions, as well as, with small adjustments, the global well-posedness result in Theorem 7. For simplicity we will work with the problem (QNLS); the corresponding arguments for (DQNLS) are identical, and are omitted.

Our starting point here is the local well-posedness result for regular data in Theorem 1. For these solutions we will prove the frequency envelope bounds in Theorems 9, 10[(i),(iii)], which in turn allow us to continue the solutions, with uniform bounds, as follows:

- (1) globally in time if  $n \geq 3$ ,
- (2) globally in time if  $n = 2$  for restricted  $s$  i.e.,  $s \geq \frac{7}{4}$ .

(3) up to time  $\epsilon^{-6}$  if  $n = 2$  at low regularity  $s > \frac{3}{2}$ .

We will then prove Theorems 3,4,6,7 by constructing rough solutions as unique limits of smooth solutions. We proceed in several steps:

**9.1. A-priori bounds for regular solutions: Proof of Theorems 9, 10[(i)(iii)].** We recall that we can freely make the bootstrap assumption that the bounds (3.14) and (3.15) hold with a large universal constant  $C$ . Then we seek to apply Theorem 12 with  $v_\lambda = u_\lambda$ . To achieve this we write the equation for  $u_\lambda$  in the paradifferential form (4.4), using Proposition 7.2 for the source term bound.

This concludes the proof of the above Theorems. For later use, we note that in particular the bounds for the paradifferential source terms in Proposition 7.2 are also valid for our solutions.

**9.2. Higher regularity.** Here we consider regular solutions as above, with initial data  $u_0$  which is not only  $\epsilon$  small in  $H^s$ , but also belongs to a smaller space  $H^\sigma$ , say

$$\|u_0\|_{H^s} \leq \epsilon, \quad \|u_0\|_{H^{s_1}} \leq M, \quad \sigma > s,$$

where  $M$  is possibly very large. Then we claim that we have a uniform bound for the solution in  $H^\sigma$ , precisely

$$(9.1) \quad \|u\|_{L_t^\infty H_x^\sigma} \lesssim M.$$

To prove this, for the initial data  $u_0$  we consider a minimal  $H^s$  frequency envelope  $\epsilon c_\lambda$  with the unbalanced slowly varying condition as in Remark 3.1. Then by construction we must also have

$$\sum_\lambda (\epsilon c_\lambda \lambda^{\sigma-s})^2 \lesssim M^2.$$

By Theorems 9, 10 the frequency envelope  $\epsilon c_\lambda$  is propagated along the flow, and we obtain

$$\|u(t)\|_{H^\sigma}^2 \lesssim \sum_\lambda (\epsilon c_\lambda \lambda^{\sigma-s})^2 \lesssim M^2, \quad t \in [0, T].$$

We remark that not only the uniform  $L^2$  higher regularity bounds are propagated along the flow, but also the corresponding bilinear  $L^2$  and Strichartz bounds.

**9.3. Continuation of regular solutions.** Given a regular initial data  $u_0 \in H^\sigma$  where  $\sigma > \frac{n}{2} + \frac{3}{2}$ , and which is also small in  $H^s$ , we ask whether a regular solution exists and for how long it can be continued:

**Proposition 9.1.** *For every regular initial data  $u_0 \in H^\sigma$  and which also satisfies the smallness condition*

$$(9.2) \quad \|u_0\|_{H^s} \leq \epsilon \ll 1,$$

*there exists a local solution  $u \in C_t H_x^\sigma$ . Furthermore, this local solution can be continued for as long as*

$$\|u\|_{L_t^\infty H_x^s} \lesssim \epsilon.$$

Combining this result with Theorems 9, 10[(i)(iii)], we obtain the following lifespan bounds:

**Corollary 9.2.** *Given initial data as in Proposition 9.1, a regular solution exists*

- (i) *globally in time in dimension  $n \geq 3$ ,*
- (ii) *up to time  $O(\epsilon^{-6})$  in dimension  $n = 2$ .*

*Proof of Proposition 9.1.* For local solutions we want to apply Theorem 1. We remark that even if we have the data  $H^s$  smallness in (9.2), there is no guarantee that the its  $H^\sigma$  size is small. Hence we need the large data version of local well-posedness from [52], which in turn requires verifying the nontrapping assumption. Indeed, we claim that

**Lemma 9.3.** *Let  $\epsilon \ll 1$ . Then the initial data  $u_0$  satisfying the smallness condition (9.2) are uniformly nontrapping.*

Here by uniformity we mean that the intersection of any geodesic for the metric  $g$  with a ball  $B(x_0, r)$  has length  $\lesssim r$ , with a uniform constant.

Assuming that the lemma holds, the large data local well-posedness theorem in [52] guarantees a local  $H^\sigma$  solution. Let  $[0, T)$  be the maximal interval on which an  $H^\sigma$  solution exists. Our apriori bounds guarantee that the  $H^\sigma$  size of the solution remains bounded,

$$\|u(t)\|_{H^\sigma} \lesssim \|u(0)\|_{H^\sigma}, \quad t \in [0, T),$$

as well as that we have a corresponding  $H^\sigma$  frequency envelope bound, uniformly in time. This does not directly guarantee that the limit

$$u(T) = \lim_{t \rightarrow T} u(t)$$

exists in  $H^\sigma$ . However, by examining the (QNLS) equation one easily sees that the limit exists in  $H^{\sigma-2}$ , which combined with the frequency envelope bound implies that the limit exists in  $H^s$ . This in turn contradicts the maximality of  $T$ , and concludes the proof of the proposition.

It remains to prove the nontrapping lemma:

*Proof of Lemma 9.3.* The equations for the Hamilton flow have the form

$$\dot{x}^j = 2g^{jk}\xi_k, \quad \dot{\xi}_j = \partial_{x_j} g^{kl}\xi_k\xi_l,$$

where  $g = g(u_0)$ . Here the full symbol  $g^{jk}\xi_j\xi_k$  is a conserved quantity, so once we fix  $|\xi(0)| = 1$  by scaling, we have the global uniform bound  $|\xi(t)| \approx 1$ .

We will prove that the bicharacteristics are nearly straight, in the sense that

$$(9.3) \quad |\xi(t) - \xi(0)| \lesssim \epsilon, \quad |\dot{x}(t) - \dot{x}(0)| \lesssim \epsilon.$$

Given that  $|g - I| \lesssim \epsilon$ , the second bound clearly implies the first, so it remains to prove the second. For this we make a bootstrap assumption

$$(9.4) \quad |\xi(t) - \xi(0)| \leq C\epsilon,$$

with a large universal constant  $C$ . Then we also have

$$|\dot{x}(t) - \dot{x}(0)| \lesssim C\epsilon \ll 1,$$

which in turn indicates that our bicharacteristic, call it  $\gamma$ , is nearly straight. It remains to estimate

$$|xi(T) - \xi(0)| \lesssim \int_0^T |\nabla g(x(t))| ds \lesssim \int_0^t |u_0(x(t))| |\nabla u_0(x(t))| ds.$$

Since  $\gamma$  is nearly straight and  $s > \frac{n+1}{2}$ , by the trace theorem it follows that

$$\|u\|_{L^2(\gamma)} + \|\nabla u\|_{L^2(\gamma)} \lesssim \|u_0\|_{H^s} \lesssim \epsilon.$$

Then by Cauchy-Schwarz inequality we have

$$|\xi(T) - \xi(0)| \lesssim \epsilon^2,$$

which closes our bootstrap and thus the proof of the nontrapping lemma.  $\square$

This in turn completes the proof of the continuation proposition.  $\square$

**9.4. A-priori  $L^2$  bounds for the linearized equation: Proof of Theorem 11.** Here we seek again to apply Theorem 12 by writing the linearized equation in the form (4.5). Correspondingly, we take  $d_\lambda$  to be an  $L^2$  frequency envelope for the initial data.

It suffices then to make a bootstrap assumption for  $v_\lambda$ , which is the same as in the proof of Theorem 12, and to prove the following bounds on the source terms  $N_\lambda^{lin}v$ , which represent the linearized equation counterpart of Proposition 7.2:

**Proposition 9.4.** *Let  $s \geq \frac{n+1}{2}$ . Assume that the function  $u$  satisfies the bounds (3.1), (3.2) and (3.3) in a time interval  $[0, T]$ , where  $T \leq \epsilon^{-6}$  in dimension  $n = 2$ . Assume also that  $v$  satisfies the bootstrap bounds (7.13)-(7.16). Then for  $\epsilon$  small enough, the functions  $N_\lambda^{lin}v$  satisfy*

$$(9.5) \quad \|N_\lambda^{lin}v \cdot \bar{v}_\lambda^{x_0}\|_{L^1_{t,x}} \lesssim \epsilon^2 d_\lambda^2.$$

*The similar bound holds if we replace  $\bar{v}_\lambda^{x_0}$  by  $\bar{w}_\lambda$ , where  $w_\lambda$  is a solution for the linear flat Schrödinger flow (8.3).*

*Proof.* This proof is virtually identical to the proof of Proposition 7.2, so the details are left for the interested reader.  $\square$

**9.5. Rough solutions as limits of smooth solutions.** Here we show that the  $H^s$  energy estimates for the full equation, combined with the  $L^2$  energy estimates for the linearized equation, both in the frequency envelope formulation, imply the local well-posedness results in Theorems 3, 4, as well as the global well-posedness results in Theorems 6 and 7. A standard argument applies here, and for which we only outline the steps. For more details we refer the reader to the same argument in our prior 1D paper [36], as well as the expository paper [32], where the strategy of the proof is presented in detail. Here it is more convenient to use the dyadic notation for frequencies, allowing also for an easier comparison with [36], [32]. The steps are as follows:

**I. Initial data regularization:** given  $u_0 \in H^s$ , with frequency envelope  $\epsilon c_k$ , we consider the regularized data

$$u_0^h = P_{<h} u_0,$$

where  $h \in \mathbb{R}^+$ , for which we can use an improved frequency envelope

$$c_j^h = \begin{cases} c_j & j \leq h, \\ c_h 2^{-N(j-h)} & j > h, \end{cases}$$

with decay beyond frequency  $2^h$ .

For this data we consider the corresponding smooth solutions  $u^h$ . Using the continuation result for regular solutions in Proposition 9.1 and the bounds in Theorems 9 and 10, a continuity argument shows that these solutions extend as regular solutions up to times  $T$  as in Theorems 9 and 10 with similar frequency envelope bounds

$$(9.6) \quad \|P_j u^h\|_{L_t^\infty L_x^2} \lesssim \epsilon c_j^h 2^{-sj}.$$

**II. Difference bounds:** Defining

$$v^h = \frac{d}{dh} u^h,$$

which solves the linearized equation around  $u^h$ , with initial data

$$v^h(0) = P_h u_0,$$

satisfying

$$\|v^h(0)\|_{L_x^2} \lesssim \epsilon 2^{-sh} c_h,$$

we use the linearized energy estimates in Theorem 11 to obtain the  $L^2$  bounds

$$(9.7) \quad \|v^h\|_{L_t^\infty L_x^2} \lesssim \epsilon 2^{-sh} c_h.$$

**III. Convergence:** Here we define the rough solution  $u$  as the limit of  $u^h$  via the telescopic sum

$$u = u^0 + \sum_{k \geq 0} u^{k+1} - u^k.$$

By (9.7) we have

$$(9.8) \quad \|u^{k+1} - u^k\|_{L_t^\infty L_x^2} \lesssim \epsilon 2^{-sk} c_k,$$

which implies rapid convergence of the series in  $L^2$ , and

$$(9.9) \quad \|u - u^k\|_{L_t^\infty L_x^2} \lesssim \epsilon 2^{-sk} c_k.$$

On the other hand by (9.6) we have

$$(9.10) \quad \|u^{k+1} - u^k\|_{L_t^\infty H_x^{s+m}} \lesssim \epsilon 2^{sm} c_k.$$

Combining the last two relations, it follows that the summands in the series are almost orthogonal in  $H^s$ , and we have

$$(9.11) \quad \|u^k - u^j\|_{L_t^\infty H_x^s} \lesssim \epsilon c_{[j,k]}, \quad j < k.$$

This shows convergence in  $L_t^\infty H_x^s$ , and

$$(9.12) \quad \|u - u^j\|_{L_t^\infty H_x^s} \lesssim \epsilon c_{\geq j}.$$

Now it is easily verified that the limit  $u$  solves the equation (QNLS), and also that it satisfies all the bounds in Theorems 9 and 10.

**IV. Continuous dependence:** This again follows the argument in [32]. The weak Lipschitz dependence implies convergence in a weaker topology, which is then upgraded to strong convergence using the uniform decay of the high frequency tails derived from frequency envelope bounds.



**9.6. Scattering.** Our aim here is to prove the scattering part of our global well-posedness theorems. For clarity we begin with the simpler higher dimensional case in Theorem 6, and then discuss the differences in the 2D case.

We begin by writing the equation for each  $u_\lambda$  as a constant coefficient Schrödinger flow,

$$(i\partial_t + \Delta)u_\lambda = f_\lambda := N_\lambda(u) - (g_{[\cdot, \lambda]} - I)\partial^2 u_\lambda.$$

Now we estimate the source term  $f_\lambda$ , taking our cue from Section 8. For  $N_\lambda$  we have the bounds from Proposition 7.2, while for the second term on the right we repeat the proof of (8.14). This yields

$$(9.13) \quad \|f_\lambda\|_{N^\sharp} \lesssim \epsilon^3 c_\lambda^3 \lambda^{-s+1},$$

which has a one derivative loss. This bound holds globally in time, but if we restrict the time interval we further obtain the decay

$$(9.14) \quad \lim_{T \rightarrow \infty} \|f_\lambda\|_{N^\sharp[T, \infty)} = 0.$$

This is a consequence of the fact that the proof of (9.13) uses at least one translation free bilinear  $L^2$  bound, for which we can directly use the absolute continuity of the  $L^2$  norm.

We remark that we could also simply use  $L^{\frac{4}{3}}$  bounds, but that would be non-optimal as it would involve a larger loss of derivatives (two in 3D). The bound (9.13) and the above decay property allow us to estimate

$$(9.15) \quad \|u_\lambda(t) - e^{i(t-s)\Delta} u_\lambda(s)\|_{L_x^2} \lesssim \epsilon^3 c_\lambda^3 \lambda^{-s+1},$$

with the continuity property

$$(9.16) \quad \lim_{t, s \rightarrow \infty} \|u_\lambda(t) - e^{i(t-s)\Delta} u_\lambda(s)\|_{L_x^2} = 0.$$

This implies that the limit

$$u_\lambda^\infty = \lim_{t \rightarrow \infty} e^{-it\Delta} u_\lambda(t)$$

exists in  $L^2$ , and

$$(9.17) \quad \|u_\lambda^\infty - u_{0\lambda}\|_{L^2} \lesssim \epsilon^3 c_\lambda^3 \lambda^{-s+1}.$$

On the other hand, passing to the limit in the energy bounds for  $u_\lambda$  we also get

$$(9.18) \quad \|u_\lambda^\infty\|_{L_x^2} \lesssim \epsilon c_\lambda \lambda^{-s}.$$

Then we can set

$$u^\infty = \sum_\lambda u_\lambda^\infty \in H^s,$$

where, combining the above properties, we have the strong limit

$$(9.19) \quad \lim_{t \rightarrow \infty} e^{-it\Delta} u(t) = u^\infty \quad \text{in } H^s,$$

i.e. in the strong topology. This concludes the proof of the scattering result in dimensions three and higher.

The proof of the scattering result in two dimensions is similar, with only two differences:

- The space  $N^\sharp$  is now replaced by  $DV_\Delta^2$ , exactly as in the proof of the Strichartz estimates 8.

- The bound for  $N_\lambda$  now separates into two parts: the unbalanced part  $F_\lambda$ , for which we can still use the estimates in Proposition 7.2, and the cubic balanced part  $C_\lambda$ , for which we simply use the  $L^4$  bound (3.7). This allows us to estimate  $C_\lambda$  in  $L^{\frac{4}{3}} \subset DV^2$ . We remark that the  $L^4$  bound (3.7) has a loss of  $1/2$  derivative, but this is acceptable since our target is a bound with a one derivative loss, and we are at least one half derivative above scaling.

## 10. THE 2D GLOBAL RESULT AT LOW REGULARITY

The aim of this section is to complete the proof of the global well-posedness result in Theorem 8. In view of the local well-posedness result in Theorem 3 and its associated frequency envelope bounds in Theorem 10, it suffices to show that the frequency envelope bounds in Theorem 10 hold globally in time. Further, it suffices to do this under appropriate bootstrap assumptions, see Proposition 3.2.

For the proof of the local well-posedness result in Theorem 3, as well as the global result, except for the low regularity regime in dimension  $n = 2$ , it was enough to interpret the (QNLS) evolution as a paradifferential equation with a perturbative source term. Here, on the other hand, we also single out the balanced cubic terms. At higher regularity  $s \geq \frac{7}{4}$ , these balanced cubic terms were estimated using the  $L^4$  Strichartz estimates (3.7) with a half derivative loss. At lower regularity, this loss turns out to be too large, and instead we need a more complex argument. For clarity, we begin with a brief description of our strategy for balanced cubic terms:

- (i) Low frequency terms: for these we simply establish and use an  $L^4$  Strichartz bound perturbatively.
- (ii) High frequency terms without phase rotation symmetry. These may be further divided into two terms
  - transversal, where the three input frequencies and the output frequency may be split into two pairs of separate frequencies, and then we can perturbatively use two bilinear  $L^2$  bounds.
  - nonresonant, which heuristically may be eliminated using a normal form correction, which in our setting corresponds to quartic density and flux corrections for the mass and the momentum.
- (iii) High frequency terms with phase rotation symmetry, where we use the conservative assumption to construct a suitable flux correction modulo a perturbative part.

Our goal is to prove the bounds (3.1), (3.2), (3.3) and (3.7). In doing this, without any loss of generality we can make the bootstrap assumptions (3.14), (3.15), (3.16) and (3.17).

We consider the localized mass and momentum densities

$$M_\lambda = M_\lambda(u, \bar{u}),$$

$$P_\lambda^j = P_\lambda^j(u, \bar{u})$$

which satisfy the conservation laws

$$\partial_t M_\lambda(u) = \partial_j P_\lambda^j(u) + C_{\lambda,m}^{4,bal}(u) + C_{\lambda,m}^{4,nr}(u) + F_{\lambda,m}^4(u),$$

respectively

$$\partial_t P_\lambda^j(u) = \partial_x E_\lambda^{jk}(u) + C_{\lambda,p}^{4,bal}(u) + C_{\lambda,p}^{4,nr}(u) + F_{\lambda,p}^4(u),$$

where  $C^{4,bal}$  represents the contribution of balanced cubic terms with phase rotation symmetry and  $C^{4,nr}$  represent the contribution of balanced cubic nonresonant terms without the phase rotation symmetry. Here we remark that these terms are needed only at high frequency  $\lambda \gg 1$ , as at low frequency  $\lambda \ll 1$  we can place all balanced cubic contributions in  $F_\lambda^4$  due to the  $L^4$  bootstrap bound (3.17).

From Lemma 7.3 we have bounds for the source terms  $F_\lambda^4$ , namely

$$(10.1) \quad \|F_{\lambda,m}^4(u)\|_{L_{t,x}^1} \lesssim \epsilon^4 C^4 c_\lambda^4 \lambda^{-2s}, \quad \|F_{\lambda,p}^4(u)\|_{L_{t,x}^1} \lesssim \epsilon^4 C^4 c_\lambda^4 \lambda^{-2s+1}.$$

Unfortunately, the contributions of  $C_\lambda^{4,bal}$  and  $C_\lambda^{4,nr}$  cannot be also estimated directly in  $L^1$ , and will be treated separately.

*A. The  $C_\lambda^{4,bal}$  term.* Here we need to use the conservative assumption. This guarantees that the quartic forms  $C^{4,bal}$  have the additional structural property

$$(10.2) \quad c_{\lambda,m}^{4,bal}(\xi, \xi, \xi, \xi) = 0, \quad c_{\lambda,p}^{4,bal}(\xi, \xi, \xi, \xi) = 0, \quad \xi \in \mathbb{R}^2.$$

At the symbol level, this allows us to obtain a decomposition

$$\lambda^{-1} c_{\lambda,m}^{4,bal}(\xi_1, \xi_2, \xi_3, \xi_4) = (\xi_{odd} - \xi_{even}) q_{\lambda,m}^4(\xi_1, \xi_2, \xi_3, \xi_4) + \Delta^4 \xi r_{\lambda,m}^4(\xi_1, \xi_2, \xi_3, \xi_4),$$

where all symbols on the right are smooth and bounded. Separating variables in  $c_{\lambda,m}^{4,bal}$ , we arrive at the following

**Lemma 10.1.** *The quartic form  $C_m^{4,bal}$  admits a representation of the form*

$$(10.3) \quad \lambda^{-1} C_{\lambda,m}^{4,bal}(u, \bar{u}, u, \bar{u}) = \partial_x R_m^4(u, \bar{u}, u, \bar{u}) + \sum \partial_x Q_{j,a}^2(u, \bar{u}) Q_{j,b}^2(u, \bar{u}),$$

where the sum is rapidly convergent in  $j$ .

We remark that one corollary of this is the bound

$$(10.4) \quad \left| \int C_m^{4,bal}(u, \bar{u}, u, \bar{u}) dx dt \right| \lesssim \sum_j \| |D|^{\frac{1}{2}} Q_{j,a}^2(u, \bar{u}) \|_{L_{t,x}^2} \| |D|^{\frac{1}{2}} Q_{j,b}^2(u, \bar{u}) \|_{L_{t,x}^2},$$

where the right hand side may be estimated using the bilinear balanced bound (3.16). This suffices for energy estimates, but not for interaction Morawetz.

Instead, there we use a paraproduct decomposition for the terms in the series, writing

$$\begin{aligned} \partial_x Q_{j,a}^2(u, \bar{u}) Q_{j,b}^2(u, \bar{u}) &= T_{\partial_x Q_{j,a}^2(u, \bar{u})} Q_{j,b}^2(u, \bar{u}) + \partial_x Q_{j,a}^2(u, \bar{u}) Q_{j,b}^2(u, \bar{u}) - T_{\partial_x Q_{j,b}^2(u, \bar{u})} Q_{j,a}^2(u, \bar{u}) \\ &\quad + \partial_x T_{Q_{j,b}^2(u, \bar{u})} Q_{j,a}^2(u, \bar{u}). \end{aligned}$$

The first three terms can be estimated in  $L^1$  by the right hand side in (10.4), while the last term together with  $R_m^4$  go into a flux correction. Precisely, we have proved the following

**Proposition 10.2.** *Assume that  $c_{\lambda,m}^{4,bal}$  satisfies (10.2). Let  $u$  satisfy (3.14) and (3.16). Then there exists a decomposition*

$$(10.5) \quad C_{\lambda,m}^{4,bal}(u) = \partial_x Q_{\lambda,m}^{4,bal}(u) + F_{\lambda,m}^{4,bal}(u),$$

so that

$$(10.6) \quad \|Q_{\lambda,m}^{4,bal}(u)\|_{L_t^{\frac{6}{5}} L_x^{\frac{4}{3}}} \lesssim C^4 \epsilon^4 c_\lambda^4 \lambda^{2+\frac{5}{6}-4s},$$

while

$$(10.7) \quad \|F_{\lambda,m}^{4,bal}(u)\|_{L_{t,x}^1} \lesssim C^4 \epsilon^4 c_\lambda^4 \lambda^{3-4s}.$$

We note that in (10.6) we have in effect a full range of estimates between  $L_t^1 L_x^2$  and  $L_t^{\frac{3}{2}} L_x^1$ , of which the endpoints are not so useful to us, so we chose the middle. To prove it, we interpolate between the energy bound (3.14) and the balanced bilinear  $L^2$  bound (3.16).

The  $C^{4,nr}$  term. For this term we use a normal form type correction to both the densities and the fluxes:

**Proposition 10.3.** *Let  $u$  satisfy (3.14) and (3.16). Then there exists a decomposition*

$$(10.8) \quad C_{\lambda,m}^{4,nr}(u) = -\partial_t B_{\lambda,m}^{4,nr}(u) + \partial_x Q_{\lambda,m}^{4,nr}(u) + F_{\lambda,m}^{4,nr}(u),$$

so that we have the fixed time bound

$$(10.9) \quad \|B_{\lambda,m}^{4,nr}(u)\|_{L_x^1} \lesssim C^4 \epsilon^4 c_\lambda^4 \lambda^{2-4s},$$

and the space-time bounds

$$(10.10) \quad \|Q_{\lambda,m}^{4,nr}(u)\|_{L_t^{\frac{6}{5}} L_x^{\frac{4}{3}}} \lesssim C^4 \epsilon^4 c_\lambda^4 \lambda^{2+\frac{5}{6}-4s},$$

respectively

$$(10.11) \quad \|F_{\lambda,m}^{4,nr}(u)\|_{L_{t,x}^1} \lesssim C^4 \epsilon^4 c_\lambda^4 \lambda^{3-4s}.$$

*Proof.* The corrections are computed as if the linear part were the flat Schrödinger flow. This yields quartic forms with smooth symbols

$$B_{\lambda,m}^{4,bal}(u) = L(u_\lambda, u_\lambda, u_\lambda, u_\lambda), \quad Q_{\lambda,m}^{4,bal}(u) = \lambda L(u_\lambda, u_\lambda, u_\lambda, u_\lambda).$$

Then the bound (10.9) follows from the energy estimate and Bernstein's inequality, while (10.10) is exactly as in the previous Proposition. It remains to estimate the error term, which has the form

$$F_{\lambda,m}^{4,nr}(u) = \lambda^2 L(g_{<\lambda} - I, u_\lambda, u_\lambda, u_\lambda, u_\lambda) + L(N_\lambda(u), u_\lambda, u_\lambda, u_\lambda) := F_{\lambda,m}^{4,nr,1}(u) + F_{\lambda,m}^{4,nr,2}(u).$$

In the first term we can use two bilinear  $L^2$  bounds and two  $L^\infty$  bounds from energy and Bernstein's inequality,

$$\|F_{\lambda,m}^{4,nr,1}(u)\|_{L_{t,x}^1} \lesssim C^6 \epsilon^6 c_\lambda^4 \lambda^{-4s+n+1},$$

which suffices.

In the second term we separate  $N_\lambda = F_\lambda + C_\lambda$ , where for  $F_\lambda$  we use Proposition 7.2 and two  $L^\infty$  bounds from energy and Bernstein's inequality, while for  $C_\lambda$  we obtain

$$\lambda^2 L(u_\lambda, u_\lambda, u_\lambda, u_\lambda, u_\lambda, u_\lambda),$$

which can be estimated directly by combining (3.14) with (3.16) and Bernstein's inequality. This yields a frequency factor  $\lambda^{-6s+4}$ , which is more than sufficient.  $\square$

To summarize our findings, we have the modified densities

$$(10.12) \quad M_\lambda^\sharp(u) = M_\lambda(u) + B_{m,\lambda}^{4,nr}(u),$$

respectively

$$(10.13) \quad P_\lambda^\sharp(u) = P_\lambda(u) + B_{p,\lambda}^{4,nr}(u),$$

and the associated density-flux identities

$$(10.14) \quad \partial_t M_\lambda^\sharp(u) = \partial_j (P_\lambda^j(u) + Q_{\lambda,m}^{4,bal,j} + Q_{\lambda,m}^{4,nr,j}) + F_{\lambda,m}^{4,bal}(u) + F_{\lambda,m}^4(u),$$

respectively

$$(10.15) \quad \partial_t P_\lambda^{\sharp,j}(u) = \partial_x (E_\lambda^{jk}(u) + Q_{\lambda,p}^{4,bal,jk} + Q_{\lambda,p}^{4,nr,jk}) + F_{\lambda,m}^{4,bal,j}(u) + F_{p,\lambda}^{4,j}(u),$$

where the flux corrections  $Q_{\lambda,m}^{4,bal}$ ,  $Q_{\lambda,m}^{4,nr}$  are as in (10.6), respectively (10.10), and all the  $F^4$  terms are estimated in  $L^1$  as in (10.1), (10.7) and (10.9).

**10.1. The uniform energy bounds.** Here we use the bounds in Proposition 10.2 and Proposition 10.3 in order to prove the dyadic energy estimates (3.1) in our main result in Theorem 8, under the appropriate bootstrap assumptions as stated in Proposition 3.2.

**10.2. The interaction Morawetz identities.** Here we prove the bilinear  $L^2$  bounds (3.3) and (3.2), thereby concluding the proof of Theorem 8.

Overall, we will seek to pair a frequency  $\lambda$  portion of one solution  $u$  with a frequency  $\mu$  portion of another solution  $v$ , which will eventually be taken to be  $v = u^{x_0}$ , i.e. a translate of  $u$ . The argument follows the computations done earlier in Section 7, with the only difference that we need to add in the contribution of our density and flux corrections.

We will write here all identities in the general case. Then, in order to prove the global result in Theorem 8 we will specialize to three cases:

- (1) The diagonal case  $\lambda = \mu$ ,  $u = v$  (and thus  $x_0 = 0$ ).
- (2) The balanced shifted case  $\lambda \approx \mu$ , with  $x_0$  arbitrary.
- (3) The unbalanced case  $\mu < \lambda$ , with  $x_0$  arbitrary.

We define the interaction Morawetz functional for the functions  $(u, v)$  and the associated pair of frequencies  $(\lambda, \mu)$  as

$$(10.16) \quad \mathbf{I}_{\lambda\mu}(u, v) := \iint a_j(x - y) (M_\lambda^\sharp(u)(x) P_\mu^{\sharp,j}(v)(y) - P_\lambda^{\sharp,j}(u)(x) M_\mu^\sharp(v)(y)) dx dy.$$

We use the modified density-flux identities for the mass and for the momentum in order to compute the time derivative of  $\mathbf{I}_{\lambda\mu}$  as

$$(10.17) \quad \frac{d}{dt} \mathbf{I}_{\lambda\mu} = \mathbf{J}_{\lambda\mu}^4 + \mathbf{J}_{\lambda\mu}^6 + \mathbf{K}_{\lambda\mu}.$$

Compared to the computations done in the case of the paradifferential equation, all terms are similar except for  $\mathbf{J}_{\lambda\mu}^6$ , which contains the contribution of our density and flux corrections, namely it has the form

$$\begin{aligned} \mathbf{J}_{\lambda\mu}^6 := & \iint a_{jk}(x - y) \left( (Q_{\lambda,m}^{4,bal,k}(u) + Q_{\lambda,m}^{4,nr,k}(u))(x) P_\mu^{\sharp,j}(v)(y) - (Q_{\lambda,p}^{4,bal,jk}(u) + Q_{\lambda,p}^{4,nr,jk}(u))(x) M_\mu^\sharp(v)(y) \right. \\ & \left. - M_\lambda^\sharp(u)(x) (Q_{\mu,p}^{4,bal,jk}(v) + Q_{\mu,p}^{4,nr,jk}(v))(y) + P_\lambda^{\sharp,j}(u)(x) (Q_{\mu,m}^{4,bal,k}(v) + Q_{\mu,m}^{4,nr,k}(v))(y) \right) dx dy. \end{aligned}$$

To obtain the same outcome as in the paradifferential case, it will suffice to estimate the contribution of  $\mathbf{J}_{\lambda\mu}^{6,bal}$  perturbatively, i.e. to show that

$$(10.18) \quad \left| \int_0^T \mathbf{J}_{\lambda\mu}^{6,bal} dt \right| \lesssim \epsilon^4 c_\lambda^2 c_\mu^2 \lambda^{1-2s} \mu^{-2s}.$$

For simplicity we consider one of the second term expressions, which is the worst if  $\mu \lesssim \lambda$ . Since  $|a_{jk}| \leq \frac{1}{|x-y|}$ , we can use Young's inequality to estimate

$$\left| \int_0^T \iint Q_{\lambda,p}^{4,bal,jk}(u)(x) M_\mu^\sharp(v)(y) dx dy dt \right| \lesssim \|Q_{\lambda,p}^{4,bal,jk}(u)\|_{L_t^{\frac{6}{5}} L_x^{\frac{4}{3}}} \|M_\mu(v)\|_{L_t^6 L_x^{\frac{4}{3}}},$$

and similarly for  $Q_{\lambda,p}^{4,bal,jk}$ . For  $M_\mu^\sharp(v)$  we can interpolate the  $L_t^\infty L_x^1$  bound provided by (3.14) and the  $L_t^2 L_x^4$  bound derived from (3.16) to obtain

$$\|M_\mu^\sharp(v)\|_{L_t^6 L_x^{\frac{4}{3}}} \lesssim C^2 \epsilon^2 c_\mu^2 \mu^{\frac{1}{6}-2s},$$

where the quartic correction is better than the leading quadratic term.

Combining this with (10.6) or (10.10) we get

$$\lesssim C^6 \epsilon^6 c_\lambda^4 c_\mu^2 \lambda^{3+\frac{5}{6}-2s} \mu^{\frac{1}{6}-2s}$$

which is better than (10.18) provided that  $s > \frac{3}{2}$ . This concludes the proof of (10.18), and thus the proof of Theorem 8.

## REFERENCES

- [1] Thomas Alazard and Jean-Marc Delort. Global solutions and asymptotic behavior for two dimensional gravity water waves. *Ann. Sci. Éc. Norm. Supér. (4)*, 48(5):1149–1238, 2015.
- [2] Ioan Bejenaru and Daniel Tataru. Large data local solutions for the derivative NLS equation. *J. Eur. Math. Soc. (JEMS)*, 10(4):957–985, 2008.
- [3] Fernando Bernal-Vilchis, Nakao Hayashi, and Pavel I. Naumkin. Quadratic derivative nonlinear Schrödinger equations in two space dimensions. *NoDEA Nonlinear Differential Equations Appl.*, 18(3):329–355, 2011.
- [4] Jean Bourgain. A remark on normal forms and the “*I*-method” for periodic NLS. *J. Anal. Math.*, 94:125–157, 2004.
- [5] Jean-Yves Chemin and Delphine Salort. Wellposedness of some quasi-linear Schrödinger equations. *Sci. China Math.*, 58(5):891–914, 2015.
- [6] Hiroyuki Chihara. Local existence for semilinear Schrödinger equations. *Math. Japon.*, 42(1):35–51, 1995.
- [7] M. Colin. On the local well-posedness of quasilinear Schrödinger equations in arbitrary space dimension. *Comm. Partial Differential Equations*, 27(1-2):325–354, 2002.
- [8] M. Colin. On the local well-posedness of quasilinear Schrödinger equations in arbitrary space dimension. *Comm. Partial Differential Equations*, 27(1-2):325–354, 2002.
- [9] J. Colliander, M. Keel, G. Staffilani, H. Takaoka, and T. Tao. Almost conservation laws and global rough solutions to a nonlinear Schrödinger equation. *Math. Res. Lett.*, 9(5-6):659–682, 2002.
- [10] J. Colliander, M. Keel, G. Staffilani, H. Takaoka, and T. Tao. Global well-posedness and scattering for the energy-critical nonlinear Schrödinger equation in  $\mathbb{R}^3$ . *Ann. of Math. (2)*, 167(3):767–865, 2008.
- [11] Anne de Bouard, Nakao Hayashi, Pavel I. Naumkin, and Jean-Claude Saut. Scattering problem and asymptotics for a relativistic nonlinear Schrödinger equation. *Nonlinearity*, 12(5):1415–1425, 1999.
- [12] Jean-Marc Delort. Semiclassical microlocal normal forms and global solutions of modified one-dimensional KG equations. *Ann. Inst. Fourier (Grenoble)*, 66(4):1451–1528, 2016.
- [13] Benjamin Dodson. Global well-posedness and scattering for the mass critical nonlinear Schrödinger equation with mass below the mass of the ground state. *Adv. Math.*, 285:1589–1618, 2015.

- [14] Shin-ichi Doi. On the Cauchy problem for Schrödinger type equations and the regularity of solutions. *J. Math. Kyoto Univ.*, 34(2):319–328, 1994.
- [15] P. Germain, N. Masmoudi, and J. Shatah. Global solutions for 2D quadratic Schrödinger equations. *J. Math. Pures Appl. (9)*, 97(5):505–543, 2012.
- [16] J. Ginibre and N. Hayashi. Almost global existence of small solutions to quadratic nonlinear Schrödinger equations in three space dimensions. *Math. Z.*, 219(1):119–140, 1995.
- [17] Nakao Hayashi and Pavel I. Naumkin. Asymptotics for large time of solutions to the nonlinear Schrödinger and Hartree equations. *Amer. J. Math.*, 120(2):369–389, 1998.
- [18] Nakao Hayashi and Pavel I. Naumkin. Global existence of small solutions to the quadratic nonlinear Schrödinger equations in two space dimensions. *SIAM J. Math. Anal.*, 32(6):1390–1403, 2001.
- [19] Nakao Hayashi and Pavel I. Naumkin. Large time asymptotics for the fractional nonlinear Schrödinger equation. *Adv. Differential Equations*, 25(1-2):31–80, 2020.
- [20] Nakao Hayashi and Tohru Ozawa. Remarks on nonlinear Schrödinger equations in one space dimension. *Differential Integral Equations*, 7(2):453–461, 1994.
- [21] Sebastian Herr, Daniel Tataru, and Nikolay Tzvetkov. Global well-posedness of the energy-critical nonlinear Schrödinger equation with small initial data in  $H^1(\mathbb{T}^3)$ . *Duke Math. J.*, 159(2):329–349, 2011.
- [22] Jiaxi Huang, Ze Li, and Daniel Tataru. Global Regularity of Skew Mean Curvature Flow for Small Data in  $d \geq 4$  Dimensions. *Int. Math. Res. Not. IMRN*, (5):3748–3798, 2024.
- [23] Jiaxi Huang and Daniel Tataru. Local well-posedness of skew mean curvature flow for small data in  $d \geq 4$  dimensions. *Comm. Math. Phys.*, 389(3):1569–1645, 2022.
- [24] Jiaxi Huang and Daniel Tataru. Local well-posedness of the skew mean curvature flow for small data in  $d \geq 2$  dimensions. *Arch. Ration. Mech. Anal.*, 248(1):Paper No. 10, 79, 2024.
- [25] John K. Hunter and Mihaela Ifrim. Enhanced life span of smooth solutions of a Burgers-Hilbert equation. *SIAM J. Math. Anal.*, 44(3):2039–2052, 2012.
- [26] John K. Hunter, Mihaela Ifrim, Daniel Tataru, and Tak Kwong Wong. Long time solutions for a Burgers-Hilbert equation via a modified energy method. *Proc. Amer. Math. Soc.*, 143(8):3407–3412, 2015.
- [27] Wataru Ichinose. On  $L^2$  well posedness of the Cauchy problem for Schrödinger type equations on the Riemannian manifold and the Maslov theory. *Duke Math. J.*, 56(3):549–588, 1988.
- [28] Mihaela Ifrim and Daniel Tataru. Global bounds for the cubic nonlinear Schrödinger equation (NLS) in one space dimension. *Nonlinearity*, 28(8):2661–2675, 2015.
- [29] Mihaela Ifrim and Daniel Tataru. Two dimensional water waves in holomorphic coordinates II: Global solutions. *Bull. Soc. Math. France*, 144(2):369–394, 2016.
- [30] Mihaela Ifrim and Daniel Tataru. The lifespan of small data solutions in two dimensional capillary water waves. *Arch. Ration. Mech. Anal.*, 225(3):1279–1346, 2017.
- [31] Mihaela Ifrim and Daniel Tataru. Well-posedness and dispersive decay of small data solutions for the Benjamin-Ono equation. *Ann. Sci. Éc. Norm. Supér. (4)*, 52(2):297–335, 2019.
- [32] Mihaela Ifrim and Daniel Tataru. Local well-posedness for quasilinear problems: a primer. *arXiv e-prints*, page arXiv:2008.05684, August 2020.
- [33] Mihaela Ifrim and Daniel Tataru. Global solutions for 1D cubic defocusing dispersive equations: Part I. *arXiv e-prints*, page arXiv:2205.12212, May 2022.
- [34] Mihaela Ifrim and Daniel Tataru. Long time solutions for 1D cubic dispersive equations, Part II: the focusing case. *arXiv e-prints*, page arXiv:2210.17007, October 2022.
- [35] Mihaela Ifrim and Daniel Tataru. Testing by wave packets and modified scattering in nonlinear dispersive pde’s. *arXiv e-prints*, page arXiv:2204.13285, April 2022.
- [36] Mihaela Ifrim and Daniel Tataru. Global solutions for 1d cubic dispersive equations, part iii: the quasilinear schrödinger flow, 2023.
- [37] Mihaela Ifrim and Daniel Tataru. The global well-posedness conjecture for 1D cubic dispersive equations. *arXiv e-prints*, page arXiv:2311.15076, November 2023.
- [38] In-Jee Jeong and Sung-Jin Oh. Wellposedness of the electron MHD without resistivity for large perturbations of the uniform magnetic field. *arXiv e-prints*, page arXiv:2402.06278, February 2024.
- [39] Jun Kato and Fabio Pusateri. A new proof of long-range scattering for critical nonlinear Schrödinger equations. *Differential Integral Equations*, 24(9-10):923–940, 2011.
- [40] Markus Keel and Terence Tao. Endpoint Strichartz estimates. *Amer. J. Math.*, 120(5):955–980, 1998.

- [41] C. E. Kenig, G. Ponce, C. Rolvung, and L. Vega. Variable coefficient Schrödinger flows for ultrahyperbolic operators. *Adv. Math.*, 196(2):373–486, 2005.
- [42] Carlos E. Kenig, Gustavo Ponce, Christian Rolvung, and Luis Vega. The general quasilinear ultrahyperbolic Schrödinger equation. *Adv. Math.*, 206(2):402–433, 2006.
- [43] Carlos E. Kenig, Gustavo Ponce, and Luis Vega. Small solutions to nonlinear Schrödinger equations. *Ann. Inst. H. Poincaré C Anal. Non Linéaire*, 10(3):255–288, 1993.
- [44] Carlos E. Kenig, Gustavo Ponce, and Luis Vega. Smoothing effects and local existence theory for the generalized nonlinear Schrödinger equations. *Invent. Math.*, 134(3):489–545, 1998.
- [45] Carlos E. Kenig, Gustavo Ponce, and Luis Vega. The Cauchy problem for quasi-linear Schrödinger equations. *Invent. Math.*, 158(2):343–388, 2004.
- [46] Herbert Koch and Daniel Tataru. Energy and local energy bounds for the 1-d cubic NLS equation in  $H^{-\frac{1}{4}}$ . *Ann. Inst. H. Poincaré Anal. Non Linéaire*, 29(6):955–988, 2012.
- [47] Wee Keong Lim and Gustavo Ponce. On the initial value problem for the one dimensional quasi-linear Schrödinger equations. *SIAM J. Math. Anal.*, 34(2):435–459, 2002.
- [48] Hans Lindblad, Jonas Lührmann, and Avy Soffer. Asymptotics for 1D Klein-Gordon equations with variable coefficient quadratic nonlinearities. *Arch. Ration. Mech. Anal.*, 241(3):1459–1527, 2021.
- [49] Hans Lindblad and Avy Soffer. Scattering and small data completeness for the critical nonlinear Schrödinger equation. *Nonlinearity*, 19(2):345–353, 2006.
- [50] Jeremy L. Marzuola, Jason Metcalfe, and Daniel Tataru. Quasilinear Schrödinger equations I: Small data and quadratic interactions. *Adv. Math.*, 231(2):1151–1172, 2012.
- [51] Jeremy L. Marzuola, Jason Metcalfe, and Daniel Tataru. Quasilinear Schrödinger equations, II: Small data and cubic nonlinearities. *Kyoto J. Math.*, 54(3):529–546, 2014.
- [52] Jeremy L. Marzuola, Jason Metcalfe, and Daniel Tataru. Quasilinear Schrödinger equations III: Large data and short time. *Arch. Ration. Mech. Anal.*, 242(2):1119–1175, 2021.
- [53] Sigeru Mizohata. *On the Cauchy problem*, volume 3 of *Notes and Reports in Mathematics in Science and Engineering*. Academic Press, Inc., Orlando, FL; Science Press Beijing, Beijing, 1985.
- [54] Ben Pineau and Mitchell A. Taylor. Low regularity solutions for the general Quasilinear ultrahyperbolic Schrödinger equation. *arXiv e-prints*, page arXiv:2310.19221, October 2023.
- [55] Fabrice Planchon and Luis Vega. Bilinear virial identities and applications. *Ann. Sci. Éc. Norm. Supér. (4)*, 42(2):261–290, 2009.
- [56] Markus Poppenberg. On the local well posedness of quasilinear Schrödinger equations in arbitrary space dimension. *J. Differential Equations*, 172(1):83–115, 2001.
- [57] Jie Shao and Yi Zhou. Local wellposedness for the quasilinear Schrödinger equations via the generalized energy method. *arXiv e-prints*, page arXiv:2311.02556, November 2023.
- [58] Jalal Shatah. Normal forms and quadratic nonlinear Klein-Gordon equations. *Comm. Pure Appl. Math.*, 38(5):685–696, 1985.
- [59] Jiro Takeuchi. Le problème de Cauchy pour certaines équations aux dérivées partielles du type de Schrödinger. VIII. Symétrisations indépendantes du temps. *C. R. Acad. Sci. Paris Sér. I Math.*, 315(10):1055–1058, 1992.
- [60] Terence Tao. Global regularity of wave maps. II. Small energy in two dimensions. *Comm. Math. Phys.*, 224(2):443–544, 2001.
- [61] Terence Tao. Global well-posedness of the Benjamin-Ono equation in  $H^1(\mathbf{R})$ . *J. Hyperbolic Differ. Equ.*, 1(1):27–49, 2004.
- [62] Daniel Tataru. On global existence and scattering for the wave maps equation, unpublished original version, 1999.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF WISCONSIN, MADISON  
 Email address: ifrim@wisc.edu

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA AT BERKELEY  
 Email address: tataru@math.berkeley.edu