

Strong-weak symmetry and quantum modularity of resurgent topological strings on local \mathbb{P}^2

Veronica Fantini¹ and Claudia Rella²

¹*IHÉS, 91440 Bures-sur-Yvette, France*

²*Département de Physique Théorique, Université de Genève,
CH-1211 Genève, Switzerland*

Email: fantini@ihes.fr, claudia.rella@unige.ch

Abstract

Quantizing the mirror curve to a toric Calabi–Yau threefold gives rise to quantum operators whose fermionic spectral traces produce factorially divergent formal power series in the Planck constant and its inverse. These are conjecturally captured by the Nekrasov–Shatashvili and standard topological string free energies, respectively, via the TS/ST correspondence. The resurgent structures of the first fermionic spectral trace of local \mathbb{P}^2 in both weak and strong coupling limits were solved exactly by the second author in [1]. Here, we take the perspective of the Stokes constants and their generating functions. We prove that a full-fledged *strong-weak resurgent symmetry* is at play, exchanging the perturbative/nonperturbative contributions to the holomorphic and anti-holomorphic blocks in the factorization of the spectral trace. This relies on a global net of relations connecting the perturbative series and the discontinuities in the dual regimes, which is built upon the analytic properties of the L -functions with coefficients given by the Stokes constants and the q -series acting as their generating functions. Then, we show that the latter are holomorphic quantum modular forms for $\Gamma_1(3)$ and are reconstructed by the median resummation of their asymptotic expansions.

Contents

1	Introduction	2
2	Resurgence of the spectral trace of local \mathbb{P}^2	4
2.1	Geometric setup and the fermionic spectral traces	4
2.2	Resurgent structures in the weak and strong coupling regimes	8
2.3	Exact solutions for the spectral trace	11
3	Resurgent L-functions	16
3.1	Generating functions of the Stokes constants	16
3.2	Strong-weak resurgent symmetry	18
3.3	Functional equation from a unified perspective	23
3.4	Arithmetic twist of the Stokes constants	24

4	Summability and quantum modularity	27
4.1	Borel–Laplace sums	27
4.2	Median resummation	30
4.3	Quantum modularity of the generating functions	36
5	Conclusions	40
A	Quantum dilogarithms	42
B	Basics of resurgence	44
	References	47

1 Introduction

The A-model topological string theory compactified on a local Calabi–Yau (CY) threefold X is defined perturbatively by a worldsheet genus expansion in the string coupling constant g_s and can be expressed in terms of the enumerative invariants of the background geometry. The fixed-genus topological string free energies $F_g(\mathbf{t})$, $g \geq 0$, are well-defined expansions in a common region of convergence near the large radius limit $\Re(t_i) \gg 1$ in the moduli space of Kähler structures of X . Yet, at a fixed point \mathbf{t} in the convergence region, $F_g(\mathbf{t})$ diverges factorially as $(2g)!$ [2], signaling the presence of exponentially small corrections in g_s .

A non-perturbative definition of the topological string on X has been proposed in [3, 4]. By mirror symmetry, the B-model topological string theory on the mirror \hat{X} is described by the theory of deformations of its complex structures [5, 6]. This is encoded in algebraic equations parametrizing a family of Riemann surfaces Σ of genus g_Σ embedded in $\mathbb{C}^* \times \mathbb{C}^*$, whose quantization leads naturally to quantum operators acting on the real line. A Planck constant $\hbar \in \mathbb{R}_{>0}$ is introduced as a quantum deformation parameter. The conjectural statement of [3, 4], known as Topological String/Spectral Theory (TS/ST) correspondence, implies a strong-weak coupling duality $\hbar \propto g_s^{-1}$ and leads to exact formulae for the fermionic spectral traces $Z_X(\mathbf{N}, \hbar)$, $\mathbf{N} \in \mathbb{N}^{g_\Sigma}$, of these quantum-mechanical operators. These are well-defined functions of \hbar and are expressed in terms of the total grand potential of the topological string on X . In particular, the A-model total free energies of both the conventional and Nekrasov–Shatashvili (NS) topological strings on X , that is, the generating functions of the Gromov–Witten and refined BPS invariants, are needed to define the total grand potential and can be regarded as non-perturbative corrections of one another in the appropriate regimes [7, 8]. Thus, the TS/ST correspondence provides a way to access the non-perturbative effects associated with the factorial divergence of the topological string perturbation series in the spirit of large- N gauge/string dualities. After the original proposal of [5, 9, 10], growing evidence indicates that the theory of resurgence [11] can be applied to obtain a systematic understanding of the hidden non-perturbative sectors of topological string theory.

The machinery of resurgence uniquely associates a divergent formal power series with a collection of non-analytic, exponential-type corrections paired with a non-trivial set of complex numbers, known as Stokes constants, which capture information about the large-order behaviour of the original asymptotic series and its additional non-perturbative sectors in a mathematically precise way. See, e.g., [12–14] for reviews on resurgence and [2, 15] for its application to gauge and string theories. For instance, following the work of [16, 17], the resurgent structure of the

topological string has recently been investigated by means of an operator formulation of the BCOV holomorphic anomaly equations [18, 19] satisfied by the conventional closed topological string free energies on toric CY threefolds [20], which has later been extended to arbitrary CY threefolds [21, 22] and Walcher’s real topological string [23]. These works produced formal solutions for the multi-instanton trans-series extensions of the free energies that are conjecturally related to the resurgent structure of their perturbative genus expansion in a specific way, although a determination of which of the possible Borel singularities are realized and the values of their Stokes constants are missing. The same methods have been applied to the refined topological string and its NS limit on toric CY threefolds [24–27]. The Stokes constants in these resurgent structures are generally unknown, and only a few have been computed numerically. Yet, growing evidence supports a conjectural identification between them and enumerative invariants counting BPS states of the topological string. The case of the resolved conifold has been studied in [28–31].

Simultaneously, the alternative approach of [1, 32], which we follow and promote in this paper, investigates the resurgent structure of the topological string on a toric CY threefold X via the factorially divergent numerical power series obtained by asymptotically expanding the fermionic spectral traces $Z_X(\mathbf{N}, \hbar)$ at fixed \mathbf{N} in the limits of $\hbar \rightarrow 0$ and $\hbar \rightarrow \infty$. Crucially, this leads, in particular, to peacock patterns of singularities in the Borel plane and infinite sets of generally calculable and rational Stokes constants. By means of the TS/ST correspondence, the dual \hbar -regimes of the fermionic spectral traces above are explicitly related to the free energies of the NS and conventional topological string theories on X , respectively.

The power of studying the resurgence of the fermionic spectral traces is manifest in the work of the second author on the local \mathbb{P}^2 geometry [1], which is the benchmark example among all non-compact CY threefolds. Indeed, the full resurgent structures of the logarithm of the first fermionic spectral trace of local \mathbb{P}^2 in both weak and strong limits in \hbar have been exactly solved, leading to proven analytic formulae for the Stokes constants. These have a transparent and strikingly simple arithmetic meaning as divisor sum functions, while their generating series are known in closed form in terms of q -Pochhammer symbols. In addition, the Stokes constants are the coefficients of two explicit L -functions, whose analytic number-theoretic properties underlie a global net of relations connecting the resurgent structures at weak and strong coupling. These results are the starting ground for the investigation performed in this paper. The arithmetic duality discovered in [1] is completed and upgraded into a beautiful full-fledged symmetry. We call it the *strong-weak resurgent symmetry*. In a nutshell, it is an exact symmetry between the resurgent structures in the strong and weak coupling regimes building upon the interplay of q -series and L -functions and revolving around the central role played by the Stokes constants. This newly found symmetry acts at the level of the perturbative/non perturbative contributions to the holomorphic and anti-holomorphic blocks in the factorization of the spectral trace in a precise way. As a side effect, we shed some light on one of the formal dissimilarities between topological strings and complex Chern–Simons theory pointed out in [32].

The richness of the resurgence of the spectral trace of local \mathbb{P}^2 leads us to investigate further the properties of the generating functions of the weak and strong coupling Stokes constants, which we denote by f_0 and f_∞ , respectively. On the one hand, we prove that f_0 and f_∞ and their images under Fricke involution are *holomorphic quantum modular forms* of weight zero for the congruence subgroup $\Gamma_1(3) \subset \mathrm{SL}_2(\mathbb{Z})$ (see Theorems 4.7 and 4.8). Notice that the group $\Gamma_1(3)$ is deeply related to the geometry of the problem since the moduli space parametrizing the complex structures of the mirror of local \mathbb{P}^2 is the compactification of the quotient $\mathbb{H}/\Gamma_1(3)$, where \mathbb{H} denotes the upper half of the complex plane. Moreover, the generating functions of

Gromov–Witten invariants of local \mathbb{P}^2 are known to be quasi modular forms for $\Gamma_1(3)$ [33–35]. On the other hand, we study the summability properties of the asymptotic expansions of f_0 and f_∞ , which are explicitly governed by the perturbative series of the logarithm of the spectral trace in the two regimes in \hbar . We prove that the asymptotic expansion of f_0 is resumable and specifically reconstructs the generating function by means of the *median resummation* (see Theorem 4.6). A similar result is only conjectured for f_∞ (see Conjecture 1) on the basis of numerical evidence.

In the companion paper [36], we lift the paramount features of the spectral trace of local \mathbb{P}^2 to a generic paradigm of *modular resurgence*, which revolves around the functional equations satisfied by the resurgent L -functions. This leads us to present and discuss new perspectives on the resurgence of divergent formal power series and quantum modularity and to introduce the notion of a *modular resurgent structure*.

This paper is organized as follows. In Section 2, we review the basic ingredients in the construction of the fermionic spectral traces of local \mathbb{P}^2 and their conjectural resurgent structures in the weak and strong coupling regimes in \hbar . We then focus on the logarithm of the first fermionic spectral trace and revise the exact results obtained by the second author in [1]. In Section 3, we delve into the analytic number-theoretic characterization of the dual resurgent structures discovered in [1] and complete it into a full-fledged strong-weak resurgent symmetry. As we take the perspective of the Stokes constants, their generating functions, and their L -functions, we find new exact relations intertwining the dual regimes in \hbar . In Section 4, we study the quantum modularity properties of the generating functions of the Stokes constants and the summability of their asymptotic expansions. Finally, in Section 5, we conclude and mention further perspectives to be addressed in future works. There are two Appendices.

2 Resurgence of the spectral trace of local \mathbb{P}^2

In this section, we revise the construction of the fermionic spectral traces from the quantization of the mirror curve of the local \mathbb{P}^2 geometry and their connection to the topological string theory compactified on the same background via the conjectural TS/ST correspondence [3, 4]. We refer to [37] for a thorough overview. We describe the recent proposal of [1, 32] on the resurgence of the asymptotic series obtained from the weak and strong coupling expansions of the fermionic spectral traces. Finally, we summarize the exact results of [1] on the dual resurgent structures of the first fermionic spectral trace and their startling number-theoretic properties.

2.1 Geometric setup and the fermionic spectral traces

The local \mathbb{P}^2 geometry is the simplest example of a toric del Pezzo Calabi–Yau (CY) threefold.¹ It is defined as the total space of the canonical line bundle on the projective surface \mathbb{P}^2 , that is,

$$X = \mathcal{O}(-3) \rightarrow \mathbb{P}^2. \quad (1)$$

Mirror symmetry pairs X with a mirror CY threefold \hat{X} in such a way that the theory of variations of complex structures of the mirror \hat{X} is encoded in the one-parameter family of algebraic equations

$$O_{\mathbb{P}^2}(x, y) + \kappa = e^x + e^y + e^{-x-y} + \kappa = 0, \quad x, y \in \mathbb{C}, \quad (2)$$

¹A toric del Pezzo CY threefold is defined as the total space of the canonical line bundle on a toric del Pezzo surface. Examples of toric del Pezzo surfaces are the projective surface \mathbb{P}^2 , the Hirzebruch surfaces \mathbb{F}_n for $n = 0, 1, 2$, and the blowups of \mathbb{P}^2 at n points, usually denoted by \mathcal{B}_n , for $n = 1, 2, 3$.

where κ is the complex deformation parameter of \hat{X} . Note that there are no mass parameters. Eq. (2) describes a genus-one Riemann surface Σ embedded in $\mathbb{C}^* \times \mathbb{C}^*$, which is known as the mirror curve of X and determines the B-model topological string theory on \hat{X} [5, 6].

Remark 2.1. *Toric del Pezzo surfaces are classified by reflexive polyhedra in two dimensions. The polyhedron $\Delta_{\mathbb{P}^2}$ associated with the projective surface \mathbb{P}^2 is the convex hull of the origin together with the two-dimensional vectors*

$$\nu^{(1)} = (1, 0), \quad \nu^{(2)} = (0, 1), \quad \nu^{(3)} = (-1, -1), \quad (3)$$

as shown in Fig. 1. Note that the function $O_{\mathbb{P}^2}(x, y)$ in Eq. (2) can be written as

$$O_{\mathbb{P}^2}(x, y) = \sum_{i=1}^3 \exp\left(\nu_1^{(i)} x + \nu_2^{(i)} y\right), \quad (4)$$

where $\nu^{(i)} = (\nu_1^{(i)}, \nu_2^{(i)})$, $i = 1, 2, 3$.




Figure 1: Toric diagram of the local \mathbb{P}^2 geometry. We show the vectors in Eq. (3) and the polyhedron $\Delta_{\mathbb{P}^2}$. The complex modulus κ corresponds to the internal vertex of the diagram.

The Batyrev coordinate $z = 1/\kappa^3$ is related to the Kähler parameter of X via the classical mirror map

$$-t(z) = \log z + \tilde{S}(z), \quad (5)$$

where $\tilde{S}(z)$ is a power series in z with finite radius of convergence. Following a choice of symplectic basis $\{A, B\}$ of one-cycles on the spectral curve Σ , the classical periods of the meromorphic differential one-form $\lambda = y(x)dx$, where the function $y(x)$ is locally defined by Eq. (2), satisfy

$$t(z) \propto \oint_A \lambda, \quad \partial_t F_0(z) \propto \oint_B \lambda, \quad (6)$$

where the function $F_0(z)$ is the classical prepotential of local \mathbb{P}^2 [38], representing the genus-zero amplitude of the B-model topological string on \hat{X} – or, equivalently, the generating function of the genus-zero Gromov–Witten invariants of X convoluted with the mirror map. In the parametrization of the moduli space given by the Batyrev coordinate, the Picard–Fuchs differential equation associated with the mirror of local \mathbb{P}^2 is

$$\mathcal{L}_z \Pi = (\Theta^3 - 3z(3\Theta + 1)(3\Theta + 2)\Theta) \Pi = 0, \quad (7)$$

where $\Theta = zd/dz$ and Π is the full period vector of λ . The Picard–Fuchs differential operator \mathcal{L}_z has three singular points, which are the large radius point at $z = 0$, the conifold point at $z = -1/27$, and the orbifold point at $1/z = 0$. In addition, by looking at the monodromy data of the Picard–Fuchs equation, the moduli space of complex structures on \hat{X} can be identified with the compactified quotient $\overline{\mathbb{H}}/\Gamma_1(3)$, where \mathbb{H} denotes the upper half-plane and $\Gamma_1(3)$ is the congruence subgroup

$$\Gamma_1(3) := \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \mathrm{SL}_2(\mathbb{Z}) \mid a, d \equiv_3 1, c \equiv_3 0 \right\}, \quad (8)$$

which is generated by the elements

$$T = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}, \quad \gamma_3 = \begin{pmatrix} 1 & 0 \\ 3 & 1 \end{pmatrix}. \quad (9)$$

We refer to [34, Section 10.1] for a detailed discussion.

The canonical Weyl quantization scheme of [3, 4] applied to the function $O_{\mathbb{P}^2}(x, y)$ in Eq. (2), together with an appropriate choice of reality conditions for the variables $x, y \in \mathbb{C}$, produces the Hermitian quantum operator²

$$\mathbf{O}_{\mathbb{P}^2}(\mathbf{x}, \mathbf{y}) = e^{\mathbf{x}} + e^{\mathbf{y}} + e^{-\mathbf{x}-\mathbf{y}}, \quad (10)$$

acting on $L^2(\mathbb{R})$, where \mathbf{x}, \mathbf{y} are self-adjoint Heisenberg operators on the real line satisfying the commutation relation $[\mathbf{x}, \mathbf{y}] = i\hbar$. A Planck constant $\hbar \in \mathbb{R}_{>0}$ is introduced as a quantum deformation parameter. Consequently, the classical mirror map $t(z)$ in Eq. (5) is promoted to the quantum mirror map

$$-t(z, \hbar) = \log z + \tilde{S}(z, \hbar), \quad (11)$$

which reproduces the conventional mirror map in the semiclassical limit $\hbar \rightarrow 0$ and is determined as an A-period of a quantum-corrected version of the differential λ obtained via the all-orders perturbative WKB approximation [40, 41]. It was conjectured in [3, 4] and later proven in [42] that the inverse operator

$$\rho_{\mathbb{P}^2} = \mathbf{O}_{\mathbb{P}^2}^{-1} \quad (12)$$

is positive-definite and of trace class. Its spectral (or Fredholm) determinant

$$\Xi_{\mathbb{P}^2}(\kappa, \hbar) = \det(1 + \kappa \rho_{\mathbb{P}^2}) \quad (13)$$

is an entire function of κ [4, 43], while its fermionic spectral traces $Z(N, \hbar)$, where N is a non-negative integer, are well-defined functions of $\hbar \in \mathbb{R}_{>0}$ and can be computed explicitly. In particular, they appear in the convergent power series expansion of the analytically continued spectral determinant around the orbifold point $\kappa = 0$, that is,

$$\Xi_{\mathbb{P}^2}(\kappa, \hbar) = 1 + \sum_{N=1}^{\infty} Z(N, \hbar) \kappa^N. \quad (14)$$

Besides, classical results in Fredholm theory provide explicit determinant expressions for the fermionic spectral traces, which can be regarded as multi-cut matrix model integrals [44, 45]. In particular, we have the multi-dimensional integral representation

$$Z(N, \hbar) = \frac{1}{N!} \sum_{\sigma \in S_N} (-1)^{\epsilon(\sigma)} \int \prod_{i=1}^N \rho_{\mathbb{P}^2}(x_i, x_{\sigma(i)}) d^N x = \frac{1}{N!} \int \det(\rho_{\mathbb{P}^2}(x_i, x_j)) d^N x, \quad (15)$$

²The spectral theory of $\mathbf{O}_{\mathbb{P}^2}$ has been studied numerically in [39].

where \mathcal{S}_N is the N -th permutation group, $\epsilon(\sigma)$ is the signature of the element $\sigma \in \mathcal{S}_N$, and $\rho_{\mathbb{P}^2}(x_i, x_j)$ denotes the integral kernel of the operator $\rho_{\mathbb{P}^2}$. This is known explicitly as

$$\rho_{\mathbb{P}^2}(x, y) = \frac{e^{\pi \mathbf{b}(x+y)/3}}{2\mathbf{b} \cosh(\pi(x-y)/\mathbf{b} + i\pi/6)} \frac{\Phi_{\mathbf{b}}(y + i\mathbf{b}/3)}{\Phi_{\mathbf{b}}(x - i\mathbf{b}/3)}, \quad (16)$$

where \mathbf{b} is related to \hbar by

$$2\pi \mathbf{b}^2 = 3\hbar \quad (17)$$

and $\Phi_{\mathbf{b}}$ denotes the Faddeev's quantum dilogarithm (see Appendix A for its definition and some of its fundamental properties). Note that the integral kernel in Eq. (16) is well-defined for

$$\hbar \in \mathbb{C}' = \mathbb{C} \setminus \mathbb{R}_{\leq 0}, \quad (18)$$

since $\Phi_{\mathbf{b}}$ can be analytically continued to all values of \mathbf{b} such that $\mathbf{b}^2 \notin \mathbb{R}_{\leq 0}$. As a consequence, the same holds for the fermionic spectral traces in Eq. (15).

The conjectural statement of the TS/ST correspondence of [3, 4], which is now supported by an increasing amount of evidence obtained in applications to concrete examples, implies an exact formula for the fermionic spectral traces of local \mathbb{P}^2 in terms of the topological string amplitudes on the same background. Namely, we have that³

$$Z(N, \hbar) = \frac{1}{2\pi i} \int_{-i\infty}^{i\infty} d\mu \exp(J(\mu, \hbar) - \mu N), \quad N \in \mathbb{N}, \quad (19)$$

where the integration contour along the imaginary axis can be suitably deformed to guarantee convergence, the chemical potential μ is related to the complex modulus by $\kappa = e^\mu$, and $J(\mu, \hbar)$ is the total grand potential of topological string theory on X . We recall that this is given by the sum

$$J(\mu, \hbar) = J^{\text{WS}}(\mu, \hbar) + J^{\text{WKB}}(\mu, \hbar), \quad (20)$$

where the worldsheet component $J^{\text{WS}}(\mu, \hbar)$ encodes the non-perturbative corrections in \hbar due to complex instantons, while the WKB component $J^{\text{WKB}}(\mu, \hbar)$ contains the perturbative contributions in \hbar to the quantum-mechanical spectral problem in the mirror B-model [47]. Precisely, $J^{\text{WS}}(\mu, \hbar)$ can be expressed in terms of the all-genus generating functional of Gopakumar–Vafa invariants of X [48, 49], which is captured by the total free energy of the A-model standard topological string, while $J^{\text{WKB}}(\mu, \hbar)$ is a function of the all-order generating functional of refined BPS invariants of X [50–52], which is captured by the total free energy of the A-model Nekrasov–Shatashvili (NS) topological string⁴. As we will discuss in Section 2.2, only one of the two components of the total grand potential above contributes to the each of the dual perturbative regimes of $\hbar \rightarrow 0$ and $\hbar \rightarrow \infty$. Moreover, and importantly, the TS/ST correspondence implies the identification

$$g_s = \frac{4\pi^2}{\hbar}, \quad (21)$$

where g_s is the conventional topological string coupling constant. The statement above can be interpreted as a strong-weak coupling duality between the spectral theory of the operator $\rho_{\mathbb{P}^2}$ arising from the quantization of the mirror curve Σ and the standard topological string theory on X . We will return to this in Section 3.2.

³Although being initially defined for non-negative integer values of N , the Airy-type integral in Eq. (19) naturally extends the fermionic spectral traces to entire functions of $N \in \mathbb{C}$ [46].

⁴The standard and NS topological string partition functions can be engineered as two one-parameter specializations of the partition function of the refined topological string theory compactified on a toric CY background [53].

Remark 2.2. In its original formulation, the TS/ST correspondence of [3, 4], and therefore also the statement in Eq. (19), applies to $\hbar \in \mathbb{R}_{>0}$. The issue of the complexification of \hbar , or, equivalently, of g_s , in the context of the TS/ST correspondence has been addressed in various studies since [54–57]. As we mentioned above, the integral kernel of local \mathbb{P}^2 in Eq. (16) admits a well-defined analytic continuation to $\hbar \in \mathbb{C}'$. Moreover, the analytically continued fermionic spectral traces obtained in this way match the natural analytic continuation of the total grand potential of local \mathbb{P}^2 to complex values of \hbar such that $\Re(\hbar) > 0$. The TS/ST correspondence is, then, still applicable.

2.2 Resurgent structures in the weak and strong coupling regimes

In light of the strong-weak duality between the topological string coupling g_s and the quantum deformation parameter \hbar in Eq. (21), the regimes of $\hbar \rightarrow 0$ and $\hbar \rightarrow \infty$ of the spectral theory correspond to the strong and weak coupling limits of the dual topological string theory on local \mathbb{P}^2 , respectively. Besides, due to the functional forms of the worldsheet and WKB grand potentials appearing in Eq. (20), there are appropriate scaling regimes in the coupling constants in which only one of the two components effectively contributes to the total grand potential. In the standard double-scaling limit [58–60]

$$\hbar \rightarrow \infty, \quad \mu \rightarrow \infty, \quad \frac{\mu}{\hbar} = \zeta \text{ fixed}, \quad (22)$$

the quantum mirror map $t(z, \hbar)$ in Eq. (11) becomes trivial, and the total grand potential in Eq. (20) reduces to its worldsheet component, which has the all-genus asymptotic expansion

$$J^{\text{WS}}(\zeta, \hbar) \sim \sum_{g=0}^{\infty} J_g(\zeta) \hbar^{2-2g}, \quad (23)$$

where $J_g(\zeta)$ is essentially the genus g free energy of the conventional topological string in the large radius limit $\Re(t) \gg 1$ of the moduli space of X after the so-called B-field has been turned on. Similarly, in the dual semiclassical regime

$$\hbar \rightarrow 0, \quad \mu \text{ fixed}, \quad (24)$$

the quantum mirror map $t(z, \hbar)$ in Eq. (11) becomes classical by construction, and the total grand potential in Eq. (20) retains only the WKB contribution, which can be formally expanded in powers of \hbar as

$$J^{\text{WKB}}(\mu, \hbar) \sim \sum_{n=0}^{\infty} J_n^{\text{NS}}(\mu) \hbar^{2n-1}, \quad (25)$$

where the fixed-order WKB grand potential $J_n^{\text{NS}}(\mu)$ is a function of the order n free energy of the NS topological string at large radius and its derivative in the Kähler parameter t . Evidence suggests that both asymptotic series in Eqs. (23) and (25) diverge factorially [47]. Thus, their non-perturbative completions will contain exponentially small effects in \hbar and $1/\hbar$, respectively. See the discussion in Section 3.2.

As a consequence of the TS/ST correspondence, there are related 't Hooft limits for the fermionic spectral traces that extract the perturbative all-genus and all-order expansions of the conventional and NS topological strings on local \mathbb{P}^2 , respectively. In the standard 't Hooft limit associated with Eq. (22), that is,

$$\hbar \rightarrow \infty, \quad N \rightarrow \infty, \quad \frac{N}{\hbar} = \lambda \text{ fixed}, \quad (26)$$

the saddle-point evaluation of the integral in Eq. (19) represents a symplectic transformation from the large radius point in moduli space to the so-called maximal conifold point⁵ [4, 43]. Specifically, it follows from the geometric formalism of [33] that the 't Hooft parameter λ is the flat coordinate corresponding to the maximal conifold frame of local \mathbb{P}^2 . In the recent work of [1], a dual WKB double-scaling regime associated with Eq. (24) is studied, that is,

$$\hbar \rightarrow 0, \quad N \rightarrow \infty, \quad N\hbar = \sigma \text{ fixed}, \quad (27)$$

so that the symplectic transformation of the total grand potential at large radius encoded in the integral in Eq. (19) can be interpreted again as a suitable change of frame in moduli space. The dual 't Hooft coupling σ is shown to be a simple function of the periods in Eq. (6) and the modular coordinate $\partial_t^2 F_0(z)$ which appears in the study of the modular properties and BPS spectrum of local \mathbb{P}^2 [34, 35, 61]. We expect further progress in this direction [62, 63].

Let us go back to the fermionic spectral traces $Z(N, \hbar)$, $N \in \mathbb{Z}_{>0}$, in Eq. (15) and perturbatively expand them in the limits $\hbar \rightarrow 0$ and $\hbar \rightarrow \infty$ with N fixed. Following [1, 32], we construct the two families of asymptotic series

$$\log Z(N, \hbar) \sim \phi_N(\hbar) \quad \text{for } \hbar \rightarrow 0, \quad (28a)$$

$$\log Z(N, \hbar) \sim \psi_N(\hbar^{-1}) \quad \text{for } \hbar \rightarrow \infty, \quad (28b)$$

indexed by the non-negative integer N . Under the assumption that the formal power series in Eq. (28) are Gevrey-1 and simple resurgent, which is explicitly verified in the examples considered in this paper, the theory of resurgence can be applied to give us access to the non-analytic sectors that are hidden in perturbation theory. We briefly recall some of the basic notions of resurgence in Appendix B. In order to perform a resurgent analysis of the fermionic spectral traces, we consider their analytic continuation to $\hbar \in \mathbb{C}'$. A conjectural proposal for the resurgent structure of the asymptotic series in Eq. (28) has been recently formulated⁶ in [1, 32]. Let us briefly recall its main statements. For fixed $N \in \mathbb{Z}_{>0}$, each of the formal power series in Eq. (28a), which emerge in the semiclassical limit $\hbar \rightarrow 0$ of the spectral theory, is associated with a minimal resurgent structure \mathfrak{B}_{ϕ_N} of the particular form

$$\mathfrak{B}_{\phi_N} = \{\Phi_{\sigma,n;N}(\hbar)\}_{\sigma=0,\dots,l_0; n \in \mathbb{Z}}, \quad \Phi_{\sigma,n;N}(\hbar) = \phi_{\sigma;N}(\hbar) e^{-n \frac{\mathcal{A}_0}{\hbar}}, \quad (29)$$

where the non-negative integer l_0 and the complex constant \mathcal{A}_0 depend on the choice of N . Namely, for each value of $\sigma \in \{0, \dots, l_0\}$, a Gevrey-1 asymptotic series $\phi_{\sigma;N}(\hbar)$ resurges from the original perturbative expansion $\phi_N(\hbar) = \phi_{0;N}(\hbar)$ and the singularities of its Borel transform $\hat{\phi}_{\sigma;N}(\zeta)$ are located along infinite towers in the Borel ζ -plane so that every two singularities in the same tower are spaced by an integer multiple of \mathcal{A}_0 . The global arrangement of the complete set of Borel singularities of \mathfrak{B}_{ϕ_N} is known as a peacock pattern⁷. In this way, each asymptotic series $\phi_{\sigma;N}(\hbar)$ gives rise to an infinite family of basic trans-series $\Phi_{\sigma,n;N}(\hbar)$, $n \in \mathbb{Z}$. The corresponding infinite-dimensional matrix of Stokes constants \mathcal{S}_{ϕ_N} can be written as

$$\mathcal{S}_{\phi_N} = \{S_{\sigma,\sigma',n;N} \in \mathbb{Q}\}_{\sigma,\sigma'=0,\dots,l_0; n \in \mathbb{Z}}, \quad (30)$$

⁵The maximal conifold point can be defined as the unique point in the conifold locus of moduli space where its connected components intersect transversally.

⁶The conjecture of [1, 32] describes a universal structure underlying the resurgence of the strong and weak coupling perturbative expansions of the fermionic spectral traces for all local CY threefolds. Explicit results in support of the conjecture are obtained in the examples of local \mathbb{P}^2 and local \mathbb{F}_0 .

⁷Peacock patterns are typically observed in theories controlled by a quantum curve in exponentiated variables, including complex Chern–Simons theory on the complement of a hyperbolic knot [64, 65].

after fixing a canonical normalization of the asymptotic series $\phi_{\sigma;N}(\hbar)$. It is conjectured that the Stokes constants $S_{\sigma,\sigma',n;N}$ are closely related to non-trivial sequences of integers, thus representing a new class of enumerative invariants of the topological string on local \mathbb{P}^2 . Furthermore, they can be naturally organized as the coefficients of generating functions that are expressible in the form of q -series and are uniquely determined by the original perturbative expansion $\phi_N(\hbar)$. Schematically,

$$S_{\sigma\sigma';N}(q) = \sum_{n \in \mathbb{Z}} S_{\sigma\sigma',n;N} q^n. \quad (31)$$

The conjectural proposal of [1, 32] for the formal power series in Eq. (28b), which emerge in the strongly-coupled limit $\hbar \rightarrow \infty$ of the spectral theory, is entirely analogous. At fixed $N \in \mathbb{Z}_{>0}$, the minimal resurgent structure \mathfrak{B}_{ψ_N} is given by

$$\mathfrak{B}_{\psi_N} = \{\Psi_{\sigma,n;N}(\hbar^{-1})\}_{\sigma=0,\dots,l_\infty; n \in \mathbb{Z}}, \quad \Psi_{\sigma,n;N}(\hbar^{-1}) = \psi_{\sigma;N}(\hbar^{-1}) e^{-n\mathcal{A}_\infty \hbar}, \quad (32)$$

where again $l_\infty \in \mathbb{Z}_{>0}$ and $\mathcal{A}_\infty \in \mathbb{C}$ depend on the choice of N , resulting in a dual peacock arrangement of singularities in the complex Borel plane. The infinite-dimensional matrix of Stokes constants \mathcal{S}_{ψ_N} is instead

$$\mathcal{S}_{\psi_N} = \{R_{\sigma,\sigma',n;N} \in \mathbb{Q}\}_{\sigma,\sigma'=0,\dots,l_\infty; n \in \mathbb{Z}}, \quad (33)$$

after fixing the normalization of the asymptotic series $\psi_{\sigma;N}(\hbar^{-1})$. Once more, the Stokes constants $R_{\sigma,\sigma',n;N}$ provide a new non-trivial class of integer invariants of the geometry and can be naturally organized into the generating q -series

$$R_{\sigma\sigma';N}(q) = \sum_{n \in \mathbb{Z}} R_{\sigma\sigma',n;N} q^n, \quad (34)$$

which are uniquely determined by the original perturbative expansion $\psi_N(\hbar^{-1})$.

Remark 2.3. *The work of [32] refers to the exponential of the series in Eq. (28b), that is, the perturbative expansion in g_s of the fermionic spectral traces directly. As pointed out in [1], however, the arithmetic properties underlying the resurgent structures of the series in Eq. (28) for $N = 1$ become less easily accessible after exponentiation. For example, the Stokes constants for $\exp(\phi_1(\hbar))$ are complex numbers with no manifest interpretation beyond the fact that they can be expressed in terms of the Stokes constants of $\phi_1(\hbar)$ by means of a closed partition-theoretic formula [1]. Similarly, the exact number-theoretic duality between the weak and strong coupling regimes emerges when looking at the logarithm of the spectral trace – which is akin to considering the free energy instead of the partition function. Let us stress that, as shown explicitly in [1], the analytic solution to the resurgent structures of the series in Eq. (28) can be translated into results on the corresponding exponentiated series by applying the tools of alien calculus [2, 12–14].*

Remark 2.4. *Let us stress that the TS/ST statement in Eq. (19) and the strong-weak coupling duality in Eq. (21) map the semiclassical \hbar -series in Eq. (28a) into a dual perturbative \hbar -expansion that is described by the NS limit of the refined topological string on local \mathbb{P}^2 . Analogously, the asymptotic series for $\hbar \rightarrow \infty$ in Eq. (28b) is associated with a dual perturbative expansion in the weakly coupled regime $g_s \rightarrow 0$ that is captured by the conventional topological string on the same geometry. Notably, the standard and NS topological string free energies can be regarded as non-perturbative corrections of one another in the appropriate regimes [7, 8]. The interplay of the two one-parameter specializations of the refined topological string theory on a local CY threefold with the perturbative and non-perturbative contributions to the fermionic spectral traces in the dual limits in \hbar is discussed in Section 3.2.*

2.3 Exact solutions for the spectral trace

In the rest of this paper, unless explicitly stated, we will focus on the spectral trace

$$Z_{\mathbb{P}^2}(1, \hbar) = \text{Tr}(\rho_{\mathbb{P}^2}), \quad (35)$$

whose resurgent structures in the semiclassical limit $\hbar \rightarrow 0$ and the dual strongly-coupled limit $\hbar \rightarrow \infty$ have been obtained analytically in closed form in [1]. This is the only example of a fermionic spectral trace of a local CY threefold to have been solved exactly. Under the assumption that $\Re(\mathbf{b}) > 0$, it has the integral representation [44]

$$\text{Tr}(\rho_{\mathbb{P}^2}) = \frac{1}{\sqrt{3}\mathbf{b}} \int_{\mathbb{R}} e^{2\pi \mathbf{b} x/3} \frac{\Phi_{\mathbf{b}}(x + \mathbf{i}\mathbf{b}/3)}{\Phi_{\mathbf{b}}(x - \mathbf{i}\mathbf{b}/3)} dx, \quad (36)$$

which is an analytic function of $\hbar \in \mathbb{C}'$. The integral above can be evaluated using the integral Ramanujan formula, or analytically continuing x to the complex domain, completing the integration contour from above, and summing over residues, yielding the closed formula [42]

$$\text{Tr}(\rho_{\mathbb{P}^2}) = \frac{1}{\sqrt{3}\mathbf{b}} e^{-\frac{\pi \mathbf{i}}{36}(12c_{\mathbf{b}}^2 + 4\mathbf{b}^2 - 3)} \frac{\Phi_{\mathbf{b}}(c_{\mathbf{b}} - \frac{\mathbf{i}\mathbf{b}}{3})^2}{\Phi_{\mathbf{b}}(c_{\mathbf{b}} - \frac{2\mathbf{i}\mathbf{b}}{3})} = \frac{1}{\mathbf{b}} \left| \Phi_{\mathbf{b}}\left(c_{\mathbf{b}} - \frac{\mathbf{i}\mathbf{b}}{3}\right) \right|^3, \quad (37)$$

where $c_{\mathbf{b}} = \mathbf{i}(\mathbf{b} + \mathbf{b}^{-1})/2$. Moreover, following [32], the expression in Eq. (37) can be factorized into a product of q, \tilde{q} -series by applying the infinite product representation in Eq. (183). Namely, we have that

$$\Phi_{\mathbf{b}}\left(c_{\mathbf{b}} - \frac{\mathbf{i}\mathbf{b}}{3}\right) = \frac{(q^{2/3}; q)_{\infty}}{(w^{-1}; \tilde{q})_{\infty}}, \quad \Phi_{\mathbf{b}}\left(c_{\mathbf{b}} - \frac{2\mathbf{i}\mathbf{b}}{3}\right) = \frac{(q^{1/3}; q)_{\infty}}{(w; \tilde{q})_{\infty}}, \quad (38)$$

where $(xq^{\alpha}; q)_{\infty}$ is the quantum dilogarithm defined in Eq. (176). Therefore,

$$\text{Tr}(\rho_{\mathbb{P}^2}) = \frac{1}{\sqrt{3}\mathbf{b}} e^{-\frac{\pi \mathbf{i}}{36}\mathbf{b}^2 + \frac{\pi \mathbf{i}}{12}\mathbf{b}^{-2} + \frac{\pi \mathbf{i}}{4}} \frac{(q^{2/3}; q)_{\infty}^2}{(q^{1/3}; q)_{\infty}} \frac{(w; \tilde{q})_{\infty}}{(w^{-1}; \tilde{q})_{\infty}^2}, \quad (39)$$

where we have introduced

$$q = e^{2\pi \mathbf{i} \mathbf{b}^2} = e^{3\mathbf{i} \hbar}, \quad \tilde{q} = e^{-2\pi \mathbf{i} / \mathbf{b}^2} = e^{-4\pi^2 \mathbf{i} / (3\hbar)}, \quad w = e^{2\pi \mathbf{i} / 3}. \quad (40)$$

Note that the factorization into holomorphic/anti-holomorphic blocks in Eq. (39) is not symmetric in q, \tilde{q} . We further assume that $\Im(\mathbf{b}^2) > 0$, and therefore $\Im(\hbar) > 0$, which implies $|q|, |\tilde{q}| < 1$, so that the q, \tilde{q} -series converge.

Let us now recall some of the results of [1] on the exact resurgent structures of the perturbative expansions of $\text{Tr}(\rho_{\mathbb{P}^2})$ in both weak and strong \hbar -regimes and the analytic number-theoretic duality relating them, which is the starting point for the investigation performed in this paper. In the following, we will summarize part of [1, Section 4]. We invite the interested reader to consult the original reference for a complete and detailed description with proofs.

The limit $\hbar \rightarrow 0$

The all-orders perturbative expansion in the limit $\hbar \rightarrow 0$ of the spectral trace of local \mathbb{P}^2 in Eq. (39) is obtained by applying the asymptotic expansion formula for the quantum dilogarithm

in Eq. (193b) to the holomorphic q -series component⁸. Namely,

$$\mathrm{Tr}(\rho_{\mathbb{P}^2}) \sim \frac{\Gamma(\frac{1}{3})^3}{6\pi\hbar} \exp\left(3 \sum_{n=1}^{\infty} (-1)^{n-1} \frac{B_{2n}B_{2n+1}(2/3)}{2n(2n+1)!} (3\hbar)^{2n}\right), \quad (41)$$

where $B_n(x)$ is the n -th Bernoulli polynomial, $B_n = B_n(0)$ is the n -th Bernoulli number, and $\Gamma(x)$ is the gamma function. Let us denote by $\phi(\hbar)$ the formal power series appearing in the exponent in Eq. (41), that is,

$$\phi(\hbar) = \sum_{n=1}^{\infty} a_{2n} \hbar^{2n} \in \mathbb{Q}[[\hbar]], \quad a_{2n} = (-1)^{n-1} \frac{B_{2n}B_{2n+1}(2/3)}{2n(2n+1)!} 3^{2n+1} \quad n \geq 1. \quad (42)$$

Its perturbative coefficients satisfy the factorial growth

$$|a_{2n}| \leq (2n)! \mathcal{A}_0^{-2n} \quad n \gg 1, \quad \mathcal{A}_0 = \frac{4\pi^2}{3}, \quad (43)$$

and $\phi(\hbar)$ is thus a Gevrey-1 asymptotic series. The Borel transform $\hat{\phi}(\zeta)$ can be explicitly resummed into a well-defined, exact function of ζ which is manifestly simple resurgent. Its singularities are simple poles⁹ located along the imaginary axis at

$$\zeta_n = \mathcal{A}_0 i n, \quad n \in \mathbb{Z}_{\neq 0}, \quad (44)$$

tracing two Stokes lines at the angles $\pm\pi/2$. This is the simplest occurrence of the peacock pattern described in Section 2.2.

The local expansion of the Borel transform $\hat{\phi}(\zeta)$ at $\zeta = \zeta_n$, $n \in \mathbb{Z}_{\neq 0}$, is given by

$$\hat{\phi}(\zeta) = -\frac{S_n}{2\pi i(\zeta - \zeta_n)} + \text{regular in } \zeta - \zeta_n, \quad (45)$$

where S_n is the corresponding Stokes constant. It follows that all the Stokes constants can be accessed analytically. In particular, the normalized Stokes constant S_n/S_1 , where $S_1 = 3\sqrt{3}i$, is given by the divisor sum function

$$\frac{S_n}{S_1} = \sum_{\substack{d|n \\ d \equiv 1 \pmod{3}}} \frac{1}{d} - \sum_{\substack{d|n \\ d \equiv 2 \pmod{3}}} \frac{1}{d} = \sum_{d|n} \frac{1}{d} \chi_{3,2}(d) \in \mathbb{Q}_{>0}, \quad n \in \mathbb{Z}_{\neq 0}, \quad (46)$$

where $d|n$ denotes the positive integer divisors d of n and $\chi_{3,2}(n)$ is the unique non-principal Dirichlet character modulo 3, that is,

$$\chi_{3,2}(n) = \begin{cases} 0 & \text{if } n \equiv 0 \pmod{3} \\ 1 & \text{if } n \equiv 1 \pmod{3} \\ -1 & \text{if } n \equiv 2 \pmod{3} \end{cases}, \quad n \in \mathbb{Z}. \quad (47)$$

It follows that S_n/S_1 is a multiplicative arithmetic function and satisfies $S_{-n} = S_n$. Note that we can introduce the sequence of integers

$$\alpha_n = \frac{S_n}{S_1} n, \quad n \in \mathbb{Z}_{\neq 0}, \quad (48)$$

which satisfies $\alpha_n \in \mathbb{Z}_{>0}$ for $n \in \mathbb{Z}_{>0}$ and $\alpha_{-n} = -\alpha_n$. The pattern of singularities in the Borel plane and the associated α_n , $n \in \mathbb{Z}_{\neq 0}$ are shown in Fig. 2 on the left.

⁸The \tilde{q} -series giving the anti-holomorphic block in the factorized expression in Eq. (39) converge for $\hbar \rightarrow 0$.

⁹We apply the conventions of Appendix B, which differ slightly from those adopted in [1] due to a change in the definition of the Borel and Laplace transforms. In particular, with the conventions of [1], the singularities of the Borel transforms of the series $\phi(\hbar), \psi(\tau)$ are logarithmic branch points.




Figure 2: On the left, the first few singularities of the Borel transform of the asymptotic series $\phi(\hbar)$, defined in Eq. (42), and the associated integer constants α_n , $n \in \mathbb{Z}_{\neq 0}$, defined in Eq. (48). On the right, the first few singularities of the Borel transform of the asymptotic series $\psi(\hbar)$, defined in Eq. (51), and the associated integer constants β_n , $n \in \mathbb{Z}_{\neq 0}$, defined in Eq. (56).

The limit $\hbar \rightarrow \infty$

The all-orders perturbative expansion in the limit $\hbar \rightarrow \infty$ of the spectral trace of local \mathbb{P}^2 in Eq. (39) is obtained by applying the asymptotic expansion formula for the quantum dilogarithm in Eq. (193a) to the anti-holomorphic \tilde{q} -series component¹⁰. Namely,

$$\text{Tr}(\rho_{\mathbb{P}^2}) \sim \sqrt{\frac{2\pi}{3^{5/2}\hbar}} e^{-\frac{3V}{4\pi^2}\hbar} \exp \left(\sqrt{3} \sum_{n=1}^{\infty} (-1)^{n-1} \frac{B_{2n} B_{2n-1}(2/3)}{(2n)!(2n-1)} \left(\frac{4\pi^2}{\hbar} \right)^{2n-1} \right), \quad (49)$$

where $V = 2\Im(\text{Li}_2(e^{\pi i/3}))$. Let us introduce the parameter

$$\tau = -\frac{1}{b^2} = -\frac{2\pi}{3\hbar} \quad (50)$$

and denote by $\psi(\tau)$ the formal power series appearing in the exponent in Eq. (49), that is,

$$\psi(\tau) = \sum_{n=1}^{\infty} b_{2n} \tau^{2n-1} \in \mathbb{Q}[\pi, \sqrt{3}][[\tau]], \quad b_{2n} = (-1)^n \sqrt{3} \frac{B_{2n} B_{2n-1}(2/3)}{(2n)!(2n-1)} (6\pi)^{2n-1} \quad n \geq 1. \quad (51)$$

Its perturbative coefficients satisfy the factorial growth

$$|b_{2n}| \leq (2n)! \mathcal{A}_{\infty}^{-2n} \quad n \gg 1, \quad \mathcal{A}_{\infty} = \frac{2\pi}{3} = \frac{\mathcal{A}_0}{2\pi}, \quad (52)$$

¹⁰The q -series giving the holomorphic block in the factorized expression in Eq. (39) converge for $\hbar \rightarrow \infty$.

and $\psi(\tau)$ is thus a Gevrey-1 asymptotic series. Analogously to the case of $\hbar \rightarrow 0$, the Borel transform $\hat{\psi}(\zeta)$ can be explicitly resummed into an exact function of ζ which is manifestly simple resurgent. Its singularities are simple poles located along the imaginary axis at

$$\eta_n = \mathcal{A}_\infty i n, \quad n \in \mathbb{Z}_{\neq 0}, \quad (53)$$

again tracing the only two Stokes lines at the angles $\pm\pi/2$.

The local expansion of the Borel transform $\hat{\psi}(\zeta)$ at $\zeta = \eta_n$, $n \in \mathbb{Z}_{\neq 0}$, is given by

$$\hat{\psi}(\zeta) = -\frac{R_n}{2\pi i(\zeta - \eta_n)} + \text{regular in } \zeta - \eta_n, \quad (54)$$

where R_n is the corresponding Stokes constant. Once more, all the Stokes constants can be derived analytically in closed form. Indeed, the normalized Stokes constant R_n/R_1 , where $R_1 = 3$, is given by the divisor sum function

$$\frac{R_n}{R_1} = \sum_{\substack{d|n \\ d \equiv_3 1}} \frac{d}{n} - \sum_{\substack{d|n \\ d \equiv_3 2}} \frac{d}{n} = \sum_{d|n} \frac{d}{n} \chi_{3,2}(d) \in \mathbb{Q}_{\neq 0}, \quad n \in \mathbb{Z}_{\neq 0}, \quad (55)$$

which is a multiplicative arithmetic function satisfying the property $R_{-n} = -R_n$. Note the strikingly simple arithmetic symmetry between the formulae in Eqs. (46) and (55). Moreover, we can introduce a dual sequence of integers

$$\beta_n = \frac{R_n}{R_1} n, \quad n \in \mathbb{Z}_{\neq 0}, \quad (56)$$

which satisfies $\beta_n \in \mathbb{Z}_{\neq 0}$ for $n \in \mathbb{Z}_{>0}$ and $\beta_{-n} = \beta_n$. The pattern of singularities in the Borel plane and the associated β_n , $n \in \mathbb{Z}_{\neq 0}$, are shown in Fig. 2 on the right.

Exact formulae for the discontinuities

A notable consequence of the closed formulae in Eqs. (46) and (55) is that the Stokes constants S_n, R_n , $n \in \mathbb{Z}_{>0}$, can be naturally organized as coefficients of exact generating functions given by the discontinuities of the corresponding asymptotic series $\phi(\hbar), \psi(\tau)$ across the positive imaginary axis. Explicitly,

$$\text{disc}_{\frac{\pi}{2}} \phi(\hbar) = \sum_{n=1}^{\infty} S_n \tilde{q}^n = -3 \log \frac{(w; \tilde{q})_\infty}{(w^{-1}; \tilde{q})_\infty} - i\pi, \quad (57a)$$

$$\text{disc}_{\frac{\pi}{2}} \psi(\tau) = \sum_{n=1}^{\infty} R_n q^{n/3} = 3 \log \frac{(q^{2/3}; q)_\infty}{(q^{1/3}; q)_\infty}. \quad (57b)$$

Observe that the quantum dilogarithms appearing in the generating function in Eq. (57a) associated with the weak coupling asymptotic expansion $\phi(\hbar)$ are the same \tilde{q} -series that occur in the anti-holomorphic block of the spectral trace in Eq. (39). Similarly, the q -series in the generating function in Eq. (57b) associated with the strong coupling asymptotic expansion $\psi(\tau)$ are those appearing in the dual holomorphic block of the spectral trace in Eq. (39)¹¹.

¹¹Disregarding constant prefactors, the q, \tilde{q} -series in Eq. (57) differ from the logarithms of the holomorphic/anti-holomorphic blocks in Eq. (39) by powers of two acting on the numerator/denominator, respectively.

Remark 2.5. *The q, \tilde{q} -series in the factorized expression of the spectral trace of local \mathbb{P}^2 encode the perturbative information in one asymptotic limit in \hbar , while being invisible to perturbation theory in the dual limit. Simultaneously, the same q, \tilde{q} -series supply the discontinuities in the opposite regimes. Thus, the holomorphic block resurges from the asymptotics of the anti-holomorphic one, and vice versa. This two-way exchange of perturbative/non-perturbative content between the holomorphic/anti-holomorphic blocks is one of several exact cross-relations connecting the weak and strong coupling resurgent structures and paving the way towards the formulation of a global strong-weak resurgent symmetry, which we will describe in Section 3.2.*

A number-theoretic resurgent duality

The standard large- n asymptotics of the perturbative coefficients a_{2n}, b_{2n} , $n \in \mathbb{Z}_{>0}$, in Eqs. (42) and (51) can be upgraded by systematically including the contributions from all sub-dominant singularities in the Borel plane to give the exact large-order relations

$$a_{2n} = \frac{\Gamma(2n)}{\pi i (\mathcal{A}_0 i)^{2n}} \sum_{m=1}^{\infty} \frac{S_m}{m^{2n}}, \quad (58a)$$

$$b_{2n} = \frac{\Gamma(2n-1)}{\pi i (\mathcal{A}_\infty i)^{2n-1}} \sum_{m=1}^{\infty} \frac{R_m}{m^{2n-1}}, \quad (58b)$$

that is, up to the simple prefactors above, the perturbative coefficients of the series $\phi(\hbar), \psi(\tau)$ are given by the Dirichlet series encoding the corresponding weak and strong coupling Stokes constants evaluated at even and odd integer points, respectively.

Amazingly, the Dirichlet series defined by the Stokes constants satisfy an Euler product expansion indexed by the set of prime numbers, are absolutely convergent in the right half-plane $\Re(s) > 1$, and can be analytically continued to meromorphic functions on the whole complex s -plane. The resulting weak and strong coupling L -functions $L_0(s), L_\infty(s)$, $s \in \mathbb{C}$, further admit the factorization

$$L_0(s) = \sum_{m=1}^{\infty} \frac{S_m}{m^s} = S_1 L(s+1, \chi_{3,2}) \zeta(s), \quad (59a)$$

$$L_\infty(s) = \sum_{m=1}^{\infty} \frac{R_m}{m^s} = R_1 L(s, \chi_{3,2}) \zeta(s+1), \quad (59b)$$

where $L(s, \chi_{3,2})$ is the Dirichlet L -function of the primitive character $\chi_{3,2}$ and $\zeta(s)$ is the Riemann zeta function. Observe that the arithmetic duality¹² relating the weak and strong coupling Stokes constants S_n, R_n , $n \in \mathbb{Z}_{>0}$, translates at the level of the L -functions encoded in the perturbative coefficients into a symmetric unitary shift in the arguments of the factors in the RHS (right-hand-side) of Eq. (59).

Remark 2.6. *If we extend the discrete index $n \in \mathbb{Z}_{>0}$ of the sequences of perturbative coefficients $\{a_{2n}\}, \{b_{2n}\}$ to a continuous variable $s \in \mathbb{C}$, then the exact large-order relations in Eq. (58) allow us, in principle, to analytically continue the perturbative coefficients to meromorphic functions throughout the complex s -plane as*

$$a_s = \frac{\Gamma(s)}{\pi i (\mathcal{A}_0 i)^s} L_0(s), \quad b_s = \frac{\Gamma(s-1)}{\pi i (\mathcal{A}_\infty i)^{s-1}} L_\infty(s-1). \quad (60)$$

¹²The symmetric decomposition of the Stokes constants into the Dirichlet convolution of simple arithmetic functions dictates the dual factorizations of the corresponding L -functions [1, Section 4.4].

A similar observation has been made in the recent work of [66] on the exact large order relations for general resurgent trans-series.

3 Resurgent L -functions

In this section, we complete the analytic number-theoretic duality of [1], which connects the exact resurgent structures of $\log \text{Tr}(\rho_{\mathbb{P}^2})$ in the dual regimes of $\hbar \rightarrow 0$ and $\hbar \rightarrow \infty$, and upgrade it into a full-fledged exact *strong-weak resurgent symmetry*. This unique global construction rests on the interplay of q -series and L -functions and revolves around the central role played by the weak and strong coupling Stokes constants.

3.1 Generating functions of the Stokes constants

As it will be useful in the rest of this paper, let us now write the generating series of the weak coupling Stokes constants S_n , $n \in \mathbb{Z}_{\neq 0}$, in Eq. (46) in the form

$$f_0(y) := \begin{cases} \sum_{n>0} S_n e^{2\pi i n y} & \text{if } \Im(y) > 0 \\ -\sum_{n<0} S_n e^{2\pi i n y} & \text{if } \Im(y) < 0 \end{cases}, \quad (61)$$

which defines a holomorphic function on $\mathbb{C} \setminus \mathbb{R}$ with the periodicity and parity properties

$$f_0(y+1) = f_0(y), \quad f_0(-y) = -f_0(y). \quad (62)$$

It follows that $f_0(y)$, $y \in \mathbb{C} \setminus \mathbb{R}$, need only be specified in the upper half of the complex y -plane, where it is known in closed form by means of the exact discontinuity formula in Eq. (57a). Namely, we have that [1, Corollary 4.8]

$$f_0(y) = -3 \log \frac{(w; e^{2\pi i y})_{\infty}}{(w^{-1}; e^{2\pi i y})_{\infty}} - i\pi, \quad y \in \mathbb{H}, \quad (63)$$

where $w = e^{2\pi i/3}$ as before and therefore also

$$f_0\left(-\frac{2\pi}{3\hbar}\right) = \text{disc}_{\frac{\pi}{2}} \phi(\hbar), \quad \hbar \in \mathbb{H}. \quad (64)$$

Analogously, we write the generating series of the strong coupling Stokes constants R_n , $n \in \mathbb{Z}_{\neq 0}$, in Eq. (55) in the form

$$f_{\infty}(y) := \begin{cases} \sum_{n>0} R_n e^{2\pi i n y} & \text{if } \Im(y) > 0 \\ -\sum_{n<0} R_n e^{2\pi i n y} & \text{if } \Im(y) < 0 \end{cases}, \quad (65)$$

which defines a holomorphic function on $\mathbb{C} \setminus \mathbb{R}$ with the periodicity and parity properties¹³

$$f_{\infty}(y+1) = f_{\infty}(y), \quad f_{\infty}(-y) = f_{\infty}(y). \quad (66)$$

¹³Note the exchange of even/odd parity between the Stokes constants and the corresponding generating functions in both weak and strong coupling limits.

Again, it follows that $f_\infty(y)$, $y \in \mathbb{C} \setminus \mathbb{R}$, need only be specified in the upper half of the complex y -plane, where it is known in closed form through the exact discontinuity formula in Eq. (57b). Explicitly, we have that [1, Corollary 4.19]

$$f_\infty(y/3) = 3 \log \frac{(e^{4\pi i y/3}; e^{2\pi i y})_\infty}{(e^{2\pi i y/3}; e^{2\pi i y})_\infty}, \quad y \in \mathbb{H}, \quad (67)$$

and therefore also

$$f_\infty\left(-\frac{1}{3\tau}\right) = \text{disc}_{\frac{\pi}{2}} \psi(\tau), \quad \tau \in \mathbb{H}. \quad (68)$$

Remark 3.1. Let us now consider the discontinuities $\text{disc}_{\frac{\pi}{2}} \phi(\hbar)$ and $\text{disc}_{\frac{\pi}{2}} \psi(\tau)$ as functions of the variable $\tau \in \mathbb{H}$ using Eqs. (57a), (57b), and (50). Namely, we take the functions

$$\left(\text{disc}_{\frac{\pi}{2}} \phi\right)(\tau) = -3 \log \frac{(w; \tilde{q})_\infty}{(w^{-1}; \tilde{q})_\infty} - i\pi, \quad \left(\text{disc}_{\frac{\pi}{2}} \psi\right)(\tau) = 3 \log \frac{(q^{2/3}; q)_\infty}{(q^{1/3}; q)_\infty}, \quad (69)$$

where $\tilde{q} = e^{2\pi i \tau}$ and $q = e^{-2\pi i/\tau}$ as before. It follows that

$$\left(\text{disc}_{\frac{\pi}{2}} \phi\right)(\tau + 1) = \left(\text{disc}_{\frac{\pi}{2}} \phi\right)(\tau), \quad \left(\text{disc}_{\frac{\pi}{2}} \psi\right)\left(\frac{\tau}{3\tau + 1}\right) = \left(\text{disc}_{\frac{\pi}{2}} \psi\right)(\tau), \quad (70)$$

that is, the discontinuities are invariant under the action of the generators $T : \tau \mapsto \tau + 1$ and $\gamma_3 : \tau \mapsto \tau/(3\tau + 1)$ of the congruence subgroup $\Gamma_1(3)$, respectively. Recall that we have introduced $\Gamma_1(3)$ in Eq. (8) as it dictates the symmetries of the moduli space of complex structures of the mirror of local \mathbb{P}^2 . The same $\Gamma_1(3)$ -structure crucially occurs in Section 4.3, where the modular considerations above are upgraded to full quantum modularity statements on the generating functions of the Stokes constants $f_0(y)$ and $f_\infty(y)$, $y \in \mathbb{H}$.

Asymptotic series for the generating functions

Let us compute the all-orders perturbative expansions in the limit $y \rightarrow 0$ of the weak and strong coupling generating functions $f_0(y)$, $f_\infty(y)$, $y \in \mathbb{H}$, in Eqs. (63) and (67), which we denote by $\tilde{f}_0(y)$, $\tilde{f}_\infty(y)$, respectively. We apply the known asymptotic expansion formula for the quantum dilogarithm in Eq. (193a) to the closed expression for the generating function of the weak coupling Stokes constants in Eq. (63) and explicitly evaluate the special functions that appear. We find that

$$\tilde{f}_0(y) = -\frac{\pi i}{2} - \frac{3}{2\pi i y} (\text{Li}_2(w) - \text{Li}_2(w^{-1})) - 2\sqrt{3}i \sum_{n=1}^{\infty} (6\pi i y)^{2n-1} \frac{B_{2n} B_{2n-1}(2/3)}{(2n)!(2n-1)!}, \quad (71)$$

which simplifies into

$$\tilde{f}_0(y) = -\frac{\pi i}{2} - \frac{3\mathcal{V}}{2\pi i y} - 2\psi(y), \quad (72)$$

where¹⁴ $\mathcal{V} = 2\Im(\text{Li}_2(w))$ and $\psi(y)$ is the asymptotic series occurring in the strong coupling perturbative expansion of $\text{Tr}(\rho_{\mathbb{P}^2})$ in Eq. (51). Similarly, we apply the known asymptotic expansion formula for the quantum dilogarithm in Eq. (193b), with the choice of $\alpha = 1/3, 2/3$, to the

¹⁴Observe that $\frac{27}{8\pi^2}\mathcal{V} \simeq 0.462758$ is the quantum volume of local \mathbb{P}^2 , which is given by the value of its Kähler parameter at the conifold [67]. This is related to the constant V in Eq. (49) by $\mathcal{V} = -2\pi^2/3 + 2iV/3$.

closed expression for the generating function of the strong coupling Stokes constants in Eq. (67) and explicitly evaluate the special functions that appear. We derive in this way that

$$\tilde{f}_\infty(y) = -3 \log \frac{\Gamma(2/3)}{\Gamma(1/3)} - \log(-6\pi i y) - 6 \sum_{n=1}^{\infty} (6\pi i y)^{2n} \frac{B_{2n} B_{2n+1}(2/3)}{2n(2n+1)!}, \quad (73)$$

which equals to

$$\tilde{f}_\infty(y) = -3 \log \frac{\Gamma(2/3)}{\Gamma(1/3)} - \log(-6\pi i y) + 2\phi(2\pi y), \quad (74)$$

where $\phi(y)$ is the asymptotic series occurring in the weak coupling perturbative expansion of $\text{Tr}(\rho_{\mathbb{P}^2})$ in Eq. (42). As a consequence of Eqs. (72) and (74), the exact resurgent structures of the asymptotic series $\tilde{f}_0(y)$ and $\tilde{f}_\infty(y)$ follow straightforwardly from the work of [1] on the perturbative expansions $\psi(\tau)$ and $\phi(\hbar)$, respectively, which we have summarized in Section 2.3.

Remark 3.2. Putting together Eqs. (64) and (68) with Eqs. (72) and (74), we obtain the asymptotic expansions of the discontinuities of the original perturbative series $\phi(\hbar)$ and $\psi(\tau)$ in the large limits of the variables \hbar and τ , respectively. Namely, we have that

$$\text{disc}_{\frac{\pi}{2}} \phi(\hbar) \sim -\frac{\pi i}{2} - \frac{9\mathcal{V}}{4\pi^2 i} \hbar - 2\psi\left(-\frac{2\pi}{3\hbar}\right), \quad \hbar \rightarrow \infty, \quad (75a)$$

$$\text{disc}_{\frac{\pi}{2}} \psi(\tau) \sim -3 \log \frac{\Gamma(2/3)}{\Gamma(1/3)} - \log\left(\frac{2\pi i}{\tau}\right) + 2\phi\left(-\frac{2\pi}{3\tau}\right), \quad \tau \rightarrow \infty. \quad (75b)$$

3.2 Strong-weak resurgent symmetry

As we described in Section 2.3, following the work of [1], and in Section 3.1, the resurgent structures of the logarithm of the spectral trace of local \mathbb{P}^2 in both weak and strong coupling limits display a rich analytic number-theoretic fabric centered around the key role played by the Stokes constants and their generating functions. The interaction of q -series and L -functions governs a network of relations encompassing and intertwining the regimes of $\hbar \rightarrow 0$ and $\hbar \rightarrow \infty$. Let us give a concise overview.

- The discontinuity $\text{disc}_{\frac{\pi}{2}} \psi(\tau)$, which equals the generating function $f_\infty(-\frac{1}{3\tau})$ in Eq. (67) of the strong coupling Stokes constants R_n , $n \in \mathbb{Z}_{\neq 0}$, in Eq. (55), reproduces the weak coupling perturbative series $\phi(\hbar)$ in Eq. (42) when expanded in the limit $\tau \rightarrow \infty$, as we have seen in Eq. (75b).
- Resurgence associates $\phi(\hbar)$ with the collection $\{\zeta_n, S_n\}$, $n \in \mathbb{Z}_{\neq 0}$, of simple poles and Stokes constants, which satisfy $\zeta_n = \mathcal{A}_0 i n$ and $S_{-n} = S_n$.
- The exact large- n relations in Eq. (58a) for the perturbative coefficients a_{2n} , $n \in \mathbb{Z}_{>0}$, of $\phi(\hbar)$ uniquely determine the weak coupling L -function $L_0(s)$ in Eq. (59a). The latter reproduces in turn the perturbative coefficients when evaluated at $s = 2n$.
- The discontinuity $\text{disc}_{\frac{\pi}{2}} \phi(\hbar)$, which equals the generating function $f_0(-\frac{2\pi}{3\hbar})$ in Eq. (63) of the weak coupling Stokes constants S_n , $n \in \mathbb{Z}_{\neq 0}$, in Eq. (46), reproduces the strong coupling perturbative series $\psi(\tau)$ in Eq. (51) when expanded in the limit $\hbar \rightarrow \infty$, as we have seen in Eq. (75a).
- Resurgence associates $\psi(\tau)$ with the collection $\{\eta_n, R_n\}$, $n \in \mathbb{Z}_{\neq 0}$, of simple poles and Stokes constants, which satisfy $\eta_n = \mathcal{A}_\infty i n$ and $R_{-n} = -R_n$.

- The exact large- n relations in Eq. (58b) for the perturbative coefficients b_{2n} , $n \in \mathbb{Z}_{>0}$, of $\psi(\tau)$ uniquely determine the strong coupling L -function $L_\infty(s)$ in Eq. (59b). The latter reproduces in turn the perturbative coefficients when evaluated at $s = 2n - 1$.

Assembling the above relations together, we find that the resurgent structures of the dual perturbative expansions of $\log \text{Tr}(\rho_{\mathbb{P}^2})$ embed into a unique global construction as they compose the symmetric diagram below.

$$\begin{array}{ccccc}
 & \text{disc}_{\frac{\pi}{2}} \psi(\tau) & \xrightarrow{\hbar \rightarrow 0} & \phi(\hbar) & \\
 & \updownarrow \begin{array}{c} \text{inverse Mellin} \\ \text{Mellin} \end{array} & & \updownarrow \begin{array}{c} \text{evaluation} \\ \text{exact large-}n \end{array} & \\
 \begin{array}{c} \{\eta_n, R_n\} \end{array} & \xrightarrow{\text{Dirichlet series}} & \begin{array}{c} \{\mathcal{A}_\infty, L_\infty\} \end{array} & & \begin{array}{c} \{\mathcal{A}_0, L_0\} \end{array} \xleftarrow{\text{Dirichlet series}} \begin{array}{c} \{\zeta_n, S_n\} \end{array} \\
 & \nwarrow \text{resurgence} & & \nwarrow \text{resurgence} & \\
 & \psi(\tau) & \xleftarrow{\tau \rightarrow 0} & \text{disc}_{\frac{\pi}{2}} \phi(\hbar) & \\
 & \updownarrow \begin{array}{c} \text{exact large-}n \\ \text{evaluation} \end{array} & & \updownarrow \begin{array}{c} \text{Mellin} \\ \text{inverse Mellin} \end{array} & \\
 & \begin{array}{c} \{\eta_n, R_n\} \end{array} & & \begin{array}{c} \{\mathcal{A}_0, L_0\} \end{array} & \begin{array}{c} \{\zeta_n, S_n\} \end{array}
 \end{array} \quad (76)$$

Notably, starting from any fixed vertex, all other vertices of this commutative diagram are spanned by following its directed arrows – that is, the information contained in each vertex reconstructs the whole diagram and is therefore equivalent to the information content of every other vertex. Moreover, two additional arrows are drawn in the diagram in Eq. (76) that were not previously stated. They originate from the well-established relation between power series and Dirichlet series sharing the same coefficients, which makes use of the Mellin transform.

Lemma 3.1. *The weak and strong coupling L -functions $L_0(s)$, $L_\infty(s)$, $s \in \mathbb{C}$, in Eqs. (59a) and (59b) are given by the Mellin transform of the generating functions $f_0(y)$, $f_\infty(y)$, $y \in \mathbb{H}$, of the corresponding Stokes constants, that is,*

$$L_0(s) = \frac{(2\pi)^s}{\Gamma(s)} \int_0^\infty t^{s-1} f_0(it) dt, \quad (77a)$$

$$L_\infty(s) = \frac{(2\pi)^s}{\Gamma(s)} \int_0^\infty t^{s-1} f_\infty(it) dt. \quad (77b)$$

Proof. Let us start with the weak coupling regime of $\hbar \rightarrow 0$ and explicitly compute the Mellin transform of the generating function of the Stokes constants in Eq. (61). We find that

$$\int_0^\infty t^{s-1} f_0(it) dt = \int_0^\infty t^{s-1} \left(\sum_{k=1}^\infty S_k e^{-k2\pi t} \right) dt = \sum_{k=1}^\infty S_k \int_0^\infty t^{s-1} e^{-k2\pi t} dt, \quad (78)$$

where we have permuted sum and integration due to absolute convergence. The simplified integral in the RHS is then easily computed via the well-known formula

$$\int_0^\infty x^{s-1} e^{-kx} dx = \frac{\Gamma(s)}{k^s}, \quad k \in \mathbb{Z}_{>0}, \quad s \in \mathbb{C}, \quad (79)$$

which gives the desired expression in Eq. (77a). The same procedure above is straightforwardly applied in the strong coupling regime of $\hbar \rightarrow \infty$ to explicitly compute the Mellin transform of the generating function of the Stokes constants in Eq. (65), yielding the expression in Eq. (77b). \square

By the Mellin inversion theorem, we recover the generating functions of the Stokes constants via the inverse Mellin transform of the corresponding L -functions. Composing Lemma 3.1 with the exact large-order relations, we find a direct way of obtaining the perturbative expansion of the generating function in one asymptotic limit from the generating function in the dual limit, which amounts to performing an operation that is the formal inverse of taking the discontinuity of the perturbative series.

Proposition 3.2. *The perturbative coefficients a_{2n} , b_{2n} , $n \in \mathbb{Z}_{>0}$, in Eqs. (42) and (51) are given by the Mellin transform of the discontinuities $\text{disc}_{\frac{\pi}{2}} \phi(\hbar)$, $\text{disc}_{\frac{\pi}{2}} \psi(\tau)$, respectively, that is,*

$$a_{2n} = \frac{(-1)^n}{\pi i} \int_0^\infty \hbar^{-2n-1} \text{disc}_{\frac{\pi}{2}} \phi(-i\hbar) d\hbar, \quad (80a)$$

$$b_{2n} = \frac{(-1)^n}{\pi} \int_0^\infty \tau^{-2n} \text{disc}_{\frac{\pi}{2}} \psi(-i\tau) d\tau. \quad (80b)$$

Proof. Using Eq. (64) and applying the change of variable $2\pi t = \mathcal{A}_0/\hbar$, the integral in the RHS of Eq. (80a) becomes

$$\int_0^\infty \hbar^{-2n-1} f_0\left(\frac{\mathcal{A}_0 i}{2\pi \hbar}\right) d\hbar = \frac{(2\pi)^{2n}}{\mathcal{A}_0^{2n}} \int_0^\infty t^{2n-1} f_0(it) dt. \quad (81)$$

Using Eq. (77a) for $s = 2n$ and the exact large- n relations for the weak coupling perturbative coefficients in Eq. (58a), we conclude. Similarly, using Eq. (68) and applying the change of variable $2\pi t = \mathcal{A}_\infty/\tau$, the integral in the RHS of Eq. (80b) becomes

$$\int_0^\infty \tau^{-2n} f_\infty\left(\frac{\mathcal{A}_\infty i}{2\pi \tau}\right) d\tau = \frac{(2\pi)^{2n-1}}{\mathcal{A}_\infty^{2n-1}} \int_0^\infty t^{2n-2} f_\infty(it) dt. \quad (82)$$

Using Eq. (77b) for $s = 2n-1$ and the exact large- n relations for the strong coupling perturbative coefficients in Eq. (58b) yields the desired expression. \square

Consider the commutative diagram in Eq. (76) once more. The internal vertices and arrows composing the left-side sub-diagram that connects the q -series $\text{disc}_{\frac{\pi}{2}} \psi(\tau)$ with the strong coupling perturbative series $\psi(\tau)$ reduce to one vertical two-headed arrow according to the formulae in Eqs. (57b) and (80b). Analogously, the set of internal vertices and arrows composing the right-side sub-diagram that connects the \tilde{q} -series $\text{disc}_{\frac{\pi}{2}} \phi(\hbar)$ with the weak coupling perturbative series $\phi(\hbar)$ is equivalent to one vertical two-headed arrow according to the dual formulae in Eqs. (57a) and (80a). Thus, the two-way exchange of perturbative/non-perturbative information between the dual regimes in \hbar takes the form of a mathematically precise mechanism, which we refer to as *strong-weak resurgent symmetry*. We represent it schematically in the box diagram below,

where we stress the contribution of the Stokes constants to all quantities of interest.

$$\begin{array}{ccc}
\text{disc}_{\frac{\pi}{2}}\psi(\tau) = \sum_{n=1}^{\infty} R_n q^{n/3} & \xrightarrow{\hbar \rightarrow 0} & \phi(\hbar) = \sum_{n=1}^{\infty} \left(\frac{\Gamma(2n)}{\pi i (\mathcal{A}_0 i)^{2n}} \sum_{m=1}^{\infty} \frac{S_m}{m^{2n}} \right) \hbar^{2n} \\
\uparrow \text{strong-weak} & & \uparrow \text{strong-weak} \\
& \text{symmetry} & \text{symmetry} \\
\psi(\tau) = \sum_{n=1}^{\infty} \left(\frac{\Gamma(2n-1)}{\pi i (\mathcal{A}_{\infty} i)^{2n-1}} \sum_{m=1}^{\infty} \frac{R_m}{m^{2n-1}} \right) \tau^{2n-1} & \xleftarrow{\tau \rightarrow 0} & \text{disc}_{\frac{\pi}{2}}\phi(\hbar) = \sum_{n=1}^{\infty} S_n \tilde{q}^n
\end{array} \tag{83}$$

Remark 3.3. *The strong-weak resurgent symmetry of the spectral trace of local \mathbb{P}^2 , which we have presented here, fundamentally arises from the interplay of the Stokes constants, their generating functions in the form of q , \tilde{q} -series, and their L -functions. This is the starting point in a wider mathematical program that aims at linking the resurgent properties of q -series with the analytic number-theoretic properties of L -functions and makes contact with the notion of quantum modularity. We will present this new paradigm of resurgence in detail and full generality in the companion paper [36].*

As pointed out in [32, 68], the TS/ST correspondence for a toric CY threefold X shares many formal similarities with complex Chern–Simons theory on the complement of a hyperbolic knot \mathcal{K} in the three-sphere. In particular, the spectral determinant $\Xi_X(\kappa, \hbar)$, which is an entire function of the complex deformation parameter κ of the mirror \hat{X} , corresponds to the state integral, or Andersen–Kashaev invariant, of the knot $Z_{\mathcal{K}}(u, \mathbf{b}^2)$ [69, 70], which is an entire function of the holonomy u around the knot. Here, \mathbf{b}^2 is a complex coupling parameter. Thus, the spectral traces $Z_X(N, \hbar)$ are dual to the coefficients of $Z_{\mathcal{K}}(u, \mathbf{b}^2)$ in a Taylor expansion around $u = 0$. Importantly, the state integral can be expressed as a sum of products of holomorphic and anti-holomorphic blocks given by q , \tilde{q} -series [71, 72], where q and \tilde{q} are related to \mathbf{b}^2 by the same formulae in Eq. (40), and the blocks are exchanged under the transformation $\mathbf{b}^2 \mapsto -\mathbf{b}^{-2}$, which swaps q and \tilde{q} . In the TS/ST correspondence, a similar factorization property is observed in closed expressions for the fermionic spectral traces $Z_X(N, \hbar)$, although the holomorphic and anti-holomorphic blocks are generally different functions [32]. Yet, the resurgent study of the spectral trace of local \mathbb{P}^2 shows how such different functions are nonetheless deeply connected. Indeed, as we have already remarked in Section 2.3, the holomorphic block resurges from the perturbative expansion of the anti-holomorphic block, and vice versa. This statement assumes a concrete meaning in light of the proven, exact results on the generating functions of the weak and strong coupling Stokes constants assembled into the strong-weak resurgent symmetry. To sum up, we display a minimal scheme below.

$$\text{disc}_{\frac{\pi}{2}}\psi(\tau) = \sum_{n=1}^{\infty} R_n q^{n/3} \xleftrightarrow{\text{red arrows}} \text{disc}_{\frac{\pi}{2}}\phi(\hbar) = \sum_{n=1}^{\infty} S_n \tilde{q}^n \tag{84}$$

Here, the red arrows represent the diagonals joining the top-left and bottom-right vertices of the diagram in Eq. (83).

On the physics of the strong-weak resurgent symmetry

The strong-weak coupling duality in Eq. (21), which is at the heart of the TS/ST correspondence, allows us to describe the strongly coupled conventional topological string compactified on local \mathbb{P}^2 in terms of the semiclassical regime of the quantum-mechanical spectral problem defined by the quantization of its mirror curve. Analogously, the semiclassical limit in the topological string coupling constant determines the strong dynamics of the spectral theory. Namely, the strong regime of each theory is approachable through the weak regime of the other. We point out that our strong-weak symmetry relating the exact resurgent structures of the spectral trace of local \mathbb{P}^2 in the limits of $\hbar \rightarrow 0$ and $\hbar \rightarrow \infty$ is a mathematically precise realization of this intuition on the exchange of perturbative and non-perturbative contributions.

The discussion on the physical mechanism underlying the strong-weak resurgent symmetry can be pushed further. Indeed, recall the definition of the total grand potential $J(\mu, \hbar)$ in Eq. (20). As we have briefly described in Section 2, the worldsheet generating functional $J^{\text{WS}}(\mu, \hbar)$, which determines the perturbative expansion of $J(\mu, \hbar)$ in the weakly coupled limit $g_s \rightarrow 0$, encodes the non-perturbative contributions in \hbar from the conventional topological string partition function. At the same time, the WKB generating functional $J^{\text{WKB}}(\mu, \hbar)$, which determines the perturbative expansion of $J(\mu, \hbar)$ in the semiclassical limit $\hbar \rightarrow 0$ instead, contains the non-perturbative, exponentially small effects in g_s from the NS limit of the refined topological string partition function. By means of the TS/ST statement in Eq. (19) and the strong-weak coupling duality in Eq. (21), we can translate the above argument into the following. The non-perturbative \hbar -corrections to the semiclassical perturbative expansion of the fermionic spectral trace $Z(N, \hbar)$, $N \in \mathbb{Z}_{>0}$, in Eq. (28a), determine its perturbative expansion in the strong coupling regime $\hbar \rightarrow \infty$. Conversely, the non-perturbative \hbar^{-1} -corrections to the strongly coupled perturbative expansion of the fermionic spectral trace $Z(N, \hbar)$, $N \in \mathbb{Z}_{>0}$, in Eq. (28b), determine its semiclassical perturbative \hbar -expansion.

Take now $N = 1$ and consider the first fermionic spectral trace $Z(1, \hbar)$, as we have done in this paper. We know that the non-analytic, exponential-type contributions at strong and weak coupling are governed by two dual sequences of Stokes constants S_n, R_n , $n \in \mathbb{Z}_{\neq 0}$. Equivalently, we can repackage the same information in the corresponding generating functions $f_0(y), f_\infty(y)$, $y \in \mathbb{C} \setminus \mathbb{R}$, or the corresponding L -functions $L_0(s), L_\infty(s)$, $s \in \mathbb{C}$. Thus, the argument above implies that the Stokes constants/generating functions/ L -functions in one regime must dictate the perturbative coefficients in the other, which we have proven true. Yet, substantially more information is discovered by analytically determining the complete resurgent structures of the spectral trace of local \mathbb{P}^2 at strong and weak coupling, as we have summarized in the commutative diagram in Eq. (76) and will further detail in the rest of this work. Finally, we note that our results recall the traditional notion of S-duality in string theory [73]¹⁵.

Remark 3.4. *As the ideas above apply in full generality, our conceptual argument underlying the strong-weak resurgent symmetry of the spectral trace of local \mathbb{P}^2 supports the existence of an appropriate generalization of our results to other toric CY threefolds and higher-order fermionic spectral traces. The detailed exploration of the physical principles of the strong-weak resurgent symmetry will be the subject of a subsequent publication [62].*

¹⁵In the work of [73], it was shown that the S-duality of type IIB superstrings in ten dimensions implies the existence of an S-duality relating the A-model and B-model topological string theories on the same CY background. In particular, the D-instantons of one model correspond to the perturbative amplitudes of the other.

3.3 Functional equation from a unified perspective

Let us go back to the L -functions $L_0(s)$, $L_\infty(s)$, $s \in \mathbb{C}$, which are defined by the meromorphic continuation to the complex s -plane of the Dirichlet series with coefficients given by the weak and strong coupling Stokes constants S_n , R_n , $n \in \mathbb{Z}_{>0}$, respectively, as we have briefly recalled in Section 2.3. These weak and strong coupling L -functions satisfy the factorizations in Eqs. (59a) and (59b) in terms of the Dirichlet L -function $L(s, \chi_{3,2})$ and the Riemann zeta function $\zeta(s)$, whose meromorphic completions

$$\Lambda_{3,2}(s) = \frac{3^{\frac{s}{2}}}{\pi^{\frac{s+1}{2}}} \Gamma\left(\frac{s+1}{2}\right) L(s, \chi_{3,2}), \quad \Lambda_\zeta(s) = \frac{1}{\pi^{\frac{s}{2}}} \Gamma\left(\frac{s}{2}\right) \zeta(s), \quad s \in \mathbb{C}, \quad (85)$$

satisfy the well-known functional equations

$$\Lambda_{3,2}(s) = \Lambda_{3,2}(1-s), \quad \Lambda_\zeta(s) = \Lambda_\zeta(1-s), \quad (86)$$

which are centered at $s = 1/2$ and relate s with $1-s$. Following the factorization formulae in Eqs. (59a) and (59b) and using the completed L -functions in Eq. (85), we define

$$\Lambda_0(s) = \Lambda_{3,2}(s+1)\Lambda_\zeta(s) = \frac{3^{\frac{s+1}{2}}}{S_1 \pi^{s+1}} \Gamma\left(\frac{s}{2}\right) \Gamma\left(\frac{s}{2} + 1\right) L_0(s), \quad (87a)$$

$$\Lambda_\infty(s) = \Lambda_{3,2}(s)\Lambda_\zeta(s+1) = \frac{3^{\frac{s}{2}}}{R_1 \pi^{s+1}} \Gamma\left(\frac{s+1}{2}\right)^2 L_\infty(s), \quad (87b)$$

which are meromorphic functions for $s \in \mathbb{C}$. Observe, however, that they do not individually satisfy a functional equation of the form in Eq. (86). Indeed, these completions of the weak and strong coupling L -functions are analytically continued to the whole complex s -plane through each other. The following statement is a consequence of the remarkable symmetry that relates the factorizations of $L_0(s)$ and $L_\infty(s)$ via a unitary cross-shift in the arguments of the factors.

Theorem 3.3. *The weak and strong coupling completed L -functions $\Lambda_0(s)$ and $\Lambda_\infty(s)$, $s \in \mathbb{C}$, in Eqs. (87a) and (87b), respectively, satisfy the functional equation*

$$\Lambda_0(s) = \Lambda_\infty(-s). \quad (88)$$

Proof. Using the definitions in Eqs. (87a) and (87b) and the functional equations in Eq. (86), the statement follows from the explicit computation

$$\Lambda_0(1-s) = \Lambda_{3,2}(2-s)\Lambda_\zeta(1-s) = \Lambda_{3,2}(s-1)\Lambda_\zeta(s) = \Lambda_\infty(s-1), \quad (89)$$

after the change of variable $s \rightarrow 1-s$. \square

As the weak and strong coupling L -functions are naturally paired, their combined functional equation in Eq. (88) provides an additional two-headed arrow joining them directly in the global

commutative diagram in Eq. (76), which we explicitly reproduce below.¹⁶

$$\begin{array}{ccccc}
 & \text{disc}_{\frac{\pi}{2}} \psi(\tau) & \xrightarrow{h \rightarrow 0} & \phi(\hbar) & \\
 & \updownarrow \text{Mellin} & & \updownarrow \text{evaluation} & \\
 & \text{inverse Mellin} & & \text{exact large-}n & \\
 \begin{array}{c} \nearrow \text{generating series} \\ \text{Dirichlet series} \end{array} & \begin{array}{c} \{\eta_n, R_n\} \\ \xrightarrow{\text{Dirichlet series}} \end{array} & \{\mathcal{A}_\infty, L_\infty\} & \xleftarrow{\text{functional equation}} & \{\mathcal{A}_0, L_0\} & \xleftarrow{\text{Dirichlet series}} & \{\zeta_n, S_n\} \\
 & \updownarrow \text{Mellin} & & \updownarrow \text{inverse Mellin} & \\
 & \text{exact large-}n & & \text{evaluation} & \\
 \begin{array}{c} \nwarrow \text{resurgence} \\ \text{resurgence} \end{array} & \begin{array}{c} \psi(\tau) \\ \xleftarrow{\tau \rightarrow 0} \end{array} & \text{disc}_{\frac{\pi}{2}} \phi(\hbar) & \nwarrow \text{generating series} & \\
 & \updownarrow \text{Mellin} & & \updownarrow \text{evaluation} & \\
 & \text{inverse Mellin} & & \text{exact large-}n &
 \end{array} \quad (90)$$

3.4 Arithmetic twist of the Stokes constants

Recall that the weak and strong coupling Stokes constants $S_n, R_n, n \in \mathbb{Z}_{\neq 0}$, are explicitly given by the divisor sum functions in Eqs. (46) and (55), respectively. As we have already observed, an enticingly simple arithmetic symmetry relates them. Namely, the n -th Stokes constants in the two asymptotic limits are obtained from one another by exchanging the positive integer divisors d and n/d that multiply the Dirichlet character inside the sums. We can say something more. Let us start by introducing the notation

$$n = \text{sign}(n) \prod_{p \in \mathbb{P}} p^{n_p} \in \mathbb{Z}_{\neq 0}, \quad n_p \in \mathbb{N}, \quad (91)$$

where \mathbb{P} is the set of prime numbers. We then define the arithmetic function¹⁷

$$\mathfrak{f}(n) = 3^{n_3} \chi_{3,2}(n/3^{n_3}), \quad n \in \mathbb{Z}_{\neq 0}, \quad (92)$$

where $\chi_{3,2}(n)$ is the non-principal Dirichlet character modulo 3 in Eq. (47). In particular, $\mathfrak{f}(n) = \chi_{3,2}(n)$ when $3 \nmid n$. Note that $\mathfrak{f}(n)$ is completely multiplicative and satisfies $\mathfrak{f}(-n) = -\mathfrak{f}(n)$.

Proposition 3.4. *The normalized weak and strong coupling Stokes constants $S_n/S_1, R_n/R_1, n \in \mathbb{Z}_{\neq 0}$, in Eqs. (46) and (55) are related by the arithmetic twist*

$$\frac{S_n}{S_1} = \mathfrak{f}(n) \frac{R_n}{R_1}, \quad (93)$$

where $\mathfrak{f}(n)$ is the arithmetic function in Eq. (92).

Proof. Let us fix $n \in \mathbb{Z}_{\neq 0}$ and write $n = 3^{n_3}m$, where $n_3 \in \mathbb{N}$ and $m \in \mathbb{Z}_{\neq 0}$ with $3 \nmid m$. Using the closed formulae in Eqs. (46), (55), and (92), we find that

$$\mathfrak{f}(n) \frac{R_n}{R_1} = \sum_{d|n} \frac{d 3^{n_3}}{n} \chi_{3,2}(dm) = \sum_{d|m} \frac{d}{m} \chi_{3,2}(m/d) = \sum_{d|m} \frac{1}{d} \chi_{3,2}(d) = \frac{S_m}{S_1}, \quad (94)$$

¹⁶For completeness, we remind that the fundamental constants $\mathcal{A}_0, \mathcal{A}_\infty$ are related by $\mathcal{A}_0 = 2\pi\mathcal{A}_\infty$.

¹⁷The function in Eq. (92) can be interpreted as the Dirichlet convolution $\mathfrak{f}(n) = \sum_{d|n} f_3(d) \chi_{3,2}(n/d)$, where $f_3(n) = 3^{n_3}, n \in \mathbb{Z}_{\neq 0}$.

where we have applied the property

$$\chi_{3,2}(dm) = \chi_{3,2}(d)^2 \chi_{3,2}(m/d) = \chi_{3,2}(m/d), \quad (95)$$

which makes use of $\chi_{3,2}(d)^2 = 1$ for $3 \nmid d \mid m$. We conclude by substituting $S_m = S_n$. \square

The arithmetic twist above plays a role in the global net of relations among the dual resurgent structures of $\text{Tr}(\rho_{\mathbb{P}^2})$ that is analogous to the one played by the functional equation linking the two L -functions. Indeed, Proposition 3.4 supplies a two-headed arrow joining the weak and strong coupling Stokes constants directly in the commutative diagram in Eq. (76). We reproduce it below in an equivalent form that allows us to visualize the contribution from Eq. (93).¹⁸

$$\begin{array}{ccccc}
 & \text{disc}_{\frac{\pi}{2}} \psi(\tau) & \xrightarrow{\hbar \rightarrow 0} & \phi(\hbar) & \\
 & \uparrow \text{generating series} & & \downarrow \text{resurgence} & \\
 \{ \mathcal{A}_\infty, L_\infty \} & \xleftarrow{\text{Dirichlet series}} \{ \eta_n, R_n \} & \xleftrightarrow{\text{arithmetic twist}} & \{ \zeta_n, S_n \} & \xrightarrow{\text{Dirichlet series}} \{ \mathcal{A}_0, L_0 \} \\
 & \downarrow \text{evaluation} & & \downarrow \text{evaluation} & \\
 & \psi(\tau) & \xleftarrow{\tau \rightarrow 0} & \text{disc}_{\frac{\pi}{2}} \phi(\hbar) & \\
 & \uparrow \text{resurgence} & & \downarrow \text{generating series} & \\
 & \text{exact large-}n & & \text{exact large-}n & \\
 & \uparrow \text{Mellin} & & \downarrow \text{inverse Mellin} &
 \end{array} \quad (96)$$

L -functions and character twists

The arithmetic twist of the Stokes constants described in Eq. (93) appropriately translates in the language of the weak and strong coupling L -functions $L_0(s)$, $L_\infty(s)$, $s \in \mathbb{C}$, in Eqs. (59a) and (59b). In particular, Eq. (93) straightforwardly implies that

$$L_0(s) = \sum_{n=1}^{\infty} \frac{S_n}{n^s} = \frac{S_1}{R_1} \sum_{n=1}^{\infty} \frac{R_n}{n^s} \mathfrak{f}(n). \quad (97)$$

We can reformulate this arithmetic twist in a way that makes its meaning and implications more explicit. Before doing so, we introduce the principal Dirichlet character modulo 3, that is,

$$\chi_{3,1}(n) = \begin{cases} 0 & \text{if } n \equiv_3 0 \\ 1 & \text{else} \end{cases}, \quad n \in \mathbb{Z}, \quad (98)$$

which leads us to the following corollary to Proposition 3.4.

Corollary 3.5. *The normalized weak and strong coupling Stokes constants S_n/S_1 , R_n/R_1 , $n \in \mathbb{Z}_{\neq 0}$, in Eqs. (46) and (55) satisfy*

$$\chi_{3,1}(n) \frac{S_n}{S_1} = \chi_{3,2}(n) \frac{R_n}{R_1}, \quad (99)$$

where $\chi_{3,1}(n)$, $\chi_{3,2}(n)$ are the Dirichlet characters modulo 3 in Eqs. (47) and (98), respectively.

¹⁸For completeness, we remind that the locations of the singularities ζ_n , η_n , $n \in \mathbb{Z}_{\neq 0}$, are related by $\zeta_n = 2\pi\eta_n$.

Proof. Observe that Eq. (93) implies

$$\frac{S_m}{S_1} = \chi_{3,2}(m) \frac{R_m}{R_1}, \quad m \in \mathbb{Z}_{\neq 0}, \quad 3 \nmid m. \quad (100)$$

Multiplying both sides of Eq. (100) by the Dirichlet character $\chi_{3,1}(m)$ in Eq. (98) and applying

$$\chi_{3,1}(m)\chi_{3,2}(m) = \chi_{3,2}(m), \quad (101)$$

we obtain the desired statement. \square

Note that Corollary 3.5 can be equivalently stated as

$$\chi_{3,2}(n) \frac{S_n}{S_1} = \chi_{3,1}(n) \frac{R_n}{R_1}. \quad (102)$$

Indeed, multiplying both sides of Eq. (99) by $\chi_{3,2}(n)$ and using that $\chi_{3,2}(n)^2 = \chi_{3,1}(n)$, we find the expression in Eq. (102). Finally, let us introduce the following notation. We denote by $L(s, \chi)$ the L -function that results from twisting an arbitrary L -function $L(s)$, $s \in \mathbb{C}$, by an arbitrary Dirichlet character $\chi(n)$, $n \in \mathbb{Z}$. Namely,¹⁹

$$L(s, \chi) = \sum_{n=1}^{\infty} \frac{A_n}{n^s} \chi(n), \quad (103)$$

where the complex numbers A_n are the coefficients in the L -series representation of $L(s)$.

Theorem 3.6. *The weak and strong coupling L -functions $L_0(s)$, $L_{\infty}(s)$, $s \in \mathbb{C}$, in Eqs. (59a) and (59b) are obtained from each other by applying a character twist. Specifically, they satisfy*

$$L_0(s) = \frac{S_1}{R_1} \frac{1}{1 - 3^{-s}} L_{\infty}(s, \chi_{3,2}), \quad (104a)$$

$$L_{\infty}(s) = \frac{R_1}{S_1} \frac{1}{1 - 3^{-(s+1)}} L_0(s, \chi_{3,2}), \quad (104b)$$

where $\chi_{3,2}(m)$ is the unique non-principal Dirichlet character modulo 3 in Eq. (47).

Proof. We write an arbitrary integer $n \in \mathbb{Z}_{>0}$ in the form $n = 3^{n_3}m$, where $n_3 \in \mathbb{N}$ and $m \in \mathbb{Z}_{>0}$ with $3 \nmid m$, as before. As a straightforward consequence of their closed formulae as divisor sum functions, the weak and strong coupling Stokes constants satisfy the simple properties

$$S_n = S_m, \quad R_n = 3^{-n_3} R_m. \quad (105)$$

It follows from the definition of the weak coupling L -function in Eq. (59a) that it factorizes as

$$L_0(s) = \sum_{n_3=0}^{\infty} 3^{-sn_3} \sum_{\substack{m=1 \\ 3 \nmid m}}^{\infty} \frac{S_m}{m^s} = \frac{1}{1 - 3^{-s}} \sum_{\substack{m=1 \\ 3 \nmid m}}^{\infty} \frac{S_m}{m^s}, \quad (106)$$

¹⁹Note that the Dirichlet L -function $L(s, \chi_{3,2})$, which appears in the factorization of the weak and strong coupling L -functions in Eqs. (59a) and (59b), is obtained by twisting the Riemann zeta function $\zeta(s)$ by the Dirichlet character $\chi_{3,2}$ in Eq. (47).

where we have used the first formula in Eq. (105) and resummed the geometric series over the index n_3 . As a consequence of Corollary 3.5, the series in the RHS of Eq. (106) can then be written as

$$\sum_{\substack{m=1 \\ 3 \nmid m}}^{\infty} \frac{S_m}{m^s} = \sum_{n=1}^{\infty} \frac{S_n}{n^s} \chi_{3,1}(n) = \frac{S_1}{R_1} \sum_{n=1}^{\infty} \frac{R_n}{n^s} \chi_{3,2}(n). \quad (107)$$

Substituting Eq. (107) into Eq. (106) and using the notation introduced in Eq. (103) for $L_{\infty}(s)$ yields the statement in Eq. (104a). Analogously, the definition of the strong coupling L -function in Eq. (59b) implies the factorization

$$L_{\infty}(s) = \sum_{n_3=0}^{\infty} 3^{-(s+1)n_3} \sum_{\substack{m=1 \\ 3 \nmid m}}^{\infty} \frac{R_m}{m^s} = \frac{1}{1 - 3^{-(s+1)}} \sum_{\substack{m=1 \\ 3 \nmid m}}^{\infty} \frac{R_m}{m^s}, \quad (108)$$

where we have used the second formula in Eq. (105) and resummed the geometric series over the index n_3 . As a consequence of Corollary 3.5 in the equivalent form of Eq. (102), the series in the RHS of Eq. (108) can then be written as

$$\sum_{\substack{m=1 \\ 3 \nmid m}}^{\infty} \frac{R_m}{m^s} = \sum_{n=1}^{\infty} \frac{R_n}{n^s} \chi_{3,1}(n) = \frac{R_1}{S_1} \sum_{n=1}^{\infty} \frac{S_n}{n^s} \chi_{3,2}(n). \quad (109)$$

Substituting Eq. (109) into Eq. (108) and using the notation introduced in Eq. (103) for $L_0(s)$ yields the statement in Eq. (104b). \square

4 Summability and quantum modularity

In this section, we present our results on the summability and modularity properties of the generating functions f_0 , f_{∞} of the Stokes constants at weak and strong coupling, defined in Eqs. (61) and (65), respectively. In particular, we show how the median resummation allows us to effectively reconstruct $f_0(y)$ and (conjecturally) $f_{\infty}(y)$ from their asymptotic expansions in the limit $y \rightarrow 0$. Moreover, we prove that they are holomorphic quantum modular forms of weight zero under the congruence subgroup $\Gamma_1(3) \subset \mathrm{SL}_2(\mathbb{Z})$.

4.1 Borel–Laplace sums

As a preliminary step toward the summability study we perform in Section 4.2, we explicitly compute the Borel–Laplace sums of the asymptotic series $\phi(\hbar)$ and $\psi(\tau)$ in Eqs. (42) and (51) capturing the perturbative expansions of the spectral trace of local \mathbb{P}^2 in the limits $\hbar \rightarrow 0$ and $\hbar \rightarrow \infty$, respectively. We start by introducing the following notation. Let $\mathbf{e}_1: \mathbb{C} \setminus i\mathbb{R}_{\geq 0} \rightarrow \mathbb{C}$ be the function defined by²⁰

$$\mathbf{e}_1(y) := \frac{1}{2\pi i} \int_0^{\infty} e^{-2\pi t} \frac{dt}{t + iy}. \quad (110)$$

Note that, applying the simple change of coordinates $t \mapsto t/(2\pi) - iy$, the exponential integral $\mathbf{e}_1(y)$ can be written in the equivalent form

$$\mathbf{e}_1(y) = \frac{1}{2\pi i} e^{2\pi iy} \Gamma(0, 2\pi iy), \quad (111)$$

²⁰The notation in Eq. (110) was suggested by M. Kontsevich as the function \mathbf{e}_1 plays a similar role in other examples (e.g., the case of σ, σ^* in [74, from min. 21.55] and the case of even/odd Maass cusp forms in [36]).

where $\Gamma(s, 2\pi iy)$ denotes the upper incomplete gamma function. It follows from the analyticity properties of $\Gamma(0, 2\pi iy)$ that the function $\mathbf{e}_1(y)$ is analytic for $y \in \mathbb{C} \setminus i\mathbb{R}_{\geq 0}$ and its jump across the branch cut along the positive imaginary axis is given by

$$\lim_{\delta \rightarrow 0^+} \mathbf{e}_1(ix - \delta) - \mathbf{e}_1(ix + \delta) = e^{-2\pi x}, \quad (112)$$

where we have fixed $x \in \mathbb{R}_{\geq 0}$. Let us then go back to the Gevrey-1 asymptotic series $\phi(\hbar)$, $\psi(\tau)$ in Eqs. (42) and (51). As recalled in Section 2.3, their perturbative coefficients a_{2n} , b_{2n} , $n \in \mathbb{Z}_{>0}$, satisfy the exact large-order relations in Eq. (58), which can be straightforwardly written as

$$a_{2n} = \frac{\Gamma(2n)}{\pi i} \sum_{m=1}^{\infty} \frac{S_m}{\zeta_m^{2n}}, \quad (113a)$$

$$b_{2n} = \frac{\Gamma(2n-1)}{\pi i} \sum_{m=1}^{\infty} \frac{R_m}{\eta_m^{2n-1}}, \quad (113b)$$

where the Stokes constants S_m , R_m are given explicitly in Eqs. (46) and (55), while the locations ζ_m , η_m of the singularities in the complex ζ -plane are in Eqs. (44) and (53). We now prove that both $\phi(\hbar)$ and $\psi(\tau)$ are Borel–Laplace summable at the angles 0 and π and their Borel–Laplace sums can be expressed in terms of the exponential integral in Eq. (110).

Proposition 4.1. *The Borel–Laplace sum of the weak coupling perturbative series $\phi(\hbar)$, defined in Eq. (42), along the positive real axis is*

$$s_0(\phi)(\hbar) = - \sum_{m \in \mathbb{Z}_{\neq 0}} S_m \mathbf{e}_1 \left(-\frac{2\pi m}{3\hbar} \right), \quad (114)$$

which is analytic for $\Re(\hbar) > 0$.

Proof. As summarized in Appendix B, the Borel–Laplace resummation requires two successive steps. First, we compute the Borel transform $\hat{\phi}(\zeta)$, which gives

$$\begin{aligned} \hat{\phi}(\zeta) &= \sum_{n=1}^{\infty} \frac{a_{2n}}{\Gamma(2n)} \zeta^{2n-1} = \frac{1}{\pi i} \sum_{m=1}^{\infty} S_m \zeta^{-1} \sum_{n=1}^{\infty} \left(\frac{\zeta}{\zeta_m} \right)^{2n} = \frac{1}{\pi i} \sum_{m=1}^{\infty} S_m \frac{\zeta}{\zeta_m^2 - \zeta^2} = \\ &= -\frac{1}{2\pi i} \sum_{m=1}^{\infty} \frac{S_m}{\zeta_m + \zeta} + \frac{1}{2\pi i} \sum_{m=1}^{\infty} \frac{S_m}{\zeta_m - \zeta} = \frac{1}{2\pi i} \sum_{m \in \mathbb{Z}_{\neq 0}} \frac{S_m}{\zeta_m - \zeta}, \end{aligned} \quad (115)$$

where we have substituted Eq. (113a), applied the identity

$$\frac{1}{(x-y)(x+y)} = \frac{1}{2x(x-y)} + \frac{1}{2x(x+y)}, \quad (116)$$

and used the properties $\zeta_{-m} = -\zeta_m$ and $S_{-m} = S_m$, $m \in \mathbb{Z}_{\neq 0}$. Then, assuming $\hbar \in \mathbb{R}_{\geq 0}$, we perform the Laplace transform along the positive real axis, that is,

$$\begin{aligned} s_0(\phi)(\hbar) &= \int_0^{\infty} d\zeta e^{-\zeta/\hbar} \hat{\phi}(\zeta) = \frac{1}{2\pi i} \int_0^{\infty} d\zeta \sum_{m \in \mathbb{Z}_{\neq 0}} S_m \frac{e^{-\zeta/\hbar}}{\zeta_m - \zeta} = \\ &= \frac{1}{2\pi i} \int_0^{\infty} dt \sum_{m \in \mathbb{Z}_{\neq 0}} S_m \frac{e^{-2\pi t}}{\frac{\zeta_m}{2\pi\hbar} - t} = - \sum_{m \in \mathbb{Z}_{\neq 0}} S_m \mathbf{e}_1 \left(\frac{i\zeta_m}{2\pi\hbar} \right), \end{aligned} \quad (117)$$

where we have substituted Eq. (115), applied the change of variable $2\pi t = \zeta/\hbar$, permuted sum and integral due to absolute convergence, and used the notation introduced in Eq. (110). Finally, applying Eq. (44) in Eq. (117), we find the desired expression for the Borel–Laplace sum of $\phi(\hbar)$, which can be analytically continued to $\Re(\hbar) > 0$ thanks to the properties of the function \mathbf{e}_1 . \square

Proposition 4.2. *The Borel–Laplace sum of the strong coupling perturbative series $\psi(\tau)$, defined in Eq. (51), along the positive real axis is*

$$s_0(\psi)(\tau) = - \sum_{m \in \mathbb{Z}_{\neq 0}} R_m \mathbf{e}_1 \left(-\frac{m}{3\tau} \right), \quad (118)$$

which is analytic for $\Re(\tau) > 0$.

Proof. We follow the same steps of the proof of Proposition 4.1. First, we compute the Borel transform $\hat{\psi}(\tau)$ and obtain

$$\begin{aligned} \hat{\psi}(\tau) &= \sum_{n=1}^{\infty} \frac{b_{2n}}{\Gamma(2n-1)} \zeta^{2n-2} = \frac{1}{\pi i} \sum_{m=1}^{\infty} R_m \eta_m^{-1} \sum_{n=1}^{\infty} \left(\frac{\zeta}{\eta_m} \right)^{2n-2} = \frac{1}{\pi i} \sum_{m=1}^{\infty} R_m \frac{\eta_m}{\eta_m^2 - \zeta^2} = \\ &= \frac{1}{2\pi i} \sum_{m=1}^{\infty} \frac{R_m}{\eta_m + \zeta} + \frac{1}{2\pi i} \sum_{m=1}^{\infty} \frac{R_m}{\eta_m - \zeta} = \frac{1}{2\pi i} \sum_{m \in \mathbb{Z}_{\neq 0}} \frac{R_m}{\eta_m - \zeta}, \end{aligned} \quad (119)$$

where we have substituted Eq. (113b), applied the fractional identity in Eq. (116), and used the properties $\eta_{-m} = -\eta_m$ and $R_{-m} = -R_m$, $m \in \mathbb{Z}_{\neq 0}$. Then, assuming $\tau \in \mathbb{R}_{\geq 0}$, we compute the Laplace transform along the positive real axis, which gives

$$\begin{aligned} s_0(\psi)(\tau) &= \int_0^{\infty} d\zeta e^{-\zeta/\tau} \hat{\psi}(\zeta) = \frac{1}{2\pi i} \int_0^{\infty} d\zeta \sum_{m \in \mathbb{Z}_{\neq 0}} R_m e^{-\zeta/\tau} \frac{1}{\eta_m - \zeta} = \\ &= \frac{1}{2\pi i} \int_0^{\infty} dt \sum_{m \in \mathbb{Z}_{\neq 0}} R_m e^{-2\pi t} \frac{1}{\frac{\eta_m}{2\pi\tau} - t} = - \sum_{m \in \mathbb{Z}_{\neq 0}} R_m \mathbf{e}_1 \left(\frac{i\eta_m}{2\pi\tau} \right), \end{aligned} \quad (120)$$

where we have substituted Eq. (119), performed the change of variable $2\pi t = \zeta/\tau$, permuted sum and integral due to absolute convergence, and again applied the notation in Eq. (110). We conclude by substituting Eq. (53) into Eq. (120). Notice that $s_0(\psi)(\tau)$ can be analytically continued to $\Re(\tau) > 0$ thanks to the properties of the function \mathbf{e}_1 . \square

In addition, following the same steps in the proofs of Propositions 4.1 and 4.2, we obtain the analogous expressions for the Borel–Laplace sums of $\phi(\hbar)$ and $\psi(\tau)$ at angle π . Namely,

$$s_{\pi}(\phi)(\hbar) = - \sum_{m \in \mathbb{Z}_{\neq 0}} S_m \mathbf{e}_1 \left(-\frac{2\pi m}{3\hbar} \right), \quad (121a)$$

$$s_{\pi}(\psi)(\tau) = - \sum_{m \in \mathbb{Z}_{\neq 0}} R_m \mathbf{e}_1 \left(-\frac{m}{3\tau} \right), \quad (121b)$$

which are analytic for $\Re(\hbar) < 0$ and $\Re(\tau) < 0$, respectively.

Recall that the asymptotic behaviour in the limit $y \rightarrow 0$ of the generating functions $f_0(y)$ and $f_{\infty}(y)$ of the weak and strong coupling Stokes constants in Eqs. (61) and (65) is dictated by the original perturbative series $\psi(\tau)$ and $\phi(\hbar)$, respectively, according to the formulae in

Eqs. (72) and (74). Therefore, the Borel–Laplace sums of the asymptotic series $\tilde{f}_0(y)$ and $\tilde{f}_\infty(y)$ easily follow from Propositions 4.1 and 4.2. For completeness, we write their explicit formulae in the two corollaries below.

Corollary 4.3. *The Borel–Laplace sums of the perturbative expansion $\tilde{f}_0(y)$ in Eq. (71) along the positive and negative real axes are written in terms of the function \mathbf{e}_1 in Eq. (110) as*

$$s_0(\tilde{f}_0)(y) = -\frac{\pi i}{2} - \frac{3\mathcal{V}}{2\pi i y} + 2 \sum_{m \in \mathbb{Z}_{\neq 0}} R_m \mathbf{e}_1 \left(-\frac{m}{3y} \right), \quad \Re(y) > 0, \quad (122a)$$

$$s_\pi(\tilde{f}_0)(y) = -\frac{\pi i}{2} - \frac{3\mathcal{V}}{2\pi i y} + 2 \sum_{m \in \mathbb{Z}_{\neq 0}} R_m \mathbf{e}_1 \left(-\frac{m}{3y} \right), \quad \Re(y) < 0, \quad (122b)$$

where $\mathcal{V} = 2\Im(\text{Li}_2(e^{2\pi i/3}))$ as before.

Proof. The proof of the statement follows directly from Eq. (72) and Proposition 4.2. \square

Corollary 4.4. *The Borel–Laplace sums of the perturbative expansion $\tilde{f}_\infty(y)$ in Eq. (73) along the positive and negative real axes are written in terms of the function \mathbf{e}_1 in Eq. (110) as*

$$s_0(\tilde{f}_\infty)(y) = -3 \log \frac{\Gamma(2/3)}{\Gamma(1/3)} - \log(-6\pi i y) - 2 \sum_{m \in \mathbb{Z}_{\neq 0}} S_m \mathbf{e}_1 \left(-\frac{m}{3y} \right), \quad \Re(y) > 0, \quad (123a)$$

$$s_\pi(\tilde{f}_\infty)(y) = -3 \log \frac{\Gamma(2/3)}{\Gamma(1/3)} - \log(-6\pi i y) - 2 \sum_{m \in \mathbb{Z}_{\neq 0}} S_m \mathbf{e}_1 \left(-\frac{m}{3y} \right), \quad \Re(y) < 0. \quad (123b)$$

Proof. The proof of the statement follows directly from Eq. (74) and Proposition 4.1. \square

4.2 Median resummation

We will now show that the median resummation of the asymptotic expansion $\tilde{f}_0(y)$ of the generating function of the weak coupling Stokes constants S_n , $n \in \mathbb{Z}_{\neq 0}$, appropriately reconstructs the generating function $f_0(y)$ itself. As a consequence, our results showcase the effectiveness of the median resummation as a summability tool. We begin with the preliminary Lemma 4.5, where we prove that the Borel–Laplace sums at angles 0 and π of the asymptotic expansion $\tilde{f}_0(y)$ in Eq. (71) reproduce the original function $f_0(y)$ with a correction that is suitably encoded in the dual function $f_\infty(y)$.

Lemma 4.5. *Let $\tilde{f}_0(y)$ be the asymptotic expansion in the limit $y \rightarrow 0$ of the generating function $f_0(y)$ in Eq. (61) with $\Im(y) > 0$, which we have written explicitly in Eq. (71). Its Borel–Laplace sums along the positive and negative real axes can be expressed as*

$$s_0(\tilde{f}_0)(y) = f_0(y) + f_\infty \left(-\frac{1}{3y} \right), \quad \Re(y) > 0, \quad (124a)$$

$$s_\pi(\tilde{f}_0)(y) = f_0(y) - f_\infty \left(-\frac{1}{3y} \right), \quad \Re(y) < 0. \quad (124b)$$

Proof. Applying Eq. (193a) to expand the two quantum dilogarithms in Eq. (63) separately, $\tilde{f}_0(y)$ can be written as

$$\begin{aligned} \frac{1}{3}\tilde{f}_0(y) = & -\frac{\pi i}{3} - \frac{1}{2} (\log(1-w) - \log(1-w^{-1})) \\ & - \frac{1}{2\pi i y} (\text{Li}_2(w) - \text{Li}_2(w^{-1})) - (\tilde{\varphi}_+(y) - \tilde{\varphi}_-(y)), \end{aligned} \quad (125)$$

where $w = e^{2\pi i/3}$ as before and we have introduced the two formal power series

$$\tilde{\varphi}_{\pm}(y) := \sum_{k=1}^{\infty} (2\pi i y)^{2k-1} \frac{B_{2k}}{(2k)!} \text{Li}_{2-2k}(w^{\pm 1}). \quad (126)$$

Let us compute the Borel transform of $\tilde{\varphi}_{\pm}(y)$ in the variable ζ conjugate to y . We find that

$$\begin{aligned} \mathcal{B}[\tilde{\varphi}_{\pm}](\zeta) &= \sum_{k=1}^{\infty} (2\pi i)^{2k-1} \frac{B_{2k}}{(2k)!} \text{Li}_{2-2k}(w^{\pm 1}) \frac{\zeta^{2k-2}}{(2k-2)!} \\ &= \left(\sum_{k=1}^{\infty} \frac{B_{2k}}{(2k)!} \zeta^{2k-2} \right) \diamond \left(\sum_{k=1}^{\infty} (2\pi i)^{2k-1} \frac{\text{Li}_{2-2k}(w^{\pm 1})}{(2k-2)!} \zeta^{2k-2} \right) =: \tilde{g}(\zeta) \diamond \tilde{f}_{\pm}(\zeta), \end{aligned} \quad (127)$$

where \diamond denotes the Hadamard product [75]. The formal power series $\tilde{g}(\zeta)$, $\tilde{f}_{\pm}(\zeta)$ have finite radius of convergence at $\zeta = 0$ and can be resummed explicitly into the functions²¹

$$g(\zeta) = -\frac{1}{\zeta^2} + \frac{1}{2\zeta} \coth\left(\frac{\zeta}{2}\right), \quad |\zeta| < 2\pi, \quad (128a)$$

$$f_{\pm}(\zeta) = -\pi i \left(\frac{1}{1 + e^{\pm \frac{\pi i}{3} - 2\pi i \zeta}} + \frac{1}{1 + e^{\pm \frac{\pi i}{3} + 2\pi i \zeta}} \right), \quad |\zeta| < \frac{1}{3}, \quad (128b)$$

respectively. After being analytically continued to the whole complex ζ -plane, the function $g(\zeta)$ has poles of order one along the imaginary axis at

$$\mu_m = 2\pi i m, \quad m \in \mathbb{Z}_{\neq 0}, \quad (129)$$

while the functions $f_{\pm}(\zeta)$ have poles of order one along the real axis at

$$\nu_k^{(+,1)} = \frac{1}{3} + k, \quad \nu_k^{(+,2)} = -\frac{1}{3} + 2k, \quad \nu_k^{(+,3)} = \frac{2}{3} + 2k, \quad (130a)$$

$$\nu_k^{(-,1)} = \frac{1}{3} + 2k, \quad \nu_k^{(-,2)} = -\frac{1}{3} + 2k, \quad \nu_k^{(-,3)} = \frac{2}{3} + 2k, \quad \nu_k^{(-,4)} = -\frac{2}{3} + 2k, \quad (130b)$$

for $k \in \mathbb{Z}$, respectively. We consider a circle γ in the complex s -plane with center $s = 0$ and radius $0 < r < 2\pi$ and apply Hadamard's multiplication theorem. The Borel transform of $\tilde{\varphi}_{\pm}(y)$ can be written as the integral

$$\begin{aligned} \mathcal{B}[\tilde{\varphi}_{\pm}](\zeta) &= \frac{1}{2\pi i} \int_{\gamma} g(s) f_{\pm}\left(\frac{\zeta}{s}\right) \frac{ds}{s} \\ &= \frac{1}{2} \int_{\gamma} \left(\frac{1}{s} - \frac{1}{2} \coth\left(\frac{s}{2}\right) \right) \left(\frac{1}{1 + e^{\pm \frac{\pi i}{3} - 2\pi i \frac{\zeta}{s}}} + \frac{1}{1 + e^{\pm \frac{\pi i}{3} + 2\pi i \frac{\zeta}{s}}} \right) \frac{ds}{s^2}, \end{aligned} \quad (131)$$

for $|\zeta| < r/3$. For such values of ζ , the function $s \mapsto f_{\pm}(\zeta/s)$ has singular points at $s = \zeta/\nu_k^{(\pm,i)}$, $k \in \mathbb{Z}$, which sit inside the contour of integration γ and accumulate at the origin, and no singularities for $|s| > r$. The function $g(s)$ has simple poles at the points $s = \mu_m$ with residues

$$\text{Res}_{s=\pm 2\pi i m} g(s) = \pm \frac{1}{2\pi i}, \quad m \in \mathbb{Z}_{>0}. \quad (132)$$

²¹We impose that $g(0) = 1/12$ in order to eliminate the removable singularity of $g(\zeta)$ at the origin.

By Cauchy's residue theorem, the integral in Eq. (131) can be evaluated by summing the residues at the poles of the integrand which lie outside γ , allowing us to express the Borel transform as an exact function of ζ . More precisely, we find that

$$\begin{aligned}\mathcal{B}[\tilde{\varphi}_{\pm}](\zeta) &= - \sum_{m \in \mathbb{Z} \setminus \{0\}} \text{Res}_{s=2\pi i m} \left[g(s) f_{\pm} \left(\frac{\zeta}{s} \right) \frac{1}{s} \right] \\ &= \sum_{m \in \mathbb{Z} \setminus \{0\}} \frac{1}{4\pi i m^2} \left(\frac{1}{1 + e^{\pm \frac{\pi i}{3} - \frac{\zeta}{m}}} + \frac{1}{1 + e^{\pm \frac{\pi i}{3} + \frac{\zeta}{m}}} \right) \\ &= \frac{1}{2\pi i} \sum_{m=1}^{\infty} \frac{1}{m^2} \left(\frac{1}{1 + e^{\pm \frac{\pi i}{3} - \frac{\zeta}{m}}} + \frac{1}{1 + e^{\pm \frac{\pi i}{3} + \frac{\zeta}{m}}} \right),\end{aligned}\tag{133}$$

which is a well-defined, exact function of ζ for $|\zeta| < 2\pi/3$. In fact, the Borel transform in Eq. (133) converges when $\frac{\zeta}{2\pi i} \neq \frac{m}{3} + n$ and $\frac{\zeta}{2\pi i} \neq \frac{2m}{3} + n$ with $m \in \mathbb{Z} \setminus \{0\}$ and $n \in \mathbb{Z}$, respectively.²²

We can now compute the Laplace transform of $\mathcal{B}[\tilde{\varphi}_{\pm}](\zeta)$ along the positive and negative real axes. When $\Re(y), \Im(y) > 0$, we have that

$$\begin{aligned}s_0(\tilde{\varphi}_{\pm})(y) &= \frac{1}{2\pi i} \int_0^{\infty} e^{-\zeta/y} \left[\sum_{m=1}^{\infty} \frac{1}{m^2} \left(\frac{1}{1 + e^{\pm \frac{\pi i}{3} - \frac{\zeta}{m}}} + \frac{1}{1 + e^{\pm \frac{\pi i}{3} + \frac{\zeta}{m}}} \right) \right] d\zeta \\ &= \frac{1}{2\pi i} \int_0^{\infty} \left(\sum_{m=1}^{\infty} \frac{e^{-mt/y}}{m} \right) \left(\frac{1}{1 + e^{\pm \frac{\pi i}{3} - t}} + \frac{1}{1 + e^{\pm \frac{\pi i}{3} + t}} \right) dt,\end{aligned}\tag{134}$$

where we have applied the change of variable $\zeta = mt$. Resumming the series in m , we find that

$$\begin{aligned}s_0(\tilde{\varphi}_{\pm})(y) &= -\frac{1}{2\pi i} \int_0^{\infty} \log(1 - e^{-t/y}) \left(\frac{1}{1 + e^{\pm \frac{\pi i}{3} - t}} + \frac{1}{1 + e^{\pm \frac{\pi i}{3} + t}} \right) dt \\ &= -\frac{1}{2\pi i} \int_{-\infty}^{\infty} \frac{\log(1 - e^{-t/y})}{1 + e^{\pm \frac{\pi i}{3} - t}} dt + \frac{1}{2\pi i} \int_0^{-\infty} \frac{\log(-e^{t/y})}{1 + e^{\pm \frac{\pi i}{3} - t}} dt,\end{aligned}\tag{135}$$

where we have divided the integral into the sum of two contributions for simplicity. Indeed, we observe that the second term in the RHS of Eq. (135) gives

$$\frac{1}{2\pi i} \int_0^{-\infty} \frac{\log(-e^{t/y})}{1 + e^{\pm \frac{\pi i}{3} - t}} dt = -\frac{\text{Li}_2(w^{\pm 1})}{2\pi i y} - \frac{1}{2} \log(1 - w^{\pm 1}),\tag{136}$$

while the first term gives

$$-\frac{1}{2\pi i} \int_{-\infty}^{\infty} \frac{\log(1 - e^{-t/y})}{1 + e^{\pm \frac{\pi i}{3} - t}} dt = \log \frac{(w; e^{2\pi i y})_{\infty}}{(e^{-4\pi i/(3y)}; e^{-2\pi i/y})_{\infty}},\tag{137a}$$

$$-\frac{1}{2\pi i} \int_{-\infty}^{\infty} \frac{\log(1 - e^{-t/y})}{1 + e^{-\frac{\pi i}{3} - t}} dt = \log \frac{(w^{-1}; e^{2\pi i y})_{\infty}}{(e^{-2\pi i/(3y)}; e^{-2\pi i/y})_{\infty}}.\tag{137b}$$

The last two formulae follow from [76, Theorem A-29] as shown in [77, Eq. (5.108)].²³ Therefore, substituting Eqs. (136), (137a), and (137b) into the formula for the Borel–Laplace sum in

²²The convergence of the infinite sum in the RHS of Eq. (133) can be easily verified by, e.g., the limit comparison test. Indeed, where defined, the generic term of the series is dominated by $1/m^2$.

²³We thank C. Wheeler for sharing the preliminary version of his book in preparation [77].

Eq. (135), we find that

$$s_0(\tilde{\varphi}_+)(y) = -\frac{\text{Li}_2(w)}{2\pi iy} - \frac{1}{2}\log(1-w) + \log \frac{(w; e^{2\pi iy})_\infty}{(e^{-4\pi i/(3y)}; e^{-2\pi i/y})_\infty}, \quad (138a)$$

$$s_0(\tilde{\varphi}_-)(y) = -\frac{\text{Li}_2(w^{-1})}{2\pi iy} - \frac{1}{2}\log(1-w^{-1}) + \log \frac{(w^{-1}; e^{2\pi iy})_\infty}{(e^{-2\pi i/(3y)}; e^{-2\pi i/y})_\infty}. \quad (138b)$$

Now, it follows directly from Eqs. (125), (138a), and (138a) that

$$\begin{aligned} s_0(\tilde{f}_0)(y) &= -\pi i - 3 \log \frac{(w; e^{2\pi iy})_\infty}{(e^{-4\pi i/(3y)}; e^{-2\pi i/y})_\infty} + 3 \log \frac{(w^{-1}; e^{2\pi iy})_\infty}{(e^{-2\pi i/(3y)}; e^{-2\pi i/y})_\infty} \\ &= f_0(y) + f_\infty(-1/3y), \end{aligned} \quad (139)$$

where we have applied the closed formulae for the generating functions in Eqs. (63) and (67).

Similarly, when $\Re(y) < 0$ and $\Im(y) > 0$, the Borel-Laplace sum of the formal power series $\tilde{\varphi}_\pm(y)$ along the negative real axis is given by

$$\begin{aligned} s_\pi(\tilde{\varphi}_\pm)(y) &= \frac{1}{2\pi i} \int_0^{-\infty} e^{-\zeta/y} \left[\sum_{m=1}^{\infty} \frac{1}{m^2} \left(\frac{1}{1 + e^{\pm \frac{\pi i}{3} - \frac{\zeta}{m}}} + \frac{1}{1 + e^{\pm \frac{\pi i}{3} + \frac{\zeta}{m}}} \right) \right] d\zeta \\ &= \frac{1}{2\pi i} \int_0^{-\infty} \left(\sum_{m=1}^{\infty} \frac{e^{-mt/y}}{m} \right) \left(\frac{1}{1 + e^{\pm \frac{\pi i}{3} - t}} + \frac{1}{1 + e^{\pm \frac{\pi i}{3} + t}} \right) dt, \end{aligned} \quad (140)$$

where we have applied the change of variable $\zeta = mt$. Resumming the series in m , we find that

$$\begin{aligned} s_\pi(\tilde{\varphi}_\pm)(y) &= -\frac{1}{2\pi i} \int_0^{-\infty} \log(1 - e^{-t/y}) \left(\frac{1}{1 + e^{\pm \frac{\pi i}{3} - t}} + \frac{1}{1 + e^{\pm \frac{\pi i}{3} + t}} \right) dt \\ &= \frac{1}{2\pi i} \int_0^{-\infty} \log(1 - e^{t/y}) \left(\frac{1}{1 + e^{\pm \frac{\pi i}{3} + t}} + \frac{1}{1 + e^{\pm \frac{\pi i}{3} - t}} \right) dt \\ &= \frac{1}{2\pi i} \int_{-\infty}^{\infty} \frac{\log(1 - e^{t/y})}{1 + e^{\pm \frac{\pi i}{3} - t}} dt - \frac{1}{2\pi i} \int_0^{-\infty} \frac{\log(-e^{-t/y})}{1 + e^{\pm \frac{\pi i}{3} - t}} dt, \end{aligned} \quad (141)$$

where we have divided the integral into the sum of two contributions for simplicity. Let us now set $x = -1/y$, so that $\Re(x) > 0$ and $\Im(x) > 0$, then the second term in the RHS of Eq. (141) becomes

$$-\frac{1}{2\pi i} \int_0^{-\infty} \frac{\log(-e^{tx})}{1 + e^{\pm \frac{\pi i}{3} - t}} dt = \frac{x}{2\pi i} \text{Li}_2(w^{\pm 1}) + \frac{1}{2} \log(1 - w^{\pm 1}), \quad (142)$$

while the first term gives

$$\frac{1}{2\pi i} \int_{-\infty}^{\infty} \frac{\log(1 - e^{-tx})}{1 + e^{\frac{\pi i}{3} - t}} dt = -\log \frac{(w; e^{2\pi i/x})_\infty}{(e^{-4\pi ix/3}; e^{-2\pi ix})_\infty}, \quad (143a)$$

$$\frac{1}{2\pi i} \int_{-\infty}^{\infty} \frac{\log(1 - e^{-tx})}{1 + e^{-\frac{\pi i}{3} - t}} dt = -\log \frac{(w^{-1}; e^{2\pi i/x})_\infty}{(e^{-2\pi ix/3}; e^{-2\pi ix})_\infty}, \quad (143b)$$

following [77, Eq. (5.108)]. Therefore, substituting Eqs. (142), (143a), and (143b), after we change back x to $-1/y$, into the formula for the Borel–Laplace sum in Eq. (141), we find that

$$s_\pi(\tilde{\varphi}_+)(y) = -\frac{\text{Li}_2(w)}{2\pi iy} + \frac{1}{2} \log(1-w) - \log \frac{(w; e^{-2\pi iy})_\infty}{(e^{4\pi i/(3y)}; e^{2\pi i/y})_\infty}, \quad (144a)$$

$$s_\pi(\tilde{\varphi}_-)(y) = -\frac{\text{Li}_2(w^{-1})}{2\pi iy} + \frac{1}{2} \log(1-w^{-1}) - \log \frac{(w^{-1}; e^{-2\pi iy})_\infty}{(e^{2\pi i/(3y)}; e^{2\pi i/y})_\infty}. \quad (144b)$$

Putting together Eqs. (125), (144a), and (144b) yields

$$\begin{aligned} s_\pi(\tilde{f}_0)(y) &= 3 \log \frac{(w; e^{-2\pi iy})_\infty}{(e^{4\pi i/(3y)}; e^{2\pi i/y})_\infty} - 3 \log \frac{(w^{-1}; e^{-2\pi iy})_\infty}{(e^{2\pi i/(3y)}; e^{2\pi i/y})_\infty} \\ &= -3 \log \frac{(w e^{2\pi iy}; e^{2\pi iy})_\infty}{(w^{-1} e^{2\pi iy}; e^{2\pi iy})_\infty} - 3 \log \frac{(e^{4\pi i/(3y)}; e^{2\pi i/y})_\infty}{(e^{2\pi i/(3y)}; e^{2\pi i/y})_\infty}, \end{aligned} \quad (145)$$

where we have applied the relation in Eq. (177). Let us now recall that

$$(w^{\pm 1} e^{2\pi iy}; e^{2\pi iy})_\infty = (w^{\pm 1}; e^{2\pi iy})_\infty - (1 - w^{\pm 1}), \quad (146)$$

which we substitute into Eq. (145) to get

$$\begin{aligned} s_\pi(\tilde{f}_0)(y) &= -\pi i - 3 \log \frac{(w; e^{2\pi iy})_\infty}{(w^{-1}; e^{2\pi iy})_\infty} - 3 \log \frac{(e^{4\pi i/(3y)}; e^{2\pi i/y})_\infty}{(e^{2\pi i/(3y)}; e^{2\pi i/y})_\infty} \\ &= f_0(y) - f_\infty(1/3y) \\ &= f_0(y) - f_\infty(-1/3y), \end{aligned} \quad (147)$$

where we have first applied the closed formulae for the generating functions in Eqs. (63) and (67) and then used the parity property in Eq. (66).²⁴ \square

We can now prove that the median resummation of the asymptotic expansion $\tilde{f}_0(y)$ reproduces the generating function $f_0(y)$ itself.

Theorem 4.6. *Let $\tilde{f}_0(y)$ be the asymptotic expansion in the limit $y \rightarrow 0$ of the generating function $f_0(y)$ in Eq. (61) with $\Im(y) > 0$, which we have written explicitly in Eq. (71). Then, the median resummation of $\tilde{f}_0(y)$ reconstructs the original function $f_0(y)$, that is,*

$$\mathcal{S}_{\frac{\pi}{2}}^{\text{med}} \tilde{f}_0(y) = f_0(y), \quad y \in \mathbb{H}. \quad (148)$$

Proof. Following the definition in Eq. (215), the median resummation of $\tilde{f}_0(y)$ along the positive imaginary axis is given by

$$\mathcal{S}_{\frac{\pi}{2}}^{\text{med}} \tilde{f}_0(y) = \begin{cases} s_0(\tilde{f}_0)(y) + \frac{1}{2} \text{disc}_{\frac{\pi}{2}} \tilde{f}_0(y), & \Re(y) > 0, \\ s_\pi(\tilde{f}_0)(y) - \frac{1}{2} \text{disc}_{\frac{\pi}{2}} \tilde{f}_0(y), & \Re(y) < 0, \end{cases} \quad (149)$$

²⁴We note that the proof in the case of $\Re(y) > 0$ can be adapted to allow for an arbitrary choice of $w = \exp(z)$, $z \in \mathbb{C}^*$. However, the proof in the case of $\Re(y) < 0$ makes explicit use of the symmetry properties of the Stokes constants and their generating functions, which cannot be straightforwardly generalized.

where we take $\Im(y) > 0$. As a consequence of Eqs. (68) and (72), we have that

$$f_\infty\left(-\frac{1}{3y}\right) = -\frac{1}{2}\text{disc}_{\frac{\pi}{2}}\tilde{f}_0(y), \quad (150)$$

and substituting this in Eq. (149) yields

$$\mathcal{S}_{\frac{\pi}{2}}^{\text{med}}\tilde{f}_0(y) = \begin{cases} s_0(\tilde{f}_0)(y) - f_\infty\left(-\frac{1}{3y}\right), & \Re(y) > 0, \\ s_\pi(\tilde{f}_0)(y) + f_\infty\left(-\frac{1}{3y}\right), & \Re(y) < 0, \end{cases} \quad (151)$$

where again $\Im(y) > 0$. Then, the conclusion follows from Lemma 4.5. \square

Let us now consider the strong coupling asymptotic expansion $\tilde{f}_\infty(y)$ in Eq. (73). Despite its similarities with the dual formal series $\tilde{f}_0(y)$ in Eq. (71), the analogue to the analytic argument of Lemma 4.5 is still missing. Therefore, relying on the support of numerical tests, we limit ourselves to conjecture the effectiveness of the median resummation for the generating functions of the strong coupling Stokes constants.

Conjecture 1. *Let $\tilde{f}_\infty(y)$ be the asymptotic expansion in the limit $y \rightarrow 0$ of the generating function $f_\infty(y)$ in Eq. (65) with $\Im(y) > 0$, which we have written explicitly in Eq. (73). Then, the median resummation of $\tilde{f}_\infty(y)$ reconstructs the original function $f_\infty(y)$, that is,*

$$\mathcal{S}_{\frac{\pi}{2}}^{\text{med}}\tilde{f}_\infty(y) = f_\infty(y), \quad y \in \mathbb{H}. \quad (152)$$

Remark 4.1. *As a consequence Eqs. (64) and (74), we have that*

$$f_0\left(-\frac{1}{3y}\right) = \frac{1}{2}\text{disc}_{\frac{\pi}{2}}\tilde{f}_\infty(y), \quad y \in \mathbb{H}. \quad (153)$$

Substituting this into the definition of the median resummation in Eq. (215) for the asymptotic expansion $\tilde{f}_\infty(y)$ at the angle $\pi i/2$, Conjecture 1 straightforwardly implies that the Borel–Laplace sums along the positive and negative real axes of $\tilde{f}_\infty(y)$ reproduce the original function $f_\infty(y)$ with a correction that is suitably encoded in the dual function $f_0(y)$, and viceversa. More precisely,

$$s_0(\tilde{f}_\infty)(y) = f_\infty(y) - f_0\left(-\frac{1}{3y}\right), \quad \Re(y) > 0, \quad (154a)$$

$$s_\pi(\tilde{f}_\infty)(y) = f_\infty(y) + f_0\left(-\frac{1}{3y}\right), \quad \Re(y) < 0, \quad (154b)$$

which represent the dual statement to Lemma 4.5, are equivalent to Eq. (152).

Theorem 4.6 and Conjecture 1 can be placed within the context of the strong-weak resurgent symmetry discussed in Section 3.2 as they provide additional arrows in the commutative diagram in Eq. (83). In particular, taking the median resummation of the asymptotic expansions represents the formal inverse of perturbatively expanding the generating functions. We show

below the resulting completed diagram in terms of f_0 , f_∞ and their asymptotic expansions.

$$\begin{array}{ccc}
 f_\infty(y) & \xrightleftharpoons[\text{median resummation}]{y \rightarrow 0} & \tilde{f}_\infty(y) \\
 \uparrow \text{strong-weak symmetry} & & \uparrow \text{strong-weak symmetry} \\
 \tilde{f}_0(y) & \xrightleftharpoons[y \rightarrow 0]{\text{median resummation}} & f_0(y)
 \end{array} \tag{155}$$

Note that all four arrows tracing the edges of the box are now invertible. We display Conjecture 1 with a dashed arrow to distinguish it from the proven results that compose the rest of the diagram.

Remark 4.2. *The full content of the q , \tilde{q} -series in the block factorization of the spectral trace of local \mathbb{P}^2 cannot be reconstructed by a simple Borel–Laplace resummation. Yet, and remarkably, the missing information is not lost. Instead, it is collected by the discontinuities that are then recovered by the median resummation. The effectiveness of the median resummation as a summability tool is known in quantum Chern–Simons theory, as it was conjectured for the quantum invariants of knots and 3-manifolds in [78, Conjecture 1.1] and proven for the special cases of the trefoil knot [78] and the Seifert fibered homology sphere [79]. We expect more evidence to be discovered. Furthermore, as we discuss in the companion paper [36], the effectiveness of the median resummation appears to be closely related to quantum modularity.*

4.3 Quantum modularity of the generating functions

We will now present our results on the quantum modularity properties of the generating functions of the Stokes constants in both \hbar -regimes, that is, the periodic holomorphic functions $f_0(y)$ and $f_\infty(y)$ defined in Eqs. (61) and (65). Let us start our discussion by recalling the definition of a quantum modular form. We refer to [76, Chapter IV] for an overview. A first description of quantum modular forms was given by Zagier in [80]. Namely, a function $f: \mathbb{Q} \setminus \{0\} \rightarrow \mathbb{C}$ is called a weight- ω quantum modular form with respect to a subgroup $\Gamma \subseteq \mathrm{SL}_2(\mathbb{Z})$, where ω is a fixed integer or half-integer, if the cocycle

$$h_\gamma[f](y) := (cy + d)^{-\omega} f\left(\frac{ay + b}{cy + d}\right) - f(y) \tag{156}$$

is *better behaved* than $f(y)$ for all choices of $\gamma = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \Gamma$. Conceptually, being better behaved means having better analyticity properties than the function $f(y)$ itself – for instance, being a real analytic function over $\mathbb{R} \setminus \{0, -d/c\}$. Quantum modular forms of weight zero are called quantum modular functions. Some examples of quantum modular forms are built from the usual holomorphic modular forms. In these cases, the cocycle in Eq. (156) is expressed in terms of the underlying modular forms (see [80, Examples 0 to 4] and [81–83]). Besides, sometimes, the function f is defined and analytic in the upper half-plane \mathbb{H} , implying that it already has good analyticity properties. In this case, the cocycle $h_\gamma[f]$ in Eq. (156) is requested to be analytic

in some domain that is larger than \mathbb{H} – for instance, in $\mathbb{C}' = \mathbb{C} \setminus \mathbb{R}_{\leq 0}$. The quantum modular forms in this class are called *holomorphic quantum modular forms*. The following definition was proposed by Zagier [84, min 21:45].

Definition 4.1. *A function $f: \mathbb{H} \rightarrow \mathbb{C}$ is a weight- ω holomorphic quantum modular form for a subgroup $\Gamma \subseteq \mathrm{SL}_2(\mathbb{Z})$, where ω is integer or half-integer, if the cocycle $h_\gamma[f]: \mathbb{H} \rightarrow \mathbb{C}$ in Eq. (156) extends to a holomorphic function in $\mathbb{C}_\gamma := \{y \in \mathbb{C}: cy + d \in \mathbb{C}'\}$ ²⁵ for every $\gamma = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \Gamma$.*

Note that an analogous definition of holomorphic quantum modular form can be written for functions $f: \mathbb{H}_- \rightarrow \mathbb{C}$, where \mathbb{H}_- denotes the lower half of the complex plane. Remarkably, the resurgent analysis of the spectral trace of local \mathbb{P}^2 in both weak and strong coupling regimes is a new source of examples of holomorphic quantum modular forms. As we will show in Theorem 4.7, the generating functions of the Stokes constants are holomorphic quantum modular functions for the congruence subgroup $\Gamma_1(3)$, which we have defined in Eq. (8).

Let us start by defining the holomorphic functions $F_R, F_S: \mathbb{H} \rightarrow \mathbb{C}$ as

$$F_R(y) := f_\infty\left(-\frac{1}{3y}\right) + f_0(y), \quad (157a)$$

$$F_S(y) := f_\infty(y) - f_0\left(-\frac{1}{3y}\right). \quad (157b)$$

Note that the function $F_R(y)$ is equal to the Borel–Laplace sum $s_0(\tilde{f}_0)(y)$ in Eq. (124a) for $\Re(y) > 0$, while the function $-F_S(-\frac{1}{3y})$ is equal to the Borel–Laplace sum $s_\pi(\tilde{f}_0)(y)$ in Eq. (124b) for $\Re(y) < 0$. They can, therefore, be expressed explicitly as functions of the strong coupling Stokes constants according to Eqs. (122a) and (122b), respectively. We find in this way that

$$F_R(y) = -\frac{\pi i}{2} - \frac{3\mathcal{V}}{2\pi i y} + 2 \sum_{m \in \mathbb{Z}_{\neq 0}} R_m \mathbf{e}_1\left(-\frac{m}{3y}\right), \quad \Re(y) > 0, \quad (158a)$$

$$F_S(y) = \frac{\pi i}{2} - \frac{9\mathcal{V}y}{2\pi i} - 2 \sum_{m \in \mathbb{Z}_{\neq 0}} R_m \mathbf{e}_1(my), \quad \Re(y) > 0, \quad (158b)$$

where $\mathcal{V} = 2\Im(\mathrm{Li}_2(e^{2\pi i/3}))$ and \mathbf{e}_1 is the exponential integral in Eq. (110) as before. Indeed, Theorem 4.6 can be equivalently restated in terms of the functions above as

$$\frac{F_R(y) - F_S\left(-\frac{1}{3y}\right)}{2} = f_0(y), \quad y \in \mathbb{H}. \quad (159)$$

Remark 4.3. *Assuming the validity of Conjecture 1, we can complete the above discussion as follows. We observe that the function $F_R(-\frac{1}{3y})$ is equal to the Borel–Laplace sum $s_\pi(\tilde{f}_\infty)(y)$ in Eq. (154b) for $\Re(y) < 0$, while the function $F_S(y)$ is equal to the Borel–Laplace sum $s_0(\tilde{f}_\infty)(y)$ in Eq. (154a) for $\Re(y) > 0$. They can, therefore, be expressed explicitly as functions of the weak coupling Stokes constants according to Eqs. (123b) and (123a), respectively. Namely,*

$$F_R(y) = -3 \log \frac{\Gamma(2/3)}{\Gamma(1/3)} - \log(2\pi i/y) - 2 \sum_{m \in \mathbb{Z}_{\neq 0}} S_m \mathbf{e}_1(my), \quad \Re(y) > 0, \quad (160a)$$

$$F_S(y) = -3 \log \frac{\Gamma(2/3)}{\Gamma(1/3)} - \log(-6\pi i y) - 2 \sum_{m \in \mathbb{Z}_{\neq 0}} S_m \mathbf{e}_1\left(-\frac{m}{3y}\right), \quad \Re(y) > 0, \quad (160b)$$

²⁵Note that $\mathbb{C}_\gamma = \mathbb{C} \setminus (-\infty; -d/c]$ when $c > 0$ and $\mathbb{C}_\gamma = \mathbb{C} \setminus [-d/c; +\infty)$ when $c < 0$.

where \mathbf{e}_1 is again the exponential integral in Eq. (110). Indeed, Conjecture 1 is equivalent to

$$\frac{F_S(y) + F_R\left(-\frac{1}{3y}\right)}{2} = f_\infty(y), \quad y \in \mathbb{H}. \quad (161)$$

In addition, the functions $F_R(y)$ and $F_S(y)$ in Eqs. (157a) and (157b) can be simply written as the logarithm of a ratio of Faddeev's quantum dilogarithms. Indeed, setting $\mathbf{b}^2 = y$, it follows from the infinite product representations in Eqs. (63), (67), and (183) that

$$F_R(y) = 3 \log \frac{\Phi_{\mathbf{b}}\left(\frac{2i}{3\mathbf{b}} - c_{\mathbf{b}}\right)}{\Phi_{\mathbf{b}}\left(\frac{i}{3\mathbf{b}} - c_{\mathbf{b}}\right)} - i\pi, \quad (162a)$$

$$F_S(y/3) = 3 \log \frac{\Phi_{\mathbf{b}}\left(\frac{2i\mathbf{b}}{3} - c_{\mathbf{b}}\right)}{\Phi_{\mathbf{b}}\left(\frac{i\mathbf{b}}{3} - c_{\mathbf{b}}\right)}, \quad (162b)$$

where $c_{\mathbf{b}}$ is defined in Eq. (180). Thus, thanks to the analyticity properties of the Faddeev's quantum dilogarithm $\Phi_{\mathbf{b}}(x)$ reviewed in Appendix A, $F_R(y)$ and $F_S(y)$ can be analytically continued to the whole cut complex plane \mathbb{C}' . We can now prove our statements on the quantum modularity of the generating functions.

Theorem 4.7. *The generating functions $f_0, f_\infty: \mathbb{H} \rightarrow \mathbb{C}$, defined in Eqs. (61) and (65), respectively, are holomorphic quantum modular functions for $\Gamma_1(3)$.*

Proof. Recall that the generators of $\Gamma_1(3)$ are the matrices T and γ_3 in Eq. (9). It is sufficient to prove that the corresponding cocycles, that is,

$$h_T[f](y) = f(y+1) - f(y), \quad (163a)$$

$$h_{\gamma_3}[f](y) = f\left(\frac{y}{3y+1}\right) - f(y), \quad (163b)$$

for $f = f_0, f_\infty$, are analytic in \mathbb{C}' . It follows from the periodicity properties in Eqs. (62) and (66) that the cocycles for T are trivial, i.e., $h_T[f_0](y) = 0$ and $h_T[f_\infty](y) = 0$.

Applying the definition of $F_R(y)$ in Eq. (157a) and the first formula in Eq. (66), we find that

$$\begin{aligned} f_0\left(\frac{y}{3y+1}\right) &= F_R\left(\frac{y}{3y+1}\right) - f_\infty\left(-\frac{1}{3y}\right) \\ &= F_R\left(\frac{y}{3y+1}\right) - F_R(y) + f_0(y). \end{aligned} \quad (164)$$

Therefore, we obtain that the cocycle of f_0 for γ_3 is

$$h_{\gamma_3}[f_0](y) = F_R\left(\frac{y}{3y+1}\right) - F_R(y), \quad (165)$$

which is analytic in \mathbb{C}' following Eq. (162a). Analogously, applying the definition of $F_S(y)$ in Eq. (157b) and the first formula in Eq. (62), we find that

$$\begin{aligned} f_\infty\left(\frac{y}{3y+1}\right) &= F_S\left(\frac{y}{3y+1}\right) + f_0\left(-\frac{1}{3y}\right) \\ &= F_S\left(\frac{y}{3y+1}\right) - F_S(y) + f_\infty(y). \end{aligned} \quad (166)$$

Therefore, the cocycle of f_∞ for γ_3 is

$$h_{\gamma_3}[f_\infty](y) = F_S\left(\frac{y}{3y+1}\right) - F_S(y), \quad (167)$$

which is analytic on \mathbb{C}' following Eq. (162b). \square

Let us stress that the generating functions $f_0(y)$ and $f_\infty(y)$, $y \in \mathbb{H}$, have trivial cocycles for the generator T , while their cocycles for the generator γ_3 in Eqs. (165) and (167) are given by the functions $F_R(y)$ and $F_S(y)$ in Eq. (157a) and (157b), respectively. Specifically, we have that

$$h_{\gamma_3}[f_0](y) = h_{\gamma_3}[F_R](y), \quad h_{\gamma_3}[f_\infty](y) = h_{\gamma_3}[F_S](y), \quad (168)$$

which are determined by the strong and weak coupling Stokes constants R_n, S_n , $n \in \mathbb{Z}_{>0}$, as a consequence of Eqs. (158a) and (160b). Once more, note the exchange of information between the two regimes in \hbar . The non-trivial cocycle of the generating function of the Stokes constants in one regime is controlled by the Stokes constants in the dual regime.

Remark 4.4. *As we learn from the proof of Theorem 4.7, the holomorphic quantum modular functions f_0, f_∞ are deeply related to the Faddeev's quantum dilogarithm $\Phi_{\mathfrak{b}}$ in Eq. (183), which enters the formulae for the cocycles. This is indeed a general feature of the q -Pochhammer symbols, as discussed in [76, Section 8.11].*

Quantum modularity and Fricke involution

As we will show, a pair of “dual” holomorphic quantum modular functions for $\Gamma_1(3)$ is constructed by acting on the generating functions $f_0(y)$ and $f_\infty(y)$ with the transformation $y \mapsto -\frac{1}{3y}$, which is known as the Fricke involution of $\mathbb{H}/\Gamma_1(3)$. In particular, we define the holomorphic functions $f_0^*, f_\infty^*: \mathbb{H} \rightarrow \mathbb{C}$ as

$$f_0^*(y) := f_0\left(-\frac{1}{3y}\right), \quad f_\infty^*(y) := f_\infty\left(-\frac{1}{3y}\right), \quad (169)$$

where we have used the symbol $*$ to denote the action of the Fricke involution.

Theorem 4.8. *The functions $f_0^*, f_\infty^*: \mathbb{H} \rightarrow \mathbb{C}$, defined in Eq. (169) as the images of the generating functions under Fricke involution, are holomorphic quantum modular functions for $\Gamma_1(3)$.*

Proof. Recall that the generators of $\Gamma_1(3)$ are the matrices T and γ_3 in Eq. (9). It is sufficient to prove that the corresponding cocycles, that is,

$$h_T[f^*](y) = f^*(y+1) - f^*(y) = f\left(-\frac{1}{3y+3}\right) - f\left(-\frac{1}{3y}\right), \quad (170a)$$

$$h_{\gamma_3}[f^*](y) = f^*\left(\frac{y}{3y+1}\right) - f^*(y) = f\left(-\frac{1}{3y} - 1\right) - f\left(-\frac{1}{3y}\right), \quad (170b)$$

for $f = f_0, f_\infty$, are analytic in \mathbb{C}' . It follows from the periodicity properties in Eqs. (62) and (66) that the cocycles for γ_3 are trivial, i.e., $h_{\gamma_3}[f_0^*](y) = 0$ and $h_{\gamma_3}[f_\infty^*](y) = 0$.

Applying the definition of $F_S(y)$ in Eq. (157b) and the first formula in Eq. (66), we find that

$$\begin{aligned} f_0^*(y+1) &= f_0\left(-\frac{1}{3y+3}\right) = f_\infty(y+1) - F_S(y+1) \\ &= F_S(y) - F_S(y+1) + f_0\left(-\frac{1}{3y}\right) \\ &= F_S(y) - F_S(y+1) + f_0^*(y). \end{aligned} \quad (171)$$

Therefore, we obtain that the cocycle of f_0^* for T is

$$h_T[f_0^*](y) = F_S(y) - F_S(y+1), \quad (172)$$

which is analytic on \mathbb{C}' following Eq. (162b). Analogously, applying the definition of $F_R(y)$ in Eq. (157a) and the first formula in Eq. (62), we find that

$$\begin{aligned} f_\infty^*(y+1) &= f_\infty\left(-\frac{1}{3y+3}\right) = F_R(y+1) - f_0(y+1) \\ &= F_R(y+1) - F_R(y) + f_\infty\left(-\frac{1}{3y}\right) \\ &= F_R(y+1) - F_R(y) + f_\infty^*(y), \end{aligned} \quad (173)$$

Therefore, the cocycle of f_∞^* for T is

$$h_T[f_\infty^*](y) = F_R(y+1) - F_R(y), \quad (174)$$

which is analytic on \mathbb{C}' following Eq. (162a). \square

The Fricke involution has exchanged the roles played by the generators of the modular group $\Gamma_1(3)$ and, simultaneously, by the weak and strong coupling Stokes constants. Indeed, the dual functions $f_0^*(y)$ and $f_\infty^*(y)$, $y \in \mathbb{H}$, have trivial cocycles for the generator γ_3 , while their cocycles for the generator T in Eqs. (172) and (174) are given by the functions $F_S(y)$ and $F_R(y)$ in Eq. (157b) and (157a), respectively. Namely, we have that

$$h_T[f_0^*](y) = -h_T[F_S](y), \quad h_T[f_\infty^*](y) = h_T[F_R](y), \quad (175)$$

which are determined by the weak and strong coupling Stokes constants S_n , R_n , $n \in \mathbb{Z}_{>0}$, as a consequence of Eqs. (160b) and (158a).

Remark 4.5. *Our results on the quantum modularity of the generating functions and the role of the Fricke involution pave the way for a geometric interpretation of the Stokes constants. Indeed, as briefly discussed in Section 2.1, the group $\Gamma_1(3)$ is deeply related to the geometry of local \mathbb{P}^2 . The moduli space $\mathcal{M}_{\text{cpx}}(\hat{X})$ parametrizing the complex structures of the mirror \hat{X} is the compactification of the quotient $\mathbb{H}/\Gamma_1(3)$. Furthermore, the generating functions of the Gromov–Witten invariants are known to be quasi modular functions in the modular coordinate $\tau \in \mathcal{M}_{\text{cpx}}(\hat{X})$ at the conifold point [34]. This enticing connection opens a new direction of investigation aimed at studying an explicit relationship between the Stokes constants appearing in TS/ST correspondence and the Gromov–Witten invariants of toric CY threefolds, which we plan to address in future work [63].*

Finally, a large class of examples of quantum modular forms comes from the study of quantum invariants of 3-manifolds and knots [80, 85, 86]. Thus, our results on the quantum modularity of the generating functions of the Stokes constants in Theorems 4.7 and 4.8 provide new evidence in addition to the many formal similarities that are shared by topological string theory and complex Chern–Simons theory, which we have briefly discussed in Section 3.2.

5 Conclusions

The Stokes constants are analytic invariants capturing information about the non-perturbative corrections to a resurgent asymptotic series. Yet, in some remarkable cases, they are rational or integer numbers and possess an interpretation as enumerative invariants based on the counting of BPS states [87–89]. Examples come from several quantum theories, including $4d \mathcal{N} = 2$ supersymmetric gauge theory in the NS limit of the Omega-background [90], complex Chern–Simons theory on Seifert fibered homology spheres [79] and on complements of hyperbolic knots [64, 65],

the standard and NS topological string theories on (toric) CY threefolds [1, 20–22, 24, 26, 28–32] and their refinement [25, 27], and Walcher’s real topological string [23].

In this paper, following the exact results of [1], we completed our understanding of the unique set of relations connecting the resurgent structures of the logarithm of the spectral trace of local \mathbb{P}^2 in the weak and strong coupling limits. In particular, we uncovered a direct way of obtaining the perturbative expansion of the generating function of the Stokes constants in one regime from the generating function in the other, which amounts to performing an operation that is the formal inverse of taking the discontinuity of the asymptotic series. Thus, we proved that a *strong-weak resurgent symmetry* is at the heart of the analytic number-theoretic duality of [1]. Through this newly discovered exact symmetry, the two-way exchange of perturbative/non-perturbative information between the holomorphic and anti-holomorphic blocks in the factorization of the spectral trace takes a mathematically precise form. This is a realization of underlying physical mechanisms that can be intuitively traced back to the duality between the worldsheet and WKB contributions to the total grand potential of the topological string on local \mathbb{P}^2 . Moreover, the weak and strong coupling Stokes constants are related by an arithmetic twist, while the corresponding L -functions analytically continue each other by means of a functional equation.

Finally, we proved that the q -series acting as generating functions of the Stokes constants are holomorphic quantum modular functions for the group $\Gamma_1(3)$ and showed the effectiveness of the median resummation in the weakly coupled regime, while the analogous statement at strong coupling is conjectured in light of numerical evidence.

Multiple questions and open problems follow from the investigation performed in this paper. The geometric and physical meaning of the enumerative invariants and non-perturbative sectors of [1, 32] is yet to be understood. However, the remarkable arithmetic fabric underpinning the resurgent properties of the dual \hbar -expansions of $\log \text{Tr}(\rho_{\mathbb{P}^2})$ discovered in [1], and developed here into a global strong-weak resurgent symmetry, paves the way for new insights on the physical interpretation of the Stokes constants to be explored in upcoming work [62].

Along the same lines, a study of the relation between the formalism of [20, 24] and the framework proposed in [1, 32] and advanced here might help us achieve a more comprehensive understanding of the non-perturbative structure of topological string theory. Let us also mention that other techniques, which do not employ resurgence, have been recently used to address the problem of identifying and counting BPS states of type IIA compactifications on toric CY threefolds [91]. These include generalizations of the WKB approach [92–95] and techniques based on attractor flows [61].

Simultaneously, our results on the quantum modularity of the generating functions suggest a new direction of research as they naturally prompt us to investigate the geometric content of the weak and strong coupling Stokes constants in connection to the Gromov–Witten invariants of local \mathbb{P}^2 . We plan to address these ideas in future work [63].

Finally, it would be interesting to extend our results on the first fermionic spectral trace of local \mathbb{P}^2 to other toric CY threefolds and possibly study higher-order fermionic spectral traces, where we expect that similar structures might show up, guiding us toward a potential generalization of the strong-weak resurgent symmetry.

Acknowledgements

We thank Alba Grassi, Maxim Kontsevich, Marcos Mariño, Campbell Wheeler, and Don Zagier for many useful discussions. This work has been supported by the ERC-SyG project “Recur-sive and Exact New Quantum Theory” (ReNewQuantum), which received funding from the

European Research Council (ERC) within the European Union's Horizon 2020 research and innovation program under Grant No. 810573 and by the Swiss National Centre of Competence in Research SwissMAP (NCCR 51NF40-141869 The Mathematics of Physics).

A Quantum dilogarithms

In this section, we recall some of the main properties of Faddeev's quantum dilogarithm [96]. We refer to [70, Appendix A] and [76, Section 8.10] for detailed expositions. We call *quantum dilogarithm* the function of two variables defined by the series

$$(xq^\alpha; q)_\infty = \prod_{n=0}^{\infty} (1 - xq^{\alpha+n}), \quad \alpha \in \mathbb{R}, \quad (176)$$

which is analytic in $x, q \in \mathbb{C}$ with $|q| < 1$ and has asymptotic expansions around q a root of unity. It satisfies the relation

$$(x; q^{-1})_\infty = (xq; q)_\infty^{-1}. \quad (177)$$

The *q-Pochhammer symbols*, also known as *q-shifted factorials*, are denoted by

$$(x; q)_m = \prod_{n=0}^{m-1} (1 - xq^n), \quad (x; q)_{-m} = \frac{1}{(xq^{-m}; q)_m}, \quad m \in \mathbb{Z}_{>0}, \quad (178)$$

with $(x; q)_0 = 1$. Equivalently, we can write

$$(x; q)_m = \frac{(x; q)_\infty}{(xq^m; q)_\infty}, \quad m \in \mathbb{Z}. \quad (179)$$

The *Faddeev's quantum dilogarithm* $\Phi_{\mathbf{b}}(x)$ is defined in the strip $|\Im(x)| < |\Im(c_{\mathbf{b}})|$, where

$$c_{\mathbf{b}} = i(\mathbf{b} + \mathbf{b}^{-1})/2, \quad (180)$$

by the integral representation

$$\Phi_{\mathbf{b}}(x) = \exp \left(\int_{\mathbb{R}+i\epsilon} \frac{e^{-2ixz}}{4 \sinh(z\mathbf{b}) \sinh(z\mathbf{b}^{-1})} \frac{dz}{z} \right), \quad (181)$$

which implies the symmetry properties

$$\Phi_{\mathbf{b}}(x) = \Phi_{-\mathbf{b}}(x) = \Phi_{\mathbf{b}^{-1}}(x). \quad (182)$$

When $\Im(\mathbf{b}^2) > 0$, the formula in Eq. (181) is equivalent to

$$\Phi_{\mathbf{b}}(x) = \frac{(e^{2\pi\mathbf{b}(x+c_{\mathbf{b}})}; q)_\infty}{(e^{2\pi\mathbf{b}^{-1}(x-c_{\mathbf{b}})}; \tilde{q})_\infty} = \prod_{n=0}^{\infty} \frac{1 - e^{2\pi\mathbf{b}(x+c_{\mathbf{b}})} q^n}{1 - e^{2\pi\mathbf{b}^{-1}(x-c_{\mathbf{b}})} \tilde{q}^n}, \quad (183)$$

where

$$q = e^{2\pi i \mathbf{b}^2}, \quad \tilde{q} = e^{-2\pi i \mathbf{b}^{-2}}. \quad (184)$$

Note that the function in Eq. (183) can be extended to the region $\Im(\mathbf{b}^2) < 0$ by means of Eq. (182) and further admits an analytic continuation to all values of \mathbf{b} such that $\mathbf{b}^2 \notin \mathbb{R}_{\leq 0}$.

Moreover, $\Phi_{\mathbf{b}}(x)$ can be extended to the whole complex x -plane as a meromorphic function with an essential singularity at infinity, poles at the points

$$x = c_{\mathbf{b}} + i m \mathbf{b} + i n \mathbf{b}^{-1}, \quad (185)$$

and zeros at the points

$$x = -c_{\mathbf{b}} - i m \mathbf{b} - i n \mathbf{b}^{-1}, \quad (186)$$

for $m, n \in \mathbb{N}$. It satisfies the inversion formula

$$\Phi_{\mathbf{b}}(x)\Phi_{\mathbf{b}}(-x) = e^{\pi i x^2} \Phi_{\mathbf{b}}(0)^2, \quad \Phi_{\mathbf{b}}(0) = \left(\frac{q}{\tilde{q}}\right)^{1/48} = e^{\pi i (\mathbf{b}^2 + \mathbf{b}^{-2})/24}, \quad (187)$$

the complex conjugation formula

$$\overline{\Phi_{\mathbf{b}}(x)} = \frac{1}{\Phi_{\bar{\mathbf{b}}}(\bar{x})}, \quad (188)$$

and the quasi-periodicity relations

$$\Phi_{\mathbf{b}}(x \pm i \mathbf{b}) = \Phi_{\mathbf{b}}(x) \left(1 + e^{2\pi \mathbf{b} x \pm \pi i \mathbf{b}^2}\right)^{\mp 1}, \quad (189a)$$

$$\Phi_{\mathbf{b}}(x \pm i \mathbf{b}^{-1}) = \Phi_{\mathbf{b}}(x) \left(1 + e^{2\pi \mathbf{b}^{-1} x \pm \pi i \mathbf{b}^{-2}}\right)^{\mp 1}. \quad (189b)$$

In addition, when $\mathbf{b}^2 = M/N \in \mathbb{Q}$ with $M, N \in \mathbb{Z}_{>0}$ coprime, the expression for $\Phi_{\mathbf{b}}(x)$ in Eq. (183) simplifies into [97, Theorem 1.9]

$$\Phi_{\mathbf{b}}\left(\frac{x}{2\pi\sqrt{MN}} - c_{\mathbf{b}}\right) = \frac{e^{\frac{i}{2\pi MN} \text{Li}_2(e^x)} (1 - e^x)^{1 + \frac{ix}{2\pi MN}}}{D_N(e^{x/N}; e^{2\pi i M/N}) D_M(e^{x/M}; e^{2\pi i N/M})}, \quad (190)$$

where

$$D_N(z; q) = \prod_{k=1}^{N-1} (1 - z q^k)^{\frac{k}{N}} \quad (191)$$

is the cyclic quantum dilogarithm.²⁶

Finally, we recall the known asymptotic behaviour of Faddeev's quantum dilogarithm. We refer to Appendix B for an introduction to resurgence. In the limit of $\mathbf{b} \rightarrow 0$ with $\Im(\mathbf{b}^2) > 0$, $\Phi_{\mathbf{b}}(x)$ gives the asymptotic series [70]

$$\log \Phi_{\mathbf{b}}\left(\frac{x}{2\pi \mathbf{b}}\right) \sim \sum_{k=0}^{\infty} (2\pi i \mathbf{b}^2)^{2k-1} \frac{B_{2k}(1/2)}{(2k)!} \text{Li}_{2-2k}(-e^x), \quad (192)$$

where x is kept fixed, $\text{Li}_n(z)$ is the polylogarithm of order n , and $B_n(z)$ is the n -th Bernoulli polynomial. Similarly, for $\mathbf{b} \rightarrow 0$ with $\Im(\mathbf{b}^2) > 0$ and x fixed, the following special cases of the quantum dilogarithm in Eq. (176) have the asymptotic expansions [98]

$$\log(x; q)_{\infty} \sim \frac{1}{2} \log(1 - x) + \sum_{k=0}^{\infty} (2\pi i \mathbf{b}^2)^{2k-1} \frac{B_{2k}}{(2k)!} \text{Li}_{2-2k}(x), \quad (193a)$$

$$\begin{aligned} \log(q^{\alpha}; q)_{\infty} &\sim -\frac{\pi i}{12 \mathbf{b}^2} - B_1(\alpha) \log(-2\pi i \mathbf{b}^2) - \log \frac{\Gamma(\alpha)}{\sqrt{2\pi}} \\ &\quad - B_2(\alpha) \frac{\pi i \mathbf{b}^2}{2} - \sum_{k=2}^{\infty} (2\pi i \mathbf{b}^2)^k \frac{B_k B_{k+1}(\alpha)}{k(k+1)!}, \quad \alpha > 0, \end{aligned} \quad (193b)$$

²⁶We use the simplified formula for Faddeev's quantum dilogarithm in Eq. (190) to perform numerical checks at rational points of the results presented in Section 4.2.

where $B_n = B_n(0)$ is the n -th Bernoulli number and $\Gamma(\alpha)$ is the gamma function. Besides, $\Phi_b(x)$ is Borel–Laplace summable [99, Theorem 1.3].

B Basics of resurgence

The resurgent analysis of factorially divergent formal power series unveils a universal mathematical structure involving a set of numerical data called Stokes constants. See [12–14] for a formal introduction to the theory of resurgence and [2, 15] for its application to gauge and string theories. In this section, we review the basics of resurgence. Let z be a formal variable. The Borel transform is a map acting on z -monomials as

$$\mathcal{B}[z^{n-\alpha}] := \frac{\zeta^{n-\alpha-1}}{\Gamma(n-\alpha)}, \quad n \in \mathbb{Z}_{\geq 0}, \quad \alpha \in \mathbb{R} \setminus \mathbb{Z}_{\geq 0}, \quad (194)$$

where ζ is a new formal variable and $\Gamma(n-\alpha)$ is the gamma function. The action of the Borel transform above is extended by linearity to all formal power series in $z^{-\alpha}\mathbb{C}[[z]]$, while the Borel transform of the identity is conventionally denoted with the symbol $\delta := \mathcal{B}[1]$. Let $\phi(z)$ be a Gevrey-1 asymptotic series of the form

$$\phi(z) = z^{-\alpha} \sum_{n=0}^{\infty} a_n z^n \in z^{-\alpha}\mathbb{C}[[z]], \quad |a_n| \leq \mathcal{A}^{-n} n! \quad n \gg 1, \quad (195)$$

where $\alpha \in \mathbb{R} \setminus \mathbb{Z}_{\geq 0}$ and $\mathcal{A} \in \mathbb{R}_{>0}$. Its Borel transform, which we denote for simplicity by $\hat{\phi}(\zeta) := \mathcal{B}\phi(\zeta)$, is by definition

$$\hat{\phi}(\zeta) = \sum_{k=0}^{\infty} \frac{a_k}{\Gamma(k-\alpha)} \zeta^{k-\alpha-1} \in \zeta^{-\alpha}\mathbb{C}\{\zeta\} \quad (196)$$

and is a holomorphic function in an open neighborhood of $\zeta = 0$ of radius $|\mathcal{A}|$. When extended to the complex ζ -plane, known as Borel plane, $\hat{\phi}(\zeta)$ shows a (possibly infinite) set of singularities $\zeta_\omega \in \mathbb{C}$, which we label by the index $\omega \in \Omega$. A ray in the Borel plane of the form

$$\mathcal{C}_{\theta_\omega} = e^{i\theta_\omega} \mathbb{R}_+, \quad \theta_\omega = \arg(\zeta_\omega), \quad (197)$$

which starts at the origin and passes through a singularity ζ_ω , is called a Stokes ray. The Borel plane is partitioned into sectors bounded by the Stokes rays in such a way that the Borel transform converges to a generally different holomorphic function in each of these sectors.

We recall that the Gevrey-1 asymptotic series $\phi(z)$ is called *resurgent* if its Borel transform $\hat{\phi}(\zeta)$ can be endlessly analytically continued. Namely, for every $L > 0$, there is a finite set of points Ω_L in the Riemann surface of ζ^α such that $\hat{\phi}(\zeta)$ can be analytically continued along any path that avoids Ω_L and has length at most L . If, additionally, its Borel transform has only simple poles and logarithmic branch points, then it is called *simple resurgent*. Let us assume that the formal power series $\phi(z)$ in Eq. (195) is simple resurgent. If the singularity ζ_ω is a simple pole, the local expansion of the Borel transform in Eq. (196) around it has the form

$$\hat{\phi}(\zeta) = -\frac{S_\omega}{2\pi i(\zeta - \zeta_\omega)} + \text{regular in } \zeta - \zeta_\omega, \quad (198)$$

where $S_\omega \in \mathbb{C}$ is the Stokes constant at ζ_ω . Whereas, if the singularity ζ_ω is a logarithmic branch point, the local expansion of the Borel transform in Eq. (196) around it has the form

$$\hat{\phi}(\zeta) = -\frac{S_\omega}{2\pi i} \log(\zeta - \zeta_\omega) \hat{\phi}_\omega(\zeta - \zeta_\omega) + \text{regular in } \zeta - \zeta_\omega, \quad (199)$$

where again $S_\omega \in \mathbb{C}$ is the corresponding Stokes constant. If we introduce the variable $\xi = \zeta - \zeta_\omega$, the function

$$\hat{\phi}_\omega(\xi) = \sum_{k=0}^{\infty} \hat{a}_{k,\omega} \xi^{k-\beta} \in \xi^{-\beta} \mathbb{C}\{\xi\}, \quad (200)$$

where $\beta \in \mathbb{R} \setminus \mathbb{Z}_+$, is locally analytic at $\xi = 0$ and can be regarded as the Borel transform of the Gevrey-1 asymptotic series

$$\phi_\omega(z) = z^{-\beta} \sum_{n=0}^{\infty} a_{n,\omega} z^n \in z^{-\beta} \mathbb{C}[[z]], \quad a_{n,\omega} = \Gamma(n - \beta + 1) \hat{a}_{n,\omega}. \quad (201)$$

Note that the value of the Stokes constant S_ω depends on the normalization of $\phi_\omega(z)$.

If the analytic continuation of the Borel transform $\hat{\phi}(\zeta)$ in Eq. (196) does not grow too fast at infinity²⁷, its Laplace transform at an arbitrary angle θ in the Borel plane gives the Borel–Laplace sum of the original, divergent formal power series $\phi(z)$, denoted by $s_\theta(\phi)(z)$. Explicitly,

$$s_\theta(\phi)(z) = \int_0^{e^{i\theta}\infty} e^{-\zeta/z} \hat{\phi}(\zeta) d\zeta = z \int_0^{e^{i\theta}\infty} e^{-\zeta} \hat{\phi}(\zeta z) d\zeta, \quad (202)$$

whose asymptotics near the origin reconstructs $\phi(z)$. If the Borel–Laplace sum in Eq. (202) exists in some region of the complex z -plane, we say that the series $\phi(z)$ is Borel–Laplace summable along the direction θ . Note that the Borel–Laplace sum inherits the sectorial structure of the Borel transform. It is a locally analytic function with discontinuities across the special rays identified by

$$\arg(z) = \arg(\zeta_\omega), \quad \omega \in \Omega. \quad (203)$$

The discontinuity across an arbitrary ray $\mathcal{C}_\theta = e^{i\theta} \mathbb{R}_+$ in the complex z -plane is the difference between the Borel–Laplace sums along two rays in the complex ζ -plane that lie slightly above and slightly below \mathcal{C}_θ . Namely,

$$\text{disc}_\theta \phi(z) = s_{\theta_+}(\phi)(z) - s_{\theta_-}(\phi)(z) = \int_{\mathcal{C}_{\theta_+} - \mathcal{C}_{\theta_-}} e^{-\zeta/z} \hat{\phi}(\zeta) d\zeta, \quad (204)$$

where $\theta_\pm = \theta \pm \epsilon$ for some small positive angle ϵ . A standard contour deformation argument shows that the two lateral Borel–Laplace sums differ by exponentially small terms. More precisely, if the Borel transform $\hat{\phi}(z)$ has only simple poles, then we have that

$$\text{disc}_\theta \phi(z) = \sum_{\omega \in \Omega_\theta} S_\omega e^{-\zeta_\omega/z}, \quad (205)$$

where the index $\omega \in \Omega_\theta$ labels the singularities ζ_ω such that $\arg(\zeta_\omega) = \theta$ and the complex numbers S_ω are the same Stokes constants that appear in Eq. (198). Similarly, if there are only logarithmic branch points, we have that

$$\text{disc}_\theta \phi(z) = \sum_{\omega \in \Omega_\theta} S_\omega e^{-\zeta_\omega/z} s_{\theta_-}(\phi_\omega)(z), \quad (206)$$

where again the complex numbers S_ω are the Stokes constants appearing in Eq. (199), while $\phi_\omega(z)$ is the formal power series in Eq. (201).

²⁷Roughly, we require that the Borel transform grows at most exponentially in an open sector of the Borel plane containing the angle θ .

Let us now focus on the case of logarithmic singularities. If we regard the lateral Borel–Laplace sums as operators acting on formal power series, the Stokes automorphism \mathfrak{S}_θ at an arbitrary angle θ is defined by the composition

$$s_{\theta+} = s_{\theta-} \circ \mathfrak{S}_\theta, \quad (207)$$

and the discontinuity formula in Eq. (206) has the equivalent form

$$\mathfrak{S}_\theta(\phi) = \phi + \sum_{\omega \in \Omega_\theta} S_\omega e^{-\zeta_\omega/z} \phi_\omega. \quad (208)$$

An interesting feature of resurgence is that it allows us to build a whole family of new asymptotic series ϕ_ω , $\omega \in \Omega$, from the knowledge of the singularity structure of the Borel transform of one single resurgent series ϕ . We can then repeat the procedure above with each new series obtained this way, assuming they are simple resurgent. For each $\omega \in \Omega$, let us define the basic trans-series

$$\Phi_\omega(z) = e^{-\zeta_\omega/z} \phi_\omega(z), \quad (209)$$

such that its Borel–Laplace sum at the angle θ is given by

$$s_\theta(\Phi_\omega)(z) = e^{-\zeta_\omega/z} s_\theta(\phi_\omega)(z). \quad (210)$$

The corresponding Stokes automorphism acts on the basic trans-series as

$$\mathfrak{S}_\theta(\Phi_\omega) = \Phi_\omega + \sum_{\omega' \in \Omega_\theta} S_{\omega\omega'} \Phi_{\omega'}, \quad (211)$$

where we have denoted by $S_{\omega\omega'} \in \mathbb{C}$ the Stokes constants of the secondary asymptotic series $\phi_\omega(z)$. The minimal resurgent structure associated with $\phi(z)$ is the smallest collection of basic trans-series that resurge from it and form a closed set under Stokes automorphisms. It is denoted by [32]

$$\mathfrak{B}_\phi = \{\Phi_\omega(z)\}_{\omega \in \bar{\Omega}}, \quad \bar{\Omega} \subseteq \Omega. \quad (212)$$

We stress that \mathfrak{B}_ϕ does not necessarily include all the basic trans-series arising from $\phi(z)$. Finally, we construct the (possibly infinite-dimensional) matrix of Stokes constants

$$\mathcal{S}_\phi = \{S_{\omega\omega'}\}_{\omega, \omega' \in \bar{\Omega}}, \quad (213)$$

which is indexed by the distinct basic trans-series in the minimal resurgent structure of $\phi(z)$ and incorporates additional information about the non-analytic corrections to the original asymptotic series. In the case of simple poles, the same definitions and considerations above apply after imposing that $\phi_\omega(z) = 1$, $\omega \in \Omega$.

In the study of the summability properties of factorially divergent formal power series that arise as asymptotic limits of well-defined holomorphic functions, it is natural to ask the question of which summability method is efficient in giving back the original function. In quantum Chern–Simons theory, Costin and Garoufalidis conjectured that the asymptotic expansions at roots of unity of quantum invariants, such as the WRT invariant for knots and the Kashaev invariant for 3-manifolds, are Gevrey-1 series whose median resummation reconstructs the non-perturbative invariants [78, Conjecture 1]. As we show in this paper, more evidence of the efficacy of the median resummation comes from the study of the spectral traces of the quantum operators obtained by quantizing the mirror curve to a toric CY threefold. We recall that the

median resummation of the Gevrey-1 asymptotic series $\phi(z)$ in Eq. (195) across an arbitrary ray \mathcal{C}_θ in the complex z -plane is the average of the two lateral Borel–Laplace sums $s_{\theta_\pm}(\phi)(z)$, that is,

$$\mathcal{S}_\theta^{\text{med}}\phi(z) = \frac{s_{\theta_+}(\phi)(z) + s_{\theta_-}(\phi)(z)}{2}, \quad (214)$$

which is an analytic function for $\arg z \in (\theta - \frac{\pi}{2}, \theta + \frac{\pi}{2})$. Equivalently, we can write

$$\mathcal{S}_\theta^{\text{med}}\phi(z) = \begin{cases} s_{\theta_-}(\phi)(z) + \frac{1}{2} \text{disc}_\theta\phi(z), & \Re(e^{-i\theta_-}z) > 0, \\ s_{\theta_+}(\phi)(z) - \frac{1}{2} \text{disc}_\theta\phi(z), & \Re(e^{-i\theta_+}z) > 0, \end{cases} \quad (215)$$

where $\Re(e^{-i\theta}z) > 0$. The expression above highlights how the discontinuities can be suitably interpreted as corrections to the standard Borel–Laplace transform. In addition, by varying the contour of integration of the Laplace transform, the domain of analyticity of the median resummation can be extended, which is helpful in some applications.

References

- [1] C. Rella, *Resurgence, Stokes constants, and arithmetic functions in topological string theory*, *Commun. Num. Theor. Phys.* **17** (2023) 709–820, [2212.10606].
- [2] M. Mariño, *Lectures on non-perturbative effects in large N gauge theories, matrix models and strings*, *Fortsch. Phys.* **62** (2014) 455–540, [1206.6272].
- [3] A. Grassi, Y. Hatsuda and M. Mariño, *Topological Strings from Quantum Mechanics*, *Ann. Henri Poincaré* **17** (2016) 3177–3235, [1410.3382].
- [4] S. Codesido, A. Grassi and M. Mariño, *Spectral Theory and Mirror Curves of Higher Genus*, *Ann. Henri Poincaré* **18** (2017) 559–622, [1507.02096].
- [5] M. Mariño, *Open string amplitudes and large order behavior in topological string theory*, *JHEP* **03** (2008) 060, [hep-th/0612127].
- [6] V. Bouchard, A. Klemm, M. Mariño and S. Pasquetti, *Remodeling the B-model*, *Commun. Math. Phys.* **287** (2009) 117–178, [0709.1453].
- [7] K. Sun, X. Wang and M.-x. Huang, *Exact Quantization Conditions, Toric Calabi-Yau and Nonperturbative Topological String*, *JHEP* **01** (2017) 061, [1606.07330].
- [8] A. Grassi and J. Gu, *BPS relations from spectral problems and blowup equations*, *Lett. Math. Phys.* **109** (2019) 1271–1302, [1609.05914].
- [9] M. Mariño, R. Schiappa and M. Weiss, *Nonperturbative Effects and the Large-Order Behavior of Matrix Models and Topological Strings*, *Commun. Num. Theor. Phys.* **2** (2008) 349–419, [0711.1954].
- [10] M. Mariño, *Nonperturbative effects and nonperturbative definitions in matrix models and topological strings*, *JHEP* **12** (2008) 114, [0805.3033].
- [11] J. Écalle, *Les fonctions résurgentes. Tome I*, vol. 5 of *Publications Mathématiques d’Orsay* 81. Université de Paris-Sud, Département de Mathématiques, Orsay, 1981.

- [12] I. Aniceto, G. Basar and R. Schiappa, *A Primer on Resurgent Transseries and Their Asymptotics*, *Phys. Rept.* **809** (2019) 1–135, [1802.10441].
- [13] C. Mitschi and D. Sauzin, *Divergent Series, Summability and Resurgence I*, vol. 2153 of *Lecture Notes in Mathematics*. Springer, 2016, 10.1007/978-3-319-28736-2.
- [14] D. Dorigoni, *An Introduction to Resurgence, Trans-Series and Alien Calculus*, *Annals Phys.* **409** (2019) 167914, [1411.3585].
- [15] M. Mariño, *Instantons and Large N: An Introduction to Non-Perturbative Methods in Quantum Field Theory*. Cambridge University Press, 2015, 10.1017/CBO9781107705968.
- [16] R. Couso-Santamaría, J. D. Edelstein, R. Schiappa and M. Vonk, *Resurgent Transseries and the Holomorphic Anomaly*, *Ann. Henri Poincaré* **17** (2016) 331–399, [1308.1695].
- [17] R. Couso-Santamaría, J. D. Edelstein, R. Schiappa and M. Vonk, *Resurgent Transseries and the Holomorphic Anomaly: Nonperturbative Closed Strings in Local \mathbb{CP}^2* , *Commun. Math. Phys.* **338** (2015) 285–346, [1407.4821].
- [18] M. Bershadsky, S. Cecotti, H. Ooguri and C. Vafa, *Holomorphic anomalies in topological field theories*, *Nucl. Phys. B* **405** (1993) 279–304, [hep-th/9302103].
- [19] M. Bershadsky, S. Cecotti, H. Ooguri and C. Vafa, *Kodaira-Spencer theory of gravity and exact results for quantum string amplitudes*, *Commun. Math. Phys.* **165** (1994) 311–428, [hep-th/9309140].
- [20] J. Gu and M. Mariño, *Exact multi-instantons in topological string theory*, *SciPost Phys.* **15** (2023) 179, [2211.01403].
- [21] J. Gu, A.-K. Kashani-Poor, A. Klemm and M. Mariño, *Non-perturbative topological string theory on compact Calabi-Yau 3-folds*, *arXiv e-prints* (5, 2023) , [2305.19916].
- [22] K. Iwaki and M. Mariño, *Resurgent Structure of the Topological String and the First Painlevé Equation*, *arXiv e-prints* (7, 2023) , [2307.02080].
- [23] M. Mariño and M. Schwick, *Non-perturbative real topological strings*, *arXiv e-prints* (9, 2023) , [2309.12046].
- [24] J. Gu and M. Mariño, *On the resurgent structure of quantum periods*, *SciPost Phys.* **15** (2023) 035, [2211.03871].
- [25] S. Alexandrov, M. Mariño and B. Pioline, *Resurgence of refined topological strings and dual partition functions*, *arXiv e-prints* (11, 2023) , [2311.17638].
- [26] J. Gu, *Relations between Stokes constants of unrefined and Nekrasov-Shatashvili topological strings*, *arXiv e-prints* (7, 2023) , [2307.02079].
- [27] J. Gu and G. Guo, *Resurgent Wilson loops in refined topological string*, *arXiv e-prints* (3, 2024) , [2403.04528].
- [28] S. Pasquetti and R. Schiappa, *Borel and Stokes Nonperturbative Phenomena in Topological String Theory and $c=1$ Matrix Models*, *Ann. Henri Poincaré* **11** (2010) 351–431, [0907.4082].

- [29] M. Alim, A. Saha, J. Teschner and I. Tulli, *Mathematical Structures of Non-perturbative Topological String Theory: From GW to DT Invariants*, *Commun. Math. Phys.* **399** (2023) 1039–1101, [2109.06878].
- [30] A. Grassi, Q. Hao and A. Neitzke, *Exponential Networks, WKB and the Topological String*, *arXiv e-prints* (1, 2022) , [2201.11594].
- [31] M. Alim, L. Hollands and I. Tulli, *Quantum curves, resurgence and exact WKB*, *arXiv e-prints* (3, 2022) , [2203.08249].
- [32] J. Gu and M. Mariño, *Peacock patterns and new integer invariants in topological string theory*, *SciPost Phys.* **12** (2022) 058, [2104.07437].
- [33] M. Aganagic, V. Bouchard and A. Klemm, *Topological Strings and (Almost) Modular Forms*, *Commun. Math. Phys.* **277** (2008) 771–819, [hep-th/0607100].
- [34] T. Coates and H. Iritani, *Gromov–Witten invariants of local \mathbb{P}^2 and modular forms*, *J. Math. (Kyoto)* **61** (2021) 543 – 706, [1804.03292].
- [35] P. Bousseau, H. Fan, S. Guo and L. Wu, *Holomorphic anomaly equation for (\mathbb{P}^2, E) and the Nekrasov–Shatashvili limit of local \mathbb{P}^2* , *Forum Math. Pi* **9** (2021) e3, [2001.05347].
- [36] V. Fantini and C. Rella, “Modular resurgent structures.” *In preparation*.
- [37] M. Mariño, *Spectral Theory and Mirror Symmetry*, *Proc. Symp. Pure Math.* **98** (2018) 259, [1506.07757].
- [38] P. Candelas, X. C. De La Ossa, P. S. Green and L. Parkes, *A Pair of Calabi-Yau manifolds as an exactly soluble superconformal theory*, *Nucl. Phys. B* **359** (1991) 21–74.
- [39] M.-x. Huang and X.-f. Wang, *Topological Strings and Quantum Spectral Problems*, *JHEP* **09** (2014) 150, [1406.6178].
- [40] A. D. Mironov and A. Morozov, *Nekrasov Functions and Exact Bohr-Zommerfeld Integrals*, *JHEP* **04** (2010) 040, [0910.5670].
- [41] M. Aganagic, M. C. N. Cheng, R. Dijkgraaf, D. Krefl and C. Vafa, *Quantum Geometry of Refined Topological Strings*, *JHEP* **11** (2012) 019, [1105.0630].
- [42] R. M. Kashaev and M. Mariño, *Operators from mirror curves and the quantum dilogarithm*, *Commun. Math. Phys.* **346** (2016) 967–994, [1501.01014].
- [43] S. Codesido, J. Gu and M. Mariño, *Operators and higher genus mirror curves*, *JHEP* **02** (2017) 092, [1609.00708].
- [44] M. Mariño and S. Zakany, *Matrix models from operators and topological strings*, *Ann. Henri Poincaré* **17** (2016) 1075–1108, [1502.02958].
- [45] R. M. Kashaev, M. Mariño and S. Zakany, *Matrix models from operators and topological strings*, *2, Ann. Henri Poincaré* **17** (2016) 2741–2781, [1505.02243].
- [46] S. Codesido, A. Grassi and M. Mariño, *Exact results in $\mathcal{N} = 8$ Chern-Simons-matter theories and quantum geometry*, *JHEP* **07** (2015) 011, [1409.1799].

- [47] Y. Hatsuda, M. Mariño, S. Moriyama and K. Okuyama, *Non-perturbative effects and the refined topological string*, *JHEP* **09** (2014) 168, [1306.1734].
- [48] R. Gopakumar and C. Vafa, *M-theory and topological strings. 2.*, *arXiv e-prints* (12, 1998) , [hep-th/9812127].
- [49] R. Pandharipande and R. P. Thomas, *Curve counting via stable pairs in the derived category*, *Invent. Math.* **178** (2009) 407–447, [0707.2348].
- [50] N. A. Nekrasov and S. L. Shatashvili, *Quantization of Integrable Systems and Four Dimensional Gauge Theories*, in *16th ICMP*, pp. 265–289, 8, 2009. 0908.4052. DOI.
- [51] J. Choi, S. Katz and A. Klemm, *The refined BPS index from stable pair invariants*, *Commun. Math. Phys.* **328** (2014) 903–954, [1210.4403].
- [52] N. A. Nekrasov and A. Okounkov, *Membranes and Sheaves*, *arXiv e-prints* (4, 2014) , [1404.2323].
- [53] N. A. Nekrasov, *Seiberg-Witten prepotential from instanton counting*, *Adv. Theor. Math. Phys.* **7** (2003) 831–864, [hep-th/0206161].
- [54] A.-K. Kashani-Poor, *Quantization condition from exact WKB for difference equations*, *JHEP* **06** (2016) 180, [1604.01690].
- [55] R. M. Kashaev and S. M. Sergeev, *Spectral equations for the modular oscillator*, *Rev. Math. Phys.* **30** (2018) 1840009, [1703.06016].
- [56] R. M. Kashaev and S. Sergeev, *On the Spectrum of the Local \mathbb{P}^2 Mirror Curve*, *Ann. Henri Poincaré* **21** (2020) 3479–3497, [1904.12315].
- [57] A. Grassi and M. Mariño, *The complex side of the TS/ST correspondence*, *J. Phys. A* **52** (2019) 055402, [1708.08642].
- [58] M. Mariño and P. Putrov, *ABJM theory as a Fermi gas*, *J. Stat. Mech.* **1203** (2012) P03001, [1110.4066].
- [59] V. A. Kazakov, I. K. Kostov and N. A. Nekrasov, *D particles, matrix integrals and KP hierarchy*, *Nucl. Phys. B* **557** (1999) 413–442, [hep-th/9810035].
- [60] A. Grassi and M. Mariño, *M-theoretic matrix models*, *JHEP* **02** (2015) 115, [1403.4276].
- [61] P. Bousseau, P. Descombes, B. Le Floch and B. Pioline, *BPS Dendroscopy on Local \mathbb{P}^2* , *arXiv e-prints* (10, 2022) , [2210.10712].
- [62] V. Fantini and C. Rella, “On the physics of the strong-weak resurgence symmetry.” *In preparation*.
- [63] V. Fantini and C. Rella, “Geometric interpretation of the Stokes invariants for local \mathbb{P}^2 .” *In preparation*.
- [64] S. Garoufalidis, J. Gu and M. Mariño, *The Resurgent Structure of Quantum Knot Invariants*, *Commun. Math. Phys.* **386** (2021) 469–493, [2007.10190].

- [65] S. Garoufalidis, J. Gu and M. Mariño, *Peacock patterns and resurgence in complex Chern-Simons theory*, *arXiv e-prints* (11, 2020) , [2012.00062].
- [66] C. Marinissen, A. van Spaendonck and M. Vonk, *Resurgence of large order relations*, *arXiv e-prints* (12, 2023) , [2312.12497].
- [67] T. M. Chiang, A. Klemm, S.-T. Yau and E. Zaslow, *Local mirror symmetry: Calculations and interpretations*, *Adv. Theor. Math. Phys.* **3** (1999) 495–565, [hep-th/9903053].
- [68] M. Mariño and C. Rella, *On the structure of wave functions in complex Chern-Simons theory*, *arXiv preprint* (12, 2023) , [2312.00624].
- [69] T. Dimofte, S. Gukov, J. Lenells and D. Zagier, *Exact Results for Perturbative Chern-Simons Theory with Complex Gauge Group*, *Commun. Num. Theor. Phys.* **3** (2009) 363–443, [0903.2472].
- [70] J. Ellegaard Andersen and R. M. Kashaev, *A TQFT from Quantum Teichmüller Theory*, *Commun. Math. Phys.* **330** (2014) 887–934, [1109.6295].
- [71] C. Beem, T. Dimofte and S. Pasquetti, *Holomorphic Blocks in Three Dimensions*, *JHEP* **12** (2014) 177, [1211.1986].
- [72] T. Dimofte, *Complex Chern–Simons Theory at Level k via the 3d–3d Correspondence*, *Commun. Math. Phys.* **339** (2015) 619–662, [1409.0857].
- [73] N. A. Nekrasov, H. Ooguri and C. Vafa, *S duality and topological strings*, *JHEP* **10** (2004) 009, [hep-th/0403167].
- [74] V. Fantini, “Mini school on Wall-Crossing Structures, Analyticity, and Resurgence: q -Series, Resurgence, and Modularity.”
https://www.youtube.com/watch?v=x_B5C_QNiQc.
- [75] J. Hadamard, *Théorème sur les séries entières*, *Acta Math.* **22** (1900) 55 – 63.
- [76] C. Wheeler, *Modular q -difference equations and quantum invariants of hyperbolic three-manifolds*. PhD thesis, Rheinischen Friedrich–Wilhelms–Universität Bonn, 2023.
- [77] C. Wheeler, “Quantum modular forms and 3-manifolds.” *In preparation*.
- [78] O. Costin and S. Garoufalidis, *Resurgence of the Kontsevich–Zagier series*, *Annales de l’Institut Fourier* **61** (2011) 1225–1258, [math/0609619].
- [79] J. Ellegaard Andersen and W. Elbæk Mistegård, *Resurgence Analysis of Quantum Invariants of Seifert Fibered Homology Spheres*, *J. Lond. Math. Soc.* **105** (2022) 709–764, [1811.05376].
- [80] D. Zagier, *Quantum modular forms*, *Quanta of Maths* **11** (2010) 659–675.
- [81] K. Bringmann, J. Lovejoy and L. Rolin, *On Some Special Families of q -hypergeometric Maass Forms*, *Int. Math. Res. Not.* **2018** (2018) 5537–5561, [1603.01783].
- [82] A. Goswami and R. Osburn, *Quantum modularity of partial theta series with periodic coefficients*, *Forum Math.* **33** (2021) 451–463, [2012.02457].

- [83] D. Zagier, *Vassiliev invariants and a strange identity related to the Dedekind eta-function*, *Topology* **40** (2001) 945–960.
- [84] D. Zagier, “HIM Lectures. Trimester Programm on Dynamics: Topology and Numbers.” https://www.youtube.com/watch?v=2Rj_xh3UKrU.
- [85] R. Lawrence and D. Zagier, *Modular forms and quantum invariants of 3-manifolds*, *Asian J. Math.* **3** (1999) 93–107.
- [86] T. Dimofte and S. Garoufalidis, *Quantum modularity and complex Chern-Simons theory*, *Commun. Num. Theor. Phys.* **12** (2018) 1–52, [1511.05628].
- [87] M. Kontsevich and Y. Soibelman, *Analyticity and resurgence in wall-crossing formulas*, *Lett. Math. Phys.* **112** (2022) Paper No. 32, 56, [2005.10651].
- [88] K. Iwaki and O. Kidwai, *Topological Recursion and Uncoupled BPS Structures I: BPS Spectrum and Free Energies*, *Adv. Math.* **398** (2022) 108191, [2010.05596].
- [89] K. Iwaki and O. Kidwai, *Topological Recursion and Uncoupled BPS Structures II: Voros Symbols and the τ -Function*, *Commun. Math. Phys.* **399** (2023) 519–572, [2108.06995].
- [90] A. Grassi, J. Gu and M. Mariño, *Non-perturbative approaches to the quantum Seiberg-Witten curve*, *JHEP* **07** (2020) 106, [1908.07065].
- [91] M. R. Douglas, B. Fiol and C. Romelsberger, *The Spectrum of BPS branes on a noncompact Calabi-Yau*, *JHEP* **09** (2005) 057, [hep-th/0003263].
- [92] D. Gaiotto, G. W. Moore and A. Neitzke, *Wall-crossing, Hitchin systems, and the WKB approximation*, *Adv. Math.* **234** (2013) 239–403, [0907.3987].
- [93] R. Eager, S. A. Selmani and J. Walcher, *Exponential Networks and Representations of Quivers*, *JHEP* **08** (2017) 063, [1611.06177].
- [94] P. Longhi, *Instanton Particles and Monopole Strings in 5D $SU(2)$ Supersymmetric Yang-Mills Theory*, *Phys. Rev. Lett.* **126** (2021) 211601, [2101.01681].
- [95] F. Del Monte and P. Longhi, *The threefold way to quantum periods: WKB, TBA equations and q -Painlevé*, *SciPost Phys.* **15** (2023) 112, [2207.07135].
- [96] L. D. Faddeev and R. M. Kashaev, *Quantum Dilogarithm*, *Mod. Phys. Lett. A* **9** (1994) 427–434, [hep-th/9310070].
- [97] S. Garoufalidis and R. M. Kashaev, *Evaluation of state integrals at rational points*, *Commun. Num. Theor. Phys.* **9** (2015) 549–582, [1411.6062].
- [98] M. Katsurada, *Asymptotic expansions of certain q -series and a formula of Ramanujan for specific values of the riemann zeta function*, *Acta Arith.* **107** (2003) 269–298.
- [99] S. Garoufalidis and R. M. Kashaev, *Resurgence of Faddeev’s quantum dilogarithm*, *arXiv e-prints* (8, 2020) , [2008.12465].